

Exploring Ozone-climate Interactions in Idealized CMIP6 DECK Experiments

Jingyu Wang^{1,2}, Gabriel Chiodo^{1,3}, Blanca Ayarzagüena⁴, William T. Ball[†], Mohamadou Diallo⁵, Birgit Hassler⁶, James Keeble⁷, Peer Nowack⁸, Clara Orbe⁹, Sandro Vattioni¹, and Timofei Sukhodolov¹⁰

¹Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

²Lunar & Planetary Laboratory/Department of Planetary Sciences, University of Arizona, Tucson AZ, USA

³Instituto de Geociencias (IGEO), CSIC-UCM, Madrid, Spain

⁴Facultad de CC Fisicas, Universidad Complutense de Madrid, Madrid, Spain

⁵Forschungszentrum Juelich, Germany

⁶Deutsches Zentrum für Luft- und Raumfahrt, Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

⁷Lancaster Environment Center, Lancaster University, Lancaster, UK

⁸Institute of Theoretical Informatics & Institute of Meteorology and Climate Research (IMK-ASF), Karlsruhe Institute of Technology, Germany

⁹NASA Goddard Institute for Space Studies, New York, USA

¹⁰Physikalisch-Meteorologisches Observatorium Davos and World Radiation Center, Davos, Switzerland

[†]deceased

Correspondence: Jingyu Wang (wangjingyu@arizona.edu), Gabriel Chiodo (gabriel.chiodo@csic.es)

Abstract

Under climate change driven by increased carbon dioxide (CO₂) concentrations, stratospheric ozone will respond to temperature and circulation changes, leading to chemistry-climate feedback by modulating large-scale atmospheric circulation and Earth's energy budget. However, there is a significant model uncertainty since many processes are involved and few models have a detailed chemistry scheme. This work employs the latest data from Coupled Model Intercomparison Project Phase 6 (CMIP6) to investigate the ozone response to increased CO₂. We find that in most models, ozone increases in the upper stratosphere (US) and extratropical lower stratosphere (LS), and decreases in the tropical LS, thus the total column ozone (TCO) response is small in the tropics. The ozone response is mainly driven by the slower chemical destruction cycles in the US and enhanced upwelling in the LS, with a highly model-dependent Arctic ozone response to polar vortex strength changes. We then explore the ozone-climate feedback, by combining offline calculations and comparisons between models with ("chem") and without ("no-chem") interactive chemistry. We find that the stratospheric temperature response is substantial, with a global negative radiative forcing ranging from -0.03 W m⁻² to -0.19 W m⁻². We find that chem models consistently simulate less tropospheric warming and strong weakening of the polar stratospheric vortex, which results in a larger increase of sudden stratospheric warming (SSW) frequency than in most no-chem models. Our findings show that ozone-climate feedback is essential for the climate system and should be considered in the development of Earth System Models.

1 Introduction

Stratospheric ozone abundances are sensitive to temperature and circulation variations and, thus, respond to climate change (Shepherd, 2008). In turn, ozone variations can also affect temperature and the large-scale atmospheric circulation, as ozone is a radiatively active gas that absorbs solar and terrestrial radiation, and radiatively heats the stratosphere (Brasseur and Solomon, 2005). Therefore, under externally-forced climate change, stratospheric ozone will respond to and in turn feedback onto climate. Understanding the mechanisms driving the ozone response and its implications for climate, and in particular the uncertainty across models in this feedback, is critical for future climate projections.

The distribution of ozone is primarily determined by production and loss from chemical reactions and transport processes (Brasseur and Solomon, 2005). Chemistry-climate models are often employed to simulate the ozone distribution and analyze the response of ozone and its feedback on climate. From a chemical perspective, two major factors influence the changes in production and destruction of ozone due to changes in atmospheric abundances of greenhouse gases (GHGs). Firstly, an increase in the abundance of GHGs, most importantly CO₂, leads to cooling of the stratosphere. In particular in the upper stratosphere (US) where the role of transport is comparatively less important due to the short chemical lifetime of ozone, this cooling slows down the destruction of ozone due to the positive temperature dependence of the catalytic cycles (Barnett et al., 1975; Haigh and Pyle, 1982) and also reduces odd oxygen loss. Therefore, local (radiative) cooling leads to a net increase in ozone mixing ratios. Secondly, since the loss rate of odd oxygen is proportional to the abundance of atomic oxygen, the increased efficiency of the three-body reaction $O_2 + O + M \rightarrow O_3 + M$ reduces the atomic oxygen abundance and consequently reduces odd oxygen loss (Jucks and Salawitch, 2000; Jonsson et al., 2004). Conversely, in the lower stratosphere (LS), the larger ozone column abundance overhead acts as a shield, reducing the sunlight responsible for ozone production. As a result of this self-shielding effect, the abundance of ozone decreases in this region (Haigh and Pyle, 1982; Jonsson et al., 2004; Meul et al., 2014; Keeble et al., 2017).

From a transport perspective, the expansion of the tropopause due to warming at the surface will effectively replace ozone-rich stratospheric air with ozone-poor tropospheric air, leading to a decrease of ozone near the tropopause. The warming in the tropical troposphere also accelerates the subtropical jets and results in a faster Brewer-Dobson Circulation (BDC), which means faster tropical upwelling and poleward transport, resulting in more efficient transport of ozone from the tropics to the extratropics in the LS (Butchart, 2014; Abalos et al., 2021). This results in a change in the latitudinal distribution of ozone. Such transport effects on ozone are more pronounced in the lower/middle stratosphere, while chemical effects dominate in its upper part (Oman et al., 2010; Zubov et al., 2012). Because of the different sign of the ozone change in the US and LS, the total column ozone (TCO) response in the tropics depends on the opposing impacts from different dominant mechanisms. When modeled, the representation and relative importance of these processes can be model dependent, resulting in different sign of the low-latitude TCO response (Oman et al., 2010; Chiodo et al., 2018; Keeble et al., 2021). The stratospheric polar vortex is also an important feature that shapes the distribution of ozone, since it creates a transport barrier between mid-latitudes and the

poles of the winter hemisphere (Shepherd, 2008; Seinfeld et al., 1998). The response of the polar vortex to increased CO₂ level
50 is very uncertain, with a large divergence in the projection of polar vortex strength in winter across climate model ensembles,
which will affect the mixing of ozone-poor polar air with ozone-rich air in lower latitudes (Ayarzagüena et al., 2020; Karpechko
et al., 2022).

Many factors can influence the ozone response to climate change, making it difficult to separate the influence of individual
forcing agents. First, because of their role as sources of radical species in the stratosphere, the different concentrations of CH₄
55 and N₂O used in inter-model comparisons may potentially offset the effects of CO₂ (Revell et al., 2012). Second, comparing
different future scenarios may be non-trivial due to the non-linearity from the combined effects of ozone-depleting substances
(ODSs), GHGs, and ozone precursors (Meul et al., 2015). CO₂ is the only forcing considered for diagnosing transient and
equilibrium climate model sensitivity (TCR and ECS), and, thus, a large amount of data is available for this individual forcing
in the past multi-model comparison.

60 Previous studies comparing simulations of a small set of models found that under increasing CO₂, ozone mixing ratios
will increase in the US, decrease in the tropical LS and increase in the extratropical LS (Oman et al., 2010; Chiodo et al.,
2018). Aside from documenting changes in ozone, previous research has also highlighted another crucial element in the ozone-
climate interaction: changes in ozone can potentially lead to a chemistry-climate feedback, with implications for the modeled
changes in the tropospheric and surface climate. It has been suggested that the ozone-climate feedback may affect both climate
65 sensitivity (Dietmüller et al., 2014; Muthers et al., 2014, 2015; Nowack et al., 2018; Hardiman et al., 2019) and dynamical
sensitivity (Chiodo and Polvani, 2017; Nowack et al., 2017, 2018; Chiodo and Polvani, 2019; Orbe et al., 2024). In addition,
it may play a role in modulating the modeled response of El Niño–Southern Oscillation (ENSO) to global warming (Nowack
et al., 2017) and in dampening the climate system response to solar forcing (Chiodo and Polvani, 2016; Muthers et al., 2016).
Furthermore, the importance of chemistry feedback has been shown for the stratosphere-troposphere coupling (Haase and
70 Matthes, 2019), in particular for the Arctic climate (Friedel et al., 2022) and the Southern annular mode (Morgenstern, 2021).
Consequently, implementing interactive chemistry, i.e. an online scheme (either a full chemical submodel or a linearized ozone
scheme) that allows to capture the feedbacks between chemistry and temperature/dynamics, in Earth System Models (ESM) is
thought to be important for climate projections.

However, there remains a large inter-model discrepancy in both the ozone response and its climate feedback, and reasons
75 for this uncertainty are still unclear. For instance, the magnitude and peak location of stratospheric ozone response have a
notable inter-model discrepancy, which can also lead to a spread in stratospheric cooling (Chiodo et al., 2018). Simulated
tropical TCO shows a significant inter-model spread in its magnitude and sign of the response, which stems mainly from
the tropical lower stratospheric ozone (LSO3). The one possible source of the inter-model discrepancy in the tropics is the
spread of the strengthening of the ascending branch of the BDC (Chiodo et al., 2018). The magnitude of the ozone-climate
80 feedback, quantified as the impact of interactive ozone chemistry on the global-mean surface air temperature response to
abrupt quadrupling of CO₂, ranges from 20% (Nowack et al., 2015), to 7%-8% (Dietmüller et al., 2014; Muthers et al., 2016),
to nil (Marsh et al., 2016; Chiodo and Polvani, 2019). However, these previous studies have relied either on individual model
simulations or a maximum of three models.

Considering the limited number of models analyzed in the past, it is necessary to compare the ozone response and associated
85 climate feedback across more models and additional idealized scenarios (including transient experiments). This is now possible
given that the number of models employing interactive chemistry schemes has substantially increased (by factor of three) in the
last Coupled Model Intercomparison Project Phase 6 (CMIP6) (Keeble et al., 2021) compared to the previous phase 5 (Chiodo
et al., 2018).

This work analyzes the latest CMIP6 dataset, which has more and updated models and a larger range of ECS compared to
90 CMIP5 (Flynn and Mauritsen, 2020). Specifically, three 150 year-long Diagnostic, Evaluation and Characterization of Klima
(DECK) experiments which are piControl, abrupt-4 $\times$ CO₂ and 1pctCO₂ (increase up to 4 $\times$ CO₂), are used to analyze the
response and potential drivers of ozone response to elevated CO₂ (Eyring et al., 2016). The linearity of ozone response to
CO₂ increase can be investigated by comparing results from the two increased CO₂ scenarios. We also use the data from the
same three 150 year-long time-slice experiments from the new version of our in-house model, SOLar Climate Ozone Links
95 v4.0 (SOCOLv4), under piControl (1850) CMIP6 boundary conditions. The climate impact is then investigated by conducting
offline calculations of radiative transfer using Parallel Offline Radiative Transfer (PORT) for one year and prescribing the ozone
mixing ratio the same as that from CMIP6 abrupt-4 $\times$ CO₂ experiments to get the radiative perturbation. Lastly, by grouping
models based on whether interactive chemistry is employed, comparison between chem and no-chem models is conducted
through the investigation of the response in temperature and circulation between these two categories.

100 The structure of this paper is as follows: Section 2 introduces the data and models used in this work, Section 3 presents the
main findings, and Section 4 concludes and discusses the findings and broader implications.

2 Data and Models

2.1 Data

Data analyzed in this research are from the DECK experiments of CMIP6, including piControl, abrupt-4×CO₂ and 1pctCO₂ experiments. There are in total 20 models which have ozone data for piControl, 20 models for abrupt-4×CO₂ and 19 models for 1pctCO₂. Among the 20 models, there are 13 chem models that performed abrupt-4×CO₂ and 12 chem models which performed 1pctCO₂ (see Table 1). We only use 7 no-chem models that have chem counterpart (see below). Data within the time-period of 150 years from the start of each experiment are extracted. The number of ensemble members in each experiment of each model varies, thus we only use one ensemble member with the same physics version (“p”) for the analysis within each model. Note that zonal wind and vertical velocity data from MRI-ESM2-0 are not included because of the large departure from the data of other models with no reasonable trend. Furthermore, data of the first 15 years of ozone mixing ratio from the GISS-E2-1-G piControl experiment are not used due to an existing trend in this period, indicating that the model hadn’t finished the spin-up phase. Brief descriptions of CMIP6 models are proved in Keeble et al. (2021) with exceptions of SOCOL and GISS-E2-1-G, which we provide below.

SOCOLv4 (Sukhodolov et al., 2021) is based on the combination of the MPIMET (Hamburg, Germany) Earth System model (MPI-ESM1-2LR, Mauritsen et al. (2019)) consisting of ECHAM6 for atmosphere and MPIOM (Jungclaus et al., 2013) for the ocean as well as JSBACH for terrestrial biosphere and HAMOCC for the ocean’s biogeochemistry with the latest versions of the chemical (MEZON) (Egorova et al., 2003) and microphysical (AER) (Sheng et al., 2015) modules. SOCOLv4 uses the low-resolution (LR) configuration of the MPI-ESM model, which corresponds to a spectral truncation at T63, providing an approximate horizontal grid spacing of $1.9^\circ \times 1.9^\circ$. The vertical resolution of the atmosphere is set to 47 levels from the surface to 0.01 hPa, using a hybrid sigma-pressure coordinate system. Besides SOCOLv4, we also use the results from its earlier CMIP5 version SOCOL-MPIOM (Muthers et al., 2014) to compare the ozone response within the SOCOL model family. Boundary conditions for SOCOLv4 simulations follow the recommendations of Eyring et al. (2016), with the exception being the volcanic forcing, which is set to the quiet conditions of year 2000 instead of the 1850-2014 mean. Initial conditions for the coupled atmosphere-ocean system have been taken from the piControl simulation of MPI-ESM1-2-LR.

GISS-E2-1 (Kelley et al., 2020) consists of an atmosphere component and an ocean component. The horizontal and vertical resolution of the atmospheric component is 2° latitude by 2.5° longitude with 40 vertical layers from the surface to 0.1 hPa in the lower mesosphere. Among the two versions used in this work, atmospheric composition is prescribed in the first version, denoted in the CMIP6 archive as physics-version=1 (“p1”). In the other version (“p3”), ozone is calculated prognostically using the One-Moment Aerosol (OMA) model. Each of the two physics versions of the GISS-E2-1 atmospheric component is coupled to the ocean general circulation models GISS Ocean version 1 (GO1). It has a horizontal resolution of 1° latitude by 1.25° longitude and 40 vertical layers. We also examine results from the high vertical resolution version of the GISS CMIP6 climate model submission, GISS-E2-2 (Rind et al., 2020; Orbe et al., 2020). Though identical in horizontal resolution to E2-1, E2-2 has more than twice the number of vertical levels (102) and a higher model top (0.002 hPa). This, in combination a non-

Table 1. Summary of the data from CMIP6 and SOCOL experiments used in this research (Keeble et al., 2021). For models without interactive chemistry, ozone fields in most models are prescribed using CMIP6 dataset, with the exception of CESM2 and CESM-FV2, which use simulations performed with the CESM2-WACCM model; for GISS-E2-1-G (p1) and GISS-E2-2-G (p1), it is prescribed with the offline ozone fields from GISS-E2-1-G (p3) and GISS-E2-2-G (p3), respectively (Kelley et al., 2020); for HadGEM3-GC31-LL in increased CO₂ experiments, the ozone field is vertically redistributed from the piControl field to account for the tropopause shift (Hardiman et al., 2019).

Model	Horizontal Resolution (lon×lat)	Vertical Resolution	Interactive Chemistry (Y/N)	Ozone Scheme	piControl/ abrupt-4×CO ₂ / 1pctCO ₂ (year)
CESM2	288×192	32 levels; top level 2.25 hPa	N	Prescribed	150/150/150
CESM2-FV2	144×96	32 levels; top level 2.25 hPa	N	Prescribed	150/150/150
CESM2-WACCM	288×192	70 levels; top level 4.5×10^{-6} hPa	Y	Interactive chemistry	150/150/150
CESM2-WACCM-FV2	144×96	70 levels; top level 4.5×10^{-6} hPa	Y	Interactive chemistry	150/150/150
CNRM-CM6-1	256×128	91 levels; top level 78.4 km	Y	Simplified online scheme	150/150/150
CNRM-CM6-1-HR	720×360	91 levels; top level 78.4 km	Y	Simplified online scheme	150/150/150
CNRM-ESM2-1	256×128	91 levels; top level 78.4 km	Y	Interactive chemistry	150/150/150
E3SM-1-0	360×180	72 levels; top level 0.1 hPa	Y	Simplified online scheme	150/150/150
GFDL-CM4	360×180	33 levels; top level 1 hPa	N	Prescribed (CMIP6 dataset)	150/150/150
GFDL-ESM4	288×180	49 levels; top level 1 Pa	Y	Interactive chemistry	150/150/150
GISS-E2-1-G (p1)	144×90	40 levels; top level 0.1 hPa	N	Prescribed	150/150/150
GISS-E2-1-G (p3)	144×90	40 levels; top level 0.1 hPa	Y	Interactive chemistry	150/150/150
GISS-E2-2-G (p1)	144×90	102 levels; top level 0.002 hPa	N	Prescribed	150/150/150
GISS-E2-2-G (p3)	144×90	102 levels; top level 0.002 hPa	Y	Interactive chemistry	150/150/150
HadGEM3-GC31-LL	192×144	85 levels; top level 85km	N	Prescribed	150/150/150
MPI-ESM1-2-LR	192×96	47 levels; top level 0.01 hPa	N	Prescribed (CMIP6 dataset)	150/150/150
MRI-ESM2-0	128×64	80 levels; top level 0.01 hPa	Y	Interactive chemistry	150/150/150
SOCOL-MPIOM	96×48	47 levels; top level 0.01 hPa	Y	Interactive chemistry	150/150/–
SOCOLv4	192×96	47 levels; top level 0.01 hPa	Y	Interactive chemistry	150/150/150
UKESM1-0-LL	192×144	85 levels; top level 85km	Y	Interactive chemistry	150/150/150

135 orographic gravity wave drag scheme that is directly tethered to parameterized convection, produces in E2-2 more credible middle atmosphere dynamical and transport circulations, compared to observations (Orbe et al., 2020).

Table 2. Chem vs. No-chem pairs chosen to compare chem and no-chem models

Chem	No-chem
CESM2-WACCM	CESM2
CESM2-WACCM-FV2	CESM2-FV2
GFDL-ESM4	GFDL-CM4
UKESM1-0-LL	HadGEM3-GC31-LL
SOCOLv4	MPI-ESM1-2-LR
GISS-E2-1-G (p3)	GISS-E2-1-G (p1)
GISS-E2-2-G (p3)	GISS-E2-2-G (p1)

2.2 CESM-PORT

Parallel Offline Radiative Transfer in the Community Earth System Model (CESM-PORT) (Conley et al., 2013) is driven by model-generated datasets that can be used for any radiation calculation that the underlying radiative transfer schemes can perform, such as diagnosing radiative forcing. The inclusion of stratospheric temperature adjustment under the assumption of fixed dynamical heating is necessary to compute radiative forcing in addition to the more straightforward instantaneous radiative forcing.

The CESM-PORT experiment includes two parts: verification and perturbation. We validate CESM-PORT using the ozone data from the Whole Atmosphere Community Climate Model (WACCM) (Marsh et al., 2013) under piControl configurations with interactive chemistry. For the perturbation process, another series of CESM-PORT experiments are carried out using ozone mixing ratio from abrupt-4×CO₂ experiment of CMIP6 models with interactive chemistry. This is done by first computing the ozone fraction ratio of abrupt-4×CO₂ to piControl, and then multiplying the background ozone mixing ratio of the CESM-PORT piControl experiment above its tropopause by the corresponding fraction ratio obtained for each model, which has been interpolated onto the CESM grid.

The radiative forcing of the surface-troposphere system due to the perturbation or the introduction of an agent is defined as “the change in net (down minus up) irradiance (solar plus long-wave; in Wm^{−2}) at the tropopause after allowing for stratospheric temperatures to re-adjust to radiative equilibrium, but with surface and tropospheric temperatures and state held fixed at the unperturbed values” (Ramaswamy et al., 2001). It can be investigated by looking at the difference of radiative fluxes and temperature adjustment between the CESM-PORT reference and perturbation runs.

2.3 Pairs of chem and no-chem models

To investigate the impact of interactive chemistry, we compare models that simulate interactive ozone with those that impose a fixed pre-industrial climatological ozone forcing data-set (i.e. the "no-chem" models) from Checa-Garcia et al. (2018), following the method discussed in Morgenstern et al. (2022). We identified seven such "pairs" (Table 2)

3 Results

160 In this section, we examine the response of ozone mixing ratio and column ozone abundance from all the chem models except for GISS-E2-2-G since it has similar behavior with GISS-E2-1-G. We then investigate potential drivers of these responses. The responses are assessed by taking the difference between the climatology obtained from the last 100 years for abrupt- $4\times\text{CO}_2$ ($4\times\text{CO}_2$ hereafter) or years 135 to 145 for 1pct CO_2 experiment (when it reaches the same CO_2 level as abrupt- $4\times\text{CO}_2$) and the climatology of the 150-year-long piControl experiments.

165 3.1 Annual-mean zonal-mean ozone response




Figure 1. Annual-mean zonal-mean ozone response to $4\times\text{CO}_2$ of each chem model. Tropopause for piControl and $4\times\text{CO}_2$ is denoted using black and red dotted line, respectively. Regions that are not stippled are statistically significant (at 99% level), according to the t-test.

Figure 1 shows the annual-mean zonal-mean ozone response to $4\times\text{CO}_2$. We assume the timeseries of ozone concentration under piControl and $4\times\text{CO}_2$ are independent samples with the same variance, then we compute the t statistic to see if the two samples have same mean value. This also applies to other variables we analyze hereafter. Ozone increases in the tropical middle troposphere in most models, and decreases in the upper troposphere (although not the case for CNRM-ESM2-1, GFDL-

170 ESM4 and UKESM1-0-LL). The increase is likely due to enhanced NO_x emission from lightning, which can increase ozone abundance by the cycling of NO_x and HO_x (Banerjee et al., 2014; Revell et al., 2015; Iglesias-Suarez et al., 2018), and increased stratosphere-troposphere exchange through isotropic mixing (Abalos et al., 2020; Hegglin and Shepherd, 2009; Wang and Fu, 2023). A similar pattern was simulated in some of the CCMI1 models (Morgenstern et al., 2018), even though not all those models fully represent NO_x production changes under climate change. The decrease near the tropical tropopause is linked to
175 the expansion of the tropopause due to tropospheric warming, which replaces ozone-rich stratospheric air with tropospheric air that has a lower ozone abundance. Note that E3SM-1-0 does not employ interactive chemistry in the troposphere, which explains the lack of response in this region for this model. When the CO_2 forcing is transient rather than abrupt (1pct CO_2), we find a largely similar response (see Fig. B1).

In the stratosphere, the ozone mixing ratio increases in the US due to the reduction in chemical destruction caused by cooling.
180 There is a decrease in tropical LS ozone of about 40%, which is caused mostly by dynamical processes. But partly, it also results from the increased US ozone that absorbs more ozone-producing UV radiation (200-240 nm), thus less ozone will form in the LS (self-shielding, Dütsch et al., 1991). The relative effects of both processes could be partially isolated by using the $4\times\text{CO}_2$ experiment with prescribed sea surface temperatures (SSTs) from the piControl experiment, as has been done in Match and Gerber (2022) and Chrysanthou et al. (2020), suggesting that the self-shielding effect would be responsible for up to one third
185 of the total, though prescribing the SSTs doesn't fully prevent tropical upwelling from changing. The strengthened transport by the BDC of the tropical ozone to the extratropics and the expansion of the tropopause will also lead to the replacement of ozone-rich air with ozone-poor air. In the extratropics, the enhanced ozone transport from the tropics and upper levels is more significant than the change of the efficiency of ozone formation by photolysis, resulting in a net increase in the LS.

Comparing the ozone response of these models with CMIP5 (Chiodo et al., 2018), we see that the pattern in CMIP6 models
190 generally agrees well, indicating consistency between the two generations of models. However, having more models for our analysis allows us to test whether the mechanisms of ozone response based on CMIP6 simulations are more robust.




Figure 2. 150-year-long annual-mean ozone response to temperature change in stratosphere at different pressure levels and latitude bands based on the 1pctCO₂ experiment.

Next, we explore the relationship between ozone and local temperature, to identify how chemistry and transport affect ozone in different stratospheric regions. We look at this relationship by plotting the annual mean temperature change against the ozone response at different pressure levels and latitude bands (Fig. 2). Here, we use the data from 1pctCO₂ experiment instead of the 4×CO₂ experiment, since the linearly increasing forcing allows tracking the regional dependencies in individual models.

In the US, there is an inverse relationship between the ozone and temperature response for most models: this is expected from the slow-down of ozone destruction with radiative cooling from increased CO₂. On the other hand, more ozone enhances UV radiation absorption, heating up the US, and reducing the cooling from increased CO₂. The competition between the two processes determines the cooling trend in the US, and differences in the photochemistry and radiation schemes of the models explain the different sensitivities of ozone to temperature change across models, defined by the slope in the scatter plots. For instance, the ozone increase at 1 hPa in SOCOLv4 is the largest by the end of the 1pctCO₂ experiment, but it does not correspond to the largest cooling, which might be caused by more efficient heating by ozone. For UKESM1-0-LL, we find the

opposite behavior, indicating a lower temperature-dependence of gas-phase ozone chemistry in this model. The general trend is consistent among all latitude bands at 1 hPa and 5 hPa.

205 The correlation is the opposite in the tropical LS (30 hPa and 70 hPa) since dynamics and related changes in ozone transport play a dominant role. Here, tropical LS cooling is the result of strengthened upwelling caused by the acceleration of BDC (Abalos et al., 2021). The stronger upwelling results in enhanced transport of ozone out of the tropical pipe and ozone-poor air from the troposphere to the LS, leading (locally) to a decrease (Oman et al., 2010). Hence, the colder the upper troposphere and lower stratosphere (UTLS) gets over the course of the 1pctCO₂ experiment, the more ozone is transported out of the tropics, locally
210 reducing ozone abundances (i.e. a positive relationship). Again, this relationship is model-dependent, and in some models it becomes less linear (e.g. UKESM1-0-LL). In the extratropical LS, the relation is less evident since dynamics and chemistry are equally important.

Overall, the response of ozone to elevated CO₂ is consistent with previous research, and the mechanisms of the responses are examined using CMIP6 data and a larger number of models. The ozone response pattern in the stratosphere can therefore
215 be described as follows: ozone increase in the US dominated by the chemistry response to temperature and ozone heating, ozone decrease in the tropical LS caused by ozone self-shielding and transport, and increase in extratropical LS as a response of both chemistry and dynamics.

3.1.1 Polar vortex vs. ozone response




Figure 3. 150-year-long seasonal-mean ozone response in 60-90°N to zonal wind (u) change in 50-70°N at 70hPa in DJF and MAM for 1pctCO₂. Fitting lines retrieved from linear regression are plotted as dash-dotted lines with the corresponding color for each model. The thick black line is fitted using data from all models with the corresponding R² denoted in the upper right corner of the plot. R² values for each model are denoted in the legend for DJF and MAM respectively.

Figure 3 shows that in winter, for most models, a weaker NH polar vortex correlates with an increase of ozone in the Arctic. Antarctic vortex shows similar behavior (see Figure B2). It might be caused by the more efficient wave forcing from surface warming, and results in enhanced mixing of ozone-rich air masses into the polar vortex. Also, the heating from more ozone could partly contribute to polar warming and weakening of the polar vortex throughout the extended winter season, and there is more ozone in the surf zone available to be transported into the polar region. Conversely, during the deep winter months, LW cooling effect from ozone anomalies can be dominant at the polar-most latitudes (London, 1980). Distinguishing the relative importance of individual factors would require additional experiments, where ozone would respond to dynamical changes but would not in turn affect the radiation and, thus, the circulation. The relation in spring (R² = 0.01) is not as evident as in winter (R² = 0.10). The breakup of the polar vortex may lead to enhanced transport of ozone to polar region, but averaging over MAM may mask this relationship. Investigation of the breakup time of polar vortex and how it changes under climate change would need to be considered for each model, which is out of scope, but which merits further investigation. Note that in the 1pctCO₂ simulations analyzed here, ODSs are set to pre-industrial values, and heterogeneous chemistry is less

important than under present-day conditions. Hence, polar ozone abundances are mostly determined by mid-winter transport, and thus correlate better with the vortex strength in winter than in spring. Under near-present day ODSs concentrations, the springtime ozone/vortex relationship gets magnified by the combined feedback of heterogeneous chemistry, temperature and dynamics (Kult-Herdin et al., 2023).

235 3.1.2 Tropical upwelling vs. ozone response

In the models used here, surface temperature increases in most regions, with some exceptions over the North Atlantic in some models (see Fig. B3), which will lead to more efficient wave generation and propagation, thus enhancing the upwelling in the tropics, which is consistent with previous studies (Chrysanthou et al., 2020). It is thus pertinent to ask whether the climate model sensitivity correlates with tropical upwelling, and thus ozone. We explore this potential linkage by examining the relationship
240 between upwelling and tropical ozone, using grid-scale vertical velocity ($\overline{w}$, Fig. 4), and residual upwelling ($\overline{w^*}$) diagnosed via the Transformed Eulerian Mean (TEM, see Fig. B4), provided via the DynVar initiative for some of the CMIP6 models (Gerber and Manzini, 2016). Both metrics indicate strengthened tropical upwelling with warming (panels (a)-(b) in Fig. 4 and Fig. B4), consistent with previous work (Abalos et al., 2021). Again as expected, we find an inverse relationship between upwelling and ozone changes (panels (c)-(d) in Fig. 4 and Fig. B4), indicating that more tropical upwelling decreases tropical ozone.
245 Climatological responses (panels (e)-(f) in Fig. 4 and Fig. B4) are also consistent between the models so that they all show a negative lower-stratospheric tropical ozone change and a positive one in tropical upwelling under higher GHGs. However, we don't find a consistent relationship in the inter-model differences in the responses in the two variables, which seems to be dominated by "outliers" (e.g. SOCOL-MPIOM in panel (e)) and the regression slope there is insignificant. The sensitivity of ozone to upwelling may differ across models, as it also depends on the vertical ozone gradient, which differs across models
250 (e.g. in UKESM1-0-LL, not shown) and the background climatology of stratospheric transport and dynamics.




Figure 4. (a)-(b): Annual-mean tropical (15°S-15°N) upwelling (w) response to global mean surface temperature for the 4×CO₂ and 1pctCO₂ experiments for individual year. (c)-(d): Same as (a)-(b), but contrasting upwelling changes with the ozone response. (e)-(f): Same as (c)-(d), but averaged over the last 100 years for 4×CO₂ and for all the 150 years for 1pctCO₂ experiment. Fitting lines retrieved from a linear regression are plotted with the corresponding color of each model.

3.2 Column ozone response




Figure 5. Annual-mean column ozone response to $4\times\text{CO}_2$ change. (a) TCO, (b) tropospheric (TRO3), (c) lower-stratosphere (LSO3), and (d) upper-stratosphere (USO3) partial ozone columns. The lower stratosphere is defined as the atmospheric layer between the tropopause and 20 hPa, and the upper stratosphere is defined as the layer between 20 hPa and 1 hPa.

One of the key ozone metrics of interest is total column ozone (TCO), as it affects the amount of UV reaching the Earth's surface: this is shown in Figure 5. TCO increases by up to 50-75 DU in the polar regions, while it's around zero in the tropics. In the tropics, the multi-model uncertainty is smaller, though UKESM1-0-LL and SOCOL-MPIOM show a more negative TCO response and GISS-E2-1-G_p3 is an outlier in the opposite direction. In the extratropics, the spread among models is increasing and is the strongest in the polar regions, with values ranging between 5 and 80 DU in SH and 30 and 80 DU in NH. Decomposing the response of TCO into three parts, we see that for tropospheric ozone (TRO3), the response is relatively small and slightly positive in the middle latitudes. The lower-stratospheric ozone (LSO3) response dominates the uncertainty of the model spread in TCO with a negative response in the tropics and a mostly positive but highly model-dependent response in the high latitudes; and for upper-stratospheric ozone (USO3), there is an uniform increase. Therefore, the compensation between LSO3 and USO3 leads to small response in the tropics for TCO. These results are consistent with the analysis of the data from four CMIP5 models (Chiodo et al., 2018), including also the response in the NH being larger than that in the SH due to a stronger BDC (Butchart, 2014), and are similar with previous study (Morgenstern et al., 2018). When looking at the TCO seasonal cycle, there is a larger seasonal variation at high latitudes than in the tropics with the peak in both hemispheres occurring in the late boreal winter and early spring (not shown). The seasonal variability in high latitudes is consistent with that of BDC, which is stronger in the winter-spring hemisphere (Shepherd, 2008).




Figure 6. 150-year-long annual-mean tropical (20°S-20°N) LSO3, USO3 and TCO response to a 4×CO₂ change with time.

Figure 6 shows that tropical LSO3 decreases quickly in the first ~40 years, followed by a slight trend potentially due to the slow response of the deep ocean, while tropical USO3 reaches equilibrium almost instantaneously. This confirms again the different dominant mechanisms in the two layers. In the LS, the change in transport dominates, while in the US, the cooling-induced change in the efficiency of ozone destruction in the Chapman mechanism dominates. Since the response of dynamics is much slower, as it is linked to the surface temperature changes, it takes longer for LSO3 to reach equilibrium. As a result of this lag in the LSO3 response, the tropical TCO response to the 4×CO₂ increase is slightly positive for the first several decades and then gets mostly negative over the subsequent 80 years. For 1pctCO₂ (see Fig. B5), we can see the lag of the response since the tropical TCO response around year 140, when the CO₂ level in 1pctCO₂ reaches that in 4×CO₂, is smaller than the equilibrated value from 4×CO₂. This is also revealed in the slightly larger decrease of ozone in tropical LS for 4×CO₂ (see Fig. 1) compared to 1pctCO₂ (see Fig. B1).

3.3 Climate feedback

3.3.1 Temperature adjustment




Figure 7. Temperature adjustment to ozone for the $4\times\text{CO}_2$ experiment from the chem models.

So far, we have examined how ozone is affected by CO_2 -driven warming, via changes in circulation and temperature. However, ozone also largely affects stratospheric climate, via changes in radiative heating. One way to disentangle the radiative effects of ozone is by performing offline radiative transfer calculations to quantify the temperature adjustments needed to achieve radiative equilibrium, by using the Fixed Dynamical Heating (FDH) approximation with fixed tropospheric temperature (Fels et al., 1980). We achieve this in CESM-PORT for all the ozone perturbations derived from each of the models displayed in Figure 1 (see Section 2.3), and plot the corresponding temperature adjustments in Figure 7. We find that the temperature adjustments broadly correspond to the pattern of ozone responses to $4\times\text{CO}_2$ change (Fig. 1), with a cooling in the tropical LS and warming elsewhere.

By comparing the temperature adjustment to ozone with the actual zonal-mean temperature response in the coupled experiments (displayed in Fig. B6), we can see the relative contribution of the radiative heating induced by ozone to stratospheric temperature changes. In the LS, we find a slight warming in the NH middle and high-latitudes, contributing to the total tem-

290 perature response in this region, competing with a similar level of cooling from CO₂, which results in differences in the sign of the total temperature response across models. Over the tropical tropopause and LS, we find cooling, indicating that reduced ozone in the UTLS amplifies the stratospheric cooling from increased CO₂ and can explain about half of the total temperature change, whereas in the US, heating from increased ozone is outweighed by radiative cooling from increased CO₂, consistent with previous work on historical temperature trends (Chiodo and Polvani, 2022; McLandress et al., 2011). At high latitudes

295 in the LS, the temperature response depends on the opposing influences of warming induced by increased ozone abundances and downwelling, and radiative cooling from CO₂ (Chiodo et al., 2023; Kult-Herdin et al., 2023). For most of the models, the warming dominates at high latitudes, while the cooling dominates close to the tropics.

3.3.2 Radiative impacts of ozone




Figure 8. Annual-mean zonal-mean response of radiative fluxes at the tropopause to ozone from chem models for the 4×CO₂ experiment. The longwave, shortwave and net radiative flux response are denoted with red (dash-dotted), blue (dotted) and black lines, respectively. The value of the global mean net flux change is shown as number in each subplot.

Changes in ozone can have a sizable impact on the radiative balance of the stratosphere, via changes in shortwave (SW) heating,

300 but also in the longwave (LW), by changing the trapping of LW radiation coming from the troposphere, and via the stratospheric temperature adjustments. Hence, changes in ozone that are induced by CO₂ lead to a radiative forcing (RF), potentially altering tropospheric and surface climate response to CO₂ in some models (Dietmüller et al., 2014; Muthers et al., 2014; Nowack

et al., 2015, 2018). We quantify the ozone-induced RF across all CMIP6 models, by calculating the stratospherically-adjusted changes in the LW and SW at the tropopause. By inspecting these quantities in Figure 8, we see that the latitudinal structure of the LW flux changes is consistent with the ozone changes near the tropopause. The ozone changes determine the temperature adjustment and modulate the LW flux by increasing the local LW absorption and emission. The net LW flux is reduced (meaning more LW escaping the troposphere) in the tropics, and is increased in the high latitudes. These opposing effects are due to changes in the LW absorption as well as in the local temperatures induced by ozone. The reduced ozone abundances near the UTLS radiatively cool the UTLS, leading to reduced LW emission from the stratosphere towards the troposphere; conversely, increased ozone abundances in the LS at high latitudes lead to warming, increasing the LW emission from the stratosphere towards the troposphere. In the SW range, the resulting RF mirrors the changes in TCO and is due to the "shielding" effect of the ozone columns; the SW flux barely changes in the tropics (due to small changes in TCO there), while it's reduced in the extratropics (where TCO increases). At high latitudes, the thicker TCO absorbs more SW, reducing the SW flux reaching the tropopause. The net flux change is the sum of that of LW and SW, and is negative in the tropics and positive in the extratropics, with the LW generally dominating over the SW forcing. Due to the large area of the tropics, the global mean net RF is negative in all models, varying between -0.03 (E3SM-1-0) and -0.19 Wm^{-2} (UKESM-1-0-LL). Compared to CMIP5, the range in the ozone-induced RF is larger (Fig. 2, e.g. Chiodo and Polvani, 2019) in CMIP6, possibly due to the larger number of models considered in this study. Taken together, the negative RF at the tropopause implies a reduction in tropospheric and surface warming from ozone, consistent with some previous studies (Dietmüller et al., 2014; Muthers et al., 2014; Nowack et al., 2015). However, the ozone-induced RF does not tell the full story, as it does not consider other physical feedbacks or large-scale circulation changes, as well as it can be dependent on the background model climatologies, while in the PORT calculations we used a single prescribed background. Therefore, we examine the overall feedback by comparing models with and without interactive chemistry in the next section.

3.3.3 Impact of interactive ozone chemistry on the coupled response

In this section, our aim is to examine the overall climate feedback in the context of coupled experiments. The cleanest way to achieve this is by running pairs of experiments within the same model system, with and without interactive ozone, as done in previous work (Dietmüller et al., 2014; Muthers et al., 2014; Nowack et al., 2015; Chiodo and Polvani, 2016; Marsh et al., 2016; Chiodo and Polvani, 2019). As an alternative approach, we compare the seven pairs of chem and no-chem models from CMIP6 (see Table 2), and look at the differences in the $4\times\text{CO}_2$ response among them. The caveat is that the comparison of such pairs will not only isolate the impact of interactive ozone, but also other effects, such as differences in the model physics, as discussed in Morgenstern et al. (2022). More specifically, the pairs also differ in other components like gravity wave drag, and convective parameterization.




Figure 9. Comparison of the 100-year long annual-mean air temperature response to $4\times\text{CO}_2$ between seven pairs of chem and no-chem CMIP6 models. The left column shows the response from chem models, the middle column shows the response from no-chem models and the right column shows the difference between chem and no-chem models.

First, we start by looking at zonal-mean temperature in Figure 9. The seven pairs of models share the same pattern of zonal-mean temperature response. Compared with no-chem models, chem models have significantly less warming in the troposphere, more cooling in the tropical LS and less cooling in the extratropical LS and US, which is consistent with previous findings (Dietmüller et al., 2014; Nowack et al., 2015; Marsh et al., 2016). The negative stratospherically-adjusted RF at the tropopause (Fig. 8) might partly explain the reduced tropospheric warming in chem models. Another coupling mechanism could be related to the transport of water vapor from the upper troposphere to the LS being reduced due to the cooling of LS, which would lead to less tropospheric warming due to the GHG effects of the water vapor (Nowack et al., 2018; Banerjee et al., 2019; Nowack et al., 2023). In the stratosphere, the temperature pattern is coherent with the ozone response, with decreased ozone in the tropical LS leading to cooling and increased ozone in the US and extratropical LS leading to a warming (Fig. 1). Among these pairs, UKESM1-0-LL/HadGEM3-GC31-LL exhibits the largest difference in the tropical LS, while SOCOLv4/MPI-ESM1-2-LR has the largest difference in the US. In the multi-model mean (Fig. 10), chem models are about ~ 2 K cooler in the troposphere in their $4\times\text{CO}_2$ response, ~ 3 K cooler in the tropical LS and ~ 4 K warmer in the US. In the stratosphere, the temperature pattern is broadly consistent with the temperature adjustments calculated with CESM-PORT (Fig. 7), suggesting that the chem vs no-chem differences are indeed indicative of a true "ozone effect". These temperature changes have implications for the zonal mean zonal wind, as discussed in the next paragraph. In the tropical stratosphere, we explore if the inclusion of interactive ozone also affects the degree of BDC acceleration, as suggested by (Hufnagl et al., 2023). Out of the three pairs of experiments that provide tropical upwelling ($\overline{w^*}$, Table B1), for all of them we find a reduction in the increase due to CO_2 in the chem version of the models. Changes in tropical upwelling critically depend on zonal wind changes near the tropical tropopause layer (TTL), as these modulate the propagation and dissipation of tropospheric waves. Ozone decreases the meridional temperature gradient near the TTL, altering the BDC as a consequence of changes in zonal winds, as discussed next.




Figure 10. Similar to Figure 9, but average of 100-year long annual mean air temperature response to $4\times\text{CO}_2$ of seven pairs of chem and no-chem models.




Figure 11. Comparison of the 100-year long seasonal mean zonal wind response to a $4\times\text{CO}_2$ change in DJF between seven pairs of chem and no-chem models. Black contour lines depict the zonal wind climatology from the piControl experiment.

We then look at the response of polar vortex in boreal winter (Fig. 11) since this is the season when the strength of the polar vortex is at its peak. All the individual pair comparisons of zonal wind in DJF reveal a weaker polar vortex compared to no-chem models (Fig. 11) because chem models have a smaller latitudinal temperature gradient (Fig. 9). The one exception is the GISS-E2-2-G model, in which interactive composition produces a more dramatic weakening of the Atlantic Meridional Overturning Circulation (AMOC); this, on longer timescales, accelerates the NH jet, obfuscating the initial weakening of the NH polar vortex (Orbe et al., 2024). On shorter timescales, however, the initial response of the polar vortex in the interactive chemistry simulation is consistent with the response observed in the other models. In terms of the frequency of sudden stratospheric warmings (SSWs), models widely differ in terms of their background (piControl) SSW frequencies, as well as their effects of global warming, consistent with previous work (Ayarzagüena et al., 2020). Interestingly, chem models tend to show larger and more consistent (4 out of 5 pairs) increases in the SSW frequency compared to no-chem models (Table. B1), in agreement with the zonal wind weakening (Fig. 11). In some models, the non-chem configuration has different sign in the frequency change under abrupt-4xCO₂ compared to the chem counterpart (e.g. SOCOLv4/MPI-ESM1-1-2-LR and UKESM1-0-LL/HadGEM3-GC31-LL). In some pairs, the easterly wind anomalies extend to the surface (CESM2-WACCM/CESM2, GFDL-ESM4/GFDL-CM4 and UKESM1-0-LL/HadGEM3-GC31-LL). The positive zonal wind anomalies in the tropical US could be related to the expansion of the polar vortex area due to its weakening, as discussed in Section 3.1.1. Averaged over all pairs (Fig. 12), we find that zonal wind weakens in both hemispheres in chem models with respect to non-chem models. Most remarkably, these anomalies extend to the troposphere in the SH, and are indicative of an equatorward shift of the middle latitude jet, thus opposing the effect of increasing CO₂. This result is consistent with previous work using individual models, suggesting that models with prescribed ozone may over-estimate the tropospheric circulation response to CO₂ (Chiodo and Polvani, 2017; Nowack et al., 2018; Li and Newman, 2023). The crucial new addition is that here, we confirm this finding with more models, and find that it is a robust feature among all models in the stratosphere. This effect could be one of the reasons for uncertainty across some of the CMIP6 models in terms of NH polar vortex response to future CO₂ scenarios (Karpechko et al., 2022, 2024). In addition, we confirm the finding of Chiodo and Polvani (2019) concerning the role of stratospheric ozone in reducing the poleward shift of the tropospheric mid-latitude jets, as we show in Figure B7 as a difference between the chem and no-chem multi-model mean pairs in the 850 hPa zonal winds.




Figure 12. Similar to Figure 11, but a model average.

Lastly, we examine the differences in the global-mean surface air temperature response in $4\times\text{CO}_2$ and 1pctCO_2 experiments as well as the difference of the two (see Table 3). Chem models tend to exhibit less surface warming than no-chem models in both types of CO_2 experiments, consistent with the tropospheric cooling shown in Figure 9. In the multi-model mean, we find that chem models have approx. 9.0% less warming under $4\times\text{CO}_2$ than no-chem models, while for 1pctCO_2 , the difference is smaller ($\sim 7.0\%$). This leads to an overall larger response of global mean surface temperature for chem version than no-chem version. Taken together, the negative RF induced by ozone is consistent with the reduction in surface warming in chem models, which is in agreement with the initial work by Dietmüller et al. (2014) and in qualitative agreement with Nowack et al. (2015). However, we still also report a large uncertainty across models in this effect, partly also confirming the previous difference between the single-model studies (Dietmüller et al., 2014; Nowack et al., 2015; Muthers et al., 2016; Marsh et al., 2016; Chiodo and Polvani, 2019). Also, models with a large RF from ozone (e.g. UKESM-1-0-LL and SOCOLv4) are not the models showing the largest reduction in surface warming, suggesting that other factors such as differences in model physics and other climate feedbacks (e.g. clouds and water vapor) may contribute to the differences between both pairs of models. The bias arising from inconsistencies between the thermal and chemical tropopauses has been eliminated in the HadGEM3-GC31-LL $4\times\text{CO}_2$ and 1pctCO_2 experiments, as described in Hardiman et al. (2019). Hence, the UKESM-1-0-LL/HadGEM3-GC31-LL pair is not affected by this, but it may be of importance in other pairs.

Table 3. Comparison of the global-mean surface air temperature of the seven chem/no-chem CMIP6 model pairs.

Pair	4×CO ₂ (K)			1pctCO ₂ (K)		
	Chem	No-chem	Chem minus	Chem	No-chem	Chem minus
			No-chem			No-chem
CESM2-WACCM/CESM2	6.07	6.84	-0.77	5.03	5.49	-0.46
CESM2-WACCM-FV2/CESM2-FV2	5.65	6.02	-0.37	4.96	4.96	-0.0015
GFDL-ESM4/GFDL-CM4	4.12	5.48	-1.36	3.78	5.06	-1.29
GISS-E2-1-G (r1i1p3f1)/GISS-E2-1-G (r1i1p1f1)	3.63	3.81	-0.18	3.68	3.70	-0.0167
GISS-E2-2-G (r1i1p3f1)/GISS-E2-2-G (r1i1p1f1)	3.15	3.66	-0.51	-	-	-
UKESM1-0-LL/HadGEM3-GC31-LL	7.3	7.47	-0.17	6.54	6.64	-0.10
SOCOLv4/MPI-ESM1-2-LR	4.46	4.74	-0.27	4.05	4.3	-0.26
Multi-model Mean	4.91	5.43	-0.52±0.39 (7.2%)	4.67	5.03	-0.35±0.45 (8.9%)

4 Conclusions

In this study, we investigate the ozone response to elevated CO_2 levels, by analyzing data from 20 models from the CMIP6 DECK experiments. We assess the role of potential drivers of ozone changes, by exploring the relationships between ozone and parameters like near-surface and stratospheric temperature, zonal wind, and residual upwelling (w^*). We find that most stratospheric ozone changes can be explained by these drivers, but we also find large inter-model differences in the ozone response in some regions, that cannot be explained by any of them. The larger number of models enables us to have a more robust comparison than previous studies, which employed three models at most (Chiodo and Polvani, 2019).

The main findings of this paper can be concluded as follows:

1. The ozone response to $4\times\text{CO}_2$ and 1pctCO_2 is very similar, although the ozone decrease in the tropical lower stratosphere is smaller in 1pctCO_2 due to the lag of response to transient forcing.
2. The analyzed models exhibit a broadly similar pattern of zonal-mean ozone response, with an increase in the upper stratosphere and extratropical lower stratosphere and a decrease in the tropical lower stratosphere.
3. The ozone response in the upper stratosphere is dominated by changes in gas-phase chemistry, while in the lower stratosphere, chemistry and transport changes both play a role. Therefore, the timescale of the response in the lower stratosphere and upper stratosphere is different. In the upper stratosphere, it reaches equilibrium almost instantaneously because of the quick response of gas-phase chemistry. In the lower stratosphere, the response is much slower because of the slower chemical time-scales, as well as the sizable role of tropical upwelling, which is influenced by SSTs and thus slow equilibration timescale of the ocean.
4. In the lower stratosphere, there is a decrease of ozone dominated by stronger upwelling, while in the middle and high latitudes, the ozone anomalies are positive, because of the combination of increased transport by BDC and overall higher ozone abundance due to cooling from CO_2 . Additionally, the weakening of westerlies also leads to more in-mixing of ozone-rich air into the polar region during winter.
5. In the multi-model mean sense, the total column ozone response is negligible in the tropics because of the cancellation between decreases in the lower stratosphere and increases in the upper stratosphere. Total column ozone increases at high latitudes, but with a larger inter-model discrepancy compared to the tropics, which is related to the differences in the response of residual ozone transport in the models and, to some extent, uncertainty in the polar vortex response. Overall, the pattern (and its uncertainty) in the total column ozone response is dominated by lower stratospheric ozone.
6. The response of upwelling to surface warming, as well as the ozone response to strengthened upwelling, is strongly model dependent. These two sensitivities combined determine how tropical total column ozone responds to increased CO_2 concentrations in different models. Disentangling the two effects for ozone response uncertainty would require additional experiments with prescribed SSTs, to constrain the tropical upwelling, and having ozone calculated online but radiatively decoupled.

- 425 7. Models with interactive chemistry show less warming under increased CO₂ in the troposphere and tropical lower strato-
sphere, and more warming in the extratropical lower and upper stratosphere, consistent with previous studies. These
temperature changes cause the weakening of the stratospheric polar vortex during boreal winter, with this signal extend-
ing to the troposphere. Compared to the model uncertainty in the total response of polar vortex to increased CO₂, the
models are more consistent in terms of the sign of the response. Also, due to the weakening in the polar vortex, chem
430 models tend to show larger and more consistent increases in the SSW frequency compared to no-chem models.
8. Models with interactive chemistry simulate, on average, have about $\sim 9.6\% \pm 7.2\%$ smaller surface warming than models
without chemistry under $4 \times \text{CO}_2$, and about $\sim 7.0\% \pm 8.9\%$ less warming under 1pctCO_2 .

Previous studies also show that the ozone response to a $4 \times \text{CO}_2$ change has a considerable impact on the tropospheric
circulation in NH, and will induce an equatorward shift of the North Atlantic jet during boreal winter (Chiodo and Polvani,
435 2019). This shift of the North Atlantic jet may induce a rapid weakening of the AMOC, and in turn, might result in an
eastward acceleration and poleward shift of the Atlantic jet (Orbe et al., 2024). Although this is out of the scope of this work,
it emphasizes the extensive effect that the ozone response may have and thus the importance of including ozone interactive
chemistry in climate sensitivity studies. Therefore, it highlights the need for using the ozone field simulated in chem models as
the forcing for no-chem models in future model intercomparison projects such as CMIP7.

440 A caveat of this work is that the abundance of ODSs in the three experiments is fixed at pre-industrial level to separate the
effect of CO₂, thus the effect of anthropogenic halogens cannot be simulated. The present-day ODSs level is high, which will
lead to ozone depletion through heterogeneous chemistry in polar stratospheric clouds (PSCs) with CO₂ induced stratospheric
cooling, which may counteract the positive ozone response. However, we fix the ODSs level in this work following the standard
approach of studying climate feedback (Gregory and Webb, 2008; Andrews et al., 2012). Future analyses are needed to study
445 the ozone response considering present-day and future ODSs levels.

The negative (damping) climate feedback from stratospheric ozone changes is in agreement with previous single model
studies (Dietmüller et al., 2014; Nowack et al., 2015) utilizing the current available data from CMIP6. Although three out of
the seven pairs we chose to conduct the chem/no-chem comparison have other differences other than chemistry such as height of
model top, they share similar patterns with those that only differ in chemistry scheme. This indicates that the different chemistry
450 scheme contributes the most to the chem/no-chem comparison and other model differences play a minor role. However, to
isolate the feedback of ozone responses, future experiments which compare the same model system with and without interactive
chemistry directly should be included in future model intercomparison projects such as CMIP7. Lastly, the large difference of
global mean surface air temperature between chem and no-chem models of $\sim 10\%$ cannot be explained by the direct radiative
effect of ozone response alone ($\sim -0.1 \text{ W m}^{-2}$). Other feedbacks, such as via cloud and water vapor changes, as well as biases
455 induced by, e.g., inconsistencies between the chemical and thermal tropopause, might contribute: further studies are needed to
explain the cause of these large differences.

Code and data availability. CMIP6 model datasets used in this study are available through the Earth System Grid Federation (ESGF; <https://esgf-index1.ceda.ac.uk/projects/cmip6-ceda/>). Data and code to reproduce the figures in this work can be found on <https://doi.org/10.5281/zenodo.14545386>.

460 **Appendix A**

Before analyzing the data, some pre-processing processes are necessary for the cohesion of data format and computation of required variables.

A1 Calculation of tropopause

Since the variable for tropopause height is not available for all models in all three experiments, we compute the tropopause
465 height following WMO definition¹. The 19 pressure levels of CMIP6 models are not sufficient to compute the tropopause
height, thus we first interpolate the zonal-mean annual-mean geopotential height and air temperature to 40 pressure levels to
get a better vertical resolution. Then, the annual-mean zonal-mean tropopause height in Pa is calculated. The annual-mean
tropopause height for a time period is derived by averaging the annual-mean values. Note that for the abrupt-4×CO₂ experi-
ment, we only average over the last 100 years to make sure the system reaches equilibrium, and for 1pctCO₂, the time average
470 is done from year 135 to year 145, which is the time series selected around the year 140 when the CO₂ concentration reaches
4×CO₂.

A2 Remap of MRI-ESM2-0

The horizontal resolution of air temperature of MRI-ESM2-0 is remapped so that it is the same as that of its ozone mixing
ratio.

475 **A3 Vertical wind velocity for UKESM1-0-LL and GFDL-ESM4**

Since the vertical wind velocity in Pa s⁻¹ (wap) variable for UKESM1-0-LL piControl is not available, We average over the
first two decades (1850-1870) from historical runs of five ensemble members (r9-r13) to represent the equilibrium state of
piControl experiment.

Similarly, since the vertical velocity of the residual mean meridional circulation ($\overline{w^*}$) variable for GFDL-ESM4 piControl
480 is not available, We average over the first three decades (1850-1880) from historical run to represent the equilibrium state of
piControl experiment.

¹It is defined as the lowest level at which the lapse rate decreases to 2k km⁻¹ or less, provided also that the average lapse rate between this level and all
higher levels within 2 km does not exceed 2K⁻¹.

A4 Conversion from ω to w

When analyzing vertical velocity, the only available variable is omega in Pa s^{-1} , thus to better represent the vertical motion, we convert ω to w in mm s^{-1} using Eq.(A1):

$$w_j = -1000\omega_j \frac{H}{P_j} \quad (\text{A1})$$

In which w_j is the vertical velocity in mm s^{-1} on j-th pressure level; ω_j is the vertical velocity in Pa s^{-1} on j-th pressure level; H is the scale height and P_j is the pressure of j-th pressure level.

A5 Computation of $\overline{w^*}$

Transferred-Eulerian-Mean (TEM) circulation combines the contribution of eddy and mean transport (Butler et al., 2016) and group the eddy fluxes of heat and momentum into the zonal momentum equation (David G. Andrews, 1987). It keeps the benefit of the Eulerian view, but also includes eddy fluxes to understand particles transport from the Lagrangian view, and it is usually used in stratosphere.

Since w^* for SOCOL-MPIOM is not provided, we compute them following Eq.(A2):

$$\overline{w^*} = \overline{w} + (a \cos \phi)^{-1} \frac{\partial(\cos \phi \overline{v' \theta'}) / (\partial \overline{\theta} / \partial z)}{\partial \phi} \quad (\text{A2})$$

In which $\overline{w^*}$ is the mean vertical velocity of the residual mean meridional circulation, $\overline{w}$ is the mean vertical velocity, a is the mean radius of Earth, ϕ is latitude, v' is the deviation of meridional velocity from zonal mean value, $\overline{\theta}$ is the zonal mean potential temperature and θ' is the deviation of potential temperature from zonal mean value.

A6 Computation of global-mean surface temp

The global mean surface temperature is computed using weighted mean of surface temperature as Eq.(A3):

$$T_{glbm} = \sum_{i=1}^{i=N} \sigma_i T_i \quad (\text{A3})$$

In which, T_{glbm} is the global mean surface temperature, N is the number of grid cells, σ_i is the area of the i-th grid cell, and T_i is the average surface temperature of the i-th grid cell, which is represented by the temperature at the corresponding grid point.

A7 Computation of column ozone

Column ozone is computed by integrating the number of ozone molecule between certain pressure levels, and then converting it to Dobson unit (DU) following Eq.(A4):

$$Col_{ozone} = \frac{1}{2.69 \times 10^{16}} \int_{P_1}^{P_2} \frac{10vmr}{M_{air}g} dP \quad (A4)$$

In which, Col_{ozone} is the column ozone value in DU, P_1 and P_2 are the starting and ending pressure level in Pa, vmr is the ozone volume mixing ratio, M_{air} is the mass of one air molecule in gram.

510 A8 Comparison of chem and no-chem models

For the comparison of chem and no-chem models, We first remap all models' horizontal grid to the same resolution, which is chosen as 288×192 . Since all CMIP6 models have 19 pressure levels, vertical interpolation is done on SOCOLv4 so that the data are on the same pressure levels.

Appendix B




Figure B1. Annual-mean ozone response to 1pctCO₂ of each chem model. Tropopause for piControl (1pctCO₂) is denoted using black (red dotted) curve. Regions that are not stippled are statistically significant (at the 99% level), according to the t test.




Figure B2. 150-year-long seasonal-mean ozone response in 60-90°S to zonal wind (u) change in 50-70°S at 70hPa in JJA and SON for 1pctCO₂. Fitting lines retrieved from linear regression are plotted as dash-dotted lines with the corresponding color for each model. The thick black line is fitted using data from all models with the corresponding R^2 denoted in the upper right corner of the plot. R^2 values for each model are denoted in the legend for JJA and SON respectively.




Figure B3. Annual-mean surface air temperature response to $4\times\text{CO}_2$. Regions that are not stippled are statistically significant.




Figure B4. (a)-(b): Annual-mean tropical (15°S-15°N) residual upwelling ($\overline{w^*}$) response to global mean surface temperature for 4×CO₂ and 1pctCO₂. (c)-(d): annual-mean tropical (15°S-15°N) ozone response at 50hPa to upwelling (w) change at 70hPa for 4×CO₂ and 1pctCO₂. (e)-(f): same as (c)-(d), but averaged over the last 100 years for 4×CO₂ and for all the 150 years for 1pctCO₂ experiment. Fitting lines retrieved from linear regression are plotted with the corresponding color of each model.




Figure B5. 150-year-long annual-mean tropical (20°S-20°N) LSO3, USO3 and TCO response to 1pctCO₂ with time.




Figure B6. Annual-mean air temperature response to $4\times\text{CO}_2$ of each chem model.




Figure B7. Multi-model average of annual-mean zonal wind response to $4\times\text{CO}_2$ at 850hPa in DJF for chem and no-chem models, along with the difference between the two categories of models.

Table B1. Residual tropical (20°S-20°N) upwelling ($\overline{w^*}$) change due to 4×CO₂ at 70 hPa (as provided on ESGF via the DynVarMip initiative) and average SSW frequency (identified based on Charlton and Polvani (2007) and calculated as number of SSW events per year) in piControl and 4×CO₂ in Chem vs. No-chem pairs.

Pair (Chem/No-chem)	SSW (Chem/No-chem)		$\overline{w^*}$ (Chem/No-chem) (mm s ⁻¹)
	piControl	4×CO ₂	4×CO ₂ - piControl
CESM2-WACCM/CESM2	0.43/0.23	0.82/0.37	0.23/0.49
CESM2-WACCM-FV2/CESM2-FV2	0.52/0.4	0.71/0.32	-/-
GFDL-ESM4/GFDL-CM4	0.52/0.29	0.94/0.34	0.11/0.19
UKESM1-0-LL/HadGEM3-GC31-LL	0.59/0.83	0.42/0.29	0.13/0.16
SOCOLv4/MPI-ESM1-2-LR	0.72/0.95	1.02/0.65	0.10/-

515 *Author contributions.* JW processed, analyzed the CMIP6 model datasets, run the SOCOLv4 model experiments, analyzed the SOCOLv4 data and drafted the manuscript. GC supervised JW in this project, helped with data analysis, and also discussion and edition of the manuscript. BA conducted the SSW analysis. MD, BH, JK, PN, CO, SV and BA gave suggestions on the data and results analysis. TS prepared the SOCOLv4 model experiments, discussed and edited the manuscript. All authors contributed to the preparation of the manuscript.

Competing interests. At least one of the (co-)authors is a member of the editorial board of the Atmospheric Chemistry and Physics journal.
520 Besides this, we have no other competing interests to declare.

Acknowledgements. We acknowledge the World Climate Research Programme (WCRP) for coordinating and promoting CMIP6 through its Working Group on Coupled Modelling. We thank the climate modelling groups, who produced and made available their model output, the Earth System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies who support CMIP6 and ESGF. We thank DynVarMIP for providing the TEM analysis metrics. We also thank the GFDL model development team, who led the
525 development of GFDL-CM4 and GFDL-ESM4, as well as the numerous scientists and technical staff at GFDL who contributed to the development of these models and conducted the CMIP6 simulations. SOCOL simulations have been performed at the ETH cluster EULER and Swiss National Supercomputing Centre (CSCS) under projects s1191 and s1144. We thank Urs Beyerle (IAC ETH) for his help with acquiring and postprocessing the CMIP6 data. Jingyu Wang acknowledges support from the University of Arizona startup funds (PI: S. Ranjan). Support for Gabriel Chiodo was provided by the Swiss National Science Foundation within the Ambizione grant no. PZ00P2_180043, the
530 European Research Council within the ERC StG project no. 101078127, and the Spanish Ministry of Science and Innovation via the Ramon y Cajal grant no. RYC2021-033422-I. Support for Sandro Vattioni was provided by the ETH Research grant no. ETH-1719-2 as well as by the Harvard Geoengineering Research Program. Sandro Vattioni received also funding from the Simons Foundation (grant no. SFI-MPS-SRM-00005217). Timofei Sukhodolov acknowledges the support from the Swiss National Science Foundation (grant no. 200020E_219166) and the Karbacher Fonds, Graubünden, Switzerland. Timofei Sukhodolov and Gabriel Chiodo also acknowledge the support from the Si-
535 mons foundation (SFI-MPS-SRM-00005208). Birgit Hassler acknowledges the support from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No. 101003536 (ESM2025—Earth System Models for the Future). Blanca Ayarza-güena acknowledges support from the project Stratospheric Ozone recovery in the Northern Hemisphere under climate change (RecO3very): PID2021-124772OB-I00 from the Spanish Ministry of Science and Innovation.

References

- 540 Abalos, M., Orbe, C., Kinnison, D. E., Plummer, D., Oman, L. D., Jöckel, P., Morgenstern, O., Garcia, R. R., Zeng, G., Stone, K. A., and Dameris, M.: Future trends in stratosphere-to-troposphere transport in CCMI models, *Atmospheric Chemistry and Physics*, 20, 6883–6901, <https://doi.org/10.5194/acp-20-6883-2020>, 2020.
- Abalos, M., Calvo, N., Benito-Barca, S., Garny, H., Hardiman, S. C., Lin, P., Andrews, M. B., Butchart, N., Garcia, R., Orbe, C., Saint-Martin, D., Watanabe, S., and Yoshida, K.: The Brewer–Dobson circulation in CMIP6, *Atmospheric Chemistry and Physics*, 21, 13 571–13 591, <https://doi.org/10.5194/acp-21-13571-2021>, 2021.
- 545 Andrews, T., Gregory, J. M., Webb, M. J., and Taylor, K. E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, *Geophysical Research Letters*, 39, <https://doi.org/https://doi.org/10.1029/2012GL051607>, 2012.
- Ayarzagüena, B., Charlton-Perez, A. J., Butler, A. H., Hitchcock, P., Simpson, I. R., Polvani, L. M., Butchart, N., Gerber, E. P., Gray, L., Hassler, B., Lin, P., Lott, F., Manzini, E., Mizuta, R., Orbe, C., Osprey, S., Saint-Martin, D., Sigmond, M., Taguchi, M., Volodin, E. M., and Watanabe, S.: Uncertainty in the Response of Sudden Stratospheric Warmings and Stratosphere-Troposphere Coupling to Quadrupled CO₂ Concentrations in CMIP6 Models, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032 345, <https://doi.org/https://doi.org/10.1029/2019JD032345>, 2020.
- 550 Banerjee, A., Archibald, A. T., Maycock, A. C., Telford, P., Abraham, N. L., Yang, X., Braesicke, P., and Pyle, J. A.: Lightning NO_x, a key chemistry–climate interaction: impacts of future climate change and consequences for tropospheric oxidising capacity, *Atmospheric Chemistry and Physics*, 14, 9871–9881, <https://doi.org/10.5194/acp-14-9871-2014>, 2014.
- Banerjee, A., Chiodo, G., Previdi, M., Ponater, M., Conley, A. J., and Polvani, L. M.: Stratospheric water vapor: an important climate feedback, *Climate Dynamics*, 53, 1697–1710, <https://doi.org/10.1007/s00382-019-04721-4>, 2019.
- Barnett, J. J., Houghton, J. T., and Pyle, J. A.: The temperature dependence of the ozone concentration near the stratopause, *Quarterly Journal of the Royal Meteorological Society*, 101, 245–257, <https://doi.org/https://doi.org/10.1002/qj.49710142808>, 1975.
- 560 Brasseur, G. and Solomon, S.: *Aeronomy of the Middle Atmosphere: Chemistry and Physics of the Stratosphere and Mesosphere*, <https://doi.org/10.1007/1-4020-3824-0>, 2005.
- Butchart, N.: The Brewer-Dobson circulation, *Reviews of Geophysics*, 52, 157–184, <https://doi.org/https://doi.org/10.1002/2013RG000448>, 2014.
- Butler, A. H., Daniel, J. S., Portmann, R. W., Ravishankara, A. R., Young, P. J., Fahey, D. W., and Rosenlof, K. H.: Diverse policy implications for future ozone and surface UV in a changing climate, *Environmental Research Letters*, 11, <https://doi.org/10.1088/1748-9326/11/6/064017>, 2016.
- 565 Charlton, A. J. and Polvani, L. M.: A New Look at Stratospheric Sudden Warmings. Part I: Climatology and Modeling Benchmarks, *Journal of Climate*, 20, 449–469, <https://doi.org/https://doi.org/10.1175/JCLI3996.1>, 2007.
- Checa-Garcia, R., Hegglin, M. I., Kinnison, D., Plummer, D. A., and Shine, K. P.: Historical Tropospheric and Stratospheric Ozone Radiative Forcing Using the CMIP6 Database, *Geophysical Research Letters*, 45, 3264–3273, <https://doi.org/https://doi.org/10.1002/2017GL076770>, 2018.
- 570 Chiodo, G. and Polvani, L. M.: Reduction of climate sensitivity to solar forcing due to stratospheric ozone feedback, *Journal of Climate*, 29, 4651–4663, <https://doi.org/10.1175/JCLI-D-15-0721.1>, 2016.
- Chiodo, G. and Polvani, L. M.: Reduced Southern Hemispheric circulation response to quadrupled CO₂ due to stratospheric ozone feedback, *Geophysical Research Letters*, 44, 465–474, <https://doi.org/10.1002/2016GL071011>, 2017.
- 575

- Chiodo, G. and Polvani, L. M.: The Response of the Ozone Layer to Quadrupled CO₂ Concentrations: Implications for Climate, *Journal of Climate*, 32, 7629–7642, <https://doi.org/10.1175/JCLI-D-19-0086.1>, 2019.
- Chiodo, G. and Polvani, L. M.: New Insights on the Radiative Impacts of Ozone-Depleting Substances, *Geophysical Research Letters*, 49, e2021GL096783, <https://doi.org/https://doi.org/10.1029/2021GL096783>, 2022.
- 580 Chiodo, G., Polvani, L. M., Marsh, D. R., Stenke, A., Ball, W., Rozanov, E., Muthers, S., and Tsigaridis, K.: The Response of the Ozone Layer to Quadrupled CO₂ Concentrations, *Journal of Climate*, 31, 3893–3907, <https://doi.org/10.1175/JCLI-D-17-0492.1>, 2018.
- Chiodo, G., Friedel, M., Seeber, S., Domeisen, D., Stenke, A., Sukhodolov, T., and Zilker, F.: The influence of future changes in springtime Arctic ozone on stratospheric and surface climate, *Atmospheric Chemistry and Physics*, 23, 10451–10472, <https://doi.org/10.5194/acp-23-10451-2023>, 2023.
- 585 Chrysanthou, A., Maycock, A. C., and Chipperfield, M. P.: Decomposing the response of the stratospheric Brewer–Dobson circulation to an abrupt quadrupling in CO₂, *Weather and Climate Dynamics*, 1, 155–174, <https://doi.org/10.5194/wcd-1-155-2020>, 2020.
- Conley, A. J., Lamarque, J.-F., Vitt, F., Collins, W. D., and Kiehl, J.: PORT, a CESM tool for the diagnosis of radiative forcing, *Geoscientific Model Development*, 6, 469–476, <https://doi.org/10.5194/gmd-6-469-2013>, 2013.
- 590 David G. Andrews, James R. Holton, C. B. L.: Chapter 3 - Basic Dynamics, in: *Middle Atmosphere Dynamics*, edited by Andrews, D. G., Holton, J. R., and Leovy, C. B., vol. 40 of *International Geophysics*, pp. 113–149, Academic Press, <https://doi.org/https://doi.org/10.1016/B978-0-12-058575-5.50008-6>, 1987.
- Dietmüller, S., Ponater, M., and Sausen, R.: Interactive ozone induces a negative feedback in CO₂-driven climate change simulations, *Journal of Geophysical Research: Atmospheres*, 119, 1796–1805, <https://doi.org/https://doi.org/10.1002/2013JD020575>, 2014.
- 595 Dütsch, H., Bader, J., and Staehelin, J.: Separation of Solar Effects on Ozone from Anthropogenically Produced Trends, *Journal of geomagnetism and geoelectricity*, 43, 657–665, https://doi.org/10.5636/jgg.43.Supplement2_657, 1991.
- Egorova, T., Rozanov, E., Zubov, V. A., and Karol, I.: Model for investigating ozone trends (MEZON), *Izvestiya - Atmospheric and Ocean Physics*, 39, 277–292, 2003.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 600 Fels, S. B., Mahlman, J. D., Schwarzkopf, M. D., and Sinclair, R. W.: Stratospheric Sensitivity to Perturbations in Ozone and Carbon Dioxide: Radiative and Dynamical Response, *Journal of Atmospheric Sciences*, 37, 2265 – 2297, [https://doi.org/10.1175/1520-0469\(1980\)037<2265:SSTPIO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2265:SSTPIO>2.0.CO;2), 1980.
- 605 Flynn, C. M. and Mauritsen, T.: On the climate sensitivity and historical warming evolution in recent coupled model ensembles, *Atmospheric Chemistry and Physics*, 20, 7829–7842, <https://doi.org/10.5194/acp-20-7829-2020>, 2020.
- Friedel, M., Chiodo, G., Stenke, A., Domeisen, D. I. V., Fueglistaler, S., Anet, J. G., and Peter, T.: Springtime arctic ozone depletion forces northern hemisphere climate anomalies, *Nature Geoscience*, 15, 541–547, <https://doi.org/10.1038/s41561-022-00974-7>, 2022.
- Gerber, E. P. and Manzini, E.: The Dynamics and Variability Model Intercomparison Project (DynVarMIP) for CMIP6: assessing the stratosphere–troposphere system, *Geoscientific Model Development*, 9, 3413–3425, <https://doi.org/10.5194/gmd-9-3413-2016>, 2016.
- 610 Gregory, J. and Webb, M.: Tropospheric Adjustment Induces a Cloud Component in CO₂ Forcing, *Journal of Climate*, 21, 58–71, <https://doi.org/https://doi.org/10.1175/2007JCLI1834.1>, 2008.

- Haase, S. and Matthes, K.: The importance of interactive chemistry for stratosphere–troposphere coupling, *Atmospheric Chemistry and Physics*, 19, 3417–3432, <https://doi.org/10.5194/acp-19-3417-2019>, 2019.
- 615 Haigh, J. D. and Pyle, J. A.: Ozone perturbation experiments in a two-dimensional circulation model, *Quarterly Journal of the Royal Meteorological Society*, 108, 551–574, <https://doi.org/https://doi.org/10.1002/qj.49710845705>, 1982.
- Hardiman, S. C., Andrews, M. B., Andrews, T., Bushell, A. C., Dunstone, N. J., Dyson, H., Jones, G. S., Knight, J. R., Neining, E., O’Connor, F. M., Ridley, J. K., Ringer, M. A., Scaife, A. A., Senior, C. A., and Wood, R. A.: The Impact of Prescribed Ozone in Climate Projections Run With HadGEM3-GC3.1, *Journal of Advances in Modeling Earth Systems*, 11, 3443–3453, <https://doi.org/10.1029/2019MS001714>, 2019.
- 620 Hegglin, M. I. and Shepherd, T. G.: Large climate-induced changes in ultraviolet index and stratosphere-to-troposphere ozone flux, *Nature Geoscience*, 2, 687–691, <https://doi.org/10.1038/ngeo604>, 2009.
- Hufnagl, L., Eichinger, R., Garny, H., Birner, T., Kuchař, A., Jöckel, P., and Graf, P.: Stratospheric Ozone Changes Damp the CO₂-Induced Acceleration of the Brewer–Dobson Circulation, *Journal of Climate*, 36, 3305–3320, [https://doi.org/https://doi.org/10.1175/JCLI-D-22-](https://doi.org/https://doi.org/10.1175/JCLI-D-22-0512.1)
- 625 0512.1, 2023.
- Iglesias-Suarez, F., Kinnison, D. E., Rap, A., Maycock, A. C., Wild, O., and Young, P. J.: Key drivers of ozone change and its radiative forcing over the 21st century, *Atmospheric Chemistry and Physics*, 18, 6121–6139, <https://doi.org/10.5194/acp-18-6121-2018>, 2018.
- Jonsson, A. I., de Grandpré, J., Fomichev, V. I., McConnell, J. C., and Beagley, S. R.: Doubled CO₂-induced cooling in the middle atmosphere: Photochemical analysis of the ozone radiative feedback, *Journal of Geophysical Research D: Atmospheres*, 109, 1–18, <https://doi.org/10.1029/2004JD005093>, 2004.
- 630 Jucks, K. W. and Salawitch, R. J.: Future Changes in Upper Stratospheric Ozone, in: *Atmospheric Science Across the Stratopause*, Geophysical Monograph Series, pp. 241–255, <https://doi.org/https://doi.org/10.1029/GM123p0241>, 2000.
- Jungclaus, J. H., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and von Storch, J. S.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *Journal of Advances in Modeling Earth Systems*, 5, 422–446, <https://doi.org/https://doi.org/10.1002/jame.20023>, 2013.
- Karpechko, A. Y., Afargan-Gerstman, H., Butler, A. H., Domeisen, D. I. V., Kretschmer, M., Lawrence, Z., Manzini, E., Sigmond, M., Simpson, I. R., and Wu, Z.: Northern Hemisphere Stratosphere-Troposphere Circulation Change in CMIP6 Models: 1. Inter-Model Spread and Scenario Sensitivity, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036992, <https://doi.org/https://doi.org/10.1029/2022JD036992>, 2022.
- 640 Karpechko, A. Y., Wu, Z., Simpson, I. R., Kretschmer, M., Afargan-Gerstman, H., Butler, A. H., Domeisen, D. I. V., Garny, H., Lawrence, Z., Manzini, E., and Sigmond, M.: Northern Hemisphere Stratosphere-Troposphere Circulation Change in CMIP6 Models: 2. Mechanisms and Sources of the Spread, *Journal of Geophysical Research: Atmospheres*, 129, e2024JD040823, <https://doi.org/https://doi.org/10.1029/2024JD040823>, 2024.
- Keeble, J., Bednarz, E. M., Banerjee, A., Abraham, N. L., Harris, N. R. P., Maycock, A. C., and Pyle, J. A.: Diagnosing the radiative and chemical contributions to future changes in }tropical column ozone with the UM-UKCA chemistry–climate model, *Atmospheric Chemistry and Physics*, 17, 13801–13818, <https://doi.org/10.5194/acp-17-13801-2017>, 2017.
- Keeble, J., Hassler, B., Banerjee, A., Checa-Garcia, R., Chiodo, G., Davis, S., Eyring, V., Griffiths, P. T., Morgenstern, O., Nowack, P., Zeng, G., Zhang, J., Bodeker, G., Burrows, S., Cameron-Smith, P., Cugnet, D., Danek, C., Deushi, M., Horowitz, L. W., Kubin, A., Li, L., Lohmann, G., Michou, M., Mills, M. J., Nabat, P., Olivie, D., Park, S., Seland, Ø., Stoll, J., Wieners, K. H., and Wu, T.: Evaluating
- 650

- stratospheric ozone and water vapour changes in CMIP6 models from 1850 to 2100, *Atmospheric Chemistry and Physics*, 21, 5015–5061, <https://doi.org/10.5194/acp-21-5015-2021>, 2021.
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezzuman, K., Miller, R. L., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 025, <https://doi.org/10.1029/2019MS002025>, 2020.
- Kult-Herdin, J., Sukhodolov, T., Chiodo, G., Checa-Garcia, R., and Rieder, H. E.: The impact of different CO₂ and ODS levels on the mean state and variability of the springtime Arctic stratosphere, *Environmental Research Letters*, 18, <https://doi.org/10.1088/1748-9326/acb0e6>, 2023.
- Li, F. and Newman, P. A.: Prescribing stratospheric chemistry overestimates southern hemisphere climate change during austral spring in response to quadrupled CO₂, *Climate Dynamics*, 61, 1105–1122, <https://doi.org/10.1007/s00382-022-06588-4>, 2023.
- London, J.: Radiative energy sources and sink in the stratosphere and mesosphere, in: *Proceeding of the NATO Advanced Study Institute on Atmospheric Ozone: Its Variation and Human Influences*, 1980.
- Marsh, D. R., Mills, M. J., Kinnison, D. E., Lamarque, J.-F., Calvo, N., and Polvani, L. M.: Climate Change from 1850 to 2005 Simulated in CESM1(WACCM), *Journal of Climate*, 26, 7372 – 7391, <https://doi.org/10.1175/JCLI-D-12-00558.1>, 2013.
- Marsh, D. R., Lamarque, J. F., Conley, A. J., and Polvani, L. M.: Stratospheric ozone chemistry feedbacks are not critical for the determination of climate sensitivity in CESM1(WACCM), *Geophysical Research Letters*, 43, 3928–3934, <https://doi.org/10.1002/2016GL068344>, 2016.
- Match, A. and Gerber, E. P.: Tropospheric Expansion Under Global Warming Reduces Tropical Lower Stratospheric Ozone, *Geophysical Research Letters*, 49, <https://doi.org/10.1029/2022GL099463>, 2022.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E., Nam, C. C., Notz, D., Nyawira, S. S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J. S., Tian, F., Voigt, A., Vrese, P., Wieners, K. H., Wilkenskeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.
- McLandress, C., Shepherd, T. G., Scinocca, J. F., Plummer, D. A., Sigmond, M., Jonsson, A. I., and Reader, M. C.: Separating the Dynamical Effects of Climate Change and Ozone Depletion. Part II: Southern Hemisphere Troposphere, *Journal of Climate*, 24, 1850 – 1868, <https://doi.org/10.1175/2010JCLI3958.1>, 2011.
- Meul, S., Langematz, U., Oberländer, S., Garny, H., and Jöckel, P.: Chemical contribution to future tropical ozone change in the lower stratosphere, *Atmospheric Chemistry and Physics*, 14, 2959–2971, <https://doi.org/10.5194/acp-14-2959-2014>, 2014.
- Meul, S., Oberländer-Hayn, S., Abalichin, J., and Langematz, U.: Nonlinear response of modelled stratospheric ozone to changes in greenhouse gases and ozone depleting substances in the recent past, *Atmospheric Chemistry and Physics*, 15, 6897–6911, <https://doi.org/10.5194/acp-15-6897-2015>, 2015.

- Morgenstern, O.: The Southern Annular Mode in 6th Coupled Model Intercomparison Project Models, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034161, <https://doi.org/https://doi.org/10.1029/2020JD034161>, 2021.
- 690 Morgenstern, O., Stone, K. A., Schofield, R., Akiyoshi, H., Yamashita, Y., Kinnison, D. E., Garcia, R. R., Sudo, K., Plummer, D. A., Scinocca, J., Oman, L. D., Manyin, M. E., Zeng, G., Rozanov, E., Stenke, A., Revell, L. E., Pitari, G., Mancini, E., DI Genova, G., Visioni, D., Dhomse, S. S., and Chipperfield, M. P.: Ozone sensitivity to varying greenhouse gases and ozone-depleting substances in CCMI-1 simulations, *Atmospheric Chemistry and Physics*, 18, 1091–1114, <https://doi.org/10.5194/acp-18-1091-2018>, 2018.
- 695 Morgenstern, O., Kinnison, D. E., Mills, M., Michou, M., Horowitz, L. W., Lin, P., Deushi, M., Yoshida, K., O'Connor, F. M., Tang, Y., Abraham, N. L., Keeble, J., Dennison, F., Rozanov, E., Egorova, T., Sukhodolov, T., and Zeng, G.: Comparison of Arctic and Antarctic Stratospheric Climates in Chemistry Versus No-Chemistry Climate Models, *Journal of Geophysical Research: Atmospheres*, 127, <https://doi.org/10.1029/2022JD037123>, 2022.
- Muthers, S., Anet, J. G., Stenke, A., Raible, C. C., Rozanov, E., Brönnimann, S., Peter, T., Arfeuille, F. X., Shapiro, A. I., Beer, J., Steinhilber, F., Bruhnara, Y., and Schmutz, W.: The coupled atmosphere–chemistry–ocean model SOCOL-MPIOM, *Geoscientific Model Development*, 7, 2157–2179, <https://doi.org/10.5194/gmd-7-2157-2014>, 2014.
- 700 Muthers, S., Arfeuille, F., Raible, C. C., and Rozanov, E.: The impacts of volcanic aerosol on stratospheric ozone and the Northern Hemisphere polar vortex: separating radiative-dynamical changes from direct effects due to enhanced aerosol heterogeneous chemistry, *Atmospheric Chemistry and Physics*, 15, 11461–11476, <https://doi.org/10.5194/acp-15-11461-2015>, 2015.
- 705 Muthers, S., Raible, C. C., Rozanov, E., and Stocker, T. F.: Response of the AMOC to reduced solar radiation – the modulating role of atmospheric chemistry, *Earth System Dynamics*, 7, 877–892, <https://doi.org/10.5194/esd-7-877-2016>, 2016.
- Nowack, P., Ceppi, P., Davis, S. M., Chiodo, G., Ball, W., Diallo, M. A., Hassler, B., Jia, Y., Keeble, J., and Joshi, M.: Response of stratospheric water vapour to warming constrained by satellite observations, *Nature Geoscience*, 16, 577–583, <https://doi.org/10.1038/s41561-023-01183-6>, 2023.
- 710 Nowack, P. J., Luke Abraham, N., Maycock, A. C., Braesicke, P., Gregory, J. M., Joshi, M. M., Osprey, A., and Pyle, J. A.: A large ozone-circulation feedback and its implications for global warming assessments, *Nature Climate Change*, 5, 41–45, <https://doi.org/10.1038/nclimate2451>, 2015.
- Nowack, P. J., Braesicke, P., Luke Abraham, N., and Pyle, J. A.: On the role of ozone feedback in the ENSO amplitude response under global warming, *Geophysical Research Letters*, 44, 3858–3866, <https://doi.org/https://doi.org/10.1002/2016GL072418>, 2017.
- 715 Nowack, P. J., Abraham, N. L., Braesicke, P., and Pyle, J. A.: The Impact of Stratospheric Ozone Feedbacks on Climate Sensitivity Estimates, *Journal of Geophysical Research: Atmospheres*, 123, 4630–4641, <https://doi.org/https://doi.org/10.1002/2017JD027943>, 2018.
- Oman, L. D., Plummer, D. A., Waugh, D. W., Austin, J., Scinocca, J. F., Douglass, A. R., Salawitch, R. J., Canty, T., Akiyoshi, H., Bekki, S., Braesicke, P., Butchart, N., Chipperfield, M. P., Cugnet, D., Dhomse, S., Eyring, V., Frith, S., Hardiman, S. C., Kinnison, D. E., Lamarque, J.-F., Mancini, E., Marchand, M., Michou, M., Morgenstern, O., Nakamura, T., Nielsen, J. E., Oliv  , D., Pitari, G., Pyle, J., Rozanov, E., Shepherd, T. G., Shibata, K., Stolarski, R. S., Teyss  dre, H., Tian, W., Yamashita, Y., and Ziemke, J. R.: Multimodel assessment of the factors driving stratospheric ozone evolution over the 21st century, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/https://doi.org/10.1029/2010JD014362>, 2010.
- 720 Orbe, C., Rind, D., Jonas, J., Nazarenko, L., Faluvegi, G., Murray, L. T., Shindell, D. T., Tsigaridis, K., Zhou, T., Kelley, M., and Schmidt, G. A.: GISS Model E2.2: A Climate Model Optimized for the Middle Atmosphere—2. Validation of Large-Scale Transport and Evaluation of Climate Response, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD033151, <https://doi.org/https://doi.org/10.1029/2020JD033151>, 2020.

- Orbe, C., Rind, D., Waugh, D. W., Jonas, J., Zhang, X., Chiodo, G., Nazarenko, L., and Schmidt, G. A.: Coupled Stratospheric Ozone and Atlantic Meridional Overturning Circulation Feedbacks on the Northern Hemisphere Midlatitude Jet Response to 4xCO₂, *Journal of Climate*, 37, 2897–2917, <https://doi.org/https://doi.org/10.1175/JCLI-D-23-0119.1>, 2024.
- 730 Ramaswamy, V., Boucher O., Haigh J., Hauglustaine D., Haywood J., Myhre, G., Nkajima T., Shi G. Y., and Solomon S.: Climate Change 2001: The Scientific Basis, in: Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change (IPCC), pp. 350–416, 2001.
- Revell, L. E., Bodeker, G. E., Huck, P. E., Williamson, B. E., and Rozanov, E.: The sensitivity of stratospheric ozone changes through the 21st century to N₂O and CH₄, *Atmospheric Chemistry and Physics*, 12, 11 309–11 317, <https://doi.org/10.5194/acp-12-11309-2012>, 2012.
- 735 Revell, L. E., Tummon, F., Stenke, A., Sukhodolov, T., Coulon, A., Rozanov, E., Garny, H., Grewe, V., and Peter, T.: Drivers of the tropospheric ozone budget throughout the 21st century under the medium-high climate scenario RCP 6.0, *Atmospheric Chemistry and Physics*, 15, 5887–5902, <https://doi.org/10.5194/acp-15-5887-2015>, 2015.
- Rind, D., Orbe, C., Jonas, J., Nazarenko, L., Zhou, T., Kelley, M., Lacis, A., Shindell, D., Faluvegi, G., Romanou, A., Russell, G., Tausnev, N., Bauer, M., and Schmidt, G.: GISS Model E2.2: A Climate Model Optimized for the Middle Atmosphere—Model
- 740 Structure, Climatology, Variability, and Climate Sensitivity, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032 204, <https://doi.org/https://doi.org/10.1029/2019JD032204>, 2020.
- Seinfeld, J. H., Pandis, S. N., and Noone, K. J.: Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, *Physics Today*, 51, 88–90, <https://api.semanticscholar.org/CorpusID:98232768>, 1998.
- Sheng, J.-X., Weisenstein, D. K., Luo, B.-P., Rozanov, E., Stenke, A., Anet, J., Bingemer, H., and Peter, T.: Global atmospheric sulfur budget
- 745 under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *Journal of Geophysical Research: Atmospheres*, 120, 256–276, <https://doi.org/https://doi.org/10.1002/2014JD021985>, 2015.
- Shepherd, T. G.: Dynamics, stratospheric ozone, and climate change, *Atmosphere-Ocean*, 46, 117–138, <https://doi.org/10.3137/ao.460106>, 2008.
- Sukhodolov, T., Egorova, T., Stenke, A., Ball, W. T., Brodowsky, C., Chiodo, G., Feinberg, A., Friedel, M., Karagodin-Doyennel, A., Peter, T.,
- 750 Sedlacek, J., Vattioni, S., and Rozanov, E.: Atmosphere-ocean-aerosol-chemistry-climate model SOCOLv4.0: Description and evaluation, *Geoscientific Model Development*, 14, 5525–5560, <https://doi.org/10.5194/gmd-14-5525-2021>, 2021.
- Wang, M. and Fu, Q.: Changes in Stratosphere-Troposphere Exchange of Air Mass and Ozone Concentration in CCM1 Models From 1960 to 2099, *Journal of Geophysical Research: Atmospheres*, 128, e2023JD038 487, <https://doi.org/https://doi.org/10.1029/2023JD038487>, 2023.
- 755 Zubov, V., Rozanov, E. V., Egorova, T. A., Karol, I. L., and Schmutz, W. K.: Role of external factors in the evolution of the ozone layer and stratospheric circulation in 21st century, *Atmospheric Chemistry and Physics*, 13, 4697–4706, <https://api.semanticscholar.org/CorpusID:54635161>, 2012.