

AGILE v0.1: The Open Global Glacier Data Assimilation Framework

Patrick Schmitt¹, Fabien Maussion^{1,2}, Daniel N. Goldberg³, and Philipp Gregor⁴

¹Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria

²School of Geographical Sciences, University of Bristol, Bristol, United Kingdom

³School of Geosciences, University of Edinburgh, Edinburgh, United Kingdom

⁴Meteorologisches Institut, Ludwig-Maximilians-Universität München, München, Germany

Correspondence: Patrick Schmitt (patrick.schmitt@uibk.ac.at)

Received: 15 July 2025 – Discussion started: 11 August 2025

Revised: 16 December 2025 – Accepted: 16 December 2025 – Published:

Abstract. The growing availability of glacier observations poses a challenge for models to integrate this heterogeneous information in a dynamically consistent way. At the same time, estimates of current glacier volume and area remain uncertain, as many global inventories and thickness datasets date back to the early 2000s. We present the Open Global Glacier Data Assimilation Framework, named AGILE, a time-dependent variational method inspired by 4D-Var data assimilation. AGILE is built on a reimplementation of the OGGM flowline glacier evolution model in PyTorch, enabling full differentiability through automatic differentiation (AD). We test AGILE v0.1 in a series of idealized experiments designed to reflect common initialization and calibration scenarios in global glacier modeling. The goal is to recover glacier bed topography and distributed ice volume in 2020 through transient calibration, based on dynamical simulations starting in 1980. In these experiments, we assume a perfectly known mass balance and fixed ice dynamics parameters. While this setup simplifies real-world complexity, it allows us to isolate and evaluate the core functionality of the approach. Our results show that AGILE efficiently optimizes multiple control variables by leveraging AD-derived gradients, requiring only a few iterations to substantially improve upon initial guesses. We also examine the potential to reconstruct earlier glacier states (e.g., in 1980) without direct observations and find that this is fundamentally limited because glacier dynamics are governed by a diffusion equation, which leads to a loss of information about past states over time, even in an idealized setting. Overall, our experiments demonstrate AGILE’s potential as a flexible and ef-

ficient data assimilation framework. Its ability to integrate diverse datasets in a dynamically consistent manner makes it a promising tool for future real-world glacier modeling applications.

1 Introduction

Mountain glaciers are retreating globally (The GlAMBIE Team, 2025; Hugonnet et al., 2021; Zemp et al., 2015), causing significant impacts on sea level rise (e.g., Marzeion et al., 2020; Rounce et al., 2023; Slangen et al., 2022; Zemp et al., 2019), freshwater resources (e.g., Aguayo et al., 2024; Huss and Hock, 2018; Ultee et al., 2022; Wimberly et al., 2025), and ecosystems (e.g., Bosson et al., 2023; Cannone et al., 2008; Gobbi et al., 2021). Predicting these impacts requires dynamic glacier evolution models that rely on past observations to simulate glacier behaviour into the future. However, many glaciers are located in remote and hard-to-access regions, making direct measurements difficult to obtain and scarce. As a result, global models are highly dependent on Earth observation (EO) data from satellites.

Early EO datasets mainly included glacier outlines (e.g., RGI 6 Consortium, 2017; RGI 7 Consortium, 2023) and digital elevation models (e.g., COP-DEM, 2022; NASA JPL, 2020). Recently, more glacier-specific datasets have become available (e.g., Hugonnet et al., 2021; Millan et al., 2022), allowing global glacier models to shift towards using glacier-specific observations for calibration rather than relying on regional averages or data from a few well-studied glaciers

(Zekollari et al., 2024; Marzeion et al., 2012). This shift is reflected in the design choices of the eight large-scale glacier models participating in the recent Glacier Model Intercomparison Project 3 (GlacierMIP3; Zekollari et al., 2025).

The GlacierMIP3 models adopt different calibration and initialization strategies, but share many common features and rely on the same datasets. The initial glacier surface geometry and corresponding start date are defined using RGI version 6 outlines (RGI 6. Consortium, 2017) and their associated timestamps. Mass balance is calibrated separately by matching to a geodetic mass balance observation over a period of approximately 20 years, using data from Hugonnet et al. (2021) (seven models) or Shean et al. (2020) (one model), while assuming a fixed glacier geometry during this period. All models rely on the consensus ice thickness estimates from Farinotti et al. (2019), either by directly using the distributed fields or by matching total glacier volumes on a glacier-specific or regional basis.

Although widely used, the consensus ice thickness estimate from Farinotti et al. (2019) is not a direct observation but rather the mean result from an ensemble of models. In large-scale glacier modelling, however, it is often treated as an observed quantity (i.e. a target to match via calibration) because glacier volume is essential for estimating a plausible initial glacier state. Improving glacier-specific volume estimates in absence of observations remains an active area of research, with several methods currently being developed and tested in selected regions, aiming for future global applications (e.g., Cook et al., 2023; van Pelt and Frank, 2025).

To better approximate initial conditions and account for glacier evolution prior to the outline date, three GlacierMIP3 models include a dynamic spin-up during initialization. In these cases, a past glacier state is defined, and the model is run forward in time to match specific targets. The Community Ice Sheet Model v2.1 (CISM2; Lipscomb et al., 2019) matches the ice thickness field, the Global Glacier Evolution Model ice flow (GloGEMflow; Zekollari et al., 2019) targets the glacier-specific total volume and glacier length from the outline, and the Open Global Glacier Model (OGGM; Mausson et al., 2019) uses the glacier total area, total volume and geodetic mass balance to additionally refine its previously calibrated mass balance by accounting for evolving surface geometry (Aguayo et al., 2024).

Outside GlacierMIP3, an adapted version of OGGM's dynamic spin-up was extended to incorporate a second glacier outline in a small region of the Alps, enabling the model to simultaneously match observed area and volume changes at regional scale for the first time (Hartl et al., 2025). This enhancement allowed performance improvements when tested against additional validation data and increased confidence in the model's projections through 2100 compared to the setup relying solely on globally available datasets.

Another recent approach in the Alps is presented by Cook et al. (2023), who use the Instructed Glacier Model (IGM; Jouvét, 2022), a deep learning based 3D ice flow model. IGM

assimilates various observations such as distributed thickness data (GlaThiDa; Welty et al., 2020) and surface velocity fields (Millan et al., 2022) to initialize glaciers in a dynamically consistent state. The transient simulation then begins from this point, removing the need for a spin up. However, all inputs must correspond to the same timestamp, such as the outline date, since the method is limited to snapshot inversions. IGM also focuses only on ice dynamics and bed reconstruction and does not include a dedicated mass balance model.

Despite these efforts, current approaches face several limitations. Most models neglect past glacier evolution during initialization and do not account for surface geometry changes during calibration. They are tailored to currently available global datasets and struggle to incorporate new observation types or repeated measurements in a seamless way, limiting their flexibility to use all available data on a glacier-specific scale. It is also often implicitly assumed that all observations apply at the outline timestamp. Furthermore, many methods rely on computationally intensive optimization schemes where only one parameter is adjusted at a time, or they apply steady state assumptions to manage the general problem of overparameterization in global glacier modelling (e.g., Rounce et al., 2020).

To overcome these limitations, an ideal calibration method should ensure dynamic consistency, be able to use temporally distributed observations, avoid assumptions about the glacier's dynamic state, and allow the simultaneous optimization of multiple model parameters and state variables. It must also remain computationally efficient to be applicable at regional and global scales.

To move closer to the transient calibration of large scale glacier models, we present a proof-of-concept for the Open Global Glacier Data Assimilation Framework, named AGILE. AGILE is based on a time-dependent variational assimilation approach, inspired by 4D-Var methods (Lorenç, 1997), which extend the spatial domain (x, y, z) of 3D-Var or snapshot approaches by adding time as a fourth dimension. Following this concept, AGILE treats time as a coordinate during assimilation, enabling the consistent integration of temporally distributed observations.

In line with this concept, AGILE iteratively adjusts all control variables of a dynamic glacier evolution model during a transient (forward) simulation to minimize a cost function that quantifies the mismatch between model output and available observations. To ensure computational efficiency, AGILE is implemented in PyTorch, a machine learning framework (Paszke et al., 2019), which enables the use of automatic differentiation (AD). AD-based methods have previously been applied to snapshot inversions in regional glacier modeling (Cook et al., 2023) and in ice sheet modeling (Brinkerhoff and Johnson, 2013), as well as to transient assimilation in ice sheet modeling (Goldberg and Heimbach, 2013; Recinos et al., 2023). To our knowledge, AGILE rep-

resents the first application of a transient AD-based assimilation approach in the context of global glacier modeling.

In this study, we introduce AGILE v0.1, which reimplements the dynamic flowline model of OGGM in PyTorch to make it fully differentiable through AD (Sect. 2). We evaluate the implementation using idealized experiments with synthetic glaciers and observations designed to reflect globally available datasets (Sect. 3). The objective is to initialize a dynamically consistent glacier by optimizing bed topography and ice volume distribution as control variables, assuming a perfectly known mass balance and fixed dynamic parameters. While these assumptions simplify real-world complexity, they allow us to isolate and assess how effectively AD guides the optimization of control variables during a transient model run (Sect. 4). We conclude by discussing the broader implications of our results and outlining future directions for real-world applications (Sect. 5).

2 Methods

AGILE is developed as an open-source extension of the Open Global Glacier Model (OGGM) framework (Maussion et al., 2019), with the code publicly available at <https://github.com/OGGM/AGILE> (last access: 21 January 2026), and version 0.1 archived on Zenodo (Schmitt et al., 2025). The following sections provide a detailed description of the underlying methodology and its individual components.

2.1 General principles

The general approach of AGILE is illustrated in Fig. 1. A set of initial control variables, referred to as the first guess and which may represent a combination of model parameters, state variables or boundary conditions, is iteratively adjusted to minimize the mismatch between the output of a glacier evolution model and available observations. This mismatch is quantified by a cost function, which combines the discrepancies across all observational targets into a single value, with each contribution normalized by its associated uncertainty. In addition, regularization terms are included in the cost function to address the ill-posed nature of the problem and to reduce the risk of overfitting. The optimization aims to minimize this cost function by systematically adjusting the control variables.

To support this process efficiently, AGILE leverages automatic differentiation (AD) through PyTorch. This enables flexible definition of control variables while still allowing the calculation of their gradients with respect to the cost function. These gradients guide the optimization and allow for efficient and simultaneous updates of all variables. More technical details about the AD implementation in PyTorch are provided in Appendix A1.




Figure 1. Principle workflow of AGILE: Starting with a First Guess at the top, an initial glacier evolution model run is performed (blue line). The mismatch between the model output and observations (depicted in green), in particular one total volume, one volume change and one total area, is calculated using a cost function. If the cost function has not reached a minimum, all control variables (i.e., the unknown variables we aim to estimate) are updated simultaneously using AD, and a new glacier evolution model run is conducted (orange line).

2.2 Glacier representation and forward model

AGILE re-implements the glacier evolution model from OGGM (Maussion et al., 2019) in PyTorch, making it fully differentiable. The model is based on the shallow ice approximation to compute depth-integrated ice velocity (Eq. A3) and

a mass conservation equation, where changes in ice thickness at a point equal the mass balance input minus the divergence of the ice flux (Eq. A4). Since version 1.6.0, OGGM uses a semi-implicit numerical scheme instead of the scheme described in Maussion et al. (2019). Details on this scheme are provided in Appendix A2.

AGILE uses the same 1.5D flowline representation as OGGM. In this approach, glacier dynamics are computed along a flowline with variable surface widths. The flowlines are derived from geographical input data, including glacier outlines and a digital elevation model (DEM), using the 'elevation band flowlines' method (e.g. Huss and Farinotti, 2012; Huss and Hock, 2015; Werder et al., 2019), which preserves the glacier's area-height distribution. Each grid point along the flowline is assigned a trapezoidal bed shape with a 45° wall-slope, and the bottom width depends on the glacier bed height (see Sect. 2.3.1 for more details). The 1.5D flowline setup reduces spatial complexity compared to full 2D or 3D models, making it computationally feasible to track the full computational graph during model runs to be used for AD, without running into memory limitations (see Appendix A1 for details about AD in PyTorch).

To integrate height-dependent Mass Balance (MB) forcing into the computation of gradients, AGILE includes a PyTorch wrapper (see Appendix A3 for details). This accounts for changes in MB forcing caused by dynamically evolving glacier surface heights. The wrapper allows the use of any OGGM-compatible MB model for dynamic simulations without re-implementing it in PyTorch. However, this approach does not support gradient computation for MB model parameters (e.g., the melt factor in temperature index models), but this can be added in the future.

2.3 Cost function

The cost function is defined as

$$\mathcal{J}(\Theta) = \mathcal{J}_{\text{obs}}(\Theta) + \lambda \mathcal{J}_{\text{reg}}(\Theta), \quad (1)$$

where \mathcal{J}_{obs} represents the target observational component (see Sect. 2.3.1), and \mathcal{J}_{reg} is the regularization component (see Sect. 2.3.2). The control variables are denoted with Θ (see Sect. 2.4), and λ determines the weight of the regularization term relative to the observational term.

2.3.1 Target observational cost component

The target observational component, \mathcal{J}_{obs} , measures the alignment of the model with the target observations using squared differences. For a single observed quantity x , it is defined as

$$\mathcal{J}_{\text{obs},x} = \frac{(x_{\text{mdl}} - x_{\text{obs}})^2}{\sigma_x^2}, \quad (2)$$

where x_{mdl} is the modeled quantity, x_{obs} is the corresponding target observation, and σ_x represents the uncertainty of

the target observation. This ensures that discrepancies within the uncertainty bounds have values smaller than one and also scales different types of target observations. For multiple observations, \mathcal{J}_{obs} is averaged over all normalized discrepancies.

$$\mathcal{J}_{\text{obs}} = \frac{1}{n_{\text{obs}}} \sum_j^{n_{\text{obs}}} \mathcal{J}_{\text{obs},j}, \quad (3)$$

where n_{obs} equals the number of considered target observations.

In this study, we focus on target observations that are readily available from global datasets. Specifically, a single area-height distribution, one geodetic mass balance value (ΔM), and one total glacier volume estimate (V). The choice of these variables and the implications for real-world applications were discussed in Sect. 1, these are the currently available global datasets used by the glacier models participating in GlacierMIP3.

The area-height distribution is derived from a glacier outline and a DEM, and is also used to define the elevation-band flowlines (see Sect. 2.2). As a result, we aim to match the surface heights (S) at each grid point along the flowline at the outline year. By doing so, we also match the area-height distribution, since it was used to construct the flowlines. To maintain this relationship during minimization, the bottom width of the trapezoidal bed shape is adjusted at each iteration based on the current bed height estimate (see control variables in Sect. 2.4). This preserves the link between surface height and the corresponding elevation-band area at each grid point.

As an example, the target observational cost for volume is given by

$$\mathcal{J}_{\text{obs},V} = \frac{(V_{\text{mdl}} - V_{\text{obs}})^2}{\sigma_V^2}, \quad (4)$$

while for surface height along the flowline it is

$$\mathcal{J}_{\text{obs},S} = \frac{\sum_{i=1}^{n_x} (S_{\text{mdl}}^i - S_{\text{obs}}^i)^2}{\sigma_S^2 n_x}, \quad (5)$$

where, n_x is the number of grid points and S^i is the surface height at grid point i . Here, the volume represents a glacier-integrated observation, while the surface heights are an observation along the flowline, incorporating the observed area-height distribution. AGILE's flexible cost function design allows for expanding the range of supported target observations as long as corresponding model counterparts exist.

2.3.2 Regularization cost component

While \mathcal{J}_{obs} ensures alignment with target observations, the problem can remain ill-posed or prone to overfitting to observations. Regularization, \mathcal{J}_{reg} , introduces constraints to address these issues, typically focusing on smoothness (e.g.

Goldberg and Heimbach, 2013; Fürst et al., 2017; Jouvét, 2022). The regularization term is scaled to ensure compatibility with \mathcal{J}_{obs} , similar to the observation uncertainties.

In this study, we apply regularization solely to enforce a smooth glacier bed. This is defined as

$$\mathcal{J}_{\text{reg,bed}} = \frac{1}{\gamma_{\text{bed}}} \sum_i^{n_x-1} \left(\frac{b^{i+1} - b^i}{\Delta x} \right)^2, \quad (6)$$

where b^i is the glacier bed height at grid point i , Δx is the grid spacing, and γ_{bed} is the scaling factor based on the smoothness of the first guess bed height b_{fg} ($\gamma_{\text{bed}} = \sum_i^{n_x-1} \left(\frac{b_{\text{fg}}^{i+1} - b_{\text{fg}}^i}{\Delta x} \right)^2$). At the start of the flowline (the highest point), an additional grid point is introduced to preserve the bed slope. Without this extra grid point, the algorithm would lower the bed height at the highest point to match the height of the second-highest point, minimizing the smoothness term. This effect gets stronger with larger values of λ . The extra grid point acts as an additional regularization to counteract this effect, regardless of the value of λ .

For our experiments we defined $\mathcal{J}_{\text{reg}} = \mathcal{J}_{\text{reg,bed}}$. However, this regularization framework could be extended in future studies, for instance, to include smoothness constraints on the initial flux. Such extensions will become increasingly important when working with real-world target observations, which are subject to measurement uncertainties. In these cases, regularization plays a crucial role in preventing the model from overfitting to noisy data.

2.4 Control variables

Control variables, denoted as $\Theta = [\Theta_1, \Theta_2, \dots, \Theta_n]$, are adjusted at each iteration to minimize the cost function (see Sect. 2.3). They represent the unknown variables to be estimated or optimized. In theory, these can include any unknown aspect of the system, such as flowline-geometry variables (e.g. glacier bed height), dynamic parameters (e.g. the deformation-sliding parameter A of Eq. A3), MB model parameters (e.g. melt factor), or a combination of these. The flexibility of AD enables the simultaneous calculation of gradients for multiple control variables, providing essential information to guide the minimization algorithm (see Sect. 2.5). However, this is constrained by observational data availability and the risk of overfitting.

In this study, we set the control variables to be the glacier bed elevation and the distributed ice volume in 1980 at each grid point. This choice reflects our primary goal: to demonstrate, as a proof-of-concept, that variables with different types, meanings, and magnitudes can be optimized simultaneously within each iteration. We also aim to explore whether AGILE can (i) improve upon the current OGGM bed inversion, which assumes equilibrium and may introduce biases for non-steady-state glaciers, and (ii) reconstruct glacier conditions in 1980. The second objective is of secondary im-

portance and mainly serves to enable transient simulations from 1980 to 2020. This allows us to incorporate target observations at their respective timestamps, such as the area-height distribution and the total volume both in 2000, as well as the geodetic mass balance from 2000 to 2020. The goal is to obtain a dynamically consistent glacier state in 2020, which is not part of the optimization but is important for initializing future projections.

To handle control variables with varying magnitudes, we apply min-max normalization. For a given control variable Θ_x , this scaling is defined as

$$\Theta_{x,\text{scaled}} = \frac{\Theta_x - \Theta_{x,\text{min}}}{\Theta_{x,\text{max}} - \Theta_{x,\text{min}}}, \quad (7)$$

where $\Theta_{x,\text{min}}$ and $\Theta_{x,\text{max}}$ represent the defined minimum and maximum bounds, respectively. This ensures all scaled variables are within the range $[0, 1]$, making changes proposed by the minimization algorithm approximately proportional across all variables. The minimum and maximum values for the bed heights are defined such that the resulting ice thickness at the time of the observed outline and DEM remains within $\pm 60\%$ of the first-guess estimate. Similarly, the 1980 ice volume at each grid point is constrained within $\pm 40\%$ of the first guess. Additionally, ten extra grid points are added beyond the terminus for the ice volume, allowing the glacier to be initialized with a larger extent than observed. For these extra points, the scaling boundaries are set to 0 and the maximum value at the terminus grid point (140% of the first guess volume).

We also must account for the non-uniform grid used in the 1.5D flowline representation (see Sect. 2.2), where grid-cell areas vary along the flowline. These variations influence glacier-wide properties when control variables are changed. For example, a change in the bed height affects the glacier area-height distribution more in a larger grid cell than in a smaller one. To account for this, we multiply the glacier bed height by the initial surface width at each grid point and use the resulting values as our control variables. During each iteration, these control variables are divided by the same initial surface width to recover the estimated glacier bed, which is then used in the dynamic simulation. The initial surface width is set during flowline initialization to preserve the observed area-height distribution and remains constant throughout minimization (see Sect. 2.2). We use width instead of grid-cell area because the spacing along the flowline is the same for all grid cells, making it a constant factor. The 1980 glacier volume at each grid point naturally accounts for varying grid-cell sizes. For the actual control variables, we divide the volume by the constant grid-cell spacing along the flowline and use the resulting volume per unit length as the control. This corresponds to the cross section area at each grid point.

2.5 First guess and minimization

To initiate the algorithm, an initial estimate of the control variables, specifically the glacier bed height and the 1980 volume distribution, is required. In our experiments, we used two methods to generate these estimates, allowing us to assess AGILE’s sensitivity to the first guess. The first method relies on the default OGGM bed inversion, a mass-conserving approach based on an equilibrium assumption and an apparent mass balance, as described in Maussion et al. (2019). The second uses the shear-stress-based GlabTop method (Linsbauer et al., 2012), which depends solely on surface geometry.

In both cases, the estimated bed geometry directly provides the initial values for both the bed height and the 1980 volume distribution. However, it is important to note that the volume estimates from both methods correspond to the outline year (2000), and we introduce a temporal mismatch by using them as the first guess 1980 volume distribution.

The minimization process is conducted using the L-BFGS-B algorithm, as implemented in the Python package SciPy (Virtanen et al., 2020, <https://www.scipy.org/>, last access: 21 January 2026). L-BFGS-B is a limited-memory quasi-Newton optimization algorithm that, given a smooth objective function, its gradient, an initial guess, and simple bound constraints on the control variables, iteratively approximates the inverse Hessian to efficiently find a constrained minimum (Byrd et al., 1995; Zhu et al., 1997; Morales and Nocedal, 2011). Similar algorithms have been used in comparable geophysical studies (e.g., Goldberg and Heimbach, 2013; Fürst et al., 2017). In our case, we set the bounds for the control variables to the minimum and maximum values defined in Sect. 2.4.

3 Experiments

The goal of these experiments is to demonstrate that AGILE can recover both the glacier bed heights along the flowline and the dynamic glacier state in 2020 by performing a transient simulation from 1980 to 2020. A key objective is to show that AGILE improves upon OGGM’s bed inversion, which assumes glacier equilibrium and may introduce errors for retreating or advancing glaciers, while at the same time producing a dynamically consistent glacier state in 2020 that could serve as the initial condition for future projections. To achieve this, we define the control variables as the glacier bed height and the 1980 ice volume at each grid point (see Sect. 2.4).

The target observations are based on globally available datasets, used here in an idealized setting. Specifically, we use an area–height distribution and a total volume estimate for the year 2000, along with a geodetic mass balance between 2000 and 2020. This setup is considered idealized because we exclude uncertainties in both mass balance forc-

ing and ice dynamics. While this simplification does not reflect real-world complexity, it serves as a proof of concept, demonstrating that AGILE is correctly implemented and that gradients obtained with AD can be used to optimize multiple control variables simultaneously.

To create a controlled test environment, we generate synthetic glaciers using OGGM and treat the model output as observations. This gives us complete knowledge of the system, allowing us to directly assess the accuracy of our methodology. This type of setup, often referred to as an “inverse crime” (Colton and Kress, 2013), offers an optimistically favourable testing environment. However, demonstrating robust and reliable performance under such idealized conditions is an essential first step for validating a new inversion method (Goldberg and Heimbach, 2013).

3.1 Creation of synthetic glaciers and measurements

The idealized experiments are designed to replicate realistic glacier geometries. We selected the glaciers Aletsch in the Alps (Europe), Artesonraju in the Cordillera Blanca (South America), Baltoro in the Karakoram (Asia), and Peyto in the Canadian Rockies (North America) because they represent a range of different climates and glacier area size. For each glacier, a flowline is constructed using the RGI v6 outline (RGI 6. Consortium, 2017), DEM data from NASADEM (NASA JPL, 2020), and the consensus ice thickness estimate (Farinotti et al., 2019). Based on these flowlines, we generate dynamic glacier simulations from 1980 to 2020 for three different dynamic states: retreating, advancing, and equilibrium.

In particular, the retreating and advancing cases allow us to evaluate whether AGILE can improve upon OGGM’s bed inversion, which assumes a glacier is in equilibrium. The resulting glacier geometries, shown in Fig. 2 panels a and b for Aletsch, and in Fig. S1 panels a and b for Artesonraju, panels e and f for Baltoro, and panels i and j for Peyto, serve as the ground truth we aim to invert for.

The driving mass balance is defined using a simple degree-day model (see Eq. 1 in Schuster et al., 2023), with precipitation and temperature inputs taken from the W5E5 dataset (Lange, 2019) for each glacier’s location. Precipitation factors (ranging from 1.4 to 5.2), degree-day factors (3.1 to $6.5 \text{ kg m}^{-2} \text{ }^\circ\text{C}^{-1} \text{ d}^{-1}$), and temperature biases (-5.6 to $0.9 \text{ }^\circ\text{C}$) follow the calibration values from OGGM v1.6. The deformation-sliding parameter A (ranging from 0.1 to $7.9 \times 10^{-24} \text{ s}^{-1} \text{ Pa}^{-3}$) is also taken from OGGM v1.6, leading to different dynamic behaviors for each synthetic glacier.

For the creation of advancing and retreating cases, we first performed a 60-year dynamic simulation using climate data from 1920 to 1980, followed by a second simulation from 1980 to 2020. For each glacier, dynamic state, and simulation period, temperature biases were applied to ensure that the total volume changes from 1980 to 2020 were of similar magnitude for both retreating and advancing conditions.

For the equilibrium state, we defined a constant mass balance profile based on the average over specific historical periods (Aletsch: 1988–1998, Arensonraju: 1981–1983, Baltoro: 1981–1993, Peyto: 1948–1966). This profile was applied for 120 years before 1980 to allow the glacier to reach equilibrium, and the same mass balance was then used from 1980 to 2020. Figure 2 panels c and d show the resulting mass balance and volume evolution for the different dynamic states of the Aletsch glacier. For the other glaciers, see Fig. S1 panels c and d for Artesonraju, panels g and h for Baltoro, and panels k and l for Peyto.

During the creation of the synthetic glaciers, we “measure” the target observations used in our experiments. Specifically, we record the surface elevation at each grid point in the year 2000 (S_{2000}) for capturing the area height distribution (for details see Sect. 2.3.1), the total glacier volume in 2000 (V_{2000}), and the geodetic mass balance from 2000 to 2020 ($\Delta M_{2000/2020}$). We define the associated measurement uncertainties to reflect the typical order of magnitude found in reported real-world data, setting them as $\sigma_S = 10$ m (Ueema et al., 2020), $\sigma_{\Delta M} = 100 \text{ kg m}^{-2} \text{ yr}^{-1}$ (Hugonnet et al., 2021) and $\sigma_V = 10\%$ of V_{2000} (Farinotti et al., 2019). These target observations are designed to reflect the type of readily available global datasets and represent the only inputs provided to both the first-guess methods and AGILE in our experiments.

3.2 Performance measurements

To evaluate how well AGILE can reconstruct the glacier bed and the dynamic glacier state in 2020, we track the mean absolute difference (MAD) of key variables throughout the optimization process. Specifically, we compute the MAD of the glacier bed elevation (MAD_BED) and the distributed ice volume in 2020 (MAD_V_2020). Since the simulation runs dynamically from 1980 to 2020 and the distributed volume in 1980 is one of the control variables, we also calculate the MAD of the distributed ice volume in 1980 (MAD_V_1980). Although we do not expect to accurately reconstruct the 1980 glacier state, given the lack of target observations before 2000 and the inherently diffusive nature of glacier dynamics, tracking MAD_V_1980 still offers valuable insight into how strongly diffusion limits the ability to recover past glacier states.

4 Results and Discussion

We begin by evaluating the performance of the two first guess methods in Sect. 4.1, with a particular focus on the glacier bed. As noted in Sect. 2.5, a temporal mismatch is introduced in the first guess of the 1980 volume distribution, which limits the interpretability at this stage.

This is followed by an evaluation of AGILE’s core functionality (Sect. 4.2), beginning with the synthetic Aletsch




Figure 2. Synthetic glacier geometry inspired by Aletsch, shown for retreating, equilibrium, and advancing glacier states. Panel (a) displays the defined glacier bed, ice thickness along the flowline, and area-height distribution for the year 2000. Panel (b) presents the volume distribution along the flowline for the year 2000. Panel (c) illustrates the driving mass balance used during the simulation from 1980 to 2020, and panel (d) shows the corresponding total glacier volume over the same period.

glacier geometry in a retreating dynamic state (Sect. 4.2.1). We then expand the analysis across different glacier geometries and dynamic states (Sect. 4.2.2), and finally examine the recovery of control variables along the flowline (Sect. 4.2.3). Next, we evaluate how different first guess methods affect AGILE’s performance (Sect. 4.3). Finally, we assess the impact of varying cost function settings, including different values of λ and numbers of target observations (Sect. 4.4).

4.1 First guess performance

Using our synthetic glacier measurements, we evaluated the two first guess methods described in Sect. 2.5. At this stage, we focus only on the glacier bed, as we acknowledge the temporal mismatch introduced by using a volume distribution estimate from 2000 as the first guess for the 1980 state (see Sect. 2.5). In later analyses, however, we return to the 1980 volume distribution to assess AGILE’s ability to reconstruct past glacier states prior to the first available observation.

Starting from OGGM’s first guess glacier bed, Fig. 3 shows the best performance in the equilibrium state, with a maximum absolute difference of 39.5 m and a MAD of

1.5 m. This is expected, as OGGM’s inversion assumes the glacier is in equilibrium. In contrast, bed elevations are systematically underestimated in the retreating state and overestimated in the advancing state, especially near the terminus. These states show maximum absolute differences of 56.1 m (retreating) and 70.5 m (advancing), with MAD values of 8.7 and 12.7 m, respectively. This pattern holds across all glacier geometries and reflects previous findings (e.g., Fig. 5, panel d in Maussion et al., 2019), highlighting the importance of assessing performance under varying dynamic conditions.

In contrast, the first guess glacier bed of the GlabTop method shows similar behavior across all dynamic states, as it relies solely on surface geometry and does not incorporate any mass balance forcing. The maximum absolute differences for Aletsch in Fig. 3 are 96.0 m (retreating), 100.9 m (equilibrium), and 162.2 m (advancing). This behavior is consistent across all tested geometries.

For all other glacier geometries and dynamic states the performance metrics of the two first guess methods are summarized in Table S1, using mean absolute differences for bed height (MAD_BED), distributed volume in 1980 (MAD_V_1980), and in 2020 (MAD_V_2020). The MAD_V_2020 metric is computed after running the glacier evolution model from 1980 to 2020 with the prescribed mass balance. The OGGM values listed in the table are later used to normalize performance metrics, allowing for easier assessment of whether AGILE improves upon the OGGM first-guess results.

Across all cases, Table S1 shows that OGGM generally performs as well as or better than GlabTop, with the most accurate results seen in equilibrium states. This trend is particularly pronounced for the Baltoro glacier, where OGGM outperforms GlabTop by nearly an order of magnitude. Consequently, Baltoro represents an especially relevant case for evaluating AGILE’s ability to improve upon poor initial guesses that deviate strongly from the synthetic truth.

4.2 Proof of concept for AGILE functionality

4.2.1 Aletsch retreating

The first experiment showcases the proper functionality of AGILE and illustrates that gradient calculations are working as expected. For this we selected our synthetic Aletsch-retreating case and set λ to 0.01 (see Eq. 1). Further we use all three target observations we took during the glacier creation, as defined in Sect. 3.1 (S_{2000} , V_{2000} and $\Delta M_{2000/2020}$) and our first guess is coming from OGGM. Our control variables consist of the distributed volume in 1980 as well as the bed height at each grid point, which gives in total 120 control variables.

Figure 4 panel a shows the evolution of the cost function over 20 iterations, with the largest decrease occurring in the first two iterations, reducing the cost from 2.3 to 0.2. This demonstrates that the gradient-informed updates to the 120




Figure 3. Difference between the synthetic glacier bed for Aletsch and the glacier bed for the two first guess methods, OGGM and GlabTop (see Sect. 4.1), shown for all three dynamic states: retreating, equilibrium, and advancing.

control variables are working as intended. The correctness of the gradient calculations, along with the simultaneous updating of all control variables, is further highlighted by AGILE requiring only 21 forward model runs for 20 iterations (see Fig. 4 panel c), with most iterations needing just one run to find a smaller cost value. This indicates that the minimization algorithm rarely performs ‘search’ runs, where updates to control variables fail to reduce the cost.

Examining individual cost components (Fig. 4, panels a and b), the largest initial contributor is the mismatch to observed surface heights along the flowline S_{2000} , followed by the volume V_{2000} and the geodetic mass balance $\Delta M_{2000/2020}$ terms. The S_{2000} mismatch significantly decreased after two iterations, becoming smaller than $\Delta M_{2000/2020}$, but later increasing slightly again. This highlights that not all observation mismatches decrease monotonically and may temporarily increase as the total cost continues to decrease. The regularization term, starting at 0.005, remained almost constant throughout the iterations. However, as the total cost decreases, its relative importance grows, eventually becoming the dominant cost component after 13 iterations (Fig. 4, panel b).

Analyzing the performance metrics (MAD_BED, MAD_V_1980, and MAD_V_2020) over iterations (Fig. 4, panel c) confirms that reductions in the cost function led to improved agreement with the synthetic truth. Significant improvements were observed for both the bed height and the distributed volume in 2020 compared to the first guess. While the distributed volume for 1980 initially worsened during the early iterations, it began improving relative to the first guess after the sixth iteration.

This experiment demonstrates AGILE’s ability to adapt 120 control variables in just a few iterations, achieving significant improvements over the OGGM-provided first guess. Additionally, the computational demand is minimal, with each iteration (forward model run and gradient calculation) taking approximately 1.3 s for the Aletsch geometry on a laptop equipped with an 11th Gen Intel(R) Core(TM) i7-1165G7 CPU (2.8 GHz) and 16 GB RAM (see Table S2 for other computing times of other dynamic states and geometries). The gradient calculation, specifically the backward propagation through the computing graph (see Appendix A1), requires only around 0.1 s, or roughly ten percent of the total iteration. This efficiency makes the method highly promising for regional to global-scale applications.

4.2.2 Performance across glacier geometries and dynamic states

To evaluate the performance of AGILE across a range of conditions, we generated 12 synthetic glaciers representing four geometries (Aletsch, Artesonraju, Baltoro, and Peyto), each in three dynamic states: retreating, equilibrium, and advancing, as described in Sect. 3.1. In this section, we further examine the core functionality of the method by assessing its ability to minimize the cost function under these varying configurations. We focus here only on experiments relying on the OGGM first guess.

In the equilibrium experiments, the OGGM first guess performs very well, which is expected given its underlying equilibrium assumption. The only exception is Baltoro, where the first guess for the equilibrium case closely resembles that of the retreating case. As a result, the minimization process also mirrors the retreating scenario.

As an example of typical behavior in the equilibrium case, Fig. 4 panels d to f show the results for the Aletsch geometry starting from the OGGM first guess. The strong performance of the OGGM first guess is evident in the low initial total cost of just 0.18 (panel d), indicating that mismatches for all observations fall within their defined uncertainty ranges, suggesting that further optimization is not strictly necessary. Nonetheless, AGILE is able to refine the solution and improve upon this already strong first guess, as shown by the decreasing values of the performance metrics in panel f.

Beyond the equilibrium case, a consistent pattern in the performance metrics across many experiments is that MAD_BED and MAD_V_2020 improve more than MAD_V_1980, highlighting the limited ability to recover earlier glacier states. For example, Fig. 5, panel c (Artesonraju, advancing) and panel f (Peyto, retreating), both starting from the OGGM first guess, show similar behavior. This highlights the broader challenge of reconstructing past glacier states in the absence of direct observations, regardless of glacier geometry or dynamic state. This limitation stems from the diffusive nature of glacier dynamics and the resulting loss of information over time.




Figure 4. Evolution of the cost function and performance metrics for Aletsch retreating (a to c) and equilibrium (d to f), starting from the OGGM first guess, with a λ value of 0.01. Panels (a) and (d) show the evolution of the cost function over minimization iterations, with individual cost terms represented in different colors. The relative contribution of the cost terms to the total cost is shown in panels (b) and (e). Panels (c) and (f) illustrate the evolution of performance metrics over the same iterations. The numbers in boxes along the x-axis indicate the total number of forward model runs required for each corresponding iteration.

For all glacier geometries and dynamic states, AGILE required 21 to 27 model runs over 20 iterations, demonstrating its ability to efficiently inform the minimization algorithm with accurate gradients. An exception was the retreating Peyto experiment (Fig. 5, panel f), where the number of model runs increase substantially from Iteration 14 onwards.




Figure 5. Same as Fig. 4, for Artesonraju advancing (a, b and c) and Peyto retreating (d, e and f) starting from the OGGM first guess.

This hints that the minimization algorithm has difficulties in further minimizing the cost as probably a local minimum is reached. For real-world applications, AGILE includes an option to limit the number of forward model runs (default: 100) to reduce computational effort. Forward model run times varied between 0.7 and 2.5 s (see Table S2), depending on the maximum glacier velocity and corresponding time-step constraints imposed by the stability criterion (Eq. A19).

4.2.3 Distributed differences along the flowline

Figure 6 shows the differences between the modeled results and the synthetic truth along the flowline across iterations for the Aletsch-retreating and Baltoro-advancing cases. Specifi-

cally, it shows the difference in bed height (panels a and d), distributed volume in 1980 (panels b and e), and distributed volume in 2020 (panels c and f).

For Aletsch, improvements are visible along most of the flowline, except near the highest points (start of the flowline, at 0 distance). At these points, a persistent noisy pattern is observed in DIF_BED, which appears in several of the 12 synthetic experiments. As iterations increase, these noisy patterns either dampen or remain stable in DIF_V_2020, but can sometimes become more pronounced in DIF_V_1980. This behaviour depends on the glacier volume distribution along the flowline (Fig. 2, panel b; Fig. S1, panels b, f, and j). These patterns likely reflect the diffusive nature of glacier dynamics and the inherent limitations in accurately inverting for distributed glacier volume when relying on glacier-integrated observations.

Overall, improvements across most grid points are seen regardless of the dynamic state or first-guess method. One notable exception is the Baltoro advancing case (Fig. 6, panels d to f), where AGILE was unable to correct an overly high glacier bed at the terminus. Despite this local limitation, clear improvements are observed between 5 and 25 km along the flowline. Since a large portion of the glacier’s total volume is located within this section (Fig. S1, panel f), the overall distributed volume is still substantially improved.

4.3 Influence of first guess

This section compares AGILE’s performance when initialized with either the OGGM or GlabTop first guess (see Sect. 2.5). Looking at the Aletsch-retreating case, shown panel a of both Figs. 4 and 7, the initial cost value at iteration 0 is significantly higher for GlabTop (10.1) than for OGGM (2.3), due to larger mismatches with the target observations (S_{2000} , V_{2000} , and $\Delta M_{2000/2020}$). Despite this, AGILE effectively minimizes the cost function, reaching a comparable value to OGGM’s initial cost after three iterations and continuing to decrease thereafter.

Examining the performance metrics (Fig. 7, panel c), we find that at Iteration 0, GlabTop has a slightly better MAD_BED compared to OGGM, while MAD_V_1980 and MAD_V_2020 are approximately twice as large. With further iterations, all three metrics improve. After three iterations, MAD_V_2020 improves enough to outperform the OGGM first guess (value < 1). However, MAD_V_1980 shows only minor improvement after 20 iterations and remains roughly twice as large as the value achieved with OGGM.

To further assess AGILE’s performance under more challenging conditions, we examine the Baltoro-retreating case initialized with the GlabTop first guess (Fig. 7, panels d to f). The poor initial performance is evident from the high total cost of 82.5 (panel d) and large values of the performance metrics, around 10, which means they are approximately ten times higher than those for the OGGM first guess.




Figure 6. Differences between the synthetic truth and AGILE guess after every fifth iteration along the flowline for Aletsch retreating (a to c) and Baltoro advancing (d to f), starting from the OGGM first guess (0 Iteration) with a λ value of 0.01. Panels (a) and (d) show the differences in bed height, panels (b) and (e) shows the differences in volume for 1980, and panels (c) and (f) show the differences in volume for 2020. All differences are displayed for every grid point along the flowline.

Despite this challenging starting point, AGILE is able to substantially improve the solution. After 20 iterations, it reaches performance levels close to those of the OGGM first guess, with values around 1.4 (where 1 indicates equal performance to the OGGM first guess). When AGILE is allowed to continue beyond 20 iterations, it further improves the results, achieving values around 0.5 for both MAD_BED and MAD_V_2020 after 40 iterations (not shown in panel f). However, MAD_V_1980 does not improve further beyond 20 iterations.

These results highlight that starting from a worse first guess, such as GlabTop, can reduce the accuracy of reconstructing the glacier state in 1980. This limitation is partly due to the diffusive nature of glacier dynamics, which leads to a gradual loss of information over time during dynamic simulations. Nonetheless, AGILE demonstrates its ability to improve even a poor first guess, successfully refining the glacier bed and simultaneously initialize the model with the 2020 dynamic glacier state.




Figure 7. Same as Fig. 4, for Aletsch (a, b and c) and Baltoro (d, e and f) retreating using GlabTop first guess.

4.4 Different cost function settings

Since the cost function is central to AGILE, we analyzed here the impact of different cost function settings (λ values and number of provided target observations) for retreating and advancing cases, for individual glacier geometries using the OGGM first guess. Specifically, we explored 30 different values of λ , including 0 (no regularization) and 29 values ranging logarithmically from 10^{-4} to 10^3 . Additionally, we varied the number of provided target observations (S_{2000} , V_{2000} and $\Delta M_{2000/2020}$) across seven configurations: using only one of the three target observations, all combinations of two target observations, and all three target observations together. We evaluated the performance in Fig. 8 by looking

if the first guess of MAD_BED and MAD_V_2020 could be improved after 20 iterations and the range of λ values that result in improved performance.

For Aletsch (Fig. 8, panels a to g), in both the retreating and advancing cases, surface height observations (S_{2000}) had the strongest influence in improving MAD_BED and MAD_V_2020 compared to the OGGM first guess. This is expected, as surface height provides distributed information about the glacier's slope along the flowline, while V_{2000} and $\Delta M_{2000/2020}$ are integrated over the entire glacier. Interestingly, even with only providing V_{2000} , small λ values improved MAD_V_2020 .

Combining two target observations consistently outperformed single target observations, especially when S_{2000} were included. Notably, combining V_{2000} and $\Delta M_{2000/2020}$ produced significant improvements compared to using either target observation alone. Furthermore, using two target observations broadened the range of λ values that yielded improvements over the first guess, especially in the advancing case. Using all three target observations simultaneously did not provide substantial benefits compared to configurations that included two target observations, particularly when S_{2000} was one of them. This suggests that two target observations may already provide sufficient information for the control variables for this particular experiment.

For Artesonraju (Fig. 8, panels h to n), S_{2000} was by far the most important target observation. Even when used alone, it achieved results comparable to using two or all three observations. Combining V_{2000} and $\Delta M_{2000/2020}$ did not outperform using these observations individually.

For Baltoro (Fig. 8, panels o to u), in the retreating case (solid lines) single observations alone were insufficient to improve upon the first guess, with V_{2000} showing the best performance. Combining V_{2000} with either S_{2000} or $\Delta M_{2000/2020}$ resulted in substantial improvements, with the best performance achieved using all three target observations. For the advancing case (dotted lines), S_{2000} alone was sufficient to improve MAD_BED and MAD_V_2020 , with further gains observed when combining target observations.

In the retreating case for Peyto (solid lines in Fig. 8, panels v to z, aa, ab), simultaneous improvement of MAD_BED and MAD_V_2020 was only achieved by using all three target observations. For other configurations, improving one metric often led to no improvement or degradation in the other. In the advancing case (dotted lines), S_{2000} alone improved both metrics, with further improvements observed when combining target observations, though using all three added little additional value.

The importance of specific target observations varies with glacier geometry and dynamic state. While it is challenging to disentangle the roles of geometry and state, increasing the number of target observations consistently improves performance. All cases benefit from using all three target observations, though in some instances, performance with one or two target observations is nearly equivalent.

Furthermore, the results demonstrate that the range of effective λ values is relatively broad, with many retreating and advancing cases performing well for λ values between 10^{-4} and 10^{-1} . We observed small to no improvements when transitioning from no regularization ($\lambda = 0$) to small values of λ . However, as λ increased further, performance worsened. This indicates that overly large regularization weights cause AGILE to prioritize smoothing the bed over matching target observations. It is important to note that in this experiment, we used perfectly accurate target observations. With real, imperfect data, regularization becomes more crucial to prevent overfitting, which explains why $\lambda = 0$ performed well under these idealized conditions.

5 Conclusion and future work

Our experiments show that AGILE can adjust many control variables in just a few iterations, proving that the gradient calculations using AD work well. Adding more target observations makes the inversion more reliable, but in some cases, good results can be achieved with fewer target observations. Reconstructing the glacier volume in 1980 was harder than in 2020 because no direct target observations were used before 2000, and the diffusive nature of glacier dynamics, which leads to a gradual loss of information over time during dynamic simulations.

These results are promising, but AGILE has not yet been tested on real-world data. Its flexibility offers many possibilities, but these need careful exploration. For example, with new glacier bed estimates becoming available (e.g., Cook et al., 2023; van Pelt and Frank, 2025), AGILE could focus on other aspects, like combining the calibration of glacier dynamics and mass balance models. Setting up test cases with plenty of observations for validation will be key to understanding what AGILE can do in practical applications.

In our idealized experiments, we assumed the target observations were perfect, so regularization was less important. In real-world cases, where observations are uncertain, regularization will be critical to avoid overfitting. Finding the right balance for the regularization weight (λ) will be necessary. Techniques like L-curves (see, e.g., Hansen, 1992; Gillet-Chaulet et al., 2012; Recinos et al., 2023; Wolovick et al., 2023), could help in deciding this balance.

The next step for applying AGILE to real-world problems would be to include a differentiable mass balance model. A simple temperature index model (e.g., Marzeion et al., 2012) could be a good starting point. For more complex mass balance models (e.g. PyGEM Rounce et al., 2020) it will be important to check if their equations work with AD. Similarly, adding dynamic processes like calving or debris cover will require compatibility with AGILE's AD framework. Therefore, AD may limit how complex the models can get, which is further restricted by the availability of target observations. On the other hand, AGILE's ability to work with diverse




Figure 8. MAD_BED (orange color) and MAD_V_2020 (blue color), normalized by the OGGM first guess values, for retreating (dot and solid line) and advancing (cross and dotted line) experiments after 20 iterations, starting from the OGGM first guess. Results are shown for Aletsch (a to g), Artesonraju (h to n), Baltoro (o to u), and Peyto (v to z, aa, ab). The y-axis shows the number of provided target observations, and the x-axis shows different values of λ ($\lambda = 0$ is shown with a dot or a cross to the left of the axis). The gray shaded area indicates the region where the OGGM first guess could be improved after 20 iterations.

datasets could support more complex models as more target observations become useable in a consistent way.

AGILE provides a promising way to integrate new datasets as they become available or to use all available glacier-specific observations in a consistent way. This could help address the issue of equifinality in global glacier modeling (e.g. Rounce et al., 2020) by combining all available information. Additionally, AGILE could generate consistent glacier histories, filling in data gaps from the past (e.g., creating a reanalysis dataset for glaciers) or providing a solid starting point for future projections.

Appendix A: Model implementations using PyTorch

A1 Automatic Differentiation in PyTorch

PyTorch implements automatic differentiation (AD) using a system called Autograd, which is designed for deep learning optimization tasks. This system enables the construction of a dynamic computation graph and efficient gradient computation. We use this system in our forward model by replacing NumPy arrays with PyTorch tensors.

During a forward pass, PyTorch constructs a dynamic computational graph that keeps track of all operations on tensors requiring gradients, in our case the control variables. For those, all operations involving this tensor are tracked for later differentiation.

After the forward pass, we call PyTorch's `.backward()` function on the cost function tensor. This triggers PyTorch to compute the gradient of that tensor with respect to all tensors requiring gradients, in our case the partial derivatives of our control variables with respect to the cost function. This computation is done by propagating the computational graph backward and applying the chain rule

$$\frac{dy}{dz} = \frac{dy}{dx} \frac{dx}{dz} \quad (\text{A1})$$

for each operation applied during the forward pass. To achieve this, PyTorch stores the derivatives of all standard operations.

PyTorch also allows users to add custom operations by defining both a forward computation and a corresponding backward computation (the derivative of the forward computation). This can be useful for optimizing parts of the code where the derivatives are known. We applied this approach in our semi-implicit solver (see Sect. A2).

A2 Semi-Implicit Solver including AD

With the idea of a global application in mind, AGILE uses the same 1.5D flowline representation as OGGM. In particular, AGILE uses one flowline with changing widths. The bed shape is trapezoidal, with a constant wall angle of 45° . The flowlines are generated from the geographical input data using the “elevation band flowlines” method (e.g. Huss and Farinotti, 2012; Huss and Hock, 2015; Werder et al., 2019).

The glacier evolution model from OGGM (originally introduced by Oerlemans, 1997)

$$\frac{\partial C}{\partial t} = w\dot{m} - \nabla \cdot q \quad (\text{A2})$$

is adapted in AGILE and re-implemented with PyTorch. This advection equation for the cross-section area C (m^2) provides flexibility for varying surface widths w (m) and allows the use of mixed bed shapes with the same equation. Further \dot{m} is the mass balance ($\text{kg m}^{-2} \text{s}^{-1}$), $q = Cu$ is the ice flux ($\text{m}^3 \text{s}^{-1}$) and u is the depth-integrated shallow-ice velocity (m s^{-1}). This velocity is defined as

$$u = \left(\frac{2A}{n+2} h + f_s \frac{1}{h} \right) \left(-\rho g h \frac{\partial s}{\partial x} \right)^n \quad (\text{A3})$$

where A is the ice creep parameter ($\text{s}^{-1} \text{Pa}^{-3}$), n is the exponent of Glen's flow law ($n = 3$), h is the ice thickness (m), f_s is a sliding parameter ($\text{s}^{-1} \text{Pa}^{-3}$), ρ is the ice density (900 kg m^{-3}), g is the gravitational acceleration (9.81 m s^{-2}) and $\frac{\partial s}{\partial x}$ is the surface slope, where s is the surface height (m). By default, the sliding parameter f_s is set to $0 \text{ s}^{-1} \text{Pa}^{-3}$ because there are currently no methods/observations available on a global scale to distinguish the contributions of ice deformation (defined by A) and sliding to the total velocity.

Besides the explicit forward finite difference approximation scheme of OGGM also a semi-implicit scheme is included in AGILE. The reason for this is that numeric instabilities can occur (a known problem, see e.g. <https://oggm.org/2020/01/18/stability-analysis/> and <https://oggm.org/2020/07/08/numerics/>, last access: 21 January 2026) which cause problems when using AD. In particular, during the backward pass for the gradient calculation, these instabilities are amplified and dominate the resulting gradients. The semi-implicit scheme was derived by starting from Eq. (A2) and rearranging it from an advection equation into a diffusion equation

$$\frac{\partial C}{\partial t} = w\dot{m} + \nabla \cdot \left(D \frac{\partial s}{\partial x} \right) \quad (\text{A4})$$

with Diffusivity D

$$D = \left(\frac{2A}{n+2} h + f_s \frac{1}{h} \right) (\rho g h)^n \left| \frac{\partial s}{\partial x} \right|^{n-1} C. \quad (\text{A5})$$

In the following the derivation of the semi-implicit scheme using a rectangular cross-section $C = wh$ is shown. This simplifies the problem because for a rectangular bed shape the surface width is not changing over time ($\frac{\partial C}{\partial t} = \frac{\partial wh}{\partial t} = w \frac{\partial h}{\partial t}$). Afterwards, this solution is generalized for a trapezoidal cross-section. First, we modify Eq. (A4) by using the rectangular cross-section area

$$\frac{\partial h}{\partial t} = \dot{m} + \frac{1}{w} \nabla \cdot \left(D \frac{\partial s}{\partial x} \right) \quad (\text{A6})$$

with Diffusivity D

$$D = \left(\frac{2A}{n+2} h^2 + f_s \right) (\rho g h)^n \left| \frac{\partial s}{\partial x} \right|^{n-1} w. \quad (\text{A7})$$

For the discretization, the ice flux q is defined on a staggered grid denoted with indices $i \pm 1/2$. Further using $(\nabla q)_i = \frac{q_{i+1/2} - q_{i-1/2}}{\Delta x}$, $\left(\frac{\partial s}{\partial x} \right)_{i+1/2} = \frac{s_{i+1} - s_i}{\Delta x}$ and $q^t = D^t \left(\frac{\partial s}{\partial x} \right)^{t+1}$ we get

$$\begin{aligned} \frac{h_i^{t+1} - h_i^t}{\Delta t} &= \dot{m} \\ &+ \frac{1}{w_i} \frac{D_{i+1/2}^t \left(\frac{s_{i+1}^{t+1} - s_i^{t+1}}{\Delta x} \right) - D_{i-1/2}^t \left(\frac{s_i^{t+1} - s_{i-1}^{t+1}}{\Delta x} \right)}{\Delta x}. \end{aligned} \quad (\text{A8})$$

For spatial interpolation of variables from the unstaggered to the staggered grid an arithmetic mean value is used. This is needed in the calculation of D for h and w (e.g. $h_{i+1/2} = \frac{h_i + h_{i+1}}{2}$).

Next, we rearrange the equation to put all terms involving the future time-step $t + 1$ on the left side [TS1](#)

$$\begin{aligned} h_i^{t+1} - \frac{\Delta t}{\Delta x^2 w_i} \\ \cdot \left(D_{i-1/2}^t s_{i-1}^{t+1} - (D_{i+1/2}^t + D_{i-1/2}^t) s_i^{t+1} + D_{i+1/2}^t s_{i+1}^{t+1} \right) \\ = h_i^t + \Delta t \dot{m}. \end{aligned} \quad (\text{A9})$$

and use the definition of the surface height $s = b + h$ (where the glacier bed height b is constant over time) together with a vector notation

$$\begin{aligned} & \left[-\frac{\Delta t}{\Delta x^2 w_i} D_{i-1/2}^t + \frac{\Delta t}{\Delta x^2 w_i} (D_{i+1/2}^t + D_{i-1/2}^t) - \frac{\Delta t}{\Delta x^2 w_i} D_{i+1/2}^t \right] \\ & \cdot \begin{bmatrix} h_{i-1}^{t+1} \\ h_i^{t+1} \\ h_{i+1}^{t+1} \end{bmatrix} \\ & = h_i^t + \Delta t \dot{m} + \frac{\Delta t}{\Delta x^2 w_i} \left[D_{i-1/2}^t - (D_{i+1/2}^t + D_{i-1/2}^t) D_{i+1/2}^t \right] \\ & \cdot \begin{bmatrix} b_{i-1} \\ b_i \\ b_{i+1} \end{bmatrix}. \end{aligned} \quad (\text{A10})$$

Finally, this equation can be used to set up a final linear equation for all grid points. We include the boundary conditions $D_{-1/2} = D_{nx+1/2} = 0$ for all time steps, where nx denotes the last grid point of the unstaggered grid and the unstaggered grid starts with index 0. With this, we define two $nx \times nx$ matrices

$$\begin{aligned} \mathbf{M}_h = & \delta_{i+1,j} \cdot \begin{bmatrix} -\frac{\Delta t}{\Delta x^2 w_0} D_{-1/2} \\ -\frac{\Delta t}{\Delta x^2 w_1} D_{1/2} \\ \vdots \\ -\frac{\Delta t}{\Delta x^2 w_{nx}} D_{nx-1/2} \end{bmatrix} + \\ & + \delta_{i,j} \cdot \begin{bmatrix} 1 + \frac{\Delta t}{\Delta x^2 w_0} (D_{-1/2} + D_{1/2}) \\ 1 + \frac{\Delta t}{\Delta x^2 w_1} (D_{1/2} + D_{3/2}) \\ \vdots \\ 1 + \frac{\Delta t}{\Delta x^2 w_{nx}} (D_{nx-1/2} + D_{nx+1/2}) \end{bmatrix} + \quad (\text{A11}) \\ & + \delta_{i,j+1} \cdot \begin{bmatrix} -\frac{\Delta t}{\Delta x^2 w_0} D_{1/2} \\ -\frac{\Delta t}{\Delta x^2 w_1} D_{3/2} \\ \vdots \\ -\frac{\Delta t}{\Delta x^2 w_{nx}} D_{nx+1/2} \end{bmatrix} \end{aligned}$$

and

$$\begin{aligned} \mathbf{M}_b = & \delta_{i+1,j} \cdot \begin{bmatrix} \frac{\Delta t}{\Delta x^2 w_0} D_{-1/2} \\ \frac{\Delta t}{\Delta x^2 w_1} D_{1/2} \\ \vdots \\ \frac{\Delta t}{\Delta x^2 w_{nx}} D_{nx-1/2} \end{bmatrix} + \\ & + \delta_{i,j} \cdot \begin{bmatrix} -\frac{\Delta t}{\Delta x^2 w_0} (D_{-1/2} + D_{1/2}) \\ -\frac{\Delta t}{\Delta x^2 w_1} (D_{1/2} + D_{3/2}) \\ \vdots \\ -\frac{\Delta t}{\Delta x^2 w_{nx}} (D_{nx-1/2} + D_{nx+1/2}) \end{bmatrix} + \quad (\text{A12}) \\ & + \delta_{i,j+1} \cdot \begin{bmatrix} \frac{\Delta t}{\Delta x^2 w_0} D_{1/2} \\ \frac{\Delta t}{\Delta x^2 w_1} D_{3/2} \\ \vdots \\ \frac{\Delta t}{\Delta x^2 w_{nx}} D_{nx+1/2} \end{bmatrix} \end{aligned}$$

where δ is the Kronecker delta defined as

$$\delta_{i,j} = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases} \quad (\text{A13})$$

and

$$\begin{aligned} \delta_{i+1,j} = & \begin{bmatrix} 0 & 0 & \dots & 0 & 0_{nx} \\ 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0_{nx} \end{bmatrix}, \\ \delta_{i,j+1} = & \begin{bmatrix} 0 & 1 & 0 & \dots & 0_{nx} \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & 0 & \dots & 0_{nx} \end{bmatrix}. \end{aligned}$$

The final linear equation we solve is defined as

$$\mathbf{M}_h \cdot \begin{bmatrix} h_0^{t+1} \\ h_1^{t+1} \\ \vdots \\ h_{nx}^{t+1} \end{bmatrix} = \begin{bmatrix} h_0^t \\ h_1^t \\ \vdots \\ h_{nx}^t \end{bmatrix} + \Delta t \dot{m} + \mathbf{M}_b \cdot \begin{bmatrix} b_0 \\ b_1 \\ \vdots \\ b_{nx} \end{bmatrix} \quad (\text{A14})$$

which can be solved by using the function `scipy.linalg.solve_banded` from the Python package `SciPy` (Virtanen et al., 2020).

This solution for a rectangular cross-section can be generalized to a trapezoidal cross-section by using the definition of the area

$$C = \frac{1}{2} (w + w_0) h, \quad (\text{A15})$$

where w_0 is the bottom width (m) and the surface width w (m) is defined as

$$w = w_0 + \eta h, \quad (\text{A16})$$

where η defines the angle of the side wall (e.g. $\eta = 2$ is a 45° wall angle). Now we can use this definition of the cross-section area in the definition of the diffusivity (Eq. A5)

$$D = \left(\frac{2A}{n+2} h^2 + f_s \right) (\rho g h)^n \left| \frac{\partial s}{\partial x} \right|^{n-1} \frac{w_0 + w}{2}. \quad (\text{A17})$$

Further we can rewrite $\frac{\partial C}{\partial t}$ by inserting the trapezoidal cross-section (Eq. A15) together with the surface width definition (Eq. A16) to get

$$\begin{aligned} \frac{1}{2} \frac{\partial}{\partial t} (h(2w_0 + \eta h)) &= \frac{1}{2} \frac{\partial h}{\partial t} (2w_0 + \eta h) + \frac{1}{2} h \eta \frac{\partial h}{\partial t} \\ &= \frac{1}{2} (2w_0 + 2\eta h) \frac{\partial h}{\partial t} = w \frac{\partial h}{\partial t}. \end{aligned} \quad (\text{A18})$$

With this we are able to rewrite Eq. (A4) again into Eq. (A6) and use our solution derived before. The only difference is

that we need to use the diffusivity of the trapezoidal cross-section defined in Eq. (A17).

For the time-stepping the stability criterion [TS2](#)

$$\Delta t \leq \text{cfl_nr} \frac{\Delta x^2}{\max(D_{i+1/2}/w_{i+1/2})} \quad (\text{A19})$$

is used. This criterion derives from a linearised form of Eq. (A6) (assuming D is constant), which will become a heat equation with diffusivity D/w . Therefore the stability criterion of the heat equation is adapted here. The `cfl_nr` was set to 0.5 following Hindmarsh (2001) equation 91.

To use this semi-implicit solver now in AGILE we further need a differentiable version of the linear equation solver `scipy.linalg.solve_banded`. Otherwise, the linear solve would need too many operations and the computational graph (see Sect. A1 more infos about the computational graph) will become very large and hence the memory consumption when using AD. For this, a new function is defined which uses the original `SciPy` solver during the forward pass, which solves Eq. (A14) for \mathbf{h}^{t+1} by

$$\mathbf{h}^{t+1} = \mathbf{M}_h^{-1} \cdot \mathbf{rhs} \quad (\text{A20})$$

where

$$\mathbf{h}^{t+1} = \begin{bmatrix} h_0^{t+1} \\ h_1^{t+1} \\ \vdots \\ h_{nx}^{t+1} \end{bmatrix} \quad (\text{A21})$$

and

$$\mathbf{rhs} = \begin{bmatrix} h_0^t \\ h_1^t \\ \vdots \\ h_{nx}^t \end{bmatrix} + \Delta t \dot{m} + \mathbf{M}_b \cdot \begin{bmatrix} b_0 \\ b_1 \\ \vdots \\ b_{nx} \end{bmatrix}. \quad (\text{A22})$$

For the backward pass of this newly defined function, the Eqs. (7) and (8) from Goldberg and Heimbach (2013) are used to calculate the adjoints $\delta^* \mathbf{M}_h$ and $\delta^* \mathbf{rhs}$ with [TS3](#)

$$\delta^* \mathbf{rhs} = \mathbf{M}_h^{-T} \cdot \delta^* \mathbf{h}^{t+1} \quad (\text{A23})$$

and

$$\delta^* \mathbf{M}_h = -\delta^* \mathbf{rhs} \cdot (\mathbf{h}^{t+1})^T \odot (\delta_{i,j+1} + \delta_{i,j} + \delta_{i+1,j}), \quad (\text{A24})$$

where \odot is an element-wise matrix multiplication. This matrix multiplication ensures that we only get gradients of the non-zero elements of \mathbf{M}_h . The δ^* notation is used for the adjoints and how they are connected to the gradients is explained in more detail in Goldberg and Heimbach (2013). The implementation of the gradient calculation of this new function was tested against a finite-difference approximation using `torch.autograd.gradcheck` from Pytorch.

A3 Mass-Balance Model wrapper

The idea of the wrapper is to circumvent the need to re-implementing part of the modelling chain with PyTorch, but still incorporate its influence on the gradient calculation. The downside is that with this you can not obtain gradients of variables used in the wrapped part.

In our case, we decided to put the MB model forcing into a wrapper. The MB model gives us for each year a value for the climatic MB depending on the height $\text{mb}_{\text{orig}}(\text{height})$. The MB height profile is defined each year by the temperature and precipitation input.

The idea is to utilize the differentiable 1D interpolation tool from <https://github.com/aliutkus/torchinterp1d> (last access: 21 January 2026). With this, we define constant height bins h_{bins} which cover the vertical glacier extension. The differentiable MB wrapper finally looks like

$$\text{mb}_{\text{torch},i}(h) = \text{torchinterp1d}(h_{\text{bins}}, \text{mb}_{\text{orig},i}(h_{\text{bins}}))(h) \quad (\text{A25})$$

for a year i .

This method works best if the desired modelling part is smooth and does not change too much depending on the input. The spacing of bins (h_{bins} in our case) should be determined by the variability of the output of the wrapped function.

Code and data availability. AGILE [TS4](#) is written in Python and openly available on GitHub (<https://github.com/OGGM/AGILE>, last access: 21 January 2026) under a BSD 3-Clause License. Version 0.1 is also archived on Zenodo with a permanent DOI (<https://doi.org/10.5281/zenodo.17774989>, Schmitt et al., 2025). Under this DOI, all scripts used to run the experiments and generate the figures presented in this study are included in the folder `agile1d/sandbox/paper_v01_code`, along with a README that provides instructions on using a bash script to reproduce the results. In addition, we provide Docker images of the computing environment used, publicly accessible via GitHub Packages (<https://github.com/OGGM/AGILE/pkgs/container/agile>, last access: 21.01.2026). All results presented in this work were produced using the Docker image <https://ghcr.io/oggm/agile:20230525> [TS5](#), archived on Zenodo with a permanent DOI (<https://doi.org/10.5281/zenodo.18314100>, Schmitt et al., 2026).

Supplement. The supplement related to this article is available online at [the link will be implemented upon publication].

Author contributions. P.G. initiated the development of AGILE during his master's thesis, after an idea of F.M. P.S. continued this work in his own master's thesis and subsequently implemented the methods presented in this study. D.G. designed the semi-implicit numeric scheme and P.S. implemented it. P.S. conceptualized the study, designed the idealized experiment setup, and carried out the analysis with continuous input from all co-authors. P.S. wrote the manuscript with contributions from all authors. Large language

models were used to assist with grammar, wording, and style during the preparation of the manuscript.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Geoscientific Model Development*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. P.S. and F.M. received funding from the European Union's Horizon 2020 research and innovation programme (grant agreement no. 101003687) (PROVIDE), from the Austrian Climate Research Programme (ACRP) – 14th call, under grant agreement no. KR21KB0K00001 (HyMELT-CC), and from ESA's "Digital Twin Component for Glaciers" project (4000146160/24/I-KE). D.N.G. acknowledges funding from NERC grant NE/X005194/1. D.N.G. acknowledges funding from NERC grant NE/X005194/1 and NE/T001607/1. The Open Access Publication Fund of the University of Innsbruck contributed to publication costs.

Financial support. This research has been supported by the Horizon 2020 (grant no. 101003687), the European Space Agency (grant no. 4000146160/24/I-KE), the Natural Environment Research Council, British Antarctic Survey (grant nos. NE/X005194/1 and NE/T001607/1), and the Austrian Climate Research Programme (grant no. KR21KB0K00001). ^{TS6}

Review statement. This paper was edited by Ludovic Räss and reviewed by two anonymous referees.

References

- Aguayo, R., Maussion, F., Schuster, L., Schaefer, M., Caro, A., Schmitt, P., Mackay, J., Ultee, L., Leon-Muñoz, J., and Aguayo, M.: Unravelling the sources of uncertainty in glacier runoff projections in the Patagonian Andes (40–56° S), *The Cryosphere*, 18, 5383–5406, <https://doi.org/10.5194/tc-18-5383-2024>, 2024.
- Bosson, J. B., Huss, M., Cauvy-Fraunié, S., Clément, J. C., Costes, G., Fischer, M., Poulenard, J., and Arthaud, F.: Future emergence of new ecosystems caused by glacial retreat, *Nature*, 620, 562–569, <https://doi.org/10.1038/s41586-023-06302-2>, 2023.
- Brinkerhoff, D. J. and Johnson, J. V.: Data assimilation and prognostic whole ice sheet modelling with the variationally derived, higher order, open source, and fully parallel ice sheet model Var-
- GlaS, *The Cryosphere*, 7, 1161–1184, <https://doi.org/10.5194/tc-7-1161-2013>, 2013.
- Byrd, R. H., Lu, P., Nocedal, J., and Zhu, C.: A Limited Memory Algorithm for Bound Constrained Optimization, *SIAM Journal on Scientific Computing*, 16, 1190–1208, <https://doi.org/10.1137/0916069>, 1995.
- Cannone, N., Diolaiuti, G., Guglielmin, M., and Smiraglia, C.: Accelerating Climate Change Impacts on Alpine Glacier Forefield Ecosystems in the European Alps, *Ecological Applications*, 18, 637–648, <https://doi.org/10.1890/07-1188.1>, 2008.
- Colton, D. and Kress, R.: *Inverse Acoustic and Electromagnetic Scattering Theory*, Springer New York, <https://doi.org/10.1007/978-1-4614-4942-3>, 2013.
- Cook, S. J., Jouvet, G., Millan, R., Rabatel, A., Zekollari, H., and Dussaillant, I.: Committed Ice Loss in the European Alps Until 2050 Using a Deep-Learning-Aided 3D Ice-Flow Model With Data Assimilation, *Geophysical Research Letters*, 50, <https://doi.org/10.1029/2023gl105029>, 2023.
- COP-DEM: Copernicus DEM, CDSE [data set], <https://doi.org/10.5270/esa-c5d3d65>, 2022.
- Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nature Geoscience*, 12, 168–173, <https://doi.org/10.1038/s41561-019-0300-3>, 2019.
- Fürst, J. J., Gillet-Chaulet, F., Benham, T. J., Dowdeswell, J. A., Grabiec, M., Navarro, F., Pettersson, R., Moholdt, G., Nuth, C., Sass, B., Aas, K., Fettweis, X., Lang, C., Seehaus, T., and Braun, M.: Application of a two-step approach for mapping ice thickness to various glacier types on Svalbard, *The Cryosphere*, 11, 2003–2032, <https://doi.org/10.5194/tc-11-2003-2017>, 2017.
- Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, <https://doi.org/10.5194/tc-6-1561-2012>, 2012.
- Gobbi, M., Ambrosini, R., Casarotto, C., Diolaiuti, G., Ficetola, G. F., Lencioni, V., Seppi, R., Smiraglia, C., Tampucci, D., Valle, B., and Caccianiga, M.: Vanishing permanent glaciers: climate change is threatening a European Union habitat (Code 8340) and its poorly known biodiversity, *Biodiversity and Conservation*, 30, 2267–2276, <https://doi.org/10.1007/s10531-021-02185-9>, 2021.
- Goldberg, D. N. and Heimbach, P.: Parameter and state estimation with a time-dependent adjoint marine ice sheet model, *The Cryosphere*, 7, 1659–1678, <https://doi.org/10.5194/tc-7-1659-2013>, 2013.
- Hansen, P. C.: Analysis of Discrete Ill-Posed Problems by Means of the L-Curve, *SIAM Review*, 34, 561–580, <http://www.jstor.org/stable/2132628> (last access: 21 January 2026), 1992.
- Hartl, L., Schmitt, P., Schuster, L., Helfricht, K., Abermann, J., and Maussion, F.: Recent observations and glacier modeling point towards near-complete glacier loss in western Austria (Ötztal and Stubai mountain range) if 1.5 °C is not met, *The Cryosphere*, 19, 1431–1452, <https://doi.org/10.5194/tc-19-1431-2025>, 2025.
- Hindmarsh, R. C. A.: *Notes on Basic Glaciological Computational Methods and Algorithms*, Springer Berlin Heidelberg, Berlin, Heidelberg, ISBN 978-3-662-04439-1, 222–249, https://doi.org/10.1007/978-3-662-04439-1_13, 2001.

- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., Brun, F., and Käab, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Huss, M. and Farinotti, D.: Distributed ice thickness and volume of all glaciers around the globe, *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2012jf002523>, 2012.
- Huss, M. and Hock, R.: A new model for global glacier change and sea-level rise, *Frontiers in Earth Science*, 3, <https://doi.org/10.3389/feart.2015.00054>, 2015.
- Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8, 135–140, <https://doi.org/10.1038/s41558-017-0049-x>, 2018.
- Jouvet, G.: Inversion of a Stokes glacier flow model emulated by deep learning, *Journal of Glaciology*, 69, 13–26, <https://doi.org/10.1017/jog.2022.41>, 2022.
- Lange, S.: WFDE5 over land merged with ERA5 over the ocean (W5E5), GFZ Data Services [data set], <https://doi.org/10.5880/PIK.2019.023>, 2019.
- Linsbauer, A., Paul, F., and Haerberli, W.: Modeling glacier thickness distribution and bed topography over entire mountain ranges with GlabTop: Application of a fast and robust approach, *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011jf002313>, 2012.
- Lipscomb, W. H., Price, S. F., Hoffman, M. J., Leguy, G. R., Bennett, A. R., Bradley, S. L., Evans, K. J., Fyke, J. G., Kennedy, J. H., Perego, M., Ranken, D. M., Sacks, W. J., Salinger, A. G., Vargo, L. J., and Worley, P. H.: Description and evaluation of the Community Ice Sheet Model (CISM) v2.1, *Geosci. Model Dev.*, 12, 387–424, <https://doi.org/10.5194/gmd-12-387-2019>, 2019.
- Lorenc, A. C.: Development of an Operational Variational Assimilation Scheme (gtSpecial Issue: Data Assimilation in Meteorology and Oceanography: Theory and Practice), *Journal of the Meteorological Society of Japan. Ser. II*, 75, 339–346, https://doi.org/10.2151/jmsj1965.75.1b_339, 1997.
- Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers, *The Cryosphere*, 6, 1295–1322, <https://doi.org/10.5194/tc-6-1295-2012>, 2012.
- Marzeion, B., Hock, R., Anderson, B., Bliss, A., Champollion, N., Fujita, K., Huss, M., Immerzeel, W. W., Kraaijenbrink, P., Malles, J., Maussion, F., Radić, V., Rounce, D. R., Sakai, A., Shannon, S., van de Wal, R., and Zekollari, H.: Partitioning the Uncertainty of Ensemble Projections of Global Glacier Mass Change, *Earth's Future*, 8, <https://doi.org/10.1029/2019ef001470>, 2020.
- Maussion, F., Butenko, A., Champollion, N., Dusch, M., Eis, J., Fourteau, K., Gregor, P., Jarosch, A. H., Landmann, J., Oesterle, F., Recinos, B., Rothenpieler, T., Vlug, A., Wild, C. T., and Marzeion, B.: The Open Global Glacier Model (OGGM) v1.1, *Geosci. Model Dev.*, 12, 909–931, <https://doi.org/10.5194/gmd-12-909-2019>, 2019.
- Millan, R., Mouginot, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world's glaciers, *Nature Geoscience*, 15, 124–129, <https://doi.org/10.1038/s41561-021-00885-z>, 2022.
- Morales, J. L. and Nosedal, J.: Remark on “algorithm 778: L-BFGS-B: Fortran subroutines for large-scale bound constrained optimization”, *ACM Transactions on Mathematical Software*, 38, 1–4, <https://doi.org/10.1145/2049662.2049669>, 2011.
- NASA JPL: NASADEM Merged DEM Global 1 arc second V001, NASA Land Processes Distributed Active Archive Center [data set], https://doi.org/10.5067/MEASURES/NASADEM/NASADEM_HGT.001, 2020.
- Oerlemans, J.: A flowline model for Nigardsbreen, Norway: projection of future glacier length based on dynamic calibration with the historic record, *Annals of Glaciology*, 24, 382–389, <https://doi.org/10.3189/s0260305500012489>, 1997.
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein, N., Antiga, L., Desmaison, A., Kopf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., Bai, J., and Chintala, S.: PyTorch: An Imperative Style, High-Performance Deep Learning Library, in: *Advances in Neural Information Processing Systems* 32, edited by: Wallach, H., Larochelle, H., Beygelzimer, A., d’Alché-Buc, F., Fox, E., and Garnett, R., Curran Associates, Inc., 8024–8035, <http://papers.neurips.cc/paper/9015-pytorch-an-imperative-style-high-performance-deep-learning-library.pdf> (last access: 21 January 2026), 2019.
- Recinos, B., Goldberg, D., Maddison, J. R., and Todd, J.: A framework for time-dependent ice sheet uncertainty quantification, applied to three West Antarctic ice streams, *The Cryosphere*, 17, 4241–4266, <https://doi.org/10.5194/tc-17-4241-2023>, 2023.
- RGI 6 Consortium: A Dataset of Global Glacier Outlines: Version 6.0: Technical Report, Global Land Ice Measurements from Space, <http://www.glims.org/RGI/randolph60.html> (last access: 21 January 2026), 2017.
- RGI 7 Consortium: Randolph Glacier Inventory – A Dataset of Global Glacier Outlines, Version 7, National Snow and Ice Data Center [data set], <https://doi.org/10.5067/F6JMOVY5NAVZ>, 2023.
- Rounce, D. R., Khurana, T., Short, M. B., Hock, R., Shean, D. E., and Brinkerhoff, D. J.: Quantifying parameter uncertainty in a large-scale glacier evolution model using Bayesian inference: application to High Mountain Asia, *Journal of Glaciology*, 66, 175–187, <https://doi.org/10.1017/jog.2019.91>, 2020.
- Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., Farinotti, D., Menounos, B., and McNabb, R. W.: Global glacier change in the 21st century: Every increase in temperature matters, *Science*, 379, 78–83, <https://doi.org/10.1126/science.abo1324>, 2023.
- Schmitt, P., Gregor, P., and Maussion, F.: OGGM/AGILE: AGILE v0.1.2, Zenodo [code], <https://doi.org/10.5281/zenodo.17774989>, 2025.
- Schmitt, P., Gregor, P., and Maussion, F.: OGGM/AGILE: Container build 20230525 used in Publication, Zenodo [code], <https://doi.org/10.5281/zenodo.18314100>, 2026.
- Schuster, L., Rounce, D. R., and Maussion, F.: Glacier projections sensitivity to temperature-index model choices and calibration strategies, *Annals of Glaciology*, 64, <https://doi.org/10.1017/aog.2023.57>, 2023.
- Shean, D. E., Bhushan, S., Montesano, P., Rounce, D. R., Arendt, A., and Osmanoglu, B.: A Systematic, Regional Assessment of High Mountain Asia Glacier Mass Balance, *Frontiers in Earth Science*, 7, <https://doi.org/10.3389/feart.2019.00363>, 2020.

- Slangen, A. B., Palmer, M. D., Camargo, C. M., Church, J. A., Edwards, T. L., Hermans, T. H., Hewitt, H. T., Garner, G. G., Gregory, J. M., Kopp, R. E., Santos, V. M., and van de Wal, R. S.: The evolution of 21st century sea-level projections from IPCC AR5 to AR6 and beyond, *Cambridge Prisms: Coastal Futures*, 1, <https://doi.org/10.1017/cft.2022.8>, 2022.
- The GlaMBIE Team: Community estimate of global glacier mass changes from 2000 to 2023, *Nature*, 639, 382–388, <https://doi.org/10.1038/s41586-024-08545-z>, 2025.
- Ultee, L., Coats, S., and Mackay, J.: Glacial runoff buffers droughts through the 21st century, *Earth Syst. Dynam.*, 13, 935–959, <https://doi.org/10.5194/esd-13-935-2022>, 2022.
- Uuemaa, E., Ahi, S., Montibeller, B., Muru, M., and Kmoch, A.: Vertical Accuracy of Freely Available Global Digital Elevation Models (ASTER, AW3D30, MERIT, TanDEM-X, SRTM, and NASADEM), *Remote Sensing*, 12, 3482, <https://doi.org/10.3390/rs12213482>, 2020.
- van Pelt, W. and Frank, T.: New glacier thickness and bed topography maps for Svalbard, *The Cryosphere*, 19, 1–17, <https://doi.org/10.5194/tc-19-1-2025>, 2025.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., and SciPy 1.0 Contributors: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Welty, E., Zemp, M., Navarro, F., Huss, M., Fürst, J. J., Gärtner-Roer, I., Landmann, J., Machguth, H., Naegeli, K., Andreassen, L. M., Farinotti, D., Li, H., and GlaThiDa Contributors: Worldwide version-controlled database of glacier thickness observations, *Earth Syst. Sci. Data*, 12, 3039–3055, <https://doi.org/10.5194/essd-12-3039-2020>, 2020.
- Werder, M. A., Huss, M., Paul, F., Dehecq, A., and Farinotti, D.: A Bayesian ice thickness estimation model for large-scale applications, *Journal of Glaciology*, 66, 137–152, <https://doi.org/10.1017/jog.2019.93>, 2019.
- Wimberly, F., Ultee, L., Schuster, L., Huss, M., Rounce, D. R., Maussion, F., Coats, S., Mackay, J., and Holmgren, E.: Inter-model differences in 21st century glacier runoff for the world's major river basins, *The Cryosphere*, 19, 1491–1511, <https://doi.org/10.5194/tc-19-1491-2025>, 2025.
- Wolovick, M., Humbert, A., Kleiner, T., and Rückamp, M.: Regularization and L-curves in ice sheet inverse models: a case study in the Filchner–Ronne catchment, *The Cryosphere*, 17, 5027–5060, <https://doi.org/10.5194/tc-17-5027-2023>, 2023.
- Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble, *The Cryosphere*, 13, 1125–1146, <https://doi.org/10.5194/tc-13-1125-2019>, 2019.
- Zekollari, H., Huss, M., Schuster, L., Maussion, F., Rounce, D. R., Aguayo, R., Champollion, N., Compagno, L., Hugonnet, R., Marzeion, B., Mojtabavi, S., and Farinotti, D.: Twenty-first century global glacier evolution under CMIP6 scenarios and the role of glacier-specific observations, *The Cryosphere*, 18, 5045–5066, <https://doi.org/10.5194/tc-18-5045-2024>, 2024.
- Zekollari, H., Schuster, L., Maussion, F., Hock, R., Marzeion, B., Rounce, D. R., Compagno, L., Fujita, K., Huss, M., James, M., Kraaijenbrink, P. D. A., Lipscomb, W. H., Minallah, S., Oberrauch, M., Van Tricht, L., Champollion, N., Edwards, T., Farinotti, D., Immerzeel, W., Leguy, G., and Sakai, A.: Glacier preservation doubled by limiting warming to 1.5 °C versus 2.7 °C, *Science*, 388, 979–983, <https://doi.org/10.1126/science.adu4675>, 2025.
- Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Hoelzle, M., Paul, F., Haerberli, W., Denzinger, F., Ahlstrøm, A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R., Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Ove Hagen, J., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V., Severskiy, I., Sigurdsson, O., Soruco, A., Usabaliyev, R., and Vincent, C.: Historically unprecedented global glacier decline in the early 21st century, *Journal of Glaciology*, 61, 745–762, <https://doi.org/10.3189/2015jog15j017>, 2015.
- Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568, 382–386, <https://doi.org/10.1038/s41586-019-1071-0>, 2019.
- Zhu, C., Byrd, R. H., Lu, P., and Nocedal, J.: Algorithm 778: L-BFGS-B, *ACM Transactions on Mathematical Software*, 23, 550–560, <https://doi.org/10.1145/279232.279236>, 1997.

Remarks from the typesetter

TS1 Please confirm the equation.

TS2 Please give an explanation of why the equation needs to be changed. We have to ask the handling editor for approval.
Thanks.

TS3 Please confirm the equation.

TS4 Please confirm adjusted section header and add data set information, since there are only codes currently.

TS5 Please confirm URL.

TS6 Please confirm both Acknowledgements and Financial support sections, and make sure that the funding information is identical in both sections. Please also confirm updated funding information in the MS records.

TS7 Should the name remain as is then?