

# Mineral-bound organic carbon exposed by hillslope thermokarst terrain: case study in Cape Bounty, Canadian High Arctic

Maxime Thomas<sup>1</sup>, Julien Fouché<sup>2</sup>, Hugues Titeux<sup>1</sup>, Charlotte Morelle<sup>1</sup>, Nathan Bemelmans<sup>1</sup>,  
Melissa J. Lafrenière<sup>3</sup>, Joanne K. Heslop<sup>3,4</sup>, Sophie Opfergelt<sup>1</sup>

5 <sup>1</sup>Earth and Life Institute, Université catholique de Louvain, Louvain-la-Neuve, Belgium

<sup>2</sup>LISAH, Univ Montpellier, INRAE, IRD, Institut Agro, Montpellier, France

<sup>3</sup>Department of Geography and Planning, Queen's University, Kingston, ON, K7L 3N6, Canada

<sup>4</sup>now at: The Alaska Earthquake Center, University of Alaska, Fairbanks, Alaska, USA

*Correspondence to:* Maxime Thomas (maxime.thomas@uclouvain.be)

10 **Abstract.** Arctic landscapes could add 55 - 230 Pg of carbon (in CO<sub>2</sub> equivalent) to the atmosphere, through CO<sub>2</sub> and CH<sub>4</sub>  
emissions, by the end of this century. These estimates could be quantified more accurately by constraining the contribution of  
rapid thawing processes such as thermokarst landscapes to permafrost carbon loss, and by investigating the exposed organic  
carbon (OC) interacting with mineral surfaces or metallic cations, i.e., the nature of these interactions and what controls their  
relative abundance. Here, we investigate two contrasted types of hillslope thermokarst landscapes: an Active Layer  
15 Detachment (ALD) which is a one-time event, and a Retrogressive Thaw Slump (RTS) which repeats annually during summer  
months in the Cape Bounty Arctic Watershed Observatory (Melville Island, Canada). We analyzed mineralogy, total and  
soluble element concentrations, total OC and mineral-OC interactions within the headwalls of both disturbances, and within  
corresponding undisturbed profiles. Our results show that **OC-small fragments of biopolymers** stabilized by chemical bonds  
account for 13 ± 5% of total OC in the form of organo-metallic complexes and up to 6 ± 2% associated with poorly crystalline  
20 iron oxides. If we add the mechanisms of physical protection of particulate organic matter in aggregates and larger molecules  
stabilized by chemical bonds, we reach 64 ± 10% of the total OC being stabilized. Importantly, we observe a decrease in the  
proportion of mineral-bound OC in the deeper layers exposed by the retrogressive thaw slump: the proportion of organo-  
metallic complexes drops from ~18% in surface samples to ~1 % in the deepest samples. These results therefore suggest that  
the OC exposed by thermokarst disturbances at Cape Bounty is protected by interactions with minerals to a certain extent, but  
25 that deep thaw features could expose OC more readily accessible to degradation.


## 1 Introduction

In the Arctic, recent studies show that air temperatures are rising 3 to 4 times faster than the global average (AMAP, 2021;  
Rantanen et al., 2022). This polar amplification exposes permafrost – soil, sediment or rock frozen for at least two years – to  
thawing. The latest IPCC predictions therefore forecast a decrease in near-surface permafrost area of 2-66% for IPCC scenario  
30 RCP2.6 and 30-99% for RCP8.5, by 2100 (Fox-Kemper et al., 2021; Meredith et al., 2019). This degradation could induce a

permafrost-carbon-climate feedback (e.g., Abbott and Jones, 2015; Koven et al., 2011; McGuire et al., 2018; Schuur et al., 2015), since northern circumpolar permafrost contains ~1000 Pg of carbon in the first three meters, roughly half the soil carbon present in all other terrestrial biomes (Hugelius et al., 2014; Strauss et al., 2021, 2024). It is estimated that CO<sub>2</sub> and CH<sub>4</sub> emissions from the Arctic could add 55 - 230 Pg of carbon (in CO<sub>2</sub> equivalent) to the atmosphere by the end of this century - a level comparable to emissions of industrialized countries - and must therefore be considered in climate models (Schuur et al., 2022). These assessments are based on simulated volumes of OC that will be exposed by the increase in thickness of the seasonally thawing active layer (i.e., gradual thawing). In order to refine these estimates, rapid thawing processes such as thermokarst, thermo-denudation and thermo-erosion – which could represent an additional release of greenhouse gases – should benefit from further studies (e.g., Natali et al., 2021).

Thermokarst landforms (e.g., Heginbottom et al., 2012; Kokelj and Jorgenson, 2013) are punctual events, both in space and time, and occur when the ground collapses due to the melting of the excess ice in the soil or sediment. These physical disturbances of permafrost are sometimes referred to as ‘abrupt thaw’. In particular, hillslope thermokarst landscapes occur where physical degradation leads to slumps (Fig. 1a), active layer detachments (Fig. 1b) and gullies. These events potentially expose deep and ancient materials to thawing, including OC that would never have re-entered the modern carbon cycle if these disturbances had not occurred. The only existing simulations of thermokarst terrain impacts on permafrost carbon balance indicate that these could provide a similar climate feedback as gradual thaw emissions (Turetsky et al., 2020). Active layer detachments (ALD), are landslides in which the active layer slides down onto the permafrost table over tens of meters (e.g., Lamoureux and Lafrenière, 2009). They thus expose a vertical headwall of the height of the active layer at the location where it has detached. This type of event is punctual in time and typically triggered by deep thaw combined with heavy precipitation.

Retrogressive thaw slumps (RTS) also consist in the detachment of materials, but generally expose greater depths, where permafrost is part of the exported mass. RTSs expand through the years as frozen deposits thaw and ground ice melts at a headwall. The retreat of the head scarp each summer can reach tens of meters per year (e.g., Brooker et al., 2014; Günther et al., 2015; Kokelj et al., 2021; Kunitsky et al., 2013; Lacelle et al., 2015; Leibman et al., 2021; Séjourné et al., 2015; Vadakkedath et al., 2020; van der Sluijs et al., 2023, 2018). These structures therefore expose and move large volumes of material (Kokelj et al., 2015, 2021; Shakil et al., 2020; van der Sluijs et al., 2018; Tanski et al., 2017), involving previously perennially frozen masses of OC, until they stabilize. The area susceptible to be affected by hillslope thermokarst landforms is indeed projected to increase by 250,000 km<sup>2</sup> by the end of the 21st century (Turetsky et al., 2020). The soils and sediments exposed by ALDs and RTSs are generally diverse-varied depending on their headwall heights, lithology and historical permafrost thaw dynamics, and contain organic matter of variable composition and interactions with minerals (e.g., Jongejans et al., 2022; Thomas et al., 2024, 2023).



**Figure 1: Conceptual model of a retrogressive thaw slump (a) and an active layer detachment (b).** Panel (a) modified from (Thomas et al., (2023)).

65 In soils and sediments, organic matter as a whole can be separated into fractions differing in their interactions with minerals and resistance to decay (i.e., stabilization). The free particulate organic matter, composed of plant fragments non associated with minerals, is considered as prone to microbial degradation (labile) displaying a high turnover (Cotrufo and Lavellee, 2022). The fraction of organic matter interacting with mineral surfaces or metallic cations is considered as stabilized, being protected against microbial processing and therefore more persistent (Lavellee et al., 2020). Although different stabilization mechanisms

70 often act simultaneously, they can be classified into different categories, providing very different levels of stabilization (Fig. 12a): (i) physical protection of OC trapped in the aggregates, with no chemical binding to mineral surfaces. This enables stabilization over a period of 10 to ~100 years and depends on the size of the aggregate (von Lützow et al., 2006); (ii) the formation of cationic bridges between negatively charged clay mineral surfaces (e.g., smectite) and negatively charged organic molecules. These are electrostatic interactions between the organic molecule, the cation and the clay, which allows reversible

75 binding; (iii) the formation of cationic bridges between organic molecules of various sizes. These are organo-metallic complexes, where metals (i.e.,  $\text{Fe}^{3+/2+}$ ,  $\text{Al}^{3+}$ ,  $\text{Mn}^{3+/2+}$  or  $\text{Ca}^{2+}$ ) act as polyvalent cations and coordinate to the functional groups of organic acids through electrostatic interactions or coordinate covalent bond (e.g., Boudot et al., 1989; Courchesne and Turmel, 2008; von Lützow et al., 2006). This complexation leads to the coagulation of soluble molecules into insoluble “pseudo-polymers”; (iv) the adsorption of OC onto mineral surfaces such as poorly crystalline iron oxides, via the


80 attachment – by ligand exchange – of organic molecules to metal atoms (Al, Fe) structurally present in the minerals (Kleber et al., 2015). Such poorly-crystalline surfaces are among the most effective sorbents for dissolved OC (e.g., Kaiser et al., 1997). Overall, the turnover time of the mineral- interacting pool of OC ranges from decades to millennia (Kleber et al., 2015 and

references therein), although a substantial portion of mineral-interacting OC may cycle at relatively fast timescales (minutes to years); (Jilling et al., 2025). Quantifying the share of different mechanisms for OC stability requires a combination of physical methods, which enable to isolate mineral-associated OC from the most labile OC pool, and chemical extraction methods which are designed to target specific pools such as complexes or OC associated with poorly-crystalline surfaces (Fig. 1b2b).

The nature, quantities and environmental factors controlling the different organo-mineral interactions in sediments exposed by hillslope thermokarst features are still poorly understood (Opfergelt, 2020). Recent studies show that the proportion of stabilized OC in permafrost sediments is sometimes minor (i.e., averaging 15%; Patzner et al., 2020; Salvadó et al., 2015) or, on the contrary, very significant (i.e., reaching up to 90%; Dutta et al., 2006; Martens et al., 2023; Monhonval et al., 2022). Besides, approaches to characterizing mineral-OC interactions using physical and chemical methods are often applied separately. A comprehensive understanding of the different mechanisms is, however, necessary for realistic assessments of the fate of OC exposed by these features, particularly from RTSs and ALDs, which expose potentially heterogeneous sediments and mobilize materials from varying depths.

In this study, we investigate the nature and controls of different mechanisms of mineral-OC interactions in soils exposed by two contrasted types of hillslope thermokarst landscapes and within profiles of corresponding undisturbed sites. We hypothesize that organo-mineral interactions may be more prominent in surface layers that are subject to recurrent positive temperatures, but that it may be limited at depth if buried sediments have not experienced pedological development~~will depend on historical permafrost thaw dynamics~~. We analyzed mineralogy, total and soluble element concentrations, total OC and mineral-OC interactions within the headwalls of two disturbances: an ALD and a RTS at Cape Bounty (south-central coast of the Melville Island, Nunavut, Canada). We investigate whether these contrasted disturbances expose mineral-interacting OC of different natures (i.e., OC extracted by physical or chemical techniques), estimate the level of stabilization of the different pools and discuss their potential fate.

105



**Figure 21:** Organic carbon (OC) pools with their respective target protocol of extraction and associated stabilization mechanisms; (a) Mechanisms of stabilization of organic matter by mineral surfaces or elements in soils or sediments, i.e. free particulate organic matter; physical protection of organic matter trapped in the aggregates – no chemical link; formation of cation bridges between organic matter and negatively-charged mineral surfaces (such as smectites) – electrostatic interactions; formation of cation bridges between molecules of organic matter of various sizes – electrostatic interactions or coordinate covalent bond; binding of organic matter to metal ions (Al and Fe) contained in mineral surfaces - coordinate covalent bond; (b) Physical and chemical extraction methods and associated target of OC pool;

## 2 Materials & Methods


### 115 2.1 Study site

The Cape Bounty Arctic Watershed Observatory (CBAWO), located on the south-central coast of the Melville Island (Nunavut, Canada; **Fig. 32a-b**), was established in 2003 to examine Arctic ecosystem processes that would be impacted by climate warming and permafrost degradation (Lamoureux and Lafrenière, 2018; Treitz et al., 2024). The Melville Island is characterized by a polar desert climate (mean annual air temperature:  $-14.8 \pm 1.3$  °C) with limited annual precipitation and runoff ( $< 150$  mm year<sup>-1</sup>; Beel et al., 2020, 2018; Lamoureux and Lafrenière, 2018) and a short thaw season between June and August (Lamoureux and Lafrenière, 2009). Bedrock in this region is composed of upper Devonian near-shore marine sandstone and siltstone units (Hodgson et al., 1984). Glacial and early Holocene marine sediments later overlaid the region resulting in a landscape characterized by incised low elevation plateaus and gentle hills (Hodgson et al., 1984). Permafrost is continuous and the thickness of the active layer is most often confined to the 0.75-1 meter of surface material (Lamoureux and Lafrenière, 2009, 2018; Rudy et al., 2013). Exceptionally warm temperatures and clear skies during July 2007 combined with a major

rainfall event (5-year return period), caused extensive thawing of ground ice and degradation of permafrost in Cape Bounty, in the form of hillslope thermokarst structures (Lamoureux and Lafrenière, 2009). This degradation manifested locally as RTSs and ALDs.

## 2.2 Sampling

130 The two different types of hillslope thermokarsts were sampled the 7<sup>th</sup> and 10<sup>th</sup> of August, 2018, i.e., a RTS (**Fig. 2e3c**) and an  
ALD (**Fig. 2d3d**), both for so-called "disturbed" and "undisturbed" locations. Samples were taken at 10 cm increments in the  
active layer until the top of the frozen layer was reached (where only one sample was taken). A location is defined as disturbed  
(D) when disturbance (ALD or RTS) is present and sampling was conducted within the headwall (**Fig. 3e-f**). The corresponding  
undisturbed (UD) site is located at 29 m (ALD) and 234 m (RTS) from the disturbance, in an area where the collapse had not  
135 (yet) occurred. The ALD sampled in this study is within the Ptarmigan sub-catchment and formed in 2007 (Lamoureux and  
Lafrenière, 2009). The catchment hosted other ALDs in 2007, collectively covering 12% of the total catchment area (21.3 ha;  
Lafrenière and Lamoureux, 2013). Field observations showed that, while the ALD was not active in summer 2018 at the time  
of sampling, the RTS was still very active (up to 1-meter annual retreat rate) and exposed fresh sediments from that summer.



140

**Figure 32: Study site and sampling location.** (a) Location of the Melville island; (b) Location of the Cape Bounty Arctic Watershed Observatory (CBAWO), on the south-central coast of the Melville island; (c-d) Locations of the sampled profiles. RTS = retrogressive thaw slump; ALD = active layer detachment; D = disturbed; UD = undisturbed.; (e-f) Photographs and profile locations of the disturbed sites at the time of sampling for the RTS-D (e) and the ALD-D (f). Profile locations on panels (e) and (f) are represented by colored vertical rectangles. -Map created in ArcMap® 10.8. Basemap source: World Imagery (Esri et al., 2024). -Photos from panels e-f: M. Lafrenière, J. Heslop.

145

### 2.3 Mineralogy

150 We used X-ray diffraction (XRD) to characterize the crystalline mineral phases in the samples. The mineralogy of the bulk samples was determined on non-oriented powder finely ground in a mortar (Cu K $\alpha$ , Bruker Advance D8 diffractometer, detection limit 5% by weight) in the sample closest to the surface and in the deepest sample for the four modalities (n = 8).

### 2.4 Bulk Mineral Element Concentrations (Fe, Al, Mn, Ca, K)

155 We measured the total concentrations of Fe, Al, Mn, Ca and K in all samples (n = 33) using a portable X-ray fluorescence (XRF) device (*Niton XL3t GOLDD* + pXRF; ThermoFisher Scientific, Waltham, the United States). Measurements were carried out in laboratory conditions (ex-situ) on air-dried samples to prevent the introduction of additional variability (e.g. water content, sample heterogeneity). Briefly, the samples were placed on a circular plastic cap (2.5 cm in diameter), its base covered with a thin transparent film (prolene 4  $\mu$ m). To ensure that detected intensities were not underestimated, the minimum sample thickness in the cap was set at 2 cm (Ravansari et al., 2020), and the total analysis time was set at 90 s to standardize  
160 each measurement. Concentrations measured by pXRF were calibrated using a method following Monhonval et al. (2021a). Linear regressions were used to correct pXRF concentrations for trueness on all samples (n = 33). These regressions were obtained from element concentrations measured by pXRF and by inductively coupled plasma optical emission spectrometry (ICP-OES) after alkaline fusion on samples from different permafrost environments (n = 172), including 8 samples from the present study. Those 8 samples were the top and bottom samples for each modality (ALD-D, ALD-UD, RTS-D, RTS-UD).  
165 The regressions had good coefficients of determination (robust  $R^2 \geq 0.9$  for Fe, Ca, K ; robust  $R^2 \geq 0.8$  for Mn; robust  $R^2 \geq 0.6$  for Al; **Fig. A. 1**). In the following, the total element concentration measured by XRF and corrected for trueness will be referred to as Fe<sub>t</sub>, Al<sub>t</sub>, Mn<sub>t</sub>, Ca<sub>t</sub>, K<sub>t</sub>.

### 2.5 Water extractions

170 The water-soluble concentration in Ca and K were determined by water extraction on all samples (n=33). This measurement is used to assess to what extent a soil or sediment has undergone leaching of water-extractable soluble elements. The water extraction protocol follows that in Thomas et al. (2023), consisting of placing 2 g of soil in 20 mL of ultrapure water (resistivity range = 10 - 18 M $\Omega$ ·cm) for 2 h at room temperature and targets the elements released in the soluble phase (Ca, K). The element concentrations are measured in solution after filtration at 0.2  $\mu$ m by ICP-OES. We also measured the pH (pH Electrode Inlab micro) and conductivity (probe Inlab 715-4mm) on the water extract (Mettler Toledo SevenCompact DuoS213). In the  
175 following, elements extracted with ultrapure water will be referred to as the symbol of the corresponding element followed by a subscript letter 'w' (Ca<sub>w</sub>, K<sub>w</sub>).

## 2.6 Total organic carbon concentrations

The total organic carbon (TOC) content was determined on finely ground sediments by subtracting the total inorganic carbon from the total carbon on all samples ( $n = 33$ ) and is reported as the mass of OC as a percentage of the total mass of dry soil (weight percent; %<sub>w</sub>t). The total carbon was measured after dry combustion via a Vario El Cube Elemental Analyser in accordance with the NF ISO 10694 standard (LOD = 0.1%<sub>w</sub>t). The measurement of total inorganic carbon was carried out by measuring the release of carbon dioxide (CO<sub>2</sub>) following acid attack of 1 g of dry soil with hydrochloric acid, following the XP CEN/TS 16375 standard. Briefly, the CO<sub>2</sub> released generates a pressure related to the inorganic carbon content and the relationship is established from a series of standards of 10, 20, 50, 100 and 200 mg CaCO<sub>3</sub> (see also Baize, 2018). For the samples analyzed ( $n = 33$ ), inorganic carbon averaged  $0.04 \pm 0.02$  %<sub>w</sub>t. Inorganic carbon concentration is therefore not significant.

## 2.7 Selective extractions (Fe, Al, Mn, Ca, organic carbon)

Two approaches of selective extraction from soil were used as indicators of complexed and poorly crystalline oxide phases (Rennert, 2019). Extraction of Fe, Al, Mn and Ca by sodium pyrophosphate targets organometallic complexes (Bascomb, 1968; Parfitt and Childs, 1988). We recognize a possible contribution from oxide nanoparticles in addition to organically bound metals (Courchesne and Turmel, 2008; Jeanroy and Guillet, 1981; Kaiser and Guggenberger, 2007), but limited by centrifugation as well as filtration of the extract. Extraction of Fe, Al and Mn by ammonium oxalate in the dark targets poorly crystalline oxides and organometallic complexes (Blakemore et al., 1981). The pool of mineral elements that form organometallic complexes or associations with OC is sometimes referred to as “reactive”. This reactive pool combines all poorly crystalline, amorphous and complexed forms of Fe, Mn, Al (and Ca), and corresponds here to extraction with ammonium oxalate (Fig. 2b). The selective extractions were carried out on all samples ( $n = 33$ ). Concentrations in Fe, Al, Mn and Ca were measured in solution by ICP-OES after each extraction. In the following, elements extracted by the pyrophosphate and oxalate methods will be referred to as the symbol of the corresponding element followed by a subscript letter indicating the type of extraction, i.e. 'p' for pyrophosphate extraction (Fe<sub>p</sub>, Al<sub>p</sub>, Mn<sub>p</sub>, Ca<sub>p</sub>) and 'o' for oxalate extraction (Fe<sub>o</sub>, Al<sub>o</sub>, Mn<sub>o</sub>). The concentration of poorly crystalline oxides is calculated as the difference between the concentration in the oxalate extract and the pyrophosphate extract (Fe<sub>o</sub>-Fe<sub>p</sub>; Al<sub>o</sub>-Al<sub>p</sub>; Mn<sub>o</sub>-Mn<sub>p</sub>).

The pool of OC selectively extracted with sodium pyrophosphate (Bascomb, 1968; Jeanroy and Guillet, 1981; Parfitt and Childs, 1988) was measured on the same solutions as those used for the selective extraction of metals ( $n = 33$ ). Briefly, we measured dissolved OC released after dispersion by pyrophosphate using a Shimadzu TOC-L analyzer (measuring non-purgeable OC ; LOD = 50 µg/L). In the following, this carbon extracted by pyrophosphate will be referred to as C<sub>p</sub>.


For oxalate extracted carbon, direct measurement is impractical as ammonium oxalate ((NH<sub>3</sub>)<sub>2</sub>C<sub>2</sub>O<sub>4</sub>) and oxalic acid (H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>) are organic reagents. As a proxy, we measured the absorbance at 430 nm in the oxalate extract (via a Genesys 10 S VIS

spectrophotometer, with the extractant solution as a blank) to evaluate the organic acid concentration. The optical density of the oxalate extract (ODOE) is mainly influenced by the extracted fulvic acids present in the oxalate extract (Daly, 1982).

210 The selective extractions presented here target organic carbon in the form of small fragments of biopolymers stabilized by chemical bonds. Some larger fragments of biopolymers—also stabilized by chemical bonds—will not be completely dissolved by selective extractions and will therefore not be included in this pool. These must therefore be quantified using density fractionation (see section 2.8).

## 2.8 Soil aggregate and density fractionation

215 The samples followed a two-stage fractionation protocol, based on aggregate size and density (**Fig. 4**) adapted from and extensively described in Burgeon et al. (2021). This protocol was applied to the sample closest to the surface and to the deepest sample for the four modalities (n = 8) to separate free particulate organic matter from mineral-associated organic matter, representing organic matter occluded in aggregates and sharing chemical bonds with minerals. In the first stage, we performed a series of wet sieving (in a bath of 2 cm of distilled water over the mesh, we rotated and tilted the sieve for 50 rotations during  
220 2 min) to separate the soil into 3 aggregate size classes: the coarse sand size fraction (2000–250  $\mu\text{m}$ ), the fine sand size fraction (250–50  $\mu\text{m}$ ) and the silt and clay size (S&C) fraction (<50  $\mu\text{m}$ ). Particles larger than 2 mm were eliminated. The second stage used sodium polytungstate (SPT – 1.85  $\text{g}/\text{cm}^3$ ) to differentiate light fractions non occluded in aggregates considered as particulate organic matter (< 1.85  $\text{g}/\text{cm}^3$ ) from heavy fractions (> 1.85  $\text{g}/\text{cm}^3$ ): aggregate of coarse sand size, aggregate of fine sand size and heavy fractions of free silt and clay size, aggregated or not. Briefly, we submerged 5 g of 60°C oven dried  
225 fractions obtained from the first stage in 50 mL of SPT. These were then gently shaken head over head 10 times to soak the sample and remove air bubbles and placed in a vacuum chamber (140 kPa) for 10 min then left to rest under atmospheric pressure for 20 min. All samples were then centrifuged at 4700 g for 10 min and the supernatant, considered as the light fraction, was collected on glass fiber filters (performed thrice). The light fraction was thoroughly rinsed with distilled water and dried at 60°C. The pellet was shaken into suspension using distilled water, then centrifuged. The distilled water was  
230 discarded in order to remove the SPT. This step was performed multiple times to recover heavy fraction dominated by mineral associated organic matter. On each fraction, the total carbon was measured by dry combustion via a Vario El Cube Elemental Analyser (LOD = 0.1%<sub>wt</sub>).



235 **Figure 4: Two-stage aggregate and density fractionation protocol** (adapted from Burgeon et al., 2021).

## 2.9 Statistical analyses

240 We performed computations for statistical analysis using R software version R.3.6.14.5.1 (R Core Team, 2025). Robust mixed-effect linear regression models (robust  $R^2$ ) presented in this study were fitted are implemented with an alpha of 0.95 using the `rlmer` function from the `robustlmm` package v3.3-3 (Koller, 2016). The variable ‘profile’ (RTS-D, RTS-UD, ALD-D, ALD-UD) has been included as a random factor, with random intercept and slope. We report the marginal  $R^2$  ( $R^2_m$ ), representing the population-level trend (variance explained by fixed effects) in the main text. The conditional  $R^2$  ( $R^2_c$ ), which includes both fixed and random effects and reflects total model performance, is additionally shown on the plots for reference.

245 When numerical statistics are presented in the text for dataset descriptions, the mean  $\pm$  standard deviation of the distribution is presented. For comparing two datasets, we performed nonparametric statistical Wilcoxon test.

## 3 Results


### 3.1 Mineralogy, Bulk Mineral Element Concentrations and Water Extractions

250 The diffractograms of the samples characterized indicate the presence in all samples of primary silicate minerals (quartz, feldspars and micas) and secondary silicate minerals (kaolinite, illite, vermiculite, chlorite) (**Fig. C. 1 – Fig. C. 4**) with similar diffractogram patterns in all samples.

Total concentrations in major elements (Ca, K, Fe, Al, Mn) do not vary significantly between the RTS and the ALD for the disturbed and undisturbed modalities (**Fig. B. 1**). Total iron concentration is  $36 \pm 4$  g/kg, total aluminum concentration is  $71 \pm 11$  g/kg and total manganese concentration is  $0.53 \pm 0.09$  g/kg. The concentrations are homogeneous along the depths. However, total calcium ( $\text{Ca}_t$ ;  $2.3 \pm 0.4$  g/kg) and potassium ( $\text{K}_t$ ;  $15 \pm 2$  g/kg) concentrations increase slightly with depth for the sediments exposed by the retrogressive thaw slump (RTS-D) compared to the undisturbed (RTS-UD) location (**Fig. B. 1d,e**).

The proportion of highly soluble water extractable Ca and K shows a clear increase with depth for sediments exposed by the retrogressive thaw slump (RTS-D ;  $Ca_w/Ca_t$  ;  $K_w/K_t$ ; **Fig. 5a-b**), likewise for electrical conductivity of the solution (**Fig. 5c**) and pH (**Fig. 5d**). In contrast, there is no trend with depth for active layer detachment samples (**Fig. 5e-h**).

260



**Figure 5: Results of water extraction measurements** for (a,e) the ratio between water extracted (w) and total (t) concentrations in calcium ( $Ca_w/Ca_t$ ), (b,f) the ratio between water extracted (w) and total (t) concentrations in potassium ( $K_w/K_t$ ), (c,g) electrical conductivity and (d,h) pH. RTS = retrogressive thaw slump; ALD = active layer detachment; D = disturbed; UD = undisturbed.

265


### 3.2 Selective extractions of metals

Four cations were initially examined for the formation of organo-metallic complexes, i.e., extracted by pyrophosphate: iron, aluminum, manganese and calcium. Calcium extracted by pyrophosphate ( $Ca_p$ ) was not further investigated, as it showed no correlation with pyrophosphate-extracted carbon (robust  $R^2_m = 0.0301$ ). We acknowledge that analytically, calcium extracted with pyrophosphate or other extractants reflects both calcium bound to OC and calcium released by dissolution of highly soluble mineral phases, and the pool of organically bound calcium cannot be isolated. Yet, calcium can sometimes play an important role in OC stabilization (e.g., Rowley et al., 2018). Among the three other metals involved in complexation, aluminum is dominant ( $Al_p/(Fe_p+Al_p+Mn_p) = 55 \pm 6 \%$  when concentrations are expressed on a molar basis: mmol/kg), followed by iron ( $Fe_p/(Fe_p+Al_p+Mn_p) = 40 \pm 7 \%$ ) and manganese ( $Mn_p/(Fe_p+Al_p+Mn_p) = 5 \pm 3 \%$ ). We observe a distinct decrease in the concentration of metals forming complexes in the deepest samples of the profile exposed by the thaw slump

270

275

(RTS-D; **Fig. 6a**), while the concentration of all metal complexes are homogeneous along the depth for the three other profiles (RTS-UD; ALD-D; ALD-UD; **Fig. 6b-d**). Aluminum, besides being the dominant metal in terms of molar concentration, is also the metal that best explains the distribution of complexed OC (linear-mixed-effect model regression between  $Al_p$  and  $C_p$ ; robust  $R^2_m = 0.92-89$ ; **Fig. D. 1b**), followed by iron ( $Fe_p$ ; robust  $R^2_m = 0.758$ ; **Fig. D. 1a**) and manganese ( $Mn_p$ ; robust  $R^2_m \approx 0.270$ ; **Fig. D. 1c**). The proportions of these metals in complex form in relation to total concentrations are low, with  $0.3 \pm 0.2\%$  for aluminum ( $Al_p/Al_t$ ),  $0.9 \pm 0.6\%$  for iron ( $Fe_p/Fe_t$ ) and  $6 \pm 4\%$  for manganese ( $Mn_p/Mn_t$ ).



**Figure 6: Evolution with depth of the metals forming organo-metallic complexes in (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD).**

For oxalate extraction, the proportion extracted relative to total concentration is 5 to 20 times higher than for pyrophosphate (**Fig. D. 2; Fig. D. 3; Fig. D. 4**), with  $1.4 \pm 0.2\%$  for aluminum ( $Al_o/Al_t$ ),  $10 \pm 2\%$  for iron ( $Fe_o/Fe_t$ ) and  $63 \pm 14\%$  for manganese ( $Mn_o/Mn_t$ ). As for the proportion of poorly crystalline oxides, it stands at  $1.1 \pm 0.2\%$  for aluminum ( $(Al_o-Al_p)/Al_t$ ),  $9 \pm 2\%$  for iron ( $(Fe_o-Fe_p)/Fe_t$ ) and  $56 \pm 14\%$  for manganese ( $(Mn_o-Mn_p)/Mn_t$ ). Concentrations and proportion of poorly crystalline oxides are homogeneous along depth, except in the headwall exposed by the retrogressive thaw slump (RTS-D), where there is a clear increase with depth in the proportion of poorly crystalline iron oxides (**Fig. D. 2a**) and poorly crystalline aluminum oxides (**Fig. D. 3a**), relative to the total.


### 295 3.3 Total and selectively extracted organic carbon

Total organic carbon concentrations are low along the profiles, i.e., at  $1.1 \pm 0.2\%$ . For the profile exposed by the thaw slump (RTS-D), we see an increase to  $2\%$  at 20 cm depth, before dropping to  $0.9\%$  for the deepest samples (**Fig. 7a**). The concentration of carbon forming complexes with metals is  $0.14 \pm 0.07\%$  on average ( $C_p$ ; **Fig. 7**). Relative to the TOC

concentration, this means that  $13 \pm 5\%$  of TOC is in the form of complexes with metals ( $C_p/\text{TOC}$ ; **Fig. D. 6**), and trends along  
300 the depth follow those of metal complexes (see **Fig. 6**).

Assuming a maximal sorption capacity of  $0.22 \text{ gOC/gFe}$  as ferrihydrite (Wagai and Mayer, 2007), we can calculate a  
concentration of up to  $0.08 \pm 0.02\%_{\text{wt}}$  of OC bound to poorly crystalline iron oxides ( $\text{Fe}_o\text{-Fe}_p$ ; **Fig. 7**). We acknowledge that  
the samples could potentially also contain Al-bearing amorphous silica (e.g., allophanes, imogolite; Rennert, 2019) extracted  
with oxalate, yet these cannot be quantified and singled out with confidence. Besides, the concentration in Al-bearing  
305 amorphous silica or oxides ( $\text{Al}_o\text{-Al}_p$ ) is on average 4 times lower than those containing iron ( $\text{Fe}_o\text{-Fe}_p$ ; **Fig. D. 5**), as are  
concentrations in poorly crystalline manganese oxides ( $\text{Mn}_o\text{-Mn}_p$ ; 12 times lower; **Fig. D. 5**). These Al- or Mn-bearing  
amorphous silica and poorly crystalline oxides have therefore not been considered for associations with OC. In the following,  
the calculation of OC bound to poorly crystalline iron oxides will be referred to as  $C_{\text{amorph}}$  and refer to a maximal sorption  
capacity of OC to poorly crystalline iron oxides only. Relative to TOC, this means that up to  $6 \pm 2\%$  of the TOC is bound to  
310 poorly crystalline iron oxides in these profiles (**Fig. D. 6**). We estimate accordingly that the combination of the mechanisms  
of complexation and adsorption to poorly crystalline iron oxides results in  $20 \pm 4\%$  of the TOC pool being chemically bound  
with mineral surfaces or metallic cations ( $(C_p + C_{\text{amorph}})/\text{TOC}$ ).

The optical density of the oxalate extract (ODOE) averages  $0.039 \pm 0.012$ . This value correlates well with the concentration  
of OC forming complexes with metals (linear regression between ODOE and  $C_p$ ; robust  $R^2_{\text{m}} = 0.8884$ ; **Fig. D. 7a**) and TOC  
315 (robust  $R^2_{\text{m}} = 0.6659$ ; **Fig. D. 7e**), but shows poor to no correlation with oxalate extracted metals (robust  $R^2_{\text{m}} = 0.05$  with  $\text{Fe}_o$ ;  
**Fig. D. 7b**; robust  $R^2_{\text{m}} = 0.38\text{-}23$  with  $\text{Al}_o$ ; **Fig. D. 7c** and robust  $R^2_{\text{m}} = 0.25\text{-}28$  with  $\text{Mn}_o$ ; **Fig. D. 7d**) or with the proportions  
in poorly crystalline oxides (**Fig. D. 7f-h**). On the other hand, ODOE correlates very well with the sum  $C_p + C_{\text{amorph}}$   
(robust  $R^2_{\text{m}} = 0.9492$ ), supporting our approach to calculate  $C_{\text{amorph}}$ .



320

**Figure 7: Evolution with depth of pyrophosphate-extracted carbon ( $C_p$ ) and maximum proportion of organic carbon (OC) bound to poorly crystalline Fe oxides ( $C_{amorph}$ ) in (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD). The values of OC bound to crystalline Fe oxides correspond to a calculation with a maximum sorption capacity of  $0.22 \text{ gOC/gFe}$  as ferrihydrite ( $\text{Fe}_o\text{-Fe}_p$ ). %wt = mass of OC as a percentage of the total mass of dry soil, i.e., weight percent.**

325

Sediments exposed by the retrogressive thaw slump (RTS-D) exhibit a significant decrease in OC forming complexes with metals ( $C_p$ ) along depth (Wilcoxon tests comparing the 5 surface samples with the 4 deepest samples; p-value < 0.05; see also **Fig. 7a**). In contrast, no clear trend is observed for the other three modalities (p-value > 0.3 for RTS-UD, ALD-D and ALD-UD; **Fig. 7b-d**). The OC associated with poorly crystalline Fe oxides ( $C_{amorph}$ ) in RTS-D sediments shows an opposite pattern, with concentrations increasing slightly with depth (p-value = 0.2), although  $C_{amorph}$  concentrations are, on average, half of those for  $C_p$  (**Fig. 7**). When both mechanisms ( $C_p$  and  $C_{amorph}$ ) are combined, there is still a significant decrease in overall stabilized OC concentration with depth in RTS-D sediments (p-value < 0.05; **Fig. 7a**).

330


### 3.4 Particulate and mineral associated organic matter pools

335

The mass fractions of 2000-250  $\mu\text{m}$ , 250-50  $\mu\text{m}$  and <50  $\mu\text{m}$  aggregates are  $15 \pm 4\%$ ,  $37 \pm 9\%$  and  $48 \pm 10\%$ , respectively, with no trend with depth or between RTS and ALD. The particulate organic matter resulting from density fractionation represents  $1.1 \pm 0.2\%$  on average of the total soil mass. It is divided into  $0.24 \pm 0.06\%$  for 2000-250  $\mu\text{m}$  size,  $0.4 \pm 0.1\%$  for 250-50  $\mu\text{m}$  size and  $0.5 \pm 0.1\%$  for <50  $\mu\text{m}$  size. The mineral-associated fraction, which represents  $98.3 \pm 0.2\%$  of the total

mass, is divided into  $15 \pm 4\%$  for 2000-250  $\mu\text{m}$ ,  $36 \pm 8\%$  for 250-50  $\mu\text{m}$  size and  $47 \pm 10\%$  for  $<50 \mu\text{m}$  (Tab. E. 1). The OC concentration is much higher for the particulate organic matter fractions ( $33 \pm 10 \text{ g}_{\text{OC}}/100\text{g}_{\text{fraction}}$ ) than for the mineral associated organic matter fractions ( $0.7 \pm 0.3 \text{ g}_{\text{OC}}/100\text{g}_{\text{fraction}}$ ; Tab. E. 1). When we recalculate these OC contents taking into account the mass proportion of each fraction in the soil, we obtain that the free particulate organic matter fraction represents a content of  $0.4 \pm 0.1\%_{\text{wt}}$  on average (Tab. E. 1; Fig. 8); divided in  $0.06 \pm 0.03\%_{\text{wt}}$ ,  $0.12 \pm 0.03\%_{\text{wt}}$  and  $0.20 \pm 0.07\%_{\text{wt}}$  for 2000-250  $\mu\text{m}$ , 250-50  $\mu\text{m}$  and  $<50 \mu\text{m}$  sizes, respectively (Fig. E. 1). The mineral associated organic matter fraction represents a content of  $0.7 \pm 0.1\%_{\text{wt}}$  (Tab. E. 1; Fig. 8), divided in  $0.08 \pm 0.04\%_{\text{wt}}$ ,  $0.20 \pm 0.09\%_{\text{wt}}$  and  $0.4 \pm 0.1\%_{\text{wt}}$  for 2000-250  $\mu\text{m}$ , 250-50  $\mu\text{m}$  and  $<50 \mu\text{m}$  sizes, respectively (Fig. E. 1). The free particulate organic matter represents on average  $36 \pm 10\%$  of the bulk soil TOC content, and the mineral-associated organic matter fraction represents  $64 \pm 10\%$  of TOC. For sediment exposed by the RTS, the content of mineral-associated OC decreases with depth, i.e., from  $0.84\%_{\text{wt}}$  mineral associated organic C for the shallowest sample to  $0.41\%_{\text{wt}}$  at depth (Fig. 8a), as it is the case for the chemical extractions (Fig. 8a; Fig. 7a).

350



355 **Figure 8: Comparative results of organic carbon (OC) stabilized through organo-mineral interactions with the selective chemical extraction protocol and the size and density fractionation protocol for the shallowest and deepest samples in (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD). POC = particulate organic carbon.**

## 4 Discussion

### 4.1 The proportion of organic carbon stabilized by chemical interactions is in the low range at Cape Bounty relative to other Arctic environments

360 Our estimate is that  $20 \pm 4\%$  of the TOC ( $(C_p + C_{\text{amorph}})/\text{TOC}$ ) in the soil and sediments from Cape Bounty is chemically bound  
with mineral surfaces or metallic cations, which is in the low range of values reported elsewhere. Indeed, it compares with  
 $43 \pm 20\%$  from other studies on hillslope thermokarst or Yedoma sediments in Siberia and Canada (Monhonval et al., 2021a,  
2022; Thomas et al., 2023, 2024), all of which report fairly low TOC content. Indeed, it compares with  $34 \pm 8\%$  for sediments  
from the Batagay thaw slump in Siberia (Thomas et al., 2024),  $37 \pm 12\%$  from thaw slump headwalls in the Peel Plateau,  
365 western Canada (Thomas et al., 2023),  $60 \pm 20\%$  in Yedoma sediments in Central Yakutia (Monhonval et al., 2022) and  $42$   
 $\pm 26\%$  in circum-Arctic Yedoma sediments (Monhonval et al., 2021a). In all these sites, the TOC content is quite low, i.e. on  
average  $1.6 \pm 1.5\%$ <sub>wt</sub>, which is in the same range as the samples from this study ( $1.1 \pm 0.2\%$ <sub>wt</sub>; section 3.3). This confirms that  
– for sites with comparable TOC contents – the level of stabilization through chemical interactions is appears to be about half  
as highlower at Cape Bounty than-in comparison to in-sites from other studies in the Arctic. This could be partially explained  
370 by the particular location of Cape Bounty, which is at the highest latitude ( $74^\circ 55' \text{N}$ ) compared to the study sites referenced  
here above ( $66^\circ 45' \text{N} - 67^\circ 34' \text{N}$ ). In addition, Cape Bounty landscapes consists of Holocene syngenetic permafrost developed  
in marine sediments (England et al., 2009; Hodgson et al., 1984; Lajeunesse and Hanson, 2008; Paquette et al., 2023) compared  
to the Pleistocene age of sites referenced above. This means that Cape Bounty is roughly 800-900 km farther north than the  
comparative sites, with a polar desert climate (mean annual air temperature:  $-14.8 \pm 1.3$  °C and low precipitations: <  
375 150 mm/year; Beel et al., 2018; Lamoureux and Lafrenière, 2018). Such past permafrost history and climatic conditions (i.e.,  
negative temperature and low precipitation) allow only limited pedological development and constitute no favorable conditions  
for forming mineral-OC bounds. This could partially explain why the level of OC chemical stabilization is generally low at  
Cape Bounty. Similar low proportions have also been reported in sediments from the Eurasian arctic-Arctic shelf, with mineral-  
interacting proportion of TOC of 0.5 – 22 % (Salvadó et al., 2015; TOC content =  $1.4 \pm 0.4\%$ <sub>wt</sub>) with samples collected between  
380  $70^\circ \text{N}$  and  $80^\circ \text{N}$ . García-Palacios et al. (2024) have shown conversely that higher concentrations of mineral-associated OC are  
found at lower temperatures, arguing a role of temperature limitation for persistence of the mineral-protected fraction of OC.  
This suggests that, once the mineral-OC interactions have been formed, cold can help to maintain them over time.  
Permafrost peatland environments also show fairly low levels of mineral-interacting proportion of TOC, namely in  $20 \pm 10\%$   
in lowland thermokarst degradations in Eight Mile Lake, Alaska (Monhonval et al., 2023) and 10 – 15 % in a palsa mire in  
385 Sweden (Patzner et al., 2020). In peatland ecosystems, however, the concentration of TOC is much higher, (e.g.  $21 \pm 19\%$ <sub>wt</sub>  
from Monhonval et al., 2023) and it can be assumed that the surfaces and metallic cations available for mineral-OC interactions  
are saturated and unable to establish further bonds (Thomas et al., 2024).  
The results of aggregate and density fractionation at Cape Bounty, i.e.,  $64 \pm 10\%$  of the TOC being both physically and  
chemically stabilized, show on the other hand proportions that are in the same range compared with the literature. By

390 comparison, the same method applied to Siberian permafrost sediments showed a stabilization  $73 \pm 17\%$  of the TOC (Dutta et  
al., 2006; Gentsch et al., 2015; Martens et al., 2023). These proportions are extremely variable across the Arctic and show no  
latitudinal trend; these include  $74 \pm 13\%$  in Yedoma sediments, northeast Siberia (Dutta et al., 2006), a range of 33–74% (OC  
associated with mineral particles  $<6.3 \mu\text{m}$ ) in the Pleistocene permafrost of Bol'shoy Lyakhovsky Island, Siberia (Martens et  
395 al., 2023),  $54 \pm 16\%$  in Kolyma permafrost soils in Siberia (Gentsch et al., 2015). These estimates are extremely variable  
across the Arctic, showing no latitudinal trend. Future work needs to address a systematic analysis of past and present  
environmental controls (e.g., soil and air temperature, precipitation, soil moisture) and edaphic controls (e.g., lithology, pH,  
pyrophosphate- & oxalate-extractable Fe and Al, silt and clay content) of the concentration and proportion of mineral-  
interacting OC. Total precipitation and effective soil moisture appear promising since they has been shown to be significant  
positive drivers of soil carbon accumulation in dry tundra (Klaminder et al., 2009) and mineral-interacting OC at the global  
400 scale (Kramer and Chadwick, 2018), as are silt and clay content (e.g., Georgiou et al., 2022).


#### **4.2 Organic carbon exposed by the retrogressive thaw slump in deeper horizons is less protected by organo-mineral interactions**

The retrogressive thaw slump at Cape Bounty exposes sediments at depth whose OC is less protected by organo-mineral interactions (**Fig. 7a; Fig. 8a**). In particular, organometallic complexescomplexation, an efficient mechanisms for OC  
405 protection (Kleber et al., 2015; Mikutta et al., 2006), are virtually non-existent in deep sediments exposed by the RTS (RTS-D). As it happens, deep sediments in the RTS-D are also those that are rich in soluble mineral elements (**Fig. 4a-c**; see also Lamhonwah et al., 2017) with higher pH (**Fig. 4d**). For the other profiles studied (i.e., RTS-UD, ALD-D & ALD-UD), however, there was no change with depth, either for stabilized carbon (**Fig. 7b-d**) or for soluble elements, conductivity and pH (**Fig. 4**). Yet all four profiles have a similar mineralogy (**Fig. C. 1 – Fig. C. 4**) supporting a similar lithology of the parent  
410 material. We attribute the observed trends in the RTS-D to the fact that it probably exposes sediments that were previously perennially frozen at lower depths (as they are rich in solutes) and result from a syngenetic permafrost aggradation (French, 2007; Paquette et al., 2023), thereby limiting pedological development and the potential to form organo-mineral interactions. The retrogressive nature of a thaw slump guarantees exposure to this type of fresh sediment every summer until the slump is not active (**Fig. 9a**).

415 This is in agreement with the conclusion drawn at the Peel Plateau in Canada (Thomas et al., 2023) and in the Batagay thaw slump in Siberia (Thomas et al., 2024), where shallowest soils and sediments show significantly higher  $C_p$  concentrations than deep sediments, for which pedological development has been limited. Historical permafrost thaw dynamics therefore appear to be a first-order factor dictating the concentration of stabilized OC in a given sediment type. Data gathered in this study therefore confirm that thaw slumps, which are likely to develop more intensively in the future (Kokelj et al., 2015, 2021), will  
420 potentially further expose masses of deep OC that is less well protected by organo-mineral interactions. We can point out that this limited protection in deep horizons is driven by  $C_p$  and, at Cape Bounty, is partially offset by a higher concentration of

$C_{amorph}$  (section 3.3). As  $C_{amorph}$  constitutes only a minor fraction of the TOC pool however, its offsetting effect is negligible (Fig. 7a; Fig. 8a).

We would argue that materials exposed by the ALD are sediments that have undergone multiple freeze-thaw cycles since the ALD development in 2007 up to the sampling in 2018. Therefore, the exposed materials in the ALD headwall have thawed for several summers in a row. In addition, in 2018, the active layer thickness at the time of the sampling was lower (i.e., ~0.6 m on average) due to low summer temperatures (Beel et al., 2021) than average active layer thickness (i.e., 0.75 - 1.0 m) from previous years (Lamoureux and Lafrenière, 2018) (Fig. 9b-c).



430

**Figure 9: Conceptual model of materials exposed by hillslope thermokarst structures and mineral-organic carbon (OC) interactions in surface and deep samples.** (a) an active retrogressive thaw slump, whose headwall exposes fresh material every summer; (b) a 'fresh' active layer detachment, i.e. in the days/months following its formation; (c) an "old" active layer detachment, i.e. several years after its formation and whose headwall has undergone freeze/thaw cycles for several years. The white dots on the outcrops (a) and (c) represent the sampling depths of the top and bottom samples.

435

Using a mass balance approach, we can derive a first order estimate of the OC stock exposed by the ALD and the RTS per square meter of exposed horizontal surface area and for the entire headwall. Assuming an excess ice content of 50% for depths  $\geq 60$  cm (Lamhonwah et al., 2016) and dry bulk densities of 1.53 g/cm<sup>3</sup> for depths of 0-7 cm, 1.55 g/cm<sup>3</sup> for depths of 7-26 cm, 1.67 g/cm<sup>3</sup> for depths of 26-46 cm, and 1.71 g/cm<sup>3</sup> for depths greater than 46 cm (Stanton, 2023), it appears that the RTS exposes a greater stock of OC than the ALD (~13.5 kg/m<sup>2</sup> for RTS-D versus ~9.8 kg/m<sup>2</sup> for the ALD-D; Fig. D.8). This besides supports evidence that the proportion of OC stabilized in sediments exposed by the RTS is lower than that of the ALD (~17.2% for the RTS-D versus ~18.4% for the ALD-D; Fig. D.8). It is worth pointing out that OC stocks exposed at Cape Bounty are quite low compared to OC stocks exported by thaw slumps such as at the Peel Plateau (~150 to ~400 kg/m<sup>2</sup>; Thomas et al., 2023) or Batagay (~460 kg/m<sup>2</sup>; Thomas et al., 2024), which is mainly due to the shallow headwall heights in Cape Bounty.

440

### 445 4.3 Physical and chemical mechanisms of mineral-organic carbon interactions at Cape Bounty: implications for the organic carbon stability

Measurements of the proportion of OC stabilized through mineral-OC interactions greatly differ between the selective extractions protocol ( $20 \pm 4\%$ ; **Fig. 7**) and the aggregate and density fractionation protocol ( $64 \pm 10\%$ ; **Tab. E. 1; Fig. 8**). One could therefore argue that chemical extractions (pyrophosphate and oxalate) greatly underestimate the proportion of OC that is stabilized in the sediments at the Cape Bounty. Yet, these two protocols do not target the same pools of OC. Chemical extractions target (i) organometallic complexes via pyrophosphate and (ii) organometallic complexes & OC bound through adsorption onto mineral surfaces such as poorly crystalline iron oxides via oxalate (**Fig. 12**). These two chemical extractions only include small fragments of biopolymers, which are fully dissolved during extraction. Large fragments of biopolymers can also be stabilized by complexation or adsorption, but will not be soluble and therefore not measured in the oxalate or pyrophosphate extract, due to centrifugation. Density fractionation, on the other hand, targets both (i) the same pool as chemical extractions, but also (ii) the OC occluded within the aggregates of various sizes as well as all organic molecules sharing chemical bounds with minerals (even the large fragments of biopolymers that are not completely dissolved by the extractants). The different pools of OC at Cape Bounty and the associated stabilization mechanisms can be summarized as follows:

- 460 (i) up to  $6 \pm 2\%$  of the TOC is stabilized through adsorption of organic molecules onto poorly crystalline iron oxides ( $C_{\text{amorph}}$ ; **Fig. 10**). This is calculated based on a maximum sorption capacity of  $0.22 \text{ g}_{\text{OC}}/\text{g}_{\text{Fe}}$  as ferrihydrite (Wagai and Mayer, 2007;  $\text{Fe}_o\text{-Fe}_p$ ) and therefore reflects the difference between oxalate and pyrophosphate extractions (**Fig. 1a2b**). The mechanism involves the attachment – by ligand exchange – of organic molecules to metal atoms (Al, Fe) structurally present in the minerals (**Fig. 1b2a**). The chemical links are coordinate covalent bonds, which provide the longest residence times in soils (Cui et al., 2014; von Lützow et al., 2006; Mikutta et al., 2006), i.e., more than 100 years, or even millennia (Kleber et al., 2015) under stable physico-chemical conditions (pH, redox), as poorly-crystalline surfaces are among the most effective sorbents for dissolved OC (e.g., Kaiser et al., 1997).
- 470 (ii)  $13 \pm 5\%$  of the TOC is stabilized in the form of complexes ( $C_p$ ; **Fig. 10**), and represents the pool extracted with pyrophosphate (**Fig. 1a2b**). This mechanism involves the formation of cation bridges between organic matter and negatively-charged mineral surfaces (such as smectites) via electrostatic interactions; or the formation of cation bridges between molecules of organic matter of various sizes via electrostatic interactions or coordinate covalent bond (**Fig. 1b2a**). The mechanism of interaction with polymeric metal species is also one of the most important for preservation of OC (Cui et al., 2014; Kleber et al., 2015; von Lützow et al., 2006; Mikutta et al., 2006) and provides residence time in soils up to hundreds of years (Kleber et al., 2015; von Lützow et al., 2006) under stable pH and redox conditions.
- 475 (iii)  $45 \pm 8\%$  of the TOC is either occluded in aggregates, or in the form of chemically stabilized large fragments of biopolymers that are not completely dissolved by the pyrophosphate and oxalate extractants (**Fig. 10; Fig. 1a2b**). The distinction between the two cannot be made here. This is calculated as the difference between the mineral-

interacting OC obtained by the protocol of soil aggregate and density fractionation and the mineral-interacting OC obtained via chemical extractions ( $C_p + C_{\text{amorph}}$ ). It should be pointed out that the portion of OC occluded in aggregates is stabilized by being not easily accessible to microorganisms, but there are no chemical links between organic molecules and mineral surfaces or metallic cations (**Fig. 2a1b**). This pool is therefore stabilized, but to a lesser extent than in the chemically-extracted pools. Specifically, the physically occluded pool appears to be highly sensitive to physical disturbance in cultivated soils (Bischoff et al., 2017; Cui et al., 2014; von Lützow et al., 2006) and of relatively low importance in Pleistocene permafrost samples (Martens et al., 2023), although the number of relevant studies in permafrost environments is limited and further research is essential.

- (iv)  $36 \pm 10\%$  of the TOC is a pool of free particulate OC, not being stabilized through interactions with minerals but could be recalcitrant to microbial decomposition, therefore being persistent, depending on intrinsic properties of organic molecules, diversity of microbial communities and functional traits, stoichiometry constrains and abiotic factors (Derrien et al., 2023; **Fig. 10**; **Fig. 21b-e**).

Chemical stabilization mechanisms targeting small fragments of biopolymers therefore account for a relatively small proportion of the TOC ( $20 \pm 4\%$ ) but likely support residence times spanning hundreds to thousands of years with no drastic environmental changes (e.g., pH, redox, organic matter inputs, etc.). In contrast, stabilization of larger molecules via chemical bonds or physical protection within aggregates involve a comparatively larger fraction ( $45 \pm 8\%$ ) of TOC, though the residence times of aggregates could be expected to be much shorter, typically on the scale of decades.

The average figures presented above do not reflect the signature of the deep horizon exposed by the thaw slump (RTS-D). For this horizon, the OC bound to poorly crystalline oxides contributes twice as much as the average (12% of TOC; **Fig. 10**), but this is offset by the virtual absence of OC forming complexes with metals (1% of TOC; **Fig. 10**) and by the low contribution of OC occluded in aggregates or in stable large fragments of biopolymers (32% of TOC; **Fig. 10**). Consequently, more than half of the TOC is in the form of free particulate organic carbon in this horizon, meaning that the thaw slump exposes potentially more labile OC (**Fig. 10**). We attribute this to the limited pedological development in this horizon (**section 4.2**), yet the actual fate of this OC after material export from the thaw slump headwall remains uncertain. Thomas et al. (2023) showed that the pools of  $C_p + C_{\text{amorph}}$  were well preserved in materials exported by thaw slumps in the Peel Plateau in Canada. This supports that, although limited, this mineral-interacting OC can be expected to be well protected over the long term if the physico-chemical conditions do not change. It should be noted, however, that pH or redox changes – e.g. caused by water saturation – would be likely to cause a release of mineral-bound OC (Monhonval et al., 2023; Patzner et al., 2020). Yet, hillslope thermokarst landforms are generally located on well-drained terrains leading to redox conditions considered, arguably, as oxic and relatively stable (Abbott and Jones, 2015). Since thaw slumps expose deep poorly weathered material at depth, a substantial decrease in pH, which would induce dissolution of Al-OC bonds, is unlikely at Cape Bounty, where pH levels are close to neutral (Fig. 5d,h). The fate of the sediments carried downstream from the headwall and into the debris tongues remains yet unclear, and ~~We could also hypothesize that the materials, once exported will benefit from favorable conditions for pedological development and could therefore form new organo-mineral interactions in the debris tongues in the~~

decades to come. Further research needs to evaluate mineral-OC interactions along the sediment cascade of materials exported by thaw slumps.

515




Figure 10: Individual organic carbon (OC) pools at Cape Bounty. Average concentrations and standard deviations (SD) of the physically and chemically extracted OC pools on all samples (n=8) compared to the deepest sediments exposed by the retrogressive thaw slump (RTS-D). The color-coding of the OC targeted pools matches those of the selective extractions shown in Fig. 21.

520

## 5 Conclusion


Our study leads to the following conclusions:

- At the Cape Bounty, chemical stabilization mechanisms targeting small fragments of biopolymers account for  $13 \pm 5\%$  of total organic carbon in the form of complexes and up to  $6 \pm 2\%$  associated with poorly crystalline iron oxides. These results are in the low range compared to other studies conducted on organo-mineral interactions in the Arctic, possibly attributable to temperature and humidity conditions at the Cape Bounty that are not favorable to the formation of organo-mineral interactions at this latitude.
- Deep organic carbon exposed by the retrogressive thaw slump is significantly less stabilized by organo-mineral interactions. The retrogressive thaw slumps are therefore expected to expose sediments in which organic carbon is more prone to decomposition.
- Only about one-fifth of the total organic carbon ( $20 \pm 4\%$ ) is in the form of small fragments of biopolymers stabilized through strong associations with minerals, yet this fraction likely persists the longest in soils. Physical protection, which traps carbon within aggregates or large, chemically stable molecules, accounts for a larger portion ( $45 \pm 8\%$ ) and spans a wider variety of carbon forms. Chemical stabilization mechanisms represent a minor fraction of the total organic carbon ( $20 \pm 4\%$ ), yet likely enable longer residence times, while the mechanisms of physical protection in aggregates and larger molecules stabilized by chemical bonds target comparatively wider pools ( $45 \pm 8\%$ )

535

While the data presented in this study have been put into perspective with the available literature, the conclusions are limited to four distinct profiles. Such studies on headwalls and materials exported by thaw slumps or active layer detachments should receive more attention from the scientific community, and should include structures of various sizes and morphologies, at different latitudes and with contrasting development dynamics.


## Appendix A | Bulk elements concentrations assessment



545

**Figure A. 1: Total concentrations in Fe (a), Al (b), Mn (c), Ca (d) and K (e) measured by inductively coupled plasma optical-emission spectrometry (ICP-OES) as a function of those measured by the portable X-ray fluorescence (pXRF).** Cross-arctic measurements include data from the yedoma domain (Monhonval et al., 2021b) and include samples from Khardang Island (n = 1), Kurungnakh Island (n = 2), Sobo sise Island (n = 58), Bykovsky peninsula (n = 2), Buor Khaya peninsula (n = 44), Stolbovoy Island (n = 1), Bol'shoy lyakhovsky Island (n = 3), Oyogos Yar coast (n = 1), Kytalyk (n = 4), Duvanny Yar (n = 5), Yukechi (n = 2), Kitluk (n = 2), Baldwin peninsula (n = 1), Colville (n = 8) and Itkilik (n = 10). Other samples originate from Zackenberg (n = 4) (Opfergelt et al., 2018), Peel Plateau (n = 3) (Thomas et al., 2023), Batagay (n = 14) (Thomas et al., 2024) and from Cape Bounty (n = 8; this study). 95% prediction intervals are presented in grey.

550



**Figure B. 1: Total concentrations in Fe, Al, Mn, Ca, and K as a function of depth for (a-e) the retrogressive thaw slump (RTS; top row, filled symbols) and (f-j) the active layer detachment (ALD; bottom row, outline symbols). The modalities ‘disturbed’ (D) are drawn in red and the modalities ‘undisturbed’ (UD) are drawn in blue.**




Figure C. 1: Diffractograms for Cape Bounty thaw slump in the disturbed profile (RTS-D).




Figure C. 2: Diffractograms for Cape Bounty thaw slump in the undisturbed profile (RTS-UD).




Figure C. 3: Diffractograms for Cape Bounty active layer detachment in the disturbed profile (ALD-D).






Figure C. 4: Diffractograms for Cape Bounty active layer detachment in the undisturbed profile (ALD-UD).


570




575 **Figure D. 1: Robust linear regression plots** between carbon extracted by pyrophosphate ( $C_p$ ) and (a) Fe extracted by pyrophosphate ( $Fe_p$ );  
 (b) Al extracted by pyrophosphate ( $Al_p$ ); (c) Mn extracted by pyrophosphate ( $Mn_p$ ); (d) sum of metal complexes ( $Fe_p + Al_p + Mn_p$ ).  
 RTS = retrogressive thaw slump; ALD = active layer detachment; D = disturbed; UD = undisturbed. Color code as in Fig. 5. The variable  
 580 'profile' (RTS-D, RTS-UD, ALD D, ALD UD) have been included as a random factor, with random intercept and slope. The marginal  $R^2$   
( $R^2_m$ ), represents the population-level trend (variance explained by fixed effects). The conditional  $R^2$  ( $R^2_c$ ) have been included to account for  
both fixed and random effects and reflects total model performance.




585 **Figure D. 2: Evolution with depth of Fe in the form of organo-metallic complexes ( $Fe_p$ ) and poorly crystalline oxides ( $Fe_o-Fe_p$ )** for (a)  
 sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed  
 by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD).



590 **Figure D. 3:** Evolution with depth of Al in the form of organo-metallic complexes ( $Al_p$ ) and poorly crystalline oxides ( $Al_o-Al_p$ ) for (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD).




595 **Figure D. 4:** Evolution with depth of Mn in the form of organo-metallic complexes ( $Mn_p$ ) and poorly crystalline oxides ( $Mn_o-Mn_p$ ) for (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD).




**Figure D. 5: Evolution with depth of the metals forming poorly crystalline oxides in (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD).**

600




**Figure D. 6: Evolution with depth of the proportion of pyrophosphate-extracted carbon relative to the total ( $C_p/TOC$ ) and maximum proportion of organic carbon (OC) bound to poorly crystalline Fe oxides, relative to the total ( $C_{amorph}/TOC$ ) for (a-b) sediments exposed by the retrogressive thaw slump (RTS-D) and the corresponding undisturbed profile (RTS-UD); (c-d) sediments exposed by the active layer detachment (ALD-D) and the corresponding undisturbed profile (ALD-UD). The values of OC bound to crystalline Fe oxides correspond to a calculation with a maximum sorption capacity of  $0.22 \text{ goC}/\text{grFe}$  as ferrihydrite ( $Fe_o-Fe_p$ ). Color code as in Fig. 5.**

605



610 **Figure D. 7: Robust linear regression plots** between optical density of oxalate extract (ODOE) and (a) carbon extracted by pyrophosphate ( $C_p$ ); (b) Fe extracted by oxalate ( $Fe_o$ ); (c) Al extracted by oxalate ( $Al_o$ ); (d) Mn extracted by oxalate ( $Mn_o$ ); (e) total organic carbon (TOC);  
615 (f) the concentration in poorly crystalline iron oxides ( $Fe_o - Fe_p$ ); (g) the concentration in poorly crystalline aluminum oxides ( $Al_o - Al_p$ ); (h) the concentration in poorly crystalline manganese oxides ( $Mn_o - Mn_p$ ). RTS = retrogressive thaw slump; ALD = active layer detachment; D = disturbed; UD = undisturbed. Color code as in Fig. 5. The variable 'profile' (RTS-D, RTS-UD, ALD-D, ALD-UD) have been included as a random factor, with random intercept and slope. The marginal  $R^2$  ( $R^2_m$ ), represents the population-level trend (variance explained by fixed effects). The conditional  $R^2$  ( $R^2_c$ ) have been included to account for both fixed and random effects and reflects total model performance.



**Figure D. 8: Evolution with depth of the stock of pyrophosphate-extracted carbon ( $C_p$ ) and maximum proportion of organic carbon (OC) bound to poorly crystalline Fe oxides ( $C_{amorph}$ ) in (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) sediments exposed by the active layer detachment (ALD-D); Stocks are estimated by assuming an excess ice content of 50% for depths  $\geq 60$  cm (Lamhonwah et al., 2016) and dry bulk densities of 1.53 g/cm<sup>3</sup> for depths of 0–7 cm, 1.55 g/cm<sup>3</sup> for depths of 7–26 cm, 1.67 g/cm<sup>3</sup> for depths of 26–46 cm, and 1.71 g/cm<sup>3</sup> for depths greater than 46 cm (Stanton, 2023). OC concentrations as in Figure 7. The bars with the gray background represent the integral over the entire depth.**

620


Table E. 1: Results of soil aggregate and density fractionation and carbon measurements on fractions

RTS/ALD	disturbed/ undisturbed (D - UD)	Depth (cm)	mass proportion of each fraction <2 mm (gF <sub>x</sub> /100g dry soil)						OC content in each fraction (gC/100gF <sub>x</sub> )					
			F2 <i>coarse sand size</i>		F3 <i>fine sand size</i>		F4 <i>silt and clay size</i>		F2 <i>coarse sand size</i>		F3 <i>fine sand size</i>		F4 <i>silt and clay size</i>	
			F5 <i>LF</i>	F6 <i>HF</i>	F7 <i>LF</i>	F8 <i>HF</i>	F26 <i>LF</i>	F27 <i>HF</i>	F5 <i>LF</i>	F6 <i>HF</i>	F7 <i>LF</i>	F8 <i>HF</i>	F26 <i>LF</i>	F27 <i>HF</i>
RTS	D	3	0,25	18,97	0,44	46,29	0,61	32,97	31,19	0,78	40,70	0,65	38,44	1,17
RTS	D	69	0,28	19,01	0,28	21,89	0,57	57,29	29,65	0,55	49,28	0,42	49,82	0,38
RTS	UD	3	0,16	14,99	0,30	33,30	0,57	50,26	30,37	0,66	44,30	0,56	37,59	0,97
RTS	UD	61,5	0,18	11,20	0,28	31,29	0,78	55,68	16,18	0,24	43,50	0,51	36,11	0,82
ALD	D	2	0,23	14,15	0,23	40,49	0,32	43,95	12,23	0,39	31,65	0,33	36,89	1,26
ALD	D	66	0,29	9,35	0,30	33,30	0,51	55,47	20,19	0,41	35,45	0,60	33,14	0,69
ALD	UD	3	0,34	17,64	0,34	47,54	0,37	33,47	29,12	0,33	27,33	0,72	31,39	1,26
ALD	UD	60	0,20	11,34	0,68	37,36	0,48	49,58	19,78	0,88	21,35	0,41	35,05	0,87

RTS/ALD	disturbed/ undisturbed (D - UD)	Depth (cm)	OC content in soil from each fraction (gC/g dry soil) or (% <sub>wt</sub> )						OC content (gC/g dry soil) or (% <sub>wt</sub> )		
			F2 <i>coarse sand size</i>		F3 <i>fine sand size</i>		F4 <i>silt and clay size</i>		f-POC LF	OC-mineral interactions HF	TOC
			F5 <i>LF</i>	F6 <i>HF</i>	F7 <i>LF</i>	F8 <i>HF</i>	F26 <i>LF</i>	F27 <i>HF</i>	F5 + F7 + F26	F6 + F8 + F27	
RTS	D	3	0,08%	0,15%	0,18%	0,30%	0,23%	0,39%	0,49%	0,84%	1,33%
RTS	D	69	0,08%	0,10%	0,14%	0,09%	0,28%	0,22%	0,51%	0,41%	0,92%
RTS	UD	3	0,05%	0,10%	0,13%	0,18%	0,22%	0,49%	0,40%	0,77%	1,17%
RTS	UD	61,5	0,03%	0,03%	0,12%	0,16%	0,28%	0,46%	0,43%	0,65%	1,08%
ALD	D	2	0,03%	0,06%	0,07%	0,13%	0,12%	0,55%	0,22%	0,74%	0,96%
ALD	D	66	0,06%	0,04%	0,10%	0,20%	0,17%	0,38%	0,33%	0,62%	0,95%
ALD	UD	3	0,10%	0,06%	0,09%	0,34%	0,12%	0,42%	0,31%	0,82%	1,13%
ALD	UD	60	0,04%	0,10%	0,14%	0,15%	0,17%	0,43%	0,35%	0,69%	1,04%

F<sub>x</sub> = fraction as defined in Burgeon et al. (2021); LF = light fraction; HF = heavy fraction; OC = organic carbon; f-POC = free particulate organic carbon; TOC = total organic carbon



635 **Figure E. 1: Comparative results of organic carbon (OC) stabilized through organo-mineral interactions with the selective chemical extraction protocol and the soil aggregate and density fractionation protocol for the shallowest and deepest samples in (a) sediments exposed by the retrogressive thaw slump (RTS-D); (b) the corresponding undisturbed profile (RTS-UD); (c) sediments exposed by the active layer detachment (ALD-D); (d) the corresponding undisturbed profile (ALD-UD). POC = particulate organic carbon. F<sub>x</sub> = fraction as defined in Burgeon et al. (2021). These results are the same as in Fig. 8, but with details of the contribution of each fraction.**

### **Data availability**

A spreadsheet containing concentrations in total Ca, K, Fe, Mn, Al and organic carbon (Ca, K, Fe, Mn, Al and TOC); Ca and K extracted with ultrapure water ( $C_{a_w}$ ,  $K_w$ ); Fe, Al, Mn, Ca and carbon selectively extracted by pyrophosphate ( $Fe_p$ ,  $Al_p$ ,  $Mn_p$ ,  $Ca_p$  and  $C_p$ ); Fe, Al and Mn selectively extracted by ammonium oxalate ( $Fe_o$ ,  $Al_o$ ,  $Mn_o$ ); optical density of the oxalate extract (ODOE) is available at <https://doi.org/10.14428/DVN/5O6FJ3>. Raw data on soil aggregate and density fractionation have been presented in the appendices of the paper.

### **Author contribution**

MT and SO conceived and planned the experimental work. ML and JKH carried out the sampling campaign. JF supervised the size-density fractionation protocol. Water extractions were carried out by MT and NB. Selective extractions and total organic carbon measurements were carried out by MT and CM. Total and mineralogical analyses were carried out by MT under the supervision of SO. ML, JF and JKH contributed their expertise on the study site and Arctic environments. HT contributed with his expertise on selective extractions and organic carbon stability. MT wrote the manuscript under the supervision of SO with inputs from all co-authors.

### **Competing interests**

The authors declare that they have no conflict of interest.

### **Acknowledgements**

Thomas Rochereau is acknowledged for carrying out the size-density fractionation. Matthieu Leclerc and Brice de Saint-Aubert are acknowledged for the total inorganic carbon and total carbon measurements. Claudine Givron is acknowledged for the carbon measurements on fractions. Elodie Devos and Laurence Monin are acknowledged for element analyses on the selective extractions. MT thanks the Mineral and Organic Chemical Analysis (MOCA) platform at UCLouvain for support in the lab as well as members of the WeThaw project and the ELIe research department for useful critical comments. MT further thanks Sandra Arndt, Veerle Vanacker & François Jonard for their constructive feedbacks.

### **Financial support**

This project received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No.714617 to SO (WeThaw), and SO acknowledges funding from the Fund for Scientific Research FNRS in Belgium (FC69480).

## References

- Abbott, B. W. and Jones, J. B.: Permafrost collapse alters soil carbon stocks, respiration, CH<sub>4</sub>, and N<sub>2</sub>O in upland tundra, *Global Change Biology*, 21, 4570–4587, <https://doi.org/10.1111/gcb.13069>, 2015.
- AMAP: Arctic Climate Change Update 2021: Key Trends and Impacts. Summary for Policy-makers, Arctic Monitoring and Assessment Programme (AMAP), Tromsø, Norway, 2021.
- Baize, D.: Guide des analyses en pédologie, 3. ed., Quae Editions, Versailles, 328 p. pp., 2018.
- Bascomb, C. L.: Distribution of Pyrophosphate-Extractable Iron and Organic Carbon in Soils of Various Groups, *Journal of Soil Science*, 19, 251–268, <https://doi.org/10.1111/j.1365-2389.1968.tb01538.x>, 1968.
- Beel, C. R., Lamoureux, S. F., and Orwin, J. F.: Fluvial Response to a Period of Hydrometeorological Change and Landscape Disturbance in the Canadian High Arctic, *Geophysical Research Letters*, 45, 10,446-10,455, <https://doi.org/10.1029/2018GL079660>, 2018.
- Beel, C. R., Lamoureux, S. F., Orwin, J. F., Pope, M. A., Lafrenière, M. J., and Scott, N. A.: Differential impact of thermal and physical permafrost disturbances on High Arctic dissolved and particulate fluvial fluxes, *Scientific Reports*, 10, 11836, <https://doi.org/10.1038/s41598-020-68824-3>, 2020.
- Beel, C. R., Heslop, J. K., Orwin, J. F., Pope, M. A., Schevers, A. J., Hung, J. K. Y., Lafrenière, M. J., and Lamoureux, S. F.: Emerging dominance of summer rainfall driving High Arctic terrestrial-aquatic connectivity, *Nat Commun*, 12, 1448, <https://doi.org/10.1038/s41467-021-21759-3>, 2021.
- Bischoff, N., Mikutta, R., Shibistova, O., Puzanov, A., Silanteva, M., Grebennikova, A., Fuß, R., and Guggenberger, G.: Limited protection of macro-aggregate-occluded organic carbon in Siberian steppe soils, *Biogeosciences*, 14, 2627–2640, <https://doi.org/10.5194/bg-14-2627-2017>, 2017.
- Blakemore, L. C., Searle, P. L., and Daly, B. K.: Methods for chemical analysis of soils, New Zealand Soil Bur. Scientific Rep., 10A, <https://doi.org/10.7931/DL1-SBSR-10A>, 1981.
- Boudot, J. P., Bel Hadj^Brahim, A., Steiman, R., and Seigle-Murandi, F.: Biodegradation of synthetic organo-metallic complexes of iron and aluminium with selected metal to carbon ratios, *Soil Biology and Biochemistry*, 21, 961–966, [https://doi.org/10.1016/0038-0717\(89\)90088-6](https://doi.org/10.1016/0038-0717(89)90088-6), 1989.
- Brooker, A., Fraser, R. H., Olthof, I., Kokelj, S. V., and Lacelle, D.: Mapping the Activity and Evolution of Retrogressive Thaw Slumps by Tasselled Cap Trend Analysis of a Landsat Satellite Image Stack, *Permafrost and Periglacial Processes*, 25, 243–256, <https://doi.org/10.1002/ppp.1819>, 2014.
- Burgeon, V., Fouché, J., Leifeld, J., Chenu, C., and Cornélis, J.-T.: Organo-mineral associations largely contribute to the stabilization of century-old pyrogenic organic matter in cropland soils, *Geoderma*, 388, 114841, <https://doi.org/10.1016/j.geoderma.2020.114841>, 2021.
- Cotrufo, M. F. and Lavalée, J. M.: Chapter One - Soil organic matter formation, persistence, and functioning: A synthesis of current understanding to inform its conservation and regeneration, in: *Advances in Agronomy*, vol. 172, edited by: Sparks, D. L., Academic Press, 1–66, <https://doi.org/10.1016/bs.agron.2021.11.002>, 2022.

- Courchesne, F. and Turmel, M.-C.: Extractable Al, Fe, Mn, and Si, in: Soil sampling and methods of analysis, edited by: Carter, M. R. and Gregorich, E. G., Canadian Society of Soil Science ; CRC Press, [Pinawa, Manitoba] : Boca Raton, FL, 307–315, 2008.
- 700 Cui, J., Li, Z., Liu, Z., Ge, B., Fang, C., Zhou, C., and Tang, B.: Physical and chemical stabilization of soil organic carbon along a 500-year cultivated soil chronosequence originating from estuarine wetlands: Temporal patterns and land use effects, *Agriculture, Ecosystems & Environment*, 196, 10–20, <https://doi.org/10.1016/j.agee.2014.06.013>, 2014.
- Daly, B. K.: Identification of podzols and podzolised soils in New Zealand by relative absorbance of oxalate extracts of A and B horizons, *Geoderma*, 28, 29–38, [https://doi.org/10.1016/0016-7061\(82\)90038-6](https://doi.org/10.1016/0016-7061(82)90038-6), 1982.
- 705 Derrien, D., Barré, P., Basile-Doelsch, I., Cécillon, L., Chabbi, A., Crème, A., Fontaine, S., Henneron, L., Janot, N., Lashermes, G., Quénéa, K., Rees, F., and Dignac, M.-F.: Current controversies on mechanisms controlling soil carbon storage: implications for interactions with practitioners and policy-makers. A review, *Agron. Sustain. Dev.*, 43, 21, <https://doi.org/10.1007/s13593-023-00876-x>, 2023.
- Dutta, K., Schuur, E. a. G., Neff, J. C., and Zimov, S. A.: Potential carbon release from permafrost soils of Northeastern Siberia, *Global Change Biology*, 12, 2336–2351, <https://doi.org/10.1111/j.1365-2486.2006.01259.x>, 2006.
- England, J. H., Furze, M. F. A., and Doupé, J. P.: Revision of the NW Laurentide Ice Sheet: implications for paleoclimate, the northeast extremity of Beringia, and Arctic Ocean sedimentation, *Quaternary Science Reviews*, 28, 1573–1596, <https://doi.org/10.1016/j.quascirev.2009.04.006>, 2009.
- Esri, Maxar, Earthstar Geographics, and GIS User Community: “World Imagery” [basemap]., [https://services.arcgisonline.com/ArcGIS/rest/services/World\\_Imagery/MapServer](https://services.arcgisonline.com/ArcGIS/rest/services/World_Imagery/MapServer), 2024.
- 715 Fox-Kemper, B., Hewitt, H. T., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S. S., Edwards, T. L., Golledge, N. R., Hemer, M., Koop, R. E., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I. S., Ruiz, L., Sallée, J.-B., Slangen, A. B. A., and Yu, Y.: Ocean, Cryosphere and Sea Level Change, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1211–1362, <https://doi.org/10.1017/9781009157896.011>, 2021.
- French, H. M.: *The Periglacial Environment*, Third Edition., John Wiley & Sons, Ltd, Chichester, England, 2007.
- 725 García-Palacios, P., Bradford, M. A., Benavente-Ferraces, I., de Celis, M., Delgado-Baquerizo, M., García-Gil, J. C., Gaitán, J. J., Goñi-Urtiaga, A., Mueller, C. W., Panettieri, M., Rey, A., Sáez-Sandino, T., Schuur, E. A. G., Sokol, N. W., Tedersoo, L., and Plaza, C.: Dominance of particulate organic carbon in top mineral soils in cold regions, *Nat. Geosci.*, 1–6, <https://doi.org/10.1038/s41561-023-01354-5>, 2024.
- 730 Gentsch, N., Mikutta, R., Shibistova, O., Wild, B., Schnecker, J., Richter, A., Urich, T., Gittel, A., Šantrůčková, H., Bárta, J., Lashchinskiy, N., Mueller, C. W., Fuß, R., and Guggenberger, G.: Properties and bioavailability of particulate and mineral-associated organic matter in Arctic permafrost soils, Lower Kolyma Region, Russia, *European Journal of Soil Science*, 66, 722–734, <https://doi.org/10.1111/ejss.12269>, 2015.
- Georgiou, K., Jackson, R. B., Vindušková, O., Abramoff, R. Z., Ahlström, A., Feng, W., Harden, J. W., Pellegrini, A. F. A., Polley, H. W., Soong, J. L., Riley, W. J., and Torn, M. S.: Global stocks and capacity of mineral-associated soil organic carbon, *Nat Commun*, 13, 3797, <https://doi.org/10.1038/s41467-022-31540-9>, 2022.

- 735 Günther, F., Grosse, G., Wetterich, S., Jones, B. M., Kunitsky, V. V., Kienast, F., and Schirrmeister, L.: The Batagay mega thaw slump, Yana Uplands, Yakutia, Russia: permafrost thaw dynamics on decadal time scale, in: PAST Gateways - Palaeo-Arctic Spatial and Temporal Gateways - Third International Conference and Workshop, PAST Gateways - Palaeo-Arctic Spatial and Temporal Gateways - Third International Conference and Workshop, Potsdam, Germany, 18 May 2015 - 22 May 2015, 2015.
- 740 Heginbottom, J. A., Brown, J., Humlum, O., and Svensson, H.: Permafrost and Periglacial Environments, in: State of the Earth's cryosphere at the beginning of the 21st century: glaciers, global snow cover, floating ice, and permafrost and periglacial environments, U.S. Geological Survey, Reston, VA, A425–A496, 2012.
- Hodgson, D. A., Vincent, J.-S., and Fyles, J. G.: Quaternary geology of central Melville Island, Northwest Territories, Canada, 1984.
- 745 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. a. G., Ping, C.-L., Schirrmeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.
- Jeanroy, E. and Guillet, B.: The occurrence of suspended ferruginous particles in pyrophosphate extracts of some soil horizons, *Geoderma*, 26, 95–105, [https://doi.org/10.1016/0016-7061\(81\)90078-1](https://doi.org/10.1016/0016-7061(81)90078-1), 1981.
- 750 Jilling, A., Grandy, A. S., Daly, A. B., Hestrin, R., Possinger, A., Abramoff, R., Annis, M., Cates, A. M., Dynarski, K., Georgiou, K., Heckman, K., Keiluweit, M., Lang, A. K., Phillips, R. P., Rocci, K., Shabtai, I. A., Sokol, N. W., and Whalen, E. D.: Evidence for the existence and ecological relevance of fast-cycling mineral-associated organic matter, *Commun Earth Environ*, 6, 690, <https://doi.org/10.1038/s43247-025-02681-8>, 2025.
- 755 Jongejans, L. L., Mangelsdorf, K., Karger, C., Opel, T., Wetterich, S., Courtin, J., Meyer, H., Kizyakov, A. I., Grosse, G., Shepelev, A. G., Syromyatnikov, I. I., Fedorov, A. N., and Strauss, J.: Molecular biomarkers in Batagay megaslump permafrost deposits reveal clear differences in organic matter preservation between glacial and interglacial periods, *The Cryosphere*, 16, 3601–3617, <https://doi.org/10.5194/tc-16-3601-2022>, 2022.
- Kaiser, K. and Guggenberger, G.: Sorptive stabilization of organic matter by microporous goethite: sorption into small pores vs. surface complexation, *European Journal of Soil Science*, 58, 45–59, <https://doi.org/10.1111/j.1365-2389.2006.00799.x>, 2007.
- 760 Kaiser, K., Guggenberger, G., Haumaier, L., and Zech, W.: Dissolved organic matter sorption on sub soils and minerals studied by <sup>13</sup>C-NMR and DRIFT spectroscopy, *European Journal of Soil Science*, 48, 301–310, <https://doi.org/10.1111/j.1365-2389.1997.tb00550.x>, 1997.
- 765 Klaminder, J., Yoo, K., and Giesler, R.: Soil carbon accumulation in the dry tundra: Important role played by precipitation, *Journal of Geophysical Research: Biogeosciences*, 114, <https://doi.org/10.1029/2009JG000947>, 2009.
- Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., and Nico, P. S.: Mineral–Organic Associations: Formation, Properties, and Relevance in Soil Environments, in: *Advances in Agronomy*, vol. 130, edited by: Sparks, D. L., Academic Press, 1–140, <https://doi.org/10.1016/bs.agron.2014.10.005>, 2015.
- 770 Kokelj, S. V. and Jorgenson, M. T.: Advances in Thermokarst Research, *Permafr. Periglac. Process.*, 24, 108–119, <https://doi.org/10.1002/ppp.1779>, 2013.

- Kokelj, S. V., Tunnicliffe, J., Lacelle, D., Lantz, T. C., Chin, K. S., and Fraser, R.: Increased precipitation drives mega slump development and destabilization of ice-rich permafrost terrain, northwestern Canada, *Global and Planetary Change*, 129, 56–68, <https://doi.org/10.1016/j.gloplacha.2015.02.008>, 2015.
- 775 Kokelj, S. V., Kokoszka, J., van der Sluijs, J., Rudy, A. C. A., Tunnicliffe, J., Shakil, S., Tank, S. E., and Zolkos, S.: Thaw-driven mass wasting couples slopes with downstream systems, and effects propagate through Arctic drainage networks, *The Cryosphere*, 15, 3059–3081, <https://doi.org/10.5194/tc-15-3059-2021>, 2021.
- Koller, M.: *robustlmm: An R Package for Robust Estimation of Linear Mixed-Effects Models*, *Journal of Statistical Software*, 75, 1–24, <https://doi.org/10.18637/jss.v075.i06>, 2016.
- 780 Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: Permafrost carbon-climate feedbacks accelerate global warming, *PNAS*, 108, 14769–14774, <https://doi.org/10.1073/pnas.1103910108>, 2011.
- Kramer, M. G. and Chadwick, O. A.: Climate-driven thresholds in reactive mineral retention of soil carbon at the global scale, *Nature Climate Change*, 8, 1104–1108, <https://doi.org/10.1038/s41558-018-0341-4>, 2018.
- 785 Kunitsky, V. V., Syromyatnikov, I., Schirrmeyer, L., Skachov, Y. B., Grosse, G., Wetterich, S., and Grigoriev, M. N.: Ice-rich Permafrost and thermal denudation in the Batagay area (Yana Upland, East Siberia), *Earth Cryosphere (Kriosfera Zemli)*, 17, 56–58, 2013.
- Lacelle, D., Brooker, A., Fraser, R. H., and Kokelj, S. V.: Distribution and growth of thaw slumps in the Richardson Mountains–Peel Plateau region, northwestern Canada, *Geomorphology*, 235, 40–51, <https://doi.org/10.1016/j.geomorph.2015.01.024>, 2015.
- 790 Lafrenière, M. J. and Lamoureux, S. F.: Thermal Perturbation and Rainfall Runoff have Greater Impact on Seasonal Solute Loads than Physical Disturbance of the Active Layer, *Permafr. Periglac. Process.*, 24, 241–251, <https://doi.org/10.1002/ppp.1784>, 2013.
- Lajeunesse, P. and Hanson, M. A.: Field observations of recent transgression on northern and eastern Melville Island, western Canadian Arctic Archipelago, *Geomorphology*, 101, 618–630, <https://doi.org/10.1016/j.geomorph.2008.03.002>, 2008.
- 795 Lamhonwah, D., Lafreniere, M., Lamoureux, S., and Wolfe, B.: Multi-year impacts of permafrost disturbance and thermal perturbation on High Arctic stream chemistry, *Arctic Science*, 3, 254–276, <https://doi.org/10.1139/AS-2016-0024>, 2016.
- Lamhonwah, D., Lafrenière, M. J., Lamoureux, S. F., and Wolfe, B. B.: Evaluating the hydrological and hydrochemical responses of a High Arctic catchment during an exceptionally warm summer, *Hydrological Processes*, 31, 2296–2313, <https://doi.org/10.1002/hyp.11191>, 2017.
- 800 Lamoureux, S. F. and Lafrenière, M. J.: Fluvial Impact of Extensive Active Layer Detachments, Cape Bounty, Melville Island, Canada, *Arctic, Antarctic, and Alpine Research*, 41, 59–68, <https://doi.org/10.1657/1523-0430-41.1.59>, 2009.
- Lamoureux, S. F. and Lafrenière, M. J.: More than just snowmelt: integrated watershed science for changing climate and permafrost at the Cape Bounty Arctic Watershed Observatory, *WIREs Water*, 5, e1255, <https://doi.org/10.1002/wat2.1255>, 2018.
- 805 Lavallee, J. M., Soong, J. L., and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century, *Global Change Biology*, 26, 261–273, <https://doi.org/10.1111/gcb.14859>, 2020.

- Leibman, M., Kizyakov, A., Zhdanova, Y., Sonyushkin, A., and Zimin, M.: Coastal Retreat Due to Thermodenudation on the Yugorsky Peninsula, Russia during the Last Decade, Update since 2001–2010, *Remote Sensing*, 13, 4042, <https://doi.org/10.3390/rs13204042>, 2021.
- von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Matzner, E., Guggenberger, G., Marschner, B., and Flessa, H.: Stabilization of organic matter in temperate soils: mechanisms and their relevance under different soil conditions – a review, *European Journal of Soil Science*, 57, 426–445, <https://doi.org/10.1111/j.1365-2389.2006.00809.x>, 2006.
- 815 Martens, J., Mueller, C. W., Joshi, P., Rosinger, C., Maisch, M., Kappler, A., Bonkowski, M., Schwamborn, G., Schirrmeister, L., and Rethemeyer, J.: Stabilization of mineral-associated organic carbon in Pleistocene permafrost, *Nat Commun*, 14, 2120, <https://doi.org/10.1038/s41467-023-37766-5>, 2023.
- McGuire, A. D., Lawrence, D. M., Koven, C., Klein, J. S., Burke, E., Chen, G., Jafarov, E., MacDougall, A. H., Marchenko, S., Nicolsky, D., Peng, S., Rinke, A., Ciais, P., Gouttevin, I., Hayes, D. J., Ji, D., Krinner, G., Moore, J. C., Romanovsky, V., Schädel, C., Schaefer, K., Schuur, E. A. G., and Zhuang, Q.: Dependence of the evolution of carbon dynamics in the northern permafrost region on the trajectory of climate change, *PNAS*, 115, 3882–3887, <https://doi.org/10.1073/pnas.1719903115>, 2018.
- Meredith, M., Sommerkorn, M., Cassotta, S., Derksen, C., Ekaykin, A., Hollowed, A., Kofinas, G., Mackintosh, A., Melbourne-Thomas, J., Muelbert, M. M. C., Ottersen, G., Pritchard, H., and Schuur, E. A. G.: Polar Regions, in: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, edited by: Pörtner, H.-O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegria, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 203–320, 2019.
- Mikutta, R., Kleber, M., Torn, M. S., and Jahn, R.: Stabilization of Soil Organic Matter: Association with Minerals or Chemical Recalcitrance?, *Biogeochemistry*, 77, 25–56, <https://doi.org/10.1007/s10533-005-0712-6>, 2006.
- 830 Monhonval, A., Strauss, J., Mauclet, E., Hirst, C., Bemelmans, N., Grosse, G., Schirrmeister, L., Fuchs, M., and Opfergelt, S.: Iron Redistribution Upon Thermokarst Processes in the Yedoma Domain, *Front. Earth Sci.*, 9, 703339, <https://doi.org/10.3389/feart.2021.703339>, 2021a.
- Monhonval, A., Mauclet, E., Pereira, B., Vandeuren, A., Strauss, J., Grosse, G., Schirrmeister, L., Fuchs, M., Kuhry, P., and Opfergelt, S.: Mineral Element Stocks in the Yedoma Domain: A Novel Method Applied to Ice-Rich Permafrost Regions, *Frontiers in Earth Science*, 9, 773, <https://doi.org/10.3389/feart.2021.703304>, 2021b.
- Monhonval, A., Strauss, J., Thomas, M., Hirst, C., Titeux, H., Louis, J., Gilliot, A., du Bois d’Aische, E., Pereira, B., Vandeuren, A., Grosse, G., Schirrmeister, L., Jongejans, L. L., Ulrich, M., and Opfergelt, S.: Thermokarst processes increase the supply of stabilizing surfaces and elements (Fe, Mn, Al, and Ca) for mineral–organic carbon interactions, *Permafr. Periglac. Process.*, 33, 452–469, <https://doi.org/10.1002/ppp.2162>, 2022.
- 840 Monhonval, A., Mauclet, E., Hirst, C., Bemelmans, N., Eekman, E., Schuur, E. A. G., and Opfergelt, S.: Mineral organic carbon interactions in dry versus wet tundra soils, *Geoderma*, 436, 116552, <https://doi.org/10.1016/j.geoderma.2023.116552>, 2023.
- Natali, S. M., Holdren, J. P., Rogers, B. M., Treharne, R., Duffy, P. B., Pomerance, R., and MacDonald, E.: Permafrost carbon feedbacks threaten global climate goals, *Proc. Natl. Acad. Sci.*, 118, e2100163118, <https://doi.org/10.1073/pnas.2100163118>, 2021.
- 845 Opfergelt, S.: The next generation of climate model should account for the evolution of mineral-organic interactions with permafrost thaw, *Environ. Res. Lett.*, 15, 091003, <https://doi.org/10.1088/1748-9326/ab9a6d>, 2020.

- Opfergelt, S., Mauclet, E., Lefebvre, B., Monhonval, A., Bertouille, N., Vandeuren, A., Pereira, B., Hirst, C., Kuhry, P., Strauss, J., Grosse, G., and Conley, D. J.: The permafrost mineral reserve: identify potential mineral nutrient hotspots upon thawing, 5th European Conference on Permafrost, 5th European Conference on Permafrost, Chamonix Mont-Blanc, France, 780–781, 2018.
- Paquette, M., Lafrenière, M. J., and Lamoureux, S. F.: Landscape influence on permafrost ground ice geochemistry in a polar desert environment, Resolute Bay, Nunavut, *Arctic Science*, 9, 465–482, <https://doi.org/10.1139/as-2021-0049>, 2023.
- Parfitt, R. L. and Childs, C. W.: Estimation of forms of Fe and Al - a review, and analysis of contrasting soils by dissolution and Mossbauer methods, *Soil Res.*, 26, 121–144, <https://doi.org/10.1071/sr9880121>, 1988.
- Patzner, M. S., Mueller, C. W., Malusova, M., Baur, M., Nikeleit, V., Scholten, T., Hoeschen, C., Byrne, J. M., Borch, T., Kappler, A., and Bryce, C.: Iron mineral dissolution releases iron and associated organic carbon during permafrost thaw, *Nat Commun*, 11, 6329, <https://doi.org/10.1038/s41467-020-20102-6>, 2020.
- R Core Team: R: A Language and Environment for Statistical Computing, 2025.
- 860 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Commun Earth Environ*, 3, 1–10, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Ravansari, R., Wilson, S. C., and Tighe, M.: Portable X-ray fluorescence for environmental assessment of soils: Not just a point and shoot method, *Environment International*, 134, 105250, <https://doi.org/10.1016/j.envint.2019.105250>, 2020.
- 865 Rennert, T.: Wet-chemical extractions to characterise pedogenic Al and Fe species – a critical review, *Soil Res.*, 57, 1–16, <https://doi.org/10.1071/SR18299>, 2019.
- Rowley, M. C., Grand, S., and Verrecchia, É. P.: Calcium-mediated stabilisation of soil organic carbon, *Biogeochemistry*, 137, 27–49, <https://doi.org/10.1007/s10533-017-0410-1>, 2018.
- Rudy, A. C. A., Lamoureux, S. F., Treitz, P., and Collingwood, A.: Identifying permafrost slope disturbance using multi-temporal optical satellite images and change detection techniques, *Cold Regions Science and Technology*, 88, 37–49, <https://doi.org/10.1016/j.coldregions.2012.12.008>, 2013.
- Salvadó, J. A., Tesi, T., Andersson, A., Ingri, J., Dudarev, O. V., Semiletov, I. P., and Gustafsson, Ö.: Organic carbon remobilized from thawing permafrost is resequenced by reactive iron on the Eurasian Arctic Shelf, *Geophysical Research Letters*, 42, 8122–8130, <https://doi.org/10.1002/2015GL066058>, 2015.
- 875 Schuur, E. a. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.
- Schuur, E. A. G., Abbott, B. W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., Grosse, G., Jones, M., Koven, C., Leshyk, V., Lawrence, D., Lorant, M. M., Mauritz, M., Olefeldt, D., Natali, S., Rodenhizer, H., Salmon, V., Schädel, C., Strauss, J., Treat, C., and Turetsky, M.: Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic, *Annual Review of Environment and Resources*, 47, 343–371, <https://doi.org/10.1146/annurev-environ-012220-011847>, 2022.
- 880 Séjourné, A., Costard, F., Fedorov, A., Gargani, J., Skorve, J., Massé, M., and Mège, D.: Evolution of the banks of thermokarst lakes in Central Yakutia (Central Siberia) due to retrogressive thaw slump activity controlled by insolation, *Geomorphology*, 241, 31–40, <https://doi.org/10.1016/j.geomorph.2015.03.033>, 2015.

- 885 Shakil, S., Tank, S. E., Kokelj, S. V., Vonk, J. E., and Zolkos, S.: Particulate dominance of organic carbon mobilization from thaw slumps on the Peel Plateau, NT: Quantification and implications for stream systems and permafrost carbon release, *Environ. Res. Lett.*, 15, 114019, <https://doi.org/10.1088/1748-9326/abac36>, 2020.
- van der Sluijs, J., Kokelj, S. V., Fraser, R. H., Tunnicliffe, J., and Lacelle, D.: Permafrost Terrain Dynamics and Infrastructure Impacts Revealed by UAV Photogrammetry and Thermal Imaging, *Remote Sensing*, 10, 1734, <https://doi.org/10.3390/rs10111734>, 2018.
- 890 van der Sluijs, J., Kokelj, S. V., and Tunnicliffe, J. F.: Allometric scaling of retrogressive thaw slumps, *The Cryosphere*, 17, 4511–4533, <https://doi.org/10.5194/tc-17-4511-2023>, 2023.
- Stanton, T.: Soil Properties and Trace Gas Fluxes in a Chronosequence of Permafrost Disturbances, Cape Bounty, Melville Island, Nunavut, Master Thesis, Queen’s University, Kingston, Ontario, Canada, 128 pp., 2023.
- 895 Strauss, J., Abbott, B. W., Hugelius, G., Schuur, E. A. G., Treat, C., Fuchs, M., Schädel, C., Ulrich, M., Turetsky, M., Keuschnig, M., Biasi, C., Yang, Y., and Grosse, G.: Permafrost, in: Recarbonizing global soils – A technical manual of recommended management practices, vol. Volume 2: Hot spots and bright spots of soil organic carbon, FAO, Rome, Italy, <https://doi.org/10.4060/cb6378en>, 2021.
- 900 Strauss, J., Fuchs, M., Hugelius, G., Miesner, F., Nitze, I., Opfergelt, S., Schuur, E., Treat, C., Turetsky, M., Yang, Y., and Grosse, G.: Organic matter storage and vulnerability in the permafrost domain, in: Reference Module in Earth Systems and Environmental Sciences, Elsevier, <https://doi.org/10.1016/B978-0-323-99931-1.00164-1>, 2024.
- Tanski, G., Lantuit, H., Ruttor, S., Knoblauch, C., Radosavljevic, B., Strauss, J., Wolter, J., Irrgang, A. M., Ramage, J., and Fritz, M.: Transformation of terrestrial organic matter along thermokarst-affected permafrost coasts in the Arctic, *Sci. Total Environ.*, 581–582, 434–447, <https://doi.org/10.1016/j.scitotenv.2016.12.152>, 2017.
- 905 Thomas, M., Monhonval, A., Hirst, C., Bröder, L., Zolkos, S., Vonk, J. E., Tank, S. E., Keskitalo, K. H., Shakil, S., Kokelj, S. V., van der Sluijs, J., and Opfergelt, S.: Evidence for preservation of organic carbon interacting with iron in material displaced from retrogressive thaw slumps: Case study in Peel Plateau, western Canadian Arctic, *Geoderma*, 433, 116443, <https://doi.org/10.1016/j.geoderma.2023.116443>, 2023.
- 910 Thomas, M., Jongejans, L. L., Strauss, J., Vermeylen, C., Calcus, S., Opel, T., Kizyakov, A., Wetterich, S., Grosse, G., and Opfergelt, S.: A Third of Organic Carbon Is Mineral Bound in Permafrost Sediments Exposed by the World’s Largest Thaw Slump, Batagay, Siberia, *Permafr. Periglac. Process.*, 35, 278–293, <https://doi.org/10.1002/ppp.2230>, 2024.
- Treitz, P. M., Atkinson, D. M., Blaser, A., Bonney, M. T., Braybrook, C. A., Buckley, E. C., Collingwood, A., Edwards, R., van Ewijk, K., Freemantle, V., Gregory, F., Holloway, J., Hung, J. K. Y., Lamoureux, S. F., Liu, N., Ljubicic, G., Robson, G., Rudy, A. C. A., Scott, N. A., Shang, C., and Wall, J.: Remote sensing of biogeophysical variables at the Cape Bounty Arctic Watershed Observatory, Melville Island, Nunavut, Canada, *Arctic Science*, 10, 281–304, <https://doi.org/10.1139/as-2023-0043>, 2024.
- 915 Turetsky, M. R., Abbott, B. W., Jones, M. C., Anthony, K. W., Olefeldt, D., Schuur, E. A. G., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., Lawrence, D. M., Gibson, C., Sannel, A. B. K., and McGuire, A. D.: Carbon release through abrupt permafrost thaw, *Nature Geoscience*, 13, 138–143, <https://doi.org/10.1038/s41561-019-0526-0>, 2020.
- 920 Vadakkedath, V., Zawadzki, J., and Przędziecki, K.: Multisensory satellite observations of the expansion of the Batagaika crater and succession of vegetation in its interior from 1991 to 2018, *Environ Earth Sci*, 79, 150, <https://doi.org/10.1007/s12665-020-8895-7>, 2020.

Wagai, R. and Mayer, L. M.: Sorptive stabilization of organic matter in soils by hydrous iron oxides, *Geochimica et Cosmochimica Acta*, 71, 25–35, <https://doi.org/10.1016/j.gca.2006.08.047>, 2007.

925