

Upscaling of soil methane fluxes from topographic attributes derived from a digital elevation model in a cold temperate mountain forest

Sumonta Kumar Paul¹, Keisuke Yuasa¹, Masako Dannoura¹, Daniel Epron¹

¹ Graduate School of Agriculture, Kyoto University, 606-8502, Japan

5

Correspondence to: Sumonta Kumar Paul (paulsumonta@gmail.com), Daniel Epron (daniel.epron.3a@kyoto-u.ac.jp)

ORCID ID

SKP: 0009-0000-7293-7255

MD: 0000-0003-0389-871X

10 DE: 0000-0001-9451-3437

Abstract. Forest soils are generally considered a sink for atmospheric methane (CH_4), but their uptake rate can vary considerably in space and time. This study **investigated the role of terrain attributes (topography, vegetation)** on soil CH_4 fluxes and the temporal patterns **of spatially distributed soil CH_4 fluxes** in a topographically complex cold-temperate mountain forest in central Japan. We measured soil CH_4 fluxes nine times during the snow-free season at multiple locations within a 40-ha area in a forested watershed. We used a machine-learning approach to upscale the measured soil CH_4 fluxes to the landscape scale at each date of measurement, using topographic and vegetation attributes derived from a digital elevation model and aerial images. **Upland** soils were a sink of CH_4 , while small wetland patches emitted CH_4 consistently throughout the study period.

Unfortunately, the quantile random forest models were unable to accurately reproduce the measured fluxes when wetland measurements were included in the training dataset. The accuracy of predicted **upland** fluxes varied seasonally, with the highest

20 model performance observed in early autumn ($R^2 = 0.67$) and the lowest in mid-summer ($R^2 = 0.31$). **Within the study landscape**, predicted **upland** CH_4 fluxes varied significantly across topographic positions, with greater uptake on ridges and slopes than on the plain and foot slopes. **Topography played a predominant role compared to vegetation in the spatial variability of CH_4 fluxes.** **Landscape-scale predicted** CH_4 fluxes in the **upland** area ranged from -0.34 to -0.60 g $\text{CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$ in spring, -0.39 to -1.28 g $\text{CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$ in summer, and -0.48 to -0.89 g $\text{CH}_4 \text{ ha}^{-1} \text{ h}^{-1}$ in autumn. Seasonal **upland** fluxes were highly correlated with the 20-day antecedent precipitation index ($R^2 = 0.70$), revealing the importance of seasonal moisture conditions in regulating CH_4 flux dynamics. **This study highlighted the importance of topography in controlling the soil CH_4 fluxes and the efficiency of remote sensing and machine learning approaches in scaling field measurements to the landscape level, enabling visualization of spatial patterns of fluxes across the landscape over time.**

1 Introduction

30 Methane (CH_4), the second most important anthropogenic greenhouse gas, contributes substantially to the anthropogenic radiative forcing and is responsible for approximately 0.5°C of current global warming compared to 1850 - 1900 (IPCC, 2023). Natural wetlands (149 Tg $\text{CH}_4 \text{ yr}^{-1}$) and rice cultivation (30 Tg $\text{CH}_4 \text{ yr}^{-1}$) are important sources of CH_4 ; in contrast, **upland** soils are considered a biological sink of atmospheric CH_4 , with an estimated uptake of 25-45 Tg yr^{-1} , contributing 5-7% to the global CH_4 sink (Saunois et al., 2020). **Among the upland ecosystems**, forest soils account for approximately 60% of global soil CH_4 uptake (Dutaur and Verchot, 2007), and soil uptake rates are particularly high in Japanese **mountainous** forests due to their high porosity (Ishizuka et al., 2000). CH_4 uptake by forest soils is driven by methane-oxidizing bacteria in oxic soil layers, whereas anaerobic environments such as wetland soils are usually dominated by methanogenic archaea producing CH_4 (Christiansen et al., 2016). CH_4 production can also occur in **upland** soils, either in deeper soil layers or in microsites located in otherwise well-aerated soil layers, if anaerobic conditions prevail (Angel et al., 2012). Hence, CH_4 oxidation and production can occur simultaneously at the same location, **determining** the net flux.

40 Net soil CH_4 fluxes depend mainly on the soil air-filled porosity (AFP), which in turn depends on total porosity and soil water content. A high AFP enhances gas diffusion in soil and, consequently, microbial CH_4 oxidation (Kruse et al., 1996). Soil organic

matter at the soil surface can act as a physical barrier to atmospheric CH₄ diffusion and reduce the CH₄ uptake rate (Yu et al., 2017). Conversely, carbon substrates released by the decomposition of soil organic matter can increase CH₄ oxidation activity either by directly stimulating the growth of methanotrophs or by promoting CH₄ production in anaerobic microsites and indirectly supporting the growth of methanotrophs (West and Schmidt, 1999). Additionally, soil nutrients can influence soil CH₄ fluxes by regulating the soil microbial community. The activity of methanotrophic microorganisms is affected by the availability of inorganic nitrogen (Bodelier and Laanbroek, 2004). Although methanotrophic activity can be nitrogen-limited in forest soils (Veldkamp et al., 2013), increasing ammonium (NH₄⁺) concentration often reduces CH₄ uptake due to competitive inhibition by NH₄⁺ of the enzyme methane mono-oxygenase, which can oxidize both CH₄ and NH₄⁺. Nitrate (NO₃⁻) can also be a potent inhibitor of CH₄ oxidation in some soils (Mochizuki et al., 2012). Although temperature affects microbial activities, including methanogenesis and methanotrophy (Luo et al., 2013; Praeg et al., 2017), CH₄ uptake is generally less sensitive to changes in soil temperature than in soil moisture (Epron et al., 2016).

Topography and vegetation cover can create a predictable distribution of soil moisture and nutrients across topographically complex landscapes (Jeong et al., 2017; Murphy et al., 2011). In Japan, forests cover 68% of the land, mostly in mountainous areas. Conifers account for 44% of the total forest area (Lundbäck et al., 2021; Nakamura and Krestov, 2005). Topography is a critical determinant of soil hydrological conditions, from well-drained slopes to waterlogged riparian areas (Kaiser et al., 2018). Topography can also impact soil nutrient availability by altering leaf litter accumulation and the movement of soil nutrients (Osborne et al., 2017; Tateno and Takeda, 2003). The spatial distribution of trees, differences in species abundance across the landscape, and variation in litter chemistry often create heterogeneity in soil nitrogen cycling (Osborne et al., 2017). Furthermore, differences in stem flow and throughfall related to differences in canopy structure between tree species can indirectly influence spatial patterns of soil moisture (Holwerda et al., 2006).

In situ chamber measurements have long been the dominant method for studying CH₄ fluxes in forests, providing insight into the processes that drive them (Brumme and Borken, 1999; Guckland et al., 2009; Itoh et al., 2009). Until recently, most studies reported spatially average flux values measured at several locations (Gomez et al., 2016; Itoh et al., 2009). This method is acceptable for small patches of homogeneous landscapes, such as crops or single-species tree plantations in flat terrain. However, it is inappropriate for more complex landscapes, as the number of sampling points required to obtain an accurate spatially-averaged flux would increase considerably.

In complex terrains, measurement locations can be grouped into several distinct categories according to landforms (Courtois et al., 2018; Gomez et al., 2016; Itoh et al., 2009; Kagotani et al., 2001; Kaiser et al., 2018; Warner et al., 2018), soil microtopographic features (Epron et al., 2016), vegetation characteristics (Guckland et al., 2009), or land uses (Jacinthe et al., 2015). However, as Vainio et al. (2021) pointed out, aggregation assumes spatial homogeneity of fluxes within each category or requires a large number of sampling points to capture the spatial heterogeneity, and this approach ignores the spatially continuous nature of soil processes and their drivers.

More recently, regressions with multiple landscape attributes derived from remote sensing-based maps have been successfully applied to upscale CH₄ to a catchment scale (Kaiser et al., 2018). Recent studies conducted on a 12-ha forested watershed (Warner et al., 2019), a 10-ha boreal forest plot (Vainio et al., 2021), two northern peatland-forest-mosaic catchments of 450 ha and 790 ha, respectively (Räsänen et al., 2021), and a 450-ha subarctic tundra (Virkkala et al., 2024) have demonstrated the effectiveness of machine-learning modelling approaches for upscaling CH₄ fluxes from remote sensing data.

Soil CH₄ fluxes exhibit strong spatiotemporal variability in temperate mountain forests, and robust large-scale estimates remain scarce despite their importance for consolidating the global methane budget because upscaling fine-scale chamber-measured CH₄ fluxes requires an explicit understanding of their spatial and seasonal heterogeneity. We assessed the role of terrain attributes (topography, vegetation) on methane fluxes throughout the snow-free season in a topographically complex mountainous landscape, and how the spatial heterogeneity of predicted fluxes and the aggregated fluxes at the landscape level vary over time. We measured soil CH₄ fluxes several times during the snow-free season at multiple locations within a 40-ha

area in a forested watershed. We applied a random forest machine-learning approach in combination with terrain attributes from remotely sensed data, i.e., a digital elevation model (DEM) and a vegetation map derived from aerial images, to upscale measured soil CH₄ fluxes to the landscape level. Our strategy was to directly predict CH₄ fluxes using topographic and vegetation variables as proxies for soil moisture and chemistry, because incorporating soil moisture and chemistry as predictors, which would need to be extrapolated to the landscape level, would introduce additional layers of uncertainty. We hypothesized that (1) terrain attributes related to water accumulation are reliable predictors of soil CH₄ fluxes, (2) spatial patterns of uncertainties in predicted soil CH₄ fluxes vary seasonally due to a wet early summer influenced by the East Asian monsoon, (3) predicted soil CH₄ fluxes vary within the landscape depending on topography and vegetation, and (4) seasonal variations of CH₄ flux at the landscape scale are explained by recent past precipitations.

90

95

2 Materials and methods

2.1 Description of the study site and experimental design

This study was conducted in the forested upper Yura River watershed (520 ha, 35.34 N; 135.76 E) located at the Ashiu Experimental Forest of Kyoto University in northeastern Kyoto Prefecture, Japan (Fig. 1). The mean annual temperature and precipitation were 10.3°C and 2,732 mm, respectively, between 2011 and 2020 and the ground was covered by snow (2-3 m depth) from mid-December to mid-April (Epron et al., 2023). The study area is characterized by a cool-temperate monsoon climate, with a very humid early summer (520 mm in June and July on average between 2011 and 2020) and occasionally heavy precipitation caused by typhoons in late summer. The soils in the study area are classified as brown forest soils according to the Classification of Forest Soils in Japan, with a relatively thick brownish-black A horizon with a crumb structure and a brown B horizon with a blocky structure (Hirai et al., 1988; Ueda et al., 1993). The forest is primarily dominated by *Cryptomeria japonica* D. Don (Japanese cedar, 73% of the basal area in four 1-ha census plots), mixed with more than 50 broadleaved species (Ishihara et al., 2011).




Figure 1: Map of the upper Yura River watershed with its location in Japan on the left and an enlargement of the 40.2 ha study area on the right. The green triangles represent the 52 flux measurement locations on unsaturated soils (upland) and the red dots the 3 measurement locations on waterlogged wetlands. (Japan map: http://www.gsi.go.jp/ENGLISH/page_e30286.html)

The study site covered an area of 40.2 hectares and included 55 sampling points for CH₄ flux measurements and soil sampling (Fig. 1). The sampling points were chosen along three transects perpendicular to the main river, from the plain to the ridges covering two slopes (south-facing and north-facing), as well as in a lateral canyon, and along transects parallel to the main

115 river, on the plain, above the foot slope and on a ridge. The sampling was designed to encompass the landscape heterogeneity, while being constrained by the geography of the site and safety considerations. We recorded the positions of all sampling locations using a portable GPS tracker (Garmin, eTrex® Touch 35) with an accuracy of less than 5 m.

2.2 Soil sampling and analysis

120 Soil cores were collected using a sampling cylinder at 0-10 cm depth at approximately 0.3 m of the flux measurement points. Samples were sieved at 2 mm and separated into stones and fine earth. The fresh weight of the fine earth fraction was measured before being air-dried. Bulk density of this fraction was determined as the ratio of oven-dried soil (subsample dried at 105°C) to the soil volume. Soil texture was analysed using the micro-pipette method, following Burt et al. (1993). Total soil carbon (C) and nitrogen (N) contents were measured using a Macro Corder JM 1000CN (J-SCIENCE LAB Co., Ltd., Japan). The soil pH was measured in a suspension (10 g of soil in 25 ml distilled H₂O) after shaking for 1 hour.

125 2.3 Topographic characterization

To characterize and process the terrain attributes related to soil CH₄ fluxes, we used a 0.5 m grid digital elevation model (DEM) based on airborne laser surveys conducted throughout the upper Yura River watershed in 2012 by the Ashiu Experimental Forest staff. The DEM was further processed and conditioned into a 5 m grid DEM image according to the GPS tracker's accuracy (≤ 5 m) that was used to locate each collar position, enabling us to identify the corresponding pixels on the terrain attribute grids. We derived several topographic attributes from the DEM using SAGA Next Generation in QGIS (v3.34.5-Prizren). The calculated attributes included slope, profile curvature (PrC), topographic position index (TPI), saga wetness index (SWI), and vertical distance to channel network (VDCN). Among the many attributes that can be derived from a DEM, we avoided selecting those that would be redundant to limit collinearities and overparameterization. Our preselection was motivated by the fact that methane fluxes result from the activity of methanotrophic and methanogenic communities, which are controlled by soil moisture and chemistry (C, N, pH), and, to a lesser extent, temperature. All the preselected attributed were correlated with soil moisture and chemistry (Table A1) and can potentially serve as a proxy for the spatial distribution of soil moisture and nutrient availability (Jeong et al., 2017; Kemppinen et al., 2018).

130 Slope, and profile curvature were calculated following the 9-parameter 2nd order polynom method (Zevenbergen and Thorne, 1987). Negative values of profile curvature indicate a convex surface where the flow of water accelerates as it moves downslope; in contrast, positive values suggest a concave surface where the flow slows down (Pachepsky et al., 2001).

135 SWI is a refined version of the topographic wetness index (TWI) (Beven and Kirkby, 1979), which indicates that the spatial distribution of soil moisture is defined as $TWI = \ln(SCA/\tan \Theta)$, where SCA refers to the specific catchment area and Θ is the local slope. SWI considers small differences in elevation values by using an iterative modification of the specific catchment area, assuming rather homogenous hydrologic conditions in the flat areas. The SWI was calculated using the Saga wetness index algorithm, which is available in the SAGA library and integrated within QGIS (Conrad et al., 2015). Prior to computation, the DEM was hydrologically corrected by filling sinks to ensure continuous flow routing (Wang and Liu, 2006).

140 TPI describes the relative position of a location within a landscape, indicating whether it is on a ridge, slope, or valley based on the elevation compared to the surrounding terrain at a specified radius (Ågren et al., 2014). Positive values indicate ridges; negative values indicate depressions, and zero or near-zero values indicate slopes or flat areas. TPI is a highly scale-dependent variable and was calculated at the centre of circular areas with radii of 20 m, 30 m, and 50 m using the unfilled DEM. In our final model, we used TPI calculated with a 30 m radius, as it had the highest Spearman correlations with soil physical and chemical properties that influence soil CH₄ fluxes (Table A1).

145 VDCN was calculated as the elevation difference between each grid cell and the baseline of the nearest stream channel. This parameter serves as a proxy for groundwater depth, with lower VDCN values typically corresponding to areas with shallower groundwater and higher water tables, and higher values indicating deeper groundwater levels often found in upland positions (Bock and Köthe, 2008). To calculate VDCN, the filled DEM was first used to create a flow accumulation layer using the

multiple flow direction method (Freeman, 1991). The resulting flow accumulation raster was then used to create topographically defined flow channel networks, applying flow initiation thresholds of 0.5, 2.5, and 5 ha. VDCN then subsequently calculated for each threshold. In our final model, we used VDCN calculated with a 5-ha initiation threshold, as it has the highest Spearman correlations with soil physical and chemical properties that influence soil CH₄ fluxes (Table A1).
160 The site was classified into uplands (including ridges, slopes, foot slopes and plains as topographic positions where the soil is almost always unsaturated), wetlands (small patches with water-saturated soil year-round in the plain), and rivers. To distinguish wetland and upland areas, we collected additional GPS positions at the edges and within the three wetland patches, in addition to the positions of the 55 sampling points. We then used SWI, PrC, slope, and VDCN to predict the locations of
165 wetlands using a machine learning approach described in the supplementary file (Note S1). Finally, the boundaries between wetlands and uplands were refined by visual inspection. We acknowledged that using a fixed boundary between "upland" and "wetland" conditions, although these boundaries may vary seasonally depending on the balance between precipitation and evaporation, may increase uncertainties in CH₄ flux prediction. Predicting the temporal variations of these boundaries was beyond the scope of this work, and, at our site, wetlands represent only 1% of the pixels (see below), and their boundaries even
170 less. A posteriori, pixels classified as wetland had SWI values above 8.1, profile curvature between -0.003 and 0.001, slope values below 6.8, and VDCN values below 2.2 (Fig. S1). For river mapping, pixels corresponding to rivers were identified in the channel network raster, which was calculated using a 5-ha initiation threshold. Slope angle and TPI at 30 m radius were used to partition the upland into ridges, slopes, foot slopes, and the plain. Locations with TPI values of 5 or greater were defined as ridges, representing locally elevated, convex surfaces. Locations with TPI values ≤ -5 were defined as foot slopes, concave surface. Areas with intermediate TPI values ($-5 < \text{TPI} < 5$) were further divided according to slope angle: sites with slope $> 18^\circ$ were defined as slopes, and those with slope $\leq 18^\circ$ were defined as plains. Uplands, wetlands, and rivers, accounted for 94%, 1%, and 5% of the total study area, with respectively 52 sampling points located in upland areas, including 14 in plains, 9 in foot slopes, 16 in slopes, and 13 in ridges, while 3 were situated in wetland areas.
175

2.4 Vegetation classification

180 Tree inventory was conducted during the flux measurement period to classify the vegetation surrounding the flux measurement points. A circular plot with a 10-meter radius was established, centred at each flux measurement point. Within the plot, all trees were identified at the species level, and their diameter at breast height (DBH) was measured. We calculated the plot basal area (BA) as the sum of the cross-sectional areas (CSA) at breast height of all tree trunks in each plot, and subsequently determined the relative basal area of coniferous trees (RBA_{CON}) in each plot. Then, we predicted the BA and RBA_{CON} for the entire study area using SWI, TPI, VDCN, and the normalized vegetation index (NDVI) using a machine learning approach described in the supplementary file (Note S2, Fig. S2). Vegetation density was classified into three categories based on the quantile distribution of BA: high (BA > 2.6 , upper quartile), medium ($0.9 < \text{BA} < 2.6$, interquartile range), and low (BA > 2.6 , lower quartile). High, medium, and low vegetation density accounted for 37%, 37% and 26% of the total study area (Fig. S3), represented by 14, 28 and 10 sampling points, respectively. Vegetation types were classified based on RBA_{CON}. Three types
185 were defined: coniferous when RBA_{CON} was higher than 0.75, broadleaf when it was lower than 0.25, or mixed (comprising both conifers and broadleaved trees). These three types accounted 6%, 22% and 72% of the total study area (Fig. S3), represented by 11, 19 and 22 sampling points, respectively.
190

2.5 Flux measurements

195 Soil CH₄ fluxes were measured using a static, non-steady-state, non-flow-through system composed of a dark acrylic chamber (20 cm diameter and 12.5 cm height) connected to a cavity-enhanced absorption spectroscopy gas analyser (Li 7810, Licor; Lincoln, USA) with two PTFE tubes, each 1.8 m long and 4 mm in inner diameter. One week before the first measurements, a 20 cm diameter, 9 cm tall PVC collar was inserted approximately 5 cm into the soil at each sampling point. Flux from each collar was measured on nine occasions in 2023: in early spring after snowmelt (4/27), mid-spring (5/12), late spring (5/31),
5

early summer (7/06), mid-summer (7/26), late summer (9/04), early autumn (10/07), mid-autumn (11/07), and late autumn (11/30). When measuring fluxes from the three small wetland patches, we took care to avoid trampling the soil near the collars, taking advantage of the abundant presence of stones and coarse woody debris.

To measure soil CH₄ flux, the chamber was placed on the collar, and changes in CH₄ and CO₂ concentrations inside were recorded for 4 minutes at a frequency of 1 Hz. The slope of the linear regression of CH₄ concentration over time was used to calculate the soil CH₄ flux:

$$F_{CH_4} = \frac{\Delta[CH_4]}{\Delta t} \times \frac{V \times P}{A \times R \times T}$$

where F_{CH_4} is the soil CH₄ flux, $\frac{\Delta[CH_4]}{\Delta t}$ is the slope of the linear change in CH₄ concentrations over time, V is the system volume (chamber, collar above the ground, tubes, and analyzer), A is the soil area covered by the collar, and R is the ideal gas constant (8.314 J K⁻¹ mol⁻¹). A constant value of 93,525 Pa for an elevation of 650 m was used for the atmospheric pressure (P). The slope was calculated over 90 seconds following Epron et al. (2023). The R^2 of the linear variation of CH₄ concentration was less than 0.9 for a single measurement, and for this measurement, the R^2 of the linear variation of CO₂ concentration was 0.99, indicating that the low R^2 for CH₄ was due to the near-zero CH₄ flux and not to an erroneous measurement.

Soil moisture content and soil temperature near each collar were recorded on each measurement date using a soil moisture probe (SM150-T Device, Cambridge, UK) and a digital thermometer.

2.6 Climatic data

Air temperature and rainfall were measured every 10 minutes at a nearby weather station operated by the Field Science Education and Research Centre of Kyoto University. The antecedent precipitation index (API), an indicator of soil moisture conditions, was calculated using the following equation:

$$API_n = \sum_{t=1}^n P_t \times k^t$$

where, P_t is the precipitation during day t , k is the recession coefficient, and n is the number of antecedent days. The parameter k accounts for the water removed from the soil by evapotranspiration and drainage.

2.7 Modeling

We applied quantile regression forests (QRF) introduced by Meinshausen (2006), an extension of the random forests (RF) algorithm. RF is an ensemble learning method that builds a set of regression trees, and the final prediction is the average of all the regression trees, which are evaluated using out-of-bag cross-validation (Breiman, 2001). The QRF algorithm estimates the full conditional distribution of the response variable as a function of its predictors, not just the mean as with the original RF algorithm. Therefore, it is possible to extract the prediction interval for each pixel across the landscape for each measurement period. We followed three steps to develop models for predicting soil CH₄ fluxes at each measurement period. We used the five terrain attributes (slope, PrC, TPI at 30-m radius, SWI, and VDCN at 5-ha initiation threshold), basal area (BA), and relative basal area of coniferous trees to BA (RBAcon) as predictors. Unfortunately, the machine learning model was unable to accurately reproduce the measured fluxes when wetland measurements were included in the training dataset, likely due to the imbalance between the 52 upland and only 3 wetland sampling points. The comparison of models including and not including wetland data is shown in Table A2. Only permanent wetlands in the plain area were excluded (3 of 55 collars for data, less than 1% of the landscape pixels). Wet patches, which had temporarily water-saturated soils, were not excluded.

Before applying QRFs, we eliminated the less important variables and identified the most relevant predictors for each measurement date, using a variable selection algorithm for random forest models proposed by Genuer et al. (2010) and implemented in the “VSURF” package for R (Genuer et al., 2015). This approach systematically uses a repeated cross-validation procedure to rank variables by their importance index and iteratively eliminates the least informative ones to

minimize model error. The result is a refined subset of predictors that enhances model interpretation and predictive performance. The predictor reduction approach has previously been used to map CH₄ fluxes (Räsänen et al., 2021; Warner et al., 2019) and soil properties (Jeong et al., 2017; Miller et al., 2015).

After selecting the relevant predictor variables, the QRF models were trained to predict CH₄ fluxes using the R-packages “caret” (Kuhn and Johnson, 2013) and “quantregForest” (Meinshausen, 2017). The mtry parameter, which determines the number of randomly selected predictor variables at each node, was tested from 2 to n-1 (n being the total number of predictors) using leave-one-out cross-validation to minimize prediction error and maximize the variance explained by the model. The ntree parameter was set to 500, ensuring the model constructed an ensemble of 500 decision trees. Furthermore, we calculated the variable's importance scores using the “vip” R-package (Greenwell and Boehmke, 2020). Variable importance scores were estimated using a permutation-based approach, in which the values of each predictor in the training data were randomly permuted to assess the resulting change in model performance, as quantified by the adjusted R-squared value. A greater reduction in adjusted R² indicated a higher importance of the predictor variable. We generated the accumulated local effect (ALE) plots to visualize the response of CH₄ fluxes to the predictor variables, accounting for the effect of the predictors in the model (Apley and Zhu, 2020). In ALE plots, an ALE value of zero on the y-axis corresponds to the mean predicted CH₄ flux, with positive values indicating higher and negative values indicating lower flux under the specific predictor on the x-axis. ALE reduces a complex machine learning function to depend on only one or, in some cases, two input variables, and visualizes the effect of a selected variable on the predicted CH₄ flux. The method removes the confounding effects of other input variables, computes the partial derivatives (local effects) of the prediction function with respect to the variable of interest, and integrates (accumulates) these effects across the range of that variable.

For each of the nine measurement dates, model accuracy was evaluated based on the root mean square error (RMSE) and coefficient of determination (R²). The output of the QRFs was a set of conditional prediction distributions of CH₄ fluxes for each landscape pixel and measurement dates. Because these prediction distributions were often not normally distributed, the median of the conditional prediction distribution at each pixel was used as the final prediction, and the interquartile range of the distribution was used to quantify the uncertainty in the prediction (Warner et al., 2019). Prediction uncertainties were expressed as a percentage (i.e., interquartile range of the conditional prediction distribution divided by the median). Modelling was conducted independently for each of the nine measurement dates, without including meteorological data, as in previous studies (Vainio et al., 2021; Warner et al., 2019). Indeed, since all pixels would exhibit the same values for meteorological variables at a given date, it would be pointless to include them in RF models to predict the spatial heterogeneity of fluxes at the landscape scale.

2.8 Statistical analysis

We used analysis of variance (ANOVA) to test the differences in soil properties across the topographic positions and vegetation types and densities. Interactions were not included because the model would be rank-deficient as there were no “pure” broadleaved plots on the ridge. We examined the relationships between soil properties and topographic and vegetation variables using Spearman's rank correlation analysis using the ‘Hmisc’ package (Harrell Jr, 2003). Linear mixed-effect models (LMM) were used to test the relationship between the predicted fluxes at pixel levels and measured fluxes (fixed effect), with flux measurement dates as a random effect and between the predicted soil CH₄ fluxes and measured soil CH₄ fluxes aggregated by landscape units (topographical position, vegetation types, and vegetation density), which were included as the random factor on both slope and intercept. The root mean square error (RMSE) was used to evaluate model performance at each date, and the marginal and conditional coefficients of the determination (R_m² and R_c²) were used to determine the strength of the relationship between the predicted and measured fluxes. LMM was carried out using the ‘lmerTest’ package (Bates et al., 2015; Kuznetsova et al., 2017), and R_m² and R_c² were calculated using the ‘MUMIn’ package (Bartoń, 2010). To test the effects of topographic positions, vegetation types, and densities on predicted CH₄ fluxes while accounting for spatial autocorrelation, we

280 also used a linear mixed-effect model. Topographic positions, vegetation types, and densities were included in the model as
fixed effects, and pixel ID as a random effect. Interactions were not included as no pixel contains “pure” broadleaved vegetation
on the ridge. To eliminate spatial autocorrelation among residuals, we incorporated an exponential spatial correlation structure
based on each pixel coordinate nested within each measurement date. This was performed using the ‘nlme’ package (Pinheiro
et al., 1999). The semi-variogram of the residuals confirmed that the residuals were not spatially correlated. To quantify the
285 effect size that indicates the relative contribution of each factor to the total variance in the response variable, we calculated
eta-squared (η_p^2) values using the ‘effectsize’ package (Ben-Shachar et al., 2019). A pairwise comparison across the topographic
positions, vegetation types, and densities was performed using the ‘emmeans’ package (Lenth, 2017). Linear regression models
290 were used to examine the relationship between landscape-scaled predicted soil CH₄ fluxes and API. The recession coefficient
(k) and the number of antecedent days (n) were not fixed a priori but optimized to maximize R² while ensuring the best
distribution of the residuals, allowing parameters k and n to vary iteratively from 0.6 to 0.9 with an increment of 0.01 and from
0 to 30 with an increment of 0.01, respectively. Using a more complex bivariate model with an exponential function of air
temperature did not improve the quality of the fit and returned Q₁₀ values that were not significantly different from 1, as
previously reported (Epron et al., 2016). Calculations, modelling, and statistical analyses were performed using the R statistical
programming environment (R Core Team, 2024).

295 **3 Results**

3.1 Environmental conditions and soil properties across upland topographic and vegetation features

The total rainfall in the study area during the snow-free period of 2023 was 1578.5 mm, with relatively high rainfall in late-
May to mid-June and a peak on August 15 due to the typhoon Lan (Fig. 2a). The monthly mean air temperature ranged from
7.5 to 24.2 °C during the study period (Fig. 2b). Mean soil moisture content varied seasonally, with the highest (47.7 ± 1.1 %;
300 mean ± standard error) observed in the early summer (07/06) and the lowest (32.9 ± 1.2 %) in the late summer (09/04) (Fig.
2c). Mean soil temperature followed a similar trend to air temperature across the study period (Fig. 2d). Upland soils
consistently uptake CH₄ (negative fluxes, Fig. 2e), while soils in the three small wetland patches emitted CH₄ (positive flux,
Fig. A1). Variation in upland CH₄ fluxes across the measurement dates was consistent with the seasonal patterns of rainfall
and air temperature. The fluxes measured on two collars located in the upland area, but in wet patches, were positive on one
305 occasion each.




Figure 2: Seasonal variation in (a) daily rainfall and (b) daily air temperature from April to November in 2023, and (c) mean soil moisture content, (d) mean soil temperature, and (e) mean soil CH₄ fluxes from upland sampling points, including all topographic positions (n = 52). Vertical bar indicating the standard error.

310

Topographic positions significantly influenced several soil properties (bulk density, pH, total carbon and nitrogen, and mean temperature), whereas vegetation type and vegetation density significantly influenced soil temperature and soil moisture, respectively (Table 1). The bulk density of the fine earth fraction was relatively low due to the presence of stones, highest in the plain ($0.42 \pm 0.04 \text{ g cm}^{-3}$, mean \pm SE) and significantly lowest in the ridge ($0.26 \pm 0.04 \text{ g cm}^{-3}$). Soil pH differed significantly across topographic positions ($p < 0.001$), with more acidic conditions observed at higher elevations (ridge: 4.0 ± 0.14) compared to the plain (5.1 ± 0.73). Similarly, total carbon (C) and total nitrogen contents (N) were significantly higher

315

on the ridges ($16.7 \pm 2.2\%$ C and $0.8 \pm 0.10\%$ N) and lower in the plain ($5.1 \pm 0.73\%$ C and $0.3 \pm 0.04\%$ N). Mean soil temperature also significantly varied across topographic positions, with the highest in the plain ($14.3 \pm 0.18^\circ\text{C}$) and the lowest in the foot slope ($13.4 \pm 22^\circ\text{C}$). In contrast, the soil texture of the fine earth fraction (clay, silt, and sand) and mean soil moisture content did not vary significantly with topographic positions.

Vegetation type significantly influenced soil temperature, with the highest values observed under broadleaved stands and the lowest under coniferous stands. Vegetation density significantly affected soil moisture content, which was highest in low-density areas and lowest in medium-density areas.

Table 1. Mean (\pm standard error) of soil bulk density (BD), texture (clay, silt and sand), total carbon (C) and nitrogen (N) concentration, pH, mean soil water content (SWC) and temperature (T_{soil}) according to topographic position, vegetation type, and vegetation density. SWC and T_{soil} are the average of the 9 measurement dates. Different lowercase letters indicate significant differences among topographic positions, vegetation types, and vegetation densities ($p < 0.05$). The p-values of the ANOVA are shown in the last rows. The number of independent replicates in each factor level is indicated in the first column.

Factor	Bulk density (g cm^{-3})	Clay (%)	Silt (%)	Sand (%)	pH	Total C (%)	Total N (%)	Mean SWC (%)	Mean T_{soil} ($^\circ\text{C}$)
Position									
Plain (n=14)	0.42 ± 0.04 a	8 ± 2	27 ± 3	65 \pm 4	4.9 ± 0.1 a	5.1 ± 0.7 a	0.3 ± 0.0 a	44.2 ± 2.0	14.3 ± 0.2 a
Foot slope (n=9)	0.36 ± 0.06 ab	8 ± 2	33 ± 4	60 ± 5	4.3 ± 0.1 b	8.3 ± 2.2 a	0.5 ± 0.1 ab	42.1 ± 2.0	13.4 ± 0.2 b
Slope (n=16)	0.29 ± 0.03 ab	10 ± 2	29 ± 3	60 ± 4	4.3 ± 0.1 b	10.0 ± 1.1 a	0.6 ± 0.1 ab	39.2 ± 1.4	13.9 ± 0.1 a
Ridge (n=13)	0.26 ± 0.04 b	9 ± 2	23 ± 4	68 ± 4	4.0 ± 0.1 b	16.7 ± 2.2 b	0.8 ± 0.1 b	40.1 ± 2.0	13.9 ± 0.1 ab
Vegetation type									
Broadleaved (n=19)	0.38 ± 0.04	10 ± 2	29 ± 3	61 ± 4	4.7 ± 0.1	6.8 ± 1.0	0.4 ± 0.1	43.3 ± 1.6	14.2 ± 0.1 a
Coniferous (n=11)	0.32 ± 0.04	7 ± 1	27 ± 4	66 ± 5	4.3 ± 0.1	11.1 ± 2.4	0.6 ± 0.1	40.8 ± 1.7	13.4 ± 0.2 b
Mixed (n=22)	0.29 ± 0.03	9 ± 1	26 ± 2	64 ± 3	4.2 ± 0.1	12.4 ± 1.6	0.6 ± 0.1	39.7 ± 1.3	13.9 ± 0.1 a
Vegetation density									
High (n=14)	0.32 ± 0.04	9 ± 2	28 ± 3	64 ± 4	4.4 ± 0.1	10.6 ± 2.02	0.57 ± 0.1	40.4 ± 0.1 a	13.7 ± 0.2
Medium (n=28)	0.30 ± 0.03	10 ± 1	28 ± 3	62 ± 3	4.3 ± 0.1	11.3 ± 1.34	0.59 ± 0.1	39.5 ± 1.0 a	13.9 ± 0.1
Low (n=10)	0.43 ± 0.05	8 ± 2	26 ± 3	66 ± 4	4.7 ± 0.2	6.1 ± 1.27	0.40 ± 0.1	47.6 ± 2.1 b	14.2 ± 0.2
Anova results									
Position	p < 0.05	p = 0.55	p = 0.33	p = 0.49	p < 0.001	p < 0.001	p < 0.01	p = 0.12	p < 0.001
Vegetation type	p = 0.81	p = 0.30	p = 0.92	p = 0.66	p = 0.75	p = 0.90	p = 0.99	p = 0.52	p < 0.001
Vegetation density	p = 0.58	p = 0.45	p = 0.58	p = 0.43	p = 0.99	p = 0.93	p = 0.98	p < 0.05	p = 0.91


3.2 Selected variables and performance of the upland CH₄ flux models

The topographic position index (TPI) was consistently selected in all seasons, with high importance scores, ranging from 0.54 to 0.88, depending on the measurement dates (Table 2). The saga wetness index (SWI) was selected for most measurement dates, except for two, where the vertical distance to the channel network (VDCN) was selected instead. SWI importance scores were higher in summer than in the other seasons. VDCN and profile curvature (PrC) were occasionally selected along with TPI and TWI. VDCN showed moderate to low importance scores, contributing mostly in mid-spring (0.66) and early autumn (0.58). PrC, although less consistently selected, played a role in specific seasons, particularly in early spring and mid to late autumn. **Accumulated local effect (ALE) plots showed the direction of the variables' effects on soil CH₄ fluxes for each measurement date (Fig. A2).** For the two most influential predictors, low CH₄ uptake rates were associated with low TPI values, while they were associated with high SWI values. The vegetation density (BA) was selected only on two dates in 2023/04/27 and 2023/10/07, without improving the model accuracy, so we did not include it the final models (Appendix Table A3).

Table 2. Selected variables for the quantile regression forest (QRF) models at each measurement date, along with the R² and root mean square error (RMSE) values to evaluate the accuracy of the models. Importance scores of the selected variables are shown in parentheses, indicating their contribution to predicting soil CH₄ fluxes.

Measurement dates	Selected variables	R ²	RMSE (nmol m ⁻² s ⁻¹)
2023/04/27	SWI (0.57), TPI (0.67), PrC (0.58)	0.53	0.52
2023/05/12	TPI (0.80), VDCN (0.66)	0.31	0.82
2023/05/31	SWI (0.55), TPI (0.57), VDCN (0.42)	0.43	0.48
2023/07/06	SWI (0.73), TPI (0.60)	0.40	0.50
2023/07/26	SWI (0.80), TPI (0.69)	0.37	1.02
2023/09/04	SWI (0.74), TPI (0.85)	0.40	1.18
2023/10/07	TPI (0.88), VDCN (0.58)	0.67	0.81
2023/11/07	SWI (0.32), TPI (0.84), VDCN (0.12), PrC (0.45)	0.59	0.66
2023/11/30	SWI (0.32), TPI (0.54), VDCN (0.21), PrC (0.28)	0.51	0.56

Model accuracy showed seasonal variation, with the highest obtained in early autumn ($R^2 = 0.67$; RMSE = 0.81 nmol m⁻² s⁻¹) and the lowest in mid-spring ($R^2 = 0.31$; RMSE = 0.82 nmol m⁻² s⁻¹; Table 2). The relationship between measured and predicted fluxes for each measurement date showed that estimated fluxes were close to the observed fluxes (Fig. 3a-i).



355

Figure 3: Comparison of predicted (median of the quartile predictions from QRFs) and measured CH₄ fluxes for each measurement date. Vertical bars indicate the interquartile ranges of the prediction distribution. Intercepts and slopes are estimated using a linear mixed-effect model with measurement dates as a random effect (full statistics are shown in Table A4). The diagonals are the identity (1:1) lines.

360

Overall, the slope of the relationship between measured and predicted fluxes (fixed effects) was not significantly different from 1 and was similar at all dates. The marginal (R^2_m) and conditional (R^2_c) coefficients of determination were 0.93 and 0.94, respectively, highlighting the consistency of the prediction for all measurement dates (linear mixed model, Table A4). To validate the fluxes at the landscape level, predicted fluxes were aggregated by landscape unit (i.e., topographic position, vegetation type, and vegetation density) and compared with the aggregated measured fluxes, which were consistent with the measured fluxes (Fig. 4, Table A5).




Figure 4: Comparison between aggregated mean predicted and measured CH₄ fluxes for (a) topographic position, (b) vegetation type, and (c) vegetation density (full statistics are shown in Table A5). The diagonals are the identity (1:1) lines.


370

3.3 Predicted upland soil CH₄ fluxes

We predicted that upland soils consistently uptake CH₄ across the seasons (negative fluxes, Fig. 5). Predicted median CH₄ fluxes showed significant spatial heterogeneity, which was consistent across the seasons (Fig. 5). The highest net CH₄ uptake was predicted on ridges and the steepest parts of the slopes and decreased toward the foot slopes near streams and the flat plain

375

(Fig. 6a). Coniferous and mixed stands showed the highest uptake compared to the broadleaved stands (Fig. 6b). **Vegetation density (BA)** also influenced the soil CH₄ uptake with higher uptake in the high and medium density areas. Although substantial variation was observed within each landscape unit, topographic position exerted the strongest control on CH₄ fluxes ($\eta_p^2 = 0.43$), followed by vegetation density ($\eta_p^2 = 0.11$) and vegetation type ($\eta_p^2 = 0.006$) (Table A6).



380

Figure 5: Maps of predicted soil CH_4 fluxes at each pixel of the study area for each measurement date. Values represent the median of the conditional prediction distribution for each pixel.




Figure 6: Predicted landscape-scale soil CH₄ fluxes averaged over all measurement dates, aggregated by (a) upland topographic positions, (b) vegetation type, and (c) vegetation density (Full statistics with significance of the differences between each landscape unit shown in Table A6). Green squares indicate the mean of the measured fluxes with standard errors, blue triangles indicate the mean of the predicted fluxes at the landscape level and grey boxplots indicate the distribution of the predicted fluxes.

3.4 Uncertainty of predicted upland soil CH₄ fluxes

The spatial distribution of the percentage of predicted uncertainty varied across seasons (Fig. 7). The percentage was consistently low to moderate (less than 100%) for pixels on ridges and steep slopes, but extremely high uncertainties (more than 500%) was observed at some dates for low-elevation pixels when predicted fluxes were close to zero. However, low predicted fluxes were often associated with equally low predicted uncertainty (Fig A3). The proportion of pixels with low uncertainty (<50%) was highest in early autumn (39.7% of the total upland pixels) and lowest in early spring (5.7% of the total upland pixels). In contrast, moderate uncertainty (50-100%) was predominant in most seasons, particularly in spring and autumn, accounting for approximately 50% of the landscape. Moderate to high uncertainty (101-500%) was also predominant at some measurement dates, reaching its highest contribution to the landscape in late spring (49.8%). Extreme uncertainty (>500%) was very rare in all seasons, except for a small peak in late autumn (0.26%, Table A7).




Figure 7. Uncertainty map of predicted soil CH₄ fluxes at each pixel of the study area for each measurement date. Values represent the ratio of the interquartile range to the median of the prediction distribution for each pixel.

3.5 Predicted seasonal upland fluxes at the landscape level

The predicted upland CH₄ flux per hectare was calculated as the sum of the predicted fluxes at each pixel multiplied by pixel area (25 m²), and the sum divided by the upland area. Across the landscape, the average CH₄ uptake by upland soils during the snow-free season was -0.66 (interquartile range: -0.94 to -0.44) g CH₄ ha⁻¹ hr⁻¹. Predicted median seasonal fluxes ranged from -0.34 to -0.60 g CH₄ ha⁻¹ hr⁻¹ in spring, from -0.39 to -1.28 g CH₄ ha⁻¹ hr⁻¹ in summer, and from -0.48 to -0.89 g CH₄ ha⁻¹ hr⁻¹ in autumn (Fig. 8a). CH₄ uptake was low across the landscape in early (April 27) and late spring (May 31), while higher uptake was predicted in mid-spring (May 12). CH₄ uptake remained low in the early wet summer (July 6) and increased toward the mid (July 26) to late dry summer (Sep 4) when it reached its peaks. Net CH₄ uptake then decreased from early autumn (Oct 7) and reached its lowest rate in late autumn (Nov 30).




Figure 8: Predicted soil CH_4 fluxes, calculated as the mean of all pixels in the study area, and antecedent precipitation index (API). (a) Seasonal variations in landscape-scaled predicted soil CH_4 fluxes and (b) relationship between landscape-scaled predicted soil CH_4 fluxes and the 20-day API. Vertical bars indicate the uncertainty of the predicted fluxes.

415

This seasonal variation in predicted upland median fluxes was well explained by the 20-day antecedent precipitation index ($R^2 = 0.70, p < 0.01$) with a recession coefficient of 0.69 (Fig. 8b), followed closely by the 30-day ($R^2 = 0.69$) and 7-day ($R^2 = 0.68$) API (Table A8).

4. Discussion

420

4.1 Selected variables

We employed quantile regression forest (QRF) models, **driven only by topographic attributes**, to upscale *in-situ* soil CH_4 flux measurements from sampling points to the landscape level for each measurement date in all upland topographic positions, but excluding wetlands (1% of our study area). **Although selected twice, the inclusion of BA did not improve the accuracy and performance of the model and was eventually not retained in any final models, while RBA_{CON} was never selected.**

425

Among all tested topographic variables derived from the DEM, SWI, TPI, PrC, and VDCN were consistently selected in different models across all measurement periods, emphasizing their importance in upscaling CH_4 fluxes. Overall, the results validated our first hypothesis, as the selected topographic attributes were related to water circulation and accumulation.

430

Among these variables, SWI, **which positively influence CH_4 fluxes (low uptake in areas with high TWI)**, represents water accumulation potential and is a common surrogate for soil moisture in mountainous regions. This key factor controls CH_4 fluxes by affecting gas diffusion and microbial activity (Kaiser et al., 2018; Vainio et al., 2021; Warner et al., 2019), as SWI integrates potential inflows and discharges through runoff and drainage (Ågren et al., 2014; Beven and Kirkby, 1979). SWI was selected in seven out of nine measurement periods but not on May 12 and October 7. These two periods correspond to transitional seasons, i.e., mid-spring and early autumn, when the landscape is generally drier, and water does not accumulate. TPI describes the elevation of a location relative to those of the surrounding terrain within a given radius, allowing the identification of landform positions such as ridges, slopes, and valleys (Ågren et al., 2014). TPI is generally calculated using a non-filled DEM, which is also more representative of local-scale moist depression that SWI doesn't capture, as SWI is

calculated using the filled DEM (Kemppinen et al., 2018). In our study, TPI was consistently selected in all measurement periods, and clearly related, highlighting that localized moisture, and potentially soil chemistry, are more influential parameters in controlling the CH₄ fluxes at the landscape level. Areas with negative TPI values (e.g., valleys or depressions) typically function as convergence zones, where water and nutrients accumulate due to gravitational flow and reduced drainage. In contrast, positive TPI values (e.g., ridges and convex upper slopes) are more divergent, often characterized by increased drainage and runoff, and limited water and nutrient retention. **TPI negatively affected soil CH₄ fluxes (high uptake in areas with high TPI).**

PrC refers to the curvature of the land surface in the direction of the slope (along a flow line) which was selected three times (April 27, Nov 7 and Nov 30) across the measurement dates. It influences the acceleration or deceleration of surface and subsurface water flow (Ågren et al., 2014). Negative values (concave slopes) tend to slow water movement, promoting water and nutrient accumulation in soils. Conversely, positive values (convex slopes) accelerate flow, often reducing water retention time and lowering nutrient accumulation due to leaching or erosion. Excluding PrC from the list of available variables for selection decreased the model performance for these three dates, probably because PrC helps discriminate between plains and slopes, both of which have near-zero TPI values.

VDCN is another important variable reflecting groundwater level conditions. Lower values typically observed near stream channels with higher groundwater level (Bock and Köthe, 2008). When the landscape was drier (May 12 and October 7), and SWI was not selected, TPI and VDCN had more substantial explanatory power. VDCN was also selected several times with SWI. Interestingly, VDCN has been shown to be useful in distinguishing well-drained from poorly drained soils (Bell et al., 1992; Kravchenko et al., 2002). It may explain why excluding VDCN from the list of variables available for selection decreased model performance. This highlights that SWI and TPI alone were not sufficient to reflect local soil moisture conditions, as drainage conditions can potentially vary across the landscape, which controls soil microhabitat conditions and thus influences CH₄ fluxes.

QRF modelling is non-parametric machine learning approach is particularly suited for handling non-linear relationships and complex interactions among predictors (Meinshausen, 2006). However, although the topographic predictors have successfully predicted CH₄ fluxes, the QRF method, like other statistical methods, does not provide a mechanistic understanding of the underlying biogeochemical processes, and the existence of confounding factors cannot be ruled out.

4.2 Model performance and uncertainty

Soil CH₄ fluxes predicted by QRF models were close to the measured fluxes for all measurement periods (Fig. 3; Table A4). We recognize that our models, by forcing pixel-scale predictors (5 m resolution) to explain finer-scale chamber measurements (20 cm diameter), may actually overestimated the predictive accuracy of the models at coarser scales. However, the predicted soil CH₄ fluxes not only closely matched the individual measured fluxes, but also when the two were aggregated by topographic position classes (ridge, slope, foot slope, and plain), vegetation density classes (high, medium, and low) or vegetation type classes (coniferous, mixed, and broadleaf, Fig. 4; Table A5). This confirmed that our models did not only predict point-level flux heterogeneity but was also able to capture the landscape-scale flux patterns and indicated that topographic attributes could be used for upscaling CH₄ fluxes in mountainous landscapes. Overall, the performance of the models developed for scaling CH₄ fluxes was comparable to previous studies using topographic data for similar purposes (Kaiser et al., 2018; Vainio et al., 2021; Virkkala et al., 2024; Warner et al., 2019). However, it is important to note that direct comparisons between studies are difficult due to variations in cross-validation approaches, as the choice of cross-validation technique can significantly influence model performance (Roberts et al., 2017).

Unfortunately, it was not possible to accurately predict CH₄ fluxes when measurements collected in wetland patches were included in the training data, as the model accuracy decreased at all dates (Table A9). As a consequence, the marginal and conditional coefficients of determination of the relationship between the predicted and measured fluxes decreased from 0.93

480 and 0.94 respectively to 0.70 when wetland data were included. This is probably because neither the topographic features nor
481 the vegetation differed sufficiently between the large areas functioning as CH₄ sinks and the small wetland patches in the plain
482 area functioning as CH₄ sources (Fig. A1). Räsänen et al. (2021) noticed that spatial patterns of CH₄ fluxes could be accurately
483 predicted in a northern peatland-forest-mosaic landscape when they were modelled for sinks and sources separately. This
484 separation was not possible in our study due to the low number of measurement locations in wetlands, related to their small
485 extent (1%) in our upland-dominated landscape. **Wetland exclusion, although acceptable in our 40-ha study area due to their**
small extent, would overestimate CH₄ uptake if incorrectly applied at larger scales, i.e., to the entire upper Yura River
catchment in our case or to other hydrologically complex forest landscapes.

486 One advantage of the QRF approach is its ability to estimate prediction intervals (Meinshausen, 2006), thus offering insights
487 into the uncertainty associated with the predicted flux value at each pixel. The spatial distribution of the uncertainty associated
488 with the predicted soil CH₄ fluxes varied seasonally (Fig. 7; Table A7) in agreement with our second hypothesis, reflecting
489 both spatial heterogeneity and temporal changes in model confidence. In our study, the spatial patterns of QRF-derived
490 uncertainties were consistently related to topographic position and flux magnitude. Predictions in ridge and steep slope pixels
491 generally exhibited low percentage uncertainties (often below 100%), likely because these well-drained upland areas were
492 well represented in the training data and exhibited relatively stable and high CH₄ uptake across seasons. In contrast, extremely
493 high percentage uncertainties (exceeding 500%) were observed in some low-lying pixels during specific seasons, especially
494 where predicted CH₄ fluxes were close to zero. **Our models did not predict median positive fluxes although positive fluxes**
were occasionally measured. However, the possibility of positive fluxes is reflected in the large uncertainties associated with
near-zero fluxes. A crucial methodological point is that percentage uncertainty is a relative measure; even a small absolute
495 uncertainty around a near-zero prediction can yield a very large percentage (Warner et al., 2019). In addition, large absolute
496 uncertainties can result from large differences in fluxes measured at locations with similar topographic characteristics.

500 The lowest uncertainty was obtained in late summer and early autumn, i.e., under warm and dry conditions, indicating better
501 model performance when hydrological conditions were less variable. In contrast, larger uncertainties were produced by the
502 models in early spring and late autumn, as well as in late spring and early summer, when measured and predicted soil CH₄
503 fluxes were lowest. The East Asian monsoon flow bringing warm and humid air mass and resulting in the rainy season in late
504 spring and early summer, as well as low evapotranspiration in early spring and late autumn, may have introduced greater
505 variability in soil hydrology, contributing to higher uncertainties. Nevertheless, low to moderate uncertainty (<100%) was the
506 most prevalent class across all seasons, consistently accounting for more than half the landscape—up to 80% in late summer
507 and early autumn—while extreme uncertainties (>500%) were very rare in all seasons. This suggests that the models performed
508 well overall. Although some areas remain challenging to model, the QRF approach provides generally reliable spatial
509 predictions of soil CH₄ fluxes with quantifiable and interpretable uncertainties.

510 **4.3 Spatial patterns of predicted soil CH₄ fluxes**

511 The models revealed clear spatial patterns in soil CH₄ fluxes that were consistent across measurement dates, even though the
512 models selected different variables at each date. Predicted soil CH₄ fluxes closely matched topographic gradients, consistent
513 with our third hypothesis. Ridges and upper slopes exhibited the highest net CH₄ uptake, functioning as strong sinks for CH₄
514 across all seasons, whereas CH₄ uptakes were lowest in plain and foot slope positions. These topographic patterns of CH₄
515 uptake are consistent with previous studies. In a temperate forest in central Ontario, Canada, the highest CH₄ uptake was
516 observed on slopes and ridges (Wang et al., 2013). Similarly, in a temperate forest in Maryland, USA, transition zones were
517 identified as hotspots for CH₄ uptake (Warner et al., 2018). In a tropical forest in China, hillslopes exhibited the highest CH₄
518 uptake, while lower uptake was observed at the foot slopes and in groundwater discharge areas (Yu et al., 2021). Similarly,
519 CH₄ uptake was greater on ridges than at valley bottoms in a subtropical forest in Puerto Rico (Quebbeman et al., 2022).

520 In our studied landscape, we observed lower soil bulk density on ridges and slopes than on the plain area, indicating that ridge
and slope soils have higher porosity, which is consistent with higher soil CH₄ oxidation rates due to higher diffusion rates of
O₂ and CH₄ from the atmosphere through soil pores (Ishizuka et al., 2009). Although we did not assess the methanotroph
community structure, the greater atmospheric CH₄ uptake on slopes and ridges is consistent with the community structure
observed in a subalpine forest, with type I methanotrophs dominating in riparian soils, whereas type II methanotrophs were
525 more prevalent in upland soils (Du et al., 2015). The higher soil carbon (C) and nitrogen (N) contents observed on ridges and
slopes at our site may contribute to higher soil CH₄ uptake, as soil CH₄ uptake has been found to be positively correlated with
soil organic matter content in subtropical and temperate forests (Lee et al., 2023). Possible explanations are that higher soil
carbon may increase the availability of labile substrates that stimulate methanotrophic activity by increasing CH₄ supply
530 through enhanced methanogenesis in anoxic microsites or by directly providing substrate for facultative methane-oxidizing
bacteria, thereby increasing their abundance (Jensen et al., 1998; Semrau et al., 2011; West and Schmidt, 1999). Soil nitrogen
was probably predominantly in organic form, and therefore the soil concentration of nitrate and ammonium, known to inhibit
CH₄ oxidation by methanotrophs at high concentration (King and Schnell, 1994; Mochizuki et al., 2012), likely remained low
(Aronson and Helliker, 2010; Bodelier and Laanbroek, 2004). Nitrogen is an essential nutrient for the growth of methanotrophs,
535 whose activity has been shown to be nitrogen-limited in forest soils (Börjesson and Nohrstedt, 2000; Martinson et al., 2021;
Veldkamp et al., 2013). Therefore, mineralization of these low levels of organic nitrogen could alleviate the nitrogen limitation
of CH₄ oxidation and partly explain the higher soil CH₄ uptake observed on ridges and slopes, where total nitrogen
concentration was higher than at the foot slopes and in the plain.

540 Although the effect-size of vegetation density was much smaller than that of topographic position, the predicted soil CH₄
uptake was significantly lower in areas with low basal area. Vegetation density can also potentially be related to local moisture
conditions, as dense vegetation likely consume more water, thus increasing the soil air-filled porosity (Hakamada et al., 2020;
Vanclay, 2009). Unexpectedly, although a very small effect-size, our models predicted higher soil CH₄ uptake in conifer-
545 dominated areas and lower uptake in broadleaf-dominated areas, contrary to previous evidence of greater soil CH₄ uptake in
plots containing only deciduous broadleaved tree species than in plots containing evergreen coniferous trees, either alone or in
mixture (Jevon et al., 2023). The discrepancy between this previous study and our results may be related to the fact that their
study area was ten times smaller and more topographically homogeneous than ours (4 *versus* 40 ha). Moreover, soil properties
550 that could explain the lower rate of CH₄ oxidation in coniferous than in broadleaved stands, such as higher acidity (Borken et
al., 2003; Hütsch, 1998; Ishizuka et al., 2000) did not differ significantly among the three types of vegetation cover at our site,
whereas they differed according to topographic position. However, vegetation types and density were not randomly distributed
among topographic positions (Table A9), meaning that the confounding effects of vegetation and DEM-derived variables on
555 the prediction soil CH₄ uptake could make it difficult to separate the influence of vegetation and topography in our complex
mountain landscape.

4.4 Landscape-scaled soil CH₄ fluxes and seasonal variation

555 The upland CH₄ fluxes per hectare were calculated by aggregating pixel-level predictions and normalizing them to the total
upland area, allowing for standardized comparison across sites, although there are still very few comparable data available,
making it difficult to analyse the causes of differences across sites. Our highest CH₄ uptake in late summer was -1.28 g CH₄
ha⁻¹ hr⁻¹ (interquartile range -1.70 to -0.89), 2.6 times higher in absolute value than in a forested watershed in Maryland, USA
(-0.47 g CH₄ ha⁻¹ hr⁻¹, Warner et al. 2019), but slightly lower than in a boreal pine forest in Finland (-1.59 g CH₄ ha⁻¹ hr⁻¹,
Vainio et al. 2021).

560 Consistent with our fourth hypothesis, the seasonal variation in landscape-scaled soil CH₄ fluxes in the upland areas reflects
strong sensitivity to soil moisture dynamics, which were effectively captured using the Antecedent Precipitation Index (API).
The API, serving as a proxy for soil moisture dynamics, integrates precipitation over a defined period and includes a recession

factor to account for evapotranspiration and drainage. Short durations (e.g., 7 days) reflect surface moisture, while longer durations (e.g., 30 days) capture deeper soil moisture conditions (Schoener and Stone, 2020; Sidle et al., 2000; Yamao et al., 2016). Among the API durations tested, the 20-day API with a recession coefficient of 0.69 showed the highest explanatory power ($R^2 = 0.70$), although using either a 30-day or a 7-day API would provide similar goodness of fit with similar recession coefficients, indicating that soil moisture conditions across different depths had similar influence on CH_4 flux variability. The consistently low recession coefficient (Kohler and Linsley, 1951) suggested that rainwater does not accumulate in our watershed. High API values indicate wetter antecedent conditions, which can suppress CH_4 uptake by reducing oxygen availability and thus limiting methanotrophic activity, and by temporarily turning the subsoil condition to anoxic, promoting methane production and reducing net CH_4 uptake (Angel et al., 2012; Hu et al., 2023; Kruse et al., 1996). Conversely, drier periods with low API values were observed in mid and late summer and earlier autumn, when soils were better aerated, creating favourable conditions for atmospheric CH_4 oxidation and leading to greater CH_4 uptake.

5 Conclusion

In conclusion, our study showed the dominant role of topography, **compared to that of vegetation**, on the spatial variation of soil CH_4 fluxes in upland forest landscapes **throughout the snow-free season**. The quantile regression forest models successfully captured these ridge-to-plain spatial gradients in the upland area where the soil is almost always unsaturated, with strong performance. However, our modelling approach was unable to accurately predict CH_4 fluxes when including measurements collected in three wetland patches functioning as CH_4 sources in the plain area (1% of the total landscape). CH_4 uptake was consistently highest on ridges and slopes, where well-drained soils with lower bulk density and higher porosity supported enhanced methanotrophic activity. Furthermore, the seasonal dynamics of the **landscape-scaled predicted** soil CH_4 flux was well-captured by the 20-day Antecedent Precipitation Index (API), with a significant positive relationship between API and CH_4 uptake, emphasizing the sensitivity of CH_4 uptake by upland soils to seasonal fluctuations in soil moisture conditions. The integration of terrain-based predictors and moisture history provides a reliable framework for scaling soil CH_4 fluxes across complex landscapes, highlighting the importance of considering both static (topography, vegetation) and dynamic (climate) controls in future assessments of CH_4 flux.

Appendix A

Table A1. Spearman's rank correlation test between soil properties and topographic and vegetation attributes. Significant coefficients are shown in bold and italic.

	Mean SWC	Mean T _{soil}	Total C	Total N	pH	BD	Sand	Silt	Clay
PrC	-0.21	0.14	<u>0.27</u>	0.23	-0.27	-0.14	0.24	-0.26	-0.06
Slope	-0.17	-0.24	0.25	0.22	<u>-0.51</u>	-0.11	0.00	-0.02	0.01
TPI 20m	<u>-0.35</u>	-0.11	<u>0.45</u>	<u>0.39</u>	<u>-0.34</u>	-0.26	0.05	-0.04	0.03
TPI 30m	<u>-0.31</u>	-0.08	<u>0.64</u>	<u>0.55</u>	<u>-0.46</u>	<u>-0.36</u>	0.11	-0.14	0.03
TPI 50m	<u>-0.33</u>	-0.08	<u>0.59</u>	<u>0.50</u>	<u>-0.42</u>	<u>-0.31</u>	0.11	-0.13	0.02
SWI	<u>0.43</u>	0.27	<u>-0.62</u>	<u>-0.55</u>	<u>0.65</u>	<u>0.41</u>	-0.03	0.04	-0.01
VDCN 0.5ha	-0.24	-0.16	<u>0.61</u>	<u>0.52</u>	<u>-0.51</u>	<u>-0.36</u>	0.16	-0.20	0.04
VDCN 2.5ha	-0.24	-0.20	<u>0.71</u>	<u>0.62</u>	<u>-0.65</u>	<u>-0.42</u>	0.18	-0.22	-0.02
VDCN 5ha	<u>-0.31</u>	<u>-0.29</u>	<u>0.65</u>	<u>0.56</u>	<u>-0.71</u>	<u>-0.47</u>	0.16	-0.23	0.09
BA	<u>-0.37</u>	<u>-0.29</u>	0.22	0.13	<u>-0.25</u>	-0.19	-0.03	0.08	0.00
RBA _{CON}	-0.17	<u>-0.56</u>	<u>0.30</u>	0.19	<u>-0.41</u>	-0.17	0.03	0.02	-0.11

SWC: soil water content; T_{soil}: soil temperature; BD: soil bulk density

Table A2. Selected variables for each measurement date, along with the R² and root mean square error (RMSE) values of the quantile regression forest models without wetland and with wetland. Note that the same variables were selected at all dates in both cases.

Measurement dates	Selected variables	R ² (RMSE, nmol m ⁻² s ⁻¹)	R ² (RMSE, nmol m ⁻² s ⁻¹)
		Without wetland	With wetland
2023/04/27	SWI, TPI, PrC	0.53 (0.52)	0.37 (0.86)
2023/05/12	TPI, VDCN	0.31 (0.82)	0.22 (1.27)
2023/05/31	SWI, TPI, VDCN	0.43 (0.48)	0.25 (1.07)
2023/07/06	SWI, TPI	0.40 (0.50)	0.34 (1.99)
2023/07/26	SWI, TPI	0.37 (1.02)	0.30 (1.60)
2023/09/04	SWI, TPI	0.40 (1.18)	0.38 (1.27)
2023/10/07	TPI, VDCN	0.67 (0.81)	0.42 (1.07)
2023/11/07	SWI, TPI, VDCN, PrC	0.59 (0.66)	0.47 (1.25)
2023/11/30	SWI, TPI, VDCN, PrC	0.51 (0.56)	0.40 (0.57)

600 **Table A3: Comparison of the accuracy of the quantile regression forest (QRF) models without and with vegetation at the two dates where BA was selected. Selected variables and their importance scores in parentheses, along with the R² and root mean square error (RMSE) values (model with vegetation is in bold italic letters).**

Date	Selected variables			R ²	RMSE	605
2023/04/27	SWI (0.57)	PrC (0.58)	TPI (0.67)	0.53	0.52	
	<u>SWI (0.64)</u>	<u>BA (0.42)</u>	<u>TPI (0.50)</u>	<u>0.51</u>	<u>0.52</u>	
2023/10/07	TPI (0.88)	VDCN (0.58)		0.67	0.81	
	<u>TPI (0.77)</u>	<u>VDCN (0.39)</u>	<u>BA (0.27)</u>	<u>0.55</u>	<u>0.80</u>	

610

615 **Table A4. Summary of the linear mixed model (LMMs) analysing the relationship between the predicted soil CH₄ fluxes and measured soil CH₄ fluxes, where measurement periods were included as the random factor on both slope and intercept. The p-values of the fixed effect were for testing if the intercept was different from zero and the slope different from 1. The statistics panel at the bottom left shows the marginal (R²_m) and conditional (R²_c) coefficients of determination, the root mean square error of the model, and the overall significance of the model (p-value).**

Fixed effect: predicted CH ₄ flux			Random effects: measurement dates		
Estimate ± SE		p-values		Intercept	Slope
Intercept	0.15 ± 0.03	from 0: 0.003	2023/04/27	-0.05	0.03
Slope	1.16 ± 0.02	from 1: 0.92	2023/05/12	-0.02	0.02
			2023/05/31	-0.05	0.02
			2023/07/06	-0.06	0.02
Statistics			2023/07/26	0.07	0.00
n	467		2023/09/04	0.14	-0.03
R ² _m	0.94		2023/10/07	0.02	-0.04
R ² _c	0.95		2023/11/07	-0.01	-0.04
RMSE	0.26		2023/11/30	-0.03	0.01
p-value	2.5 × 10 ⁻⁹				

620

Table A5. Summary of the linear mixed model (LMMs) analysing the relationship between the predicted soil CH₄ fluxes and measured soil CH₄ fluxes, where landscape units (positions, vegetation types, and vegetation density) were included as the random factor on both slope and intercept. The p-values of the fixed effect were for testing if the intercept was different from zero and the slope different from 1. The statistics panel at the bottom left shows the marginal (R^2_m) and conditional (R^2_c) coefficients of determination, the root mean square error of the model, and the overall significance of the model.

Fixed effect: Predicted CH ₄ flux		Random effect: Positions		
Estimate \pm SE	p values		Intercept	Slope
Intercept	-0.17 \pm 0.04	from 0: 0.006	Plain	0.01
Slope	0.94 \pm 0.05	from 1: 0.99	Foot slope	0.01
Statistics			Slope	-0.05
n	36		Ridge	0.05
R^2_m	0.93			-0.09
R^2_c	0.98			
RMSE	0.20			
p-value	7.8×10^{-5}			
Fixed effect: Predicted CH ₄ flux		Random effect: Vegetation type		
Intercept	-0.19 \pm 0.07	from 0: 0.08	Broadleaf	0.02
Slope	0.82 \pm 0.09	from 1: 0.99	Coniferous	-0.09
Statistics			Mixed	0.08
n	27			0.12
R^2_m	0.91			
R^2_c	0.96			
RMSE	0.17			
p-value	0.01			
Fixed effect: Predicted CH ₄ flux		Random effect: Vegetation density		
Intercept	-0.13 \pm 0.05	from 0: 0.08	High	-0.01
Slope	0.81 \pm 0.08	from 1: 0.99	Medium	0.06
Statistics			Low	-0.05
n	27			0.11
R^2_m	0.80			
R^2_c	0.95			
RMSE	0.19			
p-value	0.01			

Table A6. Summary of the LMM analysing the effects of topographic position, vegetation type, and density on predicted soil CH₄ fluxes. Pixel ID was included as a random effect, and spatial autocorrelation among residuals was eliminated. η_p^2 was calculated as the effect size of each explanatory variable. Letters indicated the significance within each landscape unit.

Explanatory variables	p-value	Effect size (η_p^2) [with 95% CI]
Position [df=3]	<0.001	0.43 [0.42, 0.44]
<i>Plain (a), Foot slope (b), Slope (c), Ridge (c)</i>		
Vegetation type [df=2]	<0.001	0.006 [0.00, 0.01]
<i>Broadleaf (a), Coniferous (b) and Mixed (c)</i>		
Vegetation density [df=2]	<0.001	0.11 [0.10, 0.12]
<i>High (c), Medium (b), Low (a)</i>		

Table A7. Percentage of upland pixels in the study area distributed among four levels of predicted relative uncertainty for soil CH₄ fluxes.

Measurement date	Uncertainty			
	< 50 %	50 - 99 %	100- 500 %	> 500 %
2023/04/27	5.72%	53.29%	40.99%	0.01%
2023/05/12	19.93%	54.01%	26.06%	-
2023/05/31	10.90%	39.29%	49.81%	-
2023/07/06	27.76%	38.32%	33.92%	-
2023/07/26	21.12%	40.66%	38.13%	0.09%
2023/09/04	30.19%	51.24%	18.58%	-
2023/10/07	39.68%	38.13%	22.19%	-
2023/11/07	17.50%	46.10%	36.4%	-
2023/11/30	7.83%	43.65%	48.26%	0.26%

Table A8. Statistics of the linear relationship between landscape-scaled soil CH₄ fluxes and antecedent precipitation indexes (API).

665


20 antecedent days provided the best fit. 30 and 7 antecedent days are shown as common metrics in hydrology. Adjusted recession coefficients (k) and determination coefficients (R²) are shown.

Antecedent days	k	R ²
20	0.69	0.70
30	0.69	0.69
7	0.67	0.68

670

Table A9. Proportion of vegetation density and type associated with the different topographic positions across the landscape.

Position	Vegetation density	Proportion (%)	Vegetation type	Proportion (%)
Plain	High	19.6	Broadleaf	91.8
	Medium	8.4	Coniferous	2.7
	Low	72.0	Mixed	5.5
Foot slope	High	4.9	Broadleaf	12.9
	Medium	38.5	Coniferous	0.5
	Low	56.6	Mixed	86.6
Slope	High	33.4	Broadleaf	4.5
	Medium	46.3	Coniferous	20.1
	Low	20.3	Mixed	75.4
Ridge	High	86.0	Coniferous	13.6
	Medium	14.0	Mixed	86.4



675

Fig A1. Seasonal variation of soil CH_4 fluxes from wetlands (means and standard error, $n = 3$).





Fig A2. Accumulated local effect plots for each model on each measurement date.



680

Fig A3. Relationship between predicted uncertainty and predicted soil CH_4 fluxes. The highest uncertainty is observed for a near-zero prediction.

Data availability

The data used in this study are available at The Kyoto University Research Information Repository (KURANAI, DOI: to be

685

added)

Author contribution

DE had the original idea of this research. DE and SKP designed the research framework with suggestions from MD. SKP, DE, and KY conducted the vegetation survey and flux measurements. SKP analyzed and performed the modeling under the supervision of DE. SKP wrote the manuscript that was critically reviewed and edited by all co-authors.

690

Competing interest

The authors declare that they have no conflict of interest.

Acknowledgement

We would like to thank the staff of the Ashiu Forest Station of the Field Science Education and Research Centre, Kyoto University for enabling this research, providing the access to the forest, and for sharing the Digital Elevation Model (DEM) 695 and climate data. We are grateful to Takumi Mochidome for his helpful discussion about GIS and Lucie Bivaud and Makoto Nagasawa for their help during the field survey.

Funding

The research was supported by grants from the Research Institute for Sustainable Humanosphere (RISH), Kyoto University, the Japan Society for the Promotion of Science (KAKENHI Grant no. JP24K01797) and the SPRING fellowship program 700 from Japan Science and Technology (JST SPRING Grant no. JPMJSP2110).

References

Ågren, A. M., Lidberg, W., Strömgren, M., Ogilvie, J., and Arp, P. A.: Evaluating digital terrain indices for soil wetness mapping – a Swedish case study, *Hydrol. Earth Syst. Sci.*, 18, 3623–3634, <https://doi.org/10.5194/hess-18-3623-2014>, 2014.

705 Angel, R., Claus, P., and Conrad, R.: Methanogenic archaea are globally ubiquitous in aerated soils and become active under wet anoxic conditions, *ISME J.*, 6, 847–862, <https://doi.org/10.1038/ismej.2011.141>, 2012.

Apley, D. W. and Zhu, J.: Visualizing the Effects of Predictor Variables in Black Box Supervised Learning Models, *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 82, 1059–1086, <https://doi.org/10.1111/rssb.12377>, 2020.

Aronson, E. L. and Helliker, B. R.: Methane flux in non-wetland soils in response to nitrogen addition: a meta-analysis, *Ecology*, 91, 3242–3251, <https://doi.org/10.1890/09-2185.1>, 2010.

710 Bartoń, K.: MuMIn: Multi-Model Inference, <https://doi.org/10.32614/CRAN.package.MuMIn>, 2010.

Bates, D., Mächler, M., Bolker, B., and Walker, S.: Fitting linear mixed-effects models using lme4, *J. Stat. Soft.*, 67, <https://doi.org/10.18637/jss.v067.i01>, 2015.

Bell, J. C., Cunningham, R. L., and Havens, M. W.: Calibration and validation of a soil-landscape model for predicting soil 715 drainage class, *Soil Science Soc of Amer J*, 56, 1860–1866, <https://doi.org/10.2136/sssaj1992.03615995005600060035x>, 1992.

Ben-Shachar, M. S., Makowski, D., Lüdecke, D., Patil, I., Wiernik, B. M., Thériault, R., and Waggoner, P.: effectsize: Indices of Effect Size, <https://doi.org/10.32614/CRAN.package.effectsize>, 2019.

720 Beven, K. J. and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology / Un modèle à base physique de zone d'appel variable de l'hydrologie du bassin versant, *Hydrol. Sci. Bull.*, 24, 43–69, <https://doi.org/10.1080/02626667909491834>, 1979.

Bock, M. and Köthe, R.: Predicting the depth of hydromorphic soil characteristics influenced by ground water, *SAGA-Seconds Out*, 19, 13–22, 2008.

Bodelier, P. L. E. and Laanbroek, H. J.: Nitrogen as a regulatory factor of methane oxidation in soils and sediments, *FEMS Microbiol. Ecol.*, 47, 265–277, [https://doi.org/10.1016/S0168-6496\(03\)00304-0](https://doi.org/10.1016/S0168-6496(03)00304-0), 2004.

725 Börjesson, G. and Nohrstedt, H.-Ö.: Fast recovery of atmospheric methane consumption in a Swedish forest soil after single-shot N-fertilization, *For. Ecol. Manage.*, 134, 83–88, [https://doi.org/10.1016/S0378-1127\(99\)00249-2](https://doi.org/10.1016/S0378-1127(99)00249-2), 2000.

Borken, W., Xu, Y., and Beese, F.: Conversion of hardwood forests to spruce and pine plantations strongly reduced soil methane sink in Germany, *Glob Change Biol.*, 9, 956–966, <https://doi.org/10.1046/j.1365-2486.2003.00631.x>, 2003.

Breiman, L.: Random Forest, *Machine learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.

730 Brumme, R. and Borken, W.: Site variation in methane oxidation as affected by atmospheric deposition and type of temperate forest ecosystem, *Glob. Biogeochem. Cycles*, 13, 493–501, <https://doi.org/10.1029/1998GB900017>, 1999.

Burt, R., Reinsch, T. G., and Miller, W. P.: A micro-pipette method for water dispersible clay, *Commun. Soil Sci. Plant Anal.*, 24, 2531–2544, <https://doi.org/10.1080/00103629309368975>, 1993.

735 Christiansen, J. R., Levy-Booth, D., Prescott, C. E., and Grayston, S. J.: Microbial and environmental controls of methane fluxes along a soil moisture gradient in a pacific coastal temperate rainforest, *Ecosystems*, 19, 1255–1270, <https://doi.org/10.1007/s10021-016-0003-1>, 2016.

Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., Wehberg, J., Wichmann, V., and Böhner, J.: System for Automated Geoscientific Analyses (SAGA) v. 2.1.4, *Geosci. Model Dev.*, 8, 1991–2007, <https://doi.org/10.5194/gmd-8-1991-2015>, 2015.

740 Courtois, E. A., Stahl, C., Van Den Berge, J., Bréchet, L., Van Langenhove, L., Richter, A., Urbina, I., Soong, J. L., Peñuelas, J., and Janssens, I. A.: Spatial Variation of Soil CO₂, CH₄ and N₂O Fluxes Across Topographical Positions in Tropical Forests of the Guiana Shield, *Ecosystems*, 21, 1445–1458, <https://doi.org/10.1007/s10021-018-0232-6>, 2018.

Du, Z., Riveros-Iregui, D. A., Jones, R. T., McDermott, T. R., Dore, J. E., McGlynn, B. L., Emanuel, R. E., and Li, X.: Landscape position influences microbial composition and function via redistribution of soil water across a watershed, *Appl Environ Microbiol*, 81, 8457–8468, <https://doi.org/10.1128/AEM.02643-15>, 2015.

745 Dutaur, L. and Verchot, L. V.: A global inventory of the soil CH₄ sink, *Glob. Biogeochem. Cycles*, 21, 2006GB002734, <https://doi.org/10.1029/2006GB002734>, 2007.

Epron, D., Plain, C., Ndiaye, F.-K., Bonnaud, P., Pasquier, C., and Ranger, J.: Effects of compaction by heavy machine traffic on soil fluxes of methane and carbon dioxide in a temperate broadleaved forest, *For. Ecol. Manage.*, 382, 1–9, <https://doi.org/10.1016/j.foreco.2016.09.037>, 2016.

750 Epron, D., Mochidome, T., Tanabe, T., Dannoura, M., and Sakabe, A.: Variability in stem methane emissions and wood methane production of different tree species in a cold temperate mountain forest, *Ecosystems*, 26, 784–799, <https://doi.org/10.1007/s10021-022-00795-0>, 2023.

Freeman, T. G.: Calculating catchment area with divergent flow based on a regular grid, *Computers & Geosciences*, 17, 413–422, [https://doi.org/10.1016/0098-3004\(91\)90048-i](https://doi.org/10.1016/0098-3004(91)90048-i), 1991.

Genuer, R., Poggi, J.-M., and Tuleau-Malot, C.: Variable selection using random forests, *Pattern Recognit. Lett.*, 31 (14), 2225–2236, 2010.

Genuer, R., Poggi, J.-M., and Tuleau-Malot, C.: VSURF: An R package for variable selection using random forests, *The R Journal*, 7, 19, <https://doi.org/10.32614/RJ-2015-018>, 2015.

760 Gomez, J., Vidon, P., Gross, J., Beier, C., Caputo, J., and Mitchell, M.: Estimating greenhouse gas emissions at the soil–atmosphere interface in forested watersheds of the US Northeast, *Environ. Monit. Assess.*, 188, 295, <https://doi.org/10.1007/s10661-016-5297-0>, 2016.

Greenwell, B., M. and Boehmke, B., C.: Variable importance plots—an introduction to the vip package, *The R Journal*, 12, 343, <https://doi.org/10.32614/RJ-2020-013>, 2020.

765 Guckland, A., Flessa, H., and Prenzel, J.: Controls of temporal and spatial variability of methane uptake in soils of a temperate deciduous forest with different abundance of European beech (*Fagus sylvatica* L.), *Soil Biol. Biochem.*, 41, 1659–1667, <https://doi.org/10.1016/j.soilbio.2009.05.006>, 2009.

Hakamada, R. E., Hubbard, R. M., Moreira, G. G., Stape, J. L., Campoe, O., and Ferraz, S. F. D. B.: Influence of stand density on growth and water use efficiency in Eucalyptus clones, *For. Ecol. Manage.*, 466, 118125, <https://doi.org/10.1016/j.foreco.2020.118125>, 2020.

770 Harrell Jr, F. E.: Hmisc: Harrell Miscellaneous, 5.2–4, <https://doi.org/10.32614/CRAN.package.Hmisc>, 2003.

Hirai, H., Araki, S., and Kyuma, K.: Characteristics of brown forest soils developed on the paleozoic shale in northern Kyoto with special reference to their pedogenetic process, *Soil Sci. Plant Nutr.*, 34, 157–170, <https://doi.org/10.1080/00380768.1988.10415670>, 1988.

775 Holwerda, F., Scatena, F. N., and Bruijnzeel, L. A.: Throughfall in a Puerto Rican lower montane rain forest: A comparison of sampling strategies, *J. Hydrol.*, 327, 592–602, <https://doi.org/10.1016/j.jhydrol.2005.12.014>, 2006.

Hu, R., Hirano, T., Sakaguchi, K., Yamashita, S., Cui, R., Sun, L., and Liang, N.: Spatiotemporal variation in soil methane uptake in a cool-temperate immature deciduous forest, *Soil Biol. Biochem.*, 184, 109094, <https://doi.org/10.1016/j.soilbio.2023.109094>, 2023.

780 Hütsch, B. W.: Tillage and land use effects on methane oxidation rates and their vertical profiles in soil, *Biol. Fertil. Soils*, 27, 284–292, <https://doi.org/10.1007/s003740050435>, 1998.

IPCC: Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, 1st ed., Cambridge University Press, <https://doi.org/10.1017/9781009157896>, 2023.

785 Ishihara, M. I., Suzuki, S. N., Nakamura, M., Enoki, T., Fujiwara, A., Hiura, T., Homma, K., Hoshino, D., Hoshizaki, K., Ida, H., Ishida, K., Itoh, A., Kaneko, T., Kubota, K., Kuraji, K., Kuramoto, S., Makita, A., Masaki, T., Namikawa, K., Niiyama, K., Noguchi, M., Nomiya, H., Ohkubo, T., Saito, S., Sakai, T., Sakimoto, M., Sakio, H., Shibano, H., Sugita, H., Suzuki, M., Takashima, A., Tanaka, N., Tashiro, N., Tokuchi, N., Yakushima Forest Environment Conservation Center, Yoshida, T., and Yoshida, Y.: Forest stand structure, composition, and dynamics in 34 sites over Japan, *Ecol. Res.*, 26, 1007–1008, <https://doi.org/10.1007/s11284-011-0847-y>, 2011.

Ishizuka, S., Sakata, T., and Ishizuka, K.: Methane oxidation in Japanese forest soils, *Soil Biol. Biochem.*, 32, 769–777, [https://doi.org/10.1016/S0038-0717\(99\)00200-X](https://doi.org/10.1016/S0038-0717(99)00200-X), 2000.

Ishizuka, S., Sakata, T., Sawata, S., Ikeda, S., Sakai, H., Takenaka, C., Tamai, N., Onodera, S., Shimizu, T., Kan-na, K., Tanaka, N., and Takahashi, M.: Methane uptake rates in Japanese forest soils depend on the oxidation ability of topsoil, with a new estimate for global methane uptake in temperate forest, *Biogeochemistry*, 92, 281–295, <https://doi.org/10.1007/s10533-009-9293-0>, 2009.

790 Itoh, M., Ohte, N., and Koba, K.: Methane flux characteristics in forest soils under an East Asian monsoon climate, *Soil Biol. Biochem.*, 41, 388–395, <https://doi.org/10.1016/j.soilbio.2008.12.003>, 2009.

Jacinthe, P. A., Vidon, P., Fisher, K., Liu, X., and Baker, M. E.: Soil methane and carbon dioxide fluxes from cropland and riparian buffers in different hydrogeomorphic settings, *J. Environ. Qual.*, 44, 1080–1090, <https://doi.org/10.2134/jeq2015.01.0014>, 2015.

Jensen, S., Priemé, A., and Bakken, L.: Methanol improves methane uptake in starved methanotrophic microorganisms, *Appl. Environ. Microbiol.*, 64, 1143–1146, <https://doi.org/10.1128/AEM.64.3.1143-1146.1998>, 1998.

Jeong, G., Oeverdieck, H., Park, S. J., Huwe, B., and Ließ, M.: Spatial soil nutrients prediction using three supervised learning methods for assessment of land potentials in complex terrain, *Catena*, 154, 73–84, <https://doi.org/10.1016/j.catena.2017.02.006>, 2017.

805 Jevon, F. V., Gewirtzman, J., Lang, A. K., Ayres, M. P., and Matthes, J. H.: Tree species effects on soil CO₂ and CH₄ fluxes in a mixed temperate forest, *Ecosystems*, <https://doi.org/10.1007/s10021-023-00852-2>, 2023.

Kagotani, Y., Hamabata, E., and Nakajima, T.: Seasonal and spatial variations and the effects of clear-cutting in the methane absorption rates of a temperate forest soil, *Nutr. Cycl. Agroecosystems*, 59, 169–175, <https://doi.org/10.1023/A:1017554031367>, 2001.

810 Kaiser, K. E., McGlynn, B. L., and Dore, J. E.: Landscape analysis of soil methane flux across complex terrain, *Biogeosciences*, 15, 3143–3167, <https://doi.org/10.5194/bg-15-3143-2018>, 2018.

Kemppinen, J., Niittynen, P., Riihimäki, H., and Luoto, M.: Modelling soil moisture in a high-latitude landscape using LiDAR and soil data, *Earth Surf. Processes Landf.*, 43, 1019–1031, <https://doi.org/10.1002/esp.4301>, 2018.

815 King, G. M. and Schnell, S.: Ammonium and nitrite inhibition of methane oxidation by *Methylobacter albus* BG8 and *Methylosinus trichosporium* OB3b at low methane concentrations, *Appl. Environ. Microbiol.*, 60, 3508–3513, <https://doi.org/10.1128/aem.60.10.3508-3513.1994>, 1994.

Kohler, M. A. and Linsley, R. K.: Predicting the Runoff from Storm Rainfall, US department of commerce, Weather Bureau, 820 20, 1951.

Kravchenko, A. N., Bollero, G. A., Omonode, R. A., and Bullock, D. G.: Quantitative mapping of soil drainage classes using topographical data and soil electrical conductivity, *Soil Sci. Soc. Am. J.*, 66, 235–243, <https://doi.org/10.2136/sssaj2002.2350>, 2002.

Kruse, C. W., Moldrup, P., and Iversen, N.: Modeling diffusion and reaction in soils: II. atmospheric methane diffusion and consumption in a forest soil, *Soil Sci.*, 161, 355–365, <https://doi.org/10.1097/00010694-199606000-00002>, 1996.

825 Kuhn, M. and Johnson, K.: *Applied Predictive Modeling*, Springer New York, New York, NY, <https://doi.org/10.1007/978-1-4614-6849-3>, 2013.

Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: lmerTest package: tests in linear mixed effects models, *J. Stat. Soft.*, 82, <https://doi.org/10.18637/jss.v082.i13>, 2017.

830 Lee, J., Oh, Y., Lee, S. T., Seo, Y. O., Yun, J., Yang, Y., Kim, J., Zhuang, Q., and Kang, H.: Soil organic carbon is a key determinant of CH₄ sink in global forest soils, *Nat. Commun.*, 14, 3110, <https://doi.org/10.1038/s41467-023-38905-8>, 2023.

Lenth, R. V.: emmeans: Estimated Marginal Means, aka Least-Squares Means, <https://doi.org/10.32614/cran.package.emmeans>, 2017.

Lundbäck, M., Persson, H., Häggström, C., and Nordfjell, T.: Global analysis of the slope of forest land, *Forestry*, 94, 54–69, <https://doi.org/10.1093/forestry/cpaa021>, 2021.

835 Luo, G. J., Kiese, R., Wolf, B., and Butterbach-Bahl, K.: Effects of soil temperature and moisture on methane uptake and nitrous oxide emissions across three different ecosystem types, *Biogeosciences*, 10, 3205–3219, <https://doi.org/10.5194/bg-10-3205-2013>, 2013.

Martinson, G. O., Müller, A. K., Matson, A. L., Corre, M. D., and Veldkamp, E.: Nitrogen and phosphorus control soil methane uptake in tropical montane forests, *JGR Biogeosciences*, 126, e2020JG005970, <https://doi.org/10.1029/2020JG005970>, 2021.

840 Meinshausen, N.: Quantile Regression Forests, *J. Mach. Learn. Res.*, 7, 983–999, 2006.

Meinshausen, N.: quantregForest: Quantile Regression Forests, <https://doi.org/10.32614/CRAN.package.quantregForest>, 2017.

Miller, B. A., Koszinski, S., Wehrhan, M., and Sommer, M.: Impact of multi-scale predictor selection for modeling soil properties, *Geoderma*, 239–240, 97–106, <https://doi.org/10.1016/j.geoderma.2014.09.018>, 2015.

845 Mochizuki, Y., Koba, K., and Yoh, M.: Strong inhibitory effect of nitrate on atmospheric methane oxidation in forest soils, *Soil Biol. Biochem.*, 50, 164–166, <https://doi.org/10.1016/j.soilbio.2012.03.013>, 2012.

Murphy, P. N. C., Ogilvie, J., Meng, F.-R., White, B., Bhatti, J. S., and Arp, P. A.: Modelling and mapping topographic variations in forest soils at high resolution: A case study, *Ecol. Model.*, 222, 2314–2332, <https://doi.org/10.1016/j.ecolmodel.2011.01.003>, 2011.

850 Nakamura, Y. and Krestov, P. V.: Coniferous forests of the temperate zone of asia, in: *Ecosystems of the world. Coniferous forests*, vol. 6, edited by: Andersson, F. A., Elsevier, 163–220, 2005.

Osborne, B. B., Nasto, M. K., Asner, G. P., Balzotti, C. S., Cleveland, C. C., Sullivan, B. W., Taylor, P. G., Townsend, A. R., and Porder, S.: Climate, topography, and canopy chemistry exert hierarchical control over soil N cycling in a neotropical lowland forest, *Ecosystems*, 20, 1089–1103, <https://doi.org/10.1007/s10021-016-0095-7>, 2017.

855 Pachepsky, Ya. A., Timlin, D. J., and Rawls, W. J.: Soil Water Retention as Related to Topographic Variables, *Soil Science Soc of Amer J*, 65, 1787–1795, <https://doi.org/10.2136/sssaj2001.1787>, 2001.

Pinheiro, J., R Core Team, and Bates, D.: nlme: Linear and Nonlinear Mixed Effects Models, <https://doi.org/10.32614/cran.package.nlme>, 1999.

Praeg, N., Wagner, A. O., and Illmer, P.: Plant species, temperature, and bedrock affect net methane flux out of grassland and 860 forest soils, *Plant Soil.*, 410, 193–206, <https://doi.org/10.1007/s11104-016-2993-z>, 2017.

Quebbeman, A. W., Menge, D. N. L., Zimmerman, J., and Uriarte, M.: Topography and tree species improve estimates of spatial variation in soil greenhouse gas fluxes in a subtropical forest, *Ecosystems*, 25, 648–660, <https://doi.org/10.1007/s10021-021-00677-x>, 2022.

R Core Team: A language and environment for statistical computing, R: A language and environment for statistical computing, 865 2024.

Räsänen, A., Manninen, T., Korkiakoski, M., Lohila, A., and Virtanen, T.: Predicting catchment-scale methane fluxes with multi-source remote sensing, *Landscape Ecol.*, 36, 1177–1195, <https://doi.org/10.1007/s10980-021-01194-x>, 2021.

Roberts, D. R., Bahn, V., Ciuti, S., Boyce, M. S., Elith, J., Guillera-Arroita, G., Hauenstein, S., Lahoz-Monfort, J. J., Schröder, B., Thuiller, W., Warton, D. I., Wintle, B. A., Hartig, F., and Dormann, C. F.: Cross-validation strategies for data with temporal, 870 spatial, hierarchical, or phylogenetic structure, *Ecography*, 40, 913–929, <https://doi.org/10.1111/ecog.02881>, 2017.

Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carroll, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, 875 G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., Van Weele, M., Van Der Werf, G. R., Weiss, R. F., Worthy, 880 D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth Syst. Sci. Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.

Schoener, G. and Stone, M. C.: Monitoring soil moisture at the catchment scale – A novel approach combining antecedent precipitation index and radar-derived rainfall data, *J. Hydrol.*, 589, 125155, <https://doi.org/10.1016/j.jhydrol.2020.125155>, 2020.

885 Semrau, J. D., DiSpirito, A. A., and Vuilleumier, S.: Facultative methanotrophy: false leads, true results, and suggestions for future research: Facultative methanotrophy, *FEMS Microbiol. Lett.*, 323, 1–12, <https://doi.org/10.1111/j.1574-6968.2011.02315.x>, 2011.

890 Sidle, R. C., Tsuboyama, Y., Noguchi, S., Hosoda, I., Fujieda, M., and Shimizu, T.: Stormflow generation in steep forested headwaters: a linked hydrogeomorphic paradigm, *Hydrol. Process.*, 14, 369–385, [https://doi.org/10.1002/\(SICI\)1099-1085\(20000228\)14:3<369::AID-HYP943>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1099-1085(20000228)14:3<369::AID-HYP943>3.0.CO;2-P), 2000.

Tateno, R. and Takeda, H.: Forest structure and tree species distribution in relation to topography-mediated heterogeneity of soil nitrogen and light at the forest floor, *Ecol. Res.*, 18, 559–571, <https://doi.org/10.1046/j.1440-1703.2003.00578.x>, 2003.

Ueda, S., Ando, M., and Kanzaki, K.: Forest soil surveys of the Kyoto University Forest in Ashiu. II. Soil types, grain size, and chemical and physical properties of soils, *Bulletin of the Kyoto University Forests*, 65, 94–112, 1993.

895 Vainio, E., Peltola, O., Kasurinen, V., Kieloaho, A.-J., Tuittila, E.-S., and Pihlatie, M.: Topography-based statistical modelling reveals high spatial variability and seasonal emission patches in forest floor methane flux, *Biogeosciences*, 18, 2003–2025, <https://doi.org/10.5194/bg-18-2003-2021>, 2021.

Vanclay, J. K.: Managing water use from forest plantations, *For. Ecol. Manage.*, 257, 385–389, <https://doi.org/10.1016/j.foreco.2008.09.003>, 2009.

900 Veldkamp, E., Koehler, B., and Corre, M. D.: Indications of nitrogen-limited methane uptake in tropical forest soils, *Biogeosciences*, 10, 5367–5379, <https://doi.org/10.5194/bg-10-5367-2013>, 2013.

Virkkala, A.-M., Niittynen, P., Kemppinen, J., Marushchak, M. E., Voigt, C., Hensgens, G., Kerttula, J., Happonen, K., Tyystjärvi, V., Biasi, C., Hultman, J., Rinne, J., and Luoto, M.: High-resolution spatial patterns and drivers of terrestrial ecosystem carbon dioxide, methane, and nitrous oxide fluxes in the tundra, *Biogeosciences*, 21, 335–355, <https://doi.org/10.5194/bg-21-335-2024>, 2024.

905 Wang, J. M., Murphy, J. G., Geddes, J. A., Winsborough, C. L., Basiliko, N., and Thomas, S. C.: Methane fluxes measured by eddy covariance and static chamber techniques at a temperate forest in central Ontario, Canada, *Biogeosciences*, 10, 4371–4382, <https://doi.org/10.5194/bg-10-4371-2013>, 2013.

910 Wang, L. and Liu, H.: An efficient method for identifying and filling surface depressions in digital elevation models for hydrologic analysis and modelling, *Int. J. Geogr. Inf. Sci.*, 20, 193–213, <https://doi.org/10.1080/13658810500433453>, 2006.

Warner, D. L., Vargas, R., Seyfferth, A., and Inamdar, S.: Transitional slopes act as hotspots of both soil CO₂ emission and CH₄ uptake in a temperate forest landscape, *Biogeochemistry*, 138, 121–135, <https://doi.org/10.1007/s10533-018-0435-0>, 2018.

915 Warner, D. L., Guevara, M., Inamdar, S., and Vargas, R.: Upscaling soil-atmosphere CO₂ and CH₄ fluxes across a topographically complex forested landscape, *Agric. For. Meteorol.*, 264, 80–91, <https://doi.org/10.1016/j.agrformet.2018.09.020>, 2019.

West, A. E. and Schmidt, S. K.: Acetate stimulates atmospheric CH₄ oxidation by an alpine tundra soil, *Soil Biol. Biochem.*, 31, 1649–1655, [https://doi.org/10.1016/S0038-0717\(99\)00076-0](https://doi.org/10.1016/S0038-0717(99)00076-0), 1999.

920 Yamao, M., Sidle, R. C., Gomi, T., and Imaizumi, F.: Characteristics of landslides in unwelded pyroclastic flow deposits, southern Kyushu, Japan, *Nat. Hazards Earth Syst. Sci.*, 16, 617–627, <https://doi.org/10.5194/nhess-16-617-2016>, 2016.

Yu, L., Huang, Y., Zhang, W., Li, T., and Sun, W.: Methane uptake in global forest and grassland soils from 1981 to 2010, *Sci. Total Environ.*, 607–608, 1163–1172, <https://doi.org/10.1016/j.scitotenv.2017.07.082>, 2017.

925 Yu, L., Zhu, J., Ji, H., Bai, X., Lin, Y., Zhang, Y., Sha, L., Liu, Y., Song, Q., Dörsch, P., Mulder, J., and Zhou, W.: Topography-related controls on N₂O emission and CH₄ uptake in a tropical rainforest catchment, *Sci. Total Environ.*, 775, 145616, <https://doi.org/10.1016/j.scitotenv.2021.145616>, 2021.

Zevenbergen, L. W. and Thorne, C. R.: Quantitative analysis of land surface topography, *Earth Surf. Processes Landf.*, 12, 47–56, <https://doi.org/10.1002/esp.3290120107>, 1987.