

Supplementary information for:

Mesozoic to Cenozoic bathymetric and gateway changes reduced Atlantic Ocean sensitivity to deoxygenation

^{1,2}Nina M. Papadomanolaki, ^{3,4}Anta-Clarisse Sarr, ¹Anthony Gramoullé, ¹Marie Laugie,
⁵Jean-Baptiste Ladant and ¹Yannick Donnadieu

¹CEREGE, Aix Marseille Univ, CNRS, IRD, INRAE, France

²Institute of Geology and Paleontology, University of Münster, Münster, Germany

³Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IRD, Univ. Gustave Eiffel, ISTERre, Grenoble, France

⁴Department of Earth Sciences, University of Oregon, Eugene, 97403 USA

⁵Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Univ. Paris-Saclay, Gif-sur-Yvette, France

Content:

Figure S1: Time series.

Figure S2: Northern Hemisphere mixed layer depth.

Figure S3: Southern Ocean salinity and wind velocity.

Figure S4: Ideal age of bottom waters.




Figure S1. Time series for (a) sea surface salinity (top 50m of ocean), (b) deep ocean salinity (below 2500 meters of water depth), (c) maximum intensity of the meridional stream function (negative for counterclockwise motion), (d) sea surface temperature (top 50m), (e) deep ocean temperatures (below 2500m) and (f) maximum (winter) mixed layer depth.




Figure S2. Maximum (winter) mixed layer depth (m) in the Southern Hemisphere. *DeepCAS* results from (Laugié et al., 2021).

Water age at seafloor >2500m (yr)




Figure S3. Ideal age (top row) and age versus oxygen concentrations for each of the major basins (bottom row), at 60Ma.




Figure S3. Surface salinity (psu) (left column) and changes 850hPa wind velocity (m s^{-1}) (right column) in the Southern Ocean. From top to bottom, salinity is shown for TG_{CLOSED} , TG_{OPEN} and $4 \times TG_{\text{OPEN}}$. Differences in wind velocity are shown as a difference between TG_{OPEN} and TG_{CLOSED} , $4 \times TG_{\text{OPEN}}$ and TG_{OPEN} , and $4 \times TG_{\text{OPEN}}$ and TG_{CLOSED} .

Table S1. Freshwater Flux (Sv) into the Southern Ocean sectors and corresponding average salinity values for the top 50m of these sectors. Italics indicate the sector where deep-waters form.

Simulation	Southern ocean sector	Freshwater Flux (Sv)	Average salinity (0-50m)
<i>TG_{CLOSED}</i>	Atlantic/Indian	0.5927	33.82
	Pacific	<i>0.5668</i>	<i>34.67</i>
<i>TG_{OPEN}</i>	Atlantic/Indian	<i>0.6059</i>	<i>34.41</i>
	Pacific	<i>0.6071</i>	<i>34.51</i>
<i>4×TG_{OPEN}</i>	Atlantic/Indian	<i>0.7417</i>	<i>35.08</i>
	Pacific	0.8031	33.22

References

Laugié, M., Donnadieu, Y., Ladant, J. B., Bopp, L., Ethé, C., and Raison, F.: Exploring the impact of Cenomanian paleogeography and marine gateways on oceanic oxygen, *Paleoceanography and Paleoclimatology*, 36, e2020PA004202, 2021.