

Estimating Particulate Organic Matter Flux from In-Situ Optics: A Framework for Correcting for Suspended Particles and Incorporating Depth-Dependent Degradation

Nasrollah Moradi^{1,2}, Lili Hufnagel^{1,2,*}, Simon Ramondenc^{1,2,§}, Clara M. Flintrop^{1,†}, Rainer Kiko^{3,4,5}, Tim Fischer⁶, Helena Hauss^{7,8}, Arne Körtzinger^{9,5}, Gerhard Fischer² and Morten H. Iversen^{1,2}

¹Alfred Wegener Institute Helmholtz-Center for Polar and Marine Research Bremerhaven, Bremerhaven, Germany

²MARUM and University of Bremen, Bremen, Germany

³Laboratoire d'Océanographie de Villefranche, LOV Institut de la Mer de Villefranche, IMEV, Villefranche-sur-Mer, France

⁴Department of Biological Oceanography, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁵Faculty of Mathematics and Natural Sciences, Kiel University, Kiel, Germany.

⁶Department of Physical Oceanography, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁷Division of Climate & Environment, Norwegian Research Centre (NORCE), Bergen, Norway

⁸Department of Ocean Ecosystems Biology, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

⁹Department of Chemical Oceanography, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

*Current address: Helmholtz Centre for Environmental Research - UFZ, Department of River Ecology, Magdeburg, Germany

§Current address: Sorbonne University, CNRS, Laboratoire d'Océanographie de Villefranche, Villefranche-sur-Mer, France

†Current address: The Fredy & Nadine Herrmann Institute of Earth Sciences, Hebrew University of Jerusalem, Jerusalem, Israel

Correspondence to: Nasrollah Moradi (nasrollah.moradi@awi.de) and Morten Iversen (morten.iversen@awi.de)

Abstract. Accurate quantification of particulate organic matter (POM) flux from settling particles is crucial for understanding the efficiency of oceanic CO₂ sequestration. Recent advancements in in-situ camera systems (ISCs) have enabled high-resolution estimates of the size distribution of particle concentration (PSDc) in the global ocean, both vertically and horizontally. When calibrated against corresponding sediment trap flux observations, these PSDc data serve as a basis for estimating downward POM flux. However, these estimates are subject to several uncertainties. A central unresolved issue lies in differentiating settling particles from suspended particles in ISC data. Additionally, conventional approaches for converting PSDc to organic matter flux—which typically estimate a particle's contribution to the total flux based on a power-law relationship with its size—do not explicitly integrate particle sinking velocities and degradation rates, two key factors regulating POM export. Overcoming these limitations requires incorporating spatiotemporally aligned supplementary data and refined methodological approaches.

Here, we develop and validate a process-based, three-step framework to address these uncertainties. First, to correct for the contribution of non-sinking particles, we calibrate ISC data against particle flux measurements from gel traps using a measured size-velocity relationship, thereby yielding a robust estimate of the sinking particle flux. Second, leveraging the output from the first step, we build upon the conventional power-law approach by developing a mechanistic POC flux model that incorporates size-specific sinking velocities and a depth-dependent degradation term modulated by water temperature and oxygen concentration. Third, with the POC flux estimates from the second step as input, we extend the framework to estimate particulate organic nitrogen (PON) flux using a dynamic model for particulate organic C:N stoichiometry. By

Style Definition: Heading 5

Deleted: A Novel Approach to Estimate Carbon and Nitrogen

Deleted:

Deleted: Application to Cyclonic Eddies off the Cape Verde Islands A Framework for Correcting for Suspended Particles and

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: English (US)

Deleted: *

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US), Not Superscript/ Subscript

Formatted: Font: 10 pt, English (US)

Formatted: English (US)

Formatted: Font: Not Bold, English (US)

Formatted: Affiliation

Deleted: Abstract. Mesoscale eddies are pervasive in the global ocean and are suggested to play a significant role in enhancing carbon export from the surface to the deep ocean. However, their dynamic nature and associated challenges of in-situ sampling have limited direct measurements of carbon flux within these features, leaving their contribution to carbon export uncertain. In-situ optical devices offer a promising solution by providing high-resolution data on particle abundance and size distribution (PSD) within eddies, both vertically and horizontally. Yet, converting PSD data into reliable carbon flux estimates is particularly challenging in dynamic features like eddies, and is further complicated by the need to account for key factors regulating carbon export. To address this, we present a novel mechanistic framework that optimally integrates PSD data with flux measurements, settling velocities and respiration rates of in-situ collected aggregates, together with water temperature and oxygen concentration to estimate both particulate carbon and nitrogen fluxes in the water column. This framework tackles primary challenges by incorporating contributions from suspended particles, size-dependent sinking velocities, and the degradation rates of settling particles with depth. Applied to a multi-instrument dataset from a high-resolution survey of eddies off the Cape Verde Islands, the presented framework reveals distinct funnel-shaped flux patterns for particulate organic carbon and nitrogen, with a twofold enhancement of flux within the examined cyclonic eddy. Furthermore, it identifies lower carbon-to-nitrogen ratios in settling organic matter at the deep eddy core, indicating the transport of fresher organic matter to the deep sea. These findings provide new insights into eddy-driven carbon export, highlighting the roles of settling velocity, water temperature, and oxygen concentration in shaping carbon flux and organic matter stoichiometry. While observed in a single eddy, this doubling of carbon flux underscores the potential for mesoscale eddies to locally enhance carbon export and, given their widespread occurrence, contribute significantly to the ocean's capacity to sequester atmospheric CO₂.

linking the C:N ratio to particle age, the model reproduces the increase in C:N with depth—a key biogeochemical trend driven by preferential nitrogen remineralization over carbon that is not resolved by conventional, static-ratio modeling approaches.

1 Introduction

The oceanic biological carbon pump (BCP) plays a pivotal role in Earth's climate system by transporting large quantities of photosynthetically fixed carbon from the sunlit surface ocean to the deep sea and, thereby, maintaining the vertically increasing gradient in dissolved organic carbon in the water column (Volk and Hoffert, 1985; Sarmiento and Gruber, 2006; Iversen, 2023). This BCP is driven by the downward flux of particulate organic matter (POM) via settling marine particles, which delivers particulate organic carbon (POC) to depths where it can be sequestered on long-term (Stukel et al., 2018; Boyd et al., 2019; Buesseler et al., 2020). Model estimates for annual carbon export to the mesopelagic currently range between 4 and 12 Pg (DeVries and Weber, 2017; Henson et al., 2011). Therefore, more accurate quantification of POC flux is needed to reduce uncertainties in global carbon budgets, to improve projections of climate–ocean feedbacks, and to assess the efficacy of oceanic carbon storage in a rapidly changing ocean (Siegel et al., 2023).

Direct measurement of POC flux has traditionally relied on sediment traps (ST)—moored, drifting, or neutrally buoyant—which collect settling particles over defined deployment periods. While STs provide invaluable ground-truth data, they inherently sample a limited area and depth, and the subsequent sample processing is labor-intensive and time-consuming (Buesseler et al., 2007, 2008). These limitations constrain the spatiotemporal resolution of ST-based flux datasets. Consequently, capturing the full spectrum of POC flux variability—especially in highly dynamic oceanic features such as mesoscale eddies, which can significantly influence regional carbon export (McGillicuddy et al., 2007; Siegel et al., 2011)—remains challenging with STs alone, as the biogeochemical properties within these features can evolve rapidly over timescales of days to weeks.

To enhance observational capacity, in situ camera systems (ISCs) have emerged as valuable tools (Picheral et al., 2010; Giering et al., 2020). These instruments capture high-resolution spatiotemporal observations of marine particles and zooplankton (Jackson et al., 1997; Kiko et al., 2017). After aggregating particle counts into discrete, often logarithmically spaced size bins to reduce sampling noise, these observations yield the depth-resolved size distribution of particle concentration (PSDc)—that is, the number of particles per unit water volume within each size bin. Once calibrated against corresponding sediment trap (ST) flux measurements, these PSDc data can serve as a credible proxy for estimating high-resolution POC flux profiles throughout the water column. The most common approach for this calibration, hereafter referred to as the conventional method, models each particle's contribution to the total POC flux as a power-law function of

Formatted: Font: +Headings (Times New Roman), English (US)

Formatted: Font: 12 pt, Bold, English (US)

Formatted: Font: 12 pt, Font colour: Auto, English (US)

Formatted: Font: 12 pt, Bold, English (US)

Formatted: Justified, Line spacing: 1,5 lines

Formatted: Font: +Headings (Times New Roman), Font colour: Black, English (US), Kern at 16 pt

Formatted: Line spacing: 1,5 lines

Moved (insertion) [1]

Formatted: Font: +Headings (Times New Roman), English (US)

Moved (insertion) [2]

Formatted: Font: +Headings (Times New Roman), English (US)

115 its size (Guidi et al., 2008). The total flux at a given depth, z_ℓ , is then calculated by summing the contributions of all particles:

$$F_{\text{conv}}(z_\ell) = \sum n_i \times A_{\text{conv}} \times d_i^{B_{\text{conv}}} \quad (1)$$

120 where n_i is the ISC-derived particle concentration in size bin i at depth z_ℓ , d_i is the midpoint particle diameter for that size bin, and A_{conv} and B_{conv} are free parameters optimized empirically using corresponding ST-based flux measurements over a predefined particle size range. This approach is similarly applied to estimate other biogeochemical fluxes, such as PON flux. The straightforward implementation of this conventional method has led to its widespread use (Guidi et al., 2008; Iversen et al., 2010; Ramondenc et al., 2016); however, it has inherent limitations that may introduce uncertainty into flux estimates. The first limitation arises from the nature of ISC observations: ISCs detect all particles within their field of view, including both sinking particles that contribute to vertical flux and non-sinking particles (i.e., suspended particles and zooplankton). Because non-sinking particles do not directly contribute to downward POM flux, their inclusion in flux calculations may introduce uncertainty, particularly in situations where the concentration of suspended particles varies vertically due to changes in the pycnocline structure.

130 The second limitation stems from the assumption that a particle's contribution to the flux depends solely on its size (Iversen and Lampitt, 2023). However, as particles settle, their organic carbon content progressively decreases due to microbial degradation (Ploug et al., 1999; Simon et al., 2002). Since the physical structure of settling aggregates is often maintained by mineral components, this degradation of organic matter can occur without a significant change in the particle's apparent size as detected by an ISC.

135 A third limitation arises when extending this method to multiple elements. Its formulation—applying separate power-law relationships to POC and PON—implicitly yields a C:N ratio that is effectively static with depth. This static behavior directly contradicts the well-documented principle of preferential nitrogen remineralization (Grossart and Ploug, 2001; Hach et al., 2020; Iversen, 2023 and references therein), a process known to cause the C:N ratio of sinking organic matter to increase with depth (Romankevich, 1984; Schneider et al., 2003, 2004). Consequently, the conventional method can lead to a misrepresentation of C:N stoichiometry throughout the water column. These limitations become more pronounced when ISC particle data are calibrated using flux measurements confined to either the upper or lower water column (e.g., drifting ST or moored ST data, respectively), potentially leading to systematic over- or underestimation of POC and PON flux at unmeasured depths.

140 Overcoming these limitations poses a dual challenge: it requires both the integration of ISC data with a suite of supplementary measurements and the development of a refined methodological framework to synthesize these diverse data into mechanistically coherent models. Leveraging such a spatiotemporally aligned dataset, we develop and validate an enhanced framework aimed at mitigating the uncertainties associated with the conventional method. The datasets integrated

Moved (insertion) [3]

Formatted: English (US)

in our study include: (i) profiles of ISC-derived PSD; (ii) sediment trap measurements of POC and PON flux and gel trap data on the flux of sinking particles; (iii) direct, size-specific measurements of particle sinking velocities and particle respiration rates; and (iv) corresponding profiles of temperature and oxygen concentration from CTD casts. The framework is implemented through three sequential methods (Fig. 1) that systematically tackle each of the outlined limitations.

Method I yields reliable profiles of the size distribution of particle flux (PSDf) from ISC data. It achieves this by implementing a calibration against gel trap flux data and size-specific sinking velocities, which corrects for the contribution of non-sinking particles. Building upon the corrected particle flux, **Method II** enhances POC flux estimation by incorporating a mechanistic model of depth-dependent carbon degradation. This model moves beyond a purely size-based relationship by explicitly accounting for the degradation of a particle's organic carbon as a function of ambient oxygen, temperature, and travel time during its descent. Finally, with **Method III**, we extend this framework to address the issue of static C:N stoichiometry. To account for this, a new model for PON flux estimation is introduced that incorporates a time-dependent relationship between the PON and POC mass of sinking particles. When combined with the outputs from the preceding methods, this refinement effectively captures preferential nitrogen remineralization over carbon (Grossart and Ploug, 2001; Hach et al., 2020) and, consequently, the increase in the C:N ratio of organic matter flux with increasing depth (Schneider et al., 2004).




Figure 1. Methodological framework and notations. (a) Schematic diagram illustrating the inputs, outputs, and interconnections within the proposed framework for integrating in situ observations (see *Materials and Methods* for details). (b) Definitions of the primary notations used in the methodology. The abbreviations used are as follows. DST: Drifting Sediment Trap; UVP: Underwater Vision Profiler; MSC: Marine Snow Catcher; CTD: Conductivity, Temperature, and Depth profiler system.

2 Materials and Methods

2.1 In-situ observations

The data used in this study were from a high-resolution, multi-instrument survey of mesoscale eddies conducted aboard the RV *Meteor* cruise (M160) in November–December 2019 around the Cape Verde archipelago (Fig. A1). While the cruise had a broader biogeochemical focus, the work presented here is concerned primarily with methodological development, not an analysis of the eddies themselves. Below, we provide a brief overview of the equipment deployed during the cruise and the shipboard measurements used to collect these data. A detailed description is provided in the Appendix. We leveraged a subset of the M160 dataset because its co-located and concurrent Underwater Vision Profiler (UVP), Drifting Sediment Trap (DST) equipped with a gel trap, and Marine Snow Catcher (MSC) measurements provided a complementary dataset well-suited to developing our methodological framework.

DST. Six DST deployments were carried out—four southwest and two northeast of the Cape Verde Islands. The DSTs collected settling particles at depths of 100 m, 200 m, and 400 m. At each depth, one trap cylinder was equipped with a gel-filled collection cup to preserve particles for size-distribution analysis and flux calculations, while the remaining cylinders were used to measure biogeochemical fluxes, including POC and PON. Each gel sample was imaged using a high-resolution camera, and the size of the collected particles was accurately determined using Gel-PISA, an in-house software package developed for image analysis. Collected particles were subsequently sorted into logarithmically spaced size bins, and counts per bin were determined. These counts, together with the deployment time and collection cup area, were used to derive the gel trap-based particle flux (F_{Gel}) for each size bin—yielding a representative PSDf for each cup. At each deployment station and depth, POC and PON fluxes were also quantified from samples collected in the remaining cylinders; these samples were processed and measured in the laboratory.

UVP. A UVP5 camera system was mounted on the CTD-Rosette and deployed at 71 stations, including parallel deployment at all stations where a DST was deployed and recovered, to image particles in the water column inside and outside the studied eddies. We applied the same logarithmically spaced size bins used for F_{Gel} . Particle images from each water-depth layer (see Appendix for layer definitions) were analyzed to determine the UVP-based particle concentration (n_{UVP}) for each size bin, yielding a representative PSDc for that layer.

MSC. To measure the size, sinking velocities and respiration rates of in-situ collected aggregates, eight MSC deployments were conducted. Aggregates were collected from 10 m below the chlorophyll maximum, and a total of 88 were analyzed onboard using a flow chamber to measure their size (d) and sinking velocity (W). These measurements were fitted to a power-law function:

$$W(d) = A_{\text{vel}} \times d^{B_{\text{vel}}} \quad (2)$$

to establish a representative size–velocity relationship for settling particles in the study area, with the free parameters A_{vel} and B_{vel} determined from the data. Oxygen concentration profiles were also measured for a randomly selected subset of 31 aggregates. Using the same flow chamber and a microsensor system, these profiles were determined across the particle–water interface. These measurements were used to calculate the interfacial oxygen flux following the method of Moradi et al. (2021) and, subsequently, the oxygen uptake rates of the settling aggregates (Q_{O_2}).




Figure 2. Parameterization and comparison of gel trap-based and UVP-derived PSDf. (a) An example gel trap PSDf with its log-log linear fit (blue line). (b) The corresponding, spatiotemporally aligned UVP-derived PSDf (Eq. 3), parameterized as in (a). (c) Comparison of the best-fit lines from the gel trap-based and UVP-derived PSDf data shown in panels (a) and (b). (d) Schematic of the trapezoidal area used for optimizing correction factors in Method I. Note: "part." on the axis labels denotes the number of particles.

2.2 Data Integration Methods

2.2.1 Method I: Correcting UVP-derived particle flux for non-sinking particles

The UVP-based particle flux for each size bin ($F_{\text{UVP},i}$), was calculated by multiplying the corresponding UVP-based particle concentration ($n_{\text{UVP},i}$), by the sinking velocity associated with that bin. This calculation is expressed as:

$$F_{\text{UVP},i} [\text{particles m}^{-2} \text{ day}^{-1}] = n_{\text{UVP},i} [\text{particles m}^{-3}] \times W(d_i) [\text{m day}^{-1}] \quad (3)$$

where, i and d_i indicate the i -th size bin and its midpoint, respectively, and $W(d_i)$ is the sinking velocity for particles of size d_i , obtained from the size–velocity relationship. The resulting UVP-based PSDf was further characterized by a pair of intercept and slope values (a_{UVP} and b_{UVP}), derived from the linear regression of $\log_{10}(F_{\text{UVP}})$ versus $\log_{10}(d)$ over the considered particle size range (Clements et al., 2023). The logarithmic transformation was used to capture contributions across a wide range of particle sizes, from small to large.

It is important to note that the calculated F_{UVP} values generally tend to overestimate actual particle fluxes, as the UVP camera detects both settling and non-sinking particles. Therefore, adjustments to the characteristic intercept-slope pairs were necessary to reliably use them as a proxy for downward particle flux estimates in the water column.

Moved (insertion) [4]

Formatted: Font: 12 pt, English (US)

Formatted: Left, Line spacing: single

230 Correction factors α and β were introduced to adjust each original intercept-slope pair as $\alpha \cdot a_{\text{UVP}}$ and $\beta \cdot b_{\text{UVP}}$, respectively. We assumed that F_{Gel} provided reliable flux estimates of sinking particles, since suspended particles are unlikely to enter sediment traps and zooplankton swimmers are manually removed during image analysis. The F_{Gel} data were processed in the same manner to derive a set of intercept-slope pairs (a_{Gel} and b_{Gel}), which served as the reference for the optimization of the correction factors α and β . These correction factors were then optimized by minimizing the discrepancy between each paired gel trap and UVP-based PSDf in log-space. For each gel trap-based PSDf, its spatiotemporally aligned UVP counterpart was identified.

235 A trapezoidal area (S) was then calculated for each distribution, defined by its respective characteristic fitted line (see Fig. 2 for an illustration). The optimization was achieved by minimizing the sum of squared errors (Err_S) between the areas of the paired gel trap and UVP-based trapezoids across all k samples

$$240 \quad \text{Err}_S = \sum_k (S_{\text{Gel},k} - S_{\text{UVP},k})^2 \quad (4)$$

where $S_{\text{Gel},k}$ is the reference area from the k -th gel trap sample and $S_{\text{UVP},k}$ is the area from its UVP counterpart. This procedure finds the optimal α and β that best align the corrected UVP-derived areas with the gel trap references. The areas of the resulting trapezoids were calculated as follows:

$$245 \quad S_{\text{Gel}} = L \cdot (y_{1,\text{Gel}} + y_{2,\text{Gel}}) / 2 \quad \text{where } L = x_2 - x_1, \quad y_{1,\text{Gel}} = a_{\text{Gel}} + b_{\text{Gel}} \cdot x_1, \quad \text{and } y_{2,\text{Gel}} = a_{\text{Gel}} + b_{\text{Gel}} \cdot x_2. \quad (5)$$

Similarly,

$$S_{\text{UVP}} = L \cdot (y_{1,\text{UVP}} + y_{2,\text{UVP}}) / 2 \quad \text{where } y_{1,\text{UVP}} = \alpha \cdot a_{\text{UVP}} + \beta \cdot b_{\text{UVP}} \cdot x_1, \quad \text{and } y_{2,\text{UVP}} = \alpha \cdot a_{\text{UVP}} + \beta \cdot b_{\text{UVP}} \cdot x_2. \quad (6)$$

The size range boundaries x_1 and x_2 were fixed to $\log_{10}(144)$ and $\log_{10}(2315)$, respectively, defining an overlapping size range where both the UVP5 and the gel trap provide reliable spectral slopes. These sizes correspond to the midpoints of the 128–161 μm and 2050–2580 μm size bins, respectively.

250 Prior to the logarithmic transformation, particle sizes (d) were normalized by 1 μm and particle fluxes (F_{Gel} and F_{UVP}) by 1 particle $\text{day}^{-1} \text{m}^{-2}$, making the resulting intercepts, slopes, and trapezoid areas unitless for the optimization procedure. Note that these correction factors were applied to the intercepts and slopes originally derived from $\log_{10}(F_{\text{UVP}})$ data. Accordingly, the UVP-derived particle flux corrected for non-sinking particles (\tilde{F}_{UVP}) was calculated for each size bin i as:

$$255 \quad \tilde{F}_{\text{UVP},i} = 10^a \times d_i^b \quad (7)$$

with $\alpha = a \cdot a_{\text{UVP}}$ and $\tilde{b} = \beta \cdot b_{\text{UVP}}$.

If there are no particles in a size bin, \tilde{F}_{UVP} is set to zero for that bin. The size bins considered in this study are shown in Fig. 3b. In Eq. 7 using particle size d_i in μm yields a final flux in the units of particles $\text{day}^{-1} \text{m}^{-2}$.

Moved (insertion) [5]

Formatted: Font: +Body (Times New Roman), English (US)

Moved (insertion) [6]

Formatted: Font: +Body (Times New Roman), English (US)

Moved (insertion) [7]

Formatted: Font: +Body (Times New Roman), English (US)

260 2.2.2 Method IIa: Incorporating depth-dependent carbon degradation into POC flux estimates from UVP particle data

265 In this method, we develop a simplified mechanistic model that accounts for variations in the organic carbon content of settling particles as a function of both particle size and depth. The depth dependency stems from progressive carbon degradation during particle descent, which is modulated by the temperature and oxygen concentration of the ambient water at depth. To accurately estimate the organic carbon content of a settling particle at a given depth, the model incorporates the cumulative degradation it experiences during its descent. Each UVP-detected particle in size bin i is treated as a spherical particle with diameter d_i (units: mm) originating at the sea surface (i.e., at depth $z = 0$) and sinking through the water column at a constant velocity $W(d_i)$. The initial organic carbon content of the particle ($M_{\text{POC},i}(0)$, units: mg C) is assumed to follow a power-law function of particle size:

$$270 \quad M_{\text{POC},i}(0) = A_{\text{POC}} \times d_i^{B_{\text{POC}}} \quad (8)$$

where A_{POC} and B_{POC} are parameters to be optimized using flux measurements as described below.

To account for organic carbon degradation during particle descent, the water column is divided into a series of discrete vertical layers, each with thickness Δz_j , where j denotes the layer index. The layer thickness increases with depth, consistent with the applied depth averaging of the UVP data: 5 m in the upper 100 m, 10 m from 100–200 m, 15 m from 200–400 m, and 25 m below 400 m. As a particle settles through each layer, it undergoes carbon loss due to microbial respiration. The carbon loss of the particle within a layer is determined by the effective carbon degradation rate in that layer, scaled by the time the particle spends transiting through it. This residence time is given by:

$$275 \quad \Delta t_j = \Delta z_j / W(d_i) \quad (9)$$

280 where $W(d_i)$ is the sinking velocity of the particle calculated from the size-velocity relationship (Eq. 2). To model POC loss in sinking particles, we employ a zero-order degradation model. This choice is mechanistically grounded in our experimental approach, which derives carbon degradation rates from oxygen fluxes measured at the particle-water interface (detailed in Eqs. 14–16). This formulation treats remineralization as a surface-area-limited process and assumes that the microbial enzymes responsible for degradation are saturated with substrate, resulting in a constant degradation rate under constant ambient conditions.

285 The effective carbon degradation rate of a particle in size bin i while in layer j , denoted as $\Lambda_{\text{eff},i}(z_j)$ (units: mg C day⁻¹), is derived from a reference degradation rate ($\Lambda_{\text{ref},i}$), which represents the rate under reference conditions. This reference rate is then modulated by the mean temperature (T) and mean oxygen concentration (O_2) (Iversen and Ploug, 2013; Ploug and Bergkvist, 2015) within that layer:

$$290 \quad \Lambda_{\text{eff},i}(z_j) = \Lambda_{\text{ref},i} \times \theta_j(T, O_2) \quad (10)$$

where the modulation term (θ_j) is modeled using a Q_{10} temperature response and Michaelis-Menten kinetics for oxygen limitation:

$$\theta_j(T, O_2) = (Q_{10}^{(T(z_j) - T_{\text{ref}})/10}) \times \left(\frac{O_2(z_j)}{K_m + O_2(z_j)} \right) \quad (11)$$

Here, T_{ref} is the reference temperature and the temperature coefficient (Q_{10}) and the half-saturation oxygen constant (K_m) are set to the optimized values of $Q_{10} = 2.5 \pm 0.2$ and $K_m = 19 \pm 7 \mu\text{mol O}_2 \text{ kg}^{-1}$ from DeVries and Weber (2017). The mass of carbon lost while the particle transits layer j ($\Delta M_{\text{POC},i}(z_j)$) is therefore:

$$\Delta M_{\text{POC},i}(z_j) = \Lambda_{\text{eff},i} \times \Delta t_j = \Lambda_{\text{eff},i} \times \Delta z_j / W(d_i) \quad (12)$$

The carbon content of the particle at the base of layer l ($M_{\text{POC},i}(z_\ell)$) is calculated by subtracting the cumulative carbon losses from its initial surface mass. To ensure physical realism, a non-negativity constraint is applied:

$$M_{\text{POC},i}(z_\ell) = \max \left(A_{\text{POC}} \times d_i^{B_{\text{POC}}} - \sum_{j=1}^{\ell} \Delta M_{\text{POC},i}(z_j), 0 \right) \quad (13)$$

where the sum is over the ℓ layers from the sea surface down to depth z_ℓ . The non-negativity constraint reflects the natural scenario where, once a settling marine particle's organic carbon is entirely consumed, no further carbon degradation of that particle can occur. The reference degradation rate ($\Lambda_{\text{ref},i}$) was parameterized using microbial respiration rates measured on in-situ collected particles (see *Results* for MSC). It is defined based on the total oxygen consumption rate per particle ($Q_{O_2,i}$) as:

$$\Lambda_{\text{ref},i} = Q_{O_2,i} \times \Phi \times \sigma \quad (14)$$

where $Q_{O_2,i}$ itself is estimated from the measured oxygen flux at the particle-water interface (J_{O_2}) and the particle's surface area (A_i) (Ploug et al., 2002; Moradi et al., 2018):

$$Q_{O_2,i} = J_{O_2} \times A_i \quad (15)$$

Here, J_{O_2} (units: $\mu\text{mol O}_2 \text{ mm}^{-2} \text{ day}^{-1}$) is the mean oxygen flux measured at the interface of in-situ collected particles, and A_i represents the surface area of the particle, πd_i^2 (units: mm^2). The term Φ represents the stoichiometric ratio for oxygen-to-carbon conversion, assumed to be 1, which corresponds to a respiratory quotient of 1 mol CO_2 produced per 1 mol O_2 consumed (Iversen and Ploug, 2013). The term σ is the molar mass of carbon (12 g mol^{-1}). With this formulation, a surface-area-normalized organic carbon degradation rate can be defined as $\Lambda_0 \equiv J_{O_2} \cdot \Phi \cdot \sigma$. This rate quantifies the organic carbon degradation driven by microbial respiration, normalized to the particle surface area. Thus, $\Lambda_{\text{ref},i}$ (Eq. 14) can be expressed

as:

$$\Lambda_{\text{ref},i} = \Lambda_0 \times A_i \quad (16)$$

The total POC flux estimate at depth z_ℓ ($F_{\text{POC}}(z_\ell)$) is calculated by summing the carbon content of all particles passing through that depth:

$$F_{\text{POC}}(z_\ell) = \sum_{i=1}^N F_{\text{UVP},i}(z_\ell) \times M_{\text{POC},i}(z_\ell) \quad (17)$$

where $F_{UVP,i}$ is the corrected UVP-derived particle flux for the i -th size bin from Method I. The free parameters (A_{POC} and B_{POC}) were optimized by minimizing the sum of squared logarithmic errors between the available sediment trap measurements ($F_{POC,k}$) and their corresponding model estimates ($\hat{F}_{POC,k}$) (units: $\text{mg C day}^{-1} \text{ m}^{-2}$):

$$\text{Err}_{POC} = \sum [\log_{10}(F_{POC,k}) - \log_{10} \hat{F}_{POC,k}]^2. \quad (18)$$

2.2.3 Method IIb: Alternative formulation for estimating POC flux using first-order degradation model

For comparison, we also present an alternative formulation based on a first-order degradation model—a commonly used approach in large-scale biogeochemical models, where the degradation rate constant is typically treated as an adjustable parameter to be optimized rather than as a directly measured input (DeVries and Weber, 2017). The first-order degradation model is based on the assumption that the rate of change of a substance is proportional to its current amount. Under this assumption, the POC mass of a particle as a function of time is described by:

$$M_{POC,i}(t) = M_{POC,i}(0) \times e^{-\lambda^* t} \quad (19)$$

where λ^* is the degradation rate constant (units: day^{-1}). By applying this model to our discretized vertical layer framework, the POC mass of a sinking particle at the base of layer ℓ is calculated by multiplying its initial mass by a decay factor derived from the sum of layer-specific decay terms:

$$M_{POC,i}(z_\ell) = A_{POC}^* \times d_i^{B_{POC}^*} \times \exp\left(-\lambda^* \sum_{j=1}^{\ell} \theta_j(T, O_2) \frac{\Delta z_j}{W(d_i)}\right) \quad (20)$$

where the starred superscripts indicate that the free parameters (A_{POC}^* , B_{POC}^* and λ^*) are specific to the first-order model. These parameters were then optimized simultaneously by minimizing the error function (Eq. 18), subject to the constraint that $\lambda^* > 0$.

2.2.4 Method III: A dynamic model for PON flux: accounting for preferential nitrogen degradation

To account for the preferential remineralization of nitrogen over carbon in sinking particles (Grossart and Ploug, 2001; Hach et al., 2020), an alternative modeling approach for estimating PON flux is presented. This approach begins by assuming a correlation between the particulate organic nitrogen ($M_{PON,i}$) and particulate organic carbon mass of a particle, an assumption supported by our in-situ flux observations (see Sect. 3.1.2). This relationship is initially expressed as:

$$M_{PON,i} = M_{POC,i} \gamma_i \quad (21)$$

where γ_i is a dimensionless correlation factor representing the C:N ratio for particles in the i -th size bin. To incorporate the preferential degradation of nitrogen, we model γ_i as a function of particle age, defined as its travel time (in days), t_i , from the sea surface. Since preferential nitrogen degradation causes the C:N ratio to increase with particle age, a power-law relationship can serve as a baseline model to capture this variation:

$$\gamma_i = A_{\text{PON}} \times t_i^{B_{\text{PON}}}. \quad (22)$$

350 However, this straightforward formulation presents two conceptual inconsistencies in its asymptotic behavior. First, at short travel times, it yields biogeochemically implausible low C:N ratios, including a theoretically incorrect value of zero at the surface ($t_i = 0$). Second, at long travel times, its unbounded growth is inconsistent with the expectation that the C:N ratio should approach a constant, maximum value as the organic matter becomes refractory. While the first issue can be pragmatically managed by applying a minimum depth constraint, the second issue of unbounded growth is a more

355 fundamental limitation that requires a different model formulation. To resolve this asymptotic behavior while maintaining a parsimonious two-parameter structure to avoid over-fitting, we propose a more robust model based on the Michaelis-Menten formulation:

$$\gamma_i = A_{\text{PON}} \times t_i / (B_{\text{PON}} + t_i) \quad (23)$$

where A_{PON} and B_{PON} are positive parameters to be optimized. In this formulation, the dimensionless parameter A_{PON} represents the asymptotic C:N mass ratio of refractory organic matter as $t_i \rightarrow \infty$, corresponding to a maximum molar C:N ratio of $(14/12) \times A_{\text{PON}}$. The parameter B_{PON} is the half-saturation constant, representing the time required to reach half of this maximum ratio (in days). To avoid the implausibly low C:N values this model produces near the surface, a practical constraint is applied by setting a minimum depth, such as 50 m.

Using the particle's sinking velocity $W(d_i)$ to determine its travel time to a given depth z_ℓ , the nitrogen content of the particle at that depth ($M_{\text{PON},i}(z_\ell)$) is then given by:

$$M_{\text{PON},i}(z_\ell) = M_{\text{POC},i}(z_\ell) / [A_{\text{PON}} \times (z_\ell / W(d_i)) / (B_{\text{PON}} + (z_\ell / W(d_i)))] \quad (24)$$

With this, the estimated PON flux at a given position and depth is calculated by summing the nitrogen content of all particles, weighted by their corrected UVP-derived flux:

$$F_{\text{PON}}(z_\ell) = \sum F_{\text{UVP},i}(z_\ell) \times M_{\text{PON},i}(z_\ell) \quad (25)$$

370 where the index i refers to the i -th size bin and the summation is over all considered size bins. Similar to the POC flux estimation, the parameters A_{PON} and B_{PON} were determined by minimizing the sum of squared errors between measured and estimated PON fluxes:

$$\text{Err}_{\text{PON}} = \sum [\log_{10}(F_{\text{PON},k}) - \log_{10} F_{\text{PON},k}^*]^2 \quad (26)$$

where $F_{\text{PON},k}$ and $F_{\text{PON},k}^*$ are the k -th sediment trap-based PON flux measurement and its corresponding model-estimated PON flux (Eq. 25), respectively, and the summation is over all available PON flux observations.

2.3 Uncertainty Analysis

A comprehensive uncertainty analysis was performed using a sequential Monte Carlo procedure to quantify and propagate uncertainties through the entire modeling chain. This approach was essential to correctly account for the statistical

dependencies between parameters estimated in successive model stages and to incorporate parametric, sampling, and measurement uncertainties. The procedure was repeated for $\mathcal{M}=1000$ simulations to generate an ensemble representing the full joint probability distribution of all model parameters. This resulting ensemble allows for the robust calculation of the median and 95% confidence intervals for any model parameter or derived quantity.

2.3.1 Step 1: Uncertainty in the particle flux estimates

A nested bootstrap procedure was implemented to determine the uncertainty in the corrected particle flux, F_{UVP} . For each of the $\mathcal{M}=1000$ iterations, the following two-part process was executed:

Velocity uncertainty. A plausible size-velocity relationship was generated via a parametric bootstrap, drawing a sample pair of A_{vel} and B_{vel} from the bivariate normal distribution defined by the best-fit velocity parameters and their covariance matrix. This preserves the inherent correlation between A_{vel} and B_{vel} .

Sampling uncertainty & dependent optimization. To account for sampling uncertainty and the dependent optimization, a reference particle flux dataset was created by resampling with replacement from the six DST deployments ($\mathcal{R}=6$). Each deployment was treated as a single block to preserve the vertical structure of the observations at 100 m, 200 m, and 400 m. The correction factors α and β were then re-optimized using this bootstrapped dataset in conjunction with the specific velocity model (generated for this iteration).

This initial stage yielded an ensemble of \mathcal{M} self-consistent parameter sets (A_{vel} , B_{vel} , α , and β) and their corresponding corrected particle flux models, F_{UVP} .

2.3.2 Step 2: Uncertainty in the POC flux estimates

The uncertainty from the particle flux model was propagated into the POC flux model. For each of the $\mathcal{M}=1000$ iterations the following process was executed:

Input condition assembly. For each iteration, a self-consistent set of plausible input conditions was assembled. This set included: (i) the corrected particle flux model, F_{UVP} , from the corresponding iteration of the previous step; (ii) three independent biogeochemical parameters— Λ_{0z} , Q_{10z} , and K_m —each drawn from its respective normal distribution defined by its mean and standard error; and (iii) a bootstrap sample of the reference POC flux measurements, created by resampling the six DST deployments ($\mathcal{R}=6$) with replacement. To ensure a consistent propagation of sampling uncertainty across the modeling chain, the same random bootstrap indices used in Step 1 were re-applied to generate this sample.

Dependent optimization. The POC flux model parameters, A_{POC} and B_{POC} , were then re-optimized for each of these \mathcal{M} scenarios. The output of this stage was an expanded ensemble of \mathcal{M} parameter sets, A_{vel} , B_{vel} , α , β , A_{POC} , and B_{POC} , now including the POC flux model parameters, fully conditioned on the results of the previous stage.

2.3.3 Step 3: Uncertainty in the PON flux estimates

The analysis was extended in a third sequential step to propagate all preceding uncertainties into the PON flux model. For each iteration the following process was executed:

Input condition assembly. All previously determined plausible models and parameters for that iteration were utilized, including the velocity model (A_{vel} , B_{vel} , α , and β) and the POC flux model (A_{POC} and B_{POC}). A bootstrap sample of reference PON flux measurements was created by applying the same resampling indices from the prior stages to the six DST deployments ($\mathcal{R}=6$).

Dependent optimization. The PON flux model parameters (A_{PON} and B_{PON}) were then re-optimized based on this complete and self-consistent set of inputs.

This step finalized the generation of the full ensemble, resulting in \mathcal{M} complete sets of all eight model parameters (A_{vel} , B_{vel} , α , β , A_{POC} , B_{POC} , A_{PON} , and B_{PON}) that capture the entire cascade of correlated uncertainties.


2.3.4 Step 4: Uncertainty in the estimated C:N ratio

The uncertainty in the final estimated ratio of POC to PON flux (C:N) was determined as a derived quantity from the full Monte Carlo simulation. For each of the $\mathcal{M} = 1000$ iterations, the total POC flux and the total PON flux were calculated using the complete parameter set for that iteration. The ratio of these two fluxes was then computed. This process yielded a final distribution of 1000 plausible ratio values, from which the median and 95% CI were determined to represent the final estimate and its propagated uncertainty.

We determined the optimal model parameters by minimizing the defined error terms using the “fmin” function from Python's SciPy library. To robustly search for the global minimum and avoid local optima, this optimization was performed 300 times, each initiated with a random starting point within the specified parameter space. The parameter set yielding the lowest final error was selected as the best-fit solution.

Throughout this study, optimized parameters are reported as the median—a robust statistic for potentially skewed distributions—and the associated 95% CI, derived from their optimized bootstrap distributions.

For all power-law relationships of the form $Y = A \cdot d^B$ used in this study, the particle diameter (d) is first expressed in millimeters (mm) and then normalized by a reference size of 1 mm, yielding a dimensionless size term ($d [\text{mm}] / 1 [\text{mm}]$). This normalization results in units for the pre-factor (A) that are the same as the units of the dependent variable (Y), while the exponent (B) is always dimensionless.



440 **Figure 3. In-situ observations.** (a) Measured particle sinking velocity W versus particle size d (equivalent spherical diameter). The blue
 445 curve shows the best-fit power law $W(d) = A_{vel}d^{B_{vel}}$, with $A_{vel}=121.0$ (m day $^{-1}$) and $B_{vel}=0.5$ ($R^2=0.11$); the shaded area denotes the
 95% confidence interval (CI). (b) Sinking velocities for the defined size bins, derived from the fitted size-velocity relationship. (c)
 Comparison of size-velocity relationships for sinking marine particles from this study with those reported by Cael et al. (2021). (d)
 Calculated diffusive O $_2$ flux across the aggregate-water interface as a function of particle size, with the dashed line indicating the mean
 flux. (e-g) PSDf from individual gel traps across all six DST deployments at 100 m, 200 m, and 400 m, respectively. (h-j) Corresponding
 PSDf derived from UVP data for the same stations and depths, calculated using Eq. 3. (Mean DST-based fluxes (N=6): (k) POC flux, (l)
 PON flux, (m) Corresponding molar C:N calculated as $14/12 \times (POC/PON)$. (n) Individual POC fluxes vs. their corresponding PON fluxes.
 The black regression line shows the correlation between measured POC and PON fluxes.

Formatted: Caption, Left, Space Before: 0 pt, After: 0 pt, Add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Moved (insertion) [8]

Formatted: Font: +Body (Times New Roman), 9 pt, English (US)

Formatted: Font: +Body (Times New Roman), 9 pt, Not Bold, English (US)

3. Results

3.1 Observational data

3.1.1 Settling velocity and respiration

The size-specific sinking velocities of in-situ collected aggregates from the study area were parameterized as $W(d) = A_{\text{vel}} d^{B_{\text{vel}}}$, with the best-fit parameters $A_{\text{vel}} = 121.0 [99.9, 141.3]$ (m day⁻¹) and $B_{\text{vel}} = 0.50 [0.21, 0.81]$ determined via least-squares regression (Fig. 3a). This implies that the average sinking velocity for particles sized 1 mm in the study area was approximately 121 m day⁻¹. Applying this relationship to the midpoint diameter of each particle size bin used in this study yielded sinking velocities ranging from 33 m day⁻¹ to 232 m day⁻¹ (Fig. 3b). This range represents the full particle size spectrum considered, from the UVP5 detection limit (64–80 μm) to the size bin encompassing the largest sinking particles observed in our gel traps (3.25–4.1 mm). Notably, the size-velocity relationship observed in the region aligns with results from a comprehensive size-velocity dataset (N=5655) compiled from 54 different studies (Fig. 3c), which presents an overall size-velocity relationship by increasing the degree of variability (Cael et al., 2021).

Figure 3d shows the microbial respiration-induced diffusive oxygen flux at the water–aggregate interface of the collected sinking aggregates. No evident correlation was found between the interfacial diffusive oxygen flux and particle size. The mean measured oxygen flux was $J_{\text{O}_2} = 1.15 \pm 0.16$ nmol O₂ mm⁻² day⁻¹, resulting in a surface-area–normalized particle carbon degradation rate of $A_0 = (1.38 \pm 0.19) \times 10^{-5}$ mg C mm⁻² day⁻¹.

3.1.2. Gel trap-based and UVP-derived PSDf

The PSDf data obtained from all gel traps mounted on DST are shown for all studied stations in Fig. 3e–g, with the corresponding spatiotemporally aligned UVP-derived PSDf, calculated using Eq. 3, shown in Fig. 3h–j. As a general trend, after log–log transformation, the PSDf data appeared approximately linear for the considered particle size ranges, which justifies applying a linear fit to characterize their shape. The intercept and slope of the fitted lines were within the ranges $8.6 \leq a_{\text{Gel}} \leq 11.1$ and $-2.7 \leq b_{\text{Gel}} \leq -1.6$ for the gel trap-based PSDf, and $9.6 \leq a_{\text{UVP}} \leq 12.4$ and $-2.8 \leq b_{\text{UVP}} \leq -1.9$ for the UVP-derived PSDf (Table A1). On average, the a_{UVP} (mean value: 10.5 ± 0.15) were higher than a_{Gel} (mean value: 9.8 ± 0.14), and b_{UVP} (mean value: -2.25 ± 0.05) were slightly steeper than b_{Gel} (mean value: -2.12 ± 0.06).

3.1.3 DST-based POC, PON fluxes, and C:N ratios

The measured POC and PON fluxes ranged from 35.5 to 105.4 mg C m⁻² day⁻¹ and 5.1 to 14.9 mg N m⁻² day⁻¹, respectively, with corresponding molar C:N ratios ranging from 5.7 to 11.4 (Table A1). Across all stations, the highest POC and PON fluxes occurred at 100 m depth and were significantly higher than fluxes at 200 m (Wilcoxon paired rank-sum test, $p = 0.0064$), indicating pronounced attenuation between these depths (Fig. 3k). Below 200 m, flux attenuation was weak, and

Moved (insertion) [9]

Formatted: English (US)

Moved (insertion) [10]

Formatted: Font: Not Bold, Font colour: Auto, English (US)


Formatted: Justified, Space Before: 0 pt, After: 0 pt, Add space between paragraphs of the same style, Line spacing: 1,5 lines, Don't keep with next, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Formatted: Font: Not Bold, English (US)

Moved (insertion) [11]

Formatted: English (US)

480 average POC fluxes at 200 m and 400 m were not significantly different (Wilcoxon paired rank-sum test, $p = 0.7488$). A
 485 similar depth-dependent pattern was observed for PON fluxes (Fig. 3I). On average, molar C:N ratios increased with depth
 490 (Fig. 3m), and POC and PON fluxes were positively correlated across individual measurements (Fig. 3n).



485 **Figure 4.** Correcting UVP-derived particle fluxes for the influence of non-sinking particles with Method I, using gel trap flux data
 as a reference. (a) Ratios of uncorrected UVP-derived particle flux (which is biased high by non-sinking particles) to the corresponding
 gel trap particle flux across particle size bins. Each point represents the ratio for one UVP–gel trap pair within a given size bin. The dashed
 black line shows the mean ratio per bin, and the dashed grey line indicates the 1:1 line (ratio = 1). (b–d) Mean PSDs from gel traps (pink
 circles) and corresponding uncorrected UVP flux estimates (blue circles) at 100 m (b), 200 m (c), and 400 m (d). The shaded blue region
 denotes the 95% confidence interval of the UVP-derived PSDf, propagated from uncertainties in the size–velocity relationship. (e–h)
 490 Ratios of log–log linear fit parameters from UVP-derived PSDf relative to the gel trap-based PSDf data: intercepts before correction (e),
 slopes before correction (f), intercepts after correction (g), and slopes after correction (h). Ratios of UVP- to gel trap-derived fluxes across
 size bins before (i) and after (j) correction. (k–m) Box-and-whisker plots showing (k) the distribution of intercept ratios and (l) the
 distribution of slope ratios (both derived from the bootstrap analysis), and (m) the distribution of log-transformed flux ratios corresponding
 to the data in panels (i) and (j).

Moved (insertion) [12]

Formatted: Font: 9 pt, English (US)

3.2 Particle flux estimate from Method I

The uncorrected UVP-derived particle fluxes were systematically higher than those measured by the spatiotemporally aligned gel traps. The ratio of the uncorrected UVP flux to the gel trap flux averaged ca. 2.6 across all size bins and UVP-gel trap pairs (Fig. 4a). This consistent overestimation was also visually apparent when comparing the mean PSDf from both instruments at depths of 100 m, 200 m, and 400 m (Fig. 4b-d), even when including the uncertainties in the size-specific settling velocity relationship. These results confirm that UVP-based estimates of carbon flux may have a high bias if the PSDs are not corrected for the presence of non-sinking particles. It is of course important to note that the magnitude of such bias is dependent on the contribution of suspended particles versus settling particles, which is likely to be both regionally and seasonally dependent.

To address this bias, Method I was applied to adjust the log-log linear fits of the UVP-derived PSDfs. By optimizing against the gel trap data, the correction factors for the intercept and slope were determined to be $\alpha = 0.92$ (CI: 0.84, 1.02) and $\beta = 0.94$ (CI: 0.82, 1.10), respectively. The effectiveness of this correction is demonstrated in Fig. 4e-h, which shows the distribution of the slope and intercept ratios before and after the adjustment. The application of these factors using Eq. 7 yields corrected UVP-based particle flux estimates that are in much closer agreement with the reference measurements (Fig. 4i-j). As summarized in Fig. 4k-m, the correction procedure successfully centers both the mean and median of the UVP-to-gel trap ratios for slope, intercept, and total flux near the ideal value of 1.0.




Figure 5. Bootstrap analysis of optimized model parameters for particulate organic matter. The panels show the joint distributions for the parameters of: (a) the zero-order degradation model for POC (Eq. 13); (b) the PON model with dynamic C:N stoichiometry (Eq. 23); and (c, d) the first-order degradation model for POC (Eq. 20).

Moved (insertion) [13]

Formatted: Font: 9 pt, Font colour: Auto, English (US)

Formatted: Font: +Body (Times New Roman), 9 pt, Not Bold, Font colour: Auto, English (US)




Figure 6. POC and PON flux estimates, and resulting C:N ratios, from the mechanistic framework compared with DST measurements. Vertical profiles show model estimates for POC flux (a–f), PON flux (g–l), and the resulting molar C:N ratio (m–r) at the six DST stations. POC fluxes were estimated using the zero-order degradation model (Method IIa), while PON fluxes were estimated using the dynamic C:N stoichiometry model (Method III). Solid blue lines represent the median of the model estimates from the optimized bootstrap distribution, with blue shading indicating the 95% confidence interval. Black circles represent the discrete measurements from the DSTs. The errors shown indicate the relative deviation of the model estimates (blue lines) from the DST measurements. All model estimates are based on the average of two UVP profiles acquired during the deployment and recovery of each DST.

3.3 Model Performance: POC, PON, and C:N Stoichiometry

3.3.1 Performance of the mechanistic framework (Methods II & III)

The zero-order degradation model for estimating POC flux (Method IIa) was optimized against DST-based POC flux measurements. This procedure proved to be highly robust, converging for 100% of the 1000 bootstrap resamples and allowing for the reliable estimation of parameter uncertainties (Fig. 5a-b). The optimization yielded a scaling exponent of


Moved (insertion) [14]

Formatted: Font: 12 pt, Not Bold, Font colour: Auto, English (US)

Formatted: Left, Space Before: 0 pt, After: 0 pt, Add space between paragraphs of the same style, Don't keep with next, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Formatted: Font: 9 pt, Font colour: Auto, English (US)

535 $B_{\text{POC}} = 1.86$ (CI: 1.28, 2.18) and a scaling coefficient of $A_{\text{POC}} = 1.1 \times 10^{-3}$ (CI: 4.5×10^{-4} , 1.4×10^{-3}) (mg C). With an average relative error of 20%, the optimized model provided a close fit to the measured POC fluxes across all studied stations and depths (Fig. 6a-f). This indicates that the model structure is well-suited to capture the observed vertical flux attenuation patterns. The estimated profiles exhibit a classic attenuation pattern consistent with the Martin curve, showing a sharp flux decrease in the euphotic zone followed by a more gradual decline at greater depths (Martin et al., 1987).
 540 Similarly, the dynamic PON flux model (Method III), which accounts for changes in C:N stoichiometry as a function of particle age, was optimized against DST-based PON flux measurements. This yielded the parameters $A_{\text{PON}} = 9.6$ (CI: 8.3, 12) and $B_{\text{PON}} = 0.49$ (CI: 0.071, 2.1) (days). The PON model effectively captured the vertical and spatial variability in the measured flux data (24% average relative error) (Fig. 6g-l). As expected from its direct link to the carbon model, the resulting PON flux profiles exhibited a similar vertical attenuation pattern to their POC counterparts. The model's robustness is further supported by its emergent prediction of particulate C:N stoichiometry. Remarkably, despite not being directly
 545 optimized against C:N data, the model-derived C:N ratios matched DST measurements well (17% average relative error). Furthermore, they robustly captured the gradual increase with depth (Schneider et al., 2004), reflecting the biogeochemical signature of preferential nitrogen remineralization (Fig. 6m-r).



555 **Figure 7. Comparison of POC flux estimates from the mechanistic zero-order and first-order degradation models with the conventional method at two contrasting stations. (a, b) Particle size distribution and corresponding vertical POC flux profiles for a station at the core of a cyclonic eddy (M160 expedition, zone A2 in Fig. A1), characterized by an abundance of larger particles. (c, d) The same comparison for the CVOO observatory station (see Fig. A1), which had a lower relative abundance of large particles. All flux profiles represent the median estimate for each model.**

Moved (insertion) [15]

Formatted: Font: 9 pt, Font colour: Auto, English (US)

3.3.2 Comparison with alternative approaches

560 For comparison, the first-order degradation model for POC flux was likewise optimized. Similar to the zero-order model, it was numerically stable, with the optimization converging for 99% of the bootstrap ensemble (Fig. 5c-d). It yielded a scaling coefficient of $A_{\text{POC}}^* = 1.1 \times 10^{-3}$ (CI: 4.6×10^{-4} , 1.5×10^{-3}) (mg C), a scaling exponent of $B_{\text{POC}}^* = 1.88$ (CI: 1.24, 2.20), and a

565 first-order degradation rate constant of $\lambda^* = 2.3 \times 10^{-2}$ (CI: 2.7×10^{-3} , 2.0×10^{-1}) (day^{-1}), with a comparable average relative error of $\sim 19\%$. This model produced POC flux profiles and uncertainties very similar to those produced using the zero-order model in the upper 1000 m (Fig. 7).

570 Finally, the conventional method was applied to our dataset to provide a performance baseline. This approach proved to be numerically unstable during optimization. Convergence was achieved for ca. 70% of the bootstrap ensemble for POC and only 9% for PON flux. After removing the large proportion of non-converged runs, the optimization yielded the parameters $A_{\text{conv}}(\text{POC}) = 1.3 \times 10^{-2}$ (CI: 4.4×10^{-4} , 3.8×10^{-2}) (mg C m day^{-1}) and $B_{\text{conv}}(\text{POC}) = 1.59$ (CI: 0.019, 2.2) for POC flux. For PON flux, the results were $A_{\text{conv}}(\text{PON}) = 6.2 \times 10^{-5}$ (CI: 4.8×10^{-5} , 5.9×10^{-3}) (mg N m day^{-1}) and $B_{\text{conv}}(\text{PON}) = 4.9 \times 10^{-2}$ (CI: 6.8×10^{-4} , 2.4). The estimated fluxes using the conventional method are shown in Fig. A3.

575 The apparently lower flux uncertainties of the conventional method (Fig. A3), when compared to our mechanistic approach (Fig. 6), must be interpreted with caution for two reasons. First, the conventional method exhibited poor bootstrap convergence. Consequently, its reported uncertainty does not reflect the full range of sampling variability, particularly for PON, and is therefore artificially narrow. Second, the lower scaling exponent in the conventional model, inherently suppresses the contribution of larger particles to the total flux (Fig. 7a–b). Since these large particles are the most variable in abundance, this suppression also dampens their propagated uncertainty, leading to a significant underestimation of the true total flux uncertainty.

580 Despite achieving reasonably low average relative errors (18.7% for POC, 18.0% for PON, 14.6% for C:N), the conventional model's predictions exhibited two limitations. First, it yielded consistently lower flux estimates in areas of high abundance of large particles, such as the epipelagic zone, when compared to estimations from the mechanistic framework (Fig. A3a–f, Fig. 7). Second, it failed to reproduce the biogeochemically expected trend of increasing C:N ratios with depth (Fig. A3m–r), a trend also supported by the measured DST flux data (Fig. 3m). Instead, it produced higher C:N ratios in the epipelagic zone, where POM is typically freshest and C:N ratios are expected to be lowest. This was followed by a nearly constant C:N ratio at deeper depths, where a gradual increase in the C:N ratio is expected (Schneider et al., 2004).

590 **4. Discussion**

595 **Method I.** A fundamental challenge in estimating particle and biogeochemical fluxes from in-situ cameras (ISCs) is the inability to distinguish sinking aggregates from the background pool of suspended or neutrally buoyant particles (McDonnell and Buesseler, 2012; Cael and White, 2020). This issue can be exacerbated by mesoscale ocean features that locally concentrate suspended particles (Sweeney et al., 2003). Furthermore, while the removal of zooplankton from ISC datasets is possible, it remains a laborious task, whether performed manually on large datasets or through artificial intelligence models

500 that require extensive training data (Giering et al., 2020). Our study introduces a novel approach (Method I) to overcome these challenges directly. By calibrating UVP-derived PSDf (Eq. 3) against those obtained from gel traps—which selectively capture sinking particles and enable the straightforward removal of swimmers during image analysis—we developed a quantitative correction to account for the contribution of non-sinking particles (suspended particles and zooplankton) when estimating particle flux from ISC data. Our results clearly show that including non-sinking particles in ISC-derived flux estimates leads to a systematic overestimation of the sinking particle flux (Fig. 4a). Specifically, the characteristic regression line of the uncorrected UVP-derived PSDf exhibited a higher intercept and a slightly steeper slope in log-log space, an effect primarily driven by the high concentration of small suspended particles in our study area (Fig. 4e-f).

505 A critical consideration in Method I is the inherent uncertainty of the size-velocity relationship. While the heterogeneous nature of marine aggregates makes particle size an imperfect predictor of sinking velocity (Iversen and Lampitt, 2020), it remains the most practical proxy that can be readily measured in situ (Cael et al., 2021). Crucially, our analysis reveals a key insight: the bias introduced by non-sinking particles is a far more dominant source of error than the uncertainty in the sinking velocity itself. As shown in Fig. 4b-d, the uncorrected UVP-derived particle flux (Eq. 3) systematically overestimates the sinking particle flux, a discrepancy that persists even when the full uncertainty of the size-velocity model is propagated. This result demonstrates that, despite the inherent uncertainty in size-velocity relationships (Fig. 3a), using a locally-derived, spatiotemporally aligned size-velocity relationship to calibrate ISC and gel trap observations is not only justified but also crucial for accurately isolating and correcting the primary bias from suspended particles.

515 Quantifying the standing stock of suspended particles is essential for understanding ocean particle dynamics, as it represents a vast reservoir of organic matter that fuels microbial metabolism and serves as the building blocks for larger, fast-sinking aggregates (Allredge & Jackson, 1995; Kiorboe, 2001). It can also act as an additional carbon sequestration mechanism to the sinking flux, where carbon within suspended particles is captured at depth away from exchange with the atmosphere (Boyd et al., 2019; Rogge et al. 2023). Beyond its primary role of providing corrected particle fluxes for our mechanistic POC and PON flux models (Methods II and III), Method I may also serve as a practical tool for estimating the concentration of the background suspended particle pool. For example, our data showed that uncorrected UVP-derived particle fluxes were, on average, ca. 2.6 times higher than the true sinking particle flux measured by gel traps across all particle size bins (Fig. 4a). If this overestimation in smaller size classes is assumed to be primarily from suspended particles, it implies that the suspended (and extremely slow-sinking) pool is substantial, potentially constituting up to ca. 70% of the total small particles detected by the UVP for the study area. Although this inference hinges on the assumption that gel trap measurements provide a reliable estimate of the true sinking flux, it nevertheless highlights the high abundance of suspended particles relative to sinking particles in the small size classes (e.g., Cael & White, 2020; Baker et al., 2017; Guidi et al., 2008). Indeed, such high amounts of suspended particles have been seen in the study area off Cape Verde during previous studies with pronounced intermediate nepheloid layers (e.g., Valiente et al., 2022).

Moved (insertion) [16]

Formatted: ng-star-inserted1, Font: +Body (Times New Roman), Font colour: Custom Colour (RGB(26,28,30)), English (US)

530

Method II & III. The primary contribution of this study is a mechanistic framework that moves beyond traditional statistical approaches to estimate POM fluxes from ISC data. A key advantage of this framework is its ability to integrate locally measured, size-specific parameters—such as sinking velocities and respiration rates—as direct inputs. This allows flux estimations to be tailored to the specific biogeochemical conditions of a study region, leading to more accurate and regionally relevant quantifications of carbon export.

535

The presented models for POC flux, based on both zero- and first-order degradation, accurately reproduced the observed vertical and spatial variability in the DST flux data (Fig. 6a) while also yielding physically meaningful parameters. The models' scaling exponents ($B_{\text{POC}} = 1.86$ and $B_{\text{POC}}^* = 1.88$) provide an empirical estimate of the fractal dimension (D). Crucially, these values are highly consistent with a morphology where particle carbon content scales more with surface area ($D \approx 2$) than with volume—a fundamental characteristic of porous marine snow aggregates (Alldredge and Gotschalk, 1988; Ploug and Passow, 2007; Burd and Jackson, 2009; Cael and White, 2020). By independently converging on a fractal dimension that reflects this key scaling characteristic, our mechanistic framework demonstrates its ability to capture the underlying physics of particle structure, a significant advantage over purely statistical approaches.

540

545

The first-order kinetic model produced flux estimates and scaling exponents that were nearly identical to those from the primary zero-order model in the upper 1000 m (Fig. 7). This convergence is likely explained by the modest overall impact of degradation, as suggested by the low optimized degradation rate ($\lambda^* = 0.023 \text{ day}^{-1}$). The POC flux estimates from the two models may, however, differ at greater depths, particularly where large particles are sparse and the flux is dominated by smaller particles with long residence times (Fig. A4). While both models performed similarly, we argue that the zero-order model is mechanistically superior as it is directly parameterized by our respiration measurements, although the first-order model remains a practical alternative when such data are unavailable.

550

It is also worth noting that, although the zero-order model uses a non-negativity constraint (see Eq. 13), in practice, it is unlikely to result in zero POC mass even at great depths, as illustrated in Fig. A4. This is due to the combined effects of the low overall degradation rate and the temperature-driven suppression of microbial respiration at depth (Iversen and Ploug, 2013), an effect governed by the Q_{10} temperature response.

555

A low λ^* does not diminish the importance of degradation; rather, it highlights its context-dependency. While the effect may be minor for fast-sinking particles due to their much shorter travel times, degradation becomes a critical flux regulator for the smaller, slow-sinking particles responsible for most deep-ocean export. The optimized λ^* in our study represents a bulk degradation rate for the entire natural particle assemblage, including both organic-rich aggregates and more refractory,

560

Moved (insertion) [17]

Deleted: The eastern tropical North Atlantic (ETNA) is known for its high-productivity and low-oxygen eddies, which typically are anticyclonic mode-water and cyclonic eddies (Karstensen et al., 2015, 2017; Fiedler et al., 2016; Schütte et al., 2016a, 2016b; Pietri and Karstensen, 2018; Dilmahamad et al., 2022). When these eddies form in coastal productive waters and migrate to oligotrophic regions, they become oases of elevated biological activities and may result in high trophic transfer (Christiansen et al., 2018) and carbon export to the deep ocean (Fiedler et al., 2016; Romero et al., 2016; Hauss et al. 2015; Fischer et al., 2016, 2021). However, the spatial variability of carbon export within eddies necessitates high-resolution flux measurements of organic matter, both horizontally and vertically, to better understand their contribution to carbon export to the deep ocean. While sediment traps provide direct measurements of organic matter flux, they collect bulk samples of sinking particles from a limited area and depth. This is due to the combined effects of the low

Formatted: ng-star-inserted1, Font: +Body (Times New Roman), Font colour: Custom Colour (RGB(26,28,30)), English (US)

580 mineral-ballasted particles. It is therefore possible that carbon-specific degradation from more refractory in situ aggregates is
lower than rates from aggregates formed from laboratory-grown phytoplankton cultures, for which reported values average
~0.15 day⁻¹ (e.g., Iversen and Ploug, 2013). Indeed, our results align with findings showing that in-situ marine particles
degrade more slowly than lab-formed ones (Belcher et al., 2016), although the upper range of the confidence interval of the
585 optimized λ^* still includes these higher lab-based rates (Fig. 5d). Further support for a low λ^* value comes from a recent
study by Bressac et al. (2024), which demonstrated that microbial degradation is a secondary driver of flux attenuation
(contributing only 7–29%), subordinate to the more significant impact of zooplankton grazing in the upper water column
(Stemmann et al., 2004; Iversen et al., 2010; van der Jagt et al., 2020)—an effect inherently captured by the ISC-detected
attenuation of particle concentration with depth.

590 A notable feature of our framework is the dynamic modeling of C:N stoichiometry (Method III). Traditional biogeochemical
models often rely on static stoichiometric ratios (Moore et al., 2004; Galbraith and Martiny, 2015), a practice that fails to
account for the preferential nitrogen remineralization. In contrast, our PON model links C:N stoichiometry to particle age
(Eq. 23), providing a numerically robust representation that not only reproduces measured C:N ratios but also captures the
trend of increasing C:N with depth (Fig. 6m–r), consistent with the trend observed across a large compilation of sediment
695 trap data (see Fig. 1 in Schneider et al., 2004). This emergent property, achieved without explicit optimization against C:N
observations, validates the mechanistic integrity of the model and its potential to represent POM transformation of sinking
particles (Grossart and Ploug, 2001). Furthermore, the model's parameters, A_{PON} and B_{PON} , provide additional informative
metrics for comparing POM transformation in sinking particles across different regions. Theoretically, A_{PON} represents a
representative the asymptotic deep-ocean C:N mass ratio of a particle's refractory POM (after labile material is fully
700 degraded), while B_{PON} is the time required for the particle's C:N ratio to reach half of its saturating value. The optimized
value of B_{PON} (0.49 days) implies that it takes approximately half a day for particles to reach a molar C:N ratio of 5.6 (half
of the asymptotic value). This suggests that the bulk molar C:N ratio of fresh particles in the study area was less than the
Redfield ratio of 6.6, but that it increased quickly as the particles aged.

705 **4.3 Limitations of the conventional method and the prospect for mechanistic models.**

The conventional method is often a practical compromise between sampling effort and flux prediction and benefits from a
710 simple implementation. However, unlike our mechanistic framework, it functions mainly as an unconstrained statistical fit.
This distinction reveals several limitations, particularly when applied to the smaller datasets typical of regional or local
studies. The most immediate issue is numerical instability. This was highlighted by our bootstrap analysis, in which the
optimization converged for approximately 70% of the bootstrap ensemble for the POC flux and only 9% for the PON flux.
This suggests the conventional method requires larger sample sizes and is therefore better suited for large-scale studies than
for smaller regional studies where measured flux data are often limited. Indeed, the conventional method has been shown to

715 be effective for deriving global-scale relationships from vast datasets (Guidi et al., 2008). Although adjusting the particle
size range (i.e., applying a larger lower bound and/or a smaller upper bound) might improve numerical stability, as
720 demonstrated by Guidi et al. (2008) who limited their analysis to a 250 μm to 1.5 mm size range, this approach is subjective
and potentially problematic. It risks excluding key particle populations in areas where POM flux is primarily driven by small
(and slow-sinking) particles (Dall’Omo and Mork, 2014; Omand et al., 2015; Baker et al., 2017) or in cases where large
particles are abundant (Allredge et al., 1993). In contrast, this study applied a much wider particle size range (0.72 μm to
3.7 mm); its lower bound was set by the smallest UVP5 size bin with particle data and its upper bound was defined by the
size bin containing the largest particles found in our gel traps, thereby accounting for the contributions of both small and
large particles.

725 Beyond numerical instability, an inherent limitation lies in the conventional method’s use of a single, lumped scaling
exponent, B_{conv} . This parameter is forced to statistically account for the combined and implicit effects of particle POM mass,
sinking velocity, and the bias introduced by non-sinking particles. This approach can produce a compromised and potentially
physically implausible scaling relationship. In our dataset, this resulted in a weak scaling exponent for POC flux ($B_{\text{conv}(\text{POC})}$
= 1.59), significantly lower than the scaling exponent derived from our mechanistic framework ($B_{\text{vel}} + B_{\text{POC}} = 2.36$). Such a
low exponent down-weights the contribution of large, fast-sinking particles to the flux calculation, potentially failing to
capture the high flux variability driven by any patchy distribution of large aggregates (Fig. 7a–b). The limitation of this
730 lumped-parameter approach is particularly evident in the PON flux optimization for our dataset. Despite achieving a good
statistical fit to the PON flux observations (mean error of 16%), the median of scaling exponents from the converged
bootstrap runs was close to zero ($B_{\text{conv}(\text{PON})} = 0.05$), implying that particle size has a negligible influence on the estimated
PON flux and that only particle count is relevant. This demonstrates that the conventional method may yield a statistically
convenient fit at the expense of physical meaning—a key limitation when the goal is to understand the underlying
735 mechanisms of the biological carbon pump.

740 Another issue is that the conventional method tends to overestimate POC flux at greater depths when the flux is primarily
driven by small particles, particularly if it is optimized with only upper water column flux data (Fig. 7c–d). This occurs
because the small-particle pool includes a large fraction of suspended material and is also more subject to degradation due to
longer residence times—two factors the conventional method overlooks. Furthermore, the method lacks a direct mechanism
to model changes in POM stoichiometry. By treating POC and PON fluxes with the same functional form (Eq. 1), it cannot
capture the preferential nitrogen remineralization, resulting in a C:N ratio that is effectively static with depth (Fig. A3m–r).

745 Hence, by incorporating mechanistic processes into the conventional method, we can not only more reliably estimate POM
fluxes but also better understand the driving mechanisms of the biological carbon pump and examine processes such as

settling and microbial degradation individually for specific regional and seasonal datasets. Process-based approaches, however, require additional sampling effort, supplementary data, and more sophisticated implementation.

5. Conclusion

750 This study presents a novel, numerically stable methodological framework that represents a significant step forward in
estimating POM fluxes from in-situ camera systems. By incorporating key mechanistic processes, our approach moves
beyond the limitations of conventional statistical methods to improve the accuracy and physical grounding of flux estimates.
755 The core contributions of this framework are: (i) a robust method for correcting for the bias of non-sinking particles when
estimating particle flux; (ii) a mechanistic model for POC flux that explicitly incorporates size-specific sinking velocities and
depth-dependent carbon degradation; and (iii) a dynamic model for PON flux that accounts for preferential nitrogen
remineralization by linking C:N stoichiometry to particle age.

By integrating locally measured parameters for particle dynamics and respiration, our framework enables reliable
estimations of POC and PON fluxes and their spatiotemporal variability. This capability is particularly valuable for
760 quantifying export fluxes locally or within dynamic mesoscale features such as eddies—where flux measurements are often
limited—and can help improve the representation of these fluxes in biogeochemical models (Turner, 2015).

With the growing number of ISC profiles being collected across the global ocean (Kiko et al., 2022), the adoption of such
mechanistic approaches has the potential to significantly improve the accuracy of regional and global carbon export
765 estimates. This could help constrain the large uncertainties that currently exist in large-scale models of carbon sequestration
(Siegel et al., 2016; Giering et al., 2017; Lombard et al., 2019; Henson et al., 2022) and ultimately enhance our
understanding of the efficiency of the biological carbon pump.

770

Appendix

1. Shipboard-based measurements

775 **Drifting Sediment Traps (DST).** A total of six DST deployments were conducted, four (DST1, DST4-6) were southwest and two (DST2 and DST3) northeast of the Cape Verde Islands (see Fig. A1 for the position of examined stations). The DST array consisted of a top buoy equipped with an Iridium satellite sender, an array of 14 two-liter buoyancy spheres that acted as wave attenuators, two benthos spheres (glass floats) that provided 25 kg of buoyancy each, and three sediment trap arrays each equipped with four gyroscopically mounted trap cylinders (of diameter 10 cm) deployed at 100 m, 200 m, and 400 m water depth, respectively. One trap cylinder per collection depth was fitted with a gel-filled collection cup with an ethanol-based viscous cryo-gel (Tissue-Tek, O.C.T.TM COMPOUND, Sakura) to collect and preserve settling aggregates, allowing determination of the physical structure, type, and size of individual aggregates. The three remaining trap cylinders at each collection depth were used to quantify biogeochemical fluxes (Pauli et al. 2021), including particulate organic carbon (POC) and particulate organic nitrogen (PON).

785 **Underwater Vision Profiler (UVP).** During the expedition a UVP5 camera system (Picheral et al., 2010) was mounted on the CTD-Rosette system and deployed at 71 stations to quantify the vertical abundance and size distribution of particles and aggregates (Fig. 2). This included parallel deployments with each deployment and recovery of DST, allowing for simultaneous observations by both the DST and UVP5. The UVP5 provides data on particles larger than 64 μm in equivalent spherical diameter (ESD). However, due to its pixel size of $\sim 30 \mu\text{m}$, counts of particles from ca. 120 μm upwards are generally considered to be reliable. The majority of the profiles were done down to 1000 m depth and the size-specific particle concentrations were calculated for 5 m depth bins. The UVP5 particle data and the corresponding CTD measurements of water temperature and oxygen concentration have been published in the PANGAEA data repository by Kiko et al. (2022) and Dengler et al. (2022), respectively.

790 **Marine Snow Catcher (MSC).** Eight MSC (OSIL, UK) deployments were conducted to collect in-situ formed marine aggregates, which were used to measure size-specific sinking velocity of settling particles. The in-situ formed aggregates were collected from 10 m below the chlorophyll maximum. After deployment, aggregates were allowed to sink in the MSC for three hours before they were gently collected from the base of the MSC. A total of 95 individual particles were collected, and their size-specific sinking velocities were measured on board using a flow chamber system equipped with an oxygen microsensor (Ploug and Jørgensen, 1999). The flow chamber was filled with the filtered sea water sampled by the MSC. A heating system was used to adjust the water temperature of the flow chamber to match the in-situ temperature, as indicated

800

Moved up [2]: (Buesseler et al., 2007, 2008).

Moved up [3]: 2008; Iversen et al., 2010; Ramondenc et al., 2016

Deleted: Furthermore, flux measurements using drifting sediment traps (DST) or neutrally buoyant sediment traps in deeper layers of eddies are rare. In contrast, in-situ camera systems (hereafter ISC) are operationally simple and can obtain high-resolution spatiotemporal observations of particle abundance and size distribution (PSD), including settling aggregates and zooplankton (Guidi et al., 2008; Iversen et al., 2010; Picheral et al., 2010; Kiko et al., 2022). ISC-derived depth-specific particle concentrations can serve as proxies for local particle flux in the water column. As such, calibration of the ISC-derived PSD from depths with corresponding DST flux allows for estimates of size-specific carbon flux associated with aggregates, which can be upscaled to have high vertical resolution of particulate organic matter (POM) flux through the water column (Guidi et al., ...

Deleted:). In a well-established statistical approach proposed by Guidi et al. (2008) for this calibration, hereafter referred to as the conventional method, particulate organic matter (POM) fluxes in the water column are parameterize as $\sum n_i A d_i^B$. Here, n denotes the ISC-derived depth-specific particle concentration as a function of particle size (d), with the summation performed over all considered particle size bins, indexed by i . The free parameters A and B are empirically optimized through related DST-based flux measurements. A common challenge when using ISC-derived particle profiles to estimate biogeochemical fluxes is distinguishing between settling and suspended particles in ISC images (McDonnell and Buesseler, 2012; Cael and White, 2020), as including suspended particles leads to an inaccurate estimate of sinking POM flux. Sinking velocity and carbon degradation rates of settling aggregates are crucial factors that, along with zooplankton grazing, influence the strength and efficiency of the biological carbon pump and, consequently, the flux of POM through the water column (Asper et al., 1992; Simon et al., 2002; Iversen et al., 2010, 2013, 2015; Stukel et al., 2018; Moradi et al., 2018, 2021; Iversen, 2023). However, besides the contribution of suspended particles, these two factors are not explicitly considered in the conventional method, either as input parameters to be measured regionally or as free parameters to be adjusted. Furthermore, in regions lacking DST-based flux measurements from either the upper or lower parts of the water column, the conventional method may not accurately optimize the parameters A and B . This can lead to overestimation of carbon flux at depth if only shallow DSTs are available, and vice versa. In this study, we address all these issues to more accurately determine particulate organic carbon (POC) and nitrogen (PON) fluxes through the entire water column. We achieve this by developing a novel approach that effectively integrates in-situ camera particle profiles, associated DST-based flux observations, measured particle settling velocities and respiration rates of in-situ collected aggregates, together with CTD measurements of water temperature and oxygen concentration (Fig. 1). Our approach is particularly useful for quantifying POC and PON flux fields across oceanic dynamic features, such as mesoscale eddies, where flux ... [1]

Formatted: Font: +Headings (Times New Roman)

Deleted: done

Formatted: Font: Bold, Not Italic

Deleted: 2

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold, Not Italic

by the CTD measurement of temperature profile from the collection depth for each station. The size-specific sinking velocities of 88 (out of 95) aggregates were fitted by a power law function. Seven particles were excluded: one particle (6.1 mm) was larger than the upper size-range considered in this study (64 μm - 4.1mm) for calculating POC flux, and the remaining six were excluded due to a lack of replicate measurements of sinking velocity. During the measurements of the sinking velocities of the collected aggregates in the flow chamber, the oxygen concentration profile across the water-aggregate interface of 31 randomly selected aggregates was measured using oxygen microsensor, following the procedure described in Iversen and Ploug (2013). The microsensor used was a Clark-type oxygen microelectrode with a guard cathode (Revsbech, 1989) and a tip diameter of 8-12 μm , and was calibrated in oxygen-saturated and oxygen-depleted water before mounting on a micromanipulator. The concentration was measured in increments of 50 μm with a 3 sec waiting period before and a 3 sec measuring period during each measurement. The current was measured on a multimeter (Unisense) with a 90% response time of <35 msec. The interfacial oxygen flux for each aggregate was then calculated following Moradi et al. (2021).

Deleted: of the form $W = A_w d^{B_w}$ where A_w and B_w are the fitting parameters, d is the equivalent spherical diameter (ESD, mm) of particles, and W is the settling velocity (m d^{-1}).

Gel trap image analysis. Aggregates collected in the gel cups were imaged using a high-resolution camera (2.8 $\mu\text{m px}^{-1}$) to determine the size distribution of particle flux at depths 100, 200, and 400 m for each station. To ensure clear imaging of both small and large particles, we imaged smaller subsections of each cup area at three prespecified depths of camera's focus. This process was automated using a pre-programmed XYZ-stage equipped with the camera and a motor. A custom Python package, Gel-Trap Particle Image Size Analyzer (Gel-PISA), was developed to reconstruct a single high-resolution image of the entire gel cup area. This tool facilitates precise stacking and stitching of subsection images from each gel cup (Fig. A2) and is specifically tailored for analyzing images captured by our camera system. Gel-PISA also removes background noise from each image frame, producing clean, high-quality images of individual particles. The removal of background image was achieved by analysing the grayscale value of each pixel at the three focus depths. For that, we leveraged the fact that the variation in grayscale values of pixels in the uncovered areas on the image (devoid of particles) across the three focus depths is different from those in the covered areas. Subsequently, the final gel trap image underwent conversion into a binary black and white format. Particle size was calculated from its image area, determined using the *contour* module of *opencv-python* software. This allowed for accurate quantification of particle sizes and counts within the gel trap samples.

Deleted: The average water temperature in the flow chamber during microsensor measurements across different stations was 18.10 ± 0.18 °C. An example of a measured O_2 concentration profile and its calculated corresponding diffusive flux is shown in Fig. A1....

Formatted: Font: Bold, Not Italic

Deleted: 3

Gel trap-based particle fluxes. The equivalent spherical diameter (ESD) of all particles that were collected in the gel traps was calculated using the formula $2\sqrt{A/\pi}$, where A represents the pixel area of the particle multiplied by the pixel size of the image. The swimmers, i.e., zooplankton that actively entered the trap, were visually annotated and excluded from the calculations. Given the rarity of larger particles, particularly at greater water depths, it was necessary to group them into larger size bins. The particles were sorted into logarithmically spaced size bins (Jackson et al., 1997) such that the equivalent spherical volume corresponding to the particle sizes at the left and right edges of each bin increases by 50% from one bin to

Deleted: "

Deleted: "

Deleted: "

Deleted: "

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Bold, Not Italic

Formatted: Font colour: Text 1

Deleted: $2\sqrt{A/\pi}$.

the next. Specifically, the bins were defined so that the right edge of each bin (D_{i+1}) is $\sqrt[3]{2}$ times larger than its left edge (D_i), i.e., $D_{i+1} = \sqrt[3]{2}D_i$ with $D_1 = 32 \mu\text{m}$. The index i refers to the i^{th} size bin represented by $d_i = 0.5 \times (D_{i+1} + D_i)$, the midpoint of the size bin i . The size bin i contains the number of particles whose diameters are within the range $D_i \leq d < D_{i+1}$. Using the deployment time of the sediment trap and the area of the gel trap cup (ca. 50 cm^2), the corresponding gel trap-based particle flux (F_{Gel}) was calculated in units of number of particles per square meter per day (particles $\text{m}^{-2} \text{day}^{-1}$). The obtained size distribution of particle flux (PSDF) for each gel trap was characterized by the best linear fit of $\log_{10}(F_{\text{Gel}})$ versus $\log_{10}(d)$, represented by an associated pair of intercept as exemplified. For the fitting procedure, a particle size-range from $64 \mu\text{m}$ to 2.58 mm was considered. The midpoints of the smallest ($64\text{--}80 \mu\text{m}$) and largest ($2.05\text{--}2.58 \mu\text{m}$) size bins within this range are $72 \mu\text{m}$ and 2.315 mm , respectively. Particle counts for sizes larger than 2.58 mm were considered unreliable due to the rarity of large sinking particles, especially at greater depths, and the small sampling area by the trap. If a size bin in the considered particle size range had $\log_{10}(F_{\text{Gel}}) \leq 0$ (or simply no particles in that size bin), that size bin was excluded from fitting procedure.

UVP-based particle concentration. To determine UVP-based size distribution of particle concentration (φ_{UVP}) the same size bins used for the gel traps were applied, but the first three bins were skipped and D_1 was set to $64 \mu\text{m}$ because of lower resolution of UVP camera. Due to the UVP's shorter observation time and the notable decrease in particle counts with increasing water depth, particularly for larger particles at greater water depth, a depth-specific averaging approach was implemented to statistically enhance the reliability of aggregate abundance and the size distribution. Averaging was conducted within specific vertical water layers: 5-meter layers for depths shallower than 50 m , 10-meter layers for depths between 50 m and 200 m , 15-meter layers for depths between 200 m and 400 m , and 25-meter layers for greater depths. To associate a spatiotemporally consistent φ_{UVP} counterpart to each F_{Gel} , the average of two φ_{UVP} were used. These two were extracted from the UVP profiles obtained at the times of deployment and recovery of the corresponding DST, aligning with the associated depth of respective F_{Gel} .

2. Supplementary Figures and Tables

Deleted: (Jackson et al., 2005).

Deleted:

Deleted: size distribution of

Deleted: $n_{\text{gel}}^{\text{F}}$

Deleted: (#

Deleted: d

Deleted: Each $n_{\text{gel}}^{\text{F}}$

Deleted: $\log_{10}(n_{\text{gel}}^{\text{F}})$

Deleted: (a^{gel}) and slope (b^{gel}),

Deleted: in Fig. 4a

Deleted: $n_{\text{gel}}^{\text{F}}$

Deleted: Using the characteristic a^{gel} and b^{gel} for each gel trap, the estimate of $n_{\text{gel}}^{\text{F}}$ as a function of the size bin is given by $n_{\text{gel}}^{\text{F}}(d_i) \approx 10^{a^{\text{gel}}} \times d_i^{b^{\text{gel}}}$.

Deleted: $n_{\text{UVP}}^{\text{C}}$

Formatted: Font: Bold, Not Italic

Formatted: Font: Bold

Deleted: $n_{\text{UVP}}^{\text{C}}$

Deleted: $n_{\text{gel}}^{\text{F}}$

Deleted: $n_{\text{UVP}}^{\text{C}}$

Deleted: $n_{\text{gel}}^{\text{F}}$

Deleted: *Eddy location and 3D-structure.* To locate survey stations and ship tracks relative to eddy coordinates, and to characterize the eddy dynamical regime, we used in-situ observed currents from the ship's vessel-mounted Acoustic Doppler Current Profiler (vmADCP). During Meteor cruise M160, an RDI Ocean Surveyor 75 kHz and an RDI Ocean Surveyor 38kHz (the latter only for the first half of the cruise) continuously recorded currents down to 600 m and 1200 m .

Moved up [4]:

Deleted: integration methods

Moved up [5]: The areas of the resulting trapezoids were

Formatted: Font: +Body (Times New Roman)

Deleted:

Moved up [6]: Similarly,

Formatted: Font: +Body (Times New Roman)

Deleted:

Moved up [7]: These sizes correspond to the midpoints of the

Formatted: Font: +Body (Times New Roman)

Deleted: Note that the particle fluxes ($n_{\text{gel}}^{\text{F}}$ and $n_{\text{UVP}}^{\text{F}}$) and pa

Moved up [9]: Notably, the size-velocity relationship observed in

Deleted: 5c), which presents an overall size-velocity relation




Figure A1 (a) Map of station locations within the Cape Verde archipelago region, sampled during the M160 expedition (Nov.–Dec. 2019). Arrows indicate the dominant flow directions of major current systems in the area: Canary Current (CC), North Equatorial Current (NEC), Mauritanian Current (MC), Cape Verde Current (CVC), and Poleward Undercurrent (PUC). The depiction of primary current systems is adapted from Fig. 1 in Romero and Ramondenc (2022). CVOO marks the position of the Cape Verde Ocean Observatory. (b–d) Enlarged views of stations within the three main sampling zones. The chlorophyll maps represent the average satellite-derived chlorophyll *a* concentration (cds.climate.copernicus.eu) for the sampling period within each zone: A1 (25/11–01/12/2019), A2 (25/11–03/12/2019), and A3 (10–19/12/2019). The satellite-derived surface currents are shown for the midpoint of each period.





Figure A2 Image analysis of gel trap sample using Gel-PISA. (a) High-resolution reconstruction of the entire gel trap cup image, achieved by stacking and stitching subsection frames. (b) Conversion of the final image to grayscale, followed by background removal. (c) Transformation of the grayscale image into a binary format for particle counting and size calculation. (d) Examples of particle images extracted from the gel trap sample image.

- Formatted [8]
- Moved up [11]: (mean value: -2.12 ± 0.06).
- Formatted [9]
- Deleted: To account for these deviations, primarily due to [10]
- Moved up [1]: 2023).
- Formatted [11]
- Deleted: As such, when combined with DSTs, ISCs facilit [12]
- Moved down [18]: 2021).
- Formatted [13]
- Deleted: These considerations also apply to estimating PC [14]
- Moved up [16]: As shown in Fig.
- Moved up [10]: ¶
- Deleted: 8a–f, this leads the conventional method to estim [16]
- Deleted: 2. (a) Map of station locations in the ETNA,
- Formatted [15]
- Formatted [17]
- Formatted [18]
- Formatted [19]
- Deleted: In zones A1, A2, and A3, we sampled a frontal z [20]
- Formatted [21]
- Deleted: ¶ [22]
- Formatted [23]
- Formatted [24]
- Deleted: 3.
- Formatted [25]
- Deleted:
- Formatted [26]
- Deleted:Page Break..... [27]
- Moved up [12]: Figure 4.
- Formatted [28]
- Deleted: Characterization of particle flux size distribution. [29]
- Moved up [13]: Figure 5.
- Formatted [30]
- Formatted [31]
- Deleted: (a) Measured particle sinking velocity *W* versus [32]
- Moved up [14]: ¶
- Formatted [33]
- Deleted: Mean DST-based fluxes (N=6): (a) POC flux, ([34]
- Moved up [8]:) Individual POC fluxes vs. their corresponding
- Formatted [35]
- Formatted [36]



545 **Figure A3.** POC and PON flux estimates, and resulting C:N ratios, from the conventional method (Eq. 1 in the main text) compared with
DST measurements. Vertical profiles show model estimates for POC flux (a–f), PON flux (g–l), and the resulting molar C:N ratio (m–r) at
the six DST stations. Solid black lines represent the median of the model estimates from the optimized bootstrap distribution, with grey
shading indicating the 95% confidence interval. Black circles represent the discrete measurements from the DSTs. The errors shown
550 indicate the relative deviation of the model estimates (black lines) from the DST measurements. All model estimates are based on the
average of two UVP profiles acquired during the deployment and recovery of each DST.





Figure A4. Comparison of POC flux estimation using zero-order and first-order degradation models at the CVOO observatory station (see Fig. A1). (a) Relative POC mass loss of modelled particles plotted against particle size at three depths: 100 m, 1,000 m, and 3,000 m. (b) Estimated POC flux as a function of water depth for both models. The low POC flux values estimated at depth by the zero-order model are consistent with measurements from a deep moored sediment trap at the CVOO station (see Fig. 5 in Fischer et al., 2021).

Table
 Measured DST-based POC and PON fluxes with corresponding molar C:N alongside the intercept and slope of the characteristic lines fitted to the log-transformed gel trap-based size distributions of particle flux and their UVP counterparts. Note that the values for a_{UVP} and b_{UVP} are presented without correction. Both particle flux (F_{Gel} and F_{UVP}) as well as particle size d were initially normalized by $1 \text{ m}^2 \text{ day}^{-1}$ and $1 \mu\text{m}$ respectively, prior to log-transformation. Consequently, the intercepts and slopes are unitless.

DST sample reference				Measured Biogeochemical fluxes			Gel trap-based particle flux		UVP-derived particle flux	
DST label	Station	Depth (m)	Index k	POC flux ($\text{mg m}^{-2} \text{ day}^{-1}$)	PON flux ($\text{mg m}^{-2} \text{ day}^{-1}$)	Molar C:N	a_{Gel}	b_{Gel}	a_{UVP}	b_{UVP}
DST_1	D: M160-4-1	100	1	78.69	9.20	9.97	9.85	-2.11	10.49	-2.20
	R: M160-12-1	200	2	47.76	5.81	9.58	9.45	-2.03	10.29	-2.18
		400	3	48.97	5.19	10.99	9.81	-2.14	10.80	-2.38
DST_2	D: M160-87-1	100	4	65.48	13.20	5.78	9.29	-1.90	10.87	-2.37
	R: M160-87-1	200	5	35.68	6.68	6.22	9.80	-2.15	10.74	-2.35
		400	6	38.86	5.34	8.48	9.51	-2.00	10.48	-2.20
DST_3	D: M160-110-1	100	7	105.43	12.67	9.70	10.02	-2.21	12.44	-2.83
	R: M160-116-1	200	8	49.41	5.31	10.84	NA	NA	10.67	-2.27
		400	9	39.25	4.11	11.13	9.54	-2.03	10.78	-2.36
DST_4	D: M160-146-1	100	10	68.03	14.92	5.31	9.43	-1.95	10.50	-2.22
	R: M160-164-1	200	11	58.60	5.94	11.49	10.23	-2.29	9.78	-1.95
		400	12	60.29	7.94	8.84	10.51	-2.47	11.26	-2.56
DST_5	D: M160-170-1	100	13	52.03	7.39	8.20	9.96	-2.16	10.37	-2.18
		200	14	42.81	5.54	9.00	11.13	-2.69	10.83	-2.41

Moved (insertion) [18]

Deleted: -----Page Break-----



Formatted: Font: +Body (Times New Roman), 9 pt, Not Bold

Moved up [15]: Figure 7.

Moved up [17]: A4.

Formatted: Caption, Left, Line spacing: single, Keep with next

Deleted: C:N.

Formatted: Font: +Body (Times New Roman), 9 pt, Not Bold

Formatted: ... [37]

Formatted: ... [38]

Deleted: a_{UVP}

Formatted: ... [39]

Deleted: b_{UVP}

Deleted: n_{Gel}^F

Deleted: n_{UVP}^F

Deleted: 2d

Formatted: ... [40]

Formatted: ... [41]

Formatted: ... [42]

Formatted: Font: 9 pt, English (US)

Formatted: ... [43]

Formatted: ... [44]

Formatted: Font: Times New Roman, 9 pt

Deleted: 1 ... [45]

Deleted: DST2

Deleted: DST3

Deleted: DST4

Deleted: DST5

DST 6	R: M160-182-1	400	15	59.11	6.29	10.95	10.47	-2.44	10.50	-2.28
	D: M160-189-1	100	16	76.17	9.51	9.33	8.61	-1.65	9.60	-1.88
	R: M160-198-1	200	17	46.73	5.57	9.78	9.11	-1.85	10.0	-2.07
	R: M160-198-1	400	18	44.00	5.28	9.70	9.51	-2.04	9.66	-1.95

Deleted: DST6

Table A2. Particle size and sinking velocity data. Size (equivalent spherical diameter, ESD) and sinking velocity of in-situ formed aggregates collected during the M160 expedition. Measurements were conducted using the flow chamber system on board.

Station	Size (mm)	Sinking Vel. (m day ⁻¹)	Station	Size (mm)	Sinking Vel. (m day ⁻¹)	Station	Size (mm)	Sinking Vel. (m day ⁻¹)
M160-14-1	1.34	153.30	M160-71-1	0.71	88.06	M160-144-1	0.69	56.08
M160-14-1	0.95	347.90	M160-71-1	0.35	79.27	M160-144-1	0.43	59.15
M160-14-1	1.55	70.54	M160-71-1	0.87	0.63	M160-144-1	0.96	132.01
M160-14-1	1.23	197.52	M160-71-1	1.43	19.65	M160-144-1	1.26	189.87
M160-14-1	1.37	496.35	M160-71-1	0.78	32.31	M160-144-1	0.44	95.64
M160-14-1	1.42	55.58	M160-71-1	0.57	32.01	M160-144-1	0.44	174.60
M160-14-1	0.81	102.67	M160-71-1	0.98	18.75	M160-144-1	0.48	102.21
M160-14-1	1.45	124.99	M160-71-1	0.34	7.36	M160-144-1	0.58	52.43
M160-14-1	0.80	144.85	M160-91-1	0.61	67.29	M160-144-1	0.43	62.11
M160-14-1	1.89	346.97	M160-91-1	0.72	48.23	M160-144-1	0.46	25.68
M160-53-1	0.77	206.31	M160-91-1	0.37	36.90	M160-144-1	0.29	207.17
M160-53-1	0.59	324.50	M160-91-1	0.69	32.11	M160-177-1	0.54	38.12
M160-53-1	0.65	92.43	M160-91-1	0.39	37.37	M160-177-1	0.54	46.16
M160-53-1	1.04	84.47	M160-91-1	0.47	39.60	M160-177-1	0.26	41.06
M160-53-1	0.61	186.74	M160-91-1	0.82	149.51	M160-177-1	2.23	4.96
M160-53-1	0.55	262.25	M160-91-1	0.71	90.78	M160-177-1	0.46	19.18
M160-53-1	1.21	186.54	M160-91-1	0.55	25.61	M160-177-1	0.68	53.05
M160-53-1	0.54	94.52	M160-91-1	0.49	85.63	M160-177-1	1.22	20.73
M160-53-1	0.76	114.95	M160-91-1	0.62	102.32	M160-177-1	0.50	56.17
M160-53-1	0.99	179.77	M160-91-1	0.42	77.11	M160-186-1	1.00	62.64
M160-53-1	0.63	581.61	M160-112-1	0.78	145.04	M160-186-1	0.26	77.35
M160-53-1	0.58	114.42	M160-112-1	0.59	126.65	M160-186-1	0.61	24.03
M160-53-1	1.90	62.06	M160-112-1	0.27	18.77	M160-186-1	0.54	101.25
M160-71-1	1.84	193.39	M160-112-1	0.36	23.64	M160-186-1	0.28	32.26
M160-71-1	1.65	110.22	M160-112-1	0.36	176.53	M160-186-1	0.52	105.43
M160-71-1	2.17	86.74	M160-112-1	0.42	84.11	M160-186-1	0.40	25.15
M160-71-1	0.98	31.70	M160-112-1	0.33	86.00	M160-186-1	0.30	28.73
M160-71-1	1.52	261.87	M160-112-1	0.60	132.20	M160-186-1	0.39	43.21
M160-71-1	2.09	36.13	M160-112-1	0.19	29.17			
M160-71-1	0.39	11.72	M160-112-1	0.36	41.97			

Deleted: Page Break

Deleted: A3

Formatted: Font: 9 pt

Formatted: Font: 9 pt

Deleted: d

Deleted: d

Deleted: d

Table A3. Particle size and oxygen flux data. Size (equivalent spherical diameter, ESD) and interfacial diffusive flux of oxygen for in-situ formed aggregates collected during the M160 expedition. Measurements were conducted on board using the microsensor in the flow chamber.

Station	Size (mm)	Diff. O ₂ flux (nmol O ₂ cm ⁻² h ⁻¹)	Station	Size (mm)	Diff. O ₂ flux (nmol O ₂ cm ⁻² h ⁻¹)	Station	Size (mm)	Diff. O ₂ flux (nmol O ₂ cm ⁻² h ⁻¹)
M160-53-1	0.77	4.5327	M160-53-1	0.55	4.8873	M160-53-1	1.52	7.0449
M160-53-1	1.04	3.1428	M160-53-1	1.60	4.285	M160-71-1	1.84	7.2257
M160-53-1	0.61	2.3854	M160-53-1	1.25	8.8444	M160-71-1	1.65	6.9982
M160-71-1	2.17	1.1999	M160-71-1	0.98	4.8523	M160-71-1	1.52	7.5312
M160-71-1	2.09	0.8756	M160-91-1	0.61	0.9358	M160-91-1	0.72	1.4843
M160-91-1	0.82	0.7346	M160-91-1	0.71	0.2925	M160-91-1	0.49	19.5579
M160-91-1	0.62	1.5165	M160-112-1	0.78	4.4208	M160-112-1	0.59	3.5465
M160-112-1	0.27	0.8509	M160-144-1	0.69	8.0996	M160-144-1	0.43	2.4215

Deleted: Page Break

Formatted: Font: Bold, Not Italic, Font colour: Black

Deleted: A4

Formatted: Font: 9 pt

Formatted: Line spacing: single

Deleted: d

M160-144-1	0.96	5.1427	M160-144-1	1.26	6.7506	M160-144-1	0.48	8.9725
M160-177-1	0.54	4.4655	M160-177-1	0.54	6.834	M160-186-1	1.0	3.0204
M160-186-1	1.66	5.596						

605

Deleted: ¶
.....Page Break.....

Formatted: Default Paragraph Font

Formatted: Heading 1

Data availability

All data used in this study are freely accessible. The UVP, CTD, ADCP, and drifter datasets can be respectively found at the following links:

<https://doi.org/10.1594/PANGAEA.924375>, <https://doi.org/10.1594/PANGAEA.943432>,

610 <https://doi.org/10.1594/PANGAEA.943409>, and <https://doi.org/10.1594/PANGAEA.918612>.

DST-based flux measurements, MSC-based size-sinking rates, and diffusive oxygen fluxes are available in the Appendix (Tables A1–3). The computer scripts used to produce the results are available from the corresponding authors upon request.

Deleted: A2–4

Author contribution

NN analysed the data, developed and implemented the presented methods and Gel-PISA package, prepared and visualized the results, and wrote the manuscript. LH processed DST samples and measured biogeochemical fluxes. SR prepared the map of study region. CMF, LH, and NM performed in-situ sampling and measurements (MSC and DST) on board. RK contributed to processing of UVP profiles. HH prepared the UVP5 for the cruise and instructed its use on board, downloaded and pre-processed the UVP5 data and contributed to postprocessing. AK conceptualized, planned and organized the scientific programme of the multidisciplinary, multi-platform eddy studies during RV Meteor Cruise M160 and led the 620 cruise. MHI planed the sampling strategy and supervised the study. All co-authors contributed to the discussion of results, critically reviewed and approved the manuscript.

Deleted: TF contributed to the deployment of vmADCP on board, analysed the ADCP data and provided the position of the studied eddy. HH prepared the UVP5 for the cruise and instructed its use on

Deleted: preprocessed

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

625 The authors extend gratitude to the captain and crew of RV Meteor M160, as well as to Juri Knudsen for operating the UVP5 on board.

Financial support

635 Funding from the German Federal Ministry of Education and Research (BMBF) through the following projects is greatly
acknowledged: REEBUS-WP6 (Grant 03F0815D), REEBUS-WP1 (Grant 03F0815A), Project 03F0629A, CUSCO-WP5
(Grant 03F0813A), and CO2MESO (Grant 03F0876A). Additional financial support was provided by the DFG-Research
Center/Cluster of Excellence "The Ocean in the Earth System" (EXC-2077-390741603), and the Alfred Wegener Institute
640 Helmholtz Center for Polar Marine Research through the PoF IV program "Changing Earth - Sustaining our Future," Topic
6.3 of the German Helmholtz Association. NM and MHI acknowledge additional funding from the EU HORIZON-CL6
projects OceanICU (Grant 101083922) and SEA-Quester (Grant 101136480). CMF was funded by the AWI internal strategy
fund EcoPump. RK acknowledges support from the French National Research Agency through the Make Our Planet Great
Again program (ANR, #ANR-19-MPGA-0012) and funding from the German Science Foundation's Heisenberg Programme
(#KI 1387/5-1).

645

References

[Aldredge, A. L., and Jackson, G. A.: Aggregation in marine systems, Deep-Sea Res. II, 42, 1–7,
https://doi.org/10.1016/0967-0645\(95\)90003-9,1995.](https://doi.org/10.1016/0967-0645(95)90003-9)

650 [Aldredge, A. L. and Gotschalk, C.: In situ settling behavior of marine snow, Limnol. Oceanogr., 33, 339–351,
https://dx.doi.org/10.4319/lo.1988.33.3.0339, 1998.](https://dx.doi.org/10.4319/lo.1988.33.3.0339)

[Aldredge, A.L., Passow, U. and Logan, B.E.: The abundance and significance of a class of large, transparent organic
particles in the ocean. Deep Sea Research Part I: Oceanographic Research Papers, 40, .1131–1140,
655 https://doi.org/10.1016/0967-0637\(93\)90129-Q, 1993.](https://doi.org/10.1016/0967-0637(93)90129-Q)

[Baker, C. A., Henson, S. A., Cavan, E. L., Giering, S. L. C., Yool, A., Gehlen, M., Belcher, A., Riley, J. S., Smith, H. E. K.,
and Sanders, R.: Slow-sinking particulate organic carbon in the Atlantic Ocean: magnitude, flux, and potential controls,
Global Biogeochem. Cycles, 31, 1051–1065, https://dx.doi.org/10.1002/2017gb005638, 2017.](https://dx.doi.org/10.1002/2017gb005638)

660 [Belcher, A., Iversen, M., Manno, C., Henson, S. A., Tarling, G. A., and Sanders, R.: The role of particle associated microbes
in remineralization of fecal pellets in the upper mesopelagic of the Scotia Sea, Antarctica, Limnol. Oceanogr. 61, 2016,
1049–1064, https://doi: 10.1002/lno.10269, 2016.](https://doi.org/10.1002/lno.10269)

Deleted: Archibald, K. M., Siegel, D. A., and

Moved down [19]: Doney, S.

Deleted: C.: Modeling the impact of zooplankton diel vertical migration on the carbon export flux of the biological pump. Global Biogeochem. Cycles, 33, 188–199, https://dx.doi.org/10.1029/2018gb005983, 2019. Asper, V. L., Deuser, W. G., Knauer, G. A., and Lohrenz, S. E.: Rapid coupling of sinking particle fluxes between surface and deep ocean waters, Nature, 357, 670–672, https://dx.doi.org/10.1038/357670a0, 1992....

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Deleted: Briggs, N., Dall'Olmo

Deleted: Claustre, H.: Major

Deleted: fragmentation

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Deleted: regulating biological sequestration

Deleted: CO2 by

Deleted: oceans, Science, 367, 6479

Formatted: English (US)

Formatted: English (US)

Deleted: dx.

Formatted: English (US)

685 Boyd, P. W., Claustre, H., Levy, M., Siegel D. A., and Weber, T.: Multi-faceted particle pumps drive carbon sequestration in the ocean. *Nature*, 568, 327–335, <https://doi.org/10.1038/s41586-019-1098-2>, 2019.

Formatted: English (US)

685 Bressac, M., Laurenceau-Cornec, E. C., Kennedy, F., Santoro, A. E., Paul, N. L., Briggs, N., Carvalho, F., and Boyd, P. W.: Decoding drivers of carbon flux attenuation in the oceanic biological pump. *Nature*, 633, 587–593, <https://doi.org/10.1038/s41586-024-07850-x>, 2024.

Moved (insertion) [20]

Deleted: 1126/science.aay1790

Formatted: English (US)

690 Buesseler, K. O., Boyd, P. W., Black, E. E., and Siegel, D. A.: Metrics that matter for assessing the ocean biological carbon pump. *Proc. Natl. Acad. Sci. U.S.A.*, 117, 9679–9687, <https://doi.org/10.1073/pnas.1918114117>, 2020.

Formatted: English (US)

Formatted: English (US)

695 Buesseler, K. O., Antia, A. N., Chen, M., Fowler, S. W., Gardner, W. D., Gustafsson, O., Harada, K., Michaels, A. F., Rutgers van der Loeff, M., Sarin, M., Steinberg, D. K., and Trull, T.: An assessment of the use of sediment traps for estimating upper ocean particle fluxes, *J. Mar. Res.*, 65, 345–416, <https://dx.doi.org/10.1357/002224007781567621>, 2007.

695 Buesseler, K. O., Lamborg, C., Cai, P., Escoube, R., Johnson, R., Pike, S., Masque, P., McGillicuddy, D., and Verdeny, E.: Particle fluxes associated with mesoscale eddies in the Sargasso Sea, *Deep Sea Res. Part II*, 55, 1426–1444, <https://dx.doi.org/10.1016/j.dsr2.2008.02.007>, 2008.

Formatted: English (US)

700 Burd, A. B. and Jackson, G. A.: Particle Aggregation, *Annual Review of Marine Science*, 1, <https://doi.org/10.1146/annurev.marine.010908.163904>, 65-90, 2009.

Moved (insertion) [21]

Formatted: English (US)

Formatted: English (US)

705 Cael, B. B., Cavan, E. L., and Britten, G. L.: Reconciling the size-dependence of marine particle sinking speed, *Geophys. Res. Lett.*, 5, e2020GL091771, <https://dx.doi.org/10.1029/2020gl091771>, 2021.

Moved up [21]: Cael, B. B., Cavan, E. L., and Britten, G. L.: Reconciling the size-dependence of marine particle sinking speed, *Geophys. Res. Lett.*, 5, e2020GL091771, <https://dx.doi.org/10.1029/2020gl091771>, 2021.

705 Cael, B. B. and White, A. E.: Sinking versus suspended particle size distributions in the North Pacific Subtropical Gyre, *Geophys. Res. Lett.*, 47, e2020GL087825, <https://dx.doi.org/10.1029/2020gl087825>, 2020.

Moved down [22]: J.,

Deleted: Carrasco, R., Horstmann,

710 Clements, D. J., Yang, S., Weber, T., McDonnell, A. M. P., Kiko, R., Stemmann, L., and Bianchi, D.: New estimate of organic carbon export from optical measurements reveals the role of particle size distribution and export horizon, *Global Biogeochem. Cycles*, 37, e2022GB007633, <https://dx.doi.org/10.1029/2022gb007633>, 2023.

Deleted: and Körtzinger, A.: Surface drifter data from Meteor cruise M160, PANGAEA, <https://doi.org/10.1594/PANGAEA.918612>, 2020.

Formatted: English (US)

Formatted: English (US)

715 Dall’Olmo, G. and Mork, K. A.: Carbon export by small particles in the Norwegian Sea, *Geophys. Res. Lett.*, 41, 2921–2927, <https://dx.doi.org/10.1002/2014gl059244>, 2014.

Deleted: Christiansen, S., Hoving, H.-J., Schütte, F., Hauss, H., Karstensen, J., Körtzinger, A., Schröder, S.-M., Stemmann, L., Christiansen, B., Picheral, M., Brandt, P., Robison, B., Koch, R., and Kiko, R.: Particulate matter flux interception in oceanic mesoscale eddies by the polychaete *Poecobius* sp., *Limnol. Oceanogr.*, 63, 2093–2109, <https://dx.doi.org/10.1002/lno.10926>, 2018...

Formatted: English (US)

- Dengler, M., Körtzinger, A., and Krahnemann, G.: Physical oceanography (CTD) during METEOR cruise M160, PANGAEA, <https://doi.pangaea.de/10.1594/PANGAEA.943432>, 2022.
- 735 DeVries, T. and Weber, T.: The export and fate of organic matter in the ocean: New constraints from combining satellite and oceanographic tracer observations, *Global Biogeochem. Cycles*, 31, 535–555, <https://dx.doi.org/10.1002/2016gb005551>, 2017.
- 740 Fischer, G., Romero, O. E., Karstensen, J., Baumann, K.-H., Moradi, N., Iversen, M., Ruhland, G., Klann, M., and Körtzinger, A.: Seasonal flux patterns and carbon transport from low-oxygen eddies at the Cape Verde [Ocean Observatory](#): lessons learned from a time series sediment trap study (2009–2016). *Biogeosciences*, 18, 6479–6500, <https://dx.doi.org/10.5194/bg-18-6479-2021>, 2021.
- 745 [Galbraith, E.D. and Martiny, A.C.: A simple nutrient-dependence mechanism for predicting the stoichiometry of marine ecosystems. *Proc. Natl. Acad. Sci. U.S.A.*, 112, 8199–8204. <https://doi.org/10.1073/pnas.1423917112>, 2015.](#)
- Giering, S. L. C., Cavan, E. L., Basedow, S. L., Briggs, N., Burd, A. B., Darroch, L. J., Guidi, L., Irisson, J.-O., Iversen, M. H., Kiko, R., Lindsay, D., Marcolin, C. R., McDonnell, A. M. P., Möller, K. O., Passow, U., Thomalla, S., Trull, T. W., and Waite, A. M.: Sinking organic particles in the ocean—flux estimates from in situ optical devices, *Front. Mar. Sci.*, 6, 834, <https://dx.doi.org/10.3389/fmars.2019.00834>, 2020.
- 750 Giering, S. L. C., Sanders, R., Martin, A. P., Henson, S. A., Riley, J. S., Marsay, C. M., and Johns, D. G.: Particle flux in the oceans: challenging the steady state assumption, *Global Biogeochem. Cycles*, 31, 159–171, <https://dx.doi.org/10.1002/2016gb005424>, 2017.
- 755 Guidi, L., Jackson, G. A., Stemmann, L., Miquel, J. C., Picheral, M., and Gorsky, G.: Relationship between particle size distribution and flux in the mesopelagic zone, *Deep Sea Res. Part I*, 55, 1364–1374, <https://dx.doi.org/10.1016/j.dsr.2008.05.014>, 2008.
- 760 Grossart, H.-P. and Ploug, H.: Microbial degradation of organic carbon and nitrogen on diatom aggregates, *Limnol. Oceanogr.*, 46, 267–277, <https://dx.doi.org/10.4319/lo.2001.46.2.0267>, 2001.
- 765 [Hach P. F., Marchant H. K., Krupke A., Riedel T., Meier D. V., Lavik G., Holtappels M., Dittmar T., and Kuypers M. M. M.: Rapid microbial diversification of dissolved organic matter in oceanic surface waters leads to carbon sequestration. *Sci. Rep.*, 10, 13025, <https://doi.org/10.1038/s41598-020-69930-y>, 2020.](#)

Formatted: English (US)

Deleted: 2022a

Formatted: English (US)

Deleted: Dengler, M., Fischer, T., Körtzinger, A., and Krahnemann, G.: ADCP current measurements (38 and 75 kHz) during METEOR cruise M160, PANGAEA, <https://doi.org/10.1594/PANGAEA.943409>, 2022b.

Formatted: English (US)

Deleted: Dilmahamad, A. F., Karstensen, J., Dietze, H., Löptien, U., and Fennel, K.: Generation mechanisms of mesoscale eddies in the Mauritanian upwelling region, *J. Phys. Oceanogr.*, 52, 161–182, <https://dx.doi.org/10.1175/jpo-d-21-0092.1>, 2022.

Deleted: Early, J. J., Samelson, R. M., and Chelton, D. B.: The evolution and propagation of quasigeostrophic ocean eddies, *J. Phys. Oceanogr.*, 41, 1535–1555, <https://doi.org/10.1175/2011JPO4601.1>, 2011.

Deleted: Fiedler, B., Grundle, D. S., Schütte, F., Karstensen, J., Löscher, C. R., Hauss, H., Wagner, H., Loginova, A., Kiko, R., Silva, P., Tanhua, T., and Körtzinger, A.: Oxygen utilization and downward carbon flux in an oxygen-depleted eddy in the eastern tropical North Atlantic, *Biogeosciences*, 13, 5633–5647, <https://dx.doi.org/10.5194/bg-13-5633-2016>, 2016.

Deleted: Fischer, G., Karstensen, J., Romero, O., Baumann, K.-H., Donner, B., Heffer, J., Mollenhauer, G., Iversen, M., Fiedler, B., Monteiro, I., and Körtzinger, A.: Bathypelagic particle flux signatures from a suboxic eddy in the oligotrophic tropical North Atlantic: production, sedimentation, and preservation, *Biogeosciences*, 13, 3203–3223, <https://dx.doi.org/10.5194/bg-13-3203-2016>, 2016....

Formatted: English (US)

Deleted: OceanObservatory

Formatted: English (US)

795 Henson, S. A., Sanders, R., Madsen, E., Morris, P. J., Le Moigne, F., and Quartly, G. D.: A reduced estimate of the strength of the ocean's biological carbon pump, *Geophys. Res. Lett.*, **38**, L04606, doi:10.1029/2011GL046735, 2011.

800 Iversen, M. H., Laufkötter, C., Leung, S., Giering, S. L. C., Palevsky, H. I., and Cavan, E. L. C.: Uncertain response of ocean biological carbon export in a changing world, *Nat. Geosci.*, **15**, 248–254, <https://dx.doi.org/10.1038/s41561-022-00927-0>, 2022.

805 Iversen, M. H. and Ploug, H.: Ballast minerals and the sinking carbon flux in the ocean: carbon-specific respiration rates and sinking velocity of marine snow aggregates, *Biogeosciences*, **7**, 2613–2624, <https://dx.doi.org/10.5194/bg-7-2613-2010>, 2010.

810 Iversen, M. H. and Ploug, H.: Temperature effects on carbon-specific respiration rate and sinking velocity of diatom aggregates – potential implications for deep ocean export processes, *Biogeosciences*, **10**, 4073–4085, <https://dx.doi.org/10.5194/bg-10-4073-2013>, 2013.

815 Iversen, M. H.: Carbon export in the ocean: A biologist's perspective, *Annu. Rev. Mar. Sci.*, **15**, 357–381, <https://dx.doi.org/10.1146/annurev-marine-032122-035153>, 2023.

Iversen, M. H. and Lampitt, R. S.: Size does not matter after all: No evidence for a size-sinking relationship for marine snow, *Prog. Oceanogr.*, **189**, 102445, <https://dx.doi.org/10.1016/j.pocan.2020.102445>, 2020.

Jackson, G. A., Maffione, R., Costello, D. K., Alldredge, A. L., Logan, B. E., and Dam, H. G.: Particle size spectra between 1 μm and 1 cm at Monterey Bay determined using multiple instruments. *Deep-Sea Res.*, **144**, 1739–1767, [https://doi.org/10.1016/S0967-0637\(97\)00029-0](https://doi.org/10.1016/S0967-0637(97)00029-0), 1997.

Kiko, R., Picheral, M., Antoine, D., Babin, M., Berline, L., Biard, T., Boss, E., Brandt, P., Carlotti, F., Christiansen, S., Coppola, L., de la Cruz, L., Diamond-Riquier, E., Durrieu de Madron, X., Elineau, A., Gorsky, G., Guidi, L., Hauss, H., Irisson, J.-O., Karp-Boss, L., Karstensen, J., Kim, D.-g., Lekanoff, R. M., Lombard, F., Lopes, R. M., Marec, C., McDonnell, A. M. P., Niemeyer, D., Noyon, M., O'Daly, S. H., Ohman, M. D., Pretty, J. L., Rogge, A., Searson, S., Shibata,

Moved (insertion) [22]

Formatted: English (US)

Moved (insertion) [23]

Deleted: Hauss, H., Christiansen, S., Schütte, F., Kiko, R., Lima, M. E., Rodrigues, 38, L04606, doi:10.1029/2011GL046735, 2011.

Formatted: English (US)

Moved up [20]: E.,

Deleted: Karstensen, J., Löscher, C. R., Körtzinger, A., and Fiedler, B.: Dead zone or oasis in the open ocean? Zooplankton distribution and migration in low-oxygen medowater eddies, *Biogeosciences*, **13**, 1977–1989, <https://dx.doi.org/10.5194/bg-13-1977-2016>, 2016....

Formatted: English (US)

Moved down [24]: Mar.

Deleted: Iversen, M. H. and Robert, M. L.: Ballasting effects of smectite on aggregate formation and export from a natural plankton community, *Mar. Chem.*, **175**, 18–27,

Deleted: Chem., **175**, 18–27, <https://dx.doi.org/10.1016/j.marchem.2015.04.009>, 2015....

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Deleted: Waite

Deleted: M

Deleted: Boyd, P. W.: Role of algal aggregation in vertical carbon export during SOIREE Dam, H. G.: Particle size spectra between

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Deleted: in other low biomass environments. *Geophys.*

Deleted: Res., **13**, L13607

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Deleted: 1029/2005GL023180, 2005

Formatted: English (US)

Deleted: Karstensen, J., Fiedler, B., Schütte, F., Brandt, P., Körtzinger, A., Fischer, G., Zantopp, R., Hahn, J., Visbeck, M. ... [46]

Formatted: English (US)

860 M., Tanaka, Y., Tanhua, T., Taucher, J., Trudnowska, E., Turner, J. S., Waite, A., and Stemmann, L.: A global marine particle size distribution dataset obtained with the Underwater Vision Profiler 5, *Earth Syst. Sci. Data*, 14, 4315–4337, <https://doi.org/10.5194/essd-14-4315-2022>, 2022.

Formatted: English (US)

865 Kiko, R., Biastoch, A., Brandt, P., Cravatte, S., Hauss, H., Hummels, R., Kriest, I., Marin, F., McDonnell, A. M. P., Oschlies, A., Picheral, M., Schwarzkopf, F. U., Thurnherr, A. M., Stemmann, L.: Biological and physical influences on marine snowfall at the equator, *Nat. Geosci.*, 10, 852, <https://dx.doi.org/10.1038/ngeo3042>, 2017.

Formatted: English (US)

870 Kiko, R., Picheral, M., Antoine, D., Babin, M., Berline, L., Biard, T., Boss, E., Brandt, P., Carlotti, F., Christiansen, S., Coppola, L., de la Cruz, L., Diamond-Riquier, E., Durrieu de Madron, X., Elineau, A., Gorsky, G., Guidi, L., Hauss, H., Irisson, J.-O., Karp-Boss, L., Karstensen, J., Kim, D.-g., Lekanoff, R. M., Lombard, F., Lopes, R. M., Marec, C., McDonnell, A. M. P., Niemeyer, D., Noyon, M., O'Daly, S. H., Ohman, M. D., Pretty, J. L., Rogge, A., Searson, S., Shibata, M., Tanaka, Y., Tanhua, T., Taucher, J., Trudnowska, E., Turner, J. S., Waite, A., Stemmann, L.: A global marine particle size distribution dataset obtained with the Underwater Vision Profiler 5, *PANGAEA*, <https://doi.org/10.1594/PANGAEA.924375>, 2022.

Formatted: English (US)

875 Kjørboe, T.: Formation and fate of marine snow: small-scale processes with large-scale implications, *Sci. Mar.*, 65, 57–71, <https://dx.doi.org/10.3989/scimar.2001.65s257>, 2001.

Formatted: English (US)

880 Lombard, F., Boss, E., Waite, A. M., Vogt, M., Uitz, J., Stemmann, L., Sosik, H. M., Schulz, J., Romagnan, J.-B., Picheral, M., Pearlman, J., Ohman, M. D., Niehoff, B., Möller, K. O., Miloslavich, P., Lara-López, A., Kudela, R., Lopes, R. M., Kiko, R., Karp-Boss, L., Jaffe, J. S., Iversen, M. H., Irisson, J.-O., Fennel, K., Hauss, H., Guidi, L., Gorsky, G., Giering, S. L. C., Gaube, P., Gallagher, S., Dubelaar, G., Cowen, R. K., Carlotti, F., Briseño-Avena, C., Berline, L., Benoit-Bird, K., Bax, N., Batten, S., Ayata, S. D., Artigas, L. F., and Appeltans, W.: Globally consistent quantitative observations of planktonic ecosystems, *Front. Mar. Sci.*, 6, 196, <https://dx.doi.org/10.3389/fmars.2019.00196>, 2019.

Formatted: English (US)

885 McDonnell, A. M. P. and Buesseler, K. O.: A new method for the estimation of sinking particle fluxes from measurements of the particle size distribution, average sinking velocity, and carbon content, *Limnol. Oceanogr. Methods*, 10, 329–346, <https://dx.doi.org/10.4319/lom.2012.10.329>, 2012.

Formatted: English (US)

890 McGillicuddy, D. J. Jr., Anderson, L. A., Bates, N. R., Bibby, T., Buesseler, K. O., Carlson, C. A., Davis, C. S., Ewart, C., Falkowski, P. G., Goldthwait, S. A., Hansell, D. A., Jenkins, W. J., Johnson, R., Kosnyrev, V. K., Ledwell, J. R., Li, Q. P., Siegel, D. A., and Steinberg, D. K.: Eddy/wind interactions stimulate extraordinary mid-ocean plankton blooms. *Science* 316, 1021–1026, <https://dx.doi.org/10.1126/science.1136256>, 2007.

895 [McGillicuddy Jr., D. J.:](#) Mechanisms of Physical-Biological-Biogeochemical Interaction at the Oceanic Mesoscale, *Annu. Rev. Mar. Sci.*, 8, 125–159, <https://dx.doi.org/10.1146/annurev-marine-010814-015606>, 2016.

Formatted: English (US)

[Moore, J. K., Donev, S. C., and Lindsay, K.:](#) Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model, *Global Biogeochem. Cycles*, 18, GB4028, <https://doi.org/10.1029/2004GB002220>, 2004.

Moved (insertion) [19]

Formatted: English (US)

900 [Moradi, N., Klawonn, I., Iversen, M. H., Wenzhöfer, F., Grossart, H.-P., Ploug, H., Fischer, G., and Khalili, A.:](#) A novel measurement-based model for calculating diffusive fluxes across substrate-water interfaces of marine aggregates, sediments and biofilms. *Front. Mar. Sci.*, 8, 68997, <https://doi.org/10.3389/fmars.2021.689977>, 2021.

Formatted: English (US)

905 [Moradi, N., Liu, B., Iversen, M., Kuypers, M. M., Ploug, H., and Khalili, A.:](#) A new mathematical model to explore microbial processes and their constraints in phytoplankton colonies and sinking marine aggregates, *Science Advances*, 4, eaat1991, <https://dx.doi.org/10.1126/sciadv.aat1991>, 2018.

Formatted: English (US)

910 [Omand, M. M., D'Asaro, E. A., Lee, C. M., Perry, M. J., Briggs, N., Cetinić, I., and Mahadevan, A.:](#) Eddy-driven subduction exports particulate organic carbon from the spring bloom, *Science*, 348, 222–225, <https://dx.doi.org/10.1126/science.1260062>, 2015.

Formatted: English (US)

915 [Pauli, N.-C., Flintrop, C. M., Konrad, C., Pakhomov, E. A., Swoboda, S., Koch, F., Wang, X.-L., Zhang, J.-C., Brierley, A. S., Bernasconi, M., Meyer, B., and Iversen, M. H.:](#) Krill and salp faecal pellets contribute equally to the carbon flux at the Antarctic Peninsula, *Nature Communications*, 12, 7168, <https://doi.org/10.1038/s41467-021-27436-9>, 2021.

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

[Picheral, M., Guidi, L., Stemmann, L., Karl, D. M., Iddaoud, G., and Gorsky, G.:](#) The underwater vision profiler 5: An advanced instrument for high spatial resolution studies of particle size spectra and zooplankton, *Limnol. Oceanogr.*, 8, 462–473, <https://dx.doi.org/10.4319/lom.2010.8.462>, 2010.

920 [Ploug, H. and Jørgensen, B. B.:](#) A net-jet flow system for mass transfer and microsensor studies of sinking aggregates. *Mar. Ecol. Prog. Ser.*, 176, 279–290, <https://dx.doi.org/10.3354/meps176279>, 1999.

Deleted: Pietri, A. and Karstensen, J.: Dynamical characterization of a low oxygen submesoscale coherent vortex in the Eastern North Atlantic Ocean, *J. Geophys. Res. Ocean.*, 123, 2049–2065, <https://dx.doi.org/10.1002/2017jc013177>, 2018....

Formatted: English (US)

925 [Ploug, H., Hietanen, S., and Kuparinen, J.:](#) Diffusion and advection within and around sinking, porous diatom aggregates. *Limnology and oceanography*, 47, 1129–1136, <https://doi.org/10.4319/lo.2002.47.4.1129>, 2002.

930 [Ploug, H. and Passow, U.: Direct measurement of diffusivity within diatom aggregates containing transparent exopolymer particles, *Limnol. Oceanogr.*, 52, 1–6, <https://dx.doi.org/10.4319/lo.2007.52.1.0001>, 2007.](#)

Formatted: English (US)

[Ploug, H. and Bergkvist, J.: Oxygen diffusion limitation and ammonium production within sinking diatom aggregates under hypoxic and anoxic conditions, *Mar. Chem.*, 176, 142–149, <https://dx.doi.org/10.1016/j.marchem.2015.08.012>, 2015.](#)

Formatted: English (US)

935 [Ramondenc, S., Madeleine, G., Lombard, F., Santinelli, C., Stemmann, L., Gorsky, G., and Guidi, L.: An initial carbon export assessment in the Mediterranean Sea based on drifting sediment traps and the Underwater Vision Profiler data sets, *Deep Sea Res. Part I*, 117, 107–119, <https://dx.doi.org/10.1016/j.dsr.2016.08.015>, 2016.](#)

Formatted: English (US)

940 [Revsbech, N. P.: An oxygen microsensor with a guard cathode. *Limnol. Oceanogr.*, 34, 474–478, <https://doi.org/10.4319/lo.1989.34.2.0474>, 1989.](#)

Formatted: English (US)

[Rogge, A., Janout, M., Loginova, N., Trudnowska, E., Hörstmann, C., Wekerle, C., et al.: Carbon dioxide sink in the Arctic Ocean from cross-shelf transport of dense Barents Sea water. *Nature Geoscience*, 16, 82–88.](#)

945 <https://doi.org/10.1038/s41561-022-01069-z>, 2023.

Romankevich, E. A.: Organic carbon in late Quaternary sediments of seas and oceans, in: *Geochemistry of Organic Matter in the Ocean*, edited by: Romankevich, E. A., Springer, Berlin, pp. 105–160, 1984, https://dx.doi.org/10.1007/978-3-642-49964-7_6.

950 [Romero, O. E. and Ramondenc, S.: A 17-year time-series of diatom populations' flux and composition in the Mauritanian coastal upwelling, *Front. Mar. Sci.*, 9, 1006345, <https://dx.doi.org/10.3389/fmars.2022.1006345>, 2022.](#)

Deleted: Romero, O. E., Fischer, G., Karstensen, J., and Cermeño, P.: Eddies as trigger for diatom productivity in the open-ocean Northeast Atlantic, *Prog. Oceanogr.*, 147, 38–48, <https://dx.doi.org/10.1016/j.pocean.2016.07.011>, 2016....

Formatted: English (US)

[Sarmiento, J. L. and Gruber, N.: *Ocean Biogeochemical Dynamics*, Princeton University Press, Princeton, NJ, 2006.](#)

955 [Simon, M., Grossart, H.-P., Schweitzer, B., and Ploug, H.: Microbial ecology of organic aggregates in aquatic ecosystems, *Aquat. Microb. Ecol.*, 28, 175–211, <https://dx.doi.org/10.3354/ame028175>, 2002.](#)

Deleted: Schütte, F., Karstensen, J., Krahnemann, G., Hauss, H., Fiedler, B., Brandt, P., Visbeck, M., and Körtzinger, A.: Characterization of “dead-zone” eddies in the eastern tropical North Atlantic, *Biogeosciences*, 13, 5865–5881, <https://doi.org/10.5194/bg-13-5865-2016>, 2016. [Schütte, F., Brandt, P., and Karstensen, J.: Occurrence and characteristics of mesoscale eddies in the tropical northeastern Atlantic Ocean, *Ocean Sci.*, 12, 663–685, <https://doi.org/10.5194/os-12-663-2016>, 2016....](#)

Formatted: English (US)

960 [Schneider, B., Engel, A., and Schlitzer, R.: Effects of depth- and CO₂-dependent C:N ratios of particulate organic matter \(POM\) on the marine carbon cycle. *Global Biogeochemical Cycles*, 18, 1, <https://doi.org/10.1029/2003GB002184>, 2004](#)

[Schneider, B., Schlitzer, R., Fischer, G., and Nöthig, E.-M.: Depth-dependent elemental compositions of particulate organic matter \(POM\) in the ocean, *Global Biogeochem. Cycles*, 17, 1032, <https://dx.doi.org/10.1029/2002gb001871>, 2003.](#)

Formatted: English (US)

980 Siegel, D. A., DeVries, T., Cetinić, I., and Bisson K., M.: Quantifying the ocean's biological pump and its carbon cycle
 impacts on global scales, *Annu. Rev. Mar. Sci.*, 15,18.1–18.28, <https://doi.org/10.1146/annurev-marine-040722-115226>,
 2023.

985 Siegel, D. A., Buesseler, K. O., Behrenfeld, M. J., Benitez-Nelson, C. R., Boss, E., Brzezinski, M. A., Burd, A., Carlson, C.
 A., D'Asaro, E. A., Doney, S. C., Perry, M. J., Stanley, R. H. R., and Steinberg, D. K.: Prediction of the export and fate of
 global ocean net primary production: The EXPORTS science plan, *Front. Mar. Sci.*, 3, 22,
<https://dx.doi.org/10.3389/fmars.2016.00022>, 2016.

990 ~~Stemmann, L., Jackson, G. A., and Gorsky, G.: A vertical model of particle size distributions and fluxes in the midwater
 column that includes biological and physical processes—Part II: application to a three year survey in the NW Mediterranean
 Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 51, 885–908, <https://doi.org/10.1016/j.dsr.2004.03.002>,
 2004.~~

995 Stukel, M. R., Song, H., Goericke, R., and Miller, A. J.: The role of subduction and gravitational sinking in particle export,
 carbon sequestration, and the remineralization length scale in the California Current Ecosystem, *Limnol. Oceanogr.*, 63,
 363–383, <https://dx.doi.org/10.1002/lno.10636>, 2018.

Sweeney, E. N., McGillicuddy, D. J., and Buesseler, K. O.: Biogeochemical impacts due to mesoscale eddy activity in the
 Sargasso Sea as measured at the Bermuda Atlantic Time Series (BATS) site, *Deep Sea Res. Part II*, 50, 3017–3039,
<https://dx.doi.org/10.1016/j.dsr2.2003.07.008>, 2003.

1000 Turner, J. T.: Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's biological pump, *Prog. Oceanogr.*, 130,
 205–248, <https://dx.doi.org/10.1016/j.pocean.2014.08.005>, 2015.

1005 ~~Valiente, S., Fernández-Castro, B., Campanero, R., Marrero-Díaz, A., Rodríguez-Santana, A., Gelado-Cabellero, M. D.,
 Nieto-Cid, M., Delgado-Huertas, A., Aristegui, J., and Álvarez-Salgado, X.A.: Dissolved and suspended organic matter
 dynamics in the Cape Verde Frontal Zone (NW Africa), *Progress in Oceanography*, 201, 102727,
<https://doi.org/10.1016/j.pocean.2021.102727>, 2022.~~

1010 van der Jagt, H., Wiedmann, I., Hildebrandt, N., Barbara, N., Iversen M. H.: Aggregate feeding by the copepods *Calanus*
 and *Pseudocalanus* controls carbon flux attenuation in the arctic shelf sea during the productive period, *Front. Mar. Sci.*, 7,
[543124](https://doi.org/10.3389/fmars.2020.543124), <https://doi.org/10.3389/fmars.2020.543124>, 2020.

Moved (insertion) [24]

Formatted: English (US)

Formatted: English (US)

Moved (insertion) [25]

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Deleted: 2002.

Formatted: English (US)

Deleted: Waite, A. M.,

015

Volk, T. and Hoffert, M. I.: Ocean carbon pumps: Analysis of relative strengths and efficiencies in ocean-driven atmospheric CO₂ changes, in: *The Carbon Cycle and Atmospheric CO₂: Natural Variations Archean to Present*, edited by: Sundquist, E. T. and Broecker, W. S., American Geophysical Union, Washington, D.C., pp. 99–110, <https://doi.org/10.1029/GM032p0099, 1985>.

Moved up [25]: Stemmann, L.,

Moved up [23]: Res. Lett.,

Deleted: Guidi, L., Calil, P. H. R., Hogg, A. Mc C., Feng, M., Thompson, P. A., Picherl, M., and Gorsky, G.: The wineglass effect shapes particle export to the deep ocean in mesoscale eddies, *Geophys. Res. Lett.*, 43, 9791–9800,

Deleted: 43, 9791–9800, <https://dx.doi.org/10.1002/2015gl066463, 2016>.

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Page 26: [1] Deleted NM 14/10/2025 08:49:00

Page 28: [2] Deleted NM 14/10/2025 08:49:00

Page 28: [3] Deleted NM 14/10/2025 08:49:00

Page 28: [4] Deleted NM 14/10/2025 08:49:00

Page 28: [5] Deleted NM 14/10/2025 08:49:00

Page 28: [6] Deleted NM 14/10/2025 08:49:00

Page 28: [7] Deleted NM 14/10/2025 08:49:00

Page 29: [8] Formatted NM 14/10/2025 08:49:00

Line spacing: single

Page 29: [9] Formatted NM 14/10/2025 08:49:00

English (US)

Page 29: [10] Deleted NM 14/10/2025 08:49:00

Page 29: [11] Formatted NM 14/10/2025 08:49:00

Font: +Headings (Times New Roman), English (US)

Page 29: [12] Deleted NM 14/10/2025 08:49:00

Page 29: [13] Formatted NM 14/10/2025 08:49:00

Font: +Body (Times New Roman), 12 pt, English (US)

Page 29: [14] Deleted NM 14/10/2025 08:49:00

Page 29: [15] Formatted NM 14/10/2025 08:49:00

ng-star-inserted1, Font: +Body (Times New Roman), Font colour: Custom Colour (RGB(26,28,30)), English (US)

Page 29: [16] Deleted NM 14/10/2025 08:49:00

Page 29: [17] Formatted NM 14/10/2025 08:49:00

Font: Not Bold, Font colour: Auto, English (US)

Page 29: [18] Formatted NM 14/10/2025 08:49:00

Font: Not Bold, English (US)

Page 29: [19] Formatted NM 14/10/2025 08:49:00

Font: 9 pt

Page 29: [20] Deleted NM 14/10/2025 08:49:00

Page 29: [21] Formatted NM 14/10/2025 08:49:00

Font: 9 pt

Page 29: [22] Deleted NM 14/10/2025 08:49:00

Page 29: [23] Formatted NM 14/10/2025 08:49:00

Font: 9 pt

Page 29: [24] Formatted NM 14/10/2025 08:49:00

Caption, Left, Line spacing: single

Page 29: [25] Formatted NM 14/10/2025 08:49:00

Font: 9 pt

Page 29: [26] Formatted NM 14/10/2025 08:49:00

Font: 9 pt

Page 29: [27] Deleted NM 14/10/2025 08:49:00

Page 29: [28] Formatted NM 14/10/2025 08:49:00

Font: 9 pt, English (US)

Page 29: [29] Deleted NM 14/10/2025 08:49:00

Page 29: [30] Formatted NM 14/10/2025 08:49:00

Font: 9 pt, Font colour: Auto

Page 29: [31] Formatted NM 14/10/2025 08:49:00

Font: +Body (Times New Roman), 9 pt, Font colour: Auto

Page 29: [32] Deleted NM 14/10/2025 08:49:00

Page 29: [33] Formatted NM 14/10/2025 08:49:00

Font: 9 pt, Font colour: Auto

Page 29: [34] Deleted NM 14/10/2025 08:49:00

Page 29: [35] Formatted NM 14/10/2025 08:49:00

Font: +Body (Times New Roman), 9 pt, English (US)

Page 29: [36] Formatted NM 14/10/2025 08:49:00

Font: +Body (Times New Roman), 9 pt, Not Bold, English (US)

Page 31: [37] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [38] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [39] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [40] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [41] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [42] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [43] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [44] Formatted NM 14/10/2025 08:49:00

Font: Times New Roman, 9 pt, Not Bold, English (US)

Page 31: [45] Deleted NM 14/10/2025 08:49:00

Page 37: [46] Deleted NM 14/10/2025 08:49:00

▼

▲