

Characteristics of ocean mesoscale eddies in the Canadian Basin from a high resolution pan-Arctic model

Noémie Planat¹, Carolina O. Dufour^{1,2}, Camille Lique², Jan K. Rieck¹, Claude Talandier², and L. Bruno Tremblay¹

¹McGill University, Departement of Atmospheric and Oceanic Sciences, Montréal, Québec, Canada

²University of Brest, CNRS, Ifremer, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, F29280, Plouzané, France

Correspondence: Noémie Planat (noemie.planat@mail.mcgill.ca)

Abstract. Mesoscale eddies are ubiquitous in the Arctic Ocean and are expected to become more numerous and energetic as sea ice continues to decline. Yet, the spatio-temporal characteristics of these eddies are poorly documented. Here, we apply an eddy detection and tracking method to the output of a high resolution ($1/12^\circ$) regional model of the Arctic - North Atlantic in order to investigate mesoscale eddies in the Canadian Basin over the period 1995-2020. Over that period, about 6000 eddies

5 per year are detected in the surface layer, while about 9000 eddies per year are detected in the pycnocline layer, and about 5500 eddies per year in the Atlantic Waters (AW) layer. The eddy population is generally distributed about equally between cyclones and anticyclones. Yet, within the pycnocline and surface layer, a clear dominance of anticyclones over cyclones is found at the centre of the Beaufort Gyre, in line with observations from Ice Tethered Profilers (ITPs). The observed dominance of anticyclonic eddies reported by ITPs thus likely partially arises from the regional focus of the ITPs. On average, eddies last
10 10 days, travel 11 km and have a radius of 12.1 km. These statistics hide strong regional and temporal disparities within the eddy population. In the surface layer, the seasonal, decadal and interannual variability in the number of eddies and in their mean characteristics follow that of the sea ice cover. In contrast, within the pycnocline layer and below, the number and properties of eddies show a weakened seasonality. At all depths, the characteristics and density of the eddy population show a strong
15 asymmetry between the slope and the centre of the Canadian Basin. Over the slope, the upper 85 m show an increase in the number of eddies, while the pycnocline layer shows a net diminution of the number of eddies. Within the AW layer, an increased number of eddies is generated in the vicinity of the cyclonic boundary current. The vast majority of eddies have no temperature nor salinity signature with respect to their environment, although a significant portion of long-lived eddies, located along
20 the Chukchi shelf break, have a non-negligible temperature anomaly and penetrate into the Beaufort Gyre, thus suggesting a mechanism for the penetration of heat into the gyre. Over the 26 years analysed here, the number of eddies generated within the upper 85 m increases by 34%, with the largest increase occurring in the open ocean and Marginal Ice Zone. Within the pycnocline layer, the number of eddies increases by 45%, with a strong year-long increase in 2008, presumably in response to the Beaufort Gyre spin-up in 2007-2008 associated with the record low in sea ice extent. The number of eddies in the Atlantic
25 Waters (AW) layer shows an overall increase of 41%, but little interannual variability. We suggest that this model-based eddy census can thus help investigate the recent changes in the dynamical equilibrium of the Beaufort Gyre by providing a consistent spatio-temporal characterization of mesoscale eddies in the Canadian basin along the past two decades.

1 Introduction

Observations and numerical models reveal that mesoscale eddies are ubiquitous in the Arctic Ocean, including under sea ice (e.g. Manley and Hunkins, 1985; Cassianides et al., 2023; Liu et al., 2024). These eddies are thought to play an important role in the transport of heat, salt and nutrients from the shelves to the deep basins (Watanabe, 2011; Watanabe et al., 2014; 30 Spall et al., 2008; Pickart et al., 2005) and possibly in the modulation of the Marginal Ice Zone (MIZ; Gupta et al., 2024; Martínez-Moreno et al., 2025; Manucharyan and Thompson, 2022). In the Canadian Basin, mesoscale eddies are also hypothesized to be a key component of the dynamical equilibrium of the large scale circulation through the dissipation of potential energy that accumulates within the anticyclonic Beaufort Gyre (BG, Manucharyan et al., 2016; Manucharyan and Spall, 2016; Meneghelli et al., 2020; Armitage et al., 2020). Additionally, eddies possibly play a role in the build-up of the subsurface heat 35 reservoir by driving the penetration of relatively warm summer Pacific Waters (sPW) within the gyre (MacKinnon et al., 2021; Spall et al., 2018; Planat et al., 2025). However, despite their possible role in the thermo-dynamical equilibrium of the BG, characteristics of the mesoscale eddy field and its evolution through time remain largely unknown in the Arctic, for one part due to the sparsity of observations, in particular under ice, and for another part due to the high resolution needed for models to represent the mesoscale at high latitude.

40

Within the Canadian Basin, measurements with Ice-Tethered Profilers (ITPs; Toole et al., 2011) and moorings which were deployed as part of the Beaufort Gyre Exploration Project (<https://www.whoi.edu/beaufortgyre>) have enabled the detection of $\mathcal{O}(400)$ eddies between 2004 and 2019 (Zhao et al., 2014; Cassianides et al., 2023). The majority of these eddies were found to lie within the halocline (50-300 m), with a few detected at greater depth. Analyses of synthetic-aperture radar data in 2007, 45 2011 and 2016 in the Western Arctic identified more than 7,500 eddies within the seasonally ice-free and MIZ regions (Kozlov et al., 2019). Similarly, altimetry-based detection within the seasonally ice-free region reported 2,000 eddies between 1993 and 2018 (Kubryakov et al., 2021), an order of magnitude difference from in situ observations, likely due to the better spatial coverage of the ice free region and MIZ. Analyses of rotating ice floes with optical satellite images were also used to provide information about the eddy population within the MIZ of the Beaufort Gyre, revealing thousands of eddy-like signatures over 50 the last two decades (Manucharyan et al., 2022). Even though no consensus was found in the eddy count across the different observational datasets, all satellite observations have shown regions densely populated with eddies over the continental shelf and slope, and in the open ocean and MIZ (Kozlov et al., 2019; Kubryakov et al., 2021), while in situ observations have demonstrated the presence of numerous eddies in the central basin below sea ice and at depth (Carpenter and Timmermans, 2012; Zhao et al., 2014; Cassianides et al., 2023).

55

The density of eddy population as well as their spatial extents (both lateral and vertical), and polarity, have provided hints at the processes driving eddy generation. In particular, the polarity has been scrutinized to better understand the hydrography and dynamics of eddies, but reconciling the picture provided by the different observation datasets has proven difficult. While in situ observations show a predominantly anticyclonic eddy field ($> 95\%$ Zhao et al., 2014; Cassianides et al., 2023), in line with op-

60 tical satellite imagery which finds twice as many anticyclonic as cyclonic floes (Manucharyan et al., 2022), altimetry shows an
equally distributed polarity (Kubryakov et al., 2021), and synthetic-aperture radar imagery shows cyclones twice as numerous
cyclones as anticyclones (Kozlov et al., 2019). Mechanisms of eddy generation were proposed to sustain the strong asymmetry
documented from in-situ observations, such as subduction processes at outcropping fronts (Manucharyan and Timmermans,
2013) and baroclinically unstable coastal boundary currents (D'Asaro, 1988; Hunkins, 1974; Zhao et al., 2014). Furthermore,
65 part of the anticyclone to cyclone asymmetry may be attributed to the stronger coherency and a slower decay of anticyclones,
a characteristic reported for eddies at lower latitudes (Chelton et al., 2011), probably leading to an over-representation of anti-
cyclones in eddy censuses (Stegner et al., 2021; Giulivi and Gordon, 2006). Contrasts between surface intensified and at depth
eddies may also impact the statistics of eddy polarity, but this is yet to be shown.

Investigating eddy sizes, in-situ observations have reported eddies across both the submeso- and meso-scales, with radii rang-
70 ing from 3 to 15 km when detected from ITP profiles and from 3 to 80 km when detected from mooring profiles (Cassianides
et al., 2023). The difference in the eddy size range may be attributed to the detection methods used; while ITPs flow "across"
eddies, moorings observe eddies passing by (Cassianides et al., 2023). On the other hand, satellite observations have typically
observed eddies with diameters ranging from $\mathcal{O}(10)$ km up to $\mathcal{O}(100)$ km (Kubryakov et al., 2021; Manucharyan et al., 2022),
75 with synthetic-aperture radar images capturing features down to $\mathcal{O}(1)$ km (Kozlov et al., 2019). Only recently has the resolu-
tion reached by realistic models become fine enough to resolve at least part of the mesoscale spectrum in the Arctic, where the
first Rossby radius of deformation varies between ≈ 15 km in the Canadian Basin and ≈ 8 km in the Eurasian Basin, down to
1 – 2 km on the shelves (Nurser and Bacon, 2013; Hu et al., 2019; Wang et al., 2020). An analysis of the Eddy Kinetic Energy
(EKE) in the entire Arctic Ocean in a 1 km resolution model shows peaks of EKE at 400 m depth at spatial scales of around
60 km. Further, this model shows that about half of the EKE is contained at scales smaller than 30 km (Liu et al., 2024).

80 On the vertical, eddies are found to form respectively at the surface, within the pycnocline layer, and at depth. The shallow
eddies are confined to the upper surface layer by the strong stratification and have a vertical extent of typically 100 m, while
eddies at depth can span up to $\mathcal{O}(1)$ km and are located around 1,200 m (Carpenter and Timmermans, 2012). In between,
double core eddies have been detected with a shallow core at the base of the pycnocline and a deep core within the Atlantic
Water layer (Zhao and Timmermans, 2015). Idealized model configurations of the BG have shown vertical modes of baroclinic
85 instabilities with similar vertical structure (Meneghelli et al., 2021). Overall, while observations have revealed different types
and origins of eddies based on their dimensions and repartitions, the number of detected features has remained relatively low,
hence preventing a systematic documentation of their spatial characteristics and geographical distribution that would allow for
more robust statistics of the eddy population. The recent advent of fine resolution ocean-sea ice models has enabled such an
investigation. Yet, it remains to be done.

90

In the Canada Basin, mesoscale eddy activity displays a strong seasonal cycle at the surface that is directly linked to that of
sea ice (Hunkins, 1974; Meneghelli et al., 2021; Manucharyan and Thompson, 2022; Rieck et al., 2025b). In ice-free regions,
thus mostly during summer, a vigorous mesoscale eddy activity is reported in both observations and models. In contrast, below
sea ice, or more generally in winter, a quiescent surface layer is observed with eddies that last as short as a few days (Meneghelli

95 et al., 2021). The short lifetime of under-ice eddies highlights the role of sea ice in dissipating eddy energy through friction. At subsurface though, eddies, shielded from the effect of sea ice by the strong stratification, may persist beyond months (D'Asaro, 1988; Hunkins, 1974). Subsurface eddy lifetime cannot be precisely estimated from observations, though, as both ITPs and moorings only capture a portion of the eddy trajectory.

100 As sea ice shrinks and the gyre intensifies in the Canada Basin, the number of mesoscale eddies is expected to increase. Satellite observations of spinning ice floes hinting at the eddy field have suggested such a trend over the past two decades (Manucharyan et al., 2022). Likewise, the number of eddies has been found to vary on interannual time scales with the intensity and freshwater content in the Beaufort Gyre (Kubryakov et al., 2021; Manucharyan et al., 2022; Zhang et al., 2016). These observations tend to confirm the suggested role of mesoscale eddies in the gyre equilibration through the conversion of potential energy, which accumulates within the freshwater reservoir at the centre of the anticyclonic BG, into eddy kinetic energy. In the same line, 105 modelling showed an enhancement of the EKE concurrent with the intensification of the gyre following increased wind forcing and sea ice retreat in 2007 (Regan et al., 2020). However, EKE was shown to only increase for a couple years in the model of Regan et al. (2020), thus questioning the mechanisms at play in the long-term equilibration of the gyre. In addition, to fulfil their role in the dynamical equilibrium of the gyre, Manucharyan and Stewart (2022) argue that eddies should be generated from baroclinic instabilities within the gyre, which cannot lead to strong polarity asymmetry in contrast to what is documented from 110 ITP measurements. To reconcile this dynamical constraint with observations, Manucharyan and Stewart (2022) further suggest that both types of eddies exist in the BG. For one hand, small and cold anticyclones travelling freely from the shelfbreak, where they are generated through coastal boundary current instabilities or outcropping fronts (Manucharyan and Timmermans, 2013; D'Asaro, 1988; Zhao et al., 2014), to the centre of the Gyre. For the other hand, larger and weaker eddies formed from baroclinic instability in the interior of the gyre that are yet to be observed from in-situ measurements.

115 Finally, the shrinking and thinning of sea ice that has been observed over the past decades and is projected to continue into the future (Meredith et al., 2001; Meier and Stroeve, 2022) will inevitably reduce the frictional dissipation of eddies, thus allowing more eddies to survive in the surface layer. The projections of the future Arctic with eddy-rich models show an increasingly energetic ocean with enhanced eddy activity in ice-free regions but also under sea ice (Rieck et al., 2025b; Li et al., 2024). The enhanced eddy activity at the surface may drive more lateral mixing of heat with potential feedback on the ice. Likewise, 120 changes in stratification may affect the eddy activity and characteristics. As suggested by Meneghelli et al. (2021), if the upper layer stratification were to weaken as the sea ice reduces, subsurface eddies that persist all year long shielded from sea ice by the strong vertical stratification may extend across reaching the surface. The evolution of the eddy characteristics over the Arctic in transition are yet to be investigated to foresee the upcoming changes in the eddy field and possible feedbacks on the ice cover.

125 Overall, no consensus has yet emerged on eddy characteristics in the Canadian Basin leaving large uncertainties that hamper robust projections of a future more energetic Arctic. In this paper, we propose a census of mesoscale eddies (i.e. vortex-like features) that develop in the Canadian Basin using a high-resolution regional model of the Arctic. To do so, we detect and track eddies to extract key properties such as size, lifetime, polarity and thermohaline anomalies. The resulting eddy dataset comprises $O(10^3)$ eddies/year thus enabling us to derive robust statistics on eddy properties. Besides, the dataset, which is

130 fully coherent in space and time, is used to document changes of eddy characteristics between 1995 and 2020, hence covering
a period of changes in strong ocean dynamical and sea ice state in the Canadian Basin. We perform a detection and tracking
of eddies, the Lagrangian framework allowing to discuss the displacement of eddies within the basin. The paper is organized
as follows. The model and the eddy detection and tracking algorithm are described in Section 2. The spatio-temporal eddy
census is presented in Section 3. A discussion of key differences with observations is offered in Section 4 together with the
135 main findings of this study and future perspectives.

2 Methods

In the rest of this paper, the Canadian basin is defined as the region between 69–85°N and 108–180°W, thus fully encompassing
the BG and its surrounding area. Note that we also define the Canada Basin, abbreviated CB, between 73–77°N and 135–
152°W, as a specific region of the BG for analysis purposes (see Fig. 1).

140 2.1 The pan-Arctic high-resolution model CREG12

2.1.1 Model and simulation

We use an updated version of the 1/12° regional Arctic-North Atlantic configuration CREG12 (Canadian Regional; Dupont
et al., 2015). CREG12 is based on the ocean modelling platform Nucleus for European Modelling of the Ocean (NEMO)
version 4.2.2 (Madec et al., 2023) and the Sea Ice modelling Integrated Initiative 3 (SI3) sea ice model, with levitating sea ice,
145 five categories of ice and two layers of snow (Vancoppenolle et al., 2023). The model is run on an ORCA12 seamless regional
grid with horizontal resolution $\approx 3 - 4$ km in the central Arctic (Barnier et al., 2014). It uses a z^* vertical coordinate with 75
levels spaced by 1 m at the surface and 150 m at 1500 m. This relatively fine horizontal grid size allows for an explicit resolution
of most of the mesoscale spectrum within the deep basins where the first Rossby radius of deformation R_o is $\approx 10 - 15$ km,
but not over the continental slope and shelf where $R_o < 7$ km (Nurser and Bacon, 2013, see also Fig. S1). Higher resolution
150 simulations of the Arctic Ocean (≈ 1 km) have shown that the EKE spectrum peaks around 50 km (Li et al., 2024) and that
more than 80% (resp. 65%) of the EKE is contained in scales larger than 10 km (resp. 20 km; Liu et al., 2024). Therefore,
we argue that 1/12° is a resolution fine enough to represent most of the mesoscale features in the Beaufort Gyre and along
its margins (but not over the shelves), while it runs at a cost that allows for decadal integration. The configuration includes a
third-order momentum flux formulation, a second-order scheme for tracers advection, with an additional bi-Laplacian viscosity,
155 diffusivity formulation depending on the local velocity, and a turbulence closure scheme for vertical mixing. The representation
of tidal mixing effects is included in the comprehensive parameterization of mixing by breaking internal tides and lee waves
(De Lavergne et al., 2016).

The simulation is initialized in 1979 from the World Ocean Atlas 2009 for temperature (Levitus et al., 2010) and salinity
(Antonov et al., 2010) with the ocean at rest and is run until 2020. Sea ice conditions are initialized from the Pan-Arctic
160 Ice Ocean Modeling and Assimilation System (PIOMAS) (Zhang and Rothrock, 2003). The ocean and sea ice are forced

with hourly atmospheric fields from the European Centre for Medium-Range Weather Forecasts Reanalysis version 5 (ERA5, Hersbach et al., 2020). To compensate for the known warm biases of ERA5 at the sea surface (e.g. Batrak and Müller, 2019), the snow conductivity is set to 0.5 W/m/K , the ice-ocean drag coefficient to $7 \cdot 10^{-3}$, the atmosphere-ocean drag coefficient to $1.2 \cdot 10^{-3}$, and the ice strength to $2 \cdot 10^{-4} \text{ N m}^{-2}$. The open boundary conditions at Bering Strait and along 27°N in the Atlantic 165 are specified daily from the output of GLORYS12V1, a global reanalysis at $1/12^\circ$ resolution run from 1993 to 2020 (Lellouche et al., 2018). Prior to 1993, output of GLORYS12V1 between 1993 and 2021 is used to build a climatology and force the open boundaries of CREG12. At Bering Strait, meridional velocities are adjusted to constrain the inflow to about 1.1 Sv, matching 170 observation estimates (Woodgate, 2018). The river run-off and Greenland melting are specified following Weiss-Gibbons et al. (2024). An additional sea surface salinity restoring with piston velocity of 167 mm day^{-1} is implemented in ice-free regions at monthly frequency using the World Ocean Atlas 2009 (Antonov et al., 2010). For additional details on the run, the reader is referred to Talandier and Lique (2024).

2.1.2 Evaluation of the simulation

We present here a brief evaluation of the model's representation of the hydrography, circulation and sea ice conditions in the Canadian Basin. For a more in-depth assessment of the model's performance, the reader is referred to Regan et al. (2020) and 175 Barton et al. (2022) who use similar configurations. In this study, we focus on the period 1995-2020 to let the model equilibrate between 1979 and 1994. Over the period of analysis, the mean September sea ice concentration is comparable to that derived from satellite observations with small differences on the Eurasian shelf and a low bias in the western CB (Fig. 1a,b). On average across the Arctic and along the simulation, the sea ice extent deviates from that derived from satellite observations by -7% in September and -16% in March. When compared to the PIOMAS Arctic Sea Ice Volume Reanalysis (Zhang and 180 Rothrock, 2003), the sea ice thickness is 35 cm thinner in September and 20 cm thinner in Mars (Fig. S2b, S3b). The interannual variability of sea ice extent is well captured by the model across the 26 years of simulation. A strong decline in sea ice starting around 2000 and persisting in time appears in the model in agreement with observations (see Fig. S2c, S3c in supplementary). The corresponding location of ice loss is generally well represented despite some biases in the ice concentration along the Eurasian shelf in summer and high biases along Yermack plateau and Greenland eastern shelf in winter (see Fig. S2a,b, S3a,b 185 in supplementary).

The geostrophic circulation of the model at the surface, estimated through the gradients of Sea Surface Height (SSH, Fig. 1c,d), shows comparable pattern and intensity as observations for the BG (anticyclonic) and for the circulation in the Nansen Basin (cyclonic). Within the CB, CREG12 successfully presents the vertical distribution of temperature extrema associated with the three main water masses present in this region (Fig. 1e,f), namely the summer Pacific Waters (sPW; temperature maximum at 190 100 m), the winter Pacific Water (wPW; temperature minimum at 200 m), and the Atlantic Water (AW; temperature maximum at 550 m). Small biases in the magnitude of the temperature extrema themselves (warm bias for the wPW, and cold bias for the sPW and AW) are noted. Despite a high salinity bias at the surface in CREG12, the modelled stratification, that allows and sustains this vertical temperature structure, displays the so-called "bowl shape" of the BG visible through the tilted isopycnals along the edges of the gyre, although slightly weaker in the northernmost side of the BG in CREG12. The overall fresh water

195 content, referenced to 34.8 psu, shows a strong increase between 2003-2009 in the Canadian Basin as documented from the Beaufort Gyre Exploration Project (Proshutinsky et al., 2009) followed by a plateau (see Fig. S4 in supplementary). Overall, the model offers a realistic representation of the main circulation features, with the anticyclonic BG extending down to ≈ 250 m and intensifying along the Chukchi shelf break (see the Mean Kinetic Energy (MKE) in Fig. S5). The cyclonic boundary current within the AW layer is found around the CB at 500 m with a returning branch of weaker intensity along the 200 Canadian Archipelago (Fig. S5c). Upper outflows through the Canadian Archipelago are similar to observation-based derived circulation (see Fig. S5, see also Planat et al., 2025). Climatologies of EKE computed relative to monthly means show larger values along the shelf break and along topographic features such as Northwind Ridge, both at the surface (not shown) and within the pycnocline (Fig. 2a). In contrast, the deep basin is more quiescent, with EKE one to two orders of magnitude lower than on the shelves (Fig. 2a,b,). The shelf-deep basin contrast in EKE magnitude is a typical feature of the mooring-based estimates 205 (von Appen et al., 2022). Yet, the intensity of EKE is about one order of magnitude smaller in our model than that derived from observations (von Appen et al., 2022), as documented previously in Regan et al. (2020). The MKE, which captures the location of the main currents, is of similar order of magnitude as in observations (von Appen et al., 2022), with discrepancies being partly attributed to the difference in the exact locations of the main currents between models and observations (Fig. S5). Finally, the vertical structure of the total kinetic energy is similar to that derived from the Beaufort Gyre Exploration Project 210 Moorings (compare Fig. S6 with for instance Fig. A1 from Meneghelli et al., 2020) with sub-surface intensified structures between 30-200 m, and deeper (although weaker) structures between 400-2000 m, as evidenced in observations by Carpenter and Timmermans (2012).

2.2 Detection and tracking of mesoscale eddies

2.2.1 Detection

215 We perform an offline detection and tracking of mesoscale eddies within the Canadian Basin over 1995-2020. Our definition of *eddies* spans here a broad range of mesoscale rotating features, from the evanescent vortices quickly dissipated by sea ice to the more persistent features that may eventually evolve into materially coherent vortices. This broad definition thus includes parts of the "turbulent soup" that is expected to develop at the surface in response to the atmospheric and ice forcings and should be captured by the model. Though short-lived, these features which are characteristic of the surface ocean, deserve an 220 investigation as they allow to examine the energy dissipation exerted by sea ice and participate to the dynamical equilibrium of the basin. In the following, we focus on features with characteristic sizes from $R_0 \approx 10$ km to $2\pi R_0 \approx 60$ km (defining the mesoscale, e.g. Tulloch et al., 2011).

To identify eddies, we use the *eddytools* python package documented in Rieck et al. (2025a). Eddies are detected using the Okubo-Weiss parameter (OW; Okubo, 1970; Weiss, 1991), which measures the relative importance of shear and strain to 225 vorticity (Fig. 3a) in the velocity field:

$$\text{OW} = (\partial_x u - \partial_y v)^2 + (\partial_x v + \partial_y u)^2 - (\partial_x v - \partial_y u)^2 \quad (1)$$




Figure 1. Mean sea ice concentration in September (background color) and in March (80 % contour in red) over 1995-2020 from (a) CREG12 and (b) National Snow and Ice Data Center (NSIDC) Climate Data Record (DiGirolamo et al., 2022), a blend between the NASA-Team algorithms (Cavalieri et al., 1984) and the NASA Bootstrap algorithm (Comiso, 1986). Mean Sea Surface Height (SSH) anomaly with respect to the mean over 2011-2020 and above 65°N from (c) CREG12 and (d) the updated altimetry-based product of Armitage et al. (2016). Black contours are evenly spaced every 0.1 m between -0.75 m and 0.75 m. Mean conservative temperature (background) and potential density referenced to surface (dashed contours) along a transect at -145°E over 2005-2014 from (e) CREG12 and (f) World Ocean Atlas 2023 climatology (Locarnini et al., 2024; Reagan et al., 2024). Note the different periods displayed for each variable to match that of the observation datasets. Boxes on panel a) represent the regions used for our analyses corresponding to the Alaskan shelf area (red) and the Canada Basin (CB, green). The thick yellow box indicates the Canadian Basin i.e. the entire domain analysed in this study. The pink line is the section used for Fig. S9. In panel b), CS and LS stand from Chukchi and Laptev Sea, respectively, and NWR, PB, AM and MK for Northwind Ridge, Pt. Barrow, Alpha-Mendeleev Ridge and McKenzie River respectively. Thin gray lines show the bathymetry, respectively 100, 500 and 1000 m depth isobaths on (a), (c), (d) and 100, 500, 1000, 2000 and 3000 m depth isobaths on (b).




Figure 2. Eddy Kinetic Energy (EKE) computed from velocity anomalies with respect to the monthly means in CREG12 and averaged over the 26 years of simulation (a) at 69 m (within the halocline) and (b) at 508 m (within the AW layer). Super-imposed are mooring-based estimates of EKE from von Appen et al. (2022), computed with fourth-order Butterworth filter with 2-day to 30-day cutoffs. The reader is referred to von Appen et al. (2022) for exact calculation method.

where u, v denote the velocities along the x and y directions of the grid, locally orthogonal. The resulting OW field (Fig. 3b) is compared to the local OW standard deviation ($\sigma_{\text{OW}}(x, y)$) averaged over the full time period (Fig. 3c). σ_{OW} is computed over a $L_\sigma \times L_\sigma$ box, L_σ being chosen small enough to capture the regional differences between e.g. the centre of the gyre and the boundary currents, but large enough so that σ_{OW} is not impacted by individual eddies. To be retained, eddies have to meet the following condition :

$$\text{OW}(x, y, t) < -\alpha \sigma_{\text{OW}}(x, y) \quad (2)$$

As we aim to detect any vortex-like features that may develop in the Canadian Basin, including those which are not materially coherent, we choose a Eulerian over a Lagrangian approach for detection. The OW-method is based on velocities (u, v) and thus preferable over SSH-based methods for detection in sea ice-covered areas where SSH-based detections are known to miss objects that do not have a surface expression. Additionally, the OW-method has the advantage to be computationally efficient and thus seems well-suited for a detection run for 26 years at each model level between the surface and 1200 m. A comparison of our OW-based detection with those from Nencioli et al. (2010, u, v - based) and Chelton et al. (2011, SSH-based) was performed by Rieck et al. (2025a, see their Fig. S3). They show that the OW-based method detects higher numbers of eddies compared to the other methods, mostly due to its capability to detect weak eddies, i.e. eddies with small rotational velocities and SSH anomaly. This detection bias towards weak eddies is commented in the discussion section.

The detection is implemented using daily averaged output in the Canadian Basin and is run for each vertical level of the model independently above 1200 m (which represents a lower bound of the AW layer), totalling 49 levels. No 3D representation of eddies is attempted here as connecting the results between the vertical layers is not trivial, and would require a substantial 245 development of the detection and tracking algorithms. A brief evaluation of the vertical structure of eddies is however proposed in Section 3.1.1. Note that because σ_{OW} is computed independently for each depth level, the minimum OW used to identify an eddy also varies with depth. In other words, at depths of intense mesoscale activity, the OW an individual eddy needs to be identified is higher. For each eddy, we estimate its radius with $R = \sqrt{area/\pi}$ even though the eddies might have an elliptic 250 shape. We set the smallest eddies that the algorithm detects to occupy 20 grid points, which correspond to equivalent circular eddies with a minimum of 5 grid points across the diameter. A 5-grid point diameter circular structure corresponds to an eddy of 7.5 to 10 km radius, depending locally on the grid size of the model, which is the lower bound of R_0 over the deep part of the Canadian Basin (see Nurser and Bacon, 2013). Over the shelf, where R_0 is smaller ($\approx 2 - 5$ km), we only detect the largest 255 of the mesoscale rotating features. Statistics presented here include all detected features, but remain valid when filtering out eddies on the shelf, as the vast majority of eddies are detected over the continental slope and within the basins (not shown).

Sensitivity tests for α show that the vertical distribution of the metrics investigated, that are mean eddy radius, duration, 255 polarity ($r_{C/T}$, ratio of cyclones to total number of eddies) and a proxy for the vorticity $|\Omega|$ (see Sect. 2.2.3), are robust to changes of α from 0.1 to 0.5. Yet, we note changes in the total number of detected features with slightly larger, weaker and shorter eddies for smaller α (Fig. S7). For our analyses, we choose $\alpha = 0.3$ as an intermediate value. The box length L_σ over 260 which to compute σ_{OW} needs to be tuned to the main spatial scales of dynamical regimes in the basin. In other words, L_σ should be small enough to capture the jet-like circulation along the Alaskan and Chukchi slopes that are about 200 – 300 km 265 large, and large enough to allow statistically relevant values of σ_{OW} . We found that $L_\sigma = 200$ km is a reasonable value to resolve the different dynamical regimes within the Canadian Basin. Similarly to α , sensitivity tests indicate that changing L_σ within the range [50, 400] km does not modify the vertical distribution of the mean eddy radius, duration, vorticity and 270 polarity although the total number of detected features vary (not shown). Overall, modifications of α and L_σ impact the precise definition of particular eddies but not the statistical properties of the eddy field.

2.2.2 Tracking

Eddies are tracked over consecutive days using three main criteria: (i) their speed of propagation, (ii) their polarity, and (iii) 270 their radius R (Fig. 3c). For each eddy detected on day t , if an eddy with similar radius (within $[0.5R, 1.5R]$) and same polarity lies within a search radius R_s on day $t+1$, it is chosen as a continuation of the track. In case there is more than one eddy matching these criteria, the one with the centre located the closest to the original eddy's centre is chosen. Results do not appear sensitive to the choice of a search radius $R_s \in [15, 53]$ km, and we choose a search radius $R_s = 22$ km corresponding to a propagation speed of 25 cm/s, which is approximately the speed of the fastest simulated current within the domain, located along the Chukchi slope in summer.

The radius, location and grid point occupied by each individual eddy are detected every day. We assume eddies to be born 275 (generated) the first time they are detected and to die (dissipated) the last time they are detected. However, the algorithm may




Figure 3. Example of detection and tracking of mesoscale eddies along the south-eastern edge of the CB at 30 m on Sept. 16th, 1996. Shown are: (a) the vorticity ($= \partial_x v - \partial_y u$), (b) the OW parameter, and (c) the OW standard deviation σ_{OW} . Superimposed on each panel are contours indicating the eddies detected by the algorithm, coloured according to (a) their duration, (b) their polarity (red indicating anticyclones and blue cyclones), and (c) their intensity i.e. the absolute value of the difference between the vorticity in the centre of the eddy and the average vorticity along the edge of the eddy. Plain contours on (a), (b) and (c) indicate eddies lasting more than 2 days, dotted red and blue contours on (b) indicate eddies with a duration of one day. Black thin lines on (c) indicate the eddy trajectories. Thin gray lines indicate the 100, 500, 1000 and 1500 m isobaths.

occasionally lose track of eddies, leading to a given eddy being counted as two successive eddies of similar properties. This interruption in the tracking generally occurs with weak features that are not well developed and thus not detected as eddies over consecutive days. We remove these eddies by filtering out any eddy that does not persist over at least two consecutive days. While this filtering does not fully overcome the interruption in the eddy detection – in particular if one eddy splits into two different ones, or equivalently if two eddies merge – it removes most of the issue and enables us to focus more on well-developed eddies. Still, the majority of eddies we detect are relatively weak and have a duration shorter than their turnaround time scale, defined as the time it takes for a water parcel to do a full revolution, $\tau = 2\pi/|\Omega|$ (i.e. an approximation of the expression suggested by Smith and Vallis, 2001, that is $\tau = 2\pi/\zeta_{rms}$, where ζ_{rms} is the root-mean-square vorticity.). A discussion of the characteristics of the more vigorous and persistent eddies is proposed in Section 4.

285 **2.2.3 Properties**

For each detected and tracked eddy, we extract its intensity, that we define as the absolute vorticity amplitude of the eddy, i.e. the difference between the vorticity in the centre of the eddy and the average vorticity along the edge of the eddy ($|\Omega|$). We equivalently report its relative intensity $|\Omega|/f(\lambda)$ where f is the Coriolis parameter computed as a function of latitude λ . By 290 spatially averaging over the eddy area, we also extract its absolute salinity (S), conservative temperature (T), and the mean sea ice concentration A and thickness h above each eddy for each day.

Eddy properties are computed and tracked along the eddy pathway. Except where mentioned, properties are extracted at the eddy generation time. Locations of eddies are obtained from their centre of mass - thus, the distance travelled by a given eddy between two consecutive days is possibly smaller than the grid resolution if an eddy is very slow. The properties of the 295 eddy environment are defined by spatially averaging over a box that we take to be $n = 3$ times larger than the eddy width and length (thus not of identical size along both x and y directions) and from which we remove the eddy area. We note $\Delta X = X_i^{eddy} - X_i^{env}$ the anomaly of property X at the time of eddy generation i . If two eddies develop next to each other, they will become each other's environment as we do not use a 2D eddy mask. To increase the robustness of the results, only *significant* anomalies are reported, except where mentioned otherwise. We first define the deviation δX from the environment noise using the standard deviation of X across the environment (i.e., excluding the eddy area), σ_X^{env} :

300

$$\delta X = \Delta X + \begin{cases} +\sigma_X^{env} & \text{if } \Delta X < 0 \\ -\sigma_X^{env} & \text{if } \Delta X > 0. \end{cases} \quad (3)$$

Then, the anomaly ΔX is said *significant* if it is of the same sign as δX , that is if the anomaly is larger in absolute than the standard deviation over the area.

Finally, the normalized amplitude of the seasonal cycle is defined for each property X as:

$$305 \quad SC_X = \frac{X_{max} - X_{min}}{\bar{X}} \quad (4)$$

where the maximum, minimum and mean are taken along the seasonal cycle.

3 Results

3.1 Characteristics of mesoscale eddies at annual and seasonal scales

3.1.1 Across the Canadian Basin

310 Over 1995-2020, and on average along the vertical, we detect and track about 6,000 eddies per year in the Canadian Basin. This large number opposes the very few vortices detected from in-situ observations below the ice ($\mathcal{O}(10)$ eddies per year, Cassianides et al., 2023; Zhao et al., 2014). It is, however, closer to numbers reported from satellite observations in the MIZ or

	Radius R	Lifetime dt	Distance travelled D	Intensity $ \Omega $	Temperature anomaly ΔT	Salinity anomaly ΔS
10^{th} perc.	9.8 km	2 day	0.6 km	$5.7 \cdot 10^{-7} \text{ s}^{-1}$	-0.10°C	-0.25 psu
90^{th} perc.	15 km	21 days	26 km	$1.1 \cdot 10^{-5} \text{ s}^{-1}$	0.20°C	0.24 psu

Table 1. Eddy characteristics defined from the 10^{th} and 90^{th} percentiles of the distribution for all eddies at all depths above 1,200 m. Percentiles of the temperature and salinity anomalies are computed on $\approx 15\%$ of the total eddy population.

the Open Ocean (up to $\mathcal{O}(1000)$ eddies per year, Kubryakov et al., 2021; Kozlov et al., 2019; Manucharyan et al., 2022). Most of the eddies detected in the model have a radius similar to the Rossby radius of deformation ($\bar{R} = 12.1$ km), are short lived 315 with an average duration $\bar{dt} = 10$ days and do not travel far with an average distance travelled $\bar{D} = 11.1$ km (Fig. 4a-c). Of all eddies detected, 49% are cyclones. Cyclones and anticyclones have a similar intensity ($|\bar{\Omega}| = 4.6 \cdot 10^{-6} \text{ s}^{-1}$ corresponding to a relative intensity of $|\bar{\Omega}|/\bar{f} = 0.03$; Fig 4b). The eddy intensity, lifetime and travelled distance have a standard deviation of the same order of magnitude as the mean. In particular, the distribution of the distance travelled shows three peaks in the 320 distribution (Fig. 4b): a first one corresponding to quasi-stationary eddies, and two secondary ones centered around 4 km and 8 km. 15% of the eddies show temperature and salinity anomalies with respect to their environment (Fig. 4e,f). The narrow and 325 short tail of the statistical distribution of ΔS indicates that the overwhelming majority of detected eddies have properties close to the mean, while the wider distribution of ΔT indicate relatively large temperature anomalies for a significant portion of the eddy population (see box whiskers on Fig. 4 and the 10^{th} and 90^{th} percentiles in Table 1). Interestingly, eddies with properties at the tail of the distributions do not represent a distinct population of eddies. For instance, larger eddies do not systematically live longer (see Fig. S8).

So far, we have presented the statistics of eddy properties aggregated over the whole 1995-2020 period and over all depth levels above 1,200 m, hence accounting for the same eddy several times if that eddy spans several depth levels. Yet, eddies may span over tens of meters to kilometres in the CB (Carpenter and Timmermans, 2012; Zhao and Timmermans, 2015). We observe some vertical coherency when looking at a few structures individually, in particular for structures spanning the 330 pycnocline between 70-250 m, or surface intensified eddies, or eddies spanning the whole water column below 200 m (Fig. 5). This vertical structure is similar to vertical structure obtained from observations (Carpenter and Timmermans, 2012; Zhao and Timmermans, 2015) or predicted from baroclinic instability estimate (Meneghelli et al., 2021) in the CB. Therefore, statistics computed over the whole eddy population hide significant differences across depth (see the coloured plain lines and \blacktriangleleft , \blacktriangleright and $+$ in Fig. 4). Within the top 1200 m, the total number of eddies generated at each model depth level remains roughly constant 335 between the surface and 85 m, and below 225 m, but increases by two thirds between 85 m and 225 m (Fig. 6a). We also note important transitions in the ratio of anticyclones versus cyclones, radius and eddy durations around these depths suggesting different mechanisms of formation and dissipation. On average across the basin and along the 26 years, these transition depths




Figure 4. Histogram of the properties of eddies generated at all depths in the model: (a) radius, (b) distance travelled, (c) duration, (d) relative intensity for cyclones (blue) and anticyclones (orange), and anomalies in (e) salinity and (f) temperature with respect to the surrounding environment (see Sect. 2). All variables are estimated at the time of eddy generation, that is the first time an eddy is detected. Number of eddies are reported in million along the left axis (indigo). Anomalies are only accounted for when significant (see Sect. 2), that is only $\approx 15\%$ of the eddy population is considered for panels (e) and (f). Box plots indicate the quartiles Q1 and Q3, the median (plain line) and mean (dotted line), and the 10^{-th} and 90^{-th} percentiles in the whiskers. Plain lines correspond to the histogram of properties at specific depths (11 m, 30 m, 69 m, 147 m and 508 m), reported along the right axis in tens of thousands of eddies (orange). On panel d), plain lines report the histogram of absolute relative intensity, that is, for both cyclones and anticyclones together. Coloured \blacktriangleleft , \triangleright and \times respectively indicate the 10^{-th}, 90^{-th} and median at the corresponding depth.




Figure 5. Daily averaged total kinetic energy (upper row) of a *virtual* mooring located at $77.94^{\circ}\text{N}, -155.34^{\circ}\text{W}$ (first column), at $70.9^{\circ}\text{N}, -134.92^{\circ}\text{W}$ (second column) and $74.28^{\circ}\text{N}, -147.94^{\circ}\text{W}$ (last column). Dashed white lines correspond to surface referenced isopycnals (σ_0). Polarity of eddies passing by the virtual mooring (second row) and identified by our algorithm with cyclones in blue and anticyclones in red. White means no eddy detected. Dots (.) indicates an interruption in the tracking, meaning that the algorithm identifies a newly born eddy.

correspond the depth at which are found the summer Pacific Waters (sPW, ~ 85 m), and the winter Pacific Waters (wPW, ~ 225 m, see Fig. 6g,h), forming together the pycnocline layer.

340 Based on the evolution of the statistical properties with depth, together with the observation of the coherent structures with finite depth extent, we thus define three layers : the upper layer (0-85 m), the pycnocline layer (85-225 m) and the AW layer (225-1200 m). Next, we describe the eddy properties and discuss their formation and dissipation within each of these three layers. The results presented in the following are robust to the exact definition of the layer boundaries (± 2 depth levels).

3.1.2 Within the upper layer

345 Within the top 85 m, about 6000 eddies are detected every year. The properties of these eddies show a marked seasonal cycle (Fig. 6) that mainly follows that of the sea ice cover. From winter to summer, the number of eddies increases by a factor of 10 with a minimum in April, just before the onset of sea ice melting, and a maximum in September when sea ice is at its minimum (Fig. 6a). The Mixed Layer (ML) depth – computed as the depth at which the potential density has increased by 0.01 kg m^{-3} compared to the potential density at 1 m – decreases from 35 m in January-May to 3 m in July, and increases from August 350 to December, when it recovers 30 m (see also the stratification on Fig. 6h). The change in stratification associated to the ML depth delimits different regimes of variations for the mean radius, polarity and intensity. Within the ML, the averaged radius of eddies increases from 12 km in early summer to 14 km in fall, while below the ML, the averaged radius of eddies barely changes comparatively (increase from 12 km to 12.5 km; Fig. 6b). In winter, a dominance of anticyclones is found at the surface (0-10 m) and of cyclones just below (10-40 m, i.e. to the base of the ML), while at the very surface eddies are essentially 355 anticyclonic year long (Fig. 6d). Below the base of the ML, the proportion between cyclones and anticyclones remains more equally distributed throughout the year, with about 55% anticyclones. Within the top 85 m, the most intense eddies are found during the stratifying and de-stratifying periods, corresponding to the onsets of sea ice growth (October and November) and to the melt season (May-July), respectively. In winter, intense eddies are also found at the base of the ML (Fig. 6c). Eddies persist longer in summer (7 – 8 days) than in winter when their lifetime is reduced by about half (Fig. 6e). Lifetimes likely 360 influence distance travelled, with eddies propagating over 15 – 16 km in summer and 7 – 8 km in winter on average (Fig. 6f). Lifetimes for the vast majority of eddies (85%) are significantly shorter than the theoretical mean turnaround time. Thus, part of the detected eddies are likely not fully developed, so belong to the "turbulent soup" that is generated in response to the surface density gradients and gets quickly dissipated by sea ice in winter. These eddy lifetimes are similar to the characteristic times of spin-down through sea ice dissipation when sea ice is taken as the main drag (e.g. $\lesssim 4$ days Meneghelli et al., 2021; 365 Pedlosky, 1982). We come back to this point in Sect. 4.

370 Eddy properties present large spatial variations across the surface layer of the Canadian Basin (Fig. 7). In particular, there is a strong contrast between the continental slope and the deep basin, with up to 10 times larger density of eddy population over the slope (≈ 150 km wide, Fig. 7a). The greater generation of eddies over the slope peaks in October, when sea ice extent is close to its minimum and winds start to increase (see Fig. S9). While the production of eddies in the deep basin remains low on average, it becomes similar to the production over the slope when sea ice concentration drops below $\approx 80\%$, that is between July and November depending on the latitude (see Fig. S9a). Simultaneously, eddy lifetime increases from evanescent ($dt = 1 – 3$ days) below the pack in winter to about 15 days in summer on average (Fig. S9b). Thus, over the domain, eddy lifetime are mainly enhanced where the ice concentration is lower than 15% (not shown).

375 Over the slope, eddies have, on average, a positive temperature anomaly ($\approx 0.3^\circ\text{C}$) with the exception of some anomalously cold eddies forming over the Chukchi shelf. Where the mean September sea ice concentration is higher than 15%, eddies do not have a temperature anomaly, aligning with Cassianides et al. (2023)'s detection of a majority of vortices with no significant




Figure 6. Seasonal cycle of eddy and basin properties with depth. (a) The number of eddies generated at each model depth level and month summed over the 26 years of simulation, and associated averaged properties: (b) radius, (c) intensity, (d) polarity $r_{C/T}$ i.e. the ratio of the number of cyclones to the total number of eddies, (e) lifetime, (f) distance travelled. Basin-averaged (g) potential temperature and (h) stratification (N^2). For panels b-h, properties are averaged at each model depth level and month over the 26 years of simulation. For panels a-f, dotted, dashed and plain lines indicate the isocontours corresponding to 5,000 10,000 and 20,000 eddies as calculated in (a), thus indicating where the statistics might be less robust due to the lower number of eddies. Note the use of a non-linear vertical axis to highlight the variability in the upper layer. Dotted horizontal grey lines delineate the three layers introduced in Section 3: upper layer, pycnocline layer and Atlantic Waters layer. The red line indicates the depth of the base of the mixed layer (ML) computed from a potential density threshold referenced to 1 m of 0.01 kg m^{-3} .

temperature anomaly. We also find a contrast in radius between the eastern and western sides of the gyre, especially off-shore the Chukchi shelf break and above Northwind ridge, where eddies are found to be about 60% larger (Fig. 7b). These eddies 380 are carried within the intense anticyclonic BG circulation (Fig. S5) and therefore travel up to 40 km throughout their lifetime , a distance much larger than the averaged distance travelled of 8 km within the CB.

Anticyclones are predominant over the centre of the gyre, while over the slope, a greater proportion of cyclones are found (Fig. 7d). Vorticity anomalies within the CB does not indicate any preference for the generation of anticyclones (not shown).

One hypothesis that explains this cyclone/anticyclone asymmetry, which has been formulated for the Mediterranean Sea and 385 more generally in other contexts of turbulent flows, is that anticyclones are more persistent than cyclones that tend to split into smaller objects, leading to anticyclones being more systematically identified in eddy censuses (Stegner et al., 2021; Giulivi and Gordon, 2006). Beech et al. (2025) further suggest the role of sea ice in preferentially dissipating small cyclones. Whether this applies to the BG is worth future investigation. We speculate that such a filtering mechanism might mainly apply in the centre of the gyre where mean currents are negligible and the turbulent field can freely develop, while strong mean currents 390 that generate and destroy eddies are likely to be the dominant factor in determining eddy polarity near the gyre's edges.

Up to 300 km off the shelf, high density in the eddy population is accompanied by intensities in the eddy field up to one order of magnitude higher than in the deep basin (Fig. 7c). This is visible all along the shelf break of the domain, from the Chukchi shelf break to the Canadian Arctic Archipelago shelf break. The most intense eddies are found at the mouth of the McKenzie River and at Pt. Barrow (not shown), being respectively fresher and saltier than their environment (Fig. 7e,f). Along 395 the Alaskan and Chukchi slopes, on both sides of Pt. Barrow, eddies with positive salinity anomalies are detected in the inner part of the slope, while eddies with negative anomalies are detected in the outer part. This pattern illustrates the penetration of the Pacific Waters from Pt. Barrow along the baroclinically unstable Alaskan coastal and Chukchi Slope currents (Corlett and Pickart, 2017; Spall et al., 2008) and supports observations of the penetration of eddies associated with a salty anomaly into the CB at Pt. Barrow (MacKinnon et al., 2021, in the submesoscale range). Additionally, eddies associated with a fresh anomaly 400 found along the outer part of the slope confirm the role of fresh water input from McKenzie River in generating instabilities that develop into eddies propagating downstream along the anticyclonic circulation (Kubryakov et al., 2021).

3.1.3 Within the pycnocline layer

Over the 26 years of simulation, there are about 30% more eddies detected in the pycnocline layer (\sim 85-250 m) than in the upper layer (9,000 on average in the pycnocline layer vs 6,000 eddies on average in the upper layer per year; Fig. 6a). Eddies 405 detected within the pycnocline layer are evenly distributed between cyclones and anticyclones (Fig. 6d) and are found to be smaller and weaker than in the upper layer on average (mean radius is decreased from 12.4 km to 11.6 km and intensity is decreased from $7.4 \cdot 10^{-6} \text{ s}^{-1}$ in the upper layer to $3.3 \cdot 10^{-6} \text{ s}^{-1}$, Fig. 6b,c). Although weaker, eddies in the pycnocline layer last about 6 days longer than in the upper layer, likely due to the absence of ice or air drag to dissipate eddies through friction (Fig. 6e). Despite this increased longevity, the mean distance travelled by eddies in the pycnocline layer is only increased by 410 about 1.1 km compared to the upper layer (Fig. 6f), presumably because of the weaker background mean flow advecting these eddies. Therefore, of the eddies generated over the slope, only the strongest and longest-lived eddies may be able to travel far




Figure 7. Eddy properties at 30 m (i.e. within the upper layer) over the 26 years of simulation. (a) Density of the eddy population (i.e. number of individual eddies detected per km^2) and associated properties: (b) averaged radius, (c) intensity, (d) polarity, and anomalies of (e) temperature and (f) salinity with respect to the environment of eddies. Temperature anomalies are calculated with respect to the local freezing temperature. Temperature and salinity anomalies are only accounted for when anomalies are significant (which represents about 15% of all eddies, see Sect. 2). All variables are extracted at nominal depth 30 m, and summed (panel a) or averaged (panels b-f) over the 26 years of simulation. Note that all fields show similar structures at all depths between 0 and 70 m except for the radii that are significantly larger within the ML (see Fig. 6c). Plain, dashed, dotted and loosely dotted black lines show respectively the 90%, 80%, 50% and 15% contours of the climatological September sea ice concentration. Gray lines show the 100, 500, 1000, 1500 m isobaths.

enough to reach the gyre and could thus participate in the transport of heat, salt and nutrients from the continental shelf to the deep basins (see Sect. 4).

In the pycnocline layer, eddy characteristics show a weaker seasonality compared to the upper layer. Quantitatively, the normalized amplitude of the seasonal cycle of the number of detected eddies diminishes with depth, from $SC_N = 2.3$ at 30 m to $SC_N = 0.5$ at 150 m (see Sect. 2 for a definition of SC_N). The other properties also show a decreased normalized amplitude of their seasonal cycle compared to the upper layer, by $\approx 60\%$ for the radius, 50% for the intensity and distance travelled, and by 40% for the lifetime (refer to Table 3 for detailed seasonal cycles). This damped seasonality is expected as the pycnocline shields eddies from dissipation by sea ice.

For most eddy characteristics (radius, intensity, duration, distance and polarity), the spatial distribution within the pycnocline layer is generally similar to that of the upper layer (compare Fig. 7 with Fig. 8) but persists throughout the year due to the absence of seasonal variability. The similarity in spatial distribution with the upper layer is expected as the anticyclonic circulation that dominates most of the region investigated extends down to the pycnocline associated to the wPW (see Fig. S5, see also Planat et al., 2025). However, the spatial distribution of the density in eddy population is notably different between the pycnocline and the upper layers along the southern edge of the BG (Fig. 8a). There, a strong reduction in the density of the eddy population is found compared to the shelf and deep basin (Fig. 8a). This reduced density compared to the upper layer occurs despite the eddies being relatively intense, long-lived and travelling relatively far along the anticyclonic flow (Fig. 8c,e,f). We suggest that the inner part of this local reduction in eddy generation is linked to a stabilizing effect of the continental slope. The growth of instabilities is known to be hampered over regions where the ratio of the continental slope to the isopycnal slopes is greater than 1 (Manucharyan and Isachsen, 2019), as is the case for the slope of the CB in the model (not shown, see also Regan et al., 2020). However, this reduction occurs up to 250 km away from the shelfbreak. There, we observed diminished background PV gradients (not shown, see Fig. 9 from Meneghelli et al., 2021) associated to diminished baroclinic instabilities, which offer an alternative explanation for the extended area with diminished eddy density.

3.1.4 Within the AW layer

Below the pycnocline layer and down to 1,200 m, within the AW layer, the total number of eddies over the 26 years of simulation decreases by 37% compared to the pycnocline layer (from on average 9000 per year in the pycnocline layer to 5,500 eddies per year in the AW layer; Fig. 6a). This decrease is reduced to -20% if one compares eddy density, as the area where eddies developed is reduced due to the bathymetry. Because the layer is located below the pycnocline, the seasonal variability in eddy properties is almost completely shut down (Fig. 6). In that layer, we find eddies similar in radius (≈ 11.8 km, Fig. 6b) but weaker in intensity than in the pycnocline layer ($1.5 \cdot 10^{-6} \text{ s}^{-1}$ compared to $3.3 \cdot 10^{-6} \text{ s}^{-1}$, Fig. 6c). The distance travelled by eddies decreases from 12.2 km to 7 km (Fig. 6f), and the polarity remains equally shared between cyclones and anticyclones (Fig. 6d). We note that the averaged lifetime of eddies is longer than in the pycnocline layer (14 days compared to 11 days) but remains small despite the few processes that could dissipate eddies at this depth. This relatively short lifetime may point to the fact that most of the eddies detected in this layer are not well-developed according to their turnaround time, and hence bound to disappear quickly. Of all the eddies detected in that layer, only 6% persist for longer than their turnaround time. These




Figure 8. Eddy properties at 150 m (i.e. within the pycnocline layer) over the 26 years of simulation. (a) Density of the eddy population (i.e. number of individual eddies detected per km^2) and associated properties: (b) averaged radius, (c) intensity, (d) polarity, (e) lifetime and (f) distance travelled. All variables are extracted at nominal depth 150 m, and summed (panel a) or averaged (panels b-f) over the 26 years of simulation. Note that all fields show similar structures at all depths between 85 and 225 m. Plain, dashed, dotted and loosely dotted black lines show respectively the 90%, 80%, 50% and 15% contours of the climatological September sea ice concentration. Gray lines show the 100, 500, 1000, 1500 m isobaths.

long-lasting eddies may live up to 150 days (99th percentile), which surpasses all the maximum durations detected in the other layer and matches estimates from observations of weeks to years (Hunkins, 1974; Timmermans et al., 2008). We refer the reader to Sect. 4 for a discussion on long-lasting eddies.

Within the AW layer, eddy properties show different patterns compared to the layers above. Significant differences are 450 expected given that the mean circulation of that layer departs strongly from that above (Fig. S5). In particular, eddies are predominantly generated over the continental slope along the path of the cyclonic boundary current carrying AW (Fig. 9a). A smaller density of eddies is generated in the rest of the domain, with some hot spots of high density in eddy population located close to the northern boundary of our domain. The latter correspond to short-lived eddies, and we discuss more extensively the 455 "turbulent soup" form by short-lived eddies in Sect. 4. Along the shelfbreak and boundary currents, the eddy intensity is larger by up to one order of magnitude compared to the rest of the domain (Fig. 9c). Throughout the rest of the domain, eddies are notably weaker. No clear spatial pattern in polarity arise at the scale of the basin, except along the shelf breaks of the Chukchi Sea and Canadian Archipelago, where anticyclones dominate in the inshore part of the current while cyclones dominate in the offshore part of the current (Fig. 9d). Off the western flank of Northwind Ridge are found the largest (up to 20 km), farthest-reaching (up to 40 km) and longest-lived (up to 60 days) eddies (Fig. 9b,e,f). The EKE is one order of magnitude larger in 460 that area than within the deep basin and displays hotspots in the form of large structures detaching from the cyclonic boundary current that hugs Northwind Ridge (not shown). In this region, large uncertainties exist among the literature on the exact path of the AW. The AW are thought to intermittently detach from the slope-intensified cyclonic boundary current, or alternatively flow along double boundary currents (McLaughlin et al., 2009; Li et al., 2020; Planat et al., 2025; Karcher et al., 2012; Lique et al., 2015). We suggest here some instabilities in the cyclonic boundary currents associated to the generation of large eddies.

465 3.2 Evolution of the population of eddies over 1995-2020

Over the last two decades, both the sea ice cover and the anticyclonic mean circulation of the Canadian Basin have drastically changed. Indeed, the sea ice extent has shrunk with a trend of -12.7% (Meier and Stroeve, 2022), while an acceleration of the anticyclonic circulation of the BG has been found to occur around 2007 in both observations (Giles, 2012; Regan et al., 2019) and models (Regan et al., 2020) associated with a decrease in the ice cover and increased Ekman pumping, with intensified 470 winds (Meneghelli et al., 2018). Both this decadal change in ice cover and this fast accelerating period of the BG are represented in our model (Fig. 10a) : the sea ice extent of the Canadian Basin decreases by -55% over the 26 years of simulation, while the mean gradients of SSH over the CB, which is a proxy for the intensity of the mean circulation at the surface, shows a




Figure 9. Eddy properties at 500 m (i.e. within the AW layer) over the 26 years of simulation. (a) Density of eddy population (i.e. number of individual eddies detected per km^2) and associated properties: (b) averaged radius, (c) intensity, (d) polarity, (e) duration and (f) distance travelled. All variables are extracted at nominal depth 500 m, and summed (panel a) or averaged (panels b-f) over the 26 years of simulation. Note that all fields show similar structures at all depths between 225 and 1200 m. Plain, dashed, dotted and loosely dotted black lines show respectively the 90%, 80%, 50% and 15% contours of the climatological September sea ice concentration. Gray lines show the 100, 500, 1000, 1500 m isobaths.

step increase of 0.15mm/km between 2006 and 2008, for an overall increase of +16% (% changes are computed using averages over the first and last 5 years of the simulation). We report in the paragraphs below how the population of eddies has changed
475 between 1995 and 2026 in light of the modifications of these two forcing fields.

Over 1995-2020, the number of eddies generated in the Canadian Basin increases in all layers, by +34% in the upper layer, +45% in the pycnocline layer and +40% in the AW layer (Fig. 10). Increases in eddy number are also found when looking regionally at the CB and the Alaskan shelf area at all depths; with the exception of the AW layer that shows an overall decrease
480 in the eddy number in the CB (-26%). However, a rebound in the eddy number is seen around 2017, thus suggesting a possible lagged increase in the eddy number in that layer. A key difference between the Alaskan Shelf and CB is that the latter is mostly energized above the mixed layer (+94% above 30 m vs -15% between 30 and 85 m) while the former presents an increase in the number of eddies that is roughly constant with depth throughout the upper layer. This difference may be explained by the greater energy input in the Alaskan area that becomes seasonally ice-free earlier than the interior of the basin, where little
485 additional energy linked to the sea ice decline can thus penetrate in the water column.

Along with the increase in the eddy population, eddies become bigger (+0.7 km), travel further (+2.2 km) and carry relatively warmer waters (+0.0027°C; Table. 2). These changes are in line with an increased stratification, which increases the Rossby radius. In the upper layer, eddies last longer (+0.6 days), most probably in relation to the reduced impact of sea ice. In both the AW and pycnocline layers, the eddy intensity is increased. This is presumably due to the fastened mean circulation, with
490 increased MKE and EKE in all layers (not shown). Polarity remains unchanged through the 3 decades. A detailed comparison of the histograms for each property between the first and last 5 years of the simulation is offered in supplementary Fig. S10.

An increase in the number of eddies is expected as sea ice shrinks, and is in line with the recent literature reporting on an enhancement of the energy over the Arctic (Regan et al., 2020; Li et al., 2024; Manucharyan et al., 2022; Armitage et al.,
495 2020). Yet, it remains unclear whether this increase is due only to the expansion of the open ocean and the MIZ, or also to increased levels of energy within the MIZ and pack ice linked to the sea ice becoming less concentrated, but also thinner and more mobile (Kinnard et al., 2011; Kwok, 2018; Rampal et al., 2009) and to changes in atmospheric forcing, in particular. If the former applies, then the density of the eddy population within the open ocean should remain constant. To investigate this question, we look at the density of the eddy population within three sea ice regions in the upper layer of our domain : the pack
500 ice (where the ice concentration is $\geq 80\%$), the open ocean (where the ice concentration is $\leq 15\%$), and the MIZ which lies in between. For a given year and region, the density is computed as the total number of individual eddies detected for that year over the mean sea ice area for that year in that region. Over the 26 years, we find an increase of +10% and +20% of the density in eddy population in the open ocean and MIZ respectively, and no change on average below the pack ice (Fig. S11). In the MIZ and open ocean, the increase in eddy generation is mainly a step increase in 2008 with reduced (shut down) interannual
505 variability in the MIZ (Open Ocean) in the following years. This enhancement in the density of the eddy population presumably results from the additional energy penetrating into the ocean in the recent state of the BG, in line with future projections of Li et al. (2024). In the open ocean, this accumulation of energy could be attributed to the acceleration of the BG from atmospheric

forcings (Giles, 2012), and in the MIZ to a combination of less compacted ice (Martin et al., 2016) or to the thinner ice cover (Muilwijk et al., 2024). A quantification of these different drivers is beyond the scope of this paper and left for future analysis.

510 In the pack ice, the constant density results from opposing changes with an increase above the ML of +10% and a decrease of similar amplitude below, with strong year to year variability (see e.g. 1999 and 2008) associated with low sea ice concentration years.

Overall, our results suggest that, in the upper layer, the number of eddies do not only increase because of an expansion of the open ocean area at the expense of sea ice, but also because of an energy surplus in the Canadian Basin in the MIZ and open 515 ocean, in line with conclusions from other modelling studies (e.g., Rieck et al., 2025b). The increase of energy below the pack ice, suggested in future projections of the Arctic (Rieck et al., 2025b), is only seen in the upper mixed layer in the current climate.

On interannual time scales, the CB, Alaskan shelf area and whole Canadian Basin show important variability in the number 520 of eddies detected in the upper layer. Significant correlations with the ice cover, either with the September sea ice extent, or with the yearly cumulated area with ice less concentrated than 80%, are only visible at the surface (down to ≈ 20 m, that is approximately down to the basis of the mixed layer), at the exception of the Alaskan area where they are found significant deeper (not shown).

In both the upper and pycnocline layers, the number of eddies starts rising around 2000 before culminating in 2008 (+54% 525 between 2006 and 2008 in the pycnocline layer, Fig. 10b,c). The increase in the number of eddies that peaks in 2008 in our model bears similarity to the EKE increase that was reported by Regan et al. (2020) to occur over one year following the gyre acceleration in 2007 and low sea ice record of that year. The authors suggest that beyond 2007-2008, the BG is able to expand spatially over NorthWind Ridge and thus the need for eddies to release the accumulated potential energy is reduced. The transient increase in the number of eddies reported in our analysis thus tends to confirm this hypothesis, with a slightly 530 longer equilibration time. In the pycnocline layer of the CB, the increase in the eddy number in 2008 persists for an additional couple of years in contrast to the Alaskan shelf area and basin average where the increase remains punctual. Further down, the AW layer also displays an increase, yet weaker, in the number of eddies in the CB (Fig. 10d). This increase at depth suggests a direct top-down coupling between the BG and the AW layers, despite the insulation of the pycnocline, as documented by Lique and Johnson (2015) and Lique et al. (2015) for the mean circulation of the AW layer. The processes sustaining the increase 535 in the number of eddies within both the pycnocline and AW layer for the 5 years following the increase of the BG remain to be explored. Additionally, we note that the number of eddies generally keeps increasing after 2016, while the gyre intensity declines. Our simulation lacks a few more years to interpret changes.

Within the AW layer, little interannual variability in the number of eddies is found in contrast to the layers above (Fig. 10d). However, when looking at the long-lasting eddies (6% of the total population), we do see important year-to-year variability (see 540 Fig. S12). These more persistent features show a large relative increase (+300%) starting around 2011, which is mostly due to an increase in the number of eddies located east of Northwind Ridge. These structures, which are large (30-50 km), develop in particular in the late years of the simulation (2012-2020) when the cyclonic boundary current on the Northwest flank of

	R [km]	D [km]	dt [days]	$ \Omega [10^{-7} \text{ s}^{-1}]$	ΔS	ΔT	$r_{C/T} [\%]$
Upper layer (0 - 85 m)	1 ; 8%	2.4 ; 21%	0.6 ; 10%	-2.1 ; -3%	-0.008	0.004	-0.7
Pycnocline layer (85 - 225 m)	0.5 ; 4%	2.0 ; 18%	-1.3 ; -11%	7.9 ; 26%	0.003 ;	0.004 ;	-0.06
AW layer (225 - 1200 m)	0.4 ; 4%	2.0 ; 30%	-1.4 ; -9%	7.4 ; 57%	-2.5e-5	0.0009	3
All (0-1200 m)	0.7 ; 6%	2.2 ; 22 %	-0.5 ; -5%	3 ; 7 %	0.0003	0.0027	0.8

Table 2. Change in mean eddy properties for each layer when comparing the last 5 years of the simulation and first 5 years. Red indicate an increase, blue a decrease. Also reported are relative increases in % for the mean radius, distance, duration and intensity.

Northwind Ridge reverses to an anticyclonic flow with increased horizontal shear (not shown). What drives these changes in the AW mean pathways, and how the latter influence the generation of large eddies is beyond the scope of this paper and is left for future analysis.

4 Discussion and conclusion

In this study, we apply an eddy detection and tracking algorithm to the output of a high-resolution regional model of the Arctic, in order to document the characteristics of mesoscale eddies in the Canadian Basin and examine the evolution in the number and characteristics of eddies over 1995-2020. Over that period, we report an average of $\approx 6,000$ eddies generated per year in the surface layer, $\approx 9,000$ eddies in the pycnocline layer, and $\approx 5,500$ in the AW layer of the Canadian Basin. Most of these eddies are found to be the size of the Rossby radius of deformation (mean eddy radius of about 12.1 km), stationary (distance travelled of about 11.1 km) and short lasting (lifetime of about 10 days; Fig. 4). The distribution between cyclones and anticyclones is about equal in the investigated domain. In addition, the majority of eddies do not have a temperature nor salinity anomaly relative to their environment, although some significant anomalies are visible along the shelf in the surface layer (Fig. 7e,f). All the documented properties (radius, polarity, intensity, lifetime, distance travelled and temperature and salinity anomaly) vary significantly across space and time (see Table 3), pointing to the role of the environment, such as the stratification, sea ice or main currents, in setting different processes important for the generation and dissipation of eddies in the Canadian basin.

Our analysis highlights three layers that arise from the stratification of the Canadian Basin and show consistent characteristics for eddies across the vertical. It is important to note that the definition of the three layers relies on the statistical properties of the whole eddy field over 26 years and hence does not account for temporal variability of the mean circulation, the location of the gyre, and the mean isopycnal depths and slopes. Thus, the depths used as delimiters of the three layers do not necessarily correspond to the actual, instantaneous pycnoclines that they are assumed to represent at all times. This is particularly true where the isopycnal surfaces are strongly tilted or even outcrop over the slope and the shelf break. There, fixed depth layers ensure that there is no averaging done between regions in contact with sea ice and those away from the sea ice influence that would otherwise complicate the interpretation of the results. The results presented in this study, and in particular the spatial structure of the eddy properties and their temporal variability, remain consistent when slightly varying the layers' upper and




Figure 10. Time series of (a) the September sea ice extent over the Canadian Basin (red, total area of the domain is $2.7 \cdot 10^{12} \text{ km}^2$) and norm of the gradient of SSH calculated at every location of the domain and averaged over the CB. (b), (c) and (d) show the layered-averaged number of eddies relative to the 26-year average for (b) the upper layer, (c) the pycnocline layer, and (d) the AW layer, further divided into the whole Canadian Basin (plain black line), the Alaskan shelf area (dashed black line) and the Canada Basin (dashed dotted line). Reporting numbers of eddies relative to the 1995-2020 average for each depth level permits a comparison of the temporal evolution of depth levels with different absolute numbers of eddies.

lower boundaries indicating the robustness of the key features reported for the three layers especially in the centre of the BG where the isopycnals are relatively flat.

570 In the upper layer (top 85 m), which lies on average above the pycnocline, eddy properties display a significant seasonal cycle generally in phase with that of sea ice. Eddy population is the densest when sea ice concentration decreases below 80%, a threshold in line with the results of Manucharyan and Thompson (2022), and over the continental slope. At about \sim 70 m, the pycnocline insulates the eddy field from dissipation by sea ice. Therefore, weaker seasonality is detected in the eddy characteristics below this depth. Between 85 m and 225 m depth, a reduced density in the eddy population is found within a 250 km wide area along the Alaskan shelf break that is attributed to the smaller gradients of background potential vorticity. Deeper 575 down, the AW layer shows a muted seasonal cycle in eddy properties and little to no similarities to the layers above, due to the efficient insulation of this region by the pycnocline. In particular, while the upper and pycnocline layers of the Canadian Basin show anticyclones forming preferably at the centre of the Beaufort Gyre, in the AW layer, a symmetry is found along the slope, with anticyclones forming in-shore and cyclones off-shore.

580 The 1995-2020 period is marked by an overall rise in eddy density at all depths (+35 – 45%), in line with predictions of an increasingly energetic Arctic Ocean (Rieck et al., 2025b; Li et al., 2024) with reduced ice cover (Meredith et al., 2001; Meier and Stroeve, 2022). A smaller increase is visible in the upper open ocean and upper MIZ (+10%), and limited to the mixed layer of the pack ice area. We argue that the higher density in the eddy population is the result of an increasing penetration of energy in the upper layers of the basin where the ice concentration is small enough. Therefore, in the MIZ and open ocean, 585 these results confirm findings from Rieck et al. (2025b) of an increased energy input associated to more energy conversion toward EKE, and from Li et al. (2024) who associate this increased EKE to higher baroclinic instabilities, despite increased eddy killing with more mobile ice, in simulations of the future Arctic. Large interannual variability in the number of eddies is visible in the upper and pycnocline layers, with in particular a peak in the eddy population between 2007-2009, when the BG is known to have accelerated. Post 2010, the eddy population is similar in the upper and pycnocline layer to that prior to 2007, 590 which tend to confirm Regan et al. (2020) hypothesis that the gyre is able to expand above Northwind Ridge and thus diminish the baroclinic instabilities. We however report longer equilibration time for the number of eddies, especially in the CB. The upper layer additionally shows the imprint of the ice cover on yearly time scale with significant correlations between the ice area and the number of eddies along the Alaskan shelf break and at the very surface of the CB. Overall, the interannual and decadal variability of the pycnocline and surface layers result from both local changes in sea ice dissipation and large scale 595 changes in energy input.

One of the striking and most intriguing characteristic of the eddy field reported so far in the literature is the eddy polarity, ranging from $r_{C/A} = 5\%$ to $r_{C/A} = 70\%$ (see Sect. 1 Cassianides et al., 2023; Kozlov et al., 2019). At the scale of the Canadian Basin, as many anticyclones as cyclones are found in the upper and pycnocline layers of our model. This partition is maintained 600 for small eddies (< 15 km), or long-lived eddies (not shown). Yet, the polarity of eddies shows a marked spatial pattern over the Canadian Basin with a larger proportion of anticyclones than cyclones ($\approx 70\%$ anticyclones) in the centre of the BG (see Figures 7d, and 8d). This predominance of anticyclones aligns with that estimated using dataset based on ITPs (Cassianides

	Nmb. of eddies	R [km]		dt [days]		D [km]		$r_{C/T}$ [%]		$ \Omega $ [s $^{-1}$]		ΔT [°C]	ΔS		
Upper layer (0 - 85 m)	6000	2.4	12.4	0.1	5.4	0.6	11.3	0.7	48.1	0.09	7.410^{-6}	0.4	0.08	1.7	-0.01
Pycnocline layer (85 - 225 m)	9000	0.6	11.6	0.04	10.8	0.1	12.2	0.3	49.5	0.05	3.310^{-6}	0.3	0.01	5.1	0.008
AW layer (225 - 1200 m)	5500	0.5	11.8	0.02	14.2	0.3	7.1	0.1	50.8	0.06	1.510^{-6}	0.2	0.003	1.1	-0.001
Total	6000	2.6	12.1	0.06	9.9	0.2	11.1	0.4	49	0.05	4.610^{-6}	0.04	0.04	1.6	-0.001

Table 3. Mean (left) and seasonal cycle amplitudes (right) of the eddy properties reported along this manuscript for each layer and for the Canadian Basin as a whole.

et al., 2023; Zhao et al., 2014), though observations suggest a much higher proportion of anticyclones in that region (95% versus 65% in our model when applying our eddy detection following the temporal and spatial sampling of the ITPs, see Fig. 605 S13). Although the comparison between the ITP dataset and our model is somewhat limited by the fact that the observed eddy field may be dominated by sub-mesoscale features, which our model does not resolve, our results suggest that part of the anticyclonic dominance documented by the ITP dataset is simply linked to the ITP sampling location, as was already suggested by Beech et al. (2025). Yet, a proper and detailed comparison with observations would require using a model at higher-resolution given that most features identified with moorings, ITPs or satellite fall between the meso- and the submeso-610 scales. Such a comparison would also benefit from an adequate model subsampling, through an Observing System Simulation Experiments, to take into account the observations sampling biases.

Our algorithm identifies as "eddy" a broad range of features, from the ephemeral ones that last a couple of days to the more persistent ones that are likely more coherent. When separating between short and long-lasting eddies based on their duration being respectively shorter or longer than their turnaround time, we find that the bulk of the eddy dataset consists of short-lasting 615 eddies which we refer to as a "turbulent soup". Within this ephemeral eddy population, it is likely that some short features are artefact of the tracking algorithm, that may lose track of the weakest eddies, or of eddies splitting/merging. The OW detection algorithm is known to be biased towards weak eddies. Yet, most of the features are likely actual, short-lasting eddies that are evanescent by nature. Within the upper layer, the very short lifetime of these eddies could be attributed to the presence of sea ice (spindown time scale of eddies due to ice friction is estimated around 4 days Meneghelli et al., 2021). At depth, weak 620 vortices have been suggested to form from the stirring of interior PV gradients (Manucharyan and Stewart, 2022). Alternatively, weak eddies may arise from the relatively low EKE in our model due to its resolution (Sect. 2). Overall, these relatively weak and short-lasting eddies that form a turbulent soup are not captured by observational dataset, which may explain some of the important differences found between our census and the observation-based literature. Nonetheless, these eddies may play an 625 important role in the transfer of energy, and we leave for future analysis to quantify their integrated role in the penetration of heat, salt and nutrients into the deep basin.

Long-lasting eddies may resemble more the eddy population captured by observations. We find that these eddies represent 15% of the population in the upper layer, 10% in the pycnocline layer and 6% in the AW layer (see also the mean statistical properties of long-lasting eddies in supplementary Fig. S14). A fraction of these eddies display temperature anomalies, in particular along the shelf break, while little salinity anomaly is observed within this eddy subset. These anomalies are mostly

630 positive at the surface where eddies are formed either within the mixed layer or the warm Near Surface Temperature Maximum; and in the upper part of the pycnocline layer, where they form within the warm summer Pacific Waters (not shown). In contrast, these anomalies are mostly negative in the lower part of the pycnocline layer where they form in the cold winter Pacific Waters (not shown). Note that because these long-lasting eddies are mostly generated in regions of high density in eddy population, such as Pt. Barrow, it is difficult to quantify the associated anomaly in temperature and salinity. Still, a visual inspection seems
635 to support the presence of strong anomalies at Pt. Barrow (see Fig. 11). Because these long-lasting eddies travel for a few tens of kilometres, we hypothesize that the most coherent eddies actively play a role in transporting heat, in line with previous observations of warm eddies directly penetrating into the CB from Point Barrow (MacKinnon et al., 2021). One example of such an eddy is given on Fig. 11a, with an anticyclone carrying warm water off-shore when leaving Pt Barrow. This eddy is seen to subduct at depth from mi-September, with a colder and fresher layer developing above (Fig. 11b). However, a robust
640 quantification of the heat transport associated with these eddies is not trivial as it requires computing the temperature anomaly which is highly dependent upon the definition of the environment. This analysis is thus left for future work.

To conclude, we present a first characterization of the spatio-temporal properties of mesoscale eddies in the Canadian Basin. By doing so, we reveal strong differences in eddy properties across space and time, as well as important variability in the
645 number of eddies generated over 1995-2020 in relation to the loss of sea ice and acceleration of the BG. Our eddy census can thus provide a benchmark against which censuses from other models could be compared, and could form a starting point to explore questions that remain on the BG dynamical and thermodynamical equilibrium, as well as on the transport and mixing of nutrients, salt, or other tracers.

Code and data availability. The *eddytools* python package used to perform the eddy detection and tracking along with its documentation is available at <https://github.com/jk-rieck/eddytools@N-tracking-properties>. The scripts used in this study to detect, track and analyse the mesoscale eddies are available at https://github.com/noemieplanat/Eddies_CB/releases/tag/submission. The documentation of the CREG12 experiment can be found in Talandier and Lique (2024). The detection and tracking for 30, 69, 147 and 508 m depth are available here 10.5281/zenodo.17713586. Fresh water content estimates were obtained from <https://www2.whoi.edu/site/beaufortgyre/data/freshwater-content-gridded-data/>, accessed on October 1st, 2025. The World Ocean Atlas (WOA) 2023 climatology was downloaded from
655 <https://www.ncei.noaa.gov/access/world-ocean-atlas-2023/>, accessed in January 2025 (Locarnini et al., 2024; Reagan et al., 2024). Arctic dynamic topography/geostrophic currents data were provided by the Centre for Polar Observation and Modelling, University College London (Armitage et al., 2016, 2017). Sea ice concentration were obtained from the National Snow and Ice Data Center (NSIDC, Di-Girolamo et al., 2022). PIOMAS Reanalysis (Zhang and Rothrock, 2003) was downloaded from <https://psc.apl.uw.edu/research/projects/arctic-sea-ice-volume-anomaly/>

660 *Author contributions.* NP designed and conducted the study with input from COD, CL and LBT. CT and CL ran CREG12 model. JKR built the eddytools python package and NP applied it on the output of CREG12 with appropriate modifications. NP processed and analyzed the




Figure 11. (a) Conservative Temperature at 30 m in September 1996 overlaid with contours corresponding to all long-lasting eddies detected (not necessarily for the first time) on September 16th 1996. Colors of contours indicate the temperature anomaly of eddies, and thin black lines their trajectory. Eddies are coloured (filled) by their mean temperature. (b) Conservative temperature along the eddy track identified on (a) with a thicker black line. At a given depth, intense color shading indicates that the algorithm detects an eddy at that depth, location and time while pale color shading indicates that no eddy is detected. Thick black lines show isopycnals along the eddy track.

eddy database. NP, JKR, COD, CL and LBT contributed to the interpretation. NP wrote the manuscript, with contribution from COD, CL, LBT, JKR and CT for editing.

Competing interests. The authors declare no conflict of interest.

665 *Acknowledgements.* NP was supported by the Fonds de recherche du Québec - Nature et Technologie (FRQNT) through a Doctoral Training Scholarship, a Natural Sciences and Engineering Research Council of Canada (NSERC) Accelerator Supplements awarded to COD (grant no. RGPAS/2018-522502), and a NSERC Discovery Grant awarded to BT (grant no. RGPIN/2018-04838). NP also received a scholarship from ISblue (Interdisciplinary graduate school for the blue planet - ANR-17-EURE-0015) and financial support from Québec-Océan for this work. COD and JKR acknowledge funding from NSERC through the Accelerator Supplements (grant no. RGPAS/2018-522502) and a
670 Canada Research Chair (grant no. 252794), both awarded to COD. NP, COD, CL, JKR and LBT acknowledge the financial support from the Fonds de recherche du Québec – Nature et technologies (FRQNT) and the French Ministry of Europe and Foreign Affairs through the Samuel-de-Champlain grant (<https://doi.org/10.69777/329860>). CL and CT were supported by funding from the CLIMArcTIC project funded by the “PPR Océan et Climat—France 2030” (contract ANR-22-POCE-0005). The pan-Arctic simulations were performed using HPC resources from the French GENCI-CINES center (Grant 2023-A0130107420). The authors also acknowledge the technical and scientific contributions
675 of Benjamin Valette and Sacha Coez to the preparatory phase of the study.

References

Antonov, J. I., Seidov, D., Boyer, T. P., Locarnini, R. A., Mishonov, A. V., Garcia, H. E., Baranova, O. K., Zweng, M. M., and Johnson, D. R.: World Ocean Atlas 2009, Volume 2: Salinity., Tech. rep., U.S. Government Printing Office, Washington, D.C., 2010.

Armitage, T. W. K., Bacon, S., Ridout, A. L., Thomas, S. F., Aksenov, Y., and Wingham, D. J.: Arctic sea surface height variability and
680 change from satellite radar altimetry and GRACE, 2003–2014, *Journal of Geophysical Research*, [Dataset]http://www.cpom.ucl.ac.uk/dynamic_topography/, 2016.

Armitage, T. W. K., Bacon, S., Ridout, A. L., Petty, A. A., Wolbach, S., and Tsamados, M.: Arctic Ocean surface geostrophic circulation
2003–2014, *The Cryosphere*, 2017.

Armitage, T. W. K., Manucharyan, G. E., Petty, A. A., Kwok, R., and Thompson, A. F.: Enhanced eddy activity in the Beaufort Gyre in
685 response to sea ice loss, *Nature Communications*, 11, 761, <https://doi.org/10.1038/s41467-020-14449-z>, 2020.

Barnier, B., Blaker, A., Biastoch, A., Boening, C., Coward, A. C., Deshayes, J., Duche, A., Hirschi, J., Sommer, J. L., Madec, G., Maze,
G., Molines, J. M., New, A., Penduff, T., Scheinert, M., Talandier, C., and Treguier, A. M.: DRAKKAR:developing high resolution ocean
components for European Earth system models, 2014.

Barton, B. I., Lique, C., Lenn, Y., and Talandier, C.: An Ice-Ocean Model Study of the Mid-2000s Regime Change in the Barents Sea, *Journal
690 of Geophysical Research: Oceans*, 127, e2021JC018280, <https://doi.org/10.1029/2021JC018280>, 2022.

Batrak, Y. and Müller, M.: On the warm bias in atmospheric reanalyses induced by the missing snow over Arctic sea-ice, *Nature Communications*, 10, 4170, <https://doi.org/10.1038/s41467-019-11975-3>, 2019.

Beech, N., Rackow, T., Semmler, T., and Jung, T.: High-latitude Southern Ocean eddy activity projected to evolve with anthropogenic climate
695 change, *Communications Earth & Environment*, 6, <https://doi.org/10.1038/s43247-025-02221-4>, publisher: Springer Science and Business Media LLC, 2025.

Carpenter, J. R. and Timmermans, M.: Deep mesoscale eddies in the Canada Basin, Arctic Ocean, *Geophysical Research Letters*, 39,
2012GL053025, <https://doi.org/10.1029/2012GL053025>, 2012.

Cassianides, A., Lique, C., Tréguier, A., Meneghelli, G., and De Marez, C.: Observed Spatio-Temporal Variability of the Eddy-Sea Ice
700 Interactions in the Arctic Basin, *Journal of Geophysical Research: Oceans*, 128, e2022JC019469, <https://doi.org/10.1029/2022JC019469>, 2023.

Cavalieri, D. J., Gloersen, P., and Campbell, W. J.: Determination of sea ice parameters with the NIMBUS 7 SMMR, *Journal of Geophysical
Research: Atmospheres*, 89, 5355–5369, <https://doi.org/10.1029/JD089iD04p05355>, 1984.

Chelton, D. B., Schlax, M. G., and Samelson, R. M.: Global observations of nonlinear mesoscale eddies, *Progress in Oceanography*, 91,
167–216, <https://doi.org/10.1016/j.pocean.2011.01.002>, 2011.

705 Comiso, J. C.: Characteristics of Arctic winter sea ice from satellite multispectral microwave observations, *Journal of Geophysical Research*,
91, 975, <https://doi.org/10.1029/JC091iC01p00975>, 1986.

Corlett, W. B. and Pickart, R. S.: The Chukchi slope current, *Progress in Oceanography*, 153, 50–65,
<https://doi.org/10.1016/j.pocean.2017.04.005>, 2017.

D'Asaro, E. A.: Generation of submesoscale vortices: A new mechanism, *Journal of Geophysical Research: Oceans*, 93, 6685–6693,
710 <https://doi.org/10.1029/JC093iC06p06685>, 1988.

De Lavergne, C., Madec, G., Capet, X., Maze, G., and Roquet, F.: Getting to the bottom of the ocean, *Nature Geoscience*, 9, 857–858,
<https://doi.org/10.1038/ngeo2850>, 2016.

DiGirolamo, N., Parkinson, C., Cavalieri, D., Gloersen, P., and Zwally, H.: Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data, <https://doi.org/10.5067/MPYG15WAA4WX>, 2022.

715 Dupont, F., Higginson, S., Bourdallé-Badie, R., Lu, Y., Roy, F., Smith, G. C., Lemieux, J.-F., Garric, G., and Davidson, F.: A high-resolution ocean and sea-ice modelling system for the Arctic and North Atlantic oceans, *Geoscientific Model Development*, 8, 1577–1594, <https://doi.org/10.5194/gmd-8-1577-2015>, 2015.

Giles, K. A.: Western Arctic Ocean freshwater storage increased by wind-driven spin-up of the Beaufort Gyre, *NATURE GEOSCIENCE*, 5, 2012.

720 Giulivi, C. F. and Gordon, A. L.: Isopycnal displacements within the Cape Basin thermocline as revealed by the Hydrographic Data Archive, *Deep Sea Research Part I: Oceanographic Research Papers*, 53, 1285–1300, <https://doi.org/10.1016/j.dsr.2006.05.011>, 2006.

Gupta, M., Gürcan, E., and Thompson, A. F.: Eddy-Induced Dispersion of Sea Ice Floes at the Marginal Ice Zone, *Geophysical Research Letters*, 51, e2023GL105 656, <https://doi.org/10.1029/2023GL105656>, 2024.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, 725 A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

730 Hu, X., Myers, P. G., and Lu, Y.: Pacific Water Pathway in the Arctic Ocean and Beaufort Gyre in Two Simulations With Different Horizontal Resolutions, *Journal of Geophysical Research*, 2019.

Hunkins, K. L.: Subsurface eddies in the Arctic Ocean*, Tech. rep., Pergamon Press, volume: 21, 1974.

Karcher, M., Smith, J. N., Kauker, F., Gerdes, R., and Smethie, W. M.: Recent changes in Arctic Ocean circulation revealed by iodine-129 735 observations and modeling, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2011JC007513>, publisher: Blackwell Publishing Ltd, 2012.

Kinnard, C., Zdanowicz, C. M., Fisher, D. A., Isaksson, E., de Vernal, A., and Thompson, L. G.: Reconstructed changes in Arctic sea ice over the past 1,450 years, *Nature*, 479, 509–512, <https://doi.org/10.1038/nature10581>, publisher: Nature Publishing Group, 2011.

Kozlov, I. E., Artamonova, A. V., Manucharyan, G. E., and Kubryakov, A. A.: Eddies in the Western Arctic Ocean From Space- 740 borne SAR Observations Over Open Ocean and Marginal Ice Zones, *Journal of Geophysical Research: Oceans*, 124, 6601–6616, <https://doi.org/10.1029/2019JC015113>, 2019.

Kubryakov, A. A., Kozlov, I. E., and Manucharyan, G. E.: Large Mesoscale Eddies in the Western Arctic Ocean From Satellite Altimetry Measurements, *Journal of Geophysical Research: Oceans*, 126, e2020JC016 670, <https://doi.org/10.1029/2020JC016670>, 2021.

Kwok, R.: Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958–2018), *Environmental Research Letters*, 13, 105 005, <https://doi.org/10.1088/1748-9326/aae3ec>, 2018.

745 Lellouche, J.-M., Greiner, E., Le Galloudec, O., Garric, G., Regnier, C., Drevillon, M., Benkiran, M., Testut, C.-E., Bourdalle-Badie, R., Gasparin, F., Hernandez, O., Levier, B., Drillet, Y., Remy, E., and Le Traon, P.-Y.: Recent updates to the Copernicus Marine Service global ocean monitoring and forecasting real-time 112° high-resolution system, *Ocean Science*, 14, 1093–1126, <https://doi.org/10.5194/os-14-1093-2018>, 2018.

Levitus, S., Locarnini, R. A., Boyer, T. P., Mishonov, A. V., Antonov, J. I., Garcia, H. E., Baranova, O. K., Zweng, M. M., Johnson, D. R., 750 and Seidov, D.: World ocean atlas 2009, Volume 1: Temperature, Tech. rep., U.S. Government Printing Office, Washington, D.C, 2010.

Li, J., Pickart, R. S., Lin, P., Bahr, F., Arrigo, K. R., Juraneck, L., and Yang, X.: The Atlantic Water Boundary Current in the Chukchi Borderland and Southern Canada Basin, *Journal of Geophysical Research: Oceans*, 125, e2020JC016197, <https://doi.org/10.1029/2020JC016197>, 2020.

Li, X., Wang, Q., Danilov, S., Koldunov, N., Liu, C., Müller, V., Sidorenko, D., and Jung, T.: Eddy activity in the Arctic Ocean projected to surge in a warming world, *Nature Climate Change*, 14, 156–162, <https://doi.org/10.1038/s41558-023-01908-w>, publisher: Nature Publishing Group, 2024.

755 Lique, C. and Johnson, H. L.: Is there any imprint of the wind variability on the Atlantic Water circulation within the Arctic Basin?, *Geophysical Research Letters*, 2015.

760 Lique, C., Johnson, H. L., and Davis, P. E. D.: On the Interplay between the Circulation in the Surface and the Intermediate Layers of the Arctic Ocean, *JOURNAL OF PHYSICAL OCEANOGRAPHY*, 45, 2015.

Liu, C., Wang, Q., Danilov, S., Koldunov, N., Müller, V., Li, X., Sidorenko, D., and Zhang, S.: Spatial Scales of Kinetic Energy in the Arctic Ocean, *Journal of Geophysical Research: Oceans*, 129, e2023JC020013, <https://doi.org/10.1029/2023JC020013>, 2024.

765 Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Reagan, J. R., Boyer, T. P., Seidov, D., Wang, Z., Garcia, H. E., Bouchard, C., Cross, S. L., Paver, C. R., and Dukhovskoy, D.: World Ocean Atlas 2023, Volume 1: Temperature, <https://doi.org/10.25923/54BH-1613>, publisher: NOAA National Centers for Environmental Information, 2024.

770 MacKinnon, J. A., Simmons, H. L., Hargrove, J., Thomson, J., Peacock, T., Alford, M. H., Barton, B. I., Boury, S., Brenner, S. D., Couto, N., Danielson, S. L., Fine, E. C., Graber, H. C., Guthrie, J., Hopkins, J. E., Jayne, S. R., Jeon, C., Klenz, T., Lee, C. M., Lenn, Y.-D., Lucas, A. J., Lund, B., Mahaffey, C., Norman, L., Rainville, L., Smith, M. M., Thomas, L. N., Torres-Valdés, S., and Wood, K. R.: A warm jet in a cold ocean, *Nature Communications*, 12, 2418, <https://doi.org/10.1038/s41467-021-22505-5>, 2021.

775 Madec, G., Bell, M., Blaker, A., Bricaud, C., Bruciaferri, D., Castrillo, M., Calvert, D., Jérôme Chanut, Clementi, E., Coward, A., Epicoco, I., Éthé, C., Ganderton, J., Harle, J., Hutchinson, K., Iovino, D., Lea, D., Lovato, T., Martin, M., Martin, N., Mele, F., Martins, D., Masson, S., Mathiot, P., Mocavero, S., Müller, S., Nurser, A. G., Paronuzzi, S., Peltier, M., Person, R., Rousset, C., Rynders, S., Samson, G., Téchené, S., Vancoppenolle, M., and Wilson, C.: NEMO Ocean Engine Reference Manual, <https://doi.org/10.5281/ZENODO.1464816>, publisher: Zenodo Version Number: v4.2.1, 2023.

780 Manley, T. O. and Hunkins, K.: Mesoscale eddies of the Arctic Ocean, *Journal of Geophysical Research: Oceans*, 90, 4911–4930, <https://doi.org/10.1029/JC090iC03p04911>, 1985.

Manucharyan, G. E. and Isachsen, P. E.: Critical Role of Continental Slopes in Halocline and Eddy Dynamics of the Ekman-Driven Beaufort Gyre, *Journal of Geophysical Research: Oceans*, 124, 2679–2696, <https://doi.org/10.1029/2018JC014624>, publisher: Blackwell Publishing Ltd, 2019.

785 Manucharyan, G. E. and Spall, M. A.: Wind-driven freshwater buildup and release in the Beaufort Gyre constrained by mesoscale eddies, *Geophysical Research Letters*, 43, 273–282, <https://doi.org/10.1002/2015GL065957>, 2016.

Manucharyan, G. E. and Stewart, A. L.: Stirring of interior potential vorticity gradients as a formation mechanism for large subsurface-intensified eddies in the Beaufort Gyre, *Journal of Physical Oceanography*, <https://doi.org/10.1175/jpo-d-21-0040.1>, publisher: American Meteorological Society, 2022.

790 Manucharyan, G. E. and Thompson, A. F.: Heavy footprints of upper-ocean eddies on weakened Arctic sea ice in marginal ice zones, 2022.

Manucharyan, G. E. and Timmermans, M.-L.: Generation and Separation of Mesoscale Eddies from Surface Ocean Fronts, *Journal of Physical Oceanography*, 43, 2545–2562, <https://doi.org/10.1175/JPO-D-13-094.1>, 2013.

Manucharyan, G. E., Spall, M. A., and Thompson, A. F.: A Theory of the Wind-Driven Beaufort Gyre Variability, *Journal of Physical Oceanography*, 46, 3263–3278, <https://doi.org/10.1175/JPO-D-16-0091.1>, 2016.

790 Manucharyan, G. E., Lopez-Acosta, R., and Wilhelmus, M. M.: Spinning ice floes reveal intensification of mesoscale eddies in the western Arctic Ocean, *Scientific Reports*, 12, 7070, <https://doi.org/10.1038/s41598-022-10712-z>, 2022.

Martin, T., Tsamados, M., Schroeder, D., and Feltham, D. L.: The impact of variable sea ice roughness on changes in Arctic Ocean surface stress: A model study, *Journal of Geophysical Research: Oceans*, 121, 1931–1952, <https://doi.org/10.1002/2015JC011186>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015JC011186>, 2016.

795 Martínez-Moreno, J., Lique, C., and Talandier, C.: Sea Ice Heterogeneity as a Result of Ocean Eddy Activity During the Ice Growth Season, *Geophysical Research Letters*, 52, e2024GL113645, <https://doi.org/10.1029/2024GL113645>, 2025.

McLaughlin, F. A., Carmack, E. C., Williams, W. J., Zimmermann, S., Shimada, K., and Itoh, M.: Joint effects of boundary currents and thermohaline intrusions on the warming of Atlantic water in the Canada Basin, 1993–2007, *Journal of Geophysical Research: Oceans*, 114, 2008JC005001, <https://doi.org/10.1029/2008JC005001>, 2009.

800 Meier, W. and Stroeve, J.: An Updated Assessment of the Changing Arctic Sea Ice Cover, *Oceanography*, <https://doi.org/10.5670/oceanog.2022.114>, 2022.

Meneghelli, G., Marshall, J., Timmermans, M.-L., and Scott, J.: Observations of Seasonal Upwelling and Downwelling in the Beaufort Sea Mediated by Sea Ice, *JOURNAL OF PHYSICAL OCEANOGRAPHY*, 48, 2018.

Meneghelli, G., Doddridge, E., Marshall, J., Scott, J., and Campin, J.-M.: Exploring the Role of the “Ice–Ocean Governor” and Mesoscale 805 Eddies in the Equilibration of the Beaufort Gyre: Lessons from Observations, *JOURNAL OF PHYSICAL OCEANOGRAPHY*, 50, 2020.

Meneghelli, G., Marshall, J., Lique, C., Isachsen, L. E., Doddridge, E., Campin, J.-M., Regan, H., and Talandier, C.: Genesis and Decay of Mesoscale Baroclinic Eddies in the Seasonally Ice-Covered Interior Arctic Ocean, *JOURNAL OF PHYSICAL OCEANOGRAPHY*, 51, 2021.

Meredith, M., Heywood, K., Dennis, P., Goldson, L., White, R., Fahrbach, E., Schauer, U., and Østerhus, S.: Freshwater fluxes through the 810 Western Fram Strait, *Geophysical Research Letters*, 28, 1615–1618, <https://doi.org/10.1029/2000GL011992>, 2001.

Muilwijk, M., Hattermann, T., Martin, T., and Granskog, M. A.: Future sea ice weakening amplifies wind-driven trends in surface stress and Arctic Ocean spin-up, *Nature Communications*, 15, 6889, <https://doi.org/10.1038/s41467-024-50874-0>, 2024.

Nencioli, F., Dong, C., Dickey, T., Washburn, L., and McWilliams, J. C.: A Vector Geometry-Based Eddy Detection Algorithm and Its 815 Application to a High-Resolution Numerical Model Product and High-Frequency Radar Surface Velocities in the Southern California Bight, *Journal of Atmospheric and Oceanic Technology*, 27, 564–579, <https://doi.org/10.1175/2009JTECHO725.1>, 2010.

Nurser, A. J. G. and Bacon, S.: Arctic Ocean Rossby radius Eddy length scales and the Rossby radius in the Arctic Ocean Arctic Ocean Rossby radius, *Ocean Sci. Discuss*, 10, 1807–1831, <https://doi.org/10.5194/osd-10-1807-2013>, 2013.

Okubo, A.: Horizontal dispersion of floatable particles in the vicinity of velocity singularities such as convergences, *Deep Sea Research and Oceanographic Abstracts*, 17, 445–454, [https://doi.org/10.1016/0011-7471\(70\)90059-8](https://doi.org/10.1016/0011-7471(70)90059-8), 1970.

820 Pedlosky, J.: *Geophysical Fluid Dynamics*, Springer edn., 1982.

Pickart, R. S., Weingartner, T. J., Pratt, L. J., Zimmermann, S., and Torres, D. J.: Flow of winter-transformed Pacific water into the Western Arctic, *Deep Sea Research Part II: Topical Studies in Oceanography*, 52, 3175–3198, <https://doi.org/10.1016/j.dsr2.2005.10.009>, 2005.

Planat, N., Tremblay, L. B., Dufour, C. O., and Straub, D.: Seasonal and Decadal Geostrophic Pathways of Pacific and Atlantic Waters in the Arctic Amerasian Basin From Observations, *Journal of Geophysical Research: Oceans*, 130, e2024JC021560, 825 <https://doi.org/10.1029/2024JC021560>, 2025.

Proshutinsky, A., Krishfield, R., Timmermans, M.-L., Toole, J., Carmack, E., McLaughlin, F., Williams, W. J., Zimmermann, S., Itoh, M., and Shimada, K.: Beaufort Gyre freshwater reservoir: State and variability from observations, 2009.

Rampal, P., Weiss, J., and Marsan, D.: Positive trend in the mean speed and deformation rate of Arctic sea ice, 1979–2007, *Journal of Geophysical Research: Oceans*, 114, <https://doi.org/10.1029/2008JC005066>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2008JC005066>, 2009.

830 Reagan, J. R., Seidov, D., Wang, Z., Dukhovskoy, D., Boyer, T. P., Locarnini, R. A., Baranova, O. K., Mishonov, A. V., Garcia, H. E., Bouchard, C., Cross, S. L., and Paver, C. R.: World Ocean Atlas 2023, Volume 2: Salinity, <https://doi.org/10.25923/70QT-9574>, publisher: NOAA National Centers for Environmental Information, 2024.

Regan, H. C., Lique, C., and Armitage, T. W. K.: The Beaufort Gyre Extent, Shape, and Location Between 2003 and 2014 From Satellite Observations, *Journal of Geophysical Research*, 2019.

835 Regan, H. C., Lique, C., Talandier, C., and Meneghelli, G.: Response of Total and Eddy Kinetic Energy to the Recent Spinup of the Beaufort Gyre, *JOURNAL OF PHYSICAL OCEANOGRAPHY*, 50, 2020.

Rieck, J. K., Dufour, C. O., Nadeau, L.-P., and Thompson, A. F.: Heat Transport toward Sea Ice by Transient Processes and Coherent Mesoscale Eddies in an Idealized Southern Ocean, *Journal of Physical Oceanography*, 55, 377–396, <https://doi.org/10.1175/JPO-D-24-0073.1>, 2025a.

840 Rieck, J. K., Martínez Moreno, J., Lique, C., Dufour, C. O., and Talandier, C.: Mean Kinetic Energy and Its Projected Changes Dominate Over Eddy Kinetic Energy in the Arctic Ocean, *Geophysical Research Letters*, 52, e2025GL117957, <https://doi.org/10.1029/2025GL117957>, 2025b.

Smith, K. S. and Vallis, G. K.: The Scales and Equilibration of Midocean Eddies: Freely Evolving Flow, *Journal of Physical Oceanography*, 31, 554–571, [https://doi.org/10.1175/1520-0485\(2001\)031<0554:TSAEOM>2.0.CO;2](https://doi.org/10.1175/1520-0485(2001)031<0554:TSAEOM>2.0.CO;2), 2001.

845 Spall, M. A., Pickart, R. S., Fratantoni, P. S., and Plueddemann, A. J.: Western Arctic Shelfbreak Eddies: Formation and Transport, *Journal of Physical Oceanography*, 38, 1644–1668, <https://doi.org/10.1175/2007JPO3829.1>, 2008.

Spall, M. A., Pickart, R. S., Li, M., Itoh, M., Lin, P., Kikuchi, T., and Qi, Y.: Transport of Pacific Water Into the Canada Basin and the Formation of the Chukchi Slope Current, *Journal of Geophysical Research: Oceans*, 123, 7453–7471, <https://doi.org/10.1029/2018JC013825>, publisher: Blackwell Publishing Ltd, 2018.

850 Stegner, A., Le Vu, B., Dumas, F., Ghannami, M. A., Nicolle, A., Durand, C., and Faugere, Y.: Cyclone-Anticyclone Asymmetry of Eddy Detection on Gridded Altimetry Product in the Mediterranean Sea, *Journal of Geophysical Research: Oceans*, 126, e2021JC017475, <https://doi.org/10.1029/2021JC017475>, 2021.

Talandier, C. and Lique, C.: CREG12.L75-REF12, <https://doi.org/10.5281/ZENODO.13491948>, 2024.

855 Timmermans, M.-L., Toole, J., Proshutinsky, A., Krishfield, R., and Plueddemann, A.: Eddies in the Canada Basin, Arctic Ocean, Observed from Ice-Tethered Profilers, *Journal of Physical Oceanography*, 38, 133–145, <https://doi.org/10.1175/2007JPO3782.1>, 2008.

Toole, J., Krishfield, R., Timmermans, M.-L., and Proshutinsky, A.: The Ice-Tethered Profiler: Argo of the Arctic, *Oceanography*, 24, 126–135, <https://doi.org/10.5670/oceanog.2011.64>, 2011.

Tulloch, R., Marshall, J., Hill, C., and Smith, K. S.: Scales, Growth Rates, and Spectral Fluxes of Baroclinic Instability in the Ocean, *Journal 860 of Physical Oceanography*, 41, 1057–1076, <https://doi.org/10.1175/2011JPO4404.1>, 2011.

Vancoppenolle, M., Rousset, C., Blockley, E., Aksenov, Y., Feltham, D., Fichefet, T., Garric, G., Guémas, V., Iovino, D., Keeley, S., Madec, G., Massonnet, F., Ridley, J., Schroeder, D., and Tietsche, S.: SI3, the NEMO Sea Ice Engine, <https://doi.org/10.5281/ZENODO.7534900>, publisher: Zenodo Version Number: 4.2release_doc1.0, 2023.

von Appen, W.-J., Baumann, T. M., Janout, M. A., Koldunov, N. V., Lenn, Y.-D., Pickart, R. S., and Wang, Q.: Eddies and the Distribution
865 of Eddy Kinetic Energy in the Arctic Ocean, 35, 42–51, 2022.

Wang, Q., Koldunov, N. V., Danilov, S., Sidorenko, D., Wekerle, C., Scholz, P., Bashmachnikov, I. L., and Jung, T.: Eddy Kinetic Energy in the
Arctic Ocean From a Global Simulation With a 1-km Arctic, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL088550>,
870 publisher: Blackwell Publishing Ltd, 2020.

Watanabe, E.: Beaufort shelf break eddies and shelf-basin exchange of Pacific summer water in the western Arctic Ocean detected by satellite
and modeling analyses, *Journal of Geophysical Research*, 116, C08 034, <https://doi.org/10.1029/2010JC006259>, 2011.

875 Watanabe, E., Onodera, J., Harada, N., Honda, M. C., Kimoto, K., Kikuchi, T., Nishino, S., Matsuno, K., Yamaguchi, A.,
Ishida, A., and Kishi, M. J.: Enhanced role of eddies in the Arctic marine biological pump, *Nature Communications*, 5, 3950,
<https://doi.org/10.1038/ncomms4950>, 2014.

Weiss, J.: The dynamics of enstrophy transfer in two-dimensional hydrodynamics, *Physica D: Nonlinear Phenomena*, 48, 273–294,
880 [https://doi.org/10.1016/0167-2789\(91\)90088-Q](https://doi.org/10.1016/0167-2789(91)90088-Q), 1991.

Weiss-Gibbons, T., Tefs, A., Hu, X., Stadnyk, T., and Myers, P. G.: Sensitivity of Simulated Arctic Ocean Salinity and Strait Trans-
port to Interannually Variable Hydrologic Model Based Runoff, *Journal of Geophysical Research: Oceans*, 129, e2023JC020536,
<https://doi.org/10.1029/2023JC020536>, 2024.

Woodgate, R. A.: Increases in the Pacific inflow to the Arctic from 1990 to 2015, and insights into seasonal trends and driving mechanisms
885 from year-round Bering Strait mooring data, *Progress in Oceanography*, 160, 124–154, <https://doi.org/10.1016/j.pocean.2017.12.007>,
publisher: Elsevier Ltd, 2018.

Zhang, J. and Rothrock, D. A.: Modeling Global Sea Ice with a Thickness and Enthalpy Distribution Model in Generalized Curvilinear
Coordinates, *Monthly Weather Review*, 131, 845–861, [https://doi.org/10.1175/1520-0493\(2003\)131<0845:MGSIWA>2.0.CO;2](https://doi.org/10.1175/1520-0493(2003)131<0845:MGSIWA>2.0.CO;2), 2003.

Zhang, J., Steele, M., Runciman, K., Dewey, S., Morison, J., Lee, C., Rainville, L., Cole, S., Krishfield, R., Timmermans, M.-L., and Toole,
885 J.: The Beaufort Gyre intensification and stabilization: A model-observation synthesis, *Journal of Geophysical Research*, 2016.

Zhao, M. and Timmermans, M. L.: Vertical scales and dynamics of eddies in the Arctic Ocean’s Canada Basin, *Journal of Geophysical
Research: Oceans*, 120, 8195–8209, <https://doi.org/10.1002/2015JC011251>, publisher: Blackwell Publishing Ltd, 2015.

Zhao, M., Timmermans, M.-L., Cole, S., Krishfield, R., Proshutinsky, A., and Toole, J.: Characterizing the eddy field in the Arctic Ocean
halocline, *Journal of Geophysical Research: Oceans*, 119, 8800–8817, <https://doi.org/10.1002/2014JC010488>, 2014.