

Natural wetland methane emissions simulated by ICON-XPP

Stiig Wilkenskjeld¹, Thomas Kleinen¹, Tobias Stacke¹, and Victor Brovkin^{1,2}

¹Max Planck Institute for Meteorology, Hamburg, Germany

²CEN, University of Hamburg, Hamburg, Germany

Correspondence: Thomas Kleinen (thomas.kleinen@mpimet.mpg.de)

Abstract. ~~Methane emissions from natural wetlands account for about 1/3 of~~ Natural wetlands emit roughly one third of the
global methane emissions, ~~and thus have a significant climatic impact due to methane's high global warming potential. Among~~
~~the sources of methane, those from natural wetlands have the highest uncertainty, and it is thus of key importance to understand~~
~~and reduce the uncertainties in the estimates of the wetland emissions to be able to close the global methane budget. Using~~
5 ~~a coupled land and atmosphere setup, a new implementation of interactive wetlands and wetland methane emissions yet their~~
~~contribution remains highly uncertain. To reduce this uncertainty, we incorporated a fully coupled wetland-hydrology and~~
~~wetland-methane module~~ into the ICON-XPP Earth System Model was used to test the sensitivity of the wetland methane
~~emissions to a number of hydrological and biogeochemical model assumptions of which some are mimicing anthropogenic~~
~~influences on the Earth System. Averaged over the historical period~~ (Earth-system model and performed a suite of AMIP-type
10 ~~experiments spanning 1855-2014) the simulated emissions are 166.2 (156.3–181.2) Tg(CH₄) yr⁻¹. For the 2000-2012 period~~
~~the equivalent numbers 182.3 (154.3–205.1. Wetland extent is diagnosed online with a TOPMODEL-based wetland scheme~~
~~that exploits high-resolution topographic indices, while methane production follows a temperature-dependent, anaerobic decomposition~~
~~formulation calibrated to recent global budgets.~~

The baseline simulation yields a time-averaged natural wetland methane flux of 182 (154-205) Tg(CH₄) yr⁻¹ are in good
15 ~~agreement with estimates from other studies. Wetland methane emissions rise by about 12% during the historical period, mainly~~
~~since 1980, an increase which is due to an enhanced carbon cycle caused by the~~ for the recent period 2000-2012, in line with
the multi-model mean of Saunio et al. (2016). Sensitivity experiments reveal that:

1. ~~CO₂ fertilization associated with rising atmospheric CO₂ concentrations. The modeled emissions are very sensitive to~~
~~changes and assumptions in the model hydrology, some dependencies only revealed through the interaction with the~~
20 ~~atmosphere (changes in the moisture recycling patterns). Therefore offline land models are only of limited value to test~~
~~the influence of changes in model hydrology, which is also influenced by changes in terrestrial vegetation~~ fertilisation
is the dominant driver of the observed 12% increase in emissions since the late 19th century; fixing terrestrial CO₂ at
preindustrial levels reverses the trend, producing a net decline in methane flux.
2. Anthropogenic drainage of croplands curtails potential wetland area, especially in the northern extratropics, and offsets
25 part of the CO₂-driven rise in emissions. Experiments that relax drainage or freeze historical land-use increase wetland
area by 0.2-0.6 Mm² and raise emissions.

3. Surface-water-retention (SWR) strongly modulates regional precipitation recycling in a coupled model setup, thus significantly altering precipitation in comparison to uncoupled offline experiments.

Overall, the coupled model reproduces observed wetland extents and methane emissions, but suggests that current Earth-system models likely underestimate human-induced hydrological alterations, particularly drainage and water-management practices. Our results underscore the necessity of interactive land-atmosphere coupling and refined hydrological representations for robust projections of future wetland methane feedbacks.

1 Introduction

Methane is ~~—~~ after CO₂ ~~—~~ the second most important greenhouse gas on which humans have a significant direct impact (Zhang et al., 2017; Saunois et al., 2016, 2020, 2024). Since the mid ~~1980's~~ 1980s, the atmospheric methane concentration has been rising by about 7 ppb per year (Lan et al., 2025), with an increasing trend since 2005. Due to its much higher global warming potential compared to CO₂, methane currently contributes almost ~~1/3~~ one third of the anthropogenic climate change (Forster et al., 2021). In combination with the rather short atmospheric lifetime of methane (about 9 years, Saunois et al. (2024)) this could make methane emission reductions an important ~~contributor to reach~~ contribution to reaching the goals of the Paris agreement on a decadal time scale (UNEP, 2021).

Methane is emitted to the atmosphere from a number of marine and terrestrial sources, of which some are anthropogenic, some natural and some natural but influenced by human activities. Emissions from natural wetlands are the largest of the latter type, as they contribute about ~~1/3~~ one third of the total emissions (Saunois et al., 2016, 2020, 2024). Despite estimate uncertainties, it is commonly agreed, that ~~at least in the recent years,~~ anthropogenic sources have grown larger than the natural ones (Saunois et al., 2024), at least in recent years.

Modeling of wetland methane emissions includes many poorly quantified parameters and processes, which need to be either estimated from observations or modeled interactively. Since methane production takes place under anaerobic conditions, it is closely coupled to the hydrology of the soils: ~~—~~ which is used to determine areas where soil conditions are anaerobic. The performance of the current generation of hydrology models is ~~—positively stated—moderately good~~ (Krysanova et al., 2020; Melton et al., 2013) ~~—With—, however, less than perfect~~ (Krysanova et al., 2020; Melton et al., 2013). Furthermore, with few exceptions (e.g., Veldkamp et al. (2018)) land surface models used in global Earth System Models ignore active human water management. ~~Human activities such—, including such human activities~~ as dam construction, drainage, irrigation and changes in land use ~~may however~~ that might severely impact the hydrology on local to regional scales (de Vrese et al., 2018; Xu et al., 2024; Chen et al., 2025). ~~Such measures have historically drained~~ Historically, however, such measures have led to the drainage of large wetland areas (Davidson, 2014; Gardner and Finlayson, 2018) ~~(Davidson, 2014; Gardner and Finlayson, 2018; Fluet-Chouinard et al., 2023)~~ mainly since the year 1900.

In the high latitudes the presence of (thawing) permafrost further adds uncertainties in wetland estimates (Hagemann et al., 2016; Andresen et al., 2020). The state of the permafrost highly impacts the hydrology of the soils (de Vrese et al., 2023, 2024). Due to small evapotranspiration and the sub-surface runoff partly being blocked by frozen soils, a significant part of global

60 wetlands is located in permafrost areas. Many permafrost-related processes take place at scales far smaller than the resolution of global models (e.g., Rehder et al. (2023)) and depend on subsurface developments, which are presently poorly understood and only possible to observe on local scale ~~-(Patel et al., 2024).~~ Upscaling these processes to pan-Arctic or even to typical global model grid point scale to obtain robust estimates of soil moisture and wetland area is even more challenging.

Observational estimates of wetland area comes with their own issues and challenges. Zhang et al. (2017) reported a range in 65 the global, ~~yearly annual~~ mean wetland area from five commonly used different observational products from 5.3 to 10.2 Mm² (~~1Mm² = 10¹²m² = 10⁶km²~~ 1 Mm² = 10¹² m² = 10⁶ km²). Also, large disagreement on the global patterns and annual cycle was reported due to differences in methods, observed periods and original purpose of the data set.

Soil methane production requires available soil organic matter (SOM), anaerobic conditions and sufficiently high temperatures, ~~all parameters which.~~ All of these parameters are estimated by various model components, all of which ~~comes come~~ 70 with their own uncertainties. Over decades, atmospheric and land models have gained good skills in simulating temperatures and carbon content, whereas the ~~oxie oxygen~~ state of the soils have until recently only received little attention.

Here we present the results of a new implementation of wetland methane emissions based on interactively calculated wetlands in the ICON-XPP Earth System Model and the sensitivity to different model assumptions about hydrological management and biogeochemical influences. This will hopefully contribute to a better understanding of the controls on the modeled natural 75 methane ~~emission~~emissions, and thus aims to identify and understand uncertainties in the current estimates of wetland methane emissions.

2 Model and development

This study ~~is utilizing~~ utilizes the ICON Earth System Model, using the ICON-XPP atmosphere model (Müller et al., 2025) and the land surface model JSBACH4 (Schneck et al., 2022). To enable the model to estimate wetland areas and methane 80 emissions, a number of extensions and modifications to the model code ~~has have~~ been implemented. These are described in the following subsections.

2.1 Wetlands

~~Wetland area fraction was estimated~~ In global land-surface modelling the term “wetland” is commonly used in the sense of a hydrological diagnostic, as opposed to an ecosystem characterization: a grid cell (or sub-grid fraction) is labelled a 85 wetland when the simulated water table reaches or exceeds the soil surface, which creates anaerobic conditions that allow methanogenesis. This definition does not imply any change in vegetation type: often, the vegetation cover is prescribed from boundary condition data sets, in our case the from the LUH2 land use dataset (Hurtt et al., 2020). This then remains unchanged, regardless of the water-table status. Consequently, a cell that is classified as a wetland in our model may still be labelled “cropland” in the land use map and retain the same crop vegetation. The term “wetland” therefore refers only to the soil-water 90 state that controls methane production, not to the ecological concept of a wetland ecosystem.

We estimated the wetland area fraction using a diagnostic TOPMODEL approach (Beven and Kirkby, 1979), where the compound topographic index (CTI), which describes the relation between the (hydrological) upstream area and the local slope for any point, is used to relate the ~~grid-mean-grid-mean~~ water table to the ~~water-table-water-table~~ distribution on the sub-grid-scale topography. The wetland fraction is then assumed to be the fraction of the grid cell where the water table is at or above the soil surface.

The CTI index was obtained ~~as-the-maximum-value-of-from~~ the data set presented in Marthews et al. (2015) ~~and~~. However, ~~in order to avoid unrealistically low CTI values that would generate spurious wetlands, we enforce a minimum CTI value of 5.~~

The calculation of the wetland area does not directly feed back to the soil hydrology. ~~It-Instead~~, is assumed that the part of the soils, which are diagnosed as inundated using this approach, are anaerobic. Thereby the wetland ~~extend-extent~~ has an influence on the carbon cycle, which again influences the hydrology (see ~~See-Section~~ 2.2).

We extended the scheme presented in detail in Kleinen et al. (2020) with two improvements:

- The simple piecewise linear estimate of the ice fraction of soil water used in Kleinen et al. (2020) has been replaced by calculations internally in JSBACH’s soil hydrology (Ekici et al., 2014; Hagemann and Stacke, 2015). Thus, the freezing and melting of soil water now also influences both the hydraulic conductivity, the energy calculations of the soil and the soil carbon handling.
- In flat areas with shallow soils (e.g., the Sahara), TOPMODEL tends to produce spurious wetlands as a consequence of individual precipitation events, since the shallow soils are rapidly “filled up”. To exclude such artificial wetlands while preserving TOPMODEL’s capability to react to changing climatic conditions, an additional dynamic condition to exclude wetlands from very dry areas was introduced: We require that

$$\frac{\overline{PR}}{\overline{AET}} > 0.3 \quad (1)$$

~~were-where~~ PR is precipitation, AET is the equilibrium evapotranspiration, defined by ~~Eq-Equation~~ (2) and an ~~overline~~ ~~overbar~~ indicates a temporal mean over a “long” period (in this study defined as a running average with an e-folding time of 1 year). This ~~criteria-criterion~~ excludes the driest areas while still letting the wetlands develop freely in the rest of the world.

The equilibrium evaporation AET is, following Prentice et al. (1993), defined as:

$$AET = \frac{1}{\lambda} \frac{s}{s + \gamma} NetRad \quad (2)$$

with the rate of change of the saturated vapor pressure: $s = 2.503 \cdot 10^6 e^{\frac{17.269}{t_{ref}} \frac{t}{t_{ref}}}$, the psychrometer constant: $\gamma = 65.05 + 0.064t$, the latent heat of evaporation: $\lambda = 2.495 \cdot 10^6 - 2380t$, $t_{ref} = t + 237.3$, $t_{ref} = t + 273.3$, t is the 2m air temperature in °C and the $NetRad$ is the net radiative balance in $W m^{-2}$.

The diagnostic wetland fraction is evaluated at every model time step (7.5 min). The atmospheric component (ICON-XPP) runs on a triangular C-grid with an R2B4 configuration, i.e., the spacing between grid-cell centers is of the order of 160 km.

125 The CTI field that drives the TOPMODEL calculation is obtained from the global dataset of Marthews et al. (2015) at 15 arc seconds ($\approx 500\text{m}$) resolution. Thus, the hydrological diagnostic benefits from a high-resolution topographic input, but the underlying climate and land-surface fields are resolved only on the coarse 160 km grid. Consequently, fine-scale features, such as polygonal tundra or palsas complexes in the permafrost region (which have characteristic lengths of a few hundred metres to a few kilometres), are not explicitly represented.

2.2 Wetland methane emissions

The wetland methane production model implementation is adopted from Kleinen et al. (2020), which is based on the approach by Riley et al. (2011). ~~The basic assumption is, that In this model, the decomposition of SOM in anaerobic soils both produces~~ soil organic matter (SOM) is assumed to occur under anaerobic conditions in the diagnosed grid cell wetland fraction (section 2.1). Due to the anaerobic conditions, decomposition products are not just CO_2 , as under aerobic conditions, but both CO_2 and methane. ~~The anaerobic decomposition pathway is much slower than the aerobic and thus the SOM is changed differently depending on the oxic state of the soil. Since the soil properties depend on the SOM content, this constitute an indirect feedback to the hydrological cycle. Following Kleinen et al. (2020, 2021); Wania et al. (2010), we assume a time and space independent~~ CH_4 . The model then transports CO_2 , CH_4 and O_2 vertically through the soil column by considering diffusion, ebullition, and plant-mediated transport through aerenchyma, also determining the oxidation of CH_4 in soil layers where sufficient O_2 is present.

140 In laboratory incubations under strictly anaerobic conditions, SOM mineralisation yields a 50% CO_2 : 50% CH_4 split (Conrad, 1999). However, in actual wetland soils this ratio is usually much greater than 1:1, and can vary significantly among different kinds of wetlands (Bridgham et al., 2013). In our model, we therefore assume a significantly lower fraction of CH_4 production, which we furthermore assume to be temperature-dependent in a Q_{10} formulation. We calibrated this to give a 24% CH_4 fraction at 295 K and a $Q_{10}=1.2$ (Riley et al., 2011). This formulation does not distinguish between acetoclastic and hydrogenotrophic methanogenesis, which is the standard approach in most global methane models. Distinguishing acetoclastic from hydrogenotrophic methanogenesis would require explicit representation of substrate availability (acetate vs. H_2/CO_2) and is not currently implemented. A recent intercomparison (Xu et al., 2016) reported that 37 out of 40 global wetland-methane models do not separate the two pathways, highlighting that this simplification is common practice, albeit a known source of uncertainty.

145 Following Wania et al. (2010), we furthermore assume a time- and space-independent reduction of SOM decomposition rate, so that wetland SOM is decomposed with 35%-% of the aerobic efficiency. ~~The fraction of total decomposed SOM converted to methane (the rest is assumed to become) is determined by a Q_{10} formulation, with 24% methane fraction at 295K and a Q_{10} value of 1.2.~~

~~The produced methane decomposition rate.~~

155 The produced methane, along with CO_2 and O_2 , is transported vertically through the soil column via diffusion, ebullition and plant aerenchyma. ~~On it's way it,~~ with process descriptions and parameters adopted from Riley et al. (2011). Depending on the availability of O_2 , which can be supplied either via diffusion or via plant aerenchyma, CH_4 may be oxidized partly or

completely ~~using oxygen which is diffused downward through the soil column. The oxidation of methane is,~~ also parameterized using a ~~Q10 formulation with a Q10~~ Q₁₀ formulation with a Q₁₀ value of 1.9.

Finally, ~~uncertainty ranges for parameters used in the wetland emission model are sufficiently large that the total emissions (but not the spatial distribution) can be calibrated over a range of wetland area estimates. Thus, we used the CO₂:CH₄ ratio in methane production to calibrate our model to yield global total emissions similar to the Sauniois et al. (2016) estimate.~~

2.3 Anthropogenic drainage of croplands

~~To maximize agricultural gains, it has long been practiced to regulate the soil hydrology of crop areas (e.g. Valipour et al. (2020)). This involved draining excess water away from crop areas. This practice has been build into the model by prohibiting wetlands on crops. In the current implementation, the drainage affects only the wetland area and not the basic soil hydrology below crops.~~

~~To include drained crops furthermore~~ Historically, many agricultural regions have been drained to remove excess water from fields (Valipour et al., 2020). In the model, this practice is represented by suppressing the diagnostic wetland fraction on all grid cell fractions that are classified as cropland in the LUH2 land use map. Importantly, this drainage flag only suppresses the wetland diagnostics (i.e., the calculation that would label the cell as anaerobic and allow methane production). It does not modify the underlying soil-hydrology equations (soil moisture, infiltration, runoff) nor the carbon-cycle fluxes beneath the crops. The vegetation type therefore remains the prescribed crop (or pasture) throughout the simulation, no transition to natural wetland vegetation (e.g., emergent macrophytes) occurs. Furthermore, the inclusion of drained crops has the advantage, that methane emissions from rice fields (Sauniois et al., 2020) are unequivocally attributed as an anthropogenic source and thus ~~falls~~ fall out of the scope of this study.

2.4 Surface water retention (SWR)

The standard JSBACH model uses the ARNO-scheme (Hagemann and Gates, 2003; Hagemann and Stacke, 2015) to separate rain and melting snow into surface runoff and infiltration into the soil, based on the slope distribution of the grid cell. This scheme ignores that water might pile up in ~~small-scale~~ small-scale topographic depressions forming ponds or puddles, from where the water is available for ~~evapotranspiration, evaporation~~ evapo(transpi)ration or later infiltration into the soils. ~~Thus~~ Thus, it may suppress important water and energy feedbacks between surface and atmosphere. It is assumed, that in addition to the micro-topographical effects, this water retention takes place ~~were~~ where local soil surface hydraulic conductivity is low. Therefore, the conductivity of these areas is assumed to be equivalent to that of clay, which is the natural soil type with the lowest conductivity. To account for these effects, we implemented the SWR scheme described in detail in ~~de Vrese et al. (2021); Stacke and Hagemann (2012)~~ Stacke and Hagemann (2012); de Vrese et al. (2021).

It is assumed that the local SWR is a temporary, though possibly repeated, phenomenon compared to the wetlands arising due to saturated soils (~~See~~ Section 2.1). The ~~area~~ areas inundated by SWR is therefore not counted as wetlands. Due to the lower infiltration in areas with SWR, the soil is comparatively dry, and ~~thus~~ we assume that these soils stay oxic and ~~do thus~~ therefore do not contribute to the wetland methane production.

3 Setup, experiments and boundary conditions

The experiments were ~~done~~ performed with ICON-XPP (Müller et al., 2025) with the extensions discussed in ~~See~~ Section
190 2. ICON uses a triangular grid and was in this study run at the R2B4 resolution, which has a cell size of about 25,000 km²,
roughly corresponding to a resolution of 160 km in a grid with square cells. The atmospheric model used 90 vertical layers,
where the upper layer has a pressure of about 1 Pa, and the land model contained 5 soil layers with ~~in-creasing~~ increasing
thickness down to a total soil depth of at most ≈ 9.5 m, though generally the soil is much shallower.

The model was run using the concentration driven AMIP configuration ~~—~~ that is: land-coupled-to-the the model describes
195 a land surface coupled to an atmosphere, while ocean sea surface temperature, sea ice and greenhouse gas concentrations
are prescribed (sea surface temperature and ice using data described in Taylor et al. (2000)). Due to the prescription of the
greenhouse gases ~~—~~ mainly CO₂ and methane ~~—~~ there is no direct feedback from the terrestrial carbon cycle on to the
atmosphere. Thus, this study does not deliver any estimates of atmospheric concentration or lifetime of the methane.

The model was spun up from an initial state without carbon for 160 years, then the slow humus soil pools of the YASSO
200 soil carbon model (Goll et al., 2015; Tuomi et al., 2009) of JSBACH were equilibrated, and the model was spun up for another
50 years. The spinup was done using the vegetation distribution from the LUH2 data set (Hurtt et al., 2020) and greenhouse
gas concentrations (GHGs) for the year 1855. During the spinup, the SWR scheme was switched on and crop areas drained as
described above.

From this spinup, five experiments (~~Tab.~~ Table 1) were ~~started, all branched off, each~~ running from 1855 to 2014, to get
205 obtain a best estimate of the natural wetland methane emissions and to explore the influence of different model assumptions:
~~The “Base” experiment is a direct continuation of the spinup, using the SWR scheme and crop drainage, but it switches to~~
~~transient GHGs and annually prescribed land-use maps derived from the~~ :

Base Fully coupled experiment with SWR active, cropland drainage applied, and transient land-use (LUH2 data. Other experiments
switch-off the SWR scheme (“DrySurf”), and CO₂ concentrations.

210 DrySurf Same as Base, but surface-water-retention (SWR) disabled.

ConstCO2 Same as Base, but the CO₂ concentration in the terrestrial carbon cycle model is fixed at the 1855 value (while the
atmospheric CO₂ used for radiative forcing remains transient).

WetCrops Same as Base, but cropland drainage is disabled; the prescribed LUH2 crop fraction can become wetland (the
drainage (“WetCrops”), vegetation type remains the LUH2 crop, i.e., no shift to wetland plant functional types).

215 ConstVeg Same as Base, but the vegetation map is frozen at the 1855 LUH2 distribution; land-use changes (including cropland
expansion) are therefore ignored, while the land-cover changes (“ConstVeg”) or the transient CO₂ concentration seen by
the vegetation model, while preserving the transient GHGs used in the atmospheric model (“ConstCO2”) respectively.
hydrology (precipitation, runoff, SWR) remains fully interactive.

The experiments are stopped in 2014 due to ~~a~~ lack of newer boundary ~~condition~~ data for the atmospheric model. Parts of the
220 analysis therefore concentrate on the period 2000-2012, which was the main focus of Saunois et al. (2016, 2017).

In this study only the data from the terrestrial areas are analyzed, and thus the term “global” hereafter only refers to the
terrestrial surface. Furthermore, often the results are discussed separately for a number of regions (~~Fig-~~Figure S1). The most
often used regions are the northern extratropics (NXT), defined as the area north of 30°N, and the tropics, which (unless
otherwise explicitly stated) ~~—following Saunois et al. (2017)—~~ ~~following Saunois et al. (2017) –~~ covers the rest of the
225 world. The latter definition is justified since the southern extratropics only contribute an insignificant fraction of the global
wetland ~~area~~ and wetland methane emissions.

The LUH2 data set prescribes ~~5.8Mm²~~ ~~5.8Mm²~~ of crops in 1855 (~~3.6Mm²~~ ~~3.6Mm²~~ located in the NXT area). Globally
the crop area increase almost linearly to ~~15.1Mm²~~ ~~15.1Mm²~~ in 2014. However, the NXT contribution stabilizes at about
~~7.9Mm²~~ ~~7.9Mm²~~ around 1960 and decreases after around 1990 to about ~~7.1Mm²~~ ~~7.1Mm²~~ in 2014. The increase in crop area
230 is wide-spread in non-mountainous areas of all continents, the exceptions being the north eastern US and Western Europe,
where crop areas decrease in the last part of the experiment period. The crop distribution and its changes are important for a
proper interpretation of the results from several of the experiments.

4 Results

4.1 “Base” experiment including SWR

235 The average July–August (JJA) wetland distribution (~~Fig-~~Figure 1a) shows the main wetland areas in the tropical rain forests
of South America and Africa, as well as in the Hudson Bay area and central northern Siberia. Smaller, but still important, areas
are found in India, eastern North America and larger parts of northern Siberia. In addition, some wetlands are found in central
western Europe, south-eastern US, Indonesia and eastern China. Roughly, the majority of the northern hemisphere summer
wetland area is concentrated in two latitudinal bands: ~~The-the~~ boreal zone (55°N - 70°N) ~~contributing with,~~ ~~contributing~~ 34%
240 of the global area ~~and the in,~~ ~~and the~~ tropics (20°S - 15°N) with 38%. The global JJA wetland area is ~~≈5.2Mm²~~ ~~≈5.2Mm²~~,
with a slight decrease over time (2% per century). This trend over the ~~entire~~ experiment period, however, hides ~~the fact~~ that the
wetlands are only diminishing ~~in-during~~ the first part of the experiment, while they are growing again ~~after-around-1980~~ at a
rate of 7% per century ~~-after 1980~~. This increase is mainly driven by changes in the NXT area, though ~~also-in-the-tropics-the~~
~~trend-turns-the trend also turns the tropics~~ from negative to positive over the ~~experiment-Actually~~ ~~course of the experiment~~.
245 ~~In fact,~~ the wetland area in ~~the~~ NXT region is slightly ~~higher-in-the-latter~~ ~~larger in the later~~ part of the experiment than in the
~~experiment-long-long-term~~ average, and is thus opposing the global decreasing trend (~~Fig-~~Figure 4a), ~~so-that-leading-to-an~~
~~increase in~~ the NXT fraction of the global wetland areas ~~increase~~ from < 30% in the ~~1940's~~ ~~1940s~~ to 33% in the last decade
of the experiment (2005-2014). ~~In-At~~ the start of the experiment, this fraction was about 31.6%.

The geographical trend distribution (~~Fig-~~Figure 1b), shows areas of both increasing and decreasing wetlands: Decreasing
250 wetlands are found in the east-central US, eastern Europe, southern Brazil and India/Bangladesh, while increasing wetlands

are found in eastern US and western Europe. In general there is a close relation between the development of cropland (and thus drainage, ~~Fig-Figure~~ S3) fraction and the development of the wetlands.

The average simulated wetland area on non-snow-covered areas in “Base” is $\approx 3.8\text{Mm}^2$ with a pronounced annual cycle (~~Fig-Figure~~ 2), ranging from $\approx 2.8\text{Mm}^2$ to $\approx 5.4\text{Mm}^2$. The year to year variation on the other hand is small ($1\sigma = 0.06\text{Mm}^2$).
255 The annual cycle stems only from the NXT areas, with maximum in NH summer, June to September, where the northern extratropics are snow free. Though the NXT wetland area in the non-NH-summer increase by up to almost 1Mm^2 (November) if no snow masking is applied, the shape of the ~~yearly-annual~~ cycle remains. In the ~~yearly-annual~~ average, snow is covering $\approx 0.5\text{Mm}^2$ of wetlands. The lack of an annual cycle in the tropics is to some degree a result of a cancellation of the annual cycles on the difference continents being out of phase (~~Fig-Figure~~ S2). The rather weak South American wetland annual cycle
260 peaks in April–July and has its minimum in October–January, quite opposite to ~~the~~ somewhat more pronounced annual cycle in Africa, which is the other main tropical wetland region. South East Asia and tropical North America ~~contributes-contribute~~ only with very small wetland areas, leaving their influence on the annual cycle diminutive.

Global wetland methane emissions averaged over the entire experiment period ~~amounts-amount~~ to $166.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$, of which $35.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$, or about 21%, are emitted in the NXT area. The rest stems almost exclusively from the tropical rain
265 forest areas in South America, Africa and Indonesia, together with India (~~Tab.-3 and Fig-Table 3 and Figure~~ 3a). The relation between the boreal and tropical wetland latitudinal bands is thus quite different with respect to methane emissions: The boreal band contributes only 8% while the tropical band emits 67% of global wetland methane. Nzotungicimpaye et al. (2021) report a similar distribution of wetland methane emissions between the boreal and tropical regions using a more microbiologically detailed methane model.

270 For the period 2000-2012 the global average wetland methane emission is $182.7 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$, in close agreement with the multi-model mean presented in Saunois et al. (2016)~~-, who report $185\text{Tg}(\text{CH}_4) \text{ yr}^{-1}$ for 2003-2012, with a 1σ standard deviation of $21\text{Tg}(\text{CH}_4) \text{ yr}^{-1}$~~ .

However, our emissions are more scattered, most pronounced in the Eurasian boreal areas. Higher methane emissions are found in western Europe and southeastern US and lower emissions are found in South East Asia, eastern Africa and the Hudson
275 Bay area. These differences in patterns are quite similar to the multi-model mean of wetland fractions presented in Hardouin et al. (2024).

The emissions are increasing over time with a trend over the entire experiment of about $20 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$ (8% per century). After around 1980, the trend is about ~~4-four~~ times as large compared to the entire experiment (~~Tab.-3, Fig-Table 3, Figure~~ 4d). The global increase is mainly driven by increases in the large production areas in South America and Africa as well as Western
280 Europe (~~Fig-Figure~~ 3b). Decreasing emissions are found in the Central US.

~~The standard deviation (obtained from-To obtain the standard deviation of~~ the time series~~detrended-by-, we detrended it by separately~~ subtracting the linear regressions in the time periods 1855-1980 and 1981-2014~~separately. The result-. The results~~ show very little effect ~~on-of~~ the exact point of ~~partition-of-partitioning~~ the regression periods and ~~is-are~~ virtually identical between the two periods. ~~)-of-the-yearly-average-emissions-The standard deviation of the annual average emissions thus~~
285 ~~obtained~~ is $\approx 3.4\text{Tg}(\text{CH}_4) \text{ yr}^{-1}$.

The emission fraction (the fraction of produced methane that is not ~~oxidized~~oxidised in the soil and thus reaches the atmosphere) is 44.5% and ~~very~~-constant over the experiment period. The emission fraction is geographically very unevenly distributed in complex patterns, where the ~~fraction~~fractions range from near 0 to almost 80%. In general, the major methane emitting areas have moderate emission fractions of 30-40%, while hardly any methane escapes from dry areas.

290 4.2 No surface water retention (“DrySurf”)

~~Switching-off~~Disabling SWR lowers the average wetland area by about ~~0.1Mm²~~(Fig. 0.1 Mm² (Figure 4a)), with the entire reduction taking place in the NXT area, ~~mainly~~. This mainly happens in the period July – October, ~~where~~and the reduction is almost twice the annual mean. Geographically more specific, the reduction is confined to a narrowing stripe reaching from ~~the northeastern Europe~~turning northeastern Europe into central Siberia and the Mackenzie River basin (~~Fig.~~Figure 5a).

295 The areas of reduced wetlands are within ~~huge~~large regions where the precipitation is much lower in “DrySurf” than in “Base” (~~Fig.~~Figure 5b). In parts of central Siberia the precipitation reduction is ~~more~~larger than 30%. Most of the areas with large and consistent precipitation differences are in the areas of the NH jet stream. This indicates a ~~huge~~significant shift in the magnitude of moisture recycling through the atmosphere, which is supported by a large decrease of evapotranspiration in “DrySurf” compared to “Base” (not shown).

300 In “DrySurf” the NXT wetland area decreases by about 7% per century and thus 5 times faster than in “Base” (~~Fig.~~Figure 4c), whereas the tropical wetlands ~~changes~~change a bit less in “DrySurf” than in “Base”. The geographical patterns of changing trends between “Base” and “DrySurf” ~~is~~coincidingcoincide almost exactly with those of the absolute wetland area difference. The average wetland methane emissions (~~Fig.~~Figure 4c) from “DrySurf” behave ~~very similar~~rather similarly to the wetland area: Slightly lower (~~$\approx 2.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$~~ less emissions ($\approx 2.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$)) from the NXT than in “Base” ~~is~~are partly
305 compensated by an even smaller increase (~~$\approx 0.8 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$~~ $\approx 0.8 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$) from the Tropics.

In the NXT, the decreasing wetland area trend dominates over the generally increasing trend in methane emissions when looking at the entire experiment: A trend of -3% per century opposes the tropical trend of +8% per century. The result is that the global trend is slightly below that of “Base”. Since 1980 this picture is reversed: NXT emissions increase by 42% per century compared to 30% in the tropics.

310 4.3 Neglecting CO₂ ~~fertilization~~fertilisation (“ConstCO2”)

In AMIP ~~runs~~experiments with prescribed CO₂, there is no direct feedback from the terrestrial carbon cycle ~~on~~-to the atmosphere. Thus, it could be expected that the wetlands in “ConstCO2” would be identical to those in “Base”. This seems to be confirmed by the average wetland area (~~Fig.~~Figure 4a), where the two experiments are indeed very close. The trends over the 1855-2014 period, however, are somewhat different, with slightly smaller ~~trend~~(less negative) trends all over the world.

315 In the recent period, the NXT region and the tropics behave very differently: In the tropics the wetland area ~~is~~stagnatingstagnates with hardly any trend, while the positive NXT trend in “ConstCO2” exceeds that of “Base”. These changes arise as a feedback through changed ~~water-use~~water-use efficiency of the plants at different CO₂ levels, changing the transpiration. Thus, both water and energy fluxes to the atmosphere are changed, which then ~~triggers~~trigger different atmospheric responses.

~~Though~~ Although the global numbers of evapotranspiration are very similar in the two experiments, both the differences in the
320 evapotranspiration itself and its trend show complex regional patterns (~~Fig.~~ Figure S5).

Keeping the CO₂ concentration constant at pre-industrial levels for the carbon model has a huge impact on the trend of the methane emissions (~~Fig.~~ Figure 4d). The increasing methane emissions in “Base” are turned into a decrease in the “ConstCO2” and the trend is thus in the ~~other directions opposite direction~~ compared to all other conducted experiments. Only the NXT area has a positive trend in the latter part of the experiment, and this is still smaller than in all other experiments except “ConstVeg”
325 (discussed below). The global carbon litter is ~~—~~ opposite to any other experiment ~~—~~ decreasing after around the year 1900 (~~Fig.~~ Figure S6). There is still a slight increase in the decomposition of the soil carbon, but only about 5% of that in the other experiments. That less carbon respire ~~more~~ is due to a warmer climate increasing the respiration rate. The reversed trend in the methane emissions in “ConstCO2” reveals, that the trend found in the other experiments ~~is solely~~ is largely due to an enhanced carbon cycle ~~due to CO₂ fertilization. Different~~ caused by CO₂ fertilisation. Furthermore, different effects on the
330 wetland methane emissions from rising temperatures seems to be canceling each other out, at least on the large scale.

4.4 Increasing potential wetland area: “WetCrops” and “ConstVeg”

Both experiments increase the potential wetland area, but they do so in different ways:

- WetCrops: the drainage flag is disabled, so any cropland cell whose water table reaches the surface is diagnosed as a wetland (hydrologically). The vegetation remains the prescribed crop; no ecological conversion to wetland plants occurs.
- ConstVeg: the LUH2 land use map is frozen at its 1855 state, preventing the expansion of cropland (and the associated drainage) over time. Because the vegetation is fixed, the only driver of wetland change is the hydrological response to climate (precipitation, temperature, SWR). Thus, even with a static vegetation field, the diagnostic wetland fraction can grow or shrink, leading to corresponding changes in methane emissions.

340 In “ConstVeg” the mean JJA wetland area is 0.2 Mm² larger than in “Base” (Figure 4a). The increase is entirely due to hydrological inundation of cells that are, for example, forest or grassland in the 1855 land use map. Consequently, the global methane flux rises to 176.2 Tg yr⁻¹ (Table 3), even though the vegetation does not change.

The effect on wetland area of ~~switching off the drainage from crop~~ disabling the drainage flag on cropland areas and keeping the vegetation distribution constant at the level of the early industrialization is conceptually the same: The area on which
345 wetlands can ~~be created~~ exist is increased. In ~~ease that the drainage is switched off~~ the case of disabled drainage, all crop areas are made available for wetlands, and in the case of constant vegetation distribution, the ~~erops~~ crop areas are kept smaller, as discussed in ~~See~~ Section 3. This theoretical ~~behavior~~ behaviour is confirmed by ~~Fig.~~ Figure 4, where average “WetCrops” wetlands exceeds those of “Base” by ~~≈ 0.6 Mm²~~ ≈ 0.6 Mm² while “ConstVeg” ~~is adding ≈ 0.2 Mm²~~ adds ≈ 0.2 Mm² to the
“Base” wetland area. This increase is essentially equally distributed between NXT and tropics. These two experiments are
350 the only ones having a positive trend in the global wetland area throughout the experiments, somewhat lower in “WetCrops” than in “ConstVeg” and mostly from the tropical area. In the 1980–2014 period, the trends in “WetCrops” are much larger than

those of “ConstVeg” in all regions. In the NXT ~~regions~~region, the wetlands are even decreasing in “ConstVeg” since they do not profit from the ~~abandoning~~abandonment of croplands in Western Europe and ~~the USA~~North America. In general, the major difference to “Base” is that “WetCrops” and “ConstVeg” do not participate in the downward wetland area trend, where
355 crops are increasing over the experiment period.

The average methane emissions are also increased compared to “Base”. However, the emissions from “ConstVeg” ($176.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$) are closer to “WetCrops” ($181.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$) than to those of “Base” ($166.2 \text{ Tg}(\text{CH}_4) \text{ yr}^{-1}$). The explanation can be found in the methane emission trends, where “ConstVeg” is having the largest trend of all conducted experiments. The reason for the higher trend is that “WetCrops” only changes the wetland area, while no direct changes to the carbon cycle
360 ~~compared are made in comparison~~ to “Base”~~are done~~. “ConstVeg” changes the vegetation distribution in a way ~~leaving that~~leaves more space for the most productive vegetation type: ~~The~~the forests. Therefore, more carbon is available for methane production in “ConstVeg” compared to “WetCrops”.

5 Discussion

5.1 Impact of including surface water retention

~~The large~~The significant differences between the “Base” and “DrySurf” ~~highlights~~experiments highlight the sensitivity of the estimates of wetland ~~extend~~extent and thus wetland methane emissions to changes in the soil hydrology. Including SWR ~~regionally~~ dramatically increases the evapotranspiration and thus atmospheric water recycling in some regions. This also has major effects on the fluxes of latent and sensible heat and thus on regional temperature. These feedbacks emphasize the importance of also considering coupled (in this case land-atmosphere) simulations, since such additional water recycling (or
370 ~~lack of it~~thereof) is not accounted for in offline, land only, setups. ~~Thus~~As a result, offline sensitivity studies to changes in, e.g., soil hydrology characterizations may come to wrong conclusions: ~~In equivalent~~. In similar offline experiments (not shown), forced with the GSWP3 data set (Dirmeyer et al., 2006), the ~~major~~dominating effect of enabling SWR was to decrease the water infiltration into the soil, ~~which again led to a~~thus leading to slightly reduced wetland ~~area~~areas and thus lower wetland methane emissions. ~~Our assumption that the soil below water retention areas is as impermeable as clay likely is an exaggeration,~~
375 ~~so that the differences between~~, completely hiding the changes in moisture recycling dominating the coupled experiments.

However, since model estimates of wetland area do not differ by a very large amount between the “Base” and “DrySurf”
~~may be an extreme case.~~ experiments, and the increase in precipitation tends to occur in regions, where the difference in precipitation between model and observations (Dirmeyer et al., 2006) is relatively small, there is no clear conclusion, whether inclusion of the SWR parameterisation, as in the “Base” experiment, leads to an improvement in model results in terms of
380 wetland areas and wetland methane emissions.

Compared to the TOPMODEL wetlands, the SWR scheme is another way of allowing super-surface water to form, which can be interpreted as wetlands. Both SWR and TOPMODEL-style wetlands arise in areas with high water availability and since both approaches are based on statistical methods, it is impossible to completely separate the two ways of obtaining surface water bodies.

385 The correlation between the SWR and ~~wetland~~TOPMODEL-wetland areas in the entire “Base” experiment is ≈ 0.3 without trend. The assumption that soils below SWR areas are oxic may thus be too simplistic. Since SWR almost exclusively influences North America north-east of the Mackenzie River, north eastern Europe and north western Siberia (~~Fig.~~Figure S7), a (partial) lifting of this assumption would enhance the Arctic and Sub-Arctic wetland methane emissions compared to the tropical ones. By assuming that there is no overlap between the SWR and TOPMODEL wetland areas and that annual methane emissions are linearly dependent on the annual mean area with a wet surface, a rough estimate of the effect of the assumption, that the SWR areas do not contribute to methane production, can be obtained. In this case global wetland methane emissions would increase by $42 \text{ Tg}(\text{CH}_4)\text{yr}^{-1}$ or $\approx 25\%$. This includes an additional $27 \text{ Tg}(\text{CH}_4)\text{yr}^{-1}$ of emissions from the NXT area, increasing the NXT fraction of the global emissions from 21% to 30%. Since the assumptions for this estimate are rather unrealistic, these numbers are~~however,~~ however, regarded as the upper extreme of the consequences of excluding the SWR areas from the methane production.

5.2 AMIP setup versus offline land simulations

Using an AMIP setup introduces an atmospheric variability, which is different from the observed one. Therefore, comparisons of model results to observations and ~~observation-based~~observation-based model results are only possible in a statistical sense over longer periods. ~~Additionally~~In addition, in AMIP setups the atmospheric model may ~~in AMIP setups introduce less~~
400 ~~well-known~~introduce less well-known biases compared to observational data sets. For this study, the most important known atmospheric bias is an underestimation of the tropical precipitation by about 29% compared to GSWP3 (~~Tab.~~Table 2). The tropical bias is unevenly distributed between the continents, and is by far strongest over Indonesia and South East Asia, which explains our comparatively low wetland areas and thus methane emissions from this region. The bias distribution agrees with findings of Müller et al. (2025) and seems thus to be a general property of the ICON-XPP atmospheric model. Common
405 wetland products based on observations (GIEMS-2 (Prigent et al., 2020), WAD2M (Zhang et al., 2021)) find a significant seasonal cycle in the tropical wetland area more or less in phase with the northern boreal wetland areas. That in this study essentially no seasonal cycle is found can be attributed to the severe underestimation of south Asian and Indonesian rainfall and thus diminished wetlands in these regions. Therefore, the weak cycle in the Amazonian region can ~~out-weight~~out-weigh the out-of-phase cycles in Asia and Africa, producing a net-zero cycle for the tropics.

410 5.3 Anthropogenic land use changes including drainage of croplands

The use of land use maps compared to land use transitions (which are currently not implemented in JSBACH4) is known to underestimate the influence of anthropogenic land use changes on the carbon cycle (Wilkenskjeld et al., 2014; Stocker et al., 2014). Specifically the release of soil carbon is underestimated. Therefore, especially in the tropics, where shifting cultivation is a common agricultural practice, the carbon pools may be overestimated. This could also lead to an overestimate of the methane
415 emissions from these regions. Since farmers generally try to avoid wet areas (Valipour et al., 2020), we~~however,~~ however, assume the effect on the methane emissions to be small.

Assuming that all cropland areas in the model are drained is a strong simplification, especially due to the large-scale dynamics and development of agricultural practices. Former ~~crop-cropland~~ areas may be abandoned and become either pastures or completely be given back to ~~the~~-nature. This, of course, would not mean that drainage systems are removed, and thus the area drained is likely underestimated by the approach of this study. Also, managed forests, which in JSBACH are regarded as natural vegetation, pastures, and areas of urban expansion may be drained to enhance tree growth, making the land more suitable as grassland or to make the ground supportive for technospheric constructions (Fluet-Chouinard et al., 2023). Again these effects will lead to an underestimate of drained area. On the other hand, not all croplands are drained, either because they are dry enough by nature or because of limited local resources to establish adequate drainage systems. As an example, Valipour et al. (2020) states that in ~~present-day-present-day~~ India, only about 10% of croplands are drained. This will lead to an overestimate of ~~our-the~~ drained area, but likely not change the simulated wetlands much, since the extra drained area will ~~anyway-largely~~ be too dry to regularly become wetlands, ~~anyway~~. Though Valipour et al. (2020) reports a history of several thousand years of anthropogenic drainage, Fluet-Chouinard et al. (2023) argue that ~~the~~ main wetland area loss due to drainage takes place after 1900 and is still ongoing at a more or less constant rate. Thus, ~~our~~ assumption of drained crops may be most representative for the ~~latter-later~~ part of the simulation period. With the exception ~~of~~ the NXT region in the “ConstVeg” experiment, wetland areas are increasing ~~since-after~~ 1980 in all ~~of~~ our experiments for both NXT and ~~the~~ tropics. One reason may be different developments in the precipitation and precipitation patterns. GSWP3 and the ICON-XPP atmosphere agree that in the 1980–2014 period, the global (land) precipitation is on average enhanced ($\approx 0.48, 0.43$ and $0.32\text{mm}\cdot\text{yr}^{-2}$ ~~0.32mm yr⁻²~~ for GSWP3, “Base” and “DrySurf”, respectively). However, the global patterns of increasing precipitation in ICON-XPP correlates better with our simulated wetlands (somewhat better in “~~BestGuessBase~~” than in “DrySurf”), which may thus be enlarged. Nevertheless, it is unlikely that this better pattern correlation alone can explain the trends in wetland area going in different directions in our simulations and observations. It is more likely that we, despite the drainage assumption, largely underestimate the influence of human water management (dams, reservoirs, ~~drainage—also-of-non-croplandand drainage, also of non-croplands~~). In total we expect ~~that~~ our simulated drained area to be underestimated (and the wetland areas therefore overestimated) in regions with intense and long-lasting anthropogenic influence.

5.4 Wetland area estimates

~~Our simulated wetland is approximately in line with those presented in Prigent et al. (2007) when accounting for their data set including also anthropogenic wetlands, mainly rice fields (Kleinen et al., 2020). The estimates from Zhang et al. (2021); Prigent et al. (2020) are however much larger. The new version 2.0 of the WAD2M wetland data set (Zhang et al., 2021) reduced their wetland extent estimate by 0.5~~ Data-based estimates of global methane emitting areas are scarce, though a few map-based estimates of wetland extent and remote-sensing estimates of surface inundation are available. Lehner and Döll (2004) estimated the global annual maximum extent of lakes and wetlands based on maps, and show a global extent of methane-relevant wetlands of between 4.5 and 7 Mm² compared to the previous version (Saunois et al., 2024). One reason for our smaller wetland area might be, that the available observational data sets still do not distinguish small lakes from wetlands, whereas the TOPMODEL approach only counts, depending on which categories of wetlands are considered relevant for methane. Prigent et al. (2012)

developed a remote-sensing-based data set of surface inundation, estimating an annual mean extent of 2.5 Mm², after removal of rice-growing areas based on Monfreda et al. (2008), and a JJA extent of 4.1 Mm². This estimate includes the areas of lakes, reservoirs, and rivers, though, which are not considered methane-emitting areas in our model. Excluding those areas, Zhang et al. (2021) estimate a wetland extent of 4.4 Mm² in the annual mean, with a JJA extent of 5.2 Mm². Finally, Bernard et al. (2025) estimate an annual mean extent of 3.5 Mm² for inundated and saturated wetlands and peatlands, with an annual maximum extent of 7.7 Mm².

Our model estimate yields between 5.3 Mm² in the “DrySurf” experiment and 6 Mm² in “WetCrops”, both for the JJA extent and the 1980–2014 period (Figure 4), thus determining wetland extents generally in line with data-based estimates. However, a large uncertainty in the estimates of wetlands, and especially methane-emitting areas, has to be acknowledged. The TOPMODEL-based approach we are using determines the extent of (super-) saturated soils as wetland. Another reason is, and considers those as methane-emitting. This may, however, underestimate the extent, as some categories, for example peatlands, cannot be considered well in this approach.

Furthermore, the ICON-XPP model shows a precipitation bias over the tropics—specifically, specifically in South East Asia, likely leading to too low an estimate of wetland extent for this region in our experiments. Also, JSBACH is not equipped with a detailed river flow model capable of simulating temporary flooding of river flood plains as a consequence of heavy precipitation or snow melt events. Since such wetlands tend to be short-term phenomena, it is questionable if, however, whether the soils have time to become anaerobic and thus they would likely not contribute much significantly to the wetland methane emissions.

Excluding short-term wetlands from short-term wetlands in dry areas by applying the dynamic application of the criterion in Eq. criterion in Equation (1), instead of a static mask, allows the model to catch capture the influence of major climatic shifts. On This exclusion, the other hand it may, might also cause spurious trends in wetland area and thus in wetland methane emissions by changes in the wetland allowance changing the wetland allowance mask, either temporary temporarily or as part of a long-term long-term trend. By selecting the required ratio as low as 0.3 (Eq. (1)) Equation 1), the bounds of where wetlands are allowed are however permitted are located far away from where major wetlands are predicted. Thus, the wetland methane emissions and their trends are likely not influenced by the details of this masking.

5.5 Wetland methane emissions

In the “ConstCO2” experiment the trend in wetland methane emissions is much smaller than in “BestGuess”. Over the entire experiment period it is even negative, following the downward trend in wetland area. This suggest suggests that the main driver of the trend found in all other experiments is an accelerated carbon cycle stemming from CO₂ fertilization fertilisation. This contradicts the findings of both McNorton et al. (2016) and the WETCHIMP model ensemble (Melton et al., 2013), who attributes attribute their trend to temperature driven a temperature-driven increase of wetland area and microbial methane production rates.

For the period 2000–2012 the trend in global wetland methane emissions is about 0.4 Tg(CH₄) yr⁻², of which 75% stems stem from the tropics. This is higher than the numbers presented in Saunois et al. (2017) who reports, who report a multi-

485 model mean of $0.2 \text{ Tg}(\text{CH}_4)\text{yr}^{-2}$, and about 10% of a trend of $2.2 \text{ Tg}(\text{CH}_4)\text{yr}^{-2}$ from top-down inversions attributed to non-anthropogenic sources. Since the productivity in JSBACH is known to be comparatively sensitive to changes in ~~CO_2 atmospheric concentration~~ the atmospheric concentration of CO_2 , which is the driver of our trend in methane emissions, it ~~seems logical~~ appears likely that our predicted wetland emission trend is at the high end.

The basic TOPMODEL assumption defining wetlands as areas with super-saturated soils (i.e., the water level is at least reaching the surface) and using the wetland fraction as a measure of the soil oxygen content are simplifications, which can only serve as a first guess. For instance soils, where the root zone of macrophytes is saturated, may be anoxic. Thus, though the soil column as a whole is not saturated, such areas could be a source of methane, mainly since produced methane may bypass the upper (oxic) soil layers through aerenchyma. On the other hand, soils ~~takes~~ take time to become anoxic after being inundated. Depending on the state of the soil at inundation, this may take hours to weeks ~~Patel et al. (2024)~~ (Patel et al., 2024), and the anaerobic area could thus potentially be overestimated by assuming all wetland soils to be anoxic. ~~Also~~ Further, wetland areas may have very different and highly heterogeneous methane production and emission rates, as shown for Arctic ponds by Rehder et al. (2023).

~~The~~ Of course, the tropical precipitation bias reducing our tropical wetland area ~~of course~~ tends to decrease the importance of the tropical wetland methane emissions. There are ~~however~~, however, also effects which ~~draws~~ draw in the other direction: In this study, we accounted only for the conversion of contemporary organic matter to methane, while in reality an additional source may be added from the huge amount of ancient organic carbon stored in the thawing permafrost (Hugelius et al., 2014), as well as from degrading peatlands (Hugelius et al., 2020), since we include no explicit peatland model such as ~~HIMMELI~~ HIMMELI (Raivonen et al., 2017). Ekici et al. (2019) found that incorporating micro-scale ground subsidence following permafrost thaw will increase the wetland fraction and thus methane emissions. Both these effects would enhance the importance of the Arctic regions in the global methane budget. Even though our wetlands are in general smaller, the modeled NXT fraction agrees with the more detailed wetland methane model of Nzotungicimpaye et al. (2021).

5.6 Offline versus online simulations

Offline (stand-alone) land-surface simulations are valuable for isolating the response of wetland hydrology to prescribed climate, but they cannot capture feedbacks such as (i) moisture recycling that modifies precipitation patterns, and (ii) vegetation-mediated changes in surface energy fluxes. Conversely, online coupled simulations may inherit atmospheric biases (e.g., the tropical precipitation deficit in ICON-XPP) that propagate into the wetland response. Therefore, both experimental designs are complementary: offline experiments help to diagnose process sensitivities, while online experiments are required to assess the full climate-wetland feedback loop.

6 Conclusions

515 Sub-models for ~~interactive~~ coupled wetland dynamics and wetland methane emissions have been ~~build~~ built into the ICON-XPP Earth System Model. The resulting wetland distributions, when accounting for precipitation biases from the atmospheric

model, align very well with observations. Additionally, the estimates of wetland methane emissions agree well with published values. Sensitivity studies were done to quantify effects of certain model assumptions and anthropogenic influences on methane emissions from natural wetlands.

520 Agricultural development encompassing both the changes in farmed area and cultivation practices, is the dominant anthropogenic factor controlling the natural wetland methane emissions by controlling the potential wetland area. Until around 1980~~this result~~, this resulted in a global decrease in wetland area, counteracting the general trend of increasing wetland methane emissions. In contrast to the simulations ~~here presented~~presented here, observations suggest that global wetland area is still decreasing as a consequence of human actions, indicating that anthropogenic influences on the global soil hydrology are

525 underestimated in the current generation of Earth System Models.

The ~~second-most significant humanly-influenced~~second-most significant human-influenced factor affecting wetland area and their methane emissions is the increase in atmospheric CO₂ concentrations. This rise accelerates the carbon cycle through CO₂ ~~fertiliazation~~fertilisation. This, however, does not significantly impact the wetland area. The temperature effect of rising atmospheric CO₂ concentration does not substantially affect wetland area and methane emissions.

530 ~~Effects of changing how the~~ Changes in the model description of the surface hydrology~~is modelled~~, including the implementation of surface water retention, ~~has~~have the potential to significantly alter land-atmosphere water and energy fluxes on the regional scale and thus ~~needs~~need to be evaluated using coupled setups.

Code and data availability. Code, scripts and data used for this manuscript as available in the Edmond repository, Wilkenskield (2025).

Author contributions. S. Wilkenskield ported the wetland and methane models to the ICON-ESM and developed the model extensions,

535 designed and conducted the experiments as well as the output analysis and did the main work on writing this paper. T. Kleinen implemented the original wetland and methane models in MPI-ESM and provided much insights into wetland and methane cycling in the Earth System. He also revised the paper after the initial reviews. T. Stacke implemented the SWR scheme and contributed his knowledge on soil hydrology and its interplay with the atmosphere. V. Brovkin contributed the idea of the project and gave it directions during the process. All authors contributed to the writing process of the paper.

540 *Competing interests.* The authors declare that they have no competing interests.

Acknowledgements. This study was funded by the EU-Horizon 2020 projects ESM2025, Grant-no: 101003536 and Q-Arctic (ERC-grant-no: 951288) and used resources of the Deutsches Klimarechenzentrum (DKRZ) granted by its Scientific Steering Committee (WLA) under project ID bm1255.

545 We thank Goran Georgievski for comments on an earlier version of this manuscript. Furthermore, we thank two anonymous reviewers for
their valuable suggestions.

References

- Andresen, C. G., Lawrence, D. M., Wilson, C. J., McGuire, A. D., Koven, C., Schaefer, K., Jafarov, E., Peng, S., Chen, X., Gouttevin, I., Burke, E., Chadburn, S., Ji, D., Chen, G., Hayes, D., and Zhang, W.: Soil moisture and hydrology projections of the permafrost region – a model intercomparison, *The Cryosphere*, 14, 445–459, <https://doi.org/10.5194/tc-14-445-2020>, 2020.
- 550 Bernard, J., Prigent, C., Jimenez, C., Fluet-Chouinard, E., Lehner, B., Salmon, E., Ciais, P., Zhang, Z., Peng, S., and Saunois, M.: The GIEMS-MethaneCentric database: a dynamic and comprehensive global product of methane-emitting aquatic areas, *Earth System Science Data*, 17, 2985–3008, <https://doi.org/10.5194/essd-17-2985-2025>, 2025.
- Beven, K. J. and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology, *Hydrological Sciences Bulletin*, 24, 43–69, <https://doi.org/10.1080/02626667909491834>, 1979.
- 555 Bridgman, S. D., Cadillo-Quiroz, H., Keller, J. K., and Zhuang, Q.: Methane emissions from wetlands: biogeochemical, microbial, and modeling perspectives from local to global scales, *Global Change Biology*, 19, 1325–1346, <https://doi.org/https://doi.org/10.1111/gcb.12131>, 2013.
- Chen, Y., Li, H., Song, S., Zhou, Z., Chen, C., Guo, C., and Zheng, F.: Fast Expansion of Surface Water Extent in Coastal Chinese Mainland from the 1980s to 2020 Based on Remote Sensing Monitoring, *WATER*, 17, <https://doi.org/10.3390/w17020194>, 2025.
- 560 Conrad, R.: Contribution of hydrogen to methane production and control of hydrogen concentrations in methanogenic soils and sediments, *FEMS Microbiology Ecology*, 28, 193–202, <https://doi.org/10.1111/j.1574-6941.1999.tb00575.x>, 1999.
- Davidson, N. C.: How much wetland has the world lost? Long-term and recent trends in global wetland area, *Marine and Freshwater Research*, 65, 934 – 941, <https://doi.org/10.1071/MF14173>, 2014.
- de Vrese, P., Stacke, T., and Hagemann, S.: Exploring the biogeophysical limits of global food production under different climate change scenarios, *Earth System Dynamics*, 9, 393–412, <https://doi.org/10.5194/esd-9-393-2018>, 2018.
- 565 de Vrese, P., Stacke, T., Kleinen, T., and Brovkin, V.: Diverging responses of high-latitude CO_2 and CH_4 emissions in idealized climate change scenarios, *The Cryosphere*, 15, 1097–1130, <https://doi.org/10.5194/tc-15-1097-2021>, 2021.
- de Vrese, P., Georgievski, G., Gonzalez Rouco, J. F., Notz, D., Stacke, T., Steinert, N. J., Wilkenskjeld, S., and Brovkin, V.: Representation of soil hydrology in permafrost regions may explain large part of inter-model spread in simulated Arctic and subarctic climate, *The Cryosphere*, 17, 2095–2118, <https://doi.org/10.5194/tc-17-2095-2023>, 2023.
- 570 de Vrese, P., Stacke, T., Gayler, V., and Brovkin, V.: Permafrost Cloud Feedback May Amplify Climate Change, *Geophysical Research Letters*, 51, e2024GL109034, <https://doi.org/https://doi.org/10.1029/2024GL109034>, e2024GL109034 2024GL109034, 2024.
- Dirmeyer, P. A., Gao, X., Zhao, M., Guo, Z., Oki, T., and Hanasaki, N.: GSWP-2 - Multimodel analysis and implications for our perception of the land surface, *BULLETIN OF THE AMERICAN METEOROLOGICAL SOCIETY*, 87, 1381+, [https://doi.org/10.1175/BAMS-87-](https://doi.org/10.1175/BAMS-87-10-1381)
- 575 10-1381, 2006.
- Ekici, A., Beer, C., Hagemann, S., Boike, J., Langer, M., and Hauck, C.: Simulating high-latitude permafrost regions by the JSBACH terrestrial ecosystem model, *Geoscientific Model Development*, 7, 631–647, <https://doi.org/10.5194/gmd-7-631-2014>, 2014.
- Ekici, A., Lee, H., Lawrence, D. M., Swenson, S. C., and Prigent, C.: Ground subsidence effects on simulating dynamic high-latitude surface inundation under permafrost thaw using CLM5, *Geoscientific Model Development*, 12, 5291–5300, [https://doi.org/10.5194/gmd-12-5291-](https://doi.org/10.5194/gmd-12-5291-2019)
- 580 2019, 2019.
- Fluet-Chouinard, E., Stocker, B. D., Zhang, Z., Malhotra, A., Melton, J. R., Poulter, B., Kaplan, J. O., Goldewijk, K. K., Siebert, S., Minayeva, T., Hugelius, G., Joosten, H., Barthelmes, A., Prigent, C., Aires, F., Hoyt, A. M., Davidson, N., Finlayson, C. M., Lehner, B., Jackson, R. B.,

- and McIntyre, P. B.: Extensive global wetland loss over the past three centuries, *Nature*, 614, 281–286, <https://doi.org/10.1038/s41586-022-05572-6>, 2023.
- 585 Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., p. 923–1054, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021.
- 590 Gardner, R. C. and Finlayson, C.: Global Wetland Outlook: State of the World's Wetlands and Their Services to People, Tech. Rep. 2020-5, Stetson University College of Law Research, <https://ssrn.com/abstract=3261606>, accessed: 2025-04-09, 2018.
- Goll, D. S., Brovkin, V., Liski, J., Raddatz, T., Thum, T., and Todd-Brown, K. E. O.: Strong dependence of CO₂ emissions from anthropogenic land cover change on initial land cover and soil carbon parametrization, *GLOBAL BIOGEOCHEMICAL CYCLES*, 29, 1511–1523, <https://doi.org/10.1002/2014GB004988>, 2015.
- 595 Hagemann, S. and Gates, L. D.: Improving a subgrid runoff parameterization scheme for climate models by the use of high resolution data derived from satell, *Climate Dynamics*, 21, 349–359, <https://doi.org/10.1007/s00382-003-0349-x>, 2003.
- Hagemann, S. and Stacke, T.: Impact of the soil hydrology scheme on simulated soil moisture memory, *Climate Dynamics*, 44, 1731–1750, <https://doi.org/10.1007/s00382-014-2221-6>, 2015.
- 600 Hagemann, S., Blome, T., Ekici, A., and Beer, C.: Soil-frost-enabled soil-moisture-precipitation feedback over northern high latitudes, *EARTH SYSTEM DYNAMICS*, 7, 611–625, <https://doi.org/10.5194/esd-7-611-2016>, 2016.
- Hardouin, L., Decharme, B., Colin, J., and Delire, C.: Climate-Driven Projections of Future Global Wetlands Extent, *Earth's Future*, 12, e2024EF004553, <https://doi.org/https://doi.org/10.1029/2024EF004553>, e2024EF004553 2024EF004553, 2024.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C. L., Schirrmeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *BIOGEOSCIENCES*, 11, 6573–6593, <https://doi.org/10.5194/bg-11-6573-2014>, 2014.
- 605 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw, *Proceedings of the National Academy of Sciences*, 117, 20438–20446, <https://doi.org/10.1073/pnas.1916387117>, 2020.
- 610 Hurtt, G. C., Chini, L., Sahajpal, R., Frolking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J. O., Kennedy, J., Krisztin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.: Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6, *Geoscientific Model Development*, 13, 5425–5464, <https://doi.org/10.5194/gmd-13-5425-2020>, 2020.
- Kleinen, T., Mikolajewicz, U., and Brovkin, V.: Terrestrial methane emissions from the Last Glacial Maximum to the preindustrial period, *CLIMATE OF THE PAST*, 16, 575–595, <https://doi.org/10.5194/cp-16-575-2020>, 2020.
- Kleinen, T., Gromov, S., Steil, B., and Brovkin, V.: Atmospheric methane underestimated in future climate projections (vol 16, 094006, 2021), *ENVIRONMENTAL RESEARCH LETTERS*, 16, <https://doi.org/10.1088/1748-9326/ac2f66>, 2021.

- 620 Krysanova, V., Zaherpour, J., Didovets, I., Gosling, S. N., Gerten, D., Hanasaki, N., Müller Schmied, H., Pokhrel, Y., Satoh, Y., Tang, Q., and Wada, Y.: How evaluation of global hydrological models can help to improve credibility of river discharge projections under climate ch, *Climatic Change*, 163, 1353–1377, <https://doi.org/10.1007/s10584-020-02840-0>, 2020.
- Lan, X., K.W., T., and Dlugokencky, E.: Trends in globally-averaged CH₄, N₂O, and SF₆ determined from NOAA Global Monitoring Laboratory measurements. Version 2025-01, <https://doi.org/10.15138/P8XG-AA10>, 2025.
- 625 Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *Journal of Hydrology*, 296, 1–22, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2004.03.028>, 2004.
- Marthews, T. R., Dadson, S. J., Lehner, B., Abele, S., and Gedney, N.: High-resolution global topographic index values for use in large-scale hydrological modelling, *Hydrology and Earth System Sciences*, 19, 91–104, <https://doi.org/10.5194/hess-19-91-2015>, 2015.
- McNorton, J., Gloor, E., Wilson, C., Hayman, G. D., Gedney, N., Comyn-Platt, E., Marthews, T., Parker, R. J., Boesch, H., and Chipperfield, M. P.: Role of regional wetland emissions in atmospheric methane variability, *GEOPHYSICAL RESEARCH LETTERS*, 43, 11 433–11 444, <https://doi.org/10.1002/2016GL070649>, 2016.
- 630 Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, <https://doi.org/10.5194/bg-10-753-2013>, 2013.
- 635 Monfreda, C., Ramankutty, N., and Foley, J. A.: Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Global Biogeochemical Cycles*, 22, GB1022, <https://doi.org/10.1029/2007GB002947>, 2008.
- Müller, W. A., Früh, B., Korn, P., Potthast, R., Baehr, J., Bettems, J.-M., Bölöni, G., Brienens, S., Fröhlich, K., Helmert, J., Jungclaus, J., Köhler, M., Lorenz, S., Schneidereit, A., Schnur, R., Schulz, J.-P., Schlemmer, L., Sgoff, C., Pham, T. V., Pohlmann, H., Vogel, B., Vogel, H., Wirth, R., Zaehle, S., Zängl, G., Stevens, B., and Marotzke, J.: ICON: Towards vertically integrated model configurations for numerical weather prediction, climate predictions and projections, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-24-0042.1>, 2025.
- 640 Nzotungicimpaye, C.-M., Zickfeld, K., MacDougall, A. H., Melton, J. R., Treat, C. C., Eby, M., and Lesack, L. F. W.: WETMETH 1.0: a new wetland methane model for implementation in Earth system models, *Geoscientific Model Development*, 14, 6215–6240, <https://doi.org/10.5194/gmd-14-6215-2021>, 2021.
- Patel, K. F., Rod, K. A., Zheng, J., Regier, P., Machado-Silva, F., Bond-Lamberty, B., Chen, X., Day, D. J., Doro, K. O., Kaufman, M. H., Kovach, M., McDowell, N., McKever, S. A., Megonigal, J. P., Norris, C. G., O'Meara, T., Peixoto, R. B., Rich, R., Thornton, P., Kemner, K. M., Ward, N. D., Weintraub, M. N., and Bailey, V. L.: Time to anoxia: Observations and predictions of oxygen drawdown following coastal flood events, *Geoderma*, 444, 116 854, <https://doi.org/https://doi.org/10.1016/j.geoderma.2024.116854>, 2024.
- 650 Prentice, I. C., Sykes, M. T., and Cramer, W.: A Simulation-Model for the Transient Effects of Climate Change on Forest Landscapes, *ECOLOGICAL MODELLING*, 65, 51–70, 1993.
- Prigent, C., Papa, F., Aires, F., Rossow, W. B., and Matthews, E.: Global inundation dynamics inferred from multiple satellite observations, 1993–2000, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2006JD007847>, 2007.
- 655 Prigent, C., Papa, F., Aires, F., Jimenez, C., Rossow, W. B., and Matthews, E.: Changes in land surface water dynamics since the 1990s and relation to population pressure, *Geophysical Research Letters*, 39, <https://doi.org/https://doi.org/10.1029/2012GL051276>, 2012.

- Prigent, C., Jimenez, C., and Bousquet, P.: Satellite-Derived Global Surface Water Extent and Dynamics Over the Last 25 Years (GIEMS-2), *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030711, <https://doi.org/https://doi.org/10.1029/2019JD030711>, 2020.
- Raivonen, M., Smolander, S., Backman, L., Susiluoto, J., Aalto, T., Markkanen, T., Mäkelä, J., Rinne, J., Peltola, O., Aurela, M., Lohila, A., Tomasic, M., Li, X., Larmola, T., Juutinen, S., Tuittila, E.-S., Heimann, M., Sevanto, S., Kleinen, T., Brovkin, V., and Vesala, T.: HIMMELI v1.0: Helsinki Model of Methane build-up and emission for peatlands, *Geoscientific Model Development*, 10, 4665–4691, <https://doi.org/10.5194/gmd-10-4665-2017>, 2017.
- Rehder, Z., Kleinen, T., Kutzbach, L., Stepanenko, V., Langer, M., and Brovkin, V.: Simulated methane emissions from Arctic ponds are highly sensitive to warming, *Biogeosciences*, 20, 2837–2855, <https://doi.org/10.5194/bg-20-2837-2023>, 2023.
- Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., and Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, *Biogeosciences*, 8, 1925–1953, <https://doi.org/10.5194/bg-8-1925-2011>, 2011.
- Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J., Bergamaschi, P., Blake, D. R., Brailsford, G., Brovkin, V., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Curry, C., Frankenberg, C., Gedney, N., Höglund-Isaksson, L., Ishizawa, M., Ito, A., Joos, F., Kim, H.-S., Kleinen, T., Krummel, P., Lamarque, J.-F., Langenfelds, R., Locatelli, R., Machida, T., Maksyutov, S., McDonald, K. C., Marshall, J., Melton, J. R., Morino, I., Naik, V., O'Doherty, S., Parmentier, F.-J. W., Patra, P. K., Peng, C., Peng, S., Peters, G. P., Pison, I., Prigent, C., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J., Spahni, R., Steele, P., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., van Weele, M., van der Werf, G. R., Weiss, R., Wiedinmyer, C., Wilton, D. J., Wiltshire, A., Worthy, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z., and Zhu, Q.: The global methane budget 2000–2012, *Earth System Science Data*, 8, 697–751, <https://doi.org/10.5194/essd-8-697-2016>, 2016.
- Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Frankenberg, C., Gedney, N., Höglund-Isaksson, L., Ishizawa, M., Ito, A., Joos, F., Kim, H.-S., Kleinen, T., Krummel, P., Lamarque, J.-F., Langenfelds, R., Locatelli, R., Machida, T., Maksyutov, S., Melton, J. R., Morino, I., Naik, V., O'Doherty, S., Parmentier, F.-J. W., Patra, P. K., Peng, C., Peng, S., Peters, G. P., Pison, I., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J., Spahni, R., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., Weiss, R., Wilton, D. J., Wiltshire, A., Worthy, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z., and Zhu, Q.: Variability and quasi-decadal changes in the methane budget over the period 2000–2012, *Atmospheric Chemistry and Physics*, 17, 11 135–11 161, <https://doi.org/10.5194/acp-17-11135-2017>, 2017.
- Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguía-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang,

- 695 Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgersson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K.,
700 Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguía-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., van der Werf, G., Worthy, D. E., Xi, Y., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000–2020, *Earth System Science Data Discussions*, 2024, 1–147, <https://doi.org/10.5194/essd-2024-115>, 2024.
- 705 Schneck, R., Gayler, V., Nabel, J. E. M. S., Raddatz, T., Reick, C. H., and Schnur, R.: Assessment of JSBACHv4.30 as a land component of ICON-ESM-V1 in comparison to its predecessor JSBACHv3.2 of MPI-ESM1.2, *Geoscientific Model Development*, 15, 8581–8611, <https://doi.org/10.5194/gmd-15-8581-2022>, 2022.
- Stacke, T. and Hagemann, S.: Development and evaluation of a global dynamical wetlands extent scheme, *Hydrology and Earth System Sciences*, 16, 2915–2933, <https://doi.org/10.5194/hess-16-2915-2012>, 2012.
- 710 Stocker, B., Feissli, F., Strassmann, K., Spahni, R., and Joos, F.: Past and future carbon fluxes from land use change, shifting cultivation and wood harvest, *Tellus B*, 66, <http://www.tellusb.net/index.php/tellusb/article/view/23188>, 2014.
- Taylor, K., Williamson, D., and Zwiers, F.: The sea surface temperature and sea ice concentration boundary conditions for AMIP II simulations, Tech. Rep. 60, Program for Climate Model Diagnosis and Intercomparison (PCMDI), Lawrence Livermore National Laboratory, <https://pcmdi.llnl.gov/report/pdf/60.pdf>, 2000.
- 715 Tuomi, M., Thum, T., Jarvinen, H., Fronzek, S., Berg, B., Harmon, M., Trofymow, J. A., Sevanto, S., and Liski, J.: Leaf litter decomposition-Estimates of global variability based on Yasso07 model, *ECOLOGICAL MODELLING*, 220, 3362–3371, <https://doi.org/10.1016/j.ecolmodel.2009.05.016>, 2009.
- UNEP, U. N. E. P.: Global Methane Assessment: Benefits and Costs of Mitigating Methane Emissions - Summary for Decision Makers, <https://wedocs.unep.org/20.500.11822/35917>, 2021.
- 720 Valipour, M., Krasilnikof, J., Yannopoulos, S., Kumar, R., Deng, J., Rocco, P., Mays, L., Grismer, M. E., and Angelakis, A. N.: The Evolution of Agricultural Drainage from the Earliest Times to the Present, *SUSTAINABILITY*, 12, <https://doi.org/10.3390/su12010416>, 2020.
- Veldkamp, T. I. E., Zhao, F., Ward, P. J., de Moel, H., Aerts, J. C. J. H., Schmied, H. M., Portmann, F. T., Masaki, Y., Pokhrel, Y., Liu, X., Satoh, Y., Gerten, D., Gosling, S. N., Zaherpour, J., and Wada, Y.: Human impact parameterizations in global hydrological models
725 improve estimates of monthly discharges and hydrological extremes: a multi-model validation study, *ENVIRONMENTAL RESEARCH LETTERS*, 13, <https://doi.org/10.1088/1748-9326/aab96f>, 2018.
- Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHyMe v1.3.1, *GEOSCIENTIFIC MODEL DEVELOPMENT*, 3, 565–584, <https://doi.org/10.5194/gmd-3-565-2010>, 2010.
- Wilkenskjeld, S.: Data and scripts for “Natural wetland methane emissions simulated by ICON-XPP”, Edmond, draft version,
730 <https://doi.org/10.17617/3.YADS3P>, 2025.

- Wilkenskjeld, S., Kloster, S., Pongratz, J., Raddatz, T., and Reick, C. H.: Comparing the influence of net and gross anthropogenic land-use and land-cover changes on the carbon cycle in the MPI-ESM, *Biogeosciences*, 11, 4817–4828, <https://doi.org/10.5194/bg-11-4817-2014>, 2014.
- 735 Xu, X., Yuan, F., Hanson, P. J., Wullschleger, S. D., Thornton, P. E., Riley, W. J., Song, X., Graham, D. E., Song, C., and Tian, H.: Reviews and syntheses: Four decades of modeling methane cycling in terrestrial ecosystems, *Biogeosciences*, 13, 3735–3755, <https://doi.org/10.5194/bg-13-3735-2016>, 2016.
- Xu, Y., Feng, L., Fang, H., Song, X.-P., Gieseke, F., Kariryaa, A., Oehmcke, S., Gibson, L., Jiang, X., Lin, R., Woolway, R. I., Zheng, C., Brandt, M., and Fensholt, R.: Global mapping of human-transformed dike-pond systems, *Remote Sensing of Environment*, 313, 114 354, <https://doi.org/https://doi.org/10.1016/j.rse.2024.114354>, 2024.
- 740 Zhang, B., Tian, H., Lu, C., Chen, G., Pan, S., Anderson, C., and Poulter, B.: Methane emissions from global wetlands: An assessment of the uncertainty associated with various wetland extent data sets, *ATMOSPHERIC ENVIRONMENT*, 165, 310–321, <https://doi.org/10.1016/j.atmosenv.2017.07.001>, 2017.
- Zhang, Z., Fluet-Chouinard, E., Jensen, K., McDonald, K., Hugelius, G., Gumbrecht, T., Carroll, M., Prigent, C., Bartsch, A., and Poulter, B.: Development of the global dataset of Wetland Area and Dynamics for Methane Modeling (WAD2M), *Earth System Science Data*, 13, 2001–2023, <https://doi.org/10.5194/essd-13-2001-2021>, 2021.
- 745




Figure 1. “Base” average distribution of (a) wetland area (% of grid cell area) in northern hemisphere summer (JJA) and (b) 1855-2014 wetland area trend of yearly-annual means (% of grid cell per century). Black dots mark cells with trends significant at the 95% level according to Matlab’s fitlm-fitlm function.




Figure 2. Wetland area yearly-annual cycle for the “Base” experiment. Solid lines only include cells with less than 10% snow whereas dashed lines includes all cells. Blue lines are global values, reddish include only the NXT area (north of 30°N) and the gray line is the “tropics” (everything south of 30°N). For the tropical area, snow masking does not make any difference.




Figure 3. Global distribution of (a) wetland methane emissions in the “Base” experiment, 1855-2014 average and (b) trend of wetland methane emissions in the “Base” experiment. Black dots mark cells with trends significant at the 95% level according to Matlab’s fitlm function.




Figure 4. Top panels: NH summer (JJA) Wetland area (in $[Mm^2]$) for areas with $\leq 10\%$ snow (a) and annual wetland methane emissions (b, $[Tg(CH_4) yr^{-1}]$). Bottom panels: 100 year trend ($[\% \text{ of } 1855\text{--}2014 \text{ average}]$) of (c) annual wetland area and (d) annual wetland methane emissions. Dark (colors exactly matching the legend) partly covered bars represent the entire experiment period (1855–2014), while brighter overlaid bars represent the 1980–2014 period. Colors indicate the different experiments.




Figure 5. Mean difference in wetland area (a) and annual precipitation (b) between “DrySurf” and “Base”. Black dots mark cells significantly different at the 95% level between the experiments according to Matlab’s ttest function.

Table 1. List of conducted experiments. The “Drainage” column indicates whether the diagnostic wetland fraction is suppressed on cropland cells (Yes = drainage active, No = drainage disabled). The CO₂ column only applies to the CO₂ ~~seen-by~~ applied to the terrestrial carbon cycle model. ~~For~~, while for the radiation calculations, transient CO₂ concentration is always applied. Soil-hydrology and carbon-cycle processes are identical in all experiments.

Name	Type-Ponds	Drainage	Vegetation	CO ₂ (terrestrial)
Base	AMIP -Yes	Yes	Transient	Transient
DrySurf	AMIP -No	Yes	Transient	Transient
ConstCO2	AMIP -Yes	Yes	Transient	1855
WetCrops	AMIP -Yes	No	Transient	Transient
ConstVeg	AMIP -Yes	Yes	1855	Transient

Table 2. GSWP3 and model precipitation [mm yr⁻¹]. Annual means from year 1901 to 2014. “Tropical”: latitudes between 30°S and 30°N, “SA”: Tropical South America (latitudes 30°S to 12°N), “AF”: Africa south of 30°N, “EA”: East Asia (latitudes 15°S to 20°N, east of 90°E).

Dataset	Global	Tropical	NXT	SA	AF	EA
GSWP3	776.0	1130.4	553.1	1743.6	768.9	2376.3
Base	651.4	809.6	588.5	1316.9	633.6	964.4
DrySurf	640.2	808.0	563.9	1316.8	634.1	944.7

Table 3. Global and northern extratropics (NXT) wetland methane emissions for the entire experiments, the period 2000-2012 (global only, compare to Saunois et al. (2016)) and wetland methane emissions trends for the entire experiments and the last 30 years of the experiments. Trends are given in Tg(CH₄) per year per century.

Experiment	Global mean		NXT mean	Global trend		NXT trend	
	1855-2014	2000-2012	1855–2014	1855-2014	1980-2014	1855-2014	1980-2014
	[Tg(CH ₄) yr ⁻¹]			[Tg(CH ₄) yr ⁻² x 100]			
Base	166.2	182.7	35.2	12.8	55.6	0.2	15.1
DrySurf	164.8	179.9	32.9	11.0	57.0	-1.1	14.3
ConstCO2	156.3	154.6	34.7	-4.6	6.9	-0.9	9.9
WetCrops	181.2	204.5	42.6	21.2	70.7	2.1	15.5
ConstVeg	176.2	205.5	39.5	28.4	79.8	4.6	7.1