

GHGPSE-Net: A method towards spaceborne automated extraction of greenhouse-gas point sources using point-object-detection deep neural network

Yiguo Pang^{1,2}, Denghui Hu¹, Longfei Tian¹, Shuang Gao¹, and Guohua Liu^{1,2}

¹Innovation Academy for Microsatellites of Chinese Academy of Sciences, Shanghai, China

²University of Chinese Academy of Sciences, Beijing, China

Correspondence: Guohua Liu (liugh@microsate.com)

Abstract. Point sources account for a large portion of anthropogenic greenhouse gas (GHG) emissions. Timely detection, localization, and quantification of these emissions are critical for supporting carbon neutrality efforts. Spaceborne monitoring satellites can provide essential concentration data for identifying point sources. However, existing methods often require human intervention and typically detect plume masks instead of source locations, limiting their utility for regulatory applications.

5 In this study, we present GHGPSE-Net, a deep learning method for greenhouse gas point source extraction. GHGPSE-Net simultaneously performs detection, localization, and quantification of emissions, eliminating the need for traditional segmentation steps. To train and evaluate the model, we construct synthetic datasets using an atmospheric transport model and validate its accuracy against radiosonde profiles and satellite observations. GHGPSE-Net demonstrates desirable performance in the simulation data across detection (F_1 -score of 0.96), subpixel-level localization and quantification (Pearson's correlation of

10 0.99, root mean square error of 89.9 tCO₂ hr⁻¹), tested on ideal instrument of ~~2-0.5~~ km × ~~2-0.5~~ km resolution with retrieval noise of 1.5 parts per million (ppm). The results also demonstrate considerable generalization of the proposed model when tested using two independent datasets. On the identified sources from OCO-3 spaceborne observations, GHGPSE-Net achieves a detection precision of ~~0.60-0.89~~, localization accuracy of ~~2.47-3.02~~ km, and a Pearson's R of ~~0.89-0.59~~ for quantification. The proposed method and datasets provide a valuable foundation for future research towards rapid and automated GHG point

15 source extraction, offering critical data to support swift responses to abnormal emission events.

1 Introduction

In response to global climate change, major economies worldwide have reached climate agreements such as the Paris Agreement, which aims to reduce greenhouse gas (GHG) emissions and limit long-term global warming to well below 2°C, striving for 1.5°C above pre-industrial levels. Spaceborne satellite remote sensing offers high-coverage and objective data to support

20 climate policy making and evaluation. Its unique advantage is highlighted by the Intergovernmental Panel on Climate Change (IPCC) (~~??~~)([Calvo Buendia et al., 2019](#)). According to various emission inventories (~~??~~[Xu et al., 2024](#))([Janssens-Maenhout et al., 2019](#); [O](#) and observation campaigns (~~??~~)([Duren et al., 2019](#); [Thorpe et al., 2023](#)), a significant portion of anthropogenic GHGs is emit-

ted by spatially concentrated facilities, or point sources. Spaceborne GHG monitoring can track emission rates for these point sources, as well as reveal abnormal emissions, providing valuable reference data for environmental administration departments.

25 Most current carbon monitoring satellites measure the backscatter solar radiation in the CO₂ or CH₄ absorption band with fine spectral resolution. GHG concentrations are then retrieved using full physics algorithms (Yokota et al., 2009; Yang et al., 2020b; Crisp
(Yoshida et al., 2013; Yang et al., 2020a; O'Dell et al., 2018) or alternative algorithms such as the IMAP-DOAS (Frankenberg
et al., 2005) and WFM-DOAS (Klings et al., 2011). The concentration map can reveal high-value pixels from the emission
plume of a point source, which can be used for source detection and quantification. Exploratory research uses sparse-spatial-
30 resolution instruments, which are initially for global observation to support assimilation systems, to perform point source mon-
itoring. For example, OCO-2/3 satellites are employed for global power station CO₂ emission monitoring using the Gaussian
plume inversion method (Nassar et al., 2017, 2021; Lin et al., 2023a)(Nassar et al., 2017, 2021; Lin et al., 2023b). Following
OCO-2/3, fine-spatial-resolution and narrow-swath (FSR-NS) instruments such as GHGSat(?, PRISMA(Guanter et al., 2021a)
, AHSI(?(Varon et al., 2018), PRISMA(Guanter et al., 2021b), AHSI(He et al., 2024), and EMIT(?(Thorpe et al., 2023) are
35 of particular interest due to their ability to resolve plume structure for precise localization and quantification. However, these in-
struments are limited by their narrow swath and coverage. To address this, sparse-spatial-resolution and wide-swath (SSR-WS)
satellites represented by TROPOMI are used to identify regions with super emitters, guiding further investigations with FSR-SS
satellites (Maasakkers et al., 2021; ?)(Irakulis-Loitxate et al., 2022; Maasakkers et al., 2021; Schuit et al., 2023). New-generation
carbon monitoring satellites, such as CO2M(?(Durand et al., 2023), GOSAT-GW(?, and The (Tanimoto et al., 2025), and the
40 new generation of Chinese Carbon Dioxide Observation Satellite Mission (TanSat-2) (Wu et al., 2023; ?)(Wu et al., 2023; Fan et al., 2025)
, typically have a spatial resolution of 0.5 to 4 km and a swath width ranging from several hundred to several thousand kilome-
ters, balancing spatial resolution and swath and enhancing point source tracking ability and guiding investigations with FSR-
NS satellites. TanSat-2 is equipped with two GHG monitoring spectrometers, including the Ultra-wide-field Carbon Pollution
collaborative monitoring Instrument (UCPI) with 2 km nadir spatial resolution, and the Hotspot Greenhouse gas Emission
45 Tracker (HGET) with 0.5 km nadir spatial resolution. TanSat-2 is also equipped with a Onboard intelligent Hotspot Extraction
and Distribution Instrument (OHEDI), which aims to verify onorbit GHG point source extraction and global distribution in
near-real-time.

These new spaceborne GHG monitoring platforms require more efficient and automatic point source extraction algorithms
for swift response across departments and missions for abnormal emission events. Ongoing work is enhancing end-to-end con-
50 centration retrieval algorithms (Chen et al., 2025b; Reuter et al., 2025), laying the foundation for automatic point source
extraction. However, current point source extraction methods still largely depend on expert analysis. For plume or source detec-
tion, Nassar et al. (2017) manually labels the emission points and assigns the plume and background pixels for quantification.
Statistical test methods (Kuhlmann et al., 2019a) and classic computer vision techniques (Varon et al., 2018) are also
introduced for more automatic plume pixel identification. Though these methods reduce manual intervention, expert interpre-
55 tation is still needed to identify plume regions. Furthermore, the subsequent quantification relies on these explicitly extracted
plume pixels. Common quantification methods include Gaussian plume fitting (Bovensmann et al., 2010; Nassar et al., 2017),

cross-sectional flux, and the integrated mass enhancement (IME) method (??)([Varon et al., 2018](#)). Therefore, new methods need to be considered for automatic source extraction.

In recent years, deep learning methods have been introduced for spaceborne GHG point source quantification (??)([Jongaramrungruang et al.](#)) and plume segmentation (??)([Schuit et al., 2023](#)). Some studies ~~futher~~ further combine retrieval and plume segmentation (????)([Joyce et al., 2023](#); [Růžička et al., 2023](#); [Vaughan et al., 2024](#); [Chen et al., 2025a](#)). However, most of these approaches treat source extraction primarily as a segmentation task, which only provides plume pixel masks, instead of the point source location and the emission strength simultaneously. Explicit source location and emission rate, rather than just a plume mask, would offer more actionable data for environmental management and other observational tasks, especially when pixel size is large.

To meet the demand for fast and automatic point source extraction in current and future missions, a deep learning point-object-detection model referred GHGPSE-Net, based on the convolutional neural network (CNN) and Gaussian kernel fitting (GKF), was proposed to simultaneously detect, locate, and quantify sources in a unified framework. Observation simulation datasets over Shanghai, a large city with complex emission distributions, were constructed using the Weather Research and Forecasting model in GHG mode (WRF-GHG; ?-?[Beck et al., 2011](#)). To increase the global applicability of models trained on this dataset, a data augmentation strategy is also proposed. Synthesized observation snapshots were generated for OCO-3 and the two instruments of TanSat-2, respectively. The performance of GHGPSE-Net was then evaluated separately on each of these synthesized datasets. To further evaluate the zero-shot generalization of the proposed model, we trained GHGPSE-Net on the synthetic dataset and tested on the independent SMARTCARB simulation dataset (??[Kuhlmann et al., 2020](#))([Kuhlmann et al., 2019b, 2020](#)) and OCO-3 observations (OCO-2/OCO-3 Science Team et al., 2022). Section 2 illustrates the construction of datasets, the design of GHGPSE-Net, the experiment setup, and the evaluation approach. Section 3 describes the model performance and evaluation results. Section 4 provides discussions and concludes the study.


2 Data and method

2.1 Synthesized observation dataset

Due to the limited coverage of current OCO-2/3 observations and the inherent retrieval noise from instrumental and prior uncertainties, the current observation data are insufficient for deep learning training, especially for future missions. In this regard, simulation-based approaches are widely adopted in the spaceborne GHG monitoring domain, as demonstrated by studies such as ?-?-?, and?[Varon et al. \(2018\)](#), [Jongaramrungruang et al. \(2022\)](#), [Radman et al. \(2023\)](#), and [Dumont Le Brazidec et al. \(2024\)](#). The Weather Research and Forecasting (WRF) model is commonly used to simulate city-scale atmospheric CO₂ transport ([Zheng et al., 2019](#); ?; [Nerobellov et al., 2023](#))([Zheng et al., 2019](#); [Lei et al., 2022](#); [Nerobellov et al., 2023](#)). WRF-GHG (??)([Beck et al., 2011](#)) is a specialized branch of WRF for GHG simulation. Comparisons with observations show that high-resolution WRF-GHG simulations can accurately capture CO₂ plumes from power plants (??)([Brunner et al., 2023](#)). We synthesize pseudo-observation datasets using WRF-GHG to train and evaluate the deep learning method. [The overall workflow for dataset construction is](#)

90 illustrated in Fig.1. (1) WRF-GHG is used to simulate the three-dimensional CO₂ concentration for multiple tracers over Shanghai from January to April 2020, with hourly snapshots (Section 2.1.1). (2) The column-averaged dry air mole fraction, denoted as XCO₂, is derived from the three-dimensional CO₂ concentration, and the pseudo-observation images are then synthesized for different instrument configurations (Section 2.1.2). (3) A data augmentation method is proposed to improve the generalization of the model trained on the spatial-temporal limited dataset (Section 2.1.3).


Figure 1. Workflow for generating the synthetic XCO₂ observation dataset.



95 2.1.1 CO₂ concentration simulation using WRF-GHG

We implement the WRF-GHG model by modifying the WRF 4.6.0 source code (<https://github.com/wrf-model/WRF/releases>, last access: 30 Sep. 2024). The physical schemes, based on [Beck et al. \(2011\)](#), include the RRTM scheme for longwave radiation, the Dudhia scheme for shortwave radiation, the Kain-Fritsch cumulus parameterization for the outermost domain, the YSU scheme for the planetary boundary layer (PBL) parameterization, the Monin-Obukhov scheme for the surface layer, the WSM5 scheme for microphysics, and the NOAH scheme for land surface model (LSM). The simulation domain focuses on Shanghai, China, using a 3-layer one-way nesting setup. The innermost domain has a resolution of 0.5 km × 0.5 km, to match the ground pixel size of HGET of TanSat-2, while the outermost domain has a resolution of 12.5 km × 12.5 km given a grid ratio of 5. The innermost domain spans 110 km × 150 km, encompassing most of the land area of Shanghai. The vertical grid has 50 layers, with the top at 5 kPa. ~~A four-month simulation continuously runs to generate 2400 3-dimensional CO₂ snapshots, each with diverse~~ The model runs continuously from January 1, 2020, to April 30, 2020, and produces snapshots at 1-hour intervals. In each snapshot, the plumes and background CO₂ concentrations are stored as different tracers.

Figure 2. WRF-GHG simulation domain settings. The left panel outlines the three nested domains, while the right panel demonstrates the distribution of major point sources (indicated by circles) and the Baoshan radiosonde station (indicated by a triangle) within the innermost domain.



The primary driving data for the simulation are summarized in Table 1. The Meteorological inputs are derived from the NCEP Final (NCEP-FNL; National Centers for Environmental Prediction et al., 2015) dataset, with a temporal resolution of 6 hours and a spatial resolution of $0.25^\circ \times 0.25^\circ$. Lateral boundary conditions for background CO_2 are derived from CarbonTracker 2022 (Jacobson et al., 2023). Power plant emissions are treated as point sources, with locations, emission rates and temporal profiles obtained from the CoCO2 dataset (Guevara et al., 2024). Power plants within 0.5 km of each other are merged, and the top 10 facilities (accounting for 79.13% of total energy-related emissions in Shanghai) are modeled as individual tracers for analysis and data augmentation. Remaining anthropogenic emissions are derived from the EDGAR Community GHG Emissions database (Crippa et al., 2024), with temporal profiles from Crippa et al. (2021) to better capture the CO_2 variability. The EDGAR dataset, at 0.1° resolution, is relatively coarse compared to our simulation grid of 0.5 km. Higher-resolution emission data could potentially improve spatial details, as suggested by Kuik et al. (2016), to better capture local concentration patterns. In this regard, similar to Bisht et al. (2023), we utilize local population, road, and land-use data as proxies to redistribute the downscale EDGAR emissions to the model grid. The biomass CO_2 fluxes are pre-calculated using the VPRM model (Mahadevan et al., 2008) and the ocean CO_2 fluxes data are derived from the GONGGA inversion dataset (Jin et al., 2024).

Table 1. Configuration of driving data for WRF-GHG simulations

Item	Source	
Meteorology	NCEP-FNL 0.25° Global Analysis data (National Centers for Environmental Prediction et al., 2015)	
Lateral boundary	Carbon Tracker 2022 (Jacobson et al., 2023)	
Emissions	Point source	CoCO2 (?) (Guevara et al., 2024)
	Other anthropogenic emissions	EDGAR v8.0 GHG (?) (Crippa et al., 2024)
	Biogenic CO2 fluxes	VPRM (?) (Mahadevan et al., 2008)
	Ocean CO2 fluxes	GONGGA CO ₂ fluxes (?) (Jin et al., 2024)

2.1.2 Synthetic XCO₂ observation dataset

The column-averaged dry air mole fraction, denoted as XCO_2 with unit parts per million (ppm), is derived from the three-dimensional CO₂ concentration of the tracers, and is given by

$$XCO_2 = \frac{VCD_{CO_2}}{VCD_{dryair}} = \frac{\sum_i VCD_{dryair,i} \times XCO_{2i}}{VCD_{dryair,i}}, \quad (1)$$

125 where VCD_{CO_2} denotes the total vertical column density of CO₂, VCD_{dryair} denotes the total vertical column density of dry air. XCO_{2i} and $VCD_{dryair,i}$ denote the CO₂ volume mixing ratio and vertical column density of dry air in i -th layer of the model, respectively, and are provided by the model output.

Pseudo-observation XCO_2 images are synthesized using WRF-GHG output in the innermost domain at a 500 m × 500 m horizontal resolution, after applying shift, rotation, and downsampling procedures as described in [?-The Pang et al. \(2025\)](#).
 130 [The downsampling process simulates the ground pixel size of different instruments and the](#) resulting images are then cropped to approximately 100 km × 100 km. Retrieval [noises are noise is](#) then added to the images, with a data augmentation approach for tracers outlined in Section 2.1.3. The retrieval noise is determined by instrumental noise and illumination, which is further defined by observation geometry, surface reflectance, aerosol optical depth, etc., such as analysed by [?-Galli et al. \(2014\)](#) and Jongaramrungruang et al. (2021). A detailed noise formulation is beyond the scope of this work, so it is treated as uncorrelated
 135 Gaussian noise. [The observation datasets with-](#)


[Here, we consider five](#) different instrument configurations, including [low-retrieval-noise \(1.5 ppm\) and high-retrieval-noise \(3 ppm\) scenarios for both HGET and UCPI of TanSat-2, as well as the OCO-3 instrument. The instrument configurations, including ground pixel size and retrieval noise, image size and noise level,](#) are shown in Table 2. Five datasets are synthesized using different instrument configurations, [respectively](#). Each dataset contains 24,000 ~~XCO~~ XCO_2 concentration maps along
 140 with the corresponding source locations and emission strengths. Examples of the synthesized XCO_2 observations for the HGET instrument are shown in Fig.3, and more examples for other instruments are shown in the Supplement.

Table 2. Instrumental configurations for pseudo-observation

Scenario	Ground pixel size (km ²)	Image size	XCO ₂ 1- σ noise (ppm)
HGET	0.5 × 0.5	192 × 192	1.5
HGET-3ppm	0.5 × 0.5	192 × 192	3.0
UCPI	2.0 × 2.0	48 × 48	1.5
UCPI-3ppm	2.0 × 2.0	48 × 48	3.0
OCO-3 ^a	2.2 × 1.6	48 × 64	1.0

^a Eldering et al. (2019).

Figure 3. Snapshots of synthesized observation by HGET across multiple footprints. Each map shows the deviation of XCO₂ from the regional average (ΔXCO_2 , in ppm). It is worth noting that ΔXCO_2 is only used for visualization purposes, while the inputs to GHGPSE-Net are the original XCO₂ values.



2.1.3 Data augmentation

We propose a data augmentation method to improve the generalization of models trained on spatial-temporal limited datasets by rescaling tracer concentrations, both directly and in proportion to emission rates. CO₂ is chemically inactive in the atmosphere, as a result, its concentration is proportional to the emission rate, based on mass conservation. This relationship allows CO₂ to scale concentrations by emission rates in spaceborne GHG monitoring simulations, such as [? and ?](#) [Jongaramrungruang et al. \(2022\)](#) and [Sánchez-García et al. \(2022\)](#).

Similar to [?Dumont Le Brazidec et al. \(2024\)](#), the concentration map can ~~is~~ be expressed as a linear combination of various tracers, given by

$$150 \quad I = \alpha_{\text{bg}} I_{\text{bg}} + \sum_i^N \alpha_{\text{ps},i} s_i I_{\text{ps},i} + \alpha_{\text{anth}} I_{\text{anth}} + \epsilon. \quad (2)$$

Here I_{bg} , $I_{\text{ps},i}$ and I_{anth} denote the background CO₂ concentration, CO₂ from i -th of total N point sources and CO₂ from other anthropogenic sources, respectively. α_{bg} , $\alpha_{\text{ps},i}$ and α_{anth} denote the corresponding scaling factors. The emission rates are scaled accordingly. s_i denotes a binary switch variable for the i -th point source. The scaling factors and switching variables are derived from random distributions. ϵ denotes the observation noise, which is simplified as Gaussian noise.

155 We introduce three scaling factors, given by

$$\alpha_{\text{bg}} = \frac{c_{\text{bg}}}{|I_{\text{bg}}|}, \alpha_{\text{ps},i} = \frac{e_{\text{ps},i}}{\bar{e}_{\text{ps},i}}, \alpha_{\text{anth}} = \frac{e_{\text{anth}}}{\bar{e}_{\text{anth}}} \quad (3)$$

Here, $|I_{\text{bg}}|$ denote the mean concentration of the background CO₂ tracer from the WRF-GHG output; $\bar{e}_{\text{ps},i}$ and \bar{e}_{anth} denote the reference emission rate for the i -th point source and the non-point-source anthropogenic sources, respectively. c_{bg} , $e_{\text{ps},i}$ and e_{anth} are random scalars drawn from several distributions. To account for the observed correlation between emissions and background concentrations, as noted by such as [Hakkarainen et al. \(2016\)](#), c_{bg} and e_{anth} are sampled from their empirical joint distribution, estimated using EDGAR emission data [?\(Crippa et al., 2024\)](#) and CarbonTracker XCO₂ data [Jacobson et al. \(2023\)](#) ([Jacobson et al., 2023](#)) in global CO₂ hotspot [areaareas](#), specifically, 100 km × 100 km areas surrounding power plants emitting over 5 MtCO₂ yr⁻¹. c_{bg} is then adjusted to track the predicted global mean XCO₂ under SSP1-2.6 (Meinshausen et al., 2020), i.e., the 2 °C scenario of the Paris Agreement, spanning the full lifetime of the TanSat-2 mission.

165 Each $\bar{e}_{\text{ps},i}$ is sampled from the major power plants in the CARMA v3.0 inventory (Ummel, 2012). Finally, we introduce binary switch variables, $s_i \sim \text{Bernoulli}(p)$, where $N \times p$ is the expectation of the quantities of power plants within the sampling area.

2.2 Deep neural network for GHG point source extraction (GHGPSE-Net)

A key goal of spaceborne GHG point source monitoring is the efficient and accurate detection, localization, and quantification of emission sources. This is essential for source attribution (e.g., [?, ?Rafiq et al., 2020](#)) and for coordinating with other observational missions (e.g., [?, ? and ?, ?Irakulis-Loitxate et al., 2022](#) and [Chiba et al., 2019](#)), particularly when the satellite spatial resolution is sparse. Traditional segmentation-based methods, however, cannot extract both source locations and emission rates automatically from a single concentration map.

Over the past decade, the remote-sensing community has widely adopted the deep learning based object-detection techniques, which focus on object localization and classification rather than pixel-wise segmentation (Zhang et al., 2023). There are two main paradigms, anchor-based network and anchor-free network. The anchor-based networks, such as SSD (Liu et al., 2016) and Mask R-CNN (He et al., 2017), predict bounding boxes directly. In contrast, anchor-free networks, such as CornerNet and CenterNet (Law and Deng, 2019) and CenterNet (Duan et al., 2019), extract object centers as key points. CenterNet, in particular, generates a heatmap with peaks corresponding to object centers whose intensities represent attributes such as length, width and orientation (Zhou et al., 2019).

Inspired by CenterNet (Duan et al., 2019), we propose GHGPSE-Net (shown in Fig.4), a CNN-based model that converts an XCO_2 concentration map into an emission heatmap composed of Gaussian kernels, as to mitigate human interventions. Source locations and emission rates are then derived using Gaussian kernel fitting (GKF).

2.2.1 Deep learning model for heatmap prediction

We represent GHG point-source emissions as a heatmap generated by the summation of a series of two-dimensional Gaussian kernels, serving as the neural network’s learning target. Each kernel’s center and amplitude correspond to the source location and the emission rate. At pixel coordinate $\mathbf{x} = [x, y]^T$, the heatmap is given by

$$I(\mathbf{x}) = \sum_{i=1}^N a_i G_i(\mathbf{x}; \boldsymbol{\mu}_i), \quad (4)$$

where $a_i \geq 0$ denotes the scale of the i -th kernel centered at $\boldsymbol{\mu}_i = [\mu_{x,i}, \mu_{y,i}]^T$. The Gaussian function G_i is given by


$$G_i(\mathbf{x}; \boldsymbol{\mu}_i) = \exp\left(-\frac{1}{2} \frac{(\mathbf{x} - \boldsymbol{\mu}_i)^T (\mathbf{x} - \boldsymbol{\mu}_i)}{\sigma^2}\right), \quad (5)$$

where σ defines the spatial extent.

We train a CNN deep learning model using supervised learning to infer emission heatmaps from XCO_2 concentration maps. Deep neural network is a class of out-of-box machine learning algorithms, which has been mathematically proved to approximate any continuous real function within a hypercube (Cybenko, 1989). A CNN primarily consists of convolutional layers, activation functions, pooling layers and linear layers, allowing it to extract compact feature representations from complex, sparse inputs. Popular image-to-image CNN architectures, such as UNet (Ronneberger et al., 2015) and HourglassNet (Newell et al., 2016), preserve spatial resolution with output feature maps similar in size to the input images, making them suitable for emission heatmap inference.

We select UNet as the CNN model. UNet features a symmetric encoder-decoder structure with skip connections (shown in Fig.4), which preserves spatial information and enables multi-scale feature fusion. UNet has been proven effective for spaceborne GHG plume segmentation tasks (Dumont Le Brazidec et al., 2023; Vaughan et al., 2024). To adapt the original UNet from classification to heatmap prediction, the final softmax layer is replaced with a convolution layer. The UNet model is implemented using the PyTorch framework (Paszke et al., 2019).

Figure 4. Overview of the proposed GHGPSE-Net architecture. Point sources are represented by a heatmap generated using Gaussian kernels, and a UNet is trained to predict this heatmap from the concentration map. Source locations and emission strengths are then inferred from the predicted heatmap through Gaussian kernel fitting. The lower panel demonstrates the UNet design used in this work. The architecture of UNet is visualized using PlotNeuralNet (Iqbal, 2018).



2.2.2 Gaussian kernel fitting

We infer source locations and emission rates by fitting predicted heatmaps using a Gaussian kernel function. Let the parameter vector be denoted as $\theta = [\mathbf{a}^T, \boldsymbol{\mu}_x^T, \boldsymbol{\mu}_y^T]^T$, where $\mathbf{a} = [a_1, a_2, \dots, a_N]^T$; $\boldsymbol{\mu}_x = [\mu_{x,1}, \mu_{x,2}, \dots, \mu_{x,N}]^T$; $\boldsymbol{\mu}_y = [\mu_{y,1}, \mu_{y,2}, \dots, \mu_{y,N}]^T$. Let P denote the number of pixels and \mathbf{x}_p denote the location of the p th pixel. We can then estimate the parameters θ by fitting the modeled image \hat{I} to the heatmap I using constrained least squares, given by

$$\theta^* = \operatorname{argmin}_{\theta} \mathcal{L}(\theta; I, \hat{I}), \quad (6)$$

where the cost function is given by

$$\mathcal{L}(\theta; I, \hat{I}) = \sum_{p=1}^P \left(I(\mathbf{x}_p) - \hat{I}(\mathbf{x}_p; \theta) \right)^2 \quad \text{s.t.} \quad a_i \geq 0. \quad (7)$$

We transform the constrained problem into an unconstrained least squares formulation through a penalty term and solve it using the BFGS method (Nocedal and Wright, 2006). The BFGS method is a quasi-Newton method that has the advantage of not requiring expensive second-order derivative computations because it approximates the Hessian matrix, leading to faster convergence. The estimated emission rates are adjusted by the ratio of local pressure to the average reference pressure in the dataset. Implementation details are provided in the Supplement.

2.2.3 Experiment setup

The models are trained from scratch using supervised learning on 24,000 simulated observation snapshots for each instrument scenario (Section 2.1.2), respectively. Each dataset is divided into training, validation, and testing sets in a 3:1:1 ratio. The model is trained using the training sets, where the model weights are updated by minimizing the mean ~~square squared~~ error (MSE) between the CNN-predicted and true heatmaps (modeled by Eq.4) ~~using the Adam optimizer over~~. The Adam optimizer is used for training with an initial learning rate of 0.001 and a batch size of 16. The models generally converge within 30 epochs, by which point the loss has generally converged. The model with the lowest validation loss at each epoch is selected for final evaluation on the test set. In the evaluation stage, the model not only predicts the heatmap but also generates source locations and emission strengths using GKF. Evaluation details are described in Section 2.3.

2.3 Evaluation

2.3.1 WRF-GHG simulation evaluation

We evaluate the accuracy of our WRF-GHG simulation by comparing the meteorological variables and XCO_2 against independent observations. The modeled meteorological outputs, including temperature and wind, are compared against radiosonde profiles from the Integrated Global Radiosonde Archive (IGRA) Version 2 ~~(?)~~ (Durre et al., 2016) at Baoshan station, which locates within the $0.5 \text{ km} \times 0.5 \text{ km}$ innermost domain of the WRF-GHG simulation (Fig.2). The simulated XCO_2 is evaluated against OCO-3 retrievals (v10.4r) in snapshot area maps (OCO-2/OCO-3 Science Team et al., 2022). To avoid discrepancies

in XCO_2 caused by differences in vertical weighting methods, XCO_2 are calculated using the modeled profiles, a priori profiles, and column averaging kernels from the OCO-3 standard product. This approach is used for the evaluation, following the method described by Connor et al. (2008); Zheng et al. (2019), rather than the direct synthesis approach described in Section 2.1.2. The accuracy is assessed using root mean square error (RMSE) to measure the overall error; and mean absolute error (MAE), which is less sensitive to large anomalies compared to RMSE.

2.3.2 Source extraction evaluation

We evaluate the performance of the proposed GHGPSE-Net in three aspects, detection, localization and quantification. Firstly, the predicted and ground-truth sources are paired by solving a linear sum assignment problem with Euclidean distance. Pairs within 4 km are counted as true positives (TP); the unmatched predictions are counted as false positives (FP); and the unmatched ground-truth sources are counted as false negatives (FN). The detection performance is then evaluated by precision, recall and F_1 -score, given by

$$\text{Precision} = \frac{TP}{TP+FP}, \text{Recall} = \frac{TP}{TP+FN}, F_1 = \frac{2TP}{2TP+FP+FN}. \quad (8)$$

Secondly, for all true positives, we evaluate the Euclidean-distance errors as the localization accuracy with their mean and median. Finally, we compare the predicted emission rates against the ground truth using the coefficient of determination (R^2) to evaluate the overall fitness; Pearson’s correlation coefficient (R) to evaluate linear correlation between predictions and actual values, regardless of scale; RMSE, MAE and mean absolute percent error (MAPE) to quantify the errors.

2.3.3 Zero-shot generalization to SMARTCARB simulations and OCO-3 observations

We evaluate the zero-shot generalization of the proposed GHGPSE-Net on two independent datasets. The models are trained using the synthetic datasets and evaluated using the following external datasets.

(1) OCO-3 SAM observations over U.S. power plants.

OCO-3 observations in SAM mode provides finer spatial resolution and denser coverage of large point sources, such as power plants and urban areas, than its predecessor OCO-2 (Kiel et al., 2021), making it suited for quantifying CO_2 emissions from large point sources globally (Yang et al., 2024; Lin et al., 2023a)(Yang et al., 2024; Lin et al., 2023b). We use the emission inventory from the Clean Air Markets Program Data (CAMPD) (EPA, 2021) as the ground truth. The OCO-3 observations are interpolated onto the regular Cartesian grid (described in Table 2) as the model inputs. Discontinuous missing data points in OCO-3 sampling, caused by cloud coverage or quality control, are filled using the mean value. From the OCO-3 dataset (OCO-2/OCO-3 Science Team et al., 2022), we select 9-8 observations covering 4 different power plants-major power plants (> 1 ktCO₂ hr⁻¹) with clean backgrounds for evaluation.

(2) SMARTCARB simulations.

To supplement the relatively sparse OCO-3 observations, we also evaluate GHGPSE-Net on the SMARTCARB simulation dataset (Kuhlmann et al., 2019b), which is generated using the Consortium for Small-scale Modelling (COSMO)-GHG model rather than WRF-GHG. The SMARTCARB dataset provides one year of hourly XCO_2 simulations at 1.1 km spatial

265 resolution, covering Berlin and surrounding major power plants, where the emission rates are derived from the TNO-MACC II inventory (Kuenen et al., 2014). We evaluate the source extraction performance of GHGPSE-Net using 998 snapshots covering major power plants, including Boxberg, Jänschwalde, Lippendorf, Schwarze Pumpe, and Turów.


2.4 Results

3 Results

3.1 WRF-GHG simulation accuracy

270 First, we compare the WRF-GHG simulated temperature, wind speed, wind direction radiosonde profiles from IGRA v2.0. As the dispersion tracers are largely confined to the planetary boundary layer (Al-Hemoud et al., 2019), we focus the comparison of meteorological variables to near-surface conditions, approximated by the 1,000 hPa pressure level. As shown in Fig.5, the WRF-GHG simulation demonstrates reasonable agreement with observations for most meteorological variables. Temperature errors are mostly within ± 3 K, with an RMSE of 4.4 K and MAE of 3.2 K, indicating that the temperature is well reproduced by
275 WRF-GHG. Wind speed errors show an RMSE of 2.9 m^{-1} and MAE of 2.3 m^{-1} , suggesting that the WRF-GHG effectively captures wind magnitude. Wind direction errors, however, exhibit larger discrepancies, with an RMSE of 82.49° and an MAE of 62.98° , reflecting notable discrepancies with observations. For dew point temperature, the RMSE and MAE are 7.3 K and 5.6 K, respectively, demonstrating moderate accuracy in representing atmospheric moisture conditions.


Figure 5. Residual distribution of meteorological variables generated by WRF-GHG compared to IGRA v2.0 measurements at Baoshan station. In each boxplot, the orange dash denotes the median, the box denotes the range between the lower and upper quartiles, the whiskers extend from the box by 1.5 times the interquartile range (IQR), and the circles denote the outliers.



280 Second, we assess WRF-GHG's capability to model $X\text{CO}_2$ against OCO-3 retrievals. As shown in Fig.6, there are three clear-sky OCO-3 passes above Shanghai (Feb 20, Feb 24, and Apr 23, 2020, UTC) within the simulation running time range, with a total of 1326 retrieval samplings. In general, the WRF-GHG model well reproduces OCO-3 observed $X\text{CO}_2$ with an RMSE of 1.58 ppm, MAE of 1.16 ppm, and Pearson's R of 0.75. A small subset of points, most of which are located downwind

of urban or heavy-industry sources during the 20 Feb and 23 Apr passes, exhibit absolute errors > 3 ppm (Supplement), while the majority of the remaining samples fall within ± 1 ppm.

Figure 6. Comparison between XCO_2 simulated by WRF-GHG and that retrieved from OCO-3 observations. The observation timestamps are provided in UTC. [The large deviations in Feb 20, 2020 and Apr 23, 2020 are mostly observed downwind of urban or heavy-industry sources \(Supplement\).](#)




285 3.2 Overall evaluations of the proposed method

We evaluate the influence of Gaussian kernel fitting (GKF) and the choice of kernel size (σ) on the performance of the proposed GHGPSE-Net. Experiments are conducted using the HGET instrument dataset of Table 2. We compare the extraction performance of the GHGPSE-Net with GKF and without GKF, where the sources are extracted from the local peaks of the heatmap as described in Section 2.2.2. Additionally, we also compare the performance over kernel sizes, both with and without
290 GKF.

As shown in Fig.7, when the kernel size matches the instrument’s spatial resolution (500 m) and GKF is applied, the model achieves its best overall performance. Qualification metrics reach maximum with a Pearson’s R of 0.99, R^2 of 0.98, RMSE of 89.9 $tCO_2 \text{ hr}^{-1}$, MAE of 57.7 $tCO_2 \text{ hr}^{-1}$, and MAPE of 0.05. Detection metrics reach peak with precision of 0.98, recall of 0.95, and F_1 -score of 0.96, indicating that most point sources are correctly detected and accurately quantified. Deviating from
295 500 m, both quantification and detection performance deteriorate, with quantification metrics beginning to regrow at σ above 1500 m. Applying GKF brings a slight improvement in quantification at the small σ cases, where the R^2 improves from 0.36 to 0.54, and RMSE decreases from 556.6 to 473.4 $tCO_2 \text{ hr}^{-1}$. Detection metrics are not sensitive to the GKF procedure.

We further investigate the influence on localization of GKF and kernel size. As shown in Fig.8, small kernel size improves locating accuracy. Moreover, GKF largely reduces the locating errors, where the mean error decreases from 199.5 m to 61.4 m, and the median error from 200.1 m to 49.7 m. These errors are well below the 500 m spatial resolution, indicating the model is
300 capable of subpixel-level source localization.

Figure 7. Scatter plots comparing GHGPSE-Net under various settings. Each plot shows results from experiments with (top row) and without (bottom row) the Gaussian kernel fitting (GKF) process, across different kernel sizes ($\sigma=250, 500, 750,$ and 1500 m from left to right). Predicted emissions are plotted against true emissions. Each plot includes quantification performance metrics, including Pearson’s R , R^2 , RMSE, MAE, and MAPE, as well as detection indicators, including precision, recall, and F_1 -score.




3.3 Comparative assessment of different instrument configurations

We evaluate the performance of GHGPSE-Net across various instrument configurations, including low-noise (1.5 ppm) and high-noise (3 ppm) retrieval scenarios for HGET onboard TanSat-2, as well as the UCPI instrument, dedicated to GHG point source monitoring. We also test the model on simulated observations with spatial resolution and retrieval noise matching those of OCO-3, which is widely used for global point source monitoring. The following experiments are conducted with the GKF process. [The kernel size of GKF is set to match each instrument’s spatial resolution, i.e., 500 m for HGET, 2 km for UCPI, and \$2.2 \times 1.6\$ km for OCO-3, in the following text.](#)

As shown in Table 3, GHGPSE-Net achieves best performance in detection, localization, and quantification with the low-noise HGET configuration. In comparison, the UCPI scenario shows degraded performance, with localization errors increasing nearly four times and quantification errors increasing to 2.6-2.8 times. The performances also deteriorate under higher retrieval noise for both instruments. Notably, the HGET-3ppm case performs worse than UCPI-3ppm in recall (0.54 vs. 0.63) and quantification (0.50 vs. 0.72 in R^2 and 0.28 vs. 0.21 in MAPE), indicating that while HGET benefits from its fine spatial resolution, it is more sensitive to uncorrelated Gaussian noise. Although OCO-3 has retrieval noise [comparable-slightly lower](#)

Figure 8. Distribution of locating errors for GHGPSE-Net under different settings. Each curve represents the normalized probability density function (PDF) for a specific experiment, scaled to the overall maximum value. Panel (a) shows results with the Gaussian kernel fitting (GKF) process, while panel (b) shows results without GKF.



315 to UCPI and features a smaller ground pixel area ($2.2 \times 1.6 \text{ km}^2$ v.s. $2 \times 2 \text{ km}^2$), UCPI achieves better performance in both
 localization and quantification. This suggests that the smaller pixel area of OCO-3 does not compensate for the limitations
 brought by its narrow shape pixel with a longer along-track dimension.

3.4 Generalization evaluation using SMARTCARB dataset and OCO-3 observation

We first evaluated the impact of data augmentation on the generalization on the model's generalization performance. Com-
 320 pared to the baseline GHGPSE-Net with augmentation, the model trained without augmentation demonstrates substantial
 performance decline, where the recall decreases from 0.95 to 0.09, the mean location error increases from 61.4 m to 310.2 m,
 and quantification RMSE increases from $89.9 \text{ tCO}_2 \text{ hr}^{-1}$ to $1072.7 \text{ tCO}_2 \text{ hr}^{-1}$ (Supplement).

To further assess the generalization capability of GHGPSE-Net trained on spatially and temporally limited simulations,
 we evaluate its performance on the SMARTCARB dataset of Berlin, simulated by COSMO-GHG, as well as on OCO-3

Table 3. Performance comparison of GHGPSE-Nets on different instrument configurations.

Instrument	Detection			Localization		Quantification				
	Precision	Recall	F_1 -score	Mean ^a	Median ^a	Pearson's R	R^2	RMSE ^b	MAE ^b	MAPE
HGET	0.98	0.95	0.96	61.4	49.7	0.99	0.98	89.9	57.7	0.05
HGET-3ppm	0.82	0.54	0.65	315.8	225.7	0.84	0.50	493.4	372.5	0.28
UCPI	0.83	0.77	0.80	227.3	182.4	0.95	0.88	249.8	147.2	0.13
UCPI-3ppm	0.60	0.64	0.62	472.7	256.6	0.89	0.72	405.0	256.1	0.21
OCO-3	0.82	0.52	0.63	662.8	479.9	0.84	0.60	479.6	335.7	0.26


^a with unit [m]. ^b with unit [tCO₂ hr⁻¹].

325 observations of power plants in the U.S. For the SMARTCARB case, the model is trained on synthesized observations of
the UCPI scenario, but with a lower noise level (standard deviation of 0.7 ppm) to match the SMARTCARB dataset. For the
OCO-3 cases, the model is trained using datasets of the OCO-3 scenario.

As shown in Fig.9, the GHGPSE-Net exhibits considerable generalization on the SMARTCARB dataset. In terms of de-
tection performance, the model achieves a precision of 0.86, which is comparable to its performance on the UCPI scenario.
330 However, the recall and F_1 -score are only 0.48 and 0.62, respectively, indicating notable mis-detections. For localization ac-
curacy, the mean and median errors are 1.80 km and 1.49 km, respectively. These values surpass those obtained when testing
on the WRF-GHG dataset, yet they remain smaller than the 2 km ground pixel size of the input data, which is enough for
providing meaningful spatial locations for joint observation missions. The quantification performance is less robust, with a
Pearson's R of 0.26, RMSE of 2.4 ktCO₂ hr⁻¹, and MAPE of 0.77. Among the five power plant cases, the model achieves best
335 performance at the Jänschwalde case, with mean reported and predicted emission rates of 4.4 ktCO₂ hr⁻¹ and 4.5 ktCO₂ hr⁻¹,
respectively. However, the model tends to overestimate emissions at the other four plants, which have comparatively lower
emission rates than Jänschwalde, with an MAE of 1.2 to 1.6 ktCO₂ hr⁻¹. Further analysis on noise-free SMARTCARB data
using GHGPSE-Net trained without noise indicates that the different noise pattern is not the major factor contributing to the
undesirable quantification performance, as the quantification performance remains largely unchanged, with a Pearson's R of
340 0.28, RMSE of 2.5 ktCO₂ hr⁻¹, and MAPE of 0.81.

Among the OCO-3 observations, GHGPSE-Net achieves a detection precision of ~~0.60~~0.89 with a mean localization error of
~~2.47~~3.02 km. We compare the estimated emissions ~~with against~~ those reported by the EPA CAMPD inventory. ~~Table 4 shows~~
~~the identified~~As shown in Table 4, the model identifies several major power plants, including Colstrip, Four Corners Steam
Electric Station, Jeffrey Energy Center, and Oak Grove. ~~The overall quantification agrees overall, the quantified emissions agree~~
345 well with the inventory, with a Pearson's R of ~~0.89~~0.59, MAPE of ~~0.31~~0.29, RMSE of ~~0.43~~0.47 ktCO₂ hr⁻¹, and MAE of ~~0.37~~
~~0.41~~ ktCO₂ hr⁻¹. ~~At Colstrip, there are five observations with estimations agree well with the EPA inventory, with an RMSE~~
~~of 0.52 ktCO₂ hr⁻¹. At~~In these cases, Colstrip and Jeffrey Energy Center are two relatively isolated power plants with high

Figure 9. Violin plot illustrating GHGPSE-Net’s emission quantification performance across five major power plants from the SMARTCARB dataset. Each violin represents the distribution of predicted emissions. The lines extended to show $\pm 1\sigma$ uncertainty with markers at their mean value. The red lines indicate the Ground-truth values from the TNO-MACC II inventory, while the black lines represent GHGPSE-Net’s predictions. The overall performances of quantification detection and localization are summarized within the panel.



emission rates, where the model correctly captures and attributes their emissions. The Four Corners Steam Electric Station ; there are two observations with an RMSE of 0.20 lies near the San Juan and Bluffview power plants (Figure 10 (b)), while Oak Grove is located near the Twin Oaks power plant (Figure 10 (c-d)). In these cases, the emission rates of these smaller nearby power plants are generally under 0.57 ktCO₂ hr⁻¹ . On April 27, 2022, a single OCO-3 overpass covers both Oak Grove and Twin Oaks , with quantification errors of 0.39 and 0.22 ktCO₂ (i.e., approximately 5 MtCO₂ hr⁻¹ , respectively) according to the CAMPD inventory, resulting in weak signals that are nearly invisible in the OCO-3 observations.


4 Discussions and Conclusion

In this study, we proposed GHGPSE-Net, a deep learning model designed to simultaneously identify the locations and quantify emission rates of GHG point sources. To train and evaluate the model, we modeled the atmospheric CO₂ transport using

Table 4. The estimated emission of identified power plants over the U.S.

Power plant	Longitude	Latitude	Observation date	EPA reports (ktCO ₂ hr ⁻¹)	Estimations (ktCO ₂ hr ⁻¹)
Colstrip	-106.6140	45.8831	2021-08-13	1.40	0.69
			2022-08-13	1.43	1.53
Four Corners Steam Elec Station	-108.4814	36.6900	2020-03-05	0.60	0.83
			2020-08-01	1.43	1.17
Jeffrey Energy Center	-96.1153	39.2825	2020-08-11	1.62	2.22
Oak Grove	-96.4853	31.1850	2021-06-16	1.78	2.29
			2021-12-21	1.59	0.88
			2022-04-27	1.71	1.58

Figure 10. Examples of input OCO-3 observations (top row) and the corresponding GHGPSE-Net predicted heatmaps (bottom row) for power plants in the U.S. The power plants in the EPA CAMPD inventory are marked and labeled with their reported emission rates. The predicted source locations are marked with gold triangles in the heatmaps. Abbreviations used in the legend are defined as follows: FCSES, Four Corners Steam Elec Station; BPP, Bluffview Power Plant; SCES, Sandy Creek Energy Station.



WRF-GHG and synthesized pseudo-observations with different instrumental configurations. The simulations were evaluated using IGRA and OCO-3 measurements, which demonstrate considerable agreement with observations. We comprehensively evaluated GHGPSE-Net’s performance on GHG point source extraction from detection, localization, and quantification. For the proposed method, the impact of Gaussian kernel fitting and the choice of kernel size selection was first assessed. We then analyzed model performance under various idealized instrument configurations. Finally, we assess the model’s generalization capabilities using simulations from SMARTCARB and spaceborne observations from OCO-3.

In general, the WRF-GHG model effectively simulated the atmospheric transport of CO₂ over Shanghai, providing a reliable synthetic dataset for training deep learning models to extract emission source. Overall, The comparison of meteorological variables against radiosonde data suggests that the WRF-GHG model provides a robust and reliable simulation of atmospheric transport conditions, with RMSE of 4.4 K for temperature, 2.9 m s⁻¹ for wind speed, 82.5° for wind direction and 7.3 K for dew point temperature. Here, we mainly focus on the evaluation of the lower troposphere. The influence of upper-tropospheric atmospheric dynamics on anthropogenic CO₂ transport is limited, as most emissions remain within the PBL (Al-Hemoud et al., 2019), where the meteorological conditions is well reproduced by WRF-GHG. The near-surface wind direction may be affected by complex terrain effects and exhibits high uncertainty, the model still captures the general direction of the CO₂ plume at larger spatial scales. The results of WRF-GHG model agree well with OCO-3, particularly for the background field. The discrepancies are mainly observed downwind of major emission sources, indicating possible inaccuracies in the emission inventory. Comprehensive evaluations of the simulation accuracy, particularly its diurnal behavior, will require continuous CO₂ observations, which are currently unavailable in this study.

Based on the synthetic dataset, we evaluated the performance of the proposed deep learning method for greenhouse gas point source extraction (GHGPSE-Net), as well as quantify the impact of model design choices. GHGPSE-Net demonstrated strong performance in source detection (F_1 -score of 0.96), localization (mean error of 12.3% of pixel size), and quantification (Pearson’s R of 0.99) in the baseline experiment. The Gaussian kernel size in the heatmap label generation and fitting has a significant impact on the model’s performance. A kernel that is too wide tends to blur local features, reducing detection accuracy, while a kernel size that is too narrow tends to bring in overly localized gradients, hindering parameter updates in training. Our experiments indicate that the optimal performance is achieved when the kernel size closely matches the pixel size. The Gaussian kernel fitting (GKF) module, implemented using least-squares optimization and concatenated after the UNet, also plays an important role in GHGPSE-Net. While its impact on detection is limited, it improves quantification moderately (24.0% reduction in RMSE compared to without GKF for the $\sigma = 500$ m baseline case) and increases the localization performance substantially (69.2% reduction in RMSE for the same case). By incorporating global context from the predicted heatmap, the GKF module increases robustness to anomalies and spurious peaks, particularly when using a smaller kernel size.

We evaluated GHGPSE-Net over various instrument configurations with different pixel size and retrieval error. In general, finer pixel resolution yields better performance for point source monitoring, which contributes to capture plume shapes and provides more spatial context for inference. However, finer pixel resolution is also more subjected to increasing retrieval noise. It is also worth noting that, in practice, the observation noise may deviate from Gaussian assumptions (??) (Jongaramrungruang et al., 2022; Radman et al., 2023), as it is mainly driven by illumination conditions including solar zenith

and surface albedo. This requires more sophisticated noise modeling, such as the parameterized approach by [Kuhlmann et al. \(2019b\)](#)

We proposed a data augmentation strategy by scaling XCO_2 from different tracers to better accommodate real-world distributions beyond the original simulation domain and time range. The scaling factors reflect more realistic patterns, including future trend, power plant density, correlation between background XCO_2 and anthropogenic emissions, etc, extending from previous strategies such as [Dumont Le Brazidec et al. \(2024\)](#). Models trained without augmentation demonstrate poor source extraction ability due to a significant mismatch between training and test data. In contrast, GHGPSE-Net trained with augmentation demonstrates considerable generalization to both the SMARTCARB simulation and OCO-3 observations. On the SMARTCARB case, GHGPSE-Net achieves high precision (0.86) but moderate recall (0.48), suggesting it can reliably identify strong sources but may miss weaker sources, which is further confirmed by the quantification performance.

These results also reveal notable discrepancies between simulation datasets. One possible explanation is the more realistic vertical emission profiles used in SMARTCARB ([Brunner et al., 2019](#)) ([Kuhlmann et al., 2019b](#); [Brunner et al., 2019](#)), which are simplified as surface emissions in this work due to the lack of these profiles. This simplification may underestimate the role of emissions entering the free troposphere, which can form new plume shapes with different orientations from those in the PBL ([Brunner et al., 2023](#)), potentially leading to false detections (Supplement). [Other factors, such as differences in meteorological conditions, terrain effects, and the use of different transport models \(COSMO-GHG v.s. WRF-GHG\), may also contribute to the discrepancies.](#) On the OCO-3 case, the quantification results generally agree with the inventory (Pearson's R of [0.89](#)[0.59](#), MAPE of [0.31](#)[0.29](#)), indicating the model demonstrates considerable generalization to real satellite observations. However, due to the discontinuous [and limited](#) coverage of OCO-3, very few samples are identified with point sources. [The complex and Heterogeneous noise characteristics, diverse plume morphologies, and varying meteorological conditions in real observations also pose challenges to model generalization.](#) Further evaluation using continuous and large-scale XCO_2 measurements from upcoming missions such as TanSat-2, CO2M, and GOSAT-GW is needed to fully assess the model's performance in real applications. [Additionally, transfer learning techniques could be explored to further improve model generalization to real-world data.](#)

In this study, we propose GHGPSE-Net, a deep learning model for simultaneous ~~localization-detection~~, [localization](#), and quantification of GHG point sources. [The model is trained and evaluated using synthetic datasets generated from WRF-GHG simulations.](#) Synthetic datasets generated by WRF-GHG are constructed for training and evaluation. ~~A~~, [accompanied by a data augmentation strategy](#) ~~is also proposed to improve the model generalization to diverse datasets. Evaluations demonstrate the model can achieve to enhance model generalization. Results show the model achieves~~ accurate detection, localization, and quantification for GHG point sources. ~~The results demonstrate the potential of~~ [In this work, we demonstrate](#) GHGPSE-Net ~~to contribute to fully~~'s [potential for](#) automated spaceborne GHG point source monitoring, ~~supporting which is critical to support~~ swift detection, assessment, and response to abnormal emission events. Future work could ~~apply GHGPSE-Net extend the model~~ to high-resolution methane monitoring tasks, trained with large eddy simulation (LES) ([Jongaramrungruang et al., 2022](#); [Radma](#)) or real satellite observations ([Schuit et al., 2023](#)); ~~explore using raw radiance as inputs, such as~~ [Joyce et al. \(2023\)](#); [Růžička and Markham \(2024\)](#); [Marjani et al. \(2024\)](#), for more compact and efficient model designs.

Ongoing efforts should also consider the impact of: investigate robustness to real-world discontinuities such as cloud cover and explore the integration of water bodies; and explore integrating auxiliary information, such as NO₂ observations as auxiliary inputs (?) (Dumont Le Brazidec et al., 2024) to help alleviate the low contrast issue in CO₂ plumes.

430 *Code and data availability.* The codes, scripts, XCO₂ snapshots generated by WRF-GHG and other datasets used are available at <https://doi.org/10.5281/zenodo.17417618>. The WRF-GHG codes, configurations, initial conditions, boundary conditions and emission files are available at <https://doi.org/10.5281/zenodo.17337441>. The GONGGA flux inversion dataset is publicly available at <https://doi.org/10.5281/zenodo.8368846> (Jin et al., 2024). The SMARTCARB simulations are publicly available at <https://doi.org/10.5281/zenodo.4034266> (Kuhlmann et al., 2019b, 2020).

Author contributions. YP and GL conceptualized the GHGPSE-Net. YP, GL, and DH designed the experiments and analyzed the data. YP
435 designed the code, performed the experiments, and visualized the results. LT coordinated the computational resources. GL, SG supervised the project. YP and GL composed the original draft. All authors reviewed and edited the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Financial support. This research has been supported by the National Key Research and Development Program of China (grant no. 2022YFB3904800).

References

- 440 Al-Hemoud, A., Al-Sudairawi, M., Al-Rashidi, M., Behbehani, W., and Al-Khayat, A.: Temperature Inversion and Mixing Height: Critical Indicators for Air Pollution in Hot Arid Climate, *Natural Hazards*, 97, 139–155, <https://doi.org/10.1007/s11069-019-03631-2>, 2019.
- Beck, V., Thomas, K., Kretschmer, R., Marshall, J., Ahmadov, R., Gerbig, C., Pillai, D., and Heimann, M.: The WRF Greenhouse Gas Model (WRF-GHG), Tech. rep., Max-Planck-Institut für Biogeochemie 25, 2011.
- Bisht, J. S. H., Patra, P. K., Takigawa, M., Kanaya, Y., Yamaguchi, M., Machida, T., and Tanimoto, H.: CO₂ High-Resolution Simulation Using WRF-GHG over the Kanto Region in Japan, *ESS Open Archive eprints*, 941, [essoar.168626399](https://doi.org/10.22541/essoar.168626399), <https://doi.org/10.22541/essoar.168626399.94131688/v1>, 2023.
- 445 Bovensmann, H., Buchwitz, M., Burrows, J. P., Reuter, M., Krings, T., Gerilowski, K., Schneising, O., Heymann, J., Tretner, A., and Erzinger, J.: A Remote Sensing Technique for Global Monitoring of Power Plant CO₂ Emissions from Space and Related Applications, *Atmospheric Measurement Techniques*, 3, 781–811, <https://doi.org/10.5194/amt-3-781-2010>, 2010.
- 450 Brunner, D., Kuhlmann, G., Marshall, J., Clément, V., Fuhrer, O., Broquet, G., Löscher, A., and Meijer, Y.: Accounting for the Vertical Distribution of Emissions in Atmospheric CO₂ Simulations, *Atmospheric Chemistry and Physics*, 19, 4541–4559, <https://doi.org/10.5194/acp-19-4541-2019>, 2019.
- Brunner, D., Kuhlmann, G., Henne, S., Koene, E., Kern, B., Wolff, S., Voigt, C., Jöckel, P., Kiemle, C., Roiger, A., Fiehn, A., Krautwurst, S., Gerilowski, K., Bovensmann, H., Borchardt, J., Galkowski, M., Gerbig, C., Marshall, J., Klonecki, A., Prunet, P., Hanfland, R., Pattantyús-Ábrahám, M., Wyszogrodzki, A., and Fix, A.: Evaluation of Simulated CO₂ Power Plant Plumes from Six High-Resolution Atmospheric Transport Models, *Atmospheric Chemistry and Physics*, 23, 2699–2728, <https://doi.org/10.5194/acp-23-2699-2023>, 2023.
- 455 Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P., and Federici, S.: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Intergovernmental Panel on Climate Change, ISBN 978-4-88788-232-4, 2019.
- 460 Chen, C., Fan, M., Wang, Z., Liang, M., Tao, J., and Chen, L.: MPSUNet: A Deep Learning-Based Segmentation Framework for Methane Plume Detection With Space-Based Hyperspectral and Multispectral Imagery, *IEEE Transactions on Geoscience and Remote Sensing*, 63, 1–15, <https://doi.org/10.1109/TGRS.2025.3563599>, 2025a.
- Chen, W., Ren, T., Zhao, C., Wen, Y., Gu, Y., Zhou, M., and Wang, P.: Transformer-Based Fast Mole Fraction of CO₂ Retrievals from Satellite-Measured Spectra, *Journal of Remote Sensing*, 5, 0470, <https://doi.org/10.34133/remotesensing.0470>, 2025b.
- 465 Chiba, T., Haga, Y., Inoue, M., Kiguchi, O., Nagayoshi, T., Madokoro, H., and Morino, I.: Measuring Regional Atmospheric CO₂ Concentrations in the Lower Troposphere with a Non-Dispersive Infrared Analyzer Mounted on a UAV, Ogata Village, Akita, Japan, *Atmosphere*, 10, 487, <https://doi.org/10.3390/atmos10090487>, 2019.
- Connor, B. J., Boesch, H., Toon, G., Sen, B., Miller, C., and Crisp, D.: Orbiting Carbon Observatory: Inverse Method and Prospective Error Analysis, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2006JD008336>, 2008.
- 470 Crippa, M., Guizzardi, D., Pisoni, E., Solazzo, E., Guion, A., Muntean, M., Florczyk, A., Schiavina, M., Melchiorri, M., and Hutterli, A. F.: Global Anthropogenic Emissions in Urban Areas: Patterns, Trends, and Challenges, *Environmental Research Letters*, 16, 074 033, <https://doi.org/10.1088/1748-9326/ac00e2>, 2021.
- Crippa, M., Guizzardi, D., Pagani, F., Schiavina, M., Melchiorri, M., Pisoni, E., Graziosi, F., Muntean, M., Maes, J., Dijkstra, L., Van Damme, M., Clarisse, L., and Coheur, P.: Insights into the Spatial Distribution of Global, National, and Subnational Greenhouse

- 475 Gas Emissions in the Emissions Database for Global Atmospheric Research (EDGAR v8.0), *Earth System Science Data*, 16, 2811–2830, <https://doi.org/10.5194/essd-16-2811-2024>, 2024.
- Crisp, D., O'Dell, C., Eldering, A., Fisher, B., Oyafuso, F., Payne, V., Drouin, B., Toon, G., Laughner, J., Somkuti, P., McGarragh, G., Merrelli, A., Nelson, R., Gunson, M., Frankenberg, C., Osterman, G., Boesch, H., Brown, L., Castano, R., Christi, M., Connor, B., McDuffie, J., Miller, C., Natraj, V., O'Brien, D., Polonski, I., Smyth, M., Thompson, D., and Granat, R.: LEVEL 2 FULL PHYSICS
480 ALGORITHM Theoretical Basis Document, Tech. rep., 2021.
- Cybenko, G.: Approximation by Superpositions of a Sigmoidal Function, *Mathematics of Control, Signals and Systems*, 2, 303–314, <https://doi.org/10.1007/BF02551274>, 1989.
- Duan, K., Bai, S., Xie, L., Qi, H., Huang, Q., and Tian, Q.: CenterNet: Keypoint Triplets for Object Detection, in: 2019 IEEE/CVF International Conference on Computer Vision (ICCV), pp. 6568–6577, ISSN 2380-7504, <https://doi.org/10.1109/ICCV.2019.00667>, 2019.
- 485 Dumont Le Brazidec, J., Vanderbecken, P., Farchi, A., Bocquet, M., Lian, J., Broquet, G., Kuhlmann, G., Danjou, A., and Lauvaux, T.: Segmentation of XCO₂ Images with Deep Learning: Application to Synthetic Plumes from Cities and Power Plants, *Geoscientific Model Development*, 16, 3997–4016, <https://doi.org/10.5194/gmd-16-3997-2023>, 2023.
- Dumont Le Brazidec, J., Vanderbecken, P., Farchi, A., Broquet, G., Kuhlmann, G., and Bocquet, M.: Deep Learning Applied to CO₂ Power Plant Emissions Quantification Using Simulated Satellite Images, *Geoscientific Model Development*, 17, 1995–2014,
490 <https://doi.org/10.5194/gmd-17-1995-2024>, 2024.
- Durand, Y., Courrèges-Lacoste, G. B., Pachot, C., Fernandez, M. M., Cabezudo, D. S., Fernandez, V., Lesschaeve, S., Spilling, D., Dusaux, A., Serre, D., Komadina, J., te Hennepe, F., and Pasquet, A.: Copernicus CO2M: Status of the Mission for Monitoring Anthropogenic Carbon Dioxide from Space, in: International Conference on Space Optics — ICSO 2022, vol. 12777, pp. 1936–1950, SPIE, <https://doi.org/10.1117/12.2690839>, 2023.
- 495 Duren, R. M., Thorpe, A. K., Foster, K. T., Rafiq, T., Hopkins, F. M., Yadav, V., Bue, B. D., Thompson, D. R., Conley, S., Colombi, N. K., Frankenberg, C., McCubbin, I. B., Eastwood, M. L., Falk, M., Herner, J. D., Croes, B. E., Green, R. O., and Miller, C. E.: California's Methane Super-Emitters, *Nature*, 575, 180–184, <https://doi.org/10.1038/s41586-019-1720-3>, 2019.
- Durre, I., Yin, X., Vose, R. S., Applequist, S., Arnfield, J., Korzeniewski, B., and Hundermark, B.: Integrated Global Radiosonde Archive (IGRA), Version 2, <https://doi.org/10.7289/V5X63K0Q>, 2016.
- 500 Eldering, A., Taylor, T. E., O'Dell, C. W., and Pavlick, R.: The OCO-3 Mission: Measurement Objectives and Expected Performance Based on 1 Year of Simulated Data, *Atmospheric Measurement Techniques*, 12, 2341–2370, <https://doi.org/10.5194/amt-12-2341-2019>, 2019.
- EPA: Clean Air Markets Program Data (CAMPD), 2021.
- Fan, M., Chen, L., Tian, L., Yang, D., Huiqin, M., Chen, Tao, J., Jiang, F., Liu, L., Zhang, M., Liu, L., Yin, Z., Chen, C., Wang, J., Yao, L., Du, S., Yu, C., Zhang, Y., Hu, D., Zhou, G., Kong, Y., and Wu, W. U.: Requirements and Design of TanSat-2 Mission, *National Remote
505 Sensing Bulletin*, null, 1–12, <https://doi.org/10.11834/jrs.20255080>, 2025.
- Frankenberg, C., Platt, U., and Wagner, T.: Iterative Maximum a Posteriori (IMAP)-DOAS for Retrieval of Strongly Absorbing Trace Gases: Model Studies for CH₄ and CO₂ Retrieval from near Infrared Spectra of SCIAMACHY Onboard ENVISAT, *Atmospheric Chemistry and Physics*, 5, 9–22, <https://doi.org/10.5194/acp-5-9-2005>, 2005.
- Galli, A., Guerlet, S., Butz, A., Aben, I., Suto, H., Kuze, A., Deutscher, N. M., Notholt, J., Wunch, D., Wennberg, P. O., Griffith, D. W. T.,
510 Hasekamp, O., and Landgraf, J.: The Impact of Spectral Resolution on Satellite Retrieval Accuracy of CO₂ and CH₄, *Atmospheric Measurement Techniques*, 7, 1105–1119, <https://doi.org/10.5194/amt-7-1105-2014>, 2014.

- Guanter, L., Irakulis-Loitxate, I., Gorrono, J., Sanchez-Garcia, E., Cusworth, D. H., Varon, D. J., Cogliati, S., and Colombo, R.: Mapping Methane Point Emissions with the PRISMA Spaceborne Imaging Spectrometer, *Remote Sensing of Environment*, 265, 112671, <https://doi.org/10.1016/j.rse.2021.112671>, 2021a.
- 515 Guanter, L., Irakulis-Loitxate, I., Gorroño, J., Sánchez-García, E., Cusworth, D. H., Varon, D. J., Cogliati, S., and Colombo, R.: Mapping Methane Point Emissions with the PRISMA Spaceborne Imaging Spectrometer, *Remote Sensing of Environment*, 265, 112671, <https://doi.org/10.1016/j.rse.2021.112671>, 2021b.
- Guevara, M., Enciso, S., Tena, C., Jorba, O., Dellaert, S., Denier van der Gon, H., and Pérez García-Pando, C.: A Global Catalogue of CO₂ Emissions and Co-Emitted Species from Power Plants, Including High-Resolution Vertical and Temporal Profiles, *Earth System Science Data*, 16, 337–373, <https://doi.org/10.5194/essd-16-337-2024>, 2024.
- 520 Hakkarainen, J., Ialongo, I., and Tamminen, J.: Direct Space-Based Observations of Anthropogenic CO₂ Emission Areas from OCO-2, *Geophysical Research Letters*, 43, 11,400–11,406, <https://doi.org/10.1002/2016GL070885>, 2016.
- He, K., Gkioxari, G., Dollár, P., and Girshick, R.: Mask R-CNN, in: 2017 IEEE International Conference on Computer Vision (ICCV), pp. 2980–2988, <https://doi.org/10.1109/ICCV.2017.322>, 2017.
- 525 He, Z., Gao, L., Liang, M., and Zeng, Z.-C.: A Survey of Methane Point Source Emissions from Coal Mines in Shanxi Province of China Using AHSI on Board Gaofen-5B, *Atmospheric Measurement Techniques*, 17, 2937–2956, <https://doi.org/10.5194/amt-17-2937-2024>, 2024.
- Iqbal, H.: HarisIqbal88/PlotNeuralNet v1.0.0, Zenodo, <https://doi.org/10.5281/zenodo.2526396>, 2018.
- Irakulis-Loitxate, I., Guanter, L., Maasackers, J. D., Zavala-Araiza, D., and Aben, I.: Satellites Detect Abatable Super-
530 Emissions in One of the World’s Largest Methane Hotspot Regions, *Environmental Science & Technology*, 56, 2143–2152, <https://doi.org/10.1021/acs.est.1c04873>, 2022.
- Jacobson, A. R., Schuldt, K. N., Tans, P., Arlyn Andrews, Miller, J. B., Oda, T., Mund, J., Weir, B., Ott, L., Aalto, T., Abshire, J. B., Aikin, K., Aoki, S., Apadula, F., Arnold, S., Baier, B., Bartyzel, J., Beyersdorf, A., Biermann, T., Biraud, S. C., Boenisch, H., Brailsford, G., Brand, W. A., Chen, G., Huilin Chen, Lukasz Chmura, Clark, S., Colomb, A., Commane, R., Conil, S., Couret, C., Cox, A., Cristofanelli, P., Cuevas, E., Curcoll, R., Daube, B., Davis, K. J., De Wekker, S., Coletta, J. D., Delmotte, M., DiGangi, E., DiGangi, J. P., Di Sarra, A. G., Dlugokencky, E., Elkins, J. W., Emmenegger, L., Shuangxi Fang, Fischer, M. L., Forster, G., Frumau, A., Galkowski, M., Gatti, L. V., Gehrlein, T., Gerbig, C., Francois Gheusi, Gloor, E., Gomez-Trueba, V., Goto, D., Griffis, T., Hammer, S., Hanson, C., Haszpra, L., Hatakka, J., Heimann, M., Heliasz, M., Hensen, A., Hermansen, O., Hintsä, E., Holst, J., Ivakhov, V., Jaffe, D. A., Jordan, A., Joubert, W., Karion, A., Kawa, S. R., Kazan, V., Keeling, R. F., Keronen, P., Kneuer, T., Kolari, P., Kateřina Komínková, Kort, E., Kozlova, E.,
540 Krummel, P., Kubistin, D., Labuschagne, C., Lam, D. H., Lan, X., Langenfelds, R. L., Laurent, O., Laurila, T., Lauvaux, T., Lavric, J., Law, B. E., Lee, J., Lee, O. S., Lehner, I., Lehtinen, K., Leppert, R., Leskinen, A., Leuenberger, M., Levin, I., Levula, J., Lin, J., Lindauer, M., Loh, Z., Lopez, M., Luijkx, I. T., Lunder, C. R., Machida, T., Mammarella, I., Manca, G., Manning, A., Manning, A., Marek, M. V., Martin, M. Y., Matsueda, H., McKain, K., Meijer, H., Meinhardt, F., Merchant, L., N. Mihalopoulos, Miles, N. L., Miller, C. E., Mitchell, L., Mölder, M., Montzka, S., Moore, F., Moossen, H., Morgan, E., Josep-Anton Morgui, Morimoto, S., Müller-Williams, J., J. William
545 Munger, Munro, D., Myhre, C. L., Shin-Ichiro Nakaoka, Jaroslaw Necki, Newman, S., Nichol, S., Niwa, Y., Obersteiner, F., O’Doherty, S., Paplawsky, B., Peischl, J., Peltola, O., Piacentino, S., Jean-Marc Pichon, Pickers, P., Piper, S., Pitt, J., Plass-Dülmer, C., Platt, S. M., Prinzivalli, S., Ramonet, M., Ramos, R., Reyes-Sanchez, E., Richardson, S. J., Riris, H., Rivas, P. P., Ryerson, T., Saito, K., Sargent, M., Sasakawa, M., Scheeren, B., Schuck, T., Schumacher, M., Seifert, T., Sha, M. K., Shepson, P., Shook, M., Sloop, C. D., Smith, P., Stanley, K., Steinbacher, M., Stephens, B., Sweeney, C., Thoning, K., Timas, H., Torn, M., Tørseth, K., Trisolino, P., Turnbull, J., Van Den Bulk,

- 550 P., Van Dinter, D., Vermeulen, A., Viner, B., Vitkova, G., Walker, S., Watson, A., Wofsy, S. C., Worsley, J., Worthy, D., Dickon Young, Zaehle, S., Zahn, A., and Miroslaw Zimnoch: CarbonTracker CT2022, <https://doi.org/10.25925/Z1GJ-3254>, 2023.
- Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi, P., Pagliari, V., Olivier, J. G. J., Peters, J. A. H. W., van Aardenne, J. A., Monni, S., Doering, U., Petrescu, A. M. R., Solazzo, E., and Oreggioni, G. D.: EDGAR v4.3.2 Global Atlas of the Three Major Greenhouse Gas Emissions for the Period 1970–2012, *Earth System Science Data*, 11, 959–1002, <https://doi.org/10.5194/essd-11-959-2019>, 2019.
- 555 Jin, Z., Tian, X., Wang, Y., Zhang, H., Zhao, M., Wang, T., Ding, J., and Piao, S.: A Global Surface CO₂ Flux Dataset (2015–2022) Inferred from OCO-2 Retrievals Using the GONGGA Inversion System, *Earth System Science Data*, 16, 2857–2876, <https://doi.org/10.5194/essd-16-2857-2024>, 2024.
- Jongaramrungruang, S., Matheou, G., Thorpe, A. K., Zeng, Z.-C., and Frankenberg, C.: Remote Sensing of Methane Plumes: Instrument Tradeoff Analysis for Detecting and Quantifying Local Sources at Global Scale, *Atmospheric Measurement Techniques*, 14, 7999–8017, <https://doi.org/10.5194/amt-14-7999-2021>, 2021.
- 560 Jongaramrungruang, S., Thorpe, A. K., Matheou, G., and Frankenberg, C.: MethaNet - An AI-driven Approach to Quantifying Methane Point-Source Emission from High-Resolution 2-D Plume Imagery, *Remote Sensing of Environment*, 269, 112 809, <https://doi.org/10.1016/j.rse.2021.112809>, 2022.
- 565 Joyce, P., Ruiz Villena, C., Huang, Y., Webb, A., Gloor, M., Wagner, F. H., Chipperfield, M. P., Barrio Guilló, R., Wilson, C., and Boesch, H.: Using a Deep Neural Network to Detect Methane Point Sources and Quantify Emissions from PRISMA Hyperspectral Satellite Images, *Atmospheric Measurement Techniques*, 16, 2627–2640, <https://doi.org/10.5194/amt-16-2627-2023>, 2023.
- Kiel, M., Eldering, A., Roten, D. D., Lin, J. C., Feng, S., Lei, R., Lauvaux, T., Oda, T., Roehl, C. M., Blavier, J.-F., and Iraci, L. T.: Urban-Focused Satellite CO₂ Observations from the Orbiting Carbon Observatory-3: A First Look at the Los Angeles Megacity, *Remote Sensing of Environment*, 258, 112 314, <https://doi.org/10.1016/j.rse.2021.112314>, 2021.
- 570 Krings, T., Gerilowski, K., Buchwitz, M., Reuter, M., Tretner, A., Erzinger, J., Heinze, D., Pflüger, U., Burrows, J. P., and Bovensmann, H.: MAMAP – a New Spectrometer System for Column-Averaged Methane and Carbon Dioxide Observations from Aircraft: Retrieval Algorithm and First Inversions for Point Source Emission Rates, *Atmospheric Measurement Techniques*, 4, 1735–1758, <https://doi.org/10.5194/amt-4-1735-2011>, 2011.
- 575 Kuenen, J. J. P., Visschedijk, A. J. H., Jozwicka, M., and Denier van der Gon, H. a. C.: TNO-MACC_II Emission Inventory; a Multi-Year (2003–2009) Consistent High-Resolution European Emission Inventory for Air Quality Modelling, *Atmospheric Chemistry and Physics*, 14, 10 963–10 976, <https://doi.org/10.5194/acp-14-10963-2014>, 2014.
- Kuhlmann, G., Broquet, G., Marshall, J., Clément, V., Löscher, A., Meijer, Y., and Brunner, D.: Detectability of CO₂ Emission Plumes of Cities and Power Plants with the Copernicus Anthropogenic CO₂ Monitoring (CO₂M) Mission, *Atmospheric Measurement Techniques*, 12, 6695–6719, <https://doi.org/10.5194/amt-12-6695-2019>, 2019a.
- 580 Kuhlmann, G., Clément, V., Marshall, J., Fuhrer, O., Broquet, G., Schnadt-Poberaj, C., Löscher, A., Meijer, Y., and Brunner, D.: SMART-CARB – Use of Satellite Measurements of Auxiliary Reactive Trace Gases for Fossil Fuel Carbon Dioxide Emission Estimation, Tech. rep., Zenodo, <https://doi.org/10.5281/zenodo.4034266>, 2019b.
- Kuhlmann, G., Clément, V., Marshall, J., Fuhrer, O., Broquet, G., Schnadt-Poberaj, C., Löscher, A., Meijer, Y., and Brunner, D.: Synthetic XCO₂, CO and NO₂ Observations for the CO₂M and Sentinel-5 Satellites, <https://doi.org/10.5281/zenodo.4048228>, 2020.
- 585

- Kuik, F., Lauer, A., Churkina, G., Denier van der Gon, H. A. C., Fenner, D., Mar, K. A., and Butler, T. M.: Air Quality Modelling in the Berlin–Brandenburg Region Using WRF-Chem v3.7.1: Sensitivity to Resolution of Model Grid and Input Data, *Geoscientific Model Development*, 9, 4339–4363, <https://doi.org/10.5194/gmd-9-4339-2016>, 2016.
- Law, H. and Deng, J.: CornerNet: Detecting Objects as Paired Keypoints, <https://doi.org/10.48550/arXiv.1808.01244>, 2019.
- 590 Lei, R., Feng, S., Xu, Y., Tran, S., Ramonet, M., Grutter, M., Garcia, A., Campos-Pineda, M., and Lauvaux, T.: Reconciliation of Asynchronous Satellite-Based NO₂ and XCO₂ Enhancements with Mesoscale Modeling over Two Urban Landscapes, *Remote Sensing of Environment*, 281, 113 241, <https://doi.org/10.1016/j.rse.2022.113241>, 2022.
- Lin, X., van der A, R., de Laat, J., Eskes, H., Chevallier, F., Ciais, P., Deng, Z., Geng, Y., Song, X., Ni, X., Huo, D., Dou, X., and Liu, Z.: Monitoring and Quantifying CO₂ Emissions of Isolated Power Plants from Space, *Atmospheric Chemistry and Physics*, 23, 6599–6611, 595 <https://doi.org/10.5194/acp-23-6599-2023>, 2023a.
- Lin, X., van der A, R., de Laat, J., Eskes, H., Chevallier, F., Ciais, P., Deng, Z., Geng, Y., Song, X., Ni, X., Huo, D., Dou, X., and Liu, Z.: Monitoring and Quantifying CO₂ Emissions of Isolated Power Plants from Space, *Atmospheric Chemistry and Physics*, 23, 6599–6611, <https://doi.org/10.5194/acp-23-6599-2023>, 2023b.
- Liu, W., Anguelov, D., Erhan, D., Szegedy, C., Reed, S., Fu, C.-Y., and Berg, A. C.: SSD: Single Shot MultiBox Detector, in: *Computer Vision – ECCV 2016*, edited by Leibe, B., Matas, J., Sebe, N., and Welling, M., Lecture Notes in Computer Science, pp. 21–37, Springer International Publishing, Cham, ISBN 978-3-319-46448-0, https://doi.org/10.1007/978-3-319-46448-0_2, 2016.
- Maasakkers, J. D., Varon, D. J., Elfarsdóttir, A., McKeever, J., Jervis, D., Mahapatra, G., Pandey, S., Lorente, A., Borsdorff, T., Foorhuis, L. R., Schuit, B. J., Tol, P., van Kempen, T. A., van Hees, R., and Aben, I.: Using Satellites to Uncover Large Methane Emissions from Landfills, 2021.
- 605 Mahadevan, P., Wofsy, S. C., Matross, D. M., Xiao, X., Dunn, A. L., Lin, J. C., Gerbig, C., Munger, J. W., Chow, V. Y., and Gottlieb, E. W.: A Satellite-Based Biosphere Parameterization for Net Ecosystem CO₂ Exchange: Vegetation Photosynthesis and Respiration Model (VPRM), *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2006GB002735>, 2008.
- Marjani, M., Mohammadimanesh, F., Varon, D. J., Radman, A., and Mahdianpari, M.: PRISMethaNet: A Novel Deep Learning Model for Landfill Methane Detection Using PRISMA Satellite Data, *ISPRS Journal of Photogrammetry and Remote Sensing*, 218, 802–818, 610 <https://doi.org/10.1016/j.isprsjprs.2024.10.003>, 2024.
- Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The Shared Socio-Economic Pathway (SSP) Greenhouse Gas Concentrations and Their Extensions to 2500, *Geoscientific Model Development*, 13, 3571–3605, [https://doi.org/10.5194/gmd-13-3571-](https://doi.org/10.5194/gmd-13-3571-2020) 615 2020, 2020.
- Nassar, R., Hill, T. G., McLinden, C. A., Wunch, D., Jones, D. B. A., and Crisp, D.: Quantifying CO₂ Emissions From Individual Power Plants From Space, *Geophysical Research Letters*, 44, 10 045–10 053, <https://doi.org/10.1002/2017GL074702>, 2017.
- Nassar, R., Mastrogiacomo, J.-P., Bateman-Hemphill, W., McCracken, C., MacDonald, C. G., Hill, T., O’Dell, C. W., Kiel, M., and Crisp, D.: Advances in Quantifying Power Plant CO₂ Emissions with OCO-2, *Remote Sensing of Environment*, 264, 112 579, 620 <https://doi.org/10.1016/j.rse.2021.112579>, 2021.
- National Centers for Environmental Prediction, Service, N. W., NOAA, and U S. Department of Commerce: NCEP GDAS/FNL 0.25 Degree Global Tropospheric Analyses and Forecast Grids, <https://doi.org/10.5065/D65Q4T4Z>, 2015.

- Nerobelov, G., Timofeyev, Y., Foka, S., Smyshlyaev, S., Poberovskiy, A., and Sedeeva, M.: Complex Validation of Weather Research and Forecasting—Chemistry Modelling of Atmospheric CO₂ in the Coastal Cities of the Gulf of Finland, *Remote Sensing*, 15, 5757, <https://doi.org/10.3390/rs15245757>, 2023.
- 625 Newell, A., Yang, K., and Deng, J.: Stacked Hourglass Networks for Human Pose Estimation, <https://doi.org/10.48550/arXiv.1603.06937>, 2016.
- Nocedal, J. and Wright, S. J.: Quasi-Newton Methods, in: *Numerical Optimization*, pp. 135–163, Springer, New York, NY, ISBN 978-0-387-40065-5, https://doi.org/10.1007/978-0-387-40065-5_6, 2006.
- 630 OCO-2/OCO-3 Science Team, Chatterjee, A., and Payne, V.: OCO-3 Level 2 Bias-Corrected XCO₂ and Other Select Fields from the Full-Physics Retrieval Aggregated as Daily Files, Retrospective Processing V10.4r, 2022.
- Oda, T., Maksyutov, S., and Andres, R. J.: The Open-source Data Inventory for Anthropogenic CO₂, Version 2016 (ODIAC2016): A Global Monthly Fossil Fuel CO₂ Gridded Emissions Data Product for Tracer Transport Simulations and Surface Flux Inversions, *Earth System Science Data*, 10, 87–107, <https://doi.org/10.5194/essd-10-87-2018>, 2018.
- 635 O'Dell, C. W., Eldering, A., Wennberg, P. O., Crisp, D., Gunson, M. R., Fisher, B., Frankenberg, C., Kiel, M., Lindqvist, H., Mandrake, L., Merrelli, A., Natraj, V., Nelson, R. R., Osterman, G. B., Payne, V. H., Taylor, T. E., Wunch, D., Drouin, B. J., Oyafuso, F., Chang, A., McDuffie, J., Smyth, M., Baker, D. F., Basu, S., Chevallier, F., Crowell, S. M. R., Feng, L., Palmer, P. I., Dubey, M., García, O. E., Griffith, D. W. T., Hase, F., Iraci, L. T., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Petri, C., Roehl, C. M., Sha, M. K., Strong, K., Sussmann, R., Te, Y., Uchino, O., and Velazco, V. A.: Improved Retrievals of Carbon Dioxide from Orbiting Carbon Observatory-2 with the Version
- 640 8 ACOS Algorithm, *Atmospheric Measurement Techniques*, 11, 6539–6576, <https://doi.org/10.5194/amt-11-6539-2018>, 2018.
- Pang, Y., Tian, L., Denghui, H., Gao, S., and Liu, G.: Separating and Quantifying Facility-Level Methane Emissions with Overlapping Plumes for Spaceborne Methane Monitoring, *Atmospheric Measurement Techniques*, 18, 455–470, <https://doi.org/10.5194/amt-18-455-2025>, 2025.
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein, N., Antiga, L., Desmaison, A., Köpf, A., Yang, E., DeVito, Z., Raison, M., Tejani, A., Chilamkurthy, S., Steiner, B., Fang, L., Bai, J., and Chintala, S.: PyTorch: An Imperative
- 645 Style, High-Performance Deep Learning Library, <https://doi.org/10.48550/arXiv.1912.01703>, 2019.
- Radman, A., Mahdianpari, M., Varon, D. J., and Mohammadimanesh, F.: S2MetNet: A Novel Dataset and Deep Learning Benchmark for Methane Point Source Quantification Using Sentinel-2 Satellite Imagery, *Remote Sensing of Environment*, 295, 113 708, <https://doi.org/10.1016/j.rse.2023.113708>, 2023.
- 650 Rafiq, T., Duren, R. M., Thorpe, A. K., Foster, K., Patarsuk, R., Miller, C. E., and Hopkins, F. M.: Attribution of Methane Point Source Emissions Using Airborne Imaging Spectroscopy and the Vista-California Methane Infrastructure Dataset, *Environmental Research Letters*, 15, 124 001, <https://doi.org/10.1088/1748-9326/ab9af8>, 2020.
- Reuter, M., Hilker, M., Noël, S., Di Noia, A., Weimer, M., Schneising, O., Buchwitz, M., Bovensmann, H., Burrows, J. P., Bösch, H., and Lang, R.: Retrieving the Atmospheric Concentrations of Carbon Dioxide and Methane from the European Copernicus CO₂M Satellite
- 655 Mission Using Artificial Neural Networks, *Atmospheric Measurement Techniques*, 18, 241–264, <https://doi.org/10.5194/amt-18-241-2025>, 2025.
- Ronneberger, O., Fischer, P., and Brox, T.: U-Net: Convolutional Networks for Biomedical Image Segmentation, in: *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015*, edited by Navab, N., Hornegger, J., Wells, W. M., and Frangi, A. F., *Lecture Notes in Computer Science*, pp. 234–241, Springer International Publishing, Cham, ISBN 978-3-319-24574-4,
- 660 https://doi.org/10.1007/978-3-319-24574-4_28, 2015.

- Růžička, V. and Markham, A.: HyperspectralViTs: General Hyperspectral Models for On-board Remote Sensing, 2024.
- Růžička, V., Mateo-García, G., Gómez-Chova, L., Vaughan, A., Guanter, L., and Markham, A.: Semantic Segmentation of Methane Plumes with Hyperspectral Machine Learning Models, *Scientific Reports*, 13, <https://doi.org/10.1038/s41598-023-44918-6>, 2023.
- 665 Sánchez-García, E., Gorroño, J., Irakulis-Loitxate, I., Varon, D. J., and Guanter, L.: Mapping Methane Plumes at Very High Spatial Resolution with the WorldView-3 Satellite, *Atmospheric Measurement Techniques*, 15, 1657–1674, <https://doi.org/10.5194/amt-15-1657-2022>, 2022.
- Schuit, B. J., Maasackers, J. D., Bijl, P., Mahapatra, G., Van den Berg, A.-W., Pandey, S., Lorente, A., Borsdorff, T., Houweling, S., Varon, D. J., McKeever, J., Jarvis, D., Girard, M., Irakulis-Loitxate, I., Gorroño, J., Guanter, L., Cusworth, D. H., and Aben, I.: Automated Detection and Monitoring of Methane Super-Emitters Using Satellite Data, *Atmospheric Chemistry and Physics Discussions*, pp. 1–47, <https://doi.org/10.5194/acp-2022-862>, 2023.
- 670 Tanimoto, H., Matsunaga, T., Someya, Y., Fujinawa, T., Ohyama, H., Morino, I., Yashiro, H., Sugita, T., Inomata, S., Mueller, A., Saeki, T., Yoshida, Y., Niwa, Y., Saito, M., Noda, H., Yamashita, Y., Ikeda, K., Saigusa, N., Machida, T., and Sato, T.: The Greenhouse Gas Observation Mission with Global Observing SATellite for Greenhouse Gases and Water Cycle (GOSAT-GW): Objectives, Conceptual Framework and Scientific Contributions, *Progress in Earth and Planetary Science*, 12, <https://doi.org/10.1186/s40645-025-00684-9>, 2025.
- Thorpe, A. K., Green, R. O., Thompson, D. R., Brodrick, P. G., Chapman, J. W., Elder, C. D., Irakulis-Loitxate, I., Cusworth, D. H., Ayasse, A. K., Duren, R. M., Frankenberg, C., Guanter, L., Worden, J. R., Dennison, P. E., Roberts, D. A., Chadwick, K. D., Eastwood, M. L., Fahlen, J. E., and Miller, C. E.: Attribution of Individual Methane and Carbon Dioxide Emission Sources Using EMIT Observations from Space, *Science Advances*, 9, eadh2391, <https://doi.org/10.1126/sciadv.adh2391>, 2023.
- 675 Ummel, K.: CARMA Revisited: An Updated Database of Carbon Dioxide Emissions from Power Plants Worldwide - Working Paper 304, 2012.
- 680 Varon, D. J., Jacob, D. J., McKeever, J., Jarvis, D., Durak, B. O. A., Xia, Y., and Huang, Y.: Quantifying Methane Point Sources from Fine-Scale Satellite Observations of Atmospheric Methane Plumes, *Atmospheric Measurement Techniques*, 11, 5673–5686, <https://doi.org/10.5194/amt-11-5673-2018>, 2018.
- Vaughan, A., Mateo-García, G., Gómez-Chova, L., Růžička, V., Guanter, L., and Irakulis-Loitxate, I.: CH4Net: A Deep Learning Model for Monitoring Methane Super-Emitters with Sentinel-2 Imagery, *Atmospheric Measurement Techniques*, 17, 2583–2593, <https://doi.org/10.5194/amt-17-2583-2024>, 2024.
- 685 Wu, K., Yang, D., Liu, Y., Cai, Z., Zhou, M., Feng, L., and Palmer, P. I.: Evaluating the Ability of the Pre-Launch TanSat-2 Satellite to Quantify Urban CO₂ Emissions, *Remote Sensing*, 15, 4904, <https://doi.org/10.3390/rs15204904>, 2023.
- Xu, R., Tong, D., Xiao, Q., Qin, X., Chen, C., Yan, L., Cheng, J., Cui, C., Hu, H., Liu, W., Yan, X., Wang, H., Liu, X., Geng, G., Lei, Y., Guan, D., He, K., and Zhang, Q.: MEIC-global-CO₂: A New Global CO₂ Emission Inventory with Highly-Resolved Source Category and Sub-Country Information, *Science China Earth Sciences*, 67, 450–465, <https://doi.org/10.1007/s11430-023-1230-3>, 2024.
- 690 Yang, D., Boesch, H., Liu, Y., Somkuti, P., Cai, Z., Chen, X., Di Noia, A., Lin, C., Lu, N., Lyu, D., Parker, R. J., Tian, L., Wang, M., Webb, A., Yao, L., Yin, Z., Zheng, Y., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Pollard, D. F., Shiomi, K., Sussmann, R., Té, Y., Velazco, V. A., Warneke, T., and Wunch, D.: Toward High Precision XCO₂ Retrievals From TanSat Observations: Retrieval Improvement and Validation Against TCCON Measurements, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032794, <https://doi.org/10.1029/2020JD032794>, 2020a.
- 695 Yang, D., Boesch, H., Liu, Y., Somkuti, P., Cai, Z., Chen, X., Di Noia, A., Lin, C., Lu, N., Lyu, D., Parker, R. J., Tian, L., Wang, M., Webb, A., Yao, L., Yin, Z., Zheng, Y., Deutscher, N. M., Griffith, D. W. T., Hase, F., Kivi, R., Morino, I., Notholt, J., Ohyama, H., Pollard, D. F., Shiomi, K., Sussmann, R., Té, Y., Velazco, V. A., Warneke, T., and Wunch, D.: Toward High Precision XCO₂ Retrievals From TanSat

- Observations: Retrieval Improvement and Validation Against TCCON Measurements, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032794, <https://doi.org/10.1029/2020JD032794>, 2020b.
- 700 Yang, Y., Zhou, M., Wang, W., Ning, Z., Zhang, F., and Wang, P.: Quantification of CO₂ Emissions from Three Power Plants in China Using OCO-3 Satellite Measurements, *Advances in Atmospheric Sciences*, 41, 2276–2288, <https://doi.org/10.1007/s00376-024-3293-9>, 2024.
- Yokota, T., Yoshida, Y., Eguchi, N., Ota, Y., Tanaka, T., Watanabe, H., and Maksyutov, S.: Global Concentrations of CO₂ and CH₄ Retrieved from GOSAT: First Preliminary Results, *Sola*, 5, 160–163, <https://doi.org/10.2151/sola.2009-041>, 2009.
- 705 Yoshida, Y., Kikuchi, N., Morino, I., Uchino, O., Oshchepkov, S., Bril, A., Saeki, T., Schutgens, N., Toon, G. C., Wunch, D., Roehl, C. M., Wennberg, P. O., Griffith, D. W. T., Deutscher, N. M., Warneke, T., Notholt, J., Robinson, J., Sherlock, V., Connor, B., Rettinger, M., Sussmann, R., Ahonen, P., Heikkinen, P., Kyro, E., Mendonca, J., Strong, K., Hase, F., Dohe, S., and Yokota, T.: Improvement of the Retrieval Algorithm for GOSAT SWIR XCO₂ and XCH₄ and Their Validation Using TCCON Data, *Atmospheric Measurement Techniques*, 6, 1533–1547, <https://doi.org/10.5194/amt-6-1533-2013>, 2013.
- 710 Zhang, X., Zhang, T., Wang, G., Zhu, P., Tang, X., Jia, X., and Jiao, L.: Remote Sensing Object Detection Meets Deep Learning: A Meta-review of Challenges and Advances, <https://doi.org/10.48550/arXiv.2309.06751>, 2023.
- Zheng, T., Nassar, R., and Baxter, M.: Estimating Power Plant CO₂ Emission Using OCO-2 XCO₂ and High Resolution WRF-Chem Simulations, *Environmental Research Letters*, 14, 085001, <https://doi.org/10.1088/1748-9326/ab25ae>, 2019.
- Zhou, X., Wang, D., and Krähenbühl, P.: Objects as Points, <https://doi.org/10.48550/arXiv.1904.07850>, 2019.