

Beyond GRACE: Evaluating the benefits of NGGM and MAGIC for precipitation estimation over Europe

Muhammad Usman Liaqat¹, Stefania Camici¹, Francesco Leopardi^{1,2}, Jaime Gaona¹, Luca Brocca¹

¹Research Institute for Geo-Hydrological Protection, National Research Council of Italy, Via Madonna Alta, 126, 06128 Perugia, Italy.

²Department of Civil and Environmental Engineering, University of Perugia, Italy.

Correspondence: muhammadasman.liaqat@cnr.it

Abstract

The Gravity Recovery and Climate Experiment (GRACE) and its Follow-On (GRACE-FO) missions provide observations of terrestrial water storage (TWS) dynamics from regional to global scales. However, their limited spatio-temporal resolution is restricted in many hydrological applications. Their coarse spatio-temporal resolution limits their applicability in many hydrological studies. A joint collaboration between the National Aeronautics and Space Administration (NASA) and the European Space Agency (ESA), known as the Mass Change And Geosciences International Constellation (MAGIC), aims to launch a new paired mission, i.e., GRACE-C and NGGM (Next Generation Gravity Mission), to improve the monitoring of extreme events such as floods and droughts. The primary objective of this study is to examine the impact of the expected improvement in the spatial-temporal resolution and accuracy of NGGM and MAGIC on precipitation estimation by developing multiple synthetic experiments over Europe. The study employed the well-known SM2RAIN algorithm to estimate the precipitation accumulated between two consecutive TWS measurements. The total amount of water in the soil from the fifth generation ECMWF reanalysis for the land (ERA5L) is used as a proxy of TWS over the period of 2003-2012, while ERA5L precipitation serves as a reference dataset. The results showed that SM2RAIN exhibited satisfactory performance at a daily temporal resolution, with a mean correlation coefficient of 0.86. Good agreement was obtained across most of Europe except in some areas of northern Italy, northeastern Europe (Estonia, Latvia) and some coastal regions. Secondly, synthetic experiments were developed by degrading the temporal resolution (5–10, 15, and 30 days) and additively adding Gaussian errors to TWS data ranging from 1 to 40 mm. The results showed robust performance under moderate temporal degradation (5-10 days) and low measurement errors (<5 mm), with correlations generally exceeding 0.75. However, performance deteriorates significantly when errors exceed 10-20 mm, with correlations dropping below 0.40 for an error of at 42 mm error levels. These findings identify critical design thresholds for NGGM/MAGIC missions, demonstrating that achieving a measurement accuracy better than 5-10 mm is crucial for reliable precipitation estimation across diverse European hydroclimatic conditions.

Keywords: Precipitation estimation, SM2RAIN, TWS, NGGM, MAGIC, Synthetic experiments

Formatted: Font color: Dark Red

Formatted: Font color: Dark Red

Formatted: Font color: Dark Red

Formatted: Font color: Dark Red

Formatted: Font color: Dark Red

Formatted: Font color: Dark Red

1. Introduction

Precipitation is one of the primary components of the hydrological cycle, the precise estimation of spatio-temporal distribution of precipitation plays a vital role in assessing extreme events (Vrochidou et al., 2013), water resources management (Dorigo et al., 2021), ecological assessments (Yu et al., 2024) and hydrological research (McMillan et al., 2011; Meles et al., 2024).

Primary methods to estimate precipitation include ground observations (rain gauges and radar), reanalysis datasets and satellite based products (Song et al., 2020; Yuan et al., 2020). Although rain gauges are marked as one of the reliable sources of precipitation measurement, their spatial coverage, especially in mountainous and data-scarce regions, remains a primary limitation for their applicability on a global scale (Ahrens, 2006; Herrnegger et al., 2018; Kidd et al., 2017). Meteorological radar provides high-resolution spatial estimates of precipitation. However, it is constrained by limited accessibility and high operational costs (Sun et al., 2018). Reanalysis datasets such as the 5th reanalysis from the European Centre for Medium-Range Weather Forecasting (ECMWF), ERA5, have been developed and widely used from regional to global scales. Nevertheless, the accuracy of the reanalysis product depends upon the multi-source data, including in-situ observations and satellite products, which constrains its applicability in a data scarce mountainous environment (Ansari et al., 2022; Li et al., 2020; Liaqat et al., 2022). Satellite based precipitation products with their high spatial and temporal coverage, provide an alternative solution for precipitation estimation.

There are two primary methods for precipitation estimation from satellite, including top-down and bottom-up approaches (Brocca et al., 2014). In the top-down approach, precipitation is based on the emitted or reflected radiation from clouds or rain droplets derived from a combination of Geostationary (GEO) and Low Earth Orbiting (LEO) satellites sensors (Levizzani et al., 2020). In the bottom-up approach, precipitation is computed by surface soil moisture (SM) observations inverting the soil water balance in the so-called SM2RAIN algorithm (Brocca et al., 2013, 2014). A key difference between two approaches is that the top-down provides instantaneous measurements of precipitation which can be underestimated when the satellites do not pass over the precipitation event while the bottom-up approach gives accumulated precipitation between two subsequent SM measurements.

SM2RAIN has already been implemented from regional scale in South Asia (Lai et al., 2022) to global scale (Brocca et al., 2019), in Africa (Hengl et al., 2021; Islam, 2022), America (Paredes-Trejo et al., 2019; Satgé et al., 2020), Australia (Chua et al., 2022) and Europe (Moges et al., 2022). Various SM2RAIN products have been developed by forcing different SM satellite observations into the algorithm, including ASCAT (Advance Scatterometer (Brocca et al., 2019), Soil Moisture and Ocean Salinity (SMOS) (Brocca et al., 2016), Advanced Microwave Scanning Radiometer 2 (AMSR2) (Tarpanelli et al., 2017), Sentinel-1 (Filippucci et al., 2022).

In this study, SM is replaced for the first time with terrestrial water storage (TWS) to estimate precipitation between two consecutive measurements. SM2RAIN based on surface SM relies on top few centimeters of soil. This layer is quite sensitive to dense vegetation, frozen soils, and snow-covered locations, and it saturates quickly after intense precipitation. Gravimetry, on the other hand, observes TWS integrating water variations over the full soil column and shallow subsurface. This makes it less sensitive to surface saturation and vegetation impacts; however, the detection of precipitation signals in gravimetric observations is directly constrained by measurement uncertainty. As

70 precipitation represents the primary source of TWS, its variations inherently reflect cumulative water inputs and losses
at larger spatial scales (Zhong et al., 2025). These characteristics make TWS a suitable choice, potentially better than
SM, for estimating accumulated precipitation at sub monthly scales. However, this potential has remained largely
unexplored because TWS estimates derived from previous gravity missions, including the Gravity Recovery and
Climate Experiment (GRACE, Tapley et al., 2004), Gravity Field and steady-state Ocean Circulation Explorer (GOCE,
Drinkwater et al., 2003), and GRACE Follow-On (GRACE-FO, Landerer et al., 2020), are characterized by coarse
75 spatial (~400 km) and, importantly, temporal (monthly) resolutions, which limit their ability to accurately capture
rainfall variability, except in regions experiencing intense precipitation, such as the tropics. Currently, daily-scale TWS
estimates can only be obtained through the integration of gravimetric observations with hydrological or land surface
models (Croteau et al., 2020). Thus, it is indispensable to work on high temporal resolution satellite-based solutions
from daily to weekly scales to improve our capability to estimate global scale precipitation.

Formatted: Font color: Dark Red

Formatted: Font color: Dark Red

80 A joint collaboration between the National Aeronautics and Space Administration (NASA) and the European Space
Agency (ESA), known as the Mass change And Geosciences International Constellation (MAGIC, Daras et al., 2023),
was initiated to advance gravity-based monitoring of mass transport within the Earth system. The ultimate objective
of this collaboration is to enhance current observing capabilities through next-generation high-resolution gravity
missions, thereby improving the monitoring of extreme events such as natural hazards, droughts, and floods. The
85 European Next-Generation Gravity Mission (NGGM, Daras et al., 2024; Haagmans and Tsaoussi, 2020), a component
of MAGIC, is presently undergoing Phase A Extension activities as the first mission opportunity under ESA's Future
EO Program. Coordinated with the NASA-led GRACE-C component, NGGM is designed to drastically improve both
the temporal sampling and effective spatial resolution of TWS observations compared with the GRACE and GRACE-
FO missions. Current mission studies indicate a target temporal sampling on the order of sub-weekly days and an
90 effective spatial resolution approaching approximately 100 km at regional scales, for equivalent water height
accuracies of approximately 5–10 mm under target performance conditions (ESA, 2023). Combined with further
improvements in orbit and attitude modelling, these advances are expected to enable the generation of fast track gravity
products on sub-weekly time scales, reducing temporal aliasing while enhancing signal-to-noise characteristics.

Formatted: Font color: Dark Red


Based on that, the objectives of this study are: 1) to examine the performance of SM2RAIN algorithm to estimate
95 precipitation using TWS data as input, 2) to investigate the impact of the improved accuracy and spatiotemporal
resolution of NGGM and MAGIC on precipitation by using the SM2RAIN algorithm on a European scale, and 3) to
analyse the impact of different synthetic mission configurations by adding Gaussian error into simulated TWS
observations used as input to the SM2RAIN algorithm. It should be noted that the purpose of this study is not to
perform an operational validation of precipitation estimates, but rather to conduct synthetic experiments to assess the
100 feasibility of retrieving precipitation from TWS and evaluating the sensitivity of the SM2RAIN algorithm to the
temporal sampling and accuracy expected from future NGGM and MAGIC gravity missions.

2. Material and methods

2.1 Study area

Formatted: Space After: 0 pt, Outline numbered +
Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 +
Alignment: Left + Aligned at: 0" + Indent at: 0.25"

105 The study focuses on the European region, specifically the area with latitudes 30° to 60°N and longitudes 10°W to
350°E (see Fig. 1). This extensive domain is characterized by diverse climatic zones, each playing a critical role in
shaping precipitation dynamics and hydrological processes. Specifically, according to the Köppen-Geiger climate
classification (Beck et al., 2023), the European region exhibits significant climatic diversity: the Oceanic Climate
(Cfb), mainly evident in Western Europe along the Atlantic coastline and largely regulated by the proximity to the
110 prevailing westerlies and ocean. It is characterized by mild temperatures and relatively consistent high precipitation
throughout the year. In contrast, the Continental Climate (Dfb, Dfc), largely prevails in Central and Eastern Europe
where seasonal extremes especially for temperature are more pronounced, characterised by cold winters and warm
summers. Precipitation is generally moderate with peak eventevents in summer. The Mediterranean Climate (Csa,
Csb) is distinguished by hot dry summers and mild wet winters which largely dominatesdominate in Southern Europe
115 and in the Mediterranean Basin. Further, the Polar and Subarctic Climates (Dfc, ET) are experienced in Northern
Europe including Scandinavia, and high-altitude regions such as the Alps featured by long harsh winters and short
summers. Precipitation is generally low but persistent, with significant amounts of snowfall during the winter, feeding
into glacial and snowmelt-driven hydrological processes.



120 Figure 1. (a) Study domain covering the European region used in this analysis. The spatial grid corresponds to the
ERA5L dataset resampled to approximately 100 km spatial resolution. The analysis period spans 2003–2012,
corresponding to 3653 daily time steps used in the SM2RAIN experiments. The base map is derived from
OpenStreetMap data © OpenStreetMap contributors (available under the Open Database License). (b) Köppen–Geiger
climate classification across the study domain used to assess the influence of climatic conditions on the performance
125 of precipitation estimation from terrestrial water storage signals. The major climate classes include arid (B),
Mediterranean climates (Csa, Csb), temperate oceanic climates (Cfb), cold continental climates (Dfb, Dfc), and tundra
climates (ET).

2.2 ERA5L Soil Moisture and Precipitation data

The study utilized the volumetric soil moisture content and the total precipitation available from the European
Reanalysis 5th Generation ERA5L which is the land component of ERA5, the global reanalysis product developed by
130 ECMWF (Muñoz-Sabater et al., 2021). In particular, the volumetric soil moisture content (m³/m³) at four soil depth

Formatted: Font: 10 pt, Not Bold, Underline, Font color: Red

Formatted: Justified, Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt, Font color: Red

Formatted: Font: 10 pt, Not Bold

Formatted: Font:

Formatted: Font:

Formatted: Font:

layers (i.e., Layer 1, 0–7 cm, Layer 2, 7–28 cm, Layer 3, 28–100 cm, Layer 4, 100–289 cm) were utilized to estimate TWS. ERA5L based TWS is highly correlated with GRACE based TWS as shown in Fig S1, thus it is a reliable proxy for TWS estimation. Although this representation does not explicitly include other storage components such as snow water equivalent, groundwater, surface water, or canopy interception, which are part of the total storage measured by GRACE and GRACE-FO. However, for the selected European domain in this study, the contribution of some of these components is relatively limited for the purposes of the present sensitivity experiments. In particular, the contributions of canopy interception and surface water storage are generally small at the spatial scale considered (100 km). Further, snow water equivalent may drive storage variability locally in northern or mountainous areas, it likely contributes little to storage changes across most of our study domain during the study period. ERA5L does not explicitly represent groundwater storage, and thus groundwater cannot be included in the proxy data set used to drive the synthetic experiments. Since the goal of this study is to find out how possible and sensitive, it is to estimate the precipitation under future gravity mission configurations. By isolating the soil moisture driven component of TWS, it allows for a clearer attribution between precipitation forcing and storage response. The TWS from ERA5L is estimated by calculating the product, layer by layer, between the soil moisture content and the layer thickness. The product facilitates the conversion of volumetric moisture content into the depth of water stored within the soil column. The final TWS estimate in mm can be obtained by adding the water stored in all four layers. The equation used to calculate TWS is:

$$TWS = (SM_1 * 0.07) + (SM_2 * 0.21) + (SM_3 * 0.72) + (SM_4 * 1.89) * 1000$$

(1)

where SM_i depicts volumetric soil moisture content in (m^3/m^3) for the i -th layer. The constants (0.07, 0.21, 0.72, and 1.89) are the thicknesses in meters of the corresponding soil layers. The calculation provides the TWS in the soil column in meters of water equivalent. Although this ERA5L based TWS representation does not explicitly include additional storage components such as snow water equivalent, groundwater, surface water, or canopy interception - components included in the total signal observed by GRACE and GRACE-FO - it shows a strong correlation with GRACE based TWS (Fig S1). This agreement can be explained by the fact that, over the selected European domain, the omission of these components is expected to have a limited impact on the present sensitivity experiments. In particular, canopy interception and surface water storage are generally negligible at the ~100 km scale, while snow water equivalent may locally influence storage variability in northern and mountainous regions but has a limited overall contribution during the study period.

ERA5L total precipitation was considered as a benchmark for performance assessment of SM2RAIN algorithm. ERA5L is available at 0.1° and hourly temporal resolution from 1950 to present. In this study, the analysis period was constrained to the timeframe between 2003 and 2012, and both TWS and precipitation were resampled on a common 100 km grid using bilinear interpolation at a daily scale. This smooths out small scale spatial variability, particularly in mountainous regions like the Alps or the Scandinavian mountains. Note that this study is not designed to reproduce the full spatial resolution of the GRACE and GRACE-FO observations but instead explores the sensitivity of the precipitation retrieval to temporal sampling and measurement uncertainty. To support this choice, we conducted a set

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font:

Formatted: Font:

Formatted: Font:

Formatted: Justified, Line spacing: 1.5 lines, Tab stops: 6.5", Right

of scale consistency and sensitivity analyses (Fig. S1 and Fig. S2). The results demonstrate that resampling to a common analysis grid preserves gravity relevant information and does not introduce artificial small scale signals, thereby supporting the methodological approach adopted in the synthetic experiments.

2.3 The SM2RAIN algorithm

A modified version of the SM2RAIN algorithm, i.e., using TWS instead of soil moisture, was applied to estimate precipitation between two successive TWS measurements in the time interval dt , by inverting soil water balance equation as follows:

$$Z^* \frac{dTWS^*(t)}{dt} = p(t) - r(t) - e(t) - g(t) \quad (2)$$

Here Z^* [mm] represents the water capacity of the soil layer, calculated as the product between the depth of the soil layer and its porosity. The term TWS^* refers to the relative TWS [-] (i.e., ranging between 0 and 1), t [t] is the time, p is the precipitation rate [mm T^{-1}], $r(t)$ is the surface runoff rate [mm T^{-1}], $e(t)$ is the evaporation rate [mm T^{-1}] and $g(t)$ is the drainage rate [mm T^{-1}]. During precipitation events, evaporation and surface runoff are assumed negligible (Brocca et al., 2015). The drainage rate is dependent on TWS, which is derived from Famiglietti and Wood's (Famiglietti and Wood, 1994). Thus, the precipitation rate can be estimated by simplifying Eq. (2) as follows:

$$p(t) = Z^* \frac{dTWS^*(t)}{dt} + as(t)^b \quad (3)$$

Where a (mm T^{-1}) and b (-) refer to the saturated hydraulic conductivity and the exponent of the (Famiglietti and Wood, 1994) equation, respectively. Calibration is performed independently at each grid cell (point-by-point) using a gradient-based optimization function. The procedure estimates five parameters: Z^* , a , b , T_{pot} , and T_{base} . The estimation of parameters a , b , and Z^* is needed to compute the precipitation between two consecutive TWS^* measurements. The filter parameters (T_{pot} and T_{base}) have narrow interquartile ranges, indicating consistent noise reduction across grid cells. Before running the algorithm, the noise in the TWS signal is reduced by applying the exponential filter as described by Brocca et al. (2019), where the characteristic time length parameter (T) is modeled as a function of soil moisture using a two-parameter power law relationship. The values of the two parameters (T_{pot} and T_{base}) of the modified exponential filter equation, and the a , b , and Z^* parameter values are estimated through calibration against the ERA5L precipitation dataset. The parameter bounds are explicitly defined in the code as follows: Z^* ranges between 10 and 1200, a between 0 and 160, b between 1 and 50, T_{pot} between 0.05 and 0.75, and T_{base} between 0.05 and 3.0. Optimization is carried out using MATLAB's `fmincon` function with the active set algorithm, allowing a maximum of 300 iterations and 500 function evaluations. The calibration is carried out by minimizing an objective function based on the Nash–Sutcliffe efficiency (NS, Nash and Sutcliffe, 1970), computed between simulated and reference precipitation (e.g., ERA5L).

The Nash–Sutcliffe efficiency (NS) index. To evaluate the spatial consistency and the robustness of the calibration, we computed summary statistics of the calibrated parameters across the European domain. These values correspond to unrealistic parameter values across other parameterizations, e.g., mountainous or cold regions. The distribution of calibrated parameters (Table S1, Supplementary Material) supports the occurrence of substantial parameter variability and equifinality within this calibration framework. The storage capacity parameter " Z^* " ranges from 38.85 to 1008.70.

Formatted: Font:

with a median value of 506.80 and an interquartile range (IQR) of 432.67–584.10. These values are substantially larger than those typically reported for surface soil moisture based SM2RAIN applications, reflecting the deeper and more integrated nature of terrestrial water storage signals derived from gravimetry. The detail information about these calibrated parameters can be found in Table S1.

Regarding the validation strategy, the current implementation uses the full available time series at each grid cell for both calibration and performance assessment. An explicit temporal train/validation split or spatial cross validation was not applied. This design choice is consistent with the primary objective of the study, which is to conduct a controlled synthetic sensitivity assessment of precipitation retrieval performance under different NGGM/MAGIC sampling and error scenarios, rather than to provide operational out of sample validation of a precipitation product.

2.4 Setup of the synthetic experiments

The study employed the SM2RAIN algorithm to estimate the precipitation accumulated between two consecutive TWS measurements (Brocca et al., 2014). The study considered three types of experiments: i) feasibility experiments, in which ERA5L based TWS were used as a proxy for perfect storage retrievals and to evaluate the capability of SM2RAIN to reconstruct precipitation from storage variations; ii) synthetic experiments simulating the expected sampling and error characteristics of NGGM/MAGIC missions, iii) reference scenarios, which represent GRACE/GRACE-FO-like temporal sampling conditions.

As a first step, the ERA5L based TWS at daily time scale was used as input to SM2RAIN, while ERA5L precipitation was used as reference. This step was carried out to check the reliability of SM2RAIN in estimating precipitation from TWS. The simulated precipitation was aggregated to 15 and 30-day scales for comparative analysis and clearer visualization of performance. The choice to use ERA5L as a self consistent “proxy truth” follows established methodology used in gravity mission design and Phase A simulation studies for GRACE, GRACE-FO, MAGIC, and NGGM (e.g., Tapley et al., 2004; Landerer et al., 2020; Haagmans and Tsaoussi, 2020; Daras et al., 2023). Synthetic mission assessments typically rely on a single physically coherent dataset (model or reanalysis) to ensure compatibility between water balance terms and to isolate the effects of instrument accuracy and sampling strategy. Accordingly, ERA5L was used not for operational validation but to provide a controlled environment for evaluating the relative performance of different NGGM/MAGIC configurations.

In a second step, synthetic experiments were conducted by degrading daily TWS time series to 5-day intervals, representing the temporal sampling anticipated for future NGGM/MAGIC missions. Temporal resampling is implemented using a moving average aggregation scheme, followed by aggregation to 15-day and 30-day scales for clearer visualization of performance. A fixed window (e.g., five samples for the 5-day scenario) is applied to both the TWS signal and its time axis. The smoothed values are then embedded between the first and last original observations to preserve endpoints, and the resulting series is linearly interpolated back to the original daily time grid. This ensures temporal alignment while emulating reduced mission sampling frequency. This design choice makes sure that the experiments are consistent with each other and lets the analysis separate the effects of temporal sampling and measurement uncertainty in the TWS signal. As reference, the simulated precipitation generated from the first step was used. Although systematic errors from the initial step may carry over to the subsequent step, the purpose of the

Formatted: Justified, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Space After: 0 pt, Line spacing: 1.5 lines, Keep with next

Formatted: Justified, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Justified, Line spacing: 1.5 lines

synthetic experiments is to assess the comparative sensitivity of precipitation retrieval to NGGM/MAGIC sampling configurations, rather than to provide an independent validation.

Further, Gaussian errors (1.9 mm, 4.2 mm, 19 mm and 42 mm) were incorporated into the resampled TWS data to assess model performance as a function of the expected error values of the NGGM/MAGIC mission (Daras et al. 2023). The Gaussian noise we introduce is a simplified error model. In reality, gravimetric mission errors may have spatial and temporal correlations, depend on orbital geometry of the mission, and vary regionally (e.g., with topography), but the purpose of this noise model is to assess the sensitivity of the precipitation retrieval to measurement uncertainty. Thus, the Gaussian assumption must be understood as a linear approximation of the measurement uncertainty rather than the complete expected error structure of future gravity missions. Noise is generated using a normal distribution and is assumed to be temporally and spatially uncorrelated, providing a controlled baseline for sensitivity analysis (Landerer et al., 2020; Daras et al., 2023).

The exponential filter described by Brocca et al. (2019) is subsequently used within the SM2RAIN framework to the resampled and noise-perturbed TWS signal prior to precipitation inversion. To provide a reference comparable to the capability of current satellite gravimetry missions, a GRACE-FO-like synthetic configuration was defined using approximately ~25 mm equivalent water height uncertainty and 30-day temporal sampling. This uncertainty level is within the typical gravimetric error levels described in the ESA NGGM Mission Requirements Document (ESA, 2023). This configuration does not use actual GRACE observations but instead mimics present-day gravimetry performance by degrading the ERA5L derived TWS signal. These error values are the expected uncertainty values from the NGGM mission for the short-term time scales and for spatial resolutions of 400 km and 800 km (see Table 11 in Daras et al., 2023).

In a third step, the combined effects of temporal aggregation and error level were assessed by considering temporal intervals of 5, 10, 15, and 30-days for TWS measurements and error levels ranging from 0 to 40 mm. The results were evaluated over 100 randomly selected grid points across Europe. The final step was carried out to investigate the potential improvements of NGGM and MAGIC missions with respect to GRACE\FO mission. The obtained results were tested with a temporal aggregation of 15-day and 30-day, as having TWS measurements every 5-days and differentiating the signal, precipitation can be estimated reliably for a temporal aggregation 3 times the resolution of the input TWS measurements (Brocca et al., 2013).

For each experiment configuration model performance was evaluated using complementary metrics including bias (mm), correlation coefficient (R, -), and root mean square error (RMSE, mm). This approach is consistent with previous studies (e.g., Knoben et al., 2019; Williams, 2025), which recommend reserving skill scores (e.g., Nash Sutcliffe efficiency (NS) index) primarily for model calibration rather than performance evaluation.

Results and Discussion

3.1 SM2RAIN reliability using TWS measurements as input


Formatted: Justified, Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font color: Text 1

Formatted: Justified, Line spacing: 1.5 lines

275 The assessment of the SM2RAIN algorithm for estimating 15-day and 30-day precipitation using ERA5L based TWS
 is presented in Fig.2. The results ~~indicateshow~~ that SM2RAIN exhibited satisfactory performance, with mean values
 of R and RMSE equal to 0.86 and 13.31 mm, respectively, for the 15-day and equal to 0.86, 21.06 mm for the 30-day
 temporal assessment ~~against ERA5L precipitation~~. The mean value of BIAS indicated a slightly underestimation at
 15-days (-0.31 mm) with respect to 30-day (-0.23 mm). Significant negative values of BIAS were observed across
 280 most of Europe, with the strongest underestimation- noted in ~~the~~-central and northeastern ~~Europe~~~~regions~~. Although
 the Mediterranean ~~region~~ and southern Europe also exhibits negative bias ~~values~~, their ~~impact impact~~ is generally less
 intense than ~~Northern northern~~ regions. Based on the statistical analysis, SM2RAIN-simulated precipitation shows
 good agreement across most of Europe, with exceptions in the Alpine areas in northern Italy, in the northeastern states
 (Estonia, Latvia) and in the coastal regions of Norway. ~~The lower performance in these regions can be attributed~~
 285 ~~several hydrological factors. For instance, the strong orographic precipitation gradients and pronounced subgrid~~
~~variability of mountainous regions such as the Alps are hard to represent at the 100 km spatial resolution used in this~~
~~study. In northern latitudes (for example, Estonia, Latvia and coastal Norway), seasonal snow accumulation and melt~~
~~introduce lags between precipitation input and changes in terrestrial water storage that also weaken the direct~~
~~relationship assumed by the SM2RAIN inversion. In coastal areas, maritime precipitation variability and groundwater~~
 290 ~~dynamics may further complicate the precipitation storage relationship (Trautmann et al., 2018).These process based~~
~~explanations are consistent with the climate stratified analysis shown in Tables S2–S3, which shows that model skill~~
~~is lower in the colder, more continental, snow dominated and continental climate regimes.~~

The temporal plot of mean simulated and observed precipitation for 15-day aggregation during 2003-2012 further
 supports -the results discussed above, as shown in Fig.3. The consistent overlap of both low and high precipitation
 295 limbs supports the consistency of the simulations in capturing the observed variability. ~~The performance obtained in~~
~~this study (R = 0.86 for the 15-day aggregation) is comparable to, and in some cases higher than, correlations typically~~
~~reported in previous SM2RAIN applications based on satellite soil moisture observations, where correlations with~~
~~reference precipitation datasets generally range between 0.6 and 0.8 at regional to global scales (e.g., Brocca et al.,~~
~~2019).~~ Overall, these results pave the way for assessing the suitability of SM2RAIN to simulate precipitation for the
 300 first time using TWS, and they support the development of synthetic experiments ~~for precipitation estimation for under~~
 future gravity missions.



Formatted: Font color: Red




Figure 2: Performances of SM2RAIN algorithm forced with ERA5L based TWS against the ERA5L precipitation data over Europe for the period 2003–2012. The two columns refer to 15-day and 30-day aggregation period., rows indicate the correlation coefficient, R (top panel), BIAS (middle panel) and root mean square error, RMSE (lower panel).




Figure 3: Mean value of precipitation for 15-day aggregation as estimated from TWS through SM2RAIN and obtained from ERA5L precipitation

Additional analyses were conducted to examine the sensitivity of the SM2RAIN-based precipitation estimates to regional hydroclimatic conditions and rainfall intensity (Tables S2–S3). The results show that model performance varies across Köppen–Geiger climate classes, with the highest correlations observed in Mediterranean and temperate oceanic climates, where precipitation driven storage variations dominate the hydrological response. Lower

Formatted: Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt


performance is observed in colder continental and high latitude climates, likely due to the influence of snow accumulation, freeze thaw processes, and delayed storage responses.

Furthermore, the analysis improved by precipitation intensity indicates that the method performs significantly better for higher intensity precipitation events. This behavior can be explained by the fact that high intensity rainfall events generate larger and more rapid changes in terrestrial water storage, which are more clearly detectable above background noise and other hydrological processes such as evaporation and drainage. In contrast, low intensity precipitation events often produce weak storage signals that can be masked by measurement of uncertainty and competing fluxes, leading to reduced correlation with the reference dataset. These findings highlight the potential value of next generation gravity missions such as NGGM and MAGIC, whose improved temporal resolution and measurement accuracy are expected to enhance the detectability of short term storage changes and improve precipitation estimation, particularly for hydrological extremes.

3.2 Performance of NGGM/MAGIC synthetic TWS for precipitation estimation

In this section, synthetic experiments were designed to assess the performance of the NGGM/MAGIC mission configurations. SM2RAIN derived precipitation during the evaluation stage is used as a reference, while ERA5L based TWS aggregated at 5 day interval was used as a input. Measurement errors of 1.9 and 4.2 mm (target error) and 19 and 42 mm (threshold errors) were introduced following Daras et al. (2023).

The temporal resampling of data to a 5-day interval generally maintains a good agreement with the reference precipitation data as evident from all statistical measures with R, RMSE, and BIAS values of 0.95, 9.45 mm, and -0.21, respectively (see Fig. 4, No Error plot). Stronger correlation (R values above 0.88) were observed in the southern Europe and parts of Mediterranean regions. A similar pattern was noted for RMSE values with lower values in southern areas and moderate to higher RMSE values in northern and central Europe. The spatial average of BIAS exhibited widespread underestimation, with slight overestimation in the Mediterranean.



Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 8 pt, Bold, Do not check spelling or grammar

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: 8 pt




Figure 4: Performance of SM2RAIN algorithm as a proxy of the future NGGM mission forced with ERA5L based TWS at 5 days resolution over Europe for the period 2003-2012 and introducing an error from 0 mm to 42 mm for a 15-day aggregation period. The top panels shows the correlation coefficient, the middle panels the BIAS, and the lower panels the root mean square error, RMSE.

Introducing a target error from 1.9 mm to 4.2 mm reduced model's performance as reflected by correlation of 0.91 (-4%) and 0.87 (-8%) with respect to the no error configuration, indicating that the reduction is limited when target error is considered. The values of RMSE increased more significantly to 10.20 mm (+8%) and 12.43 mm (+32%) suggesting that the absolute accuracy begins to depicts measurable degradation even with smaller error values while the temporal agreement, measured by R, remains preserved. Spatially, both errors keeps higher correlations ($R > 0.80$), consistent error distribution ($RMSE < 12$ mm) across most of Europe, with slightly less performance in the Eastern European and Scandinavia. Further, the values of BIAS were found stable with mean values significantly low and equal to -0.01 mm and 0.04 mm using 1.9 mm and 4.2 mm error, respectively. The spatial average of BIAS values

indicated a tendency towards underestimation over majority of areas with slightly overestimation in the Mediterranean region. This suggests that low values of errors do not induce systematic algorithmic biases while preserving the optimal model operation to simulate precipitation.

At threshold error (19 mm to 42 mm), lead to a substantial degradation of model performance with mean R values decreasing to 0.63 (-34%) and 0.32 (-66%) and RMSE increasing to 20.99 mm (+121%) and 24.73 mm (+162%). The spatial pattern indicates a poor relationship between simulated precipitation and reference precipitation ($R < 0.60$ & $RMSE > 20$ mm) in several areas (Northern Scandinavia, scattered areas of Eastern Europe, the Mediterranean region, parts of Alps and surrounding mountainous regions). The results demonstrate that larger errors make it extremely difficult for the model to capture and represent meaningful precipitation patterns with 42 mm error level; reasonable performance were kept with 19 mm error level. Finally, the bias analysis illustrates interesting patterns at threshold error. Mean bias varies between 0.15 mm (19 mm) to 0.04 mm (42 mm). The spatial inspection demonstrates substantial positive bias under 19 mm error which is evident across southern and central Europe, while mix of both positive and negative bias can be evident under 42 mm indicating more complex patterns, suggesting that extreme errors induce spatially variable systematic distortions rather than uniform bias trends.

The model's performance was also evaluated by aggregating the simulated precipitation for a 30 day period, while keeping the same target and threshold errors, i.e. from 1.9 mm to 42 mm; results are shown in Table 1. Starting from the baseline without error, the model maintains good correlation and RMSE under low error conditions of 1.9 mm and 4.2 mm, where correlation decrease up 7% and RMSE increases of 22% (error level of 4.2 mm). As for 15-day precipitation estimation, the performance deteriorates significantly under large error (19 mm to 42 mm). It is also noted that model exhibited enhanced correlation stability under lower error range for the 30 day compared to 15 day aggregation period. This suggests that longer temporal aggregation helps to preserve the model's ability to capture precipitation dynamics by smoothing out short-term fluctuations.

Table.1 Performance of SM2RAIN algorithm as a proxies of the future NGGM/MAGIC mission forced with ERA5L based TWS at 5 days resolution over Europe for the period 2003-2012 and introducing an error of 1.9 mm to 42 mm (top to bottom in rows) for 30-day aggregation period.

	R (-)	BIAS (mm)	RMSE (mm)
No Error	0.95	-0.18	14.44
1.9 mm	0.92	-0.04	14.98
4.2 mm	0.88	0	17.62
19 mm	0.66	0.09	30.92
42 mm	0.32	-0.01	37.17


Differently from R and RMSE, the BIAS remains quite stable in the different configurations. This stability is particularly noteworthy when compared to the spatial patterns observed in the 15-day analysis, where regional bias variations were more pronounced. The 30-day aggregation appears to further stabilize the algorithm's systematic

performance, likely due to the temporal averaging effect that reduces the impact of short-term measurement inconsistencies.

3.2.3 Performance of NGGM/MAGIC mission with respect to GRACE-FO

The impact of varying temporal resolution (5, 10, 15, 30 days) and error level (from 0 mm to 40 mm) was evaluated at eight randomly selected locations in Italy, Germany, Spain, Greece, Austria, Czechia, Denmark, and Estonia. The results covering the period from 2003 to 2012 for 30-day precipitation estimation are shown in Fig.5. The findings reveal that the temporal resolution has a significant impact on model performance but with varying patterns across error levels. Overall, better temporal resolutions (5-10 days) maintain high R and low RMSE values, even with increasing error, while lower temporal resolutions exhibit sharper drops in correlation as the error grows. However, the rate of degradation in the performance of simulated precipitation varies between countries. Generally, for error values larger than 20 mm, the performance is quite similar for the different temporal resolutions, particularly for the RMSE values. For lower error levels, the differences in the performance for the different temporal resolutions are much larger. This regional representation improves the robustness of the findings by highlighting the model's applicability across diverse climatic conditions.

Formatted: Indent: Left: 0.25", No bullets or numbering






Figure 5: Evaluation of the NGGM/MAGIC synthetic experiments at four temporal resolutions (5, 10, 15 and 30) with error levels ranging from 0 mm to 40 mm for estimating 30-day precipitation over random locations in Italy, Germany, Spain, Greece, Austria, Czechia, Denmark, Estonia for the period 2003-2012 in terms of correlation coefficient (R) and root mean square error (RMSE).

To obtain more general conclusions on the impact of error level and temporal resolution, another experiment was conducted as presented in Fig.6 Fig.6 and Fig. S4, where a subset of 100 grid cells randomly sampled from all valid land pixels (i.e., grid cells with complete TWS and precipitation records, without missing values) in the full European domain were the mean of 100 randomly selected points across Europe were examined similarly to what was done in Figure 5. To ensure reproducibility of the sampling procedure, the random selection was performed using a fixed random seed. No explicit spatial or climatic stratification was applied in this step, as the experiment was intended to provide a representative domain-wide sensitivity assessment rather than a formally stratified evaluation. The analysis revealed a linear decrease in the R-values as a function of error levels and again more stable performance for error levels larger than 20 mm in terms of RMSE. By setting a threshold value for R to 0.7, above which the performance can be considered satisfactory, for 5-day temporal resolution data the results are satisfactory for error levels lower than 15 mm. For 30-day temporal resolution, the results are not satisfactory even for no error conditions. If we consider the expected error level of NGGM\MAGIC to be lower than 10 mm, we can foresee mean performance better than 0.78 (35 mm) in terms of R (RMSE). The error level of GRACE\FO was found equal to 25 mm and the temporal resolution is 30 days. The performances is equal to 0.40 (55 mm) in terms of R (RMSE). The expected improvements of NGGM\MAGIC with respect to GRACE are therefore highly significant.

Overall, this analysis demonstrated smoother degradation patterns compared to individual country analyses, suggesting that regional challenging areas (like Mediterranean regions) are moderated when averaged across diverse European conditions. The findings also support the 5-10 days temporal resolution which offers optimal performance for precipitation estimation at regional scale. The improvement in precipitation estimation with high temporal resolution aligns with previous scientific expectations that shorter revisit times reduce temporal aliasing of storage signals. Similar findings have been reported in previous gravity based hydrological studies (Save et al., 2016; Scanlon et al., 2018), highlighting the significance of sub-monthly sampling for improving hydrological fluxes.




Figure 6: Evaluation of the NGGM/MAGIC synthetic experiments at four temporal resolutions (5, 10, 15 and 30) with error levels ranging from 0 mm to 40 mm for estimating 30-day precipitation over 100 random points in Europe for the period 2003-2012 in terms of correlation coefficient (R) and root mean square error (RMSE).

3. Limitations and Future Work

This study conducts a controlled sensitivity experiment utilizing ERA5L as a self-consistent synthetic reference framework; hence, the results reflect relative performance assessment rather than independent validation against in situ observations. Importantly, these experiments provide a first, controlled synthetic assessment of the feasibility of retrieving precipitation from TWS. The added value of our study is not to validate the SM2RAIN algorithm (which has been done extensively using multiple observational precipitation datasets such as e.g., GPCC or E-OBS) but rather to assess its sensitivity to the sampling and accuracy expected from future gravity missions.

Measurement uncertainties are modeled as additive Gaussian noise with standard deviations derived from NGGM mission error specifications (Daras et al., 2023). This notice acknowledges that real satellite gravimetry errors may be temporally and spatially correlated and may be geographically dependent, with factors including latitude, topography and orbit geometry. While the Gaussian noise assumption provides a simplistic error model, it is sufficient in providing a first-order sensitivity of how measurement uncertainty may affect precipitation retrieval. In this work, our aim is not to replicate the full complexity of mission error characteristics but rather to study the relative effect of spatiotemporal sampling and measurement uncertainty on precipitation estimation from TWS. However, this comparison would demand mission specific error covariance models, the derivation of which is beyond the scope of this study.

Additionally, the SM2RAIN framework does not show evapotranspiration as independent fluxes, which could affect how storage dynamics change over weeks. Neglecting evaporation can lead to biased precipitation estimates, particularly in warm, vegetated, and semiarid regions where evapotranspiration is significant. Further, TWS changes generally include contributions from groundwater, but the current framework does not make a clear

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font color: Dark Red

Formatted: List Paragraph, Justified, Indent: Left: 0.25",
Line spacing: 1.5 lines

distinction between groundwater storage and other hydrological components, limiting process-level interpretation.

Future work will address these issues by integrating flux-consistent water balance constraints (incorporating evapotranspiration and runoff), parameter estimation using machine learning, and improving the attribution of groundwater dynamics with future gravity missions. This preserves the physical SM2RAIN structure while reducing calibration dependency, improving spatial coherence, and enhancing robustness in data scarce and nonstationary environments. The framework will also be extended to larger domains using more realistic, spatially, and temporally correlated error structures derived from advanced mission simulations.

2.4. Conclusion

Future gravity missions could play a key role in addressing the challenges of interpreting different hydrological components, which remain difficult to capture with the current Gravity Recovery and Climate Experiment (GRACE) and its Follow-On (GRACE-FO) mission. In this study, we aim to address these challenges by providing a comprehensive assessment of upcoming Next-Generation Gravity Mission (NGGM) and Mass-change And Geosciences International Constellation (MAGIC) for precipitation estimation on a European scale. The research applied the SM2RAIN algorithm by inverting the soil water balance equation and for the first time, considered Terrestrial Water Storage (derived from the four soil layers of ERA5L soil moisture as a proxy) for precipitation estimation. The following conclusions can be drawn from the present study:

SM2RAIN algorithm exhibited satisfactory performance to simulate precipitation using TWS data across most of Europe achieving a high correlation of 0.86 and a low RMSE of 13.31 mm for 15-day temporal resolution assessments. The performance of model varied regionally, with stronger correlations found in the Mediterranean region and in the southern Europe, while lower values were observed in the Alpine area in northern Italy, the northeastern states and the coastal regions of Norway, where snow processes, complex topography, and deeper storage dynamics likely affect the relationship between TWS variations and precipitation.

The verification of SM2RAIN leads to design NGGM/MAGIC configurations which clearly show how temporal resolution and error levels have an impact on the performance for precipitation estimation. The results indicate a high correlation range (0.88-0.61) and lower RMSE (5mm - 42 mm) with shorter temporal aggregation (5-days) compared to longer windows (e.g., 30-day sampling representative of GRACE-FO-like conditions 30 days), even when errors (0-20 mm) are added.

For error levels lower than 10 mm, the performance deterioration is limited; whereas for error levels larger than 15-20 mm the performance drop is highly significant. These results highlight the importance of reducing-minimizing the retrieval errors threshold in future gravity missions.

The use of TWS in respect to standard SM2RAIN applications based on satellite soil moisture observations (e.g. Brocca et al., 2014, Brocca et al., 2019), provides a more integrated representation of the terrestrial water cycle by combining contributions from soil moisture, groundwater, and surface water storage. While soil moisture measurements typically represent only the near-surface layer, gravity-based TWS observations capture deeper storage components and large-scale water mass redistribution. Moreover, precipitation estimation from TWS may be complementary to soil moisture based approaches for large spatial scales and over regions where surface soil moisture

490 observations are sparse and/or strongly impacted by local processes (e.g. ~~evaporation, vegetation dynamics~~densely vegetated areas and frozen soils).

The findings of this study provide critical implications for the development and execution of future gravity missions such as NGGM and MAGIC. The results suggest that TWS variations contain sufficient hydrological information to retrieve precipitation signals and that the TWS-based inversion approach could serve as a useful complement to soil moisture based precipitation estimation. They results strongly advocate the value of high temporal resolution (5-day or better) encompassing low error levels (less than 10 mm) to ensure precise precipitation estimation at both regional and global scale. The results also open new opportunities to estimate precipitation for monitoring hydrological processes from terrestrial water storage (TWS) measurements especially in data-scarce regions. By improving spatio-temporal resolution and accuracy of gravity missions we can enhance our capacity to estimate precipitation globally, with also complementary performance with respect to the use of soil moisture measurements. This important aspect will be further explored in future studies.

500 **Author's contribution**

LB and MUL designed the study; MUL conducted the analyses; MUL and LB interpreted the analyses; MUL wrote the manuscript draft; MUL, SC, GJ, FL and LB reviewed and edited the manuscript.

Data availability

505 All raw and processed data can be available by the corresponding authors upon request.

Competing interests

The contact author has declared that none of the authors has any competing interests

Acknowledgements

510 The authors acknowledge funding from the Italian Space Agency (ASI) “NGGM/MAGIC, a breakthrough in understanding Earth's dynamics” project (grant agreement No. 2023-22.HH.0.

515 **References**

Ahrens, B.: Distance in spatial interpolation of daily rain gauge data, Hydrol Earth Syst Sci, 10, 197–208, <https://doi.org/10.5194/hess-10-197-208>, 2006.

520 Ansari, R., Liaqat, M. U., and Grossi, G.: Evaluation of gridded datasets for terrestrial water budget assessment in the Upper Jhelum River Basin-South Asia, J Hydrol (Amst), 613, 128294, <https://doi.org/10.1016/j.ejrh.2024.102114>, 2022.

Beck, H.E., McVicar, T.R., Vergopolan, N. et al. High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections. Sci Data 10, 724 (2023). <https://doi.org/10.1038/s41597-023-02549-6>

525 Brocca, L., Moramarco, T., Melone, F., and Wagner, W.: A new method for rainfall estimation through soil moisture observations, Geophys Res Lett, 40, 853–858, <https://doi.org/10.1002/grl.50173>, 2013.

Formatted: Font:

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt, Ligatures: None

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

- Brocca, L., Ciabatta, L., Massari, C., Moramarco, T., Hahn, S., Hasenauer, S., Kidd, R., Dorigo, W., Wagner, W., and Levizzani, V.: Soil as a natural rain gauge: Estimating global rainfall from satellite soil moisture data, *Journal of Geophysical Research: Atmospheres*, 119, 5128–5141, <https://doi.org/10.1002/2014JD021489>, 2014.
- 530 Brocca, L., Massari, C., Ciabatta, L., Moramarco, T., Penna, D., Zuecco, G., Pianezzola, L., Borga, M., Matgen, P., and Martínez-Fernández, J.: Rainfall estimation from in situ soil moisture observations at several sites in Europe: an evaluation of the SM2RAIN algorithm, *Journal of Hydrology and Hydromechanics*, 201–209, <https://doi.org/10.1515/johh-2015-0016>, 2015.
- 535 Brocca, L., Pellarin, T., Crow, W. T., Ciabatta, L., Massari, C., Ryu, D., Su, C., Rüdiger, C., and Kerr, Y.: Rainfall estimation by inverting SMOS soil moisture estimates: A comparison of different methods over Australia, *Journal of Geophysical Research: Atmospheres*, 121, 12–62, <https://doi.org/10.1002/2016JD025382>, 2016.
- Brocca, L., Filippucci, P., Hahn, S., Ciabatta, L., Massari, C., Camici, S., Schüller, L., Bojkov, B., and Wagner, W.: SM2RAIN–ASCAT (2007–2018): Global daily satellite rainfall data from ASCAT soil moisture observations, *Earth Syst Sci Data*, 11, 1583–1601, <https://doi.org/10.5281/zenodo.3405563>, 2019.
- 540 Chua, Z.-W., Kuleshov, Y., Watkins, A. B., Choy, S., and Sun, C.: A comparison of various correction and blending techniques for creating an improved satellite-gauge rainfall dataset over Australia, *Remote Sens (Basel)*, 14, 261, <https://doi.org/10.3390/rs14020261>, 2022.
- Croteau, M. J., Nerem, R. S., Loomis, B. D., and Sabaka, T. J.: Development of a daily GRACE mascon solution for terrestrial water storage, *J Geophys Res Solid Earth*, 125, e2019JB018468, <https://doi.org/10.1029/2019JB018468>, 2020.
- 545 Daras, I., March, G., Pail, R., Hughes, C., Braitenberg, C., Güntner, A., Eicker, A., Wouters, B., Heller-Kaikov, B., and Pivetta, T.: Level-2a simulated gravity field solutions of ESA’s science support study to Mass change And Geosciences International Constellation (MAGIC) Phase AV 1.0., <https://doi.org/10.5880/icgem.2023.005>, 2023.
- 550 Daras, I., March, G., Pail, R., Hughes, C. W., Braitenberg, C., Güntner, A., Eicker, A., Wouters, B., Heller-Kaikov, B., and Pivetta, T.: Mass-change And Geosciences International Constellation (MAGIC) expected impact on science and applications, *Geophys J Int*, 236, 1288–1308, <https://doi.org/10.1093/gji/ggad472>, 2024.
- Dorigo, W., Dietrich, S., Aires, F., Brocca, L., Carter, S., Cretaux, J.-F., Dunkerley, D., Enomoto, H., Forsberg, R., and Güntner, A.: Closing the water cycle from observations across scales: Where do we stand?, *Bull Am Meteorol Soc*, 102, E1897–E1935, 2021.
- 555 Drinkwater, M. R., Floberghagen, R., Haagmans, R., Muzi, D., and Popescu, A.: VII: Closing session: GOCE: ESA’s first earth explorer core mission, *Space Sci Rev*, 108, 419–432, 2003.
- [ESA \(2023\). Next-Generation Gravity Mission \(NGGM\) Mission Requirements Document \(MRD\), Version 1.0., European Space Agency, Future Earth Observation Programme.](#)
- Famiglietti, J. S. and Wood, E. F.: Multiscale modeling of spatially variable water and energy balance processes, *Water Resour Res*, 30, 3061–3078, 1994.
- 560 Filippucci, P., Brocca, L., Quast, R., Ciabatta, L., Saltalippi, C., Wagner, W., and Tarpanelli, A.: High-resolution (1 km) satellite rainfall estimation from SM2RAIN applied to Sentinel-1: Po River basin as a case study, *Hydrol Earth Syst Sci*, 26, 2481–2497, <https://doi.org/10.5194/hess-26-2481-2022>, 2022.
- 565 Haagmans, R. and Tsaoussi, L.: Next Generation Gravity Mission as a Mass-change And Geosciences International Constellation (MAGIC) Mission Requirements Document, Earth and Mission Science Division, European Space Agency, <https://doi.org/10.5270/esa.nasa.magic-mrd.2020>, 2020.
- Hengl, T., Miller, M. A. E., Križan, J., Shepherd, K. D., Sila, A., Kilibarda, M., Antonijević, O., Glušica, L., Dobermann, A., and Haefele, S. M.: African soil properties and nutrients mapped at 30 m spatial resolution using two-scale ensemble machine learning, *Sci Rep*, 11, 6130, <https://doi.org/10.1038/s41598-021-85639-y>, 2021.

Formatted: Font: 10 pt

Formatted: Justified

Formatted: Font: (Default) Times New Roman, 10 pt

- 570 Hernegger, M., Senoner, T., and Nachtnebel, H.-P.: Adjustment of spatio-temporal precipitation patterns in a high
Alpine environment, *J Hydrol (Amst)*, 556, 913–921, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2016.04.068>,
2018.
- Islam, Z.: Soil loss assessment by RUSLE in the cloud-based platform (GEE) in Nigeria, *Model Earth Syst Environ*,
8, 4579–4591, <https://doi.org/10.1007/s40808-022-01467-7>, 2022.
- 575 Kidd, C., Becker, A., Huffman, G. J., Muller, C. L., Joe, P., Skofronick-Jackson, G., and Kirschbaum, D. B.: So, how
much of the Earth’s surface is covered by rain gauges?, *Bull Am Meteorol Soc*, 98, 69–78,
<https://doi.org/10.1175/BAMS-D-14-00283.1>, 2017.
- [Knoben, W.J., Freer, J.E. and Woods, R.A., 2019. Inherent benchmark or not? Comparing Nash–Sutcliffe and Kling–
Gupta efficiency scores. *Hydrology and Earth System Sciences*, 23\(10\), pp. 4323–4331. <https://doi.org/10.5194/hess-23-4323-2019>](https://doi.org/10.5194/hess-23-4323-2019)
- 580 Lai, Y., Tian, J., Kang, W., Gao, C., Hong, W., and He, C.: Rainfall estimation from surface soil moisture using
SM2RAIN in cold mountainous areas, *J Hydrol (Amst)*, 606, 127430, <https://doi.org/10.1016/j.jhydrol.2022.127430>,
2022.
- Landerer, F. W., Flechtner, F. M., Save, H., Webb, F. H., Bandikova, T., Bertiger, W. I., Bettadpur, S. V., Byun, S. H.,
585 Dahle, C., and Dobslaw, H.: Extending the global mass change data record: GRACE Follow-On instrument and
science data performance, *Geophys Res Lett*, 47, e2020GL088306, 2020.
- Levizzani, V., Kidd, C., Kirschbaum, D. B., Kummerow, C. D., Nakamura, K., and Turk, F. J.: Satellite precipitation
measurement, Springer, 2020.
- Li, Y., Wang, Q. J., He, H., Wu, Z., and Lu, G.: A method to extend temporal coverage of high quality precipitation
590 datasets by calibrating reanalysis estimates, *J Hydrol (Amst)*, 581, 124355,
<https://doi.org/10.1016/j.jhydrol.2019.124355>, 2020.
- Liaquat, M. U., Grossi, G., Hasson, S. ul, and Ranzi, R.: Characterization of interannual and seasonal variability of
hydro-climatic trends in the Upper Indus Basin, *Theor Appl Climatol*, 147, 1163–1184,
<https://doi.org/10.1007/s00704-021-03850-3>, 2022.
- 595 McMillan, H., Jackson, B., Clark, M., Kavetski, D., and Woods, R.: Rainfall uncertainty in hydrological modelling:
An evaluation of multiplicative error models, *J Hydrol (Amst)*, 400, 83–94,
<https://doi.org/10.1016/j.jhydrol.2011.01.026>, 2011.
- Meles, M. B., Goodrich, D. C., Unkrich, C. L., Gupta, H. V., Burns, I. S., Hirpa, F. A., Razavi, S., and Guertin, D. P.:
Rainfall distributional properties control hydrologic model parameter importance., *J Hydrol Reg Stud*, 51, 101662,
<https://doi.org/10.1016/j.ejrh.2024.101662>, 2024.
- 600 Moges, D. M., Kmoch, A., and Uemaa, E.: Application of satellite and reanalysis precipitation products for
hydrological modeling in the data-scarce Porijögi catchment, Estonia, *J Hydrol Reg Stud*, 41, 101070,
<https://doi.org/10.1016/j.ejrh.2022.101070>, 2022.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga,
605 M., Harrigan, S., and Hersbach, H.: ERA5-Land: A state-of-the-art global reanalysis dataset for land applications,
Earth Syst Sci Data, 13, 4349–4383, 2021. <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- ▲ Nash, J.E. and Sutcliffe, J.V., 1970. River flow forecasting through conceptual models part I—A discussion of
principles. *Journal of hydrology*, 10(3), pp.282–290. [https://doi.org/10.1016/0022-1694\(70\)90255-6](https://doi.org/10.1016/0022-1694(70)90255-6).
- ▲ Paredes-Trejo, F., Barbosa, H., and dos Santos, C. A. C.: Evaluation of the performance of SM2RAIN-derived
rainfall products over Brazil, *Remote Sens (Basel)*, 11, 1113, <https://doi.org/10.3390/rs11091113>, 2019.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

610 Satgé, F., Hussain, Y., Molina-Carpio, J., Pillco, R., Laugner, C., Akhter, G., and Bonnet, M.: Reliability of SM2RAIN precipitation datasets in comparison to gauge observations and hydrological modelling over arid regions, *International Journal of Climatology*, 41, <https://doi.org/10.1002/joc.6704>, 2020.

[Save, H., Bettadpur, S. and Tapley, B.D., 2016. High-resolution CSR GRACE RL05 mascons. *Journal of Geophysical Research: Solid Earth*, 121\(10\), pp.7547-7569, <https://doi.org/10.1002/2016JB013007>](#)

615 Song, X., Wu, P., Gao, X., Yao, J., Zou, Y., Zhao, X., Siddique, K. H. M., and Hu, W.: Rainwater collection and infiltration (RWCI) systems promote deep soil water and organic carbon restoration in water-limited sloping orchards, *Agric Water Manag*, 242, 106400, <https://doi.org/10.1016/j.agwat.2020.106400>, 2020.

Sun, Q., Miao, C., Duan, Q., Ashouri, H., Sorooshian, S., and Hsu, K.: A review of global precipitation data sets: Data sources, estimation, and intercomparisons, *Reviews of Geophysics*, 56, 79–107, <https://doi.org/10.1002/2017RG000574>, 2018.

620 [Scanlon, B.R., Zhang, Z., Save, H., Sun, A.Y., Müller Schmied, H., Van Beek, L.P., Wiese, D.N., Wada, Y., Long, D., Reedy, R.C. and Longuevergne, L., 2018. Global models underestimate large decadal declining and rising water storage trends relative to GRACE satellite data. *Proceedings of the National Academy of Sciences*, 115\(6\), pp.E1080-E1089, <https://doi.org/10.1073/pnas.1704665115>](#)

625 Tapley, B. D., Bettadpur, S., Watkins, M., and Reigber, C.: The gravity recovery and climate experiment: Mission overview and early results, *Geophys Res Lett*, 31, <https://doi.org/10.1029/2004GL019920>, 2004.

Tarpanelli, A., Massari, C., Ciabatta, L., Filippucci, P., Amarnath, G., and Brocca, L.: Exploiting a constellation of satellite soil moisture sensors for accurate rainfall estimation, *Adv Water Resour*, 108, 249–255, <https://doi.org/10.1016/j.advwatres.2017.08.010>, 2017.

630 [Trautmann, T., Koirala, S., Carvalho, N., Eicker, A., Fink, M., Niemann, C., and Jung, M.: Understanding terrestrial water storage variations in northern latitudes across scales, *Hydrol. Earth Syst. Sci.*, 22, 4061–4082, <https://doi.org/10.5194/hess-22-4061-2018>, 2018](#).

Vrochidou, A.-E., Tsanis, I. K., Grillakis, M. G., and Koutroulis, A. G.: The impact of climate change on hydrometeorological droughts at a basin scale, *J Hydrol (Amst)*, 476, 290–301, <https://doi.org/10.1016/j.jhydrol.2012.10.046>, 2013.

635 [Williams, G.P., 2025. Friends Don't Let Friends use Nash-Sutcliffe Efficiency \(NSE\) or KGE for Hydrologic Model Accuracy Evaluation: A rant with data and suggestions for better practice. *Environmental Modelling & Software*, p. 106665. <https://doi.org/10.1016/j.envsoft.2025.106665>](#)

Yu, Y., Hua, T., Chen, L., Zhang, Z., and Pereira, P.: Divergent changes in vegetation greenness, productivity, and rainfall use efficiency are characteristic of ecological restoration towards high-quality development in the Yellow River Basin, China, *Engineering*, 34, 109–119, <https://doi.org/10.1016/j.eng.2023.07.012>, 2024.

Yuan, S., Wang, Y., Quiring, S. M., Ford, T. W., and Houston, A. L.: A sensitivity study on the response of convection initiation to in situ soil moisture in the central United States, *Clim Dyn*, 54, 2013–2028, <https://doi.org/10.1007/s00382-019-05098-0>, 2020.

645 [Zhong, Y., Tian, B., Kim, H., Yuan, X., Liu, X., Zhu, E., Wu, Y., Wang, L. and Wang, L., 2025. Over 60% precipitation transformed into terrestrial water storage in global river basins from 2002 to 2021. *Communications Earth & Environment*, 6\(1\), p.53. <https://doi.org/10.1038/s43247-024-01967-7>](#)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt