

1 Sinking particle fluxes and biological carbon pump efficiency 2 in the Labrador Sea during a *Phaeocystis* bloom decline

3 Montserrat Roca-Martí^{1,2}, Madeline Healey¹, Colleen E. McBride³, Rachel Sipler^{3,4},
4 Emmanuel Devred⁵, Carolina Cisternas-Novoa⁶, Elisa Romanelli⁷, Kyoko Ohashi^{1,8}, Stephanie
5 S. Kienast¹

6 ¹Department of Oceanography, Dalhousie University, Halifax, NS, Canada

7 ²Physics Department, Universitat Autònoma de Barcelona, Cerdanyola del Vallès, Spain

8 ³Department of Ocean Sciences, Memorial University of Newfoundland, St. John's, NL, Canada

9 ⁴Bigelow Laboratory for Ocean Sciences, East Boothbay, ME, USA

10 ⁵Fisheries and Oceans Canada, Bedford Institute of Oceanography, Dartmouth, NS, Canada

11 ⁶Ocean Sciences Centre, Memorial University of Newfoundland, St. John's, NL, Canada

12 ⁷Institute of Environmental Engineering, Department of Civil, Environmental and Geomatic Engineering, ETH
13 Zurich, Zurich, Switzerland

14 ⁸Fisheries and Oceans Canada, Institute of Ocean Sciences, Sidney, BC, Canada

15 Correspondence to: Montserrat Roca-Martí (montserrat.roca.marti@uab.cat)

16 **Abstract.** The Labrador Sea is a key region for carbon dioxide uptake characterized by deep mixing during winter
17 that supplies nutrients to the upper water column and fuels extensive phytoplankton blooms in spring. Yet, the
18 efficiency by which organic carbon is exported from surface waters during these blooms, as well as their
19 contribution to carbon sequestration, remain poorly constrained. Here, we present an unprecedented number of
20 measurements of sinking export fluxes (particulate organic carbon, POC; and biogenic silica, bSi) collected in the
21 central Labrador Sea during a 2-week-long process study that observed the decline of a historically large
22 *Phaeocystis* bloom in spring 2022. This *Phaeocystis* bloom was unusually large and highly productive, extending
23 over more than half of the Labrador Sea for 6 weeks. During the late stages of the bloom, we found that POC
24 fluxes from the base of the euphotic zone to 500 m were variable but overall moderate to high (average of 8 ± 5
25 $\text{mmol C m}^{-2} \text{ d}^{-1}$). Nevertheless, evidence of shallow POC flux remineralization combined with the fact that POC
26 fluxes in the bloom were not higher than in a region sampled outside of the bloom (average of $13 \pm 3 \text{ mmol C m}^{-2}$
27 d^{-1}) suggested a limited role of *Phaeocystis* in carbon export. Large ($>51 \mu\text{m}$) particles collected using large
28 volume pumps presented relatively low bSi/POC ratios and, therefore, diatoms did not appear to have an important
29 ballasting role of *Phaeocystis*-derived material. Using in situ net primary production (NPP) rates, we determined
30 that 2 weeks after the bloom peak, only 6% of NPP was exported to 100 m below the euphotic zone. Three weeks
31 after the peak, this value increased to 30%, reflecting a decline in NPP while POC fluxes remained relatively
32 constant. However, when using satellite-derived NPP integrated from the bloom peak until its end, the overall
33 biological carbon pump (BCP) efficiency was 6%, indicating that this *Phaeocystis* bloom represented a low-
34 efficiency export system. We stress the importance of multiple observations of both NPP and POC export along
35 the bloom period for estimating meaningful BCP efficiencies. The results presented in this study provide a
36 foundation for comparisons with other datasets collected during this ship-based process study and autonomous
37 platforms present in the area during and beyond this study. These future efforts will provide the opportunity to
38 increase the observational period and further elucidate the mechanisms leading to the low BCP efficiency found
39 during the decline of this *Phaeocystis* bloom in the Labrador Sea.

Deleted: Institut de Ciència i Tecnologia Ambientals (ICTA-UAB)...

Deleted: of the bloom

Deleted: the total amount of

Deleted: that reached

Deleted: was only 6%

Deleted: However, 3

Deleted: of the bloom

Deleted: the

Deleted: had

Deleted: 29

Deleted: . The observed change was driven primarily by

Deleted: over the sampling period, as

Deleted:

Deleted: U

Deleted: of the bloom

Deleted: we obtained an

Deleted: of

Deleted: placing

Deleted: as

Deleted: BCP

Deleted: long-term

62 1. Introduction

63 The biological carbon pump (BCP) encapsulates a set of processes that remove carbon dioxide (CO₂) from the
64 atmosphere and sequester it in the deep ocean (Passow and Weber, 2025; Volk and Hoffert, 1985). In the surface
65 ocean, phytoplankton produce organic matter via photosynthesis, and a fraction of this organic matter is
66 subsequently transferred to depth through sinking and other processes. ~~Of the material exported below the~~
67 ~~euphotic zone, most is remineralized in the ocean interior, while only a small fraction reaches depths where CO₂~~
68 ~~can be stored for several centuries or longer. A very small fraction of exported carbon is buried in seafloor~~
69 sediments and thus removed from the atmosphere on geological timescales. While it is well understood that the
70 BCP plays a large role in climate regulation, there is low confidence in the magnitude and even the sign of
71 predicted near-future changes in carbon export fluxes and how these will in turn affect atmospheric CO₂ levels
72 (Henson et al., 2022). Better quantification of particulate matter fluxes in the ocean ~~and of the key parameters~~
73 ~~controlling them~~ is thus urgently needed.

74 Indeed, the various processes transforming and redistributing carbon between the surface and the deep ocean
75 remain poorly quantified. Assessments of carbon budgets below the euphotic zone typically have found that
76 organic carbon sources and sinks are not balanced (Baltar et al., 2009; Boyd et al., 1999; Burd et al., 2010;
77 Reinthaler et al., 2006; Steinberg et al., 2008). This mismatch has been attributed to methodological constraints,
78 the choice of parameters used to estimate budget terms, the exclusion of important midwater processes, and the
79 spatial and temporal scales over which measurements are integrated (Baumas et al., 2023; Giering et al., 2014;
80 Stephens and Roca-Martí et al., 2025).

81 Major uncertainties in the magnitude of biologically driven carbon ocean uptake and storage also lie in our
82 fragmented understanding of the different pathways by which the BCP exports particulate and dissolved organic
83 carbon (POC and DOC, respectively) from surface waters to the deep ocean. Advances in remote sensing, in situ
84 imagery, and the development of autonomous platforms (BGC-Argo floats and gliders) over the last 2 decades
85 have revealed that the BCP includes six major pathways for exporting organic carbon from the surface ocean to
86 the deep ocean: gravitational export (sinking), three pumps driven by physical processes (mixed layer, eddy
87 subduction and large-scale subduction), and two pumps driven by vertical migrations of zooplankton and larger
88 animals (Boyd et al., 2019; Claustre et al., 2021). Process studies and long-term observations of the BCP,
89 combining multiple approaches both ship- and autonomous-based, have been shown to be crucial for better
90 constraining ocean carbon budgets (Stephens and Roca-Martí et al., 2025).

91 The North Atlantic is responsible for a significant fraction of the global carbon export (~15%), driven by a
92 complex set of BCP processes (Sanders et al., 2014). The subpolar North Atlantic, specifically the Labrador Sea,
93 is an important region for CO₂ uptake (Arruda et al., 2024) characterized by deep vertical mixing during winter
94 that supplies nutrients to the upper water column and fuels extensive phytoplankton blooms in spring after the
95 water column has stratified (Tesdal et al., 2022). Yet, the efficiency by which organic carbon is exported from
96 surface waters during these blooms as well as their contribution to carbon sequestration remain poorly constrained,
97 partly because of the complex physical processes in the region (Baker et al., 2022) and the very limited POC flux
98 observations made so far (e.g., Lemaitre et al., 2018). In addition, impacts of global warming in the physical

Deleted: Most o

Deleted: is

Deleted: decomposed and respired

Deleted: remains

Deleted: until it is brought back to the surface ocean by physical transport

105 conditions of the Labrador Sea have already been reported, including freshening and weaker winter convection
106 (Yashayaev, 2024), which add uncertainty to the future state of the BCP and its downstream effects on climate
107 and marine ecosystems. Therefore, our new observations will provide an important baseline for constraining POC
108 export in the Labrador Sea under rapidly changing hydrographic and ecological conditions.


109 The phytoplankton assemblages in the Labrador Sea are typically dominated by diatoms, *Phaeocystis* spp. or
110 mixed populations (Devred et al., 2024). Observations of *Phaeocystis* blooms were previously restricted to shelf
111 and slope regimes, but unprecedented large *Phaeocystis* blooms have been reported in the Labrador basin in recent
112 years (in 2015 and 2022; Devred et al., 2025). Here, we present measurements of net primary production (NPP)
113 and sinking export fluxes (POC and biogenic silica, bSi) in the central Labrador Sea during the decline of a
114 historically large *Phaeocystis* bloom in spring 2022. The novelty of our ship-based process study (Biological
115 Carbon Export in the Labrador Sea, BELAS-1) lies in the high temporal resolution of in situ measurements
116 collected over 2 weeks which shed light on the efficiency of the gravitational sinking pump of *Phaeocystis* blooms.

117 2. Methods

118 Samples were collected in the Labrador Sea from 19 May to 2 June 2022 during the BELAS-1 expedition
119 (CE22009, RV *Celtic Explorer*). Three regions were targeted (Fig. 1): a grid of nine stations (hereafter "Eastern
120 Grid", ~1075 km²), where an extensive *Phaeocystis pouchetii* bloom was in decline (Devred et al., 2025); two
121 stations located outside of the major *Phaeocystis* bloom to the southwest of the Eastern Grid (hereafter "Central");
122 and one station located in the bloom area between the Eastern Grid and the central stations which was sampled
123 on two consecutive days (hereafter "Station 28-1" and "Station 28-2"). In the Eastern Grid, six stations were
124 sampled from 20 May to 25 May (hereafter "East 1") before Station 28 and the central stations were occupied,
125 while three stations were sampled after a major storm at the end of the expedition from 30 May to 2 June (hereafter
126 "East 2"). East 1 was characterized by higher *Phaeocystis* biomass levels and primary production rates compared
127 to East 2, which represented post-bloom conditions (Fig. 2 and 3).

Deleted: ; Bertrand et al., in prep

Deleted: (Bertrand et al., in prep)



130

131 **Figure 1. Map of the stations sampled during the BELAS-1 expedition (CE22009, RV *Celtic Explorer*) in different**
 132 **regions: the Eastern Grid or "East" (East 1: 20 May to 25 May; East 2: 30 May to 2 June), Station 28 (St. 28), and**
 133 **"Central". All stations were sampled for total ^{234}Th in seawater. Red dots denote stations where in situ large volume**
 134 **pumps (LVP) were deployed. Yellow squares denote stations sampled for net primary production (NPP).**

135 The mixed layer depth (MLD), defined as the depth where potential density exceeds the density at 10 m depth
 136 by 0.03 kg m^{-3} (de Boyer Montégut et al., 2004), ranged from 22 to 66 m. Stratification, defined as the squared
 137 buoyancy frequency, ranged from 5 to 88 $\text{m}^2 \text{s}^{-2}$ and was on average 20% deeper than the MLD. The base of the
 138 primary production zone (PPZ), defined as the depth at which fluorescence was 10% of its maximal value (Owens
 139 et al., 2015), ranged from 79 to 176 m and was used to operationally define the base of the euphotic zone.
 140 Fluorescence was obtained from an optical package mounted on the conductivity-temperature-depth (CTD)
 141 rosette system. The CTD was not equipped with a photosynthetic active radiation (PAR) sensor during the
 142 BELAS-1 expedition.





Figure 2. Satellite-derived chlorophyll-a (chl-a) concentrations (Devred et al., 2025) at the time of sampling of East 1 stations (left panel), Station 28 and central stations (middle panel), and East 2 stations (right panel).

2.1. Net primary production

Water was collected from six depths using a CTD-rosette fitted with 10 L Niskin bottles. While collections were dispersed throughout the upper 200 m of the water column, some sampling depths were selected based on key hydrographic features, including the surface and chlorophyll-a (chl-a) maximum (exact depths can be found in Table S1 in the Supplement). Water was transferred from the Niskin to the incubation bottles using acid-rinsed (10% HCl followed by a minimum of four rinses with type 1 (≥ 18.2 M Ω cm) reverse osmosis water) silicone tubing fitted with 150 μ m Nitex mesh and covered in electrical tape to exclude light from samples collected at depth. Incubations were performed in triplicate in acid-washed 1 L incubation bottles filled to the bottle neck (approximate volume of 1.2 L). Bottles were amended with $\text{H}^{13}\text{CO}_3^-$ (99%; Cambridge Isotope Laboratories) at an approximate addition of 330 $\mu\text{mol C L}^{-1}$. For stable isotope tracer studies, we aim for target additions of 10%, and the average atom percent enrichment for the current study was $13.8 \pm 0.5\%$.

Once the isotope was added, bottles were placed in flow-through deck incubators for 24 h with natural daylight. Incubators were covered at night to prevent the unintended impacts of deck lighting. The incubation bottles were covered with screen bags of varying thicknesses to mimic the light availability at the depth where they were collected. For deep samples, bottles were covered with foil and then multiple layers of electrical tape to eliminate light intrusion. Temperature and light in the incubators were monitored using HOBO TidbiT v2 water temperature data loggers (Onset Computer Corporation).

The experiments were terminated by filtration onto pre-combusted (450°C for 4 h) GF-75 filters with a nominal pore size of 0.3 μm . Filters were placed into 2 mL cryovials and frozen at -20°C until analysis at Bigelow Laboratory for Ocean Sciences using a CosTech ECS 4010/Thermo DELTA V Advantage Isotope Ratio Mass



Moved down [1]:
Figure 3. Profiles of chlorophyll-a from the CTD (CTD chl-a) and net primary production (NPP, absolute uptake) in the upper 200 m of the water column in East 1 (green), Central (dark blue) and East 2 (yellow) stations. The base of the primary production zone (PPZ, Owens et al., 2015) shown with a grey solid line is used to operationally define the base of the euphotic zone. The base of the mixed layer (MLD) is shown with a blue dashed line.

175 Spectrometer. Absolute carbon uptake rates were calculated according to Hama et al. (1983) using the following
176 equation:

177
$$\rho = \frac{PC \text{ at}\%xs}{DIC \text{ at}\%xs \times Time} \times [PC], \quad (1)$$

178 where ρ is the absolute uptake rate and at%xs is the atom% excess. For particulate carbon (PC), atom% excess
179 refers to the percentage of ^{13}C enrichment at the end of the incubation minus the percentage of ^{13}C at the beginning
180 of the incubation. For dissolved inorganic carbon (DIC), atom% excess refers to the proportional change in ^{13}C
181 available in the water when the $H^{13}CO_3^-$ label is added. Ambient HCO_3^- concentrations were estimated based on
182 the salinity of each sample (Parsons et al., 1984). Absolute uptake rate represents how much ^{13}C was taken up by
183 the target community over the incubation period and is presented as units of carbon per volume/area per time (i.e.,
184 $mmol C m^{-3} d^{-1}$).

185 Depth-integrated NPP was calculated using a trapezoidal integration (Knap et al., 1996) up to the base of the PPZ
186 from measured absolute uptake rates. In cases where the base of the PPZ exceeded the deepest sampling depth,
187 the absolute uptake rate from the deepest sampling depth was used for any depths up to and including the base of
188 the PPZ. Due to a limited number of sampling depths, NPP could not be integrated at the central stations.

189 Satellite NPP was calculated following the methods described in Devred et al. (2025). Briefly, the satellite product
190 is based on daily 4-km MODIS-Aqua level-3 fields of remote sensing reflectance, sea surface temperature (SST),
191 and PAR. Chlorophyll-a was derived using a regionally-tuned version of the MODIS OC3M algorithm, and gaps
192 in chl-a and SST were filled using Data Interpolating Empirical Orthogonal Functions (DINEOF).

193 2.2. Total ^{234}Th and ^{238}U in seawater

194 Water column samples were taken in the upper 500 m of all stations ($n = 12$, Fig. 1, 4) using a CTD-rosette at 15
195 discrete depths. Unfiltered seawater samples (2 L) were processed on board and analyzed for the activity of total
196 (dissolved + particulate) ^{234}Th following the method described in Clevenger et al. (2021). Samples were
197 immediately acidified after collection and spiked with a yield monitor (^{230}Th , 25 disintegrations per minute, dpm,
198 per sample). After a minimum 6 h equilibration time, sample pH was increased to ~ 8.5 using ammonium
199 hydroxide, and reagents were added to form a manganese oxide precipitate for scavenging of Th. Samples were
200 allowed to stand for at least 8 h, filtered onto 25-mm diameter quartz microfiber filters (QMA), dried and mounted
201 for beta counting. Samples were counted using three low-level Risø beta multiscintillation counters (5 detectors each) until
202 the uncertainty was typically $< 3\%$. Beta counters were calibrated using two sets of five deep samples (1500 m)
203 from two stations. At least 5 months after collection (> 6 ^{234}Th half-lives, where the half-life = 24.1 d), samples
204 were recounted at Dalhousie University to determine the non- ^{234}Th beta activity stemming from other
205 radionuclides included in the precipitate, which was subtracted from the first count. The net counting rate was
206 corrected for ^{234}Th decay and ingrowth from ^{238}U , counting efficiency and chemical recovery. The chemical
207 recovery of ^{230}Th was determined by using a Thermo Scientific iCAP quadrupole inductively coupled plasma
208 mass spectrometer (ICP-MS) following the procedure detailed in Clevenger et al. (2021). Recoveries averaged $83 \pm 10\%$ ($n = 218$). The uranium-238 (^{238}U) activity was derived from salinity (Owens et al., 2011).

Deleted: by the Bigelow Analytical Services

Deleted: ($mmol C m^{-3} d^{-1}$)

Deleted: and at%xs

Deleted: excess

Deleted: atoms in relation to the natural abundance of either particulate carbon (PC) or dissolved inorganic carbon (DIC)

Deleted: for stations located in the Eastern Grid using a midpoint box integration

Deleted: that incorporated both

Deleted: and chl-a concentrations

Deleted: Due to a limited number of sampling depths, NPP could not be integrated at the central stations. A trapezoidal integration is typically recommended for calculating depth-integrated NPP from in situ rate measurements (Knap et al., 1996), but this method can have difficulties when there are variances in sampling depth across and within stations, as was the case in the BELAS-1 expedition. Despite these variances, the midpoint box and trapezoidal integrations only differed by a maximum of 2.5% across Eastern Grid stations. However, the trapezoidal integration led to increased differences in depth-integrated NPP when incorporating chl-a concentrations (up to 18%) compared to the midpoint box integration. Therefore, the more conservative midpoint box approach was taken. Chl-a concentrations were used to calculate depth-integrated NPP to determine if a relationship between absolute uptake rates and chl-a could be used to model depth-integrated NPP at the central stations. However, no clear relationship was determined. Fluorometer profiles from the CTD were adjusted using a regression based on chl-a concentrations measured by high-performance liquid chromatography (HPLC) but were not corrected for non-photochemical quenching (NPQ) due to the presence of a subsurface chl-a maximum (Bertrand et al., in prep). Chl-a concentrations near the surface should, therefore, be considered slightly underestimated, but this is likely to have little to no impact on the rate when integrated over the water column. The difference in depth-integrated NPP with and without the inclusion of chl-a concentrations ranged from 0.1% (Station 7) to 11% (Station 4).

Deleted: error

2.3. Particles collected using large volume pumps and marine snow catchers


Size-fractionated particles were collected using battery-powered in situ large volume pumps (McLane Research Laboratories, Inc.) equipped with a three-tier 142-mm diameter filter holder (MULVFS style; Bishop et al., 2012; Lam et al., 2015). Five pumps were deployed at depths between 40 and 490 m at six stations (Fig. 1, 5). The pump deployment depths were chosen after examining the fluorescence profile from a CTD cast conducted shortly prior to the pump deployment. One pump was placed close to the base of the PPZ and another pump at 100 m below the PPZ base. Pumps were programmed to sample for 2.5 to 3.0 h at a starting flow rate of 6 L min⁻¹ and pumped an average of 460 L each. During each pump deployment, pressure sensors (RBR duet 3) were attached to selected pumps to confirm deployment depths.

The filter holders contained three filters: two Nitex screens (335 and 51 µm nominal pore size, acid-leached prior to the cruise) above a pre-combusted QMA filter (~1 µm nominal pore size, Graff et al., 2023) for size fractionation (1–51, 51–335 and >335 µm). A total of 28 samples were collected for each size fraction. Size-fractionated particles were analyzed for ²³⁴Th, particulate organic carbon (POC), and biogenic silica (bSi, only from Nitex screens, i.e., 51–335 and >335 µm size fractions). In every cast, an additional filter holder was mounted on the deepest pump to obtain a seawater process blank (“dipped blank” filters, Lam et al., 2015). Immediately after recovering the pumps, residual water from each filter holder was removed by vacuum to avoid particle loss.


Zooplankton that were not part of the passive sinking flux (“swimmers”) and were visible to the naked eye were carefully handpicked from the Nitex screens using forceps and discarded. Particles were gently rinsed off the Nitex screens onto 25-mm-diameter silver (Ag) filters (1.2 µm nominal pore size) using pre-filtered seawater. QMA filters were subsampled for ²³⁴Th and POC using a circular punch tool. Ag and QMA filters were dried, beta counted at sea for ²³⁴Th activities and recounted >5 months later at Dalhousie University. After final counting, Ag filters were split into halves by weight and analyzed for POC and bSi. POC was analyzed using high-temperature combustion (Costech Instruments Elemental Combustion System 4010) after acid fumigation. bSi was analyzed following the NaOH digestion method (40 min at 95°C) using a UV-Visible Spectrophotometer (Genesys 10S, Thermo Scientific) as in Roca-Martí et al. (2021).

The average of all dipped blanks was subtracted from total ²³⁴Th, POC, and bSi measurements. Dipped blanks contributed on average <3% to the total ²³⁴Th measured on the filters, <7% to total bSi, and <14% to total POC (Table S2 in the Supplement). All concentrations were above the limit of detection (3 x standard deviation of the dipped blanks, Lam et al., 2018). Analysis of triplicate punches from four QMA filters representing depths from 100 to 500 m yielded a relative standard deviation for POC of 1–4%, indicating a relatively homogeneous particle distribution across the filters (Maiti et al., 2012). POC and bSi data have average uncertainties of 3% and 6%, respectively, resulting from the dipped blank correction. Particulate ²³⁴Th have average uncertainties of 5% resulting from counting and the dipped blank correction.

At Station 12 in East 1, particles were also collected using marine snow catchers (MSCs) as an independent method to constrain POC/²³⁴Th ratios in sinking material. Three MSCs were deployed approximately 3.5 h prior to the large volume pump deployments at similar depths (165, 265, and 485 m; ± 5 m). MSCs are 1.5-m-tall, 100-



Deleted:
Figure 4. Profiles of total ²³⁴Th and ²³⁸U in the upper 500 m of the water column in East 1 stations (green), Station 28 (light blue), central stations (dark blue) and East 2 stations (yellow). The base of the primary production zone (PPZ) is shown with a grey solid line and the base of the mixed layer (MLD) is shown with a blue dashed line.



Moved down [2]:
Figure 5. Profiles of POC/²³⁴Th (C/Th) and bSi/²³⁴Th (bSi/Th) ratios in >51 µm particles across regions. Black

Deleted: 1

Deleted: , total volume of ~100 L

307 L water samplers equipped with a removable base section designed to separate particles according to their sinking
 308 velocity under low-turbulence conditions (Giering et al., 2016; Riley et al., 2012). Following MSC retrieval,
 309 particles were allowed to settle for 2 h, after which they were classified into three operationally defined fractions:
 310 suspended, slow-sinking, and fast-sinking (Romanelli et al., 2023, 2026). Suspended particles, with negligible
 311 sinking velocity, were collected from a central tap ~79 cm from the MSC top. Slow-sinking particles, with
 312 velocities $<18 \text{ m d}^{-1}$, were siphoned from the base. Fast-sinking particles, with velocities $>18 \text{ m d}^{-1}$, were trapped
 313 in the tray at the bottom of the base section. On average, 2.1 L (of 5.0 L) from the base and 1.4 L (of 2.6 L) from
 314 the tray were filtered through 25-mm-diameter Ag filters (0.45 μm nominal pore size) for the determination of
 315 ^{234}Th and POC in slow- and fast-sinking particles. Blank corrections were applied by subtracting the average of
 316 two filter blanks from total ^{234}Th and POC measurements, and the slow- and fast-sinking particle fractions were
 317 subsequently combined. Average uncertainties were 1% for POC and 17% for particulate ^{234}Th , resulting from
 318 the blank correction (POC and ^{234}Th) and counting (^{234}Th). The sample from 165 m was excluded from analysis
 319 due to a low signal-to-noise ratio in the ^{234}Th measurements.

320 2.4. ^{234}Th , POC and bSi export fluxes

321 Export fluxes of ^{234}Th were calculated for each water column profile by integrating the disequilibrium between
 322 ^{234}Th and ^{238}U from the surface to each sampled depth over the upper 500 m of the water column. This model
 323 implies steady-state (SS) conditions and neglects physical transport (Roca-Martí and Puigcorb , 2024; Savoye et
 324 al., 2006). We discuss the validity of this SS approach in Sect. 2.5.

325 To estimate POC and bSi export fluxes, steady-state ^{234}Th fluxes were multiplied with $\text{POC}/^{234}\text{Th}$ (C/Th) and
 326 $\text{bSi}/^{234}\text{Th}$ (bSi/Th) ratios measured in $>51 \mu\text{m}$ particles (i.e., combining the 51–335 and $>335 \mu\text{m}$ size fractions)
 327 collected using large volume pumps. Using the $>51 \mu\text{m}$ fraction for the calculation of fluxes is consistent with the
 328 majority of ^{234}Th studies (Puigcorb  et al., 2020). Furthermore, this choice is directly supported here by comparing
 329 C/Th ratios measured at Station 12 in size-fractionated particles with those measured in sinking particles collected
 330 using MSCs (Fig. S1 in the Supplement). Sinking particles from MSCs show C/Th ratios similar to all pump size
 331 fractions that are greater than $51 \mu\text{m}$ (i.e., 51–335 μm , $>335 \mu\text{m}$, and $>51 \mu\text{m}$; Fig. S1 in the Supplement). On the
 332 contrary, C/Th ratios from MSCs are clearly higher than C/Th ratios in pump samples that include particles smaller
 333 than $51 \mu\text{m}$ (i.e., the 1–51 μm and $>1 \mu\text{m}$ fractions).

334 Based on the samples collected at five depths across six individual stations, we derived average C/Th and bSi/Th
 335 ratios ($\pm$ standard deviation) for several depth horizons below the PPZ base (115–185, 210–240, 260–340, 465–
 336 490 m; Fig. 5). Ratios at depths not sampled by pump deployments were interpolated linearly between the mid-
 337 points of each depth horizon. For POC, all pump casts were combined to determine cruise-average C/Th ratios
 338 because no significant differences were found between East 1, Station 28, Central, and East 2 (one-way ANOVA,
 339 $p > 0.05$). For bSi, pump casts from East 1, Station 28, and Central were combined (one-way ANOVA, $p > 0.05$)
 340 to determine average bSi/Th ratios and estimate bSi fluxes from 20 May to 29 May. bSi/Th ratios in East 2 were
 341 used to estimate bSi fluxes from 30 May to 2 June given the higher bSi/Th ratios measured in East 2 (Kruskal-
 342 Wallis, $p < 0.01$).

Deleted: After a

Deleted: settling period

Deleted: subsamples were taken from each MSC: o

Deleted: compartment

Deleted: (located at the bottom of the base)

Deleted: Further details on MSC deployments and sample processing are available in Romanelli et al. (2023) and Cisternas-Novoa et al. (in prep).

351 2.5. Testing the assumptions of our export model

352 2.5.1. How valid is the steady state assumption?

353 The validity of the steady state (SS) assumption during a phytoplankton bloom depends on when the sampling
354 occurs with respect to the bloom peak and the duration of the bloom (Ceballos-Romero et al., 2018). Following
355 the approach by Ceballos-Romero et al. (2018), satellite-derived chl-a concentrations suggest that the peak of
356 maximum production in the Eastern Grid area occurred on 11 May, i.e., 9–22 d before our sampling period (20
357 May–2 June). Chl-a decreased quite rapidly with time after the peak (Fig. S2 in the Supplement), similar to a
358 hypothetical bloom scenario considered in Ceballos-Romero et al. (2018) (their Fig. 1b). In such conditions, these
359 authors found that the ^{234}Th SS model provides accurate flux estimates during a ~1 week-long sampling period,
360 referred to as "window of success", which commences 10 d after the peak of the bloom. Comparing the sampling
361 dates and bloom timing of the present study with the findings from Ceballos-Romero et al. (2018) suggests that
362 most of our sampling in East 1 happened within the window of success (21 May–27 May, Fig. S2 in the
363 Supplement), supporting the validity of the SS assumption. Sampling in East 2 occurred 3 to 6 d after the window
364 of success, and therefore, fluxes at the end of the sampling period might represent high end-member estimates
365 (Ceballos-Romero et al., 2018).

366 In addition, the sampling strategy of the BELAS-1 expedition was incompatible with a non-steady state (NSS)
367 model given that: 1) the sampling period was relatively short (6 d in East 1 and 4 d in East 2) relative to the 2–3
368 weeks recommended by Ceballos-Romero et al. (2018) to capture changes, if any, in ^{234}Th activities vs. time and
369 apply a NSS approach; and 2) in order to obtain accurate flux estimates with a NSS model, a Lagrangian sampling
370 strategy that tracks the same water mass must be followed (Resplandy et al., 2012; Savoye et al., 2006), which
371 was not possible during this expedition. Indeed, changes in upper water column properties occurred between the
372 sampling of East 1 and East 2 due to a storm (Erin Bertrand, [personal communication, 2025](#)), indicating that
373 different water masses might have been sampled. Further, in the Eastern Grid, where we have the highest density
374 of measurements, we did not observe significant changes in ^{234}Th inventories in the upper 150 m over time (Fig.
375 S3a in the Supplement). Thus, we consider that the SS model gives the best estimate of ^{234}Th export in this study.

376 2.5.2. Evaluating the effect of physical transport

377 *Vertical Transport:* The central Labrador Sea in 2022 experienced moderately deep convection reaching 1600 m
378 (Yashayaev, 2024) followed by water column stratification and bloom development in early April. Winter mixing
379 would homogenize vertical ^{234}Th gradients in the water column. However, sampling during BELAS-1 occurred
380 at least 50 d after convection ended (Yashayaev, 2024), a time interval that exceeds the half-life of ^{234}Th and its
381 mean life ($1/\text{decay constant} = 35$ d). Therefore, we are confident that vertical advection associated with winter
382 mixing did not influence the estimates presented in this study. The potential contribution of vertical diffusion was
383 estimated using vertical diffusivity estimates from a hindcast simulation using the model of Ohashi et al. (2024)
384 in the Eastern Grid at the time of the cruise. Vertical diffusivity decreased with depth from 10^{-4} – 10^{-2} $\text{m}^2 \text{s}^{-1}$ at the
385 surface to 10^{-6} $\text{m}^2 \text{s}^{-1}$ below 80 m. We estimate that vertical diffusion changed ^{234}Th fluxes at the base of the PPZ
386 by <10 $\text{dpm m}^{-2} \text{d}^{-1}$ only, given the small gradients in ^{234}Th activities observed across the base of the PPZ at each
387 station (Fig. 4). Therefore, the influence of vertical transport on ^{234}Th export flux estimates must have been
388 negligible.

Deleted: et al., in prep

Horizontal Transport: Horizontal advection in the Eastern Grid, derived from the hindcast simulation of the cruise period, had mean velocities of 2.8 km d⁻¹ over the top 150 m in a mostly southeastward direction. The observed ²³⁴Th activities show relatively large variability between profiles in the Eastern Grid (Fig. S4 in the Supplement), however, there are no consistent spatial trends in ²³⁴Th inventories (Fig. S3b in the Supplement). We attribute the variability in ²³⁴Th between profiles to small-scale spatial variations or patchiness rather than horizontal transport.

3. Results

Table 1 presents a summary of the integrated NPP rates down to the base of the PPZ, together with the sinking fluxes (POC, bSi) measured at different depths in the upper 500 m across all the stations sampled during BELAS-1.

Table 1. Integrated net primary production (NPP) down to the base of the primary production zone (PPZ), and particulate organic carbon (POC) and biogenic silica (bSi) fluxes at different depths across all stations.

Region	Station ID	Lat (°N)	Long (°E)	Sampling date (2022)	Base of PPZ (m)	Integrated NPP (mmol C m ⁻² d ⁻¹)	POC flux at PPZ base (mmol C m ⁻² d ⁻¹)	bSi flux at PPZ base (mmol Si m ⁻² d ⁻¹)	POC flux at PPZ base + 100 m (mmol C m ⁻² d ⁻¹)	bSi flux at PPZ base + 100 m (mmol Si m ⁻² d ⁻¹)	POC flux at 500 m (mmol C m ⁻² d ⁻¹)	bSi flux at 500 m (mmol Si m ⁻² d ⁻¹)
East 1	4	58.82	-49.11	20-May	119	166 ± 25	16.3 ± 4.0	1.8 ± 0.2	15.6 ± 3.5	2.1 ± 0.3	13.9 ± 6.1	2.1 ± 0.5
	6	58.82	-49.44	21-May	153	–	8.1 ± 3.1	0.9 ± 0.3	8.0 ± 2.7	1.1 ± 0.3	8.9 ± 5.0	1.3 ± 0.6
	9	58.65	-49.36	22-May	163	263 ± 30	6.2 ± 2.9	0.7 ± 0.3	5.2 ± 2.3	0.8 ± 0.3	3.0 ± 3.2	0.5 ± 0.4
	8	58.65	-49.61	23-May	153	–	9.3 ± 3.4	1.0 ± 0.3	9.1 ± 3.1	1.2 ± 0.4	11.8 ± 5.5	1.8 ± 0.5
	11	58.65	-49.11	24-May	141	126 ± 16	7.4 ± 2.7	0.8 ± 0.2	5.5 ± 2.3	0.7 ± 0.3	–	–
St. 28	12	58.48	-49.11	25-May	176	–	2.9 ± 3.8	0.3 ± 0.4	4.1 ± 3.1	0.6 ± 0.4	7.0 ± 4.6	1.1 ± 0.6
	28-1	57.72	-50.07	26-May	165	–	4.7 ± 3.0	0.5 ± 0.3	2.1 ± 2.5	0.3 ± 0.4	0.8 ± 3.7	0.1 ± 0.6
	28-2	57.72	-50.07	27-May	135	–	5.1 ± 2.9	0.5 ± 0.3	10.5 ± 3.6	1.4 ± 0.4	6.7 ± 4.6	1.0 ± 0.6
	24	56.65	-52.47	28-May	115	–	12.6 ± 3.3	1.4 ± 0.2	12.0 ± 3.4	1.6 ± 0.4	17.0 ± 7.4	2.6 ± 0.6
Central	29	56.82	-52.22	29-May	170	–	22.1 ± 5.8	2.4 ± 0.4	12.1 ± 3.7	1.8 ± 0.5	9.4 ± 5.5	1.4 ± 0.6
	15	58.48	-49.61	30-May	155	–	7.2 ± 3.0	1.3 ± 0.4	3.4 ± 2.7	0.9 ± 0.7	2.7 ± 4.0	0.6 ± 0.8
East 2	13	58.48	-49.36	31-May	143	38 ± 3	12.0 ± 3.6	2.1 ± 0.5	11.5 ± 3.3	3.0 ± 0.7	8.1 ± 5.3	1.8 ± 1.0
	7	58.82	-49.61	02-Jun	79	42 ± 6	13.9 ± 3.2	2.4 ± 0.2	18.7 ± 5.4	3.3 ± 0.7	9.9 ± 5.6	2.2 ± 0.9

3.1. NPP rates

The average NPP at the surface and chl-a maximum depths was higher in East 1 (2.01 and $2.56 \text{ mmol C m}^{-3} \text{ d}^{-1}$) compared to Central (1.79 and $1.74 \text{ mmol C m}^{-3} \text{ d}^{-1}$) and East 2 (0.92 and $0.75 \text{ mmol C m}^{-3} \text{ d}^{-1}$), which correlates with decreasing chl-a concentrations over the sampling period (Fig. 3). Surface NPP was highest at Station 11 ($2.36 \pm 0.47 \text{ mmol C m}^{-3} \text{ d}^{-1}$) while NPP at the chl-a maximum peaked at Station 9 ($3.38 \pm 0.83 \text{ mmol C m}^{-3} \text{ d}^{-1}$). Station 13 had the lowest surface and chl-a maximum NPP, $0.54 \pm 0.13 \text{ mmol C m}^{-3} \text{ d}^{-1}$ and $0.52 \pm 0.10 \text{ mmol C m}^{-3} \text{ d}^{-1}$, respectively. Depth-integrated NPP was highest during East 1, peaking at Station 9 with a value of $263 \pm 30 \text{ mmol C m}^{-2} \text{ d}^{-1}$ (Table 1). Between sampling for East 1 and East 2, depth-integrated NPP decreased notably as the bloom declined and transitioned into post-bloom conditions, reaching a minimum of $38 \pm 3 \text{ mmol C m}^{-2} \text{ d}^{-1}$ at Station 13 (Table 1). The measured in situ integrated NPP and satellite-derived NPP were in reasonable agreement (Fig. S5 in the Supplement), with a mean measured-satellite ratio of 1.25 ± 0.69 ($n = 5$), supporting the use of satellite-derived NPP for estimating export efficiency (see Sect. 4.3).




Figure 3. Profiles of chlorophyll-a from the CTD (CTD chl-a) and net primary production (NPP, absolute uptake) in the upper 200 m of the water column in East 1 (green), Central (dark blue) and East 2 (yellow) stations. The base of the primary production zone (PPZ, Owens et al., 2015) shown with a grey solid line is used to operationally define the base of the euphotic zone. The base of the mixed layer (MLD) is shown with a blue dashed line.

3.2. $^{234}\text{Th}/^{238}\text{U}$ profiles and ^{234}Th fluxes

In the upper water column, ^{234}Th activities were always lower than ^{238}U activities ($\sim 2.4 \text{ dpm L}^{-1}$) across all stations indicating particle export (Fig. 4). However, the magnitude and extent of the ^{234}Th deficits showed variability. The lowest ^{234}Th activities were found in surface waters of East 1, with average activities of $1.6 \pm 0.2 \text{ dpm L}^{-1}$,

Deleted: 252


Deleted: 6

Deleted: 39

Deleted: 1

Moved (insertion) [1]

427 compared to Station 28, Central and East 2, where average surface activities were 1.9–2.0 dpm L⁻¹. ²³⁴Th activities
 428 reached equilibrium with ²³⁸U at depths between 55 and 100 m, i.e., between the MLD and the base of the PPZ
 429 (Fig. 4). The only exception is Station 7 (East 2), the last station to be occupied, where equilibrium was only
 430 reached at ~150 m, below the PPZ.



431
 432 **Figure 4. Profiles of total ²³⁴Th and ²³⁸U in the upper 500 m of the water column in East 1 stations (green), Station 28**
 433 **(light blue), central stations (dark blue) and East 2 stations (yellow). The base of the primary production zone (PPZ) is**
 434 **shown with a grey solid line and the base of the mixed layer (MLD) is shown with a blue dashed line.**

435 In general, ²³⁴Th and ²³⁸U activities remained close to each other below the equilibrium depth. However, there
 436 were two notable exceptions. At some stations (i.e., East 1 Stations 9 and 12, and Station 28-1), large ²³⁴Th activity
 437 excesses relative to ²³⁸U were apparent within the PPZ, which is indicative of remineralization or disaggregation
 438 processes. In addition, we found a ²³⁴Th deficit below the PPZ at Station 28-2, which can indicate particle
 439 repackaging processes.

440 One-dimensional (1D) steady state ²³⁴Th fluxes (Fig. S6 in the Supplement) reflected the variability observed in
 441 ²³⁴Th/²³⁸U profiles. At the base of the PPZ, ²³⁴Th fluxes ranged from negligible in East 1 to 1640 dpm m⁻² d⁻¹ in
 442 Central. On average, ²³⁴Th fluxes (dpm m⁻² d⁻¹) at the base of the PPZ were 620 ± 330 in East 1, 360 ± 20 at
 443 Station 28, 1290 ± 500 in Central and 820 ± 260 in East 2. In general, considering all stations, ²³⁴Th fluxes
 444 increased slightly from the base of the PPZ to 100 m below the base (1.4-fold on average), and that increase was
 445 most pronounced (2.8-fold) at Station 28-2 due to the ²³⁴Th deficit observed below the PPZ at this station (Fig.
 446 4). At 500 m, ²³⁴Th fluxes were moderately high, with fluxes ≥1000 dpm m⁻² d⁻¹ at most stations except three
 447 stations located outside the central region (Stations 9, 28-1, 15).

Deleted: 5

3.3. C/Th and bSi/Th in particles

The C/Th ratios in >51 μm particles changed from 41 to 4.0 $\mu\text{mol dpm}^{-1}$ between near surface (40 m) and 490 m (Fig. 5). The highest C/Th ratios were found within the PPZ in East 1 where *Phaeocystis* dominated the phytoplankton community and phytoplankton biomass, and NPP rates were highest (Fig. 3). Below the PPZ, C/Th ratios decreased with depth from $13.5 \pm 2.8 \mu\text{mol dpm}^{-1}$ at 115–185 m to $7.1 \pm 2.7 \mu\text{mol dpm}^{-1}$ at 465–490 m considering all stations.




Figure 5. Profiles of POC/²³⁴Th (C/Th) and bSi/²³⁴Th (bSi/Th) ratios in >51 μm particles across regions. Black symbols denote average C/Th ratios (all stations) and average bSi/Th ratios (all stations except East 2) for the following depth horizons: 115–185, 210–240, 260–340, 465–490 m. Empty symbols denote samples collected within the primary production zone. Analytical uncertainties are represented by horizontal bars and uncertainties less than $\sim 3 \mu\text{mol dpm}^{-1}$ are not visible against filled symbols. POC = particulate organic carbon, bSi = biogenic silica.


bSi/Th ratios in >51 μm particles also showed a general decrease with depth, but not as marked as C/Th ratios (Fig. 5). The highest bSi/Th ratios throughout all depths were found in East 2, with values decreasing from 2.3–2.7 $\mu\text{mol dpm}^{-1}$ in the upper 220 m to 1.6–1.9 $\mu\text{mol dpm}^{-1}$ at deeper depths. In East 1, Station 28 and Central, bSi/Th ratios decreased from $1.5 \pm 0.1 \mu\text{mol dpm}^{-1}$ at 115–185 m to $1.1 \pm 0.1 \mu\text{mol dpm}^{-1}$ at 465–490 m.

3.4. POC and bSi fluxes

POC flux profiles show large station-to-station variability (Fig. 6a). At the base of the PPZ, POC fluxes ranged by 1 order of magnitude from 2.9 $\text{mmol C m}^{-2} \text{d}^{-1}$ in East 1 to 22.1 $\text{mmol C m}^{-2} \text{d}^{-1}$ in Central (Table 1). On average, POC fluxes ($\text{mmol C m}^{-2} \text{d}^{-1}$) at the base of the PPZ were 8.4 ± 4.5 in East 1, 4.9 ± 0.2 at Station 28, 17.3 ± 6.7 in Central and 11.1 ± 3.5 in East 2 (Fig. 6a, 7). POC fluxes at 100 m below the PPZ were not significantly different

Deleted: decreased
Deleted: 10-fold with depth from values of up to
Deleted: $\mu\text{mol dpm}^{-1}$ at 40 m to values as low as
Deleted: t

Moved (insertion) [2]



Deleted:

than those at the base of the PPZ (t test, $p > 0.05$), indicating negligible attenuation in POC fluxes below the PPZ and a transfer efficiency close to 1 (i.e., flux at 100 m below PPZ base $\sim$ flux at PPZ base; Fig. 7). At 500 m, POC fluxes were, on average, 8.9 ± 4.2 in East 1, 3.8 ± 4.1 at Station 28, 13.2 ± 5.3 in Central and 6.9 ± 3.7 $\text{mmol C m}^{-2} \text{ d}^{-1}$ in East 2. Only East 2 showed consistent flux attenuation (30–60%) between the base of the PPZ and 500 m at all stations.





Figure 6. Profiles of particulate organic carbon (POC, panel a) and biogenic silica (bSi, panel b) flux across regions. Fluxes measured at individual stations are shown in grey, whereas averages are shown in color (green for East 1, light blue for St. 28, dark blue for Central, yellow for East 2). The vertical red lines at $20 \text{ mmol C m}^{-2} \text{ d}^{-1}$ for POC and $4 \text{ mmol Si m}^{-2} \text{ d}^{-1}$ for bSi are for visual reference. The base of the primary production zone (PPZ) is shown with green shading (grey solid line indicates the average, black solid lines indicate the minimum and maximum) and the average base of the mixed layer (MLD) is shown with a blue dashed line.

bSi flux profiles also show large variability across stations (Fig. 6b). At the base of the PPZ, bSi fluxes ranged by 1 order of magnitude from $0.3 \text{ mmol Si m}^{-2} \text{ d}^{-1}$ in East 1 to $2.4 \text{ mmol Si m}^{-2} \text{ d}^{-1}$ in Central and East 2 (Table 1). On average, bSi fluxes ($\text{mmol Si m}^{-2} \text{ d}^{-1}$) at the base of the PPZ were 0.9 ± 0.5 in East 1, 0.5 ± 0.0 at Station 28, 1.9 ± 0.7 in Central and 1.9 ± 0.6 in East 2 (Fig. 6b, 7). In deeper waters at most stations, bSi fluxes were similar

or even higher than at the base of the PPZ indicating either no flux attenuation or addition of bSi at depth. The exception is East 2, where bSi flux attenuated by 10–50% from the base of the PPZ to 500 m.




Figure 7. Spatial maps of particulate organic carbon (POC) and biogenic silica (bSi) flux at the base of the primary production zone (PPZ, left panels) and transfer efficiency (flux at 100 m below PPZ base/flux at PPZ base) of POC and bSi over time in the study area (right panels). The horizontal dashed lines in the right panels indicate a transfer efficiency = 1.

4. Discussion

We begin the Discussion by providing an overview of the POC fluxes measured in this study and comparing them with those reported in other studies. We then examine the possible reasons why the massive *Phaeocystis* bloom observed here did not result in enhanced fluxes, with a particular focus on the ballasting hypothesis. Lastly, we present estimates of BCP efficiency during this *Phaeocystis* bloom using both snapshot and longer-term NPP observations. The assumptions underlying the export calculations, including the steady-state framework, are described in Sect. 2.5.

4.1. POC fluxes in the central Labrador Sea during the decline of a *Phaeocystis* bloom and comparison with other studies

This study presents an unprecedented number of measurements of sinking fluxes in the central Labrador Sea (~300 ²³⁴Th measurements) and represents one of the largest assessments of the impact of *Phaeocystis* blooms on

Moved down [3]: Comparison with other studies
Only three prior studies have used ²³⁴Th to estimate POC fluxes in the Labrador Sea: Moran et al. (2003) in July 1999 (three stations), Puigcorb  et al. (2017) in early May 2010 (one station), and Lemaitre et al. (2018) in late June 2014 (three stations), the last one during the decline of a diatom bloom. The ²³⁴Th flux results from our study are similar to those obtained at the PPZ base (or at 100 m, Moran et al., 2003) in June (700–860 dpm m⁻² d⁻¹, Lemaitre et al. 2018) and lie between the fluxes measured in May (270 dpm m⁻² d⁻¹, Puigcorb  et al. 2017) and later in the season in July (980–1400 dpm m⁻² d⁻¹, Moran et al. 2003). Regarding POC fluxes, our fluxes are 1 order of magnitude higher than those measured in early May (0.6 mmol C m⁻² d⁻¹, Puigcorb  et al., 2017), but encompass the range of POC fluxes measured in June (6.1–10 mmol C m⁻² d⁻¹, Lemaitre et al., 2018) and July (5.7–21 mmol C

Study area	Reference depth z (m)	Integrated NPP (mmol C m ⁻² d ⁻¹)	POC flux at z (mmol C m ⁻² d ⁻¹)	Export efficiency (%)	Transfer efficiency z=100 (%)
Labrador Sea	119–176 (PPZ)	125–252 (188)	2.9–16 (8.4)	2–9 (6)	74–139 (98)
	79–155 (PPZ)	39–43 (41)	7.2–14 (11)	31–33 (32)	48–134 (92)
	100	59 ± 2	78 ± 9	130 ± 30	–
Fram Strait	100	48–64 (56)	6.8–13 (9.8)	11–30 (21)	–
	60	–	14–62 (33)	–	–
Barents Sea	60	–	5.6–19 (12)	–	–
	90	–	17	–	–
Barents Sea	40 (0.1%PAR)	67–76 (71)	35–48 (42)	47–72 (59)	56–78 (92)
	35 (0.1%PAR)	68	54	79	41
	90–100	–	15–64 (40)	–	–
North of Svalbard	50	–	13–33 (23)	–	–
	60	–	15–47 (34)	–	–
Ross Sea	50	49–206 (105)	21–66 (34)	21–44 (34)	15–50 (30)
Crozet Plateau, Southern Ocean	87–248	–	1.0–17 (4.8)	–	–
	101–204	10–250 (70)	4.9–30 (17)	7–305 (54)	–
Labrador Sea	40–80 (Eq depth & PPZ; ± 5)	27–80 (54)	6.1–10 (8.0)	8–38 (20)	30–63 (60)
Porcupine Abyssal Plain (PPZ)	67–133	54–90 (75)	9.1–14 (11)	8–26 (16)	107–15 (139)
Atlantic sector Southern Ocean	100–120	66–234 (153)	11–44 (19)	7–34 (12)	–
Kerguelen Plateau, Southern Ocean	100	82	23 ± 4	28	107

Deleted:

Moved (insertion) [3]

Deleted: Table 2. Compilation of sinking fluxes and biological carbon pump efficiency metrics in *Phaeocystis* and diatom blooms. Values given in parentheses next to ranges indicate averages. PPZ = primary production zone, PAR = photosynthetic active radiation, Eq depth = ²³⁴Th and ²³⁸U equilibrium depth, NPP = net primary production, POC = particulate organic carbon, bSi = biogenic silica, ISP = in situ pump, ST = sediment trap.

POC export. Data collected by Fisheries and Oceans Canada along the Atlantic Repeat Hydrography Line 7 West (AR7W) as part of the Atlantic Zone Off-shelf Monitoring Program (AZOMP) between 2014 and 2022, revealed that the *Phaeocystis* bloom encountered in this study (spring 2022) was unusual given its large spatial extent and high biomass concentration (Devred et al., 2024, 2025). Using satellite remote sensing data combined with an ecological approach, these authors determined that the 2022 *Phaeocystis* bloom extended over more than half of the Labrador Sea, lasted for 6 weeks, and resulted in a remarkably high total primary production that accounted for 60% of the May production in the Labrador Sea (33.2 Tg C; Devred et al., 2025). Genomic analyses conducted during our expedition, 1 week later than the 2022 AZOMP mission, confirmed that the species of this bloom was *Phaeocystis pouchetii* (Romanelli et al., 2026; Stevens-Green et al., 2024).

Deleted: Bertrand et al., in prep

The sampling of the *Phaeocystis* bloom in this study mostly focused on a ~1075 km² grid (Eastern Grid) at the southern edge of the bloom about 2 to 3 weeks after its peak. The stations sampled from 20 May to 25 May (East 1) showed integrated NPP rates that were, on average, five times higher than those sampled from 30 May to 2 June (East 2) (Table 1) which we consider post-bloom conditions. Two stations located to the southwest of the Eastern Grid outside of the bloom (Central) were also sampled and represented more typical conditions of the central Labrador Sea with a mixed phytoplankton community composition that was more diatom dominated (Devred et al., 2024, 2025). Beyond the Eastern grid, another station (Station 28) was also sampled in the *Phaeocystis* bloom area, but no in situ NPP data were collected at that station. For this reason, below we focus our discussion on the Eastern Grid and Central (inside vs. outside of the major *Phaeocystis* bloom).

Deleted: re

Deleted: ed

Deleted: to represent

Deleted: (Bertrand et al., in prep)

POC fluxes at the base of the PPZ were, overall, moderate to high, with average fluxes of 8 and 11 mmol C m⁻² d⁻¹ in East 1 and East 2, respectively, and 17 mmol C m⁻² d⁻¹ in Central. This indicates that POC export was lower in the *Phaeocystis* bloom region relative to the stations sampled outside of the bloom. Our observations of large excesses of ²³⁴Th relative to ²³⁸U at around 70–100 m in the Eastern Grid (Stations 9 and 12) and Station 28-1 (Fig. 4) indicate remineralization of sinking particles within the euphotic zone in the *Phaeocystis* bloom region, and limited POC export to the mesopelagic. In general, POC fluxes did not decrease from the base of the PPZ to 500 m, except for East 2 where all stations showed a consistent decrease (30–60%). This lack of flux attenuation down to 500 m suggests that the material that was not subject to remineralization in the euphotic zone was more refractory in nature or sank at relatively fast rates. POC fluxes at 500 m were on average 9 and 7 mmol C m⁻² d⁻¹ in East 1 and East 2, respectively, and 13 mmol C m⁻² d⁻¹ in Central. Therefore, our results also show lower fluxes at depth in the *Phaeocystis* bloom region than at the stations outside of the bloom. Despite the limited flux attenuation found in the upper mesopelagic, taken together, these findings suggest a limited role of *Phaeocystis* in export.

Deleted: but POC fluxes in the *Phaeocystis* bloom were lower than those measured outside of the bloom,

Deleted: the decline of the extensive

Deleted: sampled during this study did not enhance POC export

Deleted: resulted in

Deleted: that

Deleted: ,

Deleted: ing

Deleted: also at depth

Only three prior studies have used ²³⁴Th to estimate POC fluxes in the Labrador Sea: Moran et al. (2003) in July 1999 (three stations), Puigcorb  et al. (2017) in early May 2010 (one station), and Lemaitre et al. (2018) in late June 2014 (three stations), the last one during the decline of a diatom bloom. The POC flux results from our study are 1 order of magnitude higher than those measured at the PPZ base (or at 100 m, Moran et al., 2003) in May (0.6 mmol C m⁻² d⁻¹, Puigcorb  et al., 2017), but encompass the range of POC fluxes measured in June (6.1–10 mmol C m⁻² d⁻¹, Lemaitre et al., 2018) and July (5.7–21 mmol C m⁻² d⁻¹, Moran et al., 2003). Compared to other *Phaeocystis* blooms (Table 2), our POC fluxes in the Eastern Grid compare well with those found in the Barents

Sea using sediment traps (Lalande et al., 2008), but are lower than those measured in that same study using ^{234}Th and large volume sampling. Our POC fluxes in the Eastern Grid are also lower (2 to 8-fold) than those reported in a number of other Arctic and sub-Arctic studies during *Phaeocystis* blooms (Coppola et al., 2002; Dybwad et al., 2021; Le Moigne et al., 2015; Reigstad and Wassmann, 2007). Compared to studies that reported blooms dominated by both *Phaeocystis* and diatoms (Table 2), our POC fluxes fall within the range of those measured in the Crozet Plateau (Morris et al., 2007; Salter et al., 2007), but are lower (4-fold) than those reported in the Barents Sea (Andreassen and Wassmann, 1998) and the Ross Sea (Asper and Smith, 1999).

Table 2. Compilation of sinking fluxes and biological carbon pump efficiency metrics in *Phaeocystis* and diatom blooms. Values given in parentheses next to ranges indicate averages. PPZ = primary production zone, PAR = photosynthetic active radiation, Eq depth = ^{234}Th and ^{238}U equilibrium depth, NPP = net primary production, POC = particulate organic carbon, bSi = biogenic silica, ISP = in situ pump, ST = sediment trap.

Study area	Reference depth z (m)	Integrated NPP (mmol C m ⁻² d ⁻¹)	POC flux at z (mmol C m ⁻² d ⁻¹)	Export efficiency (%)	Transfer efficiency z+100 m (%)	bSi flux at z (mmol Si m ⁻² d ⁻¹)	Molar bSi/POC at z	Bloom stage	Dominant species	Flux method	Number of stations considered	References
Labrador Sea	119–176 (PPZ)	126–263 (185)	2.9–16 (8.4)	2–10 (6)	74–139 (98)	0.3–1.8 (0.9)	0.1	Bloom decline	<i>Phaeocystis</i>	^{234}Th + ISPs	n = 6, East 1	This study
	79–155 (PPZ)	38–42 (40)	7.2–14 (11)	32–34 (33)	48–134 (92)	1.3–2.4 (1.9)	0.2	Post-bloom		^{234}Th + ISPs	n = 3, East 2	
Fram Strait	100	59 ± 2	78 ± 9	130 ± 30	–	–	–	After bloom peak	<i>Phaeocystis</i>	^{234}Th + ISP	n = 1	Le Moigne et al. 2015
	100	48–64 (56)	6.8–13 (9.8)	11–30 (21)	–	–	–	During/after bloom peak	Diatoms	^{234}Th + ISP	n = 2	
Barents Sea	60	–	14–62 (33)	–	–	–	–	Early bloom/bloom	<i>Phaeocystis</i>	^{234}Th + large volume sampling	n = 5	Lalande et al. 2008
	60	–	5.6–19 (12)	–	–	–	–	/late bloom		Drifting STs	n = 5	
Barents Sea	90	–	17	–	–	–	–	Bloom	<i>Phaeocystis</i>	Drifting STs	n = 1	Coppola et al. 2002
Barents Sea	40 (0.1%PAR)	67–76 (71)	35–48 (42)	47–72 (59)	56–78 (67)	–	–	Early bloom	<i>Phaeocystis</i> and diatoms	Drifting STs	n = 2	Andreassen and Wassmann 1998; Buesseler et al. 2020
	35 (0.1%PAR)	68	54	79	41	–	–	Well-developed bloom			n = 1	
North Norwegian fjords & Barents Sea	90–100	–	15–64 (40)	–	–	–	–	Bloom and after bloom	<i>Phaeocystis</i>	Drifting STs	Average of several stations and cruises	Reigstad and Wassmann 2007
North of Svalbard	50	–	13–33 (23)	–	–	–	–	Bloom	<i>Phaeocystis</i>	Ice-tethered STs	n = 6	Dybwad et al. 2021
	60	–	15–47 (34)	–	–	–	–				n = 4	
Ross Sea	50	49–206 (105)	21–66 (34)	21–44 (34)	15–50 (35)	–	–	After bloom peak	<i>Phaeocystis</i> and diatoms	Drifting STs	n = 5	Asper and Smith 1999
Crozet Plateau, Southern Ocean	87–248	–	1.0–17 (4.8)	–	–	0.3–29 (7.9)	0.2–2.8 (1.4)	Bloom	<i>Phaeocystis</i> and diatoms	Neutrally buoyant STs	n = 3 + reoccupations	Salter et al. 2007
	101–204	10–250 (70)	4.9–30 (17)	7–305 (54)	–	–	–	Bloom decline		^{234}Th + ISPs	n = 8 + reoccupations	
Labrador Sea	40–80 (Eq depth & PPZ ± 5)	27–80 (54)	6.1–10 (8.0)	8–38 (20)	30–63 (48)	2.7–6.6 (4.8)	0.4–0.8 (0.6)	After bloom peak	Diatoms	^{234}Th + ISPs	n = 3	Lemaître et al. 2018; Lemaître 2017
Porcupine Abyssal Plain (PPZ)	67–133	54–90 (75)	9.1–14 (11)	8–26 (16)	107–155 (139)	3.4–6.1 (4.4)	0.3–0.4 (0.4)	Bloom decline	Diatoms	^{234}Th + ISPs	n = 23	Clevenger et al. 2024
Atlantic sector Southern Ocean	100–120	66–234 (153)	11–44 (19)	7–34 (12)	–	–	–	Bloom decline	Diatoms	Drifting STs	n = 11	Roca-Martí et al. 2017
Kerguelen Plateau, Southern Ocean	100	82	23 ± 4	28	107	–	–	Bloom decline	Diatoms	^{234}Th + hose pump & drifting STs	n = 1 + reoccupations	Savoie et al. 2008

4.2. Why did this *Phaeocystis* bloom not lead to enhanced fluxes?

Our observations during the decline of a massive *Phaeocystis* bloom in the Labrador Sea do not show indications of enhanced sinking fluxes when compared to an area located outside of the major bloom. These observations cover a period of 2 weeks during which NPP in the bloom area strongly declined, but POC fluxes remained

Moved down [4]: *Phaeocystis* spp. play an exceptional role in marine ecosystems due to its high carbon content when blooming and because of its unique polymorphic life cycle, including free-living cells of <10 μm and gelatinous colonies that usually reach sizes of several mm (Schoemann et al., 2005). However, despite the large sizes of *Phaeocystis* colonies, most of the literature available on the topic shows that *Phaeocystis*-derived material is largely recycled in the upper ocean (Smith and Trimborn, 2024 and references therein). For instance, a compilation of sediment trap studies, including data ranging from polar to sub-Arctic and boreal regions, revealed that *Phaeocystis* POC fluxes strongly decline throughout the upper 100 m of the water column (Reigstad and Wassmann, 2007). Our observations of large excesses of ^{234}Th relative to ^{238}U at around 70–100 m in the Eastern Grid (Stations 9 and 12) and Station 28-1 (Fig. 4) also support shallow remineralization of sinking particles during *Phaeocystis* blooms. Yet, *Phaeocystis*-derived material can be efficiently exported to depth under certain circumstances, including: the formation of aggregates by *Phaeocystis* colonies facilitated by the release of transparent exopolymer particles (TEP) which can scavenge other particles like ballasting minerals (Andreassen and Wassmann, 1998; Passow and Wassmann, 1994; Wollenburg et al., 2018); and through physical mixing processes, such as downwelling associated with eddy activity (Lalande et al., 2011). The incorporation of *Phaeocystis* into fast-sinking zooplankton pellets has also been identified as an important export pathway (Dybwad et al., 2021; Wiedmann et al., 2020).

relatively constant. As described in Sect. 2.5, the assumptions behind the ^{234}Th export model used in this study have been assessed and validated to the best of our possibilities based on information of the dynamics specific to that bloom and a hindcast simulation of physical transport at the time of the cruise. Furthermore, the conversion from ^{234}Th to POC fluxes using $>51\ \mu\text{m}$ particles from large volume filtration in this study is supported with independent data from marine snow catchers (see Sect. 2.4). We also note that the C/Th ratios used to estimate POC fluxes in this study ($\leq 16\ \mu\text{mol dpm}^{-1}$) are far below the ratios measured by Lalande et al. (2008) of up to $>100\ \mu\text{mol dpm}^{-1}$ which were considered to be biased towards high values and in turn led to overestimated POC-derived fluxes. Hence, the comparison of our data with context information and independent methods gives confidence in the export estimates presented here.

Phaeocystis spp. play an exceptional role in marine ecosystems due to its high carbon content when blooming and because of its unique polymorphic life cycle, including free-living cells of $<10\ \mu\text{m}$ and gelatinous colonies that usually reach sizes of several mm (Schoemann et al., 2005). However, despite the large sizes of *Phaeocystis* colonies, most of the literature available on the topic shows that *Phaeocystis*-derived material is largely recycled in the upper ocean (Smith and Trimborn, 2024 and references therein). For instance, a compilation of sediment trap studies, including data ranging from polar to sub-Arctic and boreal regions, revealed that *Phaeocystis* POC fluxes strongly decline throughout the upper 100 m of the water column (Reigstad and Wassmann, 2007). In our study, the large excesses of ^{234}Th observed within the euphotic zone in the bloom region (Fig. 4), also support remineralization of sinking particles in the upper water column during *Phaeocystis* blooms. Yet, *Phaeocystis*-derived material can be efficiently exported to depth under certain circumstances, including: the formation of aggregates by *Phaeocystis* colonies facilitated by the release of transparent exopolymer particles (TEP) which can scavenge other particles like ballasting minerals (Andreassen and Wassmann, 1998; Passow and Wassmann, 1994; Wollenburg et al., 2018); and through physical mixing processes, such as downwelling associated with eddy activity (Lalande et al., 2011). The incorporation of *Phaeocystis* into fast-sinking zooplankton pellets has also been identified as an important export pathway (Dybwad et al., 2021; Wiedmann et al., 2020).

One hypothesis for the lack of enhanced sinking fluxes is that *Phaeocystis* material resulting from the decline of the bloom was not sufficiently ballasted. *Phaeocystis* colonies can release large amounts of TEP during their growth and their senescence, which, thanks to the high stickiness of TEP, can aggregate suspended particles and facilitate the formation of marine snow (Passow, 2002). However, the density of TEP is lower than that of seawater, which may cause the aggregates to remain in surface waters if not ballasted by other particles (Mari et al., 2017). In line with this, other studies have shown that below the mixed layer, marine biogels such as TEP can adhere to and accumulate on particles, reducing their sinking velocity by increasing both their buoyancy and hydrodynamic resistance, which may enhance carbon flux attenuation (Alcolombri et al., 2025; Romanelli et al., 2023). Sinking rates of *Phaeocystis* colonies and aggregates range from negligible to up to $200\ \text{m d}^{-1}$ (Schoemann et al., 2005), indicating that their capacity to sink and get transferred from the upper ocean to depth depends on their characteristics including the amount and type of particles that might have been scavenged. For instance, in the ice-covered Arctic Ocean, sinking of *Phaeocystis* aggregates throughout the water column down to the seafloor has been associated with ballasting by cryogenic gypsum which would have increased the density of such aggregates, facilitating their export to abyssal depths (Wollenburg et al., 2018). More common ballasting minerals that are known to increase the density of phytoplankton aggregates and enhance their export include continental

Moved (insertion) [4]

Deleted: Our observations of large excesses of ^{234}Th relative to ^{238}U at around 70–100 m in the Eastern Grid (Stations 9 and 12) and Station 28-1 (Fig. 4) also support shallow remineralization of sinking particles during *Phaeocystis* blooms. ...

dust, calcium carbonate, and biogenic silica (Armstrong et al., 2002). Yet, to our knowledge, the role of those minerals in controlling the sinking of *Phaeocystis* aggregates have not been explored.

If insufficient ballasting limited export, then we would expect a low contribution of dense mineral phases such as bSi to sinking particles. In line with this, molar bSi to POC ratios in particles >51 µm within the PPZ were lower in East 1 (0.02–0.08) relative to those in Central (0.13) and East 2 (0.15–0.20), which likely suggests a change in phytoplankton community composition towards more diatoms in East 2. This observation is consistent with genomic samples taken during the cruise showing a lower relative contribution of diatoms to biomass in East 1 (Romanelli et al., 2026; Stevens-Green et al., 2024). Likewise, bSi fluxes at the base of the PPZ were 2-fold lower in East 1 than in Central and East 2 (Table 1), indicating a smaller contribution of siliceous plankton to export flux in East 1. However, below the PPZ, bSi fluxes in East 1 and East 2 became more similar, with fluxes within the same range at 500 m. Overall, bSi/POC ratios were relatively low in all areas, with averages from the base of the PPZ down to 500 m of 0.14 ± 0.03 (East 1), 0.18 ± 0.06 (Central), and 0.22 ± 0.04 (East 2), supporting a limited contribution of siliceous ballast in this study.

To our knowledge, bSi/POC ratios during the decline of other *Phaeocystis* blooms have not been reported. Our bSi/POC ratios fall in the lower range of those measured in the Labrador Sea (Lemaitre et al., 2018), the Porcupine Abyssal Plain (Clevenger et al., 2024), and the Crozet Plateau (Salter et al., 2007) during the decline of either diatom or mixed diatom and *Phaeocystis* blooms (Table 2). The bSi/POC ratios measured in this study are also lower than those found in the upper mesopelagic of Ocean Station Papa during low flux conditions (0.63 ± 0.28 ; Roca-Martí et al., 2021). Taken together, these results suggest that sinking particles during the BELAS-1 expedition were not heavily ballasted with bSi or, in other words, that diatoms did not appear to play an important ballasting role, which could at least partly explain the limited export of *Phaeocystis* in this study.

Other factors that may help explain why this *Phaeocystis* bloom did not lead to enhanced fluxes will be assessed elsewhere, including the analysis of the biochemical composition and morphology of suspended and sinking particles (Cisternas-Novoa et al., 2026) or their susceptibility to be degraded by heterotrophic bacteria (Romanelli et al., 2026).

4.3. BCP efficiency

The export efficiency (i.e., flux at PPZ base/integrated NPP within PPZ) and transfer efficiency (i.e., flux at 100 m below PPZ base/flux at PPZ base) are two common metrics for assessing the efficiency of the biological carbon pump and allowing comparisons between studies across different regions and seasons (Buesseler et al., 2020; Buesseler and Boyd, 2009). Here, we obtain export efficiencies ranging from 2 to 10% in East 1 and 32 to 34% in East 2 using in situ estimates of integrated NPP throughout the PPZ in combination with POC flux estimates measured at the base of that layer (Table 1, 2). Transfer efficiencies ranged from 48 to 139% and were on average 98% in East 1 and 92% in East 2 suggesting that flux attenuation throughout the upper 100 m of the mesopelagic zone was low. Combining both metrics, we obtain an overall BCP efficiency, defined as the amount of NPP reaching 100 m below the PPZ base, of 6% in East 1 and 30% in East 2. That would place this *Phaeocystis* bloom in the Labrador Sea as either a low BCP efficiency system (when looking at East 1 results) like the oligotrophic site ALOHA near Hawaii (Buesseler and Boyd, 2009), or a high BCP efficiency system (when looking at East 2

Deleted: this study, we measured both POC and bSi fluxes. M...

Deleted: agrees with

Deleted: data from

Deleted: Bertrand et al., in prep

Deleted: in prep.

Deleted: in prep.


Deleted: 9

Deleted: 1

Deleted: 3


Deleted: 29

792 results) similar to that found in the Kerguelen Plateau (Savoye et al., 2008) and in the Barents Sea (Andreassen
793 and Wassmann, 1998; Buesseler et al., 2020) during diatom and mixed diatom and *Phaeocystis* blooms (Fig. 8,
794 Table 2).



795
796 Figure 8. Biological carbon pump efficiency (BCP) comparing bloom studies across different regions (see Table 2). The
797 x-axis depicts transfer efficiency (flux at 100 m below PPZ base/flux at PPZ base) and the y-axis export efficiency (flux
798 at PPZ base/integrated NPP within PPZ). Contour lines represent overall BCP efficiency (percentages), and symbol
799 area is proportional to NPP. The "East 1" and "East 2" symbols and the diamond correspond to this study. The
800 diamond represents the overall BCP efficiency for the Labrador Sea (LS) 2022 bloom, calculated using satellite-derived
801 NPP integrated from the bloom peak to the final day of sampling. Studies are colour-coded according to the dominant
802 phytoplankton species. The ALOHA site (Buesseler and Boyd, 2009) is also shown as a low-end member. PPZ = primary
803 production zone, NPP = net primary production.

804 Yet, these BCP metrics are based on NPP and export estimates measured during the same time window. Given the
805 similar POC fluxes measured in East 1 and East 2, the increase in export efficiencies observed with time in the
806 Eastern Grid, from East 1 to East 2, is due to a decrease in NPP with time. This increase in export efficiencies
807 may reflect either 1) a more efficient BCP during post-bloom conditions, or more likely 2) a temporal decoupling



Deleted: are

Deleted:

Deleted: corresponding

813 between primary production and export (Henson et al., 2015; Kiorboe et al., 1996; Smetacek et al., 1984) which
814 have been shown to explain the inverse relationship between NPP and export efficiency found in different oceanic
815 regions (Laws and Maiti, 2019; Roca-Marti et al., 2017). In other words, in this study, the export measured during
816 post-bloom conditions in East 2, when NPP was low, could have been fueled with higher NPP rates that occurred
817 at an earlier phase of the bloom. It is important to keep in mind that the NPP rates presented in this study were
818 measured in the late stages of the bloom during its decline (East 1) and post-bloom phase (East 2) and therefore
819 do not reflect the higher rates that occurred before the field campaign.

820 In order to estimate an overall BCP efficiency for the 2022 *Phaeocystis* bloom in the Labrador Sea (see diamond
821 in Fig. 8), where NPP and export estimates are compared over longer timescales (Laws and Maiti, 2019), we have
822 used satellite-derived NPP (Devred et al., 2025) in the Eastern Grid area from the peak of the *Phaeocystis* bloom
823 until the last sampling day of the BELAS-1 expedition (9 May–2 June, Fig. S5 in the Supplement). The generally
824 good agreement between in situ and satellite-derived NPP over the course of the expedition (Fig. S5 in the
825 Supplement) provides confidence in the use of satellite NPP for calculating the overall efficiency. We have
826 obtained an integrated NPP during that period of 3920 mmol C m⁻² which, combined with the POC fluxes
827 measured in the Eastern Grid (average of 9.3 mmol C m⁻² d⁻¹ in East 1 and 2) multiplied by 25 d (i.e., 232 mmol
828 C m⁻²), results in an overall export efficiency for the *Phaeocystis* bloom decline of 6%. This estimate assumes that
829 the POC export measured in East 1 and 2 using ²³⁴Th (half-life = 24.1 d) is representative of the export that
830 occurred over the 25 d from the peak of the bloom until the end of the bloom. Using that export efficiency
831 combined with an average transfer efficiency in the Eastern Grid of 96% (East 1 and 2), we would obtain an
832 overall BCP efficiency of 6%, which is the same as that obtained in East 1 (Fig. 8). This analysis suggests that
833 East 1 results represent better the overall BCP efficiency of the extensive *Phaeocystis* bloom occurred in the
834 Labrador Sea in spring 2022 and stresses the importance of long-term observations of the BCP. The overall BCP
835 efficiency obtained in this study is lower than that reported during other blooms in the North Atlantic and Southern
836 Ocean, dominated by either diatoms or *Phaeocystis* and diatoms (Fig. 8, Table 2). Insufficient mineral ballasting
837 may have contributed to the low BCP efficiency observed during this massive *Phaeocystis* bloom.

838 Altogether, our findings highlight the complexity of quantifying the contribution of phytoplankton blooms to
839 carbon export and underscore the importance of sustained observations to better capture their variability and
840 broader implications in a changing ocean. This study provides a foundation for future comparisons with other
841 datasets from the BELAS-1 expedition. Integrating these datasets will help elucidate the mechanisms underlying
842 low BCP efficiency events, such as the one observed during this study, thereby improving our ability to predict
843 the consequences of shifting bloom dynamics for global carbon export. Moreover, future efforts will offer the
844 opportunity to extend the observational period by incorporating data from autonomous platforms, thereby
845 shedding light on the evolution of sinking fluxes beyond the limited timeframe of ship-based campaigns.

846 5. Conclusions

847 This study represents one of the largest assessments ever of the impact of *Phaeocystis* blooms on sinking fluxes
848 using ²³⁴Th as a tracer during a 2-week-long process study in spring 2022. The *Phaeocystis* bloom encountered

Deleted: global

Deleted: 6

Deleted: global

was unusually large and highly productive extending over more than half of the Labrador Sea for 6 weeks. The main conclusions of this work are summarized below:

- During the late stages of the bloom, POC fluxes in the upper mesopelagic down to 500 m were variable but overall moderate to high (average of $8 \pm 5 \text{ mmol C m}^{-2} \text{ d}^{-1}$). Yet, evidence of shallow POC flux remineralization combined with the fact that POC fluxes in the bloom were not higher than in a region sampled outside of the bloom (average of $13 \pm 3 \text{ mmol C m}^{-2} \text{ d}^{-1}$) suggests a limited role of *Phaeocystis* in carbon export.
- Large ($>51 \mu\text{m}$) particles collected using large volume pumps presented relatively low bSi/POC ratios and, therefore, diatoms did not appear to have an important ballasting role of *Phaeocystis*-derived material during the observation period.
- 2 weeks after the peak of the bloom, the total amount of in situ NPP that reached 100 m below the euphotic zone was only 6%. However, 3 weeks after the peak of the bloom, the value had increased to 30%. This apparent change was driven primarily by a decline in NPP over the sampling period, highlighting how sensitive the BCP efficiency determination is to temporal changes in NPP.
- Using satellite-derived NPP from the peak of the bloom until its end, we obtain an overall BCP efficiency of 6% supporting a low BCP efficiency system. We stress the importance of long-term observations of both NPP and POC export for estimating meaningful BCP efficiencies. Future research including data from autonomous platforms will elucidate how sinking fluxes might have changed after this ship-based observation study.
- The BCP efficiency of this *Phaeocystis* bloom is clearly lower than that found during the decline of either diatom blooms or mixed diatom and *Phaeocystis* blooms (albeit with no bSi/POC information) in the North Atlantic and Southern Ocean.
- To elucidate under which conditions *Phaeocystis* can be a good exporter, future research should include the analysis of bSi and other ballasting minerals, if possible, in all particle size fractions. These measurements will be key to shedding light on the role of the BCP in sequestering carbon in a future ocean where small cells, such as *Phaeocystis*, will increasingly dominate phytoplankton communities (Finkel et al., 2010; Passow and Carlson, 2012).

Data availability. Thorium-234 (^{234}Th) and size-fractionated particulate data were published open access (Roca-Martí et al., 2025a, b).

Author contributions. SK contributed to funding acquisition. MRM, MH, SK contributed to the experimental conceptual design. All authors contributed to data generation and analysis. MRM, MH, CM, SK contributed to the initial manuscript draft (MH data visualization), and all co-authors contributed to the revision of the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work is dedicated to Markus Kienast, a dear colleague whose approach to both science and life deeply inspires us all. We will always remember his passion, enthusiasm, positivity, and sense of humor. This study is part of the Ocean Frontier Institute's research project "The Northwest Atlantic Biological Carbon Pump" (NWA-BCP). We are sincerely grateful to the crew and scientific party aboard the RV *Celtic Explorer* during the BELAS-1 expedition and the whole NWA-BCP team. We would like to acknowledge Maria Armstrong

Deleted: 29

(²³⁴Th team), and Britton Dempsey and Brianna Stanley (NPP team) for their invaluable help with the expedition's preparation, sample collection and/or processing. We also thank Stephanie Clay for providing satellite data, Gretchen Swarr and Claire Normandeau for measuring ²³⁰Th/²²⁹Th and POC, and [Erin Bertrand](#), Ken Buesseler and Uta Passow for providing insightful discussions about the data. We thank OBPg-NASA for making satellite ocean colour data available and the support from the DFO Atlantic Zone Off-shelf Monitoring Program to collect in situ data used to develop satellite-based data products.

Financial support. MRM acknowledges support from the Ocean Frontier Institute (OFI) International Postdoctoral Fellowship Program, the Beatriu de Pinós Fellowship (2021-BP-00109), the "la Caixa" Foundation (ID 100010434, fellowship code LCF/BQ/PI24/12040022), [the Ramón y Cajal Program \(RYC2023-045355-L\)](#), and ICTA-UAB MERS (2021 SGR-640) of the Generalitat de Catalunya. MH was supported by the Nova Scotia Graduate Scholarship (NSGS), and SK by the Natural Sciences and Engineering Research Council of Canada (NSERC) and OFI.

References

- Alcolombri, U., Nissan, A., Slomka, J., Charlton, S., Secchi, E., Short, I., Lee, K. S., Peaudecerf, F. J., Baumgartner, D. A., Sichert, A., Sauer, U., Sengupta, A., and Stocker, R.: Biogel scavenging slows the sinking of organic particles to the ocean depths, *Nat. Commun.* 2025 161, 16, 1–10, <https://doi.org/10.1038/s41467-025-57982-5>, 2025.
- Andreassen, I. J. and Wassmann, P.: Vertical flux of phytoplankton and particulate biogenic matter in the marginal ice zone of the Barents Sea in May 1993, *Mar. Ecol. Prog. Ser.*, 170, 1–14, <https://doi.org/10.3354/MEPS170001>, 1998.
- Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G.: A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 49, 219–236, [https://doi.org/10.1016/S0967-0645\(01\)00101-1](https://doi.org/10.1016/S0967-0645(01)00101-1), 2002.
- Arruda, R., Atamanchuk, D., Boteler, C., and Wallace, D. W. R.: Seasonality of pCO₂ and air-sea CO₂ fluxes in the Central Labrador Sea, *Front. Mar. Sci.*, 11, 1472697, <https://doi.org/10.3389/fmars.2024.1472697>, 2024.
- Asper, V. L. and Smith, W. O.: Particle fluxes during austral spring and summer in the southern Ross Sea, Antarctica, *J. Geophys. Res. Ocean.*, 104, 5345–5359, <https://doi.org/10.1029/1998JC900067>, 1999.
- Baker, C. A., Martin, A. P., Yool, A., and Popova, E.: Biological Carbon Pump Sequestration Efficiency in the North Atlantic: A Leaky or a Long-Term Sink?, *Global Biogeochem. Cycles*, 36, e2021GB007286, <https://doi.org/10.1029/2021GB007286>, 2022.
- Baltar, F., Aristegui, J., Gasol, J. M., Sintes, E., and Herndl, G. J.: Evidence of prokaryotic metabolism on suspended particulate organic matter in the dark waters of the subtropical North Atlantic, *Limnol. Oceanogr.*, 54, 182–193, <https://doi.org/10.4319/LO.2009.54.1.0182>, 2009.
- Baumas, C., Fuchs, R., Garel, M., Poggiale, J.-C., Memery, L., Le Moigne, F. A. C., and Tamburini, C.: Reconstructing the ocean's mesopelagic zone carbon budget: sensitivity and estimation of parameters associated with prokaryotic remineralization, *Biogeosciences*, 20, 4165–4182, <https://doi.org/10.5194/BG-20-4165-2023>, 2023.
- Bishop, J. K. B., Lam, P. J., and Wood, T. J.: Getting good particles: Accurate sampling of particles by large volume in-situ filtration, *Limnol. Oceanogr. Methods*, 10, 681–710, <https://doi.org/10.4319/lom.2012.10.681>, 2012.
- Boyd, P. W., Sherry, N. D., Berges, J. A., Bishop, J. K. B., Calvert, S. E., Charette, M. A., Giovannoni, S. J., Goldblatt, R., Harrison, P. J., Moran, S. B., Roy, S., Soon, M., Strom, S., Thibault, D., Vergin, K. L., Whitney, F. A., and Wong, C. S.: Transformations of biogenic particulates from the pelagic to the deep ocean realm, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 46, 2761–2792, [https://doi.org/10.1016/S0967-0645\(99\)00083-1](https://doi.org/10.1016/S0967-0645(99)00083-1), 1999.
- Boyd, P. W., Claustre, H., Levy, M., Siegel, D. A., and Weber, T.: Multi-faceted particle pumps drive carbon sequestration in the ocean, *Nature*, 568, 327–335, <https://doi.org/10.1038/s41586-019-1098-2>, 2019.

Deleted: the ICTA-UAB "Maria de Maeztu" Program for Units of Excellence of the Spanish Ministry of Science and Innovation (CEX2024-001506-M funded by MICIU/AEI /10.13039/501100011033)

940 de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the
941 global ocean: An examination of profile data and a profile-based climatology, *J. Geophys. Res.*, 109, C12003,
942 <https://doi.org/10.1029/2004JC002378>, 2004.

943 Buesseler, K. O. and Boyd, P. W.: Shedding light on processes that control particle export and flux attenuation
944 in the twilight zone of the open ocean, *Limnol. Oceanogr.*, 54, 1210–1232,
945 <https://doi.org/10.4319/lo.2009.54.4.1210>, 2009.

946 Buesseler, K. O., Boyd, P. W., Black, E. E., and Siegel, D. A.: Metrics that matter for assessing the ocean
947 biological carbon pump, *Proc. Natl. Acad. Sci.*, 117, 9679–9687, <https://doi.org/10.1073/PNAS.1918114117>,
948 2020.

949 Burd, A. B., Hansell, D. A., Steinberg, D. K., Anderson, T. R., Aristegui, J., Baltar, F., Beupré, S. R.,
950 Buesseler, K. O., DeHairs, F., Jackson, G. A., Kadko, D. C., Koppelman, R., Lampitt, R. S., Nagata, T.,
951 Reinthaler, T., Robinson, C., Robison, B. H., Tamburini, C., and Tanaka, T.: Assessing the apparent imbalance
952 between geochemical and biochemical indicators of meso- and bathypelagic biological activity: What the @\$#!
953 is wrong with present calculations of carbon budgets?, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 57, 1557–
954 1571, <https://doi.org/10.1016/j.dsr2.2010.02.022>, 2010.

955 Ceballos-Romero, E., De Soto, F., Le Moigne, F. A. C., García-Tenorio, R., and Villa-Alfageme, M.: ²³⁴Th-
956 derived particle fluxes and seasonal variability: when is the SS assumption reliable? Insights from a novel
957 approach for carbon flux simulation, *Geophys. Res. Lett.*, <https://doi.org/10.1029/2018GL079968>, 2018.

958 [Cisternas-Novoa, C., Romanelli, E., and Passow, U.: Differences between suspended and sinking particles
959 regulate carbon flux in the upper mesopelagic during a *Phaeocystis* Bloom,
960 <https://doi.org/10.64898/2026.06.09.731151>, 2026.](https://doi.org/10.64898/2026.06.09.731151)

961 Claustre, H., Legendre, L., Boyd, P. W., and Levy, M.: The Oceans' Biological Carbon Pumps: Framework for
962 a Research Observational Community Approach, *Front. Mar. Sci.*, 8,
963 <https://doi.org/10.3389/fmars.2021.780052>, 2021.

964 Clevenger, S. J., Benitez-Nelson, C. R., Drysdale, J., Pike, S., Puigcorb , V., and Buesseler, K. O.: Review of
965 the analysis of ²³⁴Th in small volume (2–4 L) seawater samples: improvements and recommendations, *J.*
966 *Radioanal. Nucl. Chem.*, 329, 1–13, <https://doi.org/10.1007/s10967-021-07772-2>, 2021.

967 Clevenger, S. J., Benitez-Nelson, C. R., Roca-Mart , M., Bam, W., Estapa, M., Kenyon, J. A., Pike, S.,
968 Resplandy, L., Wyatt, A., and Buesseler, K. O.: Carbon and silica fluxes during a declining North Atlantic
969 spring bloom as part of the EXPORTS program, *Mar. Chem.*, 258, 104346,
970 <https://doi.org/10.1016/J.MARCHEM.2023.104346>, 2024.

971 Coppola, L., Roy-Barman, M., Wassmann, P., Mulsow, S., and Jeandel, C.: Calibration of sediment traps and
972 particulate organic carbon export using ²³⁴Th in the Barents Sea, *Mar. Chem.*, 80, 11–26,
973 [https://doi.org/10.1016/S0304-4203\(02\)00071-3](https://doi.org/10.1016/S0304-4203(02)00071-3), 2002.

974 Devred, E., Wilson, K. L., Perry, T., Hardy, M., Brosnahan, M., and Ringuette, M.: Identification and validation
975 of phytoplankton taxonomic assemblages derived from pigment signatures using samples collected in the
976 Labrador Sea from 2014 to 2022, Dartmouth, Nova Scotia, Canada, 37 pp., 2024.

977 Devred, E., Clay, S., Ringuette, M., Perry, T., Amirian, M., Irwin, A., and Finkel, Z.: Net primary production in
978 the Labrador Sea between 2014 and 2022 derived from ocean colour remote sensing based on ecological
979 regimes, *Remote Sens. Environ.*, 323, 114713, <https://doi.org/10.1016/J.RSE.2025.114713>, 2025.

980 Dybwad, C., Assmy, P., Olsen, L. M., Peeken, I., Nikolopoulos, A., Krumpen, T., Randelhoff, A., Taterek, A.,
981 Wiktor, J. M., and Reigstad, M.: Carbon Export in the Seasonal Sea Ice Zone North of Svalbard From Winter to
982 Late Summer, *Front. Mar. Sci.*, 7, 525800, <https://doi.org/10.3389/fmars.2020.525800>, 2021.

983 Finkel, Z. V., Beardall, J., Flynn, K. J., Quigg, A., Rees, T. A. V., and Raven, J. A.: Phytoplankton in a
984 changing world: cell size and elemental stoichiometry, *J. Plankton Res.*, 32, 119–137,
985 <https://doi.org/10.1093/PLANKT/FBP098>, 2010.

986 Giering, S. L. C., Sanders, R., Lampitt, R. S., Anderson, T. R., Tamburini, C., Boutrif, M., Zubkov, M. V.,
987 Marsay, C. M., Henson, S. A., Saw, K., Cook, K., and Mayor, D. J.: Reconciliation of the carbon budget in the

988 ocean's twilight zone., *Nature*, 507, 480–483, <https://doi.org/10.1038/nature13123>, 2014.

989 [Giering, S. L. C., Sanders, R., Martin, A. P., Lindemann, C., Möller, K. O., Daniels, C. J., Mayor, D. J., and St.](#)
990 [John, M. A.: High export via small particles before the onset of the North Atlantic spring bloom, *J. Geophys.*](#)
991 [Res. Ocean., 121, 6929–6945, <https://doi.org/10.1002/2016JC012048>, 2016.](#)

992 Graff, J. R., Nelson, N. B., Roca-Martí, M., Romanelli, E., Kramer, S. J., Erickson, Z., Cetinic, I., Buesseler, K.
993 O., Passow, U., Zhang, X., Benitez-Nelson, C., Bisson, K., Close, H. G., Crockford, T., Fox, J., Halewood, S.,
994 Lam, P., Roesler, C., Sweet, J., VerWey, B., Xiong, Y., and Siegel, D. A.: Reconciliation of total particulate
995 organic carbon and nitrogen measurements determined using contrasting methods in the North Pacific Ocean as
996 part of the NASA EXPORTS field campaign, *Elem. Sci. Anthr.*, 11,
997 <https://doi.org/10.1525/elementa.2022.00112>, 2023.

998 Hama, T., Miyazaki, T., Ogawa, Y., Iwakuma, T., Takahashi, M., Otsuki, A., and Ichimura, S.: Measurement of
999 photosynthetic production of a marine phytoplankton population using a stable ^{13}C isotope, *Mar. Biol.*, 73, 31–
1000 36, <https://doi.org/10.1007/BF00396282>/METRICS, 1983.

1001 Henson, S. A., Yool, A., and Sanders, R.: Variability in efficiency of particulate organic carbon export: A model
1002 study, *Global Biogeochem. Cycles*, 29, 33–45, <https://doi.org/10.1002/2014GB004965>, 2015.

1003 Henson, S. A., Laufkötter, C., Leung, S., Giering, S. L. C., Palevsky, H. I., and Cavan, E. L.: Uncertain response
1004 of ocean biological carbon export in a changing world, *Nat. Geosci.* 2022 154, 15, 248–254,
1005 <https://doi.org/10.1038/s41561-022-00927-0>, 2022.

1006 [Kiorboe, T., Hansen, J. L. S., Alldredge, A. L., Jackson, G. A., Passow, U., Dam, H. G., Drapeau, D. T., Waite,](#)
1007 [A., and Garcia, C. M.: Sedimentation of phytoplankton during a diatom bloom: Rates and mechanisms, *J. Mar.*](#)
1008 [Res., 54, 1123–148, 1996.](#)

1009 Knap, A., Michaels, A., Close, A., Ducklow, H., and Dickson, A.: Protocols for the Joint Global Ocean Flux
1010 Study (JGOFS) Core Measurements. JGOFS Report Nr. 19, 1–210 pp., 1996.

1011 Lalande, C., Moran, S. B., Wassmann, P., Grebmeier, J. M., and Cooper, L. W.: ^{234}Th -derived particulate
1012 organic carbon fluxes in the northern Barents Sea with comparison to drifting sediment trap fluxes, *J. Mar.*
1013 *Syst.*, 73, 103–113, <https://doi.org/10.1016/j.jmarsys.2007.09.004>, 2008.

1014 Lalande, C., Bauerfeind, E., and Nöthig, E. M.: Downward particulate organic carbon export at high temporal
1015 resolution in the eastern Fram Strait: influence of Atlantic Water on flux composition, *Mar. Ecol. Prog. Ser.*,
1016 440, 127–136, <https://doi.org/10.3354/MEPS09385>, 2011.

1017 Lam, P. J., Ohnemus, D. C., and Auro, M. E.: Size-fractionated major particle composition and concentrations
1018 from the US GEOTRACES North Atlantic Zonal Transect, Deep Sea Res. Part II Top. Stud. Oceanogr., 116,
1019 303–320, <https://doi.org/10.1016/J.DSR2.2014.11.020>, 2015.

1020 Lam, P. J., Lee, J.-M., Heller, M. I., Mehic, S., Xiang, Y., and Bates, N. R.: Size-fractionated distributions of
1021 suspended particle concentration and major phase composition from the U.S. GEOTRACES Eastern Pacific
1022 Zonal Transect (GP16), *Mar. Chem.*, 201, 90–107, <https://doi.org/10.1016/j.marchem.2017.08.013>, 2018.

1023 Laws, E. A. and Maiti, K.: The relationship between primary production and export production in the ocean:
1024 Effects of time lags and temporal variability, Deep Sea Res. Part I Oceanogr. Res. Pap.,
1025 <https://doi.org/10.1016/J.DSR.2019.05.006>, 2019.

1026 Lemaitre, N.: Multi-proxy approach (Thorium-234, excess Barium) of export and remineralization fluxes of
1027 carbon and biogenic elements associated with the oceanic biological pump, Université de Bretagne Occidentale,
1028 319 pp., 2017.

1029 Lemaitre, N., Planchon, F., Planquette, H., Dehairs, F., Fonseca-Batista, D., Roukaerts, A., Deman, F., Tang, Y.,
1030 Mariez, C., and Sarthou, G.: High variability of particulate organic carbon export along the North Atlantic
1031 GEOTRACES section GA01 as deduced from ^{234}Th fluxes, *Biogeosciences*, 15, 6417–6437,
1032 <https://doi.org/10.5194/bg-15-6417-2018>, 2018.

1033 Maiti, K., Buesseler, K. O., Pike, S. M., Benitez-Nelson, C., Cai, P., Chen, W., Cochran, K., Dai, M., Dehairs,
1034 F., Gasser, B., Kelly, R. P., Masque, P., Miller, L. A., Miquel, J. C., Moran, S. B., Morris, P. J., Peine, F.,

1035 Planchon, F., Renfro, A. A., van der Loeff, M. R., Santschi, P. H., Turnewitsch, R., Waples, J. T., and Xu, C.:
 1036 Intercalibration studies of short-lived thorium-234 in the water column and marine particles, *Limnol. Oceanogr.*
 1037 *Methods*, 10, 631–644, <https://doi.org/10.4319/lom.2012.10.631>, 2012.

1038 Mari, X., Passow, U., Migon, C., Burd, A. B., and Legendre, L.: Transparent exopolymer particles: Effects on
 1039 carbon cycling in the ocean, *Prog. Oceanogr.*, 151, 13–37, <https://doi.org/10.1016/J.POCEAN.2016.11.002>,
 1040 2017.

1041 Le Moigne, F. A. C., Poulton, A. J., Henson, S. A., Daniels, C. J., Fragoso, G. M., Mitchell, E., Richier, S.,
 1042 Russell, B. C., Smith, H. E. K., Tarling, G. A., Young, J. R., and Zubkov, M.: Carbon export efficiency and
 1043 phytoplankton community composition in the Atlantic sector of the Arctic Ocean, *J. Geophys. Res. Ocean.*, 120,
 1044 3896–3912, <https://doi.org/10.1002/2015JC010700>, 2015.

1045 Moran, S. B., Weinstein, S. E., Edmonds, H. N., Smith, J. N., Kelly, R. P., Pilson, M. E. Q., and Harrison, W.
 1046 G.: Does $^{234}\text{Th}/^{238}\text{U}$ disequilibrium provide an accurate record of the export flux of particulate organic carbon
 1047 from the upper ocean?, *Limnol. Oceanogr.*, 48, 1018–1029, <https://doi.org/10.4319/lo.2003.48.3.1018>, 2003.

1048 Morris, P. J., Sanders, R., Turnewitsch, R., and Thomalla, S.: ^{234}Th -derived particulate organic carbon export
 1049 from an island-induced phytoplankton bloom in the Southern Ocean, *Deep Sea Res. Part II Top. Stud.*
 1050 *Oceanogr.*, 54, 2208–2232, <https://doi.org/10.1016/j.dsr2.2007.06.002>, 2007.

1051 Ohashi, K., Laurent, A., Renkl, C., Sheng, J., Fennel, K., and Oliver, E.: DalROMS-NWA12 v1.0, a coupled
 1052 circulation-ice-biogeochemistry modelling system for the northwest Atlantic Ocean: Development and
 1053 validation, *Geosci. Model Dev.*, 17, 8697–8733, <https://doi.org/10.5194/GMD-17-8697-2024>, 2024.

1054 Owens, S. A., Buesseler, K. O., and Sims, K. W. W.: Re-evaluating the ^{238}U -salinity relationship in seawater:
 1055 Implications for the ^{238}U – ^{234}Th disequilibrium method, *Mar. Chem.*, 127, 31–39,
 1056 <https://doi.org/10.1016/j.marchem.2011.07.005>, 2011.

1057 Owens, S. A., Pike, S., and Buesseler, K. O.: Thorium-234 as a tracer of particle dynamics and upper ocean
 1058 export in the Atlantic Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 116, 42–59,
 1059 <https://doi.org/10.1016/j.dsr2.2014.11.010>, 2015.

1060 Parsons, T. R., Maita, Y., and Lalli, C. M.: A manual of chemical and biological methods for seawater analysis,
 1061 Pergamon Press, Oxford, UK, 173 pp., <https://doi.org/10.25607/OBP-1830>, 1984.

1062 Passow, U.: Transparent exopolymer particles (TEP) in aquatic environments, *Prog. Oceanogr.*, 55, 287–333,
 1063 [https://doi.org/10.1016/S0079-6611\(02\)00138-6](https://doi.org/10.1016/S0079-6611(02)00138-6), 2002.

1064 Passow, U. and Carlson, C.: The biological pump in a high CO_2 world, *Mar. Ecol. Prog. Ser.*, 470, 249–271,
 1065 <https://doi.org/10.3354/meps09985>, 2012.

1066 Passow, U. and Wassmann, P.: On the trophic fate of *Phaeocystis pouchetii* (Hariot): IV. The formation of
 1067 marine snow by *P. pouchetii* on JSTOR, *Mar. Ecol. Prog. Ser.*, 104, 153–161, 1994.

1068 Passow, U. and Weber, T.: The biological carbon pump, in: *Treatise on Geochemistry (Third Edition)*, vol. 4,
 1069 Elsevier, 333–369, <https://doi.org/10.1016/B978-0-323-99762-1.00031-0>, 2025.

1070 Puigcorb , V., Roca-Mart , M., Masqu , P., Benitez-Nelson, C., Rutgers van der Loeff, M., Bracher, A., and
 1071 Moreau, S.: Latitudinal distributions of particulate carbon export across the North Western Atlantic Ocean,
 1072 *Deep Sea Res. Part I Oceanogr. Res. Pap.*, 129, 116–130, <https://doi.org/10.1016/J.DSR.2017.08.016>, 2017.

1073 Puigcorb , V., Masqu , P., and Le Moigne, F. A. C.: Global database of ratios of particulate organic carbon to
 1074 thorium-234 in the ocean: improving estimates of the biological carbon pump, *Earth Syst. Sci. Data*, 12, 1267–
 1075 1285, <https://doi.org/10.5194/essd-12-1267-2020>, 2020.

1076 Reigstad, M. and Wassmann, P.: Does *Phaeocystis* spp. contribute significantly to vertical export of organic
 1077 carbon?, *Biogeochemistry*, 83, 217–234, <https://doi.org/10.1007/s10533-007-9093-3>, 2007.

1078 Reinthaler, T., Van Aken, H., Veth, C., Aristegui, J., Robinson, C., Williams, P. J. L. B., Lebaron, P., and
 1079 Hemdl, G. J.: Prokaryotic respiration and production in the meso- and bathypelagic realm of the eastern and
 1080 western North Atlantic basin, *Limnol. Oceanogr.*, 51, 1262–1273, <https://doi.org/10.4319/LO.2006.51.3.1262>,

1081 2006.

1082 Resplandy, L., Martin, A. P., Le Moigne, F., Martin, P., Aquilina, A., Mémary, L., Lévy, M., and Sanders, R.:
 1083 How does dynamical spatial variability impact ^{234}Th -derived estimates of organic export?, *Deep Sea Res. Part I*
 1084 *Oceanogr. Res. Pap.*, 68, 24–45, <https://doi.org/10.1016/j.dsr.2012.05.015>, 2012.

1085 [Riley, J. S., Sanders, R., Marsay, C., Le Moigne, F. A. C., Achterberg, E. P., and Poulton, A. J.: The relative](https://doi.org/10.1029/2011GB004085)
 1086 [contribution of fast and slow sinking particles to ocean carbon export, *Global Biogeochem. Cycles*, 26,](https://doi.org/10.1029/2011GB004085)
 1087 <https://doi.org/10.1029/2011GB004085>, 2012.

1088 Roca-Martí, M. and Puigcorbó, V.: Combined Use of Short-Lived Radionuclides (^{234}Th and ^{210}Po) as Tracers of
 1089 Sinking Particles in the Ocean, *Ann. Rev. Mar. Sci.*, 16, 551–575, [https://doi.org/10.1146/annurev-marine-](https://doi.org/10.1146/annurev-marine-041923-013807)
 1090 [041923-013807](https://doi.org/10.1146/annurev-marine-041923-013807), 2024.

1091 Roca-Martí, M., Puigcorbó, V., Iversen, M. H., Rutgers van der Loeff, M. M., Klaas, C., Cheah, W., Bracher,
 1092 A., and Masqué, P.: High particulate organic carbon export during the decline of a vast diatom bloom in the
 1093 Atlantic sector of the Southern Ocean, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 138, 102–115,
 1094 <https://doi.org/10.1016/j.dsr2.2015.12.007>, 2017.

1095 Roca-Martí, M., Benitez-Nelson, C. R., Umhau, B. P., Wyatt, A. M., Clevenger, S. J., Pike, S., Horner, T. J.,
 1096 Estapa, M. L., Resplandy, L., and Buesseler, K. O.: Concentrations, ratios, and sinking fluxes of major
 1097 bioelements at Ocean Station Papa, *Elem. Sci. Anthr.*, 9, <https://doi.org/10.1525/elementa.2020.00166>, 2021.

1098 Roca-Martí, M., Healey, M., and Kienast, S. S.: Particulate Th-234, organic carbon, organic nitrogen and
 1099 biogenic silica concentrations in size-fractionated particles in the Labrador Sea in spring 2022 [dataset],
 1100 PANGAEA, <https://doi.org/10.1594/PANGAEA.983961>, 2025a.

1101 Roca-Martí, M., Healey, M., and Kienast, S. S.: Total (dissolved + particulate) Th-234 and U-238 activities in
 1102 seawater in the Labrador Sea in spring 2022 [dataset], PANGAEA, <https://doi.org/10.1594/PANGAEA.983957>,
 1103 2025b.

1104 Romanelli, E., Sweet, J., Giering, S. L. C., Siegel, D. A., and Passow, U.: The importance of transparent
 1105 exopolymer particles over ballast in determining both sinking and suspension of small particles during late
 1106 summer in the Northeast Pacific Ocean, *Elementa*, 11,
 1107 <https://doi.org/10.1525/ELEMENTA.2022.00122/197269>, 2023.

1108 [Romanelli, E., Stevens-Green, R., Cisternas-Novoa, C., LaRoche, J., Siegel, D. A., Carlson, C. A., and Passow,](https://doi.org/10.64898/2026.04.19.716305)
 1109 [U.: Particle lability drives degradation dynamics and bacterial community assembly during a *Phaeocystis* bloom](https://doi.org/10.64898/2026.04.19.716305)
 1110 [decline, *https://doi.org/10.64898/2026.04.19.716305*, 2026.](https://doi.org/10.64898/2026.04.19.716305)

1111 Salter, I., Lampitt, R. S., Sanders, R., Poulton, A., Kemp, A. E. S., Boorman, B., Saw, K., and Pearce, R.:
 1112 Estimating carbon, silica and diatom export from a naturally fertilised phytoplankton bloom in the Southern
 1113 Ocean using PELAGRA: A novel drifting sediment trap, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 54, 2233–
 1114 2259, <https://doi.org/10.1016/j.dsr2.2007.06.008>, 2007.

1115 Sanders, R., Henson, S. A., Koski, M., De La Rocha, C. L., Painter, S. C., Poulton, A. J., Riley, J., Salihoglu, B.,
 1116 Visser, A., Yool, A., Bellerby, R., and Martin, A. P.: The Biological Carbon Pump in the North Atlantic, *Prog.*
 1117 *Oceanogr.*, 129, 200–218, <https://doi.org/10.1016/j.pocean.2014.05.005>, 2014.

1118 Savoye, N., Benitez-Nelson, C., Burd, A. B., Cochran, J. K., Charette, M., Buesseler, K. O., Jackson, G. A.,
 1119 Roy-Barman, M., Schmidt, S., and Elskens, M.: ^{234}Th sorption and export models in the water column: A
 1120 review, *Mar. Chem.*, 100, 234–249, <https://doi.org/10.1016/j.marchem.2005.10.014>, 2006.

1121 Savoye, N., Trull, T. W., Jacquet, S. H. M., Navez, J., and Dehairs, F.: ^{234}Th -based export fluxes during a
 1122 natural iron fertilization experiment in the Southern Ocean (KEOPS), *Deep Sea Res. Part II Top. Stud.*
 1123 *Oceanogr.*, 55, 841–855, <https://doi.org/10.1016/j.dsr2.2007.12.036>, 2008.

1124 Schoemann, V., Becquevort, S., Stefels, J., Rousseau, V., and Lancelot, C.: *Phaeocystis* blooms in the global
 1125 ocean and their controlling mechanisms: a review, *J. Sea Res.*, 53, 43–66,
 1126 <https://doi.org/10.1016/J.SEAES.2004.01.008>, 2005.

1127 [Smetacek, V., von Bodungen, B., Knoppers, B., Peinert, R., Pollehne, F., Stegmann, P., and Zeitzschel, B.:](https://doi.org/10.1016/J.SEAES.2004.01.008)

1128 [Seasonal stages characterizing the annual cycle of an inshore pelagic system, Rapp. Proces-Verbaux des Reun.](#)
1129 [Cons. Int. pour l'Exploration la Mer, 183, 126–135, 1984.](#)

1130 Smith, W. O. and Trimborn, S.: *Phaeocystis*: A Global Enigma, *Ann. Rev. Mar. Sci.*, 16, 417–441,
1131 <https://doi.org/10.1146/annurev-marine-022223-025031>, 2024.

1132 Steinberg, D. K., Van Mooy, B. A. S., Buesseler, K. O., Boyd, P. W., Kobari, T., and Karl, D. M.: Bacterial vs.
1133 zooplankton control of sinking particle flux in the ocean's twilight zone, *Limnol. Oceanogr.*, 53, 1327–1338,
1134 <https://doi.org/10.4319/lo.2008.53.4.1327>, 2008.

1135 Stephens, B. M., Roca-Martí, M., Maas, A. E., Amaral, V. J., Clevenger, S., Traylor, S., Benitez-Nelson, C. R.,
1136 Boyd, P. W., Buesseler, K. O., Carlson, C. A., Cassar, N., Estapa, M., Fassbender, A. J., Huang, Y., Lam, P. J.,
1137 Marchal, O., Menden-Deuer, S., Paul, N. L., Santoro, A. E., Siegel, D. A., and Nicholson, D. P.: An upper-
1138 mesopelagic-zone carbon budget for the subarctic North Pacific, *Biogeosciences*, 22, 3301–3328,
1139 <https://doi.org/10.5194/BG-22-3301-2025>, 2025.

1140 [Stevens-Green, R., Romanelli, E., Cisternas-Novoa, C., Pollara, S., Jabre, L., Passow, U., Bertrand, E. M.,](#)
1141 [LaRoche, J., Stevens-Green, R., Romanelli, E., Cisternas-Novoa, C., Pollara, S., Jabre, L., Passow, U., Bertrand,](#)
1142 [E. M., and LaRoche, J.: Exploring the Microbial Communities Involved in the Biological Carbon Pump](#)
1143 [Throughout the Decline of a Large *Phaeocystis pouchetii* Bloom in the Labrador Sea, in: Ocean Sciences](#)
1144 [Meeting \(OSM\) 2024, 2024.](#)

1145 Tesdal, J. E., Ducklow, H. W., Goes, J. I., and Yashayaev, I.: Recent nutrient enrichment and high biological
1146 productivity in the Labrador Sea is tied to enhanced winter convection, *Prog. Oceanogr.*, 206, 102848,
1147 <https://doi.org/10.1016/J.POCEAN.2022.102848>, 2022.

1148 Volk, T. and Hoffert, M. I.: Ocean carbon pumps: analysis of relative strengths and efficiencies in ocean-driven
1149 atmospheric CO₂ changes, in: *The Carbon Cycle and Atmospheric CO₂: Natural Variations Archean to Present*,
1150 vol. 32, edited by: Sundquist, E. T. and Broecker, W. S., American Geophysical Union, Washington, D. C., 99–
1151 110, <https://doi.org/10.1029/GM032>, 1985.

1152 Wiedmann, I., Ceballos-Romero, E., Villa-Alfágeme, M., Renner, A. H. H., Dybwad, C., Jagt, H., Svensen, C.,
1153 Assmy, P., Wiktor, J. M., Tatarek, A., Różańska-Pluta, M., and Iversen, M. H.: Arctic Observations Identify
1154 Phytoplankton Community Composition as Driver of Carbon Flux Attenuation, *Geophys. Res. Lett.*, 47,
1155 e2020GL087465, <https://doi.org/10.1029/2020GL087465>, 2020.

1156 Wollenburg, J. E., Katlein, C., Nehrke, G., Nöthig, E.-M., Matthiessen, J., Wolf- Gladrow, D. A., Nikolopoulos,
1157 A., Gázquez-Sanchez, F., Rossmann, L., Assmy, P., Babin, M., Bruyant, F., Beaulieu, M., Dybwad, C., and
1158 Peeken, I.: Ballasting by cryogenic gypsum enhances carbon export in a *Phaeocystis* under-ice bloom, *Sci. Rep.*,
1159 8, 7703, <https://doi.org/10.1038/s41598-018-26016-0>, 2018.

1160 Yashayaev, I.: Intensification and shutdown of deep convection in the Labrador Sea were caused by changes in
1161 atmospheric and freshwater dynamics, *Commun. Earth Environ.*, 5, [https://doi.org/10.1038/S43247-024-01296-](https://doi.org/10.1038/S43247-024-01296-9)
1162 9, 2024.

1163