



The Observed Evolution of Arctic Amplification over the Past 45 Years

Mark C. Serreze¹, Elizabeth Cassano^{1,2}, Alex Crawford³, John J. Cassano^{1,2}, and Chen Zhang¹

¹Cooperative Institute for Research in Environmental Sciences, National Snow and Ice Data Center, University of Colorado Boulder CO

²Department of Atmospheric and Oceanic Sciences, University of Colorado Boulder, Boulder, CO, USA

³Centre for Earth Observation Science, Department of Environment and Geography, University of Manitoba, Winnipeg

Correspondence to: Mark Serreze (mark.serreze@colorado.edu)

Abstract. To address research gaps in understanding Arctic Amplification, we use data from ERA5 and sea ice concentration to examine the seasonal, spatial and decadal evolution of Arctic 2-meter and lower tropospheric temperatures and lower tropospheric (surface to 850 hPa) static stability over the past 45 years. A Local Amplification Anomaly (LAA) metric is used to examine how spatial patterns of Arctic 2-meter temperature anomalies compare to anomalies for the globe as a whole. Pointing to impacts of seasonally-delayed albedo feedback, growing areas of end-of-summer (September) open water largely co-locate with the strongest positive anomalies of 2-meter temperatures through autumn and winter and their growth through time; small summer trends reflect the effects of a melting sea ice cover. Because of seasonal ice growth, the association between rising 2-meter temperatures and sea ice weakens from autumn into winter, except in the the Barents Sea where there have been prominent downward trends in winter ice extent. Imprints of variable atmospheric circulation are prominent in the Arctic temperature evolution. Low-level (surface to 850 hPa) stability over the Arctic increases from autumn through winter, consistent with the greater depth of surface-based atmospheric heating seen in autumn. However, trends towards weaker static stability dominate the Arctic Ocean in autumn and winter, especially over areas of September and wintertime ice loss. Sea ice thinning, leading to increased conductive heat fluxes through the ice, likely also contributes to reduced stability.

1 Introduction

Arctic amplification (AA) refers to the observation that, over the last several decades, the rate of increase in surface air temperature over the Arctic region has been larger than for the globe as a whole. AA was predicted as a consequence of global warming even in the earliest generation of climate models, and was envisioned as far back as the 19th century (Arrhenius, 1896). Various studies have placed the ratio of Arctic to global warming from two to four, with differences relating to the definition of the Arctic region, data used, the time period examined and the season examined (Yu et al., 2021; Walsh 2014; Richter Menge and Druckenmiller, 2020; Jansen et al., 2020; AMAP, 2021; Rantanen et al., 2022). Using several observational data sets and defining the Arctic as the region poleward of the Arctic Circle, Rantanen et al. (2022) find a factor of four



warming relative to the globe over the period 1979–2021 based on annual mean temperatures. From comparisons with climate models, they conclude that this large ratio is either an extremely unlikely event, or that the models systematically underestimate AA. Zhou et al. (2024) conclude that the externally forced amplification is three-fold, with natural variability explaining the remainder.

35 Growing spring and summer sea ice loss, leading to more seasonal heat gain in the ocean mixed layer and subsequent upward heat release in autumn and winter - a seasonally-delayed expression of albedo feedback - is widely accepted as a key driver of AA (Perovich et al., 2007; Steele et al., 2008; Serreze et al., 2009; Screen and Simmonds, 2010a,b; Stammerjohn et al., 2012; Stroeve et al., 2014; Dai et al., 2019). However, based on observations and modeling studies, AA is also recognized as involving a suite of connected contributions including changes in poleward energy transport (Graversen et al., 2016; Woods and Caballero, 2016; Previdi et al., 2021), Planck feedback (Pithan and Mauritsen, 2014), positive lapse rate feedback (Pithan and Mauritsen, 2014; Stuecker et al., 2018; Previdi et al., 2021), changes in ocean heat transport (Beer et al., 2020), changes in autumn cloud cover (Kay and Gettelman, 2009; Wu and Lee, 2012) and even reduced air pollution in Europe (Navarro et al., 2016; Krishnan et al., 2020).

However, much remains to be understood about the seasonal expressions of AA, its regional expressions, and overall evolution. Here, using data from the ERA5 reanalysis and satellite-derived sea ice concentration, we focus on understanding the decadal evolution and seasonal/spatial expressions of Arctic temperature anomalies. We show how: 1) the pronounced autumn contribution to AA, through which internal energy gained by the upper ocean in spring and summer in growing open water areas is subsequently released back to the atmosphere, decays into winter as sea ice forms (the exception being in the Barents Sea sector, which has seen pronounced winter ice losses); 2) The decadal evolution of AA is modulated by variable spatial expressions of atmospheric circulation; 3) the deeper vertical extent of pronounced temperature anomalies in autumn than winter is consistent with the seasonal increase in static stability from autumn to winter; and 4) reductions in static stability in autumn point toward increasingly deep penetration of surface warming into the troposphere with continued sea ice loss, and potentially greater impacts of AA on altering weather patterns in lower latitudes (Ding et al., 2024).

2 Data Sources

55 Data from the European Centre for Medium-range Weather Forecasts (ECMWF) reanalysis (ERA5; Hersbach et al., 2020) are used for analysis. Monthly temperature (2 m and the significant levels from 1000 to 500 hPa) and sea level pressure were used on the 0.25° x 0.25° horizontal grid from 1979–2024. While ERA5 data are available since 1950, fields since 1979, the advent of the modern satellite database for assimilation, are more reliable. ERA5 is chosen because, in various comparisons of (near-)surface parameters throughout the Arctic, ERA5 performs similarly to or better than other global and regional reanalysis products (Graham et al., 2019; Barrett et al., 2020; Renfrew et al., 2021; Crawford et al., 2022). Reliance is placed on trends and anomalies. Anomalies are referenced to the 30-year period 1981–2010, but comparisons are made with different averaging





periods. To assess relationships with sea ice conditions, we use the satellite passive microwave records from the National Snow and Ice Data Center. The satellite passive microwave record provides estimates of concentration and extent from October 1978 through the present at 25-km resolution on a polar stereographic grid (the EASE2 grid) by combining data from the Nimbus-7 Scanning Multichannel Microwave Radiometer (SMMR, 1979–1987), the Defense Meteorological Satellite Program (DMSP) Special Sensor Microwave/Imager (SSM/I, 1987–2007) and the Special Sensor Microwave Imager/Sounder (SSMIS, 2007-onwards) (Fetterer et al., 2002).

3 Results


3.1 Seasonality of 2-Meter Temperature Trends

A key, but in our view, under-appreciated aspect of AA is its strong seasonality - under-appreciated in the sense that processes at work during summer over the Arctic Ocean, when AA is small, set the stage for understanding the strong imprints of AA during autumn and winter. Rantanen et al. (2022) find that the AA factor as assessed for the region poleward of the Arctic circle ranges from less than two in July to over five in November. Climate models examined in that study largely capture this seasonality but with smaller amplification factors. Figure 1 shows spatial patterns of surface air temperature trends by season based on ERA5. These plots extend down to 50°N to enable comparisons between changes in the Arctic and the higher middle latitudes.





80 **Figure 1: Linear trends in 2-meter temperatures by season from 1980 to 2024, in degrees per year. Shading is used for trends significant at $p < 0.05$.**



85 **Figure 2: Linear trends in sea ice concentration %/per decade 1980 through 2024 for September (left) and December (right). Only trends significant at $p < 0.05$ are shaded.**

The sharply smaller trends in summer compared to autumn and winter across Arctic latitudes clearly stands out. Summer trends are nevertheless largely positive and statistically significant across most of the Arctic and subarctic lands. Trends are very small and not statistically significant across the central Arctic Ocean. Since the skin temperature of a melting sea ice cover is



pegged to the melting point, it follows that surface air temperature trends must be small in this area. Over land, earlier loss of the snow cover (Mudryk et al., 2023) likely contributes to the rise in surface air temperatures seen there. Trends along the Russian and Alaska coastline are also positive. Melt onset typically starts in June in the southern margins of the ice cover and progresses poleward (Markus et al., 2009). Positive trends along the coastal seas are consistent with satellite observations of both a progressively earlier onset of melt (Stroeve et al., 2014; Stroeve and Notz, 2018). They are also consistent with progressively earlier exposure of dark open water areas, their expanding coverage through time, and associated increased internal energy in the ocean mixed layer (Perovich et al., 2007; Serreze et al., 2009; Perovich and Polashenski, 2012; Stammerjohn et al., 2012; Dai et al., 2019; Li et al., 2022; Bianco et al., 2024). However, the large specific heat of water and the depth of heating (10-30 m) will limit the rise in surface air temperature. Note also the positive trends over the northern North Atlantic, which is ice-free over the entire year. Somewhat larger trends are found over part of the Kara and Barents Seas.

The largest temperature trends for autumn, locally exceeding 0.2°C per year, lie primarily on the Eurasian side of the Arctic Ocean and north of Alaska. A comparison to the spatial pattern of September (end of summer) sea ice concentration (Figure 2), provides an understanding: the trends are largest in those areas with the sharpest downward trends in ice concentration, most notably in the Chukchi and East Siberian Seas and hence where there will be strong upward surface heat fluxes as the ocean loses the internal energy it gained in summer. The interpretation, building from the above discussion and from earlier studies (e.g., Stammerjohn et al., 2012; Stroeve et al., 2016; Lebrun et al., 2019), is that through the years, ice begins to retreat earlier and earlier in spring and summer, largely from the shores of Alaska and the Russian coast, exposing areas of dark open water, which absorbs solar energy. This means more energy gain in the ocean mixed layer, and over an increasingly large area, with time. As solar radiation declines in autumn, this energy is released upwards to the atmosphere, seen as positive temperature anomalies that grow in magnitude and spatial coverage with time. Before sea ice forms, all of the internal energy gained in summer must be depleted.

The pattern of winter temperature trends is quite different. The positive trends along the Eurasian coastline and in the Chukchi and Barents Seas are greatly reduced, and largest trends, exceeding 0.2°C per year, are now located in the Barents Sea. The reason for this is clear: By December, the areas of open water along the coast have re-frozen, limiting the ocean to atmosphere heat fluxes. The Barents Sea is, in turn, one of the few areas with a substantial downward trend in winter sea ice extent (Figure 2; right panel). Still positive 2-meter temperature trends in both autumn and winter encompass much of the Arctic Ocean away from areas of ice loss. One likely driver of this is progressive thinning of the ice cover, allowing for an increase in conductive fluxes through the ice (Liu and Zhang, 2025). Another driver is likely polar temperature advection from the areas of sea ice loss (Timmermans et al., 2018), as evidenced by the tongue of fairly large positive trends extending from the Barents Sea into the Arctic Ocean. Also of interest is that trends of much of the land area are very small, even negative, especially over Eurasia.



By spring, the magnitude of temperature trends over the Barents Sea has dropped relative to winter, but is still prominent. Through spring, downward trends in sea ice concentration (not shown) persist, but, compared to winter, air-sea temperature differences are smaller, hence ocean to atmosphere surface heat fluxes are smaller. Substantial positive trends are found along the Eurasian coast, again suggestive of the role of atmospheric heat advection. Trends over much of high-latitude North America are small.

To summarize, it is apparent that an assessment of Arctic Amplification based on comparing the Arctic trend with the trend for the globe as a whole must recognize the highly pronounced seasonal and spatial heterogeneity of Arctic trends. Summer 2-m temperature trends are mostly small, but the smallness over the Arctic Ocean is due to the melting of ice. The much larger autumn trends reflect energy transfer from the ocean to the atmosphere via upward surface heat fluxes from increasing extensive areas of open water. By winter, open water areas along the Eurasian coast and the Chukchi Sea have re-frozen and the locus of maximum temperature trends is shifted to the Barents Seas, consistent with the downward trends in sea ice concentration there. Spring trends are weaker than winter trends, but are still large in the Barents Sea sector. However, for autumn, winter and spring, there are also features in the spatial patterns of trends that point to advection and other processes, and winter trends in particular are small over much of the land area.

3.2 Local Amplification Anomaly Approach

To gain further insight into trends, we now look at the evolution of AA by decade, 1980-1989, 1990-1999, 2000-2009, and 2010-2019, as well as the last five years of the record, 2020-2024, making use of what we term a Local Amplification Anomaly (LAA) approach.

For each of these periods, we calculated the average 2-meter temperature at each ERA5 grid point across the globe, then calculated the anomalies at each grid point relative to the 1981-2010 climatology. Taking the (spatially weighted) average of all grid point anomalies yields the global temperature anomaly for each period. Then, at each grid point we subtracted this global temperature anomaly from the anomaly at that point. We then compiled maps of the anomalies for the region poleward of 50°N. Examining these LAAs gives us a sense of the spatial structure of Arctic temperature anomalies in terms of how they contribute to the overall AA evolution. In Table 1 we also provide, for each decade and season, the average of the anomalies relative to the global average poleward of 60°N and the average global anomaly. Results that follow will of course reflect the chosen 1981-2010 referencing period.

Autumn	Global Anomaly	Arctic Anomaly	Difference (Arctic – Global)
1980-1989	-0.22	-0.74	-0.52
1990-1999	-0.06	-0.45	-0.39
2000-2010	0.22	0.91	0.69



2019-2019	0.45	1.68	1.23
2020-2024	0.78	2.42	1.64
Winter			
1980-1989	-0.16	-0.24	-0.08
1990-1999	-0.03	-0.53	-0.50
2000-2009	0.16	0.71	0.55
2010-2019	0.38	1.66	1.28
2020-2024	0.62	1.38	0.76
Spring			
1980-1989	-0.14	-0.68	-0.54
1990-1999	-0.04	0.13	0.17
2000-2009	0.14	0.36	0.22
2010-2019	0.40	1.37	0.97
2020-2024	0.60	1.16	0.56
Summer			
1980-1989	-0.15	-0.29	-0.14
1990-1999	-0.01	-0.09	-0.08
2000-2009	0.13	0.28	0.15
2010-2019	0.35	0.70	0.35
2020-2024	0.63	1.04	0.41

Table 1: Average temperature anomalies for the Arctic (north of 60°N), the globe, and their difference

Results for autumn are examined first. For the first two decades, 1980-1989 and 1990-1999, both the average global anomaly and the average Arctic anomaly are small and negative, with the Arctic anomalies actually more negative than the global value. As the middle of the baseline period (1981-2010), the smallest anomalies are expressed in 1990-1999. This pattern reverses starting in the 2000-2010 decade. What this is capturing is that early in the record, the poleward gradient in 2-meter temperatures was stronger than it is today; as AA evolves, the gradient obviously weakens.

For the first decade, 1980-1989, LAAs are generally small across the Arctic, with a mix of positive and negative values, but with the negative anomalies obviously dominating (not shown). The exception is in the Chukchi Sea, where strong negative LAA values of up to 3°C are found. Based on data from 1979-1996, Parkinson et al. (1999) show downward trends in ice concentration in the Chukchi Sea of around 4% per decade. However, as the area had more sea ice in the 1980-1989 decade relative to the 1981-2010 climatology, it shows up as negative LAA values.

As noted, in the 1990-1999 decade, both the Arctic average and the global average anomaly are at their minimum, since this decade is in the middle of the 1981-2010 baseline (Table 1).

The difference between the 1990-1999 and the subsequent 2000-2009 decade is striking. Both the Arctic and global average anomalies are now positive. Positive LAA values encompass most of the Arctic. The largest positive LAA values lie in the



Chukchi and East Siberian Seas, reflecting the continuing development through this decade of extensive open waters in September (Figure 3). Note that the first clear indication of the emergence of AA related to sea ice loss was based on data extending through the end of the 2000-2009 decade (Serreze et al., 2009; Screen et al., 2010a,b). Wang et al. (2017) similarly found the emergence of amplified temperature anomalies over the Arctic (60-90°N) compared to the northern mid-latitudes (30-60°N) in this decade. By the 2010-2019 decade, autumn LAA values of 3-5°C are now prominent along the entire Eurasian coast and in the Chukchi Sea; consistent with the continued increase in open water areas in September. Much smaller AA values encompass most of the rest of the Arctic.

The most recent period, 2020-2024, sees a shift. While strongly positive anomalies relative to global average anomalies - that is, positive LAA values, remain over much of the Eurasian coastal sea, LAA anomalies over the Chukchi Sea are now smaller, and larger values have appeared in the Beaufort Sea and the Canadian Arctic Archipelago. In explanation, when Arctic sea ice extent began to decline, it was initially most prominent in the Chukchi Sea region, so LAA values there are especially large, as seen in the 2009-2009 and 2010-2019 plots. With the rise in the global temperature anomalies, these LAA values become more subdued.

The winter evolution is quite different. The Arctic-averaged anomaly and the global anomaly for the 1980-1989 are small and quite alike – AA had not yet emerged (Table 1). In terms of the LAA structure (not shown), positive values of typically 1-2°C over much of Eurasia, Alaska and Canada contrast with negative values of similar size elsewhere, the exception being negative values of 2-3°C in the Barents Sea sector. The story is similar for the 1990-1999 decade - AA had yet to clearly emerge (Table 1), and, indeed, the Arctic average anomaly was about half a degree colder than the global average anomaly. The LAA structure leading to this interesting finding is characterized by partly offsetting positive and negative values (Figure 4). Of interest in this regard is that North Atlantic Oscillation (or Arctic Oscillation) shifted from a negative to a strongly positive index phase between the 1970s and late 1990s. Numerous studies examined the strong temperature trends associated with this shift, notably warming over northern Eurasia, with cooling over northeastern Canada and Greenland (e.g., Hurrell 1995; 1996; Thompson and Wallace, 1998). There was vibrant debate over whether the shift might be in part a result of greenhouse gas forcing and an emerging signal of expected Arctic Amplification (see the review in Serreze et al., 2000).




Figure 3. Autumn 2-m temperature anomalies in °C relative to 1981-2010 for 1990-1999, 2000-2009, 2010-2019 and 2020-2024
185 minus the global average temperature anomaly for each period.



While there is some indication of a structure in LAA values for the 1990-1999 decade reminiscent of the rising phase of the NAO over this time (note that the index value subsequently regressed), looking back to Table 1, the behavior of the NAO clearly did not “boost” any emerging AA signal.

Turning to the decade 2000-2009, positive LAA values have become more dominant, and fairly large positive values have appeared over the Barents Sea sector, replacing the negative values of the previous decade. While by this decade, AA had clearly emerged (Table 1), note that the positive LAA values over northern Eurasia in 1990-1999 are replaced by negative values, indicative of a circulation shift, notably, regression of the NAO from its previous high index values.

The 2010-2019 is characterized by the emergence of large positive LAA values over the Barents Sea sector which have grown since the 2000-2009 decade, pointing to the effects of growing open water areas in this sector. Positive LAA values also cover almost all Arctic latitudes. The Barents Sea feature remains prominent in the past five years of the record (2020-2024). Note, however, the negative anomalies over Alaska and eastern Eurasia. As a result, the difference between the Arctic average temperature anomaly and the global average anomaly is actually smaller than in the 2010-2019 period, that is, AA is somewhat smaller. Note also by comparison with the decade 2010-2019, LAA values along most of the Eurasia coast are less pronounced. This is understood in that, by December, all areas along the Eurasian coast and north of the Chukchi and East Siberian seas have refrozen.

Turning back to the Barents Sea sector, it is notable that this is one of the few areas of the Arctic (along with eastern Hudson Bay/Hudson Strait and Bering Strait, see Figure 2) with substantial downward trends in winter sea ice concentration. Various studies attribute the loss of winter ice in the Barents Sea and associated temperature anomalies and trends to processes involving atmospheric circulation, promoting intrusions of warm moist air into the region with wind patterns promoting stronger transport of warm Atlantic waters into the region (Woods and Caballero, 2016; Lien et al., 2017; Siew et al., 2024). Warm and moist air advection raises temperatures, inhibits autumn and winter sea ice growth (Woods and Caballero, 2016; Crawford et al., 2025) and enhances spring and summer ice melt (Kapsch et al., 2013; Park et al., 2015). Intrusions of Atlantic-derived waters, which appear to be in part wind driven, also discourage winter ice growth. Beer et al. (2020) identify an oceanic mechanism that increases the vertical heat flux in the upper Arctic Ocean under global warming that causes increased ocean heat transport into the Arctic, which appears as a substantial contributor to Arctic Amplification.




Figure 4: Winter surface temperature anomalies in °C relative to 1981-1990 for 1980-1990, 2000-2009, 2010-2019 and 2020-2024 minus the global average temperature anomaly for each period.

215 While our primary focus is on the evolution of AA and LAAs in autumn and winter, it is warranted to briefly discuss spring and summer (not shown). The spring pattern of LAAs for the 1980-1989 decade is characterized by small and mostly negative



values across the Arctic, transitioning to a mix between small positive and negative values for the 1990-1999 decade, as well as for the 2000-2010 decade. The largest difference between the Arctic average and global average anomaly was for the 2010-2019 decade. Only for the last five years of the record, 2020-2024 do prominent positive LAA values of over 3°C appear over Eurasia, but these are partly balanced by negative LAAs elsewhere. This is consistent with the much smaller AA in this season compared to autumn and winter. The key feature of summer is that while as the decades pass, modest positive values of LAA appear over land, values remain close to zero over the Arctic Ocean, reflecting the effects of the melting sea ice surface. The last five years also show positive LAA values of up to 3°C along the shores of Eurasia, likely due to the open coastal waters in these areas.

The results just discussed are with reference to 1981-2010 averages. Use of an earlier climatology (e.g., 1951-1980) naturally yields stronger positive anomalies and weaker negative LAA values in the later part of the temperature records, while a more recent climatology (e.g., 1991-2020, the current NOAA standard) has the opposite effect. The 1981-2010 reference used on this paper, used by NOAA until the end of 2020, is an appropriate middle ground, and is the reference period use for sea ice analyses by the National Snow and Ice Data Center.

3.3 Vertical Structure

An assessment of the vertical structure of warming helps to both highlight the effects of sea ice and shed light on other processes known to be involved in Arctic Amplification, notably, static stability. To this end, we look at longitudinal cross sections of temperature anomalies for the most recent 10 years of the record, averaged between the longitudes 75-80°N, which corresponds to the latitude band with pronounced anomalies in surface air temperature. We look first at October, then turn attention to December (Figure 5).

The strongly positive anomalies located from 60-120°E and between 180°E to 120°W (these being stronger) are clearly surface-based, which makes sense as they are due to strong upward surface heat fluxes. The more prominent feature between 180°E and 120°W (centered along the East Siberian and Chukchi Seas) is notable in that anomalies of 3°C extend up to 700hPa. The December cross section shows maximum surface-based temperature anomalies focused between about 20-70°E (centered near the Barents Sea), but positive anomalies do not extend as far in the vertical compared to October.




Figure 5: Vertical cross sections by longitude across latitudes 75°N to 80°N for October (left) and December (right) of temperature anomalies for 2015-2024 minus 1981-2010.


3.4 Static Stability

While the magnitude of the surface temperature anomaly will bear on how high in the vertical positive anomalies will persist, the vertical stability will play a role. The strong stability of the lower Arctic troposphere has long been recognized (Wexler, 1936; Bradley et al., 1992; Kahl et al., 1992; Serreze et al., 1992) and is central to arguments that lapse rate feedback is a contributor to AA. Based on radiosonde observations, Serreze et al. (1992) report that temperature inversions (extremely strong stability), nearly ubiquitous over the ice-covered Arctic Ocean, tend to be surface-based from October through April, increasing in strength from October through winter in both depth and in the temperature difference from inversion base to top. For example, in October the median inversion depth is about 900 m and the temperature difference is about 9K, whereas corresponding values in March are 1200 m and 12K. In summer, inversions are shallower with often a deep mixed layer below. (There are also commonly shallow melt-induced surface-based inversions.) The seasonal cycle over Arctic land areas is similar but with temperature differences across the inversion of 14-16K (Figure 6).

Figure 7 shows a vertical cross section of potential temperature from the equator to 90°N for October. The much stronger vertical stability of the Arctic troposphere compared to lower latitudes is obvious. In turn, a larger vertical extent of warming in October compared to December would be expected given that stability increases from autumn into winter. In terms of potential temperature, at 80°N (for example) the increase in potential temperature from the surface to 850 hPa in October is 10K, versus 15K in December. From the surface to 700hPa, potential temperature increases by 20K in October versus 25K in December. The atmosphere starts to cool freely to space at around 5-6 km above the surface (roughly the 500 hPa level). While pronounced autumn warming does not extend upwards that far, the results nevertheless argue that radiative cooling to space is



more efficient in autumn than in winter, and that as AA progresses, cooling to space will become more efficient as a negative feedback.



265 **Figure 6: Monthly median inversion top (top of bars), base (bottom of bars) and temperature difference (solid lines) from (a) drifting station data from the central Arctic Ocean; (b) station Zhigansk over the Siberian tundra, taken as representative of the region [from Serreze et al., 1992, by permission of AMS].**





Figure 7: Vertical cross section of zonally averaged potential temperature from the equator to the pole for October, averaged over the period 1981-2010.

Oct Average dtheta/dp 1000-850 mb 1980-2024



Oct change in dtheta/dp /year 1000-850 mb 1980-2024




Figure 8. Climatological averages and linear trends in low-level vertical stability (from surface to the 850 hPa level) expressed as change in potential temperature over 1000-850 hPa for October. Trends are in units of K/hPa. Positive numbers in for the climatological averages mean weaker stability, positive values for trends mean a decrease in stability with time. Shading is used for trends significant at $p < 0.05$.

Figure 8 shows climatological averages of surface to 850 hPa static stability for October, along with linear trends. In a stable atmosphere, $d\theta/dp$ is negative (potential temperature increases with height while pressure decreases), so more negative values mean stronger stability. Consistent with Figure 7, there is a general increase in average stability moving polewards. However, stability is strongest north of Greenland and the Canadian Arctic Archipelago. It is likely not a coincidence that these areas have the thickest sea ice in the Arctic, implying especially small heat fluxes through the ice. Not surprisingly, large trends toward weaker static stability (positive values) dominate all the areas along the Eurasian coast, corresponding to the largest downward trends in September ice concentration, as well as in the Barents Sea which has seen downward trends in winter. Smaller trends towards weaker stability dominate most of the rest of the Arctic Ocean, likely driven by a thinning ice pack. While average October conductive heat fluxes through most of the ice cover in October are on the order of $5\text{--}10 \text{ W m}^{-2}$, Liu and Zhang (2025) find that fluxes have increased since 1979 due to thinning, which outcompetes the effect of upward trends in surface skin temperatures.


Corresponding results for December follow in Figure 9. Average stability is generally stronger than for October, with the clear exception of the Norwegian and Barents Seas and the extreme northern North Atlantic, where there is near neutral stability. The Norwegian and Barents Seas, in particular, are recognized for unstable near-surface boundary layers in winter that develop




290 during cold air outbreaks as Arctic air moves over open water surfaces, promoting strong surface heat fluxes and convective-
type precipitation (Olafsson and Okland, 1994). Trends towards weaker stability are in turn prominent in the Barents Sea, the
southern Chukchi Sea and Baffin and Hudson Bays, all areas where winter ice losses have been pronounced (especially the
Barents Sea). Interesting in this regard is that weakening winter stratification may lead to intensification of near surface winds
by increasing downward momentum transfer (Zapponini and Goessling, 2024), which will then foster stronger upward
295 turbulent heat fluxes.

We stress that assessments of atmospheric stability and trends should be viewed with some caution. Based on comparisons
with radiosonde profiles at coastal sites, Serreze et al. (2012) found that all three of the most modern reanalyses available at
the time of that study (MERRA, NOAA CFSR, ERA-Interim) have positive cold-season temperature (and humidity) biases
below the 850 hPa level and consequently do not capture observed low-level temperature and humidity and temperature
300 inversions. MERRA has the smallest biases. While they offer that the reanalyses are either not assimilating the radiosonde data
at low levels or are giving these data a low weight, problems in assimilating satellite data are likely involved. Further analysis
is warranted to assess whether ERA5 suffers from the same shortcomings. Indeed, Wang and Zhao (2024) find that the
depiction of static stability over the Arctic in summer appears to be sensitive to the reanalysis product examined.

Dec Average $d\theta/dp$ 1000-850 mb 1980-2024



Dec change in $d\theta/dp$ /year 1000-850 mb 1980-2024



305 **Figure 9. Climatological averages and linear trends in low-level vertical stability (from surface to the 850 hPa level) expressed as change in potential temperature over 1000-850 hPa for December. Trends are in units of K/hPa. Positive numbers in for the climatological averages mean weaker stability, positive values for trends mean a decrease in stability with time. Shading is used for trends significant at $p < 0.05$.**



4 Discussion and Conclusions

310 The results presented here show a clear association between patterns of autumn and winter sea ice concentration trends and
both the year-to-year evolution and seasonal expression of Arctic temperature anomalies. The link with sea ice loss can be
viewed as an expression of seasonally delayed albedo feedback. We also see signals of variable atmospheric circulation in
both temperature trends and the spatial structure of LAAs by decade. As discussed, a suite of other processes can also be linked
to Arctic Amplification. Given that any process leading to warming will tend to enhance sea ice melt (spring and summer) or
315 discourage its formation (autumn and winter), they can be viewed as serving to reinforce the key role of sea ice loss on observed
AA.

Consider in this regard studies from coupled models showing that AA can arise without the albedo feedback through the lapse
rate and Plank feedbacks (e.g., Caballero and Langen, 2005; Pithan and Mauritsen, 2014; Previdi et al., 2021). Lapse rate
feedback relates to the stronger stability of the Arctic atmosphere compared to low latitudes, focusing the temperature rise
320 closer to the surface and reducing longwave radiative cooling to space. From coupled simulations, Previdi et al. (2021) find
that through positive lapse rate feedback, AA develops in only a few months following an instantaneous quadrupling of
atmospheric CO₂, well before any significant sea ice loss, although ice loss contributes significantly to warming after the first
few months. While one can question what an instantaneous quadrupling of CO₂ teaches us about the real world, a key point is
once sea ice begins to decline, the positive lapse rate feedback, keeping the heating near the surface, will contribute to spring
325 and summer ice melt and delay seasonal ice growth. That static stability becomes stronger from autumn into winter indicates
that focusing the heating near the surface will also be more effective in winter. Conversely, ice loss, and likely also heat fluxes,
are changing the larger environment towards reduced stability at low levels.

Turning to the Planck feedback, the larger increase in Arctic temperatures required to bring the system back to radiative
equilibrium in response to a forcing can also be seen as a process augmenting summer sea ice loss and delaying autumn and
330 winter ice growth. Increased autumn cloud cover as a contributor to AA is closely tied to sea ice loss through reducing stability
in the boundary layer, promoting large upward surface heat fluxes (e.g., Kay and Gettleman, 2012).

In parting, a key message stemming from the present study is that the process of AA must consider both its strong seasonality
and that AA, which is generally assessed by comparing Arctic regional temperature trends against trends for the globe as a
whole, comes about by the integration across the Arctic of large spatial heterogeneity of temperature changes, seen both in the
335 spatial pattern of Arctic trends but especially when we look at the problem through local amplification anomalies – LAAs.
While AA is small in summer, evolving summer processes, namely, leading to the reduction of sea ice concentration and
enhanced energy gain in the mixed layer, set the stage for the strong regional expressions of AA in autumn, and the change in
spatial patterns of temperature anomalies into winter as areas of open water freeze over. In all seasons, variable atmospheric
circulations appear to be important, those in summer affecting spatial patterns of September ice extent, and those in autumn



340 and winter affecting temperature through advection as well as by their influence on sea ice concentration, such as in the Barents Sea. Static stability also changes seasonally, which will influence the vertical expression of temperature anomalies.

In short, the more we look at AA, the more we discover that it is a very complex beast. These complexities bear not only on the future evolution of AA and related impacts on permafrost warming and changes in the frequency of rain on snow events (Serreze et al., 2021), but on key issues such as potential impacts of Arctic warming on middle latitude weather patterns (Ding
345 et al., 2024).

Code and data availability: The ERA5 data were obtained from the Research Data Archive at the National Center for Atmospheric Research: DOI: 10.5065/BH6N-5N20. Sea ice data was obtained from the National Snow and Ice Data Center <https://nsidc.org/data/nsidc-0051/versions/2>. For processing code contact Elizabeth Cassano (Elizabeth.Cassano@colorado.edu)

350 *Author contributions:* Mark Serreze wrote the first draft of the paper. Elizabeth Cassano performed the bulk of the data analysis and creation of figures and assisted in writing. Alex Crawford, John Cassano and Chen Zhang provided intellectual input to the paper and contributed to the writing.

Competing interests. The contact author has declared that none of the authors has any competing interests.

355 *Acknowledgement:* This study was supported by NSF Navigating the New Arctic Grant 1928230 and the Canada-150 Chair Program.

References

- 360 AMAP, Arctic Climate Change Update 2021: Key trends and impacts. Summary for policy-makers, Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, 2021.
- Arrhenius, 1896: On the influence of carbonic acid in the air upon the temperature of the ground. *Philos. Mag. J. Sci.*, 5, 237-276.
- Barrett, A.P., Stroeve, J.C., and Serreze, M.C.: Arctic Ocean Precipitation from atmospheric reanalyses and comparisons with North Pole drifting station records, *Journal of Geophysical Research: Oceans*, 125(1). <https://doi.org/10.1029/2019jc015415>,
365 2020.
- Beer, E., Eisenman, I., and Wagner, T. J.: Polar amplification due to enhanced heat flux across the halocline, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019GL086706>, 2020.
- Bianco, E., Lovino, D., Masina, S., Materia, S., and Ruggieri, P.: The role of upper-ocean heat content in the regional variability of Arctic sea ice at sub-seasonal timescales, *The Cryosphere*, 18, 2357–2379, <https://doi.org/10.5194/tc-18-2357-2024>, 2024.



- 370 Bradley, R.S., Kemig, F.T. and Iaz, H.F.: Climatology of surface-based inversions in the North American Arctic, *J. Geophys. Res. Atmospheres*, 15699–15712, <https://doi.org/10.1029/92JD01451>, 1992.
- Caballero, R., and Langen, P.: The dynamic range of poleward energy transport in an atmospheric general circulation model, *Geophys. Res. Lett.*, 32, L02705, doi:10.1029/2004GL021581, 2005.
- 375 Crawford, A. D., Lukovich, J. V., McCrystall, M. R., Stroeve, J. C., and Barber, D. G.: Reduced sea ice enhances intensification of winter storms over the Arctic Ocean, *J. Climate*, 35(11), 3353–3370. <https://doi.org/10.1175/jcli-d-21-0747.1>, 2022.
- Crawford, A., Soriot, C., and Stroeve, J.: Autumn pauses in Arctic-wide sea-ice expansion, *J. Glaciol.*, 71, e21, <https://doi.org/10.1017/jog.2024.106>, 2025.
- Dai, A., Luo, D., Song, M., and Liu, J.: Arctic amplification is caused by sea-ice loss under increasing CO₂, *Nat. Comm.*, 10, 121, <https://doi.org/10.1038/s41467-018-07954-9>, 2019.
- 380 Ding, S.S., Chen, X., Zhang, X., Zhang, X., and Xu, P.: A review on the Arctic-midlatitudes connection: Interactive impacts, physical mechanisms and nonstationary, *Atmosphere*, 15, 20.33980/atmos15091115, 2024.
- Fetterer, F., Knowles, K., Meier, W., and Savoie, M.: Sea Ice Index, digital media, National Snow and Ice Data Center, Boulder, Colorado, 2002.
- Graham, R. M., Cohen, L., Ritzhaupt, N., Segger, B., Graversen, R. G., Rinke, A., et al.: Evaluation of six atmospheric reanalyses over Arctic sea ice from winter to early summer, *J. Climate*, 32(14), 4121–4143, <https://doi.org/10.1175/jcli-d-18-0643.1>, 2019.
- 385 Graversen, R. G., and Burtu, M. Arctic amplification enhanced by latent energy transport of atmospheric planetary waves, *Quart. J. Roy. Meteor. Soc.*, 142, 2046–2054, 2016.
- Hersbach and Coauthors: The ERA5 global reanalysis, *Quart. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 390 Hurrell, J.W.: Decadal trends in the North Atlantic Oscillation: Regional temperatures and precipitation, *Science*, 269, 676–679, 1995.
- Hurrell, J.W.: Influence of variations in extratropical wintertime teleconnections on Northern Hemisphere temperature, *Geophys. Res. Lett.*, 23, 665–668, 1996.
- 395 Jansen, E., Christensen, J.H., Dokken, T., Nisancioglu, K.H., Vintgher, B.M., Capron, E., Guo, C., Jensen, M.F., Langen, P.L., Pedersen, R.A., Yang, S., Bentsen, M., Kjaer, H.A., Sadatzki, H., Sessford, E., and Stendel, M.: Past perspectives on the present era of abrupt Arctic climate change, *Nat. Clim. Change*, 10, 714–721, 2020.
- Kahl, J.D., Serreze, M.C., and Schnell, R.C.: Tropospheric low-level temperature inversions in the Canadian Arctic, *Atmosphere-Ocean*, 30, 522–529, <https://www.tandfonline.com/doi/abs/10.1080/07055900.1992.9649453>, 1992.
- 400 Kapsch, M.-L., Graversen, R.G., and Tjernström, M.: Springtime atmospheric energy transport and the control of Arctic summer sea-ice extent, *Nat. Climate Change*, 3, 744–748, doi:10.1038/nclimate1884, 2013.
- Kay, J. E., and Gettelman, A.: Cloud influence on and response to seasonal Arctic sea ice loss, *J. Geophys. Res.*, 114, D18204, doi:10.1029/2009JD011773, 2009.
- 405 Krishnan, S. et al.: The roles of the atmosphere and ocean in driving Arctic warming due to European aerosol reductions, *Geophys. Res. Lett.*, 47, e2019GL086681, 2020.
- Lebrun, M., M. Vancoppenolle, Madec, G., and Massonnet, F.: Arctic sea-ice-free season projected to extend into autumn. *The Cryosphere*, 13, 79–96, <https://doi.org/10.5194/tc-13-79-2019>, 2019.
- Li, Z., Ding, Q., Steele, M., and Schweiger, A.: Recent upper Arctic Ocean warming expedited by summertime atmospheric processes, *Nat. Commun.*, 13, 362, <https://doi.org/10.1038/s41467-022-28047-8>, 2022.



- 410 Lien, V. S., Schlichtholz, P., Skagseth, Ø., and Vikebø, F. B.: Wind-driven Atlantic water flow as a direct mode for reduced Barents Sea ice cover, *Journal of Climate*, 30, 803–812, <https://doi.org/10.1175/jcli-d-16-0025.1>, 2017.
- Liu, Y., and Zhang, J.: Conductive heat flux over Arctic sea ice from 1979 to 2022, *J. Geophys. Res. Oceans*, <https://doi.org/10.1029/2024JC022062>, 2025.
- 415 Markus, T., J. C. Stroeve, J.C., and Miller, J.: 2009: Recent changes in Arctic sea ice melt onset, freezeup, and melt season length. *Journal of Geophysical Research*, 114, C12024, <https://doi.org/10.1029/2009jc005436>, 2009.
- Mudryk, L.R., Chereque, A.E., Derksen, C., Loujous K. and Decharme, B.: Terrestrial Snow Cover, NOAA Arctic Report Card 2023, doi: 10.25923/xqwa-h543, 2023.
- 420 Navarro, J.A., Varma, V., Riipinen, I., Seland, O., Kirkevåg, A., Struthers, H., Iversen, T., Hansson, H.-C., and Ekman, A.M.L.: Amplification of Arctic warming by past air pollution reductions in Europe, *Nature Geosci.*, 9, 277–281, <https://doi.org/10.1038/ngeo2673>, 2016.
- Olafsson, H., and Okland, E.: Precipitation from convective boundary layers in Arctic air masses, *Tellus A*, 46(, 4–13, <https://doi.org/10.3402/tellusa.v46i1.15422>, 1994.
- Park, H.-S., Lee, S., Kosaka, Y., Son, S.-W., and Kim, S.-W.: The impact of Arctic winter infrared radiation on early summer sea ice, *J. Climate*, 28, 6281–6296, doi:10.1175/JCLI-D-14-00773.1, 2015.
- 425 Parkinson, C.L., Cavalieri, D.J., Gloersen, P., Zwally, H.J., and Comison, J.C.C., Arctic sea ice extent, areas and trends, 1989–1996, *J. Geophys. Res.*, 104, 20,837–20, 856, 1999.
- Perovich, D.K., Light, B., Eicken, H., Jones, K.F. Runcimen, K. and Nghiem, S.V.: Increasing solar heating of the Arctic Ocean and adjacent seas, 1979–2005: Attribution and the role of ice-albedo feedback, *Geophys. Res. Lett.*, 34, L19505, doi: 10.1029/2007GL031480, 2007.
- 430 Perovich, D. K. and Polashenski, C.: Albedo evolution of seasonal Arctic sea ice, *Geophys. Res. Lett.*, 39, 8, <https://doi.org/10.1029/2012GL051432>, 2012.
- Pithan, F., and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nature Geosci.*, 7, 181–184, <https://doi.org/10.1038/ngeo2071>, 2014.
- 435 Previdi, M., K.L. Smith, and L.M. Polvani: Arctic amplification of climate change: a review of underlying mechanisms. *Env. Res. Lett.*, 16, 093003, doi:10.1088/1748-9326/ac1c29, 2021.
- Rantanen M., Karpechko A.Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Nature Communications Earth and Environment*, 3 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- 440 Renfrew, I. A., Barrell, C., Elvidge, A. D., Brooke, J. K., Duschka, C., King, J. C., et al.: An evaluation of surface meteorology and fluxes over the Iceland and Greenland Seas in ERA5 reanalysis: The impact of sea ice distribution, *Quart. J. Roy. Meteor. Soc.*, 147(734), 691–712, <https://doi.org/10.1002/qj.3941>, 2021.
- Richter-Menge, J., and Druckenmiller, M. L. (eds.) The Arctic [In “State of the Climate in 2019“], *Bull. Am. Meteor. Soc.*, 101, S239–S285 (2020).
- 445 Screen, J., and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464, 1334–1337, <https://doi.org/10.1038/nature09051>, 2010a.
- Screen, J.A., and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464, 1334–1337, doi:10.1038/nature09051, 2010b.
- Serreze, M.C., Barrett, A.P., and Stroeve, J.: Recent changes in tropospheric water vapor over the Arctic as assessed from radiosondes and atmospheric reanalyses. *Journal of Geophysical Research*, 117, D10104, doi: 10.1029/2011JD017421, 2012.



- 450 Serreze, M.C., Gustavson, J., Barrett, A.P., Druckenmiller, M.L., Fox, S., Voveris, J., Stroeve, J., Sheffield, B., Forbes, B.C., Rasmus, S., Laptander, R., Brook, M., Brubaker, M., Temte, J., McCrystal, M., and Bartsch, A.: Arctic rain on snow events: bridging observations to understand environmental and livelihood impacts. *Environmental Research Letters*, 16, <https://doi.org/10.1088/1748-9326/ac269b>, 2021
- Serreze, M.C., Kahl, J.D. and Schnell, R.C.: Low level temperature inversions of the Eurasian Arctic and comparisons with Soviet drifting station data, *J. Climate*, 5, 615-629, https://journals.ametsoc.org/view/journals/clim/5/6/1520-0442_1992_005_0615_ltiot_2_0_co_2.pdf, 1992.
- 455 Serreze, M.C., Walsh, J.E., Chapin, F.S., III, Osterkamp, T., Dyurgerov, M., Romonovsky, V., Oechel, W.C., Morison, J., Zhang, T., and Barry, R.G.: Observational evidence of recent change in the northern high latitude environment, *Climatic Change*, 46, 159-207, 2000.
- 460 Serreze, M., Barrett, A., Stroeve, J., Kindig, D., and Holland, M.: The emergence of surface-based Arctic amplification, *Cryosphere*, 3, <https://doi.org/10.5194/tc-3-11-2009>, 2009.
- Siew, P.Y.F., Wu, Y., Ting, M., Zheng, C., Ding, O. and Seager, R.: Significant contribution of internal variability to recent Barents–Kara sea ice loss in winter, *Comm. Earth Environ.*, 5, 411, <https://doi.org/10.1038/s43247-024-01582-6>, 2024.
- 465 Stammerjohn, S., Massom, R., Rind, D., and Martinson, D.: Regions of rapid sea ice change: An inter-hemispheric seasonal comparison, *Geophys. Res. Lett.*, 39, L06501, doi: 10.1029/2012GL080874, 2012.
- Steele, M., Ermold, W., and Zhang, J.: Arctic Ocean surface warming trends over the past 100 years, *Geophys. Res. Lett.*, 35, L02614, doi:10.1029/2007GL031651, 2008.
- Stroeve, J. C., A. D. Crawford, and S. Stammerjohn, 2016: Using timing of ice retreat to predict timing of fall freeze-up in the Arctic. *Geophysical Research Letters*, 43, 6332–6340, <https://doi.org/10.1002/2016gl069314>.
- 470 Stroeve, J.C., Markus, T., Boisvert, L., Miller, J., and Barrett, A.: Changes in Arctic melt season and implications for sea ice loss, *Geophys. Res. Lett.*, doi:10.1002/2013GL058951, 2014.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Env. Res. Lett.*, <https://doi.org/10.1088/1748-9326/aade56>, 2018.
- 475 Stuecker, M. F., Bitz, C.M., Armour, K.C., Proistosescu, Kang, S.M., Shang-Ping, Xie, Doyeon, K., McGregor, S., Zhang, W., Zhao, S., Cai, W., Dong, Y. and Jin, F-F.: Polar amplification dominated by local forcing and feedbacks, *Nat. Clim. Change*, 8, 1076–1081, 2018.
- Thompson, D.W.J, and Wallace, J.M.: The Arctic Oscillation signature in the wintertime geopotential height and temperature fields, *Geophys. Res. Lett.* 25, 1297-1300, 1998.
- 480 Timmermans, M.-L., Toole, J., and Krishfield, R.: Warming of the interior Arctic Ocean linked to sea ice losses at the basin margins, *Sci. Adv.*, 4, eaat6773, <https://doi.org/10.1126/sciadv.aat6773>, 2018.
- Walsh, J. E.: Intensified warming of the Arctic: Causes and impacts on middle latitudes, *Glob. Planet. Change* 117, 52–63, 2014.
- Wang, X. and Zhao, J.: Trends and oscillations in arctic mediterranean atmospheric static stability during recent Arctic warmin. *Climate and Atmospheric Science*, <https://www.nature.com/articles/s41612-024-00576-7>, 2024.
- 485 Wang, Y., F. Huang, and T. F.: Spatio-temporal variations of Arctic amplification and their linkage with the Arctic oscillation. *Acta Oceanol. Sin.*, 36, 42–51, <https://doi.org/10.1007/s13131-017-1025-z>, 2017.
- Wexler, H.: Cooling in the lower atmosphere and the structure of polar continental air, *Mon. Wea. Rev.*, 64, 122-136, 1936.
- Woods, C. and Caballero, R.: The role of moist intrusions in winter Arctic warming and sea ice decline, *J. Clim.*, 29, 4473–4485, <https://doi.org/10.1175/JCLI-D-15-0773.1>, 2016.



- 490 Wu, D.L., and Lee, J.N.: Arctic low cloud changes as observed by MISR and CALIOP: Implication for the enhanced autumnal warming and sea ice losses, *J. Geophys. Res.*, 117, D07107, doi: 10.1029/2011JD017040, 2012.
- Yu, L., Zhong, S., Vihma, T., and Sun, B.: Attribution of late summer early autumn Arctic sea ice decline in recent decades, *Clim. Atmos. Sci.* 4, 3, 2021.
- 495 Zappone, M., and Goessling, H.F.: Atmospheric destabilization leads to Arctic Ocean winter surface wind intensification, *Nature Communications Earth and Environment*, <https://doi.org/10.1038/s43247-024-01428-1>, 2024.
- Zhou, W., Leung, L. R., and Lu, J.: Steady threefold Arctic amplification of externally forced warming masked by natural variability. *Nat. Geosci.*, 17, 508–515, <https://doi.org/10.1038/s41561-024-01441-1>, 2024.