

Evaluation of UKESM aerosol size and composition using ATom measurements indicates missing marine aerosol formation mechanisms

Xu-Cheng He^{1,2,3,4}, Nathan Luke Abraham^{1,5}, Han Ding⁶, Maria R. Russo^{1,5}, Daniel P. Grosvenor^{5,7}, Yao Ge¹, Xuemei Wang^{3,8}, Anthony C. Jones⁹, Pedro Campuzano-Jost¹⁰, Benjamin Nault^{11,12}, Agnieszka Kupc¹³, Donald Blake¹⁴, Jose L. Jimenez¹⁰, Christina J. Williamson⁴, James Weber¹⁵, Alexander T. Archibald^{1,5}, and Hamish Gordon³

¹Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge, UK

²Institute for Atmospheric and Earth System Research (INAR), University of Helsinki, Helsinki, Finland

³Department of Chemical Engineering and Center for Atmospheric Particle Studies, Carnegie Mellon University, Pittsburgh, USA

⁴Finnish Meteorological Institute, Helsinki, Finland

⁵National Centre for Atmospheric Science, UK

⁶Department of Mechanical Engineering and Center for Atmospheric Particle Studies, Carnegie Mellon University, Pittsburgh, USA

⁷Centre for Environmental Modelling And Computation, University of Leeds, Leeds, UK

⁸Koninklijk Nederlands Meteorologisch Instituut, De Bilt, The Netherlands

⁹Met Office Hadley Centre, Exeter, UK

¹⁰Cooperative Institute for Research in Environmental Sciences (CIRES) and Department of Chemistry, University of Colorado Boulder, Boulder, USA

¹¹Department of Environmental Health and Engineering, Johns Hopkins University, Baltimore, USA

¹²Center for Aerosol and Cloud Chemistry, Aerodyne Research Inc., Billerica, USA

¹³Aerosol Physics and Environmental Physics, Faculty of Physics, University of Vienna, Vienna, Austria

¹⁴Department of Chemistry, University of California Irvine, Irvine, USA

¹⁵Department of Meteorology, University of Reading, Reading, UK

Correspondence: Nathan Luke Abraham (nla27@cam.ac.uk) and Alexander T. Archibald (ata27@cam.ac.uk)

Abstract.

Atmospheric aerosols influence climate through their interactions with radiation and clouds, yet large uncertainties remain in their simulation by global models. This study evaluates the United Kingdom Earth System Model version 1.1 (UKESM1.1) using global-scale aircraft observations from the Atmospheric Tomography (ATom) mission, focusing on aerosol lifecycle processes in the remote marine atmosphere. We assess model performance in simulating aerosol precursor vapours, number size distributions, chemical composition, and environmental conditions. Several process improvements are tested, including sulfuric ~~acid~~-~~ammonia~~-~~acid~~-~~ammonia~~ nucleation, ammonium nitrate scheme, methanesulfonic acid condensation, and low-temperature isoprene-derived secondary organic aerosol formation.

Model biases differ significantly between the upper troposphere (UT) and the marine boundary layer (MBL). In the UT, UKESM1.1 overestimates nucleation and Aitken mode particles while underestimating accumulation mode, indicating insufficient growth. In the MBL, the model overestimates primary aerosols (e.g. seasalt) and precursor gases but underestimates

nucleation and Aitken mode particles, even after incorporating updated nucleation and ammonium nitrate scheme. The persistence of low aerosol number concentrations, despite overestimated precursors, suggests missing formation pathways likely involving other species such as iodine, amines, or organic vapours.

15 These limitations result in an unbalanced cloud condensation nuclei budget that over-rep~~lies~~ on primary emissions. Sensitivity tests reveal that model outputs are strongly influenced by dimethyl sulfide emissions and vapour condensation schemes. Our results highlight the need for future model development to prioritise mechanistic representation of currently missing aerosol sources, rather than relying on empirical tuning, to improve ~~aerosol-climate~~ aerosol-climate interaction estimates.

20 **1 Introduction**

Aerosols profoundly influence the Earth's climate system (Charlson et al., 1992). Depending on their chemical composition and size, they can directly affect the planetary radiation balance by scattering (e.g. sulfate aerosols) or absorbing (e.g. black carbon aerosols) solar radiation. Furthermore, aerosols act indirectly by serving as cloud condensation nuclei (CCN), thereby altering cloud microphysical properties, lifetime, and albedo, which ultimately impacts Earth's radiation balance (Twomey, 25 1977). While satellite observations help constrain the contemporary radiation balance, understanding the net effect of aerosols on Earth's radiative balance since the preindustrial era, known as aerosol forcing, presents a significant challenge. This aerosol forcing quantification requires knowledge of the aerosol state in the preindustrial era, a period without direct observations.

Consequently, climate models are almost exclusively used to infer preindustrial aerosol states. These models are typically evaluated against the current aerosol state, under the assumption that present-day processes can be extrapolated backward in time. However, uncertainties associated with this assumption, coupled with limitations in accurately simulating even the present-day aerosol lifecycle, lead to substantial uncertainties in estimates of both direct and indirect aerosol forcings (Masson-Delmotte et al., 2021).

Despite the urgent need for reducing aerosol forcing uncertainties, they have shown little improvement over recent decades. Multiple factors contribute to these persistent uncertainties. These include an incomplete understanding of aerosol emission sources and airborne aerosol formation pathways (Kirkby et al., 2023), the intricate nature of aerosol-cloud interactions (Bellouin et al., 2020), difficulties in representing complex aerosol processes and natural feedback loops within Earth system models (ESMs) (Thornhill et al., 2021), and, critically, the poorly constrained preindustrial aerosol state (Carslaw et al., 2013, 2017).

A prerequisite for reducing these uncertainties and ultimately quantifying aerosol-cloud-climate interactions is the accurate 40 quantification of aerosol sources. Aerosols originate from two main pathways: direct emission of primary aerosols (e.g. dust, seasalt, primary organic aerosols and black carbon) and airborne aerosol formation (secondary aerosols) from precursor vapours. This airborne aerosol formation pathway, involving the nucleation of low-volatility molecules followed by growth through condensation or coagulation, is estimated to contribute over 50 % of global tropospheric aerosol number concentration

and dominates in the upper troposphere (UT) (Gordon et al., 2017). Therefore, a thorough understanding of airborne aerosol
45 formation is crucial for constraining the global aerosol number budget and improving forcing estimates.

Historically, research into vapours driving aerosol nucleation has centred on sulfuric acid (H_2SO_4), formed via the oxidation
of sulfur dioxide (SO_2) by the hydroxyl radical (OH) (Kulmala et al., 1998; Kirkby et al., 2011). Sulfur sources include
natural emissions (e.g. from dimethyl sulfide, DMS) and anthropogenic activities (SO_2), making H_2SO_4 nucleation key
to understanding both preindustrial conditions and anthropogenic aerosol forcing. Aerosol nucleation involving H_2SO_4 is
50 typically assisted by water vapour and atmospheric ions, with rates highly sensitive to temperature, humidity, and ion
concentrations (Kulmala et al., 1998; Vehkamäki et al., 2002; Lee et al., 2003). However, H_2SO_4 - H_2O nucleation, even
considering ions, often fails to explain observed aerosol nucleation rates in the boundary layer (Kirkby et al., 2011; Dunne
et al., 2016). Subsequent laboratory and field studies have established that alkaline vapours, particularly ammonia (NH_3) and
amines (e.g. dimethyl amine, DMA), dramatically enhance H_2SO_4 nucleation rates (Kirkby et al., 2011; Almeida et al., 2013;
55 Jen et al., 2014). This base-enhanced nucleation is observed across diverse environments, from polluted regions (Yan et al.,
2021) to pristine polar and marine atmospheres (Jokinen et al., 2018; Beck et al., 2021; Brean et al., 2021).

Beyond H_2SO_4 , recent research has highlighted other significant contributors to regional and potentially global new particle
formation (NPF). Oxidation products of monoterpenes are major drivers in boreal forests (Yokouchi and Ambe, 1985; Ehn
et al., 2014; Tunved et al., 2006; Kirkby et al., 2016), while isoprene oxidation products are implicated in nucleation events
60 observed in the tropical UT (Andreae et al., 2018; Shen et al., 2024; Curtius et al., 2024). Furthermore, iodine oxoacids play a
dominant role in coastal (Hoffmann et al., 2001; O'Dowd et al., 2002; Sipilä et al., 2016) and polar regions (Baccarini et al.,
2020; Price et al., 2023). Their widespread presence and rapid nucleation kinetics suggest a broader contribution to marine
aerosol formation (He et al., 2021a, b, 2023). These mechanisms can dominate regionally or act synergistically with H_2SO_4 ,
collectively influencing global aerosol populations (Lehtipalo et al., 2018; He et al., 2023; Shen et al., 2024).

The formation pathways involving organic vapours and iodine species are particularly important in pristine regions. In these
65 environments, cloud formation is highly sensitive to small changes in CCN concentrations (Carslaw et al., 2013; Koren et al.,
2014). Moreover, as global anthropogenic sulfur emissions decline, the relative contribution of H_2SO_4 -driven nucleation is
expected to decrease, while the importance of organic and iodine-based mechanisms for the global aerosol budget will likely
increase (He et al., 2021b, 2023). Accurately representing these processes in climate models is therefore essential for defining
70 the preindustrial baseline, constraining anthropogenic aerosol forcing, and projecting future climate change.

Despite their climatic importance, aerosol processes remain crudely represented in many ESMs. A significant number of
models used in initiatives like the Coupled Model Intercomparison Project Phase 6 (CMIP6) employ prescribed aerosol fields,
which are not interactively coupled with model chemistry and climate dynamics (Wang et al., 2022). This limits their ability to
simulate aerosol forcings and feedbacks realistically. Even in ESMs with interactive aerosol modules (Thornhill et al., 2021),
75 the representation of airborne aerosol formation is often restricted to simplified H_2SO_4 - H_2O nucleation schemes based on
theoretical rates (Kulmala et al., 1998; Vehkamäki et al., 2002). These schemes face a fundamental paradox: while experimental
and observational evidence clearly demonstrates that H_2SO_4 - H_2O nucleation mechanisms alone are insufficient to explain
observed aerosol number concentrations (Kirkby et al., 2011; Dunne et al., 2016; Kirkby et al., 2023), the most popular

theoretical nucleation rates currently adopted in models actually overestimate aerosol nucleation by several orders of magnitude
80 (Yu et al., 2020). This overestimation may unintentionally compensate for missing nucleation mechanisms involving other
species such as ions from cosmic rays or radon, NH₃, organics, and iodine compounds (Kirkby et al., 2011, 2016; He et al.,
2021b, 2023; Shen et al., 2024), but it prevents models from capturing the underlying physical processes and their associated
climate feedbacks related to nitrogen, carbon, and iodine cycles.

Robust evaluation against observations therefore is paramount for assessing ESMs performance and improving the
85 necessary aerosol precursor chemistry and microphysics. While satellite observations provide valuable global context, they
have limitations, particularly in detecting aerosols smaller than 100 nm and those near or below clouds (Seinfeld et al.,
2016). Aircraft campaigns offer complementary, high-resolution in-situ data on precursor gases, detailed aerosol number
size distributions, and sometimes chemical composition. Among available aircraft campaigns, the Atmospheric Tomography
(ATom) mission stands out, providing an exceptionally comprehensive dataset relevant to the aerosol lifecycle (Williamson
90 et al., 2019; Brock et al., 2019, 2021; Thompson et al., 2022; Wofsy et al., 2021). For example, the ability to measure aerosol
size distributions from below 10 nm to over 1 μm , quantify SO₂ at levels below 10 parts per trillion by volume (pptv), and
detect hydroxyl radicals sets the ATom observations apart from most other aircraft campaigns. Conducted across four seasons
(2016-2018), ATom flights spanning the Pacific and Atlantic Oceans from the Arctic to the Antarctic, sampling continuously
both vertically (0.2 - 12 km) and latitudinally over vast remote marine regions. This makes ATom uniquely suited for evaluating
95 ESMs simulations of aerosol processes, especially in pristine marine environments where aerosol-cloud interactions are most
sensitive and forcing uncertainties are largest (Carslaw et al., 2013).

ATom observations have been widely used to evaluate aerosol-related processes in global models. For instance, Koenig et al.
(2020) compared CAM-Chem simulations of iodine species with measurements. Froyd et al. (2022) evaluated the performance
of CESM/CARMA and GEOS/GOCART in simulating dust aerosols and examined the influence of dust on cirrus cloud
100 formation. Williamson et al. (2019) compared aerosol number size distributions from ATom with outputs from GEOS-Chem,
CAM5, and CESM. Yu et al. (2019) assessed the effectiveness of aerosol removal by convective systems in CESM. Hodzic
et al. (2020) evaluated the performance of GEOS5, ECHAM6-HAM, CESM, and GEOS-Chem in simulating aerosol chemical
composition, with a particular focus on organic aerosols. Nault et al. (2021) analysed the simulated inorganic aerosol acidity
in CCSM4, GISS, TM4, GEOS-Chem, GEOS5, and AM4.1 using ATom data. More recently, Gao et al. (2022) provided a
105 detailed comparison of GEOS-Chem simulations with ATom measurements across multiple aerosol species, including sulfate,
organics, black carbon, nitrate, and ammonium. Bian et al. (2024) evaluated the performance of AeroCom models in simulating
the sulfur cycle over the marine atmosphere. Together, these studies demonstrated the value of ATom data in identifying model
biases and highlight the importance of accurately representing aerosol sources, transformations, and removal processes to
improve simulations of global aerosol lifecycle.

110 In this study, we leverage the ATom dataset to evaluate the performance of atmosphere component of the UK Earth System
Model version 1.1 (UKESM1.1) (Mulcahy et al., 2023) in simulating key aspects of the aerosol lifecycle. While an earlier
version (UKESM1.0) was previously evaluated against ATom (Ranjithkumar et al., 2021), that study focused on only three
variables related to airborne aerosol formation (SO₂, total aerosol number, condensation sink). Our evaluation is significantly

more comprehensive, encompassing precursor vapours (DMS, SO₂, NH₃) and oxidants (OH and ozone, O₃); aerosol number
115 concentrations across nucleation, Aitken, accumulation, and coarse modes; aerosol chemical composition (sulfate, organic,
ammonium, nitrate, seasalt); and environmental conditions (temperature [T], relative humidity [RH] and condensation sink
[CS]). Such a multi-faceted evaluation is crucial, as tuning models to match a limited subset of variables can unconsciously
introduce biases in other areas, leading to equifinality without improving overall predictive skill (Lee et al., 2016).

Furthermore, beyond evaluating the default model configuration, we implement and test several major updated or alternative
120 process representations within UKESM1.1. These include: (1) a new H₂SO₄-NH₃ nucleation scheme (Dunne et al., 2016)
with the ammonium nitrate scheme (Jones et al., 2021), reflecting its proposed importance in pristine environments (Kirkby
et al., 2011; Dunne et al., 2016); (2) a condensation scheme for methanesulfonic acid (MSA), an important DMS oxidation
product involved in aerosol growth (Beck et al., 2021) which is currently ignored in UKESM1.1; (3) the incorporation of an
isoprene secondary organic aerosol (SOA) formation scheme, for isoprene's significance for organic aerosol mass (Weber et al.,
125 2021; Tsigaridis et al., 2014); and (4) testing the impact of using a traditional DMS climatology (Lana et al., 2011) and a new
chemistry scheme, CRI-Strat2 of UKESM1.1 (Weber et al., 2021; Archer-Nicholls et al., 2021).

It is important to emphasise that while many model evaluation studies focus on sensitivity tests to improve the model
performance metrics, the primary focus of this study is improving the aerosol microphysical processes in UKESM1.1. We
prioritise implementing experimentally and observationally verified processes over optimising agreement with observations
130 through existing parameter tuning. We aim to identify key discrepancies between the model (in its default and modified
configurations) and the ATom observations, analysing these differences from a process-understanding perspective. Finally,
we will discuss the implications of these findings for future model development, integrating insights from laboratory studies,
field observations, and model simulations to advance our mechanistic understanding of the aerosol lifecycle and its climate
role - ensuring improvements are driven by enhanced physical representation.

135 Consistent with this process-oriented approach, the manuscript is structured as follows. The Methods section provides an
overview of UKESM1.1, including its aerosol and chemistry modules, as well as the model developments and implementations
carried out in this study. The Results section is organised by individual parameters, allowing readers to focus on specific
variables of interest. The Discussion section synthesises these results and highlights their implications for aerosol processes
in the marine boundary layer and the upper troposphere. Finally, we present recommendations for further development of
140 UKESM to improve the representation of aerosols and cloud condensation nuclei.

2 Methods

2.1 UKESM description

This study utilises the UKESM1.1 in its atmosphere-only configuration (Sellar et al., 2019; Mulcahy et al., 2020, 2023),
145 wherein sea surface temperatures and sea ice are prescribed from Reynolds' database (Reynolds et al., 2007). ~~The land surface~~

~~model (Joint UK Land Environment Simulator, JULES) is run simultaneously, but without dynamic vegetation. As a result, land surface temperature, soil moisture, and heat and moisture fluxes to the atmosphere are simulated rather than prescribed.~~

UKESM1.1 builds upon the HadGEM3-GC3.1 global coupled atmosphere-ocean-ice climate model (Williams et al., 2018) and incorporates additional Earth system components, including representations of carbon and nitrogen cycles, land use change, 150 ocean biogeochemistry, and a unified troposphere-stratosphere chemistry and modal aerosol scheme ~~–The simulations in this study employ a horizontal resolution of 1.875° longitude × 1.25° latitude, corresponding to approximately 135 km. Vertically, the model utilises 85 levels extending up to 85 km from the Earth’s surface, with 50 levels concentrated between 0–(Sellar et al., 2019). The land surface model (Joint UK Land Environment Simulator, JULES) is run simultaneously, but without dynamic vegetation. As a result, land surface temperature, soil moisture, and 18 km, which are the primary region of~~
155 focus for this study. heat and moisture fluxes to the atmosphere are simulated rather than prescribed.

~~The model simulations are nudged using wind and temperature fields from the ERA5 reanalysis (Telford et al., 2008; Dee et al., 2011) corresponding to the period of ATom observation. This nudging aims to reproduce the specific meteorological conditions at the time and location of the measurements, thereby reducing model biases often present in free-running model configurations (Kipling et al., 2013). Since the temporal resolution of the ERA5 reanalysis is 6 hours, the relaxation time constant for the nudged simulations is set to 6 hours. Nudging is applied vertically between model levels 12 and 80.~~

2.2 United Kingdom Chemistry and Aerosol model (UKCA)

Atmospheric chemistry and aerosols are simulated using the UKCA model (version UM13.0), which is fully coupled to UKESM1.1 for handling tracer transport (O’Connor et al., 2014; Archibald et al., 2020). UKCA utilises emission datasets consistent with the CMIP6, incorporating anthropogenic (Hoesly et al., 2018), biomass-burning (Van Marle et al., 2017), 165 and biogenic sources (Guenther et al., 2012). As the CMIP6 emission datasets utilised in this study extend only up to 2014, emissions for the years after 2015 are prescribed using the Shared Socioeconomic Pathway (SSP) SSP3-7.0 scenario. Some biogenic emissions, including isoprene, terpenes, methanol, and acetone, are simulated using interactive emission schemes. Surface concentrations of methane (CH₄), nitrous oxide (N₂O) and other well-mixed greenhouse gases were prescribed following the SSP3-7.0 projection.

170 The aerosol scheme within UKCA is largely based on the Global Model of Aerosol Processes (GLOMAP)-mode (Mann et al., 2010; Mulcahy et al., 2020). GLOMAP employs a two-moment (tracking both number and mass) pseudo-modal approach to simulate the global distribution of sulfate, black carbon, organic matter, and seasalt aerosols, whereas mineral dust is simulated using the CLASSIC sectional dust scheme (Woodward, 2001). Nitrate and ammonium aerosols are not included in the standard UKESM1.1 configuration; however, a scheme incorporating these species has recently been developed (Jones 175 et al., 2021) and is included in some of the simulations we present. GLOMAP simulates comprehensive aerosol microphysical processes, including NPF, condensation of vapours onto existing aerosols, aerosol coagulation, dry deposition, wet scavenging, and cloud processing (Mann et al., 2010). The aerosol number size distribution is represented by four modes: nucleation (geometric mean dry diameter, $\bar{d} < 10$ nm), Aitken ($10 < \bar{d} < 100$ nm), accumulation ($100 < \bar{d} < 500$ nm), and coarse ($500 < \bar{d} < 10,000$ nm). In the model version used in this study, all four modes contain soluble components. Additionally, there is an

180 insoluble Aitken mode ($100 < \bar{d} < 500$ nm) composed solely of organic matter and black carbon. At each timestep, a fraction of insoluble Aitken-mode particles is transferred to the soluble Aitken mode. This fraction is proportional to the condensation rate of soluble material (e.g. sulfate or organics) and is scaled such that the accumulation of ten monolayers of soluble material on a fraction of insoluble aerosols would result in the conversion to the soluble mode.

185 Since comprehensive descriptions of the atmospheric chemistry and aerosol formation mechanisms in UKCA have been previously published (Mann et al., 2010; O'Connor et al., 2014; Archibald et al., 2020; Mulcahy et al., 2020), we focus here only on the key processes and parameterisations directly relevant to the analysis presented in this study.

2.2.1 Primary aerosols

Primary aerosols are aerosols directly emitted from various sources. In UKESM1.1, these include mineral dust, seasalt, black 190 carbon, organic matter, and sulfate. Mineral dust is simulated using a scheme with six size bins ranging from 0.06 to 60 μm in diameter. The dust within each bin is treated independently and assigned a density of 2.65 kg m^{-3} (Woodward, 2001).

Seasalt emissions are calculated using the bin-resolved parameterisation of Gong (2003). The emitted mass and number are distributed between the soluble accumulation and coarse modes, depending on whether the centre diameter of the source bin is below or above the diameter threshold separating these modes (approximately 500 nm). A density of 2.165 kg m^{-3} is assumed 195 for seasalt in UKESM1.1.

Primary carbonaceous aerosol emissions include black carbon and organic matter originating from both anthropogenic sources (biofuel and fossil fuel combustion) and biomass burning processes. Aerosols emitted from biomass/biofuel sources are assigned a geometric mean diameter of 150 nm, while those from fossil fuel sources are assigned 60 nm; both emission types assume standard deviation of 1.59 (Stier et al., 2005; Mann et al., 2010). Notably, although the assigned diameter for 200 biomass/biofuel aerosol emissions (150 nm) exceed the typical upper size limit of the Aitken mode (100 nm), these aerosols are nevertheless emitted into the model's insoluble Aitken mode. Anthropogenic emissions are released into the lowest model layer, whereas biomass burning emissions are distributed vertically between the surface and approximately 6 km above ground level.

Primary marine organic aerosols represent another primary aerosol source which was recently implemented in UKESM 205 (Mulcahy et al., 2020). These organic aerosols are thought to be emitted as components of organic-enriched sea spray (Rinaldi et al., 2010), and their emissions show a high correlation with marine biological activity, often indicated by chlorophyll concentrations (Rinaldi et al., 2013). Consequently, UKESM1.1 adopts the parameterisation of Gantt et al. (2012), which relates emissions to both wind speed and biological activity represented by surface chlorophyll concentration. The calculated organic mass emission flux is partitioned, with 25 % attributed to the soluble Aitken mode and 75 % to the insoluble Aitken 210 mode. An emission diameter of 160 nm is assumed for both fractions, based on experimental and observational constraints (O'Dowd et al., 2004; Prather et al., 2013).

Consistent with Mann et al. (2010), the model assumes that 2.5 % of anthropogenic SO_2 emissions by mass are directly emitted as primary sulfate aerosols. This primary sulfate mass is distributed with an initial size distribution specified by Stier

et al. (2005): 50 % is allocated to the accumulation mode (assuming $\bar{d} = 150$ nm) and 50 % to the coarse mode (assuming $\bar{d} =$
215 1500 nm).

2.2.2 Sulfur sources and chemistry

Sulfate aerosols form through the oxidation of SO_2 , either via gas-phase reactions that produce H_2SO_4 and trigger NPF and growth, or through multiphase oxidation by hydrogen peroxide (H_2O_2) and O_3 dissolved in cloud liquid water, which
220 contributes to aerosol mass only. The gas-phase oxidation of SO_2 produces gaseous H_2SO_4 , which drives NPF and growth in the model. In the aqueous phase, SO_2 dissolves into cloud droplets, where it undergoes oxidation by dissolved H_2O_2 and O_3 to form sulfate. The model does not include an explicit representation of multi-phase chemical species; instead, the sulfate formed through these pathways is treated as a direct mass flux contributing to the accumulation and coarse aerosol modes. Finally, because certain removal processes (e.g. precipitation removal) associated with aqueous sulfate formation are not represented
225 in the model, a 25 % reduction factor is applied to the calculated aqueous sulfate formation mass flux (Mulcahy et al., 2020).

In UKCA, SO_2 originates from both the oxidation of DMS and direct anthropogenic emissions. Anthropogenic emissions of SO_2 are taken from the Community Emissions Data System (CEDS) inventory (Hoesly et al., 2018). SO_2 emitted from both the energy and industrial sectors is released into the model's surface layer ([up to 35 m](#)), consistent with the treatment of other trace gas emissions in UKCA. Additionally, natural SO_2 emission from continuously degassing volcanoes is prescribed using the
230 climatology developed by Dentener et al. (2006). A major revision of the SO_2 dry deposition scheme, as described by Hardacre et al. (2021), was implemented in UKESM1.1. This update, along with several additional bug fixes and model improvements detailed in Mulcahy et al. (2023), contributed to reduced surface SO_2 concentrations in UKESM1.1 compared to the earlier UKESM1 version evaluated by Ranjithkumar et al. (2021).

Marine DMS emissions in the model are calculated using an interactive scheme. The ocean biogeochemistry component,
235 incorporating the Model of Ecosystem Dynamics, nutrient Utilisation, Sequestration and Acidification (MEDUSA) module, simulates the DMS concentration in the surface ocean. This follows the formulation modified from Anderson et al. (2001) to ensure energy balance within the coupled system (Sellar et al., 2019). Additionally, terrestrial DMS emissions are also included, based on an earlier climatology (Spiro et al., 1992).

The representation of DMS chemistry within UKCA is simplified, a common necessity in large-scale global models.
240 The primary oxidation pathway for DMS represented in the model is the reaction with OH radicals, which is the dominant atmospheric sink for DMS. Additionally, DMS is also oxidised by nitrate (NO_3) radicals and atomic oxygen ($\text{O}(^3\text{P})$), although the latter pathway is generally negligible in the troposphere. The default chemistry scheme employed in UKCA is the Strat-Trop scheme (Archibald et al., 2020), but the specific reactions of DMS oxidation pathways differ between UKESM1 and UKESM1.1 (Table 1). The reaction of DMS with OH proceeds via two channels: addition and abstraction. In both UKESM
245 versions, the abstraction channel of OH oxidation, along with DMS oxidation by NO_3 , leads directly to the formation of SO_2 . While Mulcahy et al. (2023) suggested that the UKESM1.1 included the reaction $\text{DMS} + \text{O}(^3\text{P}) \rightarrow \text{SO}_2$, it is not included in the version of UKESM1.1 used in this study. The addition channel in UKESM1.1 differs from that in UKESM1; the latter

neglects the formation of dimethyl sulfoxide (DMSO). However, DMSO is recognised as an important intermediate species subject to atmospheric transport and deposition and it is a precursor of MSA. Consequently, DMS oxidation by OH directly produces MSA in UKESM1.0, partially accounting for the neglected DMSO pathway. In UKESM1.1, the addition channel was modified to produce a mixture of SO₂ and DMSO, with molar yields of 0.6 and 0.4, respectively. Subsequent DMSO oxidation by OH in UKESM1.1 further produces SO₂ and MSA, with yields of 60 % and 40 %, respectively (Table 1).

Table 1. Comparison of DMS oxidation pathways across different chemistry schemes in UKESM versions. The table shows the key DMS oxidation reactions implemented in UKESM1.0 (Strat-Trop), UKESM1.1 (Strat-Trop), and UKESM1.1 (CRI-Strat2) chemistry schemes. UKESM1.1 incorporates dimethyl sulfoxide (DMSO) as an intermediate species with simplified oxidation pathways, while CRI-Strat2 includes a more detailed representation with methylthiomethylperoxy radical (MTMP) and methylsulfinic acid (MSIA) intermediates leading to both SO₂ and MSA formation. The numerical coefficients indicate stoichiometric yields for branching reactions. We note that while Mulcahy et al. (2023) suggested that the UKESM1.1 included $\text{DMS} + \text{O}(^3\text{P}) \rightarrow \text{SO}_2$, it is not included in the version of UKESM1.1 used in this study.

UKESM1(Strat-Trop)	UKESM1.1(Strat-Trop)	UKESM1.1(CRI-Strat2)
$\text{DMS} + \text{OH} \rightarrow \text{SO}_2$	$\text{DMS} + \text{OH} \rightarrow \text{SO}_2$	$\text{DMS} + \text{OH} \rightarrow \text{MTMP} + \text{H}_2\text{O}$
$\text{DMS} + \text{OH} \rightarrow \text{SO}_2 + \text{MSA}$	$\text{DMS} + \text{OH} \rightarrow 0.6\text{SO}_2 + 0.4\text{DMSO}$	$\text{DMS} + \text{NO}_3 \rightarrow \text{MTMP} + \text{HNO}_3$
$\text{DMS} + \text{NO}_3 \rightarrow \text{SO}_2$	$\text{DMS} + \text{NO}_3 \rightarrow \text{SO}_2$	$\text{DMS} + \text{OH} \rightarrow \text{DMSO} + \text{HO}_2$
$\text{DMS} + \text{O}(^3\text{P}) \rightarrow \text{SO}_2$	$\text{DMSO} + \text{OH} \rightarrow 0.6\text{SO}_2 + 0.4\text{MSA}$	$\text{MTMP} + \text{NO} \rightarrow \text{HCHO} + \text{CH}_3\text{S} + \text{NO}_2$
		$\text{MTMP} + \text{MTMP} \rightarrow 2\text{HCHO} + 2\text{CH}_3\text{S}$
		$\text{CH}_3\text{S} + \text{O}_3 \rightarrow \text{CH}_3\text{SO}$
		$\text{CH}_3\text{S} + \text{NO}_2 \rightarrow \text{CH}_3\text{SO} + \text{NO}$
		$\text{CH}_3\text{SO} + \text{NO}_2 \rightarrow \text{CH}_3\text{SO}_2 + \text{NO}$
		$\text{CH}_3\text{SO} + \text{NO}_2 \rightarrow \text{SO}_2 + \text{CH}_3\text{O}_2 + \text{NO}$
		$\text{CH}_3\text{SO} + \text{O}_3 \rightarrow \text{CH}_3\text{SO}_2$
		$\text{DMSO} + \text{OH} \rightarrow \text{MSIA} + \text{CH}_3\text{O}_2$
		$\text{MSIA} + \text{OH} \rightarrow \text{CH}_3\text{SO}_2 + \text{H}_2\text{O}$
		$\text{MSIA} + \text{OH} \rightarrow \text{MSA} + \text{HO}_2 + \text{H}_2\text{O}$
		$\text{MSIA} + \text{NO}_3 \rightarrow \text{CH}_3\text{SO}_2 + \text{HNO}_3$
		$\text{CH}_3\text{SO}_2 \rightarrow \text{CH}_3\text{O}_2 + \text{SO}_2$
		$\text{CH}_3\text{SO}_2 + \text{O}_3 \rightarrow \text{CH}_3\text{SO}_3$
		$\text{CH}_3\text{SO}_2 + \text{NO}_2 \rightarrow \text{CH}_3\text{SO}_3 + \text{NO}$
		$\text{CH}_3\text{SO}_3 + \text{HO}_2 \rightarrow \text{MSA}$
		$\text{CH}_3\text{SO}_3 \rightarrow \text{CH}_3\text{O}_2 + \text{H}_2\text{SO}_4$

2.2.3 Airborne formation of aerosols

255 Several key airborne aerosol production processes are simulated by default in UKCA. First, new particles are formed via the
binary $\text{H}_2\text{SO}_4\text{--H}_2\text{O}$ nucleation scheme, producing aerosols in the smallest (nucleation) mode. Second, aerosols grow via the
condensation of H_2SO_4 and organic oxidation products onto existing aerosol surfaces. In the meantime, coagulation between
aerosol particles reduces the number concentration in smaller modes (e.g. the nucleation mode) while contributing to aerosol
260 growth in larger modes (e.g. the Aitken and accumulation modes). When aerosols within a specific mode grow beyond the
upper size threshold for that mode, they are transferred to the next largest mode via mode merging (Mann et al., 2010).

The default binary $\text{H}_2\text{SO}_4\text{--H}_2\text{O}$ aerosol nucleation mechanism follows the theoretical parameterisation described by
Vehkamäki et al. (2002), which is most effective at producing new particles in the free troposphere. Mechanisms for NPF
specifically within the planetary boundary layer are not explicitly included in the default model configuration. However, the
model includes an option to use a boundary layer nucleation parameterisation based on cluster activation theory, wherein the
265 nucleation rate exhibits a power-law dependence on the H_2SO_4 concentration (Kulmala et al., 2006). Additionally, a multi-
component nucleation scheme described by Metzger et al. (2010), which considers the roles of both H_2SO_4 and organic
vapours, is also available as an option within the model. It should be noted that these optional boundary layer nucleation
schemes do not quantitatively represent the state-of-the-science atmospheric nucleation mechanisms and are therefore not
utilised in this study (Kirkby et al., 2023).

270 Condensable organic oxidation products, which contribute to aerosol growth, are represented in the model as originating
from the oxidation of monoterpenes, assuming a lumped mass yield of 13 % (Mann et al., 2010). However, to account for SOA
production from other sources (such as isoprene oxidation) and to compensate for the absence of explicit anthropogenic and
marine volatile organic compound (VOC) emissions in this model configuration, this lumped monoterpene oxidation yield is
scaled by a factor of two.

275 Besides nucleation, vapour condensation, and coagulation, other aerosol microphysical processes simulated in the model
impact the aerosol lifecycle, including dry deposition, wet scavenging, and the ageing of insoluble aerosols (representing
processes that increase aerosol solubility). Readers are referred to Mann et al. (2010) for the detailed formulation of these
microphysical processes.

280 2.3 Model development

A major hurdle in climate simulations involving aerosols and their interactions with clouds and climate is the often-poor
representation of precursor chemistry (e.g. sulfur chemistry) and aerosol nucleation mechanisms. For example, the theoretical
binary $\text{H}_2\text{SO}_4\text{--H}_2\text{O}$ nucleation mechanism adopted by UKESM1.1 dates back to around the turn of the century (Kulmala
et al., 1998; Vehkamäki et al., 2002), and this mechanism is widely recognised as mechanistically insufficient to fully explain
285 atmospheric aerosol formation (Kirkby et al., 2011, 2023). Furthermore, recent advances in understanding DMS oxidation

mechanisms point to insufficient representation of these pathways in the default UKESM1.1 for accurately simulating marine sulfur chemistry and subsequent aerosol formation (Cala et al., 2023).

In addition to employing potentially outdated mechanisms, the default UKESM1.1 configuration also lacks representations of several processes known to be important for aerosol formation and growth. For instance, the model lacks representation of ion-induced nucleation processes, which are recognised as a globally significant source of aerosols (Kirkby et al., 2011; Dunne et al., 2016). Additionally, the model treats MSA as an inert tracer, meaning it is not subject to deposition, or condensation processes, despite its demonstrated importance in aerosol growth (Beck et al., 2021). The model also lacks a representation of NPF processes involving isoprene oxidation products, which has recently been identified as important in the tropical UT (Shen et al., 2024; Curtius et al., 2024). Other key omissions include, for example, the lack of representation of iodine chemistry and the associated iodine oxoacid aerosol formation mechanisms, processes shown to have a global impact on marine aerosol formation (He et al., 2021b, 2023), and alkaline molecules such as ammonia (NH_3), which are known to enhance aerosol nucleation (Kirkby et al., 2011).

To address some of these key deficiencies, this study implements several process improvements, focusing primarily on incorporating the $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation mechanism, which includes both the ion-induced and neutral nucleation channels, and a new, preliminary MSA condensation scheme based on recent observational findings. $\text{H}_2\text{SO}_4\text{-NH}_3$ aerosol nucleation has been shown to be an important global process (Kirkby et al., 2011; Dunne et al., 2016). The implemented $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation parameterisation is one of the few derived directly from experiments utilising instrumentation designed to minimise the influence of organic vapours and other strong alkaline molecules (Kirkby et al., 2011; Almeida et al., 2013). The MSA condensation scheme is based on the observational work of Beck et al. (2021), which demonstrated that MSA can condense effectively onto pre-existing aerosols, contributing significantly to their early growth. As an initial proof-of-concept implementation, this MSA condensation scheme assumes irreversible condensation onto pre-existing aerosols (analogous to the model's treatment of H_2SO_4 and oxidised organics).

In addition to these new developments, we also test the impact of using the alternative CRI-Strat2 chemistry scheme (Weber et al., 2021; Archer-Nicholls et al., 2021), incorporate the recently developed ammonium nitrate scheme (Jones et al., 2021), and evaluate the effect of replacing the interactive DMS emission scheme with a DMS climatology (Lana et al., 2011) in the model. A full list of the simulations performed is provided in Table 2, which details the key differences between the model configurations used in this study.

2.3.1 Development of $\text{H}_2\text{SO}_4\text{-NH}_3$ aerosol nucleation mechanism

The implemented $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation scheme is based on the parameterisation developed by Dunne et al. (2016), which was derived from measurements performed in the CERN CLOUD (Cosmics Leaving OUtdoor Droplets) experiments. A key feature of the CLOUD experiments, compared with previous laboratory studies, is the extremely high standard of cleanliness. This allows the nucleation capability of individual vapours, and mixtures thereof, to be studied independently with minimal contamination. Taking advantage of this capability, experiments were performed to separately evaluate the influence of H_2SO_4 ,

Table 2. Summary of model simulations and configurations used in this study. The table shows the key differences between simulations, including DMS emission schemes (interactive vs. climatology), atmospheric chemistry schemes (Strat-Trop vs. CRI-Strat2), aerosol nucleation mechanisms ($\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ vs. $\text{H}_2\text{SO}_4\text{-NH}_3$), inclusion of ammonium nitrate chemistry, methanesulfonic acid (MSA) condensation, and secondary organic aerosol (SOA) formation pathways. MTSOA refers to monoterpene-derived SOA mass yield, while IPSOA refers to isoprene-derived SOA mass yield. The SA-NH₃ (benchmark) simulation represents the primary configuration with newly implemented $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation scheme and ammonium nitrate scheme used for comparison with other model implementations. Simulation abbreviations (e.g. SA-NH₃-IPSOA) are used throughout the manuscript, where SA denotes sulfuric acid to distinguish from the chemical formula notation (e.g. $\text{H}_2\text{SO}_4\text{-NH}_3$) used for nucleation mechanisms.

Simulations	DMS	Chemistry	Nucleation	Nitrate	MSA condensation	MTSOA	IPSOA
SA-H ₂ O(default)	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-H}_2\text{O}$	N/A	N/A	26%	N/A
SA-NH ₃ -noNit	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	N/A	N/A	26%	N/A
SA-NH ₃ -slow	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes (slow)	N/A	26%	N/A
SA-NH ₃ -Lana	Climatology	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes	N/A	26%	N/A
SA-NH ₃ (benchmark)	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes	N/A	26%	N/A
SA-NH ₃ -CS2	Interactive	CRI-Strat2	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes	N/A	26%	N/A
SA-NH ₃ -MSA	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes	Yes	26%	N/A
SA-NH ₃ -IPSOA	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes	N/A	13%	3%
SA-NH ₃ -IPSOA×10	Interactive	Strat-Trop	$\text{H}_2\text{SO}_4\text{-NH}_3$	Yes	N/A	13%	30%

NH₃, H₂O, and atmospheric ions on the aerosol nucleation rate. Readers are referred to the original paper for detailed formulations and parameter values (Dunne et al., 2016).

Experiments reported in (Dunne et al., 2016) Dunne et al. (2016) identified four distinct aerosol nucleation regimes, differentiated by the concentrations of NH₃ and the presence of atmospheric ions. The first regime is termed binary neutral nucleation of $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$; this represents the experimental version of the theoretical nucleation scheme used by default in UKESM1.1 (Kulmala et al., 1998; Vehkamäki et al., 2002). This regime’s rate depends only on the concentrations of H_2SO_4 and H₂O. The second regime is termed binary ion-induced nucleation of $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$, describing nucleation in the presence of atmospheric ions. Consequently, the nucleation rate of this regime exhibits an additional dependence on the concentration of atmospheric ions. The third regime is ternary neutral nucleation of $\text{H}_2\text{SO}_4\text{-NH}_3\text{(-H}_2\text{O)}$, which depends on the concentrations of H_2SO_4 , NH₃, and H₂O. As H₂O participation is fundamental to atmospheric nucleation, it is often implicitly assumed and omitted from the nomenclature. Finally, the fourth regime is ternary ion-induced nucleation of $\text{H}_2\text{SO}_4\text{-NH}_3$, which incorporates the enhancing effect of atmospheric ions on the ternary system.

The impact of H₂O on aerosol nucleation is represented as a relative humidity dependent multiplier (K_{RH}) applied to the nucleation rate calculated based on H_2SO_4 , NH₃, and atmospheric ion concentrations. The formula reads $K_{RH} = 1 + c_1(RH - 0.38) + c_2(RH - 0.38)^3(T - 208)^2$, where $c_1 = 1.5 \pm 1.3$, $c_2 = 0.045 \pm 0.003$, RH is the relative humidity expressed as a fraction, and T is the temperature in Kelvin. This formulation uses 38 % RH (RH = 0.38)

335 as the reference condition, reflecting the humidity level at which most of the underlying experiments were performed
(Dunne et al., 2016).

As atmospheric ion pair production rates are not explicitly simulated in UKESM1.1, they are prescribed using a climatology output from the simulations described by Gordon et al. (2017). NH₃ emissions in UKESM1.1 are taken from the CEDS dataset (Hoesly et al., 2018). However, simulated atmospheric NH₃ concentrations within the model are not well constrained. For
340 example, the uptake of gaseous ammonia by acidic aerosols (leading to ammonium formation), which is widely acknowledged as an important NH₃ sink, is not represented in the default model configuration. This leads to an overestimation of gas-phase NH₃ concentrations in the model. Therefore, the recently developed ammonium nitrate scheme (Jones et al., 2021) is incorporated in most simulations in this study, aiming to improve the representation of atmospheric NH₃ budget.

The new H₂SO₄-NH₃ nucleation scheme is implemented in all simulations listed in Table 2, except for the control
345 simulation using the default binary scheme, labelled SA-H₂O(default). The SA-NH₃(benchmark) simulation represents a key experiment in this study; it incorporates the ammonium nitrate scheme using parameter settings corresponding to the fast nitrate formation scheme (specifically, a nitric acid uptake coefficient $\gamma = 0.193$) as described by Jones et al. (2021). Another sensitivity simulation, SA-NH₃-slow, explores slower nitrate formation by utilising $\gamma = 0.001$. Additionally, the SA-NH₃-noNit simulation is performed using the H₂SO₄-NH₃ nucleation scheme but without activating the ammonium nitrate
350 module, serving as a reference to evaluate the impact of the nitrate scheme itself on aerosol formation.

~~It is worth noting that the representation of the nucleation rate enhancement by H₂O differs in this study compared to the original formulation in Dunne et al. (2016). In both studies, the impact of H₂O is represented as a relative humidity dependent multiplier (K_{RH}) applied to the nucleation rate calculated based on H₂SO₄, NH₃, and atmospheric ion concentrations. The original formula reads $K_{RH} = 1 + c_1(RH - 0.38) + c_2(RH - 0.38)^3(T - 208)^2$, where $c_1 = 1.5 \pm 1.3$, $c_2 = 0.045 \pm 0.003$,
355 RH is the relative humidity expressed as a fraction, and T is the temperature in Kelvin. This formulation uses 38 % RH (RH = 0.38) as the reference condition, reflecting the humidity level at which most of the underlying experiments were performed. Unfortunately, this formulation was derived from a relatively limited number of experiments conducted at other humidity levels, and its associated uncertainty is large (Dunne et al., 2016). We find that this RH enhancement factor likely exaggerates the humidity effect at both high (> 90 %) and low (< 20 %) relative humidities, potentially distorting the calculated nucleation rate under these conditions. Therefore, a more conservative formulation for the RH dependence is adopted in this study:~~
360 ~~$K_{RH} = (RH/0.38) \times [1 + 0.02 \times (RH - 0.38) \times (T - 208)^{1.2}]$.~~

2.3.2 Development of MSA condensation scheme

In the default UKESM1.1 configuration (Mulcahy et al., 2023), MSA is treated as an inert tracer. This treatment causes MSA
365 to accumulate indefinitely in the model, leading to unrealistically high concentrations. To address this limitation, a scheme of MSA condensation onto pre-existing aerosols, as well as its wet and dry deposition, are implemented in the model in this study.

The wet deposition of MSA is treated analogously to other soluble gas-phase species in the model, following the formulation described by Giannakopoulos et al. (1999). Consistent with recommendations by Barnes et al. (2006) and Cala et al. (2023),

an effective Henry's law constant of 10^9 M atm^{-1} is adopted for MSA. The dry deposition rate of MSA is assumed to be the same as that calculated for gaseous H_2SO_4 , for the similarity in their molecular sizes and, consequently, their diffusivities in air.

Finally, the MSA condensation scheme implemented assumes irreversible condensation onto pre-existing aerosol particles, at the same rate as H_2SO_4 . This assumption is based on the comparable molecular weights (MSA = 96 g mol^{-1} ; $\text{H}_2\text{SO}_4 = 98 \text{ g mol}^{-1}$) and bulk densities (MSA = 1.5 g cm^{-3} ; $\text{H}_2\text{SO}_4 = 1.8 \text{ g cm}^{-3}$) of the two species. This treatment is supported by recent Arctic observations which suggest that MSA can condense effectively onto pre-existing aerosols, potentially at a rate comparable to that of H_2SO_4 (Beck et al., 2021). The MSA aerosol mass is currently merged with the existing sulfate aerosol mass in the model for simplicity. The MSA condensation scheme is added to the SA- NH_3 (benchmark), hereafter referred to as SA- NH_3 -MSA.

However, it must be emphasised that this MSA condensation scheme is preliminary and has not yet been fully validated against comprehensive experimental data. For instance, a potential humidity dependence of MSA partitioning to the aerosol phase has been implicated in aircraft measurements (Mauldin et al., 1999). Other modelling studies have adopted volatility-dependent parameterisations for MSA condensation, assuming it to be a temperature- and humidity-dependent process (Hodshire et al., 2019). Unfortunately, there is currently limited experimental data available to rigorously validate MSA condensation rates predicted by either type of scheme; future experimental work is crucial for developing and constraining MSA condensation parameterisations in models. Thus, the MSA condensation scheme in this study aims to provide an initial proof-of-concept implementation, allowing for the evaluation of MSA's potential impact within the current chemistry framework.

2.3.3 Coupling H_2SO_4 – NH_3 nucleation with a DMS emission climatology

Although the default UKESM1.1 configuration employs an interactive marine DMS emission scheme coupled to the MEDUSA module, this study also evaluates the widely used DMS climatology developed by Lana et al. (2011). This comparison is motivated by the significant uncertainties associated with modelled DMS emissions. For example, Bhatti et al. (2023) demonstrated that simulated DMS emissions in the Southern Ocean are highly sensitive to the choice of emission parameterisation. Specifically, the seawater DMS concentrations predicted by the interactive MEDUSA scheme generally exhibit limited spatial variability, although they show low values surrounding the Antarctic continent. In contrast, the Lana et al. (2011) climatology predicts distinct hotspots of high seawater DMS concentration near Antarctica. While rigorously verifying the true spatial pattern of global DMS emissions is beyond the scope of this study, the distinct differences between the interactive scheme and the climatology are substantial. This provides an opportunity to evaluate the sensitivity of simulated atmospheric chemistry and aerosol properties in response to different DMS source mechanisms. The DMS climatology is implemented in the simulation labelled SA- NH_3 -Lana, which also utilises the H_2SO_4 – NH_3 nucleation scheme and the ammonium nitrate scheme.

2.3.4 Coupling H₂SO₄–NH₃ nucleation with CRI-Strat2 chemistry

The coupling of the Common Representative Intermediates mechanism (CRI) with the existing UKCA stratospheric chemistry scheme (Strat) was initially carried out by Archer-Nicholls et al. (2021) and subsequently updated by Weber et al. (2021) to create the CRI-Strat2 (CS2) chemistry scheme. The development of CS2 aimed to improve the model's representation of the oxidation of non-methane volatile organic compounds and provide traceability to the CRI2.2 scheme (Jenkin et al., 2019). For instance, compared to the default Strat-Trop scheme, the CS2 scheme includes a more detailed representation of the oxidation pathways for DMS and other key biogenic VOCs, such as isoprene and monoterpenes. It is worth noting that the original CS2 scheme is not only a new chemistry mechanism but also uses different emission inventories - distinct from those used in simulations with the Strat-Trop chemistry scheme. Details of the emission inventories employed in the CS2 scheme are provided in Archer-Nicholls et al. (2021); Weber et al. (2021). To better represent NH₃ levels in the marine atmosphere, we incorporated the marine NH₃ emissions from Bouwman et al. (1997), originally used in the Strat-Trop scheme, into the Archer-Nicholls et al. (2021) and Weber et al. (2021). For example, most anthropogenic and biogenic emissions are based on datasets spanning 2000 - 2010 or on a 2010 time-slice. In this study, we updated selected anthropogenic emissions — including SO₂, nitrogen monoxide, organic carbon, black carbon, ammonia, methane, and carbon monoxide — to use CMIP6 SSP3-7.0 emissions to be consistent with the ST simulations. Emissions of volatile organic compounds (VOCs), most of which are unique in CS2 simulation, are left unchanged and therefore remain consistent with the original implementations described by Archer-Nicholls et al. (2021) and Weber et al. (2021). This modification removes the influence of inconsistent anthropogenic emissions, such that the comparison between CS2 and ST presented in this study primarily reflects differences in the underlying chemistry schemes, as well as the impact of the additional VOCs included in CS2.

The key reaction pathways of the DMS oxidation within the CS2 version used in this study are tabulated in Table 1. Similarly to other sensitivity simulations, the CS2 scheme is coupled with the H₂SO₄–NH₃ nucleation scheme and the ammonium nitrate scheme, which is labelled SA-NH₃-CS2. It is important to note, however, that despite providing an improved representation compared to Strat-Trop, the key DMS oxidation pathways included in CS2 (primarily following those described by Von Glasow and Crutzen (2004)) do not reflect the latest scientific understanding (Cala et al., 2023). Numerous recent studies, emerging from theoretical, laboratory experiments, and field observations, have provided a more comprehensive picture of DMS oxidation mechanisms (Veres et al., 2020; Shen et al., 2022; Jacob et al., 2024). However, incorporating these more up-to-date reaction pathways into the model is beyond the scope of this study. The primary purpose of this simulation is therefore to evaluate how employing a different, more complex tropospheric chemistry scheme (CS2 vs. Strat-Trop) influences the simulated atmospheric sulfur cycle, the resultant aerosol number size distributions and composition.

2.3.5 Coupling H₂SO₄–NH₃ nucleation with isoprene secondary organic aerosol formation

The default UKESM1.1 configuration incorporates a simplified representation of SOA formation, based solely on the oxidation of monoterpenes. SOA formation from isoprene oxidation is not explicitly represented; instead, its contribution is implicitly

considered by scaling the monoterpene oxidation SOA yield by a factor of two. However, Weber et al. (2022) recently implemented a new scheme in the UKCA model to explicitly represent isoprene SOA formation. This new scheme treats SOA formed from isoprene oxidation as a distinct aerosol component, produced with a fixed mass yield of 3 % from the reaction of isoprene with the major atmospheric oxidants (O_3 , NO_3 , and OH). Other aspects of the treatment of this isoprene SOA component are assumed to be identical to those of the SOA derived from monoterpene oxidation. The choice of the 3 % yield is based on the work of Scott et al. (2014), which in turn relies on the experimental findings of Kroll et al. (2005, 2006). Concurrently with the introduction of this explicit isoprene SOA source, the scaling factor previously applied to the monoterpene SOA yield is removed. Therefore, the mass yield for SOA formation from monoterpene oxidation reverts to its base value of 13 % in simulations using this scheme (Table 2). The simulation incorporating this explicit isoprene SOA scheme is referred to as SA-NH₃-IPSOA. [After formation, the condensable oxidation products of monoterpenes and isoprene are assumed to condense irreversibly onto all particle modes, including nucleation, Aitken, accumulation, and coarse.](#)

It should be acknowledged that representing SOA formation using fixed yields is a significant simplification, as actual yields are known to be strongly influenced by factors such as temperature, oxidant concentrations, aerosol acidity and nitrogen oxide (NO_x) levels. Furthermore, the formation pathways differ: monoterpene oxidation can produce extremely low-volatility compounds that contribute effectively to SOA mass through irreversible condensation (Ehn et al., 2014), whereas isoprene SOA formation is thought to be dominated by the reactive uptake of gas-phase oxidation products, such as isoprene epoxydiols (IEPOX) (Paulot et al., 2009). This reactive uptake process is significantly affected by factors like aerosol acidity; for instance, yields as high as 28.6 % have been observed under low- NO_x conditions onto acidified sulfate seed aerosol (Surratt et al., 2010). Additionally, many previous laboratory experiments on isoprene SOA yields were conducted at room temperatures. Given that lower temperatures reduce the volatility of organic vapours, the effective yield of isoprene SOA formation is expected to be higher under colder atmospheric conditions. This expectation is supported by the recent work of Shen et al. (2024), which revealed that isoprene oxidation products can initiate both aerosol nucleation and subsequent growth at low temperatures, implying a significantly higher potential SOA yield under such conditions. Therefore, to explore the sensitivity to this uncertainty, we also perform a simulation where the mass yield for the explicit isoprene SOA formation scheme is increased tenfold (to 30 %). This simulation is referred to as SA-NH₃-IPSOA \times 10.

2.4 ATom airborne campaign

The ATom airborne campaign aimed at investigating the composition of the atmosphere over the remote Pacific and Atlantic Oceans (Thompson et al., 2022). Measurements were carried out using the NASA DC-8 aircraft from 2016 to 2018 across four major deployments, roughly corresponding to the four seasons: spring (ATom-4, April-May 2018), summer (ATom-1, July-August 2016), autumn (ATom-3, September-October 2017), and winter (ATom-2, January-February 2017). The measurements covered a wide latitudinal range from the Arctic to the Antarctic (approximately 84 °N to 86 °S) and extended vertically from near the surface to the tropopause (approximately 0.2 to 12 km), primarily over remote ocean regions. The aerosol number size

distribution data used in this study are from version 2 of the dataset by Brock et al. (2022), while the trace gas data are from
470 Wofsy et al. (2021).

One of the key scientific objectives of the ATom campaign is to understand the distribution of aerosols and the precursor vapours contributing to aerosol formation and growth. Consequently, the ATom campaign boasted a comprehensive suite of instruments measuring the atmospheric constituents relevant to the full aerosol lifecycle, from precursor vapours to aerosol number size distribution and chemical composition. This rich dataset provides a unique opportunity to evaluate the performance
475 of the UKESM1.1 model simulations presented in this study.

The aerosol number size distribution of dry aerosols, spanning the diameter range from 2.7 nm to 4.8 μm , was obtained by merging measurements from several instruments. Firstly, the nucleation-mode aerosol size spectrometer (NMASS) system features a set of five condensation particle counters (CPCs) operating in parallel at different cut-off diameters (Williamson et al., 2018). Two sets of NMASS instruments were deployed on the DC-8 aircraft: the first was used across all four ATom
480 campaigns, while the second was added for ATom-2, -3, and -4. The CPCs of the first NMASS had lower cut-off diameters set at 3.2, 8.3, 14, 27, and 59 nm, while the second NMASS used cut-offs of 5.2, 6.9, 11, 20, and 38 nm, resulting in merged data across 10 size bins below approximately 60 nm. Aerosols in the diameter range 63 nm to 1000 nm were measured by an ultra-high-sensitivity aerosol size spectrometer (UHSAS) (Kupc et al., 2018). For larger aerosols from 120 nm to 10 μm , a laser aerosol spectrometer (LAS, model 3340, TSI Inc.) was employed. However, the effective upper detection size limit for
485 the LAS was restricted to 4.8 μm due to the size cut by the aircraft inlet system (Brock et al., 2019). Furthermore, due to an instrument malfunction during ATom-2, LAS data from that deployment were only used for the 0.97 μm to 4.8 μm range, after the application of correction factors (Brock et al., 2019). It is important to note that aerosol size distribution data exclude periods when the aircraft was within clouds (Brock et al., 2019).

Regarding precursor gases, the SO_2 data from ATom-1, -2, and -3 had insufficient sensitivity (detection limit > 100 pptv)
490 for typical remote marine conditions (tens of pptv). Therefore, only the SO_2 data from ATom-4 are used in this evaluation. The instrument used during the ATom-4 campaign was a laser-induced fluorescence (LIF) instrument capable of detecting SO_2 down to approximately 2 pptv, even at pressures as low as 35 hPa, making it ideal for the ATom measurement requirements (Rollins et al., 2016). OH concentrations were measured using the Penn State Airborne Tropospheric Hydrogen Oxides Sensor (ATHOS), which reported detection limits of approximately $4.5 \times 10^5 \text{ cm}^{-3}$ near the surface and $1.5 \times 10^5 \text{ cm}^{-3}$ at
495 10 km altitude. O_3 concentrations were measured by the National Oceanic and Atmospheric Administration (NOAA) NO_yO_3 instrument (Pollack et al., 2010).

Measurements of non-refractory submicron aerosol chemical composition were provided by the University of Colorado high-resolution time-of-flight aerosol mass spectrometer (AMS) (Hodzic et al., 2020; Guo et al., 2021). For consistency with the model's aerosol scheme, the evaluated AMS composition data were limited to sulfate, ammonium, nitrate, organic matter,
500 and seasalt components. The atmospheric NH_3 concentration data used in this study were not directly measured but were derived from the AMS aerosol acidity measurements provided by Nault et al. (2021). Therefore, the estimation of gas-phase NH_3 is likely subject to greater uncertainty than that of aerosol mass measurements.

Unless otherwise noted, reported observational and model data for aerosol number concentrations, vapour condensation sink, and aerosol mass concentrations used in this study have been converted to standard temperature and pressure (STP: 273.15 K, 1000 hPa). To enable direct comparisons between the model output and observations, aerosol number size distributions from ATom were recalculated into consistent size modes: nucleation mode (< 10 nm in diameter), Aitken mode (10 nm - 100 nm), accumulation mode (100 nm - $1 \mu\text{m}$), and coarse mode ($1 \mu\text{m}$ - $10 \mu\text{m}$).

2.5 Evaluation of UKESM1.1 using ATom observations

The simulations in this study employ a horizontal resolution of 1.875° longitude \times 1.25° latitude, corresponding to approximately 135 km at the equator. Vertically, the model utilises 85 levels extending up to 85 km from the Earth's surface, with 50 levels concentrated between 0 and 18 km, which are the primary region of focus for this study.

The model simulations are nudged using horizontal wind and temperature fields from the ERA5 reanalysis (Telford et al., 2008; Dee et al., 2011) corresponding to the period of ATom observation. This nudging aims to reproduce the specific meteorological conditions at the time and location of the measurements, thereby reducing model biases often present in free-running model configurations (Kipling et al., 2013). Since the temporal resolution of the ERA5 reanalysis is 6 hours, the relaxation time constant for the nudged simulations is set to 6 hours. Nudging is applied vertically between model levels 12 and 80.

To cover the full ATom observation period, the model was run from July 2016 to June 2018, with a six-month spin-up (initialisation) period from January to June 2016. For comparison with ATom observations, model outputs are retrieved as instantaneous values at a high temporal resolution of one hour. This high frequency is intended to minimise sampling bias from the model. However, due to the substantial disk space required for these high-resolution outputs (roughly 25 Gigabytes per day), model data are saved only for the specific dates corresponding to ATom flights for subsequent offline analysis.

Since this study focuses on remote marine environments, ATom observations over the continental United States and Canada are excluded from the analysis. Most of the ATom observations used in this study have 1-minute time resolution, including aerosol number size distribution (Brock et al., 2019, 2021), environmental conditions (T, RH, CS), particle composition and NH_3 (Nault et al., 2021). DMS, O_3 , OH and SO_2 used in this study have time resolution between 120 - 200 seconds (Wofsy et al., 2021). For consistency, the ATom data are sub-sampled to a fixed 5-minute interval by selecting the nearest data point within each 5-minute window. This interval is selected based on a combined assessment of the temporal resolutions of various parameters measured during the ATom campaign, the model's output resolution, and the objective of obtaining multiple data points within each model grid box (assuming a NASA DC-8 average speed of 833 km h^{-1} , a 5-minute flight segment covers roughly 70 km). Since the aerosol number size distribution is the primary dataset that underpins the entire analysis, all other parameters are included only when corresponding aerosol data are available. This ensures that the temporal and spatial coverage of the supporting data aligns with that of the aerosol measurements. Following data preparation, the model outputs are interpolated onto the four-dimensional grid (longitude, latitude, altitude, and time) of the observational data using the In-Situ Observations Simulator (Russo et al., 2025). It is important to note that this study makes frequent use of model-to-ATom ratios, as well as their logarithmic form ($\log_{10}(\text{model}/\text{ATom})$), to assess model performance. To ensure the robustness and

interpretability of these ratios, we apply a thresholding criterion whereby only ATom measurements with positive values (> 0) are included in the analysis. This step avoids artificially large ratios or undefined logarithmic values that can arise from dividing by zero or by near-zero values — particularly those resulting from background subtraction in the ATom dataset, which may yield small positive or negative values. Prior to further analysis, only the spatial and temporal points with valid ATom measurements are retained, and the corresponding model values at those locations are also preserved to ensure consistency. Data points with invalid ATom values are excluded from both datasets. While this approach may introduce a slight positive bias in the statistical summaries (e.g. mean and median) of the observational dataset, it ensures that the ratio-based comparisons remain physically meaningful and are not dominated by noise near the detection limit. The variables most affected by this thresholding are those frequently measured near instrument detection limits — especially nitrate aerosol mass throughout the vertical column, as well as ammonium and seasalt aerosol mass above the marine boundary layer (MBL).

The aerosol number size distribution data are further processed to derive the total aerosol number concentration and the size-resolved number concentrations for the four modes: nucleation, Aitken, accumulation, and coarse, to be consistent with the processed ATom data. Similarly, the modelled aerosol chemical composition is processed to facilitate comparison with the AMS measurements of submicron aerosols. The AMS size range is defined by the performance of the aerodynamic lens used as the instrument inlet, and hence, operational transmission is defined by its vacuum aerodynamic diameter, which differs from the optical/geometric diameter used by aerosol sizers and models (Guo et al., 2021; Brock et al., 2021; Kim et al., 2025). Aerodynamic diameter is typically smaller than geometric diameter, depending on particle density. Moreover, the transmission efficiency of particles larger than approximately 500 nm in aerodynamic diameter decreases log-linearly with size and is not unity. As a result, the AMS measures only a fraction of aerosol mass with a geometric diameter below 1 μm . To enable direct comparison between model outputs and AMS observations, a real-time transmission correction based on the geometric diameter is applied following Guo et al. (2021).

For sulfate, organic, nitrate, and ammonium components, their total mass is calculated by summing the mass in the model's nucleation and Aitken modes with the mass fraction of aerosols from the accumulation and coarse modes after applying transmission correction. For seasalt, which is not present in the model's nucleation and Aitken modes, only the mass fraction of aerosols smaller than 1 μm , and corrected by AMS transmission, from the accumulation and coarse modes is included in this comparison. Therefore, the total submicron aerosol mass comparison presented in this study is based on the sum of sulfate, organic matter, nitrate, ammonium, and seasalt components only. It should be noted that black carbon and dust are not included in this chemical composition comparison, as aerosol mass spectrometer measurements do not typically quantify these refractory components. However, these species are included in the model simulations and contribute to the evaluated aerosol number size distributions.

Following the recommendation of Williamson et al. (2021), we exclude potential stratospheric air from the model-ATom comparison when either the model or observation indicated $\text{O}_3 > 250$ ppbv and $\text{RH} < 10\%$. After this stratospheric filtering, the analysis is conducted across three altitude-based layers: (1) 0 - 2 km (MBL), (2) 2 - 8 km (lower to mid free troposphere), and (3) 8 - 12 km (upper troposphere, UT).

Data during aircraft takeoff and landing are not excluded in the analysis of this study, as in the original dataset provided by Brock et al. (2019). However, this inclusion is not expected to significantly affect the results. Most quantitative analyses in this study - such as tables and vertical profiles — use median values, which are robust to outliers. The only exception is the curtain plots, which display mean values due to the limited number of data points in each grid cell, making median estimates statistically unreliable. Importantly, since takeoff and landing events are geographically localised, their influence on the global-scale patterns depicted in the curtain plots is minimal. Therefore, the overall impact on our analysis is expected to be negligible.

3 Results

This study evaluates the performance of the model simulations for many aerosol relevant parameters, as listed in Table 3, against the ATom observations. The data presented in Table 3 are median ratios of modelled to observed (ATom) values. For simplicity, these model-to-ATom ratios are referred to as "ratios" throughout the manuscript, unless otherwise specified. The values presented in this summary table are calculated using all available data points below 2 km altitude, an altitude covering the marine boundary layer (MBL, for simplicity defined as < 2 km in this study) and, depending on location, parts of the lower free troposphere. In the Appendix, we further tabulate the model-to-ATom ratios for the 2 - 8 km and 8 - 12 km altitude ranges (Tables A1 and A2, respectively).

Since a single median value is insufficient to capture the full distribution of the data, our results and discussions will also feature model-ATom comparisons presented as curtain plots (latitudinal and vertical distributions) of both the original measured/modelled fields and the ratios of modelled to observed values. The analysis is further complemented by vertical profile comparisons for specific parameters, probability density functions of aerosol distribution, and horizontal spatial maps at selected altitudes.

3.1 Vapour concentrations

The observed and modelled mixing ratios of key precursor vapours and oxidants (DMS, SO₂, NH₃, O₃, and OH) are presented in Figure 1, while the corresponding ratios of modelled to observed values are shown in Figure 2. All mixing ratio data in these figures are displayed on a logarithmic scale to better visualise the magnitude of differences between the model simulations and observations.

3.1.1 Oxidants

The measured O₃ and OH mixing ratios both exhibit a distinct inter-hemispheric asymmetry, with consistently higher values observed in the Northern Hemisphere (NH) compared to the Southern Hemisphere (SH). This asymmetry is well-understood to be driven by higher anthropogenic emissions of ozone precursors, such as nitrogen oxides (NO_x) and hydrocarbons, in the

Table 3. Model-to-ATom median ratios for environmental conditions, precursor vapours, aerosol number size distributions, and chemical composition in the marine boundary layer (0 - 2 km altitude). Ratios are calculated using all available ATom data below 2 km for comparison with different UKESM1.1 configurations implementing various SA (sulfuric acid)-based nucleation schemes (see Table 2 for simulation details). The median values are calculated from the point-by-point model-to-ATom ratios (i.e. the ratio is computed at each location or time point, followed by taking the median), and therefore do not necessarily equal the ratio of the median model value to the median ATom value. Values greater than 1 indicate model overestimation, while values less than 1 indicate model underestimation relative to ATom observations. Environmental conditions include T, RH, and CS of dry aerosols. Precursor vapours include DMS, SO₂, NH₃, O₃, and OH. Aerosol number concentrations are reported for nucleation (dry diameter < 10 nm), Aitken (10 - 100 nm), accumulation (100 - 1000 nm), and coarse (1000 - 10,000 nm) modes, along with total number concentration. Chemical composition includes sulfate, organic matter, ammonium, nitrate, and seasalt mass concentrations, with total mass representing their sum. Missing values indicate that ammonium and nitrate components are not included in simulations without the ammonium nitrate scheme (SA-H₂O[default] and SA-NH₃-noNit). Similar tables for the 2 - 8 km and 8 - 12 km altitude ranges are provided in the Appendix (Tables A1 and A2, respectively).

	H₂SO₄(SA)-based schemes								
	H ₂ O(default)	NH ₃ -noNit	NH ₃ -slow	NH ₃ -Lana	NH ₃ (benchmark)	NH ₃ -CS2	NH ₃ -MSA	NH ₃ -IPSOA	NH ₃ -IPSOA×10
Temperature	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99
RH	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08	1.08
CS dry	1.12	1.24 1.27	1.26 1.30	1.24 1.28	1.23 1.27	1.23 1.34	1.28 1.31	<u>1.26</u>	<u>1.31</u>
DMS	2.35	2.35 2.25	2.48 2.52	3.45 3.28	2.59 2.58	2.61 2.58	2.59 2.57	2.52 2.48	2.62 2.58
SO₂	1.96	1.90	1.98 2.02	2.13 2.15	2.00 1.92	0.91 0.94	<u>1.91</u>	1.89	1.99 1.962.02
NH₃	371.90	378.24 329.52	11.28 11.78	6.40 7.18	12.25 12.12	16.53 23.30	7.42 10.76	11.51 13.83	11.15 12.41
O₃	1.05	1.05	0.99	0.99	0.99	<u>1.04</u>	0.98	0.98	0.99 0.98
OH	1.39	1.38 1.39	1.31 1.30	1.31 1.32	1.31	1.32 1.39	1.30	1.29 1.33	1.29 1.27
Nucleation	0.09	0.41 0.68	0.11 0.19	0.08 0.13	0.10 0.15	0.08 0.18	0.04 0.06	0.09 0.16	0.09 0.14
Aitken	0.57	0.99 1.03	0.54 0.61	0.44 0.50	0.43 0.48	0.54 0.74	0.41 0.47	0.42 0.49	0.41 0.46
Accumulation	1.10	1.25 1.27	1.28 1.31	1.26	1.20 1.25	1.23 1.34	1.24 1.29	1.21 1.25	1.21 1.24
Coarse	0.67	0.66 0.68	0.69	0.69 0.70	0.70 0.71	<u>0.69</u>	0.70	0.70	0.72 0.71
Total number	0.58	0.93 1.01	0.62 0.69	0.52 0.59	0.53 0.59	0.49 0.76	0.52 0.54	0.49 0.58	<u>0.55</u>
Sulfate	1.08	1.12	1.04 1.05	1.03 1.02	0.98 1.00	0.92 0.96	1.08	0.99 1.02	0.93 0.95
Organic	1.40	1.50 1.54	1.44	1.36 1.37	1.18 1.42	1.28 1.24	1.34 1.32	1.73 1.35	<u>1.72</u>
Ammonium	–	–	2.93 2.90	3.13 3.21	3.17 3.20	2.91 3.26	<u>3.29</u>	3.25	3.20 2.953.00
Nitrate	–	–	3.86 2.93	9.34 10.99	9.61 11.27	6.48 9.76	7.23 10.09	9.43 8.54	<u>8.43</u>
Seasalt	2.09	2.21 2.23	0.98 0.99	1.05 1.04	1.03 1.02	1.00 0.97	1.04 1.01	1.03 1.00	<u>1.00</u>
Total mass	1.24	1.29 1.31	1.23 1.28	1.25 1.24	1.24	1.14 1.18	1.22 1.25	1.21 1.25	1.27 1.28

NH (Wang and Jacob, 1998). Regarding latitudinal distribution, observed O₃ mixing ratios are generally lowest in the tropics and increase towards higher latitudes, a pattern consistent with recent compilations of oceanic and polar O₃ data (Kanaya

et al., 2025). In contrast, observed OH concentrations show approximately the opposite latitudinal trend to O₃, with the highest values found in the tropics. This inverse latitudinal distribution is primarily attributed to the stronger solar radiation in the tropics which leads to more rapid photochemical destruction of O₃, which in turn drives higher OH production rates.

~~The model-to-ATom ratios for O₃ and OH in Table 3, show relatively little variation between the different model simulations performed in this study.~~ Comparing the benchmark SA-NH₃ simulation with the ATom measurements reveals that O₃ is generally overestimated by the model in the tropics throughout the vertical profile, transitioning to a slight underestimation in the polar regions. Interestingly, despite the asymmetry in the absolute observed concentrations, the pattern of modelled O₃ discrepancies relative to observations appears broadly symmetrical between the hemispheres. Quantitatively, the median overestimation of O₃ in the tropics (25 °S to 25 °N, full altitude range unless otherwise specified) is ~~28-29~~ % (i.e. model-to-ATom ratio of ~~1.281.29~~), while the median underestimation in the polar regions (60 - 90 °N/S) is 15 % (i.e. model-to-ATom ratio of 0.85). However, the model discrepancies for OH differ significantly from those for O₃ and exhibit a strong inter-hemispheric asymmetry. North of approximately 50 °S latitude, modelled OH concentrations are generally overestimated by around ~~7-8~~ %, whereas south of 50 °S, modelled OH is substantially underestimated, by ~~53-54~~ %.

The reason for this systematic underestimation of OH concentrations south of 50 °S by the model is currently unclear and warrants further investigation in future studies. However, some potential contributing factors can be explored using the data presented. The primary photochemical production pathway for OH involves the photolysis of O₃ followed by the reaction of the resulting O(¹D) atom with water vapour (H₂O). Our evaluation of relative humidity (Figure 3) indicates a general model overestimation south of 50 °S. Given that modelled O₃ is only slightly underestimated in this region (by 19 %), inaccuracies in the modelled concentrations of the primary precursors (O₃ and H₂O) seem unlikely to be the main drivers of the substantial OH underestimation. However, uncertainties in the modelled photolysis rate of O₃ (J(O¹D))) represent one factor to be examined, as this directly impacts the OH production rate. Additionally, the primary loss processes for OH involve reactions with methane (CH₄) and carbon monoxide (CO); therefore, the accuracy of their simulated concentrations is also important. Other factors potentially contributing to the discrepancy include inaccuracies in modelled NO_x concentrations (which influence OH recycling) or other unexpected OH loss processes in the model.

3.1.2 Sulfur species

As key precursors for atmospheric H₂SO₄, DMS and SO₂ are also evaluated against ATom observations in this study (Figure 1). The measured DMS mixing ratios during ATom are generally highest in the MBL, with concentrations decreasing with increasing altitude. Within the MBL, measured DMS mixing ratios are relatively higher in the tropics compared to mid-latitude and polar regions, and the distribution appears essentially symmetrical between the hemispheres. Median DMS mixing ratios measured by ATom were around 5.6 pptv below 2 km altitude and 0.6 pptv above 2 km. In contrast to DMS, observed SO₂ mixing ratios [in ATom4](#) exhibit a more homogeneous distribution throughout the marine atmosphere, with an overall median value of 11.9 pptv. ~~However, it must be noted that the SO₂ data presented here are solely from the ATom4 campaign (Spring 2018) and thus may not fully represent the complete annual cycle of SO₂ distribution.~~

Comparing modelled DMS with observations reveals that simulations utilising the interactive DMS emission scheme consistently overestimate observed mixing ratios by roughly a factor of ~~19.80~~ 19.32 south of 40 °S across all altitudes (SA-NH₃ simulation). The SA-NH₃-Lana simulation, which employs a climatological DMS emission scheme, shows marginally better agreement in this high southern latitude region, although it still overestimates observed DMS by a factor of ~~11.15~~ 11.24. Conversely, considering the global MBL, the median modelled overestimation factor for DMS is lower in the interactive scheme simulation (SA-NH₃, factor of ~~2.59~~ 2.58) compared to the climatology simulation (SA-NH₃-Lana, factor of ~~3.45~~ 3.28) in Table 3. Therefore, judged by median performance, the interactive DMS emission scheme appears to perform better within the MBL globally but significantly worsens the model performance at high southern latitudes compared to the climatology. ~~It is worth noting, however, that the spatial patterns of the model-ATom discrepancies for DMS are remarkably similar between simulations using the interactive and climatological emission schemes (Figure 2). This similarity suggests that resolving uncertainties in DMS emissions alone is insufficient to correct the model biases in atmospheric DMS distribution. Accurate representation of atmospheric chemical transformation and loss processes is likely of equal, or even higher, importance.~~ Similarly, modelled SO₂ mixing ratios south of 40 °S (full altitude range) are also overestimated relative to ATom4 data, by ~~32~~ 33 % in the SA-NH₃ simulation and by a lesser amount, around ~~21~~ 20 %, in the SA-NH₃-Lana simulation.

While switching the DMS emission scheme primarily impacts modelled DMS and SO₂ concentrations in a latitude-dependent manner (particularly at high southern latitudes), changing the core tropospheric chemistry scheme from Strat-Trop to CS2 exerts a global influence on simulated SO₂ mixing ratios. Simulated SO₂ mixing ratios are consistently lower in the SA-NH₃-CS2 simulation compared to simulations employing the Strat-Trop scheme (Figure 2 and Tables 3, A1, A2). Consequently, the SO₂ simulation in SA-NH₃-CS2 shows reasonable agreement with observations (slight overestimation) between approximately 40 °S and 60 °N, but tends towards underestimation outside this latitude range. This suggests that the modified DMS oxidation chemistry within the CS2 scheme significantly alters the simulated SO₂ budget compared to Strat-Trop (Table 1). However, it must be pointed out that this apparent improvement in the SO₂ simulation with CS2 is not necessarily indicative of a better overall representation of sulfur chemistry, given that DMS remains significantly overestimated by the model in most regions. Therefore, the seemingly better SO₂ agreement might arise from compensating errors within the model's sulfur cycle representation, an issue warranting further investigation.

3.1.3 Ammonia

As previously reported by Nault et al. (2021), atmospheric NH₃ mixing ratios estimated during ATom are generally low, often below 1 pptv (Figure 1). The highest NH₃ mixing ratios were observed in the tropical MBL (25 °S to 25 °N; altitude < 2 km) with a median value of 6.6 pptv. Outside this tropical MBL region, observed NH₃ mixing ratios frequently dropped below 1 pptv, at which NH₃ would typically have a negligible impact on aerosol nucleation processes involving H₂SO₄.

In contrast to observations, modelled NH₃ mixing ratios in the default SA-H₂O simulation are globally overestimated by several orders of magnitude compared to observations (Figure 2). For example, the median NH₃ mixing ratio in the tropical MBL is simulated to be 215.4 pptv in this configuration, 32.64 times higher than the observed median. In the SA-NH₃-noNit

simulation (which includes $\text{H}_2\text{SO}_4\text{-NH}_3$ aerosol nucleation but not the ammonium nitrate scheme), the median tropical MBL NH_3 mixing ratio remains high at ~~219.0~~212.0 pptv. However, after implementing the ammonium nitrate scheme developed by Jones et al. (2021) (which enables the uptake of NH_3 by acidic aerosols), the modelled median NH_3 mixing ratios in the tropical
675 MBL are significantly reduced, to ~~26.3~~26.8 pptv and ~~26.8~~28.6 pptv and 25.1 pptv in the SA- NH_3 -slow and SA- NH_3 benchmark simulations, respectively. These values are considerably closer to the observed median. This result strongly suggests that the uptake of NH_3 by acidic aerosols to form ammonium is a significant sink for gaseous NH_3 in the remote marine atmosphere, and that simulated NH_3 concentrations are highly sensitive to the representation of this process.

It should be noted, however, that even in the SA- NH_3 benchmark simulation(~~incorporating the ammonium nitrate scheme~~),
680 global atmospheric NH_3 is still overestimated by a factor of ~~12.25~~12.12 when compared to ATom data in the MBL (Table 3). Using the slow nitrate uptake coefficient, the SA- NH_3 -slow scheme does not improve the ammonia simulation, with the model-to-ATom NH_3 ratio remaining at ~~11.28~~11.78. A similar trend of atmospheric NH_3 overestimation is consistently observed in other chemical transport models as well (Nault et al., 2021). This persistent, large overestimation of gaseous NH_3 inevitably drives an even larger overestimation in the formation rate of new aerosol particles via the $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation pathway,
685 as will be discussed later. Therefore, further constraining modelled NH_3 mixing ratios, likely through improved emission inventories or more detailed representation of its uptake and loss processes, remains a critical area for model development to improve aerosol simulations of NH_3 in the remote marine atmosphere (Ge et al., 2021).




Figure 1. Curtain plots of precursor vapour and oxidant concentrations along ATom flight tracks. Mean concentrations of DMS, SO₂, NH₃, O₃, and OH are shown for ATom observations (first row) and selected model simulations (subsequent rows; see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. All data are displayed on a logarithmic scale to enhance visualisation of concentration variations spanning multiple orders of magnitude. Colour scales are consistent within each column to facilitate direct comparison between observations and model simulations.




Figure 2. Curtain plots of model-to-ATom ratios for precursor vapour and oxidant concentrations along ATom flight tracks. Mean ratios of modelled to observed concentrations for DMS, SO₂, NH₃, O₃, and OH are shown for selected model simulations (see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. Data are displayed on a logarithmic scale with a diverging colour scheme where values greater than 1 indicate model overestimation and values less than 1 indicate model underestimation. Colour scales are consistent within each column to facilitate comparison between different model configurations.

3.2 Environmental conditions

690 Besides the concentrations of precursor vapours and oxidants, environmental conditions such as T and RH are also crucial for airborne aerosol formation processes. Generally, lower temperatures and higher relative humidities are more favourable for new particle formation, thus promoting aerosol nucleation processes (Kirkby et al., 2011; Dunne et al., 2016).

Since the model simulations are nudged towards reanalysis meteorological fields, the simulated temperatures are generally in good agreement with ATom observations. For example, the median model underestimation of temperature in the SA-NH₃

695 simulation is only 0.8 K globally. This relatively small temperature bias is not expected to significantly impact calculated aerosol nucleation rates, as their temperature dependence is generally modest for temperature changes of this magnitude. Both simulated temperature and RH show negligible differences between the various model configurations tested in this study, as the implemented changes in aerosol formation schemes do not significantly affect these meteorological variables on the timescales considered. However, while temperature is generally well represented, modelled RH is consistently overestimated

700 by UKESM1.1 across most of the sampled atmosphere (Figure 4). For example, the median RH overestimation is 1.08 (model-to-ATom ratio) in the MBL, increasing to $+1.13$ – 1.15 in the UT (Table A2). The overestimation of RH in the UT will inevitably lead to an overestimation of aerosol nucleation rates from H_2SO_4 - NH_3 based pathways, given the RH-dependence of these processes (Dunne et al., 2016).




Figure 3. Curtain plots of environmental conditions along ATom flight tracks. Mean T, RH, and condensation sink (CS) of dry particles are shown for ATom observations (first row) and selected model simulations (subsequent rows; see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. All data are displayed on linear scale. Colour scales are consistent within each column to facilitate direct comparison between observations and model simulations.




Figure 4. Curtain plots of model-to-ATom differences (model - ATom) for environmental conditions along ATom flight tracks. Mean differences of modelled to observed values for T, RH, and CS are shown for selected model simulations (see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. Data are displayed on linear scale with a diverging colour scheme. Colour scales are consistent within each column to facilitate comparison between different model configurations.

705 3.3 Aerosol number size distribution

The observed aerosol number concentrations within each aerosol mode from the ATom measurements, alongside those from selected model simulations, are presented in Figure 5. Ratios of modelled to observed values are similarly shown in Figure 6. While Ranjithkumar et al. (2021) reported a comparison of modelled total aerosol number concentration vertical profiles with ATom observations, this study focuses on a mode-resolved comparison across all four modes (nucleation, Aitken, 710 accumulation, and coarse), as this modal separation is necessary to understand their distinct sources, transformation pathways, and climate impacts.

3.3.1 Nucleation and Aitken mode aerosols

The ATom measurements reveal a ubiquitous presence of nucleation mode aerosols throughout the marine atmosphere (Brock et al., 2019; Williamson et al., 2019). The concentrations of nucleation mode aerosols are largely symmetrical between the hemispheres. Another notable feature of the observed nucleation mode aerosol distribution is that its vertical variation is considerably larger in the tropics than in mid-latitudes and polar regions (Figure 7). For example, nucleation mode aerosol concentrations measured at approximately 12 km altitude in the tropics are around two orders of magnitude higher than those at the marine surface level, whereas this vertical gradient in mid-latitudes and polar regions is typically less than one order 720 of magnitude. This phenomenon has previously been associated with strong NPF events occurring in the tropical UT, as

proposed by Clarke et al. (2013); Williamson et al. (2019). Similar to the nucleation mode, Aitken mode aerosols also exhibit a comparable vertical trend in the tropics, with enhanced concentrations observed in the UT relative to the lower troposphere. ~~One notable difference between the nucleation and Aitken mode distributions is that Aitken mode aerosols show consistently lower concentrations within the MBL across all latitudes. This reduction in boundary layer Aitken mode aerosols is likely explained~~
725 ~~by enhanced coagulation of nucleation mode aerosols with larger accumulation and coarse mode aerosols. This interpretation is supported by the enhanced accumulation and coarse mode aerosol concentrations observed in the MBL (Figure 5), which are significantly higher than those in the UT, providing a substantial coagulation sink.~~

Comparing the default simulation (SA-H₂O) with ATom observations (Figures 6 and 7), the bias in nucleation mode aerosol concentrations shows a clear dipole pattern: a significant overestimation globally above approximately 4 km, and a strong
730 underestimation below this altitude. For example, the default simulation overestimates nucleation mode aerosols by roughly one order of magnitude at 12 km altitude (Figure 7), while it underestimates them with a ratio of 0.09 in the MBL (Table 3). The overestimation of the nucleation mode in the UT by the default simulation is attributed to several factors. The primary factor is that the theoretical binary H₂SO₄–H₂O nucleation mechanism from Vehkamäki et al. (2002) itself overestimates the neutral H₂SO₄–H₂O nucleation rate by up to three orders of magnitude in the UT, as highlighted by Yu et al. (2020). Other secondary
735 factors include the model's underestimation of temperature, overestimation of SO₂ and RH (discussed in previous sections), all of which contribute to an increased nucleation rate. Taking the tropical UT (8 - 12 km, 25 °S to 25 °N) as an example, the model-to-ATom ratios of SO₂ and RH are 1.46 and 1.12, respectively, in the default simulation. The only evaluated factor that might partly counter this tendency to overestimation is the condensation sink, which the model overestimates by 11 % in the tropical UT; however, this effect is outweighed by the other contributing factors.

740 Implementing the H₂SO₄–NH₃ nucleation scheme without the ammonium nitrate scheme (SA-NH₃-noNit simulation) does not improve model skill in simulating aerosol number size distribution in the tropical UT; the overestimation of nucleation mode aerosols remains high, similar to that in the default SA-H₂O simulation (Figure 6). On the other hand, the SA-NH₃-noNit simulation shows nucleation mode aerosol concentrations more than ~~4.7~~ times higher in the MBL compared to the default simulation, due to the aerosol nucleation enhancement from H₂SO₄–NH₃ nucleation mechanism (Table 3). However, despite
745 this enhancement, the simulated nucleation mode aerosol concentrations are still ~~2.41 times~~ lower than the ATom observations in the MBL. Furthermore, it must be noted that this apparent improvement in predicting MBL nucleation mode aerosols in the SA-NH₃-noNit simulation is concurrent with a enormous overprediction of NH₃ by the model, which is higher than ATom observations by a factor of ~~378.24~~ 329.52 (Table 3). After implementing the ammonium nitrate scheme, with either a slow (SA-NH₃-slow) or fast (SA-NH₃ benchmark) nitric acid uptake coefficient, the overestimation of NH₃ is reduced to factors of
750 ~~11.28 and 12.25~~ 11.78 and 12.12, respectively. However, the underestimation of nucleation mode aerosols worsens, with ratios of ~~0.11 and 0.10~~ 0.19 and 0.15 for these simulations, respectively. This represents a marginal enhancement compared with the default H₂SO₄–H₂O nucleation mechanism over pristine marine environments.

~~As expected, the introduction of additional condensable material for growth, such as MSA (SA-NH₃-MSA) and isoprene-derived SOA (SA-NH₃-IPSOA and SA-NH₃-IPSOA×10), does not significantly improve the nucleation mode aerosol number bias—these primarily contribute to increasing aerosol mass, rather than number concentrations (Table 3).~~
755

In general, the biases in modelled Aitken mode aerosol concentrations show similar patterns to those for the nucleation mode. In the MBL, Aitken mode aerosols are underestimated by a factor of approximately two for ~~all~~ most model simulations except the SA-NH₃-noNit ~~simulation~~ and SA-NH₃-CS₂ simulations (Table 3). The median Aitken mode aerosol concentration is well simulated in the SA-NH₃-noNit simulation. However, as discussed previously, this improved agreement is associated with the significant overestimation of NH₃ in this particular model configuration. Despite this major caveat, the SA-NH₃-noNit simulation is the only simulation that substantially improves the model skill in predicting both nucleation mode and Aitken mode aerosol concentrations in the MBL, with model-to-ATom ratios of ~~0.41 and 0.99~~ 0.68 and 1.03, respectively. The aerosol probability density function in the MBL for this simulation also shows much closer agreement with the ATom observations than other simulations (Figure 8). This result — where better aerosol predictions require unrealistically high precursor concentrations — indicates that the H₂SO₄-NH₃ mechanism cannot adequately explain observed MBL airborne aerosol formation. Additional or alternative aerosol formation processes are necessary to explain the high nucleation mode aerosol concentrations observed in the MBL.




Figure 5. Curtain plots of aerosol number size distribution along ATom flight tracks. Mean nucleation mode (< 10 nm), Aitken mode (10 - 100 nm), accumulation mode (100 - 1000 nm), and coarse mode (1000 - 10,000 nm) aerosol number concentrations are shown for ATom observations (first row) and selected model simulations (subsequent rows; see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. All data are displayed on a logarithmic axis to enhance visualisation of the variations in parameters. Colour scales are consistent within each column to facilitate direct comparison between observations and model simulations.




Figure 6. Curtain plots of model-to-ATom ratios for aerosol number size distribution along ATom flight tracks. Mean ratios of modelled to observed values for nucleation mode (< 10 nm), Aitken mode (10 - 100 nm), accumulation mode (100 - 1000 nm), and coarse mode (1000 - 10,000 nm) aerosol number concentrations are shown for selected model simulations (see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. Data are displayed on a logarithmic scale with a diverging colour scheme where values greater than 1 indicate model overestimation and values less than 1 indicate model underestimation. Colour scales are consistent within each column to facilitate comparison between different model configurations.




Figure 7. Vertical profiles of nucleation mode (< 10 nm) and Aitken mode (10 - 100 nm) aerosol number concentrations (cm^{-3}). ATom observations (black lines) and selected model simulations (coloured lines; see Table 2 for simulation details) are presented. Median values are shown for nucleation mode aerosols in the tropics (A, 25 °S to 25 °N), mid-latitudes (B, 25 - 60 °N/S), and high-latitudes (C, 60 - 90 °N/S), and for Aitken mode aerosols in the tropics (D), mid-latitudes (E), and high-latitudes (F). Shaded areas for solid lines represent the interquartile range (25th to 75th percentiles) for both ATom observations and model simulations, indicating the variability in aerosol concentrations. Numbers on the right edge of each panel show the sample size used to calculate median values for each altitude bin at 5-minute temporal resolution.




Figure 8. Aerosol number size distribution probability density functions in the marine boundary layer (0 - 2 km altitude) showing median values for ATom observations and selected model simulations. Probability density functions of aerosol number concentration ($dN/d\log D_p$) are shown for ATom observations (gray dots) and model simulations (coloured lines; see Table 2 for simulation details). Gray shade represents the interquartile range of the ATom observations. The x-axis represents aerosol diameter in micrometers, while the y-axis shows the probability density function of aerosol number concentration.

3.3.2 Accumulation and coarse mode aerosols

The observed global distributions of accumulation mode and coarse mode aerosols differ significantly from those of the nucleation and Aitken modes. Generally, regions with high Aitken mode aerosol concentrations are typically associated with low concentrations of accumulation mode aerosols, suggesting a somewhat inverse relationship in their spatial distributions (Figures 5 and 6). This relationship is partly regulated by the amount of condensable vapours available for growth, through which Aitken mode aerosols grow into the accumulation mode. Coarse mode aerosols exhibit significantly higher concentrations in the lower troposphere (below approximately 4 km; Figure A1). This is attributed to their larger size, which limits their atmospheric mobility and makes them more susceptible to deposition processes; thus they are concentrated nearer to their emission sources. Overall, both accumulation and coarse mode aerosol concentrations are significantly higher in the mid- to high-latitudes of the NH compared to the SH, a difference primarily attributed due to greater anthropogenic activities in the NH. Emissions of primary aerosols, as well as anthropogenic precursor vapours (such as SO₂) and oxidants (such as O₃ and OH), are all substantially higher in the NH than in the SH.

Accumulation mode aerosols are notably the only mode consistently overestimated by the presented model simulations in the MBL, with a model-to-ATom ratio of 1.20-1.25 in the SA-NH₃ benchmark simulation (Table 3). There are several potential reasons for this overestimation: 1) excessive aerosol nucleation and subsequent growth processes in the MBL, 2) excessive primary emissions of accumulation mode aerosols, or 3) inefficient aerosol removal processes. The first reason is unlikely, as both nucleation mode and Aitken mode aerosols are significantly underestimated in the MBL, as discussed previously. The second reason is likely, as the model appears to overestimate primary emissions contributing to accumulation mode aerosols, such as seasalt, in the MBL (supported by aerosol composition data discussed in the next section; see Table 3). The third reason cannot be ruled out, but it cannot be assessed with the data available in this study.

While accumulation mode aerosols are overestimated in the MBL, they are underestimated in the UT, where the availability of condensable vapours strongly limits particle growth (Figure A1). Since H₂SO₄ is likely already overestimated in the tropical UT by the model (due to overestimation of SO₂ and RH_{OH}), the condensable vapours driving this additional growth are likely to be oxidised organic compounds, such as isoprene-derived SOA, transported from rainforests to the marine UT, as suggested by Shen et al. (2024) and Curtius et al. (2024). The contribution of organics is further supported by aerosol composition analysis in the next section. The addition of isoprene SOA does reduce the underestimation of accumulation mode aerosols in the tropical UT. For example, the SA-NH₃-IPSOA×10 simulation which leads to a significant increase in modelled accumulation mode aerosol concentrations in this region, resulting in a model-to-ATom ratio of 0.65. This represents an increase of 0.16-0.19 in the ratio compared to the benchmark simulation (which has a ratio of 0.49-0.46). Simultaneously, the overestimation of Aitken mode aerosols in the tropical UT is also reduced by these additional condensable vapours, with the SA-NH₃-IPSOA×10 simulation yielding a model-to-ATom ratio of 1.33-1.35. This is a decrease of 0.24-0.19 in the ratio from the benchmark simulation (which has a ratio of 1.57-1.54).

Coarse mode aerosols are generally underestimated by the model in the MBL, with a median model-to-ATom ratio of 0.70 0.71 in the SA-NH₃ benchmark simulation (Table 3). In regions other than the MBL, coarse mode aerosols are generally overestimated by the model (Figure 6). The most severe overestimation occurs south of 40 °S above the MBL, where the model overestimation can exceed a factor of 10, although the absolute number concentrations in this region are very low (less than 0.1 cm⁻³). ~~Since aerosol mass spectrometer measurements of seasalt used in this study do not include aerosols larger than 1 μm and dust aerosols are not measured by the AMS we use for model evaluation in this study. It is, however, intriguing to note that this specific region of coarse mode overestimation coincides with the region where OH is significantly underestimated, while DMS and SO₂ are overestimated by the model (Figure 2). Finding the physical and chemical reasons for this connection is therefore an interesting avenue for future research.~~

The combined surface area of accumulation and coarse mode aerosols, and to a lesser extent Aitken mode aerosols, determines the available surface for vapour condensation, and thus defines the condensation sink (CS). The CS is calculated here following the methodology of Pirjola and Kulmala (1998), based on the dry aerosol size distribution. ~~This choice is made because calculating CS based on wet size distributions introduces additional uncertainties related to aerosol composition and hygroscopicity, and also because aerosol composition is evaluated independently in the subsequent section.~~ The CS typically exhibits a negative correlation with the aerosol nucleation rate, as a higher CS indicates a larger aerosol surface area for vapour condensation, thereby reducing the vapour concentration available for aerosol nucleation. In a large fraction of atmosphere sampled by ATom, the CS is overestimated by the model (Figure 4). In the MBL, the benchmark simulation overestimates CS by 24 27 % (Table 3). Considering that the primary production pathway for H₂SO₄ in the MBL is the reaction of SO₂ with OH, while its major loss process is condensation onto existing aerosols (governed by CS), a simplified steady-state approximation suggests $[H_2SO_4] \propto ([SO_2][OH])/CS$. Given that the model overestimation of CS in the MBL is roughly counterbalanced by the overestimation of OH (by 31 %) in the benchmark simulation (Table 3), the model-to-ATom ratio for the term [OH]/CS is close to unity. Consequently, the bias in simulated H₂SO₄ in the MBL is likely dominated by the bias in SO₂, suggesting that H₂SO₄ is probably overestimated by a factor of 2 (Table 3).

~~Vertical profiles of accumulation mode (100–1,000 nm) and coarse mode (1,000–10,000) aerosol number concentrations. ATom observations (black lines) and selected model simulations (coloured lines; see Table 2 for simulation details) are presented. Median values are shown for accumulation mode aerosols in the tropics (A, 25 °S to 25 °N), mid-latitudes (B, 25–60 °N/S), and high-latitudes (C, 60–90 °N/S), and for coarse mode aerosols in the tropics (D), mid-latitudes (E), and high-latitudes (F). Shaded areas for solid lines represent the interquartile range (25th to 75th percentiles) for both ATom observations and model simulations, indicating the variability in aerosol concentrations. Numbers on the right edge of each panel show the sample size used to calculate median values for each altitude bin at 5-minute temporal resolution.~~

3.4 Aerosol composition

The observed aerosol mass concentrations ATom measurements, alongside those from selected model simulations, are presented in Figure 9. Ratios of modelled to observed values are similarly shown in Figure 10. In general, model simulations

835 tend to overestimate the MBL submicron aerosol mass (Table 3). This overestimation of aerosol mass occurs despite the general
underestimation of total aerosol number concentrations by the model in the MBL. ~~In the benchmark SA-NH₃ simulation,
for example, the modelled median MBL mass concentrations for sulfate, organic matter, ammonium, and seasalt are 0.198,
0.118, 0.067, and 0.059 $\mu\text{g m}^{-3}$, respectively. It should be noted that across much of the MBL, nitrate mass concentrations
are extremely low, with the AMS frequently reporting zero values. As a result, the spatial coverage of valid nitrate data is
840 limited, and we therefore refrain from reporting its median value. Given that nitrate constitutes only a very small fraction of
the total measured and modelled aerosol mass in the MBL, its underestimation does not offset the overestimation of other
components, leading to an overall overestimation of total submicron aerosol mass by a factor of 1.25 in the MBL by the
benchmark simulation (Table 3).~~

This seemingly contradictory result - overestimated aerosol mass alongside underestimated aerosol number concentrations
845 in the MBL - can be reconciled by considering the modal contributions: the model overestimates aerosol concentrations in the
accumulation mode (which dominates aerosol mass), while simultaneously underestimating concentrations in the nucleation
and Aitken modes (which dominate aerosol number). It should also be noted that the aerosol size distribution in Figure 8
shows that the model's accumulation mode aerosol distribution is skewed towards larger sizes, which is consistent with the
overestimation of aerosol mass in the MBL.

850 3.4.1 Sulfate

Except in very specific regions (e.g. the UT south of 40 °S and MBL), sulfate aerosol mass is consistently overestimated by the
model, with the magnitude of this overestimation typically being a few tens of percent across much of the sampled atmosphere
(Figures 9 and 10). For example, between 2 - 8 km, the default (SA-H₂O) and benchmark (SA-NH₃) simulations overestimate
sulfate by 42 % and ~~32-34~~ % , respectively (Table A1). The only model simulations that significantly change sulfate aerosol
855 mass concentrations, between 2 - 8 km, are SA-NH₃-Lana, SA-NH₃-CS2 and SA-NH₃-MSA. By modifying the DMS emission
scheme, the SA-NH₃-Lana simulation increases the sulfate overestimation by ~~14-11~~ % , resulting in a model-to-observation
ratio of ~~1.461.45~~. This increase is likely driven by higher DMS emissions from the adopted climatological dataset, relative
to the default interactive scheme, as discussed earlier. On the other hand, the SA-NH₃-CS2 simulation reduces the sulfate
overestimation from ~~32-34~~ % (in the benchmark) to ~~6-5~~ % , while the SA-NH₃-MSA simulation increases the overestimation
860 to 45 % (Table A1). The reduction in sulfate overestimation in the SA-NH₃-CS2 simulation is attributed to the lower SO₂
concentrations resulting from the CS2 chemistry scheme (see earlier discussions). The additional ~~13-11~~ % increase in bias in
the SA-NH₃-MSA simulation is due to the inclusion of condensable MSA vapours, which are treated as part of the sulfate
aerosol component in the model for mass accounting, as MSA is not independently treated in the model.

It is important to mention that the most significant changes in sulfate aerosol mass concentrations ~~relative to ATom~~, due
865 to these scheme modifications, appear to concentrate in relatively pristine environments, such as the tropical UT and the SH
(Figure 10). This supports that these changes are primarily driven by changes to the atmospheric chemistry of DMS (in SA-
NH₃-CS2) and the inclusion of its oxidation products like MSA as condensable species (in SA-NH₃-MSA). Anthropogenic
primary sulfate aerosol emissions and direct emissions of SO₂, which are more dominant in the NH, are not significantly

affected by these specific scheme changes; therefore, sulfate aerosol mass concentrations in the NH are less impacted. This
870 ~~again~~ highlights the need to improve the representation of DMS sources and its atmospheric chemistry in the model for
simulating aerosols in the pristine regions of the atmosphere.

One primary reason for this general global overestimation of sulfate aerosol is the model's overestimation of SO₂
concentrations, a key precursor for sulfate aerosol formation. Sulfate aerosol in the model is formed either from primary
sources from anthropogenic emissions, or secondary processes such as from the gas-phase condensation of H₂SO₄, or, more
875 significantly, through aqueous-phase oxidation of SO₂ in cloud droplets by H₂O₂ and O₃ (Mulcahy et al., 2020). Consequently,
the accuracy of simulated sulfate aerosol concentrations is also sensitive to the modelled concentrations of other species
involved in these pathways, such as gas-phase OH and aqueous-phase oxidants like H₂O₂ and O₃. Furthermore, the use of
a fixed cloud pH value of 5.0 in the model may also affect the efficiency of SO₂ uptake (Mulcahy et al., 2020). Since the
model's treatment of these multi-phase processes is relatively simplified, further mechanistic implementation and evaluation
880 of these processes are needed to improve the model's representation of sulfate aerosol.

Another potential contributor to the overestimation of sulfate is the underestimation of its removal processes, leading to
an overestimated aerosol lifetime. For instance, Mulcahy et al. (2020) reported a sulfate (SO₄²⁻) lifetime of 5.57 days in
UKESM1.0, which is around two days longer than the estimate of 3.7 days by Mann et al. (2010). This value also lies
at the upper end of the range reported by AeroCom-I models, which spans from 3 to 5.4 days, suggesting that sulfate
885 removal may be too slow in UKESM. ~~It is worth highlighting that sulfate aerosol remains the dominant component of total
submicron aerosol mass in the MBL, accounting for 39 % of the total submicron aerosol mass in the benchmark simulation.
Accurately representing its sources, from both natural and anthropogenic emissions, is therefore crucial for improving the
model's simulation of MBL aerosol properties and, ultimately, CCN number size distributions.~~

3.4.2 Organic matter

890 The model-ATom discrepancy in organic aerosol mass exhibits a distinct dipole pattern: underestimation in the upper
troposphere (8 - 12 km) and the tropical lower to mid free troposphere (2 - 8 km), and moderate overestimation across the
rest of the sampled atmosphere (Figures 10 and A3). In the UT, this underestimation is most likely attributed to a lack of
significant organic aerosol sources in the model at these altitudes. A clear missing source of organic aerosol in the model's UT
is SOA derived from the oxidation products of isoprene, which are not explicitly included as a significant SOA source in the
895 default model configuration. Recent ambient observations and laboratory studies have shown that isoprene oxidation products
can contribute significantly to new particle formation and growth in the tropical UT (Shen et al., 2024; Curtius et al., 2024).

Although a comprehensive isoprene new particle formation mechanism cannot be implemented in the model at this stage
due to a lack of readily available parameterisations, this study attempts to partially account for this missing contribution of
isoprene oxidation products to aerosol growth using the SA-NH₃-IPSOA and SA-NH₃-IPSOA×10 simulations.

900 The implementation of the explicit IPSOA scheme significantly increases modelled organic aerosol mass concentrations
in the tropical UT. The median measured organic aerosol mass concentration during the ATom campaigns in this region is
0.083 μg m⁻³. The model-to-ATom ratio in the benchmark SA-NH₃ simulation for this region is ~~0.36~~0.32, indicating an

underestimation by around 3 times. With a 3 % SOA mass yield from isoprene oxidation, the SA-NH₃-IPSOA simulation improves the model-to-ATom ratio for organic aerosol mass concentration in the tropical UT to 0.41. A 30 % SOA mass yield (SA-NH₃-IPSOA×10 simulation) further improves this comparison, increasing the model-to-ATom ratio to 0.86 (Figure 10).

In the MBL, the model simulations generally overestimate organic aerosol mass concentrations, for example by a factor of ~~1.37-1.42~~ in the benchmark SA-NH₃ simulation. The SA-NH₃-IPSOA simulation ~~, which introduces a 3 % isoprene SOA yield,~~ results in a model-to-ATom ratio of ~~1.34-1.35~~ for MBL organic aerosol. This slight reduction in the MBL organic aerosol overestimation, despite adding an isoprene SOA source, is likely due to the concurrent reduction in the prescribed monoterpene SOA yield ~~. In the default model, the monoterpene SOA yield is scaled by two (effectively 26 %) to implicitly account for missing isoprene SOA. When the explicit isoprene SOA pathway is activated in the SA-NH₃-IPSOA simulation, this scaling is removed, and the monoterpene SOA yield reverts to its base value (13 %), which could result in a lower overall SOA production from these combined sources compared to the default scaled approach (Table 2 (see Table 2 and Methods)).~~ The SA-NH₃-IPSOA×10 simulation, on the other hand, significantly worsens the organic aerosol mass overestimation in the MBL, increasing the model-to-ATom ratio to ~~1.73-1.72~~. This increased overestimation likely results from an isoprene SOA yield (30 %) that is unrealistically high for the warmer MBL conditions, where higher temperatures generally lead to lower SOA yields. Therefore, accurately quantifying the SOA yield from isoprene oxidation, both in the cold UT and warm MBL, is crucial for improving the model's representation of organic aerosols.

Another notable feature is the generally stronger underestimation of submicron organic aerosol mass in the NH compared to the SH. This disparity is likely due to the model's omission of secondary organic aerosol contributions from anthropogenic precursors, a limitation that warrants further investigation.

3.4.3 Ammonium and nitrate

The ammonium and nitrate aerosol scheme, recently developed by Jones et al. (2021), is not included in the default UKESM1.1 configuration. Therefore, the default (SA-H₂O) and SA-NH₃-noNit simulations do not include these two aerosol components, while the ammonium nitrate scheme is included in all other simulations discussed in this study. ATom observations indicate that the mass concentrations of both ammonium and (inorganic) nitrate are significantly lower than those of sulfate and organic matter globally. This is particularly true for nitrate mass, which exhibits low concentrations globally (Figure 9).

For the region where the nitrate component is above the detection limit of AMS, model simulations generally show an overestimation of nitrate aerosol mass concentrations compared to ATom observations below 8 km (Tables 3 and A1). For example, the benchmark SA-NH₃ simulation reports a model-to-ATom ratio of ~~9.61-11.27~~ in the MBL, indicating a significant overestimation by the model (Tables 3). Using the slow nitric acid uptake coefficient (SA-NH₃-slow simulation) reduces this overestimation in the MBL, with the ratio reduced to ~~3.86-2.93~~. This discrepancy might suggest that the nitric acid uptake coefficient is still too fast, and using an even slower value could improve the simulation of nitrate aerosol mass concentrations in this NH. Alternatively, the overprediction may be from overestimated anthropogenic NO_x emissions, which would lead to excessive atmospheric nitric acid and, consequently, particulate nitrate. A third possible explanation is that the model

underestimates the deposition rates of NO_x and its oxidation products. Therefore, the exact cause of the nitrate aerosol overestimation in the NH remains uncertain and requires further investigation.

In contrast to nitrate, ammonium aerosol mass concentrations are ubiquitously overestimated by the model throughout the sampled atmosphere when the ammonium nitrate scheme is active (Figures 10 and A2). This overestimation of ammonium
940 likely translates into an overestimation of aerosol pH (i.e. an underestimation of aerosol acidity) globally, consistent with findings for other chemical transport models reported by Nault et al. (2021). In the MBL, the benchmark SA- NH_3 simulation overestimates ammonium aerosol mass concentrations by a factor of ~~3.17~~ 3.20 compared to ATom observations (Table 3). Furthermore, this positive model bias for ammonium is even larger above the MBL. For example, the bias increases to a factor of ~~7.68~~ 7.73 between 2 - 8 km altitude. It is worth noting that both ammonium aerosol and gas-phase NH_3 concentrations
945 are significantly overestimated by the model in the MBL (Table 3). This pervasive overestimation of ammonium is unlikely to be solely explained by flaws in the implemented ammonium nitrate scheme itself. Improvements to either or both the NH_3 emission inventories and the representation of its atmospheric removal processes are needed to enhance the simulation of both gaseous NH_3 and particulate ammonium in the model.

3.4.4 Seasalt

950 Seasalt aerosols are a major component of primary marine CCN and play a crucial role in cloud microphysics and climate. The precise relative contributions of seasalt aerosols (primary) and new particle formation processes to the total CCN population remain uncertain. These two major aerosol sources have distinct feedback mechanisms within the climate system and are projected to exhibit different responses to future environmental changes. It is therefore essential to accurately represent both processes in models to enable robust assessments of future changes in CCN, cloud microphysics, and ultimately, climate.

955 As expected, the measured seasalt concentrations in the MBL (Murphy et al., 2019). However, the simulations without the ammonium nitrate scheme (SA- H_2O and SA- NH_3 -noNit) overpredict seasalt aerosol mass concentrations in submicron aerosols within the MBL, with the default simulation predicting concentrations 2.09 times higher than observed (Table 3). This overestimation persists consistently across latitudes, from the tropics to high-latitude regions (Figure 10). Additionally, simulations that include the ammonium nitrate scheme show a reduction in ~~sea-salt~~ seasalt aerosol mass. This decrease is
960 possibly linked to chemical processing, specifically the displacement of chloride by nitrate through heterogeneous uptake of HNO_3 on seasalt particles, leading to the formation of hydrogen chloride (HCl) (Jones et al., 2021). However, given the very low nitrate concentrations in the marine boundary layer, it remains uncertain whether sufficient nitrate is available to drive appreciable chloride displacement. Further investigation is needed to better understand the implications of the ammonium nitrate scheme on seasalt conservation in marine environments.

965 Constraining seasalt emissions, particularly with geometric diameter smaller than $1 \mu\text{m}$, is crucial for accurately understanding the CCN budget in the MBL, as Aitken and accumulation mode aerosols are the major contributors to CCN concentrations. The default model's overestimation of seasalt aerosols in the MBL will likely lead to an exaggerated contribution from primary seasalt to the CCN budget, potentially masking or underestimating the CCN contribution from airborne aerosol formation processes. The default model's overprediction of seasalt aerosols in the marine atmosphere is also

970 supported by a recent study by Venugopal et al. (2025), which found that the default seasalt emission scheme in UKESM overpredicts the dependence of seasalt emissions on wind speed when validated against aerosol optical depth observations.




Figure 9. Curtain plots of aerosol composition along ATom flight tracks. Mean sulfate, organic, ammonium, nitrate and seasalt mass concentrations are shown for ATom observations (first row) and selected model simulations (subsequent rows; see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. All data are displayed on a logarithmic to enhance visualisation of the variations in parameters. Colour scales are consistent within each column to facilitate direct comparison between observations and model simulations. The SA-H₂O (default) simulation does not include ammonium or nitrate in aerosols, thus the absence of data in the corresponding plots.




Figure 10. Curtain plots of model-to-ATom ratios for aerosol composition along ATom flight tracks. Mean ratios of modelled to observed values for sulfate, organic, ammonium, nitrate and seasalt mass concentrations are shown for selected model simulations (see Table 2 for simulation details). Model outputs are interpolated to ATom flight coordinates and times. Data are displayed on a logarithmic scale with a diverging colour scheme where values greater than 1 indicate model overestimation and values less than 1 indicate model underestimation. Colour scales are consistent within each column to facilitate comparison between different model configurations. The SA-H₂O (default) simulation does not include ammonium or nitrate in aerosols, thus the absence of data in the corresponding plots.

4 Discussion

Previous sections have evaluated the model's performance in simulating individual environmental conditions, precursor vapours, aerosol number size distributions, and aerosol chemical composition against ATom observations. Addressing the identified discrepancies is crucial for improving the representation of aerosol processes in the model, with the ultimate goal

of achieving accurate predictions for the correct physical and chemical reasons. In this section, we synthesise our findings to discuss the implications of our model development endeavours and to outline future directions for improving aerosol simulations.

980 The discussions focus on two specific regions: the UT, particularly between 8 and 12 km altitude, and the MBL, below 2 km altitude. The selection of these two regions is based on several key considerations. Firstly, they are the regions where the model exhibits its most significant biases for numerous evaluated parameters. Secondly, the UT is a region where ATom observations reveal significant enhancements in aerosol number concentrations, often driven by strong new particle formation (Williamson et al., 2019). Thirdly, the MBL is critical due to the sensitivity of marine low-level clouds, a key component of the Earth's radiative balance, to the MBL aerosol population that acts as CCN.

985 4.1 Marine boundary layer processes

The systematic overestimation of aerosol nucleation precursors (SO_2 , OH and NH_3) alongside the underestimation of nucleation and Aitken mode particle concentrations in the MBL highlights a fundamental inconsistency between current model representations and the actual processes governing airborne aerosol formation in marine environments. This paradox - where abundant precursor vapours fail to generate sufficient particle numbers - suggests that $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ nucleation alone cannot
990 account for observed aerosol formation rates in the marine atmosphere.

The implementation of the $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation mechanism with the ammonium nitrate scheme produces mixed results. While this leads to modest improvements in nucleation mode particle concentrations, it worsens the underestimation of Aitken mode aerosols (Table 3) and generally reduces Aitken mode concentrations across the MBL, particularly in pristine marine regions (Figure 11). In contrast, accumulation mode aerosol concentrations exhibit widespread increases, especially
995 over populated continental regions where elevated NH_3 emissions promote both aerosol mass formation and $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation.

Model sensitivity to different schemes is substantial. Adding condensable vapours such as MSA increases accumulation mode concentrations, particularly over the Southern Ocean where DMS emission is high, while simultaneously reducing Aitken mode concentrations as aerosols grow more rapidly to the accumulation mode (Figure 12). Switching DMS emission schemes
1000 or chemistry schemes (from Strat-Trop to CS2) produces significant regional changes, with polar regions being especially sensitive due to their low background aerosol concentrations.

Beyond aerosol numbers, the default model overestimates total submicron aerosol mass in the MBL. For example, seasalt is overestimated in the default model, potentially leading to exaggerated primary CCN contributions. The concurrent overestimation of both gas-phase NH_3 and particulate ammonium in the simulations with the ammonium nitrate scheme
1005 suggests fundamental issues with NH_3 budget representation.

These findings indicate that the default model's CCN budgets in the MBL are incorrectly described, with excessive dependence on primary emissions such as seasalt masking deficient airborne aerosol formation processes. This imbalance has important implications for preindustrial baseline estimates, as the relative importance of natural versus anthropogenic

aerosol sources may be fundamentally misrepresented. The results strongly support incorporating additional aerosol formation pathways, likely involving iodine oxoacids, organic vapours, or other marine-specific precursors.




Figure 11. Global distribution of marine boundary layer aerosol changes following implementation of the H₂SO₄–NH₃ nucleation mechanism. Panels show **monthly mean** aerosol number concentrations in the marine boundary layer (0 - 2 km altitude) averaged **between over monthly means from** July 2016 **and to** June 2018: (A) Aitken mode aerosols from the default SA-H₂O simulation, (B) accumulation mode aerosols from the default SA-H₂O simulation, (C) ratio of Aitken mode concentrations between benchmark SA-NH₃ and default SA-H₂O simulations, and (D) ratio of accumulation mode concentrations between benchmark SA-NH₃ and default SA-H₂O simulations.




Figure 12. Global distribution of marine boundary layer aerosol changes following implementation of the Lana et al. (2011) DMS climatology, MSA condensation and CRI-Strat2 scheme. Panels show **monthly mean** aerosol number concentrations in the marine boundary layer (0 - 2 km altitude) averaged **between-over monthly data from** July 2016 **and-to** June 2018: (A) Aitken mode aerosols from the benchmark SA-NH₃ simulation, (B) accumulation mode aerosols from the benchmark SA-NH₃ simulation, (C) ratio of Aitken mode concentrations between SA-NH₃-Lana and benchmark SA-NH₃ simulations, and (D) ratio of accumulation mode concentrations between SA-NH₃-Lana and benchmark SA-NH₃ simulations, (E) ratio of Aitken mode concentrations between SA-NH₃-MSA and benchmark SA-NH₃ simulations, and (F) ratio of accumulation mode concentrations between SA-NH₃-MSA and benchmark SA-NH₃ simulations, (G) ratio of Aitken mode concentrations between SA-NH₃-CS2 and benchmark SA-NH₃ simulations, and (H) ratio of accumulation mode concentrations between SA-NH₃-CS2 and benchmark SA-NH₃ simulations.

4.2 Upper tropospheric processes

The model's biases for the aerosol number size distribution in the UT are essentially the opposite of those in the MBL. Specifically, the model overestimates nucleation and Aitken mode aerosol concentrations while underestimating accumulation mode aerosol concentrations. This pattern of too many small aerosols and too few large ones may suggest an imbalance in the model between new particle formation and subsequent aerosol growth processes. The UT provides a clearer reflection of the model's performance in simulating these airborne aerosol formation processes, as it is a region where most primary aerosols are not efficiently transported, minimising the confounding influence of primary source biases that are more prominent in the MBL. In the default simulation, the model overestimates nucleation and Aitken mode aerosols by factors of 6.30 and 1.88, respectively, while the model-to-ATom ratio for accumulation mode aerosols is only 0.39 (Table A2). As discussed in earlier sections, this overestimation of small aerosols primarily results from the excessive aerosol nucleation rate predicted by the Vehkamäki et al. (2002) scheme, as well as from contributing factors such as the overestimated concentrations of OH and H₂O (Figures 2 and 4).

Implementing the H₂SO₄-NH₃ nucleation mechanism and ammonium nitrate scheme in the benchmark simulation significantly reduces the overestimation of nucleation and Aitken mode aerosols, yielding improved model-to-ATom ratios of ~~1.79~~ and ~~1.39~~ 1.74 and 1.41, respectively. Simultaneously, the underestimation of accumulation mode aerosols is also reduced (i.e. improved), with the model-to-ATom ratio increasing to 0.52. This can be understood in terms of mass balance: a lower nucleation rate results in fewer nucleation and Aitken mode aerosols, but the available condensable vapour is then distributed among fewer aerosols, promoting more efficient growth of the remaining population into the accumulation mode. It is worth mentioning that since sulfate aerosol mass is reasonably simulated by the model in the UT (ratio of ~~1.01~~ 1.04 in the benchmark simulation), the missing mass needed to grow aerosols into the accumulation size range and to match observations must be accounted for by other components. As nitrate, ammonium, and seasalt aerosol mass concentrations are all too low in the UT over remote marine atmosphere (Figure 9), organic matter is the only remaining component that can account for this missing mass, consistent with the hypothesis by (Williamson et al., 2019; Kupc et al., 2020).

Indeed, all simulations without the IPSOA scheme show significantly underestimated organic aerosol mass concentrations in the UT, with the benchmark simulation underestimating it with a ratio of ~~0.44~~ 0.45. Isoprene is the most likely candidate to account for this missing organic aerosol mass, as its global emission is large (around 500 Tg yr⁻¹), and its oxidation products have recently been discovered to contribute significantly to new particle formation and growth in the UT (Shen et al., 2024; Curtius et al., 2024). Implementing the IPSOA scheme with a high mass yield of 30 % (in the SA-NH₃-IPSOA×10 simulation) significantly improves the agreement, increasing the model-to-ATom ratio for organic aerosol mass in the UT to 0.86. The inclusion of the IPSOA scheme results in the largest enhancements in accumulation mode aerosol concentrations above tropical rainforests (Figure 13), a finding consistent with satellite observations (Palmer et al., 2022). It is worth highlighting that the impact of including IPSOA stretches beyond the tropical UT, extending to the mid- and high-latitude UT. This demonstrates that the long lifetime of aerosols in the UT allows them to be transported over great distances, thereby exerting a global impact. Addressing this missing organic aerosol component, and thereby the underestimated accumulation mode aerosol concentrations

1045 in the UT, is crucial for improving the model's representation of ice nucleating particles and cirrus cloud formation, which in turn affects the Earth's longwave radiation balance.

Once again, changing the chemistry scheme from Strat-Trop to CS2 has a large impact on the modelled agreement for Aitken and accumulation mode aerosols in the UT (Figure 13). The reduction in SO₂ concentrations in the SA-NH₃-CS2 simulation is the clear cause of the reduction in these aerosol populations, as can be seen in Figure 2. This is also supported by the aerosol
1050 composition analysis, which shows a corresponding reduction in the sulfate aerosol mass bias, with insignificant changes in other aerosol components (Figure 10).




Figure 13. Global distribution of upper tropospheric aerosol changes following implementation of the CRI-Strat2 scheme and isoprene secondary organic aerosol formation scheme. Panels show **monthly-mean** aerosol number concentrations in the upper troposphere (8 - 12 km altitude) averaged **between-over monthly means from July 2016 and-to June 2018**: (A) Aitken mode aerosols from the benchmark SA-NH₃ simulation, (B) accumulation mode aerosols from the benchmark SA-NH₃ simulation, (C) ratio of Aitken mode concentrations between SA-NH₃-CS2 and benchmark SA-NH₃ simulations, and (D) ratio of accumulation mode concentrations between SA-NH₃-CS2 and benchmark SA-NH₃ simulations, (E) ratio of Aitken mode concentrations between SA-NH₃-IPSOA×10 and benchmark SA-NH₃ simulations, and (F) ratio of accumulation mode concentrations between SA-NH₃-IPSOA×10 and benchmark SA-NH₃ simulations.

4.3 Recommendations for future model improvements

The evaluation of UKESM against ATom observations in this study has identified several key areas for improving the model's representation of aerosol microphysics and chemistry in the marine atmosphere. These involve primary aerosol emissions, precursor vapours, nucleation and growth processes, and aerosol chemical composition. While each evaluated parameter has its own specific bias and uncertainties, the two main trends are 1) the overestimated aerosol mass and precursor gasses and

2) the underrepresented aerosol number. This indicates that the airborne aerosol formation rates are underestimated in the UKESM.

Many of the airborne aerosol formation mechanisms are essentially missing in the model, which requires extensive efforts to implement and evaluate these processes. Several candidate mechanisms have been proposed in the literature. The first and most prominent candidate is iodine oxoacids (iodic acid, HIO_3 , and iodous acid, HIO_2), which are widely acknowledged as important for aerosol nucleation in the MBL. Iodine oxoacids are ubiquitously measured in the global atmosphere, from pristine marine to polluted urban environments (Beck et al., 2021; He et al., 2021b, 2023; Zhang et al., 2024). Depending on the location and season, the concentrations of iodine oxoacids are often comparable to (within one order of magnitude of) or can even exceed those of H_2SO_4 . Dedicated laboratory experiments have revealed that on a per-molecule basis, iodine oxoacids are significantly more efficient than H_2SO_4 at forming new particles (He et al., 2021b). Furthermore, when iodine oxoacids are mixed with H_2SO_4 and NH_3 , the synergistic aerosol nucleation of the sulfur-iodine system is even more efficient than any of the individual components alone (He et al., 2023). A recent global model study has also shown that the inclusion of iodine oxoacids significantly improved modelled aerosol number concentrations in the MBL (Zhao et al., 2024), although the parameterisation used in that study was mostly a fit to experimental data of He et al. (2021b) and did not include the synergistic effects of the sulfur-iodine system (He et al., 2023). It is worth noting that while Zhao et al. (2024) suggested that iodine oxoacid-driven aerosol nucleation has a negligible impact above the MBL due to rapid scavenging of iodine species in the MBL, ambient observations suggest the widespread presence of iodine compounds throughout the troposphere, including the UT (Koenig et al., 2020; He et al., 2021b, 2023; Schill et al., 2025). This discrepancy is likely due to the absence of halogen recycling mechanisms in the model used by Zhao et al. (2024).

The second candidate mechanism is nucleation involving atmospheric amines, which have been shown to have strong synergistic effects with H_2SO_4 (Almeida et al., 2013). Sporadic ambient observations of the participation of amines in aerosol nucleation have been reported, for example, near the Antarctic Peninsula (Brean et al., 2021). However, direct measurements of nucleating cluster ions in polar regions typically show a stronger participation of NH_3 rather than amines (Beck et al., 2021; Jokinen et al., 2018), and the global relevance of amines in marine aerosol nucleation is yet to be determined. Global simulations along the ATom flight paths have also suggested a minor contribution of amines to aerosol nucleation in the MBL (Zhao et al., 2024).

The third candidate mechanism is airborne aerosol formation from organic vapours, such as monoterpenes and isoprene, or their synergistic nucleation with inorganic acids (e.g. H_2SO_4 and iodine oxoacids). Unfortunately, the role of organic vapours in MBL aerosol nucleation remains highly debated, as their marine source strength and subsequent atmospheric concentrations are highly uncertain. For example, Luo and Yu (2010) estimated that global oceanic emissions of α -pinene and isoprene using bottom-up estimates are $0.013 \text{ TgC yr}^{-1}$ and 0.32 TgC yr^{-1} , respectively, significantly lower than oceanic DMS emissions. However, the same study reported that using a top-down approach (forcing the model to match observed organic concentrations) resulted in much higher global emissions of 29.5 TgC yr^{-1} for α -pinene and 11.6 TgC yr^{-1} for isoprene, values comparable to global oceanic DMS emissions. More recent in-situ measurements of monoterpenes in the Arctic and Atlantic Oceans suggest concentrations below 5 pptv, supporting an estimated global marine monoterpene emission of 0.16 TgC yr^{-1} (Hackenberg

et al., 2017), which aligns better with the bottom-up estimate of Luo and Yu (2010). Therefore, further constraining marine organic vapour emissions is needed to better represent the role of organic vapours in new particle formation in the MBL. On the other hand, in the UT, the role of isoprene derivatives in new particle formation is unambiguous as reported in Shen et al. (2024); Curtius et al. (2024). The long range transport of the formed aerosols will affect the aerosol populations in the marine atmosphere.

Finally, we outline some of the most important recommendations. While not exhaustive, they provide a solid foundation for future model development efforts.

- Re-evaluate primary aerosol emissions in the MBL with a focus on seasalt and organics.
- 1100 – Re-evaluate the aerosol deposition schemes in the model.
- Re-evaluate the marine NH₃ emission inventory and investigate the bias in the atmospheric NH₃ budget both in the gas phase and in aerosols (NH₄⁺).
- Investigate and improve the representation of OH in the atmosphere, especially in the southern hemisphere.
- Improve the atmospheric DMS chemistry scheme as well as the DMS emissions, particularly in the Southern Ocean, where the model currently overestimates DMS emissions.
- 1105 – Implement comprehensive halogen chemistry in the model with a focus on representing iodine chemistry and iodine oxoacid production.
- Implement the iodine oxoacid new particle formation mechanism in the model, together with the synergistic effects of sulfur-iodine aerosol nucleation, to improve the representation of new particle formation in the MBL.
- 1110 – Implement experimentally verified MSA condensation scheme in the model.
- Improve model treatment of organic oxidation pathways, especially in the upper troposphere.
- Implement the isoprene new particle formation mechanism in the model, together with the synergistic aerosol nucleation of isoprene derivatives with H₂SO₄ and iodine oxoacids, to improve the representation of new particle formation in the upper troposphere.

1115 It should be noted that the need to implement new aerosol formation schemes is not unique to UKESM; other Earth system model evaluation studies, such as those using EC-Earth3, have reached similar conclusion (Svenhag et al., 2024). While implementing these comprehensive chemical and microphysical pathways will increase the computational demand on the model's chemical solver, such developments are essential for capturing the true sensitivity of the climate to aerosols. Once these full mechanisms are implemented and validated, they can be systematically simplified or parameterised based on their performance to reduce the computational burden for long-term climate simulations without sacrificing significant accuracy, for the eventual CMIP implementations.

1120

5 Conclusions

This study provides a detailed evaluation of the United Kingdom Earth System Model version 1.1 (UKESM1.1), using observations from the Atmospheric Tomography (ATom) mission to examine parameters relevant to the aerosol lifecycle in the remote marine atmosphere. By comparing model outputs with measurements of precursor vapours, aerosol number size distributions, and aerosol chemical composition, we identify several key shortcomings in current model's representation of aerosol microphysics and chemistry, and offer targeted recommendations for future development.

Our analysis highlights two contrasting biases in UKESM1.1's treatment of aerosols in the marine boundary layer. The default model overestimates primary submicron seasalt, while underrepresenting airborne aerosol formation processes. At the same time, the model underestimates nucleation and Aitken mode aerosols, with model-to-ATom ratios down to 0.09 and 0.57, respectively. This imbalance suggests that the current cloud condensation nuclei budget in the model is overly reliant on primary emissions, potentially misrepresenting the importance of airborne aerosol formation processes that are likely important in pristine marine regions.

In contrast, the model exhibits an opposite bias in the upper troposphere, generating excessive nucleation and Aitken mode aerosols due to the default $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ nucleation scheme, but failing to grow them efficiently to accumulation mode sizes relevant for ice cloud formation.

To address these problems, we implement a $\text{H}_2\text{SO}_4\text{-NH}_3$ nucleation scheme alongside an ammonium nitrate module developed by Jones et al. (2021). This update partly improves performance in simulating aerosol number size distributions in the upper troposphere but does little to resolve the discrepancies in the MBL. The continued underprediction of small aerosols in the MBL - despite substantial overestimation of precursors such as SO_2 , OH, and NH_3 - strongly suggest that the model's representation of airborne aerosol formation processes is incomplete.

Incorporating additional condensable vapours such as isoprene oxidation products, improve the model simulations in the upper troposphere. A modified isoprene secondary organic aerosol scheme with enhanced yields improves observed organic aerosol concentrations in the upper troposphere, consistent with emerging evidence that low temperature isoprene oxidation significantly contributes to aerosol formation and growth aloft (Shen et al., 2024; Curtius et al., 2024). The implemented methanesulfonic acid condensation scheme increases particle mass and alters the particle number size distribution; however, its quantitative impact remains uncertain pending more accurate experimental constraints.

Additionally, the model simulations in this study indicate that the concentrations of cloud condensation nuclei (CCN)-active Aitken and accumulation mode aerosols in the MBL are significantly influenced not only by the implemented aerosol nucleation and growth schemes but also the emission mechanisms for precursor vapours, and the adopted atmospheric chemistry mechanisms. For instance, switching from the interactive DMS emission scheme to the Lana et al. (2011) climatology resulted in substantial changes to both Aitken and accumulation mode aerosol distributions, with reductions south of 30°S and enhancements in tropical regions. Changing the chemistry scheme from Strat-Trop to CRI-Strat2 leads to significant reductions in SO_2 concentrations, which in turn affects aerosol nucleation rates and size distributions throughout the marine atmosphere. These sensitivity tests demonstrate that achieving accurate CCN predictions requires not only mechanistically

sound aerosol nucleation and growth parameterisations, but also reliable representations of precursor emissions and their atmospheric transformation pathways. The interconnected nature of these processes means that improvements in one area can be offset by deficiencies in another, highlighting the need for comprehensive, process-based model development rather than isolated parameter adjustments.

1160 These findings carry important implications for climate modeling. The tendency to overestimate primary submicron seasalt
in the MBL suggests that the default UKESM1.1 might underestimate the role of anthropogenic aerosol sources from e.g. SO₂
in shaping present-day CCN populations. Furthermore, the absence of key natural airborne aerosol formation pathways in the
model, potentially involving iodine oxoacids, organic vapours, or their interactions with H₂SO₄, also contribute to both the
preindustrial and present day CCN populations in the atmosphere. These factors could skew estimates of preindustrial aerosol
1165 baselines and the strength of aerosol-cloud interactions.

Our results underscore the need to incorporate comprehensive halogen chemistry, especially iodine-related aerosol
nucleation mechanisms, which are increasingly recognised in both laboratory and field studies as major contributors to marine
aerosol formation. Enhancing model treatments of organic aerosol sources and their oxidation pathways is also essential,
particularly for the upper troposphere, where long range transport of isoprene-derived aerosols can shape aerosol populations
1170 on a global scale.

Looking ahead, model improvement efforts should prioritise process-based improvement over empirical tuning. The long-
lasting uncertainties in aerosol forcing, despite decades of model development, highlight the limitations of ad-hoc adjustments
and point to the importance of mechanistic understanding. Capturing the underlying physics and chemistry of aerosol processes
is one of the prerequisite for building models capable of producing robust climate projections and informing effective mitigation
1175 strategies.

The ATom dataset offers a powerful benchmark for advancing this work, but fully leveraging its value will require sustained
investment in experimental research. Laboratory studies of nucleation chemistry and targeted field campaigns are particularly
needed to resolve the complex pathways governing natural aerosol formation. As anthropogenic sulfur emissions continue to
decline, accurately representing these natural processes will become even more crucial for predicting future climate trajectories.

1180 . The ATom aerosol number size distribution measurements are publicly accessible at <https://doi.org/10.3334/ORNLDAAAC/2111> (Brock
et al., 2022). The ATom atmospheric trace gas measurements are archived at <https://doi.org/10.3334/ORNLDAAAC/1925> (Wofsy et al., 2021).
ATom aerosol composition measurements are also available at <https://doi.org/10.3334/ORNLDAAAC/1925> (Wofsy et al., 2021); however,
the specific dataset utilised in this study, together with the real-time transmission correction curve, was provided by P.C.-J. and J.L.J. The
integrated 5-minute resolution data product combining collocated UKESM1.1 model output with subsampled ATom observational data will
1185 be deposited in the Zenodo repository upon acceptance of this manuscript. Additional UKESM1.1 model output is available from X.-C.H.
upon reasonable request.

. X.-C.H., N.L.A., A.T.A., and H.G. conceived this study. N.L.A. and A.T.A. provided the computational environment and resources on ARCHER2 and H.G. on Bridges-2. X.-C.H., H.D., X.W., J.W., and H.G. carried out model development and module incorporation. X.-C.H. carried out model simulations with assistance from N.L.A., J.W., and H.G. Additionally, P.C.-J., B.N., A.K., D.B., J.L.J., and C.J.W. provided ATom observational data. X.-C.H. carried out data analysis and wrote the manuscript. All authors have commented and edited the manuscript.

. At least one of the (co-)authors is a member of the editorial board of *Atmospheric Chemistry and Physics*.

. This work used the ARCHER2 UK National Supercomputing Service. This work used the Bridges-2 machine at the Pittsburgh Supercomputing Center (PSC) through allocation atm200005p from the Advanced Cyberinfrastructure Coordination Ecosystem: Services and Support (ACCESS) program, which is supported by National Science Foundation grants no. 2138259, 2138286, 2138307, 2137603, and 2138296. We [thank Kenneth S. Carslaw for insightful discussions on the sulfuric acid - ammonia aerosol nucleation mechanism. We](#) additionally thank Charles A. Brock, William H. Brune, Thomas B. Ryerson, Alexander B. Thames, Chelsea R. Thompson, and Steven C. Wofsy for providing data directly relevant to this study, and extend our gratitude to the entire ATom team for their extraordinary work in collecting and curating these invaluable atmospheric measurements.

. X.-C.H. acknowledges support from Research Council of Finland grant no. 359331, 349659, 371185. H.G. and H.D. acknowledge support from the NASA ROSES program via grant no. 80NSSC19K0949, and H.G. additionally acknowledges support from the US Department of Energy Atmospheric System Research program via grant no. DE-SC-0022227. A.T.A. acknowledges Natural Environment Research Council CARES project no. NE/W009412/1. N.L.A. and M.R.R. thank Natural Environment Research Council for financial support through the VISION project, NE/Z503393/1. D.B. acknowledges support from NASA grant no. NNX15AJ91A. P.C.-J., B.N., and J.L.J. are supported by NASA grant no. 80NSSC21K1451 and 80NSSC23K0828.

1205 **References**

- Almeida, J., Schobesberger, S., Kürten, A., Ortega, I. K., Kupiainen-Määttä, O., Praplan, A. P., Adamov, A., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos, M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurtén, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppä, J., Loukonen, V., Makhmutov, V.,
1210 Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petäjä, T., Riccobono, F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld, J. H., Simon, M., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Tröstl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen, Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T., Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H., and Kirkby, J.: Molecular Understanding of Sulphuric Acid–Amine Particle Nucleation in the Atmosphere, *Nature*, 502, 359–363,
1215 <https://doi.org/10.1038/nature12663>, 2013.
- Anderson, T., Spall, S., Yool, A., Cipollini, P., Challenor, P., and Fasham, M.: Global Fields of Sea Surface Dimethylsulfide Predicted from Chlorophyll, Nutrients and Light, *Journal of Marine Systems*, 30, 1–20, [https://doi.org/10.1016/S0924-7963\(01\)00028-8](https://doi.org/10.1016/S0924-7963(01)00028-8), 2001.
- Andreae, M. O., Afchine, A., Albrecht, R., Holanda, B. A., Artaxo, P., Barbosa, H. M. J., Borrmann, S., Cecchini, M. A., Costa, A., Dollner, M., Fütterer, D., Järvinen, E., Jurkat, T., Klimach, T., Konemann, T., Knote, C., Krämer, M., Krisna, T., Machado, L. A. T.,
1220 Mertes, S., Minikin, A., Pöhlker, C., Pöhlker, M. L., Pöschl, U., Rosenfeld, D., Sauer, D., Schlager, H., Schnaiter, M., Schneider, J., Schulz, C., Spanu, A., Sperling, V. B., Voigt, C., Walser, A., Wang, J., Weinzierl, B., Wendisch, M., and Ziereis, H.: Aerosol Characteristics and Particle Production in the Upper Troposphere over the Amazon Basin, *Atmospheric Chemistry and Physics*, 18, 921–961, <https://doi.org/10.5194/acp-18-921-2018>, 2018.
- Archer-Nicholls, S., Abraham, N. L., Shin, Y. M., Weber, J., Russo, M. R., Lowe, D., Utembe, S. R., O’Connor, F. M., Kerridge, B.,
1225 Latter, B., Siddans, R., Jenkin, M., Wild, O., and Archibald, A. T.: The Common Representative Intermediates Mechanism Version 2 in the United Kingdom Chemistry and Aerosols Model, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002420, <https://doi.org/10.1029/2020MS002420>, 2021.
- Archibald, A. T., O’Connor, F. M., Abraham, N. L., Archer-Nicholls, S., Chipperfield, M. P., Dalvi, M., Folberth, G. A., Dennison, F., Dhomse, S. S., Griffiths, P. T., Hardacre, C., Hewitt, A. J., Hill, R. S., Johnson, C. E., Keeble, J., Köhler, M. O., Morgenstern, O.,
1230 Mulcahy, J. P., Ordóñez, C., Pope, R. J., Rumbold, S. T., Russo, M. R., Savage, N. H., Sellar, A., Stringer, M., Turnock, S. T., Wild, O., and Zeng, G.: Description and Evaluation of the UKCA Stratosphere–Troposphere Chemistry Scheme (StratTrop vn 1.0) Implemented in UKESM1, *Geoscientific Model Development*, 13, 1223–1266, <https://doi.org/10.5194/gmd-13-1223-2020>, 2020.
- Baccarini, A., Karlsson, L., Dommen, J., Duplissy, P., Vüllers, J., Brooks, I. M., Saiz-Lopez, A., Salter, M., Tjernström, M., Baltensperger, U., Zieger, P., and Schmale, J.: Frequent New Particle Formation over the High Arctic Pack Ice by Enhanced Iodine Emissions, *Nature
1235 Communications*, 11, 4924, <https://doi.org/10.1038/s41467-020-18551-0>, 2020.
- Barnes, I., Hjorth, J., and Mihalopoulos, N.: Dimethyl Sulfide and Dimethyl Sulfoxide and Their Oxidation in the Atmosphere, *Chemical Reviews*, 106, 940–975, <https://doi.org/10.1021/cr020529+>, 2006.
- Beck, L. J., Sarnela, N., Junninen, H., Hoppe, C. J. M., Garmash, O., Bianchi, F., Riva, M., Rose, C., Peräkylä, O., Wimmer, D., Kausiala, O., Jokinen, T., Ahonen, L., Mikkilä, J., Hakala, J., He, X.-C., Kontkanen, J., Wolf, K. K. E., Cappelletti, D., Mazzola, M., Traversi,
1240 R., Petroselli, C., Viola, A. P., Vitale, V., Lange, R., Massling, A., Nøjgaard, J. K., Krejci, R., Karlsson, L., Zieger, P., Jang, S., Lee, K., Vakkari, V., Lampilahti, J., Thakur, R. C., Leino, K., Kangasluoma, J., Duplissy, E.-M., Siivola, E., Marbouti, M., Tham, Y. J., Saiz-Lopez,

- A., Petäjä, T., Ehn, M., Worsnop, D. R., Skov, H., Kulmala, M., Kerminen, V.-M., and Sipilä, M.: Differing Mechanisms of New Particle Formation at Two Arctic Sites, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL091334>, 2021.
- 1245 Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mühlenthal, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Reviews of Geophysics*, 58, <https://doi.org/10.1029/2019RG000660>, 2020.
- 1250 Bhatti, Y. A., Revell, L. E., Schuddeboom, A. J., McDonald, A. J., Archibald, A. T., Williams, J., Venugopal, A. U., Hardacre, C., and Behrens, E.: The Sensitivity of Southern Ocean Atmospheric Dimethyl Sulfide (DMS) to Modeled Oceanic DMS Concentrations and Emissions, *Atmospheric Chemistry and Physics*, 23, 15 181–15 196, <https://doi.org/10.5194/acp-23-15181-2023>, 2023.
- Bian, H., Chin, M., Colarco, P. R., Apel, E. C., Blake, D. R., Froyd, K., Hornbrook, R. S., Jimenez, J., Jost, P. C., Lawler, M., Liu, M., Lund, M. T., Matsui, H., Nault, B. A., Penner, J. E., Rollins, A. W., Schill, G., Skeie, R. B., Wang, H., Xu, L., Zhang, K., and Zhu, J.: Observationally Constrained Analysis of Sulfur Cycle in the Marine Atmosphere with NASA ATom Measurements and AeroCom Model Simulations, *Atmospheric Chemistry and Physics*, 24, 1717–1741, <https://doi.org/10.5194/acp-24-1717-2024>, 2024.
- 1255 Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van Der Hoek, K. W., and Olivier, J. G. J.: A Global High-resolution Emission Inventory for Ammonia, *Global Biogeochemical Cycles*, 11, 561–587, <https://doi.org/10.1029/97GB02266>, 1997.
- Brean, J., Dall’Osto, M., Simó, R., Shi, Z., Beddows, D. C. S., and Harrison, R. M.: Open Ocean and Coastal New Particle Formation from Sulfuric Acid and Amines around the Antarctic Peninsula, *Nature Geoscience*, 14, 383–388, <https://doi.org/10.1038/s41561-021-00751-y>, 1260 2021.
- Brock, C., Froyd, K., Dollner, M., Williamson, C., Schill, G., Murphy, D., Wagner, N., Kupc, A., Jimenez, J., Campuzano-Jost, P., Nault, B., Schroder, J., Day, D., Price, D., Weinzierl, B., Schwarz, J., Katich, J., Wang, S., Zeng, L., Webb, R., Dibb, J., Scheuer, E., Diskin, G., DiGangi, J., Bui, T., Dean-Day, J., Thompson, C., Peischl, J., Ryerson, T., Bourgeois, I., Daube, B., Commane, R., and Wofsy, S.: ATom: Comprehensive Aerosol Properties, 2016–2018, Version 2, <https://doi.org/10.3334/ORNLDAAC/2111>, 2022.
- 1265 Brock, C. A., Williamson, C., Kupc, A., Froyd, K. D., Erdesz, F., Wagner, N., Richardson, M., Schwarz, J. P., Gao, R.-S., Katich, J. M., Campuzano-Jost, P., Nault, B. A., Schroder, J. C., Jimenez, J. L., Weinzierl, B., Dollner, M., Bui, T., and Murphy, D. M.: Aerosol Size Distributions during the Atmospheric Tomography Mission (ATom): Methods, Uncertainties, and Data Products, *Atmospheric Measurement Techniques*, 12, 3081–3099, <https://doi.org/10.5194/amt-12-3081-2019>, 2019.
- 1270 Brock, C. A., Froyd, K. D., Dollner, M., Williamson, C. J., Schill, G., Murphy, D. M., Wagner, N. J., Kupc, A., Jimenez, J. L., Campuzano-Jost, P., Nault, B. A., Schroder, J. C., Day, D. A., Price, D. J., Weinzierl, B., Schwarz, J. P., Katich, J. M., Wang, S., Zeng, L., Weber, R., Dibb, J., Scheuer, E., Diskin, G. S., DiGangi, J. P., Bui, T., Dean-Day, J. M., Thompson, C. R., Peischl, J., Ryerson, T. B., Bourgeois, I., Daube, B. C., Commane, R., and Wofsy, S. C.: Ambient Aerosol Properties in the Remote Atmosphere from Global-Scale in Situ Measurements, *Atmospheric Chemistry and Physics*, 21, 15 023–15 063, <https://doi.org/10.5194/acp-21-15023-2021>, 2021.
- 1275 Cala, B. A., Archer-Nicholls, S., Weber, J., Abraham, N. L., Griffiths, P. T., Jacob, L., Shin, Y. M., Revell, L. E., Woodhouse, M., and Archibald, A. T.: Development, Intercomparison, and Evaluation of an Improved Mechanism for the Oxidation of Dimethyl Sulfide in the UKCA Model, *Atmospheric Chemistry and Physics*, 23, 14 735–14 760, <https://doi.org/10.5194/acp-23-14735-2023>, 2023.
- Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., and Pierce, J. R.: Large Contribution of Natural Aerosols to Uncertainty in Indirect Forcing, *Nature*, 503, 67–71, <https://doi.org/10.1038/nature12674>, 2013.

- 1280 Carslaw, K. S., Gordon, H., Hamilton, D. S., Johnson, J. S., Regayre, L. A., Yoshioka, M., and Pringle, K. J.: Aerosols in the Pre-industrial Atmosphere, *Current Climate Change Reports*, 3, 1–15, <https://doi.org/10.1007/s40641-017-0061-2>, 2017.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley, J. A., Hansen, J. E., and Hofmann, D. J.: Climate Forcing by Anthropogenic Aerosols, *Science*, 255, 423–430, <https://doi.org/10.1126/science.255.5043.423>, 1992.
- Clarke, A. D., Freitag, S., Simpson, R. M. C., Hudson, J. G., Howell, S. G., Brekhovskikh, V. L., Campos, T., Kapustin, V. N., and Zhou, 1285 J.: Free Troposphere as a Major Source of CCN for the Equatorial Pacific Boundary Layer: Long-Range Transport and Teleconnections, *Atmospheric Chemistry and Physics*, 13, 7511–7529, <https://doi.org/10.5194/acp-13-7511-2013>, 2013.
- Curtius, J., Heinritzi, M., Beck, L. J., Pöhlker, M. L., Tripathi, N., Krumm, B. E., Holzbeck, P., Nussbaumer, C. M., Hernández Pardo, L., Klimach, T., Barmounis, K., Andersen, S. T., Bardakov, R., Bohn, B., Cecchini, M. A., Chaboureau, J.-P., Dauhut, T., Dienhart, D., Dörich, R., Edtbauer, A., Giez, A., Hartmann, A., Holanda, B. A., Joppe, P., Kaiser, K., Keber, T., Klebach, H., Krüger, O. O., Kürten, 1290 A., Mallaun, C., Marno, D., Martinez, M., Monteiro, C., Nelson, C., Ort, L., Raj, S. S., Richter, S., Ringsdorf, A., Rocha, F., Simon, M., Sree Kumar, S., Tsokankunku, A., Unfer, G. R., Valenti, I. D., Wang, N., Zahn, A., Zauner-Wieczorek, M., Albrecht, R. I., Andreae, M. O., Artaxo, P., Crowley, J. N., Fischer, H., Harder, H., Herdies, D. L., Machado, L. A. T., Pöhlker, C., Pöschl, U., Possner, A., Pozzer, A., Schneider, J., Williams, J., and Lelieveld, J.: Isoprene Nitrates Drive New Particle Formation in Amazon’s Upper Troposphere, *Nature*, 636, 124–130, <https://doi.org/10.1038/s41586-024-08192-4>, 2024.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., Van De Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., De Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim Reanalysis: Configuration and Performance of the Data Assimilation System, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, 1300 <https://doi.org/10.1002/qj.828>, 2011.
- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of Primary Aerosol and Precursor Gases in the Years 2000 and 1750 Prescribed Data-Sets for AeroCom, *Atmospheric Chemistry and Physics*, 6, 4321–4344, <https://doi.org/10.5194/acp-6-4321-2006>, 2006.
- 1305 Dunne, E. M., Gordon, H., Kürten, A., Almeida, J., Duplissy, J., Williamson, C., Ortega, I. K., Pringle, K. J., Adamov, A., Baltensperger, U., Barmet, P., Benduhn, F., Bianchi, F., Breitenlechner, M., Clarke, A., Curtius, J., Dommen, J., Donahue, N. M., Ehrhart, S., Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Jokinen, T., Kangasluoma, J., Kirkby, J., Kulmala, M., Kupc, A., Lawler, M. J., Lehtipalo, K., Makhmutov, V., Mann, G., Mathot, S., Merikanto, J., Miettinen, P., Nenes, A., Onnela, A., Rap, A., Reddington, C. L. S., Riccobono, F., Richards, N. A. D., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Sengupta, K., Simon, M., Sipilä, M., 1310 Smith, J. N., Stozkhov, Y., Tomé, A., Tröstl, J., Wagner, P. E., Wimmer, D., Winkler, P. M., Worsnop, D. R., and Carslaw, K. S.: Global Atmospheric Particle Formation from CERN CLOUD Measurements, *Science*, 354, 1119–1124, <https://doi.org/10.1126/science.aaf2649>, 2016.
- Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, 1315 L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A Large Source of Low-Volatility Secondary Organic Aerosol, *Nature*, 506, 476–479, <https://doi.org/10.1038/nature13032>, 2014.

- Froyd, K. D., Yu, P., Schill, G. P., Brock, C. A., Kupc, A., Williamson, C. J., Jensen, E. J., Ray, E., Rosenlof, K. H., Bian, H., Darmenov, A. S., Colarco, P. R., Diskin, G. S., Bui, T., and Murphy, D. M.: Dominant Role of Mineral Dust in Cirrus Cloud Formation Revealed by
1320 Global-Scale Measurements, *Nature Geoscience*, 15, 177–183, <https://doi.org/10.1038/s41561-022-00901-w>, 2022.
- Gantt, B., Johnson, M. S., Meskhidze, N., Sciare, J., Ovadnevaite, J., Ceburnis, D., and O’Dowd, C. D.: Model Evaluation of Marine Primary Organic Aerosol Emission Schemes, *Atmospheric Chemistry and Physics*, 12, 8553–8566, <https://doi.org/10.5194/acp-12-8553-2012>, 2012.
- Gao, C. Y., Heald, C. L., Katich, J. M., Luo, G., and Yu, F.: Remote Aerosol Simulated During the Atmospheric Tomography
1325 (ATom) Campaign and Implications for Aerosol Lifetime, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036524, <https://doi.org/10.1029/2022JD036524>, 2022.
- Ge, Y., Heal, M. R., Stevenson, D. S., Wind, P., and Vieno, M.: Evaluation of Global EMEP MSC-W (Rv4.34) WRF (v3.9.1.1) Model Surface Concentrations and Wet Deposition of Reactive N and S with Measurements, *Geoscientific Model Development*, 14, 7021–7046, <https://doi.org/10.5194/gmd-14-7021-2021>, 2021.
- 1330 Giannakopoulos, C., Chipperfield, M. P., Law, K. S., and Pyle, J. A.: Validation and Intercomparison of Wet and Dry Deposition Schemes Using²¹⁰ Pb in a Global Three-dimensional Off-line Chemical Transport Model, *Journal of Geophysical Research: Atmospheres*, 104, 23 761–23 784, <https://doi.org/10.1029/1999JD900392>, 1999.
- Gong, S. L.: A Parameterization of Sea-salt Aerosol Source Function for Sub- and Super-micron Particles, *Global Biogeochemical Cycles*, 17, 2003GB002 079, <https://doi.org/10.1029/2003GB002079>, 2003.
- 1335 Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A., Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozkhov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R., Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and Importance of New Particle Formation in the Present-Day and Preindustrial Atmospheres, *Journal of Geophysical Research: Atmospheres*, 122, 8739–8760, <https://doi.org/10.1002/2017JD026844>, 2017.
- 1340 Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L. K., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature Version 2.1 (MEGAN2.1): An Extended and Updated Framework for Modeling Biogenic Emissions, *Geoscientific Model Development*, 5, 1471–1492, <https://doi.org/10.5194/gmd-5-1471-2012>, 2012.
- Guo, H., Campuzano-Jost, P., Nault, B. A., Day, D. A., Schroder, J. C., Kim, D., Dibb, J. E., Dollner, M., Weinzierl, B., and Jimenez, J. L.: The Importance of Size Ranges in Aerosol Instrument Intercomparisons: A Case Study for the Atmospheric Tomography Mission,
1345 *Atmospheric Measurement Techniques*, 14, 3631–3655, <https://doi.org/10.5194/amt-14-3631-2021>, 2021.
- Hackenberg, S. C., Andrews, S. J., Airs, R. L., Arnold, S. R., Bouman, H. A., Cummings, D., Lewis, A. C., Minaeian, J. K., Reifel, K. M., Small, A., Tarran, G. A., Tilstone, G. H., and Carpenter, L. J.: Basin-Scale Observations of Monoterpenes in the Arctic and Atlantic Oceans, *Environmental Science & Technology*, 51, 10 449–10 458, <https://doi.org/10.1021/acs.est.7b02240>, 2017.
- Hardacre, C., Mulcahy, J. P., Pope, R. J., Jones, C. G., Rumbold, S. T., Li, C., Johnson, C., and Turnock, S. T.: Evaluation of SO₂, SO₄²⁻ and an Updated SO₂ Dry Deposition Parameterization in the United Kingdom Earth System Model, *Atmospheric Chemistry and Physics*,
1350 21, 18 465–18 497, <https://doi.org/10.5194/acp-21-18465-2021>, 2021.
- He, X.-C., Iyer, S., Sipilä, M., Ylisirniö, A., Peltola, M., Kontkanen, J., Baalbaki, R., Simon, M., Kürten, A., Tham, Y. J., Pesonen, J., Ahonen, L. R., Amanatidis, S., Amorim, A., Baccarini, A., Beck, L., Bianchi, F., Brilke, S., Chen, D., Chiu, R., Curtius, J., Dada, L., Dias, A., Dommen, J., Donahue, N. M., Duplissy, J., El Haddad, I., Finkenzeller, H., Fischer, L., Heinritzi, M., Hofbauer, V., Kangasluoma, J.,
1355 Kim, C., Koenig, T. K., Kubečka, J., Kvashnin, A., Lamkaddam, H., Lee, C. P., Leiminger, M., Li, Z., Makhmutov, V., Xiao, M., Marten,

- R., Nie, W., Onnela, A., Partoll, E., Petäjä, T., Salo, V.-T., Schuchmann, S., Steiner, G., Stolzenburg, D., Stozhkov, Y., Tauber, C., Tomé, A., Väisänen, O., Vazquez-Pufleau, M., Volkamer, R., Wagner, A. C., Wang, M., Wang, Y., Wimmer, D., Winkler, P. M., Worsnop, D. R., Wu, Y., Yan, C., Ye, Q., Lehtinen, K., Nieminen, T., Manninen, H. E., Rissanen, M., Schobesberger, S., Lehtipalo, K., Baltensperger, U., Hansel, A., Kerminen, V.-M., Flagan, R. C., Kirkby, J., Kurtén, T., and Kulmala, M.: Determination of the Collision Rate Coefficient between Charged Iodic Acid Clusters and Iodic Acid Using the Appearance Time Method, *Aerosol Science and Technology*, 55, 231–242, <https://doi.org/10.1080/02786826.2020.1839013>, 2021a.
- 1360 He, X.-C., Tham, Y. J., Dada, L., Wang, M., Finkenzeller, H., Stolzenburg, D., Iyer, S., Simon, M., Kürten, A., Shen, J., Rörup, B., Rissanen, M., Schobesberger, S., Baalbaki, R., Wang, D. S., Koenig, T. K., Jokinen, T., Sarnela, N., Beck, L. J., Almeida, J., Amanatidis, S., Amorim, A., Ataei, F., Baccarini, A., Bertozzi, B., Bianchi, F., Brilke, S., Caudillo, L., Chen, D., Chiu, R., Chu, B., Dias, A., Ding, A., Dommen, J., Duplissy, J., El Haddad, I., Gonzalez Carracedo, L., Granzin, M., Hansel, A., Heinritzi, M., Hofbauer, V., Junninen, H., Kangasluoma, J., Kemppainen, D., Kim, C., Kong, W., Krechmer, J. E., Kvashin, A., Laitinen, T., Lamkaddam, H., Lee, C. P., Lehtipalo, K., Leiminger, M., Li, Z., Makhmutov, V., Manninen, H. E., Marie, G., Marten, R., Mathot, S., Mauldin, R. L., Mentler, B., Möhler, O., Müller, T., Nie, W., Onnela, A., Petäjä, T., Pfeifer, J., Philippov, M., Ranjithkumar, A., Saiz-Lopez, A., Salma, I., Scholz, W., Schuchmann, S., Schulze, B., Steiner, G., Stozhkov, Y., Tauber, C., Tomé, A., Thakur, R. C., Väisänen, O., Vazquez-Pufleau, M., Wagner, A. C., Wang, Y., Weber, S. K., Winkler, P. M., Wu, Y., Xiao, M., Yan, C., Ye, Q., Ylisirniö, A., Zauner-Wieczorek, M., Zha, Q., Zhou, P., Flagan, R. C., Curtius, J., Baltensperger, U., Kulmala, M., Kerminen, V.-M., Kurtén, T., Donahue, N. M., Volkamer, R., Kirkby, J., Worsnop, D. R., and Sipilä, M.: Role of Iodine Oxoacids in Atmospheric Aerosol Nucleation, *Science*, 371, 589–595, <https://doi.org/10.1126/science.abe0298>, 2021b.
- 1365 He, X.-C., Simon, M., Iyer, S., Xie, H.-B., Rörup, B., Shen, J., Finkenzeller, H., Stolzenburg, D., Zhang, R., Baccarini, A., Tham, Y. J., Wang, M., Amanatidis, S., Piedehierro, A. A., Amorim, A., Baalbaki, R., Brasseur, Z., Caudillo, L., Chu, B., Dada, L., Duplissy, J., El Haddad, I., Flagan, R. C., Granzin, M., Hansel, A., Heinritzi, M., Hofbauer, V., Jokinen, T., Kemppainen, D., Kong, W., Krechmer, J., Kürten, A., Lamkaddam, H., Lopez, B., Ma, F., Mahfouz, N. G. A., Makhmutov, V., Manninen, H. E., Marie, G., Marten, R., Massabò, D., Mauldin, R. L., Mentler, B., Onnela, A., Petäjä, T., Pfeifer, J., Philippov, M., Ranjithkumar, A., Rissanen, M. P., Schobesberger, S., Scholz, W., Schulze, B., Surdu, M., Thakur, R. C., Tomé, A., Wagner, A. C., Wang, D., Wang, Y., Weber, S. K., Welti, A., Winkler, P. M., Zauner-Wieczorek, M., Baltensperger, U., Curtius, J., Kurtén, T., Worsnop, D. R., Volkamer, R., Lehtipalo, K., Kirkby, J., Donahue, N. M., Sipilä, M., and Kulmala, M.: Iodine Oxoacids Enhance Nucleation of Sulfuric Acid Particles in the Atmosphere, *Science*, 382, 1308–1314, <https://doi.org/10.1126/science.adh2526>, 2023.
- 1370 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 1385 Hodshire, A. L., Campuzano-Jost, P., Kodros, J. K., Croft, B., Nault, B. A., Schroder, J. C., Jimenez, J. L., and Pierce, J. R.: The Potential Role of Methanesulfonic Acid (MSA) in Aerosol Formation and Growth and the Associated Radiative Forcings, *Atmospheric Chemistry and Physics*, 19, 3137–3160, <https://doi.org/10.5194/acp-19-3137-2019>, 2019.
- 1390 Hodzic, A., Campuzano-Jost, P., Bian, H., Chin, M., Colarco, P. R., Day, D. A., Froyd, K. D., Heinold, B., Jo, D. S., Katich, J. M., Kodros, J. K., Nault, B. A., Pierce, J. R., Ray, E., Schacht, J., Schill, G. P., Schroder, J. C., Schwarz, J. P., Sueper, D. T., Tegen, I., Tilmes, S., Tsigaridis, K., Yu, P., and Jimenez, J. L.: Characterization of Organic Aerosol across the Global Remote Troposphere: A Comparison of

- ATom Measurements and Global Chemistry Models, *Atmospheric Chemistry and Physics*, 20, 4607–4635, <https://doi.org/10.5194/acp-20-4607-2020>, 2020.
- 1395 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-i., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) Anthropogenic Emissions of Reactive Gases and Aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- 1400 Hoffmann, T., O'Dowd, C. D., and Seinfeld, J. H.: Iodine Oxide Homogeneous Nucleation: An Explanation for Coastal New Particle Production, *Geophysical Research Letters*, 28, 1949–1952, <https://doi.org/10.1029/2000GL012399>, 2001.
- Jacob, L. S. D., Giorio, C., and Archibald, A. T.: Extension, Development, and Evaluation of the Representation of the OH-initiated Dimethyl Sulfide (DMS) Oxidation Mechanism in the Master Chemical Mechanism (MCM) v3.3.1 Framework, *Atmospheric Chemistry and Physics*, 24, 3329–3347, <https://doi.org/10.5194/acp-24-3329-2024>, 2024.
- 1405 Jen, C. N., McMurry, P. H., and Hanson, D. R.: Stabilization of Sulfuric Acid Dimers by Ammonia, Methylamine, Dimethylamine, and Trimethylamine, *Journal of Geophysical Research: Atmospheres*, 119, 7502–7514, <https://doi.org/10.1002/2014JD021592>, 2014.
- Jenkin, M., Khan, M., Shallcross, D., Bergström, R., Simpson, D., Murphy, K., and Rickard, A.: The CRI v2.2 Reduced Degradation Scheme for Isoprene, *Atmospheric Environment*, 212, 172–182, <https://doi.org/10.1016/j.atmosenv.2019.05.055>, 2019.
- Jokinen, T., Sipilä, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E.-M., Junninen, H., Kangasluoma, J., Manninen, H. E., Petäjä, T., 1410 Kulmala, M., Worsnop, D. R., Kirkby, J., Virkkula, A., and Kerminen, V.-M.: Ion-Induced Sulfuric Acid–Ammonia Nucleation Drives Particle Formation in Coastal Antarctica, *Science Advances*, 4, eaat9744, <https://doi.org/10.1126/sciadv.aat9744>, 2018.
- Jones, A. C., Hill, A., Remy, S., Abraham, N. L., Dalvi, M., Hardacre, C., Hewitt, A. J., Johnson, B., Mulcahy, J. P., and Turnock, S. T.: Exploring the Sensitivity of Atmospheric Nitrate Concentrations to Nitric Acid Uptake Rate Using the Met Office's Unified Model, *Atmospheric Chemistry and Physics*, 21, 15901–15927, <https://doi.org/10.5194/acp-21-15901-2021>, 2021.
- 1415 Kanaya, Y., Sommariva, R., Saiz-Lopez, A., Mazzeo, A., Koenig, T. K., Kawana, K., Johnson, J. E., Colomb, A., Tulet, P., Molloy, S., Galbally, I. E., Volkamer, R., Mahajan, A., Halfacre, J. W., Shepson, P. B., Schmale, J., Angot, H., Blomquist, B., Shupe, M. D., Helmig, D., Gil, J., Lee, M., Coburn, S. C., Ortega, I., Chen, G., Lee, J., Aikin, K. C., Parrish, D. D., Holloway, J. S., Ryerson, T. B., Pollack, I. B., Williams, E. J., Lerner, B. M., Weinheimer, A. J., Campos, T., Flocke, F. M., Spackman, J. R., Bourgeois, I., Peischl, J., Thompson, C. R., Staebler, R. M., Aliabadi, A. A., Gong, W., Van Malderen, R., Thompson, A. M., Stauffer, R. M., Kollonige, D. E., Gómez Martin, J. C., 1420 Fujiwara, M., Read, K., Rowlinson, M., Sato, K., Kurokawa, J., Iwamoto, Y., Taketani, F., Takashima, H., Navarro Comas, M., Panagi, M., and Schultz, M. G.: Observational Ozone Data over the Global Oceans and Polar Regions: The TOAR-II Oceans Data Set Version 2024, <https://doi.org/10.5194/essd-2024-566>, 2025.
- Kim, D., Campuzano-Jost, P., Guo, H., Day, D. A., Yang, D., Dhaniyala, S., Williams, L., Croteau, P., Jayne, J., Worsnop, D., Volkamer, R., and Jimenez, J. L.: Development and Characterization of an Aircraft Inlet System for Broader Quantitative Particle Sampling at 1425 Higher Altitudes: Aerodynamic Lenses, Beam and Vaporizer Diagnostics, and Pressure-Controlled Inlets, *Aerosol Research*, 3, 371–404, <https://doi.org/10.5194/ar-3-371-2025>, 2025.
- Kipling, Z., Stier, P., Schwarz, J. P., Perring, A. E., Spackman, J. R., Mann, G. W., Johnson, C. E., and Telford, P. J.: Constraints on Aerosol Processes in Climate Models from Vertically-Resolved Aircraft Observations of Black Carbon, *Atmospheric Chemistry and Physics*, 13, 5969–5986, <https://doi.org/10.5194/acp-13-5969-2013>, 2013.
- 1430 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagné, S., Ickes, L., Kürten, A., Kupc, A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F., Breitenlechner, M., David,

- A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W., Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S., Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petäjä, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä, M., Stozhkov, Y., Stratmann, F., Tomé, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner, E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of Sulphuric Acid, Ammonia and Galactic Cosmic Rays in Atmospheric Aerosol Nucleation, *Nature*, 476, 429–433, <https://doi.org/10.1038/nature10343>, 2011.
- 1435 Kirkby, J., Duplissy, J., Sengupta, K., Frege, C., Gordon, H., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J., Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J., Dias, A., Ehrhart, S., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K., Rap, A., Richards, N. A. D., Riipinen, I., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger, S., Scott, C. E., Seinfeld, J. H., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Carslaw, K. S., and Curtius, J.: Ion-Induced Nucleation of Pure Biogenic Particles, *Nature*, 533, 521–526, <https://doi.org/10.1038/nature17953>, 2016.
- 1440 Kirkby, J., Amorim, A., Baltensperger, U., Carslaw, K. S., Christoudias, T., Curtius, J., Donahue, N. M., Haddad, I. E., Flagan, R. C., Gordon, H., Hansel, A., Harder, H., Junninen, H., Kulmala, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Lelieveld, J., Möhler, O., Riipinen, I., Stratmann, F., Tomé, A., Virtanen, A., Volkamer, R., Winkler, P. M., and Worsnop, D. R.: Atmospheric New Particle Formation from the CERN CLOUD Experiment, *Nature Geoscience*, 16, 948–957, <https://doi.org/10.1038/s41561-023-01305-0>, 2023.
- 1450 Koenig, T. K., Baidar, S., Campuzano-Jost, P., Cuevas, C. A., Dix, B., Fernandez, R. P., Guo, H., Hall, S. R., Kinnison, D., Nault, B. A., Ullmann, K., Jimenez, J. L., Saiz-Lopez, A., and Volkamer, R.: Quantitative Detection of Iodine in the Stratosphere, *Proceedings of the National Academy of Sciences*, 117, 1860–1866, <https://doi.org/10.1073/pnas.1916828117>, 2020.
- Koren, I., Dagan, G., and Altaratz, O.: From Aerosol-Limited to Invigoration of Warm Convective Clouds, *Science*, 344, 1143–1146, <https://doi.org/10.1126/science.1252595>, 2014.
- 1455 Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol Formation from Isoprene Photooxidation under high-NO_x Conditions, *Geophysical Research Letters*, 32, 2005GL023 637, <https://doi.org/10.1029/2005GL023637>, 2005.
- Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol Formation from Isoprene Photooxidation, *Environmental Science & Technology*, 40, 1869–1877, <https://doi.org/10.1021/es0524301>, 2006.
- 1460 Kulmala, M., Laaksonen, A., and Pirjola, L.: Parameterizations for Sulfuric Acid/Water Nucleation Rates, *Journal of Geophysical Research: Atmospheres*, 103, 8301–8307, <https://doi.org/10.1029/97JD03718>, 1998.
- Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Cluster Activation Theory as an Explanation of the Linear Dependence between Formation Rate of 3nm Particles and Sulphuric Acid Concentration, *Atmospheric Chemistry and Physics*, 6, 787–793, <https://doi.org/10.5194/acp-6-787-2006>, 2006.
- 1465 Kupc, A., Williamson, C., Wagner, N. L., Richardson, M., and Brock, C. A.: Modification, Calibration, and Performance of the Ultra-High Sensitivity Aerosol Spectrometer for Particle Size Distribution and Volatility Measurements during the Atmospheric Tomography Mission (ATom) Airborne Campaign, *Atmospheric Measurement Techniques*, 11, 369–383, <https://doi.org/10.5194/amt-11-369-2018>, 2018.
- Kupc, A., Williamson, C. J., Hodshire, A. L., Kazil, J., Ray, E., Bui, T. P., Dollner, M., Froyd, K. D., McKain, K., Rollins, A., Schill, G. P., Thames, A., Weinzierl, B. B., Pierce, J. R., and Brock, C. A.: The Potential Role of Organics in New Particle Formation and Initial

- 1470 Growth in the Remote Tropical Upper Troposphere, *Atmospheric Chemistry and Physics*, 20, 15 037–15 060, <https://doi.org/10.5194/acp-20-15037-2020>, 2020.
- Lana, A., Bell, T. G., Simó, R., Vallina, S. M., Ballabrera-Poy, J., Kettle, A. J., Dachs, J., Bopp, L., Saltzman, E. S., Stefels, J., Johnson, J. E., and Liss, P. S.: An Updated Climatology of Surface Dimethylsulfide Concentrations and Emission Fluxes in the Global Ocean: UPDATED DMS CLIMATOLOGY, *Global Biogeochemical Cycles*, 25, n/a–n/a, <https://doi.org/10.1029/2010GB003850>, 2011.
- 1475 Lee, L. A., Reddington, C. L., and Carslaw, K. S.: On the Relationship between Aerosol Model Uncertainty and Radiative Forcing Uncertainty, *Proceedings of the National Academy of Sciences*, 113, 5820–5827, <https://doi.org/10.1073/pnas.1507050113>, 2016.
- Lee, S.-H., Reeves, J. M., Wilson, J. C., Hunton, D. E., Viggiano, A. A., Miller, T. M., Ballenthin, J. O., and Lait, L. R.: Particle Formation by Ion Nucleation in the Upper Troposphere and Lower Stratosphere, *Science*, 301, 1886–1889, <https://doi.org/10.1126/science.1087236>, 2003.
- 1480 Lehtipalo, K., Yan, C., Dada, L., Bianchi, F., Xiao, M., Wagner, R., Stolzenburg, D., Ahonen, L. R., Amorim, A., Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S. B., Chen, D., Chen, X., Dias, A., Dommen, J., Draper, D. C., Duplissy, J., Ehn, M., Finkenzeller, H., Fischer, L., Frege, C., Fuchs, C., Garmash, O., Gordon, H., Hakala, J., He, X., Heikkinen, L., Heinritzi, M., Helm, J. C., Hofbauer, V., Hoyle, C. R., Jokinen, T., Kangasluoma, J., Kerminen, V.-M., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lawler, M. J., Mai, H., Mathot, S., Mauldin, R. L., Molteni, U., Nichman, L., 1485 Nie, W., Nieminen, T., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Piel, F., Pospisilova, V., Quéléver, L. L. J., Rissanen, M. P., Rose, C., Sarnela, N., Schallhart, S., Schuchmann, S., Sengupta, K., Simon, M., Sipilä, M., Tauber, C., Tomé, A., Tröstl, J., Väisänen, O., Vogel, A. L., Volkamer, R., Wagner, A. C., Wang, M., Weitz, L., Wimmer, D., Ye, P., Ylisirniö, A., Zha, Q., Carslaw, K. S., Curtius, J., Donahue, N. M., Flagan, R. C., Hansel, A., Riipinen, I., Virtanen, A., Winkler, P. M., Baltensperger, U., Kulmala, M., and Worsnop, D. R.: Multicomponent New Particle Formation from Sulfuric Acid, Ammonia, and Biogenic Vapors, *Science Advances*, 4, eaau5363, 1490 <https://doi.org/10.1126/sciadv.aau5363>, 2018.
- Luo, G. and Yu, F.: A Numerical Evaluation of Global Oceanic Emissions of α -Pinene and Isoprene, *Atmospheric Chemistry and Physics*, 10, 2007–2015, <https://doi.org/10.5194/acp-10-2007-2010>, 2010.
- Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and Evaluation of GLOMAP-mode: A Modal Global Aerosol Microphysics Model for the UKCA Composition-Climate 1495 Model, *Geoscientific Model Development*, 3, 519–551, <https://doi.org/10.5194/gmd-3-519-2010>, 2010.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.: IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press., United Kingdom, 2021.
- 1500 Mauldin, R. L., Tanner, D. J., Heath, J. A., Huebert, B. J., and Eisele, F. L.: Observations of H₂ SO₄ and MSA during PEM-Tropics-A, *Journal of Geophysical Research: Atmospheres*, 104, 5801–5816, <https://doi.org/10.1029/98JD02612>, 1999.
- Metzger, A., Verheggen, B., Dommen, J., Duplissy, J., Prevot, A. S. H., Weingartner, E., Riipinen, I., Kulmala, M., Spracklen, D. V., Carslaw, K. S., and Baltensperger, U.: Evidence for the Role of Organics in Aerosol Particle Formation under Atmospheric Conditions, *Proceedings of the National Academy of Sciences*, 107, 6646–6651, <https://doi.org/10.1073/pnas.0911330107>, 2010.
- 1505 Mulcahy, J. P., Johnson, C., Jones, C. G., Povey, A. C., Scott, C. E., Sellar, A., Turnock, S. T., Woodhouse, M. T., Abraham, N. L., Andrews, M. B., Bellouin, N., Browse, J., Carslaw, K. S., Dalvi, M., Folberth, G. A., Glover, M., Grosvenor, D. P., Hardacre, C., Hill, R., Johnson, B., Jones, A., Kipling, Z., Mann, G., Mollard, J., O’Connor, F. M., Palmiéri, J., Reddington, C., Rumbold, S. T., Richardson, M., Schutgens,

- N. A. J., Stier, P., Stringer, M., Tang, Y., Walton, J., Woodward, S., and Yool, A.: Description and Evaluation of Aerosol in UKESM1 and HadGEM3-GC3.1 CMIP6 Historical Simulations, *Geoscientific Model Development*, 13, 6383–6423, <https://doi.org/10.5194/gmd-13-6383-2020>, 2020.
- 1510 Mulcahy, J. P., Jones, C. G., Rumbold, S. T., Kuhlbrodt, T., Dittus, A. J., Blockley, E. W., Yool, A., Walton, J., Hardacre, C., Andrews, T., Bodas-Salcedo, A., Stringer, M., de Mora, L., Harris, P., Hill, R., Kelley, D., Robertson, E., and Tang, Y.: UKESM1.1: Development and Evaluation of an Updated Configuration of the UK Earth System Model, *Geoscientific Model Development*, 16, 1569–1600, <https://doi.org/10.5194/gmd-16-1569-2023>, 2023.
- 1515 Murphy, D. M., Froyd, K. D., Bian, H., Brock, C. A., Dibb, J. E., DiGangi, J. P., Diskin, G., Dollner, M., Kupc, A., Scheuer, E. M., Schill, G. P., Weinzierl, B., Williamson, C. J., and Yu, P.: The Distribution of Sea-Salt Aerosol in the Global Troposphere, *Atmospheric Chemistry and Physics*, 19, 4093–4104, <https://doi.org/10.5194/acp-19-4093-2019>, 2019.
- Nault, B. A., Campuzano-Jost, P., Day, D. A., Jo, D. S., Schroder, J. C., Allen, H. M., Bahreini, R., Bian, H., Blake, D. R., Chin, M., Clegg, S. L., Colarco, P. R., Crouse, J. D., Cubison, M. J., DeCarlo, P. F., Dibb, J. E., Diskin, G. S., Hodzic, A., Hu, W., Katich, J. M., Kim, M. J., Kodros, J. K., Kupc, A., Lopez-Hilfiker, F. D., Marais, E. A., Middlebrook, A. M., Andrew Neuman, J., Nowak, J. B., Palm, B. B., Paulot, F., Pierce, J. R., Schill, G. P., Scheuer, E., Thornton, J. A., Tsigaridis, K., Wennberg, P. O., Williamson, C. J., and Jimenez, J. L.: Chemical Transport Models Often Underestimate Inorganic Aerosol Acidity in Remote Regions of the Atmosphere, *Communications Earth & Environment*, 2, 93, <https://doi.org/10.1038/s43247-021-00164-0>, 2021.
- 1520 O'Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., Dalvi, M., Folberth, G. A., Sanderson, M. G., Telford, P. J., Voulgarakis, A., Young, P. J., Zeng, G., Collins, W. J., and Pyle, J. A.: Evaluation of the New UKCA Climate-Composition Model – Part 2: The Troposphere, *Geoscientific Model Development*, 7, 41–91, <https://doi.org/10.5194/gmd-7-41-2014>, 2014.
- O'Dowd, C. D., Jimenez, J. L., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hämeri, K., Pirjola, L., Kulmala, M., Jennings, S. G., and Hoffmann, T.: Marine Aerosol Formation from Biogenic Iodine Emissions, *Nature*, 417, 632–636, <https://doi.org/10.1038/nature00775>, 2002.
- 1530 O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and Putaud, J.-P.: Biogenically Driven Organic Contribution to Marine Aerosol, *Nature*, 431, 676–680, <https://doi.org/10.1038/nature02959>, 2004.
- Palmer, P. I., Marvin, M. R., Siddans, R., Kerridge, B. J., and Moore, D. P.: Nocturnal Survival of Isoprene Linked to Formation of Upper Tropospheric Organic Aerosol, *Science*, 375, 562–566, <https://doi.org/10.1126/science.abg4506>, 2022.
- Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kürten, A., St. Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, *Science*, 325, 730–733, <https://doi.org/10.1126/science.1172910>, 2009.
- 1535 Pirjola, L. and Kulmala, M.: Modelling the Formation of H₂SO₄–H₂O Particles in Rural, Urban and Marine Conditions, *Atmospheric Research*, 46, 321–347, [https://doi.org/10.1016/S0169-8095\(97\)00072-0](https://doi.org/10.1016/S0169-8095(97)00072-0), 1998.
- Pollack, I. B., Lerner, B. M., and Ryerson, T. B.: Evaluation of Ultraviolet Light-Emitting Diodes for Detection of Atmospheric NO₂ by Photolysis - Chemiluminescence, *Journal of Atmospheric Chemistry*, 65, 111–125, <https://doi.org/10.1007/s10874-011-9184-3>, 2010.
- 1540 Prather, K. A., Bertram, T. H., Grassian, V. H., Deane, G. B., Stokes, M. D., DeMott, P. J., Aluwihare, L. I., Palenik, B. P., Azam, F., Seinfeld, J. H., Moffet, R. C., Molina, M. J., Cappa, C. D., Geiger, F. M., Roberts, G. C., Russell, L. M., Ault, A. P., Baltrusaitis, J., Collins, D. B., Corrigan, C. E., Cuadra-Rodriguez, L. A., Ebben, C. J., Forestieri, S. D., Guasco, T. L., Hersey, S. P., Kim, M. J., Lambert, W. F., Modini, R. L., Mui, W., Pedler, B. E., Ruppel, M. J., Ryder, O. S., Schoepp, N. G., Sullivan, R. C., and Zhao, D.: Bringing the Ocean into the Laboratory to Probe the Chemical Complexity of Sea Spray Aerosol, *Proceedings of the National Academy of Sciences*, 110, 7550–7555, <https://doi.org/10.1073/pnas.1300262110>, 2013.
- 1545

- Price, R., Baccarini, A., Schmale, J., Zieger, P., Brooks, I. M., Field, P., and Carslaw, K. S.: Late Summer Transition from a Free-Tropospheric to Boundary Layer Source of Aitken Mode Aerosol in the High Arctic, *Atmospheric Chemistry and Physics*, 23, 2927–2961, <https://doi.org/10.5194/acp-23-2927-2023>, 2023.
- 1550 Ranjithkumar, A., Gordon, H., Williamson, C., Rollins, A., Pringle, K., Kupc, A., Abraham, N. L., Brock, C., and Carslaw, K.: Constraints on Global Aerosol Number Concentration, SO₂ and Condensation Sink in UKESM1 Using ATom Measurements, *Atmospheric Chemistry and Physics*, 21, 4979–5014, <https://doi.org/10.5194/acp-21-4979-2021>, 2021.
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.: Daily High-Resolution-Blended Analyses for Sea Surface Temperature, *Journal of Climate*, 20, 5473–5496, <https://doi.org/10.1175/2007JCLI1824.1>, 2007.
- 1555 Rinaldi, M., Decesari, S., Finessi, E., Giulianelli, L., Carbone, C., Fuzzi, S., O'Dowd, C. D., Ceburnis, D., and Facchini, M. C.: Primary and Secondary Organic Marine Aerosol and Oceanic Biological Activity: Recent Results and New Perspectives for Future Studies, *Advances in Meteorology*, 2010, 310 682, <https://doi.org/10.1155/2010/310682>, 2010.
- Rinaldi, M., Fuzzi, S., Decesari, S., Marullo, S., Santolero, R., Provenzale, A., Von Hardenberg, J., Ceburnis, D., Vaishya, A., O'Dowd, C. D., and Facchini, M. C.: Is Chlorophyll- *a* the Best Surrogate for Organic Matter Enrichment in Submicron Primary Marine Aerosol?, *Journal of Geophysical Research: Atmospheres*, 118, 4964–4973, <https://doi.org/10.1002/jgrd.50417>, 2013.
- 1560 Rollins, A. W., Thornberry, T. D., Ciciora, S. J., McLaughlin, R. J., Watts, L. A., Hanisco, T. F., Baumann, E., Giorgetta, F. R., Bui, T. V., Fahey, D. W., and Gao, R.-S.: A Laser-Induced Fluorescence Instrument for Aircraft Measurements of Sulfurdioxide in the Upper Troposphere and Lower Stratosphere, *Atmospheric Measurement Techniques*, 9, 4601–4613, <https://doi.org/10.5194/amt-9-4601-2016>, 2016.
- Russo, M. R., Bartholomew, S. L., Hassell, D., Mason, A. M., Neiningner, E., Perman, A. J., Sproson, D. A. J., Watson-Parris, D., and 1565 Abraham, N. L.: Virtual Integration of Satellite and In-situ Observation Networks (VISION) v1.0: In-Situ Observations Simulator (ISO_simulator), *Geoscientific Model Development*, 18, 181–191, <https://doi.org/10.5194/gmd-18-181-2025>, 2025.
- Schill, G. P., Froyd, K. D., Murphy, D. M., Williamson, C. J., Brock, C. A., Sherwen, T., Evans, M. J., Ray, E. A., Apel, E. C., Hornbrook, R. S., Hills, A. J., Peischl, J., Ryerson, T. B., Thompson, C. R., Bourgeois, I., Blake, D. R., DiGangi, J. P., and Diskin, G. S.: 1570 Widespread Trace Bromine and Iodine in Remote Tropospheric Non-Sea-Salt Aerosols, *Atmospheric Chemistry and Physics*, 25, 45–71, <https://doi.org/10.5194/acp-25-45-2025>, 2025.
- Scott, C. E., Rap, A., Spracklen, D. V., Forster, P. M., Carslaw, K. S., Mann, G. W., Pringle, K. J., Kivekäs, N., Kulmala, M., Lihavainen, H., and Tunved, P.: The Direct and Indirect Radiative Effects of Biogenic Secondary Organic Aerosol, *Atmospheric Chemistry and Physics*, 14, 447–470, <https://doi.org/10.5194/acp-14-447-2014>, 2014.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, 1575 R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving Our Fundamental Understanding of the Role of Aerosol-cloud Interactions in the Climate System, *Proceedings of the National Academy of Sciences*, 113, 5781–5790, <https://doi.org/10.1073/pnas.1514043113>, 2016.
- Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, 1580 S., Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.:

- UKESM1: Description and Evaluation of the U.K. Earth System Model, *Journal of Advances in Modeling Earth Systems*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.
- 1585 Shen, J., Scholz, W., He, X.-C., Zhou, P., Marie, G., Wang, M., Marten, R., Surdu, M., Rörup, B., Baalbaki, R., Amorim, A., Ataei, F., Bell, D. M., Bertozzi, B., Brasseur, Z., Caudillo, L., Chen, D., Chu, B., Dada, L., Duplissy, J., Finkenzeller, H., Granzin, M., Guida, R., Heinritzi, M., Hofbauer, V., Iyer, S., Kemppainen, D., Kong, W., Krechmer, J. E., Kürten, A., Lamkaddam, H., Lee, C. P., Lopez, B., Mahfouz, N. G. A., Manninen, H. E., Massabò, D., Mauldin, R. L., Mentler, B., Müller, T., Pfeifer, J., Philippov, M., Piedadhierro, A. A., Roldin, P., Schobesberger, S., Simon, M., Stolzenburg, D., Tham, Y. J., Tomé, A., Umo, N. S., Wang, D., Wang, Y., Weber, S. K., Welti, A., Wollesen de Jonge, R., Wu, Y., Zauner-Wieczorek, M., Zust, F., Baltensperger, U., Curtius, J., Flagan, R. C., Hansel, A., Möhler, O., Petäjä, T., Volkamer, R., Kulmala, M., Lehtipalo, K., Rissanen, M., Kirkby, J., El-Haddad, I., Bianchi, F., Sipilä, M., Donahue, N. M., and Worsnop, D. R.: High Gas-Phase Methanesulfonic Acid Production in the OH-Initiated Oxidation of Dimethyl Sulfide at Low Temperatures, *Environmental Science & Technology*, 56, 13 931–13 944, <https://doi.org/10.1021/acs.est.2c05154>, 2022.
- 1590 Shen, J., Russell, D. M., DeVivo, J., Kunkler, F., Baalbaki, R., Mentler, B., Scholz, W., Yu, W., Caudillo-Plath, L., Sommer, E., Ahongshangbam, E., Alfaouri, D., Almeida, J., Amorim, A., Beck, L. J., Beckmann, H., Berntheusel, M., Bhattacharyya, N., Canagaratna, M. R., Chassaing, A., Cruz-Simbron, R., Dada, L., Duplissy, J., Gordon, H., Granzin, M., Große Schute, L., Heinritzi, M., Iyer, S., Klebach, H., Krüger, T., Kürten, A., Lampimäki, M., Liu, L., Lopez, B., Martinez, M., Morawiec, A., Onnela, A., Peltola, M., Rato, P., Reza, M., Richter, S., Rörup, B., Sebastian, M. K., Simon, M., Surdu, M., Tamme, K., Thakur, R. C., Tomé, A., Tong, Y., Top, J., Umo, N. S., Unfer, G., Vettikkat, L., Weissbacher, J., Xenofontos, C., Yang, B., Zauner-Wieczorek, M., Zhang, J., Zheng, Z., Baltensperger, U., Christoudias, T., Flagan, R. C., El Haddad, I., Junninen, H., Möhler, O., Riipinen, I., Rohner, U., Schobesberger, S., Volkamer, R., Winkler, P. M., Hansel, A., Lehtipalo, K., Donahue, N. M., Lelieveld, J., Harder, H., Kulmala, M., Worsnop, D. R., Kirkby, J., Curtius, J., and He, X.-C.: New Particle Formation from Isoprene under Upper-Tropospheric Conditions, *Nature*, 636, 115–123, <https://doi.org/10.1038/s41586-024-08196-0>, 2024.
- 1600 Sipilä, M., Sarnela, N., Jokinen, T., Henschel, H., Junninen, H., Kontkanen, J., Richters, S., Kangasluoma, J., Franchin, A., Peräkylä, O., Rissanen, M. P., Ehn, M., Vehkamäki, H., Kurten, T., Berndt, T., Petäjä, T., Worsnop, D., Ceburnis, D., Kerminen, V.-M., Kulmala, M., and O'Dowd, C.: Molecular-Scale Evidence of Aerosol Particle Formation via Sequential Addition of HIO₃, *Nature*, 537, 532–534, <https://doi.org/10.1038/nature19314>, 2016.
- 1605 Spiro, P. A., Jacob, D. J., and Logan, J. A.: Global Inventory of Sulfur Emissions with 1° × 1° Resolution, *Journal of Geophysical Research: Atmospheres*, 97, 6023–6036, <https://doi.org/10.1029/91JD03139>, 1992.
- 1610 Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The Aerosol-Climate Model ECHAM5-HAM, *Atmospheric Chemistry and Physics*, 5, 1125–1156, <https://doi.org/10.5194/acp-5-1125-2005>, 2005.
- 1615 Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive Intermediates Revealed in Secondary Organic Aerosol Formation from Isoprene, *Proceedings of the National Academy of Sciences*, 107, 6640–6645, <https://doi.org/10.1073/pnas.091114107>, 2010.
- 1620 Svenhag, C., Sporre, M. K., Olenius, T., Yazgi, D., Blichner, S. M., Nieradzik, L. P., and Roldin, P.: Implementing Detailed Nucleation Predictions in the Earth System Model EC-Earth3.3.4: Sulfuric Acid–Ammonia Nucleation, *Geoscientific Model Development*, 17, 4923–4942, <https://doi.org/10.5194/gmd-17-4923-2024>, 2024.
- 1620 Telford, P. J., Braesicke, P., Morgenstern, O., and Pyle, J. A.: Technical Note: Description and Assessment of a Nudged Version of the New Dynamics Unified Model, *Atmospheric Chemistry and Physics*, 8, 1701–1712, <https://doi.org/10.5194/acp-8-1701-2008>, 2008.

- Thompson, C. R., Wofsy, S. C., Prather, M. J., Newman, P. A., Hanisco, T. F., Ryerson, T. B., Fahey, D. W., Apel, E. C., Brock, C. A., Brune, W. H., Froyd, K., Katich, J. M., Nicely, J. M., Peischl, J., Ray, E., Veres, P. R., Wang, S., Allen, H. M., Asher, E., Bian, H., Blake, D., Bourgeois, I., Budney, J., Bui, T. P., Butler, A., Campuzano-Jost, P., Chang, C., Chin, M., Commane, R., Correa, G., Crouse, J. D.,
1625 Daube, B., Dibb, J. E., DiGangi, J. P., Diskin, G. S., Dollner, M., Elkins, J. W., Fiore, A. M., Flynn, C. M., Guo, H., Hall, S. R., Hannun, R. A., Hills, A., Hintsa, E. J., Hodzic, A., Hornbrook, R. S., Huey, L. G., Jimenez, J. L., Keeling, R. F., Kim, M. J., Kupc, A., Lacey, F., Lait, L. R., Lamarque, J.-F., Liu, J., McKain, K., Meinardi, S., Miller, D. O., Montzka, S. A., Moore, F. L., Morgan, E. J., Murphy, D. M., Murray, L. T., Nault, B. A., Neuman, J. A., Nguyen, L., Gonzalez, Y., Rollins, A., Rosenlof, K., Sargent, M., Schill, G., Schwarz, J. P., Clair, J. M. S., Steenrod, S. D., Stephens, B. B., Strahan, S. E., Strode, S. A., Sweeney, C., Thames, A. B., Ullmann, K., Wagner,
1630 N., Weber, R., Weinzierl, B., Wennberg, P. O., Williamson, C. J., Wolfe, G. M., and Zeng, L.: The NASA Atmospheric Tomography (ATom) Mission: Imaging the Chemistry of the Global Atmosphere, *Bulletin of the American Meteorological Society*, 103, E761–E790, <https://doi.org/10.1175/BAMS-D-20-0315.1>, 2022.
- Thornhill, G., Collins, W., Olivíé, D., Skeie, R. B., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Gjermundsen, A., Horowitz, L., Lamarque, J.-F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., O’Connor, F. M., Paulot, F., Schulz, M., Scott, C. E., Séférian,
1635 R., Smith, C., Takemura, T., Tilmes, S., Tsigaridis, K., and Weber, J.: Climate-Driven Chemistry and Aerosol Feedbacks in CMIP6 Earth System Models, *Atmospheric Chemistry and Physics*, 21, 1105–1126, <https://doi.org/10.5194/acp-21-1105-2021>, 2021.
- Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur, R., Balkanski, Y., Bauer, S. E., Bellouin, N., Benedetti, A., Bergman, T., Berntsen, T. K., Beukes, J. P., Bian, H., Carslaw, K. S., Chin, M., Curci, G., Diehl, T., Easter, R. C., Ghan, S. J., Gong, S. L., Hodzic, A., Hoyle, C. R., Iversen, T., Jathar, S., Jimenez, J. L., Kaiser, J. W., Kirkevåg, A., Koch, D., Kokkola, H., Lee, Y. H., Lin, G., Liu,
1640 X., Luo, G., Ma, X., Mann, G. W., Mihalopoulos, N., Morcrette, J.-J., Müller, J.-F., Myhre, G., Myriokefalitakis, S., Ng, N. L., O’Donnell, D., Penner, J. E., Pozzoli, L., Pringle, K. J., Russell, L. M., Schulz, M., Sciare, J., Seland, Ø., Shindell, D. T., Sillman, S., Skeie, R. B., Spracklen, D., Stavrakou, T., Steenrod, S. D., Takemura, T., Tiitta, P., Tilmes, S., Tost, H., Van Noije, T., Van Zyl, P. G., Von Salzen, K., Yu, F., Wang, Z., Wang, Z., Zaveri, R. A., Zhang, H., Zhang, K., Zhang, Q., and Zhang, X.: The AeroCom Evaluation and Intercomparison of Organic Aerosol in Global Models, *Atmospheric Chemistry and Physics*, 14, 10 845–10 895, <https://doi.org/10.5194/acp-14-10845-2014>,
1645 2014.
- Tunved, P., Hansson, H.-C., Kerminen, V.-M., Ström, J., Maso, M. D., Lihavainen, H., Viisanen, Y., Aalto, P. P., Komppula, M., and Kulmala, M.: High Natural Aerosol Loading over Boreal Forests, *Science*, 312, 261–263, <https://doi.org/10.1126/science.1123052>, 2006.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TlOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TlOPOT>2.0.CO;2), 1977.
- 1650 Van Marle, M. J. E., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A.-L., Field, R. D., Arneth, A., Forrest, M., Hantson, S., Kehrwald, N. M., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Yue, C., Kaiser, J. W., and Van Der Werf, G. R.: Historic Global Biomass Burning Emissions for CMIP6 (BB4CMIP) Based on Merging Satellite Observations with Proxies and Fire Models (1750–2015), *Geoscientific Model Development*, 10, 3329–3357, <https://doi.org/10.5194/gmd-10-3329-2017>, 2017.
- Vehkamäki, H., Kulmala, M., Napari, I., Lehtinen, K. E. J., Timmreck, C., Noppel, M., and Laaksonen, A.: An Improved Parameterization
1655 for Sulfuric Acid–Water Nucleation Rates for Tropospheric and Stratospheric Conditions, *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2002JD002184>, 2002.
- Venugopal, A. U., Bhatti, Y. A., Morgenstern, O., Williams, J., Edkins, N., Hardacre, C., Jones, A., and Revell, L. E.: Constraining the Uncertainty Associated With Sea Salt Aerosol Parameterizations in Global Models Using Nudged UKESM1-AMIP Simulations, *Journal of Geophysical Research: Atmospheres*, 130, e2024JD041 643, <https://doi.org/10.1029/2024JD041643>, 2025.

- 1660 Veres, P. R., Neuman, J. A., Bertram, T. H., Assaf, E., Wolfe, G. M., Williamson, C. J., Weinzierl, B., Tilmes, S., Thompson, C. R.,
Thames, A. B., Schroder, J. C., Saiz-Lopez, A., Rollins, A. W., Roberts, J. M., Price, D., Peischl, J., Nault, B. A., Møller, K. H., Miller,
D. O., Meinardi, S., Li, Q., Lamarque, J.-F., Kupc, A., Kjaergaard, H. G., Kinnison, D., Jimenez, J. L., Jernigan, C. M., Hornbrook,
R. S., Hills, A., Dollner, M., Day, D. A., Cuevas, C. A., Campuzano-Jost, P., Burkholder, J., Bui, T. P., Brune, W. H., Brown, S. S.,
1665 Brock, C. A., Bourgeois, I., Blake, D. R., Apel, E. C., and Ryerson, T. B.: Global Airborne Sampling Reveals a Previously Unobserved
Dimethyl Sulfide Oxidation Mechanism in the Marine Atmosphere, *Proceedings of the National Academy of Sciences*, p. 201919344,
<https://doi.org/10.1073/pnas.1919344117>, 2020.
- Von Glasow, R. and Crutzen, P. J.: Model Study of Multiphase DMS Oxidation with a Focus on Halogens, *Atmospheric Chemistry and
Physics*, 4, 589–608, <https://doi.org/10.5194/acp-4-589-2004>, 2004.
- Wang, Y. and Jacob, D. J.: Anthropogenic Forcing on Tropospheric Ozone and OH since Preindustrial Times, *Journal of Geophysical
1670 Research: Atmospheres*, 103, 31 123–31 135, <https://doi.org/10.1029/1998JD100004>, 1998.
- Wang, Z., Wang, C., Yang, S., Lei, Y., Che, H., Zhang, X., and Wang, Q.: Evaluation of Surface Solar Radiation Trends over
China since the 1960s in the CMIP6 Models and Potential Impact of Aerosol Emissions, *Atmospheric Research*, 268, 105 991,
<https://doi.org/10.1016/j.atmosres.2021.105991>, 2022.
- Weber, J., Archer-Nicholls, S., Abraham, N. L., Shin, Y. M., Bannan, T. J., Percival, C. J., Bacak, A., Artaxo, P., Jenkin, M., Khan, M. A. H.,
1675 Shallcross, D. E., Schwantes, R. H., Williams, J., and Archibald, A. T.: Improvements to the Representation of BVOC Chemistry–Climate
Interactions in UKCA (V11.5) with the CRI-Strat 2 Mechanism: Incorporation and Evaluation, *Geoscientific Model Development*, 14,
5239–5268, <https://doi.org/10.5194/gmd-14-5239-2021>, 2021.
- Weber, J., Archer-Nicholls, S., Abraham, N. L., Shin, Y. M., Griffiths, P., Grosvenor, D. P., Scott, C. E., and Archibald, A. T.:
Chemistry-Driven Changes Strongly Influence Climate Forcing from Vegetation Emissions, *Nature Communications*, 13, 7202,
1680 <https://doi.org/10.1038/s41467-022-34944-9>, 2022.
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder,
P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann,
R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office
Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *Journal of Advances in Modeling Earth Systems*, 10, 357–380,
1685 <https://doi.org/10.1002/2017MS001115>, 2018.
- Williamson, C., Kupc, A., Wilson, J., Gesler, D. W., Reeves, J. M., Erdesz, F., McLaughlin, R., and Brock, C. A.: Fast Time Response
Measurements of Particle Size Distributions in the 3–60 Nm Size Range with the Nucleation Mode Aerosol Size Spectrometer,
Atmospheric Measurement Techniques, 11, 3491–3509, <https://doi.org/10.5194/amt-11-3491-2018>, 2018.
- Williamson, C. J., Kupc, A., Axisa, D., Bilsback, K. R., Bui, T., Campuzano-Jost, P., Dollner, M., Froyd, K. D., Hodshire, A. L., Jimenez,
1690 J. L., Kodros, J. K., Luo, G., Murphy, D. M., Nault, B. A., Ray, E. A., Weinzierl, B., Wilson, J. C., Yu, F., Yu, P., Pierce, J. R.,
and Brock, C. A.: A Large Source of Cloud Condensation Nuclei from New Particle Formation in the Tropics, *Nature*, 574, 399–403,
<https://doi.org/10.1038/s41586-019-1638-9>, 2019.
- Williamson, C. J., Kupc, A., Rollins, A., Kazil, J., Froyd, K. D., Ray, E. A., Murphy, D. M., Schill, G. P., Peischl, J., Thompson, C.,
Bourgeois, I., Ryerson, T. B., Diskin, G. S., DiGangi, J. P., Blake, D. R., Bui, T. P. V., Dollner, M., Weinzierl, B., and Brock, C. A.:
1695 Large Hemispheric Difference in Nucleation Mode Aerosol Concentrations in the Lowermost Stratosphere at Mid- and High Latitudes,
Atmospheric Chemistry and Physics, 21, 9065–9088, <https://doi.org/10.5194/acp-21-9065-2021>, 2021.

- Wofsy, S., Afshar, S., Allen, H., Apel, E., Asher, E., Barletta, B., Bent, J., Bian, H., Biggs, B., Blake, D., Blake, N., Bourgeois, I., Brock, C., Brune, W., Budney, J., Bui, T., Butler, A., Campuzano-Jost, P., Chang, C., Chin, M., Commane, R., Correa, G., Crouse, J., Cullis, P., Daube, B., Day, D., Dean-Day, J., Dibb, J., DiGangi, J., Diskin, G., Dollner, M., Elkins, J., Erdesz, F., Fiore, A., Flynn, C., Froyd, K., Gesler, D., Hall, S., Hanisco, T., Hannun, R., Hills, A., Hints, E., Hoffman, A., Hornbrook, R., Huey, L., Hughes, S., Jimenez, J., Johnson, B., Katich, J., Keeling, R., Kim, M., Kupc, A., Lait, L., McKain, K., Mclaughlin, R., Meinardi, S., Miller, D., Montzka, S., Moore, F., Morgan, E., Murphy, D., Murray, L., Nault, B., Neuman, J., Newman, P., Nicely, J., Pan, X., Paplawsky, W., Peischl, J., Prather, M., Price, D., Ray, E., Reeves, J., Richardson, M., Rollins, A., Rosenlof, K., Ryerson, T., Scheuer, E., Schill, G., Schroder, J., Schwarz, J., St.Clair, J., Steenrod, S., Stephens, B., Strode, S., Sweeney, C., Tanner, D., Teng, A., Thames, A., Thompson, C., Ullmann, K., Veres, P., Wagner, N., Watt, A., Weber, R., Weinzierl, B., Wennberg, P., Williamson, C., Wilson, J., Wolfe, G., Woods, C., Zeng, L., and Vieznor, N.: ATom: Merged Atmospheric Chemistry, Trace Gases, and Aerosols, Version 2, <https://doi.org/10.3334/ORNLDAAAC/1925>, 2021.
- Woodward, S.: Modeling the Atmospheric Life Cycle and Radiative Impact of Mineral Dust in the Hadley Centre Climate Model, *Journal of Geophysical Research: Atmospheres*, 106, 18 155–18 166, <https://doi.org/10.1029/2000JD900795>, 2001.
- Yan, C., Yin, R., Lu, Y., Dada, L., Yang, D., Fu, Y., Kontkanen, J., Deng, C., Garmash, O., Ruan, J., Baalbaki, R., Schervish, M., Cai, R., Bloss, M., Chan, T., Chen, T., Chen, Q., Chen, X., Chen, Y., Chu, B., Dällenbach, K., Foreback, B., He, X., Heikkinen, L., Jokinen, T., Junninen, H., Kangasluoma, J., Kokkonen, T., Kurppa, M., Lehtipalo, K., Li, H., Li, H., Li, X., Liu, Y., Ma, Q., Paasonen, P., Rantala, P., Pileci, R. E., Rusanen, A., Sarnela, N., Simonen, P., Wang, S., Wang, W., Wang, Y., Xue, M., Yang, G., Yao, L., Zhou, Y., Kujansuu, J., Petäjä, T., Nie, W., Ma, Y., Ge, M., He, H., Donahue, N. M., Worsnop, D. R., Veli-Matti Kerminen, Wang, L., Liu, Y., Zheng, J., Kulmala, M., Jiang, J., and Bianchi, F.: The Synergistic Role of Sulfuric Acid, Bases, and Oxidized Organics Governing New-Particle Formation in Beijing, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL091944>, 2021.
- Yokouchi, Y. and Ambe, Y.: Aerosols Formed from the Chemical Reaction of Monoterpenes and Ozone, *Atmospheric Environment*, 19, 1271–1276, [https://doi.org/10.1016/0004-6981\(85\)90257-4](https://doi.org/10.1016/0004-6981(85)90257-4), 1985.
- Yu, F., Nadykto, A. B., Luo, G., and Herb, J.: H₂SO₄–H₂O Binary and H₂SO₄–H₂O–NH₃ Ternary Homogeneous and Ion-Mediated Nucleation: Lookup Tables Version 1.0 for 3-D Modeling Application, *Geoscientific Model Development*, 13, 2663–2670, <https://doi.org/10.5194/gmd-13-2663-2020>, 2020.
- Yu, P., Froyd, K. D., Portmann, R. W., Toon, O. B., Freitas, S. R., Bardeen, C. G., Brock, C., Fan, T., Gao, R.-S., Katich, J. M., Kupc, A., Liu, S., Maloney, C., Murphy, D. M., Rosenlof, K. H., Schill, G., Schwarz, J. P., and Williamson, C.: Efficient In-Cloud Removal of Aerosols by Deep Convection, *Geophysical Research Letters*, 46, 1061–1069, <https://doi.org/10.1029/2018GL080544>, 2019.
- Zhang, Y., Li, D., He, X.-C., Nie, W., Deng, C., Cai, R., Liu, Y., Guo, Y., Liu, C., Li, Y., Chen, L., Li, Y., Hua, C., Liu, T., Wang, Z., Xie, J., Wang, L., Petäjä, T., Bianchi, F., Qi, X., Chi, X., Paasonen, P., Liu, Y., Yan, C., Jiang, J., Ding, A., and Kulmala, M.: Iodine Oxoacids and Their Roles in Sub-3 Nm Particle Growth in Polluted Urban Environments, *Atmospheric Chemistry and Physics*, 24, 1873–1893, <https://doi.org/10.5194/acp-24-1873-2024>, 2024.
- Zhao, B., Donahue, N. M., Zhang, K., Mao, L., Shrivastava, M., Ma, P.-L., Shen, J., Wang, S., Sun, J., Gordon, H., Tang, S., Fast, J., Wang, M., Gao, Y., Yan, C., Singh, B., Li, Z., Huang, L., Lou, S., Lin, G., Wang, H., Jiang, J., Ding, A., Nie, W., Qi, X., Chi, X., and Wang, L.: Global Variability in Atmospheric New Particle Formation Mechanisms, *Nature*, 631, 98–105, <https://doi.org/10.1038/s41586-024-07547-1>, 2024.

Appendix A

Table A1. Model-to-ATom median ratios for environmental conditions, precursor vapours, aerosol number size distributions, and chemical composition between 2 - 8 km altitude. Ratios are calculated using all available ATom data between 2 - 8 km for comparison with different UKESM1.1 configurations implementing various SA (sulfuric acid)-based nucleation schemes (see Table 2 for simulation details). The median values are calculated from the point-by-point model-to-ATom ratios (i.e. the ratio is computed at each location or time point, followed by taking the median), and therefore do not necessarily equal the ratio of the median model value to the median ATom value. Values greater than 1 indicate model overestimation, while values less than 1 indicate model underestimation relative to ATom observations. Environmental conditions include T, RH, and CS of dry aerosols. Precursor vapours include DMS, SO₂, NH₃, O₃, and OH. Aerosol number concentrations are reported for nucleation (dry diameter < 10 nm), Aitken (10 - 100 nm), accumulation (100 - 1000 nm), and coarse (1000 - 10,000 nm) modes, along with total number concentration. Chemical composition includes sulfate, organic matter, ammonium, nitrate, and seasalt mass concentrations, with total mass representing their sum. Missing values indicate that ammonium and nitrate components are not included in simulations without the ammonium nitrate scheme (SA-H₂O[default] and SA-NH₃-noNit). The extreme values in the NH₃ ratios result from extremely low NH₃ concentrations inferred from aerosol acidity, as NH₃ is virtually absent in the specified region.

	H₂SO₄(SA)-based schemes								
	H ₂ O(default)	NH ₃ -noNit	NH ₃ -slow	NH ₃ -Lana	NH ₃ (benchmark)	NH ₃ -CS2	NH ₃ -MSA	NH ₃ -IPSOA	NH ₃ -IPSC
Temperature	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
RH	1.10	1.10	1.10	1.10	1.10	1.09 <u>1.10</u>	1.10	1.09 <u>1.10</u>	1.10
CS dry	1.26	1.35 <u>1.36</u>	1.40 <u>1.41</u>	1.50	1.41 <u>1.44</u>	1.31 <u>1.29</u>	1.46 <u>1.47</u>	1.42 <u>1.44</u>	1.53 <u>1.54</u>
DMS	0.44	0.43 <u>0.49</u>	0.62 <u>0.53</u>	0.94 <u>0.89</u>	0.63 <u>0.54</u>	0.63 <u>0.60</u>	<u>0.65</u>	0.64	0.66 <u>0.67</u>
SO₂	1.02	1.02 <u>1.01</u>	1.12 <u>1.10</u>	1.15 <u>1.16</u>	0.47 <u>1.13</u>	1.14 <u>0.45</u>	<u>1.13</u>	1.15	1.11 <u>1.12</u>
NH₃	2797188.44	2846798.75 <u>2692912.45</u>	33.66 <u>54.93</u>	1.66 <u>1.04</u>	7.91 <u>7.71</u>	38.70 <u>48.76</u>	3.03 <u>3.37</u>	7.24 <u>6.21</u>	6.01 <u>4.93</u>
O₃	1.09	1.08 <u>1.09</u>	1.03	1.01 <u>1.02</u>	1.02	0.99 <u>1.05</u>	1.02 <u>1.01</u>	1.02	1.01
OH	1.13	1.13 <u>1.14</u>	1.06 <u>1.09</u>	1.04 <u>1.05</u>	1.05	1.08 <u>1.09</u>	1.06	1.06 <u>1.05</u>	<u>1.04</u>
Nucleation	0.92	1.91 <u>1.97</u>	0.57	0.30 <u>0.32</u>	0.37 <u>0.39</u>	0.22 <u>0.30</u>	0.21 <u>0.23</u>	0.35 <u>0.38</u>	0.27 <u>0.29</u>
Aitken	1.18	1.49	1.07 <u>1.09</u>	<u>0.97</u>	0.94	0.91 <u>0.80</u>	0.77 <u>0.95</u>	0.94 <u>0.92</u>	0.83 <u>0.85</u>
Accumulation	0.77	0.83 <u>0.84</u>	0.92 <u>0.94</u>	<u>1.01</u>	1.00	<u>0.92</u>	0.96	0.97 <u>1.00</u>	0.97 <u>0.98</u>
Coarse	1.30	1.28 <u>1.26</u>	1.43 <u>1.40</u>	1.50 <u>1.48</u>	1.51 <u>1.49</u>	1.47 <u>1.46</u>	1.50 <u>1.48</u>	1.46	1.65
Total number	1.03	1.39	0.87 <u>0.90</u>	0.74 <u>0.76</u>	0.74 <u>0.76</u>	0.63 <u>0.67</u>	0.70 <u>0.73</u>	0.67 <u>0.75</u>	<u>0.69</u>
Sulfate	1.42	1.48 <u>1.46</u>	1.37 <u>1.39</u>	1.46 <u>1.45</u>	1.32 <u>1.34</u>	1.06 <u>1.05</u>	1.45	1.35	1.29 <u>1.31</u>
Organic	0.95	1.01 <u>1.00</u>	0.96 <u>0.95</u>	0.93 <u>0.92</u>	0.93 <u>0.94</u>	0.82 <u>0.85</u>	0.89	0.95	1.52 <u>1.53</u>
Ammonium	–	–	7.47 <u>7.40</u>	7.97 <u>8.12</u>	7.68 <u>7.73</u>	7.04 <u>7.44</u>	7.88 <u>7.97</u>	7.77 <u>7.68</u>	7.34 <u>7.35</u>
Nitrate	–	–	4.48 <u>4.23</u>	8.90 <u>8.91</u>	9.82 <u>9.80</u>	8.33 <u>9.54</u>	9.54 <u>9.92</u>	9.10 <u>9.69</u>	<u>9.52</u>
Seasalt	2.10	2.19 <u>2.14</u>	0.66 <u>0.64</u>	0.72 <u>0.68</u>	<u>0.71</u>	0.75	0.74 <u>0.71</u>	0.70 <u>0.77</u>	0.73 <u>0.70</u>
Total mass	1.26	1.29 <u>1.28</u>	1.37 <u>1.38</u>	1.46 <u>1.47</u>	1.36 <u>1.38</u>	1.23 <u>1.19</u>	1.43	1.38 <u>1.39</u>	1.55 <u>1.56</u>

Table A2. Model-to-ATom median ratios for environmental conditions, precursor vapours, aerosol number size distributions, and chemical composition in upper troposphere (8 - 12 km altitude). Ratios are calculated using all available ATom data between 8 - 12 km for comparison with different UKESM1.1 configurations implementing various SA (sulfuric acid)-based nucleation schemes (see Table 2 for simulation details). The median values are calculated from the point-by-point model-to-ATom ratios (i.e. the ratio is computed at each location or time point, followed by taking the median), and therefore do not necessarily equal the ratio of the median model value to the median ATom value. Values greater than 1 indicate model overestimation, while values less than 1 indicate model underestimation relative to ATom observations. Environmental conditions include T, RH, and CS of dry aerosols. Precursor vapours include DMS, SO₂, NH₃, O₃, and OH. Aerosol number concentrations are reported for nucleation (dry diameter < 10 nm), Aitken (10 - 100 nm), accumulation (100 - 1000 nm), and coarse (1000 - 10,000 nm) modes, along with total number concentration. Chemical composition includes sulfate, organic matter, ammonium, nitrate, and seasalt mass concentrations, with total mass representing their sum. Missing values indicate that ammonium and nitrate components are not included in simulations without the ammonium nitrate scheme (SA-H₂O[default] and SA-NH₃-noNit). The extreme values in the NH₃ ratios result from extremely low NH₃ concentrations inferred from aerosol acidity, as NH₃ is virtually absent in the specified region.

	H₂SO₄(SA)-based schemes							
	H ₂ O(default)	NH ₃ -noNit	NH ₃ -slow	NH ₃ -Lana	NH ₃ (benchmark)	NH ₃ -CS2	NH ₃ -MSA	NH ₃ -IPSO
Temperature	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
RH	1.15	1.14	1.15	<u>1.14</u>	1.15	1.15	1.13 -1.15	1.15
CS dry	1.07	1.10 - <u>1.11</u>	1.10	1.20 - <u>1.21</u>	1.10	0.93 - <u>0.88</u>	1.14 - <u>1.13</u>	1.11 - <u>1.12</u>
DMS	0.16	0.17 - <u>0.14</u>	0.21	0.18 - <u>0.15</u>	0.23 - <u>0.16</u>	0.40 - <u>0.34</u>	0.23 - <u>0.21</u>	0.26 - <u>0.21</u>
SO₂	0.76	0.75	0.81	0.88 - <u>0.90</u>	0.83 - <u>0.80</u>	0.47 - <u>0.44</u>	0.82 -0.80	0.80 - <u>0.84</u>
NH₃	3644293052.67	3888048993.51 - <u>3748927339.80</u>	30685.18 - <u>33880.13</u>	0.00	0.03 - <u>0.00</u>	0.77 - <u>1.75</u>	0.05 -0.02	<u>0.03</u>
O₃	1.21	1.20	1.15	1.14 -1.13	1.07 - <u>1.14</u>	1.14	1.14	<u>1.14</u>
OH	1.09	1.08 - <u>1.09</u>	1.01	<u>0.99</u>	1.00	0.99	0.99	0.99
Nucleation	6.30	8.87 - <u>8.58</u>	3.30 - <u>3.11</u>	<u>1.60</u>	1.74	1.79 - <u>1.46</u>	1.37 - <u>1.39</u>	1.42 - <u>1.64</u>
Aitken	1.88	2.11 - <u>2.14</u>	1.64 - <u>1.61</u>	1.51 - <u>1.54</u>	1.39 - <u>1.41</u>	1.01 - <u>0.98</u>	1.51 - <u>1.50</u>	1.36 - <u>1.35</u>
Accumulation	0.39	0.42 - <u>0.43</u>	0.46 - <u>0.47</u>	0.57 - <u>0.56</u>	0.52	0.50 - <u>0.46</u>	0.50 - <u>0.51</u>	0.55 - <u>0.56</u>
Coarse	1.20	1.13 - <u>1.19</u>	1.23 -1.27	1.31	1.28 - <u>1.27</u>	1.29 -1.24	1.36 - <u>1.35</u>	<u>1.27</u>
Total number	2.11	2.58 - <u>2.52</u>	1.51 - <u>1.48</u>	1.14 - <u>1.12</u>	1.11 - <u>1.08</u>	0.85 - <u>0.81</u>	1.08 - <u>1.03</u>	1.08 - <u>1.07</u>
Sulfate	1.09	1.10 - <u>1.09</u>	1.04 - <u>1.02</u>	1.20 - <u>1.21</u>	1.01 - <u>1.04</u>	0.73 - <u>0.69</u>	1.13 - <u>1.12</u>	1.02 - <u>1.04</u>
Organic	0.43	0.45	0.44	<u>0.46</u>	0.45	0.44 -0.46	0.43 - <u>0.44</u>	0.49 - <u>0.50</u>
Ammonium	-	-	8.47 - <u>8.54</u>	9.70 - <u>9.78</u>	8.63 - <u>8.78</u>	7.44 - <u>7.18</u>	9.42 - <u>9.52</u>	8.68 - <u>8.93</u>
Nitrate	-	-	0.64 - <u>0.52</u>	0.46 - <u>0.49</u>	0.93 - <u>0.97</u>	2.73 - <u>1.92</u>	0.42 - <u>0.64</u>	1.06 - <u>0.82</u>
Seasalt	0.26	0.28 - <u>0.27</u>	0.13	0.15 - <u>0.13</u>	0.15 - <u>0.14</u>	0.15	0.14	0.15 - <u>0.14</u>
Total mass	0.72	0.73 - <u>0.74</u>	0.80	0.94	0.83 - <u>0.84</u>	0.77 - <u>0.69</u>	<u>0.88</u>	0.87




Figure A1. Vertical profiles of accumulation mode (100 - 1,000 nm) and coarse mode (1,000 - 10,000) aerosol number concentrations. ATom observations (black lines) and selected model simulations (coloured lines; see Table 2 for simulation details) are presented. Median values are shown for accumulation mode aerosols in the tropics (A, 25 °S to 25 °N), mid-latitudes (B, 25 - 60 °N/S), and high-latitudes (C, 60 - 90 °N/S), and for coarse mode aerosols in the tropics (D), mid-latitudes (E), and high-latitudes (F). Shaded areas for solid lines represent the interquartile range (25th to 75th percentiles) for both ATom observations and model simulations, indicating the variability in aerosol concentrations. Numbers on the right edge of each panel show the sample size used to calculate median values for each altitude bin at 5-minute temporal resolution.




Figure A2. Vertical profiles of sulfate and ammonium mass concentrations in particles smaller than $1 \mu\text{m}$ ($\mu\text{g m}^{-3}$). ATom observations (black lines) and selected model simulations (coloured lines; see Table 2 for simulation details) are presented. Median values are shown for sulfate mass concentrations in the tropics (A, 25°S to 25°N), mid-latitudes (B, $25 - 60^\circ\text{N/S}$), and high-latitudes (C, $60 - 90^\circ\text{N/S}$), and for ammonium mass concentrations in the tropics (D), mid-latitudes (E), and high-latitudes (F). Shaded areas for solid lines represent the interquartile range (25th to 75th percentiles) for both ATom observations and model simulations, indicating the variability in aerosol concentrations. Numbers on the right edge of each panel show the sample size used to calculate median values for each altitude bin at 5-minute temporal resolution.




Figure A3. Vertical profiles of organic and seasalt mass concentrations in particles smaller than $1 \mu\text{m}$ ($\mu\text{g m}^{-3}$). ATom observations (black lines) and selected model simulations (coloured lines; see Table 2 for simulation details) are presented. Median values are shown for organic mass concentrations in the tropics (A, 25°S to 25°N), mid-latitudes (B, $25 - 60^\circ\text{N/S}$), and high-latitudes (C, $60 - 90^\circ\text{N/S}$), and for seasalt mass concentrations in the tropics (D), mid-latitudes (E), and high-latitudes (F). Shaded areas for solid lines represent the interquartile range (25th to 75th percentiles) for both ATom observations and model simulations, indicating the variability in aerosol concentrations. Numbers on the right edge of each panel show the sample size used to calculate median values for each altitude bin at 5-minute temporal resolution.