



Assessment of snow model uncertainty in relation to the effect of a 1 °C warming using the snow modelling framework openAMUNDSEN

Erwin Rottler¹, Brage Storebakken¹, Michael Warscher², Florian Hanzer², Elena Bertazza¹, and Ulrich Strasser¹

Correspondence: Erwin Rottler (erwin.rottler@uibk.ac.at)

Abstract. Novel climate model data at the kilometer-scale, innovative downscaling techniques, sophisticated snow modelling frameworks and increasing computational capacities are among the elements that currently pave the way for a new phase in high resolution and physically based climate impact studies for the snow hydrology of mountain regions with complex topography. However, while the assessment of climate model uncertainty is well established, the uncertainty originating from the selection of the snow model usually only receives little attention. To investigate the uncertainty induced by the selection of the snow model configuration, we simulate the seasonal snow cover in the complex mountain area of the Berchtesgaden National Park mountains (Germany) under historical conditions (10/2013–09/2023) and for a 10-year period characterized by a 1 °C warming, using a large number of openAMUNDSEN snow model configurations (n = 108) with degree-day as well as physically based snowmelt methods and varying land cover maps and spatial resolutions. The analysis of the resulting snow cover durations and snow disappearance days suggests that differences showing up depending on the selected snowmelt method, land cover map and spatial resolution can be in the same range as the impact of a 1 °C warming, whereby uncertainties in the results are pronounced in the forest covered areas and in the high elevations of the study area. Our results support the identification of critical snow model settings that need to be considered, in particular, when using energy balance instead of degree-day snow models to investigate climate change impacts on the snow hydrology in complex mountain terrain.

15 1 Introduction

Climatic changes are expected to strongly affect the build-up and melt of seasonal snow covers in mountain regions. As the snow cover controls the seasonal rhythm of mountain ecosystems and represents a key part of the hydrological cycle, a detailed knowledge of its past, current and future state is of great significance (Gobiet et al., 2014; Beniston et al., 2018; Matiu et al., 2021; Rumpf et al., 2022; Kotlarski et al., 2023; Vormoor et al., 2025). In addition to in-situ observations and remote sensing data, snow models often are used to assess the current state of the snow cover and the effects of climatic changes. Such impacts of climatic changes commonly are assessed by forcing the snow-hydrological models with bias adjusted scenario data from ensembles of climate models (e.g. Bürger et al., 2011; Marke et al., 2015; Vormoor et al., 2015; Muelchi et al., 2021).

¹Department of Geography, University of Innsbruck, Innsbruck, Austria

²lumiosys GmbH, Innsbruck, Austria





Thereby, empirical snow models using degree day factors (DDFs) are often used in the framework of climate change impact studies, as they require little input data and are computationally very efficient. These temperature index (T-Index) approaches commonly are optimized using seasonally and/or spatially varying DDFs (Hock, 2003; Zhou et al., 2021). Hanus et al. (2021), for example, use a semi-distributed hydrological model with an improved degree-day method including a seasonally variable DDF (Girons Lopez et al., 2020) to assess runoff changes in Alpine catchments in Austria. In the SNOWGRID-CL model, which was applied to the Ötztal Alps (Austria) presented in the climate change impact study by Kotlarski et al. (2023), an extended degree-day approach accounting for air temperature and shortware radiation was used (Olefs et al., 2020). The degree-day approach is also utilized in the semi-distributed HBV model (Bergström, 1995; Seibert and Bergström, 2022) or in the GR models (Coron et al., 2017) (e.g. Vormoor et al., 2015; Adhikari et al., 2022; Dakhlaoui et al., 2022; Ten Berge et al., 2025). To better describe rain-on-snow events, the mesoscale Hydrological model (mHM), which has been applied to assess changes in low flows and river floods at different warming levels (Thober et al., 2018; Marx et al., 2018; Rottler et al., 2021), accounts for energy input from liquid rain; DDFs are thereby increased depending on the amount of liquid precipitation (Samaniego et al., 2010).

In addition to degree-day approaches, energy balance based snow models, which comprise process descriptions for all relevant energy and mass fluxes, are used to simulate the seasonal snow cover. These physically based modelling approaches require more input data and go along with a higher computational effort. Despite the increased complexity and effort, first studies have already harnessed energy balance based melt process description methods to assess the climate change impact on the seasonal snow cover. Hanzer et al. (2018), for example, use an energy balance based snowmelt modelling approach to assess cryospheric and hydrological impacts in the Ötztal Alps (Austria). They process an ensemble of climate model data using statistical bias correction methods as well as temporal disaggregation tools to conduct their energy balance based simulations. Marty et al. (2017) force the Alpine3D snow pack model with an ensemble of downscaled climate data for three emission scenarios in two different catchments in Switzerland. A semi-distributed approach using representative elementary areas and the Crocus snow model (Brun et al., 2012; Vionnet et al., 2012) is used to simulate the snow cover dynamics in French mountain massifs for historical conditions (Vernay et al., 2022) as well as to assess potential climate change impacts (Verfaillie et al., 2018; Kotlarski et al., 2023). A multi-physics ensemble approach for Crocus, which supports the representation of modelling errors, is presented in Lafaysse et al. (2017).

In the process of designing a climate change impact study for complex mountain terrain, not only the selection of the snowmelt modelling method needs to be considered, also the need for lateral snow redistribution (Freudiger et al., 2017; Hanzer et al., 2016; Mott et al., 2018; Quéno et al., 2024) and snow-canopy processes (Pomeroy et al., 2012; Sun et al., 2022; Sanmiguel-Vallelado et al., 2022; Essery et al., 2025) have to be addressed. The selection of the model configuration should be tailored to the specific research goal. However, the data availability and computational resources often are limited. Currently, newly available ensembles of kilometer-scale climate data (e.g. Ban et al., 2021; Pichelli et al., 2021; Collier et al., 2024) pave the way for a new generation of original climate change impact studies in mountain areas. However, while the uncertainty in the climate scenario data is usually addressed using an ensemble of model runs, the snow model uncertainty receives only little attention. Further studies illustrating the effect of different snow model configurations in the simulations are hence required





to support the development of robust modelling approaches for climate change impact studies in seasonally snow covered mountain areas.

In this study, we use the open source, intermediate complexity mountain snow model openAMUNDSEN (Strasser et al., 2024) to simulate the build-up and melt of the seasonal snow cover in the complex mountain terrain of the Berchtesgaden National Park (BGNP) under historical conditions (10/2013–09/2023) and for a 10-year period characterized by a 1 °C warming. The warming period is obtained by means of a stochastic block bootstrap resampler (climate generator), which is available as external pre-processing routine of openAMUNDSEN. We conduct snow simulations under historic and warming conditions using degree-day as well as energy balance based snow simulation routines, different land cover maps and grid geometries for the modelling with varying spatial resolutions (Fig. 1). For the historical period, model runs are evaluated using snow coverage maps derived from Sentinel-2 and MODIS data. For each model run, we calculate snow cover durations (SCDs) and snow disappearance days (SDDs) for the entire elevation range of the BGNP. Our main goal is to assess differences in the snow modelling results induced by different snow model configurations in relation to the effect of a 1 °C warming. The results support the need for careful consideration in the choice of snow model type, input data, and spatial resolution. The simulated effects of a climate change signal need to be interpreted having these uncertainties in mind.




Figure 1. Schematic overview of the modelling set-up depicting the conducted snow cover simulations using the mountain snow model openAMUNDSEN for the BGNP area under historical and 1 °C warming conditions using different snowmelt simulation methods, land cover maps and spatial resolutions.

2 Study area and data

We identified the BGNP as an ideal study site for our snow modelling experiment. The BGNP covers an area of 208 km² and is located in the state of Bavaria (Germany) in the North-East of the European Alps. It is the only German national park in the Alps and is characterized by a complex mountain landscape with an extreme topography covering elevations between





603–2713 m a.s.l. (Fig. 2 and 3). The open areas with bare rock and Alpine grassland in the higher elevations transition into coniferous, mixed and deciduous forest toward the lower elevations. The annual precipitation totals range from 1500 mm in the lower elevations to 2600 mm in the peak regions, and the build-up and melt of a seasonal snow cover dominates the hydrology in the area (e.g. Strasser, 2008; Warscher et al., 2013; Storebakken et al., 2025). In cooperation with the Bavarian Avalanche Warning Service, the administration of the BGNP operates a dense network of automatic weather stations (AWSs) in the area. A detailed description of the station network including station names, locations, measured variables, examples of measured time series as well as a description of the pre-processing of the AWS data is presented in Storebakken et al. (2025). The BGNP administration also provides a digital elevation model (DEM) in 10 m resolution, which we resample to resolutions between 20 and 1000 m (Fig. 3), and the BGNP habitat map (Fig. 2 a). This habitat map has been reclassified to seven classes, the three forest classes being the most important ones, as the respective forest types affect the snow-canopy interaction processes and hence the entire seasonal dynamics of the snow cover in the forested areas. In addition to the BGNP land cover map, we process five other land cover maps for the area, all of which base on data which is freely available and commonly used (Fig. 2 b-f). The motivation behind this is to capture the uncertainty that can be introduced into the distributed snow modelling with the selection of the land cover map. To facilitate comparability, we re-classify the data to similar land cover types. With regard to the forest representation, some data sources only have one forest class, while others differentiate different forest types and densities (Fig. 2). The land cover maps that we process are based on the following data: (i) CORINE Land Cover 2018 obtained from the European Union's Copernicus Land Monitoring Service (https://doi. org/10.2909/960998c1-1870-4e82-8051-6485205ebbac; last access 08 July 2025), (ii) OpenStreetMap (OSM) (https://www. openstreetmap.org/; last access 08 July 2025), which we download using the R package 'osmdata' (https://github.com/ropensci/ osmdata; last access 08 July 2025) (Padgham et al., 2017), (iii) WorldCover from the European Space Agency (ESA) (https:// esa-worldcover.org/; last access 08 July 2025) (Zanaga et al., 2022), (iv) Land Use/Cover Map for the European Alps developed by Marsoner et al. (2023) and (v) Land Cover Germany from the German Aerospace Agency (DLR) described in Weigand et al. (2020) (https://doi.org/10.15489/1ccmlap3mn39; last access 07 July 2025).

3 Methods

100

3.1 Intermediate complexity snow modelling

We use the open source snow-hydrological model openAMUNDSEN v1.0.4 to simulate the seasonal snow cover of the BGNP. openAMUNDSEN is a fully distributed snow model primarily designed to simulate the mass and energy balance of a seasonal snow cover in complex mountain terrain. On top of that, two different degree-day snowmelt modelling approaches are available as alternatives, namely the T-Index and the enhanced T-Index method. A full model description including these T-Index methods and all important process representations is presented in Strasser et al. (2024), the source code of the model being available on GitHub (https://github.com/openamundsen/openamundsen; last access: 07 July 2025). In the simulations presented here, we run the model in daily resolution with a constant DDF of 2.1 kg m⁻² d⁻¹ K⁻¹. This DDF was selected as a result of the comparison of model results with satellite-based snow data (see section 3.3). No automated calibration was conducted (since not the goal



115

120






Figure 2. Land cover maps for the BGNP based on the National Park habitat map (a; BGNP), CORINE land cover data (b; CORINE), OpenStreetMap data (c; OSM), ESA WorldCover data (d; WorldCover), the land use/land cover map for the European Alps developed by Marsoner et al. (2023) (e; EUSALP) and the Land Cover DE data from the German Aerospace Center (DLR) (f; LandCover-DE). The percentage coverage of each land cover class within 50 m elevation bands is shown right of each map.

here), but a robust value for DDF was selected following a series of model experiments. In addition to the T-Index approach, which only takes temperature into account, we also use the enhanced T-Index approach available within openAMUNDSEN, which also includes solar radiation (either measured radiation or potential clear-sky radiation) and modelled snow surface albedo (Pellicciotti et al., 2005). The incorporation of a radiation-driven melt component leads to seasonally and spatially varying snowmelt factors and enables simulations to be run at the sub-daily scale. The enhanced T-Index snow model runs are conducted in a 3-hourly time step with a DDF of 1.3 kg m⁻² d⁻¹ K⁻¹ and a short-wave radiation factor (RF) of 0.1 m² kg m⁻² W⁻¹ d⁻¹. In openAMUNDSEN, the T-Index and enhanced T-Index snowmelt modelling approaches are combined with the 'cryolayers' snow layering scheme, which enables to distinguish new snow, old snow, firn and ice. On the contrary, our 3-hourly openAMUNDSEN energy balance model runs engage a multilayer snowpack scheme following Essery (2015), distinguishing up to three snow layers (Strasser et al., 2024). For all model runs, we correct precipitation for wind-induced measurement errors following the approach presented by Kochendorfer et al. (2017). The station-based meteorological forcing data is spatially interpolated using a combined lapse rate/inverse distance weighting scheme. We account for the lateral snow redistribution process with a simple snow redistribution parameterization using snow redistribution factors (SRF), which are calculated based on the concept of negative topographic openness (Yokoyama et al., 2002). An evaluation of this approach with airborne laser scanning data is presented in Hanzer et al. (2016) and a description of an elevation-dependent adjustment of the SRF,








Figure 3. Digital elevation models for the BGNP in different resolutions including the location of automatic weather stations (AWS): 20 m (a), 50 m (b), 100 m (c), 250 m (d), 500 m (e) and 1000 m (f). The elevation distribution within 50 m elevation bands is shown right of each map.

tailored for the BGNP and also used in this study, is available in Storebakken et al. (2025). We assess SRFs for different resolutions and use them to redistribute the amount of solid precipitation from areas of erosion to areas of deposition (Fig. A1). All model input data and paremeter settings used in this study are available in the BGNP model set-up data repository (https://doi.org/10.23728/B2SHARE.530A7560A73647459969F5C21639E8CB; last access 28 July 2025).

3.2 Stochastic climate generator

125

130

To assess snow model uncertainty in relation to a climate warming signal, we use the stochastic block bootstrap resampler (climate generator) from openAMUNDSEN to construct a 10-year period characterized by a 1 °C warming compared to the historical 10-year period. A detailed description of the climate generator is presented in Strasser (2008) and Strasser et al. (2024). The source code is available along with the openAMUNDSEN model (https://github.com/openamundsen/openamundsen-climategenerator, last access: 21 July 2025). The openAMUNDSEN climate generator is designed to construct time series with predefined mean temperature and precipitation. Therefore, the observational time series are split up into blocks of a defined length (one week in our case) and then - week by week - rearranged to obtain the new time series with the desired characteristics. For the selection of a week in the new time series, the algorithm does not have all weeks from all months of the year available, but can only choose weeks observed up to two weeks before or after the week to be filled. The week 20 in the constructed time series, for example, can be selected from the weeks 18–22 in the observations. The width of this selection window is chosen by the user



140

145

150

155



and facilitates to individually define a compromise between a large data pool for selection and keeping seasonal characteristics consistent. Prior to the selection process, a random variability is added to the observations. Once an observed week has been selected by the algorithm to be used in the new time series, all measured variables (i.e., air temperature, precipitation, global radiation, relative humidity, and wind speed) are copied into the new data time series to be constructed. This approach enables the generation of high temporal resolution (hourly in our case) and physically consistent station time series that directly can be used to force the snow model simulations. However, while stochastic block bootstrapping procedures can be very useful in the framework of a snow model sensitivity study like the one presented here, we do not recommend their application in sophisticated climate change impact studies.

3.3 Model evaluation using satellite-based snow data

We evaluate the historical snow simulations using snow cover maps derived from Sentinel-2 and MODIS satellite data. The Sentinel-2 constellation allows for the determination of fractional snow cover (FSC) in high spatial resolution (i.e. $20x20 \,\mathrm{m}$). In this study, we use the Copernicus FSC obtained via the European Union's Copernicus Land Monitoring Service (https://doi.org/10.2909/3e2b4b7b-a460-41dd-a373-962d032795f3, last access 30 July 2025). Copernicus FSC data is available from September 2016 onwards. The available scenes for the study area are downloaded and pre-processed by cutting out the BGNP and re-sampling to the different model grids. We conduct a pixel-by-pixel comparison for all scenes with less than 50% cloud cover in the non-forested areas. The comparison of modelled and satellite-based data has four possible outcomes: true positive (TP), true negative (TN), false positive (FP) and false negative (FN). We determine the accuracy (ACC), overestimation error (OE) and underestimation error (UE) using the following equations:

$$ACC = \frac{TP + TN}{TP + TN + FP + FN} \tag{1}$$

$$OE = \frac{FP}{FN + FP} \tag{2}$$

$$UE = \frac{FN}{FP + FN} \tag{3}$$

In comparison to, e.g., Hao et al. (2019) and Rottler et al. (2024), we adapted the OE and UE in a way that the errors are assessed as a fraction of the total amount of non-matching pixels. In this way, OE and UE sum up to 1 and deviations of the ACC from a perfect match can more directly be linked to either an OE or UE. Alternative performance measures supporting snow model evaluation using satellite-based snow data inter alia are presented in Warscher et al. (2013) and Hofmeister et al. (2022). With regard to the model evaluation using FSC data, we present results on a monthly basis for February to July. Thereby we only take non-forested areas into account, as the satellite-based observation of snow on the ground inside forests is obstructed by the canopy (e.g. Xin et al., 2012; Bair et al., 2021; Keuris et al., 2023). The FSC and SWE values are translated to a binary





classification using a threshold of 20 % for FSC and 20 mm for SWE. A sensitivity assessment with regard to this threshold selection is presented, e.g., in Rottler et al. (2024).

In addition to FSC maps, we use MODIS-based snow cover extent (SCE) to evaluate historical snow model simulations. We download and process the daily and cloud-free 'Global SnowPack' product from the German Aeorospace Center (DLR) using DLR's GeoService (https://geoservice.dlr.de/; last access 30 July 2025). This SCE product has a spatial resolution of 500 m. A description of this SCE product as well as examples of its applications in research studies are available in Dietz et al. (2012, 2013, 2014, 2015), Rößler et al. (2021) and Roessler and Dietz (2023). We calculate the seasonal course of the snow coverage in the non-forested areas of the BGNP for the 10-year period 10/2013–09/2023 and compare the satellite-based values with openAMUNDSEN simulations.

4 Results

175

180

190

195

The evaluation of the openAMUNDSEN snow simulations using the described satellite-based snow data indicates that all model configurations can well capture the build-up and melt of the seasonal snow cover in the study area. A detailed description of the snow model evaluations is presented in the appendix B. The pixel-by-pixel comparison of simulated snow with FSC from Sentinel-2 (Fig. B1 and B2) as well as the comparison with MODIS snow cover extent data (Fig. B3 and B4) confirm the model's capabilities to capture the seasonal snow cover dynamics in the BGNP.

The stochastic block bootstrap resampler (climate generator) generated a 10-year period characterized by 1 °C warmer temperatures (Fig. 4 a). Due to the stochastic nature of the approach as well as the inherent natural variability of the observational data, not every month exhibits the exact same warming. Temperatures in December, for example, are on average 1.25 °C warmer, whereas March is characterized by a more moderate warming of 0.61 °C. With regard to precipitation amounts, slight increases/decreases show up (Fig. 4). Again, theses differences reflect the stochastic nature of the method as well as the inherent variability in the data. We consider this variability in the signal appropriate for 10-year averages. According to our experience, similar variabilities in the signal can show up in the comparison of 10-year periods of observations and data from climate models.

With regard to SWE, the snow model spread caused by the usage of different model configurations increases with progressing snow season and reaches its maximum between February and April and the average SWE in the BGNP can differ more than 100 mm. While the ensemble mean of the average SWE in warming conditions is below the ensemble mean of the historical simulations, there is a strong overlap of the shaded areas that indicated the spread caused by the usage of different snow model configurations (Fig. 4 c). All model simulations under historical as well as warming conditions suggest that the BGNP is almost fully snow-covered in January and February (Fig. 4 d). Furthermore, our snow simulations hint at an earlier disappearance of snow in the BGNP under a 1 °C warming by approx. one week. However, the strong overlap of the shaded areas indicates that the model spread in the second half of the main snowmelt season (May to June) is large compared to the magnitude of this shift. The model spread in this part of the year is characterized by values of approx. 50 mm for SWE and 12.5 % for the snow cover extent (Fig. 4 c and d).





SCDs reach the highest values of approx. 200 days in elevation zones between 1800 m and 2400 m (Fig. 4 e). Our simulations suggest that a 1 °C warming can cause an average decrease in SCD of 20.5 days in the BGNP. With regard to SDD, our modelling experiment suggests that changes due to a 1 °C warming can be small compared to the differences originating from the different model configurations (Fig. 4 f). In general, our model results suggest that for SCD and SDD small differences between model configurations show up in elevations between 1600–2200 m. A stronger spread in the snow model results can be detected below 1600 m and above 2200 m, respectively (Fig. 4 e and f).




Figure 4. Monthly average temperatures (a) and precipitation totals (b) for the BGNP based on historical observations (10/2013–09/2023) and a 10-year period characterized by a 1 °C warming generated using the openAMUNDSEN climate generator. The average seasonal cycle of simulated snow water equivalent (c) and snow cover extent (d) as well as the average snow cover duration (e) and snow disappearance day (f) for 200 m elevation bands is assessed for all openAMUNDSEN model configurations (i.e. 108 for historic and 1°C warming conditions, respectively).

Our analysis of the effect of the spatial resolution suggests that results are largely consistent for resolutions between 20 m and 250 m and up to elevations of 2000 m (Fig. 5). Simulation results for coarser spatial resolutions (i.e. 500 m or 1000 m) can deviate from the ones obtained for higher resolution model runs. The median SCD for the elevation band 2200–2400 m simulated using the T-Index approach (Fig. 5 a), for example, is 229, 218, 207 and 192 days using a spatial resolution of 20, 50, 100 and 250 m, respectively.



220






Figure 5. Simulated snow cover durations and snow disappearance days in the BGNP for different grid resolutions using the land cover map derived from the BGNP habitat data and the T-Index (a and b), enhanced T-Index (c and d) and energy balance (e and f) approaches available in the openAMUNDSEN snow model. Results are aggregated for 200 m elevation bands. The boxplots are displayed without whiskers and outliers.

The analysis of the model simulations with regard to the usage of different land cover maps indicates that the T-Index approach is largely insensitive to land cover (Fig. 6 a and b). However, when using the enhanced T-index or energy balance based snowmelt approach, differences in SCD and SDD can show up in elevation zones with forest classes (Fig. 6 c–f). The median SCD or SDD for elevations bands can be shifted by up to one week when using different land cover maps. In high elevations with no or only very small forested areas, the selection of the land cover map has no or only a negligible effect on the modelling results.

Our model experiments indicate that a 1 °C warming can cause a reduction in SCD of more than 20 days in the BGNP (Fig. 7 a, c and f). With regard to SDD, the change signal in our model experiment is smaller with values of a few days up to ten days (Fig. 9 b, d and f). The changes induced by higher temperatures and assessed for elevation bands seem consistent for spatial resolutions between 20–250 m. Deviations between model configuration using different spatial resolutions mainly show up at coarser modelling resolutions (i.e., 500 m or 1000 m) and for elevations below 800 m and above 2200 m. Our results also indicate that the magnitude of the difference can vary strongly within an elevation band. The length of the boxes, which cover 50 % of the cells within the elevation band, can easily span across a range of 10 days (Fig. 7).



235






Figure 6. Simulated snow cover durations and snow disappearance days in the BGNP for different land cover maps using a 50 m spatial grid resolution and the T-Index (a and b), enhanced T-Index (c and d) and energy balance (e and f) approach available in the openAMUNDSEN snow model. Results are aggregated for 200 m elevation bands. The boxplots are displayed without whiskers and outliers.

Our analyses of the effect of using different land cover maps on a potential climate change signal indicate that when using the T-Index approach, only small differences show up (Fig. 8 a and b). When using the enhanced T-Index or energy balance approach, the usage of different land cover maps characterized by different extents and distributions of forest type classes can be stronger and alter results for elevation bands with forest (Fig. 8 c–f).

Next, we directly compare results for SCDs and SDDs using different snowmelt simulation methods and assess results with respect to differences stemming from different aspect, land cover map and a 1 °C warming (Fig. 9). The aspects for the different model resolutions are presented in appendix A (Fig. A2). Our results point out that the selection of the snowmelt modelling method can strongly affect the results obtained for SCD and SDD. Differences between the snowmelt modelling methods are most prominent below approx. 1600 m with simulations using the T-Index approach resulting in considerable shorter SCDs (Fig. 9 a, c and e) and earlier SDDs (Fig. 9 b, d and f). Energy balance based simulations, however, result in longer SCDs and later SDDs for this elevation zone, while results from the enhanced T-Index approach tend to be in between the ones achieved with these two approaches. This signal changes for high elevations (> 2000 m) and the T-Index approach in this case suggests longer SCDs and later SDDs than the other two snowmelt modelling approaches. Our modelling results indicate that for elevations below approx. 1600 m and above approx. 2200 m the differences induced by different snowmelt



245






Figure 7. Changes in simulated snow cover durations and snow disappearance days in the BGNP induced by a 1 °C warming using the land cover map based on the BGNP habitat map for different grid resolutions using the T-Index (a and b), enhanced T-Index (c and d) and energy balance (e and f) approach available within the openAMUNDSEN snow model. Results are aggregated for 200 m elevation bands. The boxplots are displayed without whiskers and outliers.

modelling approaches are stronger than the signal of change induced by a 1 °C warming (Fig. 9 e and f). For elevation bands between 1600 m and 2200 m, the results for SCDs and SDDs are more consistent and the change induced by a 1 °C warming is larger than the spread caused by different snowmelt simulation methods. Also the usage of different land cover maps can induce changes in SCDs and SDDs that can be within the same range as the change induced by a 1 °C warming (Fig. 9 c and d), however, differences are less pronounced than differences resulting from the different snowmelt modelling methods, and are limited to elevations with differences in the forest representation in the land cover maps. While the enhanced T-Index and the energy balance approach can capture the differences in SCDs and SDDs depending on the exposition within elevation bands, the T-Index approach cannot account for such variabilities in the results (Fig. 9 a and b). Also the difference between northern and southern slopes is of the same magnitude as the signal induced by a 1 °C warming (Fig. 9).








Figure 8. Changes in simulated snow cover durations and snow disappearance days in the BGNP induced by a 1 °C warming for different different land cover maps using a 50 m spatial grid resolution and the T-index (a and b), enhanced T-index (c and d) and energy balance (e and f) approach available within the openAMUNDSEN snow model. Results are aggregated for 200 m elevation bands. The boxplots are displayed without whiskers and outliers.

5 Discussion

250

255

The open source snow model openAMUNDSEN proved to be a versatile snow modelling framework offering a wide range of configuration possibilities including degree-day and physically based snowmelt modelling approaches. The stochastic block bootstrap resampler (climate generator), an external pre-processing routine, is a useful extension of the openAMUNDSEN model and allows for the generation of physically consistent meteorological times series with pre-defined warming trends. The incorporation of the climate generator into the modelling experiment allows us to analyse the snow model uncertainty in the context of a potential climate warming. While the usage of the climate generator for our particular purpose - comparison of the results of different snow model configuration simulations in relation to a potential warming signal- is appropriate, we do not recommend the usage of a stochastic block bootstrapping procedure to produce climate change scenario data in the framework of a climate change impact study, in particular, when the goal is the investigation of snow-hydrological extremes, as no new extreme events can be produced by simple re-organization of observational time slices.



265

270






Figure 9. Simulated snow cover durations and snow disappearance days in the BGNP using the T-index, enhanced T-index and energy balance approach available in the openAMUNDSEN snow model depending on aspect (a and b), land cover maps based on National Park habitat data (BGNP) and the land use/land cover map for the European Alps developed by Marsoner et al. (2023) (EUSALP) (c and d) and a warming of 1 °C (e and f). Results are aggregated for 200 m elevation bands. The boxplots are displayed without whiskers and outliers.

In our study, we compare averages of meteorological and snow-related variables for two 10-year periods. The average monthly temperatures and precipitation totals as well as the average seasonal cycles indicate that the inter-annual variability of the data is affecting the averages investigated (Fig. 4 a–d). Our analysis shows how careful one should be in the quantification of climate change signals obtained using such short time frames. However, in our study, we specifically decided to compare 10-year periods, as this is the type of data currently being available from ensemble climate simulations at kilometer-scale resolution. These high resolution, "convection permitting" climate model runs allow for an explicit, physical description of deep convection without having to use parameterization schemes (Lucas-Picher et al., 2021). Ban et al. (2021), for example, conduct the first multi-model ensemble climate simulations for the greater Alpine region at the kilometer-scale for the ten year period 2000-2009. Multi-model 10-year kilometer-scale scenarios for the greater Alpine domain are presented in Pichelli et al. (2021). First case study simulations and results for a hydrological year for the "Third Pole", using a multi-model and multi-physics ensemble at the kilometer-scale, are presented in Prein et al. (2023) and Collier et al. (2024), respectively. On the contrary, high resolution regional climate model runs covering longer time frames (e.g. 30 years or more) often are limited to a single model only (e.g. Warscher et al., 2019; Collier and Mölg, 2020). Furthermore, depending on the complexity of



280

285

290

295

300

305



the terrain, resolutions below 1 km are required to adequately consider the processes that cause the spatial variability of the simulated snow cover, such as lateral redistribution of snow by wind or snow-canopy interaction. Hence, further dynamical or statistical downscaling of kilometer-scale climate model data is required to assess changes in the seasonal snow cover dynamics in complex mountain terrain such as the BGNP. Several studies recently demonstrated that the combination of high resolution weather model output, e.g. from the Weather Research and Forecasting (WRF) model, with physically-based snow energy balance models is feasible and can provide new insights into snow dynamics at the regional scale (e.g. Corripio and López-Moreno, 2017; Liu et al., 2019; Alonso-González et al., 2021; Raparelli et al., 2023; Zhou et al., 2023). The continuous improvement of the snow process descriptions in the weather models themselves might at some point make the coupling to an additional snow model unnecessary in the future (e.g. Luo et al., 2021; Saigger et al., 2024).

The recent progress in climate and snow modelling research suggests that step by step computational hurdles will be overcome and future climate change impact studies using models with an explicit treatment of deep convection and physically based snow models including lateral redistribution as well as snow canopy processes can adequately describe the snow cover dynamics in complex mountain terrain for potential future warming conditions. In this study, we show that the selection of the snowmelt simulation method is a very important aspect to consider when conceptualizing climate change impact studies in complex mountain areas. The presented modelling experiments suggest that the snow model uncertainty can be in the same range or even larger than changes in the snow cover induced by a 1 °C warming (Fig. 9 e and f). A particular focus is required for forested areas, as additional snow-canopy processes need to be taken into account. In contrast to degree-day approaches, which usually only consider temperature and a calibrated degree-day factor and which are largely insensitive to different land cover types, energy balance snow cover models do react sensitively to the choice of the land cover map and the forest representation therein (Fig. 6), given the meteorological forcing is properly adjusted to inside-canopy conditions. The comparison of model simulations with different spatial resolutions suggests that for a robust quantification of changes of the resulting snow patterns in the highest elevations, spatial resolutions considerably below 1 km are required. In our simulations, we use a simple snow redistribution parameterization to account for the lateral snow redistribution processes (Strasser et al., 2024) (Fig. A1). More process-oriented approaches incorporating wind fields and explicit representation of gravitational snow redistribution processes exist and need to be considered in the future development of the openAMUNDSEN model (e.g. Warscher et al., 2013; Freudiger et al., 2017; Quéno et al., 2024).

Our study contributes to the general research objective of quantifying snow model uncertainty and follows up on the investigations by Essery and Etchevers (2004), Essery et al. (2013), Günther et al. (2019) and Günther et al. (2020) who assess uncertainties in snow model results depending on the snow model complexity, and errors originating in the model parameters as well as in the forcing data. The effect of process representations on the obtained results are also presented in Magnusson et al. (2015). Particular consideration of snow in forested area was paid in the second phase of the Snow Model Intercomparison Project (SnowMIP2) where output from 33 different models was compared to each other (Essery et al., 2009). All these studies highlight the importance of the accuracy of the forcing data, more important than the selection of the parameters of the model, while the type of the model is of less importance, as are the computational requirements today. In addition, human errors are an important aspect affecting snow model performance (Menard et al., 2021). The snow modelling experiment presented in



315

320

325

330

335



our study complements existing research by defining a large set of configurations for distributed snow model simulations in complex mountain terrain with open and forested areas, resulting in a large number of openAMUNDSEN model runs (n = 108) with both degree-day and physically based snowmelt simulations, varying land cover maps and different spatial resolutions. The assessment of results in relation to the effects of a 1 °C warming supports our critical considerations of snow model uncertainty compared to the simulated climate change impact on the seasonal snow cover.

As many seasonally snow covered areas are also forest covered, further research is needed to better understand the effect of the snow-canopy processes in the context of a changing climate. A direct coupling of forest and snow models will thereby be an important next step in this regard, since not only climate will change, but also species composition and hence the snow hydrological processes inside the canopy. The complex mountain research site BGNP seems to be an ideal study site for such a task, both with respect to the dense measurement network and the existing expertise in forest processes modelling (e.g. Albrich et al., 2022; Senf et al., 2017; Thom et al., 2022; Dollinger et al., 2023; Braziunas et al., 2024) as well as snow modelling (e.g. Strasser, 2008; Strasser et al., 2011; Bernhardt et al., 2012; Kraller et al., 2012; Warscher et al., 2013; Storebakken et al., 2025). At the same time, however, the evaluation of snow models inside forests remains challenging, particularly, as forest canopies obstruct the satellite-based detection of snow on the ground. An alternative approach, the evaluation of openAMUNDSEN snow modelling results inside the forest using a network of microclimate sensors, is presented in Storebakken et al. (2025). Further research will support the development of new approaches for the evaluation of distributed snow models applied to forested areas, e.g. with respect to plant physiological simulation of LAI or improved scaling function for the meteorological variables from the location of the measurement to inside-canopy conditions. In our opinion, such new techniques for snow model evaluation are required to make a new generation of distributed and physically-based snow models ready for the application in the climate change impact context. There is simply no physical justification that DDFs used to simulate historical conditions are valid under future climatic conditions, and recent investigations of DDFs based on energy flux components (Ismail et al., 2023) show that DDFs cannot be treated as constant factors. They are changing with progressing melt season and need to be specifically adapted. Also our results suggest that degree-day snowmelt modelling approaches have their limitations, i.e. the DDF needs careful calibration and varies with elevation. At least we found that in higher elevations the best DDF is obviously larger than in the valley region where intermediate melting of the shallow snow cover in early winter can be very fast. Therefore, the most consistent results and only small differences between model configurations show up in elevations between 1600-2200 m (Fig. 4 e, f and Fig. 9 e, f). The spatially and seasonally variable snowmelt factors from the enhanced T-Index approach seem to improve results in this regard (due to the consideration of radiation which varies during the course of the season). However, the differences in SCDs and SDDs results of the enhanced T-Index and the energy balance simulation runs, respectively, can be substantial (Fig. 9). In lower elevations, the different forest representation in each particular land cover map causes additional uncertainties in the results, while SCDs and SDDs in the highest elevations react very sensitively to the selection of the spatial resolution (Fig. 6 and 5). Results are only consistent for spatial resolutions between 20 m and 250 m and up until elevations of 2000 m.



340

345

350

355

360



6 Conclusions

openAMUNDSEN is a versatile snow modelling framework, which allows for distributed snow cover simulations in complex mountain areas including snow-canopy interaction and lateral snow redistribution using degree-day as well as energy balance snowmelt methods. In this study, we use a large number of openAMUNDSEN model configurations (n = 108) characterized by different snowmelt modelling methods, land cover maps and spatial resolutions to simulate the seasonal snow cover in the complex mountain terrain of the BGNP for the historical period 10/2013-09/2023 and a 10-year period characterized by a 1 °C warming. The data describing the warming period was obtained by means of a stochastic block bootstrap resampler (climate generator), which is available as external pre-processing routine of openAMUNDSEN. The incorporation of the climate generator into our modelling experiment allows us to assess snow model uncertainty in relation to the effect of a potential warming signal. The complex and diverse mountain landscape of the BGNP proved to be an ideal study area for snow modelling research and the presented investigation of snow model uncertainty. Our analyses suggest that differences in the resulting SCDs and SDDs, originating from the selected snowmelt simulation method, the used land cover map and the chosen spatial resolution can be in the same range than the impact of a 1 °C warming. The uncertainties in the results are pronounced in the forest covered areas and in the high elevations of the study area. Our results emphasize the importance of properly investigating the uncertainty of a particular snow modelling setup when the effect of a warmer climate is to be quantified; other origins of uncertainty like the method to determine snowmelt, the used land cover map, or the chosen resolution can have effects on the results in the same order of magnitude as the climate change signal. Future snow modelling studies investigating the effect of climatic changes on the seasonal snow cover in complex mountain terrain can harness our findings and carefully consider the selection of the snow model configurations and related uncertainties.

Code and data availability. The openAMUNDSEN model source code including sample data and instruction on how to install and run the model is available on GitHub (https://github.com/openamundsen/openamundsen; last access: 28 July 2025). The source code of the stochastic block bootstrap resampler (climate generator) is available as pre-processing routine along with the openAMUNDSEN model (https://github.com/openamundsen/openamundsen-climategenerator, last access: 28 July 2025). The model set-up of the BGNP used including model configuration files with all parameter settings, different land cover maps, scripts used to prepare land cover maps as well as a script to run the model is available at: https://doi.org/10.23728/B2SHARE.530A7560A73647459969F5C21639E8CB (last access 28 July 2025).

Author contributions. ER designed and carried out the snow modelling and data analysis, produced the figures and wrote the first draft of the manuscript. All co-authors contributed to the concept of the study design and the discussion of the snow model results, and all co-authors contributed to the paper-writing process. In addition, BS supported the pre-processing of the openAMUNDSEN input data and the evaluation of the snow model performance, US developed the original versions of the openAMUNDSEN snow model and the climate generator, and FH implemented the latter in its current Python-based version into the openAMUNDSEN model framework.





Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. We thank the administration and the research department of the Berchtesgaden National Park and the colleagues of the MiKRoNP project for the provision of data and for their support. The project MiKRoNP (Project-No. 80489), which aims to identify microclimatic refugia and hotspots in a changing climate in the Bavarian National Parks, is financed by the Bavarian State Ministry of the Environment and Consumer Protection. The co-author Brage Storebakken receives support for his PhD from this project. The co-author Elena Bertazza has received funding from the Austrian Climate Research Programme (ACRP) within the project VecParClim (Project-No. 53759797), which aims to assess the impact of climate change on arthropod vectors and selected parasites in Alpine Pastures. Furthermore, we thank the European Union's Copernicus Land Monitoring Service (CLMS) for the provision of the fractional snow cover data and the colleagues from the German Aerospace Center (DLR) for the 'Global SnowPack' data. The University of Innsbruck generously supported the development and provision of the openAMUNDSEN model as an open-source project to the scientific community.





Appendix A: Snow redistribution factors and aspects




Figure A1. Snow redistribution factors for the BGNP for different spatial resolutions: 20 m (a), 50 m (b), 100 m (c), 250 m (d), 500 m (e) and 1000 m (f). The frequency of different snow redistribution factors is shown in the panel to the right of each map.








Figure A2. Aspects for the BGNP for different spatial resolutions: 20 m (a), 50 m (b), 100 m (c), 250 m (d), 500 m (e) and 1000 m (f). The frequency of different aspects is shown in the panel to the right of each map.



380

385

390



Appendix B: Model evaluation using satellite-based snow data

The pixel-by-pixel comparison based on FSC maps derived from Sentinel-2 images indicates an average accuracy for the months February to July between 0.82 and 0.93 (Fig. B2). Our results indicate that the coarser the spatial resolution of the model run and the re-sampled satellite image, the higher the accuracy obtained by this pixel-by-pixel comparison (Fig. B1 and B2). Using the energy balance approach, an overestimation of the snow cover shows up at the beginning of the snowmelt season. The deviations between simulated and observed snow cover at the end of the snowmelt seasons, however, are due to an underestimation error, i.e. too little snow covered area in the model simulations (Fig. B2 c1-c6). The comparison of the simulated build-up and melt of the seasonal snow cover with MODIS snow cover extent data for the 10-year period 10/2013–09/2023 suggests a good agreement between simulated and satellite-based data (Fig. B3 and B4). Looking at the evolution of the seasonal snow cover as the extent over the entire non-forested area, only very little differences show up between the different spatial resolutions, and the lines indicating the snow cover extent mostly overlap for all snowmelt modelling approaches (Fig. B4). For individual years (e.g., 2015 and 2016), the empirical snowmelt modelling approaches seem to produce a too late melt of the snow cover in the high elevations (Fig. B4 a and b). The calculated NSE values range from 0.86 to 0.93.








Figure B1. Simulated snow water equivalent (SWE) using the energy balance approach available in openAMUNDSEN and satellite-based fractional snow cover (FSC) maps derived from Sentinel-2 data for the BGNP on 23/04/2020 for different grid resolutions: 20 m (a), 50 m (b), 100 m (c), 250 m (d), 500 m (e) and 1000 m (f). The accuracy (ACC) assessed comparing the simulated SWE with the corresponding re-sampled satellite-based FSC observation is noted on the top-right of each SWE map.








Figure B2. Evaluation of openAMUNDSEN snow simulations in different spatial resolutions for non-forested areas in the BGNP between February and July using fractional snow cover (FSC) maps derived from Sentinel-2 data by means of accuracy (ACC), overestimation error (OE) and underestimation error (UE) for images with not more than 50% cloud coverage. The average ACC for all scenes is noted on the top-right of the plot panels.








Figure B3. Simulated snow water equivalent (SWE) using the T-Index (a), enhanced T-index (b) and energy balance (c) snowmelt simulation approaches available in the openAMUNDSEN snow model with a spatial resolution of 100 m and satellite-based snow cover extend maps (SCE) derived from MODIS snow cover data (d) for the BGNP and the three selected days: (1) 16/04/2020, (2) 01/05/2020 and (3) 15/04/2020. The total snow covered extent (SCE) is noted on the top-right of each map.








Figure B4. Evaluation of the openAMUNDSEN snow simulations for non-forested areas of the BGNP conducted using the T-index (a), enhanced T-index (b) and energy balance (c) snowmelt simulation approaches available in the openAMUNDSEN snow model by comparing the evolution of the simulated snow cover extent with the snow cover extent from MODIS satellite data for the time period 10/2013 to 09/2023. The Nash-Sutcliffe-Efficiency (NSE) for model runs with different spatial resolutions is noted on the top-right of each panel.





References

405

- Adhikari, T. R., Talchabhadel, R., Shrestha, S., Sharma, S., Aryal, D., and Pradhanang, S. M.: The evaluation of climate change impact on hydrologic processes of a mountain river basin, Theoretical and Applied Climatology, 150, 749–762, https://doi.org/10.1007/s00704-022-04204-3, 2022.
 - Albrich, K., Seidl, R., Rammer, W., and Thom, D.: From sink to source: changing climate and disturbance regimes could tip the 21st century carbon balance of an unmanaged mountain forest landscape, Forestry: An International Journal of Forest Research, 96, 399–409, https://doi.org/10.1093/forestry/cpac022, 2022.
- Alonso-González, E., Gutmann, E., Aalstad, K., Fayad, A., Bouchet, M., and Gascoin, S.: Snowpack dynamics in the Lebanese mountains from quasi-dynamically downscaled ERA5 reanalysis updated by assimilating remotely sensed fractional snow-covered area, Hydrology and Earth System Sciences, 25, 4455–4471, https://doi.org/10.5194/hess-25-4455-2021, 2021.
 - Bair, E. H., Stillinger, T., and Dozier, J.: Snow Property Inversion From Remote Sensing (SPIReS): A Generalized Multispectral Unmixing Approach With Examples From MODIS and Landsat 8 OLI, IEEE Transactions on Geoscience and Remote Sensing, 59, 7270–7284, https://doi.org/10.1109/TGRS.2020.3040328, 2021.
 - Ban, N., Caillaud, C., Coppola, E., Pichelli, E., Sobolowski, S., Adinolfi, M., Ahrens, B., Alias, A., Anders, I., Bastin, S., et al.: The first multi-model ensemble of regional climate simulations at kilometer-scale resolution, part I: evaluation of precipitation, Climate Dynamics, 57, 275–302, https://doi.org/10.1007/s00382-021-05708-w, 2021.
- Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A., Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J.-I., Magnusson, J., Marty, C., Morán-Tejéda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter, J., Strasser, U., Terzago, S., and Vincent, C.: The European mountain cryosphere: a review of its current state, trends, and future challenges, The Cryosphere, 12, 759–794, https://doi.org/10.5194/tc-12-759-2018, 2018.
 - Bergström, S.: The HBV model, p. 443-476, https://www.cabidigitallibrary.org/doi/full/10.5555/19961904773, 1995.
- Bernhardt, M., Schulz, K., Liston, G., and Zängl, G.: The influence of lateral snow redistribution processes on snow melt and sublimation in alpine regions, Journal of Hydrology, 424-425, 196–206, https://doi.org/10.1016/j.jhydrol.2012.01.001, 2012.
 - Braziunas, K. H., Geres, L., Richter, T., Glasmann, F., Senf, C., Thom, D., Seibold, S., and Seidl, R.: Projected climate and canopy change lead to thermophilization and homogenization of forest floor vegetation in a hotspot of plant species richness, Global Change Biology, 30, e17 121, https://doi.org/https://doi.org/10.1111/gcb.17121, 2024.
 - Brun, É., Vionnet, V., Morin, S., Boone, A., Martin, É., Faroux, S., Le Moigne, P., and Willemet, J.-M.: Le modèle de manteau neigeux Crocus et ses applications, La Météorologie, 2012, 44–54, https://doi.org/10.4267/2042/47245, 2012.
 - Bürger, G., Schulla, J., and Werner, A. T.: Estimates of future flow, including extremes, of the Columbia River headwaters, Water Resources Research, 47, https://doi.org/10.1029/2010WR009716, 2011.
 - Collier, E. and Mölg, T.: BAYWRF: a high-resolution present-day climatological atmospheric dataset for Bavaria, Earth System Science Data, 12, 3097–3112, https://doi.org/10.5194/essd-12-3097-2020, 2020.
- Collier, E., Ban, N., Richter, N., Ahrens, B., Chen, D., Chen, X., Lai, H.-W., Leung, R., Li, L., Medvedova, A., et al.: The first ensemble of kilometer-scale simulations of a hydrological year over the third pole, Climate Dynamics, 62, 7501–7518, https://doi.org/10.1007/s00382-024-07291-2, 2024.
 - Coron, L., Thirel, G., Delaigue, O., Perrin, C., and Andréassian, V.: The suite of lumped GR hydrological models in an R package, Environmental Modelling & Software, 94, 166–171, https://doi.org/https://doi.org/10.1016/j.envsoft.2017.05.002, 2017.





- 430 Corripio, J. G. and López-Moreno, J. I.: Analysis and Predictability of the Hydrological Response of Mountain Catchments to Heavy Rain on Snow Events: A Case Study in the Spanish Pyrenees, Hydrology, 4, https://doi.org/10.3390/hydrology4020020, 2017.
 - Dakhlaoui, H., Hakala, K., and Seibert, J.: Hydrological Impacts of Projected Climate Change on Northern Tunisian Headwater Catchments—An Ensemble Approach Addressing Uncertainties, pp. 499–519, Springer International Publishing, https://doi.org/10.1007/978-3-030-78566-6 24, 2022.
- Dietz, A. J., Wohner, C., and Kuenzer, C.: European Snow Cover Characteristics between 2000 and 2011 Derived from Improved MODIS Daily Snow Cover Products, Remote Sensing, 4, 2432–2454, https://doi.org/10.3390/rs4082432, 2012.
 - Dietz, A. J., Kuenzer, C., and Conrad, C.: Snow-cover variability in central Asia between 2000 and 2011 derived from improved MODIS daily snow-cover products, International Journal of Remote Sensing, 34, 3879–3902, https://doi.org/10.1080/01431161.2013.767480, 2013.
 - Dietz, A. J., Conrad, C., Kuenzer, C., Gesell, G., and Dech, S.: Identifying Changing Snow Cover Characteristics in Central Asia between 1986 and 2014 from Remote Sensing Data, Remote Sensing, 6, 12752–12775, https://doi.org/10.3390/rs61212752, 2014.
 - Dietz, A. J., Kuenzer, C., and Dech, S.: Global SnowPack: a new set of snow cover parameters for studying status and dynamics of the planetary snow cover extent, Remote Sensing Letters, 6, 844–853, https://doi.org/10.1080/2150704X.2015.1084551, 2015.
 - Dollinger, C., Rammer, W., and Seidl, R.: Climate change accelerates ecosystem restoration in the mountain forests of Central Europe, Journal of Applied Ecology, 60, 2665–2675, https://doi.org/https://doi.org/10.1111/1365-2664.14520, 2023.
- Essery, R.: A factorial snowpack model (FSM 1.0), Geoscientific Model Development, 8, 3867–3876, https://doi.org/10.5194/gmd-8-3867-2015, 2015.
 - Essery, R. and Etchevers, P.: Parameter sensitivity in simulations of snowmelt, Journal of Geophysical Research: Atmospheres, 109, https://doi.org/10.1029/2004JD005036, 2004.
- Essery, R., Rutter, N., Pomeroy, J., Baxter, R., Stähli, M., Gustafsson, D., Barr, A., Bartlett, P., and Elder, K.: SNOWMIP2:

 450 An Evaluation of Forest Snow Process Simulations, Bulletin of the American Meteorological Society, 90, 1120 1136, https://doi.org/10.1175/2009BAMS2629.1, 2009.
 - Essery, R., Morin, S., Lejeune, Y., and B Ménard, C.: A comparison of 1701 snow models using observations from an alpine site, Advances in Water Resources, 55, 131–148, https://doi.org/https://doi.org/10.1016/j.advwatres.2012.07.013, snow—Atmosphere Interactions and Hydrological Consequences, 2013.
- Essery, R., Mazzotti, G., Barr, S., Jonas, T., Quaife, T., and Rutter, N.: A Flexible Snow Model (FSM 2.1.1) including a forest canopy, Geoscientific Model Development, 18, 3583–3605, https://doi.org/10.5194/gmd-18-3583-2025, 2025.
 - Freudiger, D., Kohn, I., Seibert, J., Stahl, K., and Weiler, M.: Snow redistribution for the hydrological modeling of alpine catchments, WIREs Water, 4, e1232, https://doi.org/https://doi.org/10.1002/wat2.1232, 2017.
- Girons Lopez, M., Vis, M. J. P., Jenicek, M., Griessinger, N., and Seibert, J.: Assessing the degree of detail of temperature-based snow routines for runoff modelling in mountainous areas in central Europe, Hydrology and Earth System Sciences, 24, 4441–4461, https://doi.org/10.5194/hess-24-4441-2020, 2020.
 - Gobiet, A., Kotlarski, S., Beniston, M., Heinrich, G., Rajczak, J., and Stoffel, M.: 21st century climate change in the European Alps—A review, Science of The Total Environment, 493, 1138–1151, https://doi.org/https://doi.org/10.1016/j.scitotenv.2013.07.050, 2014.
- Günther, D., Marke, T., Essery, R., and Strasser, U.: Uncertainties in Snowpack Simulations—Assessing the Impact of Model Structure,

 Parameter Choice, and Forcing Data Error on Point-Scale Energy Balance Snow Model Performance, Water Resources Research, 55,

 2779–2800, https://doi.org/10.1029/2018WR023403, 2019.





- Günther, D., Hanzer, F., Warscher, M., Essery, R., and Strasser, U.: Including Parameter Uncertainty in an Intercomparison of Physically-Based Snow Models, Frontiers in Earth Science, 8, https://doi.org/10.3389/feart.2020.542599, 2020.
- Hanus, S., Hrachowitz, M., Zekollari, H., Schoups, G., Vizcaino, M., and Kaitna, R.: Future changes in annual, seasonal and monthly runoff signatures in contrasting Alpine catchments in Austria, Hydrology and Earth System Sciences, 25, 3429–3453, https://doi.org/10.5194/hess-25-3429-2021, 2021.
 - Hanzer, F., Helfricht, K., Marke, T., and Strasser, U.: Multilevel spatiotemporal validation of snow/ice mass balance and runoff modeling in glacierized catchments, The Cryosphere, 10, 1859–1881, https://doi.org/10.5194/tc-10-1859-2016, 2016.
- Hanzer, F., Förster, K., Nemec, J., and Strasser, U.: Projected cryospheric and hydrological impacts of 21st century climate change in the Ötztal Alps (Austria) simulated using a physically based approach, Hydrology and Earth System Sciences, 22, 1593–1614, https://doi.org/10.5194/hess-22-1593-2018, 2018.
 - Hao, S., Jiang, L., Shi, J., Wang, G., and Liu, X.: Assessment of MODIS-Based Fractional Snow Cover Products Over the Tibetan Plateau, IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, 12, 533–548, https://doi.org/10.1109/JSTARS.2018.2879666, 2019.
- 480 Hock, R.: Temperature index melt modelling in mountain areas, Journal of Hydrology, 282, 104–115, https://doi.org/https://doi.org/10.1016/S0022-1694(03)00257-9, mountain Hydrology and Water Resources, 2003.
 - Hofmeister, F., Arias-Rodriguez, L. F., Premier, V., Marin, C., Notarnicola, C., Disse, M., and Chiogna, G.: Intercomparison of Sentinel-2 and modelled snow cover maps in a high-elevation Alpine catchment, Journal of Hydrology X, 15, 100123, https://doi.org/j.hydroa.2022.100123, 2022.
- Ismail, M. F., Bogacki, W., Disse, M., Schäfer, M., and Kirschbauer, L.: Estimating degree-day factors of snow based on energy flux components, The Cryosphere, 17, 211–231, https://doi.org/10.5194/tc-17-211-2023, 2023.
 - Keuris, L., Hetzenecker, M., Nagler, T., Mölg, N., and Schwaizer, G.: An Adaptive Method for the Estimation of Snow-Covered Fraction with Error Propagation for Applications from Local to Global Scales, Remote Sensing, 15, https://doi.org/10.3390/rs15051231, 2023.
- Kochendorfer, J., Rasmussen, R., Wolff, M., Baker, B., Hall, M. E., Meyers, T., Landolt, S., Jachcik, A., Isaksen, K., Brækkan, R., and
 Leeper, R.: The quantification and correction of wind-induced precipitation measurement errors, Hydrology and Earth System Sciences,
 21, 1973–1989, https://doi.org/10.5194/hess-21-1973-2017, 2017.
 - Kotlarski, S., Gobiet, A., Morin, S., Olefs, M., Rajczak, J., and Samacoïts, R.: 21st Century alpine climate change, Climate Dynamics, 60, 65–86, https://doi.org/10.1007/s00382-022-06303-3, 2023.
- Kraller, G., Warscher, M., Kunstmann, H., Vogl, S., Marke, T., and Strasser, U.: Water balance estimation in high Alpine terrain by combining distributed modeling and a neural network approach (Berchtesgaden Alps, Germany), Hydrology and Earth System Sciences, 16, 1969–1990, https://doi.org/10.5194/hess-16-1969-2012, 2012.
 - Lafaysse, M., Cluzet, B., Dumont, M., Lejeune, Y., Vionnet, V., and Morin, S.: A multiphysical ensemble system of numerical snow modelling, The Cryosphere, 11, 1173–1198, https://doi.org/10.5194/tc-11-1173-2017, 2017.
- Liu, L., Ma, Y., Menenti, M., Zhang, X., and Ma, W.: Evaluation of WRF Modeling in Relation to Different Land Surface Schemes and
 Initial and Boundary Conditions: A Snow Event Simulation Over the Tibetan Plateau, Journal of Geophysical Research: Atmospheres,
 124, 209–226, https://doi.org/10.1029/2018JD029208, 2019.
 - Lucas-Picher, P., Argüeso, D., Brisson, E., Tramblay, Y., Berg, P., Lemonsu, A., Kotlarski, S., and Caillaud, C.: Convection-permitting modeling with regional climate models: Latest developments and next steps, WIREs Climate Change, 12, e731, https://doi.org/10.1002/wcc.731, 2021.



515

535



- Luo, L., Zhang, J., Hock, R., and Yao, Y.: Case Study of Blowing Snow Impacts on the Antarctic Peninsula Lower Atmosphere and Surface Simulated With a Snow/Ice Enhanced WRF Model, Journal of Geophysical Research: Atmospheres, 126, e2020JD033936, https://doi.org/https://doi.org/10.1029/2020JD033936, 2021.
 - Magnusson, J., Wever, N., Essery, R., Helbig, N., Winstral, A., and Jonas, T.: Evaluating snow models with varying process representations for hydrological applications, Water Resources Research, 51, 2707–2723, https://doi.org/https://doi.org/10.1002/2014WR016498, 2015.
- Marke, T., Strasser, U., Hanzer, F., Stötter, J., Wilcke, R. A. I., and Gobiet, A.: Scenarios of Future Snow Conditions in Styria (Austrian Alps), Journal of Hydrometeorology, 16, 261 277, https://doi.org/10.1175/JHM-D-14-0035.1, 2015.
 - Marsoner, T., Simion, H., Giombini, V., Egarter Vigl, L., and Candiago, S.: A detailed land use/land cover map for the European Alps macro region, Scientific Data, 10, 468, https://doi.org/10.1038/s41597-023-02344-3, 2023.
 - Marty, C., Schlögl, S., Bavay, M., and Lehning, M.: How much can we save? Impact of different emission scenarios on future snow cover in the Alps, The Cryosphere, 11, 517–529, https://doi.org/10.5194/tc-11-517-2017, 2017.
 - Marx, A., Kumar, R., Thober, S., Rakovec, O., Wanders, N., Zink, M., Wood, E. F., Pan, M., Sheffield, J., and Samaniego, L.: Climate change alters low flows in Europe under global warming of 1.5, 2, and 3 °C, Hydrology and Earth System Sciences, 22, 1017–1032, https://doi.org/10.5194/hess-22-1017-2018, 2018.
- Matiu, M., Crespi, A., Bertoldi, G., Carmagnola, C. M., Marty, C., Morin, S., Schöner, W., Cat Berro, D., Chiogna, G., De Gregorio,
 L., Kotlarski, S., Majone, B., Resch, G., Terzago, S., Valt, M., Beozzo, W., Cianfarra, P., Gouttevin, I., Marcolini, G., Notarnicola, C.,
 Petitta, M., Scherrer, S. C., Strasser, U., Winkler, M., Zebisch, M., Cicogna, A., Cremonini, R., Debernardi, A., Faletto, M., Gaddo, M.,
 Giovannini, L., Mercalli, L., Soubeyroux, J.-M., Sušnik, A., Trenti, A., Urbani, S., and Weilguni, V.: Observed snow depth trends in the
 European Alps: 1971 to 2019, The Cryosphere, 15, 1343–1382, https://doi.org/10.5194/tc-15-1343-2021, 2021.
- Menard, C. B., Essery, R., Krinner, G., Arduini, G., Bartlett, P., Boone, A., Brutel-Vuilmet, C., Burke, E., Cuntz, M., Dai, Y., Decharme,
 B., Dutra, E., Fang, X., Fierz, C., Gusev, Y., Hagemann, S., Haverd, V., Kim, H., Lafaysse, M., Marke, T., Nasonova, O., Nitta, T.,
 Niwano, M., Pomeroy, J., Schädler, G., Semenov, V. A., Smirnova, T., Strasser, U., Swenson, S., Turkov, D., Wever, N., and Yuan,
 H.: Scientific and Human Errors in a Snow Model Intercomparison, Bulletin of the American Meteorological Society, 102, E61 E79,
 https://doi.org/10.1175/BAMS-D-19-0329.1, 2021.
- Mott, R., Vionnet, V., and Grünewald, T.: The Seasonal Snow Cover Dynamics: Review on Wind-Driven Coupling Processes, Frontiers in Earth Science, 6, https://doi.org/10.3389/feart.2018.00197, 2018.
 - Muelchi, R., Rössler, O., Schwanbeck, J., Weingartner, R., and Martius, O.: River runoff in Switzerland in a changing climate changes in moderate extremes and their seasonality, Hydrology and Earth System Sciences, 25, 3577–3594, https://doi.org/10.5194/hess-25-3577-2021, 2021.
 - Olefs, M., Koch, R., Schöner, W., and Marke, T.: Changes in Snow Depth, Snow Cover Duration, and Potential Snowmaking Conditions in Austria, 1961–2020—A Model Based Approach, Atmosphere, 11, https://doi.org/10.3390/atmos11121330, 2020.
 - Padgham, M., Rudis, B., Lovelace, R., and Salmon, M.: osmdata, Journal of Open Source Software, 2, 305, https://doi.org/10.21105/joss.00305, 2017.
 - Pellicciotti, F., Brock, B., Strasser, U., Burlando, P., Funk, M., and Corripio, J.: An enhanced temperature-index glacier melt model including the shortwave radiation balance: development and testing for Haut Glacier d'Arolla, Switzerland, Journal of Glaciology, 51, 573–587, https://doi.org/10.3189/172756505781829124, 2005.



550

560

565

570



- Pichelli, E., Coppola, E., Sobolowski, S., Ban, N., Giorgi, F., Stocchi, P., Alias, A., Belušić, D., Berthou, S., Caillaud, C., et al.: The first multimodel ensemble of regional climate simulations at kilometer-scale resolution part 2: historical and future simulations of precipitation, Climate Dynamics, 56, 3581–3602, https://doi.org/10.1007/s00382-021-05657-4, 2021.
- Pomeroy, J., Fang, X., and Ellis, C.: Sensitivity of snowmelt hydrology in Marmot Creek, Alberta, to forest cover disturbance, Hydrological Processes, 26, 1891–1904, https://doi.org/10.1002/hyp.9248, 2012.
 - Prein, A. F., Ban, N., Ou, T., Tang, J., Sakaguchi, K., Collier, E., Jayanarayanan, S., Li, L., Sobolowski, S., Chen, X., et al.: Towards ensemble-based kilometer-scale climate simulations over the third pole region, Climate Dynamics, 60, 4055–4081, https://doi.org/10.1007/s00382-022-06543-3, 2023.
 - Quéno, L., Mott, R., Morin, P., Cluzet, B., Mazzotti, G., and Jonas, T.: Snow redistribution in an intermediate-complexity snow hydrology modelling framework, The Cryosphere, 18, 3533–3557, https://doi.org/10.5194/tc-18-3533-2024, 2024.
 - Raparelli, E., Tuccella, P., Colaiuda, V., and Marzano, F. S.: Snow cover prediction in the Italian central Apennines using weather forecast and land surface numerical models, The Cryosphere, 17, 519–538, https://doi.org/10.5194/tc-17-519-2023, 2023.
 - Roessler, S. and Dietz, A. J.: Development of Global Snow Cover—Trends from 23 Years of Global SnowPack, Earth, 4, 1–22, https://doi.org/10.3390/earth4010001, 2023.
- Rottler, E., Bronstert, A., Bürger, G., and Rakovec, O.: Projected changes in Rhine River flood seasonality under global warming, Hydrology and Earth System Sciences, 25, 2353–2371, https://doi.org/10.5194/hess-25-2353-2021, 2021.
 - Rottler, E., Warscher, M., Hanzer, F., and Strasser, U.: Spatio-temporal wet snow dynamics from model simulations and remote sensing: A case study from the Rofental, Austria, Hydrological Processes, 38, e15 279, https://doi.org/https://doi.org/10.1002/hyp.15279, 2024.
 - Rumpf, S. B., Gravey, M., Brönnimann, O., Luoto, M., Cianfrani, C., Mariethoz, G., and Guisan, A.: From white to green: Snow cover loss and increased vegetation productivity in the European Alps, Science, 376, 1119–1122, https://doi.org/10.1126/science.abn6697, 2022.
 - Rößler, S., Witt, M. S., Ikonen, J., Brown, I. A., and Dietz, A. J.: Remote Sensing of Snow Cover Variability and Its Influence on the Runoff of Sápmi's Rivers, Geosciences, 11, https://doi.org/10.3390/geosciences11030130, 2021.
 - Saigger, M., Sauter, T., Schmid, C., Collier, E., Goger, B., Kaser, G., Prinz, R., Voordendag, A., and Mölg, T.: A Drifting and Blowing Snow Scheme in the Weather Research and Forecasting Model, Journal of Advances in Modeling Earth Systems, 16, e2023MS004007, https://doi.org/10.1029/2023MS004007, 2024.
 - Samaniego, L., Kumar, R., and Attinger, S.: Multiscale parameter regionalization of a grid-based hydrologic model at the mesoscale, Water Resources Research, 46, https://doi.org/https://doi.org/10.1029/2008WR007327, 2010.
 - Sanmiguel-Vallelado, A., McPhee, J., Carreño, P. E. O., Morán-Tejeda, E., Camarero, J. J., and López-Moreno, J. I.: Sensitivity of forest–snow interactions to climate forcing: Local variability in a Pyrenean valley, Journal of Hydrology, 605, 127311, https://doi.org/10.1016/j.jhydrol.2021.127311, 2022.
 - Seibert, J. and Bergström, S.: A retrospective on hydrological catchment modelling based on half a century with the HBV model, Hydrology and Earth System Sciences, 26, 1371–1388, https://doi.org/10.5194/hess-26-1371-2022, 2022.
 - Senf, C., Pflugmacher, D., Hostert, P., and Seidl, R.: Using Landsat time series for characterizing forest disturbance dynamics in the coupled human and natural systems of Central Europe, ISPRS Journal of Photogrammetry and Remote Sensing, 130, 453–463, https://doi.org/10.1016/j.isprsjprs.2017.07.004, 2017.
 - Storebakken, B., Rottler, E., Warscher, M., and Strasser, U.: Modelling of the Seasonal Snow Cover Dynamics for Open and Forested Areas in the Berchtesgaden National Park (Germany) Using the openAMUNDSEN Mountain Snow Cover Model, Hydrological Processes, 39, e70 197, https://doi.org/10.1002/hyp.70197, 2025.





- Strasser, U.: Modelling of the Mountain Snow Cover in the Berchtesgaden National Park, ISBN 978-3-922325-62-8, https://www.nationalpark-berchtesgaden.de/service/publikationen/forschungsberichte/doc/fb_55.pdf, 2008.
 - Strasser, U., Warscher, M., and Liston, G. E.: Modeling snow–canopy processes on an idealized mountain, Journal of Hydrometeorology, 12, 663–677, https://doi.org/10.1175/2011JHM1344.1, 2011.
 - Strasser, U., Warscher, M., Rottler, E., and Hanzer, F.: openAMUNDSEN v1.0: an open-source snow-hydrological model for mountain regions, Geoscientific Model Development, 17, 6775–6797, https://doi.org/10.5194/gmd-17-6775-2024, 2024.
- Sun, N., Yan, H., Wigmosta, M. S., Lundquist, J., Dickerson-Lange, S., and Zhou, T.: Forest Canopy Density Effects on Snowpack Across the Climate Gradients of the Western United States Mountain Ranges, Water Resources Research, 58, e2020WR029194, https://doi.org/10.1029/2020WR029194, 2022.
 - Ten Berge, A. A., Booij, M. J., and Rientjes, T. H.: Robustness of hydrological models for simulating impacts of climate change on high and low streamflow, Journal of Hydrology, p. 133734, https://doi.org/10.1016/j.jhydrol.2025.133734, 2025.
- Thober, S., Kumar, R., Wanders, N., Marx, A., Pan, M., Rakovec, O., Samaniego, L., Sheffield, J., Wood, E. F., and Zink, M.: Multi-model ensemble projections of European river floods and high flows at 1.5, 2, and 3 degrees global warming, Environmental Research Letters, 13, 014 003, https://doi.org/10.1088/1748-9326/aa9e35, 2018.
 - Thom, D., Rammer, W., Laux, P., Smiatek, G., Kunstmann, H., Seibold, S., and Seidl, R.: Will forest dynamics continue to accelerate throughout the 21st century in the Northern Alps?, Global Change Biology, 28, 3260–3274, https://doi.org/https://doi.org/10.1111/gcb.16133, 2022.
 - Verfaillie, D., Lafaysse, M., Déqué, M., Eckert, N., Lejeune, Y., and Morin, S.: Multi-component ensembles of future meteorological and natural snow conditions for 1500 m altitude in the Chartreuse mountain range, Northern French Alps, The Cryosphere, 12, 1249–1271, https://doi.org/10.5194/tc-12-1249-2018, 2018.
- Vernay, M., Lafaysse, M., Monteiro, D., Hagenmuller, P., Nheili, R., Samacoïts, R., Verfaillie, D., and Morin, S.: The S2M meteorological and snow cover reanalysis over the French mountainous areas: description and evaluation (1958–2021), Earth System Science Data, 14, 1707–1733, https://doi.org/10.5194/essd-14-1707-2022, 2022.
 - Vionnet, V., Brun, E., Morin, S., Boone, A., Faroux, S., Le Moigne, P., Martin, E., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, Geoscientific Model Development, 5, 773–791, https://doi.org/10.5194/gmd-5-773-2012, 2012.
- Vormoor, K., Lawrence, D., Heistermann, M., and Bronstert, A.: Climate change impacts on the seasonality and generation processes of floods: projections and uncertainties for catchments with mixed snowmelt/rainfall regimes, Hydrology and Earth System Sciences, 19, 913–931, https://doi.org/10.5194/hess-19-913-2015, 2015.
 - Vormoor, K., Herzog, A., Francke, T., and Bronstert, A.: Patterns and Processes of Diel Streamflow Cycles Along the Longitudinal Profile of an Alpine Headwater Stream, Hydrological Processes, 39, e70 189, https://doi.org/https://doi.org/10.1002/hyp.70189, 2025.
- Warscher, M., Strasser, U., Kraller, G., Marke, T., Franz, H., and Kunstmann, H.: Performance of complex snow cover descriptions in a distributed hydrological model system: A case study for the high Alpine terrain of the Berchtesgaden Alps, Water Resources Research, 49, 2619–2637, https://doi.org/10.1002/wrcr.20219, 2013.
 - Warscher, M., Wagner, S., Marke, T., Laux, P., Smiatek, G., Strasser, U., and Kunstmann, H.: A 5 km Resolution Regional Climate Simulation for Central Europe: Performance in High Mountain Areas and Seasonal, Regional and Elevation-Dependent Variations, Atmosphere, 10, https://doi.org/10.3390/atmos10110682, 2019.





- Weigand, M., Staab, J., Wurm, M., and Taubenböck, H.: Spatial and semantic effects of LUCAS samples on fully automated land use/land cover classification in high-resolution Sentinel-2 data, International Journal of Applied Earth Observation and Geoinformation, 88, 102 065, https://doi.org/10.1016/j.jag.2020.102065, 2020.
 - Xin, Q., Woodcock, C. E., Liu, J., Tan, B., Melloh, R. A., and Davis, R. E.: View angle effects on MODIS snow mapping in forests, Remote Sensing of Environment, 118, 50–59, https://doi.org/10.1016/j.rse.2011.10.029, 2012.
- Yokoyama, R., Shirasawa, M., and Pike, R. J.: Visualizing topography by openness: A new application of image processing to digital elevation models, Photogrammetric engineering and remote sensing, 68, 257–266, https://www.asprs.org/wp-content/uploads/pers/2002journal/march/2002_mar_257-265.pdf, 2002.
 - Zanaga, D., Van De Kerchove, R., Daems, D., De Keersmaecker, W., Brockmann, C., Kirches, G., Wevers, J., Cartus, O., Santoro, M., Fritz, S., Lesiv, M., Herold, M., Tsendbazar, N.-E., Xu, P., Ramoino, F., and Arino, O.: ESA WorldCover 10 m 2021 v200, https://doi.org/10.5281/zenodo.7254221, 2022.
 - Zhou, G., Cui, M., Wan, J., and Zhang, S.: A Review on Snowmelt Models: Progress and Prospect, Sustainability, 13, https://doi.org/10.3390/su132011485, 2021.
- Zhou, X., Ding, B., Yang, K., Pan, J., Ma, X., Zhao, L., Li, X., and Shi, J.: Reducing the Cold Bias of the WRF Model Over the Tibetan Plateau by Implementing a Snow Coverage-Topography Relationship and a Fresh Snow Albedo Scheme, Journal of Advances in Modeling Earth Systems, 15, e2023MS003 626, https://doi.org/10.1029/2023MS003626, 2023.