

Spatiotemporal variation in rainfall predictability in Serbia under a changing climate

Tatijana Stosic¹, Ivana Tošić², Antonio Samuel Alves da Silva¹, Vladimir Djurdjević², Borko Stosic¹

¹Departamento de Estatística e Informática, Universidade Federal Rural de Pernambuco, 52171-900 Recife-PE, Brazil

5 ²Faculty of Physics, Institute for Meteorology, University of Belgrade, Belgrade, 11000, Serbia

Correspondence to: Ivana Tošić (itosic@ff.bg.ac.rs)

Abstract. This study examines whether the predictability of precipitation dynamics in Serbia has been influenced by climate change. We apply Generalized Weighted Permutation Entropy (GWPE) to evaluate the temporal structure of daily precipitation series using the parameter q , which filters subsets of small ($q < 0$) and large ($q > 0$) fluctuations. The analysis covers data from 10 14 weather stations between 1961 and 2020. Entropy values for $q = 0$ and $q = 2$, corresponding to Permutation Entropy and Weighted Permutation Entropy respectively, remained stable spatially and temporally. In contrast, GWPE values for $q = -10$ and $q = 10$, representing the predictability of small and large fluctuations, exhibited significant spatial and temporal variation between two 30-year subperiods. Entropy values for $q = -10$ were consistently lower, indicating that small precipitation fluctuations are more predictable than large ones. In several locations, significant changes in entropy occurred despite relatively 15 stable annual precipitation amounts. In others, annual totals varied while entropy remained constant. These findings suggest that climate change has influenced the predictability of precipitation in Serbia. By filtering fluctuations across scales, GWPE effectively reveals underlying changes that may be masked by standard statistical measures.

1 Introduction

The complexity of climate systems is widely recognized by researchers from different areas, such as physics, hydrology and 20 ecology (Rind, 1999; Mihailović et al., 2014; Dey and Mujumdar, 2022; Almeida-Ñauñay et al., 2021), and the research that addresses this aspect of climate has been intensifying over the recent years. In particular, there has been an increasing number of publications that address the complexity of climate variables by using the methods developed in complex system science, such as fractal and multifractal analysis (Lovejoy and Mandelbrot, 1985; Krzyszczak et al., 2019), methods originated in information theory (Silva et al., 2021a), and more recently complex networks (Boers et al., 2019). Precipitation and temperature 25 are used to describe the climate of certain regions (Da Silva et al., 2019), for climate classification (Peel et al., 2007), and to detect possible climate change (Alexander et al., 2006).

Climate models are widely used to predict future climate conditions, but must be validated against historical observations for different temporal and spatial scales to evaluate the models' performance for future projections (Vautard et al., 2021). Empirical results indicate that climate models perform better in reproducing temporal and spatial patterns of surface

30 temperature than those of precipitation (Trenberth, 2011; Lagos-Zuniga et al., 2022), thus other aspects such as emergent
properties that are characteristics of complex systems could contribute to better understanding of the nature of stochastic
processes that govern spatial and temporal precipitation variability, and help in evaluation of performance of climate models
and the impact of climate change. Fractal, multifractal, and complex network analyses have been extensively utilized to explore
the temporal and spatial variability of rainfall, while the potential of information theory-based methods in uncovering aspects
35 of rainfall dynamics remains less explored. Recently, entropy measures have been applied to characterize irregularities and the
rate of information flow in hydrological data, providing insights into different regimes and the effects of natural and human-
induced factors (de Carvalho Barreto et al., 2020; Agarwal et al., 2016; Rolim and de Souza Filho, 2025).

However, these entropy measures lack the ability to capture temporal relationships between successive values in a time series,
which was incorporated in the Permutation Entropy method introduced by Bandt and Pompe (2002). Rosso et al. (2007)
40 combined Permutation Entropy with a statistical complexity measure to develop the "Complexity-Entropy Causality Plane"
(CECP). This tool has proven effective in comparing systems with different degrees of complexity, especially in differentiating
between noise and chaos.

In hydrology Permutation entropy methods were used to analyze the nature of stochastic process governing streamflow
dynamics (Serinaldi et al., 2014; Mihailović et al., 2024), to optimize streamflow monitoring networks (Stosic et al., 2017), to
45 improve prediction models (Wang et al., 2024) and for detecting hydrological alterations caused by natural and human factors
(de Carvalho Barreto et al., 2023; Suriano et al., 2024). For a comprehensive review of the use of Permutation entropy-based
methods in hydrological studies see a recent article of Mihailović (2025). On the other hand, there are only a few studies that
use CECP to investigate the complexity of rainfall. Silva et al. (2021b) applied CECP method on monthly rainfall data recorded
in 133 pluviometric stations in Pernambuco, Northeast Brazil, during the period from 1950 to 2012, and showed the potential
50 of CECP to distinguish between different rainfall regimes. Tongal (2025) used CECP to improve the detection of hydrological
communities in the Lake District, Türkiye. Du et al. (2022) used PE along with other entropy measures to examine the rainfall
complexity in Beijing, China. Based on the monthly precipitation data recorded from 58 meteorological stations between 1968
and 2017, they observed significant spatial variations in rainfall complexity across the study area. They also found that rainfall
complexity has been increasing due to both climate variability and human activities. Liu et al. (2020) applied multiscale
55 permutation entropy to analyze precipitation complexity across 13 major cities in Heilongjiang Province, China. Based on
monthly data from 1967 to 2016, their findings indicate that water resources and urban living areas significantly influence
precipitation complexity, exhibiting a negative correlation.

In this study, we examine whether climate change influences the complexity of precipitation dynamics in Serbia by applying
the recently introduced "Generalized Weighted Complexity-Entropy Causality Plane" (GWCECP). the recently introduced
60 Generalized Weighted Permutation Entropy (GWPE) method (Stosic et al., 2022). GWPE extends the classical Permutation
Entropy (PE) framework through a modification of the Bandt and Pompe probability distribution, incorporating an additional

parameter $-\infty < q < \infty$. This generalization enables the evaluation of temporal complexity across multiple scales, emphasizing small fluctuations for $q < 0$ and large fluctuations for $q > 0$, thus offering novel insights into the underlying structure of precipitation variability. In particular, CECP represents a special case of GWCECP for $q = 0$. Previous studies related to temporal changes in precipitation in Serbia were based on trend analysis (Milovanović et al., 2017; Luković et al., 2014), extreme indices (Malinović-Miličević et al., 2016; Tošić et al., 2025) and changes in precipitation seasonality (Stosic et al., 2024a; Amiri and Gocić, 2025). Recently climate variability in Serbia was addressed by using the concept of complexity. Mimić et al. (2017) applied two complexity measures, Kolmogorov entropy and Sample entropy, on the daily maximum and minimum air temperature and the precipitation time series from seven stations in Serbia from the period 1951–2010. They found that both Kolmogorov complexity and Sample entropy of the maximum temperature series showed increasing trends for all stations, but it was statistically significant only for Kolmogorov entropy. The trends of complexity measures for the minimum temperature depend on the location, but without statistical significance except for locations in eastern (Negotin) and southern (Vranje) Serbia, where positive trend was found for Kolmogorov entropy and Sample entropy, respectively. For precipitation series both complexity measures showed decreasing trend which was statistically significant only for the Negotin station.

We address the influence of climate change on the complexity of rainfall dynamics in Serbia through the GWPE analysis of daily precipitation temporal series from 14 synoptic stations. We compare the GWPE curves in complexity/entropy causality plane for two 30-year sub-periods: 1961-1990 and 1991-2020. The main objective of this work is threefold, to find i) if complexity of temporal fluctuations of precipitation in Serbia varies over a wide range of scales; ii) if these properties vary spatially; iii) if precipitation complexity is affected by climate change.

2 Data and methodology

2.1 Study area and dataset

Serbia is a continental country located in the western part of the Balkan peninsula in the southeast of Europe, between latitudes $41^{\circ}50'$ and $46^{\circ}10'$ N (Fig. 1). The northern part is mostly flat with low elevation terrain, which gradually turns into hills and mountains that surround river valleys as moving towards central and southern part. In the northern part the climate type is moderate continental, changing to continental climate in the central part, and modified Mediterranean climate in the southern and the southwestern part (Bajat et al., 2015). In the lowlands the mean annual temperature is between 11 and 12 °C and mean annual precipitation between 500 and 700 mm, while in the mountains, the mean annual temperature is below 8 °C, and the mean annual precipitation is above 1000 mm (Vujadinović Mandić et al., 2022).

The data used in this work is the daily precipitation amount recorded during the period 1961-2020 at 14 stations located across Serbia, as shown in Fig. 1. The chosen stations have all the measurements during the study period. The data are provided by

the Serbian Meteorological Service, which performed technical and critical controls of these measurements. The geographic information on the 14 stations is given in Table 1.

Table 1: Stations with their geographic information (latitudes, longitudes and altitudes).

Abbreviations	Stations	Latitude	Longitude	Altitude (m)
BG	Belgrade	44°48' N	20°28' E	132
KG	Kragujevac	44°02' N	20°56' E	185
KR	Kraljevo	43°44' N	20°41' E	215
LO	Loznica	44°33' N	19°14' E	121
NE	Negotin	44°13' N	22°31' E	42
NI	Niš	43°20' N	21°54' E	201
NS	Novi Sad	45°20' N	19°51' E	84
SP	Smederevska Palanka	44°22' N	20°57' E	122
SO	Sombor	45°47' N	19°05' E	88
SR	Sremska Mitrovica	44°58' N	19°38' E	81
VG	Veliko Gradište	44°45' N	21°31' E	82
ZA	Zaječar	43°53' N	22°17' E	144
ZL	Zlatibor	43°39' N	19°41' E	1085
ZR	Zrenjanin	45°24' N	20°21' E	80




Figure 1: Location of Serbia in Europe (left, https://commons.wikimedia.org/wiki/File:Location_map_of_Serbia_in_Europe_%282006%E2%80%932008%29.gif) and orography of Serbia with locations of meteorological stations (right).

100 2.2 Generalized weighted permutation entropy

The Generalized Weighted Permutation Entropy (GWPE) method was recently introduced (Stosic et al., 2022) as the generalization of Permutation entropy (PE) (Bandt and Pompe, 2002) and Weighted permutation entropy method (WPE) (Fadlallah et al., 2013). While PE and WPE evaluate overall predictability of time series, GWPE is capable of distinguishing among the predictability of subsets of small/large fluctuations and thus provides a deeper insight into the time series behavior.

105 Along with PE, WPE and GWPE, statistical complexity can be defined, and measures of entropy and complexity can be represented and jointly analyzed in the Complexity entropy causality plane (CECP) (Rosso et al., 2007).

In the PE method (Bandt and Pompe, 2002), first overlapping segments (words) of a given size $d > 1$ are extracted along the time series $x_t, t = 1, \dots, N$, denoted as $X_s = (x_s, x_{s+1}, \dots, x_{s+d-1}), s = 1, \dots, N - (d - 1)$. The values within each segment are sorted in increasing order $x_{s+r_0} \leq x_{s+r_1} \leq \dots \leq x_{s+r_{d-1}}$, forming the vector of indices $v_s \equiv (r_0, r_1, \dots, r_{d-1})$, which represent

110 one of the $d!$ possible permutations of integers $0, 1, \dots, d - 1$, symbolically encoding the original segment. The relative frequencies of these permutations $\pi_i, i = 1, \dots, d!$, form the empirical distribution $P \equiv \{p(\pi_i), i = 1, \dots, d!\}$, which is used to calculate the permutation entropy as

$$H(P) = - \sum_{i=1}^{d!} p(\pi_i) \log p(\pi_i), \quad (1)$$

To accurately estimate $H(P)$, the length of the time series should be considerably greater than $d!$ For real-world data, selecting 115 the maximum embedding dimension that satisfies $T > 5d!$ is shown to be sufficient to get good statistics (Riedl et al., 2013).

In the CECP method (Rosso et al., 2007) the Jensen-Shannon divergence

$$J(P, U) = \left\{ H \left[\frac{P+U}{2} \right] - H \left[\frac{P}{2} \right] - H \left[\frac{U}{2} \right] \right\}, \quad (2)$$

which measures the distance of the empirical pattern distribution from a uniform distribution, is used to calculate the normalized complexity measure

$$120 \quad C(P) = \frac{H(P) J(P, U)}{d! J_{max}}, \quad (3)$$

where J_{max} is the maximum possible value of $J(P, U)$

$$J_{max} = -\frac{1}{2} \left[\frac{d!+1}{d!} \log(d! + 1) - 2 \log 2d! + \log d! \right] \quad (4)$$

obtained when only a single pattern is observed. The CECP represents a two-dimensional plot where $H(P)$ is used for the horizontal and $C(P)$ for the vertical coordinate.

125 The Weighted Permutation Entropy (WPE) (Fadlallah et al., 2013) introduced to account for the amplitude of the values in the segments. Instead of using relative pattern frequencies, WPE calculates probability as

$$p(\pi_i) = \frac{\sum_s \pi_s = \pi_i w_s}{\sum_s w_s}, \quad (5)$$

where $w_s = 1/d \sum_{i=0}^{d-1} (x_{s+1} - \langle x_s \rangle)^2$ is the variance, and $\langle x_s \rangle = 1/d \sum_{i=0}^{d-1} x_{s+1}$ is the average of the values observed in segment. The WPE and the corresponding CECP extension (WCECP) are then implemented using the above Eqs. (1-4)

130 These methods have proven valuable, with applications spanning fields such as medicine (Deng et al., 2015), finance (Zunino et al., 2010), ecology (Sippel et al., 2016), geophysics (Konstantinou et al., 2022), and hydrology (de Carvalho Barreto et al., 2023). Additionally, some studies (Xia et al., 2016) have demonstrated that the weighted variants outperform the original methods.

The Generalized Weighted Permutation Entropy (GWPE) method has been recently proposed (Stosic et al., 2022) as an extension of the earlier approaches, incorporating both PE and WPE as particular cases. It is achieved through the probability definition

$$p(\pi_i, q) = \frac{\sum'_s \pi_s = \pi_i w_s^{q/2}}{\sum'_s w_s^{q/2}}, \quad (6)$$

140 where $-\infty < q < \infty$ is a continuous scaling parameter. The prime in the summation indicates that the segments with strictly zero variance w_s are omitted. The scaling parameter q acts as a “magnifying glass”, where negative q values emphasize small fluctuations, and positive q values enhance large fluctuations. The PE method corresponds to GWPE for $q = 0$, and WPE corresponds to GWPE for $q = 2$. Although it was only recently introduced, GWPE has already found applications in data analysis in several studies (Stosic et al., 2024b; Stosic and Stosic, 2024; Duarte et al., 2025). Moreover, it was shown in ref. (Duarte et al., 2025) that GWPE outperformed PE and WPE in distinguishing between different sleep stages from EEG signals.

145 By applying formulas (2) and (3) to the choice of probability distribution (6) one obtains the Generalized Weighted Permutation Complexity (GWPEC), and combining the generalized weighted entropy and complexity the corresponding generalized weighted complexity plane (GWCECP) is obtained. [Source code in C, Python and R for application of GWCECP \(GWPE and GWPEC\) is available at https://github.com/stosicresearch/gwentropy.](https://github.com/stosicresearch/gwentropy)

3 Results and discussion

Mean annual precipitation for two subperiods for all 14 stations and the change from the first to second period is presented in Fig. 2, while spatial distribution of these values across Serbia is shown in Fig. 3. In both periods annual precipitation varies between 550 mm and 1050 mm. In the first period (1961-1990) the lowest precipitation (550-650 mm) was observed in the northern lowland part (Sombor, Zrenjanin, Novi Sad, Sremska Mitrovica), in valleys along the rivers Velika Morava in the central part (Smederevska Palanka, Kragujevac), Južna Morava in the southern part (Niš) and in the eastern part (Negotin and Zaječar). The intermediate precipitation (650-750 mm) was observed in the region along the Danube River (Belgrade, Veliko Gradište) and in Zapadna Morava valley (Kraljevo), while the highest precipitation (800-1000 mm) was observed in the western mountainous region (Loznica, Zlatibor). The lowest average precipitation was observed in Zrenjanin (555.8 mm) and the highest in Zlatibor (964.1 mm). In the second period (1991-2020) the precipitation increased in most of Serbian territory with largest increase in Novi Sad (151 mm). In Sremska Mitrovica, ~~Veliko Gradište, Negotin, and Zaječar and Kraljevo~~ the mean annual precipitation did not change, and in Veliko Gradište, Negotin, and Kraljevo a slight decrease was observed in the second period. The lowest average precipitation was observed in ~~Zrenjanin~~ Zrenjanin (597.2 mm) and the highest in Zlatibor (1030.4 mm).




Figure 2: Mean annual precipitation for two periods (1961-1990 and 1991-2020) for 14 stations in Serbia.

165





Figure 3: Spatial distribution of mean annual precipitation in Serbia, for two periods: (1961-1990 and 1991-2020) and change from first to second period. The third panel illustrates the change between these periods, expressed as the difference in mean annual precipitation (1991-2020 minus 1961-1990), highlighting regions with increasing or decreasing trends means. A slight decrease in annual precipitation observed at KR, NE and VG observed in Fig.2 falls into the “no change” (white) range of the color bar in the rightmost panel.


170

The GWPE method was applied on precipitation anomalies (deseasonalized) series $y_{i,j} = (x_{i,j} - \mu_i) / \sigma_i$ where $x_{i,j}$ is the precipitation value recorded on a given calendar day i of year j , μ_i and σ_i are mean and standard deviation calculated for each calendar day i over the years j of the observation period. GWPE curves in two-dimensional entropy complexity plane for all stations and for two considered sub-periods are presented in Fig. 4 and Fig. 5, respectively. In both periods and across all locations, entropy values are consistently lower for $q < 0$, indicating higher predictability of small fluctuations.


175




Loznica




Negotin




Niš




Novi Sad




Smederevska Palanka




Sombor




Sremska Mitrovica



Veliko Gradište




Zaječar



Zlatibor



Zrenjanin







Figure 4: GWPE curves representing entropy/complexity as a function of the magnification parameter q for 14 meteorological stations in Serbia during the first sub-period (1961–1990). The parameter q highlights small fluctuations for $q < 0$ and large fluctuations for $q > 0$, enabling scale-sensitive analysis of precipitation dynamics. Solid lines indicate theoretical bounds for a segment size of $w = 6$ days, as defined by Martin et al. (2006).

180








Figure 5: GWPE curves representing entropy/complexity as a function of the magnification parameter q for 14 meteorological stations in Serbia during the second sub-period (1991–2020). The parameter q highlights small fluctuations for $q < 0$ and large fluctuations for $q > 0$, enabling scale-sensitive analysis of precipitation dynamics. Solid lines indicate theoretical bounds for a segment size of $w = 6$ days, as defined by Martín et al. (2006).

A notable exception is observed in Sremska Mitrovica during the second period, where the curve that describes behavior of small fluctuations ($q < 0$) lies below that for large fluctuations ($q > 0$). This implies that, for subsets of fluctuations with identical entropy values, the larger fluctuations exhibit greater complexity. In all cases, Weighted permutation entropy (WPE) values are lower than their corresponding Permutation entropy (PE) values, while associated complexity values are comparatively higher. Considering statistical complexity, it is observed that for $q > 0$, the points are located in the right side of the GWCECP plane, where the decrease of entropy is followed by the increase of statistical complexity, while for $q < 0$ points are mostly located in the left side of CWCECP plane where both entropy and statistical complexity simultaneously decrease/increase. These observations are further supported by the complexity-entropy-scale causality boxes (CESCB) (Stosic et al., 2022), which plot entropy, complexity, and magnification factor q as coordinates in the three-dimensional space, as depicted in Fig. 6 and Fig. 7.


Belgrade




Kragujevac




Kraljevo




Loznica




Negotin




Niš






200 Figure 6: Three-dimensional GWPE causality box for 14 stations in Serbia for the first sub-period (1961-1990). The axes represent Generalized entropy (GWPE), Generalized complexity (GWPEC), and the parameter q that emphasizes small fluctuations for $q < 0$ and large fluctuations for $q > 0$.


Belgrade




Kragujevac




Kraljevo




Loznica




Negotin




Niš




Novi Sad



Smederevska Palanka



Sombor






Figure 7: Three-dimensional GWPE causality box for 14 stations in Serbia for the second sub-period (1991-2020). The axes represent Generalized entropy (GWPE), Generalized complexity (GWPEC), and the parameter q that emphasizes small fluctuations for $q < 0$ and large fluctuations for $q > 0$.

210

The values of GWPE and GWPEC for $q = -10$, $q = 10$, $q = 0$, and $q = 2$ for all stations and for two considered sub-periods are presented in Table 2 and Table 3, respectively. It is seen that GWPE and GWPEC values for $q = 0$ and $q = 2$ which correspond to the PE and WPE methods respectively, remain relatively stable across stations and time intervals. In contrast, the values of entropy (GWPE) and statistical complexity (GWPEC) for $q = -10$ and $q = 10$ exhibit noticeable variation both spatially and temporally between the two 30 years periods.

215

For enhanced interpretability, the GWPE and GWPEC results for all stations and both sub-periods and the change from the first to second period are shown in Fig. 8 and Fig. 9 respectively. It is seen from Table 2 and Fig. 8 that GWPE values at $q = -10$ are consistently lower across both periods, indicating greater predictability of small fluctuations of precipitation.

220

In the first sub-period small fluctuations ($q = -10$) exhibited the highest predictability (lowest GWPE values) in Sombor (0.012), Zaječar (0.014), Loznica (0.022), and Kragujevac (0.072), while the lowest predictability (highest GWPE values) was observed in Smederevska Palanka (0.419), and Niš (0.383). Between the first and second sub-periods, entropy values, and

consequently, predictability of rainfall fluctuations changed across stations. An increase in entropy (reflecting decreased predictability) was noted in Kragujevac, Negotin, Niš, Sremska Mitrovica, Veliko Gradište, and Zrenjanin. Conversely, predictability improved (entropy decreased) at Belgrade, Kraljevo, Smederevska Palanka, and Zlatibor. The remaining stations
225 Novi Sad, Sombor, Zaječar, and Loznica, showed no change in entropy values. Precipitation variability in Serbia is largely driven by atmospheric circulation patterns, notably the Arctic Oscillation (AO), East Atlantic/Western Russia Oscillation (EA/WR), and North Atlantic Oscillation (NAO) (Pavlović-Berdon, 2013; Tošić, 2004, 2014; Milošević, 2021). In addition to these teleconnections, local factors such as Serbia's complex topography and its position at the intersection of Mediterranean and continental climate systems play a critical role in shaping spatial and temporal precipitation patterns (Putnikovic et al.,
230 2016; Amiri and Gocic, 2025). Ruml et al. (2016) examined spatiotemporal variations in temperature and precipitation across Serbia using observational data from 26 meteorological stations spanning the period 1961–2010. Their findings indicated that annual precipitation did not exhibit significant changes over the full study period. However, during the sub-period 1986–2010, (which overlap with our second subperiod from 1991 to 2020) an increasing trend in annual precipitation was observed at numerous stations, including Belgrade, Kraljevo, and Smederevska Palanka. Notably, our analysis revealed enhanced
235 predictability of small-scale precipitation fluctuations at these locations, suggesting a possible link between rising precipitation and improved short-term predictability.

Among all stations, Belgrade exhibited the most substantial gain in predictability, with entropy decreasing from 0.360 to 0.138, while Negotin recorded the most significant loss, with entropy rising from 0.138 to 0.360. At these locations, although the average annual precipitation remained stable in the second sub-period, significant changes were observed in the predictability
240 of small fluctuations, suggesting that even in the absence of mean precipitation shifts, underlying complexity and short-term variability can evolve markedly. In contrast, at Novi Sad, where the most pronounced increase in precipitation occurred, predictability of small fluctuations remained unchanged, indicating a decoupling between precipitation volume and temporal structure.

The predictability of small rainfall fluctuations is crucial for estimation of hydrological impacts such as surface moisture retention and evapotranspiration dynamics (Yan et al., 2021; Cui et al., 2022). It is also important for agricultural planning as
245 predictable light rainfall helps optimize irrigation and planting time. Prolonged low intensity rainfall can trigger urban landslides and significantly contribute to soil erosion (Dunkerley, 2021). Spatial patterns reveal an increase in the predictability (lower entropy values) of small precipitation fluctuations across northern and eastern Serbia, which could be associated with increasing local atmospheric order. In contrast, the central and western regions show a decline in predictability suggesting
250 increased climatic irregularity or external forcing mechanisms that elevate entropy (Fig. 8a).

For all stations and both sub-periods, entropy values at $q = 10$ are consistently higher than those at $q = -10$, indicating that large precipitation fluctuations are less predictable than small ones. This reduced predictability poses challenges for

agricultural planning, water resource management, and the implementation of mitigation strategies for hydrological extremes such as floods and droughts (Sultan et al., 2005; Butterworth et al., 1999; Zhou et al., 2023).

255

Table 2: The values of GWPE for $q=-10$, $q=0$, and $q=2$, $q=10$, for 14 stations in Serbia for the two sub-periods (1961-1990 and 1991-2020).

City	1961-1990				1991-2020				Difference			
	$q=-10$	$q=0$	$q=2$	$q=10$	$q=-10$	$q=0$	$q=2$	$q=10$	$q=-10$	$q=0$	$q=2$	$q=10$
Belgrade	0.360	0.982	0.961	0.787	0.138	0.981	0.961	0.790	-0.222	-0.001	0	0.003
Kragujevac	0.072	0.980	0.958	0.813	0.211	0.979	0.959	0.803	0.139	-0.001	0.001	-0.01
Kraljevo	0.355	0.983	0.959	0.765	0.310	0.982	0.960	0.636	-0.045	-0.001	0.001	-0.129
Loznica	0.022	0.981	0.958	0.747	0.023	0.980	0.967	0.790	0.001	-0.001	0.009	0.043
Negotin	0.237	0.978	0.949	0.756	0.568	0.976	0.950	0.810	0.331	-0.002	0.001	0.054
Nis	0.383	0.981	0.956	0.812	0.441	0.980	0.962	0.795	0.058	-0.001	0.006	-0.017
Novi Sad	0.257	0.982	0.961	0.783	0.265	0.981	0.963	0.838	0.008	-0.001	0.002	0.055
Smed. Palanka	0.419	0.982	0.960	0.773	0.323	0.983	0.964	0.804	-0.096	0.001	0.004	0.031
Sombor	0.012	0.979	0.966	0.777	0.083	0.978	0.962	0.785	0.071	-0.001	-0.004	0.008
Srem. Mitrovica	0.331	0.982	0.961	0.780	0.403	0.981	0.965	0.428	0.072	-0.001	0.004	-0.352
Veliko Gradište	0.359	0.981	0.959	0.783	0.488	0.979	0.955	0.801	0.129	-0.002	-0.004	0.018
Zaječar	0.014	0.978	0.953	0.790	0.045	0.975	0.959	0.826	0.031	-0.003	0.006	0.036
Zlatibor	0.325	0.982	0.963	0.823	0.293	0.980	0.962	0.815	-0.032	-0.002	-0.001	-0.008
Zrenjanin	0.318	0.981	0.957	0.810	0.422	0.984	0.965	0.835	0.104	0.003	0.008	0.025

260

While entropy values at $q = 10$ show limited spatial variation across stations (Fig. 8b) there are evident temporal changes between the two sub-periods (Table 2). A slight increase in entropy values was observed at seven stations, predominantly in the eastern (Negotin, Veliko Gradište, Zaječar) and northern (Novi Sad, Zrenjanin) regions suggesting a decline in the predictability of large fluctuations, while three stations exhibited a slight entropy reduction, indicating improved predictability.

265

The most pronounced changes occurred at Kraljevo (central region) where entropy decreased from 0.765 to 0.636, and Sremska Mitrovica (northern region) which showed a substantial decrease from 0.780 to 0.428. These results indicate an improvement in the predictability of large precipitation fluctuations at these locations during the second sub-period, despite the stability in average annual precipitation. In contrast, Zlatibor and Sombor experienced an increase in precipitation over the same period, yet the predictability of large fluctuations remained unchanged, highlighting once again a decoupling between precipitation


270

volume and its temporal dynamics.

**Table 3: The values of GWPEC for $q=-10$, $q=0$, and $q=2$, $q=10$, for 14 stations in Serbia for the two sub-periods (1961-1990 and 1991-2020),
the second sub-period (1991-2020).**

Formatted: English (United States)


City	1961-1990				1991-2020				Difference			
	$q=-10$	$q=0$	$q=2$	$q=10$	$q=-10$	$q=0$	$q=2$	$q=10$	$q=-10$	$q=0$	$q=2$	$q=10$
Belgrade	0.314	0.042	0.091	0.350	0.128	0.044	0.091	0.354	-0.186	0.002	0.000	0.004
Kragujevac	0.070	0.047	0.099	0.338	0.187	0.049	0.097	0.343	0.117	0.002	-0.002	0.005
Kraljevo	0.319	0.040	0.095	0.364	0.273	0.043	0.093	0.355	-0.046	0.003	-0.002	-0.009
Loznica	0.022	0.046	0.100	0.371	0.022	0.047	0.078	0.334	0.000	0.001	-0.022	-0.037
Negotin	0.210	0.052	0.118	0.368	0.419	0.058	0.117	0.339	0.209	0.006	-0.001	-0.029
Nis	0.336	0.047	0.102	0.340	0.375	0.049	0.091	0.345	0.039	0.002	-0.011	0.005
Novi Sad	0.235	0.044	0.093	0.362	0.245	0.046	0.089	0.309	0.010	0.002	-0.004	-0.053
Smed. Palanka	0.347	0.043	0.094	0.360	0.287	0.042	0.085	0.336	-0.060	-0.001	-0.009	-0.024
Sombor	0.012	0.049	0.080	0.347	0.082	0.052	0.090	0.336	0.070	0.003	0.010	-0.011
Srem. Mitrovica	0.286	0.044	0.093	0.347	0.347	0.044	0.084	0.318	0.061	0.000	-0.009	-0.029
Veliko Gradište	0.292	0.045	0.095	0.350	0.367	0.051	0.103	0.350	0.075	0.006	0.008	0.000
Zaječar	0.014	0.053	0.109	0.366	0.045	0.059	0.098	0.320	0.031	0.006	-0.011	-0.046
Zlatibor	0.274	0.043	0.086	0.330	0.260	0.048	0.089	0.332	-0.014	0.005	0.003	0.002
Zrenjanin	0.287	0.046	0.101	0.345	0.354	0.039	0.082	0.312	0.067	-0.007	-0.019	-0.033



280

Figure 8: GWPE for $q = -10$ (a) and $q = 10$ (b) for 14 stations in Serbia for two periods (1961-1990 and 1991-2020). The third panel illustrates the change between these periods, expressed as the difference in GWPE (1991-2020 minus 1961-1990), highlighting regions with increasing or decreasing predictability of small ($q=-10$) and large ($q=10$) precipitation fluctuations.

285



290

Figure 9: GWPEC for $q = -10$ (a) and $q = 10$ (b) for 14 stations in Serbia for two periods (1961-1990 and 1991-2020). The third panel illustrates the change between these periods, expressed as the difference in GWPC (1991-2020 minus 1961-1990), highlighting regions with increasing or decreasing complexity of small ($q=-10$) and large ($q=10$) precipitation fluctuations.

4 Conclusions

Understanding of temporal and spatial patterns of precipitation and their alterations under the changing climate is crucial for climate impact studies. While statistical methods remain the primary tools for evaluation of these changes, recent findings suggest that emergent properties arising from complex interactions among processes that govern climate variability, may also be influenced by climate change.

The objective of this study was to examine whether one of such emergent property, complexity, is characteristic of precipitation time series in Serbia, and whether it has been affected by climate change. To this end, we applied the Generalized Weighted Permutation Entropy (GWPE) method to precipitation records from 14 weather stations distributed cross Serbia, spanning the period 1961-2020.

The main results of this work are: i) GWPE values for $q = 0$ and $q = 2$ which corresponding to Permutation entropy (PE) and Weighted permutation entropy (WPE) respectively, remain relatively stable across stations and time intervals; ii) GWPE values describing the predictability of small ($q = -10$) and large ($q = 10$) precipitation fluctuations exhibited noticeable spatial and temporal variation between the two 30-year periods; iii) GWPE values at $q = -10$ were consistently lower across both periods, indicating greater predictability of small precipitation fluctuations. In some cases, significant changes in entropy values occurred despite stable annual precipitation totals. Conversely, in other locations, entropy values remained steady even as annual totals varied, suggesting the decoupling between precipitation volume and its underlying temporal structure. These findings suggest that the complexity of precipitation dynamics in Serbia (measured via GWPE) has been influenced by climate change.

The GWPE shows strong potential to capture hidden variability in precipitation behavior and detect climate-related impacts that may be masked if focusing only on traditional statistical averages. Future work should compare these results with outputs from climate models under various emission scenarios, complementing classical statistical analyses to help identify the most suitable model or ensemble for climate change studies in this region. [Moreover, longer series should be used in future works to perform GWPE analysis in sliding windows, to address the potential crossovers and long-term trends in predictability.](#)

Code and Data availability

The datasets generated and/or analysed during the current study are available from the corresponding author upon reasonable request.

Author contribution

320 The conception of this study was the responsibility of TS. Material preparation, data collection, and analysis were performed
by TS, IT, ASAS, VDJ and BS. The paper was written by TS and revised by BS. All the authors have read and approved the
submission of paper.

Competing interests

The authors declare that they have no conflict of interest.

Special issue statement

325 This article is part of the special issue “Emerging predictability, prediction, and early-warning approaches in climate science”.

Acknowledgments

330 TS, ASAS and BS acknowledge support of Brazilian agency CNPq (grants No 308782/2022-4, 306590/2024-7 and
309499/2022-4). B.S. acknowledges support of Brazilian agency CAPES through grant No 88887.937789/2024-00. I.T. and
V.D. acknowledge the support of the Ministry of Science, Technological Development and Innovation of the Republic of
Serbia, grant No. 451-03-136/2025-03/200162.

References

- Agarwal, A., Maheswaran, R., Sehgal, V., Khosa, R., Sivakumar, B., and Bernhofer, C.: Hydrologic regionalization using
wavelet-based multiscale entropy method. *J. Hydrol.*, 538, 22–32, <https://doi.org/10.1016/j.jhydrol.2016.03.023>, 2016.
- Alexander, L. V., Zhang, X., Peterson, T. C., Caesar, J., Gleason, B., Klein Tank, A. M. G., ... and Vazquez-Aguirre, J. L.:
335 Global observed changes in daily climate extremes of temperature and precipitation. *J. Geophys. Res. Atmos.*, 111(D5),
<https://doi.org/10.1029/2005JD006290>, 2006.
- Almeida-Nauñay, A. F., Benito, R. M., Quemada, M., Losada, J. C., and Tarquis, A. M.: Complexity of the Vegetation-Climate
System Through Data Analysis, in: *Complex Networks & Their Applications IX: Volume 1, Proceedings of the Ninth
International Conference on Complex Networks and Their Applications COMPLEX NETWORKS 2020*, Springer, Cham,
340 609–619, https://doi.org/10.1007/978-3-030-65347-7_50, 2021.
- Amiri, M. and Gocić, M.: Identification of spatial and temporal variability of seasonality in precipitation regimes over Serbia.
Environ. Earth Sci., 84(6), 159, <https://doi.org/10.1007/s12665-025-12171-x>, 2025.

[Amiri, M. and Gocić, M.: Entropy analysis of precipitation variability in Serbia. *Theoretical and Applied Climatology*, 156\(8\), 454, <https://doi.org/10.1007/s00704-025-05708-4>, 2025.](https://doi.org/10.1007/s00704-025-05708-4)

345 Bajat, B., Blagojević, D., Kilibarda, M., Luković, J., and Tošić, I.: Spatial analysis of the temperature trends in Serbia during the period 1961–2010. *Theor. Appl. Climatol.*, 121, 289–301, <https://doi.org/10.1007/s00704-014-1243-7>, 2015.

Bandt, C. and Pompe, B.: Permutation entropy: a natural complexity measure for time series. *Phys. Rev. Lett.*, 88, 174102, <https://doi.org/10.1103/PhysRevLett.88.174102>, 2002.

350 Boers, N., Goswami, B., Rheinwalt, A., Bookhagen, B., Hoskins, B., and Kurths, J.: Complex networks reveal global pattern of extreme-rainfall teleconnections. *Nature*, 566, 373–377, <https://doi.org/10.1038/s41586-018-0872-x>, 2019.

Butterworth, J. A., Schulze, R. E., Simmonds, L. P., Moriarty, P., and Mugabe, F.: Hydrological processes and water resources management in a dryland environment IV: Long-term groundwater level fluctuations due to variation in rainfall. *HESS*, 3, 353–361, <https://doi.org/10.5194/hess-3-353-1999>, 1999.

355 Cui, Z., Wang, Y., Zhang, G. J., Yang, M., Liu, J., and Wei, L.: Effects of improved simulation of precipitation on evapotranspiration and its partitioning over land. *Geophys. Res. Lett.*, 49, e2021GL097353, <https://doi.org/10.1029/2021GL097353>, 2022.

Da Silva, P. E., Santos e Silva, C. M., Spyrides, M. H. C., and Andrade, L. D. M. B.: Precipitation and air temperature extremes in the Amazon and northeast Brazil. *Int. J. Climatol.*, 39, 579–595, <https://doi.org/10.1002/joc.5829>, 2019.

360 de Carvalho Barreto, I. D., Stosic, T., Filho, M. C., Delrieux, C., Singh, V. P., and Stosic, B.: Complexity analyses of Sao Francisco river streamflow: influence of dams and reservoirs. *J. Hydrol. Eng.*, 25, 05020036, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.00019](https://doi.org/10.1061/(ASCE)HE.1943-5584.00019), 2020.

de Carvalho Barreto, I., Stosic, T., Cezar Menezes, R. S., Alves da Silva, A. S., Rosso, O. A., and Stosic, B.: Hydrological changes caused by the construction of dams and reservoirs: The CECF analysis. *Chaos, an Interdiscip. J. Nonlin. Sci.*, 33, <https://doi.org/10.1063/5.0135352>, 2023.

365 Deng, B., Liang, L., Li, S., Wang, R., Yu, H., Wang, J., and Wei, X.: Complexity extraction of electroencephalograms in Alzheimer's disease with weighted-permutation entropy. *Chaos, an Interdiscip. J. Nonlin. Sci.*, 25, 043105, doi: 10.1063/1.4917013, 2015.

Dey, P. and Mujumdar, P.: On the statistical complexity of streamflow. *Hydrol. Sci. J.*, 67, 40–53, <https://doi.org/10.1080/02626667.2021.2000991>, 2022.

- 370 Du, L., Li, X., Yang, M., Sivakumar, B., Zhu, Y., Pan, X., ... and Sang, Y. F.: Assessment of spatiotemporal variability of precipitation using entropy indexes: A case study of Beijing, China. *Stoch. Environ. Res. Risk Assess.*, 36, 939–953, <https://doi.org/10.1007/s00477-021-02116-8>, 2022.
- Duarte, C. D., Pacheco, M., Iaconis, F. R., Rosso, O. A., Gasaneo, G., and Delrieux, C. A.: Statistical Complexity Analysis of Sleep Stages. *Entropy*, 27, 76, <https://doi.org/10.3390/e27010076>, 2025.
- 375 Dunkerley, D. L.: Light and low-intensity rainfalls: A review of their classification, occurrence, and importance in landsurface, ecological and environmental processes. *Earth-Sci. Rev.*, 214, 103529, <https://doi.org/10.1016/j.earscirev.2021.103529>, 2021.
- Fadlallah, B., Chen, B., Keil, A., and Principe, J.: Weighted-permutation entropy: A complexity measure for time series incorporating amplitude information. *Phys. Rev. E*, 87, 022911, <https://doi.org/10.1103/PhysRevE.87.022911>, 2013.
- 380 Konstantinou, K. I., Rahmalia, D. A., Nurfitriana, I., and Ichihara, M.: Permutation entropy variations in seismic noise before and after eruptive activity at Shinmoedake volcano, Kirishima complex, Japan. *EPS*, 74, 175, <https://doi.org/10.1186/s40623-022-01729-9>, 2022.
- Krzyszczak, J., Baranowski, P., Zubik, M., Kazandjiev, V., Georgieva, V., Sławiński, C., ... and Nieróbca, A.: Multifractal characterization and comparison of meteorological time series from two climatic zones. *Theor. Appl. Climatol.*, 137, 1811–1824, <https://doi.org/10.1007/s00704-018-2705-0>, 2019.
- 385 Lagos-Zuniga, M., Balmaceda-Huarte, R., Regoto, P., Torrez, L., Olmo, M., Lyra, A., ... and Bettolli, M. L.: Extreme indices of temperature and precipitation in South America: Trends and intercomparison of regional climate models. *Clim. Dyn.*, 1–22, <https://doi.org/10.1007/s00382-022-06598-2>, 2022.
- Liu, D., Yan, T., Liu, S., and Yan, G.: Variation Characteristics and Complexity Measurement of Urban Precipitation Based on Multi-scale Permutation Entropy. *J. Phys. Conf. Ser.*, 1624, 032027, doi:10.1088/1742-6596/1624/3/032027, 2020.
- 390 Lovejoy, S. and Mandelbrot, B. B.: Fractal properties of rain, and a fractal model. *Tellus A*, 37, 209–232, <https://doi.org/10.1111/j.1600-0870.1985.tb00423.x>, 1985.
- Martin, M. T., Plastino, A., and Rosso, O. A.: Generalized statistical complexity measures: Geometrical and analytical properties. *Physica A*, 369(2), 439–462, <https://doi.org/10.1016/j.physa.2005.11.053>, 2006.
- Mihailović, D. T., Mimić, G., and Arsenić, I.: Climate predictions: The chaos and complexity in climate models. *Adv. Meteorol.*, 2014, 878249, <https://doi.org/10.1155/2014/878249>, 2014.
- 395

- Mihailović, D. T., Malinović-Milićević, S., Frau, F. J., Singh, V. P., and Han, J.: Predictability of monthly streamflow by considering complexity measures. *J. Hydrol.*, 634, 131103, <https://doi.org/10.1016/j.jhydrol.2024.131103>, 2024.
- Mihailović, D. T.: Permutation Entropy and Its Niche in Hydrology: A Review. *Entropy*, 27, 598, <https://doi.org/10.3390/e27060598>, 2025.
- 400 Milovanović, B., Schuster, P., Radovanović, M., Vakanjac, V. R., and Schneider, C.: Spatial and temporal variability of precipitation in Serbia for the period 1961–2010. *Theor. Appl. Climatol.*, 130, 687–700, <https://doi.org/10.1007/s00704-017-2118-5>, 2017.
- Milošević, D., Stojavljević, R., Szabó, S., Stankov, U., Savić, S., & Mitrović, L.: Spatio-temporal variability of precipitation over the Western Balkan countries and its links with the atmospheric circulation patterns. *Journal of the Geographical Institute "Jovan Cvijic"*, SASA, 71(1), 29-42. <https://doi.org/10.2298/IJGI2101029M>. 2021.
- Mimić, G., Mihailović, D. T., and Kapor, D.: Complexity analysis of the air temperature and the precipitation time series in Serbia. *Theor. Appl. Climatol.*, 127, 891–898, <https://doi.org/10.1007/s00704-015-1677-6>, 2017.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification. *HESS*, 11, 1633–1644, <https://doi.org/10.5194/hess-11-1633-2007>, 2007.
- 410 ~~Pavlović Berdon, N.: The Impact of Teleconnection on Pressure, Temperature and Precipitation in Serbia. *Int. J. Remote Sens. Appl.*, 3(4), 185–192. <https://doi.org/10.14355/ijrsa.2013.0304.03>, 2013.~~ ~~Pavlović Berdon, N. (2013). The Impact of Teleconnection on Pressure, Temperature and Precipitation in Serbia. *International Journal of Remote Sensing Applications*, 3(4), 185–192. <https://doi.org/10.14355/ijrsa.2013.0304.03>.~~
- 415 ~~Putniković, S., Tošić, I., & Đurđević, V.: Circulation weather types and their influence on precipitation in Serbia. *Meteor. Atmos. Phys.*, 128(5), 649-662, <https://doi.org/10.1007/s00703-016-0432-6>, 2016.~~
- Riedl, M., Müller, A., and Wessel, N.: Practical considerations of permutation entropy: A tutorial review. *EPJ ST*, 222, 249-262, <https://doi.org/10.1140/epjst/e2013-01862-7>, 2013.
- Rind, D.: Complexity and climate. *Science*, 284, 105–107, doi:10.1126/science.284.5411.1, 1999.
- Rolim, L. Z. R. and de Souza Filho, F. D. A.: Multiscale Entropy Analysis of Rainfall and Streamflow Dynamics in Ceará, Brazil: Implications for Hydrological Predictability and Water Resource Management. *Water Conserv. Sci. Eng.*, 10, 27, <https://doi.org/10.1007/s41101-025-00347-2>, 2025.

- Rosso, O. A., Larrondo, H. A., Martin, M. T., Plastino, A., and Fuentes, M. A.: Distinguishing noise from chaos. *Phys. Rev. Lett.*, 99, 154102, <https://doi.org/10.1103/PhysRevLett.99.154102>, 2007.
- Serinaldi, F., Zunino, L., and Rosso, O. A.: Complexity–entropy analysis of daily stream flow time series in the continental United States. *Stoch. Environ. Res. Risk Assess.*, 28, 1685–1708, <https://doi.org/10.1007/s00477-013-0825-8>, 2014.
- Ruml, M., Gregoric, E., Matovic, G., Radovanovic, S., Vujadinovic, M., & Vukovic, A. (2017). Temperature and precipitation changes in Serbia between 1961 and 2010. *Annals of the University of Craiova-Agriculture, Montanology, Cadastre Series*, 46(2), 260-265.
- Silva, A. S. A. D., Barreto, I. D. D. C., Cunha-Filho, M., Menezes, R. S. C., Stosic, B., and Stosic, T.: Multiscale complexity analysis of rainfall in northeast Brazil. *Water*, 13, 3213, <https://doi.org/10.3390/w13223213>, 2021a.
- Silva, A. S. A., Menezes, R. S. C., Rosso, O. A., Stosic, B., and Stosic, T.: Complexity entropy-analysis of monthly rainfall time series in northeastern Brazil. *Chaos, Soliton Fractal.*, 143, 110623, <https://doi.org/10.1016/j.chaos.2020.110623>, 2021b.
- Sippel, S., Lange, H., Mahecha, M. D., Hauhs, M., Bodesheim, P., Kaminski, T., Gans, F., and Rosso, O. A.: Diagnosing the dynamics of observed and simulated ecosystem gross primary productivity with time causal information theory quantifiers. *PloS one*, 11, e0164960, <https://doi.org/10.1371/journal.pone.0164960>, 2016.
- Stosic, T., Stosic, B., and Singh, V. P.: Optimizing streamflow monitoring networks using joint permutation entropy. *J. Hydrol.*, 552, 306–312, <https://doi.org/10.1016/j.jhydrol.2017.07.003>, 2017.
- Stosic, D., Stosic, D., Stosic, T., and Stosic, B.: Generalized weighted permutation entropy. *Chaos, an Interdiscip. J. Nonlin. Sci.*, 32, <https://doi.org/10.1063/5.0107427>, 2022.
- Stosic, T. and Stosic, B.: Generalized weighted permutation entropy analysis of satellite hot-pixel time series in Brazilian biomes. *Physica A*, 636, 129548, <https://doi.org/10.1016/j.physa.2024.129548>, 2024.
- Stosic, T., Tošić, M., Lazić, I., da Silva Araújo, L., da Silva, A. S. A., Putniković, S., Djurdjević, V., and Stosic, B.: Changes in Rainfall Seasonality in Serbia from 1961 to 2020. *Theor. Appl. Climatol.*, 155(5), 4123–4138, <https://doi.org/10.1007/s00704-024-04871-4>, 2024a.
- Stosic, B., Djurdjević, V., Tošić, M., Lazić, I., Tošić, I., and Stosic, T.: Generalized weighted permutation entropy analysis of SPEI index in Serbia as a proxy of corn production. *Int. J. Climatol.*, 44, 3514–3528, <https://doi.org/10.1002/joc.8536>, 2024b.

- Sultan, B., Baron, C., Dingkuhn, M., Sarr, B., and Janicot, S.: Agricultural impacts of large-scale variability of the West African monsoon. *Agric. For. Meteorol.*, 128, 93–110, <https://doi.org/10.1016/j.agrformet.2004.08.005>, 2005.
- 450 Suriano, M., Caram, L. F., and Rosso, O. A.: Daily Streamflow of Argentine Rivers Analysis Using Information Theory Quantifiers. *Entropy*, 26, <https://doi.org/10.3390/e26010056>, 2024.
- Tongal, H.: Assessing climate vulnerability and nonlinear rainfall dynamics in complex networks. *Stoch. Environ. Res. Risk Assess.*, 39, 21–40, <https://doi.org/10.1007/s00477-024-02842-9>, 2025.
- Tošić, I. Spatial and temporal variability of winter and summer precipitation over Serbia and Montenegro. *Theoretical and Applied Climatology*, 77(1–2), 47–56. <https://doi.org/10.1007/s00704-003-0022-7>. 2004.
- 455 Tošić I, Hrnjak I, Gavrilov MB, Unkašević M, Marković SB, Lukić T: Annual and seasonal variability of precipitation in Vojvodina. *Theor Appl Climatol* 117:331–341. <https://doi.org/10.1007/s00704-013-1007-9>. 2014
- Tošić, I., da Silva, A. S. A., Filipović, L., Tošić, M., Lazić, I., Putniković, S., ... and Djurdjević, V.: Trends of Extreme Precipitation Events in Serbia Under the Global Warming. *Atmosphere*, 16, 436, <https://doi.org/10.3390/atmos16040436>, 2025.
- 460 Trenberth, K. E.: Changes in precipitation with climate change. *Clim. Res.*, 47, 123–138, doi:10.3354/cr00953, 2011.
- Vautard, R., Kadyrov, N., Iles, C., Boberg, F., Buonomo, E., Bülow, K., ... and Wulfmeyer, V.: Evaluation of the large EURO-CORDEX regional climate model ensemble. *J. Geophys. Res. Atmos.*, 126, e2019JD032344, <https://doi.org/10.1029/2019JD032344>, 2021.
- Vujadinović Mandić, M., Vuković Vimić, A., Ranković-Vasić, Z., Đurović, D., Ćosić, M., Sotonica, D., Nikolić, D., and 465 Đurđević, V.: Observed changes in climate conditions and weather-related risks in fruit and grape production in Serbia. *Atmosphere*, 13(6), 948, <https://doi.org/10.3390/atmos13060948>, 2022.
- Wang, H., Zhao, X., Guo, Q., & Wu, X.: A novel hybrid model by integrating TCN with TVFEMD and permutation entropy for monthly non-stationary runoff prediction. *Sci. Rep.*, 14, 31699, <https://doi.org/10.1038/s41598-024-81574-w>, 2024.
- 470 Xia, J., Shang, P., Wang, J., and Shi, W.: Permutation and weighted-permutation entropy analysis for the complexity of nonlinear time series. *Commun. Nonlinear Sci. Numer. Simul.*, 31, 60–68, <https://doi.org/10.1016/j.cnsns.2015.07.011>, 2016.

Yan, W., Zhou, Q., Peng, D., Wei, X., Tang, X., Yuan, E., Wang, Y., and Shi, C.: Soil moisture responses under different vegetation types to winter rainfall events in a humid karst region. *Environ. Sci. Poll. Res.*, 28, 56984–56995, <https://doi.org/10.1007/s11356-021-14620-z>, 2021.

475 Zhou, W., Liu, D., Zhang, J., Jiang, S., Xing, S., Wang, J., Cheng, Y., and Chen, N.: Identification and frequency analysis of drought–flood abrupt alternation events using a daily-scale standardized weighted average of the precipitation index. *Front. Environ. Sci.*, 11, 1142259, <https://doi.org/10.3389/fenvs.2023.1142259>, 2023.

Zunino, L., Zanin, M., Tabak, B. M., Pérez, D. G., and Rosso, O. A.: Complexity-entropy causality plane: A useful approach to quantify the stock market inefficiency. *Physica A*, 389, 1891–1901, <https://doi.org/10.1016/j.physa.2010.01.007>, 2010.