

# A theoretical framework for assessing and understanding to understand sources of error in Earth System Model emulation

Christopher B. Womack<sup>1,2</sup>, Glenn Flierl<sup>3</sup>, Shahine Bouabid<sup>3</sup>, Andre N. Souza<sup>3</sup>, Paolo Giani<sup>2,3</sup>, Sebastian D. Eastham<sup>4</sup>, and Noelle E. Selin<sup>2,3,5</sup>

<sup>1</sup>Department of Aeronautics and Astronautics, Massachusetts Institute of Technology, Cambridge, MA, United States

<sup>2</sup>Center for Sustainability Science and Strategy, Massachusetts Institute of Technology, Cambridge, MA, United States

<sup>3</sup>Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA, United States

<sup>4</sup>Brahmal Vasudevan Institute for Sustainable Aviation, Department of Aeronautics, Imperial College London, London, United Kingdom

<sup>5</sup>Institute for Data, Systems, and Society, Massachusetts Institute of Technology, Cambridge, MA, United States

**Correspondence:** Christopher B. Womack (cwomack@mit.edu) and Noelle E. Selin (selin@mit.edu)

**Abstract.** Full-scale Earth ~~system models~~ System Models (ESMs) are too computationally expensive to keep pace with the growing demand for climate projections across a large range of emissions pathways. Climate emulators, reduced-order models that reproduce the output of full-scale models, are poised to fill this niche. However, the large number of emulation techniques available and lack of a comprehensive theoretical basis to understand their relative strengths and weaknesses ~~compromises~~ <sup>5</sup> compromise fundamental methodological comparisons. Here, we present a theoretical framework that connects disparate emulation techniques ~~, using it to analyze and use it to understand potential~~ sources of emulator error focusing on memory effects, hidden variables, system noise, and nonlinearities. This framework includes popular emulation techniques such as pattern scaling and response functions, relating them to less commonly used methods, such as Dynamic Mode Decomposition and the Fluctuation Dissipation Theorem (FDT). To support our theoretical contributions, we provide practical implementation ~~details~~ <sup>10</sup> guidance for each technique, ~~, evaluating performance across a series of experiments designed to highlight different potential sources of error~~ Using pedagogical examples including idealized box models and a modified Lorenz 63 model, we illustrate the expected errors from each emulation technique considered. We find that response function-based emulators outperform other techniques, particularly pattern scaling, across all scenarios tested. ~~We additionally outline potential advantages of Potential benefits and trade-offs from~~ Incorporating statistical mechanics ~~into~~ <sup>15</sup> in climate emulation through the use of the FDT, though this technique requires greater computational resources and non-standard scenarios for training. Results highlight the relative utility of each technique are discussed, along with the importance of designing future scenarios for ~~Earth system models~~ ESMs with emulation in mind, ~~suggesting~~ . We argue that large-ensemble experiments utilizing the FDT could benefit climate modeling and impacts communities. We conclude by discussing optimal use cases for each emulator, along with implications for ESMs based on our pedagogical model results.

20 **1 Introduction**

Earth-System Models (ESMs) are our most comprehensive tool to simulate the climate system, yet their high computational cost limits the range and number of scenarios that can be investigated (Flato, 2011; Müller et al., 2018). Growing demand for high-quality climate projections which differ from the scenarios considered within the Coupled Model Intercomparison Project (CMIP) drives a need for computationally efficient alternatives (Eyring et al., 2016). Climate emulators - reduced order models that reproduce the outputs of full-scale climate models - have seen a surge in popularity as they can be many orders of magnitude faster than the parent models (Sudakow et al., 2022; Tebaldi et al., 2025). Their low computational costs also make them an appealing tool to disseminate climate information to audiences beyond the climate science community.

25 Chaotic sensitivity renders prediction of the climate state infeasible beyond short time horizons (Lorenz, 2006, 2015). Climate emulators must therefore target the statistics of climate variables, such as means, variances, or higher moments, rather than simulating chaotic dynamics (Beusch et al., 2020; Souza et al., 2024; Wang et al., 2025). Many emulation techniques exist to estimate the mean state and/or probability distribution of climate variables (Meinshausen et al., 2011; Castruccio et al., 30 2014; Herger et al., 2015; Tebaldi and Knutti, 2018; Leach et al., 2021; Watson-Parris et al., 2022; Addison et al., 2024; Bassetti et al., 2024; Bouabid et al., 2024), and in this work we explore methods that emulate the mean state of the system. In a recent 35 review, Tebaldi et al. (2025) distinguished between five main categories of climate emulators, including linear pattern scaling, statistical approaches, and machine learning algorithms. Following their categorization, we focus on linear pattern scaling and its immediate extensions along with dynamical system/impulse response theory emulators.

40 In the climate context, the most commonly used emulation technique is pattern scaling (Santer et al., 1990), a simple linear regression of local climate variables (e.g. temperature or precipitation anomaly) on the global mean temperature anomaly. Pattern scaling has been used and studied extensively since its development (Mitchell, 2003; Tebaldi and Arblaster, 2014; Wells et al., 2023; Giani et al., 2024), with variations that capture seasonal anomalies, different mixes of greenhouse-gases, and spatially heterogeneous forcings such as aerosols (Schlesinger et al., 2000; Herger et al., 2015; Mathison et al., 2024). This approach produces accurate projections assuming exponential and fixed-pattern forcing, linear feedbacks, and linear and time-independent dynamics, criteria that are roughly satisfied in a number of CMIP experiments (Giani et al., 2024). Memory 45 effects in overshoot scenarios (forcing history, rather than only instantaneous forcing, affecting a future state) violate these assumptions, causing this approach to break down for many decision-relevant scenarios.

Impulse response (~~response methods, commonly referred to as either response or~~ Green's ~~function~~) ~~methods~~ ~~functions~~, fill this memory effect gap by encoding forcing history into the emulator, rather than relying only on the instantaneous forcing. These techniques have been studied thoroughly in the contexts of dynamical systems and climate science (Joos and Bruno, 1996; Hasselmann et al., 1997; Lucarini et al., 2017; Orbe et al., 2018; Freese et al., 2024; Giorgini et al., 2024), and are an 50 active area of research (Winkler and Sierra, 2025). Response functions are popular due to their ease of interpretability and improvement in skill over pattern scaling in capturing realistic dynamics (Womack et al., 2025). Pure linear response functions cannot account for nonlinear effects, though hybrid schemes that incorporate machine learning (ML) may help resolve this issue (Winkler and Sierra, 2025).

Pattern scaling and linear response functions are prevalent in climate emulation literature, yet these approaches are only 55 two methods among a broad spectrum of emulators, with each technique offering trade-offs in terms of complexity, data requirements, and interpretability. For example, quasi-equilibrium emulation is closely related to pattern scaling, though only a handful of studies explore the utility of this principal beyond the traditional choice of global mean temperature as emulator input (Huntingford and Cox, 2000; Cao et al., 2015). Other techniques, such as Dynamic Mode Decomposition (DMD) and its variants, are generally not classified as emulators despite their potential to identify and predict modes of variability in the 60 climate system (Kutz et al., 2016; Gottwald and Gugole, 2020; Navarra et al., 2021; Mankovich et al., 2025).

We consider climate emulators as defined in Tebaldi et al. (2025), excluding Simple Climate Models (SCMs) and Earth system Models of Intermediate Complexity (EMICs), though they share similarities with emulators. We also do not examine ML 65 emulators such as FourCastNet and NeuralGCM – while these techniques are promising for weather prediction, they currently lack the stability required for reliable climate prediction (Pathak et al., 2022; Kochkov et al., 2024). Several studies have employed ML techniques to instead target the statistics of the climate, rather than weather (Lewis et al., 2017; Bassetti et al., 2024; Wang et al. 70 (Lewis et al., 2017; Bassetti et al., 2024; Wang et al., 2025; Bouabid et al., 2025), but these works focus on emulator implementation rather than theoretical analysis.

In this work, we develop a framework connecting a spectrum of emulators through the Koopman and Fokker-Planck operators, which govern the evolution of stochastic processes. In doing so, we identify a gap in the Tebaldi et al. (2025) emulator 75 typology: operator-based emulators, an area largely unexplored in existing climate emulator literature. While previous work has connected operator frameworks with the Fluctuation Dissipation Theorem and thus, linear response theory (Cooper and Haynes, 2011; Lucarini et al., 2017; Lembo et al., 2020; Zagli et al., 2024; Giorgini et al., 2025b), our contribution explicitly demonstrates its utility in the context of climate emulation. Section ??-2 first presents our theoretical framework (Seet. 2), highlighting that ~~emulators attempt the goal of many emulation techniques is~~ to simplify complex climate dynamics into a linear 80 set of modes associated with the Fokker-Planck and Koopman operators. We then apply this framework to identify potential sources of ~~emulator error (Seet. 2.3), comparing error within~~ six emulation techniques, ~~analyzing them~~ from both a theoretical and practical perspective ~~, providing a toolkit for those interested in emulating dynamical systems. We then (Sect. 2.3). In~~ Sect. 3, ~~we~~ introduce a series of experiments using simplified climate models and forcing scenarios designed to stress test and evaluate each emulator (Seet. ??-); ~~these experiments include box models and a modified version of the Lorenz 63 system.~~ Section 4 contains ~~experimental results the results of these simplified climate model experiments~~, showing that response functions consistently outperform other emulators across potential high-error scenarios. We conclude by discussing optimal use cases for each emulator ~~in~~, along with implications for ESMs based on our pedagogical model results (Sect. 5).

## 2 ~~Methods~~Theoretical framework for climate emulation

~~Section 2 outlines In this section, we outline~~ a theoretical framework for ~~analyzing emulators, while Sect. 2.3 provides theory and implementation~~ climate emulation based on the Koopman and Fokker-Planck operators. Section 2.1.1 introduces our ~~emulation target, a general, stochastic system, outlining potential sources of error when emulating this system. Section 2.2 then~~

formalizes two complementary emulation strategies: emulating the full probability distribution, or emulating a collection of statistical moments (e.g. mean, variance). We conclude this section by connecting theoretical and practical (i.e. implementation) details for the six emulators of interest. ~~We detail experiments and emulator evaluation protocol in Sect. ?? (Sect. 2.3).~~ See Fig.

90 2 for a conceptual roadmap of emulator theory and Table 1 for an overview of selected methods.

Throughout this section, we denote scalars with lowercase characters, vectors with lowercase, boldface italic characters, matrices with uppercase, boldface characters, and operators with script characters (e.g.  $\mathcal{N}$  or  $\mathcal{L}$ ). We use  $x$  and  $n_x$  to denote the spatial coordinate and its dimensionality, along with  $t$  and  $n_t$  to denote the temporal coordinate and its dimensionality. Our examples focus on climate anomalies relative to a background state, though these techniques are applicable to general chaotic 95 dynamical systems.

## 2.1 Theoretical framework for climate emulation ~~Problem setup~~

~~Section 2.1.1 introduces our emulation target, a general, stochastic system, outlining potential sources of error when emulating this system. Section 2.2 formalizes two complementary emulation strategies: emulating the full probability distribution, or emulating a collection of statistical moments (e.g. mean, variance).~~

### 100 2.1.1 Problem setup

A full-scale climate model is a deterministic, albeit chaotic, system. This chaos results in extreme sensitivity to initial conditions, requiring emulation of the system's statistics, rather than its dynamics (Lorenz, 2015). To understand the statistics of the system and how they may change over time, we follow Hasselmann (1976) in modeling the evolution of a single climate variable using a stochastic differential equation (SDE) (Fig. 2, box 1). We ~~treat assume time-scale separation between slow~~ 105 ~~climate processes (e.g., ocean, cryosphere, land vegetation) and other, faster sources of variability.~~


~~In this framework, the climate is regarded as the statistical mean of a process that appears stochastic in individual realizations. We treat variations occurring either on timescales shorter than climate change (such as short-term weather fluctuations as stochastic noise and interannual variability) or in different realizations as stationary, stochastic noise. This allows us to parameterize their influence on the long-term, chaotic behavior of the system statistics of the chaotic system:~~

$$110 \quad \frac{\partial w}{\partial t} = \mathcal{N}(w) + F(t) + \varepsilon \xi(t), \quad (1)$$

where  $w$  is the climate variable (or set of variables) of interest (e.g. temperature),  $F$  is an external forcing (e.g. CO<sub>2</sub>),  $\mathcal{N}$  is the operator governing the evolution of that variable ~~(under slow climate processes)~~,  $\xi$  is a white noise term ~~, and (aggregated fast effects, including weather and interannual variability)~~, and  $\varepsilon$  is the noise standard deviation. We consider variables of interest to be anomalies relative to some base state (e.g. temperature anomaly with respect to preindustrial conditions).  $\mathcal{N}$  may involve 115 both linear and nonlinear terms in one or several fields, and we cannot directly represent this operator; this parameterization aggregates the effects of processes such as heat and momentum transfers. The operator may also be influenced by variables we observe as well as unobserved hidden variables (e.g. aerosol forcing in a pattern scaling emulator with only global mean

temperature as an input). The noise standard deviation can also be state dependent, though we treat it as independent for this exploration.

120 Climate emulators approximate Equation 1, either implicitly (pattern scaling) or explicitly (Dynamic Mode Decomposition), rendering them vulnerable to several potential sources of error. Figure 1 provides an overview of the sources of error we consider across a range of scenarios: Errors can enter from the forcing if an emulator assumes only the instantaneous forcing is significant and not the forcing history (Fig. 1 (a) - memory effects in an overshoot scenario). The presence of hidden variables can lead to errors in some techniques (Fig. 1 (b) - localized aerosol effects when assuming well-mixed forcings), while other 125 techniques are sensitive to noise (Fig. 1 (c) - overfitting on internal variability). Finally, any linear emulation technique will break down in the presence of nonlinearities (Fig. 1 (d) - ice-albedo feedbacks).



**Figure 1.** Potential sources of emulator error by scenario; [emulator](#). Emulator errors shown here are meant for illustrative purposes only; [we introduce experiments which reproduce these errors in simplified climate models \(e.g. box models\) in Sect. 3.](#) (a) Pattern scaling emulator trained on *historical* and SSP585, tested against SSP119 in 2100; error over northern North America results from memory effects. (b) Pattern scaling emulator trained on *historical*, tested against SSP370 in 2050; error over India and SE Asia results from hidden variables (aerosols not contained in training data). (c) High-order polynomial pattern scaling emulator trained on *historical*, tested against SSP585 in 2020; error results from overfitting on internal variability. (d) Pattern scaling emulator trained on *historical*, tested against SSP585 in 2100; error results from nonlinear feedbacks in the Arctic. All ScenarioMIP data shown are taken from the MPI Grand Ensemble (O'Neill et al., 2016; Maher et al., 2019).

### 2.1.1 Operator framework for emulators

## 2.2 Operator framework for emulators

Our operator framework simplifies complex, possibly nonlinear climate dynamics into a linear set of modes with associated decay rates. We use the term operator to refer to an update rule that advances the system one timestep for a quantity of interest. An emulator attempts to approximate these modes, which are physically interpretable; for temperature, the decay rates correspond to heat-uptake timescales.

Table 1 summarizes emulation techniques discussed in this section, providing a short conceptual description of each method along with their key assumptions. We focus on linear emulation techniques that target the mean state of a climate variable: pattern scaling, the Fluctuation Dissipation Theorem (FDT), deconvolution, modal fitting, Dynamic Mode Decomposition (DMD), and Extended DMD (EDMD). The FDT, deconvolution, and modal fitting emulators are all response function-based emulators, while ~~E/EDMD~~ and DMD are operator-based emulators.

**Emulating a probability distribution.** Our governing system, Equation 1, simulates a variable of interest,  $w$ , forward in time under a stochastic forcing. The trajectory of the time evolution of  $w$  is characterized by the probability distribution,  $p(w, t)$ . We therefore focus our efforts on emulating  $p(w, t)$  via the Fokker-Planck operator. This is a mathematical tool to evolve the probability distribution of a stochastic system forward in time. As this operator is linear, emulating it is equivalent to approximating a series of eigenvalues and eigenfunctions.

As shown by Hasselmann (1976), the time evolution of  $p(w, t)$  is given by the Fokker-Planck equation corresponding to the governing SDE

$$145 \quad \frac{\partial}{\partial t} p(w, t) = -\frac{\partial}{\partial w} [p(w, t) (\mathcal{N}(w) + F(t))] + D \frac{\partial^2}{\partial w^2} p(w, t), \quad (2)$$


where  $D$  is a diffusion coefficient set by the noise term,  $D = \varepsilon^2/2$ . The Fokker-Planck equation describes how the probability density evolves in time and can be viewed as an advection-diffusion process.

Advection, which shifts the mean of  $p(w, t)$ , occurs due to the deterministic action of the governing operator and the external forcing. Because the advective term acts on the flux, it both shifts the mean and reshapes the density. Diffusion, which increases the variance in  $p(w, t)$ , is driven by system noise. Integrating Equation 2 forward diffuses the probability distribution, initially increasing the variance of  $w$  until balanced by the mean-reverting drift ( $\mathcal{N}(w) + F(t)$ ). It is common practice to write a Fokker-Planck equation directly from an SDE, as there exists a general relationship between any SDE and its corresponding Fokker-Planck equation; the full general derivation can be found in Denisov et al. (2009).

Importantly, the right hand side of Equation 2 is linear in the derivatives of  $w$ , allowing us to rewrite it in terms of the linear Fokker-Planck operator,  $\mathcal{F}$ ,

$$155 \quad \mathcal{F}(\cdot) = \frac{\partial}{\partial w} \left[ D \frac{\partial}{\partial w} (\cdot) - (\cdot) (\mathcal{N}(w) + F(t)) \right], \quad (3)$$

where the notation  $\mathcal{F}(\cdot)$  means the Fokker-Planck operator is acting on some arbitrary variable (in our case,  $p(w, t)$  in Equation 2). The Fokker-Planck operator (Fig. 2, box 3a) gives us a linear method to represent the time evolution of the probability



**Figure 2.** Conceptual flowchart for building an emulator through the joint Fokker-Planck/Koopman operator framework. Pop-outs show specific emulation techniques, while the shaded color indicates which concept a class of emulators relates to. Dashed arrows indicate conceptual/theoretical connections and solid arrows indicate a direct pathway. The overall process is as follows: (1) Select a climate variable of interest,  $w$ , such as temperature, here parameterized as the output of a stochastic differential equation. (2) Choose an emulation target, either the full probability distribution (option 1; 2a, 3a, 3c) or a statistical quantity such as the mean or variance (option 2; 2b, 3b, 3d). (3) Construct an emulator by selecting an approximation for either the Fokker-Planck or Koopman operator, including their response function representations; these options are connected through duality and are directly linked to linear response theory. (4) Given a new scenario of interest, emulate selected variable. [A summary of emulation techniques explored in this work \(right side of this figure\) can be found in Table 1.](#)

**Table 1.** Summary of emulation techniques discussed in this work including a short description and their key assumptions; a conceptual overview of these methods can be found in Fig. 2. Fluctuation Dissipation Theorem assumptions are shared with deconvolution and modal fitting emulation techniques. All techniques except the Fluctuation Dissipation Theorem additionally assume no hidden variables.

Technique	Short Description	Key Assumptions	Pros	Cons
Method I: Pattern Scaling <a href="#">(Pattern Scaling and its Immediate Extensions)</a>	Time-invariant pattern based on global mean temperature	Climate is always near equilibrium; response is instantaneous; fixed spatial pattern	<a href="#">Computationally efficient</a>	<a href="#">Structurally biased with irreducible errors</a>
Method II: Fluctuation Dissipation Theorem <a href="#">(Dynamical System/Impulse Response Theory)</a>	Response functions derived through perturbation ensemble experiments	Perturbations are small; data come from linear response regime	<a href="#">Gives interpretable physical response</a>	<a href="#">Requires nonstandard, computationally expensive scenarios</a>
Method III: Deconvolution <a href="#">(Dynamical System/Impulse Response Theory)</a>	Response functions solved for from any general experiment	Quasi-equilibrium initial condition; influence of noise is small	<a href="#">Applicable to any scenario</a>	<a href="#">Sensitive to noise, can give non-physical responses</a>
Method IV: Modal Fitting <a href="#">(Dynamical System/Impulse Response Theory)</a>	Response functions fit from any general experiment	Response is a decaying exponential; few significant modes	<a href="#">Applicable to any scenario</a>	<a href="#">Requires initial guess, can give non-physical responses</a>
Method V: Dynamic Mode Decomposition (DMD) <a href="#">(Operator-based Emulation)</a>	Approximating system dynamics with a linear operator	Dynamics are approx. linear; training data capture relevant dynamics	<a href="#">Gives interpretable spatiotemporal information</a>	<a href="#">Strong assumption of linearity</a>
Method VI: Extended DMD <a href="#">(Operator-based Emulation)</a>	Approximating system dynamics with nonlinear basis functions	Basis functions span Koopman operator; dynamics are approx. linear in new basis	<a href="#">Can theoretically reproduce any system behavior</a>	<a href="#">Requires selection of basis functions</a>

distribution. Linearity additionally allows us to decompose  $\mathcal{F}$  into eigenvalues and eigenfunctions (continuous eigenvectors).

160 These are the target of our emulator, and our emulator skill is directly proportional to how well it can approximate those eigenvalues and eigenfunctions, along with our estimate of  $p(w, 0)$ . This eigendecomposition is given by

$$\mathcal{F}f_{\mathcal{F}} = \lambda_{\mathcal{F}}f_{\mathcal{F}}, \quad (4)$$

where  $\lambda_{\mathcal{F}}$  denotes an eigenvalue and  $f_{\mathcal{F}}$  denotes an eigenfunction of the Fokker-Planck operator. The collection of  $\lambda_{\mathcal{F}}$  and  $f_{\mathcal{F}}$  fully characterizes the system's behavior. Our stochastic system evolves as a linear combination of probability distributions,

165  $f_{\mathcal{F}}$ , each decaying at rate  $\lambda_{\mathcal{F}}$ ; the real part of the eigenvalues controls the decay rate, while any imaginary components result in

oscillations over time. In the advection-diffusion analogy, each eigenfunction is a probability parcel that is carried and spread by the flow. The imaginary parts of the eigenvalues transport this parcel (shifting the mean) while the real parts act like an effective diffusivity (increasing the variance). This tells us which physical behaviors dominate and on what timescales they matter for climate prediction.

170 Unfortunately, in most cases we cannot obtain an explicit representation of the Fokker-Planck operator due to  $\mathcal{N}$  being nonlinear; see Appendix C for an analytic example of when this is possible. Because it acts on functions, the operator is infinite dimensional with infinitely many eigenpairs. This poses an immediate issue since computers have a finite amount of memory. Finite dimensional matrix approximations of the Fokker-Planck operator have been studied (often framed through the more general Perron-Frobenius operator) (Klus et al., 2016, 2018; Kaiser et al., 2019; Souza, 2024b, a; Souza and Silvestri, 175 2024), but require a large amount of data to reliably estimate the operator. For climate emulation this poses an additional issue, as generating large enough ensembles to resolve  $p(w, t)$  is prohibitively expensive. Because of these difficulties, little work exists studying the Fokker-Planck/Perron-Frobenius operator in the climate context (Navarra et al., 2021), though methods that reconstruct the full probability distribution of a climate variable using statistical methods (e.g. diffusion models and Gaussian processes) implicitly represent it (Bassetti et al., 2024; Bouabid et al., 2024; Wang et al., 2025).

180 **Emulating a statistical quantity.** In practice, it is often easier to emulate statistical quantities, such as the mean or variance of a climate variable. Many common emulation techniques (e.g. pattern scaling and response functions) target only the mean of a single variable (Herger et al., 2015; Wells et al., 2023; Freese et al., 2024), though other work extends this to approximate second order moments (Beusch et al., 2020; Wang et al., 2025). Relating these techniques requires the use of Koopman operator theory (Fig. 2, box 3b), a linear framework for propagating statistical quantities (usually referred to in the Koopman literature 185 as statistical observables) forward in time (Mezić, 2013; Otto and Rowley, 2021). Emulator studies rarely link their methods to Koopman theory, while literature that explicitly connects to the theory does not use the same emulator terminology (Slawinska et al., 2017; Navarra et al., 2021), though they accomplish similar prediction tasks. The Koopman operator allows for an exact representation of nonlinear dynamics using a linear operator, making it appealing when studying complex systems. We show how it can be used to emulate climate variables, simplifying nonlinear processes to the linear problem of emulating 190 physically-interpretable eigenvalues and eigenfunctions.

To derive the Koopman operator, we first define a general statistical quantity,  $g(w)$ , whose expectation,  $\langle \cdot \rangle$ , is given by

$$\langle g(w) \rangle = \int g(w) p(w, t) dw, \quad (5)$$

We then take the time derivative of this expression, moving the partial derivative inside the integral to act only on  $p$  since  $g(w)$  is independent of time. This allows us to substitute the resulting expression into the right hand side of Equation 2. Integrating 195 this by parts twice gives

$$\frac{\partial}{\partial t} \langle g(w) \rangle = \left\langle [\mathcal{N}(w) + F(t)] \frac{\partial}{\partial w} g(w) \right\rangle + D \left\langle \frac{\partial^2}{\partial w^2} g(w) \right\rangle, \quad (6)$$

where the diffusivity,  $D = \varepsilon^2/2$ , is identical to the Fokker-Planck case. This form allows us to define the Koopman operator,  $\mathcal{K}$ . It is linear in its derivatives of  $w$ , and we rewrite it as

$$\mathcal{K}(\cdot) = \mathcal{N}(w) \frac{\partial(\cdot)}{\partial w} + D \frac{\partial^2(\cdot)}{\partial w^2}, \quad (7)$$

200 where the notation  $\mathcal{K}(\cdot)$  means the Koopman operator is acting on some arbitrary variable ( $g(w)$  in Equation 7). Substituting this into Equation 6 gives

$$\frac{\partial}{\partial t} \langle g(w) \rangle = \langle \mathcal{K}g(w) \rangle + F(t) \left\langle \frac{\partial}{\partial w} g(w) \right\rangle, \quad (8)$$

205 This expression applies to any arbitrary statistical quantity (of which there are infinitely many), thus it can be used to integrate every statistical quantity forward in time; it is an alternate way to represent the complete probability distribution by representing each individual statistic. A useful choice is to select  $g(w) = w$ , giving

$$\frac{\partial}{\partial t} \langle w \rangle = \langle \mathcal{K}w \rangle + F(t), \quad (9)$$

which we will refer back to later.

Analogously to the Fokker-Planck operator, the Koopman operator provides a linear method to represent the time evolution of our entire collection of statistical quantities. As before, we can perform an eigendecomposition on the Koopman operator

210  $\mathcal{K}f_{\mathcal{K}} = \lambda_{\mathcal{K}} f_{\mathcal{K}}, \quad (10)$

where  $\lambda_{\mathcal{K}}$  denotes an eigenvalue and  $f_{\mathcal{K}}$  denotes an eigenfunction. The time evolution of our statistical quantity of interest is a linear combination of these eigenpairs. These can be used to identify dominant system dynamics and on what timescales they emerge. Training an emulator is equivalent to approximating eigenpairs; reproducing these pairs accurately emulates the behavior of the system.

215 However, approximations of the Koopman operator are limited by the same finite memory constraint as the Fokker-Planck case and deriving analytic solutions is dependent on the exact form of  $\mathcal{N}$ ; see Appendix C for an example of when analytic approximations are possible. Matrix approximations of the Koopman operator are nevertheless more prevalent than their Fokker-Planck counterparts (Schmid, 2010; Mezić, 2013; Williams et al., 2015; Otto and Rowley, 2021). Variants of these methods have recently been implemented in the climate context to identify dominant modes of variability in the system (e.g. 220 El Niño-Southern Oscillation or Pacific decadal oscillation) (Navarra et al., 2021, 2024; Mankovich et al., 2025), but have not been applied for the purpose of climate emulation. We outline two of these methods explicitly in Sect. 2.3.3.

**Two sides of the same coin.** The Koopman operator advances all statistical quantities of interest, and provides an alternative to the Fokker-Planck description of a distribution's time evolution. Knowing every statistic is equivalent to knowing the full distribution. Access to either operator fully characterizes our system, allowing us to emulate it. Mathematically, these operators 225 are dual (adjoint), where duality refers to two mathematical objects that contain alternate descriptions of the same information; this property is how we derived the Koopman operator in the previous section. This is analogous to, but physically and mathematically distinct from adjoint methods in climate modeling. There, adjoints to dynamics (rather than statistics as is the case for

the Koopman/Fokker-Planck approach) are exploited to calculate gradients with respect to input parameters more efficiently, which can be used to tune parameters and compute output sensitivities (Thuburn, 2005; Henze et al., 2007; Lyu et al., 2018).

230 Due to internal variability in the climate system, estimating the full probability distribution of a variable requires large initial condition ensembles, incurring significant computational cost. This is exacerbated for variables such as precipitation, where internal variability masks the forced response to a greater degree (Blanusa et al., 2023). Reliably estimating the full distribution at each timestep to approximate the Fokker-Planck operator from relatively coarse data is impractical. However, under additional assumptions of quasi-ergodicity, we bolster our sampling power by assuming that the statistics do not change 235 sufficiently quickly over a given time period. We thus focus on emulating lower-order statistical quantities, presenting those techniques in Sect. 2.3.

240 **Connecting to linear response theory.** Linear response theory states that the climate system's forced response (assuming perturbations are small) is encoded by a response function,  $R(t)$ . The response function is generated by the Koopman operator,  $\mathcal{K}$ , where each eigenpair of the operator determines the characteristic timescales of the system. Considering temperature 245 anomaly as an example variable, fast modes map to rapid atmospheric adjustments, while slow modes capture deep ocean heat uptake (Caldeira and Myhrvold, 2013). Response functions have been applied to a variety of climate problems (Joos and Bruno, 1996; Hasselmann et al., 2003; Joos et al., 2013; Orbe et al., 2018; Cimoli et al., 2023), including climate emulation (Freese et al., 2024; Womack et al., 2025; Sandstad et al., 2025), though often without addressing the formal response theory underlying these techniques. As was the case with the Koopman operator, more formal applications of response theory to 250 climate science often do not share the same language as climate emulators despite the shared goal of predicting the climate's forced response (Lucarini et al., 2017; Lembo et al., 2020; Zagli et al., 2024).

255 ~~Making~~ ~~To make~~ the relationship between response theory and the Koopman operator explicit in the context of emulation ~~requires the use of the~~, we first consider the system's dynamics to be governed by an operator,  $\mathcal{K}$ . When the system is subject to a small external perturbation, this operator can be split into an unperturbed component,  $\mathcal{K}_0$ , and the perturbation itself,  $\delta\mathcal{K}$ , such that  $\mathcal{K} = \mathcal{K}_0 + \delta\mathcal{K}$ . The expectation value of a statistical quantity  $g$  under the perturbed dynamics can be approximated to first order as the sum of its unperturbed evolution,  $\langle g \rangle_0$ , and a linear correction,  $\delta\langle g \rangle$ .

260 A general solution for this linear correction is provided by Ruelle's response theory. For systems in a statistical steady state (i.e., at equilibrium), this framework simplifies to the Fluctuation Dissipation Theorem (FDT) (Lucarini et al., 2025). The FDT describes how a system (e.g. the Earth system) responds to perturbations (anthropogenic CO<sub>2</sub> emissions) relative to some baseline state (preindustrial conditions). ~~Formally, it states that the response function associated with the statistical field  $g$~~  The change in the ensemble average field,  $\delta\langle g \rangle$ , is obtained by convolving a forcing,  $F(t)$ , with the system's response function,  $R(t)$

$$\delta\langle g \rangle = \int_{-\infty}^t R(s)F(t-s) ds. \quad (11)$$

260 Formally, the response function is calculated by computing the temporal autocorrelation with the between the statistical quantity  $g$  and the system's score function,  $s$ ,

$$R(t) = \langle g(t') = t'_s) s(t' = 0) \rangle, \quad (12)$$

where the score function of the steady-state distribution encodes how a small perturbation alters the system's statistics; see Giorgini et al. (2024, 2025b) for more details. The ~~change in the ensemble average field,  $\delta g$ , is then obtained by convolving with a forcing,  $F$ ,~~

265 
$$\langle \delta g \rangle = \int_{-\infty}^t R(s) F(t-s) ds.$$

---

The connection to Koopman operator theory is that temporal autocorrelations are expressed explicitly in terms of the Koopman operator, see [\(Zagli et al., 2024\)](#)[Zagli et al. \(2024\)](#).

270 Equation 11 is one way to state the Fluctuation Dissipation Theorem (FDT, Fig. 2, box 3d), a tool widely used in statistical mechanics and one of the main features of linear response theory (Lucarini et al., 2017; Lembo et al., 2020). The FDT predicts the first-order response of a statistical quantity due to external perturbations and is defined in terms of an ensemble average over a quantity of interest. As written, this form does not account for state- or time-dependent effects (i.e. one could consider the alternate formulation:  $R = R(w, t, t')$ ), though extensions to capture these effects and higher order statistical moments have been proposed [\(Metzler et al., 2018; Giorgini et al., 2025b; Winkler and Sierra, 2025\)](#)[\(Metzler et al., 2018; Giorgini et al., 2025a, b; Winkler and Sierra, 2025\)](#).

275 Response function emulators approximate the left hand side of Equation 12 using a variety of techniques, which we outline in more detail in Sect. 2.3.2. Their emulation goal is typically either to fit the eigenpairs which make up  $\mathcal{K}$  explicitly (Sandstad et al., 2025), or to find a direct representation of  $R(t)$  (i.e. an implicit representation of  $\mathcal{K}$ ) (Lembo et al., 2020; Freese et al., 2024; Womack et al., 2025). The former may be more easily interpretable through analyzing the explicit eigenpairs, while the latter offers flexibility in allowing for parametric forms other than a decaying exponential.

280 Response theory builds upon the operator frameworks presented in the previous sections by providing a method to illustrate how a given quantity responds to small changes in forcing. While the Fokker-Planck and Koopman perspectives offer complete characterizations of the statistics of the system over time, response theory offers a practical approach to use this information to predict how a quantity shifts under perturbations, described by the FDT.

### 2.3 Emulation under this framework [Connecting emulators to theory](#)

285 Following the framework from the previous section, we introduce several emulation techniques targeting the mean of a climate variable (Fig. 2, pop-outs on right hand side). We use the example of estimating the expected (or annual-average) temperature anomaly,  $T(\mathbf{x}, t)$ , given an external forcing,  $F(t)$  (e.g. CO<sub>2</sub> or other GHG emissions), though these techniques can be applied to any climate field. Each technique relates explicitly to the Fokker-Planck or Koopman operator and/or the Fluctuation Dissipation Theorem (FDT). We begin with methods that impose strong assumptions on the underlying data and progressively lift

290 those assumptions until we are left with the most general emulation techniques; headings follow the taxonomy of Tebaldi et al.  
 (2025) when possible.

### 2.3.1 Pattern scaling and its immediate extensions

**Method I: Pattern Scaling.** Pattern scaling is arguably the most well-known climate emulation technique (Santer et al., 1990; Mitchell, 2003; Tebaldi and Arblaster, 2014; Kravitz et al., 2017; Tebaldi and Knutti, 2018; Wells et al., 2023; Giani et al., 295 2024); it is formally derived via the Koopman operator, and is a specific case of a more general quasi-equilibrium emulation framework. It assumes that, at any given moment, the climate is in a quasi-equilibrium, rather than a transient, state and that changes in the forcing are small enough and/or the response of the system is fast enough to neglect system memory. Pattern scaling also assumes that the response does not depend on the background climate state, only the instantaneous forcing. Despite work showing that there are measurable differences between transient and quasi-equilibrium climate responses depending on 300 the transient warming rate (King et al., 2021), the success of pattern scaling has led to its continued use.

We first restate Equation 9 in terms of the quasi-equilibrium assumption and our climate variable of interest as

$$\frac{\partial}{\partial t} T(\mathbf{x}, t) = \mathcal{L}(\mathbf{x}, \mathbf{x}') T(\mathbf{x}', t) + F(t) \approx 0, \quad (13)$$

where  $\mathcal{L}$  indicates that this is no longer the true Koopman operator and  $\mathbf{x}$  and  $\mathbf{x}'$  indicate summation over spatial interactions, i.e. how one location,  $\mathbf{x}$ , is influenced by all other locations (including itself),  $\mathbf{x}'$ ; a more detailed description of the transition 305 from Equation 9 to 13 can be found in Appendix A4. We additionally assume  $T(\mathbf{x}, t)$  here refers to the ensemble mean temperature, which has the practical advantage of reducing the impact of internal variability on our emulator. Inverting this equation gives

$$T(\mathbf{x}, t) = -\mathcal{L}^{-1}(\mathbf{x}, \mathbf{x}') F(t), \quad (14)$$

which is a more general formulation of pattern scaling based on a generic forcing,  $F(t)$ . Alternate definitions of pattern 310 scaling have been explored previously, with a handful of studies developing extensions based on alternatives to global mean temperature such as radiative forcing or a combination of factors (Huntingford and Cox, 2000; Cao et al., 2015). A traditional pattern scaling formulation makes the further assumption that the forcing is the global mean temperature anomaly,  $F(t) = \bar{T}(t)$ , and replaces  $\mathcal{L}^{-1}$  with a low-order polynomial, leading to

$$T(\mathbf{x}, t) = a_0(\mathbf{x}) + a_1(\mathbf{x}) \bar{T}(t) + \frac{1}{2} a_2(\mathbf{x}) \bar{T}^2(t) + \dots, \quad (15)$$

315 where  $a_i(\mathbf{x})$  indicates the spatially varying pattern, and we typically keep only the first-order ( $a_1(\mathbf{x})$ ) term. Some work has explored the utility of higher-order terms, such as the quadratic term, but found it limited in extrapolative ability and physical justification (Herger et al., 2015).

Although pattern scaling implicitly attempts to approximate the Koopman operator - the perfect linear representation of the system - it is limited by its assumption of time-invariant, quasi-equilibrium dynamics. Truncating the operator with a 320 finite dimensional approximation and using only a single predictive field (here, annual-mean temperature) further reduces its

skill. Pattern scaling's inability to reproduce the pattern effect and other nonlinear/state-dependent feedbacks illustrates these limitations (Stevens et al., 2016; Giani et al., 2024). In Sect. 2.3.3, we explore alternative low-order approximations of the Koopman operator to resolve these issues.

Pattern scaling could be extended to the Fokker-Planck operator by shifting and rescaling the full probability distribution 325 based on global mean temperature, but this faces several limitations. Reliably estimating probability distributions requires large ensembles, which are computationally expensive. An alternate approach is to use long preindustrial control runs to generate the initial probability distribution and attempt to learn the linear scaling factor through the shorter SSP experiments. However, a simple linear shift may not capture scenario-dependent changes in the shape of the distribution; recent emulation work with Gaussian process regression suggests these distributional shifts may be complex (Wang et al., 2025). When applying pattern 330 scaling to the Fokker-Planck operator, we must also ensure the process does not violate the normalization of the distribution (i.e. the area under the curve must equal one).

We implement pattern scaling by calculating the global mean temperature anomaly and solving

$$\min_{a(\mathbf{x})} ||T(\mathbf{x}, t) - a(\mathbf{x})\bar{T}(\mathbf{x}, t)||^2. \quad (16)$$

~~Appendix A1 shows~~ In Appendix A1 we show that pattern scaling has two irreducible sources of error ~~when trained on~~ 335 ~~a ScenarioMIP-like forcing~~: (1) an equilibrium term, where pattern scaling converges to the wrong steady-state value when forcing plateaus and (2) a memory term, where pattern scaling breaks down when the system responds slowly compared to changes in the forcing. The ~~former stems from the mismatch between training pattern scaling in a transient regime and attempting to use it to project an equilibrium condition. The~~ latter cannot be accounted for within the pattern scaling framework, motivating the need for methods that explicitly capture memory.

### 340 2.3.2 Dynamical system/impulse response theory

Emulators that represent the climate system through response functions connect to fundamental principles of statistical mechanics and the Koopman/Fokker-Planck framework (Joos and Bruno, 1996; Hasselmann et al., 1997; Hasselmann, 2001; Lucarini et al., 2017; Orbe et al., 2018; Lembo et al., 2020; Fredriksen et al., 2021, 2023; Cimoli et al., 2023; Freese et al., 2024; Womack et al., 2025; Sandstad et al., 2025; Farley et al., 2025). Response function emulators relax the quasi-equilibrium 345 assumption, assuming instead that the current transient climate state is close to some baseline climate state that is in statistical equilibrium (generally preindustrial conditions). Perturbations to a field of interest are assumed to be small relative to magnitude of that field. These methods enable us to capture memory effects by integrating the entire forcing time history rather than only using the instantaneous forcing. One major benefit of this is that we can use them to represent regional shifts in surface warming patterns over time (the pattern effect) (Bloch-Johnson et al., 2024).

350 The use of different methods to derive response functions affects their utility as an emulator. A key assumption behind the Fluctuation Dissipation Theorem, for example, is that we have access to the governing equation, i.e. we are free to run large ensembles as needed. We begin this section assuming this is true, and relax this assumption later.

**Method II: The Fluctuation Dissipation Theorem.** In the case of a fully deterministic system with a zero initial condition,

simply forcing our system with a spatially explicit unit impulse ( $F(\mathbf{x}, t) = \delta(\mathbf{x}, t)$ ) is used to find the system's response function

355

$$T(\mathbf{x}, \mathbf{x}', t)|_{F(\mathbf{x}', t) = \delta(\mathbf{x}', t)} = R(\mathbf{x}, \mathbf{x}', t), \quad (17)$$

where perturbations are applied at each spatial location,  $\mathbf{x}'$ , to determine their influence on a location of interest,  $\mathbf{x}$ ; pulses can also be applied at alternate times,  $t'$ , to determine how different time lags impact the response (e.g. seasonality), but we neglect these effects to simplify our analysis.

360

In this case, we can derive our response function directly without the need for an ensemble of simulations, but real systems are not this simple. Utilizing an impulse forcing naively in a chaotic system may lead to a single realization with behavior far from the expected forced response. For our nonlinear SDE, we use the Fluctuation Dissipation Theorem (FDT), to calculate a response function from an ensemble. Our system's response to a perturbation of magnitude  $\varepsilon$  is given by

$$R(\mathbf{x}, \mathbf{x}', t) = \frac{\langle T_\varepsilon(\mathbf{x}, t) - T_0(\mathbf{x}, t) \rangle}{|\varepsilon(\mathbf{x}')|}, \quad (18)$$

365

where  $T_0(\mathbf{x}, t)$  and  $T_\varepsilon(\mathbf{x}, t)$  correspond to unperturbed and perturbed initial condition ensembles, respectively. More detail on this expression can be found in Marconi et al. (2008).

370

With this definition, we implement the Fluctuation Dissipation Theorem by first spinning up a simulation to get a steady state distribution from which we draw an ensemble of initial conditions,  $T_0(\mathbf{x}, t)$ . We then create a copy of the initial condition ensemble with an additional small perturbation,  $\varepsilon$ , applied to each member,  $T_\varepsilon(t)$ , and simulate every member from both ensembles for a scenario of interest. Applying Equation 18 then gives us the response function, which can be used to emulate a variable of interest by convolving it with a forcing from a new scenario (Equation 11).

375

Both the stochastic and deterministic approaches only yield an accurate estimate of the true response function when the system is perturbed from a quasi-equilibrium rather than a transient state. For climate models, this is typically done with step change CO<sub>2</sub> experiments after a spin-up period. This method is common in the literature around climate response functions and linear response theory (Lucarini et al., 2017; Lembo et al., 2020; Freese et al., 2024), though methods from the former two citations have not been applied to climate emulation and the latter does not reference formal response theory. Repeating this perturbation exercise at multiple background climate states can produce state-dependent response functions, but it is prohibitively expensive in practice.

380

Analogously to our discussion of using the Koopman vs. Fokker-Planck operator, there also exists an extension of the FDT to probability distributions. This relationship is given by

$$R(\mathbf{x}, \mathbf{x}', t) = -\langle T(\mathbf{x}, t) s(T(\mathbf{x}', 0)) \rangle, \quad (19)$$

where  $s(w) = \nabla \ln p(w)$  is the score function of the steady-state distribution and encodes how a small perturbation alters the system's dynamics; more details can be found in (Giorgini et al., 2024).

385

The score function captures the direction a distribution shifts in response to a perturbation, and correlating it with a climate variable explains how the expectation of that variable shifts. Appendix A5 outlines the link between this approach and the

Fokker-Planck operator. Analytical expressions for the score function are unavailable for most systems, necessitating machine learning techniques to learn the score function. This approach has achieved high skill in representing the response function for several systems (Giorgini et al., 2024), though it has not yet been applied to the full climate system. We do not explore it further in this work because of the machine learning infrastructure required to implement it.

390 The FDT faces accessibility issues in practice. First, there are high costs associated with this technique: a large ensemble of ESM runs is often prohibitively expensive. Second, there are also some configurations we simply cannot access: formal response theory assumes perturbations can be applied in a straightforward manner, which is not always the case. Because response functions are defined as a mapping from some perturbed input variable (e.g. CO<sub>2</sub> or radiative forcing) to an output variable of interest (e.g. temperature or precipitation), applying the FDT requires the ability to manually perturb a variable.

395 Climate models may not be configured to accommodate e.g. radiative forcing as an input. The FDT therefore cannot be applied to derive radiative forcing response functions, though this is possible through other methods (Womack et al., 2025).

**Method III: Deconvolution.** Without access to the true system to run specific perturbation experiments to find  $R(\mathbf{x}, \mathbf{x}', t)$ , data-driven approaches can estimate it. Deconvolution has been used to calculate response functions in the climate emulation context to derive spatially explicit response functions mapping effective radiative forcing to temperature (Womack et al., 2025).

400 It implicitly approximates the Koopman operator by deriving response functions that nominally correspond to Equation 11. To derive the deconvolution algorithm, we assume the data we have (e.g. annual temperature anomaly) are taken from an ensemble average of a general scenario. We begin from the FDT (Equation 11), assuming that our experiment begins from a quasi-equilibrium initial condition

$$T(\mathbf{x}, t) = \int_0^t R(\mathbf{x}, s) F(t - s) ds. \quad (20)$$

405 Treating this expression discretely, we rewrite it as a matrix expression and invert to solve for  $R(\mathbf{x}, t)$  from any general scenario

$$\mathbf{R} = \frac{\mathbf{F}^{-1} \mathbf{T}}{\Delta t}, \quad (21)$$

where  $\mathbf{F}$  is a lower-triangular matrix with  $F_{t=0}$  along the diagonal,  $F_{t=1}$  on the first off-diagonal, and so on (a Toeplitz matrix), and  $\mathbf{T}$  is a matrix of temperature values with rows corresponding to the time dimension and columns corresponding 410 to the spatial dimension. A more in-depth exploration of this process can be found in Womack et al. (2025). As written here, deconvolution aggregates spatial interactions (i.e. does not include an  $\mathbf{x}'$  term), cutting down on data requirements. Extensions of this procedure can account for spatial interactions, though they require additional experiments with varying spatial forcings.

In practice, noisy data require us to apply regularization to Equation 21 to ensure matrix stability. We instead solve

$$\min_{\mathbf{R}} \|\mathbf{RF} - \mathbf{T}\|^2 + \alpha \|\mathbf{R}\|^2, \quad (22)$$

415 where  $\alpha$  is the hyperparameter denoting the strength of our ridge regression. This simple ridge regression is equivalent to placing a Gaussian prior on the response function and assuming that the simulated temperature data we collect are corrupted

by Gaussian noise. We discuss the rationale of Gaussian noise further in Appendix B and outline our approach to tune the hyperparameter  $\alpha$  through *maximum a posteriori* optimization.

Deconvolution can be applied to any general scenario that begins from a quasi-equilibrium initial condition. However, since we require an explicit matrix inverse to perform deconvolution, it is sensitive to the frequency spectrum of the forcing data. If the eigenvalues of the matrix  $\mathbf{F}$  are very small (corresponding to near-zero frequencies) or the system is very noisy (corresponding to large differences in magnitudes between frequencies), the matrix becomes ill-conditioned, leading to an unstable response function. To illustrate these challenges, an explicit frequency-based derivation is included in Appendix A2. In practice, we regularize the system to avoid these issues (see Appendix B for details).

**Method IV: Modal Fitting.** Modal fitting is another data-driven technique to calculate response functions that retains some physical interpretability by explicitly representing the climate's response to a forcing as a series of decaying exponentials. The decay rates then represent the various timescales of the climate system (e.g. shallow vs. deep ocean heat uptake) and the modes represent how those timescales interact spatially. It has been used for tasks such as estimating effective radiative forcing and recently for climate emulation (Fredriksen et al., 2021, 2023; Sandstad et al., 2025).

To connect this approach to our framework, we begin from the same set of assumptions as deconvolution, but make the additional assumption that our response function is exactly a decaying exponential; in this case, our response function is exactly a Green's function as described in Appendix A3. We start from a restatement of Koopman response function definition (Equation 11)

$$G(\mathbf{x}, \mathbf{x}', t) = e^{\mathcal{L}(\mathbf{x}, \mathbf{x}')t}, \quad (23)$$

where  $\mathbf{x}$  and  $\mathbf{x}'$  track spatial interactions as before. We assume we can represent the Koopman operator with a finite, linear operator,  $\mathcal{L}$  (Appendix A4).

We then diagonalize the matrix  $\mathcal{L}$  though an eigenvalue decomposition, giving

$$G(\mathbf{x}, \mathbf{x}', t) = e^{v(\mathbf{x}, n)\Lambda(n, n)v^{-1}(n, \mathbf{x}')t}, \quad (24)$$

where  $\Lambda(n, n)$  and  $v(\mathbf{x}, n)$  are matrices containing the system's eigenvalues and eigenvectors, respectively, and  $n$  is the mode number. Since the matrix exponential respects similarity transformations, we rewrite this exactly as the summation

$$G(\mathbf{x}, \mathbf{x}', t) = \sum_{i=1}^k v(\mathbf{x}, n_i) e^{\lambda_i t} v^{-1}(n_i, \mathbf{x}'), \quad (25)$$

where  $k$  is equal to total the number of eigenvalues in the system. In the case of a climate model, the dimension of  $k$  is equivalent to the number of spatial dimensions. This may be much higher than the true number of modes that are significant in determining e.g. the temperature response of the system. Instead of the explicit form above, we typically see an alternate implementation, such as that in (Fredriksen et al., 2021, 2023) and (Sandstad et al., 2025). These show that one can fit an alternate form given simply by

$$G(t) \approx R(t) = \sum_{i=1}^3 \alpha_i e^{\lambda_i t}, \quad (26)$$

where using just three timescales (inter-annual, inter-decadal, and inter-centennial) is sufficient to represent the global mean behavior of the climate system; these methods specify a range/initial guess of timescales to initialize the optimization routine.

450 As we are implementing this at a grid cell level, we opt for a hybrid approach, given by

$$R_i(t) = \sum_{j=1}^3 \alpha_{i,j} e^{\lambda_i t}, \quad (27)$$

where  $i$  indicates the grid cell/region of interest, and  $j$  denotes the contribution from each timescale in a given region. We use the three timescales given above as the initial guess for each lambda, along with an initial guess for  $\alpha_{i,j} \forall i = j$ , assuming that one mode is dominant for each box.

455 We thus need to solve

$$\tilde{T}(\alpha_{i,j}, \lambda_i) = \int_{-\infty}^t R_i(s) F(t-s) ds, \quad (28)$$

$$\min_{\alpha_{i,j}, \lambda_i} \|T - \tilde{T}(\alpha_{i,j}, \lambda_i)\|^2. \quad (29)$$

For climate applications, the decay rates ( $\lambda_i$ ) can span several orders of magnitude, which are difficult for the optimizer to identify, even with normalization. This is exacerbated by the need to solve for the eigenvectors simultaneously, which are also

460 likely to have values that span several orders of magnitude; using more sophisticated optimization techniques than we apply in our test case could potentially resolve this issue. When implementing this algorithm, we follow Fredriksen et al. (2021), providing an initial guess of the correct order of magnitude to our optimizer.

Modal fitting has two major benefits. First, by truncating the leading modes, we reduce the dimensionality of the problem without the need for e.g. Empirical Orthogonal Functions (EOFs) or a Singular Value Decomposition (SVD). Second, we

465 require all  $\Re(\lambda_i) < 0$  (the real component of  $\lambda_i$ ) to ensure response functions to decay to zero as  $t \rightarrow \infty$ , a requirement not imposed on e.g. deconvolution and DMD. Because it is a best-fit problem, it naturally damps noise, making it well suited to systems with strong internal variability. However, this method can also be sensitive to local minima, requiring multiple iterations or a stochastic fitting procedure to alleviate this issue. Fitting may also be expensive on fine grids, since the number of eigenpairs scales with grid size, though we may not require all eigenpairs to accurately emulate the system.

#### 470 2.3.3 Operator-based emulation

The most general class of emulators are those that aim to directly approximate the Koopman operator. Every previous emulator can be thought of as a specific case of this general operator framework. Tebaldi et al. (2025) do not include operator-based emulators in their classification, as they are not typically referred to explicitly as emulators. However, we classify them as such to facilitate communication across disciplines with similar prediction goals.

475 The most common data-driven approximations of the Koopman operator are Dynamic Mode Decomposition (DMD) and Extended DMD (EDMD) (Schmid, 2010; Williams et al., 2015). Schmid (2010) developed DMD to extract dynamic information from fluid flows, and it has since been used to identify dominant modes of variability within the climate system, including

El Nino–Southern Oscillation, North Atlantic Oscillation, and Pacific Decadal Oscillation (Kutz et al., 2016; Gottwald and Gugole, 2020; Navarra et al., 2021; Franzke et al., 2022; Navarra et al., 2024; Mankovich et al., 2025). Under specific conditions, 480 DMD provides a finite-dimensional approximation of the Koopman operator (Schmid, 2022). EDMD expands this idea to approximate Koopman eigenvalues and eigenfunctions directly (Williams et al., 2015). The bulk of the work surrounding EDMD is theoretical (Haseli and Cortés, 2019; Netto et al., 2021), as in practice it has several limitations that we outline later in this section.

**Method V: Dynamic Mode Decomposition (DMD).** DMD assumes that the climate response is linear in  $w$  with respect 485 to an operator. If this is the true Koopman operator, this assumption holds by definition, provided it acts on the entire infinite space of statistical climate fields,  $g(w)$ . In practice, this leads to limitations based on how accurate the assumption of linearity is, which depends on the choice of variables; this approximation may hold better for a variable such as temperature, rather than precipitation. To derive DMD, we begin from Equation 9 applied to our variable of interest

$$\frac{\partial}{\partial t} T(\mathbf{x}, t) = \mathcal{K}(\mathbf{x}, \mathbf{x}') T(\mathbf{x}', t) + F(\mathbf{x}, t). \quad (30)$$

490 DMD assumes that we separate our data in discrete snapshots,  $T(\mathbf{x}, t_0), T(\mathbf{x}, t_1), \dots, T(\mathbf{x}, t_n)$ , which we assume are linearly related

$$T_{n+1} = \mathcal{L}T_n + F_n, \quad (31)$$

where we have used the subscript  $n$  as shorthand for  $t_n$  and omitted the spatial dimension for conciseness. By discretizing, 495 we are no longer solving for the exact Koopman operator (as in the previous case), which we now denote  $\mathcal{L}$ . This notation is standard in DMD literature. The traditional DMD algorithm assumes autonomous dynamics, omitting the forcing term. Equation 31 is referred to as DMD with control (DMDc) (Proctor et al., 2016), and has only recently been studied in the climate context (Mankovich et al., 2025).

To implement DMD, we collect our snapshots into matrices and invert this system, solving for  $\mathcal{L}$

$$\mathcal{L} = [\mathbf{T}_{n+1} - \mathbf{F}_n] \mathbf{T}_n^+, \quad (32)$$

500 where the superscript  $+$  denotes the Moore-Penrose pseudo-inverse of a matrix (required as it unlikely  $\mathbf{x}$  and  $t$  will be the same dimension, i.e. it is unlikely  $\mathbf{T}$  is a square matrix) and  $\mathbf{F}$  denotes a forcing matrix with the same dimension as our data; assuming well-mixed forcing means each row is identical in the forcing matrix. This is the simplest form of DMD, though in practice the Singular Value Decomposition (SVD) is often used to further reduce the dimensionality of the problem. This also increases the algorithm’s robustness relative to real-world systems that are subject to noise (Schmid, 2010).

505 This approach suffers mainly from its strong assumption of linear dynamics, which can break down for complex systems. Its success in identifying the dominant modes of variability in the climate suggests it may have utility as an explicit emulation technique (Kutz et al., 2016; Gottwald and Gugole, 2020; Franzke et al., 2022); future work will apply DMD to a full scale climate model to test this hypothesis. Unfortunately, DMD only provides a reliable estimate for the Koopman operator if it acts on a large set of statistical fields (more than simply the temperature anomaly when considering the full climate system)

510 and/or the dynamics governing the evolution of that quantity (or quantities) is linear, which is not the case in general. While the dynamics producing the base climate state are nonlinear, the success of methods such as pattern scaling suggest the dynamics of anomalies may be close to linear. DMD assumes all hidden variables are accounted for and the observed quantities fully describe the (linear) dynamics of our anomaly of interest. For example, the atmospheric temperature may be significantly influenced by heat uptake in the deep ocean, which, if it is not explicitly accounted for, will lead to errors when applying  
515 DMD. This motivates the need for a better algorithm for approximating the Koopman operator.

**Method VI: Extended DMD (EDMD).** As the baseline DMD algorithm is only able to approximate the Koopman operator in specific contexts, EDMD instead frames the problem such that we are deliberately trying to approximate the eigenvalues and eigenfunctions of the Koopman operator. This, ideally, leads to more reliable approximation than DMD and thus, a better emulator.

520 EDMD was introduced by Williams et al. (2015) as an explicit attempt to approximate the Koopman operator. The EDMD procedure involves projecting variables of interest into a higher dimensional space that has a richer representation of the system dynamics. As an example, we consider the problem of emulating precipitation anomaly using global mean temperature anomaly as the forcing. Precipitation may depend on the global mean temperature,  $\bar{T}(t)$ , but it also may depend on higher order or nonlinear terms, such as  $(\bar{T}(t))^2$ ,  $\cos(\bar{T}(t))$ ,  $\tanh(\bar{T}(t))$ , etc. To implement EDMD, the user must select a set of  
525 basis functions,  $\phi(\cdot)$ , such as these, that provide a better representation of the system dynamics than in the purely linear DMD case. Typical choices of basis functions as described by the original EDMD manuscript are Hermite polynomials, radial basis functions, and discontinuous spectral elements (Williams et al., 2015).

After choosing a set of basis functions, the EDMD problem statement is exactly the same as the original DMD algorithm. Solve for  $\tilde{\mathcal{K}}$  from

$$530 \quad \phi(T_{n+1}) = \tilde{\mathcal{K}}\phi(T_n) + \psi(F_n), \quad (33)$$

where  $\psi(\cdot)$  is the basis chosen for the forcing, and can be the same or different than the forcing for the quantity of interest. We use  $\tilde{\mathcal{K}}$  here as we are explicitly trying to approximate the Koopman operator. We ensure the basis includes the physical field of interest, e.g.  $\phi(\mathbf{T}) = [\mathbf{T}, \mathbf{T}^2, \mathbf{T}^3, \dots]$ , where the first entry is the physical field. As in the case with DMD, we solve this as

$$\tilde{\mathcal{K}} = [\phi(\mathbf{T}_{n+1}) - \psi(\mathbf{F}_n)] \phi^+(\mathbf{T}_n), \quad (34)$$

535 which we can use an SVD to solve more efficiently and reduce the influence of noise on the system. When applying this method, we first use Equation 33 with an appropriate initial condition to emulate the solution in our high-order basis. We then must project our solution back into physical space. Since we chose our basis to include the original physical coordinate, this is done by truncating the emulator output and keeping only the entries corresponding to  $\mathbf{T}$ .

This method has seldom been applied to climate problems (Navarra et al., 2024), likely due to the limitations acknowledged  
540 in Navarra et al. (2021), particularly the dimensionality of the problem. For a full climate model, DMD requires a matrix solve of dimension  $(N_{\text{lat}} \times N_{\text{lon}})^2$  for a single variable, which is extremely costly. In the case of EDMD, this dimension grows with every basis function used. To accurately represent the Koopman operator for the climate system, we potentially require

many more variables and many basis functions, causing the problem to rapidly increase in complexity, though this may be alleviated by emulating EOFs rather than gridded data. As with DMD, EDMD implicitly assumes no hidden variables, though 545 the choice of basis function can help alleviate this issue; e.g. if the hidden variables are higher order terms, EDMD may be able to represent them accurately. The selection of basis functions typically requires some experimentation though, as it can be difficult to predict which set of functions will be best suited for a given application; exploiting physical relationships such as the logarithmic relationship between CO<sub>2</sub> concentration and temperature may help alleviate this issue, however. More work is required to fully characterize the utility of EDMD for the climate system.

## 550 2.4 Experimental overview

### 3 Experimental overview

We conduct four experiments to evaluate these emulation techniques across a range of potential high-error scenarios. Here 555 we outline a set of experiments which reproduce the sources of error seen in Fig. 1, using them to evaluate the emulation techniques introduced in Sect. 2.3. We outline a climate box model with a simple local energy balance ODE in Sect. 3.1 and Sect. 3.2, followed by a nonlinear, cubic Lorenz system in Sect. 3.3. Experiments using these two simple models highlight the following potential sources of error: (1) memory effects, Fig. 1 (a); (2) hidden variables, Fig. 1 (b); (3) noise, Fig. 1 (c); (4) weak nonlinearities, Fig. 1 (d). We then describe forcing scenarios applied to each system in Sect. 3.4.

#### 3.0.1 Experiments 1 and 2: Climate Box Model

##### 3.1 Experiments 1 and 2: Climate Box Model

560 A classical box model is a standard, easily-interpretable model for temperature evolution. We use this idealized box model as it is the simplest system that includes the pattern effect and it is not necessarily meant to replicate CMIP experiments. We assume the form of this model is given by a simple local energy balance

$$C(\mathbf{x}) \frac{\partial T(\mathbf{x}, t)}{\partial t} = \lambda(\mathbf{x})T(\mathbf{x}, t) + R(\mathbf{x}, t) + \nabla \cdot \mathbf{F}(\mathbf{x}, t), \quad (35)$$

similar to Armour et al. (2013) and Giani et al. (2024).  $C(\mathbf{x})$  is the local effective heat capacity,  $T(\mathbf{x}, t)$  is the local temperature 565 anomaly,  $\lambda(\mathbf{x})$  is the local feedback parameter,  $R(\mathbf{x}, t)$  is the forcing function, and  $\nabla \cdot \mathbf{F}(\mathbf{x}, t)$  is the anomaly in heat flux divergence; parameters for this model are listed in Table 2. Furthermore, we assume that the forcing function can be linearly decomposed as a constant-amplitude spatial pattern and a variable time series:  $R(\mathbf{x}, t) = r(\mathbf{x})R(t)$ .

We consider two configurations for our box model. The first corresponds to a horizontally coupled three box system representing atmospheric boxes over land, low-latitude ocean, and high-latitude ocean;  $\nabla \cdot \mathbf{F}(\mathbf{x}, t) = -k(\mathbf{x})\nabla T(\mathbf{x}, t)$ . We assume a 570 constant diffusivity and discretize as,  $\nabla \cdot \mathbf{F}(\mathbf{x}, t) = -k(T_{i+1}(t) - T_i(t))$ , where  $i$  refers to the index of each box. We assume uniform forcing into each box, and use this configuration for experiments one and three (memory effects and noise; noise details can be found in Sect. 3.2). The second configuration corresponds to a vertically coupled two box system representing the atmosphere and the ocean; this has the same form as the previous case, with the caveat that there is no forcing applied

into the oceanic box. We use this configuration for experiment two (hidden variables). We begin this system from a zero initial

575 condition, aiming to simulate the temperature anomaly, rather than the absolute temperature.

**Table 2.** Parameters for the three box model, adapted from Giani et al. (2024). The heat capacity of each box is given in terms of the effective water depth,  $h(\mathbf{x})$ :  $C(\mathbf{x}) = \rho_w c_w h(\mathbf{x})$ , where  $\rho_w$  and  $c_w$  are the density and specific heat capacity of water, respectively. *Land*, *Low*, and *High* refer to atmospheric boxes over land, low-latitude ocean, and high-latitude ocean, respectively.

Parameter	Symbol	Land	Low	High
Effective Water Depth (m)	$h(\mathbf{x})$	5	150	1500
Local Feedback ( $\text{Wm}^{-2}\text{K}^{-1}$ )	$\lambda(\mathbf{x})$	-0.86	-2.0	-0.67

### 3.1.1 **Experiment 3: Noisy Box Model**

## 3.2 **Experiment 3: Noisy Box Model**

As the default configuration for our box model is purely deterministic, we add a stochastic noise term to the forcing to replicate the impact of inter-annual variability on the real climate system. To ensure the impact of this variability is similar to that of

580 the true system, we use CMIP6 *piControl* experiments to estimate the magnitude of the variability. Namely, we compute the standard deviations of *piControl* runs for three climate models (ACCESS-ESM1-5, MIROC6, MPI-ESM2-LR) and set the magnitude of the variability as the multi-model average  $\sigma = 0.117\text{K}$  (Dix et al., 2023; Tatebe and Watanabe, 2023; Wieners et al., 2023).

### 3.2.1 **Experiment 4: Cubic Lorenz System**

## 585 3.3 **Experiment 4: Cubic Lorenz System**

As the previous experiments are all defined by an operator which is linear in the quantity of interest, we additionally implement a weakly nonlinear, cubic Lorenz system. This provides a representation of the atmosphere that includes chaos, allowing us to

590 test the limits of these emulation techniques. In the standard Lorenz equations that represent a simplified model of atmospheric convection (Lorenz, 1963), the steady state is a linear function of  $\rho$ , and the mean heat flux ( $\langle XY \rangle = \langle Z \rangle$ ) is very nearly linear (Souza and Doering, 2015). We modify the system to the cubic form shown below to illustrate another failure mode of simple pattern scaling: the quasi-equilibrium value may not be a linear function of the forcing.

The cubic Lorenz equations are defined by the system

$$\frac{\partial}{\partial t} X = \sigma(Y - X), \quad (36)$$

$$\frac{\partial}{\partial t} Y = -(Z + \alpha Z^3)X + \rho(t)X - Y, \quad (37)$$

$$\frac{\partial}{\partial t} Z = XY - \beta Z, \quad (38)$$

with  $\alpha = 1/1000$ . The steady-state mean of both  $X$  and  $Y$  are zero, while the steady-state behavior of  $\langle Z \rangle$  is determined by  $\rho(t)$ . Values for  $\rho(t)$  are chosen such that nonlinearities are weak, as all linear methods are expected to break down in the presence of strong nonlinearities. These vary between experiments and are outlined in Table 4. We initialize this system through an initial condition ensemble starting from  $\rho(t) = 28$  with white noise applied to perturb the starting positions of each 600 ensemble member.


### 3.3.1 **Scenarios**

## 3.4 Scenarios

We consider four scenarios of interest for both the box model and cubic Lorenz system, focusing on scenarios which have CMIP analogues: (1) *Abrupt*, an abrupt increase in forcing, (2) *High Emissions*, an exponential increase in forcing, (3) *Plateau*, an exponentially increasing in forcing that levels off, and (4) *Overshoot*, a forcing that sharply increases and decreases. Descriptions of each scenario are given in Table 3. Figure 3 shows ODE-integrated solutions for each scenario in each experiment, and descriptions of experimental parameters can be found in Tables 4 and 5.

**Table 3.** Conceptual overview of forcing scenarios considered in this work. These scenarios are used in all experiments outlined in Sect. 223, and lists of experiment-specific parameters for each scenario can be found in Tables 4 and 5.

Scenario	Short Description
<i>Abrupt</i>	An abrupt doubling of CO <sub>2</sub> concentration; corresponds roughly to the <i>Abrupt2xCO<sub>2</sub></i> CMIP experiment.
<i>High Emissions</i>	An exponential increase of CO <sub>2</sub> concentration in time; corresponds roughly to <i>SSP585</i> .
<i>Plateau</i>	An increase in CO <sub>2</sub> concentration in time that follows a hyperbolic tangent, increasing exponentially and then tapering off; corresponds roughly to <i>SSP245</i> .
<i>Overshoot</i>	An increase in CO <sub>2</sub> concentration in time that follows a Gaussian profile, increasing and decreasingly rapidly; inspired by <i>SSP119</i> , but decreases more quickly.



**Figure 3.** ODE-integrated solutions for the three box model (top), two box model (middle), and cubic Lorenz system (bottom) for the (from left to right) *Abrupt*, *High Emissions*, *Plateau*, and *Overshoot* scenarios.  $D = 0.55 \text{ [Wm}^{-2}\text{K}^{-1}\text{]}$  for the three box experiment and  $D = 0.7 \text{ [Wm}^{-2}\text{K}^{-1}\text{]}$  for the two box experiment. For the cubic Lorenz problem we show the mean value of  $Z$  over 5,000 ensemble members as a line, and the shaded region indicates its standard deviation. Values shown are anomalies relative to a baseline of  $T = 0$  (experiments one through three) or  $\rho = 28$  (experiment four).

### 3.5 Evaluation

To evaluate each emulation technique, we utilize Normalized Root Mean Square Error (NRMSE, Equation 39) given as a

610 percentage, as our primary evaluation metric:

$$\text{NRMSE} = \frac{100}{\bar{g}(w_k)} \sqrt{\frac{\sum_{k=1}^{N_{\text{years}}} (g(w_k) - \hat{g}(w_k))^2}{N_{\text{years}}}}. \quad (39)$$

$\bar{g}(w_k)$  indicates the mean of our quantity of interest over the period error is calculated over. We calculate NRMSE with respect to the entire time series. To compare performance across training datasets, we train each emulator on one scenario at a time, testing against the others which are held out from the training (e.g. train on *Abrupt* and test on *High Emissions*).

615 We implement an alternate protocol for the cubic Lorenz system as there is no ground-truth to compare with due to chaos. Instead, we compare the skill of each emulator when training on only a subset of the ensemble members for that experiment. For example, given  $n_{\text{ensemble}}$  ensemble members for a given experiment, we construct a subset of  $n$  ensemble members without

**Table 4.** Forcing scenarios for each experiment, with the upper half of each row corresponding to the box model and the lower half of each row corresponding to the cubic Lorenz system. Parameters for the box model experiments are based on Giani et al. (2024) and (Armour et al., 2013) and parameters for the cubic Lorenz system are chosen such that the system exhibits weakly nonlinear behavior.  $H(t)$  is the Heaviside step function, and parameters for these scenarios are listed in Table 5.

Scenario	Functional Form
<i>Abrupt</i>	$F(t) = F_{abr} H(t)$
	$\rho(t) = \rho_{0,abr} + \rho_{1,abr} \tanh(t - \eta_{abr})$
<i>High Emissions</i>	$F(t) = F_{high} \exp(t/\tau_{high})$
	$\rho(t) = \rho_{0,high} + \rho_{1,high} \exp(t/\eta_{high})$
<i>Plateau</i>	$F(t) = F_{plat} + F_{plat} \tanh(\omega_{plat}(t - \tau_{plat}))$
	$\rho(t) = \rho_{0,plat} + \rho_{1,plat} \tanh(\omega_{plat}(t - \tau_{plat}))$
<i>Overshoot</i>	$F(t) = F_{over} \exp(-(t - \tau_{over})^2/(2\sigma^2))$
	$\rho(t) = \rho_{0,over} + \rho_{1,over} \exp(-(t - \eta_{over})^2/(2\sigma^2))$

**Table 5.** Scenario parameters used for the experiments in this study. Values for  $\rho_0$  are listed in the order *Abrupt*, *High Emissions*, *Plateau*, and *Overshoot*. Box-model parameters have physical units to output temperature; the cubic-Lorenz parameters are dimensionless.

Box Model		Cubic Lorenz System	
Parameter	Value	Parameter	Value
-	-	$\rho_0$	[45, 28, 40, 28]
$F_{abr}$	$3.7 \text{ W m}^{-2}$	$\eta_{abr}$	10
		$\rho_{1,abr}$	17
$F_{high}$	$\frac{8.5 \text{ W m}^{-2}}{\exp(\tau_f/\tau_{high})}$	$\rho_{1,high}$	$\frac{30}{\exp(\eta_f/\eta_{high})}$
$\tau_f$	250 yr	$\eta_f$	250
$\tau_{high}$	50 yr	$\eta_{high}$	50
$F_{0,plat}$	$2.25 \text{ W m}^{-2}$	$\rho_{1,plat}$	$\frac{12}{\tanh(5)}$
$F_{1,plat}$	$\frac{2.25 \text{ W m}^{-2}}{\tanh(\omega_{plat}\tau_{plat})}$	$\tau_{plat}$	150
$\omega_{plat}$	$1/50 \text{ yr}^{-1}$	$\omega_{plat}$	$1/50$
$F_{over}$	$4 \text{ W m}^{-2}$	$\rho_{1,over}$	30
$\tau_{over}$	200 yr	$\eta_{over}$	200
$\sigma_{over}$	42.47	$\sigma$	50

replacement, where  $n = 1 : n_{\text{ensemble}} - 1$ , and train our emulator from that subset. We then test the emulator’s skill in emulating the mean response given the ensemble average forcing. We repeat this subsampling exercise 10 times, recording the average

620 performance over those trials. For the noisy three box model, we use the same protocol, additionally presenting the ground truth of emulating the noiseless three box model.


## 4 Results

Section 4.1 presents a summary of results across each of the emulation techniques outlined in Sect. 2.3 when emulating the simplified climate systems presented in Sect. 2.3, with subsequent sections highlighting key results from individual experiments. Section 4.2 contains the results for the three box model with significant memory effects (Fig. 1 (a)); the three boxes represent atmospheric boxes over the land, low-latitude ocean and high-latitude ocean. We then report emulator performance on the restricted two box model in Sect. 4.3. In this case we highlight the issue of hidden variables (Fig. 1 (b)) by only giving the emulators access to the temperature anomaly in only one of the two boxes during training; the two boxes represent an atmospheric and oceanic box (forcing only into the atmosphere). This is followed by a version of the three box model with a stochastic forcing to test the robustness of each method to noise (Fig. 1 (c)). Finally, we showcase results for the nonlinear, cubic Lorenz system in Sect. 4.5 (Fig. 1 (d)), which tests emulator performance in the presence of chaos and weak nonlinearities. In the case of models with multiple regions (boxes), we present only a single evaluation score, as relative performance across boxes was consistent for all cases analyzed.

### 4.1 Overall emulator performance

635 Figure 4 summarizes emulator performance in terms of Normalized Root Mean Square Error (NRMSE) across all four experiments. For each experiment, there are four possible train/test scenarios (*Abrupt*, *High Emissions*, *Plateau*, and *Overshoot*). We test on one scenario and train against the remaining three, showing median NRMSE over all train/test combinations. For experiments two and four, the pattern scaling emulator is trained to map forcing to quantity of interest, as these experiments do not have a global mean temperature equivalent. Results for deconvolution are shown using the regularization presented 640 in Appendix B. Error values are calculated with a constant 40 ensemble members for experiment three and 4,000 ensemble members for experiment four.

645 Response function based emulators (the FDT, deconvolution, and modal fitting methods) generally outperform other approaches, demonstrating consistently lower NRMSE across most experiments. The FDT is particularly reliable relative to all other methods, yielding consistently low errors across all four test cases, indicating its robustness regardless of scenario; while it has higher error in the cubic Lorenz case, this is primarily a function of ensemble size (see Sect. 4.5). As FDT response functions are, in principle, equation-driven rather than data-driven, they provide the perfect solution given a linear system (experiments one through three) or enough realizations (experiment four). Deconvolution similarly performs well across all experiments, while modal fitting has high performance in experiments one, two, and three; both of these methods exhibit higher errors in experiment four. For deconvolution, this is due to its sensitivity to noise as discussed in Sect. 2.3.2, while modal fitting suffers because of an inability to reliably separate timescales and the need for an accurate initialization for its unknown parameters, which we discuss in Sect. 4.4.




**Figure 4.** Summary of emulator performance over all experiments considered in this work. For each experiment, there are four scenarios. We show the median NRMSE value across all scenario train and test combinations, excluding the trivial case of training and testing on the same dataset. Error values are calculated with 40 ensemble members for experiment three and 4,000 ensemble members for experiment four. Emulator abbreviations are as follows: PS - Pattern Scaling, FDT - Fluctuation Dissipation Theorem, Deconv. - Deconvolution, Modal - Modal Fitting, DMD - Dynamic Mode Decomposition, EDMD - Extended DMD. Diagonal hatching indicates response function emulators, while cross hatching indicates operator-based emulators.


In contrast, pattern scaling consistently underperforms, exhibiting the highest error in all experiments except for the cubic Lorenz case. This is most likely due to the presence of strong memory effects in the box models, which pattern scaling cannot capture by definition. DMD and EDMD outperform pattern scaling in experiments one and two, but exhibit much more variable 655 performance in experiments three and four. For the first three experiments, DMD and EDMD produce identical results. This is because the models in these experiments are purely linear, and the use of any higher-order basis for EDMD leads to a drop in skill. These methods struggle with the noisy three box model, and more in-depth results can be found in Sect. 4.4. While theory suggests DMD/EDMD would not be well-suited for the restricted two box problem due to the presence of hidden variables, they outperform pattern scaling in practice. This is likely due to the simplicity of the problem, and more complex dependencies 660 on hidden variables would likely lead to further decreases in skill. The main advantage of EDMD over DMD begins to become apparent in the cubic Lorenz experiment, where moving to a 3rd Hermite polynomial basis allows it to slightly outperform its linear counterpart, though the variability in the system (Fig. 3) is a greater magnitude than this improvement in skill.

## 4.2 Experiment 1: Three Box Model

The three box model experiment is meant to benchmark the baseline performance of each technique in the presence of strong 665 memory effects (Fig. 1 (a)). Figure 5 summarizes the results of four emulation techniques (pattern scaling, deconvolution, modal fitting, and DMD) when trained and tested on different scenario combinations, while Fig. 6 compares the true (ODE-integrated) solution to that obtained using the Fluctuation Dissipation Theorem.



**Figure 5.** NRMSE heatmaps for pattern scaling (a), deconvolution (b), modal fitting (c), and DMD (d) emulators trained and tested against the three box model. Results are shown in percentages, where lighter values correspond to lower error (higher performance) and darker values correspond to higher error (lower performance). Scenarios used for training are shown on the x-axis, while scenarios used for testing are shown on the y-axis. We do not include results for training and testing on the same dataset.



**Figure 6.** Fluctuation Dissipation Theorem emulator performance for three box model scenarios. Solid line shows ground truth (ODE-integrated) solution, while dot-dash shows emulated solution.

Pattern scaling (Method I) consistently underperforms relative to the other techniques presented in this section, exhibiting the highest NRMSE values for all train/test combinations. It fails across almost every scenario due to the influence of long timescales on the global mean temperature (strong memory effects). This experiment highlights pattern scaling’s brittleness when key assumptions, such as exponential forcing (Giani et al., 2024), are violated. These assumptions are consistent in most ScenarioMIP experiments however, leading to higher performance in practice relative to this simple example (Wells et al., 2023).

Applying deconvolution (Method III) leads to much higher performance than pattern scaling when trained on either *Abrupt*, 670 *Plateau*, or *Overshoot*, but sees a drop in performance when trained on *High Emissions*. This is because the true solution is an eigenfunction of the forcing (i.e. both the temperature response and forcing are exponentials), so the system is effectively characterized by a single timescale, that of the forcing. Deconvolution loses skill due to difficulties identifying all the timescales in the system, leading to extrapolation errors when training on this scenario. When trained on either *Plateau* or *Overshoot*, we 675 see errors in emulating *Abrupt*, meaning that the emulator has not learned the true system response despite relatively high performance in emulating the other scenarios. This is due to ill-conditioning of the  $\mathbf{F}$  matrix in these scenarios, leading to a response function that overfits these data; we discuss the limitations of training deconvolution with these scenarios further in Sect. 5.


Modal fitting (Method IV) exhibits two interesting properties: (1) training on *High Emissions* leads to poor extrapolative capability and (2) training on *Abrupt* leads to the highest performance overall. The first is also caused by the solution being an 680 eigenfunction of the forcing. It is difficult for the optimization routine to determine the correct timescales, even when initialized near the true values. This is true to a lesser degree in *Plateau* and *Overshoot*, which also do not display clean separation of time scales like *Abrupt*.

DMD (Method V) is able to capture all relevant timescales and interactions regardless of the scenario, with a maximum of 5.8% NRMSE across all train/test combinations; this level of error results from training on *High Emissions* and testing on 685 *Abrupt*, as was the case with the modal fitting emulator. The method’s high skill here is due to the governing dynamics being purely linear and there being no hidden variables, meaning all assumptions for applying DMD are accurate. Results for EDMD (Method VI) are omitted from this section as they are identical to DMD.

The Fluctuation Dissipation Theorem (Method II) has consistently high performance across all scenarios considered, with 690 NRMSE values of 0.80%, 0.50%, 0.75%, and 1.29% for the four scenarios shown in Fig. 6 (NRMSE values given by scenario from left to right). These values are lower than any other technique on average. These errors are due to the integration scheme 695 with which we derive the FDT response function, as we only use a first-order integrator. Since it requires us to simulate two scenarios (one perturbed and one unperturbed), error can accumulate between these simulations; decreasing the integrator time step or using a higher-order integrator (not shown) increases accuracy for this method. Despite this, the FDT gives us, up to the precision of our integrator, the system’s true response function, which is a major advantage compared to the other techniques 700 which may or may not provide a physically-interpretable solution. The full implementation of the FDT requires a spatially explicit response matrix with multiple perturbation runs, but for a more even comparison to the other techniques, we only consider the well-mixed case here.

### 4.3 Experiment 2: Restricted Two Box Model

The restricted two box model investigates the impact of hidden variables (Fig. 1 (b)). This experiment is meant to test if an 705 emulator can learn the true system response if not all information is included in the training data. Figure 7 summarizes the results of four emulation techniques (pattern scaling, deconvolution, modal fitting, and DMD) when trained and tested on different scenario combinations. Restricting the data means there is only one temperature series, rather than the three in the previous case. We therefore cannot calculate a global mean, and use a modified definition of pattern scaling in this section, 710 mapping from forcing to temperature anomaly. As the FDT (Method II) has roughly equivalent performance to the previous section and is not impacted by the introduction of hidden variables, we omit it from this section.



**Figure 7.** NRMSE heatmaps for pattern scaling (a), deconvolution (b), modal fitting (c), and DMD (d) emulators trained and tested against the restricted two box model. Results are shown in percentages, where lighter values correspond to lower error (higher performance) and darker values correspond to higher error (lower performance). Scenarios used for training are shown on the x-axis, while scenarios used for testing are shown on the y-axis. We do not include results for training and testing on the same dataset.

For all methods except deconvolution (Method III), we see a sharp drop in performance when introducing a hidden variable into the system. Deconvolution exhibits the same failure mode when training on *High Emissions* as before but to a greater degree, along with the ill-conditioning failure mode when training on *Plateau* and *Overshoot*. Because this method treats each

region as independent, it is more robust to the addition of hidden variables. It is able to capture the aggregate response of the  
715 atmospheric box that includes the influence of the ocean, but would not be able to separate those effects; i.e. the response function we derive is somewhat non-physical, though it can emulate the system effectively.

For the modal fitting emulator (Method IV), we initialize the optimization routine with guesses for both dominant modes (the fast atmospheric response and slower oceanic response). It is largely unsuccessful in identifying these modes, except in the case of training with *Abrupt*. This scenario is unique in that both modes are visible in the atmospheric box alone (see the  
720 leftmost plot in the middle row of Fig. 3). Training on either *High Emissions* or *Overshoot* appears promising at first, but neither can extrapolate to *Abrupt*, meaning it effectively overfits on these scenarios and loses extrapolative capabilities. As before, we see that training on *High Emissions* leads to the worst performance overall, as this scenario is characterized by only one effective timescale.

DMD (Method V) and by extension, EDMD (Method VI), experiences the sharpest decline in performance, with errors  
725 increasing by several orders of magnitude in some cases. Both methods see lower error in emulating scenarios similar to the training data (e.g. *High Emissions* vs. *Plateau*), but rapidly increasing error outside that regime. In addition to learning timescales like the previous two methods, DMD and EDMD are attempting to learn spatial interactions as well, meaning they are disproportionately affected by the hidden variable. We can also frame this issue theoretically by stating that hidden variables violate one of the fundamental assumptions of ~~E/EDMD and~~ DMD: the quantities we emulate are representative of  
730 all relevant system dynamics. By hiding the oceanic box, neither algorithm can learn the true physical behavior of the system. With EDMD, increases in polynomial order lead to further decreases in performance (not shown).


#### 4.4 Experiment 3: Noisy Three Box Model

Results of the noisy three box model show how noise affects each emulator (Fig. 1 (c)). Figure 8 summarizes the results of four emulation techniques (pattern scaling, deconvolution, modal fitting, and DMD) when trained only on *Abrupt* and tested  
735 against the other three scenarios; we choose to train only on *Abrupt* as it yielded high performance across all methods (except pattern scaling), and we want to isolate the impact of noise. See Fig. 4 in Sect. 4.1 for performance metrics across all train/test combinations with a constant ensemble size. Since the noise is added linearly, taking the difference between the perturbed and unperturbed ensembles effectively removes the noise when using the FDT (Method II). This leads to constant performance regardless of ensemble size, which is shown in Fig. 4. We additionally omit EDMD (Method VI) as it gives no improvements  
740 over DMD (Method V) in this linear case.

For these results, we evaluate performance relative to their noiseless baseline, rather than the absolute value of NRMSE; although *Abrupt* led to high performance for most methods, each method has a different baseline and some methods (e.g. pattern scaling) performed poorly when trained on this scenario. All methods exhibit decreased performance in the noisy case relative to the noiseless baseline.

745 Pattern scaling (Method I) experiences no change in performance as the number of ensemble members is increased, as the linear regression smooths the data, reducing the impact of noise regardless of the ensemble size. With both deconvolution (Method III) and modal fitting (Method IV), there is an almost random change in performance depending on the number of

NRMSE vs. Ensemble size by method  
Training scenario: *Abrupt*




**Figure 8.** NRMSE vs. number of ensemble members for pattern scaling (a), deconvolution (b), modal fitting (c), and DMD (d) emulators trained on *Abrupt* and tested against the three remaining scenarios. Solid lines indicate the error in training/testing with noisy data, while the dashed lines indicate error in training/testing with noiseless data.

ensemble members. This is because both methods regularize the data. Deconvolution requires extra regularization when the system is noisy, or else the algorithm overfits on the noise, leading to extremely high error ( $> \mathcal{O}(10^{10})$ ). The regularization has a similar effect to pattern scaling in making the expected performance of these algorithms more robust to noise. The variation in performance is due to the random sampling of ensemble members, with combinations that exhibit high error skewing the overall results. The error in DMD (Method V) is monotonically decreasing with ensemble size, though the presence of noise leads to a drop in performance relative to the noiseless baseline.

#### 4.5 Experiment 4: Cubic Lorenz System

The cubic Lorenz system allows us to jointly investigate the impact of chaos/noise and weak nonlinear effects on our emulators (Fig. 1 (c) and (d)). We run a 5,000 member ensemble as the variation in this experiment is much higher than the previous noisy case. As in experiment two, we use a slightly modified definition of pattern scaling, mapping from forcing to quantity of interest (the ensemble mean of  $Z$ ). Figure 9 summarizes emulator performance against the number of ensemble members, while Fig. 10 shows the response function derived using the FDT.

NRMSE vs. Ensemble size by method  
Training scenario: *Overshoot*




**Figure 9.** NRMSE vs. number of ensemble members for all emulators trained on *Overshoot* and tested against the three remaining scenarios. Emulators are shown as pattern scaling (a), FDT (b), deconvolution (c), modal fitting (d), and DMD (e), and EDMD (f). The FDT is trained on separate perturbation scenarios, and is therefore tested against all four scenarios. Unlike experiment three, there is no baseline/noiseless skill to compare against.

760 Similar to the previous noisy experiment (Sect. 4.4), pattern scaling (Method I) exhibits a constant level of performance independent of the number of ensemble members. The linear fitting process creates a strong artificial smoothing effect on the data, diminishing the potential impact of noise. This is also the case with both deconvolution (Method III) and the modal fitting (Method IV) approach, both of which have little variability based on the number of ensemble members. The modal fitting approach additionally requires an imaginary component to enforce oscillations in the response function similar to those in the 765 FDT result (Fig. 10). All approaches except DMD additionally show increased skill for smaller perturbations, i.e. higher skill in predicting *Plateau* than *Abrupt*. This is likely because smaller forcings lead to smaller deviations from the theoretical limit of response theory, which assumes small perturbations from the background state.

The performance of the FDT (Method II) is strongly dependent on the number of ensemble members. Figure 10 illustrates this point by showing how the response function derived using the FDT changes based on ensemble size. We treat the 50,000 member ensemble as our point of comparison, as further increases in ensemble size did not result in notable performance improvements. Key features, such as the initial magnitude of the response along with the time to reach that magnitude are 770

consistent across all ensemble sizes, but the three cases deviate after this initial peak. All three cases exhibit a similar frequency of oscillation over the time period tested, with noise in the 500 member ensemble influencing the longer-term behavior of that response (between years 3-5). There are deviations from the 50,000 member response in the 5,000 member case as well, though 775 it is generally more in-phase than the 500 member ensemble. The NRMSE between the 50,000 and 5,000 member ensembles is 166.22%, while the NRMSE between the 50,000 and 500 member ensembles is 546.06%. Both responses are far from the ground truth, but the 5,000 member ensemble is much closer than the 500 member ensemble. Because the 5,000 member ensemble has such high error relative to the 50,000 member ensemble, the predictive skill shown in Fig. 4 and Fig. 9 does not 780 tell the full story. By further increasing ensemble size, we expect to see commensurate increases in accuracy when emulating this system with the FDT.



**Figure 10.** Response function for the cubic Lorenz system derived using the Fluctuation Dissipation Theorem with three sets of ensemble members: 500, 5,000, and 50,000. We use  $\Delta t = 0.01$  and  $\delta = 50 \Delta t$  applied to the  $Y$  component of the system.

Despite [this experiment violating the fact that this experiment violates](#) the linearity assumption of DMD (Method V), it has relatively stable performance of a similar order to the other methods tested. Predictive skill on *High Emissions* and *Plateau* increases with the number of ensemble members, as one would expect as noise is averaged out, but skill on *Abrupt* decreases, which seems to be counterintuitive. In this case, we may not be introducing any further information about the coherent, un-785 underlying dynamics, which is supported by other methods showing consistent performance in these regimes. Increasing the ensemble size is leading to further refinement of the emulator's its parameters for *Overshoot* and its more closely related scenarios (*High Emissions* and *Plateau*). A deeper investigation is required to assess DMD's suitability for Lorenz-like systems. EDMD (Method VI) does not exhibit this behavior, instead performing with consistent skill across all combinations. This is likely because the 3rd-order Hermite polynomial used as the basis is well-suited to train on this scenario, illustrating the need 790 for careful selection of basis functions.

## 5 Discussion and Conclusions

While emulators of Earth System Models (ESMs) have recently surged in popularity, uncertainty regarding their performance under a variety of scenarios and the lack of a comprehensive theoretical framework for analysis have posed problems for efforts

at fundamental methodological comparisons. Our framework for emulator design and analysis builds on ideas from statistical mechanics and stochastic calculus, facilitating analysis of several emulation techniques from a theoretical and practical perspective. Our experiments based on simplified representations of the climate stress test a suite of emulators, including pattern scaling, response functions, and operator-based emulators, in the presence of memory effects, hidden variables, noise, and nonlinearities. Response function emulators consistently outperform other techniques, and the Fluctuation Dissipation Theorem (FDT) provides a robust method to derive them, though it also requires its own experimental ensemble. [Table 6 summarizes](#) [Section 5.1 describes emulator performance and key findings from our pedagogical examples, while Sect. 5.2 discusses the implications of our findings for ESMs. Table 6 additionally summarizes our experimental findings, focusing on](#) the robustness of different emulators to different sources of error.

**Table 6.** Summary of emulator capability by technique based on the results from Sect. 4. An 'X' indicates a technique possess the listed capability, while a '∞' indicates may meet this requirement if other conditions are met; we discuss these capabilities explicitly in Sect. 5. *Memory* refers to an emulator's ability to capture memory effects (Fig. 1 (a), experiment one), *Hidden* refers to an emulator's skill in the presence of hidden variables (Fig. 1 (b), experiment two), *Noise* refers to an emulator's robustness to simulation noise (Fig. 1 (c), experiment three), and *Nonlin.* refers to an emulator's ability to capture weak nonlinear effects (Fig. 1 (d), experiment four).

Technique	<i>Memory</i>	<i>Hidden</i>	<i>Noise</i>	<i>Nonlin.</i>
Method I: Pattern Scaling			X	
Method II: Fluctuation Dissipation Theorem	X	X	∞	∞
Method III: Deconvolution	X	X	∞	∞
Method IV: Modal Fitting	X	∞	X	∞
Method V: Dynamic Mode Decomposition (DMD)	X		∞	
Method VI: Extended DMD	X		∞	∞

## 5.1 [Emulator performance and trade-offs](#)

Each emulation technique considered in this work belongs to a spectrum of methods as defined by the joint Fokker-Planck/Koopman operator framework. Some emulators on this spectrum demand strict assumptions (quasi-linear/pattern scaling), while others are much more general (EDMD). There is a trade-off between the strictness of assumptions and emulator complexity, and relaxing these assumptions can shift the emulator's optimal use case. More general techniques may require specifically designed experiments, and decreasing structural emulator error may come at the price of increased computational costs (e.g. the Fluctuation Dissipation Theorem). Using this framework additionally identifies a gap in the current emulator typology as defined

810 by Tebaldi et al. (2025), as we need to consider the potential role operator-based emulators can play in this ecosystem; e.g. characterizing physical behavior in the system in addition to emulating it, as in Navarra et al. (2024).

815 Pattern scaling is a popular emulation technique because it is easy to implement, fast to apply, and its limits are well understood empirically (Mitchell, 2003; Tebaldi and Arblaster, 2014; Wells et al., 2023). Its efficiency makes it the method of choice particularly for assessments of mean annual temperature in monotonic forcing scenarios (e.g. SSP5-8.5, 3-7.0, or 2-4.5) and  
820 for understanding first-order trends of climate signals, even in the presence of internal variability. Previous work has shown this approach is valid only when the forcing is exponential and has a fixed spatial pattern, along with linear dynamics and feedbacks (Giani et al., 2024). Our results additionally show that pattern scaling exhibits two sources of irreducible error: a mismatch between the true and predicted patterns at equilibrium and the assumption that the climate must respond instantaneously to external forcings. If forcing history is important, such as in centennial-scale or strong overshoot experiments,  
825 the single-pattern approximation breaks down, misrepresenting shifts in regional warming over time. This is also the case with highly variable fields such as precipitation, where the first-order approximation may not capture significant trends. More general quasi-equilibrium approaches show promise (e.g. mapping from forcing to temperature in experiments two and four), but have yet to be widely explored in the context of full-scale ESMs. Pattern scaling's limitations push us towards emulation techniques that can capture more complex dynamics.

830 Response functions are increasing in popularity as they can capture many processes of interest that are missed by pattern scaling, such as the pattern and memory effects (Freese et al., 2024; Sandstad et al., 2025; Winkler and Sierra, 2025; Womack et al., 2025). This makes them ideal for representing decision-relevant, non-monotonic forcing scenarios, such as temperature overshoots. Response function approaches assume a linear relationship between the input forcing and output variable interest and that perturbations to the system are small (Lucarini et al., 2017). As a result, they are able to capture weakly nonlinear effects, so long as perturbations remain within the linear response regime. ~~Response functions also have the potential to emulate longer (post-2100) scenarios due to their ability to capture memory effects, but longer ESM runs are required to test this.~~ They must be used with caution when nonlinear effects are dominant or (depending on the technique) when internal variability is significant.

835 Despite its computational costs, deriving response functions with the Fluctuation Dissipation Theorem (FDT) offers a benefit over other response function techniques: it generates the system's exact linear response. Deconvolution and modal-fitting, by contrast, can produce non-physical output. As the FDT states, the response to small perturbations can be captured by  $R(t)$  if the system statistics are approximately stationary and the dynamics drive the weakly perturbed system back to the unperturbed state. The concept of climate is predicated on assuming the latter is true, further cementing the FDT's utility in this context. Because FDT-based response functions are physically interpretable, they support linear analyses of Earth system processes  
840 and serve as a reliable foundation for climate emulators (Lucarini and Chekroun, 2024). ~~Although the approach requires large ensembles of non-standard perturbation experiments (Lucarini et al., 2017; Lembo et al., 2020), our results show that the improvements in accuracy and interpretability may outweigh these costs, highlighting the need for further research on FDT-based emulation of full-scale ESMs.~~

Emulators that seek an explicit representation of the Koopman operator are potentially powerful tools as they are founded on 845 rigorous theory and are interpretable (Tu et al., 2014; Williams et al., 2015; Schmid, 2022). They can, in principle, reproduce any behavior the climate system might exhibit. In practice, however, their utility is constrained by several factors. Both Dynamic Mode Decomposition (DMD) and Extended DMD (EDMD) require the input and output variables of interest (e.g. radiative forcing and temperature) to completely characterize the dynamics of the system, rendering them sensitive to hidden variables. DMD additionally requires linearity between inputs and outputs, which is often violated in practice (Schmid, 2010). EDMD 850 relaxes this assumption by using a higher-dimensional space at the cost of selecting an appropriate (and often problem-specific) set of basis functions (Williams et al., 2015). The choice of basis functions is a major consideration with this method, and we may have been able to improve our implementation of EDMD further with a different choice. Solving the resulting large eigenvalue problems with either algorithm can be computationally demanding, and ~~E/EDMD and~~ DMD can be sensitive to noise, potentially overfitting to data. Despite these challenges, operator methods allow us to identify dominant modes of 855 variability in the climate system. They can also, in theory, be used to capture state-dependent and non-stationary processes, though this again requires a careful selection of basis functions and a large amount of training data. ~~Further While EDMD and DMD attempt to approximate the Koopman operator, they are simplified representations and in many cases do not closely approximate the true operator. Despite this, the Koopman and Fokker-Planck operators provide the most useful theoretical basis as they offer a way to directly link disparate forms of emulators. These techniques have the potential to be highly generalizable to scenarios beyond the training data as they can reproduce the system's true dynamics, but further~~ research is required to 860 determine the potential of using operator-based methods directly for climate emulation.

Emulator performance varies depending on the experimental setup, highlighting that emulators are often designed to be application specific and not completely general. Figure 4 provides an overview of these results, but each emulator had the potential for high performance depending on the application. For example, pattern scaling performs poorly on all experiments, 865 but shows high skill regardless of the experiment when trained and tested against *High Emissions*; this is not shown, as the case where the training and testing datasets are the same is trivial (near zero error) for all emulation techniques. However, this illustrates that pattern scaling has utility if used on scenarios with exponential forcing, more akin to ScenarioMIP (O'Neill et al., 2016); see (Giani et al., 2024) for further discussion. Future work will further examine the role training data plays in emulator development.

Whether emulators learn physically interpretable representations of the system they are emulating remains an open question, 870 though our process of testing an emulator's extrapolative capability suggests that some techniques do learn the system's true behavior. The clearest example of this is the FDT, which performed consistently well across all scenarios. This is to be expected as the theory behind the FDT shows that it calculates the physical impulse response of the system (Lucarini et al., 2017; Giorgini et al., 2024). Pattern scaling on the other hand, by definition, does not learn realistic behavior unless the system is 875 fully determined by the pattern scaling coefficients. For other techniques, the results are less clear. For example, the modal fitting approach is able to extrapolate successfully in any of the first three experiments when trained on *Abrupt*, but not when trained on *High Emissions*, further supporting the need for an effort focused on quantifying the impact of training data on climate emulators. Deconvolution and DMD also exhibit mixed levels of extrapolative skill, leading to difficulties in making

a consistent argument about interpretability from our results. This is especially the case for DMD, as the  $\mathcal{L}$  matrix we derive  
880 is not easily mappable to the true underlying parameters of e.g. the coupled three box model, as this problem is effectively underdetermined; we are solving for twelve DMD parameters, whereas the full system is determined by three heat capacities, three feedback parameters, and one diffusion coefficient. Future work will investigate the possibility of learning true system parameters from these emulated representations.

## 5.2 Implications for ESMs

885 While the lack of a common conceptual baseline has historically hindered comparisons between emulator classes, our framework takes an important step towards resolving this. Efforts such as ClimateBench, which provide a common training and evaluation benchmark, have been useful to that end (Watson-Parris et al., 2022), but emulator structural differences prevent it from being applied to all existing emulation techniques. Additionally, the high computational burden of running scenarios beyond those in the CMIP archive (for training or evaluation), prevents rigorous assessment of emulator capability (e.g., emulating the impact of  
890 individual forcings) and generalizability (accuracy beyond ScenarioMIP). Results from experiments such as the Detection and Attribution MIP (DAMIP) and Regional Aerosol MIP (RAMIP) can help fill these gaps (Gillett et al., 2016; Wilcox et al., 2023), but the field of ESM emulation is currently data-constrained. Our theoretical framework and pedagogical experiments provide value in this data-limited setting, as they allow us to evaluate the assumptions present in many common emulators. Our results illustrate the potential sources of error different emulator structural assumptions invite, giving us tools to assess and improve  
895 emulation techniques independently of ESM results. As ESMs improve, this framework can help ensure emulators are prepared to train on those new results.

Our pedagogical experiments provide a useful tool to isolate and examine individual sources of error relevant to emulating ESMs (Fig. 1). Though our simplified models are limited in that they lack much of the complexity of full-scale ESMs, our experiments highlight that emulator errors can be proactively resolved through structural changes in emulation, regardless of  
900 the parent model. For example, our results further support the growing body of literature on the utility of response functions (Freese et al., 2024; Womack et al., 2025; Winkler and Sierra, 2025). Response functions offer improvements over pattern scaling, particularly when considering memory effects in decision-relevant scenarios. They may also better emulate longer (post-2100) scenarios by accounting for regional pattern shifts, though longer ESM runs, such as the extensions proposed in ScenarioMIP for CMIP7, are required to test this (Van Vuuren et al., 2025). Existing emulators of ESMs may also benefit from incorporating  
905 response functions. For example, recent work into hybrid emulation using a generative model conditioned on pattern scaling could be extended by conditioning on response functions instead (Bouabid et al., 2025).

Several promising emulation techniques explored here, including the Fluctuation Dissipation Theorem (FDT), Dynamic Mode Decomposition (DMD), and Extended DMD (EDMD), have seen uses in climate science but have yet to be applied directly as emulators of ESM outputs as defined by Tebaldi et al. (2025). An intermediate step for either the FDT or EDMD  
910 may be to first emulate an EMIC, helping determine useful training scenarios without the cost of a full ESM. Our results suggest further research into these techniques is warranted, as they may represent more complex dynamics than other methods. In this context, the FDT stands apart as the most promising technique for emulating general dynamical systems, as evidenced

by its skill in this and other recent work (Giorgini et al., 2025b). However, using the FDT to derive response functions through perturbations requires a full initial condition ensemble for every perturbed grid cell/region (Lucarini et al., 2017; Lembo et al., 2020), similar to the Green's Function MIP (Bloch-Johnson et al., 2024), and is likely prohibitively expensive for full ESMs. The score-based FDT (Sect. 2.3) provides a remedy, using statistical learning methods to learn the score function and thus the system response (Giorgini et al., 2025b). Regardless of the derivation method, our results suggest response functions are the dominant emulation technique both in terms of accuracy and interpretability.

Most work studying climate emulation focuses on the development and implementation of developing and implementing new approaches in an application-specific manner. Our results show the utility of an operator-based framework for systematic analysis and comparison across a spectrum of climate emulation techniques. The main benefit of this framework is that it provides a toolkit for understanding the trade-offs between emulator complexity and performance while connecting emulation techniques to fundamental principles of statistical mechanics and stochastic systems. Memory We find that memory effects, internal variability, hidden variables, and nonlinearities are potential sources of emulator error, and error sources, and that response function-based emulators consistently outperform other methods, such as pattern scaling and Dynamic Mode Decomposition DMD, across all experiments. Emulator performance varies by experimental setup, particularly through the choice of training data, and further work is required to fully characterize these effects in a rigorous way. This framework currently relies on simple experiments, and further work is needed to determine if operator-based methods like EDMD can be practically realized to emulate nonlinear processes in full-scale climate models. Our analysis also highlights the potential to use the Fluctuation Dissipation Theorem (FDT) to derive FDT's potential for deriving robust, physically-interpretable response functions to use for climate emulation, though this comes at the cost of increased computational resources, though its computational cost is a potential barrier. As interpretability is an ongoing point of discussion within discussion in the emulator community, investing resources in a physically-grounded method such as methods like the FDT may go a long way towards increasing the utility of emulators not just for emulation, but for linear system analysis.

Code and data availability. All code to reproduce this work is available at <https://doi.org/10.5281/zenodo.17572065> (Womack, 2025). The raw data from CMIP6 were retrieved through the Earth System Grid Federation interface at <https://aims2.llnl.gov/search/cmip6/>. Figs. 2 - 10 were produced using scientific color maps from Crameri (2023).

## Appendix A: Additional derivations

### A1 Pattern scaling errors

To understand the potential sources of error in pattern scaling, we start from the linear equation for temperature evolution

$$\frac{\partial}{\partial t} T(\mathbf{x}, t) = \mathcal{K}(\mathbf{x}, \mathbf{x}') T(\mathbf{x}', t) + P(\mathbf{x}) F(t), \quad (\text{A1})$$

where  $T(\mathbf{x}, t)$  is the spatially explicit temperature,  $\mathcal{K}(\mathbf{x}, \mathbf{x}')$  is the Koopman operator that governs the autonomous system dynamics,  $P(\mathbf{x})$  is the spatial forcing pattern, and  $F(t)$  is the time series of the forcing.

We can examine errors in pattern scaling by considering the case in which the pattern scaled temperature,  $T_{PS}(\mathbf{x}, t)$ , is trained using an exponential forcing,  $F(t) = e^{t/\tau}$ , where  $\tau$  indicates the growth rate of the exponential. Forcing our governing equation with this yields

$$T_{PS}(\mathbf{x}, t) = \left[ \frac{1}{\tau} \delta(\mathbf{x} - \mathbf{x}') - \mathcal{K}(\mathbf{x}, \mathbf{x}') \right]^{-1} P(\mathbf{x}') e^{t/\tau}. \quad (\text{A2})$$

Here  $\delta(\mathbf{x} - \mathbf{x}')$  is the Dirac delta, so  $\frac{1}{\tau} \delta - \mathcal{K}$  plays the role of  $\frac{1}{\tau} I - \mathcal{K}$  in discretized form; we assume  $\tau$  lies outside the spectrum of  $\mathcal{K}$  so the inverse exists. Factoring out the exponential from this expression leaves us with

$$a_1(\mathbf{x}) = \left[ \frac{1}{\tau} \delta(\mathbf{x} - \mathbf{x}') - \mathcal{K}(\mathbf{x}, \mathbf{x}') \right]^{-1} P(\mathbf{x}'). \quad (\text{A3})$$

$a_1(\mathbf{x})$  is therefore the spatial scaling pattern used as our emulator. Inserting  $T(\mathbf{x}, t) = a_1(\mathbf{x})F(t)$  into the governing equation with the same exponential forcing, leaving us with

$$\frac{1}{\tau} a_1(\mathbf{x}) = \mathcal{K}(\mathbf{x}, \mathbf{x}') a_1(\mathbf{x}') + P(\mathbf{x}). \quad (\text{A4})$$

This identity expresses how the pattern,  $a_1(\mathbf{x})$ , balances internal dynamics with an external forcing.

We now consider an alternate scenario with an arbitrary forcing,  $F_{alt}$ , that is not the exponential forcing used for training. We denote the error between the true solution and our emulator as

$$T'(\mathbf{x}, t) = T_{alt}(\mathbf{x}, t) - a_1(\mathbf{x})F_{alt}(t). \quad (\text{A5})$$

We then recognize that  $T_{alt}(\mathbf{x}, t) = T'(\mathbf{x}, t) + a_1(\mathbf{x})F_{alt}(t)$ . Inserting this into our governing equation and using the identity from Equation A4 gives an equation describing the evolution of errors over time

$$\frac{\partial}{\partial t} T'(\mathbf{x}, t) = \mathcal{K}(\mathbf{x}, \mathbf{x}') T'(\mathbf{x}', t) + \frac{1}{\tau} a_1(\mathbf{x}) F_{alt}(t) - a_1(\mathbf{x}) \frac{\partial}{\partial t} F_{alt}(t) \quad (\text{A6})$$

From this expression, we see that there are two distinct sources of error in pattern scaling when trained on an exponential (ScenarioMIP-like forcing). The first corresponds to an equilibrium-offset. If  $F_{alt}(t)$  asymptotes to a constant  $F_f$ , the time derivative in Equation A6 vanishes, leaving us with

$$\lim_{t \rightarrow \infty} T'(\mathbf{x}, t) = -\frac{1}{\tau} \mathcal{K}^{-1}(\mathbf{x}', \mathbf{x}) a_1(\mathbf{x}) F_f. \quad (\text{A7})$$

Since we assume  $\mathcal{K}^{-1}$  exists, there does not exist a non-zero vector such that  $\mathcal{K}^{-1}(\mathbf{x}', \mathbf{x}) a_1(\mathbf{x}) = 0$ . Therefore the temperature produced by pattern scaling does not perfectly match the true equilibrium pattern.

The second source of error occurs in the transient case. When  $F_{alt}(t)$  varies in time, the final term in Equation A6 does not go to zero. If  $F_{alt}(t)$  changes more quickly than the training growth rate (i.e.  $\frac{\partial F_{alt}(t)}{\partial t} > \frac{1}{\tau} F_{alt}(t)$ ), then pattern scaling under-predicts the true temperature change. Conversely, very slow changes in  $F_{alt}(t)$  lead to an over-prediction of the true temperature change. A non-negligible rate of change term signals that system memory will be significant in that scenario.

Physically, the first error arises because the system's equilibrium pattern depends on its slow internal modes, whereas the second arises because those modes cannot keep pace with forcing that accelerates faster (or slower) than the training rate  $\tau$ .

## A2 Deconvolution instabilities

Deconvolution can amplify noise or in the worst case, cause the response function to blow up entirely. Here we identify where those instabilities arise. While issues with deconvolution are apparent in the time domain, they are easier to diagnose in frequency space. We use the Fourier transform (denoted by  $\mathcal{F}$ ) to rewrite convolution as multiplication:

$$\mathcal{F}[g(w_t)] = \mathcal{F} \left[ \int_{-\infty}^{\infty} d\tau R(\mathbf{x}, \tau) F(t - \tau) \right] \quad (\text{A8})$$

$$\hat{g}(w_\omega) = \hat{R}(\mathbf{x}, \omega) \hat{F}(\omega), \quad (\text{A9})$$

where  $g(w_t)$  is our statistical quantity of interest,  $R(\mathbf{x}, t)$  is the response function,  $F(t)$  is the forcing, the hat denotes the (continuous-time) Fourier transform, and  $\omega$  is the angular frequency. Recovering the response function therefore becomes division:

$$\hat{R}(\omega) = \frac{\hat{g}(w_\omega)}{\hat{F}(\omega)}, \quad (\text{A10})$$

$$R(t) = \mathcal{F}^{-1} \left[ \frac{\hat{g}(w_\omega)}{\hat{F}(\omega)} \right], \quad (\text{A11})$$

where  $\mathcal{F}^{-1}$  denotes the inverse Fourier transform. In discrete space, we use the fast Fourier transform.

If  $\hat{F}(\omega)$  has any near-zero frequencies, dividing by it causes  $\hat{R}(\omega) \rightarrow \infty$  at those frequencies. The corresponding time-domain process requires an explicit matrix inverse, where small eigenvalues translate into an ill-conditioned matrix. Additionally, if  $|\hat{F}(\omega)|$  spans several orders of magnitude, the ratio  $\hat{g}(w_\omega)/\hat{F}(\omega)$  amplifies high-frequency measurement noise and round-off error. The condition number of the corresponding matrix becomes very large, yielding an unstable estimate of  $R(\mathbf{x}, t)$ .

These issues are also encountered in signal processing, where a system is said to lack a spectral inverse (i.e. zeros in the frequency domain) if it exhibits the above issues (Yeung and Kong, 1986; Zazula and Gyergyek, 1993). Even in the absence of noise, the relatively flat spectrum of a true impulse response makes it difficult to recover directly. A dominant eigenvalue can obscure the weaker ones.

## A3 Distinction between Green's and response functions

A scalar field,  $w(t)$ , governed by the linear time-invariant equation

$$\frac{\partial}{\partial t} w(t) = \mathcal{L}w(t) + F(t), \quad (\text{A12})$$

has a corresponding Green's function,  $G(t)$ , that solves

$$\frac{\partial}{\partial t} G(t) = \mathcal{L}G(t) + \delta(t), \quad G(t < 0) = 0. \quad (\text{A13})$$

For a linear operator,  $\mathcal{L}$ , the solution is

$$1000 \quad G(t) = H(t)e^{\mathcal{L}t}, \quad (\text{A14})$$

where  $H(t)$  is the Heaviside step function and  $e^{\mathcal{L}t}$  is a matrix exponential. From this, any general forcing produces a response given by

$$w(t) = \int_0^t G(\tau)F(t - \tau)d\tau. \quad (\text{A15})$$

A response function, on the other hand, is either an empirical or equation-driven function that reproduces the system's linearized output but is not required to satisfy Equation A13. When the underlying dynamics are nonlinear, as is the case in 1005 climate models, a true Green's function does not exist. In practice however, the success of techniques such as pattern scaling illustrates that temperature response is very nearly linear for most of the globe, suggesting that data-derived response functions may closely approximate Green's functions for certain variables.

#### A4 Transitioning from $\mathcal{K}$ to $\mathcal{L}$

1010 We begin from a vector form of Equation 6, the expectation of a statistical field  $g(\mathbf{w})$ , where bold symbols are used to explicitly denote vectors. The vector  $\mathbf{w}$  represents a set of state variables,  $w_i$ , at discrete points in space. The evolution of  $\langle g(\mathbf{w}) \rangle$  is

$$\frac{\partial}{\partial t} \langle g(\mathbf{w}) \rangle = \left\langle [\mathcal{N}_i(\mathbf{w}, t) + F_i(t)] \frac{\partial}{\partial w_i} g(\mathbf{w}) \right\rangle + D \left\langle \frac{\partial^2}{\partial w_i^2} g(\mathbf{w}) \right\rangle. \quad (\text{A16})$$

We consider the case where  $g(\mathbf{w}) = w_i$  to find the evolution of the mean of the state variables themselves. Substituting this gives

$$1015 \quad \frac{\partial}{\partial t} \langle w_i \rangle = \langle \mathcal{N}_i(\mathbf{w}, t) \rangle + F_i(t). \quad (\text{A17})$$

We then define a steady baseline state,  $\bar{\mathbf{w}}$ , as

$$\langle \mathcal{N}_i(\bar{\mathbf{w}}, t) \rangle = -\bar{F}_i, \quad (\text{A18})$$

where  $\bar{F}_i$  is constant in time. Deviations from the baseline satisfy

$$\frac{\partial}{\partial t} \langle w'_i \rangle = \langle \mathcal{N}_i(\bar{\mathbf{w}} + \mathbf{w}', t) - \mathcal{N}_i(\bar{\mathbf{w}}, t) \rangle + F'_i(t), \quad (\text{A19})$$

1020 where  $F'_i(t)$  is the time-varying component of the forcing. We then use a first-order Taylor expansion around  $\bar{\mathbf{w}}$  to write

$$\frac{\partial}{\partial t} \langle w_i \rangle \simeq \left. \frac{\partial \mathcal{N}_i}{\partial w_j} \right|_{\bar{\mathbf{w}}} \langle w'_j \rangle + F'_i(t) = \mathcal{L}_{ij} \langle w'_j \rangle + F'_i(t), \quad (\text{A20})$$

where the derivative term,  $\frac{\partial \mathcal{N}_i}{\partial w_j}$ , can be pulled out of the expectation because the baseline state is not stochastic. To conclude, we rewrite this with  $\langle w'_i \rangle = T(x_i, t)$  and drop the discrete notation for space

$$\frac{\partial}{\partial t} T(\mathbf{x}, t) = \mathcal{L}(\mathbf{x}, \mathbf{x}') T(\mathbf{x}, t) + F(\mathbf{x}, t). \quad (\text{A21})$$

1025 **A5 FDT relationship to Fokker-Planck and Koopman**

Here we show how the Fluctuation Dissipation Theorem (FDT) relates to the Fokker-Planck operator. The result shows that a linear response function can be computed directly from the forward operator of the unperturbed system.

Let  $\mathbf{w}$  represent our full system state. Consider an equation of the form

$$\frac{\partial \mathbf{w}}{\partial t} = f_0(\mathbf{w}, t) + f_1(\mathbf{w}, t) + \varepsilon \xi(t), \quad (\text{A22})$$

1030 where  $f_0$  governs the unperturbed system dynamics and  $f_1$  governs the perturbed system dynamics. The Fokker-Planck equation corresponding to this is

$$\partial_t p + \nabla \cdot \left[ (f_0 + f_1)p - \frac{\varepsilon^2}{2} \nabla p \right] = 0. \quad (\text{A23})$$

Without loss of generality, we decompose  $p = p_0 + p_1$ , where  $p_0$  satisfies

$$\partial_t p_0 + \nabla \cdot \left( f_0 p_0 - \frac{\varepsilon^2}{2} \nabla p_0 \right) = 0. \quad (\text{A24})$$

1035 Then  $p_1$  must exactly satisfy,

$$\partial_t p_1 + \nabla \cdot \left( f_0 p_1 + f_1 p_0 + f_1 p_1 - \frac{\varepsilon^2}{2} \nabla p_1 \right) = 0 \quad (\text{A25})$$

The perturbation variables ( $f_1$  and  $p_1$ ) form a higher order term that we neglect, giving

$$\partial_t p_1 + \nabla \cdot \left( f_0 p_1 - \frac{\varepsilon^2}{2} \nabla p_1 \right) \approx -\nabla \cdot (f_1 p_0). \quad (\text{A26})$$

The solution to this is

$$1040 \quad p_1(\mathbf{w}, t) = -e^{-\mathcal{F}_0 t} \nabla \cdot (f_1 p_0), \quad (\text{A27})$$

assuming that  $p_1(\mathbf{w}, 0) = 0$ , i.e. there is no perturbation at  $t = 0$ , and  $\mathcal{F}_0$  is the unperturbed (time-independent) Fokker-Planck operator. Multiplying through by an arbitrary statistical quantity of the state,  $g(\mathbf{w})$ , and integrating with respect to  $\mathbf{w}$  then yields the first order perturbation in  $g(\mathbf{w})$

$$\int g(\mathbf{w}) p_1(\mathbf{w}, t) d\mathbf{w} = \int g(\mathbf{w}) e^{-\mathcal{F}_0 t} \nabla \cdot (f_1 p_0) d\mathbf{w}. \quad (\text{A28})$$

1045 The quantity on the left hand side is the expected value of the perturbed statistical quantity as a function of time. The right hand side is the cross correlation of the statistical quantity,  $g$ , with  $h \equiv \nabla \cdot (f_1 p_0)/p_0$  with respect to the unperturbed system. Noting the Koopman operator is the adjoint of the Fokker-Planck operator gives

$$(e^{-\mathcal{F}_0 t})^* = e^{-\mathcal{K}_0 t}, \quad (\text{A29})$$

where  $^*$  indicates the adjoint (conjuate transpose in finite dimensions) and  $\mathcal{F}^* = \mathcal{K}$ , giving an expression for the response

1050 function in terms of the Koopman operator.

Alternatively, we can connect the Fokker-Planck operator to the FDT through the score function. Consider the score function of the state given by

$$s(\mathbf{w}) = \nabla_{\mathbf{w}} \ln p_0(\mathbf{w}). \quad (\text{A30})$$

For a small, instantaneous perturbation applied at  $t = 0$ , the linear response of the mean field at a lag  $t$  is given by

$$1055 \quad R(t) = -\langle g(\mathbf{w}_t)s(\mathbf{w}_0) \rangle_{p_0}, \quad (\text{A31})$$

where the angle brackets denote an average over the stationary ensemble. We express this correlation with a joint probability density as

$$R(t) = - \int \int p(\mathbf{w}_0, \mathbf{w}_t) g(\mathbf{w}_t) s(\mathbf{w}_0) d\mathbf{w}_t d\mathbf{w}_0, \quad (\text{A32})$$

Using Bayes' theorem, we factor the joint probability density as

$$1060 \quad p(\mathbf{w}_0, \mathbf{w}_t) = p_0(\mathbf{w}_0) p(\mathbf{w}_t | \mathbf{w}_0), \quad (\text{A33})$$

where  $p(\mathbf{w}_t | \mathbf{w}_0)$  is the conditional probability from  $\mathbf{w}_0$  to  $\mathbf{w}_t$ . For dynamics governed by the Fokker-Planck operator,  $\mathcal{F}$ , we have

$$p(\mathbf{w}_t | \mathbf{w}_0) = e^{\mathcal{F}t} \delta(\mathbf{w}_0 - \mathbf{w}_t). \quad (\text{A34})$$

We then insert this expression into Equation A32 and integrate over  $\mathbf{w}_t$ :

$$1065 \quad R(t) = - \int p_0(\mathbf{w}_0) e^{\mathcal{F}t} g(\mathbf{w}_0) s(\mathbf{w}_0) d\mathbf{w}_0. \quad (\text{A35})$$

Therefore, the linear response function can be obtained by propagating the unperturbed field with  $e^{\mathcal{F}t}$  and correlating the result with the stationary score function.

## Appendix B: Regularization for response functions

Estimating a response function from noisy data requires using deconvolution to invert an often ill-conditioned matrix. We 1070 choose to model the noise in our field of interest,  $g(\mathbf{W})$ , with a Gaussian noise term:  $\boldsymbol{\varepsilon} \sim \mathcal{N}(0, \sigma^2 \mathbf{I})$ . Rather than applying an ad-hoc smoothing algorithm, we cast the problem in a Bayesian framework, placing a Gaussian prior on the response matrix:  $\mathbf{R} \sim \mathcal{N}(0, \lambda^2 \mathbf{I})$ . Our measurement model is therefore

$$g(\mathbf{W}) = \mathbf{F}\mathbf{R} + \boldsymbol{\varepsilon}. \quad (\text{B1})$$

We have dropped  $\Delta t$  and the spatial pattern for conciseness, but this analysis can easily be repeated including those terms.

1075 Under this probabilistic model, we frame the task of estimating  $\mathbf{R}$  as finding the vector that maximizes the response function probability given the observable data we have collected, i.e.  $p(\mathbf{R}|g(\mathbf{W}))$ . This term is called the *maximum a posteriori* (MAP). As it is more convenient to work with log probabilities, we recast this problem as

$$\max_{\mathbf{R}} \log p(\mathbf{R} | g(\mathbf{W})). \quad (\text{B2})$$

Using Bayes theorem, maximizing the log-posterior,

$$1080 \quad \log p(\mathbf{R} | g(\mathbf{W})) = -\frac{1}{2\sigma^2} \|g(\mathbf{W}) - \mathbf{F}\mathbf{R}\|^2 - \frac{1}{2\lambda^2} \|\mathbf{R}\|^2 + \text{const}, \quad (\text{B3})$$

is equivalent to solving

$$\min_{\mathbf{R}} \|g(\mathbf{W}) - \mathbf{F}\mathbf{R}\|^2 + \alpha \|\mathbf{R}\|^2, \quad \alpha = \sigma^2/\lambda^2. \quad (\text{B4})$$

Thus ridge regression is equivalent to placing a Gaussian prior on the response function and assuming that the data we collect are corrupted by Gaussian noise.

1085 To avoid making an arbitrary choice for our noise and prior variance hyperparameters parameters,  $\sigma^2$  and  $\lambda^2$ , we propose to compute their maximum likelihood estimates under the distribution of the field of interest. We maximize the marginal likelihood evidence,

$$p(g(\mathbf{W}) | \sigma^2, \lambda^2) = \int p(g(\mathbf{W}) | \mathbf{R}, \sigma^2) p(\mathbf{R}, \lambda^2) d\mathbf{R} \quad (\text{B5})$$

$$= \mathcal{N}(g(\mathbf{W}) | 0, \Sigma), \quad (\text{B6})$$

1090 with covariance  $\Sigma = \sigma^2 \mathbf{I} + \lambda^2 \mathbf{F}\mathbf{F}^T$ . Maximizing the log-evidence,

$$-\frac{1}{2} (\log |\Sigma| + g(\mathbf{W})^T \Sigma^{-1} g(\mathbf{W})) + \text{const}, \quad (\text{B7})$$

has no closed-form solution for a general  $\mathbf{F}$ , so we determine  $\sigma^2$  and  $\lambda^2$  numerically.

## Appendix C: Analytic examples

In this appendix, we use a 1D Ornstein-Uhlenbeck (OU) process to analytically derive the Fokker-Planck operator, the Koopman operator, the eigenpairs of both operators, and the linear response function for the system obtained in two ways: (1) by directly solving the forced stochastic differential equation (SDE) and (2) by correlation with the score function.

### C1 Fokker-Planck and Koopman operator derivation

We define the OU SDE as

$$dw_t = -w_t dt + \sqrt{2} dW_t, \quad (\text{C1})$$

1100 where  $w_t$  is the statistical field of interest and  $W_t$  is a Wiener process. The drift coefficient,  $-w_t$ , relaxes the state toward zero, while the diffusion coefficient,  $\sqrt{2}$ , gives a unit variance.

We write the Fokker-Planck equation corresponding to this OU process directly:

$$\frac{\partial}{\partial t} p(w, t) = \frac{\partial}{\partial w} (wp) + \frac{\partial^2}{\partial w^2} p, \quad (\text{C2})$$

1105 where  $p(w, t)$  is the probability density function of the field. The stationary solution of this expression is the standard normal probability density:

$$p_0(w) = \frac{1}{\sqrt{2\pi}} e^{-\frac{w^2}{2}}. \quad (\text{C3})$$

From the previous result, we explicitly write the Fokker-Planck operator governing the evolution of the probability density as

$$\mathcal{F}(\cdot) = \frac{\partial}{\partial w} \left[ w(\cdot) + \frac{\partial}{\partial w}(\cdot) \right]. \quad (\text{C4})$$

1110 To find the eigenfunctions,  $\phi(w)$ , with  $\mathcal{F}\phi(w) = \lambda\phi(w)$ , we introduce the ansatz  $\phi(w) = h(w)e^{-\frac{w^2}{2}}$ , giving

$$h''(w) - wh'(w) - \lambda h(w) = 0, \quad (\text{C5})$$

whose solutions are Hermite polynomials,  $H_n(w)$ , with eigenvalues  $\lambda = -n$  for  $n = 0, 1, 2, \dots$ .

## C2 Response function via direct diagnosis

Adding a deterministic forcing,  $F(t)$ , to our OU process gives

$$1115 \quad dy_t = (-y_t + F(t))dt + \sqrt{2}dW_t. \quad (\text{C6})$$

Taking the expected value of this and assuming  $\langle y(0) \rangle = 0$  gives

$$\frac{d}{dt} \langle y \rangle = -\langle y \rangle + F(t), \quad (\text{C7})$$

whose solution is given by

$$\langle y(t) \rangle = \int_0^t e^{-\tau} F(t - \tau) d\tau, \quad (\text{C8})$$

1120 where the response function is  $R(t) = e^{-t}$  for  $t \geq 0$ .

## C3 Response function via correlation with score function

The stationary score function is given by

$$s(w) = \nabla_w \ln p_0(w) = -w, \quad (\text{C9})$$

where we can make this simplification since the stationary probability distribution is given by a standard normal.

1125 The Fluctuation Dissipation Theorem predicts

$$R(t) = -\langle w(t)s(w(0)) \rangle = e^{-t}, \quad (\text{C10})$$

which agrees exactly with the direct solution above.

*Author contributions.* Conceptualization: CW, GF, SB, SE, NS. Formal analysis: CW, GF, SB, AS. Funding acquisition: NS. Investigation:

CW. Methodology: CW, GF, SB, AS, PG. Supervision: GF, SE, NS. Visualization: CW, SB. Writing - original draft: CW, GF, SB. Writing -

1130 reviewing and editing: CW, GF, SB, AS, PG, SE, NS.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This research was part of the Bringing Computation to the Climate Challenge (BC3) project and supported by Schmidt Sciences, LLC. through the MIT Grand Challenges. We also acknowledge the MIT *Svante* cluster supported by the Center for Sustainability Science and Strategy for computing resources. We are grateful for the entire BC3 team who provided insightful feedback and discussions

1135 about this work. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies who support CMIP6 and ESGF. We additionally acknowledge the use of LLMs in editing and annotating the code associated with this work.

We would also like to thank the two anonymous reviewers for their thorough comments and time spent reviewing this manuscript. Their work has helped us

1140 improve the quality of our research.

## References

Addison, H., Kendon, E., Ravuri, S., Aitchison, L., and Watson, P. A.: Machine learning emulation of precipitation from km-scale regional climate simulations using a diffusion model, <https://doi.org/10.48550/arXiv.2407.14158>, arXiv:2407.14158, 2024.

1145 Armour, K. C., Bitz, C. M., and Roe, G. H.: Time-Varying Climate Sensitivity from Regional Feedbacks, *JOURNAL OF CLIMATE*, 26, 2013.

Bassetti, S., Hutchinson, B., Tebaldi, C., and Kravitz, B.: DiffESM: Conditional Emulation of Temperature and Precipitation in Earth System Models With 3D Diffusion Models, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS004194, <https://doi.org/10.1029/2023MS004194>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023MS004194>, 2024.

1150 Beusch, L., Gudmundsson, L., and Seneviratne, S. I.: Emulating Earth system model temperatures with MESMER: from global mean temperature trajectories to grid-point-level realizations on land, *Earth System Dynamics*, 11, 139–159, <https://doi.org/10.5194/esd-11-139-2020>, publisher: Copernicus GmbH, 2020.

Blanusa, M. L., López-Zurita, C. J., and Rasp, S.: Internal variability plays a dominant role in global climate projections of temperature and precipitation extremes, *Climate Dynamics*, 61, 1931–1945, <https://doi.org/10.1007/s00382-023-06664-3>, 2023.

1155 Bloch-Johnson, J., Rugenstein, M. A. A., Alessi, M. J., Proistosescu, C., Zhao, M., Zhang, B., Williams, A. I. L., Gregory, J. M., Cole, J., Dong, Y., Duffy, M. L., Kang, S. M., and Zhou, C.: The Green's Function Model Intercomparison Project (GFMIP) Protocol, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003700, <https://doi.org/10.1029/2023MS003700>, publisher: John Wiley & Sons, Ltd, 2024.

Bouabid, S., Sejdinovic, D., and Watson-Parris, D.: FaIRGP: A Bayesian Energy Balance Model for Surface Temperatures Emulation, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003926, <https://doi.org/10.1029/2023MS003926>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023MS003926>, 2024.

1160 Bouabid, S., Souza, A. N., and Ferrari, R.: Score-based generative emulation of impact-relevant Earth system model outputs, <https://doi.org/10.48550/arXiv.2510.04358>, arXiv:2510.04358 [physics], 2025.

Caldeira, K. and Myhrvold, N. P.: Projections of the pace of warming following an abrupt increase in atmospheric carbon dioxide concentration, *Environmental Research Letters*, 8, 034039, <https://doi.org/10.1088/1748-9326/8/3/034039>, publisher: IOP Publishing, 2013.

1165 Cao, L., Bala, G., Zheng, M., and Caldeira, K.: Fast and slow climate responses to CO<sub>2</sub> and solar forcing: A linear multivariate regression model characterizing transient climate change, *Journal of Geophysical Research: Atmospheres*, 120, 12,037–12,053, <https://doi.org/10.1002/2015JD023901>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2015JD023901>, 2015.

Castruccio, S., McInerney, D. J., Stein, M. L., Crouch, F. L., Jacob, R. L., and Moyer, E. J.: Statistical Emulation of Climate Model Projections Based on Precomputed GCM Runs, *Journal of Climate*, 27, 1829–1844, <https://doi.org/10.1175/JCLI-D-13-00099.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2014.

1170 Cimoli, L., Gebbie, G., Purkey, S. G., and Smethie, W. M.: Annually Resolved Propagation of CFCs and SF<sub>6</sub> in the Global Ocean Over Eight Decades, *Journal of Geophysical Research: Oceans*, 128, e2022JC019337, <https://doi.org/10.1029/2022JC019337>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022JC019337>, 2023.

Cooper, F. C. and Haynes, P. H.: Climate Sensitivity via a Nonparametric Fluctuation–Dissipation Theorem, *Journal of Atmospheric Sciences*, 68, 937–953, <https://doi.org/10.1175/2010JAS3633.1>, section: *Journal of the Atmospheric Sciences*, 2011.

Crameri, F.: Scientific colour maps, <https://doi.org/10.5281/zenodo.8409685>, language: eng, 2023.

Denisov, S. I., Horsthemke, W., and Hänggi, P.: Generalized Fokker-Planck equation: Derivation and exact solutions, *The European Physical Journal B*, 68, 567–575, <https://doi.org/10.1140/epjb/e2009-00126-3>, 2009.

Dix, M., Bi, D., Dobrohotoff, P., Fiedler, R., Harman, I., Law, R., Mackallah, C., Marsland, S., O'Farrell, S., Rashid, H., Srbinovsky, J.,  
1180 Sullivan, A., Trenham, C., Vohralik, P., Watterson, I., Williams, G., Woodhouse, M., Bodman, R., Dias, F. B., Domingues, C. M., Hannah, N., Heerdegen, A., Savita, A., Wales, S., Allen, C., Druken, K., Evans, B., Richards, C., Ridzwan, S. M., Roberts, D., Smillie, J., Snow, K., Ward, M., and Yang, R.: CSIRO-ARCCSS ACCESS-CM2 model output prepared for CMIP6 CMIP piControl, [https://www.wdc-climate.de/ui/entry?acronym=C6\\_4381092](https://www.wdc-climate.de/ui/entry?acronym=C6_4381092), 2023.

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model  
1185 Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, publisher: Copernicus GmbH, 2016.

Farley, J., MacMartin, D. G., Visioni, D., Kravitz, B., Bednarz, E., Duffey, A., and Henry, M.: A Climate Intervention Dynamical Emulator  
(CIDER) for Scenario Space Exploration, 2025.

Flato, G. M.: Earth system models: an overview, *WIREs Climate Change*, 2, 783–800, <https://doi.org/10.1002/wcc.148>, \_eprint:  
1190 <https://onlinelibrary.wiley.com/doi/pdf/10.1002/wcc.148>, 2011.

Franzke, C. L. E., Gugole, F., and Juricke, S.: Systematic multi-scale decomposition of ocean variability using machine learning, *Chaos: An  
Interdisciplinary Journal of Nonlinear Science*, 32, <https://doi.org/10.1063/5.0090064>, publisher: AIP Publishing, 2022.

Fredriksen, H.-B., Rugenstein, M., and Graversen, R.: Estimating Radiative Forcing With a Nonconstant Feedback Parameter and Lin-  
ear Response, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034145, <https://doi.org/10.1029/2020JD034145>, \_eprint:  
1195 <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020JD034145>, 2021.

Fredriksen, H.-B., Smith, C. J., Modak, A., and Rugenstein, M.: 21st Century Scenario Forcing Increases More for CMIP6  
Than CMIP5 Models, *Geophysical Research Letters*, 50, e2023GL102916, <https://doi.org/10.1029/2023GL102916>, \_eprint:  
<https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023GL102916>, 2023.

Freese, L. M., Giani, P., Fiore, A. M., and Selin, N. E.: Spatially Resolved Temperature Response Functions to  
1200 CO2 Emissions, *Geophysical Research Letters*, 51, e2024GL108788, <https://doi.org/10.1029/2024GL108788>, \_eprint:  
<https://onlinelibrary.wiley.com/doi/pdf/10.1029/2024GL108788>, 2024.

Giani, P., Fiore, A. M., Flierl, G., Ferrari, R., and Selin, N. E.: Origin and Limits of Invariant Warming Patterns in Climate Models,  
<https://doi.org/10.48550/arXiv.2411.14183>, arXiv:2411.14183, 2024.

Gillett, N. P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., Santer, B. D., Stone, D., and Tebaldi, C.: The Detection and  
1205 Attribution Model Intercomparison Project (DAMIP v1.0) contribution to CMIP6, *Geoscientific Model Development*, 9, 3685–3697,  
<https://doi.org/10.5194/gmd-9-3685-2016>, publisher: Copernicus GmbH, 2016.

Giorgini, L. T., Deck, K., Bischoff, T., and Souza, A.: Response Theory via Generative Score Modeling, *Phys. Rev. Lett.*, 133, 267302,  
<https://doi.org/10.1103/PhysRevLett.133.267302>, 2024.

Giorgini, L. T., Bischoff, T., and Souza, A. N.: Statistical Parameter Calibration with the Generalized Fluctuation Dissipation Theorem and  
1210 Generative Modeling, <https://arxiv.org/abs/2509.19660>, 2025a.

Giorgini, L. T., Falasca, F., and Souza, A. N.: Predicting forced responses of probability distributions via the fluctua-  
tion-dissipation theorem and generative modeling, *Proceedings of the National Academy of Sciences*, 122, e2509578122,  
<https://doi.org/10.1073/pnas.2509578122>, publisher: Proceedings of the National Academy of Sciences, 2025b.

Gottwald, G. A. and Gugole, F.: Detecting Regime Transitions in Time Series Using Dynamic Mode Decomposition, *Journal of Statistical Physics*, 179, 1028–1045, <https://doi.org/10.1007/s10955-019-02392-3>, 2020.

1215 Haseli, M. and Cortés, J.: Approximating the Koopman Operator using Noisy Data: Noise-Resilient Extended Dynamic Mode Decomposition, in: 2019 American Control Conference (ACC), pp. 5499–5504, <https://doi.org/10.23919/ACC.2019.8814684>, iSSN: 2378-5861, 2019.

Hasselmann, K.: Stochastic climate models Part I. Theory, *Tellus*, 28, 473–485, <https://doi.org/10.1111/j.2153-3490.1976.tb00696.x>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.2153-3490.1976.tb00696.x>, 1976.

1220 Hasselmann, K.: 1.27 - Optimizing Long-Term Climate Management, in: *Global Biogeochemical Cycles in the Climate System*, edited by Schulze, E.-D., Heimann, M., Harrison, S., Holland, E., Lloyd, J., Prentice, I. C., and Schimel, D., pp. 333–343, Academic Press, San Diego, ISBN 978-0-12-631260-7, <https://doi.org/10.1016/B978-012631260-7/50029-7>, 2001.

Hasselmann, K., Hasselmann, S., Giering, R., Ocana, V., and Storch, H. V.: Sensitivity Study of Optimal CO<sub>2</sub> Emission Paths Using a Simplified Structural Integrated Assessment Model (SIAM), *Climatic Change*, 37, 345–386, <https://doi.org/10.1023/A:1005339625015>, 1997.

1225 Hasselmann, K., Latif, M., Hooss, G., Azar, C., Edenhofer, O., Jaeger, C. C., Johannessen, O. M., Kemfert, C., Welp, M., and Wokaun, A.: The Challenge of Long-Term Climate Change, *Science*, 302, 1923–1925, <https://doi.org/10.1126/science.1090858>, publisher: American Association for the Advancement of Science, 2003.

1230 Henze, D. K., Hakami, A., and Seinfeld, J. H.: Development of the adjoint of GEOS-Chem, *Atmospheric Chemistry and Physics*, 7, 2413–2433, <https://doi.org/10.5194/acp-7-2413-2007>, publisher: Copernicus GmbH, 2007.

Herger, N., Sanderson, B. M., and Knutti, R.: Improved pattern scaling approaches for the use in climate impact studies, *Geophysical Research Letters*, 42, 3486–3494, <https://doi.org/10.1002/2015GL063569>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2015GL063569>, 2015.

1235 Huntingford, C. and Cox, P. M.: An analogue model to derive additional climate change scenarios from existing GCM simulations, *Climate Dynamics*, 16, 575–586, <https://doi.org/10.1007/s003820000067>, 2000.

Joos, F. and Bruno, M.: Pulse response functions are cost-efficient tools to model the link between carbon emissions, atmospheric CO<sub>2</sub> and global warming, *Physics and Chemistry of the Earth*, 21, 471–476, [https://doi.org/10.1016/S0079-1946\(97\)81144-5](https://doi.org/10.1016/S0079-1946(97)81144-5), 1996.

1240 Joos, F., Roth, R., Fuglestvedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmospheric Chemistry and Physics*, 13, 2793–2825, <https://doi.org/10.5194/acp-13-2793-2013>, publisher: Copernicus GmbH, 2013.

Kaiser, E., Kutz, J. N., and Brunton, S. L.: Data-driven approximations of dynamical systems operators for control, <https://doi.org/10.48550/arXiv.1902.10239>, arXiv:1902.10239 [math], 2019.

1245 King, A. D., Borowiak, A. R., Brown, J. R., Frame, D. J., Harrington, L. J., Min, S.-K., Pendergrass, A., Rugenstein, M., Sniderman, J. M. K., and Stone, D. A.: Transient and Quasi-Equilibrium Climate States at 1.5°C and 2°C Global Warming, *Earth's Future*, 9, e2021EF002274, <https://doi.org/10.1029/2021EF002274>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021EF002274>, 2021.

Klus, S., Koltai, P., and Schütte, C.: On the numerical approximation of the Perron-Frobenius and Koopman operator, *Journal of Computational Dynamics*, 3, 1–12, <https://doi.org/10.3934/jcd.2016003>, arXiv:1512.05997 [math], 2016.

Klus, S., Nüske, F., Kolta, P., Wu, H., Kevrekidis, I., Schütte, C., and Noé, F.: Data-Driven Model Reduction and Transfer Operator Approximation, *Journal of Nonlinear Science*, 28, 985–1010, <https://doi.org/10.1007/s00332-017-9437-7>, 2018.

Kochkov, D., Yuval, J., Langmore, I., Norgaard, P., Smith, J., Mooers, G., Klöwer, M., Lottes, J., Rasp, S., Düben, P., Hatfield, S., Battaglia, P., Sanchez-Gonzalez, A., Willson, M., Brenner, M. P., and Hoyer, S.: Neural general circulation models for weather and climate, *Nature*, 1255 pp. 1–7, <https://doi.org/10.1038/s41586-024-07744-y>, publisher: Nature Publishing Group, 2024.

Kravitz, B., Lynch, C., Hartin, C., and Bond-Lamberty, B.: Exploring precipitation pattern scaling methodologies and robustness among CMIP5 models, *Geoscientific Model Development*, 10, 1889–1902, <https://doi.org/10.5194/gmd-10-1889-2017>, publisher: Copernicus GmbH, 2017.

Kutz, J. N., Fu, X., and Brunton, S. L.: Multiresolution Dynamic Mode Decomposition, *SIAM Journal on Applied Dynamical Systems*, 15, 1260 713–735, <https://doi.org/10.1137/15M1023543>, 2016.

Leach, N. J., Jenkins, S., Nicholls, Z., Smith, C. J., Lynch, J., Cain, M., Walsh, T., Wu, B., Tsutsui, J., and Allen, M. R.: FaIRv2.0.0: a generalized impulse response model for climate uncertainty and future scenario exploration, *Geoscientific Model Development*, 14, 3007–3036, <https://doi.org/10.5194/gmd-14-3007-2021>, publisher: Copernicus GmbH, 2021.

Lembo, V., Lucarini, V., and Ragone, F.: Beyond Forcing Scenarios: Predicting Climate Change through Response Operators in a Coupled General Circulation Model, *Scientific Reports*, 10, 8668, <https://doi.org/10.1038/s41598-020-65297-2>, publisher: Nature Publishing Group, 1265 2020.

Lewis, J., Bodeker, G. E., Kremser, S., and Tait, A.: A method to encapsulate model structural uncertainty in ensemble projections of future climate: EPIC v1.0, *Geoscientific Model Development*, 10, 4563–4575, <https://doi.org/10.5194/gmd-10-4563-2017>, publisher: Copernicus GmbH, 2017.

1270 Lorenz, E. N.: Deterministic Nonperiodic Flow, *Journal of the Atmospheric Sciences*, 20, 130–141, [https://journals.ametsoc.org/view/journals/atsc/20/2/1520-0469\\_1963\\_020\\_0130\\_dnf\\_2\\_0\\_co\\_2.xml](https://journals.ametsoc.org/view/journals/atsc/20/2/1520-0469_1963_020_0130_dnf_2_0_co_2.xml), section: *Journal of the Atmospheric Sciences*, 1963.

Lorenz, E. N.: Predictability – a problem partly solved, in: *Predictability of Weather and Climate*, edited by Palmer, T. and Hagedorn, R., pp. 40–58, Cambridge University Press, 1 edn., ISBN 978-0-521-84882-4 978-0-511-61765-2 978-1-107-41485-3, <https://doi.org/10.1017/CBO9780511617652.004>, 2006.

1275 Lorenz, E. U.: Predictability: Does the Flap of a Butterfly's Wings in Brazil Set Off a Tornado in Texas?, *Resonance Journal of Science Education*, 20, 260–263, <https://doi.org/10.1007/s12045-015-0174-2>, 2015.

Lucarini, V. and Chekroun, M. D.: Detecting and Attributing Change in Climate and Complex Systems: Foundations, Green's Functions, and Nonlinear Fingerprints, *Physical Review Letters*, 133, 244201, <https://doi.org/10.1103/PhysRevLett.133.244201>, 2024.

Lucarini, V., Ragone, F., and Lunkeit, F.: Predicting Climate Change Using Response Theory: Global Averages and Spatial Patterns, *Journal 1280 of Statistical Physics*, 166, 1036–1064, <https://doi.org/10.1007/s10955-016-1506-z>, 2017.

Lucarini, V., Gutierrez, M. S., Moroney, J., and Zagli, N.: A General Framework for Linking Free and Forced Fluctuations via Koopmanism, <https://arxiv.org/abs/2506.16446>, 2025.

Lyu, G., Köhl, A., Matei, I., and Stammer, D.: Adjoint-Based Climate Model Tuning: Application to the Planet Simulator, *Journal of Advances in Modeling Earth Systems*, 10, 207–222, <https://doi.org/10.1002/2017MS001194>, \_eprint: 1285 <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2017MS001194>, 2018.

Maher, N., Milinski, S., Suarez-Gutierrez, L., Botzet, M., Dobrynin, M., Kornblueh, L., Kröger, J., Takano, Y., Ghosh, R., Hedemann, C., Li, C., Li, H., Manzini, E., Notz, D., Putrasahan, D., Boysen, L., Claussen, M., Ilyina, T., Olonscheck, D., Radatz, T., Stevens, B., and Marotzke, J.: The Max Planck Institute Grand Ensemble: Enabling the Exploration of Climate Sys-

tem Variability, *Journal of Advances in Modeling Earth Systems*, 11, 2050–2069, <https://doi.org/10.1029/2019MS001639>, \_eprint: 1290 <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2019MS001639>, 2019.

Mankovich, N., Bouabid, S., Nowack, P., Bassotto, D., and Camps-Valls, G.: Analyzing climate scenarios using dynamic mode decomposition with control, *Environmental Data Science*, 4, e16, <https://doi.org/10.1017/eds.2025.8>, 2025.

Marconi, U. M. B., Puglisi, A., Rondoni, L., and Vulpiani, A.: Fluctuation–dissipation: Response theory in statistical physics, *Physics Reports*, 461, 111–195, <https://doi.org/10.1016/j.physrep.2008.02.002>, 2008.

1295 Mathison, C. T., Burke, E., Kovacs, E., Munday, G., Huntingford, C., Jones, C., Smith, C., Steinert, N., Wiltshire, A., Gohar, L., and Varney, R.: A rapid application emissions-to-impacts tool for scenario assessment: Probabilistic Regional Impacts from Model patterns and Emissions (PRIME), *EGUphere*, pp. 1–28, <https://doi.org/10.5194/egusphere-2023-2932>, publisher: Copernicus GmbH, 2024.

Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, *MAGICC6 – Part 1: Model description and calibration*, *Atmospheric Chemistry and Physics*, 11, 1417–1456, <https://doi.org/10.5194/acp-11-1417-2011>, publisher: Copernicus GmbH, 2011.

Metzler, H., Müller, M., and Sierra, C. A.: Transit-time and age distributions for nonlinear time-dependent compartmental systems, *Proceedings of the National Academy of Sciences*, 115, 1150–1155, <https://doi.org/10.1073/pnas.1705296115>, publisher: Proceedings of the National Academy of Sciences, 2018.

Mezić, I.: Analysis of Fluid Flows via Spectral Properties of the Koopman Operator, *Annual Review of Fluid Mechanics*, 45, 357–378, 1305 <https://doi.org/10.1146/annurev-fluid-011212-140652>, 2013.

Mitchell, T. D.: Pattern Scaling: An Examination of the Accuracy of the Technique for Describing Future Climates, *Climatic Change*, 60, 217–242, <https://doi.org/10.1023/A:1026035305597>, 2003.

1310 Müller, W. A., Jungclaus, J. H., Mauritzen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., Ilyina, T., Kleine, T., Kornblueh, L., Li, H., Modali, K., Notz, D., Pohlmann, H., Roeckner, E., Stemmler, I., Tian, F., and Marotzke, J.: A Higher-resolution Version of the Max Planck Institute Earth System Model (MPI-ESM1.2-HR), *Journal of Advances in Modeling Earth Systems*, 10, 1383–1413, <https://doi.org/10.1029/2017MS001217>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2017MS001217>, 2018.

Navarra, A., Tribbia, J., and Klus, S.: Estimation of Koopman Transfer Operators for the Equatorial Pacific SST, *Journal of the Atmospheric Sciences*, 78, 1227–1244, <https://doi.org/10.1175/JAS-D-20-0136.1>, 2021.

Navarra, A., Tribbia, J., Klus, S., and Lorenzo-Sánchez, P.: Variability of SST through Koopman Modes, *Journal of Climate*, 37, 4095–4114, 1315 <https://doi.org/10.1175/JCLI-D-23-0335.1>, section: *Journal of Climate*, 2024.

Netto, M., Susuki, Y., Krishnan, V., and Zhang, Y.: On analytical construction of observable functions in extended dynamic mode decomposition for nonlinear estimation and prediction, in: 2021 American Control Conference (ACC), pp. 4190–4195, <https://doi.org/10.23919/ACC50511.2021.9482747>, iSSN: 2378-5861, 2021.

O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, 1320 J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geoscientific Model Development*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, publisher: Copernicus GmbH, 2016.

Orbe, C., Yang, H., Waugh, D. W., Zeng, G., Morgenstern, O., Kinnison, D. E., Lamarque, J.-F., Tilmes, S., Plummer, D. A., Scinocca, J. F., Josse, B., Marecal, V., Jöckel, P., Oman, L. D., Strahan, S. E., Deushi, M., Tanaka, T. Y., Yoshida, K., Akiyoshi, H., Yamashita, Y., Stenke, A., Revell, L., Sukhodolov, T., Rozanov, E., Pitari, G., Visioni, D., Stone, K. A., Schofield, R., and Banerjee, A.: Large-scale tropospheric transport in the Chemistry–Climate Model Initiative (CCMI) simulations, *Atmospheric Chemistry and Physics*, 18, 7217–7235, <https://doi.org/10.5194/acp-18-7217-2018>, publisher: Copernicus GmbH, 2018.

Otto, S. E. and Rowley, C. W.: Koopman Operators for Estimation and Control of Dynamical Systems, *Annual Review of Control, Robotics, and Autonomous Systems*, 4, 59–87, <https://doi.org/10.1146/annurev-control-071020-010108>, 2021.

Pathak, J., Subramanian, S., Harrington, P., Raja, S., Chattopadhyay, A., Mardani, M., Kurth, T., Hall, D., Li, Z., Azizzadenesheli, K.,  
1330 Hassanzadeh, P., Kashinath, K., and Anandkumar, A.: FourCastNet: A Global Data-driven High-resolution Weather Model using Adaptive Fourier Neural Operators, <https://doi.org/10.48550/arXiv.2202.11214>, arXiv:2202.11214 [physics], 2022.

Proctor, J. L., Brunton, S. L., and Kutz, J. N.: Dynamic Mode Decomposition with Control, *SIAM Journal on Applied Dynamical Systems*, 15, 142–161, <https://doi.org/10.1137/15M1013857>, publisher: Society for Industrial and Applied Mathematics, 2016.

Sandstad, M., Steinert, N. J., Baur, S., and Sanderson, B. M.: METEORv1.0.1: A novel framework for emulating multi-timescale regional  
1335 climate responses, <https://doi.org/10.5194/egusphere-2025-1038>, 2025.

Santer, B., Wigley, T. L., Schlesinger, M., and Mitchell, J.: Developing climate scenarios from equilibrium GCM results, *Tech. rep.*, Max-Planck-Institute für Meteorology, 1990.

Schlesinger, M. E., Malyshev, S., Rozanov, E. V., Yang, F., Andronova, N. G., De Vries, B., Grübler, A., Jiang, K., Masui, T., Morita, T., Penner, J., Pepper, W., Sankovski, A., and Zhang, Y.: Geographical Distributions of Temperature Change for Scenarios of Greenhouse Gas and  
1340 Sulfur Dioxide Emissions, *Technological Forecasting and Social Change*, 65, 167–193, [https://doi.org/10.1016/S0040-1625\(99\)00114-6](https://doi.org/10.1016/S0040-1625(99)00114-6), 2000.

Schmid, P. J.: Dynamic mode decomposition of numerical and experimental data, *Journal of Fluid Mechanics*, 656, 5–28, <https://doi.org/10.1017/S0022112010001217>, 2010.

Schmid, P. J.: Dynamic Mode Decomposition and Its Variants, *Annual Review of Fluid Mechanics*, 54, 225–254, 2022.

1345 Slawinska, J., Szekely, E., and Giannakis, D.: Data-Driven Koopman Analysis of Tropical Climate Space-Time Variability, <https://doi.org/10.48550/arXiv.1711.02526>, arXiv:1711.02526 [physics], 2017.

Souza, A. N.: Representing turbulent statistics with partitions of state space. Part 2. The compressible Euler equations, *Journal of Fluid Mechanics*, 997, A2, <https://doi.org/10.1017/jfm.2024.657>, 2024a.

Souza, A. N.: Representing turbulent statistics with partitions of state space. Part 1. Theory and methodology, *Journal of Fluid Mechanics*,  
1350 997, A1, <https://doi.org/10.1017/jfm.2024.658>, 2024b.

Souza, A. N. and Doering, C. R.: Maximal transport in the Lorenz equations, *Physics Letters A*, 379, 518–523, <https://doi.org/10.1016/j.physleta.2014.10.050>, 2015.

Souza, A. N. and Silvestri, S.: A Modified Bisecting K-Means for Approximating Transfer Operators: Application to the Lorenz Equations, <https://arxiv.org/abs/2412.03734>, 2024.

1355 Souza, A. N., Geogdzhayev, G., Ferrari, R., and Flierl, G. R.: A Statistical Emulator Design for Averaged Climate Fields, <https://doi.org/10.22541/essoar.172779540.09973901/v1>, 2024.

Stevens, B., Sherwood, S. C., Bony, S., and Webb, M. J.: Prospects for narrowing bounds on Earth's equilibrium climate sensitivity, *Earth's Future*, 4, 512–522, <https://doi.org/10.1002/2016EF000376>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2016EF000376>, 2016.

1360 Sudakov, I., Pokojovy, M., and Lyakhov, D.: Statistical mechanics in climate emulation: Challenges and perspectives, *Environmental Data Science*, 1, e16, <https://doi.org/10.1017/eds.2022.15>, 2022.

Tatebe, H. and Watanabe, M.: MIROC MIROC6 model output prepared for CMIP6 CMIP piControl, [https://www.wdc-climate.de/ui/entry?acronym=C6\\_5208751](https://www.wdc-climate.de/ui/entry?acronym=C6_5208751), 2023.

Tebaldi, C. and Arblaster, J. M.: Pattern scaling: Its strengths and limitations, and an update on the latest model simulations, *Climatic Change*, 1365 122, 459–471, <https://doi.org/10.1007/s10584-013-1032-9>, 2014.

Tebaldi, C. and Knutti, R.: Evaluating the accuracy of climate change pattern emulation for low warming targets, *Environmental Research Letters*, 13, 055 006, <https://doi.org/10.1088/1748-9326/aabef2>, publisher: IOP Publishing, 2018.

Tebaldi, C., Tebaldi, C., Selin, N. E., Ferrari, R., and Flierl, G.: Emulators of climate model output, <https://www.authorea.com/doi/full/10.22541/essoar.174721805.59111383/v1?commit=f5346ac101507c00bb1065bb0483c7e1038bcfc7>, 2025.

Thuburn, J.: Climate sensitivities via a Fokker–Planck adjoint approach, *Quarterly Journal of the Royal Meteorological Society*, 1370 131, 73–92, <https://doi.org/10.1256/qj.04.46>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1256/qj.04.46>, 2005.

Tu, J. H., Rowley, C. W., Luchtenburg, D. M., Brunton, S. L., and Kutz, J. N.: On Dynamic Mode Decomposition: Theory and Applications, *Journal of Computational Dynamics*, 1, 391–421, <https://doi.org/10.3934/jcd.2014.1.391>, arXiv:1312.0041 [math], 2014.

Van Vuuren, D., O'Neill, B., Tebaldi, C., Chini, L., Friedlingstein, P., Hasegawa, T., Riahi, K., Sanderson, B., Govindasamy, B., Bauer, N., 1375 Eyring, V., Fall, C., Frieler, K., Gidden, M., Gohar, L., Jones, A., King, A., Knutti, R., Kriegler, E., Lawrence, P., Lennard, C., Lowe, J., Mathison, C., Mehmood, S., Prado, L., Zhang, Q., Rose, S., Ruane, A., Schleussner, C.-F., Seferian, R., Sillmann, J., Smith, C., Sörensson, A., Panickal, S., Tachiiri, K., Vaughan, N., Vishwanathan, S., Yokohata, T., and Ziehn, T.: The Scenario Model Intercomparison Project for CMIP7 (ScenarioMIP-CMIP7), <https://doi.org/10.5194/egusphere-2024-3765>, 2025.

Wang, M., Souza, A. N., Ferrari, R., and Sapsis, T.: Stochastic Emulators of Spatially Resolved Extreme Temperatures of Earth System Models, 1380 <https://www.authorea.com/doi/full/10.22541/essoar.172858084.46299070?commit=9870bad85464686e7ecaad16fcd347b357da1f15>, 2025.

Watson-Parris, D., Rao, Y., Olivié, D., Seland, Ø., Nowack, P., Camps-Valls, G., Stier, P., Bouabid, S., Dewey, M., Fons, E., Gonzalez, J., Harder, P., Jeggle, K., Lenhardt, J., Manshausen, P., Novitasari, M., Ricard, L., and Roesch, C.: ClimateBench v1.0: A Benchmark for Data-Driven Climate Projections, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002954, 1385 <https://doi.org/10.1029/2021MS002954>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021MS002954>, 2022.

Wells, C. D., Jackson, L. S., Maycock, A. C., and Forster, P. M.: Understanding pattern scaling errors across a range of emissions pathways, *Earth System Dynamics*, 14, 817–834, <https://doi.org/10.5194/esd-14-817-2023>, publisher: Copernicus GmbH, 2023.

Wieners, K.-H., Giorgetta, M., Jungclaus, J., Reick, C., Esch, M., Bittner, M., Legutke, S., Schupfner, M., Wachsmann, F., Gayler, V., 1390 Haak, H., de Vrese, P., Raddatz, T., Mauritzen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler, S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters-von Gehlen, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: MPI-M MPI-ESM1.2-LR model output prepared for CMIP6 CMIP piControl, <https://hdl.handle.net/21.14106/e8f47f002749a0c617d25fee63fd6db0b96a0c04>, 2023.

Wilcox, L. J., Allen, R. J., Samset, B. H., Bollasina, M. A., Griffiths, P. T., Keeble, J., Lund, M. T., Makkonen, R., Merikanto, J., O'Donnell, 1395 D., Paynter, D. J., Persad, G. G., Rumbold, S. T., Takemura, T., Tsigaridis, K., Undorf, S., and Westervelt, D. M.: The Regional Aerosol Model Intercomparison Project (RAMIP), *Geoscientific Model Development*, 16, 4451–4479, <https://doi.org/10.5194/gmd-16-4451-2023>, publisher: Copernicus GmbH, 2023.

Williams, M. O., Kevrekidis, I. G., and Rowley, C. W.: A Data-Driven Approximation of the Koopman Operator: Extending Dynamic Mode Decomposition, *Journal of Nonlinear Science*, 25, 1307–1346, <https://doi.org/10.1007/s00332-015-9258-5>, arXiv:1408.4408 [math], 1400 2015.

Winkler, A. J. and Sierra, C. A.: Towards a New Generation of Impulse-Response Functions for Integrated Earth System Understanding and Climate Change Attribution, *Geophysical Research Letters*, 52, e2024GL112 295, <https://doi.org/10.1029/2024GL112295>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2024GL112295>, 2025.

Womack, C.: Source code for A theoretical framework to understand sources of error in Earth System Model emulation, 1405 <https://doi.org/10.5281/zenodo.17572065>, language: eng, 2025.

Womack, C. B., Giani, P., Eastham, S. D., and Selin, N. E.: Rapid Emulation of Spatially Resolved Temperature Response to Effective Radiative Forcing, *Journal of Advances in Modeling Earth Systems*, 17, e2024MS004 523, <https://doi.org/10.1029/2024MS004523>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2024MS004523>, 2025.

Yeung, W.-K. and Kong, F.-N.: Time domain deconvolution when the kernel has no spectral inverse, *IEEE transactions on acoustics, speech, and signal processing*, 34, 912–918, 1986.

1410 Zagli, N., Colbrook, M., Lucarini, V., Mezić, I., and Moroney, J.: Bridging the Gap between Koopmanism and Response Theory: Using Natural Variability to Predict Forced Response, <https://doi.org/10.48550/arXiv.2410.01622>, arXiv:2410.01622, 2024.

Zazula, D. and Gyergyek, L.: Direct frequency-domain deconvolution when the signals have no spectral inverse, *IEEE transactions on signal processing*, 41, 977–981, 1993.