

Supplement of

Measurement report: Dust impact on hygroscopicity and volatility of submicron aerosols: Based on the observation in April of Beijing

Xinyao Hu^{1,2}, Aoyuan Yu^{1,3}, Xiaojing Shen^{1,2}, Jiayuan Lu^{1,4}, Yangmei Zhang^{1,2}, Quan
5 Liu^{1,2}, Lei Liu^{1,2}, Linlin Liang^{1,2}, Hongfei Tong^{1,2}, Qianli Ma⁵, Shuxian Zhang⁵, Bing
Qi⁶, Rongguang Du⁶, Huizheng Che^{1,2}, Xiaoye Zhang^{1,2}, and Junying Sun^{1,2*}

¹State Key Laboratory of Severe Weather Meteorological Science and Technology, Chinese Academy
of Meteorological Sciences, Beijing, 100081, China

²Key Laboratory of Atmospheric Chemistry of CMA, Chinese Academy of Meteorological Sciences,
10 Beijing, 100081, China

³Dalian Meteorological Observatory, Dalian, 116000, China

⁴Department of Atmospheric and Oceanic Sciences and Institute of Atmospheric Sciences, Fudan
University, Shanghai, 200433, China

⁵Lin'an Atmosphere Background National Observation and Research Station, Lin'an 311307, Hangzhou,
15 China

⁶Hangzhou Meteorological Bureau, Hangzhou, 310051, China

*Correspondence to: Junying Sun (jysun@cma.gov.cn)




Figure S1. Mean hygroscopic growth factor probability density function (HGF-PDF)
20 and volatile shrink factor probability density function (VSF-PDF) for particles ranging

from 50 to 300 nm during the sampling period.




Figure S2. The 48-h back trajectories on April 15 calculated using the HYSPLIT model arriving at the CAMS site with a terminal height of 500 m a.g.l.