



Measurement report: New particle formation and aerosol properties at a newly founded atmospheric observatory at the Finnish Baltic Sea coast

Maija Peltola^{1,2}, Roseline Thakur¹, Kurt Spence³, Janne Lampilahti¹, Ronja Mäkelä¹, Sasu Karttunen¹, Ekaterina Ezhova¹, Sami Haapanala⁴, Aki Vähä¹, Juha Kangasluoma¹, Tommy Chan¹, Pauli Paasonen¹, Joanna Norkko³, Alf Norkko³, Markku Kulmala¹, and Mikael Ehn¹

¹Institute for Atmospheric and Earth System Research/Physics, University of Helsinki, 00014 Helsinki, Finland.

²Now at: Finnish Meteorological Institute, P.O. Box 503, 00101 Helsinki, Finland

³Tvärminne Zoological Station, University of Helsinki, 10900 Hanko, Finland.

⁴Suvilumi Ltd., Helsinki, Finland.

Correspondence: Maija Peltola (maija.peltola@fmi.fi) and Mikael Ehn (mikael.ehn@helsinki.fi)

Abstract. Aerosol formation is an important contributor to climate in different environments. Coastal ecosystems are of special interest, since their habitat types and species can be highly diverse which can lead to high variability in fluxes of aerosol precursor vapours. Here, we present the first results from an atmospheric observatory established in 2022 at the Tvärminne Zoological Station (TZS) on the southern coast of Finland as a part of the Centre for Coastal Ecosystem and Climate Change Research (CoastClim). This work is focused on new particle formation (NPF) and how it is influenced by the coastal environment. NPF occurred at TZS roughly as often as at the well-established boreal forest site Hyytiälä, 230 km north-north-east of TZS. However, only one-third of the events occurred during the same days at both stations. Similarly to Hyytiälä, NPF events were most likely to be observed when clean air masses from the Norwegian Sea pass over the boreal forest region. To understand the special characteristics of aerosol formation at the coast, we also studied local intermediate ion formation (LIIF), which can tell us about aerosol formation potential in the vicinity of the measurement site. LIIF was more abundant over the continent compared to the marine sector and it was favoured by low condensation sink and sunny dry conditions. This study presents an overview of aerosol size distributions at TZS and provides a basis for future work, which will aim to understand the sources of coastal aerosol precursors and their links to coastal ecology and biogeochemistry.

1 Introduction

The majority of global cloud condensation nuclei (CCN) are produced by new particle formation (NPF) in the atmosphere (Gordon et al., 2017). Even though the oceans cover over 70 % of the globe, most NPF studies have been conducted over continental regions (Kerminen et al., 2018). Recent research has highlighted the role of marine aerosols for the radiative properties of clouds (Rosenfeld et al., 2019) and the importance of marine aerosol observations in constraining climate models (Regayre et al., 2020). Marine secondary aerosol has been of interest since Charlson et al. (1987) first proposed that dimethyl sulfide emitted by phytoplankton could influence the climate by forming aerosols. This hypothesis boosted research on atmosphere-



biosphere interactions in marine environments (Quinn and Bates, 2011), and recent studies have found evidence of NPF occurring within the marine boundary layer (Zheng et al., 2021; Peltola et al., 2022). More advanced measurement techniques have also uncovered new chemical mechanisms that can form aerosols in marine environments including amines (Brean et al., 2021) and ammonia (Jokinen et al., 2018) together with sulfuric acid, as well as organics (Mayer et al., 2020; Zheng et al., 2020), and iodine species (Sipilä et al., 2016; He et al., 2021). Despite these recent advances, further studies on marine NPF are still needed.

Apart from several studies from Mace Head at the Irish coast and few studies from other locations strongly influenced by tidal processes (McFiggans et al., 2004; O'Connor et al., 2008; Grose et al., 2007; Furneaux et al., 2010; Sipilä et al., 2016), most marine NPF studies have focused on open ocean or polar conditions, while coastal areas remain less studied. Coastal zones are of special interest due to their highly diverse habitat types and species as well as significant influence from human activities. Coastal zones are also often characterized by dynamic phenomena such as coastal upwelling and downwelling that can induce strong changes in the coastal water properties, including nutrient transport, temperature, and salinity (e.g., Lehmann and Myrberg, 2008). More complexity is added to the coastal atmosphere interactions by the heterogeneity of surface forcing, which can lead to e.g. sea and land breezes. The Baltic Sea coast is a great example of an anthropogenically influenced coastal environment, because it has been impacted by eutrophication for a long time (Andersen et al., 2017). Measurements at the Baltic Sea coast have already shown a high variability of greenhouse gas emissions both seasonally and between habitat types (Roth et al., 2022). Contrary to open oceans where CO₂ fluxes are driven by temperature and concentration of dissolved inorganic carbon (Gallego et al., 2018), the carbon fluxes at the coastal Baltic Sea are dominated by biogeochemical processes and can be influenced by run-off of organic matter from land (Honkanen et al., 2024).

To fully evaluate the interactions between coastal areas and climate, it is not enough to only study greenhouse gases, but we need to also understand the role of aerosol formation in these areas since aerosols can have a cooling effect on climate. This relates to the concept of CarbonSink+, which accounts for both carbon sink and potential for local aerosol formation when estimating the climate impacts of various ecosystems (Kulmala et al., 2020; Ke et al., 2025).

Coastal areas can be a mixing spot of aerosol precursors from the sea and land. For example de Jonge et al. (2024) have shown that having aerosol precursors both from the sea and the land can boost CCN production. Past research in the Baltic Sea has identified cyanobacterial blooms as a source of iodic acid that formed new particles (Thakur et al., 2021). This was however connected to air masses coming from the open Baltic Sea and coastal contribution could not be partitioned from the open sea contribution. One way to study the new particle formation potential in different locations is to focus on local aerosol production, which can be studied by examining the concentrations of freshly formed 2-2.3 nm particles which must have formed close to the measurement location since they have not yet grown to larger sizes (Kulmala et al., 2024; Tuovinen et al., 2024). This is known as local intermediate ion formation (LIIF) and we use this approach also in this study.

The best way to tackle the challenges related to understanding coastal climate interactions is through multidisciplinary collaboration and long-term measurements. As a part of the Centre for Coastal Ecosystem and Climate Change Research (CoastClim, <https://www.coastclim.org/>), we have recently established permanent atmospheric observations at the Tvärminne Zoological Station, located at the southern coast of Finland, where marine measurements have been conducted for nearly a



century (see e.g., Goebeler et al., 2022). This observatory conducts comprehensive atmospheric measurements spanning from greenhouse and trace gases to chemical composition of volatile organic compounds and their oxidation products to aerosols ranging in size from clusters to coarse mode. In the future the atmospheric parameters will also be connected to continuous measurements of seawater properties, enabling us to directly measure the interactions between the coastal ecosystem and climate.

In this work we show the first results from the aerosol measurements of a newly founded coastal station with a focus on understanding NPF and the effects that the coastal environment and meteorological conditions have on particle formation at this site. First, we study the NPF event frequency at the station and compare it to a boreal forest site. Then we focus on more local phenomena and study intermediate ions that have formed in the vicinity of the station. We compare the concentrations of these ions in marine and continental wind sectors to understand how the environment influences particle formation. We also study the effect of meteorological parameters and trace gas concentrations on particle and local intermediate ion formation.

2 Methods

2.1 Measurements

This paper is focused on aerosol and ion data measured at the Tvärminne Zoological Station, a research station founded in 1902 and located at the southern coast of Finland (TZS, Fig. 1). In 2022, we started a new atmospheric observatory at the station. The observatory for aerosol, ion, and aerosol precursor measurements consists of two trailers at the coast, less than 10 m from the sea, and one container located 160 m north-west from the trailers further inland. The observatory also includes greenhouse gas observations and flux measurements using an eddy covariance system located on a small nearby island 130 m east from the trailers (Fig. B3).

Our analysis is focused on data from a Neutral cluster and Air Ion Spectrometer (NAIS, Airel Ltd, Mirme and Mirme, 2013), located in one of the trailers at the coast (Fig. B3). The NAIS was first installed at the station on 26 February 2022. Initially the instrument was installed near the main building of the station, 300 m north-east from the final measurement location in the trailer at the coast. There it ran until 11 April 2022 and after a short break it was installed at the final location at the coast on 1 June 2022. Here we use its measurement from 26 February 2022 to 15 June 2024. The NAIS measures the size distribution of 2.5–42 nm particles and 0.8–42 nm negative and positive ions. The total particle size distribution is measured with the NAIS in both positive and negative modes, but here we used only the negative mode. We also used NAIS data from the Hyytiälä Forest Station (see e.g., Manninen et al., 2009), located 230 km north-north-east from Tvärminne, for the same time period to understand how our coastal observatory compares to this well-established boreal forest station.

In addition to the NAIS data, we use data from a custom-made Differential Mobility Particle Sizer (DMPS, e.g. Aalto et al., 2001) for the size distribution of 10–800 nm particles and a TSI Aerodynamic Particle Sizer 3321 (APS) for 500 nm – 20 µm particles. These data were available 22 May 2023 to 29 June 2023 and 1 February 2024 to 24 June 2024. For trace gas measurements, we used a Teledyne API T-400 ozone analyser, Teledyne API T300U carbon monoxide analyser, Teledyne API T200UP NO-NO₂ analyser and a Thermo Scientific 43i-TLE sulfur dioxide analyser, all located in the container further




Figure 1. Map of the location of Tvärminne Zoological Station (TZS) and Hyytiälä Forest Station (HYY). The grey shaded area in the zoomed-in figure indicates the marine wind sector used in this work. The maps are modified from maps prepared using European Union's Copernicus Land Monitoring Service information.

inland. The gas analyser data are available starting from 22 December 2022 for SO_2 and from 30 January 2022 for the other instruments. In this paper data until 24 June 2024 is used. Apart from the NO_x analyser, which had technical issues related to on-site calibration, these were all calibrated on-site. All the trace gas instruments, the DMPS, and the APS were located in the container further inland. Meteorological data, including wind, global radiation, and temperature were collected at the flux measurement island.

2.2 Data analysis

A simple way to understand how often new particle formation events (NPFE) occur is to classify each day based the particle size distributions observed during that day. For this we used the NPFE classification criteria described by Dal Maso et al. (2005). In this classification, each day is classified either as a clear new particle formation day (Classes I and II), a non-event day, or undefined day when it is not clear if particle formation occurs or not. Class I and II events require observing nucleation mode (< 25 nm) particles and that particle mode needs to grow in size. Observing this requires the meteorological conditions to be somewhat homogeneous over long distances. Because of this requirement these NPFE can also be called regional events. The difference between Classes I and II is that for Class II events, the new aerosol mode diameter or concentration is fluctuating, making it difficult to define a growth rate for the event, whereas for Class I events, growth rate determination is possible. On undefined days either nucleation mode particles are observed for more than an hour, but they do not grow in size, or there is a growing Aitken mode (25-100 nm). On non-event days, no new < 25 nm particle mode lasting more than an hour nor growth in Aitken mode is found. For the event analysis, we used NAIS size distribution data (total particle concentration measured with the negative polarity) from both Tvärminne and Hyytiälä. After removing days with missing or partially missing data



or instrument malfunctions, Tvärminne had 735 and Hyytiälä 714 days of NAIS data, with 655 days where both stations had complete data.

For rest of the data analysis in this work, we averaged data from particle and gas measurements over 30 minutes to be able to compare variables with different time resolution. In most of the analysis we divided the data into marine and continental wind sector. This was done based on the wind direction measured on the flux measurement island. This simple approach was chosen to get a picture of how aerosol formation is influenced by whether the air mass was last travelling over the sea or land. If the wind direction was from the sea (100° - 240°) the data were considered marine and if the wind direction was from the continent ($< 100^{\circ}$ or $> 240^{\circ}$) the data were considered as continental (see Fig. 1). With this division, 46% of the data were considered marine data.

Appendix Fig. B4 shows the statistical distribution of wind direction at the station during different seasons. The most notable difference between the seasons is that during the spring and summer winds from the south-west (190° - 230°) are more dominant than during the autumn and winter. In the autumn and winter the wind direction is more evenly distributed, but the wind speeds are higher when they are coming from the west and south-west compared to the opposite direction. Higher wind speeds in the marine sector are expected for this location during the winter season (Tammelin et al., 2013) and they are overall typical during the winter in Finland (Riuttanen et al., 2013).

The local wind direction can be influenced by e.g. sea or land breeze and wind channelling between the islands of the archipelago, but these effects cannot be separated with this data set. The wind sector division can indicate whether the air mass last travelled over land or sea, which provides insights about the effects of local phenomena in these different environments, as such phenomena occur over short time scales. Analysing trace gas data (see Appendix A) also indicated that the marine sector is less polluted than the continental sector, which supports this division.

Despite being simple, the wind sector division can be especially suitable for studying local effects such as local intermediate ion formation (LIIF) at the coast. It was recently highlighted that LIIF can be studied by focusing on 2–2.3 nm negative ions since in this size range the particles have been freshly formed but have not yet grown to larger sizes, meaning that they must have originated close to the measurement location (Kulmala et al., 2024; Tuovinen et al., 2024). This means that even if the wind direction is somewhat local it should still tell if the air mass last passed over the sea or the continent. This is why, in this paper we focus our analysis on 2–2.3 nm negative ions. In addition to this narrow size range that indicates local particle production, we show results for negative cluster ions (sub-2 nm), intermediate ions (2–7 nm), and larger 7–25 nm ions that are still within the nucleation mode. Negative ion data from the NAIS was used to calculate all of these concentrations. Additionally, we used data from the DMPS and APS for aerosols in nucleation (10–25 nm), Aitken (25–100 nm), accumulation (100–1000 nm), and coarse mode particles (1000–20 000 nm). APS data were used for sizes starting from 600 nm and DMPS for smaller particles.

Size distribution data were used also for the calculation of condensation sink. The sink was calculated using sulfuric acid as the condensable species as described by (Kulmala et al., 2001). For this we combined data from NAIS (3–40 nm), DMPS (40–650 nm) and APS (> 650 nm) to obtain the full size distribution.



3 Results & discussion

3.1 Occurrence of new particle formation events

To understand how aerosol formation at a coastal site differs from a continental site, we compared new particle formation event (NPFE) frequencies in Tvärminne and the well-studied boreal forest station Hyytiälä, located 230 km north-north-east from Tvärminne. For this we used the Dal Maso et al. (2005) criteria described in the Section 2.2. Figure 2 summarises the results of NPFE classification for both stations and shows the seasonal distribution of the events. On average Tvärminne had 9.3 % clear NPFE days (Class I 1.5 % and Class II 7.8 %) while Hyytiälä had 8.2 % clear NPFE days (Class I 1.1 % and Class II 7.1 %). For undefined days, the fraction was 23.8 % for Tvärminne and 17.3 % for Hyytiälä, leaving 66.9 % and 74.4 % of the days as non-event days for Tvärminne and Hyytiälä, respectively. As seen in Figure 2, both stations show similar seasonal patterns with a peak in event frequencies in April and a smaller peak in the early autumn. Although the average event frequencies we report here are lower than for example Nieminen et al. (2014) reported earlier for Hyytiälä, the spring and early autumn maxima were also observed in their 16-year data set. The lower event frequency observed here compared to previous literature can be explained by several factors: 1) classifying events manually is subjective, 2) they used DMPS, not NAIS data, and 3) different years can have different event frequencies, depending on meteorological conditions etc.

Despite the similarities in event fractions and seasonal patterns at the two stations, only 37% of the events were observed on the same days at both stations. An inspection of air mass back trajectories calculated with HYSPLIT (Stein et al., 2015; Rolph et al., 2017) for Class I and II events suggests that air mass history is likely to explain these differences. During these clear regional NPFE days, the air masses arrived primarily from north or north-west (Appendix Fig. B5) whereas on non-event days the air mass back trajectories could be from any direction. This is also in line with previous results from Hyytiälä since there events typically occur when the air masses come from the sector 280°–30° (Nieminen et al., 2014). When the air masses come from this direction, the air has typically crossed over the Norwegian Sea and the Scandinavian Mountains and it is clean compared to other directions, and this clean air favours NPF (Riuttanen et al., 2013).

3.2 Effect of marine and continental air masses on ion concentrations

To better understand NPF in Tvärminne and the influence of the coastal environment on NPF processes, we explored aerosol data of different size ranges and compared aerosol concentrations from the marine and continental wind sectors. When comparing the medians of these ion concentrations in the two wind sectors, sub-2 nm ions had 39%, and 2–2.3 nm ions 22 % higher concentrations in continental air. For larger ions the differences were smaller, with both 2–7 nm and 7–25 nm ions having 12 % higher concentrations in continental air. Observing higher concentrations of < 25 nm negative ions in the continental air is in line with the results that NPFE's were observed when the air mass back trajectories arrived from north-west.

Figure 3 shows the diurnal cycles of negative ions in different size ranges. For sub-2 nm ions, the continental air has higher median concentrations throughout the day. This is sensible, since over land radon can increase ion production (Chen et al., 2016) compared to marine conditions. In the continental wind sector highest sub-2 nm ion concentrations are observed during




Figure 2. The seasonal cycle of the contributions of different new particle formation event classes in each month for a) Tvärminne and b) Hyytiälä.

the night and early morning. This is likely due to less mixing during the night. In the marine air, no clear diurnal pattern is observed.

175 For 2–2.3 nm ions, the differences between the air masses are less clear, with the differences in the median concentrations in continental and marine air being statistically insignificant during 10.00–13.30 h and 18.00–19.30 h. During other times the continental air had higher concentrations. For the full range of intermediate ions (2–7 nm) the continental air has statistically significantly higher medians than the marine air all times but 12.00–13.30 h, when the concentrations are more variable and the confidence interval of the median is larger. In this size range the highest 90th percentiles are observed around or after midday
180 which is in line with typical NPFE times.

For the larger 7–25 nm ions the differences between the wind sectors are less clear, with continental air having statistically significantly higher concentrations only during 6.00–7.30 h and after 12 h. Having clearer differences in concentrations after midday is sensible considering that events happen mainly over land and in the afternoon particles produced by NPF have grown past 7 nm.

185 Comparing the diurnal cycles during different seasons (Appendix Figs. B6-B9) shows that the afternoon maximum in the 2–2.3 nm and 2–7 nm ion concentrations is more pronounced during the spring and summer compared to autumn and winter. This can also be expected, considering that there is more radiation in the summer and NPF occurrence peaked in the spring. Other clear differences between seasons include clearer seasonal cycle for sub-2 nm ions in the spring and less clear diurnal cycles and differences in air masses for all size ranges in the winter compared to other seasons. In the winter there is less
190 NPFE's and variations in boundary layer height are smaller so these results are expected.

Figure 4 shows the seasonal cycles for ions in the same size bins as Figure 3. The seasonal cycles are weak and vary between size ranges. For sub-2 nm ions the highest concentrations are observed in both wind sectors in the autumn (Fig. 4a) with continental concentrations higher throughout the seasons. The seasonal cycle is somewhat similar to that observed by Sulo et al. (2022) for Hyytiälä, with maximum cluster ion concentrations in the autumn.




Figure 3. Diurnal cycles of negative ions in different size ranges for marine and continental wind sectors. The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th-75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.

195 The intermediate ion size ranges (2-2.3 nm and 2-7 nm) have weak seasonal cycles with higher concentrations from late spring until early autumn compared to the rest of the year. Somewhat similar seasonal trends can be observed for the larger 7-25 nm ions. These trends are similar to the seasonal cycle that was observed for NPFE's.

These results show that more locally and regionally produced particles are observed when the wind direction is from the continent. The particle production is most intense during the day and in the spring and autumn. In the next sections we explore the effects of environmental conditions such as pre-existing particle population and meteorological parameters on NPF.

3.3 Effect of environmental conditions on particle formation

In this section we study the effects of condensation sink, meteorological parameters and trace gases on new particle formation observed in Tvärminne.

3.3.1 Effect of pre-existing particle population

205 Since particle formation depends on the relative strength of sources and sinks, we first explore the effect of condensation sink. Condensation sink is used to represent how fast condensable vapours are lost to the total surface area of particles. Figure 5 shows the average size distribution from all the particle instruments when DMPS data were available (22 May 2023 to 29 June 2023 and 2 January 2024 to 24 June 2024). From this figure and Appendix Figure B10, we can see that the number concentrations of negative ions and particles are higher in the continental wind sector in size ranges up to 30-40 nm. This is likely explained by NPF over the continent. From approximately 40 nm to 5000 nm marine air has slightly higher concentrations than continental




Figure 4. Seasonal cycles of ions in different size ranges for marine and continental wind sectors. The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.

air. The reason for this could be related to more long-range transported particles or sea spray in air masses coming from the marine sector.

One notable thing in Figure 5 is that the particle concentrations measured with NAIS are higher than those measured with the DMPS. This is expected since NAIS is known to show higher particle concentrations compared to DMPS systems (Kangasluoma et al., 2020). This discrepancy does not influence interpretation of the results in this analysis.

Figure 6 illustrates the diurnal cycles of particle concentrations measured with the DMPS and APS in nucleation (10–25 nm), Aitken (25–100 nm), accumulation (100–1000 nm), and coarse (1000–20 000 nm) modes and the differences seen between the wind sectors in different size ranges are in line with the previous results. In the nucleation mode the continental air has clearly higher concentrations than marine air during the day (8.00–17.30 h) which is in line with the results seen with negative ions. In Aitken mode continental air has lower concentrations than marine air around midday (10.00–13.30 h) while rest of the day the differences are statistically insignificant. Similar but stronger pattern is seen for accumulation mode particles, which have lower concentrations in continental air for most of the day with the differences being clearest around midday. These lower concentrations in the continental wind sector are likely related to clean air masses coming from the north-west and are one of the main reasons why particle formation is being favoured in this wind sector. This effect is the most pronounced during the day, when the mixed layer height increases and clean air from the free troposphere dilutes aerosol concentrations in the mixed layer. This has been shown to favour NPF (Wu et al., 2021) which is in line with seeing the highest nucleation mode particle concentrations around the same time. As can be expected in a location far from major pollution sources, coarse mode particle




Figure 5. Average size distribution from all instruments when DMPS data is available (22 May 2023 to 29 June 2023 and 1 February 2024 to 24 June 2024). All blue lines are for the marine wind sector and all the green lines for the continental wind sectors whereas the different line styles are for different instruments as indicated by the legend.




Figure 6. Diurnal cycles of 10–20 000 nm particles in different size ranges for coastal and continental air. The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.

concentrations were low. During the day the two wind sectors had similar concentrations whereas during the night the marine sector had higher concentrations.

230 Figure 7 shows the diurnal cycle for the calculated condensation sink. As the surface area of the particles is dominated by accumulation mode particles, it is no surprise that the CS was higher in marine air. The median CS for the period when




Figure 7. The diurnal cycle of condensation sink when data from all particle instruments is available (22.5.2023–29.6.2023 and 2.1.2024–24.6.2024). The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.

data from all particle instruments was available is $2.5 \cdot 10^{-3} \text{ s}^{-1}$ and the ratio of median CS in marine wind sector divided by median of CS in continental wind sector is 1.34. Similarly to accumulation mode particles, which likely dominate the CS, the difference in the two air masses is clearest during the early day (8.00–17.30 h) and this is likely explained by boundary layer dynamics as explained earlier.

Since DMPS and APS data were available only for part of the time, we calculated the diurnal cycle for NAIS data for this time period separately (Fig. B11). The diurnal cycles and wind sector differences of ions during this time period are similar to those observed for the whole data set in Figure 3. The differences in median intermediate ion concentrations between the two wind sectors during this period are smaller than the differences in condensation sink during daytime. Condensation sink is thus one of the key players in particle formation at this location, but not the only factor.

These results support previous research for Hyytiälä that has shown that NPFE's are favoured when clean air masses come from the north-western wind sector having crossed the Norwegian Sea and Scandinavian Mountains and later land over Finland (Riuttanen et al., 2013). The results also show that one of the factors favouring NPF during the day in the continental sector is the reduction in CS related to boundary layer dynamics.

3.3.2 Influence of meteorology and trace gases

To better understand the phenomena influencing particle formation at the observatory, we studied the effect of local meteorological conditions and trace gas levels on particle formation. Here, we focus first on the 2–2.3 nm negative ions and meteorological conditions. The general behaviour of trace gases is described in Appendix Section A in order to keep the focus of the main text on aerosols.




Figure 8. This figure illustrates how the 2–2.3 nm negative ion concentrations vary as a function of wind direction and other meteorological variables by plotting the median ion concentration in a given wind direction and a) wind speed, b) total radiation and c) relative humidity bins. The colour of each square shows the median concentration of 2–2.3 nm negative ions in that wind direction and a) wind speed, b) total radiation and c) relative humidity range.

Figure 8 shows the average 2–2.3 nm ion concentration binned with mean wind speed and direction, total incident radiation and relative humidity. All the plots show that there are more ions when the wind direction is from the north-west. This is in line with the previous results that NPF is most likely to occur in air masses from this direction. There are also more ions when wind speeds are high, total incident radiation is high and relative humidity is low. Sunny conditions favour photochemistry which can drive particle formation. Relative humidity is often anti-correlated with radiation and high relative humidity can be associated with cloudy or rainy conditions. This explains why the lowest 2–2.3 nm ion concentrations are seen when relative humidity is high. These results are in line with previous work done at Hyytiälä (e.g., Hamed et al., 2011; Dada et al., 2018) and other boreal forest sites (Garmash et al., 2024).

As noted in the previous section, the condensation sink can be one of the factors explaining higher 2–2.3 nm negative ion concentrations in continental air compared to marine air. Figure B12 shows the average condensation sink in wind direction and wind speed bins. It indicates that the condensation sink is at its highest when wind speed is low (below 5 ms^{-1}) and the wind direction from south-east. The lowest sink values are observed when wind speed is above 4 ms^{-1} and from north-west. This is largely opposite to what was seen for the 2–2.3 nm ions, indicating again that low condensation sink favours the formation of 2–2.3 nm ions.

To further study the relationship of 2–2.3 nm negative ions and different variables, Figure 9 shows the correlations of 2–2.3 nm ions with air temperature, total radiation, relative humidity, and ozone separately for the marine and continental wind sectors. The correlations are overall weak, but have similar directions in both of the wind sectors. Positive correlations are observed with temperature, total incident radiation, and ozone, whereas relative humidity has a negative correlation with ion




Figure 9. Concentration of 2–2.3 nm ions as a function of a) temperature, b) total radiation, c) relative humidity, and d) ozone in marine (blue) and continental (green) wind sectors

concentrations as can be expected from previous results. Positive correlations with radiation and temperature are in line with the previous results from the wind direction bin plots. Ozone can act as an oxidiser of VOCs and thus influence NPF, but it is also produced by radiation, so it is difficult to distinguish its role from that of photochemistry.

We also analysed these results limiting the data to only summer daytime data (Appendix Fig. B13) to limit the effect of different seasons and time of day since different processes can influence ion formation in different seasons. With this limitation, correlations between the 2-2.3 nm ions and temperature turn negative. This could be related to warmer air masses being more aged and thus having higher condensation sink that limits the production of new particles. Another reason for this could be lower volatilities of particle precursor vapours in colder air, which could favour condensation under these conditions (e.g., Stolzenburg et al., 2018).

With radiation, 2-2.3 nm ions in the continental wind sector still had a positive correlation, but in marine air there was no significant correlation. The correlations with relative humidity remained negative and were slightly stronger in both wind sectors with the limited data compared to the whole data set. With ozone there were no statistically significant correlations in either of the wind sectors.

To further understand the data, correlation coefficients were calculated for ions in different size ranges (< 2 nm, 2–2.3 nm, 2–7 nm, 7–25 nm), meteorological conditions (wind speed, temperature, radiation, relative humidity), and trace gases (CO, O₃, SO₂, NO_x). This was done separately for the whole data set (Fig. B14), only for marine air (Fig. B15), only for continental air (Fig. B16), and only for summer (June–August) day-time (10–15 h) data (Fig. B17).



285 For the whole data set (Fig. B14), the observed correlations are primarily weak. Apart from sub-2 nm and 7–25 nm ions which correlate negatively with each other, the ions in different size ranges have positive correlations with each other which is sensible since for example 2–2.3 is a subset of 2–7 ions and if these ions grow larger they can reach the 7–25 nm size range. All the size ranges have weak positive correlations with temperature and radiation and negative correlations with RH. This is in line with the previously presented results.

290 Focusing on the relationship of ions and trace gases, ions in sub-7 nm size ranges have negative correlations with NO_x , which is reasonable since the NO_x has higher concentrations in the marine air and marine CS was higher, suppressing NPF. The larger 7–25 nm ions have weak positive correlations with all the trace gases but CO. The 7–25 nm ions represent more aged particles that have sources further away from the station and they can have sources over land just like the trace gases. Ozone has weak positive correlations also with the ions in the intermediate ion classes (2–2.3 nm and 2–7 nm). This was
295 already seen in the scatter plots of 2–3.3 nm ions and ozone. Here we can see that ozone has even stronger correlations with temperature and radiation and as ozone itself is formed through photochemistry, the positive correlation with ozone and ions can also just be explained with both ozone and particles being formed in warm sunny conditions.

If the data set is limited to only marine (Fig. B15) or only continental (Fig. B16) wind sectors, the trends are largely similar to what was seen with the whole data set. When looking at only daytime data (Fig. B17), the correlations with temperature,
300 radiation, and ozone get stronger, but the trends are still similar. If the data are limited to only summertime and day data (Fig. B18), some of the correlations between particles and temperature or radiation turn weakly negative and positive correlations between ozone and particles are no longer observed. This would support the assumption that the previously observed correlation between the ions and ozone was caused by the fact that they are both produced in warm and sunny conditions. This is also in line with the results shown earlier for the 2–2.3 nm ions that indicated that particle formation is not favoured by higher
305 temperatures but just more likely to occur during the day when photochemistry plays a role and temperatures are in general higher.

To summarise, particle formation in Tvärminne is favoured by high total incident radiation and low relative humidity. The trace gas analysis suggested that higher levels of most trace gases are related to higher condensation sink that inhibited particle formation. Ozone was an exception to this since both ozone and particles are typically formed in sunny conditions.

310 4 Conclusions

Here we reported the first results of new particle formation and aerosol physical properties at a new coastal atmospheric observatory in Tvärminne, Southern Finland. We showed that new particle formation events (NPFE) occur at the coast approximately as often as at the boreal forest station Hyytiälä, located 230 km north-north-east of Tvärminne. These regional NPFE occurred when air masses arrived from the north-west and crossed over land. Northwestern air masses have also been connected to
315 regional NPFE before in Hyytiälä as these air masses are typically clean and have low concentrations of pre-existing particles that could inhibit NPF.



Further analysis focused on understanding the local effects of the coast and for this we compared two wind sectors, one where the winds were coming from the sea and another where the winds were coming from the continent. The results of this analysis were in line with the NPFE analysis and showed the concentrations of ions and particles in nucleation mode size ranges (2–25 nm) were higher when the winds were from the continent, particularly from the north-west. Studying local intermediate ion formation (LIIF) showed that in the near vicinity of the measurement site, more particles were formed over the continent than over the sea.

One factor favouring NPF in the continental wind sector was the clean air likely transported from the Norwegian Sea. This lead to lower condensation sink in the continental wind sector especially during the day when clean air was likely mixed down from the free troposphere and that favoured the occurrence of NPFE. Overall, NPF was favoured by high radiation levels and low relative humidity which is in line with previous research from other stations. These correlations were however low, so more research is needed to better understand the driving factors of NPF at the station.

Overall, our results are well in line with previous research from the thoroughly investigated Hyytiälä boreal forest site. Although clear differences were seen between the continental and marine wind sectors, more research is needed to distinguish how potential coastal sources can influence aerosol formation processes at the Baltic Sea coast and further downwind away from the coast. This will require not only deeper understanding of the local meteorology and coastal oceanographical processes, but also information about the chemical composition of aerosols and their precursors. Eventually knowledge on the coastal aerosol formation processes can be integrated with information about the coastal biogeochemistry and biodiversity. This way we can work towards fully understanding coastal climate interactions.

Code and data availability. The data are publicly available in Zenodo at <https://doi.org/10.5281/zenodo.16693683> (Peltola et al., 2025). The code used for wind rose plots was by Perreira (2024).

Appendix A: Trace gases

Trace gases can help us understand the sources and chemical make-up of air in the two wind sectors. As the Tvärminne station is situated in a nature conservation area, there are no major sources of anthropogenic pollutants in the near vicinity of the station and the levels of trace gases are overall low and similar to the levels observed in a 13 year long data set from Hyytiälä (Riuttanen et al., 2013). Figures B1 and B2 show the diurnal and seasonal cycles of carbon monoxide (CO), ozone (O₃), sulphur dioxide (SO₂), and nitrogen oxides (NO_x). For the seasonal cycles, months were included only if they had a minimum of one week of data per wind sector.

Comparing the median levels of the gases in the two wind sectors shows that while CO and SO₂ have higher levels in the continental wind sector, O₃ and NO_x are on average higher in the marine wind sector. Both CO and SO₂ are produced by anthropogenic sources and there are more of these sources over land, so having higher levels over land is reasonable. Ozone levels are lower in the continental wind sector, because other pollutants consume ozone. Higher NO_x levels in the marine



sector are likely related long-range transport as explained later in this section. Next we go through the observed trends for each gas separately.

350 CO is formed in incomplete combustion and its major sources in Finland include traffic and small-scale combustion, such as heating homes with wood burning. The main sink for CO is hydroxyl radicals, which are produced in sunny conditions and are thus most abundant in the summer (Holloway et al., 2000). This explains the observed seasonal cycle (Fig. B2a) in which we can see the lowest CO levels in the summer. The seasonal trends observed here are in line with the results observed previously in Helsinki and Sodankylä by Karppinen et al. (2024). This is expected since the lifetime of CO is relatively long, and we do
355 not expect any strong CO sources in the near vicinity of the Tvärminne station.

The diurnal cycle of CO has a maximum in the late afternoon (16–18 h) and a minimum in the morning (6–8 h) in both wind sectors. The diurnal cycle observed here is related to emissions accumulating in the boundary layer during the day and our results are somewhat similar to what has been observed in Helsinki during the weekends and less pronounced than what was observed in Helsinki during the weekdays (Järvi et al., 2009). This is reasonable since the traffic volumes in Helsinki especially
360 during the week are much higher than those in the Tvärminne region.

The diurnal cycle of ozone (Fig. B1b) is fairly similar to CO, with the highest values observed in the afternoon (14–16 h) and lowest values late at night (02–06 h). This is somewhat similar to what has been observed before in Hyytiälä (Lyubovtseva et al., 2005) and Helsinki (Järvi et al., 2009), since during the day ozone can be produced by photolysis and mixed down from higher altitudes. The differences between the two wind sectors are small, but during the day, the marine sector has statistically
365 significantly higher concentrations. The seasonal cycle of ozone (Fig. B2b) shows highest levels in both wind sectors in the late spring. This has been observed previously for example in Hyytiälä (Lyubovtseva et al., 2005; Riuttanen et al., 2013), because the radiation levels are high in the summer and ozone is produced by photolysis. The fact that ozone levels are higher in the marine sector is in line with previous work by Riuttanen et al. (2013) who observed that in Hyytiälä in the winter the ozone concentrations are higher when the air masses come from marine areas where the levels of ozone deposition and ozone
370 destroying pollutants are low. The spring ozone maximum coincided also with the maximum in NPF event frequency which is rational since both ozone and particles can be produced by photochemical reactions and ozone can oxidize many vapours so that they become less volatile and are more likely to form particles.

SO₂ has a weak diurnal cycle with both wind sectors reaching their highest concentrations during the day (Fig. B1c). The marine wind sector has lower concentrations than the continental wind sector during the night. The trend of having higher
375 SO₂ concentrations during the day is similar to what has been observed previously in Helsinki during the weekdays (Järvi et al., 2009). Looking at the seasonal cycle, SO₂ concentrations are the lowest in the late summer and early autumn (Fig B2c). This is reasonable as previous work in Hyytiälä has connected high SO₂ values in winter with more heating emissions, less photochemical losses, and temperature inversions trapping pollutants (Riuttanen et al., 2013).

For NO_x, the concentrations are in general low and in the continental wind sector the variations in concentration of each
380 month are larger than a possible diurnal cycle (Fig. B1d). In the marine sector we observe higher concentrations during the day. Previous results for Hyytiälä (Riuttanen et al., 2013) indicate that a possible reason for higher NO_x concentrations in our marine sector is long-range transport from south and south-east. In the seasonal cycle, more variability can be seen in the data



from February to June, with lowest values observed in late summer and early autumn (Fig. B2d). Similar seasonal trends have been observed before in Hyytiälä. There, the higher NO_x levels in the winter and early spring were associated with combustion
385 sources, low photochemical sink, and increased long-range transport (Riuttanen et al., 2013), and similar phenomena are likely to play a role in Tvärminne.




Figure B1. Diurnal cycles of different trace gases in coastal and continental air. Please note that no calibration factor was applied to the NO_x data due to technical difficulties. The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.




Figure B2. Seasonal cycles of different trace gases in coastal and continental air. Months were included only if they had a minimum of one week of data per wind sector. Please note that no calibration factor was applied to the NO_x data due to technical difficulties. The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.

Appendix B: Additional figures



Figure B3. The locations of 1. trailers including the NAIS, 2. the container with trace gas measurements, DMPS, and APS, 3. flux measurement island where meteorological data are measured, and 4. the main building of the station.




Figure B4. Wind roses for Tvärminne for different seasons.



Figure B5. Each black line is a 96 h air mass back trajectory for one Class I or II event day at 12 h for Tvärminne.




Figure B6. Diurnal cycles of ions in different size ranges for coastal and continental air in the spring (March-May). The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.




Figure B7. Diurnal cycles of ions in different size ranges for coastal and continental air in the summer (June-August). The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.




Figure B8. Diurnal cycles of ions in different size ranges for coastal and continental air in the autumn (September-November). The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.




Figure B9. Diurnal cycles of ions in different size ranges for coastal and continental air in the winter (December-February). The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.




Figure B10. Average marine size distribution divided by average continental size distribution.




Figure B11. Diurnal cycles of ions in different size ranges for coastal and continental air when DMPS data is available. The black dots indicate the medians, the triangles indicate the confidence interval of the medians, the boxes the 25th–75th percentiles, the crosses the 10th and 90th percentiles, and the whiskers cover approximately 99.3 % of the data.




Figure B12. Average condensation sink in wind direction and wind speed bins. Note that the data are included only if the wind speed and direction bin has at least 10 data points and the condensation sink data are available only when data from all particle instruments is available (22.5.2023–29.6.2023 and 2.1.2024–24.6.2024)




Figure B13. Concentration of 2–2.3 nm ions as a function of a) temperature, b) total radiation, c) relative humidity, and d) ozone only including data from summer and 10–15 h. The upper panel with blue data points contain data only from the marine wind sector and the lower panels with green data points data from the continental wind sector.




Figure B14. Spearman correlation coefficients between different variables for the whole dataset. Each square corresponds to the correlation coefficient between the variables on the x- and y-axes and the colour indicates the strength of the correlation with the square being white if the p-value was above 0.05.




Figure B15. Spearman correlation coefficients between different variables in marine air. Each square corresponds to the correlation coefficient between the variables on the x- and y-axes and the colour indicates the strength of the correlation with the square being white if the p-value was above 0.05.




Figure B16. Spearman correlation coefficients between different variables in air coming from the land. Each square corresponds to the correlation coefficient between the variables on the x- and y-axes and the colour indicates the strength of the correlation with the square being white if the p-value was above 0.05.




Figure B17. Spearman correlation coefficients between different variables during the day (10–15 h). Each square corresponds to the correlation coefficient between the variables on the x- and y-axes and the colour indicates the strength of the correlation with the square being white if the p-value was above 0.05.




Figure B18. Spearman correlation coefficients between different variables during the day (10–15 h) in the summer (June–August). Each square corresponds to the correlation coefficient between the variables on the x- and y-axes and the colour indicates the strength of the correlation with the square being white if the p-value was above 0.05.



Author contributions. MP conducted the data analysis and wrote the paper. MP, RT, KS, JL, SH, AV, JK, and TC were responsible for the measurements. All authors contributed to discussion of the results and provided input for the paper.

390 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This work was funded by Jane and Aatos Erkkö Foundation, Ella and Georg Ehrnrooth Foundation and Academy of Finland flagship ACCC (337549).



References

- Aalto, P., Hämeri, K., Becker, E., Weber, R., Salm, R., Mäkelä, J., Hoell, C., O'Dowd, C., Karlsson, H., Hansson, H., Väkevä, M., Koponen, I., Buzorius, G., and Kulmala, M.: Physical characterization of aerosol particles during nucleation events., *Tellus*, 53B, 344–358, 2001.
- Andersen, J. H., Carstensen, J., Conley, D. J., Dromph, K., Fleming-Lehtinen, V., Gustafsson, B. G., Josefson, A. B., Norkko, A., Villnäs, A., and Murray, C.: Long-term temporal and spatial trends in eutrophication status of the Baltic Sea, *Biological Reviews*, 92, 135–149, 2017.
- Brean, J., Dall'Osto, M., Simó, R., Shi, Z., Beddows, D., and Harrison, R. M.: Open ocean and coastal new particle formation from sulfuric acid and amines around the Antarctic Peninsula, *Nature Geoscience*, 14, 383–388, 2021.
- Charlson, R. J., Lovelock, J. E., Andreae, M. O., and Warren, S. G.: Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, 326, 655, 1987.
- Chen, X., Kerminen, V.-M., Paatero, J., Paasonen, P., Manninen, H. E., Nieminen, T., Petäjä, T., and Kulmala, M.: How do air ions reflect variations in ionising radiation in the lower atmosphere in a boreal forest?, *Atmospheric Chemistry and Physics*, 16, 14 297–14 315, <https://doi.org/10.5194/acp-16-14297-2016>, 2016.
- Dada, L., Chellapermal, R., Buenrostro Mazon, S., Paasonen, P., Lampilahti, J., Manninen, H. E., Junninen, H., Petäjä, T., Kerminen, V.-M., and Kulmala, M.: Refined classification and characterization of atmospheric new-particle formation events using air ions, *Atmospheric Chemistry and Physics*, 18, 17 883–17 893, 2018.
- Dal Maso, M., Kulmala, M., Riipinen, I., Wagner, R., Hussein, T., Aalto, P. P., and Lehtinen, K. E.: Formation and growth of fresh atmospheric aerosols: eight years of aerosol size distribution data from SMEAR II, Hyytiälä, Finland, *Boreal Environment Research*, 10, 323, 2005.
- de Jonge, R. W., Xavier, C., Olenius, T., Elm, J., Svenhag, C., Hyttinen, N., Nieradzik, L., Sarnela, N., Kristensson, A., Petäjä, T., Ehn, M., and Roldin, P.: Natural Marine Precursors Boost Continental New Particle Formation and Production of Cloud Condensation Nuclei, *Environmental Science & Technology*, 58, 10 956–10 968, <https://doi.org/10.1021/acs.est.4c01891>, 2024.
- Furneaux, K., Whalley, L., Heard, D., Atkinson, H., Bloss, W., Flynn, M., Gallagher, M., Ingham, T., Kramer, L., Lee, J., et al.: Measurements of iodine monoxide at a semi polluted coastal location, *Atmospheric Chemistry and Physics*, 10, 3645–3663, 2010.
- Gallego, M. A., Timmermann, A., Friedrich, T., and Zeebe, R. E.: Drivers of future seasonal cycle changes in oceanic $p\text{CO}_2$, *Biogeosciences*, 15, 5315–5327, <https://doi.org/10.5194/bg-15-5315-2018>, 2018.
- Garmash, O., Ezhova, E., Arshinov, M., Belan, B., Lampilahti, A., Davydov, D., Rätty, M., Aliaga, D., Baalbaki, R., Chan, T., et al.: Heatwave reveals potential for enhanced aerosol formation in Siberian boreal forest, *Environmental Research Letters*, 19, 014 047, 2024.
- Goebeler, N., Norkko, A., and Norkko, J.: Ninety years of coastal monitoring reveals baseline and extreme ocean temperatures are increasing off the Finnish coast, *Communications earth & environment*, 3, 215, 2022.
- Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M., Dunne, E. M., et al.: Causes and importance of new particle formation in the present-day and preindustrial atmospheres, *Journal of Geophysical Research: Atmospheres*, 122, 8739–8760, 2017.
- Grose, M. R., Cainey, J. M., McMinn, A., and Gibson, J. A.: Coastal marine methyl iodide source and links to new particle formation at Cape Grim during February 2006, *Environmental Chemistry*, 4, 172–177, 2007.
- Hamed, A., Korhonen, H., Sihto, S.-L., Joutsensaari, J., Järvinen, H., Petäjä, T., Arnold, F., Nieminen, T., Kulmala, M., Smith, J. N., et al.: The role of relative humidity in continental new particle formation, *Journal of Geophysical Research: Atmospheres*, 116, 2011.
- He, X.-C., Tham, Y. J., Dada, L., Wang, M., Finkenzeller, H., Stolzenburg, D., Iyer, S., Simon, M., Kürten, A., Shen, J., et al.: Role of iodine oxoacids in atmospheric aerosol nucleation, *Science*, 371, 589–595, 2021.



- 430 Holloway, T., Levy, H., and Kasibhatla, P.: Global distribution of carbon monoxide, *Journal of Geophysical Research: Atmospheres*, 105, 12 123–12 147, 2000.
- Honkanen, M., Aurela, M., Hatakka, J., Haraguchi, L., Kielosto, S., Mäkelä, T., Seppälä, J., Siiriä, S.-M., Stenbäck, K., Tuovinen, J.-P., Ylöstalo, P., and Laakso, L.: Interannual and seasonal variability of the air–sea CO₂ exchange at Utö in the coastal region of the Baltic Sea, *Biogeosciences*, 21, 4341–4359, <https://doi.org/10.5194/bg-21-4341-2024>, 2024.
- 435 Jokinen, T., Sipilä, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E.-M., Junninen, H., Kangasluoma, J., Manninen, H., Petäjä, T., et al.: Ion-induced sulfuric acid–ammonia nucleation drives particle formation in coastal Antarctica, *Science Advances*, 4, eaat9744, 2018.
- Järvi, J., Hannuniemi, H., Hussein, T., Junninen, H., Aalto, P. P., Hillamo, R., Mäkelä, T., Keronen, P., Siivola, E., Vesala, T., et al.: The urban measurement station SMEAR III: Continuous monitoring of air pollution and surface–atmosphere interactions in Helsinki, Finland, *Boreal environment research*, 14, 86, 2009.
- 440 Kangasluoma, J., Cai, R., Jiang, J., Deng, C., Stolzenburg, D., Ahonen, L. R., Chan, T., Fu, Y., Kim, C., Laurila, T. M., et al.: Overview of measurements and current instrumentation for 1–10 nm aerosol particle number size distributions, *Journal of Aerosol Science*, 148, 105 584, 2020.
- Karppinen, T., Sundström, A.-M., Lindqvist, H., Hatakka, J., and Tamminen, J.: Satellite-based assessment of national carbon monoxide concentrations for air quality reporting in Finland, *Remote Sensing Applications: Society and Environment*, 33, 101 120, 2024.
- 445 Ke, P., Lintunen, A., Kolari, P., Lohila, A., Tuovinen, S., Lampilahti, J., Thakur, R., Peltola, M., Peräkylä, O., Nieminen, T., Ezhova, E., Pihlatie, M., Laasonen, A., Koskinen, M., Rautakoski, H., Heimsch, L., Kokkonen, T., Vähä, A., Mammarella, I., Noe, S., Bäck, J., Kerminen, V.-M., and Kulmala, M.: Potential of carbon uptake and local aerosol production in boreal and hemi-boreal ecosystems across Finland and in Estonia, *Biogeosciences*, 22, 3235–3251, <https://doi.org/10.5194/bg-22-3235-2025>, 2025.
- Kerminen, V.-M., Chen, X., Vakkari, V., Petäjä, T., Kulmala, M., and Bianchi, F.: Atmospheric new particle formation and growth: review of
- 450 field observations, *Environmental Research Letters*, 13, 103 003, 2018.
- Kulmala, M., Maso, M. D., Mäkelä, J., Pirjola, L., Väkevä, M., Aalto, P., Miikkulainen, P., Hämeri, K., and O’dowd, C.: On the formation, growth and composition of nucleation mode particles, *Tellus B: Chemical and Physical Meteorology*, 53, 479–490, 2001.
- Kulmala, M., Ezhova, E., Kalliokoski, T., Noe, S., Vesala, T., Lohila, A., Liski, J., Makkonen, R., Bäck, J., Petäjä, T., and Kerminen, V.-M.: CarbonSink+—Accounting for multiple climate feedbacks from forests, *Boreal Environment Research*, 25, 1, 2020.
- 455 Kulmala, M., Ke, P., Lintunen, A., Peräkylä, O., Lohtander, A., Tuovinen, S., Lampilahti, J., Kolari, P., Schiestl-Aalto, P., Kokkonen, T., Nieminen, T., Dada, L., Ylivinkka, I., Petäjä, T., Bäck, J., Lohila, A., Heimsch, L., Ezhova, E., and Kerminen, V.-M.: A novel concept for assessing the potential of different boreal ecosystems to mitigate climate change (CarbonSink+ Potential), *Boreal Environment Research*, 29, 1–16, 2024.
- Lehmann, A. and Myrberg, K.: Upwelling in the Baltic Sea—A review, *Journal of Marine Systems*, 74, S3–S12, 2008.
- 460 Lyubovtseva, Y. S., Sogacheva, L., Dal Maso, M., Bonn, B., Keronen, P., and Kulmala, M.: Seasonal variations of trace gases, meteorological parameters, and formation of aerosols in boreal forests, *Boreal environment research*, 10, 493, 2005.
- Manninen, H. E., Petäjä, T., Asmi, E., Riipinen, I., Nieminen, T., MikkiIä, J., Hörrak, U., Mirme, A., Mirme, S., Laakso, L., et al.: Long-term field measurements of charged and neutral clusters using Neutral cluster and Air Ion Spectrometer (NAIS)., *Boreal Environment Research*, 14, 2009.
- 465 Mayer, K. J., Wang, X., Santander, M. V., Mitts, B. A., Sauer, J. S., Sultana, C. M., Cappa, C. D., and Prather, K. A.: Secondary marine aerosol plays a dominant role over primary sea spray aerosol in cloud formation, *ACS Central Science*, 6, 2259–2266, 2020.



- McFiggans, G., Coe, H., Burgess, R., Allan, J., Cubison, M., Alfarra, M. R., Saunders, R., Saiz-Lopez, A., Plane, J., Wevill, D., et al.: Direct evidence for coastal iodine particles from *Laminaria* macroalgae–linkage to emissions of molecular iodine, *Atmospheric Chemistry and Physics*, 4, 701–713, 2004.
- 470 Mirme, S. and Mirme, A.: The mathematical principles and design of the NAIS-a spectrometer for the measurement of cluster ion and nanometer aerosol size distributions, *Atmospheric Measurement Techniques*, 6, 1061, 2013.
- Nieminen, T., Asmi, A., Dal Maso, M., Aalto, P. P., Keronen, P., Petäjä, T., Kulmala, M., and Kerminen, V.-M.: Trends in atmospheric new-particle formation: 16 years of observations in a boreal-forest environment, *Boreal Environmental Research*, 19, 191–214, 2014.
- O'Connor, T., Jennings, S., and O'Dowd, C.: Highlights of fifty years of atmospheric aerosol research at Mace Head, *Atmospheric Research*, 475 90, 338–355, 2008.
- Peltola, M., Rose, C., Trueblood, J. V., Gray, S., Harvey, M., and Sellegri, K.: New particle formation in coastal New Zealand with a focus on open-ocean air masses, *Atmospheric Chemistry and Physics*, 22, 6231–6254, <https://doi.org/10.5194/acp-22-6231-2022>, 2022.
- Peltola, M., Thakur, R., Spence, K., Lampilahti, J., Mäkelä, R., Karttunen, S., Ezhova, E., Haapanala, S., Vähä, A., Kangasluoma, J., Chan, T., Paasonen, P., Norkko, J., Norkko, A., Kulmala, M., and Ehn, M.: Dataset for Measurement report: New particle formation and aerosol 480 properties at a newly founded atmospheric observatory at the Finnish Baltic Sea coast, <https://doi.org/10.5281/zenodo.16693683>, 2025.
- Perreira, D.: Wind Rose, Retrieved July 30, 2024., <https://www.mathworks.com/matlabcentral/fileexchange/47248-wind-rose>, 2024.
- Quinn, P. and Bates, T.: The case against climate regulation via oceanic phytoplankton sulphur emissions, *Nature*, 480, 51, 2011.
- Regayre, L. A., Schmale, J., Johnson, J. S., Tatzelt, C., Baccarini, A., Henning, S., Yoshioka, M., Stratmann, F., Gysel-Beer, M., Grosvenor, D. P., and Carslaw, K. S.: The value of remote marine aerosol measurements for constraining radiative forcing uncertainty, *Atmospheric 485 Chemistry and Physics*, 20, 10 063–10 072, <https://doi.org/10.5194/acp-20-10063-2020>, 2020.
- Riuttanen, L., Hulkkonen, M., Dal Maso, M., Junninen, H., and Kulmala, M.: Trajectory analysis of atmospheric transport of fine particles, SO₂, NO_x and O₃ to the SMEAR II station in Finland in 1996–2008, *Atmospheric Chemistry and Physics*, 13, 2153–2164, 2013.
- Rolph, G., Stein, A., and Stunder, B.: Real-time environmental applications and display system: READY, *Environmental Modelling & Software*, 95, 210–228, 2017.
- 490 Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 363, 2019.
- Roth, F., Sun, X., Geibel, M. C., Prytherch, J., Brüchert, V., Bonaglia, S., Broman, E., Nascimento, F., Norkko, A., and Humborg, C.: High spatiotemporal variability of methane concentrations challenges estimates of emissions across vegetated coastal ecosystems, *Global Change Biology*, 28, 4308–4322, <https://doi.org/https://doi.org/10.1111/gcb.16177>, 2022.
- 495 Sipilä, M., Sarnela, N., Jokinen, T., Henschel, H., Junninen, H., Kontkanen, J., Richters, S., Kangasluoma, J., Franchin, A., Peräkylä, O., et al.: Molecular-scale evidence of aerosol particle formation via sequential addition of HIO₃, *Nature*, 537, 532, 2016.
- Stein, A., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M., and Ngan, F.: NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bulletin of the American Meteorological Society*, 96, 2059–2077, 2015.
- Stolzenburg, D., Fischer, L., Vogel, A. L., Heinritzi, M., Schervish, M., Simon, M., Wagner, A. C., Dada, L., Ahonen, L. R., Amorim, A., 500 Baccarini, A., Bauer, P. S., Baumgartner, B., Bergen, A., Bianchi, F., Breitenlechner, M., Brilke, S., Mazon, S. B., Chen, D., Dias, A., Draper, D. C., Duplissy, J., Haddad, I. E., Finkenzeller, H., Frege, C., Fuchs, C., Garmash, O., Gordon, H., He, X., Helm, J., Hofbauer, V., Hoyle, C. R., Kim, C., Kirkby, J., Kontkanen, J., Kürten, A., Lampilahti, J., Lawler, M., Lehtipalo, K., Leiminger, M., Mai, H., Mathot, S., Mentler, B., Molteni, U., Nie, W., Nieminen, T., Nowak, J. B., Ojdanic, A., Onnela, A., Passananti, M., Petäjä, T., Quéléver, L. L. J., Rissanen, M. P., Sarnela, N., Schallhart, S., Tauber, C., Tomé, A., Wagner, R., Wang, M., Weitz, L., Wimmer, D., Xiao, M., Yan, C., Ye,



- 505 P., Zha, Q., Baltensperger, U., Curtius, J., Dommen, J., Flagan, R. C., Kulmala, M., Smith, J. N., Worsnop, D. R., Hansel, A., Donahue, N. M., and Winkler, P. M.: Rapid growth of organic aerosol nanoparticles over a wide tropospheric temperature range, *Proceedings of the National Academy of Sciences*, 115, 9122–9127, <https://doi.org/10.1073/pnas.1807604115>, 2018.
- Sulo, J., Lampilahti, J., Chen, X., Kontkanen, J., Nieminen, T., Kerminen, V.-M., Petäjä, T., Kulmala, M., and Lehtipalo, K.: Measurement report: Increasing trend of atmospheric ion concentrations in the boreal forest, *Atmospheric Chemistry and Physics*, 22, 15 223–15 242, <https://doi.org/10.5194/acp-22-15223-2022>, 2022.
- 510 Tammelin, B., Vihma, T., Atlaskin, E., Badger, J., Fortelius, C., Gregow, H., Horttanainen, M., Hyvönen, R., Kilpinen, J., Latikka, J., et al.: Production of the Finnish wind atlas, *Wind Energy*, 16, 19–35, 2013.
- Thakur, R. C., Dada, L., Beck, L. J., Quéléver, L. L. J., Chan, T., Marbouti, M., He, X.-C., Xavier, C., Sulo, J., Lampilahti, J., Lampimäki, M., Tham, Y. J., Sarnela, N., Lehtipalo, K., Norkko, A., Kulmala, M., Sipilä, M., and Jokinen, T.: An evaluation of new particle formation events in Helsinki during a Baltic Sea cyanobacterial summer bloom, *Atmospheric Chemistry and Physics Discussions*, 2021, 1–45, <https://doi.org/10.5194/acp-2021-729>, 2021.
- 515 Tuovinen, S., Lampilahti, J., Kerminen, V.-M., and Kulmala, M.: Intermediate ions as indicator for local new particle formation, *Aerosol Research*, 2, 93–105, <https://doi.org/10.5194/ar-2-93-2024>, 2024.
- Wu, H., Li, Z., Li, H., Luo, K., Wang, Y., Yan, P., Hu, F., Zhang, F., Sun, Y., Shang, D., et al.: The impact of the atmospheric turbulence-development tendency on new particle formation: a common finding on three continents, *National science review*, 8, nwaa157, 2021.
- 520 Zheng, G., Kuang, C., Uin, J., Watson, T., and Wang, J.: Large contribution of organics to condensational growth and formation of cloud condensation nuclei (CCN) in the remote marine boundary layer, *Atmospheric Chemistry and Physics*, 20, 12 515–12 525, <https://doi.org/10.5194/acp-20-12515-2020>, 2020.
- Zheng, G., Wang, Y., Wood, R., Jensen, M. P., Kuang, C., McCoy, I. L., Matthews, A., Mei, F., Tomlinson, J. M., Shilling, J. E., et al.: New particle formation in the remote marine boundary layer, *Nature communications*, 12, 1–10, 2021.
- 525