

Decoding the North Atlantic Ocean Circulation Breakthrough in the Aptian–Albian Transition

João. M. F. Ramos ^{1,2}, Jairo. F. Savian ^{1,3}, Daniel. R. Franco ⁴, Milene F. Figueiredo ⁵, Rodolfo Coccioni ⁶, Fabrizio Frontalini ⁷

¹Programa de Pós-Graduação em Geociências, Instituto de Geociências, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, 91501-970 Porto Alegre, RS, Brazil.

²Petróleo Brasileiro S.A., Rio de Janeiro (PETROBRAS), Av. Henrique Valadares 28, 20231-030 - Rio de Janeiro- RJ, Brazil.

³Instituto de Geociências, Universidade Federal do Rio Grande do Sul, Av. Bento Gonçalves 9500, 91501-970 Porto Alegre, RS, Brazil.

⁴Coordenação de Geofísica, Observatório Nacional, R. General José Cristino 77, 20921-400 Rio de Janeiro, RJ, Brazil.

⁵Centro de Pesquisas e Desenvolvimento Leopoldo Américo Miguez de Mello, Petrobras Petróleo Brasileiro S.A, Avenida Horácio Macedo 950, 21941-915 Rio de Janeiro, Brazil.

⁶Università degli Studi di Urbino “Carlo Bo”, 61029 Urbino, Italy.

⁷Dipartimento di Scienze Pure e Applicate (DiSPeA), Università degli Studi di Urbino “Carlo Bo”, 61029 Urbino, Italy.

Correspondence to: João Mauricio Figueiredo Ramos (j.m.f.ramos@petrobras.com.br)

Abstract. The Cretaceous experienced marked climatic variability driven by large igneous province volcanism, monsoonal dynamics and changes in ocean circulation, which strongly influenced the expression of oceanic anoxic events and Cretaceous Oceanic Red Beds (CORBs). Here we integrate cyclostratigraphic analysis of magnetic susceptibility records from Ocean Drilling Program Site 1049 (western North Atlantic) to evaluate the timing and synchronicity of CORB deposition between the Tethyan and North Atlantic realms. Our results show that long-lived Aptian CORBs coincided with global cooling and enhanced thermohaline circulation, whereas Albian CORBs were shorter lived and orbitally paced, reflecting greenhouse conditions dominated by circulation driven primarily by mesoscale eddies. Astrochronological tuning further constrains two short geomagnetic polarity reversals within the Cretaceous Normal Polarity Superchron, dating the M–2r reversal to 110.76 Ma (duration ~150 kyr) and placing reversed-polarity subchron 3 between 111.45 and 111.53 Ma. These ages provide robust geochronological tie points for Aptian–Albian stratigraphic correlations.

1 Introduction

35 The Aptian–Albian interval (121.4 - 100.5 Ma; Gale et al., 2020) was marked by climatic shifts and extremely warm climate (Savin, 1977; Weissert, 1989; Huber et al., 2018), which was much warmer than that of today (Friedrich et al., 2012; O'Brien et al., 2017). These significant paleoclimatic shifts driven by intense volcanism associated with the emplacement of large igneous provinces (LIPs) (Schlanger and Jenkyns, 1976; Tarduno et al., 1991; Larson and Erba, 1999; Percival et al., 2024; Li et al., 2024), elevated production of ocean crust (Larson, 1991), enhanced monsoonal activity (Matsumoto et al., 2022), and
40 changes in oceanic circulation (Weissert, 1981; Premoli Silva et al., 1989; Hay, 2008, 2009; Giorgioni et al., 2015; Bottini and Erba, 2018). These fluctuations played a crucial role in shaping the sedimentary record, contributing to both the deposition of organic-rich black shales which, with different definitions, are known in the literature as oceanic anoxic events (OAEs; Schlanger and Jenkyns, 1976, Leckie et al., 2002) and the formation of Cretaceous oceanic red beds (CORBs), which signal episodes of improved oceanic oxygenation (Premoli Silva et al., 1989; Erbacher et al., 2001; Wang et al., 2009; Li et al., 2011; Hu et al., 2012).

CORBs are fine-grained sedimentary rocks exhibiting red, pink, or brown hues, which are deposited in oxygen-rich pelagic marine settings (Hu et al., 2005a,b; Wang et al., 2005; Li et al., 2011). Their coloration is primarily attributed to iron oxides such as hematite and goethite, formed via syn-depositional oxidation and influenced by early diagenetic processes (Channell et al., 1982; Eren and Kadir, 2001; Li et al., 2011). Based on XRD and diffuse reflectance spectrophotometry (DRS) analyses,
50 Hu et al. (2006b) proposed that the red coloration of the ODP Site1049 sediments could be attributed to the presence of microscopic, finely dispersed hematite. The widespread presence of benthic foraminifera in CORBs suggests oligotrophic and oxic bottom-water conditions and shifts in their assemblages from dysoxic to oxic states further reflect stepwise improvements in ocean ventilation during their deposition (Erbacher et al., 2001; Wang et al., 2009; Kochhann et al., 2023).

Whereas OAE deposition has been linked to an intensified hydrological cycle and reduced vertical mixing (Fischer and Arthur, 1977; Sinninghe Damsté and Köster, 1998; Erbacher et al., 2001; Matsumoto et al., 2022), CORB formation reflects
55 contrasting environmental conditions, characterized by increased aridity, enhanced input of iron-rich aeolian dust, and intensified vertical mixing and bottom-current activity that promoted water-column ventilation and oxygenation (Hu et al., 2012; Giorgioni et al., 2017). Although the periodic influx of cooler deep waters is considered a key driver in CORB formation (Hu et al., 2005b), multiple factors may have contributed to the formation of CORBs (e.g., oxygen flux into pore waters, prolonged oxygen exposure, changes in sea level, basin geometry, organic matter content, and atmospheric dust fluxes; Hu et al., 2005a,b; Li et al., 2011; Hu et al., 2012), hence suggesting a complex and multifactorial origin for these deposits.

Among the proposed mechanisms for CORB formation, the presence of oxygenated bottom waters is considered the primary prerequisite for organic matter oxidation and the associated early diagenetic processes that characterize CORBs (Wang et al., 2009; Jansa and Hu, 2009). In this context, thermohaline circulation is widely recognized as the dominant mechanism
65 responsible for deep-ocean oxygenation (Yamamoto et al., 2015). A fundamental requirement for the establishment of such

circulation is the existence of strong latitudinal temperature gradients between cold polar regions and warm equatorial zones (Hay, 2008) – a condition generally considered atypical for the Cretaceous.

However, multiple lines of evidence indicate the occurrence of cooling events during the Aptian, despite the prevailing greenhouse climate, which may have enabled the development of a thermohaline circulation capable of modifying water-column stratification and promoting the oxidation of iron-bearing sediments in the presence of more oxygenated bottom waters (Hu et al., 2005a,b; Tiraboschi et al., 2009; Erbacher et al., 2001; Li et al., 2011; McAnena et al., 2013;). Accordingly, a link between CORB deposition and thermohaline circulation can be proposed for the Aptian Cold Snap (McAnena et al., 2013). Nevertheless, CORB occurrence is not restricted to the late Aptian. In the Tethyan realm, CORBs span from the Aptian to the Albian (Premoli Silva et al., 1989; Coccioni et al., 2012; Gambacorta et al., 2016), encompassing both typical Cretaceous greenhouse conditions and cooler interludes. A similar pattern is observed in the North Atlantic, where CORBs occur over the same time interval (Erbacher et al., 2001; Trabucho Alexandre et al., 2011; Li et al., 2011). The geochemical characteristics of these red beds, together with the duration of oxidation events across contrasting thermal regimes, likely hold key insights into Aptian–Albian paleoclimate evolution and ocean circulation modes.

In such context, the Ocean Drilling Program (ODP) Leg 171B, Site 1049, located at Blake Nose in the western North Atlantic, provides a unique opportunity to study short-term CORBs that appear to be orbitally-controlled (Ogg and Bardot, 2001). These CORB intervals are marked by distinctive sediment coloration and high magnetic susceptibility (MS) readings due to hematite presence (Li et al., 2011; Hu et al., 2012). Shipboard indications of two thin reversed-polarity zones in Holes from ODP Site 1049 (Norris et al., 1998) were later confirmed by Ogg and Bardot (2001), providing well-dated geomagnetic reversals based on global bioevents (Huber et al., 2011).

This study presents a magneto-cyclostratigraphic analysis based on spliced MS records from Holes A, B, and C of the ODP Site 1049 (Norris et al., 1998). The excellent preservation of planktonic foraminifera at this site (Erbacher et al., 2001) allows for precise dating of CORB intervals, as well as the estimation of sedimentation rates and durations. Astrochronological age models enable the temporal correlation of geological events across sedimentary basins, linking records from classical sections studied using different methodologies and deposited in distinct oceanic settings, such as ODP Site 1049 in the North Atlantic and the Poggio Le Guaine (PLG) section in the Tethyan realm.

By integrating orbital-cycle tuning of the MS dataset, the ages of short geomagnetic polarity reversals were constrained and updated in accordance with the latest Aptian–Albian biochronologies (Gale et al., 2020) and the ages of the key features of the carbon isotope curves (Ramos et al., 2024a,b). These reversals were subsequently correlated with other known events recorded during the Cretaceous Normal Superchron (CNPS) in multiple global locations (Baksi, 1995; Tarduno, 1990; Gilder et al., 2003; Zhang et al., 2021; Fauth et al., 2022; Ramos et al., 2024b). Our results provide new insights into the paleoclimate of the Aptian–Albian interval and offer a comprehensive review of the key mechanisms related to CORB formation under different circulation modes.

2 Geological setting

The Blake Nose, located in the western North Atlantic (Fig. 1), resembles an appendage on the eastern margin of the Blake Plateau, a gently sloping ramp that reaches a maximum depth of about 2700 m (Norris et al., 1998; Li et al., 2011). The Upper Cretaceous and Cenozoic deposits on the Blake Nose lie upon a sequence of Jurassic and Lower Cretaceous limestone. ODP Leg 171B drilled five sites along the Blake Nose transect, and three holes were drilled at Site 1049, positioned at 23°N during the Cretaceous, above the calcite compensation depth (CCD).

The Cretaceous sediments at this site consist mainly of nannofossil chalk and claystone, including clayey and organic-rich intervals, with red, white, green, and black beds (Norris et al., 1998). Abundant planktonic and benthic foraminifera with well-preserved glassy tests indicate minimal diagenetic alteration (Erbacher et al., 2001; Li et al., 2011). Several intervals at different Holes 1049A, 1049B and 1049C display color and compositional oscillations that appear to reflect a spectrum of Milankovitch orbital climate cycles (Ogg and Bardot, 2001). The laminated black shale, associated with OAE 1b (Norris et al., 1998; Erbacher et al., 2001; Huber and Leckie, 2011), is located within the *M. rischi* planktonic foraminiferal zone and near the base of the *Hayesites albiensis* nannofossil zone, linking it to the Urbino/Paquier Level (Coccioni et al., 2012, 2014; Ramos et al., 2024a).

The color of oceanic red beds, influenced by hematite and goethite, reflects syn-depositional oxidation and early diagenesis, with no evidence of late diagenetic alteration (Wang et al., 2009; Li et al., 2011). At ODP Site 1049, the formation of iron oxides under oxic bottom-water conditions is supported by the presence of a flourishing benthic community (Wang et al., 2009; Norris et al., 1998; Kochhann et al., 2023). These conditions were likely driven by low organic matter accumulation and high dissolved oxygen content in bottom waters (Li et al., 2011).

The Poggio Le Guaine (PLG) core, drilled in the Umbria–Marche Basin of central Italy, includes some of the most complete Aptian and Albian sedimentary successions known from the Tethyan Realm and provides the basis for an accurate and precise calibration of the Paleogene time scale (Coccioni et al., 2012; Leandro et al., 2022; Ramos et al., 2026). The astrochronology performed on this core, based on magnetic susceptibility, provides the most detailed zonation of OAE 1b (and its sub-events), as well as the definition of the ages of the main features of the carbon-isotope curve associated with this anoxic event (Ramos et al., 2024a), sequenced using Greek letters (Fig. 3). These C-marks (prominent features of the stable $\delta^{13}\text{C}$ curve, including excursions, peaks, troughs, and shifts in the long-term behavior of the carbon isotope records, as described in Ramos et al., 2024a), due to the global nature of the carbon-related isotopic anomalies (Weissert, 1989), incorporate chemostratigraphic tie-points that enable long-distance correlations between successions in different sedimentary basins (Ramos et al., 2024a).





Figure 1: Top: Early Albian paleogeographic reconstructions at 110 Ma with the location of the ODP Site 1049, PLG core and the other sites/outcrops used for the short geomagnetic reversed-polarity correlations. Sea Level +80 m, Mollweide Projection. Modified from CR Scotese, PALEOMAP Project. Bottom: a) The stratigraphic framework with age (Ma), planktonic foraminiferal and calcareous nannofossils zones (Huber and Leckie, 2011; Coccioni et al., 2012; Gale et al., 2020; Ramos et al., 2024a,b). b) Geomagnetic polarity and reversal identifications (Savian et al., 2016; Gale et al., 2020; Li et al., 2023; Ramos et al., 2024b, 2025). c) Tuoyun section (Gilder et al., 2003). d) Raajmahal Traps (Baksi, 1995). e) Sanbaoying, Liaoning Province (Shi et al., 2004). f) Jiaolai Basin (Li et al., 2023). g) DSDP Site 402A (Tarduno, 1990; Ramos et al., 2024b). h) DSDP Site 463 (Tarduno, 1990). i) SER-3 core (Fauth et al., 2022). j) ODP Site 317A (Tarduno, 1990). l) ODP Site 1049 (Ogg and Bardot, 2001).

3 Methods

All data used in this study (i.e., the MS dataset and the spliced instructions for cores from Holes 1049A [Core 19X, 1-4], 1049B [Core 11X, 1-2 and Core 12X, 1-5], and 1049C [Core 12X, 1-cc]) were obtained from the Laboratory Information Management System (LIMS) online repository of the International Ocean Discovery Program (available at <https://web.iodp.tamu.edu/LORE/>). Norris et al. (1998) provided a ODP Site 1049 spliced (Fig. 2d) ranging from 131.0 to 151.3 mbsf based on the shipboard low-field magnetic susceptibility (MS) data (measured on whole-round core sections at each 2 cm) from Holes A, B and C (Figs. 2a to 2c, respectively), which was used for our cyclostratigraphic analyses. The ODP Site 1049 spliced comprises the upper section from Hole A (Cores 19X-1 to 19X-4), complemented by Hole B (Core 11X-1). From 138.28 mbsf to 146.03 mbsf, the Spliced Hole is constructed using Cores 12X-1 to 12X-6 from Hole C. Finally, the basal part of the Spliced Hole is composed using Cores 12X-1 to 12X-5 from Hole C (Fig. 2d). We constrained the magnetostratigraphic evaluation of Ogg and Bardot (2001) at the 144.0 – 160.5 mbsf interval from Hole 1049A, which exhibited the higher core recovery and comprises both the top Albian unconformity and the unconformity separating the Aptian-Albian intervals.

3.1 Cyclostratigraphy

Magnetic susceptibility, CaCO_3 and other geochemical records from the Aptian/Albian intervals (Li et al., 2011; Hu et al., 2022) show pronounced short-term cyclicity linked to Milankovitch forcing (Ogg and Bardot, 2001), reflecting alternations between phases of active water bottom circulation and episodes of water-column stagnation (Fig. 3). Cyclostratigraphic analysis of the ODP Site 1049 spliced was performed using version 2.4.1 of the Acycle software (Li et al., 2019). Due to the evenly spaced acquisition of the MS data, interpolation was not required. A locally weighted scatterplot smoothing ('lowess') trend of 3.97 m window (equivalent to ~20% of the entire data length) was removed prior to spectral analysis.

To investigate potential imprints of orbital forcing in the stratigraphy, spectral analyses were conducted using the multitaper method (MTM; Thompson, 1982) with two prolate tapers. Statistical significance was assessed by fitting a first-order autoregressive [AR(1)] red-noise null hypothesis and estimating confidence levels at 95% and 99%, following the robust background estimation procedure of Mann and Lees (1996). The transience and/or persistence of astronomical frequencies throughout the dataset was evaluated using Evolutive Harmonic Analysis (EHA; Meyers et al., 2001). Candidate orbital frequencies were then compared with those predicted by the La2004 astronomical solution for Aptian-Albian times (Laskar et al., 2004). Astronomical calibration was anchored on four tie points:

- **TP.1:** an age of 111.88 Ma for the base of the *H. albiensis* calcareous nannofossil Zone at 142.4 mcd, anchored between the c-markers ν and ξ (112.20 and 111.80 Ma, respectively) (Ramos et al., 2024a);
- **TP.2:** an age of 111.52 Ma for marker *o* at 139.5 mcd;

- **TP.3:** an age of 114.91 Ma for the first occurrence (FO) of the planktonic foraminifera *Paraticinella rohri* at 147.8 mcd; and
- **TP.4:** an age of 115.64 Ma for the FO of *Hedbergella trocoidea* at 153 mcd (Huber et al., 2011; Huber and Leckie, 2011; Ramos et al., 2026) (Fig. 3).




Figure 2: a) Magnetostratigraphy of ODP Site 1049 Hole A, showing planktonic foraminiferal zones (Huber et al., 2011), cores, core photographs, and geomagnetic polarity (Ogg and Bardot, 2001). b) Section of ODP Site 1049 Hole B used to compose the spliced section (Norris et al., 1998). c) Section of ODP Site 1049 Hole C. Planktonic foraminiferal and calcareous nannofossil zones from Huber et al. (2011). d) Spliced section of magnetic susceptibility data from Holes 1049A, 1049B, and 1049C (Norris et al., 1998). The magnetic polarity curve was derived from Hole A.

180

185

190




Figure 3: a) Magnetostratigraphy of ODP Site 1049 (spliced), showing planktonic foraminiferal zones (Huber et al., 2011), cores, core photographs, and geomagnetic polarity (Ogg and Bardot, 2001). Spliced section of magnetic susceptibility data from Holes 1049A, 1049B, and 1049C (Norris et al., 1998). The magnetic polarity curve was derived from Hole A. TP = tie-points. b) Bulk carbonate samples from the upper Aptian–lower Albian interval of ODP Hole 1049C plotted alongside ages of the c-markers (Ramos et al., 2024a) and the CORBs levels. c) Correlated interval of the PLG core with stages, planktonic foraminiferal and calcareous nannofossil zones, bulk carbonate profile, c-markers, and CORBs numbers (Coccioni et al., 2012, 2014; Leandro et al., 2022). The shaded polygon represents the interval absent at ODP Site 1049. The grey square indicates the interval related to OAE 1b (Urbino/Paquier Level; Erbacher et al., 2001). d) Magnetic susceptibility and geochemical data (Li et al., 2011; Hu et al., 2012) from ODP Site 1049C, showing the cyclic variations between CORBs and non-CORBs intervals.

4 Results

An unconformity separating the Aptian and Albian intervals at ODP Site 1049 was previously reported (Huber et al., 2011; Huber and Leckie, 2011; Hu et al., 2012). The identification of orbital imprints in the MS data was carried out independently for each interval. MTM spectral analyses of the MS records (Fig. 4a,d) revealed a broad spectral content, commonly exceeding the 95%–99% confidence thresholds (Fig. 4b,e). At the Aptian section, a sequence of spectral peaks compatible to the orbital imprint was identified with ratios of 256:83:62:23:14 cm = 18.3:5.9:4.4:1.6:1. These values closely align with the expected Milankovitch cyclicity for the Aptian (405:125:95:39:23 = 17.6:5.4:4.1:1.7:1; Waltham, 2015). Similarly, a set of spectral peaks for the Albian interval (357:128:97:34:18:15 cm) corresponds to a ratio of 23.8:8.5:6.5:2.3:1.2:1, which is in strong agreement with the predicted astronomical frequencies (405:125:95:39:23:18 = 22.5:6.9:5.3:2.2:1.3:1; Waltham, 2015).

Such findings indicate that the correspondence between observed and theoretical periodicities for both intervals suggests that sedimentation at ODP Site 1049 during the Aptian–Albian was likely influenced by orbital forcing, supporting earlier interpretations by Ogg and Bardot (2001). Additionally, analysis of both the power spectra and EHA results (Fig. 4c,f) revealed a low wavelength range pattern compatible to the orbital imprint which is quite stable throughout the studied interval. For the Aptian interval, we associated the following spectral bands with eccentricity signals: 405-kyr and 125/95-kyr, corresponding to 0.1–0.5 and 0.7–1.8 cycles/m, respectively. Additionally, we attributed the spectral bands of 2.2–4.6 m and greater than 5.2 cycles/m to the obliquity (~51 to ~29 kyr) and precession (~23 to ~14 kyr) signals, respectively (Fig. 4b,c). In the Albian interval, we related the following spectral bands with eccentricity signals: 405-kyr and 125/95-kyr, corresponding to 0.1–0.6 and 0.8–3.0 cycles/m, respectively. Similarly, the obliquity (~51 to ~29 kyr) and precession (~23 to ~14 kyr) signals were linked to the spectral bands of 3.6–6.4 and greater than 6.8 cycles/m, respectively (Fig. 4e,f).

The interpreted 405-kyr long-eccentricity sinusoidal component was extracted from the MS record of ODP Site 1049 through Gaussian filtering, following the methodology outlined by Li et al. (2019). In our analysis, we correlated the minimum eccentricity phases derived from astronomical models (Laskar et al., 2004) with the lowest MS values in the dataset (Fig. 5a).

After anchoring the stratigraphic positions of the tie points—TP.1 at 111.88 Ma, TP.2 at 111.52 Ma (located in close proximity to the reversal-polarity event “3”), TP.3 at 114.91 Ma, and TP.4 at 115.64 Ma—we estimated sediment accumulation rates (SARs) ranging between 0.4 and 0.8 cm/kyr (Fig. 5b).

Our astronomical tuning provided an age of ~111.45 Ma for the onset of reversal-polarity event “3”, located at a depth of 139.5 mcd and spanning an estimated duration of ~80 kyr. Furthermore, an age of 110.61 Ma was assigned for the base of the M-2r reversed interval (~134 mcd) (Figs. 5c and 5d). We also constrained the age of the black shale interval—interpreted as a time equivalent of the Urbino Level (Coccioni et al., 2014; Ramos et al., 2024a)—to between 111.59 and 111.66 Ma. Additionally, the stratigraphic gap (Huber et al., 2011) represented by the unconformity was estimated to span approximately 2.56 million years.




Figure 4: MS dataset prior and after Lowess detrending from the Albian (a) and Aptian (d) interval ODP Site 1049. Gray line: original MS values. black line: detrended MS data. red line: 405-kyr long eccentricity component. (b, e) Multitaper spectral estimator-based spectral analysis showing the interpreted cycles and the respective bands (colored rectangles). (c, f) evolutionary spectral analysis of series. E: long-eccentricity; e: short-eccentricity; Obliq: obliquity; Prec: precession index.




Figure 5: (a) Spliced MS (dashed grey line) and Lowess detrended MS (black line) and 405-kyr component (red line) of ODP Site 1049, presenting the stages, planktonic foraminifera and calcareous nannofossils zones, and geomagnetic polarity. TP.=tiepoint. b) SAR resulting from the tuning. c) Eccentricity signal from the La2004 astronomical solution (Laskar et al., 2004) Long-eccentricity: red line; short-eccentricity: blue line) and its 405-kyr sinusoidal filtered component (red line). (d) Tuned Spliced MS (black line) and 405-kyr component (red line) of ODP Site 1049 and its associated 405-kyr component (red line). The ages of rapid reversals within the CNPS resulting from the tuning are highlighted in green.

The cyclostratigraphic evaluation of the MS record from ODP Site 1049 supports the conclusion that sedimentation during the analyzed interval was modulated by orbital forcing, with a clear imprint of both long- and short-eccentricity cycles. Based on the age model derived from the tuning process, it was also possible to evaluate the SAR, to date the CORB intervals and to correlate them with those identified in the PLG core (Coccioni et al., 2012; Ramos et al., 2026) (Fig. 6).

In the Aptian, CORB A showed a strong correlation with CORB 1 from the PLG core (Coccioni et al., 2012). Both intervals, with durations exceeding 1 Myr, can be classified as long-term CORBs (Hu et al., 2012). Although there were smaller, darker

245 layers (short-term CORBs; Hu et al., 2012) within CORB A, with durations corresponding to Milankovitch cycles, almost all the Aptian sedimentation at ODP Site 1049 between 115.8 and 114.5 Ma consists of reddened sediments. In addition to the short-term CORBs, which reflect orbital forcing, there were other distinctly reddish intervals with durations of ~350 and ~210 kyr. These intervals have durations that do not correspond to any known orbital cycle (Fig. 6, left).

The basal section of the Albian stage, which was also affected by the unconformity (absence of the *Microhedbergella renilaevius* planktonic foraminifera Zone), is characterized by the absence of CORBs. The CORB 3, associated with the Monte Nerone level, is also absent.

Similarly to the PLG core, but with distinct durations, the ~400-kyr-long interval following the first occurrence (FO) of the *H. albiensis* nannofossil Zone is composed of sediments predominantly green in color. The onset of the black shale, corresponding to the Urbino Level, is marked by a shift from yellow-greenish to black coloration. This event begins at 111.66 Ma and lasts 255 for approximately 70 kyr (Fig. 6). CORB B starts at 111.50 Ma in the ODP 1049 section and consists of at least eight main levels of reddish-brown coloration, with durations consistent with Milankovitch cycles. The high concentration of hematite in this interval (Hu et al., 2006a,b) results in a significantly larger MS amplitude compared to the rest of the studied section. After the termination of CORB B (~110.97 Ma), an interval with alternating yellowish and greenish hues predominates until the onset of CORB C at ~110.84 Ma. The onset of CORB C is gradual, allowing for a potential correlation with CORB 4. However, 260 despite the possible correlation between the onsets of CORB C (ODP Site 1049) and CORB 4 (PLG core), the latter exhibits a shorter duration (Fig. 6).

The eight main CORB levels composing CORB B (Fig. 6) have the following thicknesses and estimated durations (from oldest to youngest): ~55 cm / 125 kyr; 12 cm / 39 kyr; 7 cm / 23 kyr; 34 cm / 95 kyr; 12 cm / 39 kyr; 5 cm / 18 kyr; 6 cm / 22 kyr; and 5 cm / 18 kyr. For CORB C, from oldest to youngest, the values are: ~12 cm / 39 kyr; 5 cm / 18 kyr; 5 cm / 18 kyr; 5 cm 265 / 18 kyr; 5 cm / 18 kyr; 7 cm / 23 kyr; 12 cm / 39 kyr; 7 cm / 23 kyr; 34 cm / 95 kyr; and 12 cm / 39 kyr. It should be noted that these values are approximate, as some red beds appear (and disappear) abruptly, whereas others show more gradational boundaries.

270

275




Figure 6: a) Aptian interval from ODP Site 1049, highlighting the time spans of the CORBs layers. b) Aptian interval from the PLG core. c) Albian interval from ODP Site 1049, illustrating the time spans corresponding to the CORBs layers and the Urbino/Paquier Level (indicated by the grey rectangle). d) Albian interval from the PLG core, depicting the durations associated with the CORBs and the Urbino Level (represented by the black rectangle). Age models and CORBs boundaries follow Ramos et al. (2024a) and Coccioni et al. (2012).

5.1 Circulation regimes and CORB formation

CORBs serve as key archives of mid-Cretaceous climate and circulation dynamics (Premoli Silva et al., 1989; Hu et al., 2005; Li et al., 2011; Giorgioni et al., 2017; Gambacorta et al., 2016). Their formation is linked to episodes of cold, oxygenated deep-water formation rather than elevated productivity, as LIP-derived nutrients would have promoted organic carbon burial incompatible with oxidized conditions (Leckie et al., 2002; Browning and Watkins, 2008; Wang et al., 2009). Enhanced burial of organic carbon during OAEs, combined with LIP-induced perturbations, reduced atmospheric CO₂ and promoted global cooling, increasing the oxidizing capacity of deep waters during the Aptian Cold Snap (Arthur et al., 1988; Weissert, 1989; Hu et al., 2012).

Aptian CORBs are long-lived, regionally correlatable, and reflect sustained thermohaline circulation over ~2 Myr, supported by isotopic evidence for well-oxygenated, cooler, and low-productivity deep waters (Erbacher et al., 2001; Premoli Silva et al., 1989; Leandro et al., 2022). Enhanced vertical mixing driven by current–topography interactions may have further facilitated oxygenation (Ahmerkamp et al., 2017).

In contrast, Albian CORBs are shorter, orbitally paced, and reflect transient ventilation interspersed with stratification and episodic black shale deposition (Paquier/Urbino Level; Erbacher et al., 2001). Color, facies, and geochemical variability among Albian CORBs indicate localized differences in paleogeography and oceanography, with obliquity and short-eccentricity cycles controlling sedimentation (Li et al., 2011; Trabuco Alexandre et al., 2011). Aptian CORBs display more uniformity, consistent with sustained circulation and restricted oceanic connectivity prior to full opening of the Equatorial Atlantic Gateway (Dummann et al., 2023; Ramos et al., 2025).

One of the most frequently mentioned characteristics of CORB deposition is its association with low sedimentation rates in pelagic deep waters of oceanic basins (Wang et al., 2009; Hu et al., 2012). Previous studies have estimated sedimentation rates in the Aptian section of ODP Hole 1049C to range between 0.25 and 0.35 cm/kyr, based on the *Paraticinella eubejaouaensis* (*Paraticinella rohri*) and *Globigerinelloides algerianus* zones (Li et al., 2011). Additional estimates using paleomagnetic data and astronomical tuning have inferred sedimentation rates of 0.4 cm/kyr during the late Aptian and up to 1 cm/kyr in the early Albian (Ogg and Bardot, 2001; Ogg et al., 1999). In contrast, our study provides sedimentation rates ranging from 0.5 to 0.9 cm/kyr for the Aptian section and from 0.6 to 0.8 cm/kyr for the Albian section at ODP Site 1049 (Fig. 5). Notably, these values do not indicate significant variations that would support the hypothesis of extremely slow sediment accumulation as a primary control on CORB formation, as previously proposed (Wang et al., 2009).

5.2 Geological synchrony between the Tethyan and North Atlantic oceans

Correlations between CORBs at ODP Site 1049 and those identified in the PLG core provide insight into the temporal synchrony of these deposits and the paleoceanographic conditions under which they formed. Using first occurrences (FO) of

key planktonic foraminiferal biozones—recognized as more reliable markers for diachronism studies (Berggren and Van Couvering, 1974)—our analysis shows that the *H. trocoidea* zone spans 0.73 Myr in the Tethys (115.64 to 114.91 Ma, PLG core; Ramos et al., 2026), while its duration in the North Atlantic is slightly shorter, at 0.61 Myr (115.75 to 115.14 Ma, ODP Site 1049; this study). The FO of the nannofossil *H. albiensis* marks the base of the NC8B subzone (Bralower et al., 1993) and shows a temporal offset of only 40 kyr between basins (Huber et al., 2011; Ramos et al., 2026).

For the OAE 1b-related black shale in ODP Hole 1049C, Erbacher et al. (2001) estimated a duration of approximately 46 kyr. Our results, however, indicate a slightly longer duration of 70 kyr and a central age of 111.63 Ma. This interval correlates with the Urbino Level (111.74 to 111.65 Ma; Fig. 6). Even with marked differences in oxidation levels observed through the correlation (in geological time) between the Albian interval of ODP Site 1049 and the PLG core—differences dependent on palaeogeographic and depositional settings (Trabucho Alexandre et al., 2011)—the small age and duration offsets indicate that this event can be considered effectively synchronous between the Tethyan and North Atlantic domains. This observation suggests that the widespread anoxia associated with this particular OAE 1b sub-event transcends local physiographic barriers, supporting its interpretation as at least a Central Atlantic–western Tethys phenomenon.

The astrochronological tuning applied in this study yields a maximum error of ~200 kyr (half of the long-eccentricity cycle period of 405 kyr used in the tuning) over a 250 Myr timespan, or roughly ~1.6‰ (Laskar, 2020). Both the ~100 kyr difference in the duration of the *H. trocoidea* zone and the discrepancies in FO and LO ages fall within this error range, preventing a conclusive assessment of dispersal mechanisms (Petrizzo, 2003; Lam et al., 2022). However, the strong synchrony observed for both the Urbino Level and the FO of *H. albiensis* across the two oceanic regions was expected due to the relatively short distance between the North Atlantic and the Tethyan Ocean, which, together with the presence of persistent easterly equatorial winds (Hay, 2008), favored rapid dispersal.

5.3 Milankovitch-band orbital forcing of CORB deposition

High-frequency Milankovitch-scale cyclicity is widely recorded in Lower Cretaceous pelagic successions and is expressed in variations of CaCO_3 and SiO_2 contents (Herbert et al., 1986), carbon isotopes, and magnetic susceptibility (Leandro et al., 2022). These cycles are superimposed on a long-term perturbation of the global carbon cycle (Weissert, 1989), characterized by reduced carbon turnover and multi-million-year cold interludes (Leandro et al., 2022), potentially associated with polar ice-sheet growth (Trabucho Alexandre et al., 2011) and eustatic sea-level fall (Weissert and Lini, 1991).

Orbital-scale variability in ocean circulation reflects the interplay between seasonal contrast, hydrological balance, and water-column stratification. Humid climates with strong seasonality favor stratification and are typically associated with non-CORB intervals, whereas reduced seasonality under more arid conditions promotes weaker stratification and CORB deposition (Giorgioni et al., 2017). Variations in runoff, evaporation, saline exchange between basins, and temperature-dependent oxygen solubility further modulate surface-water density gradients and deep-water formation (Ryan and Cita, 1977; Weissert et al., 1985; MacLeod et al., 2001; Steinig et al., 2024). Episodic saline incursions and astronomically paced bottom currents may

have intermittently enhanced deep-water renewal and seafloor reoxygenation without requiring major reorganization of global circulation (Friedrich et al., 2008; Gambacorta et al., 2016).

345 At ODP Site 1049, the Albian interval displays pronounced orbital-scale cyclicity in CaCO_3 , Ba/Al, SiO_2 , Al_2O_3 , and magnetic susceptibility (Li et al., 2021; Hu et al., 2022), consistent with repeated alternations between enhanced ventilation and stratified conditions (Figs. 3 and 4). The Albian cyclic patterns are consistent with enhanced variability in bottom-water ventilation and primary productivity, in agreement with previous observations of orbitally paced changes in Early Cretaceous deep-ocean oxygenation (Herbet et al., 1986; Gambacorta et al., 2016; Leandro et al., 2022). In contrast, Aptian proxies display more
350 uniform behavior, with low-amplitude CaCO_3 variations and the absence of clear productivity cycles (Fig. 3). This suggests a more stable paleoceanographic regime during the Aptian, likely maintained by restricted oceanic connectivity prior to the full opening of the Equatorial Atlantic Gateway (EAG; Duarte et al., 2025; Ramos et al., 2025) and by the persistence of the Cold Snap climatic state. Orbital pacing, however, is still recorded in Aptian MS fluctuations, indicating sensitivity to insolation forcing even under comparatively steady circulation.

355 Albian CORBs are best explained by the combined effects of evaporation-driven deep-water formation—through episodic saline flooding—and orbitally regulated bottom-current activity that transiently reoxygenated the seafloor, enabling cyclic early diagenesis. In contrast, the Aptian interval shows more uniform proxy behavior, limited CaCO_3 variability, and weaker productivity signals, pointing to a comparatively stable circulation regime. Although orbital forcing is still expressed in magnetic susceptibility, its impact was muted, consistent with sustained deep-water ventilation during the Aptian cold snap
360 and restricted oceanic connectivity prior to the full opening of the Equatorial Atlantic Gateway.

5.4 Long-cycle-related unconformity

The unconformity observed at ODP Site 1049 is consistent with the local geological setting of the Blake Escarpment, a continental-slope environment characterized by persistent bottom-current activity and high erosional potential (Benson et al., 1978; Li et al., 2011). Along continental margins in both the Tethyan Ocean and the North Atlantic, buried contourite drifts
365 and stratigraphic gaps are common features and are widely attributed to variations in bottom-current strength and reorganizations of deep-water circulation, frequently linked to OAEs and major paleoceanographic transitions (Gambacorta et al., 2016; Liu et al., 2023).

However, hiatuses of comparable duration have been documented at multiple sites, including DSDP Sites 511, 545, 763B, and 392A (Huber and Leckie, 2011), as well as at DSDP Site 545 off Morocco and ODP Site 1276 in the Newfoundland Basin
370 (Trabucho Alexandre et al., 2011). The recurrence of similarly timed unconformities across different ocean basins indicates that the discontinuity at Site 1049 reflects a regional paleoceanographic signal rather than a purely local phenomenon. Incomplete stratigraphic records near the Aptian–Albian boundary—commonly associated with OAE 1b (Ramos et al., 2024b)—are therefore not unusual; several drill sites, including DSDP Sites 390, 392A, and 511, show partial or complete removal of upper Aptian and lower Albian successions (Huber et al., 2011).

5.5 Short geomagnetic reversed-polarity subchrons

As discussed above, a preliminary magnetostratigraphic framework for ODP Site 1049 was first proposed by Norris et al. (1998) and later re-evaluated through detailed paleomagnetic analyses of minicore suites using stepwise thermal demagnetization (Ogg and Bardot, 2001). These authors interpreted the upper Aptian reversed-polarity interval (~155 mbsf; 171B-1049A-20X-1) as a diagenetic artifact related to post-Albian iron mobilization driven by redox contrasts near an organic-rich horizon. However, a normally magnetized sample (20X-2, 30–32 cm) occurs stratigraphically between this reversed interval and the underlying black shale and exhibits pronounced yellowish to greenish staining, suggesting preservation of primary magnetization.

Independent mineralogical and biostratigraphic evidence further argues against significant late diagenetic overprinting at ODP Hole 1049C. Clay mineral assemblages show no systematic burial-related trends, with no progressive replacement of smectite by illite or chlorite, indicating the absence of burial diagenesis (Li et al., 2011). In addition, planktonic and benthic foraminifera display glassy tests with well-preserved ornamentation and no calcite infilling, supporting minimal diagenetic alteration (Erbacher et al., 2001; Li et al., 2011). The occurrence of goethite precisely at the level recording the older reversed geomagnetic field further supports the preservation of the primary magnetic mineralogy (Li et al., 2011). Accordingly, this reversal is retained as primary in our interpretation.

Within a broader context, Ryan et al. (1978) documented seven short-duration geomagnetic reversals during the Aptian–Albian interval, excluding M0r, and introduced a partial nomenclature from M-1r to M-3r. Several of these reversals were left unnamed and have often been treated as isolated events (e.g., Gale et al., 2020), despite evidence that short reversals within the CNPS commonly occur as closely spaced sets (Ryan et al., 1978; Tarduno et al., 1992; Zhang et al., 2021; Ramos et al., 2025).

Using biostratigraphic constraints and an estimated inter-event spacing of ~860 kyr, Ramos et al. (2024b) subdivided the reversal traditionally referred to as M-1r into two distinct Lower Aptian events, termed M-1r and reversal “2”. This reversal pair has been identified both at DSDP Site 402A and along the Brazilian Equatorial Margin, with comparable durations and spacing (Ramos et al., 2024b, 2025). Because reversal “2” is constrained to the Aptian and stratigraphically underlies the reversed interval documented here, we designate this newly identified short reversal as “3” (Fig. 2).

A second reversed-polarity interval at ~146.5 mbsf (171B-1049A-19X-2, 103–105 cm) is interpreted as the Albian subchron M-2r and is consistently recorded in two holes at ODP Site 1049 (Fig. 2). For both reversal “3” and M-2r, boundaries were defined at the intersections between the inclination–depth interpolation and the 0° inclination axis. On this basis, the reversal previously assigned to M-2r (*sensu* Gale et al., 2020) is subdivided into two discrete events, retaining the name M-2r for the younger reversal and assigning the older event to reversal “3”.

The M0r Chron (120.32–119.89 Ma; Ramos et al., 2026) marks the base of both the Aptian and the CNPS (Savani et al., 2016) and is widely regarded as a marker for the Barremian–Aptian boundary (Helsley and Steiner, 1968; Erba et al. 1996; Gale et al., 2020; Zhang et al., 2021). The first Aptian short reversed-polarity subchron following M0r, M-1r (or ISEA; VandenBerg

et al., 1978), occurs within the *Globigerinelloides algerianus* planktonic foraminiferal Zone and has an estimated age of 117.03 ± 0.14 Ma, with a duration of ~20 kyr (Tarduno, 1990; Ramos et al., 2024a). The subsequent reversal “2” is dated at 116.17 ± 0.14 Ma and lasted approximately 10 kyr (Ramos et al., 2024a).

A second pair of short reversals occurs ~5 Myr later within the CNPS (Ryan et al., 1978). Although ⁴⁰Ar/³⁹Ar ages of basalt flows from the Tuoyun section (113.3 ± 1.6 Ma; Gilder et al., 2003) have been interpreted as indicating an additional upper Aptian reversal, our cyclostratigraphic results place this event in the lower Albian, consistent with Ryan et al. (1978). The stratigraphically youngest reversal of this pair, reversal “3”, spans approximately 111.45–111.53 Ma. Given the low temporal resolution of earlier paleomagnetic data (Ogg and Bardot, 2001), its estimated duration of ~80 kyr may represent an upper bound.

Reversal “3” corresponds to reversal number 6 of Zhang et al. (2021) and occurs within the *Microhedbergella rischi* planktonic foraminiferal Zone, near the first occurrence of *Hayesites albiensis* (Fig. 5). The excellent preservation of planktonic and benthic foraminifera, together with the absence of stable isotope homogenization, argues against a diagenetic origin (Erbacher et al., 2001). Although minor recording artifacts cannot be entirely excluded, these biostratigraphic and sedimentological constraints strongly support the authenticity of this reversal. Future high-resolution paleomagnetic and cyclostratigraphic studies should further refine its duration.

The M-2r reversal was originally recognized as a composite interval of two reversed events within the lower Albian, associated with the *Biticinella breggiensis* planktonic foraminiferal Zone and the *Prediscosphaera cretacea* nannofossil Zone (Ryan et al., 1978). Subsequent work in the Contessa Valley section described this interval as a complex reversal set comprising multiple reversed events (Tarduno et al., 1992). Despite suggestions that seafloor oxidation may have influenced parts of the signal, M-2r has been consistently recorded in both marine sections and sedimentary basins, including the Vientiane Basin (Zhang et al., 2021).

Our data indicate that M-2r began at ~110.76 Ma and lasted approximately 150 kyr. Similar to the spacing between M-1r and reversal “2”, reversals “3” and M-2r are separated by ~800 kyr, supporting their interpretation as distinct subchrons (Fig. 7). The cyclostratigraphic framework developed here allows correlation of reversal “3” with the Tuoyun basalt age (Gilder et al., 2003), providing a robust tie point for future Aptian–Albian correlations.

5.6 Aptian–Albian paleoclimatic reorganization of ocean circulation

A major paleoclimatic transition across the Aptian–Albian boundary is recorded by long-term fluctuations in the carbon isotope record, reflecting changes in the global carbon cycle linked to ocean circulation and productivity (Weissert, 1989). At ODP Site 1049, δ¹³C values shift abruptly from ~4‰ to ~2‰ VPDB, providing clear evidence for a fundamental climatic reorganization (Huber et al., 2011). Independent proxies, including floral and faunal turnover, plant fossils indicating cooler conditions, glendonites, ice-rafted debris, and intensified hydrological cycling associated with the late Aptian Equatorial Humid Belt, corroborate this transition (Hochuli, 1981; Krassilov, 1973; Leckie, 1989; Kemper, 1987; Herrle et al., 2015;

Santos et al., 2022; Ramos et al., 2025). Tethyan records further indicate reduced siliciclastic input, turnover within the *G. algerianus* Zone, and a pronounced global sea-level fall (Weissert and Lini, 1991).


This cooling, often termed the Aptian “Cold Snap” (McAnena et al., 2013), is commonly attributed to declining atmospheric CO₂ levels, likely driven by enhanced burial of organic carbon and pyrite, which slowed carbon cycling and promoted polar ice growth (Weissert and Lini, 1991; Leandro et al., 2022). Large Igneous Province (LIP) emplacement during the Early Cretaceous, particularly the development of the Kerguelen Plateau, is widely regarded as a major driver of carbon-cycle perturbations and climatic variability (Coffin and Eldholm, 1994; Percival et al., 2024). Extensive volcanic degassing initially promoted greenhouse conditions and widespread oceanic anoxia during the early Aptian (Weissert and Lini, 1991; Jenkyns, 2003). In contrast, enhanced silicate weathering of newly emplaced basalts and increased organic carbon burial likely led to a drawdown of atmospheric CO₂, facilitating episodic to transient cooling during the late Aptian (Weissert et al., 1998; Jenkyns et al., 2012).

Geochemical proxies, including benthic and planktonic $\delta^{18}\text{O}$, indicate that Aptian bottom waters were up to $\sim 10^\circ\text{C}$ cooler than those of the Albian, highlighting a rapid warming of both surface and deep waters across the boundary (Huber et al., 2011; Kochhann et al., 2023). Under typical Cretaceous greenhouse conditions, high-latitude regions were unlikely to generate dense, oxygenated deep waters, leading to sluggish deep-ocean ventilation (Schlanger and Jenkyns, 1976; Hay, 2008, 2009). The Aptian Cold Snap temporarily disrupted this mode, enhancing thermohaline circulation through increased oxygen solubility in colder waters and possibly the presence of polar ice. This circulation regime promoted deep-water renewal and organic matter oxidation, processes that facilitated CORB formation, as evidenced by widespread hardgrounds, intensified bottom currents, and a basin-wide shift from grey-green to red-brown sediments (Premoli Silva et al., 1989; Weissert and Lini, 1991).

During the Albian, renewed greenhouse conditions suppressed deep-water formation by reducing latitudinal thermal gradients, with ocean heat transport dominated by mesoscale eddies and low-latitude sinking of warm, saline waters (Hay, 2008, 2009; Gambacorta et al., 2016). This shift is reflected in clay-mineral assemblages and geochemical proxies: Aptian Ca-rich CORBs formed under drier conditions with limited terrigenous input, whereas Albian Al-rich CORBs indicate more humid climates, greater paleoceanographic variability, and orbitally paced alternations in ventilation (Li et al., 2011; Hu et al., 2012).

The slightly higher illite concentrations in the Aptian CORBs—reflecting moderate to weak weathering under relatively dry conditions—suggest that the atypical conditions of the Aptian Cold Snap interlude were characterized by lower terrigenous input than that experienced during the Albian (Fig. 3). Although the CaCO₃ content of the white layers interbedded with the red beds is similar in the Aptian and Albian intervals, the markedly low CaCO₃ values in the Albian CORBs are noteworthy (Li et al., 2011). This observation indicates substantial paleoceanographic variability (instability) during the Albian compared to the Aptian stage. Low CaCO₃ content may be related to: (a) reduced primary productivity, (b) higher dissolved CO₂ concentrations, (c) increased terrigenous input, and (d) water-column stagnation with the absence of younger waters. Because terrigenous input at ODP Site 1049 remained stable throughout the Albian (Cheng, 2008), we propose that alternation between intervals of active water bottom oxygenation and intervals of ocean stagnation (white and green beds) provides a more plausible explanation. Variations in the Ba/Al ratio and in SiO₂ and Al₂O₃ concentrations between the Albian red and white beds (Fig.

3) further support the presence of cyclical changes in productivity (Hu et al., 2012). In contrast, the absence of such abrupt CaCO₃ variability during the Aptian suggests more stable paleoceanographic conditions, with limited fluctuations in paleoproductivity and ocean circulation.



525 **6. Conclusions**

Contrary to earlier hypotheses that associated CORBs with very low sedimentation rates, this study reveals that accumulation rates at ODP Site 1049 possibly ranged from 0.5 to 0.9 cm/kyr during Aptian-Albian times. These values demonstrate that CORBs can form under moderately low but not necessarily exceptionally slow sedimentation conditions, suggesting that other environmental or climatic factors also played a significant role in their deposition. Temporal synchrony exists between Aptian
530 CORB-related events in the Tethys and North Atlantic, reflecting their response to global cooling and sustained thermohaline circulation. The onset of long-term Aptian CORBs, such as CORB 1 in the Tethys and CORB A in the North Atlantic, is best explained by global climatic cooling that promoted thermohaline circulation during the Aptian “Cold Snap.” This cooling episode, likely initiated by the emplacement of LIPs and subsequent organic carbon burial during OAEs, promoted oxygenation of deep waters. These sustained oxic conditions, rather than high productivity, facilitated the widespread formation of red beds.
535 Unlike the longer-lasting Aptian CORBs, the Albian red beds are shorter in duration and exhibit clear orbital pacing. This suggests that Albian CORB deposition was more sensitive to high-frequency climatic oscillations and regional variability, reflecting a shift in the dominant environmental controls on red bed formation during this interval. Near the Aptian–Albian boundary, incomplete stratigraphic records linked to the OAE 1b event are common, complicating global correlations of this interval. Our astrochronological findings support recent evidence that ~2.4 Myr grand cycles also produced global hiatuses at
540 the Aptian–Albian boundary, potentially biasing correlations of deep-water sections spanning ~114.5 to 111.9 Ma.

Data availability

All data used here is public and can be accessed on the website <http://deepseadrilling.org> and https://mlp.ldeo.columbia.edu/logdb/scientific_ocean_drilling. DSDP and ODP data can be accessed at
545 <https://web.iodp.tamu.edu/OVERVIEW/>.

Author contribution

J.M.F.R.: conceptualization, methodology, validation, cyclostratigraphic analyses, astronomical tuning, and writing. J.F.S.: conceptualization, methodology, validation, and investigation, writing, project administration, and funding acquisition. D.R.F.:
550 methodology, validation, cyclostratigraphic and astronomical tuning analyses, writing. M.F.: writing. F.F.: conceptualization, and writing. R.C.: conceptualization and writing.

Competing interests

The authors declare that they have no conflict of interest.
555

Acknowledgements

560 This study was financially supported by the Specialization and Postgraduate Program of Petrobras. J.M.F.RAMOS thanks Petrobras for Ph.D. process. The authors gratefully acknowledge Editor Shiling Yang and thank Mark Leckie, Helmut Weissert, and the anonymous reviewer for their thorough and constructive reviews.

Financial support

565 The paper is a part of the projects Processamento e interpretação de dados magnetoestratigráficos do Cretáceo das Bacias Brasileiras and Magneto-cicloestratigrafia do Cretáceo na Margem Equatorial Brasileira (MEQ), both financed by Petróleo Brasileiro S.A.—Petrobras (FAURGS 8368 and 8892). J.F.S. also thanks CNPq for grants #304022/2018–7 and #311231/2021–7. D.R.F. thanks the Foundation Carlos Chagas Filho Research Support of the State of Rio de Janeiro (FAPERJ–grant #E-26/200.931/2022) and CNPq (grant #314462/2020-1).

570 References

- Ahmerkamp, S., Winter, C., Krämer, K., de Beer, D., Janssen, F., Friedrich, J., Kuypers, M. M. M., & Holtappels, M.: Regulation of benthic oxygen fluxes in permeable sediments of the coastal ocean, *Limnol. Oceanogr.*, 62, 1–14, <https://doi.org/10.1002/lno.10379>, 2017.
- 575 Arthur, M., Dean, W. & Pratt, L.: Geochemical and climatic effects of increased marine organic carbon burial at the Cenomanian/Turonian boundary, *Nature*, 335, 714–717, <https://doi.org/10.1038/335714a0>, 1988.
- Baksi, A. K.: Petrogenesis and timing of volcanism in the Rajmahal flood basalt province, northeastern India, *Chem. Geol.*, 121, 73–90, [https://doi.org/10.1016/0009-541\(94\)00124-Q](https://doi.org/10.1016/0009-541(94)00124-Q), 1995.
- 580 Benson, W. E., Sheridan, R. E., Pastouret, L., Enos, P., Freeman, T., Murdrnaa, I.O., Worstell, P., Gradstein, F., Schmidt, R.R., Weaver, F.M. et Stuermer, D.H.: Init. Rep. Deep-Sea Drill.Proj., 44:1005 pp, 1978.
- Berggren, W. A. and Van Couvering, J. A.: The Cenozoic and Mesozoic stages and their boundaries, *Bull. Geol. Soc. Am.*, 585 85, 713–725, 1974.
- Bottini, C., and Erba, E.: Mid-Cretaceous paleoenvironmental changes in the western Tethys, *Clim. Past*, 14, 1147–1163, 2018.
- Bralower, T. J., Sliter, W. V., Arthur, M., R. Leckie, R. M., Allard, D., Schlanger, S. O.: Dysoxic/Anoxic Episodes in the Aptian-Albian (Early Cretaceous), *The Mesozoic Pacific: Geology, Tectonics, and Volcanism*, Volume 77, 590 <https://doi.org/10.1029/GM077p0005>, 1993.
- Breheret, J. G.: The mid-cretaceous organic-rich sediments from the Vocontian zone of the French southeast basin. In A. Mascle (Ed.), *Hydrocarbon and petroleum geology of France*. Special publication of the European association of petroleum geoscientists, No. 4, 295–320. Springer, 1994.
- 595 Browning, E. L. and Watkins, D. K.: Elevated Primary Productivity of Calcareous Nannoplankton Associated with Ocean Anoxic Event 1b during the Aptian/Albian Transition (Early Cretaceous), *Papers in the Earth and Atmospheric Sciences*, 237, <https://digitalcommons.unl.edu/geosciencefacpub/237>, 2008.

- 600 Channell, J. E. T., Freeman, R., Heller, F., & Lowrie, W.: Timing of diagenetic hematite growth in red pelagic limestones from Gubbio (Italy), *Earth Planet. Sci. Lett.*, 58, 189–201, 1982.
- Clarke, L. J. and Jenkyns, H. C.: New oxygen isotope evidence for long-term Cretaceous climatic change in the Southern Hemisphere. *Geology*, 27: 699–702, 1999.
- 605 Coccioni, R., Jovane, L., Bancalà, G., Bucci, C., Fauth, G., Frontalini, F., Janikian, L., Savian, J., Paes de Almeida, R., Mathias, G. L., & Trindade, R. I. F.: Umbria-Marche Basin, Central Italy: A reference section for the Aptian–Albian interval at low latitudes, *Sci. Dril.*, 13, 42–46, <https://doi.org/10.2204/iodp.sd.13.07.2011>, 2012.
- 610 Coccioni, R., Sabatino, N., Frontalini, F., Gardin, S., Sideri, M., & Sprovieri, M.: The neglected history of Oceanic Anoxic Event 1b: insights and new data from the Poggio le Guaine section (Umbria–Marche Basin), *Stratigraphy*, 11, 245–282, 2014.
- Coffin and , M. F. and Eldholm, O.: Large igneous provinces: Crustal structure, dimensions, and external consequences, *Reviews of Geophysics*, Volume 32, Issue 1, 1-36, <https://doi.org/10.1029/93RG02508>, 1994.
- 615 Duarte, D., Erba, E., Bottini, C., Wagner, T., Aduomahor, B., Jones, T. D., Nicholson, U.: Early Cretaceous deep-water bedforms west of the Guinea Plateau revise the opening history of the Equatorial Atlantic Gateway, *Global and Planetary Change*, 249, 104777, ISSN 0921-8181, <https://doi.org/10.1016/j.gloplacha.2025.104777>, 2025.
- Dummann, W., Hofmann, P., Herrle, J. O., Frank, M., Wagner, T.: The early opening of the Equatorial Atlantic gateway and the evolution of Cretaceous peak warming. *Geology* 51, 476–480, 2023.
- 620 Dutkiewicz, A., Boulila, S., & Müller, R. D.: Deep-sea hiatus record reveals orbital pacing by 2.4 Myr eccentricity grand cycles, *Nat. Commun.*, 15, 1998, 2024.
- 625 Erba, E., Aguado, R., Avram, E., Baraboschkin, E. J., Bergen, J. A., Bralower, T. J., ... and Weissert, H.: The Aptian stage. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre*, 66, 31-43, 1996.
- Erbacher, J., Huber, B. T., Norris, R. D., and Markey, M.: Increased thermohaline stratification as a possible cause for an ocean anoxic event in the Cretaceous period, *Nature*, 409, 18–21, 2001.
- 630 Eren, M., & Kadir, S.: Colour genesis of Upper Cretaceous pelagic red sediments within the Eastern Pontides, NE Turkey, *Yerbilimleri*, 24, 71–79, 2001.
- 635 Fauth, G., Krah, G., Kochhann, K. G. D., Hünig Bom, M. H., Baecker-Fauth, S., Bruno, M. D. R., da Silva dos Santos, A., Ceolin, D., Villegas-Martín, J., Strohschoen Jr., O., Savian, J. F., Leandro, C. G., de Mello, R. G., & de Oliveira Lima, F. H. O.: Astronomical calibration of the latest Aptian to middle Albian in the South Atlantic Ocean, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 602, 111175, <https://doi.org/10.1016/j.palaeo.2022.111175>, 2022.
- Fischer, A. G., and Arthur, M. A.: Secular variations in the pelagic realm, *SEPM Spec. Publ.*, 25, 19–50, 1977.
- 640 Friedrich, O., Norris, R. D., & Erbacher, J.: Evolution of middle to Late Cretaceous oceans — a 55 My record of Earth's temperature and carbon cycle, *Geology*, 40, 107–110, 2012.
- Friedrich, O., J. Erbacher, K. Moriya, P. A. Wilson, and H. Kuhnert.: Warm saline intermediate waters in the Cretaceous tropical Atlantic Ocean, *Nat. Geosci.*, 1, 453–457, 2008.
- 645 Gale, A. S., Mutterlose, J., & Batenburg, S.: The Cretaceous period, in Gradstein, F. M., Ogg, J. G., Schmitz, M. D., & Ogg, G. M. (Eds.), *The Geologic Time Scale 2020*, Elsevier, 1023–1068, 2020.

- 650 Gambacorta, G., Bersezio, R., Weissert, H., and E. Erba, E.: Onset and demise of Cretaceous oceanic anoxic events: The coupling of surface and bottom oceanic processes in two pelagic basins of the western Tethys, *Paleoceanography*, 31, 732–757, doi:10.1002/2015PA002922, 2016.
- 655 Gilder, S., Chen, Y., Cogne, J. P., Tan, X. D., Courtillot, V., Sun, D. J., & Li, Y. A.: Paleomagnetism of Upper Jurassic to Lower Cretaceous volcanic and sedimentary rocks from the western Tarim Basin ..., *Earth Planet. Sci. Lett.*, 206, 587–600, [https://doi.org/10.1016/S0012-821X\(02\)01074-9](https://doi.org/10.1016/S0012-821X(02)01074-9), 2003.
- Giorgioni, M., et al.: Paleoceanographic changes during the Albian–Cenomanian in the Tethys and North Atlantic and the onset of the Cretaceous chalk, *Glob. Planet. Change*, 126, 46–61, 2015.
- 660 Giorgioni, M., Tiraboschi, D., Erba, E., Hamann, Y., Weissert, H.: Sedimentary patterns and palaeoceanography of the Albian Marne a Fucoidi Formation (Central Italy) revealed by high-resolution geochemical and nannofossil data, *Sedimentology*, 64, 111–126, 2017.
- 665 Haq, B. U., Hardenbol, J. and Vail, P. R.: Chronology of Fluctuating Sea Levels since the Triassic. *Science*, 235, 1156–1167, <http://dx.doi.org/10.1126/science.235.4793.1156>, 1987.
- Hay, W. W.: Evolving ideas about the Cretaceous climate and ocean. *Cretaceous Research*, v. 29, p. 725–753, 2008.
- Hay, W. W.: Cretaceous oceans and ocean modeling. From: SEPM Society for Sedimentary Geology 91, *Cretaceous Oceanic Red Beds: Stratigraphy, Composition, Origins, and Paleoceanographic and Paleoclimatic Significance*: ISBN 978-1-56576-135-3, 243–271, 2009.
- 670 Herbert, T. D., Stallard, R. E. and Fischer, A. G.: Anoxic Events, Productivity and Rhythms, and the Orbital Signature in a mid-Cretaceous deep-sea sequence from Central Italy, *Paleoceanography*, 1, 495–506, 1986.
- 675 Herrle, J. O., Kossler, P., Friedrich, O., Erlenkeuser, H., & Hemleben, C.: High-resolution carbon isotope records of the Aptian to lower Albian from SE France and the Mazagan Plateau (DSDP site 545): A stratigraphic tool for paleoceanographic and paleobiologic reconstruction. *Earth and Planetary Science Letters*, 218(1–2), 149–161. [https://doi.org/10.1016/S0012-821X\(03\)00646-0](https://doi.org/10.1016/S0012-821X(03)00646-0), 2004.
- 680 Herrle, J. O., Schroder-Adams, C. J., Davis, W., Pugh, A. T., Galloway, J. M., & Fath, J.: Mid-cretaceous High Arctic stratigraphy, climate, and oceanic anoxic events. *Geology*, 43(5), 403–406. <https://doi.org/10.1130/G36439.1>, 2015.
- Hochuli, E.: North Gondwanan Floral Elements in Lower to Middle Cretaceous Sediments of the Southern Alps (Southern Switzerland, Northern Italy)', *Review of Paleobotany and Palynology* 35, 337–358, 1981.
- 685 Hu, X., Jansa, L., and Sarti, M.: Mid-Cretaceous oceanic red beds in the Umbria-Marche Basin, central Italy: constraints on paleoceanography and paleoclimate, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 233, 163–186, 2005a.
- Hu, X. M., Wang, C. S., Li, X. H., & Jansa, L.: Upper Cretaceous oceanic red beds in southern Tibet: lithofacies, environments and colour origin, *Sci. China Ser. D – Earth Sci.*, 49, 785–795, 2006b.
- 690 Hu, X., Scott, R. W., Cai, Y., Wang, C., Melinte-Dobrinescu, M. C.: Cretaceous oceanic red beds (CORBs): different time scales and models of origin, *Earth Sci. Rev.*, 115, 217–248, 2012.
- 695 Huber, B. T., MacLeod, K. G., Gröcke, D., and Kucera, M.: Paleotemperature and paleosalinity inferences and chemostratigraphy across the Aptian/Albian boundary in the subtropical North Atlantic, *Paleoceanography*, v. 26, p. PA4221, 2011.

- Huber, B. T., and Leckie, M. R.: Planktic foraminiferal species turnover across deep-sea Aptian/Albian Boundary sections, *Journal of Foraminiferal Research*, 41, 53–95, 2011.
- 700 Huber, B. T., MacLeod, K. G., Watkins, D. K., and Coffin, M. F.: The rise and fall of the Cretaceous Hot Greenhouse climate, *Glob. Planet. Change*, 167, 1–23, 2018.
- Jansa, L. and Hu, X.: Cretaceous Pelagic Black Shales and Red Beds in Western Tethys: Origins, Paleoclimate, and
 705 Paleooceanographic Implications. 10.2110/sepmsp.091.059, 2009.
- Jenkyns, H. C., Schouten-Huibers, L., Schouten, S. and Sinninghe Damsté, J. S. Warm Middle Jurassic–Early Cretaceous high-latitude sea-surface temperatures from the Southern Ocean. *Clim. Past* 8, 215–226 (2012).
- 710 Kemper, E.: Das Klima der Kreidezeit, *Geol. Jb.*, A96, 5-185, 1987.
- Kochhann, K. G. D., Huber, B. T., Holbourn, A. E., and Kuhnt, W.: Benthic foraminiferal response to the Aptian–Albian carbon cycle perturbation in the Atlantic Ocean, *J. Foraminiferal Res.*, 53, 214–225, 2023.
- 715 Krassilov, V. A.: Climatic changes in Eastern Asia as indicated by fossil floras. I. Early cretaceous, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 13, Issue 4, 261-273, ISSN 0031-0182, [https://doi.org/10.1016/0031-0182\(73\)90028-X](https://doi.org/10.1016/0031-0182(73)90028-X), 1973.
- Lam, A. R., Suganuma, Y., Kuroda, J., Hasegawa, T. and O'Regan, M.: Diachroneity rules the mid-latitudes: a test case using
 720 late Neogene planktic foraminifera across the Western Pacific, *Geosciences*, 12, 1–43, 2022.
- Larson, R. L.: Geological consequences of superplumes, *Geology*, 19, 963–966, 1991.
- Larson, R. L., & Erba, E.: Onset of the mid-Cretaceous greenhouse in the Barremian-Aptian: Igneous events and the biological,
 725 sedimentary, and geochemical responses. *Paleoceanography*, 14(6), 663–678. <https://doi.org/10.1029/1999PA900040.1999>.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astron. Astrophys.*, 428, 261–285, <https://doi.org/10.1051/0004-6361:20041335>, 2004.
- 730 Laskar, J.: Astrochronology. In *Geologic time scale 2020*, 139–158, 2020.
- Leandro, C. G., Savian, J. F., Kochhann, M. V. L., Franco, D. R., Coccioni, R., Frontalini, F., Gardin, S., Jovane, F., Figueiredo, M., Tedeschi, L. R., Janikian, L., Almeida, R. P., & Trindade, R. I. F.: Astronomical tuning of the Aptian stage and its
 735 implications for age recalibrations and paleoclimatic events, *Nat. Commun.*, 13, 2941, <https://doi.org/10.1038/s41467-022-30075-3>, 2022.
- Leckie, R. M.: A paleoceanographic model for the early evolutionary history of planktonic foraminifera, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 73, Issues 1–2, 107-138, ISSN 0031-0182, [https://doi.org/10.1016/0031-0182\(89\)90048-5](https://doi.org/10.1016/0031-0182(89)90048-5), 1989.
- 740 Leckie, R. M., Bralower, T. J. and Cashman, R.: Oceanic anoxic events and plankton evolution: biotic response to tectonic forcing during the mid-Cretaceous. *Paleoceanography*, 17, 13–1 to 13–29. doi: 10.1029/2001PA000623, 2012.
- Li, M., Hinnov, L., and Kump, L.: Acycle: time-series analysis software for paleoclimate research and education, *Comput. Geosci.*, 127, 12–22, <https://doi.org/10.1016/j.cageo.2019.02.011>, 2019.
- 745

- Li, X., Hu, X., Cai, Y., and Han, Z.: Quantitative analysis of iron oxide concentrations within Aptian–Albian cyclic oceanic red beds in ODP Hole 1049C, North Atlantic, *Sediment. Geol.*, 235, 91–99, 2011.
- 750 Li, Y., Singer, B. S., Takashima, R., Schmitz, M. D., Podrecca, L. G., Sageman, B. B., Selby, D., Yamanaka, T., Mohr, M. T., Hayashi, K., Tomaru, T., Savatic, K.: Radioisotopic chronology of Ocean Anoxic Event 1a: Framework for analysis of driving mechanisms. *Sci. Adv.* 10, eadn8365, 2024.
- 755 Liu, S., Hernández-Molin, F. J., Rodrigues, S. and Rooij, D. V.: Deep-water circulation in the northeast Atlantic during the mid- and Late Cretaceous, *Geology*, 51 (6): 515–520, <https://doi.org/10.1130/G50886.1>, 2023.
- Li, Y., Qin, H., Jicha, B. R., Huyskens, M. H. et al.: Revised onset age of magnetochron M0r: Chronostratigraphic and geologic implications: *Geology*, v. 51, p. 565, 2023.
- 760 Matsumoto, H., Minoshima, K., Yamaguchi, K., & Ishikawa, T.: Mid-Cretaceous marine Os isotope evidence for heterogeneous causes of oceanic anoxic events, *Nat. Commun.*, 13, 239, 2022.
- Mann, M. E., and Lees, J. M.: Robust estimation of background noise and signal detection in climatic time series, *Clim. Change*, 33, 409–445, 1996.
- 765 McAnena, A., Flögel, S., Hofmann, P., Herrle, J.O., Griesand, A., Pross, J., Talbot, H.M., Rethemeyer, J. and Wagner, T.: Atlantic cooling associated with a marine biotic crisis during the mid-Cretaceous period, *Nat. Geosci.*, 6, 558–561, 2013.
- MacLeod, K. G., Huber, B. T., Pletsch, T., T, Röhl, U. and Kucera, M.: Maastrichtian foraminiferal and paleoceanographic changes on Milankovitch timescales, *Paleoceanography*, 16, 2, 133–154, 2001.
- 770 Meyers, S. R., Sageman, B., and Hinnov, L.: Integrated quantitative stratigraphy of the Cenomanian–Turonian Bridge Creek Limestone Member using evolutive harmonic analysis and stratigraphic modeling, *J. Sediment. Res.*, 71, 627–643, 2001.
- 775 Norris, R. D., Kroon, D., Klaus, A., et al.: Proceedings of the Ocean Drilling Program, Initial Reports, Vol. 171B, 1998.
- O’Brien, C. L., Lear, C. H., & Wade, B. S.: Cretaceous sea-surface temperature evolution: constraints from TEX86 and planktonic foraminiferal oxygen isotopes, *Earth Sci. Rev.*, 172, 224–247, 2017.
- 780 Ogg, J. G., and Bardot, L.: Aptian through Eocene magnetostratigraphic correlation of the Blake Nose Transect (Leg 171B), Florida continental margin, in Kroon, D., Norris, R. D., & Klaus, A. (Eds.), *Proc. ODP, Sci. Results*, 171B, 1–58, 2001.
- Ogg, J.G., Röhl, U., and Geib, T., :Astronomical tuning of Aptian–Albian boundary interval: oceanic anoxic event OAE 1b through Lower Albian magnetic polarity subchron M”-2r.”, *Eos*, 80:F491–492, 1999.
- 785 Percival L. M. E., Matsumoto, H., Callegaro, S., Erba, E., Kerr, A. C., Mutterlose, J. and Suzuki, K.: Cretaceous large igneous provinces: from volcanic formation to environmental catastrophes and biological crises, in: *Cretaceous Project 200 Volume 1: the Cretaceous World*, edited by: Hart, M. B., Batenburg, S. J., Huber, B. T., Price, G. D., Thibault, N., Wagreich, M. and Walaszczyk, I., Geological Society, London, Special Publications, 544, <https://doi.org/10.1144/SP544-2023-88>, 2024.
- 790 Petrizzo, M. R.: Late Cretaceous planktonic foraminiferal bioevents in the Tethys and in the Southern Ocean record: an overview, *J. Foraminiferal Res.*, 33, 330–337, 2003.
- Premoli Silva, I., Erba, E. and Tornaghi, M.E.: Paleoenvironmental signals and changes in surface fertility in mid Cretaceous Corg-Rich pelagic facies of the Fucoid Marls (Central Italy). *Geobios*, 22, 225–236, 1989.
- 795

- Ramos, J. M. F., Savian, J. F., Franco, D. R., Figueiredo, M. F., Leandro, C. G., Frontalini, F., et al.: Astronomical calibration of the Ocean Anoxic Event 1b and its implications for the cause of mid-Cretaceous events: a multiproxy record, *Paleoceanogr. Paleoclimatol.*, 39, e2024PA004860, <https://doi.org/10.1029/2024PA004860>, 2024a.
- 800 Ramos, J. M. F., Savian, J. F., Franco, D. R., Figueiredo, M. F., Leandro, C. G., & Frontalini, F.: Orbital tuning of short reversed geomagnetic polarity intervals in the Cretaceous Normal Polarity Superchron, *Geophys. Res. Lett.*, 51, e2024GL110530, <https://doi.org/10.1029/2024GL110530>, 2024b.
- 805 Ramos, J.M.F., Savian, J.F., Franco, D. R. et al.: New insights into the early opening of the Equatorial Atlantic Gateway revealed through a magneto-cyclostratigraphy framework from the Brazilian Equatorial Margin. *Commun. Earth Environ.*, 6, 958, <https://doi.org/10.1038/s43247-025-02910-0>, 2025.
- 810 Ramos, J.M.F., Savian, J.F., Franco, D.R. et al.: A novel chronostratigraphic framework for the Aptian–Albian paleoclimate events. *Sci Rep.* <https://doi.org/10.1038/s41598-026-35714-z>, 2026.
- Ryan, W. B. F., Bolli, H. M., Foss, G. N., Natland, J. H., Hottman, W. E., and Foresman, J. B.: Objectives, principle results, operations, and explanatory notes of Leg 40, South Atlantic. In Bolli, H.M., Ryan, W.B.F., et al., *Init. Repts. DSDP*, 40: Washington (U.S. Govt. Printing Office), 5–26, 1978.
- 815 Ryan, W. B. E and Cita, M. B.: Ignorance Concerning Episodes of Ocean-Wide Stagnation, *Marine Geology* 23, 197–215 , 1977.
- Santos, A., Mota, M. A. L. Kern, H. P., Fauth, G. et al.: Earlier onset of the Early Cretaceous Equatorial humidity belt. *Glob. Planet. Change* 208, 1–14, 2022.
- 820 Savin, S. M.: The History of the Earth's Surface Temperature During the Last 100 Million Years, *Annual Reviews of Earth and Planetary Science*, 5, 319–355, 1977.
- 825 Savian, J. F., Trindade, R. I. F., Janikian, L., Jovane, L., Almeida, R. P., Coccioni, R., et al.: The Barremian-Aptian boundary in the Poggio le Guaine core (central Italy): Evidence for magnetic polarity Chron M0r and oceanic anoxic event 1a. *Geological Society of America Special Paper*, 524, 57–78. [https://doi.org/10.1130/2016.2524\(05\)](https://doi.org/10.1130/2016.2524(05)), 2016.
- Schlanger, S. O., and H. C. Jenkyns.: Cretaceous oceanic anoxic events: Causes and consequence, *Geol. Mijnbouw*, 55(3–4), 179–184, 1976.
- 830 Shi, R., He, H., Zhu, R., & Pan, Y.: ISEA reversed event in the Cretaceous normal superchron (CNS): $^{40}\text{Ar}/^{39}\text{Ar}$ dating and paleomagnetic results, *Chin. Sci. Bull.*, 49, 926–930, <https://doi.org/10.1007/bf03184013>, 2004.
- 835 Sinninghe Damsté, J. S., and Köster, J. A.: A euxinic southern North Atlantic Ocean during the Cenomanian/Turonian oceanic anoxic event, *Earth Planet. Sci. Lett.*, 158, 165–173, 1998.
- Steinig, S., Dummann, W., Hofmann, P., Frank, M., Park, W., Wagner, T., and Flögel, S.: Controls on Early Cretaceous South Atlantic Ocean circulation and carbon burial – a climate model–proxy synthesis, *Clim. Past*, 20, 1537–1558, <https://doi.org/10.5194/cp-20-1537-2024>, 2024.
- 840 Tarduno, J. A.: Brief reversed polarity interval during the Cretaceous Normal Polarity Superchron, *Geology*, 18, 683–686, [https://doi.org/10.1130/0091-7613\(1990\)018<0683:BRPIDT>2.3.CO;2](https://doi.org/10.1130/0091-7613(1990)018<0683:BRPIDT>2.3.CO;2), 1990.
- 845 Tarduno, J. A., Sliter, W. V., Kroenke, L., Leckie, M., Mayer, H., Mahoney, J. J., et al.: Rapid Formation of Ontong Java plateau by Aptian mantle Plume volcanism. *Science*, 254(5030), 399–403. <https://doi.org/10.1126/science.254.5030.399>, 1991.

- 850 Tarduno, J. A., Lowrie, W., Sliter, W. V., Bralower, T. J., & Heller, F.: Reversed polarity characteristic magnetizations in the Albian Contessa sections, Umbrian Apennines, Italy: Implications for the existence of a Mid-Cretaceous mixed polarity interval. *Journal of Geophysical Research*, 97(B1), 241–271. <https://doi.org/10.1029/91jb02257>, 1992.
- 855 Tiraboschi, D., Coccioni, R., Jovane, F., and Frontalini, F.: Origin of rhythmic Albian black shales (Piobbico core, central Italy): calcareous nannofossil quantitative and statistical analyses and paleoceanographic reconstructions, *Paleoceanography*, 24, PA4206, 2009.
- Thomson, D. J.: Spectrum estimation and harmonic analysis, *Proc. IEEE*, 70, 1055–1096, 1982.
- Trabucho Alexandre, J., Gilst, R. I. V., Rodriguez-Lopez, J. P., De Boer, P. L.: The sedimentary expression of oceanic anoxic event 1b in the North Atlantic, *Sedimentology*, 58, 1217–1246, 2011.
- 860 VandenBerg, J., Klootwijk, C. T., and Wonders, A. A. H.: The late Mesozoic and Cenozoic movements of the Umbrian Peninsula: further paleomagnetic data from the Umbrian sequence, *Geol. Soc. Am. Bull.*, 89, 133–155, 1978.
- Waltham, D.: Milankovitch Period Uncertainties and their Impact on cyclostratigraphy, *Journal of Sedimentary Research*, 85(8), 990–998, <https://doi.org/10.2110/jsr.2015.66>, 2015.
- 865 Wang, C., Hu, X., Jansa, L., Li, X., & Wang, S.: Upper Cretaceous oceanic red beds in southern Tibet: a major change from anoxic to oxic, deep-sea environments, *Cretac. Res.*, 26, 21–32, 2005.
- 870 Wang, C., Hu, X., Scott, R. W., Wagreich, M., & Jansa, L.: Overview of Cretaceous Oceanic Red Beds (CORBs): a window on global oceanic and climate change, in *Cretaceous Oceanic Red Beds: Stratigraphy, Composition, Origins, and Paleoceanographic and Paleoclimatic Significance*, *SEPM Spec. Publ.* 91, 13–33, 2009.
- Weissert, H. and McKenzie, J. A.: Anoxic Sediments in the Early Cretaceous Tethys Ocean: An Oceanographic Explanation, *International Association of Sedimentologists, European Meeting, Abstracts*, 205 – 207, 1981.
- 875 Weissert, H., McKenzie, J. A., and Channell, J. E. T.: Natural Variations in the Carbon Cycle During the Early Cretaceous', in E. T. Sundquist and W. S. Broecker (eds.), *The Carbon Cycle and Atmospheric CO₂: Natural Variations Archean to the Present*, American Geophysical Union, *Geophysical Monograph*, 32, pp. 531–545, 1985.
- 880 Weissert, H.: C-Isotope stratigraphy, a monitor of paleoenvironmental change: A case study from the early cretaceous. *Surveys in Geophysics*, 10, 1–61, <https://ui.adsabs.harvard.edu/abs/1989SGeo...10....1W>, 1989.
- 885 Weissert, H., Lini, A., Föllmi, K. B., and Kuhn, O.: Correlation of Early Cretaceous carbon isotope stratigraphy and platform drowning events: a possible link?, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 137, 189–203, 1998.
- Yamamoto, A., Abe-Ouchi, A., Shigemitsu, M., Oka, A., Takahashi, K., Ohgaito, R., & Yamanaka, Y.: Global deep ocean oxygenation by enhanced ventilation in the Southern Ocean under long-term global warming, *Glob. Biogeochem. Cycles*, 29, 1801–1815, <https://doi.org/10.1002/2015GB005181>, 2015.
- 890 Zhang, D., Yan, M., Song, C., Zhang, W., Fang, X., & Li, B.: Frequent polarity reversals in the Cretaceous Normal Superchron, *Geophys. Res. Lett.*, 48, 2020GL091501, <https://doi.org/10.1029/2020GL091501>, 2021.