

Projecting changes in rainfall-induced landslide susceptibility across China under climate change

Jinqi Wang^{1,2}, Hao Fang^{2,3,4}, Kai Liu^{1,2*}, Yi Yue², Ming Wang^{1,2}, Bohao Li^{1,2}, Xiaoyi Miao^{1,2}

¹Joint International Research Laboratory of Catastrophe Simulation and Systemic Risk Governance, Beijing Normal University, Zhuhai 519087, China

²School of National Safety and Emergency Management, Beijing Normal University, Beijing 100875, China

³Zhejiang Institute of Geosciences, Hangzhou 310007, China

⁴Qiantang Institute of Geosciences, Hangzhou 310007, China

Correspondence to: Kai Liu (liukai@bnu.edu.cn)

Abstract Landslides pose a significant threat to human lives and property. Evaluating dynamic changes in landslide susceptibility under climate change can provide decision-making support for future disaster prevention. Using historical landslide inventories (2008–2023) and a random forest algorithm, this study developed an annual-scale landslide susceptibility model to assess spatiotemporal patterns of landslide susceptibility across China under different simulated scenarios. The results showed that the model achieved excellent performance (AUC = 0.96), with annual precipitation being the most influential factor (16% contribution). Compared to the baseline (1950–2014), China's landslide susceptibility is projected to increase under future climate conditions. By the late 21st century (2076–2100), the national mean annual precipitation is expected to rise by 59–111 mm, corresponding to an expansion of 1.4×10^5 – 3.4×10^5 km² in areas with median to very high susceptibility across SSP scenarios. Spatially, increased susceptibility is expected mainly in the border region between the South China bedrock hills region (SC) and the Southwest karst mountain region (SW), the western permafrost region of the Qinghai–Tibet Plateau (Tibet), and the Northwest Loess Plateau region (Loess) near the Taihang Mountains, with SSP5-8.5 further amplifying susceptibility toward the end of the century. These findings underscore the necessity of proactive risk management in these identified hotspots to mitigate escalating landslide threats.

1 Introduction

Rainfall-induced landslides rank among the world's most destructive geological disasters (Gutiérrez et al., 2015), having claimed over 48,000 documented lives globally between 2004 and 2016 (Emberson et al., 2020, 2021; Froude and Petley, 2018). China faces particularly severe impacts, with landslide-related fatalities averaging 600 annual deaths since 2000—accounting for nearly 25% of the country's natural disaster mortality (Lin et al., 2022; Lin and Wang, 2018). Climate change is expected to exacerbate this threat through increased frequency and intensity of extreme precipitation events (Gariano et al., 2017; Kirschbaum et al., 2020), potentially elevating regional landslide susceptibility and associated risks. A comprehensive

30 understanding of future susceptibility patterns is therefore critical for developing effective, climate-resilient landslide mitigation strategies.

Landslide occurrence results from the interplay between static predisposing factors and dynamic triggering conditions. Recent advances in susceptibility modelling have increasingly incorporated time-dependent variables to better capture spatiotemporal patterns of landslide risk. Currently, landslide susceptibility mapping (LSM) methodologies fall into four main categories:

35 physical models, expert-driven models, statistical models, and machine learning models (Chang et al., 2019; Li et al., 2019). Among these, machine learning algorithms have demonstrated particular effectiveness in landslide susceptibility modelling due to their capability to capture both linear and nonlinear relationships between causative factors and landslide occurrence (Merghadi et al., 2020). Commonly employed machine learning approaches include random forest (RF), support vector machines (SVM), Extreme gradient boosting (XGBoost), convolutional neural networks (CNN), logistic regression (LR), k-

40 nearest neighbours (KNN), and long short-term memory networks (LSTM) (Avand et al., 2019; Hakim et al., 2022; Huang et al., 2023; Jin et al., 2022; Zhang et al., 2023). RF algorithm has gained particular attention in this field due to its distinctive advantages. First, its parallel computing architecture enables efficient processing of large datasets, while the ensemble approach of multiple decision trees significantly enhances both training efficiency and prediction accuracy. Second, RF incorporates a built-in feature importance evaluation module that assesses variable significance through permutation-based

45 comparison with prediction outcomes (Zhao et al., 2018). Extensive studies have confirmed RF's superior performance in landslide susceptibility assessments (Chen et al., 2017; Dou et al., 2019; Hong et al., 2019; Reichenbach et al., 2018).

In recent years, LSM has increasingly been combined with climate-change scenarios. The aim is to assess how the spatial distribution of rainfall-triggered landslide susceptibility may change across future climate scenarios and time periods. Existing work can be broadly grouped into three types. First, physically coupled approaches combine outputs from global climate

50 models (GCMs) or regional climate models (RCMs) with physically based slope-stability models to examine climate-change impacts on landslides (Hürlimann et al., 2022; Peres and Cancelliere, 2018). Typical models and applications include the Fast Shallow Landslide Assessment Model (FSLAM) (Hürlimann et al., 2022), the Transient Rainfall Infiltration and Grid-Based Regional Slope-Stability model (TRIGRS) (Alvioli et al., 2018), and associated landslide warning models (Ciabatta et al., 2016). Second, threshold methods extract rainfall indicators related to triggering from climate scenarios (e.g. daily rainfall and

55 antecedent cumulative rainfall), and then apply historical or scenario-based thresholds to quantify threshold exceedance (e.g. exceedance frequency or counts); these results are used to update landslide susceptibility (Lin et al., 2022; Wang et al., 2021). Third, downscaled climate variables or rainfall-derived indicators are fed into machine-learning frameworks to produce time-slice susceptibility maps (Park and Lee, 2021; Viet Du et al., 2023). Machine learning captures both linear and non-linear relationships and handles high-dimensional data with complex interactions. It also facilitates the integration of multi-source

60 remote-sensing data and large-area samples, making it particularly suitable for regional-to-national scale susceptibility studies. Despite continued methodological advances, several key limitations remain in landslide susceptibility studies that incorporate climate-change scenarios. On the one hand, the incompleteness and spatial-temporal biases of historical inventories—due to limited event reporting and archival standards—can undermine the representativeness of training samples and inject substantial

uncertainty into model predictions (Lin et al., 2021). This issue may be more pronounced at the national scale (Du et al., 2020).
65 On the other hand, the future climate projections from CMIP6 feature relatively low spatial resolution, and inter-model differences persist among different simulation outputs (Dobler et al., 2024; John et al., 2022). This can further increase the uncertainty of future landslide susceptibility predictions (Reichenbach et al., 2018).
To mitigate these uncertainties, this study developed an annual landslide susceptibility model based on 2008–2023 geological disaster records from the Ministry of Natural Resources of China. The model employs a RF algorithm with feature engineering,
70 combined with statistically downscaled climate projections from NASA Earth Exchange Global Daily Downscaled Projections (NEX-GDDP-CMIP6) to quantify the future changes in landslide susceptibility across China under climate change scenarios. The results can provide critical insights for climate-adaptive land management and disaster risk reduction strategies in vulnerable regions.

2 Materials

75 2.1 Landslide inventory

This study uses the national geohazard records of China for 2008–2023, provided by the Ministry of Natural Resources of the People’s Republic of China, as the basic dataset for landslide susceptibility modelling and future trend analysis. The dataset was compiled annually and covers the whole of China each year, rather than only small areas affected by individual events. Therefore, the study area of each annual inventory is the whole of China. It includes the occurrence time and spatial location
80 of landslides, debris flows, and collapses, which were collectively treated as “landslides” in this study. The dataset includes both post-disaster event records and geohazard potential sites. The records of occurred hazards form an event-based post-disaster inventory, compiled through post-event field investigation and verification by professional geological survey teams organized by local governments, together with multiple sources of information such as remote-sensing interpretation and UAV surveys, and then reported level by level. In addition, the dataset also contains geohazard potential sites identified through
85 annual investigation and screening. The dataset is mainly used for disaster verification, loss assessment, hazard identification, risk management, and geohazard prevention and control. Therefore, the recorded landslide events are mainly distributed in areas where landslides may pose risks to people. The investigation and compilation procedures followed the industry technical specification, “Specification of Geological Hazard Risk Survey and Assessment (1:50,000)”. The minimum identifiable landslide area in the inventory is approximately 10 m². Because the dataset was compiled annually within a unified national
90 reporting framework, the standards for investigation, identification, recording, and compilation were generally consistent across years. As a nationwide operational geohazard dataset, it may still be affected by differences in regional accessibility, reporting practices, and investigation intensity, which may lead to some underreporting in remote areas or for small-scale events. Fig. 1 presents the spatial distribution of landslide events in the dataset. More detailed information is provided in Fig. S1 in the Supplementary Materials.




Figure. 1: Spatial distribution of rainfall-induced geological disaster events and major geological environmental regions in China. Geological environmental regions boundaries were obtained from <https://geocloud.cgs.gov.cn/>. NE: Northeastern wetland ecology region; Yangtz: Huang-Huaihai-Yangtze River Delta plain region; SC: South China bedrock hills region; Loess: Northwest Loess Plateau region; SW: Southwest karst mountain region; NW: Northwest arid desert region; Tibet: permafrost region of the Qinghai-Tibet Plateau.

2.2 Landslide influencing factors

In this study, we selected 14 factors that may influence landslide susceptibility (Table 1). All variables were resampled to a resolution of $1 \text{ km} \times 1 \text{ km}$. For continuous variables such as elevation and distance to river, we used bilinear interpolation, while for discrete variables including lithology and land use type, we employed mode sampling.

Table 1: Datasets and their corresponding landslide influencing factors. “-” indicates no time-series information.

Data Type	Dataset	Resolution	Time Period	Influencing Factor
Geological Features	GLiM	1:3750,000	-	Lithology
	GEM GAF-DB	Vectors	-	Distance to fault
	Global Seismic Hazard Map	0.04°	-	475-year return period PGA
Geomorphometric Features	SRTM Digital Elevation Data Version4	90 m	-	Elevation
				Slope TWI Curvature Plan curvature Profile curvature
Hydrological Features	1-km Monthly Precipitation Dataset for China	1 km	1950-2023	Annual total precipitation
	NEX-GDDP-CMIP6	0.25°	1950-2100	
	Five-level River Dataset of China	1:1,000,000	-	Distance to river
Environmental Features	China's Land-Use/Cover Datasets (CLCD)	30 m	2008-2023	Landcover
	MOD13Q1.006 Terra Vegetation Indices	250 m	2008-2023	NDVI
	16-DayGlobal250m			
	Open Street Map Global Primary Roads	Vectors	-	Distance to road

110

2.2.1 Geological factors

Fault structures and lithological characteristics may affect slope stability by weakening the structural integrity of rock masses. Under the combined influence of earthquakes and heavy rainfall, geologically vulnerable areas are more likely to experience slope failure. This research combines three geological elements. First, the distance to fault was calculated. This calculation
115 used the Euclidean Distance tool in ArcGIS Pro. The data were sourced from the Global Active Faults Database (GAF-DB), which contains spatial information for 13,500 faults (Styron and Pagani, 2020). The database is available at <https://github.com/GEMScienceTools/gem-global-active-faults>. Second, lithology was represented using the Global Lithological Map (GLiM), compiled by the International Union of Geological Sciences (IUGS). The GLiM geodatabase was obtained from https://www.dropbox.com/scl/fi/5v00i8op7a9brmn4qeg8b/LiMW_GIS-2015.gdb.zip. This dataset has a scale

120 of 1:3,750,000. It describes the engineering geological properties of surface rock layers (Hartmann and Moosdorf, 2012). Third,
the 475-year return period peak ground acceleration (PGA) was adopted to characterize the long-term influence of seismic
activity on slope stability. This indicator is not intended to represent earthquake-triggered landslides, but rather to describe
long-term geological structural changes induced by seismic activity. Such changes may persist over time and increase the
susceptibility of slopes to failure under intense or prolonged rainfall. Within this framework, earthquakes are regarded as
125 influencing factors that modify geological structures, rather than as direct triggering mechanisms of landslides. Therefore, the
PGA return period is incorporated as a long-term seismic background factor in rainfall-induced landslide susceptibility
modelling (Leshchinsky et al., 2021). The 475-year RP PGA spatial distribution data came from Johnson et al. (2023). This
dataset can be accessed and downloaded from <https://hazard.openquake.org/gem>.

2.2.2 Geomorphometric factors

130 Geomorphometric factors are the main factors controlling the spatial distribution of surface runoff and soil moisture movement.
They are also key triggers for landslide disasters. This study selects five geomorphological factors: elevation, slope, curvature,
plan curvature, and profile curvature. The elevation data are sourced from the fourth edition of the Shuttle Radar Topography
Mission (SRTM V4) digital elevation database provided by NASA (Jarvis et al., 2008), and the other four topographic factors
were calculated based on ArcGIS Pro. The SRTM V4 dataset is available at <https://srtm.csi.cgiar.org>.

135 **2.2.3 Hydrological factors**

We selected three hydrological factors for this study: annual total precipitation, distance to river, and the topographic wetness
index (TWI). Precipitation data were obtained from the "China 1 km Resolution Monthly Precipitation Dataset (1901–2023)"
provided by the Qinghai-Tibet Plateau Science Data Centre (Ding and Peng, 2020; Peng et al., 2017, 2018, 2019). This dataset
was obtained from <https://doi.org/10.5281/zenodo.3114194>. In this study, TWI was calculated based on SRTM V4 using the
140 hydrological analysis tools in ArcGIS Pro. The river system affects slope stability through two primary mechanisms. First,
riverbank erosion can compromise slope structural integrity. Second, the distance to river factor indirectly modulates slope
instability by influencing surface runoff energy and erosion intensity. We employed the five-level river dataset of China at a
1:1,000,000 scale, which is available from [https://open.geovisearth.com](https://open.geoviseearth.com). We then generated the distance-to-river map using
the Euclidean Distance tool in ArcGIS Pro.

145 **2.2.4 Environmental factors**

Land cover and Normalized Difference Vegetation Index (NDVI) influence the occurrence of landslides by affecting surface
runoff and infiltration. For example, areas with extensive vegetation cover have more permeable surfaces, while areas with
urban land have large impermeable surfaces and lower infiltration rates. Land cover data is sourced from China's Land-
Use/Cover Datasets (Yang and Huang, 2023), which is available at <https://zenodo.org/records/15853565>. The NDVI was
150 calculated by averaging all available MOD13Q1 V6 NDVI images from 2008 to 2023 in Google Earth Engine (Didan, 2021),

which are available at <https://doi.org/10.5067/MODIS/MOD13Q1.061>. Human activities may indirectly affect landslides by altering topography and land cover. For instance, road construction often involves cutting through slopes, weakening slope stability, and damaging soil and rock structures, which increases landslide susceptibility. In this study, road network data for China was obtained from OpenStreetMap (www.openstreetmap.org). The distance to road was then calculated using the
155 Euclidean Distance tool in ArcGIS Pro.

2.2.5 Simulation data

High-resolution climate projections from the NEX-GDDP-CMIP6 dataset (0.25° spatial resolution) were used to drive the simulation of future landslide susceptibility in China. (Thrasher et al., 2022). This dataset was obtained from <https://registry.opendata.aws/nex-gddp-cmip6>. This study comprehensively selects three typical shared socioeconomic
160 pathway (SSP) scenarios: SSP1-2.6 (sustainable development path), SSP2-4.5 (middle-of-the-road path), and SSP5-8.5 (high carbon emission path). These scenarios cover simulation output data from 31 General Circulation Models (GCMs). The study compares the spatial and temporal variation characteristics of mean annual precipitation during the historical baseline period (1950-2014) and future periods: near-term future (2026-2050), mid-term future (2051-2075), and long-term future (2076-
2100). It should be noted that this study focuses on national-scale, long-term variations in annual landslide susceptibility, rather
165 than on the detailed representation of individual landslide events or localized extreme rainfall triggering mechanisms. Within this spatial-temporal framework, the primary objective is to identify the relative spatial patterns and long-term evolution of landslide susceptibility under different climate change scenarios. Previous studies have shown that, at large spatial scales and over long-term periods, CMIP6 climate model outputs can be used to characterize overall trends in climate-change-related landslide processes (Duan et al., 2025; Lin et al., 2022). Building on this framework, this study adopts NEX-GDDP-CMIP6
170 data, which preserve long-term climate change signals while providing higher spatial resolution; therefore, data at this resolution are suitable for landslide susceptibility assessment under climate change.

3 Methodology

3.1 Framework for this work

The assessment of rainfall-triggered landslide susceptibility in China under climate change is structured into six key steps (Fig.
175 2): (1) extracting landslide points from the inventory and generating non-landslide points by random sampling outside buffer zones around landslide locations; (2) applying feature selection techniques to identify the conditioning factors from landslide conditioning variables; (3) constructing a feature matrix by extracting static and dynamic conditioning factors at each sample point, and dividing the dataset into training (70%) and testing (30%) sets; (4) training and optimizing an annual landslide susceptibility model using the RF algorithm; (5) LSM for both historical and future periods by applying downscaled climate
180 projections under different climate change scenarios; (6) analysing the spatiotemporal changes in landslide susceptibility across China under various climate change scenarios.




Figure. 2: Methodological framework of this study

185 3.2 Extracting disaster samples and generation of non-disaster samples

All landslide records were spatialized in ArcGIS Pro, resulting in 93,935 landslide points. For each landslide point, a 3 km buffer was established. Outside all buffers within China, non-landslide points were randomly generated at twice the number of landslide points, but only in areas with human presence to ensure the representativeness of the non-landslide samples. Human presence was determined using the WorldPop R2025A version v1 population data, from which the annual mean population (2015–2023) for China was calculated at 1 km resolution (Bondarenko et al., 2025). Population data are publicly available at <https://hub.worldpop.org/>. To maintain temporal consistency, each non-landslide point was randomly assigned a year between 2008 and 2023, matching the temporal distribution of the landslide point.

3.3 Feature selection

Feature selection is an essential preprocessing step for high-dimensional data analysis, visualization, and modelling. In this study, two methods were employed for feature selection to enhance computational efficiency while preserving the accuracy of the model: Pearson correlation coefficient and tolerance (TOL) and variance inflation factor (VIF).

3.3.1. Pearson correlation coefficient

To quantify the linear relationship between features, the Pearson correlation coefficient was computed. A correlation was considered strong and statistically significant if the absolute value of the correlation coefficient ($|r|$) exceeded 0.7 and the associated p-value was less than 0.05. When such highly correlated features were identified, we addressed potential multicollinearity by removing one of the redundant features. This approach provides a quantitative means of identifying and mitigating multicollinearity among input features (Li et al., 2022).

$$r = \frac{1}{n-1} \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma_x} \right) \left(\frac{Y_i - \bar{Y}}{\sigma_y} \right), \quad (1)$$

Here, r is the Pearson correlation coefficient; n is the sample size; X and Y are the variables; $\bar{X}$ and $\bar{Y}$ are the sample means of X and Y , respectively; σ_x and σ_y are the sample standard deviations.

3.3.2. Tolerance and variance inflation factor

A strong linear relationship between two influencing factors may reduce a model's predictive accuracy. To diagnose potential multicollinearity, TOL and VIF are widely used. Multicollinearity is indicated when $TOL < 0.1$ and $VIF > 10$. In such cases, variables exhibiting high collinearity should be removed (He et al., 2021; O'brien, 2007; Roy and Saha, 2019). The specific formulas are as follows.

$$TOL = 1 - R_i^2 (i = 1, 2, \dots, m), \quad (2)$$

$$VIF = \frac{1}{1 - R_i^2}, \quad (3)$$

Here, R_i^2 represents the coefficient of determination obtained by regressing the i th independent variable on the remaining $m - 1$ independent variables.

3.4 Generation of a feature matrix

Relevant features were extracted for each data point. Dynamic variables were obtained by constructing a time-series three-dimensional matrix (time $\times$ location $\times$ variable) in Python. Static variables, such as elevation and slope, were extracted using the Extract Multi Values to Points tool in ArcGIS Pro. All extracted variables were then integrated into a single feature matrix for model training.

3.5 RF algorithm

RF is an ensemble learning algorithm that improves predictive performance by building and aggregating multiple decision trees (Breiman, 2001). The algorithm employs bootstrap aggregation, whereby each tree is trained on a randomly drawn bootstrap sample from the original dataset, and at each node, a random subset of features is selected for splitting. This

randomization process promotes model diversity and robustness, enhancing accuracy and reducing sensitivity to noise and outliers (Stumpf and Kerle, 2011). By averaging the results of multiple trees, RF effectively reduces variance and the risk of overfitting, making it more robust than individual decision trees. In addition, RF can efficiently handle high-dimensional datasets and categorical variables with minimal preprocessing or encoding. Importantly, RF provides feature importance measures, which facilitate the identification of key variables influencing model predictions (Bureau et al., 2003). These strengths make RF particularly suitable for tasks such as landslide susceptibility mapping, where high accuracy, robustness, and interpretability are essential.

3.6 Model training and performance evaluation

Accurate evaluation of the RF model is essential for predicting future changes in landslide susceptibility. In this study, we employed several widely used evaluation metrics, including accuracy, precision, recall, F1-score, the receiver operating characteristic (ROC) curve, and the area under the curve (AUC). These metrics are derived from the confusion matrix (M and M.N, 2015), with values closer to 1 indicating better model performance. Specifically, the ROC curve visualizes the trade-off between the false positive rate (FPR) on the x-axis and the true positive rate (TPR) on the y-axis.

For model development and validation, we randomly split the dataset into training (70%) and testing (30%) subsets. RF model was trained on the training subset, and all performance metrics were evaluated on the independent testing subset to avoid overfitting. Model hyperparameters were optimized and set to $n_estimators = 886$, $max_depth = 20$, $max_features = "sqrt"$, and $min_samples_split = 6$, using the Gini index as the splitting criterion.

$$Accuracy = \frac{TP+TN}{TP+TN+FP+FN}, \quad (4)$$

$$Precision = \frac{TP}{TP+FP}, \quad (5)$$

$$Recall = TPR = \frac{TP}{TP+FN}, \quad (6)$$

$$F1 - score = \frac{2TP}{2TP+FP+FN}, \quad (7)$$

$$FPR = \frac{FP}{FP+TN}, \quad (8)$$

Here, TP (true positive) and TN (true negative) represent the number of landslide and non-landslide samples that are correctly classified, respectively. FP (false positive) refers to the number of non-landslide samples that are incorrectly classified as landslides, while FN (false negative) refers to the number of landslide samples that are incorrectly classified as non-landslides.

3.7 Assessment of future landslide susceptibility

We selected 31 GCMs from the NEX-GDDP-CMIP6 dataset. Each of these models includes the SSP1-2.6, SSP2-4.5, and SSP5-8.5 scenarios. Then, we evaluated the performance of these models against historical observations using the root mean

square error (RMSE) to identify those suitable for China (Table S1). Subsequently, based on the selected models, we constructed a multimodel ensemble of annual precipitation, which was used to establish the historical baseline and to project future landslide susceptibility under different climate scenarios (SSP1-2.6, SSP2-4.5, and SSP5-8.5) and future periods, including the near-term future (2026–2050), mid-term future (2051–2075), and long-term future (2076–2100). To further account for uncertainty in rainfall-induced landslides across China, we applied a 95% confidence interval derived from the T-distribution. Then, landslide susceptibility was classified into five levels using the Jenks natural breaks method. To ensure comparability of future results, we applied the classification thresholds derived from the baseline period to all subsequent assessments under different future periods and scenarios. Finally, we assessed the impact of climate change on landslide susceptibility by analysing the changes in susceptibility relative to the baseline period across the different future periods and climate change scenarios.

4. Results

4.1 Selection of conditioning factors

First, Pearson correlation coefficients were used to select the most suitable landslide influencing factors from 12 continuous variables. The correlation coefficients among these 12 factors are shown in the correlation matrix (Fig. 3(a)). A strong positive correlation was observed between plan curvature and curvature ($r = 0.85$, $p < 0.05$); therefore, curvature was removed to avoid multicollinearity. To further verify whether the remaining 11 continuous variables and 2 categorical variables were suitable for landslide susceptibility modelling, multicollinearity tests were conducted (Fig. 3(b-c)). All factors had TOL values greater than 0.1 and VIF values less than 10, indicating no multicollinearity among the 13 influencing factors. The final selected factors included annual total precipitation, TWI, elevation, 475-year RP PGA, NDVI, plan curvature, profile curvature, slope, distance to road, distance to fault, distance to river, landcover, and lithology.




Figure. 3: The results of feature selection in this study: (a) Pearson correlation coefficients between landslide influencing factors (‘×’ indicates no significant linear relationship [p value > 0.05]) (b) TOL and VIF for different landslide influencing factors. M: Lithology, N: Landcover.

4.2 Model evaluation

Model performance evaluation is a crucial component in the analysis of future rainfall-induced landslide susceptibility. In this study, 30% of the sample data were utilized as a test set to assess the performance of the RF model. As illustrated in Fig. 4, model performance was visually analysed through the ROC curve and the confusion matrix. The RF model demonstrated good predictive performance, with an AUC value as high as 0.96. In the test set, the model achieved an accuracy of 0.89, a precision of 0.87, a recall of 0.80, and an F1 score of 0.83, validating its robust predictive performance. Additionally, through the analysis of Gini coefficients in the RF model, we explored the interpretability of the model. The results revealed that annual total precipitation was the key factor influencing model predictions, with a contribution rate as high as 16%, significantly higher than other factors. Slope contributed 12% and elevation contributed 10% to the model (see Fig. S3).




Figure. 4: Performance evaluation of the RF model. (a) The ROC curve demonstrates the model’s classification capability, with an AUC of 0.96. (b) The confusion matrix provides a detailed breakdown of the prediction results, including the numbers of correctly and incorrectly classified samples in each category.

4.3 Influence of key factors on landslide susceptibility

We analysed the relationship between landslide susceptibility and key influencing factors in China during the historical baseline period (1950–2014). The top six continuous variables identified by RF importance ranking were categorized, and the area proportions of each category within different landslide susceptibility classes were calculated (Fig. 5). The results show that mean annual precipitation, slope, and NDVI are strongly positively correlated with landslide susceptibility. In areas with mean annual precipitation greater than 800 mm, the proportions of very high, high, and median susceptibility zones are 62.15%, 66.51%, and 41.65%, respectively (Fig. 5a). Landslide susceptibility also increases markedly with slope. In areas with slopes steeper than 20°, the proportions of very high, high, and median susceptibility zones reach 51.40%, 35.18%, and 27.13%, respectively (Fig. 5b). Similarly, in regions where NDVI exceeds 0.5, these proportions are 73.91%, 63.65%, and 35.18%, respectively (Fig. 5f). These findings indicate that high landslide susceptibility is closely associated with humid environments, steep terrain, and relatively high NDVI values. Elevation exhibits a nonlinear relationship with landslide susceptibility, with higher susceptibility concentrated in both low- and high-elevation regions. Specifically, 42.78%, 45.49%, and 27.40% of the very high, high, and median susceptibility zones are distributed below 1000 m, respectively, while 32.36%, 30.07%, and 51.01% are located above 3000 m (Fig. 5c). Landslide susceptibility is generally higher within 0–50 km of faults, decreases at intermediate distances, and shows a secondary increase at 200–500 km (Fig. 5e). The relationship between 475-year RP PGA and landslide susceptibility is nonlinear (Fig. 5d). In general, areas with very high and high susceptibility are more concentrated in the lowest PGA class, while their proportions decrease in the intermediate classes and increase slightly again in the highest

PGA class. This suggests that seismic background conditions may not act as a direct trigger of rainfall-induced landslides, but may indirectly increase slope sensitivity to subsequent rainfall by damaging geologic structures and reducing slope stability.




Figure. 5: Area proportions of landslide susceptibility classes across different categories of the top six continuous variables during the historical baseline period (1950–2014): (a) mean annual precipitation, (b) slope, (c) elevation, (d) 475-year RP PGA, (e) distance to fault, and (f) NDVI. Colours indicate susceptibility levels from very low to very high.

We further analysed the effects of multi-factor combinations on the spatial distribution of landslide susceptibility in China during the historical baseline period (1950–2014). Using the top three variables ranked by feature importance derived from the RF model (mean annual precipitation, slope, and elevation), we calculated the area proportion of regions with median to very high landslide susceptibility under different combinations, and visualized the results as heatmaps (Fig. 6). In the precipitation–slope combinations (Fig. 6a), the proportion of median to very high susceptibility areas generally increases with increasing slope, and this tendency becomes more pronounced under relatively high precipitation conditions. The highest value (93.61%) is observed in areas with slopes greater than 30° and mean annual precipitation of 800–1200 mm. In the precipitation–elevation combinations (Fig. 6b), high susceptibility is mainly concentrated in areas with relatively high precipitation, the effect of elevation is nonlinear; the maximum proportion (96.56%) is observed in areas above 3000 m with precipitation of 800–1200 mm. The elevation–slope combinations (Fig. 6c) further indicates that highly susceptible areas tend to occur where

steep slopes coincide with relatively low and high elevations. In particular, the 0–500 m and >3000 m elevation ranges exhibit higher landslide susceptibility under steep-slope conditions.





Figure. 6: Area proportions (%) of median to very high landslide susceptibility during the historical baseline period (1950–2014), based on combinations of key continuous variables: (a) mean annual precipitation and slope, (b) mean annual precipitation and elevation, and (c) elevation and slope. Each cell represents the proportion of areas with median to very high susceptibility (classes 3–5) within the total area corresponding to that specific combination of factor ranges.

4.4 Future precipitation changes in China

Fig. 7 shows projected changes in mean annual precipitation across China and its geological environmental zones under three climate scenarios and future periods, relative to the baseline. Overall, multimodel ensemble simulations show that future mean annual precipitation across China exhibits an increasing trend. The relative increases and associated uncertainties are greater under the SSP5-8.5 than under the SSP1-2.6 and SSP2-4.5. Compared to the near-term (2026–2050) and mid-term future (2051–2075), the long-term future (2076–2100) generally shows larger increases. According to projections from the NEX-GDDP-CMIP6 multimodel ensemble, mean annual precipitation in China is expected to increase by 59 mm, 63 mm, and 111 mm under the SSP1-2.6, SSP2-4.5, and SSP5-8.5, respectively, during the long-term future from 2076 to 2100. These correspond to increases of 10.46%, 11.18%, and 19.14% relative to the baseline.

Projected changes in mean annual precipitation exhibit substantial spatial heterogeneity across different geological environmental zones. In the NW, where the baseline mean annual precipitation is only 136 mm, the relative increase in future mean annual precipitation remains low. During the long-term future under the SSP5-8.5, the increase is 59 mm. In geological environmental zones where baseline mean annual precipitation ranges from 400 to 900 mm, the relative increases in mean annual precipitation are generally higher, particularly in the Tibet and Yangtz. For instance, during the long-term future, under the SSP5-8.5, the largest increases are projected in these regions, reaching 170 mm and 147 mm, respectively. In the Southern region including SC and SW, where baseline mean annual precipitation exceeds 1000 mm, the absolute increase is relatively

smaller compared to areas with baseline mean annual precipitation between 400 and 1000 mm. In the geological environment division, the associated uncertainties are the highest.



355

Figure. 7: Changes in multimodel mean annual precipitation under different future scenarios and time periods relative to the baseline (1950–2014). Bars represent the mean change derived from the multimodel ensemble, while error bars indicate the 95% confidence interval, reflecting the uncertainty in projected mean annual precipitation. The value in the upper right corner of each panel represents the mean annual precipitation during the baseline period.


4.5 Future landslide susceptibility in China

360

Fig. 8(a) illustrates the spatial distribution of landslide susceptibility during the baseline period (1950–2014). The results show that approximately 61.51% of the simulated area in China exhibited low to very low susceptibility, primarily located in northern Tibet, southern NW, the northwestern Loess, and central NE. Regions with moderate to very high susceptibility accounted for 38.49% of the total simulated area, mainly distributed across SC, SW, eastern Tibet, and the Loess–Taihang Mountain area. Notably, in the SC, approximately 81.11% of the area was exposed to median to very high landslide susceptibility. Fig. 9 illustrates the increase in the area of median to very high landslide susceptibility zones in China compared to the baseline period across future scenarios and periods. At the national scale, the spatial extent of median to very high susceptibility zones

is projected to expand under all future scenarios and periods. The expansion is more pronounced under SSP5-8.5 compared to
 365 SSP1-2.6 and SSP2-4.5. In addition, long-term future shows a more significant increase in susceptibility area than near-term
 and mid-term future. In the baseline period, the proportion of median to very high susceptibility areas in China is 38.49%.
 During the long-term future under the SSP5-8.5, it increases to 42.03%, corresponding to a spatial expansion of approximately
 $3.4 \times 10^5 \text{ km}^2$.

At the regional scale, all geological environmental zones show an increasing trend in the extent of median to very high landslide
 370 susceptibility areas. The Loess and the Tibet exhibit the most significant expansion (Fig. 8, Fig. 9, and Fig. S5). In the baseline
 period, the proportion of areas with median to very high susceptibility in these regions is 10.21% and 60.79%, respectively.
 During the long-term future under the SSP5-8.5, these proportions increase to 17.16% and 69.04%, corresponding to area
 increases of $6.1 \times 10^4 \text{ km}^2$ and $2.2 \times 10^5 \text{ km}^2$. These expansions mainly occur in western Tibet, and around the Taihang
 Mountains in the Loess (Fig. 8 and Fig. S4). During the long-term future under the SSP5-8.5, other regions such as SC, SW
 375 and NE also show increases. According to the multimodel ensemble averages, the proportion of median to very high
 susceptibility areas in these regions increase by approximately $1.6 \times 10^4 \text{ km}^2$, $2.3 \times 10^4 \text{ km}^2$, and $1.1 \times 10^4 \text{ km}^2$, respectively,
 compared to the baseline. The expansions are mainly located in northern SC, northern SW and southern NE. The NW and
 Yangtz show relatively small increases in median to very high susceptibility areas based on the ensemble mean. (Fig. 8).



380 **Figure. 8:** Spatial pattern of landslide susceptibility in China during (a) the baseline period (1950–2014), and (b) the long-term future (2076–2100) under the SSP5-8.5.




Figure. 9: Changes in the area of regions with median to very high landslide susceptibility under different future periods and SSPs, relative to the baseline period. Bars represent the multimodel ensemble mean changes, and error bars indicate the 95% confidence interval of the ensemble mean.

4.6 Revealing the relationship between landslides and conditioning factors

To further investigate the relationship between landslide events and conditioning factors, this study selected the top nine conditioning factors based on feature importance rankings from a random forest model for a graded statistical analysis (Fig. 10). For continuous variables, the natural break method was applied for classification, and landslide event frequencies were calculated for each interval. Landslide hazard frequency was defined as the "number of landslide samples/total sample size," where the total sample consisted of hazard points and twice the number of non-hazard points. The results showed a clear positive relationship between annual total precipitation and landslide frequency (Fig. 10a). Landslide frequency generally increased with annual precipitation and exceeded 50% when annual precipitation was greater than 1147 mm. Landslide frequency increased markedly with slope and exceeded 60% when the slope was greater than 25° (Fig. 10b). In terms of

elevation, the frequency of geohazards exhibited an M-shaped pattern with increasing altitude. The frequency was lowest in the -217–112 m elevation range, at only 16%, whereas it reached a peak in the 2338–3514 m elevation band, with the highest value of 59% (Fig. 10c). For the 475-year return period PGA, rainfall-induced landslides were mainly concentrated in the low-value intervals, while the frequency showed a secondary increase in some high-value intervals. This may suggest that seismic background conditions indirectly influence rainfall-induced landslides by weakening slope stability (Fig. 10d). Landslide events were more numerous near fault lines, especially within 0–40 km. However, landslide frequency did not decrease monotonically with fault distance and reached relatively high values in some intermediate-distance intervals (Fig. 10e). NDVI, which reflects vegetation coverage, showed an increasing trend in landslide frequency as NDVI values increased. When NDVI exceeded 0.56, landslide frequency was generally greater than 45% (Fig. 10f). Profile curvature showed a pattern of higher geohazard frequency at both ends and lower frequency in the middle: the frequency was lowest in the near-zero interval (-0.02–0.01), at only 15%, but increased markedly toward both negative and positive extreme ranges, reaching as high as 70%. This suggests that nearly planar slopes are relatively stable, whereas strongly concave or convex slopes are more susceptible to rainfall-induced instability (Fig. 10g). TWI showed a negative correlation with geohazard frequency: as TWI increased, the frequency decreased continuously from 60% in the low-value interval of 14.2–15.9 to 1.5% in the high-value interval of 25.8 and above. This indicates that geohazards are more likely to occur in areas with lower TWI (Fig. 10h). Among all lithological types, siliciclastic sedimentary rocks (ss) contained the largest number of landslide events, with 26,298 cases (Fig. 10i).

415




Figure. 10 The relationship between landslide points and non-landslide points and conditioning factors. (a) Precipitation: Annual total precipitation, (b) Slope, (c) Elevation, (d) 475-year return period PGA, (e) Distance to Fault, (f) NDVI, (g) Profile curvature, (h) TWI, and (i) Lithology.

420 5. Discussion

This study investigated the future trends of rainfall-induced landslide susceptibility under various climate change scenarios using SSP1-2.6, SSP2-4.5, and SSP5-8.5 from the NEX-GDDP-CMIP6 climate dataset. A RF model, trained with a relatively complete landslide inventory, was employed for this analysis. In the landslide susceptibility modelling, we focused on rainfall-triggered landslide events. During model construction, landslides, rockfalls, and debris flows were grouped into a single

425 “landslides” category, which is a common practice in landslide inventory compilation. To assess the influence of this merging, we compared the susceptibility maps derived from the grouped-category model with those from a landslide-only model. The two maps showed a relatively high degree of consistency (Pearson’s $r = 0.912$; $MSE = 0.198$), indicating that the merging had a relatively small impact on the overall spatial pattern of susceptibility, although minor numerical differences remained (Fig. S5). In addition, we compared susceptibility maps produced using two different non-landslide sampling strategies: random

430 sampling across the whole study area and sampling restricted to areas with human presence. The results revealed substantial differences between the two approaches. When non-landslide samples were randomly generated across the whole of China, susceptibility tended to be overestimated in densely recorded regions, such as SC and SW, while being underestimated in sparsely recorded regions, such as NW and Tibet (Fig. S6). This was also reflected in the feature-importance results, where whole-area random sampling led to lower importance of geomorphological and lithological factors (Fig. S3). This is likely

435 because the landslide inventory used in this study was compiled by integrating post-disaster investigations and annual geological hazard potential-site surveys, and these records are relatively more concentrated in areas with human settlements and activities. Under such conditions, randomly generating non-landslide samples across the entire study area may incorrectly treat poorly surveyed or uninhabited regions as true negatives, thereby amplifying the effect of spatial inventory incompleteness and increasing the uncertainty and spatial bias in susceptibility mapping results (Reichenbach et al., 2018).

440 Therefore, for inventories such as the one used in this study, in which records are more concentrated in areas with human presence, non-landslide samples may be better generated within inhabited areas to reduce spatial sampling bias.

In landslide susceptibility modelling, this study incorporated annual total precipitation as one of the precipitation-related predictors. It should be noted that monthly precipitation may better represent antecedent rainfall and soil saturation. However, in the context of this study, annual total precipitation can be regarded as a representation of the regional long-term

445 hydroclimatic background: it reflects overall precipitation regimes and their interannual variability and may influence the spatial patterns of landslide susceptibility at the regional scale through its effects on hydro-environmental conditions, including soil moisture, vegetation cover, and slope stability (Reichenbach et al., 2018; Lin et al., 2021). In addition, the landslide inventory released for this study includes only the year of occurrence; the month and day of occurrence are not available. This limits the ability to establish a temporally consistent linkage with monthly or daily antecedent precipitation indices. The use

450 of annual total precipitation helps maintain temporal comparability between the precipitation indicator and the landslide timing resolution and serves as a proxy for long-term wetness conditions. Previous studies at national or regional scales have also commonly adopted annual-scale precipitation variables (e.g., annual total precipitation or multi-year mean annual precipitation) as climatic background factors to provide a broad-scale constraint in landslide susceptibility modelling (Reichenbach et al., 2018; Lin et al., 2021). The feature importance results of this study indicated that annual total precipitation had the highest

455 contribution, which further highlighted the importance of this variable at the national scale (Fig. S3). In addition to precipitation, geomorphological factors also showed high importance, with slope, elevation, profile curvature, and TWI ranking second, third, seventh, and eighth, respectively. This suggests that terrain morphology and terrain-controlled hydrological processes play important roles in landslide occurrence. Landslides are more likely to occur on relatively steep slopes and in both low-

and high-elevation settings under different environmental backgrounds. In such areas, unfavorable terrain conditions and runoff accumulation may promote slope failure. In addition, the relatively high importance of the 475-year return period PGA suggests that seismic background conditions may act as an indirect conditioning factor for rainfall-induced landslides. Past strong ground motion may weaken geological structures and reduce slope stability. NDVI also showed relatively high importance, which may reflect humid hilly and mountainous environments where increased soil moisture, elevated pore-water pressure, and runoff convergence can further enhance landslide triggering probability (Li and Duan, 2024). Lithology is also an important conditioning factor because differences in material strength, weathering resistance, and hydrological properties among lithological units can affect slope stability and landslide susceptibility.

Based on these inputs and the RF framework, our projections indicate an overall increase in China's landslide susceptibility in the future compared to the baseline. This increase is particularly notable under the SSP5-8.5 scenario during the long-term future. These results are consistent with previous research. For example, Du et al. (2025) predicted increased rainfall-induced landslide susceptibility in southern Jiangxi Province using generalised additive models (GAMs). Similarly, Guo et al. (2023) projected an increase in shallow landslide susceptibility in Wanzhou County based on the fast shallow landslide assessment model (FSLAM). Lin et al. (2022) also found a potential increase in future landslide susceptibility and frequency across China using a generalized additive mixed effects model (GAMM). These studies collectively support our finding of a growing trend in future landslide susceptibility. Furthermore, we analysed regional variations by dividing China into seven distinct areas. Our analysis revealed specific regional changes in landslide susceptibility. In particular, the border region between SC and SW, western Tibet, and the Loess–Taihang Mountain area were identified as key regions where landslide susceptibility is likely to intensify, especially under SSP5-8.5.

The accuracy of landslide inventories is a key source of uncertainty in susceptibility mapping at the national scale. It is generally assessed along three dimensions: positional accuracy, thematic accuracy, and completeness. Regarding positional accuracy, previous studies have shown that, once the landslide extent is correctly delineated, using either the scarp or a random location within the mapped extent as the locational reference does not significantly affect the overall susceptibility results (Margottini et al., 2013; Petschko et al., 2013). However, since the scarp is more readily identifiable, adopting it as a consistent locational reference can reduce positional uncertainty and thereby improve the spatial accuracy and reliability of susceptibility mapping. Therefore, the landslide points in our study were based on the scarp as the locational reference. Thematic accuracy relates to the accuracy of classification attributes, including landslide type, size, among others. Small landslides, due to their limited spatial extent and ephemeral morphological expression, are often difficult to capture in national-scale inventories, leading to systematic omissions or under-reporting. To examine this issue, modeling and mapping were performed for China after randomly removing 50% and 100% of the small-landslide samples, respectively (Fig. S7). Relative to the results based on the complete inventory, the exclusion of small-landslide records had a pronounced effect on susceptibility: susceptibility decreased markedly in SC, whereas it increased markedly in Tibet and SW. When only 50% of small landslides were removed, the overall susceptibility pattern remained highly consistent with that from the complete inventory. With respect to completeness, in SC and SW—where landslides are frequent and research attention is high—inventory records are relatively

adequate. By contrast, in NW, NE, and Tibet, constrained by terrain conditions, data accessibility, and limited monitoring capacity, landslide samples are relatively scarce (Fig. S8) (Liu et al., 2013; Liu and Miao, 2018). To examine this issue, we randomly removed 50% and 75% of the samples in NW, NE, and Tibet and then performed modeling and mapping (Fig. S9). Relative to the complete-inventory results, these sample-poor regions exhibited an overall decrease in susceptibility as more samples were removed, whereas regions with sufficient samples showed almost no change. Based on the above experiments, in the compilation of national-scale landslide inventories, we suggest adopting the scarp as the locational reference, improving thematic classification quality, and continuously supplementing and refining inventory records to enhance the robustness and reliability of susceptibility-mapping results.

Currently, CMIP6 GCMs are widely used to explore the impact of climate change on landslides. This study integrated the latest NEX-GDDP-CMIP6 GCM dataset, encompassing SSP1-2.6, SSP2-4.5, and SSP5-8.5. These three scenarios were selected to provide a representative range of future socioeconomic and radiative forcing conditions, thereby spanning a broad range of plausible future precipitation changes. This design enabled us to examine how rainfall-induced landslide susceptibility may evolve under contrasting but plausible climate pathways and to characterize the range of possible national-scale outcomes. This dataset offers globally downscaled scenarios from CMIP6 simulations at a $0.25^\circ \times 0.25^\circ$ resolution. It employs bias correction and spatial disaggregation techniques, incorporating ground observations and raw GCM data for downscaling (Thrasher et al., 2022), thereby enhancing resolution while preserving long-term climate trends (Thrasher et al., 2012). This higher-resolution future climate dataset enabled a more precise assessment of rainfall-induced landslide susceptibility under climate change. First, we selected climate models suitable for the China region by comparing historical observed and simulated precipitation using RMSE. Following this selection, we employed a multimodel ensemble method to project future landslide susceptibility. To quantify the associated uncertainty, we used a 95% confidence interval based on the t-distribution. These steps helped reduce uncertainty in future landslide susceptibility projections. Furthermore, this study used future climate data to analyse landslide susceptibility under low, medium, and high socioeconomic development scenarios. This framework can be combined with future population and economic conditions to assess the potential impact and risks of future landslides on human society (Emberson et al., 2020; Gariano and Guzzetti, 2016). Consequently, the findings of this research provide a robust data foundation for further investigation into the potential socioeconomic risks and impacts associated with future landslides. In addition, we acknowledge that projection uncertainty generally increases toward later periods (e.g., 2051–2075 and especially 2076–2100). Our results are interpreted as scenario-dependent envelopes of plausible long-term change rather than deterministic predictions for specific years.

This study systematically assesses the spatial distribution and future trends of rainfall-induced landslide susceptibility in China and its geological environmental regions under climate change at a national scale. By integrating a relatively complete national-scale landslide inventory and employing high-resolution precipitation data from NEX-GDDP-CMIP6, this study helps reduce uncertainties associated with historical landslide records and future rainfall projections. The proposed framework helps improve the accuracy and reliability of national-scale landslide susceptibility assessment. The results provide scientific support for national-scale landslide management and disaster mitigation strategies.

The main limitations of this study are as follows: (1) This research developed an annual-scale future landslide susceptibility model for China. However, short-duration intense rainfall is a recognized critical trigger for rainfall-induced landslides, and analysing the seasonal variability of precipitation could provide further insights into future landslide susceptibility changes (Lin et al., 2021). Future advancements in global navigation satellite systems (GNSS) and aerospace remote sensing (RS) technologies, which enable the acquisition of more precise spatiotemporal information on landslide events and enhance the dynamic representation of landslide influencing factors, hold the potential to improve modelling reliability. Consequently, future work could explore monthly landslide susceptibility changes in China and its geological environmental zones under climate change, building upon improved landslide inventories and incorporating data from these advanced technologies. (2) The resolution constraints of the future climate model data introduce a limitation in predicting landslide susceptibility across various future periods and scenarios. This may limit the ability to accurately capture localized precipitation patterns. As a result, this could lead to either underestimation or overestimation of local landslide susceptibility, thereby increasing uncertainty in susceptibility projections. (3) As a national-scale analysis across China's vast territory, the modelling was conducted using influencing factors at a 1 km resolution, which may not fully capture local heterogeneity and could increase prediction uncertainty.

6. Conclusion

This study combined the latest NEX-GDDP-CMIP6 climate projections with a RF model to assess future changes in landslide susceptibility across China and its geological environmental zones under different climate scenarios. The model demonstrated good performance, with an overall accuracy of 0.89, precision of 0.87, recall of 0.80, F1-score of 0.83, and an AUC of 0.96. These results indicate that the model effectively captures spatial patterns of landslide susceptibility. Among all variables, annual total precipitation contributed the most to the model's output, with a relative importance of 16%. Under the selected scenarios, China's mean annual precipitation is projected to increase by 59 to 111 mm by the end of the 21st century. Compared to the baseline period, this change is expected to result in an increase of $1.4 \times 10^5 \text{ km}^2$ to $3.4 \times 10^5 \text{ km}^2$ in the area classified as having median to very high landslide susceptibility. These increases are more pronounced in the long-term future and under SSP5-8.5. Spatially, several regions are projected to become more vulnerable. Notably, the border region between SC and SW, western Tibet, and the Loess–Taihang Mountain area are projected to be the main hotspots of intensified landslide susceptibility, with the most pronounced expansion of median to very high susceptibility zones occurring in these areas. These regions should be prioritized in future landslide susceptibility assessments and climate change adaptation strategies.

Data availability

Data that support the findings of this study are available from the corresponding author upon reasonable request.

Author contributions

J.W. designed the research, conducted the formal analysis, and wrote the original draft. H.F. provided the data and related resources. K.L. contributed to methodology, visualization, review, and funding acquisition. Y.Y. contributed to editing the manuscript. M.W. supervised the research. B.L. was responsible for software development and validation. X.M. contributed to validation.

Competing interests

The contact author has declared that none of the authors has any competing interests.

Disclaimer

Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Special issue statement

This manuscript is submitted to the NHESS Special Issue titled “The influence of landslide inventory quality on susceptibility and hazard map reliability” (1 July 2024 – 1 April 2026), edited by Santangelo et al.

Financial support

This work was supported by the General Program of National Natural Science Foundation of China (grant number 42377467) and the Fundamental Research Funds for the Central Universities (grant number 2243300007).

References

Avand, M., Janizadeh, S., Naghibi, S. A., Pourghasemi, H. R., Khosrobeigi Bozchaloei, S., and Blaschke, T.: A Comparative Assessment of Random Forest and k-Nearest Neighbor Classifiers for Gully Erosion Susceptibility Mapping, *Water*, 11, 2076, <https://doi.org/10.3390/w11102076>, 2019.

Alvioli, M., Melillo, M., Guzzetti, F., Rossi, M., Palazzi, E., Von Hardenberg, J., Brunetti, M. T., and Peruccacci, S.: Implications of climate change on landslide hazard in Central Italy, *Science of The Total Environment*, 630, 1528–1543, <https://doi.org/10.1016/j.scitotenv.2018.02.315>, 2018

Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.

- Bureau, A., Dupuis, J., Hayward, B., Falls, K., and Van Eerdewegh, P.: Mapping complex traits using Random Forests, *BMC Genet*, 4, S64, <https://doi.org/10.1186/1471-2156-4-S1-S64>, 2003.
- Bondarenko, M., Priyatikanto, R., Tejedor-Garavito, N., Zhang, W., McKeen, T., Cunningham, A., Woods, T., Hilton, J., Cihan, D., Nosatiuk, B., Brinkhoff, T., Tatem, A., and Sorichetta, A.: The spatial distribution of population in 2015–2030 at a resolution of 30 arc (approximately 1 km at the Equator), R2025A version v1, WorldPop, School of Geography and Environmental Science, University of Southampton, <https://doi.org/10.5258/SOTON/WP00845>, 2025.
- Chang, K.-T., Merghadi, A., Yunus, A. P., Pham, B. T., and Dou, J.: Evaluating scale effects of topographic variables in landslide susceptibility models using GIS-based machine learning techniques, *Sci Rep*, 9, 12296, <https://doi.org/10.1038/s41598-019-48773-2>, 2019.
- Chen, W., Xie, X., Wang, J., Pradhan, B., Hong, H., Bui, D. T., Duan, Z., and Ma, J.: A comparative study of logistic model tree, random forest, and classification and regression tree models for spatial prediction of landslide susceptibility, *CATENA*, 151, 147–160, <https://doi.org/10.1016/j.catena.2016.11.032>, 2017.
- Ciabatta, L., Camici, S., Brocca, L., Ponziani, F., Stelluti, M., Berni, N., and Moramarco, T.: Assessing the impact of climate-change scenarios on landslide occurrence in Umbria Region, Italy, *Journal of Hydrology*, 541, 285–295, <https://doi.org/10.1016/j.jhydrol.2016.02.007>, 2016.
- Didan, K.: MODIS/Terra Vegetation Indices 16-Day L3 Global 250m SIN Grid V061, NASA Land Processes Distributed Active Archive Centre, <https://doi.org/10.5067/MODIS/MOD13Q1.061>, 2021.
- Ding, Y. and Peng, S.: Spatiotemporal Trends and Attribution of Drought across China from 1901–2100, *Sustainability*, 12, 477, <https://doi.org/10.3390/su12020477>, 2020.
- Duan, Y., Ding, M., He, Y., Zheng, H., Delgado-Téllez, R., Sokratov, S., Dourado, F., and Fuchs, S.: Global projections of future landslide susceptibility under climate change, *Geoscience Frontiers*, 16, 102074, <https://doi.org/10.1016/j.gsf.2025.102074>, 2025.
- Dobler, A., Benestad, R. E., Lussana, C., and Landgren, O.: CMIP6 models project a shrinking precipitation area, *npj Clim Atmos Sci*, 7, 239, <https://doi.org/10.1038/s41612-024-00794-z>, 2024.
- Dou, J., Yunus, A. P., Tien Bui, D., Merghadi, A., Sahana, M., Zhu, Z., Chen, C.-W., Khosravi, K., Yang, Y., and Pham, B. T.: Assessment of advanced random forest and decision tree algorithms for modeling rainfall-induced landslide susceptibility in the Izu-Oshima Volcanic Island, Japan, *Science of The Total Environment*, 662, 332–346, <https://doi.org/10.1016/j.scitotenv.2019.01.221>, 2019.
- Du, B., Wang, Y., Fang, Z., Liu, G., and Tian, Z.: Spatiotemporal modeling and projection framework of rainfall-induced landslide risk under climate change, *Journal of Environmental Management*, 373, 123474, <https://doi.org/10.1016/j.jenvman.2024.123474>, 2025.
- Du, J., Glade, T., Woldai, T., Chai, B., and Zeng, B.: Landslide susceptibility assessment based on an incomplete landslide inventory in the Jilong Valley, Tibet, Chinese Himalayas, *Engineering Geology*, 270, 105572, <https://doi.org/10.1016/j.enggeo.2020.105572>, 2020.

- 615 Emberson, R., Kirschbaum, D., and Stanley, T.: New global characterisation of landslide exposure, *Natural Hazards and Earth System Sciences*, 20, 3413–3424, <https://doi.org/10.5194/nhess-20-3413-2020>, 2020.
- Emberson, R., Kirschbaum, D., and Stanley, T.: Global connections between El Nino and landslide impacts, *Nat Commun*, 12, 2262, <https://doi.org/10.1038/s41467-021-22398-4>, 2021.
- Froude, M. J. and Petley, D. N.: Global fatal landslide occurrence from 2004 to 2016, *Natural Hazards and Earth System Sciences*, 18, 2161–2181, <https://doi.org/10.5194/nhess-18-2161-2018>, 2018.
- 620 Gariano, S. L. and Guzzetti, F.: Landslides in a changing climate, *Earth-Science Reviews*, 162, 227–252, <https://doi.org/10.1016/j.earscirev.2016.08.011>, 2016.
- Gariano, S. L., Rianna, G., Petrucci, O., and Guzzetti, F.: Assessing future changes in the occurrence of rainfall-induced landslides at a regional scale, *Science of The Total Environment*, 596–597, 417–426, <https://doi.org/10.1016/j.scitotenv.2017.03.103>, 2017.
- 625 Guo, Z., Ferrer, J. V., Hürlimann, M., Medina, V., Puig-Polo, C., Yin, K., and Huang, D.: Shallow landslide susceptibility assessment under future climate and land cover changes: A case study from southwest China, *Geoscience Frontiers*, 14, 101542, <https://doi.org/10.1016/j.gsf.2023.101542>, 2023.
- Gutiérrez, F., Linares, R., Roqué, C., Zarroca, M., Carbonel, D., Rosell, J., and Gutiérrez, M.: Large landslides associated with a diapiric fold in Canelles Reservoir (Spanish Pyrenees): Detailed geological–geomorphological mapping, trenching and electrical resistivity imaging, *Geomorphology*, 241, 224–242, <https://doi.org/10.1016/j.geomorph.2015.04.016>, 2015.
- 630 Hakim, W. L., Rezaie, F., Nur, A. S., Panahi, M., Khosravi, K., Lee, C.-W., and Lee, S.: Convolutional neural network (CNN) with metaheuristic optimization algorithms for landslide susceptibility mapping in Icheon, South Korea, *Journal of Environmental Management*, 305, 114367, <https://doi.org/10.1016/j.jenvman.2021.114367>, 2022.
- 635 Hartmann, J. and Moosdorf, N.: The new global lithological map database GLiM: A representation of rock properties at the Earth surface, *Geochemistry, Geophysics, Geosystems*, 13, <https://doi.org/10.1029/2012GC004370>, 2012.
- He, Q., Wang, M., and Liu, K.: Rapidly assessing earthquake-induced landslide susceptibility on a global scale using random forest, *Geomorphology*, 391, 107889, <https://doi.org/10.1016/j.geomorph.2021.107889>, 2021.
- Hong, H., Miao, Y., Liu, J., and Zhu, A.-X.: Exploring the effects of the design and quantity of absence data on the performance of random forest-based landslide susceptibility mapping, *CATENA*, 176, 45–64, <https://doi.org/10.1016/j.catena.2018.12.035>, 2019.
- 640 Huang, W., Ding, M., Li, Z., Yu, J., Ge, D., Liu, Q., and Yang, J.: Landslide susceptibility mapping and dynamic response along the Sichuan-Tibet transportation corridor using deep learning algorithms, *CATENA*, 222, 106866, <https://doi.org/10.1016/j.catena.2022.106866>, 2023.
- 645 Hürlimann, M., Guo, Z., Puig-Polo, C., and Medina, V.: Impacts of future climate and land cover changes on landslide susceptibility: regional scale modelling in the Val d’Aran region (Pyrenees, Spain), *Landslides*, 19, 99–118, <https://doi.org/10.1007/s10346-021-01775-6>, 2022.

- Jarvis, A., Reuter, H. I., Nelson, A., and Guevara, E.: Hole-filled SRTM for the globe Version 4, CGIAR-CSI SRTM 90m Database, <https://srtm.csi.cgiar.org/>, 2008.
- 650 Jin, J., Chen, G., Meng, X., Zhang, Y., Shi, W., Li, Y., Yang, Y., and Jiang, W.: Prediction of river damming susceptibility by landslides based on a logistic regression model and InSAR techniques: A case study of the Bailong River Basin, China, *Engineering Geology*, 299, 106562, <https://doi.org/10.1016/j.enggeo.2022.106562>, 2022.
- John, A., Douville, H., Ribes, A., and Yiou, P.: Quantifying CMIP6 model uncertainties in extreme precipitation projections, *Weather and Climate Extremes*, 36, 100435, <https://doi.org/10.1016/j.wace.2022.100435>, 2022.
- 655 Johnson, K., Villani, M., Bayliss, K., Brooks, C., Chandrasekhar, S., Chartier, T., Chen, Y.-S., Garcia-Pelaez, J., Gee, R., Styron, R., Rood, A., Simionato, M., and Pagani, M.: Global Seismic Hazard Map, <https://doi.org/10.5281/zenodo.8409647>, 2023.
- Kirschbaum, D., Kapnick, S. B., Stanley, T., and Pascale, S.: Changes in Extreme Precipitation and Landslides Over High Mountain Asia, *Geophysical Research Letters*, 47, e2019GL085347, <https://doi.org/10.1029/2019GL085347>, 2020.
- 660 Li, B., Liu, K., Wang, M., He, Q., Jiang, Z., Zhu, W., and Qiao, N.: Global Dynamic Rainfall-Induced Landslide Susceptibility Mapping Using Machine Learning, *Remote Sensing*, 14, 5795, <https://doi.org/10.3390/rs14225795>, 2022.
- Li, Y. and Duan, W.: Decoding vegetation’s role in landslide susceptibility mapping: An integrated review of techniques and future directions, *Biogeotechnics*, 2, 100056, <https://doi.org/10.1016/j.bgtech.2023.100056>, 2024.
- Li, C., Fu, Z., Wang, Y., Tang, H., Yan, J., Gong, W., Yao, W., and Criss, R. E.: Susceptibility of reservoir-induced landslides and strategies for increasing the slope stability in the Three Gorges Reservoir Area: Zigui Basin as an example, *Engineering Geology*, 261, 105279, <https://doi.org/10.1016/j.enggeo.2019.105279>, 2019.
- 665 Leshchinsky, B., Lehmann, P., and Or, D.: Enhanced Rainfall-Induced Shallow Landslide Activity Following Seismic Disturbance—From Triggering to Healing, *JGR Earth Surface*, 126, e2020JF005669, <https://doi.org/10.1029/2020JF005669>, 2021.
- 670 Lin, Q. and Wang, Y.: Spatial and temporal analysis of a fatal landslide inventory in China from 1950 to 2016, *Landslides*, 15, 2357–2372, <https://doi.org/10.1007/s10346-018-1037-6>, 2018.
- Lin, Q., Lima, P., Steger, S., Glade, T., Jiang, T., Zhang, J., Liu, T., and Wang, Y.: National-scale data-driven rainfall induced landslide susceptibility mapping for China by accounting for incomplete landslide data, *Geoscience Frontiers*, 12, 101248, <https://doi.org/10.1016/j.gsf.2021.101248>, 2021.
- 675 Lin, Q., Steger, S., Pittore, M., Zhang, J., Wang, L., Jiang, T., and Wang, Y.: Evaluation of potential changes in landslide susceptibility and landslide occurrence frequency in China under climate change, *Science of The Total Environment*, 850, 158049, <https://doi.org/10.1016/j.scitotenv.2022.158049>, 2022.
- Liu, C., Li, W., Wu, H., Lu, P., Sang, K., Sun, W., Chen, W., Hong, Y., and Li, R.: Susceptibility evaluation and mapping of China’s landslides based on multi-source data, *Nat Hazards*, 69, 1477–1495, <https://doi.org/10.1007/s11069-013-0759-y>, 2013.
- 680 Liu, X. and Miao, C.: Large-scale assessment of landslide hazard, vulnerability and risk in China, *Geomatics, Natural Hazards and Risk*, 9, 1037–1052, <https://doi.org/10.1080/19475705.2018.1502690>, 2018.

- M, H. and M.N, S.: A Review on Evaluation Metrics for Data Classification Evaluations, *IJDKP*, 5, 01–11, <https://doi.org/10.5121/ijdkp.2015.5201>, 2015.
- Merghadi, A., Yunus, A. P., Dou, J., Whiteley, J., ThaiPham, B., Bui, D. T., Avtar, R., and Abderrahmane, B.: Machine
685 learning methods for landslide susceptibility studies: A comparative overview of algorithm performance, *Earth-Science Reviews*, 207, 103225, <https://doi.org/10.1016/j.earscirev.2020.103225>, 2020.
- Margottini, C., Canuti, P., Sassa, K. (Eds.), 2013. *Landslide Science and Practice: Volume 1: Landslide Inventory and Susceptibility and Hazard Zoning*. Springer Berlin Heidelberg, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-31325-7>
- 690 Ministry of Natural Resources of the People’s Republic of China: Specification of Geological Hazard Risk Survey and Assessment (1:50,000), DZ/T 0438-2023, <https://std.cgs.gov.cn/content/7262410361846894592> (last access: 25 March 2026), 2023.
- O’Brien, R. M.: A Caution Regarding Rules of Thumb for Variance Inflation Factors, *Qual Quant*, 41, 673–690, <https://doi.org/10.1007/s11135-006-9018-6>, 2007.
- 695 Peng, S., Ding, Y., Wen, Z., Chen, Y., Cao, Y., and Ren, J.: Spatiotemporal change and trend analysis of potential evapotranspiration over the Loess Plateau of China during 2011–2100, *Agricultural and Forest Meteorology*, 233, 183–194, <https://doi.org/10.1016/j.agrformet.2016.11.129>, 2017.
- Peres, D. J. and Cancelliere, A.: Modeling impacts of climate change on return period of landslide triggering, *Journal of Hydrology*, 567, 420–434, <https://doi.org/10.1016/j.jhydrol.2018.10.036>, 2018.
- 700 Park, S.-J. and Lee, D.: Predicting susceptibility to landslides under climate change impacts in metropolitan areas of South Korea using machine learning, *Geomatics, Natural Hazards and Risk*, 12, 2462–2476, <https://doi.org/10.1080/19475705.2021.1963328>, 2021.
- Peng, S., Gang, C., Cao, Y., and Chen, Y.: Assessment of climate change trends over the Loess Plateau in China from 1901 to 2100, *International Journal of Climatology*, 38, 2250–2264, <https://doi.org/10.1002/joc.5331>, 2018.
- 705 Peng, S., Ding, Y., Liu, W., and Li, Z.: 1km monthly temperature and precipitation dataset for China from 1901 to 2017, *Earth System Science Data*, 11, 1931–1946, <https://doi.org/10.5194/essd-11-1931-2019>, 2019.
- Petschko, H., Brenning, A., Bell, R., Goetz, J., Glade, T., 2013. Assessing the quality of landslide susceptibility maps – case study Lower Austria. <https://doi.org/10.5194/nhessd-1-1001-2013>
- Reichenbach, P., Rossi, M., Malamud, B. D., Mihir, M., and Guzzetti, F.: A review of statistically-based landslide
710 susceptibility models, *Earth-Science Reviews*, 180, 60–91, <https://doi.org/10.1016/j.earscirev.2018.03.001>, 2018.
- Roy, J. and Saha, S.: Landslide susceptibility mapping using knowledge driven statistical models in Darjeeling District, West Bengal, India, *Geoenvironmental Disasters*, 6, 11, <https://doi.org/10.1186/s40677-019-0126-8>, 2019.
- Stumpf, A. and Kerle, N.: Object-oriented mapping of landslides using Random Forests, *Remote Sensing of Environment*, 115, 2564–2577, <https://doi.org/10.1016/j.rse.2011.05.013>, 2011.

- 715 Styron, R. and Pagani, M.: The GEM Global Active Faults Database, *Earthquake Spectra*, 36, 160–180, <https://doi.org/10.1177/8755293020944182>, 2020.
- Thrasher, B., Maurer, E. P., McKellar, C., and Duffy, P. B.: Technical Note: Bias correcting climate model simulated daily temperature extremes with quantile mapping, *Hydrology and Earth System Sciences*, 16, 3309–3314, <https://doi.org/10.5194/hess-16-3309-2012>, 2012.
- 720 Thrasher, B., Wang, W., Michaelis, A., Melton, F., Lee, T., and Nemani, R.: NASA Global Daily Downscaled Projections, CMIP6, *Sci Data*, 9, 262, <https://doi.org/10.1038/s41597-022-01393-4>, 2022.
- Viet Du, Q. V., Nguyen, H. D., Pham, V. T., Nguyen, C. H., Nguyen, Q.-H., Bui, Q.-T., Doan, T. T., Tran, A. T., and Petrisor, A.-I.: Deep learning to assess the effects of land use/land cover and climate change on landslide susceptibility in the Tra Khuc river basin of Vietnam, *Geocarto International*, 38, 2172218, <https://doi.org/10.1080/10106049.2023.2172218>, 2023.
- 725 Wang, N., Lombardo, L., Gariano, S. L., Cheng, W., Liu, C., Xiong, J., and Wang, R.: Using satellite rainfall products to assess the triggering conditions for hydro-morphological processes in different geomorphological settings in China, *International Journal of Applied Earth Observation and Geoinformation*, 102, 102350, <https://doi.org/10.1016/j.jag.2021.102350>, 2021.
- Yang, J. and Huang, X.: The 30 m annual land cover datasets and its dynamics in China from 1985 to 2022 (1.0.2), <https://doi.org/10.5281/zenodo.8176941>, 2023.
- 730 Zhang, J., Ma, X., Zhang, J., Sun, D., Zhou, X., Mi, C., and Wen, H.: Insights into geospatial heterogeneity of landslide susceptibility based on the SHAP-XGBoost model, *Journal of Environmental Management*, 332, 117357, <https://doi.org/10.1016/j.jenvman.2023.117357>, 2023.
- Zhao, G., Pang, B., Xu, Z., Yue, J., and Tu, T.: Mapping flood susceptibility in mountainous areas on a national scale in China, *Science of The Total Environment*, 615, 1133–1142, <https://doi.org/10.1016/j.scitotenv.2017.10.037>, 2018.

735