

# Evaluating simulations of ship tracks in a **high-resolution** km-scale model

Anna Tippett<sup>1</sup>, Paul R. Field<sup>2,3</sup>, and Edward Gryspeerdt<sup>1</sup>

<sup>1</sup>Department of Physics, Imperial College London, London, UK

<sup>2</sup>Met Office, Exeter, UK

<sup>3</sup>School of Earth and Environment, University of Leeds, Leeds, UK

**Correspondence:** Anna Tippett (a.tippett22@imperial.ac.uk)

**Abstract.** Clouds, and in particular their interactions with aerosols, remain a major source of uncertainty in climate projections, due to the wide range of scales over which cloud processes act on. This uncertainty limits our capability to simulate potential solar radiation management strategies, such as marine cloud brightening (MCB). A **good-naturalanalogue-“natural”, or “opportunistic”, experiment** for investigating MCB is analysis of ship tracks, as they mimic the intended effect and allow us to investigate time evolving aerosol perturbations. In this study, we model a real case of ship tracks, and evaluate model performance through comparisons with satellite observations. **We evaluate our model simulations against three criteria, in order** Requiring accuracy in individual cases creates a particularly challenging test for simulations of aerosol-cloud interactions, but it is necessary to ascertain whether this model is suitable for simulating MCB accurately.

Our findings highlight a key deficiency in activation parameterisations when simulating high aerosol concentrations - such as those expected in MCB scenarios. While the model can replicate the mean cloud properties within ship tracks, it struggles to capture their temporal evolution. Specifically, in precipitating clouds, enhancements in droplet number concentration ( $N_d$ ) and liquid water path (LWP) are overestimated and persist too long. This discrepancy between model and observations is linked to excessive model sensitivity to aerosol loading in precipitating conditions, leading to unrealistically **easy-strong** suppression of drizzle, and ultimately resulting in simulated ship tracks which overestimate the cooling effect in these cases. We identify scenarios in which current formulations of parameterisations are not suitable for use in simulating high-concentration aerosol perturbations, such as MCB, and scenarios in which models are more capable.

## 1 Introduction

A large portion of the uncertainty in estimates of the effective radiative forcing from aerosol-cloud interactions originates from poor model representation of clouds (Smith et al., 2020). Clouds depend on processes operating at micron scales, rendering their explicit representation unfeasible in most computationally viable resolutions of weather or climate models. Consequently, cloud processes must be parametrised. These parameterisations define the process rates, which describe how much physical properties change between each model time step. Uncertainty in these parameterisations leads to considerable uncertainty in model estimates of the radiative forcing from aerosol-cloud interactions, with a large portion of this uncertainty stemming from the amount of liquid cloud (Zelinka et al., 2014). Addressing and reducing this uncertainty is crucial for enhancing the

25 accuracy of future climate projections (Andreae et al., 2005). Better representation of clouds in regional weather forecasting  
models would improve predictability of clouds and the onset of precipitation (Field et al., 2023). This uncertainty is also  
important for marine cloud brightening (MCB), a proposed solar radiation ~~management modification~~ (SRM) strategy aimed  
at reflecting more sunlight by intentionally seeding clouds with aerosols (Diamond et al., 2022; Wood, 2021; Salter et al.,  
2008; Zhang and Feingold, 2023). Uncertainty in model representation of high-concentration aerosol-cloud processes limits  
30 our capability to simulate MCB experiments, and makes the design of suitable field trials difficult.

To assess the longer term impacts of MCB on our climate, it must be incorporated into climate model simulations. One  
suggested approach involves the use of regional models to accurately simulate the localised effects of MCB, and use this  
to parametrise its impact in coarser resolution global climate models (Feingold et al., 2024). ~~While This is a particularly~~  
challenging task for climate models. While recent models are able to adequately simulate cloud patterns of cloudiness (Tselioudis et al., 202  
35 , their simulation of aerosol-cloud interactions is more varied (Gryspeerd et al., 2020). However, these evaluations are typically  
done on large-scale ~~turbulence is resolved in these models,~~ temporal or spatial averages - assessments of MCB efficacy  
require accurate simulations of the model response to individual clouds. This is not well evaluated by current techniques.  
In addition, as many of the same parameterisations ~~are used in GCMs (e.g., for activation or for cloud microphysics are used~~  
across both coarse and finer resolution models (e.g. activation and precipitation processes). ~~Therefore, it is critical that we~~  
40 have certainty, greater confidence in our regional ~~representations of MCB representation of MCB should inform our global~~  
modelling capability, avoiding the equifinality problem in larger-scale studies (Mülmenstädt and Feingold, 2018). Robust eval-  
uation frameworks are needed to assess the realism of regional models, not only in terms of short-wave (SW) radiative forcing  
but also with respect to the representation of key processes.

Ship tracks, the narrow cloud features formed by aerosol emissions from ships, ~~and~~ offer a valuable ~~natural experiment~~  
45 "natural experiment" for studying the interactions that would occur in intentional MCB (Christensen et al., 2022; Diamond  
et al., 2022). Aerosols in the ship plume act as cloud condensation nuclei, causing the cloud to have more, smaller, droplets  
which reflect more incoming solar radiation. This is known as the "Twomey" effect, and is well documented on short timescales  
(Twomey, 1974, 1977; Ferek et al., 1998; Hobbs et al., 2000; Ackerman et al., 2000; Feingold et al., 2003; Penner et al., 2004;  
Segrin et al., 2007; Christensen et al., 2022). The brightening observed in ship tracks mimics the intended effect of MCB,  
50 offering a real-world analogue for testing the feasibility of such interventions. Ship track studies provide information about  
aerosol-cloud interactions in this marine context, and can be useful ~~they can be~~ in assessing the behaviour of models that are  
intended for use in simulating MCB.

After the initial droplet number ( $N_d$ ) perturbation, clouds respond to aerosol perturbations over longer timescales through  
effects known as 'cloud adjustments'. ~~Changes in liquid water path (LWP) - the amount of liquid in a cloud column - are~~  
55 particularly important because they influence not only cloud radiative properties but also precipitation processes. "cloud  
adjustments". A precipitation suppression effect (Albrecht, 1989; Rosenfeld, 2000), where smaller cloud droplets take longer to  
form precipitation (~~increasing,~~ increasing liquid water path (LWP), tends to result in additional cooling. Alternatively, smaller  
droplets can also enhance the mixing of dry air above cloud-top into the cloud (a process known as entrainment) and reduce  
LWP, resulting in a warming effect (Ackerman et al., 2004; Bretherton et al., 2007). This bidirectional response in the LWP

60 can depend on the initial conditions of the unperturbed cloud (Han et al., 2002; Ackerman et al., 2004; Michibata et al., 2016; Toll et al., 2017, 2019; Gryspeerdt et al., 2019a; Possner et al., 2020; Glassmeier et al., 2021; Zhang et al., 2022; Fons et al., 2023), whilst also being controlled by covarying meteorology (Goren et al., 2025).

There is considerable uncertainty in both the sign and magnitude of adjustments (Glassmeier et al., 2021), which is relevant for MCB purposes, since adjustments involving decreasing the LWP could offset the intended cooling impact. Part of this uncertainty stems from both the difficulty in making observations of aerosol-cloud processes, and isolating the impact of aerosol from background meteorology (Christensen et al., 2022). Cloud adjustments are inherently time-dependent processes (Gryspeerdt et al., 2021), yet many observational studies do not consider the time evolution of ~~an a~~ cloud response, making it difficult to observe the processes occurring. This limits our modelling capability of these interactions, since without time-resolved observations, it is difficult to capture the time dependence for instantaneous/short timescale injections (such as ~~from~~ ship aerosols or MCB).

Simulating ship tracks provides an opportunity to simulate the intended effects of MCB, and evaluate a model's ability to represent key processes realistically. By simulating ship tracks we can disentangle aerosol effects from meteorological variability because the aerosol source is known and relatively localised. This is possible since we can use the region neighbouring the ship track as an unperturbed 'control' region (Segrin et al., 2007), which is a ~~best estimate proxy~~ for what the cloud would have looked like if no ship was there. ~~This makes them~~ Ship tracks are therefore ideal for isolating the causal impact of aerosols on cloud properties and separate meteorological co-variations (Goren et al., 2025). Additionally, we can view ship tracks as linear formations of independently perturbed clouds (Kabatás et al., 2013), thereby allowing us to infer information about the time evolution of the aerosol perturbation (Gryspeerdt et al., 2021). This helps us evaluate model process representation, such as the activation of cloud droplets (which occurs on ~~short timescales~~ the order of seconds to minutes; Arabas and Shima, 2016), or the autoconversion of cloud droplets into rain droplets (occurring on longer timescales) ~~;~~ on the order of minutes to hours; Stephens and Haynes, 2007.

In order to answer ~~logistical~~ questions relating to MCB experimental design, accurate simulations of ship tracks are vital, however we must be certain that model simulations produce the correct answer for the right reasons. This calls for in depth evaluation of the representation of model processes, not just the final forcings ~~-Ship-~~ ship tracks provide an avenue to do this. Simulating MCB provides an exceptionally challenging task for the field, since accuracy is not only required in terms of large-scale / domain-wide averages, but instead at the scale of individual cloud perturbations.

Previous efforts have been made to simulate ship tracks in atmospheric models, in order to assess model performance (~~Wang and Feingold, 2009; Possner et al., 2015, 2018; Berner et al., 2015~~) (Wang and Feingold, 2009; Possner et al., 2015, 2018; Berner et al. (2015) and Wang and Feingold (2009) utilise fine resolution LES models to investigate the effect of aerosol perturbations on marine stratocumulus, but cannot make direct comparisons to observations since they are not real cases of ship tracks. Similarly, McMichael et al. (2024, 2025) investigate ship track spreading rates with LES, and Prabhakaran et al. (2024) use LES to investigate impacts on the stratocumulus-to-cumulus transition. Possner et al. (2015) use observations of a real case of ship tracks, yet the simulated ship locations are ~~prescribed and~~ not from the actual ships that caused the observed ship tracks, which limits the ability to assess the accuracy of MCB simulations.

95 In this work, we produce simulations of a real case of ship tracks, using ship emissions locations from ship Automatic Identification System (AIS) data. We utilise a double-moment cloud microphysics scheme - Cloud AeroSol Interacting Microphysics (CASIM; Field et al., 2023) in the Met Office Unified Model (UM) in regional configuration (allowing for convection to be resolved), and compare directly to satellite observations. We ~~infer-simulate~~ changes in cloud properties ~~and the processes involved from changes in~~ ( $N_d$  and LWP, ~~and the~~), investigating the processes involved as well as the time evolution of ~~these~~ ~~responses~~the response. We evaluate our simulations according to the three criteria described in Section 2.1, in order to ensure the proper change in cloud radiative effect is captured via the correct microphysical processes. In order to properly simulate MCB impacts we must simulate the correct behaviour, and for the correct reasons.


Through evaluating our simulated ship tracks against these criteria, we reveal issues in current model parameterisations when modelling high-concentration aerosol perturbations, such as those that are expected in MCB. We identify scenarios in which current formulations are more suitable for simulating MCB, and scenarios in which further work is needed before models can be considered credible. Furthermore, we make suggestions on experimental design, such that analysis and quantification of the aerosol effect is least uncertain.

## 2 Methods

### 2.1 Evaluating Criteria for evaluating ship tracks

110 In order to evaluate the model representation of aerosol perturbation in individual ship tracks, rather than a larger-scale temporal or spatially aggregated assessment, we define the following criteria.

1. ***Accurate representation of changes in CRE.*** The cloud radiative effect (CRE) in overcast scenes is largely dependent on the inside ship track  $N_d$  and LWP values (Possner et al., 2015), therefore we want to ensure our model is producing simulations with the correct numerical values inside the ship track. Due to difficulties in obtaining perfect background simulations, our first criteria considers solely the absolute values where the aerosol perturbation has been applied, since this is ultimately where it is key to get the behaviour correct.
2. ***Accurate time evolution of the CRE.*** This criteria enables us to evaluate the lifetime of the aerosol effect and therefore the associated radiative forcing. Again, this effect is driven by changes in LWP and  $N_d$ . The model sensitivity to aerosol can be evaluated, as the enhancement from the unperturbed (control) state is taken into account. Ship tracks provide a method for obtaining time evolution even in snapshot satellite imagery (Gryspeerd et al., 2021), therefore information about the time-evolution of the aerosol effect can be made.
3. ***Sensitivity in the  $N_d$  response to initial conditions.*** Previous studies have shown that the cloud response is sensitive to the initial conditions of the cloud (Ackerman et al., 2004; Michibata et al., 2016; Toll et al., 2017, 2019; Gryspeerd et al., 2019a; Possner et al., 2020; Glassmeier et al., 2021; Zhang et al., 2022; Fons et al., 2023). In order for our model representation of clouds to be accurate this sensitivity must be reflected in the model. We expect to see different responses



**Figure 1.** [Granules from Aqua and Terra overpasses on 12th July 2018 that intersect with the model domain of this study.](#)

in precipitating and non-precipitating conditions, and this is investigated in our model simulations. In order for the model to reproduce the correct response for the right reasons, this dependence on the initial cloud conditions should be satisfied. This should confirm that the model is sensitive to the correct model processes.

130 [We apply the above criteria by considering temporal evolutions \(treating the length of a ship track as a time since aerosol injection axis\) of inside track  \$N\_d\$  and LWP, and the associated enhancements from the background \(the calculation of which is explained in Section 2.6\). We discuss the “correctness” with respect to each criterion in terms of the similarities and differences between the observed responses compared to the simulated responses.](#)

## 2.2 Model configuration

Our case study is simulated using the Met Office Unified Model (UM, version 13.0; Brown et al., 2012), in the Regional Atmosphere and Land (RAL) configuration 3.1 (Bush et al., 2020, 2023, 2025). [Our domain covers the Eastern and Central Pacific, off the coast of California \(Fig. 1\).](#) The 2625 km by 1125 km domain is centred on (40°N, 135.25°W) with a horizontal resolution of 1.5 km. There are 90 levels up to 40 km, with 16 levels below 1 km. The model time step is 75 s, and the lateral boundary conditions to the nested regional model are provided hourly by the UM global model at N216 resolution ( $\approx 60$  km) in Global Atmosphere (GA) 6.1 science configuration (Walters et al., 2017). Our simulations are initialised from global model analysis at 0000Z on the 11th July 2018 and run for 48 hours.

145 To properly simulate the aerosol-cloud interactions in a cloud resolving model, a double moment cloud microphysics scheme is recommended (Morrison et al., 2009; Igel et al., 2015). Double-moment microphysics schemes prognose both the mass and number concentration of each hydrometeor species (e.g. [Ferrier, 1994; Seifert and Beheng, 2006; Morrison and Gettelman, 2008; Taufour and Ferrier, 1994; Seifert and Beheng, 2006](#)). A double-moment microphysics scheme allows  $N_d$  to be recalculated at each time step, and therefore can subsequently modify process rates (such as autoconversion), impacting the cloud evolution over time (Field et al., 2023). In single moment schemes, the number concentration of each species (from which the process rates are derived) are either constant or diagnosed from other meteorological parameters using empirical relationships. These schemes,

whilst computationally cheaper, can fail to accurately represent the indirect effect from aerosol-cloud interactions (Gordon et al., 2020).

150 For cloud microphysics, we employ the Cloud AeroSol Interacting Microphysics (CASIM) scheme (Shipway and Hill, 2012; Field et al., 2023). CASIM is a double moment scheme with five hydrometeor species (cloud liquid, rain, ice, snow, and graupel) in which both mass and number concentration mixing ratios are prognosed.

~~Our configuration of CASIM allows to be derived from aerosol (rather than prescribed~~Each cloud species' particle size distribution (PSD) is described using a generalised gamma function with constant shape parameters. The autoconversion of cloud droplets into rain droplets and accretion of cloud water by rain are parametrised in CASIM following Khairoutdinov and Kogan (2000)  
155 . CASIM also contains other warm cloud processes, such as aggregation (rain collecting rain), and we~~couple it to the~~refer the reader to Field et al. (2023) for details on these parametrised processes.

We couple CASIM to the double moment modal Global Model of Aerosol Processes aerosol microphysics scheme (GLOMAP-mode; Mann et al., 2010), within United Kingdom Chemistry and Aerosols (UKCA)~~aerosol microphysics scheme. UKCA is a full prognostic two moment aerosol microphysics scheme, based on GLOMAP (Mann et al., 2010), and provides both aerosol sub-model. Coupling to GLOMAP-mode in CASIM allows for aerosol mass and number~~information for concentration to influence the activation of cloud droplets~~, which is parametrised by Abdul-Razzak and Ghan (ARG; 2000).~~

To simplify our simulations, we only consider aerosol ~~only in the accumulation mode, in the soluble accumulation mode~~ and initialise our simulation with a constant ~~sulfate sulphate~~ aerosol number concentration of  $200 \text{ cm}^{-3}$ , characteristic mass  
165  $3 \times 10^{-18} \text{ kg}$ , and aerosol diameter of  $\approx 0.1 \mu\text{m}$ , which is typical of sulphate aerosols (Noone et al., 2000). ~~The aerosol activation scheme used is that of Abdul-Razzak and Ghan (ARG; 2000).~~

### 2.3 Control run

~~Left column: control simulations of 12th July 2018 at 20:00 without ship emissions turned on. Panel (b) contains the locations of the ships simulated during this study, up until the time step of this figure. Right column: the same simulation with shipping emissions on. Variables shown are aerosol number concentration ( $\cdot$ ), cloud droplet number concentration ( $\cdot$ ), liquid water path (LWP), surface rain rate (RR), and top of atmosphere outgoing short wave radiation (SW).~~  
170

~~Using the model configuration detailed above, our simulation is run for 48 hours from 0000Z 11th July to 0000Z 13th July. In our control run, no ship emissions are added to the model, and we use this simulation to characterise our “unpolluted” cloud. Typically, in ship track observations, we have no pure “control” cloud, and therefore we must treat a region outside of a ship track as being representative of what the cloud would have looked like if there was no aerosol perturbation present. In the model case, however, we can simply turn ships on and off to isolate the aerosol impact on the cloud~~Aerosol is allowed to evolve freely after the model simulation begins, with the boundaries determined by the global driving model. As described in Gordon et al., 2020, aerosol concentrations can vary as determined by the microphysical processes in GLOMAP-mode, however CASIM does not have the capability to track aerosol tracers within hydrometeors and therefore cloud microphysics  
175 does not alter aerosol mass/size/composition except through irreversible removal via wet-scavenging.  
180

The aerosol number concentration ( $N$ ), cloud droplet number concentration ( $N_c$ ), liquid water path (LWP), surface rain rate (RR), and top of atmosphere outgoing short wave radiation (SW) for the This assumption that we only have aerosol in the soluble accumulation mode enables us to consider differences between control run 44 hours after initialisation can be found in the left hand column of Fig. 4. Fig. 4a demonstrates how the coupled UKCA aerosol scheme allows depletion of aerosol in locations where there is precipitation and ships runs as solely a function of increasing aerosol concentration at the ship location, and the land sources of aerosol are the most significant over the course of the simulation. The coupled UM-CASIM configuration is able to reproduce open effects of aerosol chemistry are neglected. Differences in hygroscopicities in other aerosol modes are assumed to be a less important factor in droplet activation than the mass/closed cellular convection in this marine stratocumulus, with a drizzling scene (surface rain rates on average of roughly 0.5 mm/hr). Overpasses from the CloudSat Cloud Profiling Radar (Stephens et al., 2008; Tanelli et al., 2008) verify this, with observations of pockets of precipitating clouds (with maximum surface rain rate of roughly 2 mm/hr) throughout the simulation run time (see Fig. S??).

With our initialised background aerosol field of 200, we produce a relatively clean cloud droplet background of roughly 50. This is largely in line with that seen in satellite observations (Fig. 7a, c), however we note that two large sources of cloud droplets are missing from our model simulation. This is likely due to our initialisation from a constant aerosol field, and therefore we are missing these sources of aerosol in our simulation diameter of the mode (Dusek et al., 2006). Non-ship aerosol sources are described in Mann et al. (2010), with emissions from marine phytoplankton, SO<sub>2</sub> from volcanoes, fires, and industrial sources. These sources can exist across different aerosol modes (not just the accumulation mode).

### 2.3 Simulating ship tracks

In order to simulate the shipping impacts on clouds, we add five ~~container~~ ships to our simulation in our *ships* run. We use container ship locations identified in AIS data and matched to observed shiptracks from 11th-12th July 2018, to add realistic moving sources of aerosol (Smith et al., 2015). Details of ship start/end positions, ~~velocities, and ship type and velocities~~ can be found in Tab. 1. ~~These ships were selected because they produced some of the most visible ship tracks within this domain, and covered~~ Container ships are selected because of their large size and high emissions, meaning that the resulting ship tracks are the most distinct in the observations. Our five selected ships also cover both directions of travel across the domain.

the shipping route within the domain. Aerosol is added in a model level at 10 m above sea level and immediately dispersed throughout the 1.5 km by 1.5 km grid box using the following equation:

$$N_{x,y,z,t+1} = N_{x,y,z,t} + \frac{dN}{dt} \times \frac{\Delta t}{V} \quad (1)$$

where  $\frac{dN}{dt}$  is the rate of production number (the rate of ship aerosol emissions) in s<sup>-1</sup>,  $\Delta t$  is the model time step (75s), and  $V$  is the grid box volume. This describes the aerosol concentration added to the entire grid box per time step. At this model resolution, ship tracks form at a minimum width of 1.5km. Typical ship tracks have a width of on the order of 10km (Durkee et al., 2000), therefore this model resolution ~~should be is~~ suitable to capture the horizontal extent. Aerosol is almost immediately transported up to cloud base, and there is little deposition of aerosol at the bottom model level, therefore the 10m injection height (whilst an underestimation) is approximately correct.

Ship Label	Type	$lat_i$ (deg)	$lon_i$ (deg)	$lat_f$ (deg)	$lon_f$ (deg)	$v$ (knots)	$\frac{dN}{dt}$ ( $s^{-1}$ )
A	Container	38.768	234.396	44.762	219.362	19.895	$5 \times 10^{14}$
B	Container	37.798	237.677	41.639	222.035	20.231	$5 \times 10^{14}$
C	Container	37.796	237.686	44.738	222.516	20.179	$5 \times 10^{14}$
D	Container	43.156	212.417	38.228	224.687	16.427	$5 \times 10^{14}$
E	Container	43.337	210.858	36.580	227.716	22.534	$5 \times 10^{14}$

**Table 1.** Start and end locations of 5 container ships, with associated velocities, emission rates. Ships are initialised at 00:00Z 11th July 2018, and allowed to travel for the 48 hours of the simulation.

Realistic emissions of condensation nuclei (CN) from ships fall in the range  $10^{16}$  to  $10^{18}$  kg/s (Taylor and Ackerman, 1999; Hobbs et al., 2000; Berner et al., 2015). ~~In this work we use Connolly et al. (2014) find that the ARG activation parameterisation (Abdul-Razzak and Ghan, 2000) exhibits a sharp “drop off” in activated fraction at increasing aerosol concentrations (albeit, for NaCl), therefore we investigate the use of a high ship aerosol case (ship production number  $10^{16}$ ) and a lower ship aerosol case (ship production number  $5 \times 10^{14}$  due to issues with activation schemes at very high aerosol concentrations. This approximately matches the perturbation seen in the ship tracks (see-) in this study~~ (Section 3.1). We add the aerosol into the accumulation soluble mode with a mass  $3 \times 10^{-18}$  kg, the same as background aerosol, ~~so that all aerosols within the simulation are consistent~~ meaning that both our background and ship aerosols are only in the accumulation mode with the same mass. This is done to simplify simulations and isolate causal aerosol effects in our analysis, ~~since differences between the ships and control run can be attributed to the increased aerosol concentration at the ship locations.~~

## 2.4 Satellite data

To ~~constrain-evaluate~~ our model simulations, we use observations from NASA’s Aqua and Terra satellites to compare to our model output. We use Level-2 Collection 6.1 data from the Moderate Resolution Imaging Spectroradiometer (MODIS; Platnick et al., 2017) to obtain LWP and derive  $N_d$  following Quaas et al. (2006).

We collocate Aqua and Terra overpasses the occur during our simulation with our model domain, and only consider images the contain any part of a ship track, obtaining ~~6-six~~ snapshots of our domain, as shown in Fig. 1. Model data is output hourly, therefore we compare satellite data to model output from the nearest time step. When model data is compared to satellite data, the model output is masked only to consider the region contained within the associated satellite snapshot.

~~Observations of Simulated~~ precipitation in the domain ~~are confirmed with overpassed-is~~ evaluated against ERA5 surface precipitation (Hersbach et al., 2020), as well as Global Precipitation Measurement (GPM-IMERG; Huffman et al., 2015) and overpasses from the Cloud Profiling Radar (CPR) onboard CloudSat (Stephens et al., 2008) during the simulation run time (both the 2C-Precip-Column and 2C-Rain-Profile products; Stephens et al., 2008; Haynes et al., 2009; L’Ecuyer and Stephens, 2002). We investigate the probability of precipitation across  $N_d$ -LWP space using observations from the CCCM (CERES-CloudSat-CALIPSO-

combined product (~~Kato et al., 2010~~)(Kato et al., 2010, 2011). We use the CloudSat ~~precipitation-flag~~ CPR precipitation flag (from 2C-Precip-Column; Haynes et al., 2009) to define a probability of precipitation as counts of liquid precipitation or drizzle divided by all counts, per  $N_d$ -LWP bin (from MODIS Aqua). Since in 2018 CloudSat was not part of the A-train (and therefore not collocated with MODIS), we use observations from 2007-2011, as in Gryspeerdt et al. (2022).

~~Granules from Aqua and Terra overpasses on 12th July 2018 that intersect with the model domain of this study.~~

## 2.5 Ship track locations

We follow a methodology similar to that of Tippet et al. (2024); Gryspeerdt et al. (2021); Manshausen et al. (2022) to obtain our ship track locations. ~~Ship~~ Using ship positions (from AIS data) we infer the locations of clouds which experienced an aerosol perturbation (neglecting the time taken for aerosol to reach cloud base). These positions are advected in wind fields over time to ~~obtain a ship track which consists~~ predict ship tracks which consist of not only location, but the associated time since that location experienced the aerosol perturbation. This allows the length of a ship track to be treated as a time axis, and discern time evolution after an aerosol perturbation even in snapshot imagery (Kabatas et al., 2013).

In our observations of ship tracks in this case study, the hourly ship locations are advected in ERA5 reanalysis winds at 0.25° resolution and 3 hour intervals (Hersbach et al., 2020) between the surface and the boundary layer top, with vertical motion within the ship plume following Briggs (1965) (see Tippet et al., 2024 for further details). In model output, the same ship locations are advected instead in the model winds following the same methodology.

Since this study is comparing only ~~5~~ five ship tracks between model output and satellite data, it is essential that the exact ship track location is logged as to properly quantify the aerosol effects. We apply small corrections to our model and satellite output track locations from the above methodology to ensure that our track location falls exactly on each visible ship track, for both model output and observations. This is done by hand logging the nearest point of the visible track, perpendicular to the predicted track location. This is not possible in large composite studies which contain thousands of ship tracks, however due to the small number of ships considered in this study it is feasible, and ensures the most accurate measurement of the cloud responses.

Following Tippet et al. (2024), we regrid our datasets to 2D space for each ship track, at each model time step / MODIS snapshot. This 2D space consists of the “time along track” (binned to 1 hourly windows), and the perpendicular “distance away from track” (binned to 1 km). This allows us to investigate exactly how the cloud properties vary with time since aerosol perturbation and distance from the centre of the perturbation.

## 2.6 Quantifying enhancements

A key challenge in this study is determining the most appropriate method for quantifying the impact of ship emissions on clouds across our two datasets: model simulations and satellite observations.

For the UM-CASIM simulations, the impact of ship-emitted aerosols can be directly quantified by comparing the enhancement in cloud properties between the *ships* run and the *control* run. To mitigate stochastic noise between model runs, we apply Gaussian smoothing to the model fields (smoothing with a Gaussian kernel with standard deviation of 0.75 km) before

270 calculating the percentage difference to obtain enhancements in  $N_d$  and LWP ( $\epsilon_N$  and  $\epsilon_L$ ). This approach allows for a precise estimation of aerosol-induced changes while minimising the influence of small-scale variability in between model runs.

In contrast, defining a control region in observational data is less straightforward (Christensen et al., 2022). Traditional observational studies of ship tracks typically identify an “unpolluted” reference region located approximately 30 km perpendicular to the ship track (Manshausen et al., 2022). This region serves as a control from which the percentage enhancement in cloud  
275 properties is calculated. However, this methodology presents potential biases. Specifically, non-linear background gradients can introduce systematic errors in the estimated aerosol effect (Tippett et al., 2024). Additionally, in scenes containing multiple ship tracks, this approach becomes increasingly uncertain. For instance, in our case study, many “outside track” regions that are used as controls inadvertently overlap with other ship tracks. Consequently, the percentage enhancement derived from such comparisons likely underestimates the true aerosol impact (Yuan et al., 2025).

280 Despite these limitations, this method remains the most viable approach given the constraints of observational data.

In Fig. ~~S?? we demonstrate~~ 8j.1 we compare the use of (a) the control simulation and (b) the “outside” region to define the unperturbed cloud for the model simulation (solid orange and dashed orange lines, respectively). For the model, we find that both methods for calculating model enhancements produce the same results (to within -15% and -1015% and 10% for  $N_d$  and LWP enhancements in the first 15 hours of ship tracks, respectively). This is due to the model background being relatively clean, and therefore is a good representation of the control cloud since there is not much Our model background has little background  
285 variability, and any variability is much smaller than the large perturbation inside the ship track, therefore both methods provide similar results. This would not necessarily be the case in a model simulation with greater background variability. In this study, we choose to employ different methodologies for the model and observational datasets when calculating percentage enhancements, since in this clean case it does not impact the results.

## 290 2.7 Expected enhancements


Using equation (1) from Gryspeerd et al. (2019b), we can calculate expected enhancements in cloud droplet number concentration ( $\epsilon_N$ ) based on the background  $N_d$  as a function of aerosol mass emission rates

$$\epsilon_N = \frac{M_E^\gamma}{\alpha + \beta N_{cln}} + 1 \quad (2)$$

where  $M_E$  is the mass emission rate (in  $\text{kg s}^{-1}$ ),  $N_{cln}$  is the unperturbed background  $N_d$ , and the constants  $\alpha$ ,  $\beta$ , and  $\gamma$   
295 are 0.1041, 0.0038 and 0.36 as obtained from the fit in Gryspeerd et al. (2019b). This equation provides a useful method of evaluating whether or not our model is producing enhancements of the right order of magnitude, based on its background conditions.

### 3 Results

#### 3.1 Activation of aerosol




**Figure 2.** Example of how realistic ship emissions ( $10^{16}$ ) produce a ‘split’ ship track due to non-monotonic activation parameterisation from [Abdul-Razzak et al. \(1998\)](#) [Abdul-Razzak and Ghan \(2000\)](#).

300 Using Equation (1), and a realistic ship production number of  $10^{16} \text{ s}^{-1}$ , we obtain [an aerosol perturbation of aerosol number concentrations within the ship track](#) roughly  $20,000 \text{ cm}^{-3}$ . This is realistic given in-situ observations of ship tracks (Hobbs et al., 2000; Noone et al., 2000). However, with these values, our simulated ship tracks exhibit bizarre ‘split’ behaviour, with no cloud droplets activated inside the centre of the track where the aerosol concentrations are highest (see Fig. 2).

This unphysical split-shiptrack effect is due to non-monotonic behaviour of the Abdul-Razzak and Ghan (ARG) activation parameterisation scheme with increasing aerosol concentrations, beyond some critical aerosol concentration (dependent on temperature, pressure and updraft velocity) (Abdul-Razzak and Ghan, 2000). To demonstrate this, we utilise *pyrcel* (Rothenberg and Wang, 2016), a python package for implementing [a simple adiabatic cloud parcel model](#), to isolate [solely](#) the impact of the activation parameterisation.

We plot both the ARG and NS (Nenes and Seinfeld, 2003) parameterisations as a function of aerosol concentration in Fig. 310 3a, using values of temperature, pressure and updraft velocity that are representative of our domain. We find that with using our realistic ship production numbers ( $10^{16} \text{ s}^{-1}$ ) the aerosol concentrations ( $20,000 \text{ cm}^{-3}$ ) are beyond the critical concentration of ARG, producing the unrealistic split ship tracks. NS, whilst not containing this non-monotonic behaviour, would contain a large over estimation of cloud droplets at the realistic ship production number [and therefore would also be unsuitable with these emissions](#). This detail of the ARG activation parameterisation has [be-been](#) documented in Connolly et al. (2014), and 315 emphasises that commonly used aerosol activation parameterisations are not suitable for MCB applications.

Reduction of our ship production number to  $5 \times 10^{14} \text{ s}^{-1}$ , produces only a  $200 \text{ cm}^{-3}$  aerosol perturbation, but recovers  $N_d$  that match much closer to observations. This artificial reduction in ship production number is necessary to obtain more realistic ship tracks (without the splitting effect), but we must verify whether or not the resultant enhancements in these ship tracks is



**Figure 3.** (a) Number of aerosol particles ( $N_a$ ) activated to cloud droplets ( $N_d$ ) in different activation parameterisations calculated in a simple adiabatic cloud parcel model. Confidence intervals given by the range of temperature, pressure and updrafts simulated in our case study. (b) The expected enhancement in cloud droplets ( $\epsilon_N$ ) as a function of emissions aerosol mass emission rate ( $M_E$ ), based on equation (1) from Gryspeerd et al. (2019b). Vertical lines are the emissions from different ship production numbers. Horizontal lines are the enhancements produced from our simulations for each individual ship. Confidence interval is given by the range of background droplet number concentrations in our simulation.

as expected based on observations. Using equation (1) from Gryspeerd et al. (2019b), we calculate expected enhancements  
 320 ( $\epsilon_N$ ) based on the background  $N_d$  as a function of aerosol mass emission rates

$$\epsilon_N = \frac{M_E^\gamma}{\alpha + \beta N_{cln}} + 1$$

where  $M_E$  is the mass emission rate (in  $\text{kg}$ ),  $N_{cln}$  is the unperturbed background, and the constants  $\alpha$ ,  $\beta$ , and  $\gamma$  are 0.1041,  
 0.0038 and 0.36 as obtained from the fit in Gryspeerd et al. (2019b). Using our ship production of numbers of  $10^{16} \text{ s}^{-1}$  and  $5$   
 325  $\times 10^{14} \text{ s}^{-1}$ , and a mean aerosol mass of  $3 \times 10^{-18} \text{ kg}$ , we obtain mass emissions rates of  $1.5 \times 10^{-1} \text{ kg s}^{-1}$  and  $1.5 \times 10^{-3} \text{ kg}$

We find that the model enhancement obtained from our reduced emissions are consistent with what we would expect from  
 the real emissions (Fig. 3b), despite the emissions being unrealistically low. Since the enhancements obtained are as expected  
 based on observations, we assume that this reduced ship production number is sufficient to drive this realistic test case, and  
 suitable for evaluation of the model's process representation. For the remaining results of this work, we only simulate ships  
 330 with ship production number of  $5 \times 10^{14} \text{ s}^{-1}$ .

### 3.2 Evaluation of the control


Using the model configuration detailed in Section 2.2, our simulation is run for 48 hours from 0000Z 11th July 2018 to 0000Z 13th July 2018. In our *control* run, no ship emissions are added to the model, and we use this simulation to characterise our “unpolluted” cloud. Typically, in ship track observations, we have no pure “control” cloud, and therefore we must treat a region outside of a ship track as being representative of what the cloud would have looked like if there was no aerosol perturbation present. In the model case, however, we can simply turn ships on and off to isolate the aerosol impact on the cloud.

The aerosol number concentration ( $N_a$ ), cloud droplet number concentration ( $N_d$ ), liquid water path (LWP), surface rain rate (RR), and top of atmosphere outgoing short wave radiation (SW) for the *control* run 44 hours after initialisation can be found in the left-hand column of Fig. 4. Fig. 4a demonstrates how the coupled GLOMAP-mode aerosol scheme allows depletion of aerosol in locations where there is precipitation (in the Central Pacific region of the domain), and the land sources of aerosol are the most significant over the course of the simulation. The coupled UM-CASIM configuration produces open/closed cellular convection in this marine stratocumulus (Fig. 4c in the south-west of the domain), with the presence of drizzle (surface rain rates on average of roughly 0.5 mm/hr; see Fig. 4g).


We evaluate our simulated precipitation in Fig. 5. 12-hourly instantaneous surface rain rates are shown for UM-CASIM *control* simulation (regridDED to the same 0.25° resolution as ERA5), and compared against ERA5 and GPM-IMERG surface rain rates. Evidently, there is a significant disagreement between ERA5 and GPM on the presence of precipitation in the domain, with GPM being unable to capture drizzle rates smaller than 0.2 mm/hr (Skofronick-Jackson et al., 2017). UM-CASIM is in rough agreement with ERA5 on the presence, location, and magnitude of the precipitation, however ERA5 contains its own uncertainties in the realism of precipitation (Xiong et al., 2022). The UM-CASIM precipitation rates lie between those from GPM-IMERG and ERA5. There are 3 overpasses of this domain by the CloudSat Cloud Profiling Radar (CPR; Stephens et al., 2008; Tanelli et al., 2008) during our simulation run. Fig. 6a shows the location and presence of surface precipitation (in the 2C-Precip-Column product; Haynes et al., 2009) in these overpasses. There is significant disagreement in between the 2C-Precip-Column and 2C-Rain-Profile (L’Ecuyer and Stephens, 2002) products from the CPR, with UM-CASIM underestimating precipitation along this overpass when compared to 2C-Precip-Column, and having a small mean bias when compared to 2C-Rain-Profile.

Evidently, this evaluation demonstrates that there is some uncertainty in the simulated precipitation within our domain, as well as within the observed precipitation. The UM-CASIM precipitation falls between what is predicted from ERA5 and GPM, with different CloudSat products giving largely different precipitation rates. More generally, double moment CASIM is shown to have increased skill at representing precipitation in a frontal system over the UK, and a tropical storm in Darwin (Field et al., 2023), as compared to other model microphysics schemes (evaluated against GPM and radar), however there are still limitations in its representation of moderate precipitation, in the range of 4-16 mm/hr (Bush et al., 2025).


With our initialised background aerosol field of  $200 \text{ cm}^{-3}$ , we produce a relatively clean cloud droplet background of roughly  $50 \text{ cm}^{-3}$ . This is largely in line with that seen in satellite observations (Fig. 7a,c), however we note that two large sources of cloud droplets are missing from our model simulation. This is likely due to our initialisation from a constant aerosol




**Figure 4.** Left column: control simulations of 12th July 2018 at 20:00 without ship emissions turned on. Panel (b) contains the locations of the ships simulated during this study, up until the time step of this figure. Right column: difference from control for simulation with shipping emissions on. Stochastic noise between two model runs can be seen in the difference plots. Variables shown are aerosol number concentration ( $N_a$ ), cloud droplet number concentration ( $N_d$ ), liquid water path (LWP), surface rain rate (RR), and top of atmosphere outgoing short wave radiation (SW).  $N_d$  and  $N_a$  are mean values across model level numbers below 1km.



**Figure 5.** Evaluation of domain wide surface precipitation at 12-hourly intervals. a) UM-CASIM simulated surface precipitation (regridged to ERA5 resolution), b) ERA5 surface precipitation, c) GPM surface precipitation.



**Figure 6.** Evaluation of surface against CloudSat CPR products. a) Location of CloudSat overpasses during the 48 hour simulation, with bold points showing locations where surface precipitation from the 2C-Precip-Column product are greater than zero. Associated times of overpass are roughly as follows: CS.64996 at 2200Z 11th July, CS.65004 at 1200Z on 12th July, and CS.65011 at 2300Z on 12th July.



**Figure 7.** Modelled and Observed  $N_d$  and LWP for 12th July 2018 ship tracks. Plotted is a composite of the Aqua and Terra overpasses from Fig. 1, co-located with model nearest output time steps.

365 [field, and therefore we are missing these sources of aerosol in our simulation. Future work in initialising from a more realistic aerosol field would be beneficial.](#)


### 3.3 Comparison to observations

As outlined in Section 2.1, we evaluate our simulations against three criteria in order to determine the realism of our simulations, and the ability of our model to accurately simulate ship tracks / MCB. In the following sections, we present our findings  
370 with respect to each of these criteria.

#### 3.3.1 In-track values

Fig. 8 provides a comparison between the mean of all MODIS observations of our ship tracks, and the associated modelled ship tracks (at the nearest time step). The inside track values of both  $N_d$  and LWP are a close match between model and observations, with a visible peak in the  $N_d$  at early times along track, and a slow increase in LWP with time along track which  
375 is consistent with the majority of the tracks moving down a LWP gradient. [Due to the missing high  \$N\_d\$  region in the Central Pacific part of the domain, ship tracks are only considered up to 15 hours along their length, since times longer than this are within this missing high  \$N\_d\$  region and therefore accurate comparison between model and observations is not possible.](#)

The mean percentage errors in the inside track values of  $N_d$  and LWP, up to 15 hours along the track, are found to be -3.5% and +3.1%, respectively. Therefore, for both  $N_d$  and LWP the model reproduces observed values to within +/- 4%, which is  
380 well within our uncertainty from the methods ([Fig-S??Section 2.6](#)). With respect to our first evaluation criteria, this means



**Figure 8.** (a)-(d) Observed  $N_d$  and LWP across all observations of our 5 ship tracks, as well as the percentage enhancement of the “inside track” region (defined by  $<15$  km from the centre). Percentage enhancement is calculated from the mean of the region between 30-45 km away from the centre of the track. (e)-(h) The same as above, but for modelled ship tracks. The percentage enhancement is instead calculated as the percentage change from the *control* model run (solid orange line). Percentage enhancement using the lateral offset method (as in the observations) is shown in the orange dashed line. (i)-(j) Mean of the observed and modelled  $N_d$  in-track values, and enhancements, in the “inside track” region. (k)-(l) The same, but for LWP. Whilst inside-track values for both  $N_d$  and LWP are similar between observations and model output, enhancements vary significantly **due to the definition of “unpolluted” background.**

that with the artificial reduction in aerosol concentrations necessary for the activation parameterisation to produce realistic shiptracks (see Section 3.1), we are able to successfully reproduce the absolute  $N_d$  and LWP values at the location where the aerosol is injected.

### 3.3.2 Timescales of response

385 If instead of considering the raw in-track  $N_d$  and LWP values, which do not clearly demonstrate the time evolution of the aerosol effect (since we are not considering a change from an unperturbed state), we calculate the percentage enhancement of the “inside track” region, compared to the “unpolluted” control cloud.

We find that the initial  $N_d$  increase and peak of roughly 300% at 3-4 hours is the same in both model and observations. However, despite the in-track values matching very closely between model and observations, the enhancements vary dramatically  
390 at longer time scales. In Fig. 8j, the enhancement in  $N_d$  is much longer lived than that seen in observations, which instead has returned to the background beyond roughly 15 hours. Since tracks are corrected by hand, this observed lifetime is not a result of inaccuracies in ship track location prediction. This highlights the importance of considering the change from the unperturbed state.


Similarly, for the LWP enhancements we see distinctly different responses. In the observations we see the response typically  
395 associated with ship tracks in non-precipitating environments, with a decrease in LWP in the first 5 hours, which then returns to zero. Conversely, in the modelled ship track we see only a blanket increase in LWP which increases over time. Fig. 8k demonstrates that the absolute inside track LWP values are similar, therefore this difference in enhancement must stem from the model representation of the control / background state, and the sensitivity of the model to aerosol.

Both the lifetime of the  $N_d$  and LWP responses are found to be too long-lived, which suggests that aerosol is not being  
400 efficiently removed from the clouds (Wood et al., 2012). This prompts further investigation into which processes are acting unrealistically, and in Section 3.3.3 we explore the realism of the simulated ship tracks in two different cloud types: closed MCC and broken cumulus.

Overall, we find that evaluating our model against this criteria in this way reveals issues with the lifetime of the aerosol response, and that current set up and formulation of these model parameterisations is unable to capture the observed lifetime  
405 of the  $N_d$  and LWP changes inside these ship tracks. This will have important implications for the radiative forcing from these ship tracks, since a longer lived aerosol response would prolong the brightening inside the ship track, and therefore would cause an overestimated cooling effect.

### 3.3.3 Sensitivity to initial condition

In Fig. 9, we split our ship tracks into those that occur in different background conditions, as a means to investigate whether  
410 our model is able to capture varying responses, and to assess the impact on model performance. ~~Since there is precipitation within the scene, we divide our tracks into precipitating / non-precipitating cases.~~ We use direction of travel as a simple proxy for “initial condition” of the cloud before the ship passes, since each grouping of ships is always within approximately 100 km of each other, and therefore experiences largely similar meteorological conditions. Ships A-C follow routes from



**Figure 9.** Percentage enhancements from the “unpolluted” references background state. **(a)-(d)** show observations of ship tracks in non-precipitating in the Eastern Pacific region of the domain (ships **A, B and CA-C**) and precipitating the Central Pacific region (ships **D and E**) conditions for  $N_d$  (left) and LWP (right). **(e)-(h)** shows the same but for modelled ship tracks. **(i)-(l)** show the average of the “inside track” region, as defined by the red dashed lines in **(a)-(h)**.


the coast of California, westwards ~~into the cleaner ocean through the Eastern Pacific~~ (see Fig. 4b). Ships D and E travel  
415 towards the coast and are more southerly than ships A-C. ~~Ships A-C travel through~~, through the Central Pacific. The Eastern  
Pacific ships travel through classic closed mesoscale cellular convective clouds ("closed MCC"), where there are primarily  
non-precipitating clouds, ~~whereas ships D and E travel through drizzling clouds~~. In the Central Pacific, there are "broken  
cumulus" that potentially transition into trade cumulus or open MCC, and we see higher occurrence of drizzle at the surface ;  
from these clouds (see Fig. S1). Dividing our enhancements into these two ~~groupings~~ different types of marine boundary layer  
420 clouds (Fig. 9), we see that the model is sensitive to the initial condition of the cloud before the ship passes through (Fig. 9).

In ~~non-precipitating conditions~~ closed MCC (Fig. 9, 1st and 3rd column for  $N_d$  and LWP enhancements, respectively) we  
find similar enhancements in the model and observations. ~~On short timescales~~ Shortly after the ship passes there is better  
agreement, with observations and model simulations matching within the uncertainty in the first five hours. Following this, we  
see divergence between the two. This could be in part due to our tuning of the aerosol emissions to satisfy the non-monotonic  
425 behaviour of the activation parameterisation and reproduce correct  $N_d$  enhancements on short time scales. We would still  
expect the response from weaker aerosol perturbations to reduce over time, and therefore there must be some process within  
the model allowing for this increased  $N_d$  perturbation lifetime. A potential source of this long-lived  $N_d$  enhancement could be  
inefficient removal of cloud droplets by entrainment. In these clouds, we see larger simulated LWP enhancements from 10-15  
hours after the ship passes, therefore weak entrainment rates could correspond to the large  $N_d$  enhancements at these times also.  
430 Additionally, even in these closed MCC conditions where we do not observe much drizzle (Fig. S1), if cloud droplets cannot  
become rain droplets via accretion or autoconversion, or if larger droplets cannot sediment out, then the removal of aerosol will  
be underestimated and reinforce the  $N_d$  perturbation. A thorough analysis of the output process rates is needed to fully identify  
the source of this long-lived  $N_d$  perturbation.

Considering the ships that travel through ~~precipitating broken cumulus~~ clouds (Fig. 9 2nd and 4th column for  $N_d$  and LWP  
435 enhancements respectively), we see significant disagreement between the observations and the model. This suggests that the  
large disagreement in Fig. 8 is largely due to ~~these ships in precipitating conditions~~ the ship tracks in these conditions where  
precipitation is more prevalent. The  $N_d$  enhancement is too large, and shows no sign of diminishing even after 15 hours after  
the aerosol perturbation. This suggests that the model is not only overly sensitive to aerosol in ~~precipitating conditions~~ broken  
cumulus clouds, but is not effective enough at removing aerosol from the cloud either. Additionally, the LWP response is a  
440 different sign between model and observations. This suggests that the model is far too keen to suppress precipitation in these  
conditions, leading to large increases in LWP that are not observed in the satellite imagery.

There is some uncertainty that may be introduced due to the missing cloud structure in the Central Pacific region in our  
simulation, causing our background state to look cleaner than it should be. This missing cloud structure, however, will  
only influence the ship track over 10 hours along the track, and therefore is not the source of the large overestimation in  
445  $N_d$  enhancement seen before 10 hours along the track in Fig. 9j.

In Fig. 10, we show the surface rain rates inside our ship tracks, compared to the control run. Here, we see that for tracks  
D and E (~~which occur in more precipitating environments~~ in the broken cumulus region (where there are larger precipitation




**Figure 10.** Surface rain rate, averaged across two groupings of tracks (ships A-C and D+E), across 24 hours on the 12th July 2018, for the *ships* and *control* runs, as well as the difference between the two. ~~Clear~~The figure shows clear precipitation suppression in the “~~precipitating~~” ~~broken cumulus~~ case, with rain rates reduced to almost zero. Some precipitation suppression in the “~~non-precipitating~~” ~~closed MCC~~ case (ships A-C) since there is actually some precipitation towards the end of the model run (see Fig. S1).

~~rates~~), the precipitation is essentially completely shut off which is likely a far too strong response, since precipitation in high  $N_d$  conditions does occur (Gryspeerdt et al., 2022).

450 We demonstrate this in Fig. 11, where the probability of precipitation (PoP) is plotted in  $N_d$ -LWP ~~phase space for our space~~ for (a) our 48 hour model simulation and ~~collocated~~ (b) four years of collocated CloudSat-MODIS observations (following a similar methodology to Gryspeerdt et al., 2022). The ~~distribution of PoP within this  $N_d$ -LWP space is a representation of microphysical processes, and shouldn't differ significantly seasonally or day-to-day, therefore a comparison between a two-day simulation and four years of observational data should represent the same distribution (see Fig. S2).~~

455 The very sharp gradient in PoP with increases in  $N_d$  demonstrates the high-sensitivity of the autoconversion scheme to aerosol. In our ~~precipitating-broken cumulus~~ case, the addition of ship aerosol moves our “background” point in  $N_d$ -LWP phase space just over the sharp boundary in PoP to our “ship track” point. This explains the complete precipitation suppression



**Figure 11.** The probability of precipitation (PoP) in  $N_d$ -LWP space, in UM-CASIM simulations compared to CloudSat-MODIS collocated observations for the same domain. UM-CASIM has much faster transition from precipitating to non-precipitating with changes in  $N_d$  than observed. The orange point is the background (*control*)  $N_d$  and LWP for our precipitating ship (D-E), and the red point is the  $N_d$  and LWP inside the ship tracks, averaged over the simulation run time.

in these cases, since this gradient is so sharp. In observations, there would ~~perhaps~~ be a small decrease in PoP, but not to the same extent.

460 In our simulations, we do see that there is some sensitivity to the initial conditions since we obtain different responses in ~~precipitating / non-precipitating cases.~~ different cloud structures (closed MCC vs. broken cumulus). We can conclude that the model is more suitable for simulations of high-concentration aerosol perturbations in ~~non-precipitating conditions~~ closed MCC conditions where there is little precipitation, at least for up to 5 hours. However, in the case of ~~precipitating clouds~~ broken cumulus clouds with higher occurrence of precipitation, current model parameterisation formulations are not suitable  
465 to accurately simulate these scenarios, and the lifetime of the response is far too long lived.

## 4 Discussion

In this study, we produce convection permitting simulations of ship tracks using a regional model and double moment micro-physics scheme, exhibiting its capabilities and limitations in simulating high-concentration aerosol interactions with clouds. We evaluate our case study against three criteria, which are designed to ensure ship track simulations contain an accurate  
470 representation of cloud processes, and ~~would therefore may~~ be a useful tool for MCB research.

~~With some tuning of parameterisations~~ Adjusting ship emissions to  $5 \times 10^{14}$  particles per second as to avoid unphysical behaviour in the activation parameterisation, we reproduce realistic absolute values of  $N_d$  and LWP inside simulated ship tracks, and find that in closed MCC (non-precipitating conditions) the enhancement in  $N_d$  and LWP from the control “unpolluted” cloud is well modelled for the first 5 hours. However, beyond this, the lifetime of the response is too long lived. This is  
475 driven by the representation of parameterisations at very high aerosol concentrations.

## 4.1 Definition of the “control” region. Limitations

In order to quantify the enhancement of cloud properties inside ship tracks from their background state, we must first define this unperturbed “control

### 4.1.1 Definition of “precipitating” and “non-precipitating” clouds

480 In this work, we divide our ship tracks into those that occur in “closed MCC” ~~cloud~~-clouds (tracks A-C), and those in “broken cumulus” (tracks D&E). When evaluating against satellite observations, we find that our model performs better in these closed MCC cases, and worse in the broken cumulus. We attribute the difference in behaviour of the model in these different cloud types as being due to the representation of precipitation by the model. The closed MCC clouds in our model have higher background CCN concentrations (Fig. 4a) and greater  $N_a$  (Fig. 4b) with less precipitation (Fig. 4g), whereas the  
485 broken cumulus have heavier precipitation (Fig. 4g). This is in line with what is expected in these cloud types in literature (Alinejadtabrizi et al., 2024). We find that precipitation is too easily suppressed and therefore ship track lifetimes are too great in clouds where there is precipitation (such as our broken cumulus). Non-precipitating clouds (such as our closed MCC) are less impacted by these, and therefore the model produces more realistic simulations of these clouds.

In model simulations, we are able to define this region from our *control* simulation, where there are no ship emissions and  
490 we can directly measure the unperturbed cloud. In observations, however, we do not have a control run, and therefore region is more complex to define. Typically, we consider the region neighbouring the ship track as being representative of what the cloud would have looked like if there was no ship present, and use this to infer the aerosol effect on the cloud.

Whilst this methodology is useful to predict the control cloud, it is not perfect. Ship tracks often occur in large groupings and have considerable overlap, which may distort the ability to define the “clean  
495 closed MCC and broken cumulus other than amounts of precipitation that may impact this analysis. In Fig. 4 “control” background and make it difficult to observe aerosol effects at longer lifetimes. This introduces a source of uncertainty when comparing enhancements in Fig. 8 and Fig. 9. column the difference between the Eastern Pacific closed MCC and Central Pacific broken cumulus can be observed. Closed MCC have greater cloud fraction (CF), as seen in Fig. 4c, which impacts the spatial scale over which aerosol can impact the clouds. The simulated marine boundary layer (MBL) height increases with  
500 distance from the coast and is roughly 550m in the Eastern Pacific, deepening to 900m in the Central Pacific (not shown). In the deeper MBL, it is possible that aerosols may not reach cloud base as readily (Bretherton and Wyant, 1997), which could be a potential reason why satellite observations of tracks D&E do not show significant LWP enhancement - the model may be mixing aerosol vertically too efficiently in these clouds, therefore the perturbation to  $N_a$  is too great. Alternatively, broken cumulus regions can exhibit higher latent/sensible heat fluxes and weaker subsidence (Wood and Bretherton, 2004), creating a  
505 deeper MBL with weaker entrainment rates. If entrainment is too weak, however, then LWP anomalies inside the ship tracks will be allowed to exist for too long (Fig. 8j and l).

We compare these two different methodologies for defining the control in our model simulation in Fig. S???. The first method is by using the model control run, and the second is Additionally, within this single case study, we are restricted in the extent to

510 which we can divide our tracks into “different conditions”. All these tracks still occur within the same domain on the same day, therefore there is a limitation on how different the conditions can be, especially since all ships converge on the same region in the centre of the domain. A more comprehensive statistical analysis is necessary to determine fully the behaviour of the model across many cloud types and conditions, which is not possible with a single 48 hour case study.

#### 4.1.2 Aerosol scheme

515 In this work, we only consider aerosols in the accumulation mode, and neglect other aerosol size modes such as sea salt and organic matter. These other modes have different hygroscopicities, which can affect cloud droplet formation and the initiation of precipitation. For example, sea salt can modulate the activation of sulphate nuclei into cloud droplets (Fossum et al., 2020), therefore inclusion of even small concentrations of sea salt aerosol may reduce  $N_d$  concentrations. McCoy et al. (2018) also found that increasing sea salt concentrations limits  $N_d$  due to reduction in supersaturation by the larger sea salt CCN. This effect occurs because of the different hygroscopicity of sea salt, but also its larger size, making it an effective CCN. The neglecting  
520 of other aerosol modes with different hygroscopicities should not significantly impact cloud droplet number concentrations or initiation of precipitation, however the neglecting of aerosols with different sizes may impact our analysis, since the size of the aerosol mode remains the more important factor for cloud nucleating ability of aerosol particles (Dusek et al., 2006). Future work with a more comprehensive aerosol configuration would be beneficial in order to investigate the impact of different aerosol size modes on cloud droplet activation in ship tracks.

525 Additionally, our coupling between CASIM and GLOMAP-mode aerosols (as described in Gordon et al., 2020) means that the removal of aerosol from the cloud is only possible through wet-scavenging. Autoconversion and accretion rates determine the rate of removal of aerosol by precipitation, however cloud microphysical processing of the aerosol is not implemented, meaning that there is no reduction of  $N_a$  by using the “outside” ship track region (as is typically done for observations). In our model simulation, we find that this introduces 10% and 15% underestimation in the and LWP enhancements, respectively.  
530 However, since our model simulation is very clean, and does not contain the many other ship tracks seen in observations, this gives a lower bound on the expected underestimation from defining the control region using the neighbouring clouds. collision-coalescence of cloud droplets, nor an increase in  $N_a$  through droplet evaporation (Gordon et al., 2020). This severely limits the mechanisms by which aerosol can be removed from the cloud, since collision-coalescence scavenging is an important sink of CCN (Kang et al., 2022; Terai et al., 2014) and may explain our long-lived  $N_d$  enhancements within our ship tracks.  
535 Once precipitation is even slightly suppressed, aerosol removal from the cloud is also significantly reduced,  $N_d$  remains high (Wood et al., 2012), and therefore ship track lifetimes are too long. In Fig. 11 we see that very small increases in  $N_d$  are able to cause an almost complete suppression of precipitation (as a result of our autoconversion parameterisation), which in turn will feedback on the lifetime of the  $N_d$  response, since aerosol will continue to exist in the cloud. This indicates that cloud processing is important for accurately simulating ship tracks, as to avoid this precipitation suppression feedback.

540 Essentially, with

### 4.1.3 Definition of the “control” region

When there are fewer background sources of aerosols, it is easier to define the “unperturbed” cloud using the surrounding region-region adjacent to the ship track. When there are more ships, or other sources of aerosols, this definition becomes more difficult and becomes a greater source of uncertainty. Other studies have explored alternative methodologies of gaining insight into the unperturbed cloud in observations (such as through the use of ML algorithms; Diamond et al., 2020; Chen et al., 2022; Diamond, 2023), however will still struggle with increasingly complex and polluted backgrounds.

This has important implications for the design of field experiments of MCB. In order to isolate the aerosol effect with greatest certainty, a relatively simple-homogeneous background is necessary. If there are many overlapping ship tracks / interacting sources, this definition of the control becomes more nuanced, and additional uncertainty is introduced. If field trials were to occur for MCB, the easiest, and best attribution of the aerosol effect could be made if ships tracks were not allowed to interact, and the background was not too heavily polluted with sources that were difficult to define.

## 4.2 Suitability of parameterisations for high-concentration aerosol perturbations

We find-confirm that the commonly used activation parameterisation by Abdul-Razzak and Ghan (2000) is inadequate to simulate aerosol-cloud interactions at very high aerosol concentrations. We demonstrate the potential result of the non-monotonicity of the ARG activation when simulating ship tracks - ship tracks appear split (Fig. 2). In practical terms, this necessitates an artificial reduction in aerosol-input-ship aerosol emissions to obtain ship track structures that resemble those seen in satellite imagery. While this workaround may yield visually plausible results in non-precipitating conditions, it undermines the physical realism of the simulation at long timescales.'s temporal evolution. An alternative activation parameterisation from Nenes and Seinfeld (2003) would also be unsuitable at these high aerosol concentrations, as activated cloud droplet concentration would be overestimated. Further work is necessary in determining whether other parameterisations (such as that of Shipway and Abel, 2010) would be suitable, or if entirely new schemes for high aerosol concentrations are needed.

Model-observation We find that model-observation discrepancies become more pronounced in precipitating environments broken cumulus clouds, where there is greater precipitation. Under such conditions, it is possible that the autoconversion parameterisation of Khairoutdinov and Kogan (2000, KK00), which governs the conversion of cloud droplets-water to rain, is too sensitive to increases in  $N_d$ . This scheme tends to suppress production of rain droplets too readily when the cloud droplet number concentration ( $N_d$ ) is high - as demonstrated by the complete shut off of precipitation inside these ship tracks that form in precipitating scenes (see Fig. 10).

In Fig. 11, the PoP is almost binary in our simulation, with either 100% chance of precipitation above some threshold  $N_d$ -LWP line, and almost 0% chance below it. In observations, there exists a much wider space in which precipitation can occur. This means that for any conditions close to this threshold line in  $N_d$ -LWP in our simulation, an-increases-a small increase of  $N_d$  (such as through the injection of ship emissions), will move us beyond this threshold and decreases the PoP too extremely.

Consequently, the model predicts unrealistic suppression of precipitation, resulting in overestimation of the ship track lifetime and, by extension, the radiative cooling associated with the perturbation. This is not surprising, since the parameterisation

575 ~~from~~ KK00 ~~was only designed and tested on~~ autoconversion scheme was designed for mid-latitude and extratropical stratocu-  
mulus layers under typical meteorological conditions, with their evaluated “polluted” case only having an  $N_d$  of  $175 \text{ cm}^{-3}$ . This  
is aligned with the findings of Jing and Suzuki (2018) with respect to GCM parameterisations, where precipitation inhibition  
can lead to a wet scavenging feedback which can increase aerosol loading somewhat non-physically. ~~In Gryspeerd et al. (2022)~~  
~~, it is found that~~ KK00 ~~completely suppresses precipitation at high~~, ~~despite there still being some probability of precipitation~~  
~~if LWP is high. KK00 is likely~~ is possibly not well-suited to simulate very high aerosol concentration perturbations, as it tends  
580 to easily suppress precipitation - even in conditions where precipitation could still occur.

As discussed in Section 3.2, there is some uncertainty in the realism of the simulated precipitation in the domain, since it  
is difficult to obtain conclusive precipitation measurements across different satellite and reanalysis products. In order for the  
enhancement of LWP in a ship track to occur, precipitation need only be suppressed at cloud base, and therefore it is also  
possible that no impact in surface precipitation would occur. This highlights the difficulty in evaluating not only precipitation  
585 in models, but specifically the precipitation suppression effect.

In their designed configurations, these parameterisations limit the ability of models to provide meaningful insight into high  
concentration aerosol-cloud interactions over longer timescales or under precipitating conditions. This has important implica-  
tions with respect to ~~climate modelling~~, ~~where high-resolution modelling across many scales, not just regional simulations.~~  
The development of global cloud resolving models ~~are being employed.~~ would also benefit from improved cloud microphysics  
590 parameterisations (Satoh et al., 2019).

Additionally, for simulations aimed at informing MCB strategies, it is particularly critical that parameterisations are suitable  
for high-aerosol scenarios. Without modifications, models will likely overstate the cooling impacts, leading to inaccurate  
assessments of MCB efficacy. Improved autoconversion schemes that remain valid at high  $N_d$ , ~~(such as a scheme where~~  
~~autoconversion rate does not tend to zero as quickly with increases in~~  $N_d$  ~~for a given LWP, allowing for a more diffuse PoP~~  
595 ~~within~~  $N_d$ -LWP space), along with more robust activation parameterisations that avoid the need for artificial aerosol reductions,  
~~are~~ and the inclusion of cloud-processing such that precipitation suppression feedbacks are avoided, are essential for advancing  
the realism of such simulations.

## 5 Conclusions

Accurate model representation of aerosol-cloud processes is essential for reducing uncertainty in climate projections and  
600 simulating the impacts of marine cloud brightening (MCB). Ship tracks provide a “natural experiment” for isolating the aerosol  
effects on clouds, and gaining information about the time evolution of the cloud response. In order to evaluate the ~~model~~  
~~representation of these processes~~ simulated response to an aerosol perturbation, we simulate a real case of ship tracks observed  
off the coast of California on 12th July 2018, using an experimental configuration of the Met Office’s UK weather forecasting  
model.

605 We evaluate the performance of our model against three criteria, aiming to produce simulations of ship tracks which are a  
useful tool for investigating the impacts of MCB. We assess the model’s ability to reproduce observed relative and absolute

changes in cloud droplet number concentration ( $N_d$ ) and liquid water path (LWP), using MODIS satellite retrievals for validation. The ship tracks occur within a marine stratocumulus deck under ~~very~~ moderately clean background conditions, including pockets of drizzling cloud.

610 We find that the activation parameterisation of Abdul-Razzak and Ghan (2000, ARG) is not well-suited for simulating high aerosol concentrations typical of ship plumes. In particular, ARG exhibits non-monotonic behaviour at elevated concentrations, leading to non-physical results unless aerosol input is artificially reduced by a factor of ~~20-20~~ (Fig. 3). This adjustment yields raw  $N_d$  and LWP values within ship tracks that closely match observations (Fig. 8). An alternative activation parameterisation, from Nenes and Seinfeld (2003, NS), is found to overestimate expected  $N_d$  at high aerosol concentrations when evaluated in a simple adiabatic parcel model (Fig. 3). Further work is necessary to fully evaluate other activation parameterisations (such as that of Shipway and Abel, 2010) at these high concentrations and determine whether modifications can be made to existing schemes, or if novel ones must be developed.

620 ~~However, analysis of~~ Analysis of raw values inside ship tracks does not tell us about the time evolution of the aerosol effect, or how long it takes the clouds to return to the unperturbed state. In the model, this is easy to define through the use of a *control* simulation. In the observations, we consider the region just outside of the ship track as a representative control region. This highlights the usefulness of ship tracks in observing aerosol-cloud interactions, as it provides us with this control region that is otherwise so hard to define. When considering the enhancements from the control state, we find that the model fails to accurately simulate the time evolution of the response that is seen in observations ~~-(Fig. 8)~~. Specifically, model enhancements in  $N_d$  persist too long, and the LWP response diverges in sign from observations. On average, across five simulated ship tracks, 625 the model does not capture the correct time evolution of the aerosol effect. There is some uncertainty (roughly 10%) introduced by the definition of the control state in the observations, since it is difficult to define the unperturbed cloud when there are many overlapping ship tracks. Additionally, some of this discrepancy could be attributed to a distortion in the time evolution due to the ~~tuning of the activation parameterisation (such that physical ship tracks were produced~~ reduction of ship emissions (to better match the observed tracks, and avoid “split” tracks).

630 We further classify ship tracks by background precipitation conditions to better understand this discrepancy ~~-(Fig. 9)~~. In non-precipitating ~~environments~~ closed MCC clouds, the model captures the initial enhancement in cloud properties but still exhibits a response that is too long lived compared to observations (Fig. 9i,k). This suggests that while the initial tuning of the activation parameterisation produces realistic tracks, it may distort the subsequent cloud evolution, and points to the broader need for improved activation schemes capable of handling extreme aerosol conditions, especially in contexts such as MCB 635 (Connolly et al., 2014).

In precipitating environments, model-observation disagreement is more pronounced, which is critical for studies of cloud adjustments to aerosols. Model  $N_d$  enhancements are significantly overestimated and show no decay over time, whereas observations indicate a return to background values within 15 hours (Fig. 9j). Moreover, the model shows a strong positive LWP response, while observations show a weakly negative one. This suggests that the model either (a) overestimates the sensitivity of precipitation suppression to aerosol, (b) misrepresents background precipitating conditions (Fig. 11), or (c) ~~cannot be compared to potentially unrepresentative observations~~ does not sufficiently remove aerosol from cloud.

~~This Any model overestimation of the aerosol effect~~ has particular relevance for solar radiation management proposals such as MCB, and aerosol forcing estimates from ~~high-resolution models. km-scale models. Simulating aerosol perturbations accurately on the individual cloud level is an exceptionally difficult task for the field, and robust evaluation of model representation at these scales is necessary.~~ Current model parameterisations may cause an overestimation in the cloud responses to aerosol and predict an unrealistically strong cooling effect, especially under precipitating conditions. Any application of these models to unobserved phenomena like MCB must be grounded in rigorous comparison with real-world observations, and further work is necessary in modifying parameterisations such that they are more suitable for high-concentration aerosol perturbations.

*Code and data availability.* MODIS data used in this work were acquired from the Level-1 and Atmosphere Archive and Distribution System (LAADS) Distributed Active Archive Center (DAAC) (<https://ladsweb.modaps.eosdis.nasa.gov>, LAADS DAAC, 2021; Platnick et al. 2017). The ERA5 data are from the Copernicus Climate Change Service (C3S) Climate Data Store (CDS) (<https://doi.org/10.24381/cds.adbb2d47>, Copernicus Climate Change Service, 2023; Hersbach et al. 2020). Ship AIS data were obtained from exactEarth. exactEarth data are not publicly available; however, they can be accessed via a paid subscription: <https://www.marinetraffic.org/exactearth> (last access: 29 November 2024). Code and data from this analysis will be made available upon publication, via Zenodo.

*Author contributions.* All authors contributed to the design of the study. PRF provided access to the Met Office model and Monsoon, and supported AT in the set up and running the simulations. AT performed the model output and observational analysis, with EG providing observations data for Fig. 9. PRF and EG assisted with the interpretation of the results. AT drafted the manuscript, and PRF and EG provided comments and suggestions.

*Competing interests.* No competing interests are present

*Acknowledgements.* We acknowledge use of the Monsoon2 system, a collaborative facility supplied under the Joint Weather and Climate Research Programme, a strategic partnership between the Met Office and the Natural Environment Research Council. Anna Tippett and Edward Gryspeerdt acknowledge funding from the Horizon Europe programme (project CERTAINTY – Cloud–aERosol inTeractions & their impActs IN The earth sYstem; grant agreement no. 101137680) and a Royal Society University Research Fellowship (grant no. UR-F/R1/191602). ~~Paul R. Field acknowledges funding from ...~~

665 **References**

- Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types, *Journal of Geophysical Research: Atmospheres*, 105, <https://doi.org/10.1029/1999JD901161>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/1999JD901161>, 2000.
- Abdul-Razzak, H., Ghan, S. J., and Rivera-Carpio, C.: A parameterization of aerosol activation: 1. Single aerosol type, *Journal of Geophysical Research: Atmospheres*, 103, <https://doi.org/10.1029/97JD03735>, 1998.
- 670 Ackerman, A. S., Toon, O. B., Taylor, J. P., Johnson, D. W., Hobbs, P. V., and Ferek, R. J.: Effects of Aerosols on Cloud Albedo: Evaluation of Twomey's Parameterization of Cloud Susceptibility Using Measurements of Ship Tracks, *Journal of the Atmospheric Sciences*, 57, [https://doi.org/10.1175/1520-0469\(2000\)057<2684:EOAOCA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<2684:EOAOCA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2000.
- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol  
675 climate forcing, *Nature*, 432, 1014–1017, <https://doi.org/10.1038/nature03174>, 2004.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, publisher: American Association for the Advancement of Science, 1989.
- Alinejadtabrizi, T., Lang, F., Huang, Y., Ackermann, L., Keywood, M., Ayers, G., Krummel, P., Humphries, R., Williams, A. G., Siems, S. T., and Manton, M.: Wet deposition in shallow convection over the Southern Ocean, *npj Climate and Atmospheric Science*, 7, 76,  
680 <https://doi.org/10.1038/s41612-024-00625-1>, publisher: Nature Publishing Group, 2024.
- Andreae, M. O., Jones, C. D., and Cox, P. M.: Strong present-day aerosol cooling implies a hot future, *Nature*, 435, 1187–1190, <https://doi.org/10.1038/nature03671>, number: 7046 Publisher: Nature Publishing Group, 2005.
- Arabas, S. and Shima, S.-i.: On the CCN [de]activation nonlinearities, *Nonlinear Processes in Geophysics Discussions*, <https://doi.org/10.5194/npg-2016-50>, 2016.
- 685 Berner, A. H., Bretherton, C. S., and Wood, R.: Large eddy simulation of ship tracks in the collapsed marine boundary layer: a case study from the Monterey area ship track experiment, *Atmospheric Chemistry and Physics*, 15, <https://doi.org/10.5194/acp-15-5851-2015>, 2015.
- Bretherton, C. S. and Wyant, M. C.: Moisture Transport, Lower-Tropospheric Stability, and Decoupling of Cloud-Topped Boundary Layers, *Journal of the Atmospheric Sciences*, 54, 148–167, [https://doi.org/10.1175/1520-0469\(1997\)054<0148:MTLTA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<0148:MTLTA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 1997.
- 690 Bretherton, C. S., Blossey, P. N., and Uchida, J.: Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo, *Geophysical Research Letters*, 34, L03 813, <https://doi.org/10.1029/2006GL027648>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2006GL027648>, 2007.
- Briggs, G. A.: A Plume Rise Model Compared with Observations, *Journal of the Air Pollution Control Association*, 15, 433–438, <https://doi.org/10.1080/00022470.1965.10468404>, 1965.
- 695 Brown, A., Milton, S., Cullen, M., Golding, B., Mitchell, J., and Shelly, A.: Unified Modeling and Prediction of Weather and Climate: A 25-Year Journey, *Bulletin of the American Meteorological Society*, 93, 1865–1877, <https://doi.org/10.1175/BAMS-D-12-00018.1>, publisher: American Meteorological Society Section: *Bulletin of the American Meteorological Society*, 2012.
- Bush, M., Allen, T., Bain, C., Boutle, I., Edwards, J., Finnenkoetter, A., Franklin, C., Hanley, K., Lean, H., Lock, A., Manners, J., Mittermaier, M., Morcrette, C., North, R., Petch, J., Short, C., Vosper, S., Walters, D., Webster, S., Weeks, M., Wilkinson, J., Wood, N., and Zerroukat,  
700 M.: The first Met Office Unified Model–JULES Regional Atmosphere and Land configuration, RAL1, *Geoscientific Model Development*, 13, 1999–2029, <https://doi.org/10.5194/gmd-13-1999-2020>, publisher: Copernicus GmbH, 2020.

- Bush, M., Boutle, I., Edwards, J., Finnenkoetter, A., Franklin, C., Hanley, K., Jayakumar, A., Lewis, H., Lock, A., Mittermaier, M., Mohandas, S., North, R., Porson, A., Roux, B., Webster, S., and Weeks, M.: The second Met Office Unified Model–JULES Regional Atmosphere and Land configuration, RAL2, Geoscientific Model Development, 16, 1713–1734, <https://doi.org/10.5194/gmd-16-1713-2023>, publisher: Copernicus GmbH, 2023.
- 705 Bush, M., Flack, D. L. A., Lewis, H. W., Bohnenstengel, S. I., Short, C. J., Franklin, C., Lock, A. P., Best, M., Field, P., McCabe, A., Van Weverberg, K., Berthou, S., Boutle, I., Brooke, J. K., Cole, S., Cooper, S., Dow, G., Edwards, J., Finnenkoetter, A., Furtado, K., Halladay, K., Hanley, K., Hendry, M. A., Hill, A., Jayakumar, A., Jones, R. W., Lean, H., Lee, J. C. K., Malcolm, A., Mittermaier, M., Mohandas, S., Moore, S., Morcrette, C., North, R., Porson, A., Rennie, S., Roberts, N., Roux, B., Sanchez, C., Su, C.-H., Tucker, S., Vosper, S., Walters, D., Warner, J., Webster, S., Weeks, M., Wilkinson, J., Whittall, M., Williams, K. D., and Zhang, H.: The third Met Office Unified Model–JULES Regional Atmosphere and Land Configuration, RAL3, Geoscientific Model Development, 18, 3819–3855, <https://doi.org/10.5194/gmd-18-3819-2025>, publisher: Copernicus GmbH, 2025.
- 710 Chen, Y., Haywood, J., Wang, Y., Malavelle, F., Jordan, G., Partridge, D., Fieldsend, J., De Leeuw, J., Schmidt, A., Cho, N., Oreopoulos, L., Platnick, S., Grosvenor, D., Field, P., and Lohmann, U.: Machine learning reveals climate forcing from aerosols is dominated by increased cloud cover, *Nature Geoscience*, 15, 609–614, <https://doi.org/10.1038/s41561-022-00991-6>, 2022.
- 715 Christensen, M. W., Gettelman, A., Cermak, J., Dagan, G., Diamond, M., Douglas, A., Feingold, G., Glassmeier, F., Goren, T., Grosvenor, D. P., Gryspeerdt, E., Kahn, R., Li, Z., Ma, P.-L., Malavelle, F., McCoy, I. L., McCoy, D. T., McFarquhar, G., Mülmenstädt, J., Pal, S., Possner, A., Povey, A., Quaas, J., Rosenfeld, D., Schmidt, A., Schrödner, R., Sorooshian, A., Stier, P., Toll, V., Watson-Parris, D., Wood, R., Yang, M., and Yuan, T.: Opportunistic experiments to constrain aerosol effective radiative forcing, *Atmospheric Chemistry and Physics*, 22, 641–674, <https://doi.org/10.5194/acp-22-641-2022>, publisher: Copernicus GmbH, 2022.
- 720 Chun, J.-Y., Wood, R., Blossey, P., and Doherty, S. J.: Microphysical, macrophysical, and radiative responses of subtropical marine clouds to aerosol injections, *Atmospheric Chemistry and Physics*, 23, 1345–1368, <https://doi.org/10.5194/acp-23-1345-2023>, publisher: Copernicus GmbH, 2023.
- 725 Connolly, P. J., McFiggans, G. B., Wood, R., and Tsiamis, A.: Factors determining the most efficient spray distribution for marine cloud brightening, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372, <https://doi.org/10.1098/rsta.2014.0056>, publisher: Royal Society, 2014.
- Diamond, M. S.: Detection of large-scale cloud microphysical changes within a major shipping corridor after implementation of the International Maritime Organization 2020 fuel sulfur regulations, *Atmospheric Chemistry and Physics*, 23, 8259–8269, <https://doi.org/10.5194/acp-23-8259-2023>, publisher: Copernicus GmbH, 2023.
- 730 Diamond, M. S., Director, H. M., Eastman, R., Possner, A., and Wood, R.: Substantial Cloud Brightening From Shipping in Subtropical Low Clouds, *AGU Advances*, 1, e2019AV000111, <https://doi.org/10.1029/2019AV000111>, <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019AV000111>, 2020.
- 735 Diamond, M. S., Gettelman, A., Lebsock, M. D., McComiskey, A., Russell, L. M., Wood, R., and Feingold, G.: To assess marine cloud brightening’s technical feasibility, we need to know what to study—and when to stop, *Proceedings of the National Academy of Sciences*, 119, e2118379119, <https://doi.org/10.1073/pnas.2118379119>, 2022.
- Durkee, P. A., Chartier, R. E., Brown, A., Trehubenko, E. J., Rogerson, S. D., Skupniewicz, C., Nielsen, K. E., Platnick, S., and King, M. D.: Composite Ship Track Characteristics, *Journal of the Atmospheric Sciences*, 57, [https://doi.org/10.1175/1520-0469\(2000\)057<2542:CSTC>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<2542:CSTC>2.0.CO;2), 2000.

- Dusek, U., Frank, G. P., Hildebrandt, L., Curtius, J., Schneider, J., Walter, S., Chand, D., Drewnick, F., Hings, S., Jung, D., Borrmann, S.,  
740 and Andreae, M. O.: Size Matters More Than Chemistry for Cloud-Nucleating Ability of Aerosol Particles, *Science*, 312, 1375–1378,  
<https://doi.org/10.1126/science.1125261>, publisher: American Association for the Advancement of Science, 2006.
- Feingold, G., Eberhard, W. L., Veron, D. E., and Previdi, M.: First measurements of the Twomey indirect effect using ground-based remote sensors, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2002GL016633>,  
<https://onlinelibrary.wiley.com/doi/pdf/10.1029/2002GL016633>, 2003.
- 745 Feingold, G., Ghate, V. P., Russell, L. M., Blossey, P., Cantrell, W., Christensen, M. W., Diamond, M. S., Gettelman, A., Glassmeier, F.,  
Gryspeerd, E., Haywood, J., Hoffmann, F., Kaul, C. M., Lebsock, M., McComiskey, A. C., McCoy, D. T., Ming, Y., Mülmenstädt, J.,  
Possner, A., Prabhakaran, P., Quinn, P. K., Schmidt, K. S., Shaw, R. A., Singer, C. E., Sorooshian, A., Toll, V., Wan, J. S., Wood, R., Yang,  
F., Zhang, J., and Zheng, X.: Physical science research needed to evaluate the viability and risks of marine cloud brightening, *Science  
Advances*, 10, eadi8594, <https://doi.org/10.1126/sciadv.adi8594>, publisher: American Association for the Advancement of Science, 2024.
- 750 Ferek, R. J., Hegg, D. A., Hobbs, P. V., Durkee, P., and Nielsen, K.: Measurements of ship-induced tracks in clouds off the Washington coast,  
*Journal of Geophysical Research*, 103, <https://doi.org/10.1029/98JD02121>, aDS Bibcode: 1998JGR...10323199F, 1998.
- Ferrier, B. S.: A Double-Moment Multiple-Phase Four-Class Bulk Ice Scheme. Part I: Description, *Journal of the Atmospheric Sciences*, 51,  
249–280, [https://doi.org/10.1175/1520-0469\(1994\)051<0249:ADMMPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<0249:ADMMPF>2.0.CO;2), publisher: American Meteorological Society Section:  
*Journal of the Atmospheric Sciences*, 1994.
- 755 Field, P. R., Hill, A., Shipway, B., Furtado, K., Wilkinson, J., Miltenberger, A., Gordon, H., Grosvenor, D. P., Stevens, R., and Van We-  
verberg, K.: Implementation of a double moment cloud microphysics scheme in the UK met office regional numerical weather  
prediction model, *Quarterly Journal of the Royal Meteorological Society*, 149, 703–739, <https://doi.org/10.1002/qj.4414>,  
<https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.4414>, 2023.
- Fons, E., Runge, J., Neubauer, D., and Lohmann, U.: Stratocumulus adjustments to aerosol perturbations disentangled with a causal approach,  
760 *npj Climate and Atmospheric Science*, 6, 1–10, <https://doi.org/10.1038/s41612-023-00452-w>, number: 1 Publisher: Nature Publishing  
Group, 2023.
- Fossum, K. N., Ovadnevaite, J., Ceburnis, D., Preißler, J., Snider, J. R., Huang, R.-J., Zuend, A., and O’Dowd, C.: Sea-spray regulates sulfate  
cloud droplet activation over oceans, *npj Climate and Atmospheric Science*, 3, 14, <https://doi.org/10.1038/s41612-020-0116-2>, publisher:  
Nature Publishing Group, 2020.
- 765 Glassmeier, F., Hoffmann, F., Johnson, J. S., Yamaguchi, T., Carslaw, K. S., and Feingold, G.: Aerosol-cloud-climate cooling overestimated  
by ship-track data, *Science*, 371, 485–489, <https://doi.org/10.1126/science.abd3980>, publisher: American Association for the Advance-  
ment of Science, 2021.
- Gordon, H., Field, P. R., Abel, S. J., Barrett, P., Bower, K., Crawford, I., Cui, Z., Grosvenor, D. P., Hill, A. A., Taylor, J., Wilkinson, J.,  
Wu, H., and Carslaw, K. S.: Development of aerosol activation in the double-moment Unified Model and evaluation with CLARIFY  
770 measurements, *Atmospheric Chemistry and Physics*, 20, <https://doi.org/10.5194/acp-20-10997-2020>, publisher: Copernicus GmbH, 2020.
- Goren, T., Choudhury, G., Kretzschmar, J., and McCoy, I.: Co-variability drives the inverted-V sensitivity between liquid water path and  
droplet concentrations, *Atmospheric Chemistry and Physics*, 25, 3413–3423, <https://doi.org/10.5194/acp-25-3413-2025>, publisher: Coper-  
nicus GmbH, 2025.
- Gryspeerd, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., Dipu, S., Unglaub, C., Gettelman, A., and Christensen, M.: Constraining  
775 the aerosol influence on cloud liquid water path, *Atmospheric Chemistry and Physics*, 19, 5331–5347, <https://doi.org/10.5194/acp-19-5331-2019>, publisher: Copernicus GmbH, 2019a.

- Gryspeerdt, E., Smith, T. W. P., O’Keeffe, E., Christensen, M. W., and Goldsworth, F. W.: The Impact of Ship Emission Controls Recorded by Cloud Properties, *Geophysical Research Letters*, 46, <https://doi.org/10.1029/2019GL084700>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2019GL084700>, 2019b.
- 780 Gryspeerdt, E., Mülmenstädt, J., Gettelman, A., Malavelle, F. F., Morrison, H., Neubauer, D., Partridge, D. G., Stier, P., Takemura, T., Wang, H., Wang, M., and Zhang, K.: Surprising similarities in model and observational aerosol radiative forcing estimates, *Atmospheric Chemistry and Physics*, 20, <https://doi.org/10.5194/acp-20-613-2020>, 2020.
- Gryspeerdt, E., Goren, T., and Smith, T. W. P.: Observing the timescales of aerosol–cloud interactions in snapshot satellite images, *Atmospheric Chemistry and Physics*, 21, <https://doi.org/10.5194/acp-21-6093-2021>, 2021.
- 785 Gryspeerdt, E., Glassmeier, F., Feingold, G., Hoffmann, F., and Murray-Watson, R. J.: Observing short-timescale cloud development to constrain aerosol–cloud interactions, *Atmospheric Chemistry and Physics*, 22, <https://doi.org/10.5194/acp-22-11727-2022>, publisher: Copernicus GmbH, 2022.
- Han, Q., Rossow, W. B., Zeng, J., and Welch, R.: Three Different Behaviors of Liquid Water Path of Water Clouds in Aerosol–Cloud Interactions, *Journal of the Atmospheric Sciences*, 59, [https://doi.org/10.1175/1520-0469\(2002\)059<0726:TDBOLW>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<0726:TDBOLW>2.0.CO;2), 2002.
- 790 Haynes, J. M., L’Ecuyer, T. S., Stephens, G. L., Miller, S. D., Mitrescu, C., Wood, N. B., and Tanelli, S.: Rainfall retrieval over the ocean with spaceborne W-band radar, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/10.1029/2008JD009973>, <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2008JD009973>, 2009.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., 795 Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, [\\_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3803](https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3803), 2020.
- Hobbs, P. V., Garrett, T. J., Ferek, R. J., Strader, S. R., Hegg, D. A., Frick, G. M., Hoppel, W. A., Gasparovic, R. F., Russell, L. M., 800 Johnson, D. W., O’Dowd, C., Durkee, P. A., Nielsen, K. E., and Innis, G.: Emissions from Ships with respect to Their Effects on Clouds, *Journal of the Atmospheric Sciences*, 57, [https://doi.org/10.1175/1520-0469\(2000\)057<2570:EFSWRT>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<2570:EFSWRT>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2000.
- Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K., Joyce, R., Xie, P., and Yoo, S.-H.: NASA global precipitation measurement (GPM) integrated multi-satellite retrievals for GPM (IMERG), Algorithm theoretical basis document (ATBD) version, 4, 30, 2015.
- 805 Igel, A. L., Igel, M. R., and Heever, S. C. v. d.: Make It a Double? Sobering Results from Simulations Using Single-Moment Microphysics Schemes, *Journal of the Atmospheric Sciences*, 72, 910–925, <https://doi.org/10.1175/JAS-D-14-0107.1>, publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2015.
- Jing, X. and Suzuki, K.: The Impact of Process-Based Warm Rain Constraints on the Aerosol Indirect Effect, *Geophysical Research Letters*, 45, 10,729–10,737, <https://doi.org/10.1029/2018GL079956>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL079956](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL079956), 2018.
- 810 Kabatas, B., Menzel, W. P., Bilgili, A., and Gumley, L. E.: Comparing Ship-Track Droplet Sizes Inferred from Terra and Aqua MODIS Data, *Journal of Applied Meteorology and Climatology*, 52, 230–241, <https://doi.org/10.1175/JAMC-D-11-0232.1>, 2013.

- Kang, L., Marchand, R. T., Wood, R., and McCoy, I. L.: Coalescence Scavenging Drives Droplet Number Concentration in Southern Ocean Low Clouds, *Geophysical Research Letters*, 49, e2022GL097819, <https://doi.org/10.1029/2022GL097819>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2022GL097819](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2022GL097819), 2022.
- 815 Kato, S., Sun-Mack, S., Miller, W. F., Rose, F. G., Chen, Y., Minnis, P., and Wielicki, B. A.: Relationships among cloud occurrence frequency, overlap, and effective thickness derived from CALIPSO and CloudSat merged cloud vertical profiles, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012277>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JD012277](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JD012277), 2010.
- 820 Kato, S., Rose, F. G., Sun-Mack, S., Miller, W. F., Chen, Y., Rutan, D. A., Stephens, G. L., Loeb, N. G., Minnis, P., Wielicki, B. A., Winker, D. M., Charlock, T. P., Stackhouse Jr., P. W., Xu, K.-M., and Collins, W. D.: Improvements of top-of-atmosphere and surface irradiance computations with CALIPSO-, CloudSat-, and MODIS-derived cloud and aerosol properties, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016050>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2011JD016050](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2011JD016050), 2011.
- 825 Khairoutdinov, M. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Monthly Weather Review*, 128, 229–243, [https://doi.org/10.1175/1520-0493\(2000\)128<0229:ANCPPI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0229:ANCPPI>2.0.CO;2), publisher: American Meteorological Society Section: *Monthly Weather Review*, 2000.
- L'Ecuyer, T. S. and Stephens, G. L.: An Estimation-Based Precipitation Retrieval Algorithm for Attenuating Radars, *Journal of Applied Meteorology and Climatology*, 41, 272–285, [https://doi.org/10.1175/1520-0450\(2002\)041<0272:AEBPRA>2.0.CO;2](https://doi.org/10.1175/1520-0450(2002)041<0272:AEBPRA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of Applied Meteorology and Climatology*, 2002.
- 830 Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., Chipperfield, M. P., Pickering, S. J., and Johnson, C. E.: Description and evaluation of GLOMAP-mode: a modal global aerosol microphysics model for the UKCA composition-climate model, *Geoscientific Model Development*, 3, 519–551, <https://doi.org/10.5194/gmd-3-519-2010>, publisher: Copernicus GmbH, 2010.
- Manshausen, P., Watson-Parris, D., Christensen, M. W., Jalkanen, J.-P., and Stier, P.: Invisible ship tracks show large cloud sensitivity to aerosol, *Nature*, 610, <https://doi.org/10.1038/s41586-022-05122-0>, number: 7930 Publisher: Nature Publishing Group, 2022.
- 835 McCoy, D. T., Bender, F. A.-M., Grosvenor, D. P., Mohrmann, J. K., Hartmann, D. L., Wood, R., and Field, P. R.: Predicting decadal trends in cloud droplet number concentration using reanalysis and satellite data, *Atmospheric Chemistry and Physics*, 18, 2035–2047, <https://doi.org/10.5194/acp-18-2035-2018>, publisher: Copernicus GmbH, 2018.
- McMichael, L. A., Schmidt, M. J., Wood, R., Blossey, P. N., and Patel, L.: Exploring ship track spreading rates with a physics-informed Langevin particle parameterization, *Geoscientific Model Development*, 17, 7867–7888, <https://doi.org/10.5194/gmd-17-7867-2024>, publisher: Copernicus GmbH, 2024.
- 840 McMichael, L. A., Blossey, P. N., Wood, R., and Doherty, S. J.: Investigation of Ship-Induced Mesoscale Circulation Mechanics and Aerosol Plume Spreading Rates, *Geophysical Research Letters*, 52, e2025GL116904, <https://doi.org/10.1029/2025GL116904>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2025GL116904](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2025GL116904), 2025.
- 845 Michibata, T., Suzuki, K., Sato, Y., and Takemura, T.: The source of discrepancies in aerosol–cloud–precipitation interactions between GCM and A-Train retrievals, *Atmospheric Chemistry and Physics*, 16, <https://doi.org/10.5194/acp-16-15413-2016>, publisher: Copernicus GmbH, 2016.
- Morrison, H. and Gettelman, A.: A New Two-Moment Bulk Stratiform Cloud Microphysics Scheme in the Community Atmosphere Model, Version 3 (CAM3). Part I: Description and Numerical Tests, *Journal of Climate*, 21, 3642–3659, <https://doi.org/10.1175/2008JCLI2105.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2008.
- 850

- Morrison, H., Thompson, G., and Tatarskii, V.: Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes, *Monthly Weather Review*, 137, 991–1007, <https://doi.org/10.1175/2008MWR2556.1>, publisher: American Meteorological Society Section: Monthly Weather Review, 2009.
- Mülmenstädt, J. and Feingold, G.: The Radiative Forcing of Aerosol–Cloud Interactions in Liquid Clouds: Wrestling and Embracing Uncertainty, *Current Climate Change Reports*, 4, 23–40, <https://doi.org/10.1007/s40641-018-0089-y>, 2018.
- 855 Nenes, A. and Seinfeld, J. H.: Parameterization of cloud droplet formation in global climate models, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002911>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2002JD002911>, 2003.
- Noone, K. J., Johnson, D. W., Taylor, J. P., Ferek, R. J., Garrett, T., Hobbs, P. V., Durkee, P. A., Nielsen, K., Öström, E., O’Dowd, C., Smith, M. H., Russell, L. M., Flagan, R. C., Seinfeld, J. H., De Bock, L., Van Grieken, R. E., Hudson, J. G., Brooks, I., Gasparovic, R. F., and Pockalny, R. A.: A Case Study of Ship Track Formation in a Polluted Marine Boundary Layer, *Journal of the Atmospheric Sciences*, 57, [https://doi.org/10.1175/1520-0469\(2000\)057<2748:ACSOST>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<2748:ACSOST>2.0.CO;2), 2000.
- 860 Penner, J. E., Dong, X., and Chen, Y.: Observational evidence of a change in radiative forcing due to the indirect aerosol effect, *Nature*, 427, <https://doi.org/10.1038/nature02234>, 2004.
- Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and Microphysical Products: Collection 6 Updates and Examples From Terra and Aqua, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, conference Name: IEEE Transactions on Geoscience and Remote Sensing, 2017.
- 865 Possner, A., Zubler, E., Lohmann, U., and Schär, C.: Real-case simulations of aerosol–cloud interactions in ship tracks over the Bay of Biscay, *Atmospheric Chemistry and Physics*, 15, <https://doi.org/10.5194/acp-15-2185-2015>, 2015.
- 870 Possner, A., Wang, H., Wood, R., Caldeira, K., and Ackerman, T. P.: The efficacy of aerosol–cloud radiative perturbations from near-surface emissions in deep open-cell stratocumuli, *Atmospheric Chemistry and Physics*, 18, <https://doi.org/10.5194/acp-18-17475-2018>, publisher: Copernicus GmbH, 2018.
- Possner, A., Eastman, R., Bender, F., and Glassmeier, F.: Deconvolution of boundary layer depth and aerosol constraints on cloud water path in subtropical stratocumulus decks, *Atmospheric Chemistry and Physics*, 20, 3609–3621, <https://doi.org/10.5194/acp-20-3609-2020>, publisher: Copernicus GmbH, 2020.
- 875 Prabhakaran, P., Hoffmann, F., and Feingold, G.: Effects of intermittent aerosol forcing on the stratocumulus-to-cumulus transition, *Atmospheric Chemistry and Physics*, 24, 1919–1937, <https://doi.org/10.5194/acp-24-1919-2024>, publisher: Copernicus GmbH, 2024.
- Quaas, J., Boucher, O., and Lohmann, U.: Constraining the total aerosol indirect effect in the LMDZ and ECHAM4 GCMs using MODIS satellite data, *Atmospheric Chemistry and Physics*, 6, <https://doi.org/10.5194/acp-6-947-2006>, publisher: Copernicus GmbH, 2006.
- 880 Rosenfeld, D.: Suppression of Rain and Snow by Urban and Industrial Air Pollution, *Science*, 287, 1793–1796, <https://doi.org/10.1126/science.287.5459.1793>, publisher: American Association for the Advancement of Science, 2000.
- Rothenberg, D. and Wang, C.: Metamodeling of Droplet Activation for Global Climate Models, *Journal of the Atmospheric Sciences*, 73, 1255–1272, <https://doi.org/10.1175/JAS-D-15-0223.1>, publisher: American Meteorological Society Section: Journal of the Atmospheric Sciences, 2016.
- 885 Salter, S., Sortino, G., and Latham, J.: Sea-going hardware for the cloud albedo method of reversing global warming, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 366, 3989–4006, <https://doi.org/10.1098/rsta.2008.0136>, publisher: Royal Society, 2008.

- Satoh, M., Stevens, B., Judt, F., Khairoutdinov, M., Lin, S.-J., Putman, W. M., and Düben, P.: Global Cloud-Resolving Models, *Current Climate Change Reports*, 5, 172–184, <https://doi.org/10.1007/s40641-019-00131-0>, 2019.
- 890 Segrin, M. S., Coakley, J. A., and Tahnk, W. R.: MODIS Observations of Ship Tracks in Summertime Stratus off the West Coast of the United States, *Journal of the Atmospheric Sciences*, 64, 4330–4345, <https://doi.org/10.1175/2007JAS2308.1>, publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2007.
- Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteorology and Atmospheric Physics*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- 895 Shipway, B. and Abel, S.: Analytical estimation of cloud droplet nucleation based on an underlying aerosol population, *Atmospheric Research*, 96, 344–355, <https://doi.org/https://doi.org/10.1016/j.atmosres.2009.10.005>, 15th International Conference on Clouds and Precipitation, 2010.
- Shipway, B. J. and Hill, A. A.: Diagnosis of systematic differences between multiple parametrizations of warm rain microphysics using a kinematic framework, *Quarterly Journal of the Royal Meteorological Society*, 138, <https://doi.org/10.1002/qj.1913>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.1913>, 2012.
- 900 Skofronick-Jackson, G., Petersen, W. A., Berg, W., Kidd, C., Stocker, E. F., Kirschbaum, D. B., Kakar, R., Braun, S. A., Huffman, G. J., Iguchi, T., Kirstetter, P. E., Kummerow, C., Meneghini, R., Oki, R., Olson, W. S., Takayabu, Y. N., Furukawa, K., and Wilheit, T.: The Global Precipitation Measurement (GPM) Mission for Science and Society, *Bulletin of the American Meteorological Society*, 98, 1679–1695, <https://doi.org/10.1175/BAMS-D-15-00306.1>, publisher: American Meteorological Society Section: *Bulletin of the American Meteorological Society*, 2017.
- 905 Smith, C. J., Kramer, R. J., Myhre, G., Alterskjær, K., Collins, W., Sima, A., Boucher, O., Dufresne, J.-L., Nabat, P., Michou, M., Yukimoto, S., Cole, J., Paynter, D., Shiogama, H., O’Connor, F. M., Robertson, E., Wiltshire, A., Andrews, T., Hannay, C., Miller, R., Nazarenko, L., Kirkevåg, A., Olivie, D., Fiedler, S., Lewinschal, A., Mackallah, C., Dix, M., Pincus, R., and Forster, P. M.: Effective radiative forcing and adjustments in CMIP6 models, *Atmospheric Chemistry and Physics*, 20, 9591–9618, <https://doi.org/10.5194/acp-20-9591-2020>, publisher: Copernicus GmbH, 2020.
- 910 Smith, T., Jalkanen, J., Anderson, B., Corbett, J., Faber, J., Hanayama, S., O’Keeffe, E., Parker, S., Johansson, L., Aldous, L., Raucci, C., Traut, M., Ettinger, S., Nelissen, D., Lee, D., Ng, S., Agrawal, A., Winebrake, J., Hoen, M., Chesworth, S., and Pandey, A.: Third IMO Greenhouse Gas Study 2014, International Maritime Organization, United Kingdom, ISBN First presented to Marine Environment Protection Committee 67 as MEPC 67/INF.3, 2015.
- 915 Stephens, G. L. and Haynes, J. M.: Near global observations of the warm rain coalescence process, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL030259>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2007GL030259](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2007GL030259), 2007.
- Stephens, G. L., Vane, D. G., Tanelli, S., Im, E., Durden, S., Rokey, M., Reinke, D., Partain, P., Mace, G. G., Austin, R., L’Ecuyer, T., Haynes, J., Lebsock, M., Suzuki, K., Waliser, D., Wu, D., Kay, J., Gettelman, A., Wang, Z., and Marchand, R.: CloudSat mission: Performance and early science after the first year of operation, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2008JD009982>, [\\_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2008JD009982](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2008JD009982), 2008.
- 920 Tanelli, S., Durden, S. L., Im, E., Pak, K. S., Reinke, D. G., Partain, P., Haynes, J. M., and Marchand, R. T.: CloudSat’s Cloud Profiling Radar After Two Years in Orbit: Performance, Calibration, and Processing, *IEEE Transactions on Geoscience and Remote Sensing*, 46, 3560–3573, <https://doi.org/10.1109/TGRS.2008.2002030>, 2008.
- Taufour, M., Vié, B., Augros, C., Boudevillain, B., Delanoë, J., Delautier, G., Ducrocq, V., Lac, C., Pinty, J.-P., and  
925 Schwarzenböck, A.: Evaluation of the two-moment scheme LIMA based on microphysical observations from the HyMeX

- campaign, *Quarterly Journal of the Royal Meteorological Society*, 144, 1398–1414, <https://doi.org/10.1002/qj.3283>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3283>, 2018.
- 930 Taylor, J. P. and Ackerman, A. S.: A case-study of pronounced perturbations to cloud properties and boundary-layer dynamics due to aerosol emissions, *Quarterly Journal of the Royal Meteorological Society*, 125, 2643–2661, <https://doi.org/10.1002/qj.49712555915>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.49712555915>, 1999.
- Terai, C. R., Bretherton, C. S., Wood, R., and Painter, G.: Aircraft observations of aerosol, cloud, precipitation, and boundary layer properties in pockets of open cells over the southeast Pacific, *Atmospheric Chemistry and Physics*, 14, 8071–8088, <https://doi.org/10.5194/acp-14-8071-2014>, publisher: Copernicus GmbH, 2014.
- 935 Tippett, A., Gryspeerdt, E., Manshausen, P., Stier, P., and Smith, T. W. P.: Weak liquid water path response in ship tracks, *Atmospheric Chemistry and Physics*, 24, 13 269–13 283, <https://doi.org/10.5194/acp-24-13269-2024>, publisher: Copernicus GmbH, 2024.
- Toll, V., Christensen, M., Gassó, S., and Bellouin, N.: Volcano and Ship Tracks Indicate Excessive Aerosol-Induced Cloud Water Increases in a Climate Model, *Geophysical Research Letters*, 44, 12 492–12 500, <https://doi.org/10.1002/2017GL075280>, 2017.
- Toll, V., Christensen, M., Quaas, J., and Bellouin, N.: Weak average liquid-cloud-water response to anthropogenic aerosols, *Nature*, 572, 51–55, <https://doi.org/10.1038/s41586-019-1423-9>, 2019.
- 940 Tselioudis, G., Rossow, W. B., Jakob, C., Remillard, J., Tropf, D., and Zhang, Y.: Evaluation of Clouds, Radiation, and Precipitation in CMIP6 Models Using Global Weather States Derived from ISCCP-H Cloud Property Data, *Journal of Climate*, 34, 7311–7324, <https://doi.org/10.1175/JCLI-D-21-0076.1>, publisher: American Meteorological Society Section: Journal of Climate, 2021.
- Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment (1967)*, 8, 1251–1256, [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- 945 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOP>2.0.CO;2), publisher: American Meteorological Society Section: Journal of the Atmospheric Sciences, 1977.
- 950 Walters, D., Boutle, I., Brooks, M., Melvin, T., Stratton, R., Vosper, S., Wells, H., Williams, K., Wood, N., Allen, T., Bushell, A., Copsey, D., Earnshaw, P., Edwards, J., Gross, M., Hardiman, S., Harris, C., Heming, J., Klingaman, N., Levine, R., Manners, J., Martin, G., Milton, S., Mittermaier, M., Morcrette, C., Riddick, T., Roberts, M., Sanchez, C., Selwood, P., Stirling, A., Smith, C., Suri, D., Tennant, W., Vidale, P. L., Wilkinson, J., Willett, M., Woolnough, S., and Xavier, P.: The Met Office Unified Model Global Atmosphere 6.0/6.1 and JULES Global Land 6.0/6.1 configurations, *Geoscientific Model Development*, 10, 1487–1520, <https://doi.org/10.5194/gmd-10-1487-2017>, publisher: Copernicus GmbH, 2017.
- 955 Wang, H. and Feingold, G.: Modeling Mesoscale Cellular Structures and Drizzle in Marine Stratocumulus. Part II: The Microphysics and Dynamics of the Boundary Region between Open and Closed Cells, *Journal of the Atmospheric Sciences*, 66, 3257–3275, <https://doi.org/10.1175/2009JAS3120.1>, publisher: American Meteorological Society Section: Journal of the Atmospheric Sciences, 2009.
- Wood, R.: Assessing the potential efficacy of marine cloud brightening for cooling Earth using a simple heuristic model, *Atmospheric Chemistry and Physics*, 21, <https://doi.org/10.5194/acp-21-14507-2021>, publisher: Copernicus GmbH, 2021.
- 960 Wood, R. and Bretherton, C. S.: Boundary Layer Depth, Entrainment, and Decoupling in the Cloud-Capped Subtropical and Tropical Marine Boundary Layer, *Journal of Climate*, 17, 3576–3588, [https://doi.org/10.1175/1520-0442\(2004\)017<3576:BLDEAD>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<3576:BLDEAD>2.0.CO;2), publisher: American Meteorological Society Section: Journal of Climate, 2004.

- Wood, R., Leon, D., Lebsock, M., Snider, J., and Clarke, A. D.: Precipitation driving of droplet concentration variability in marine low clouds, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD018305>, \_eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2012JD018305>, 2012.
- 965 Xiong, W., Tang, G., Wang, T., Ma, Z., and Wan, W.: Evaluation of IMERG and ERA5 Precipitation-Phase Partitioning on the Global Scale, *Water*, 14, 1122, <https://doi.org/10.3390/w14071122>, publisher: Multidisciplinary Digital Publishing Institute, 2022.
- Yuan, T., Song, H., Oreopoulos, L., Wood, R., Meyer, K., Crawford, A., Smith, W., and Eastman, R.: Analyses of Virtual Ship-Tracks Systematically Underestimate Aerosol-Cloud Interactions Signals, *Geophysical Research Letters*, 52, e2024GL114356, <https://doi.org/10.1029/2024GL114356>, \_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2024GL114356>, 2025.
- 970 Zelinka, M. D., Andrews, T., Forster, P. M., and Taylor, K. E.: Quantifying components of aerosol-cloud-radiation interactions in climate models, *Journal of Geophysical Research: Atmospheres*, 119, 7599–7615, <https://doi.org/10.1002/2014JD021710>, \_eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2014JD021710>, 2014.
- Zhang, J. and Feingold, G.: Distinct regional meteorological influences on low-cloud albedo susceptibility over global marine stratocumulus regions, *Atmospheric Chemistry and Physics*, 23, <https://doi.org/10.5194/acp-23-1073-2023>, publisher: Copernicus GmbH, 2023.
- 975 Zhang, J., Zhou, X., Goren, T., and Feingold, G.: Albedo susceptibility of northeastern Pacific stratocumulus: the role of covarying meteorological conditions, *Atmospheric Chemistry and Physics*, 22, <https://doi.org/10.5194/acp-22-861-2022>, publisher: Copernicus GmbH, 2022.