

Observations of stratospheric streamers and frozen-in anticyclones in aerosol extinction

Christian Löns¹, Ronald Eixmann², Christine Pohl^{3,*}, Alexei Rozanov³, and Christian von Savigny¹

¹Institute of Physics, University of Greifswald, 17489 Greifswald, Germany

²Leibniz-Institute of Atmospheric Physics at the University of Rostock, 18225 Kühlungsborn, Germany

³Institute of Environmental Physics, University of Bremen, 28359 Bremen, Germany

* now at: Department of Physics, Lund University, Lund, 22100, Sweden

Correspondence: Christian Löns (christian.loens@uni-greifswald.de)

Abstract. When the polar vortex meanders and shifts towards the equator, air masses from the tropics and subtropics can be transported towards the pole in so-called tropical-subtropical streamers. These large-scale structures are areas of low potential vorticity and high pressure, containing dry air with a high ozone mixing ratio. The presence of these streamers can also be seen in changes in stratospheric optical properties. Satellite instruments such as OMPS-LP measuring the limb scattering of these aerosols are capable of observing an increase in the aerosol extinction coefficient in the mid-stratosphere at the edge of the vortex. The high spatial sampling of the limb instrument ensures that the trajectory of the streamer can be accurately monitored. Following a displacement and deformation of the vortex, aerosol transport to high latitudes occurred in the Northern Hemisphere in spring 2017. The additional stratospheric aerosol mass of around 1,000 t at an altitude of 24–38 km remained at middle and high latitudes for just under a month this in that year. This aerosol mass increase resulted in an estimated 70 % rise in the total mass within this altitude range at high latitudes. Frozen-in anticyclones, in which low latitude air is trapped in the circulation at high latitudes after the end of the polar vortex, can also be observed in the aerosol extinction coefficient. The observation of a particularly long-lived anticyclone in 2005, which is visible in the aerosol-extinction coefficient, is presented. This is the first study documenting streamer events and frozen-in anticyclones in stratospheric aerosols.

1 Introduction

The stratospheric part of the polar vortex (hereinafter referred to as the polar vortex) is a large area of cold rotating air. The westerly winds that shape it extend from just above the tropopause into the mesosphere (Vaugh et al., 2017). It forms in the autumn months in the polar regions after the onset of the polar night. ~~Here, the atmosphere is not heated by solar radiation.~~ A strong temperature difference arises between the equator and the pole, which creates a pressure gradient with a region of lower pressure at the pole (Schoeberl and Hartmann, 1991). The air flowing towards the pole is deflected eastwards by the Coriolis force and a circumpolar vortex is formed (Schoeberl and Newman, 2015). The maximum of these intensified westerly winds is a good approximation of the edge of the vortex (Vaugh and Polvani, 2010). Within the vortex, wind speeds are minimal. While the Antarctic vortex flows very evenly around the pole, the Arctic polar vortex is more strongly characterized by waves and disturbances (van Loon et al., 1973; Vaugh et al., 2017).

So-called sudden stratospheric warmings (SSWs) are ~~an important and a~~ frequently occurring zonal indicator of such disturbances. Their observation was first published in Scherhag (1952). During SSWs, planetary waves propagate vertically from the troposphere and transport westward angular momentum upwards (Baldwin and Dunkerton, 2001). These waves often break in the surf zone in the mid-latitudes of the stratosphere and can lead to a weakening or collapse of the vortex (Matsuno, 1971; Baldwin et al., 2021). The cold air near the pole descends and the polar temperature in the lower and mid-stratosphere rises abruptly due to the warmer air coming from above and the associated adiabatic heating. This results in a change in the usually westerly wind. There is no unambiguous standard definition for SSWs ~~and definitions often differ in the selection of characteristic criteria, threshold values or are defined for a particular latitude or altitude (Butler et al., 2015; Butler and Gerber, 2018). A frequently used and easily applicable definition was proposed (Butler et al., 2015; Butler and Gerber, 2018). According to a frequently used one~~ by Charlton and Polvani (2007), ~~which is employed in this study. According to this definition,~~ major SSWs are defined by the reversal of the mean zonal wind at 60° N and 10 hPa. ~~A distinction often made is between minor SSWs (Minor SSWs are usually defined by a~~ weakening of the wind ~~) and major SSWs (reversal of the wind direction) (Labitzke, 1981; Andrews et al., 1987). Major SSWs can be divided into two types: displacement events in which the polar vortex is deflected and splitting events in which the vortex splits into two or more vortices (Schoeberl and Hartmann, 1991; Charlton and Polvani, 2007). As shown by Butler et al. (2015), the statistical frequency of SSWs depends strongly on the respective definition. According to Charlton and Polvani (2007), in (McInturff, 1978; Labitzke, 1981). In~~ the Northern Hemisphere major mid-winter SSWs occur about six times per decade ~~(Charlton and Polvani, 2007),~~ minor SSWs about once per year (Gogoi et al., 2023). A final warming marks the end of the polar vortex season.

In order to understand the transport processes in the ~~winter~~ stratosphere, Plumb (2002) suggests dividing it into three meridional regions: the tropics, the surf zone and the polar vortex. ~~To this end, Plumb (2002) examined the relationships between long-lived tracers and reference tracers in these regions.~~ The subtropics largely serve as a transport barrier to isolate the tropics from the surf zone ~~(Plumb, 1996), as could be shown by the tape recorder signature in the tropical lower stratosphere (Mote et al., 1996)(Plumb, 1996; Mote et al., 1996).~~ As described in McIntyre and Palmer (1984), the surf zone is the region into which air masses can be transported vertically by breaking planetary waves. Similarly ~~but stronger than the subtropical barrier (Krüger et al., 2005),~~ the edge of the polar vortex serves as a transport barrier between the surf zone and the centre of the polar vortex. ~~It~~ ~~The edge of the polar vortex~~ can be characterized by the maximum gradient of potential vorticity (Nash et al., 1996) or the maximum of the circumpolar westerly winds (Waugh and Polvani, 2010). ~~It is stronger than the subtropical barrier (Krüger et al., 2005).~~

Small-scale mixing processes through transport barriers are referred to as filaments. Filaments are finger-like structures with a vertical extent of less than 2.5 km (Reid and Vaughan, 1991) and a horizontal extent of less than 1000 km (Reid et al., 1993). Such structures, which were initially found mainly in ozone profiles and are sometimes also referred to as laminae (Dobson, 1973) ~~, were explained by Appenzeller and Holton (1997) by differential advection of air masses of different origins. Filaments occur particularly frequently near the tropopause of the subtropics and mid-latitudes as well as at the edge of the polar vortex (Dobson, 1973; Reid and Vaughan, 1991; Appenzeller and Holton, 1997).~~

Sometimes the ~~The~~ term 'stratospheric streamer' is ~~sometimes~~ used to describe ~~these small-scale~~ vertical transport processes from the stratosphere to the troposphere (~~Appenzeller and Davies, 1992; Wernli and Sprenger, 2007~~)~~(e.g. Appenzeller and Davies, 1992; Wernli and Sprenger, 2007)~~. In this ~~paper study~~, the term stratospheric streamer is employed to denote large-scale tongue-like structures that transport tropical-subtropical or polar air masses towards mid-latitudes in the mid-stratosphere, as described in Offermann et al. (1999) and Krüger et al. (2005).

~~An early descriptive~~ ~~An early~~ study of such transport processes was published by Leovy et al. (1985). Using satellite measurements from the ~~Nimbus-7~~ Limb Infrared Monitor of the Stratosphere (LIMS), tongues of ozone-rich air were observed after the polar vortex at 10 hPa in the Northern Hemisphere was deflected from the pole in early 1979. These ~~ozone-rich~~ tongues, characterized by their spiral trajectory towards the pole, ~~also~~ exhibited low potential vorticity.

The occurrence of streamers has been shown to be primarily associated with the disturbance of the polar vortex, leading to the edge of the polar vortex reaching low latitudes and tongues of tropical air being drawn towards mid-latitudes (Offermann et al., 1999). It has been demonstrated that this process can occur ~~in both hemispheres especially~~ during the winter months ~~in both hemispheres~~, although it is less pronounced in the Southern Hemisphere. This is due to the typically smaller amplitude of planetary waves (Randel et al., 1993).

~~These streamers~~ ~~Streamers~~ have a width ranging from 500 to 2000 km, a length exceeding 10,000 to 20,000 km, and persist for a duration of 1 to 3 weeks (Krüger et al., 2005). Observations and model simulations have identified two primary pathways over the Atlantic and over Asia, respectively (Offermann et al., 1999; Krüger et al., 2005). Such meridional transport has been observed for various trace gases, including nitric acid (HNO_3) (~~Offermann et al., 1999~~)~~(e.g. Kouker et al., 1999; Offermann et al., 1999; Engel et al., 2005)~~, nitrous oxide (N_2O) (~~Randel et al., 1993; Offermann et al., 1999~~)~~(e.g. Randel et al., 1993; Offermann et al., 1999; Khosrawi, 2005)~~, ozone (O_3) (~~Leovy et al., 1985; Hocke et al., 2017~~)~~(e.g. Leovy et al., 1985; Hocke et al., 2017; Wüst et al., 2025)~~ and water vapour (H_2O) (~~Randel et al., 1993~~)~~(e.g. Randel et al., 1993; Hocke et al., 2017)~~.

Another phenomenon associated with the polar vortex, whereby air from lower latitudes is compressed and can be held at high latitudes for several months, is a frozen-in anticyclone (FrIAC). The first documentation of this phenomenon was by Manney et al. (2006), who observed the FrIAC using MLS data of N_2O , H_2O and ozone following the final warming of the Arctic polar vortex in March 2005. The vertical extent of this FrIAC was found to be approximately 25–45 km, and characteristic tropical signals of high N_2O and low H_2O were still identifiable ~~several months later~~ in August 2005. FrIACs were also described for other years, such as in 2003 (Lahoz et al., 2007), 2007 (Thiéblemont et al., 2011) and 2011 (Allen et al., 2013).

The ~~shape of the vortex also influences the global distribution of stratospheric aerosol~~ ~~observability of streamers and FrIACs~~ ~~has not yet been demonstrated for aerosols~~. Satellite instruments can be used to retrieve the aerosol extinction coefficient, with the two most common measurement methods being limb scattering and solar occultation.

In context of solar occultation the aerosol extinction coefficient is a measure of the attenuation of solar radiation due to its scattering or absorption by particles, such as stratospheric sulfate particles. ~~Solar occultation, in contrast to limb scattering measurements, allows for the retrieval of the extinction coefficient without the necessity of prior assumptions regarding the composition and size of the particles (e.g. McCormick, 1982).~~ In the context of limb scattering ~~the~~ aerosol extinction

is ~~predominantly an retrieved exploiting the~~ amplification of the measured signal due to ~~stronger~~ scattering by aerosols. It has the advantage of considerably higher temporal and spatial coverage (e.g. Loughman et al., 2004; von Savigny et al., 2015) and is therefore well suited for global observations of aerosol extinction.

The ~~present work is based on data products of the Institute of Environmental Physics of the University of Bremen for aerosol extinction derived from limb scattering for the instruments SCIAMACHY V3.0 (von Savigny et al., 2015; Rieger et al., 2018; Sofieva et al. and OMPS-LP V2.1 (Rozanov et al., 2024), hereafter OMPS-UBr. Aerosol extinction data from 2002 to 2023 is available for these two data products. In order to also estimate the size information of the observed aerosols, measurements and derived~~ particle size distribution (PSD) parameters from the Stratospheric Aerosol and Gas Experiment (SAGE) III solar occultation instrument on the ISS are used.

The extinction coefficient above the Junge layer decreases with altitude ~~-.The and the~~ presence of liquid aerosol particles at an altitude of 30 km to 35 km is constrained by the positive temperature gradient in the stratosphere ~~-.As the temperature increases, the and the increasing~~ saturation vapour pressure of H_2SO_4 ~~increases by several orders of magnitude. This process has been shown to impede. This impedes~~ the development of new H_2SO_4 particles ~~that are situated~~ above an altitude of around 36 km ~~-.The formation of these particles has been observed to be exclusive to regions where the partial pressure exceeds the saturation vapour pressure (Hamill et al., 1977). Furthermore, an increase in saturation vapour pressure is accompanied by an increase in the rate of evaporation. (Hamill et al., 1977).~~ An increasing extinction coefficient with altitude can therefore be explained either by dynamic effects or by material coming from above, potentially meteoric or space debris.

~~A~~ Beside that, a decrease in the extinction coefficient with increasing latitude towards the pole is to be expected. This is due to the fact that sulfate aerosols form in the tropical tropopause region and most active volcanoes are also located in the subduction zones of the tropics and subtropics. Meridional mixing is limited by the dynamic transport barriers in the subtropics and at the edge of the polar vortex.

The ~~present work is based on data products of the Institute of Environmental Physics of the University of Bremen for aerosol extinction derived from limb scattering for the instruments SCIAMACHY V3.0 (von Savigny et al., 2015; Rieger et al., 2018; Sofieva et al. and OMPS-LP V2.1 (Rozanov et al., 2024), hereafter OMPS-UBr. Aerosol extinction data from 2002 to 2023 is available for these two data products. In order to also estimate the size information of the observed aerosols, measurements and derived~~ particle size distribution (PSD) parameters from the Stratospheric Aerosol and Gas Experiment (SAGE) III instrument on the ISS are used.

~~The~~ transport of tropical aerosol towards mid-latitudes was observed in satellite measurements following the eruption of Mt. Pinatubo ~~in~~ (McCormick and Veiga, 1992; Trepte et al., 1993). The poleward transport of tropical aerosols at an altitude of 15–30 km was also described in Harvey et al. (1999). Here, SAM II measurements of the aerosol extinction coefficient revealed a reduced surface aerosol density (SAD) in the polar vortex during the winter months from 1978 to 1991 (McCormick and Veiga, 1992; Trepte et al., 1993), while the SAD in the Aleutian High was increased. The relationship between the extinction coefficient and the polar vortex was also demonstrated by Thomason et al. (2007) using the potential vorticity (PV) as a tracer for the vortex. Above an altitude of 20 km, a negative correlation between the extinction coefficient and PV is was observed, indicating that as PV increases, the extinction coefficient decreases. Therefore, the aerosol extinction within the vortex is minimal. ~~In this study, in addition to the~~

low-extinction-coefficient within the polar-vortex, an increased extinction-coefficient is observed at high and medium latitudes at the edge of the vortex.

130 The aim of ~~the work~~ this study is to describe ~~anomalies in streamers and FrIACs in the~~ aerosol extinction coefficient ~~in the~~ high latitudes that are related to a changing or collapsing polar-vortex, in particular the meridional transport of aerosol towards the pole in the Northern Hemisphere. ~~The period of spring as case studies of two events in 2017 stands out because the transport at the edge of the vortex is clearly visible and tropical air remains at middle to high latitudes for a month.~~

and 2005. Section 2 describes the data sets used of the SCIAMACHY, OMPS-UBr and SAGE III/ISS satellite measurements
135 and the ECMWF ERA5 reanalysis data. Section ~~3.1 presents results of the zonal-mean~~ 3 describes the evolution of the aerosol extinction coefficient in the Northern Hemisphere ~~for the period of OMPS measurements at different altitudes and latitude bins. Sections 3.2 and 3.4 describe the development of the polar-vortex and the aerosol extinction coefficient in early in spring 2017 and 2005. In Sect. 3.2 the methodology used to estimate the transported sulfate aerosol mass is explained. The interpretation and validity~~ 2005 and compares it with meteorological reanalysis data. The validity and interpretation of the results are discussed
140 in Sections 4 and 5.

2 Instruments and Methodology

2.1 Limb scattering instruments

The study of aerosol extinction associated with the polar vortex requires high spatial and temporal resolution. Therefore, the focus of this work is the investigation of stratospheric aerosol data sets obtained from remote sensing with a high spatial sampling,
145 such as limb scattering instruments or atmospheric lidars. However, spaceborne lidars are not well-suited to the investigation of background conditions in the mid-stratosphere in this study due to relatively low signal-to-noise ratios. Instruments in limb viewing geometry measure the intensity of limb-scattered solar radiation, by repeating line-of-sight measurements at different tangent height altitudes from approximately 0 to 100 km, or simultaneously over all heights. Limb scattering instruments are confronted with the challenge of receiving both reflected and scattered light from the Earth and atmosphere along its line of
150 sight. This complicates the methodology, with each measurement depending not only on the aerosol extinction but also on the conditions of the atmosphere and the aerosol microphysical properties such as aerosol size and composition. In addition, as is usually the case in remote sensing, aerosol extinction becomes increasingly difficult to measure with height due to a decreasing signal-to-noise ratio, and the error in the extinction coefficient increases.

The SCanning Imaging Absorption spectroMeter for Atmospheric ChartographY (SCIAMACHY) instrument was launched
155 on 1 March 2002 on board the European Environmental Satellite (Envisat) into a sun-synchronous orbit at an altitude of approximately 800 km. It was operational from August 2002 to April 2012, during which time it measured, amongst other things, the scattered solar radiation in limb and nadir geometry. A grating spectrometer recorded the radiation in eight wavelength channels in the spectral range 214–2386 nm with a spectral resolution of 0.2–1.5 nm (Bovensmann et al., 1999). The retrieval of aerosol extinction from SCIAMACHY measurements was described in von Savigny et al. (2015) and further developed in

160 Rieger et al. (2018). Level 2 version 3.0 aerosol extinction data from the University of Bremen were used for this work ~~as~~
described in Sofieva et al. (2024).

The Ozone Mapping and Profiler Suite Limb Profiler (OMPS-LP) was first launched on 28 October 2011 on board the
Suomi National Polar-orbiting Partnership (S-NPP) satellite as one of three OMPS instruments into a sun-synchronous orbit
at an altitude of approximately 800 km. The instrument has been in operation since February 2012 and remains active to
165 the present day. Additionally, OMPS-LP was launched on board the NOAA-21 satellite in December 2022. The OMPS-LP
instrument is a prism spectrometer covering a spectral range of 280-1000 nm with an increasingly coarser spectral resolution
towards infrared wavelengths (Jaross et al., 2014). OMPS-LP has three slits separated horizontally by 4.25°. The measured
tangent points are 250 km apart. The solar radiation scattered by the atmospheric limb is simultaneously recorded from the
ground up to an altitude of 80-100 km.

170 ~~The One advantage of the~~ OMPS-LP data product from the University of Bremen is ~~used for this work. One advantage of~~
~~this data set is~~ that the aerosol extinction coefficient is usually derived to a greater height than the derived aerosol product of
NASA. The retrieval of vertical profiles of aerosol extinction from limb radiance measurements is described in Rozanov et al.
(2024). In addition, atmospheric pressure profiles from the OMPS-LP data product of the Goddard Earth Sciences Data and
Information Services Centre are used (Taha, 2020) to interpolate the aerosol extinction coefficient to specific pressure levels.
175 These are forward processing data from the NASA GSFC Global Modelling Assimilation Office (GMAO) at the nearest grid
cell to each LP event, which are interpolated at the corresponding measurement time.

SCIAMACHY and OMPS-LP have approximately 14 orbits per day and it takes 3–4 days to achieve full global coverage.
SCIAMACHY covers latitudes up to 60° N throughout the year. Latitudes north of 75° N are covered from the end of February
until October. The coverage extends to a maximum of 84.4° N. OMPS-LP covers latitudes above 60° N from the end of
180 January until November and latitudes north of 75° N from mid-March until October. The coverage extends to a maximum of
81.4° N. For both data sets, only retrieved profiles with a vertical resolution greater better than 7 km ~~are filtered out to ensure~~
~~high-quality data at all heights~~were used to ensure high data quality at all altitudes.

2.2 Particle size distribution parameters

Estimating the transported sulfate aerosol mass requires assumptions about the size distribution of the aerosols responsible
185 for the increase in extinction. Previous studies have sought to derive information regarding the size of particles using limb
scattering measurements (Rieger et al., 2014; Pohl et al., 2024). However, the retrievals are dependent on many assumptions
about scattering processes in the atmosphere and aerosol properties. Therefore, the stratospheric aerosol data set from SAGE
III/ISS was used, which has been used ~~several times previously~~ to determine information about the PSD (Wrana et al., 2021;
Knepp et al., 2024). In addition to the reduced spatial and temporal coverage, SAGE III/ISS observes high latitudes later in the
190 year than the Limb instruments.

SAGE III on ISS is a solar occultation instrument (Cisewski et al., 2014) that was launched on 19 February 2017 and started
collecting data in June 2017. This instrument is an update of the SAGE III instrument on Meteor-3M (Thomason et al., 2010)
and is very similar in functionality. Version 6.0 of the SAGE III/ISS data set (NASA/LARC/SD/ASDC, 2025) is used for this

work, which contains PSD parameters derived according to the procedure described in Knepp et al. (2024). Their solution
195 algorithm for the PSD only works if the extinction coefficient is positive in all channels used in the algorithm.

~~In addition to the reduced spatial and temporal coverage, SAGE III/ISS observes high latitudes later in the year than the~~
~~Limb instruments.~~ A common challenge faced by remote sensing instruments is the signal-to-noise ratio, which decreases with
altitude. The inversion method employed for SAGE III/ISS permits negative values of the extinction coefficient, resulting in an
increase in the proportion of negative coefficients with altitude. For SAGE III/ISS, at short wavelengths such as $\lambda = 449$ nm,
200 this results in a greater proportion of negative extinction coefficients at altitudes above approximately 32 km, relative to positive
extinction coefficients, thereby causing the mean extinction coefficient to become negative. This phenomenon has the potential
to introduce biases or make a retrieval in the mid-stratosphere infeasible. Therefore, the robustness of the derived PSD also
decreases with altitude.

2.3 ~~ECMWF~~ERA5 reanalysis

205 The ECMWF ERA5 ~~Reanalysis~~ meteorological reanalysis data is used ~~to verify for this study to investigate~~ the atmospheric
dynamics in specific periods (Hersbach et al., 2020). ERA5 integrates model data and observations to create a uniform global
data set. The data set "ERA5 hourly data on pressure levels from 1940 to present" ~~is used for this study.~~, which is available on
a $0.25^\circ \times 0.25^\circ$ grid is used (Copernicus Climate Change Service, 2018).

ERA5 data products are widely used and generally provide a good representation of transport processes in the atmosphere.
210 Several studies have validated stratospheric transport in ERA5. Age of air analyses were used to investigate the representation
of the Brewer-Dobson circulation in ERA5 and to compare it with observational data (Ploeger et al., 2021). Here, in the lower
and middle stratosphere in the Northern Hemisphere, there are indications that ERA5 is at the upper edge of the observational
uncertainty in terms of mean age of air and that the Brewer-Dobson Circulation therefore appears to be low biased, also in
comparison to the previous version ERA-Interim (Diallo et al., 2021). ERA5 was also used to verify the upward transport
215 of greenhouse gases and pollution during the Asian monsoons via age of air and trajectory analyses (Vogel et al., 2024).
Comparison with in situ data showed that the transport into the lower stratosphere was well represented.

In this study, the parameters geopotential, potential vorticity, temperature, specific humidity, ozone and northward and east-
ward wind are considered at a pressure level of 10 hPa, i.e. at an altitude of approximately 30 km, during the polar vortex
season.

220 3 Results

3.1 Zonal mean aerosol extinction coefficient

Figure 1 shows the zonally averaged aerosol extinction coefficient (AEC) at 869 nm from OMPS-UBr in three distinct latitude
ranges in the Northern Hemisphere at altitudes of 25.5, 28.5, 31.5, 34.5 and 37.5 km.




Figure 1. Weekly averaged aerosol extinction coefficient (AEC) at 869 nm from OMPS-UBr for five altitudes in the Northern Hemisphere (NH). The subtropics are defined from 20–40° N, the mid-latitudes from 40–60° N and the high latitudes from 60–80° N. The first half of 2017 is shaded within the dashed lines in (a–e) and zoomed in on the right-hand side in (f–j).

During the OMPS period, ~~the tropical volcanic eruptions from 2012 to the present, there were several volcanic eruptions that~~
225 ~~had an impact on the stratosphere (e.g. Carn, 2025; Kovilakam et al., 2025). In the tropics, these were particularly the eruptions~~
~~of Kelut (2014), Ambae (2018), Ulawun (2019), Soufriere St. Vincent (2021) and Hunga (2022) are expected to influence the~~
~~aerosol extinction coefficient.~~ Moreover, two eruptions in mid-latitudes had an impact on the stratosphere. These were the
eruptions of Calbuco in the Southern Hemisphere in 2015 and of Raikoke in the Northern Hemisphere in 2019. ~~The impact~~
~~of volcanic eruptions in the stratosphere is closely associated with the altitude of the plume. Notably, the eruption of Hunga~~
230 ~~was the only one for which a significant aerosol plume was documented to reach the mid-stratosphere. This plume could be~~
~~observed up to~~ For the altitudes and latitude regions shown in Fig. 1, only a direct impact from the Hunga eruption is to be
expected, whose plume was observed at an altitude of approximately 26 km (Duchamp et al., 2023). The impact of this eruption
can be identified in Fig. 1 e. After the eruption, the aerosol extinction coefficient reaches its maximum at an altitude of 25.5 km

in the northern subtropics during the ~~OMPS period.~~ second half of 2022. The other eruptions referenced are not expected to exert a direct ~~influence upon the mid-stratosphere~~ significant influence in the Northern Hemisphere at the altitudes shown.

In general, Fig. 1 shows a decrease in the aerosol extinction coefficient towards the pole. The meridional gradient and also the minimum extinction coefficient at high latitudes are most pronounced in late summer and early autumn. In addition, the aerosol extinction coefficient in the middle and high latitudes shows a stronger variance in the polar vortex season compared to the summer months.

The occurrence of a local streamer event can be indicated by a jump in the weekly zonally averaged extinction coefficient at an altitude of 31.5 km, as observed in Fig. 1 c.

In the northern high latitudes, indicated by the blue line in Fig. 1, a sudden increase can be observed in the mid-stratosphere in the weekly zonally averaged extinction coefficient in the shaded area in the first half of 2017. The signal is strongest at an altitude of 31.5 km. At this altitude, the aerosol extinction coefficient exhibited a sixfold increase in spring 2017, marking the highest value recorded at this altitude within the northern high latitudes during the OMPS measurement period. At altitudes of 28.5 km and 34.5 km, the zonally averaged extinction coefficient increased by more than threefold. At an altitude of 28.5 km, the background level of the aerosol extinction coefficient is higher in other years than during early 2017 due to ~~a higher background aerosol level in those years~~ natural variations. At an altitude of 34.5 km, the peak in the extinction coefficient in 2017 is only exceeded by the signature of the Chelyabinsk meteor (Gorkavyi et al., 2013; Rieger et al., 2014), which exploded in the mid-stratosphere four years earlier in the spring of 2013. This is also the only period during the OMPS mission in which aerosol extinction is greater at middle and high latitudes than in the subtropics. An increase in the aerosol extinction coefficient in spring 2017 can also be observed, albeit to a lesser extent, at altitudes of 25.5 km and 37.5 km.

In the northern mid-latitudes, indicated by the red line in Fig. 1, two distinct ~~yet less pronounced~~ peaks can be identified in March and April 2017 at altitudes of 28.5 km, 31.5 km and 34.5 km, which are slightly less pronounced than in the high latitudes. At an altitude of 31.5 km, a more than twofold increase in the aerosol extinction coefficient is observed in these two peaks. The two peaks can be observed before and after the strongest increase in the high latitudes.

3.2 Aerosol extinction in the polar vortex season in early 2017

In Fig. 2, the zonal mean eastward wind at 60° N and the temperature from 65–90° N at 10 hPa for early 2017 are shown. The representation is based on Baldwin et al. (2021). Two ~~minor warming events~~ events, which are classified as minor warmings, are recorded at the beginning and end of February, which is consistent with the observations in Eswaraiah et al. 2019, Gogoi et al. 2023 and Li et al. 2023. A sudden increase in temperature can be observed at high latitudes. In addition, there is a weakening of the wind, but no reversal of the wind direction at 60° N and 10 hPa.

Figure 3 shows the development of the weekly averaged extinction coefficient interpolated to 10 hPa, i.e. approximately 30 km altitude for the initial 16 weeks of 2017. The figure shows the aerosol extinction coefficient at 869 nm in bins of 5° latitude and 15° longitude in the Northern Hemisphere.

At the beginning of 2017, the aerosol extinction coefficient in the mid-latitudes is subject to considerable variation. Figures 3 b and 3 e show tongue-like patterns of increased aerosol extinction coefficient, extending from the subtropics to the mid-




Figure 2. SSW criteria for spring 2017. The zonal mean eastward wind at 60° N is represented in red and the mean temperature from 65–90° N at 10 hPa is shown in black. Two minor SSWs in February are marked by dotted lines.

latitudes from west to central Asia. The increase in aerosol extinction that occurred subsequent to the second of these tongues in early February coincides with the first minor warming in 2017 (see Fig. 2), which caused a weakening and deflection of the vortex. At this time, the centre of the polar vortex was located at approximately 70° N. The increase in the aerosol extinction coefficient cannot be traced further in the following weeks.

On 25 February, a late minor SSW occurred (see Fig. 2). At a pressure level of 10 hPa, this had the effect of a temperature increase at the North Pole of more than 30 K. This phenomenon resulted in a weakening and separation of the polar vortex. A shift of the vortex edge into the subtropics can be observed from around 1 March. The vortex was subsequently elongated and displaced off the pole.

For a better understanding of the dynamic of the polar vortex, Fig. 4 presents in the lower five rows ERA5 reanalysis data for a four-week period in March [and April](#) 2017 following the second minor warming. The parameters under consideration are the wind, the potential vorticity with contour levels of geopotential height, the specific humidity, the ozone mass mixing ratio and the temperature, all at 10 hPa. The vortex centre itself is associated with low wind speed, geopotential height and ozone mixing ratio, and high potential vorticity and specific humidity. After the two minor SSWs, the polar vortex is no longer associated with low temperatures.

Looking at the first column of Fig. 4 for the week of 5–11 March, an increase in wind speed towards the pole of the now very uneven vortex can be observed. Following the displacement of the polar vortex towards the subtropics, the transportation of aerosols from the subtropics towards the pole can be deduced from the increased extinction coefficient at the edge of the vortex over Asia, see Figs. 4 a and 4 e. Subtropical air is transported along the edge of the polar vortex, and an area of high geopotential height or relative high pressure is formed, see Fig. 4 i. An anticyclone forms around the area of high geopotential height. Areas of an increased aerosol extinction coefficient correlate with a low potential vorticity and the air parcel is dry and initially ozone-rich.




Figure 3. Polar stereographic maps of the weekly averaged aerosol extinction coefficient at 869 nm from OMPS-UBr interpolated to 10 hPa in early 2017 in bins of 5° latitude and 15° longitude for the entire Northern Hemisphere, with 0° E as the central longitude line from bottom to centre.

As of 16 March, the ~~material is aerosols are~~ observed to depart from the edge of the polar vortex. Concurrently, an anticyclone is established over Greenland (70° N, 45° W), see Figs. 4 g and 4 k. ~~This structure persists and maintaining the material's position for nearly three weeks~~ The aerosols are kept at the same longitude in the anticyclone for about three weeks, while slowly moving towards lower latitudes. Subsequently, from approximately 6 April onwards, it can still be observed over the North Atlantic (45° N, 45° W). It is notable that the air parcel remains dry for the entire time that the signal is observable in the aerosol extinction coefficient. In the ozone mass mixing at this pressure level, the air parcel is no longer richer in ozone

295 than the ambient air after a period of one week, at which point rapid mixing occurs. There is no evident correlation between increased aerosol extinction and temperature.

After 19 March, the formation of another streamer can be observed in a substantial area of low PV, low specific humidity and high ozone mass mixing ratio at 30° N over Asia. After 26 March, Figs. 4 h, 4 l and 4 p show the transport of this subtropical air to mid-latitudes. As can be seen in 4 l, a further area of high geopotential height has formed over the Bering Sea.

300 Figure 5 shows a comparison of the vertical profiles of the aerosol extinction coefficient at middle to high latitudes, both inside and outside the anticyclone, and in the tropics over central and northern Africa at the beginning of March before the start of the streamer event. To make this comparison, profiles from the same geographical region (55–70° N and 20–45° W) were averaged for different weeks before and during the streamer event, as in Figs. 3 j and 3 l. To compare these with the vertical profile of the aerosol extinction coefficient from the region in which the streamer likely originated, measurements before the
305 start of the streamer event (see Fig. 3 i) from the tropics to subtropics (0–30° N, 0–45° E) were averaged.

~~The edge of the tropical air parcel which is transported within a high-pressure area towards high latitudes creates a strong transport barrier between the anticyclone and the ambient air.~~ At an altitude of ~~24~~26–38 km the vertical profiles in the tropics and three weeks later in the anticyclone differ only slightly. This ~~finding suggests that the mixing process is minimal for the streamer. This transport barrier is also the reason why the increase in aerosol extinction can be observed for so long~~suggests that there was little mixing of aerosols with the ambient air along the transport path. The high-pressure area, which transports tropical air towards higher latitudes, creates a barrier for mixing between the anticyclone and the surrounding air. The agreement between ~~anticyclone and~~ the profiles inside the anticyclone and from the subtropics is greatest at 30.5 km, the altitude at which the strongest signal from the streamer, i.e. the biggest percentage increase in the aerosol extinction coefficient, is visible. Up to an altitude of around 26 km, there is a clear distinction in the profiles of subtropical and high-latitude regions.
315 Conversely, a high degree of similarity is observed between the profiles both within and outside the anticyclone up to this altitude. At an altitude of 32.5 km, the aerosol extinction coefficient increases by a factor of 27 from the background level to the streamer.

In addition to 2017, anticyclones in the mid-latitudes are also evident in the aerosol extinction coefficient in other years, e.g. 2015 and 2022, albeit to a lesser extent (not shown).

320 As of 13 March, a weak polar vortex streamer becomes discernible in the reanalysis data~~through an increased potential vorticity and specific humidity~~, see the second column of Fig. 4. This is evident below 45° -N over Southern Europe ~~, see Figs. in an increased potential vorticity, see Fig. 4 j and, an increased specific humidity, see Fig. 4 n. This is not evident in the,~~ and also a reduced aerosol extinction coefficient, see Fig. 4 b.

3.3 Estimation of the transported aerosol mass in 2017

325 For the estimation of the transported aerosol mass, it is important to note that measurements with SAGE III on ISS began in June 2017, precluding the possibility of direct overlap with the observed signatures in 2017 and earlier. Consequently, the PSD parameters for comparable atmospheric conditions in subsequent years must be used. To this end, a streamer event from 2018 was selected. It began in February 2018, a few weeks earlier in the season than in 2017, and was observable with SAGE III/ISS

in early March. Below 31 km altitude, the streamer had a comparable zonal influence on the aerosol extinction coefficient as in
 330 2017 and was also briefly located in the high latitudes, see Fig. 1. Since there was no volcanic eruption between the two events
 in 2017 and 2018 that could have affected the stratosphere, the background aerosol level remained consistent. The present
 study focuses on the streamer event in early 2017 due to its greater vertical extent, its stronger influence on high latitudes, and
 the fact that there was negligible mixing taking place due to the anticyclone.

Figure 6 shows vertical profiles of the two streamers in 2017 and 2018 in the mid-latitudes (45–55° N) for different longitude
 335 regions as measured with OMPS-LP and SAGE III/ISS. Both streamer profiles originate from a time approximately 3–4 weeks
 after the start of the streamer event. Up to an altitude of 34 km, there is very good agreement between the retrieved aerosol
 extinction coefficient with OMPS-UBr and SAGE III/ISS in 2018. This indicates that the streamer event in 2018 can be
 observed with both instruments. The vertical profile from 2017 shows a higher average aerosol extinction over the entire
 altitude range of the streamer. This can be partially explained by the fact that the streamer in 2017 is more compressed within
 340 an anticyclone, thereby concentrating the aerosols in a smaller area.

Nevertheless, this period in March 2018 and this region appears to be the best possible match with the streamer event in
 2017 in the measurement period of SAGE III. For this selection, the mean median radius and geometric standard deviation of
 a monomodal log-normal distribution are obtained for every altitude step. The Mie code of the University of Oxford is used to
 calculate the extinction cross section for these PSD parameters.

345 The time period spanning 14 days from 19 March to 1 April is selected for the estimation of the transported sulfate aerosol
 mass in spring 2017. This is due to the aerosol layer exhibiting minimal zonal or meridional movement during this interval,
 thereby facilitating its observation over the North Atlantic. The mean extinction coefficient at 869 nm is calculated for each
 height level from 24.5–38.5 km for bins-grid boxes of size 3° × 3°. Furthermore, the mean extinction coefficient from 0–270° E,
 i.e. in the range in which no increase in the aerosol extinction coefficient due to transported material can be observed, is
 350 calculated for each latitude step. Bins-Grid boxes are designated as an anomaly if the extinction coefficient is higher than its
 latitudinal mean plus three times the standard deviation and if an increase is also observed in the surrounding binsgrid boxes.

For each such anomaly, the absolute number of particles $N(z)$ at height z is calculated. The height z is defined as the centre
 height of an altitude layer spanning 1 km. The volume V of the respective bin-grid box and the number density $N_0(z)$ are
 used to calculate the absolute number of particles $N(z)$. The number density is derived from the additional aerosol extinction
 355 attributable to the streamer and the PSD parameters. The mass of a log-normally distributed aerosol population can then be
 calculated using the 3rd moment of the log-normal distribution (Grainger, 2023)(Seinfeld and Pandis, 2016).

$$m(z) = \frac{4}{3} \pi \cdot N(z) \cdot r_{\text{med}}^3 \cdot \rho(z, T) \cdot \exp\left(\frac{9}{2} \ln(\sigma_g)^2\right) \quad (1)$$

In this equation, $m(z)$ denotes the total mass of the aerosol population at height z and $\rho(T)$ represents the density of the
 aerosol particles as a function of temperature T at height z . For the mass density ρ , the measured mass densities of sulfuric
 360 acid for different mass fractions and temperatures of Myhre et al. (1998) are interpolated for an H₂SO₄ mass fraction of 75 %
 and for the respective GMAO temperature at this altitude.

The aerosol mass that has been transported in 2017, as calculated with averaged PSD parameters for measurements of the streamer event in 2018, is estimated to be approximately 0.99 kt in the altitude range of 24.5–38.5 km, where the aerosol extinction signal of this air parcel is observable.

365 ~~Due to the strong mixing barrier, it can be assumed that the mass changes are negligible along the transport path.~~ Since the aerosol extinction coefficient changes only slightly along the transport path of the streamer, see Fig. 5, it is assumed that changes in mass along the path are also negligible. For the high latitudes defined as 60–90° N, the average PSD parameters from SAGE III/ISS for the same altitude range result in a total mass of 1.34 kt for background conditions. This means that the streamer event in the week of 12–18 March causes a zonal increase in the total aerosol mass in the altitude range of
370 24.5–38.5 km by about 74 %. At an altitude of 30.5 km, the aerosol mass in the high latitudes zonally increases by about 260 % during the same period. On a local scale, the mass can increase by more than one order of magnitude.

3.4 Aerosol extinction in the polar vortex season in early 2005

Manney et al. (2006) observed a strong and long-lasting FrIAC in 2005. This subsection will examine whether this can also be observed in the aerosol extinction coefficient. A notable peak in the aerosol extinction coefficient in the mid-stratosphere
375 during the SCIAMACHY measurement period was observed in spring 2005. Figure 7 shows the aerosol extinction coefficient at 750 nm in the Northern Hemisphere interpolated at an altitude of 30.5 km from spring to late-summer 2005. The northern winter of 2004/2005 was relatively cold with prolonged periods of temperatures below 200 K in the lower stratosphere and characterized by a strong and stable polar vortex (~~Sonkaew et al., 2013~~) (e.g. Sonkaew et al., 2013). On 13 March, there was an exceptionally early final warming, approximately one month before the historical median (Butler and Domeisen, 2021). As
380 can be seen in Figs. 7 a–d, there are some variations in March 2005 in the mid-latitudes. From the end of March/beginning of April an increase in the aerosol extinction coefficient at the pole is noticeable.

Figure 8 presents in the lower five rows reanalysis data for the four weeks from ~~mid-March~~ March to early April ~~following the final warming~~. Before the final warming on 13 March, a slightly increased aerosol extinction coefficient can already be observed in the mid-latitudes across a substantial geographical area extending from North America to the Pacific region, see Fig. 8 a.
385 This suggests that ~~tropical~~ subtropical air has already accumulated in a high-pressure area extending over a considerable region, see Fig. 8 i. In the weeks following the final warming, the region of low geopotential height of the collapsed polar vortex shifts towards mid-latitudes, see Fig. 8 j. Concurrently, the low-pressure area is compressed and shifts towards the pole until the end of March. By then, a strengthening anticyclone develops with low PV (Fig. 8 l), dry ozone-rich air (Figs. 8 p and 8 t) and high aerosol extinction (Fig. 8 d) over the Arctic Ocean near Siberia. The anticyclone shifts towards the pole and remains there
390 trapped in the circulation for a period of several months until August of the same year, see Fig. 7.

Similar structures to those seen in 2005 can be found in other years in the aerosol data sets examined, e.g. in 2006, 2007 and 2014–2014 (not shown).

4 ~~Discussion~~ Sources of error

395 It is conceivable that the observed signal is an artefact of the retrieval product or a specific measurement technique. However, the transport in 2017 can be observed in a very similar way in the other OMPS-LP retrieval products from NASA and the University of Saskatchewan. The FrIAC in 2005 can also be observed using aerosol extinction data collected by SCIAMACHY. This suggests that the observed phenomena are not artefacts of a single retrieval or instrument. However, both phenomena are detected using a similar measurement technique. For comparison with the aerosol extinction coefficient from a solar occultation instrument, a streamer event from 2018 was selected, which could be observed by both SAGE III/ISS and OMPS-LP, see Fig. 6.

400

Up to an altitude of 34 km, there is very good agreement between the two profiles of the different satellite instruments. The observation of increased aerosol extinction profiles up into the mid-stratosphere at mid-latitudes using two instruments with different measurement techniques and retrieval methods, indicates that this is a genuine aerosol extinction signal.

405 The likelihood of an error in retrieval due the effects of water vapour and ozone during the streamer event is negligible, as the two trace gases exhibit only very low absorption at the wavelength channels used. Increased ozone levels at an altitude of 30 km and above can only be observed during the first week of the streamer's presence. Additionally, the streamer is drier than the surrounding air.

410 The estimated amount of just below 1 kt depends heavily on the number density of the aerosols, which in turn depends on the assumptions made about the size distribution, i.e. the median radius and geometric standard deviation. For this estimation, the "weighted median PSD parameters" from the SAGE III/ISS dataset (Knepp et al., 2024) were employed for every altitude step when a streamer in 2018 was measured in mid-latitudes with SAGE III/ISS.

415 This is a very rough estimate. Therefore, the assumptions about the PSD parameters were varied. To vary the PSD parameters, a single averaged PSD combination from the altitude range of 25–30 km was applied to the entire altitude range. This is owing to the fact that PSD retrieval is more reliable at lower altitudes than at high altitudes. In addition, PSD parameters from 0–30° N from the entire SAGE III/ISS measurement period up to the Hunga eruption were averaged for each altitude step. However, both methods result only in a negligible deviation of about 5 % in aerosol mass.

420 Nevertheless, the estimate remains uncertain. No solar occultation measurements are available for spring 2017. The true PSD of aerosols above 30 km altitude is largely unknown and is subject to high bias and error when retrieved from SAGE measurements. In addition, there is general uncertainty about the true shape of the size distribution, which is assumed to be monomodal for the retrieval. However, if the true shape is different, e.g. bimodal, this can lead to significant deviations in the retrieval of PSD parameters (von Savigny and Hoffmann, 2020).

5 Discussion

Both described events in 2005 and 2017 can be explained by the combination of streamers, transporting tropical air into the middle or high latitudes and anticyclones holding material in these regions.

425 The exact extent and shape of the polar vortex and how far it reaches into the mid-latitudes appears to be decisive for the transport of aerosols towards the pole. In the majority of years in the observation period 2003–2023, a transport of tropical

air into the mid-latitudes and a variation in the aerosol extinction especially in mid-latitudes can be observed. The intensity depends largely on the background level in tropical aerosol extinction. The duration and stability of a streamer depend on the dynamics in mid-latitudes in connection with the polar vortex and ~~the formation and strength of the~~ whether the streamer is associated with an anticyclone, which, as in 2017, prevents mixing with the surrounding air.

During early 2017, two minor warmings occurred in February 2017. The impact of the former on the mesosphere was described in Eswaraiyah et al. (2020). A streamer event can also be observed in the stratospheric aerosol extinction coefficient following this first warming at the beginning of February, during which tropical air is transported to mid-latitudes. However, this signal is only observable for approximately one week, and the high latitudes are not adequately covered by OMPS-LP at the beginning of February. Consequently, the estimation of transport was only possible via surrounding latitudes on the outward or return transport from the pole. The reanalysis data indicates that no anticyclone is formed, thereby ensuring that the transported air parcel of the streamer does not remain concentrated at middle and high latitudes. Moreover, the signal in the specific humidity is no longer clearly observable after one week.

A notable characteristic of the polar vortex season 2017 is the comparatively late occurrence of a warming, i.e. the second minor warming, which was not also the final warming in the vortex season (Gogoi et al., 2023). This phenomenon enables the observation of the signal of an increased extinction coefficient above 75° N. The sequence of weakening, deflection and brief but repeated strengthening of the elongated vortex with high wind speeds in the direction of the pole and low speeds away from the pole ensures that ~~material aerosols~~ can accumulate at the pole. The vortex weakens once more, causing the material to remain in the mid-latitudes for several weeks. The formation of a high-pressure system, accompanied by the establishment of an anticyclone, results in the accumulation of material south of Greenland over the Atlantic.

Following the second warming, no correlation of low temperatures with the vortex or high temperatures with the streamer is discernible in the temperature profile. This exemplifies the significant weakening of the polar vortex, which did not regain its closed circular structure and strength following the two minor SSWs. In addition, the temperature gradient between the tropics and high latitudes is only marginal, which is why the transported air mass is no longer associated with increased temperatures.

~~The occurrence of tropical-subtropical streamers is not exclusive to a specific period during the vortex season~~ Tropical-subtropical streamers can occur at any time of the year, although streamer frequency is significantly higher during the polar vortex season (Offermann et al., 1999; Eyring et al., 2003). However, ~~as the year progresses, the Limb instrument's the~~ coverage of the high latitudes ~~diminishes, thereby impeding the comprehensive tracking of a structure's trajectory by the limb instruments used~~ decreases towards the end of the year, making it more difficult to track the course of a streamer.

It is important to note that, in contrast to the more prevalent occurrence of streamer events in the Northern Hemisphere, these events are less frequent in the Southern Hemisphere due to the presence of a more stable vortex (van Loon et al., 1973). The more-frequent observation of this phenomenon in the Northern Hemisphere may also be explained, by the elevated background level of the aerosol extinction coefficient in the mid-stratosphere in the southern mid-to-high latitudes. This results in a lower gradient of the aerosol extinction coefficient between the South pole and the tropics compared to the gradient between the North pole and the tropics. Consequently, the presence of streamer events in the Southern Hemisphere is less pronounced in the aerosol extinction coefficient. However, this observed difference in the gradient may also result from increased retrieval

uncertainties in the Southern Hemisphere (Chen et al., 2020). The scattering angles of aerosols are strongly dependent on latitude for OMPS-LP measurements and change during the orbit. The observed signal in the Southern Hemisphere originates, among other things, from aerosol scattering in the sideways and backward directions, while in the Northern Hemisphere, forward scattering predominates.

The observed ~~signals show~~ signal in 2017 shows good agreement with the characteristics expected from ~~model~~-runs for stratospheric streamers (~~Krüger et al., 2005~~) with the Freie Universität Berlin Climate Middle Atmosphere Model (FUB-CMAM) in Krüger et al. (2005). These also showed a higher probability of streamers occurring when the polar vortex shifted equatorward. The dimensions of the streamer in 2017, with a length of over 10,000 km in the development phase, a width of about 1,500 km and a vertical extent of about 14 km, correspond very well with the modelled extent of streamers. The duration of the event, spanning five weeks, is at the upper end of the expected time frame. This can be attributed to the anticyclone in 2017, which kept the tropical air in mid-latitudes for an extended period of time. The observation made in 2017 of a transport to 70–75° N also occurs in the model used in Krüger et al. (2005) in some years. In contrast to the model, no evidence of irreversible mixing at high latitudes is observed in 2017. The streamer in 2017 is located over East Asia, where the model predicts the maximum number of streamer events.

It is possible to observe different tracers of streamers and anticyclones for varying lengths of time. Ozone mixing, for instance, is a highly potent process that occurs in March 2017 a few days after the start of the streamer event. This mixing occurs more rapidly at higher altitudes (10 hPa) than at lower altitudes (20–30 hPa). At these lower altitudes, the air mass is still rich in O₃ even at the end of March. This observation is consistent with Thiéblemont et al. (2011), who also demonstrated in their description of a FrIAC in 2007 that the ozone mixing ratio decreases much faster at 10 hPa than at 20 hPa. The ozone signal is more strongly linked to gas chemistry and photochemistry than to mixing processes. The process of relaxation to lower ozone levels is even faster in spring, due to higher levels of solar radiation (Manney et al., 2006). Conversely, elevated aerosol levels and diminished levels of specific humidity persist within the anticyclone for a period of approximately one month. The correlation of anticyclone and dry air in the specific humidity is observable up to higher altitudes (5–20 hPa).

~~During the winter of 2016/17, the quasi-biennial oscillation (QBO) phase at 10 hPa shifted from westerly phase of the QBO (WQBO) to the easterly phase of QBO (EQBO). This means it was effectively neutral during the season. In general, signals from FrIACs (Thiéblemont et al., 2011) or streamers are more pronounced in years with an EQBO phase. This is due to a higher background level of aerosol extinction in the tropical mid-stratosphere in EQBO years, creating a greater gradient in the extinction coefficient between the tropics and the North Pole. The polar vortex tends to be stronger during the WQBO phase while being weaker and more unstable during the EQBO phase (Gray et al., 2018). Upward-propagating Rossby waves have a stronger orientation towards the polar vortex during the EQBO phase. A stronger interaction with the mean flow is associated with a higher probability of sudden stratospheric warming events (Baldwin et al., 2021). However, the QBO phase is not a definitive criterion, as Thiéblemont et al. (2016) show based on an observed FrIAC in 2013, while the QBO phase was westerly. The strong streamer event observed in 2017 during a neutral QBO phase also suggests that the QBO phase is not decisive for the occurrence and strength of streamer events. This is consistent with the observation in Waugh (1996) that the QBO phase has only a minor effect on winds poleward of 20° and thus on quasi-horizontal transport from the tropics.~~

The FrIAC in 2005 has already been described in detail based on observations (Manney et al., 2006) and modelling (Allen et al., 2011). As in Manney et al. (2006) for the trace gases N_2O and H_2O , the signature of the FrIAC can be observed in the extinction coefficient until August.

Similar to 2005, FrIACs of varying intensity can also be seen in the Northern Hemisphere in 2006, 2007 and 2014 (not shown), subsequent to the breakdown of the polar vortex. These circumpolar areas can be identified by a slight increase in the extinction coefficient at high latitudes.

6 Sources of error

6 Conclusions

~~It is conceivable that the observed signal is an artefact of retrieval or a specific measurement technique. The transport We have described a stratospheric tropical-subtropical streamer and a frozen-in anticyclone that can be observed in an increased aerosol extinction coefficient in two case studies in 2017 can be observed in a very similar way in the other OMPS-LP retrieval products from NASA and the University of Saskatchewan. The FrIAC in 2005 can also be observed using aerosol extinction data collected by SCIAMACHY. This suggests that the observed phenomena are not artefacts of a single retrieval or instrument. However, both phenomena are detected using a similar measurement technique. For comparison with and 2005. Streamers are horizontal transport processes by which atmospheric trace substances can be transported over long distances. Anticyclones can hold low latitude air in middle or high latitudes for periods ranging from weeks to several months. The signatures of these events were compared with ERA5 meteorological reanalysis data of the atmospheric state. While ozone mixes rapidly in both the streamer and the FrIAC, these two phenomena are evident in the aerosol extinction coefficient from a solar occultation instrument, a streamer event from 2018 was selected, which could be observed by both SAGE III/ISS and OMPS-LP, see Fig. 6 for as long as they are present in specific humidity.~~

~~Up to an altitude of 34 km, there is very good agreement between the two profiles of the different satellite instruments. The observation of increased aerosol extinction profiles up into the mid-stratosphere at mid-latitudes using two instruments with different measurement techniques and retrieval methods, indicates that this is a genuine aerosol extinction signal.~~

~~The likelihood of a error in retrieval due the effects of water vapour and ozone during the streamer event is negligible, as the two trace gases exhibit only very low absorption at the wavelength channels used. Increased ozone levels at an altitude of 30 km and above can only be observed during the first week of the streamer's presence. Additionally, Streamers are generally associated with disturbances in the polar vortex and a deflection of the vortex towards the subtropics. The streamer event of 2017 could be observed after a comparatively late second minor SSW in the streamer is drier than the surrounding air polar vortex season. This allowed tropical air to cross the subtropical transport barrier. FrIACs can occur both during the polar vortex season, such as in 2017 when the transported tropical air mass was held over the North Atlantic for several weeks, or after the polar vortex breakdown, as was the case in 2005. In 2005, a FrIAC was observed in an increased aerosol extinction coefficient in the high latitudes from March to August.~~

The estimated amount of just below 1 kt depends heavily on the number density of the aerosols, which in turn depends on the assumptions made about the size distribution, i.e. the median radius and geometric standard deviation. For this estimation, the “weighted median PSD parameters” from amount of sulphuric acid aerosols transported in the streamer towards middle and high latitudes in early 2017 was estimated to be approximately 1 kt, using aerosol properties derived from SAGE III/ISS dataset (Knepp et al., 2024) were employed for every altitude step when a streamer in 2018 was measured in mid-latitudes with SAGE III/ISS.

This is a very rough estimate. Therefore, the assumptions about the PSD parameters were varied. To vary the PSD parameters, a single averaged PSD combination from the altitude range of 25–30 km was applied to the entire altitude range. This is owing to the fact that PSD retrieval is more reliable at lower altitudes than at high altitudes. In addition, PSD parameters from 0–30° N from the entire SAGE III/ISS measurement period up to the Hunga eruption were averaged for each altitude step. However, both methods result only in PSD data for a negligible deviation comparable streamer event in 2018. A variation in the selected size parameters results in only a slight change in aerosol mass of about 5 % in aerosol mass.

Nevertheless, the estimate remains uncertain. No solar occultation measurements are available for spring 2017. The true PSD of aerosols above 30 km altitude is largely unknown and is subject to high bias and error when retrieved from SAGE measurements. In addition, there is general uncertainty about the true shape of the size distribution, which is assumed to be monomodal for the retrieval. However, if the true shape is different, e.g. bimodal, this can lead to significant deviations in the retrieval of PSD parameters (von Savigny and Hoffmann, 2020).

Stratospheric tropical-subtropical streamers are horizontal transport processes by which atmospheric trace substances can be transported over long distances. Streamers are associated with disturbances in the polar vortex and an elongation of vortex towards the subtropics. The 2017 event was the most significant poleward aerosol transport by streamers observed in the last twenty years. It almost doubled the aerosol mass in high latitudes for the altitude range of 24.5–38.5 km of the streamer. On a local scale, the aerosol extinction coefficient increased by a factor of 27.

Such streamers transport tropical air masses to middle and high latitudes, enabling them to cross the subtropical transport barrier. The transport of tropical aerosol-rich air to mid-latitudes is not an uncommon phenomenon and occurs to a small extent multiple times every year. However, the observation of a tropical-subtropical streamer reaching high latitudes is a rare occurrence. As shown for the first time in this study, the aerosol extinction coefficient also exhibits a high correlation with streamer events. The association of streamers with a low potential vorticity, high geopotential height, low specific humidity and high ozone mixing ratio has already been demonstrated in previous studies.

Satellite instruments such as OMPS only cover high latitudes at the beginning and end of winter. Therefore, the time window during which the transport of aerosols towards the pole can be tracked in detail is limited. The transport of increased aerosol concentrations in the mid-stratosphere to high latitudes is not expected to significantly impact radiative transfer.

Anticyclones can hold low latitude air in middle or high latitudes for periods ranging from weeks to several months. This phenomenon may occur both during the polar vortex season, such as In 2017, this is even evident in the zonal mean aerosol extinction at high latitudes. The correlation between streamer events and aerosol extinction coefficients was demonstrated for

~~the first time in this study. This correlation as well as the connection between streamers and anticyclones in 2017, or after the polar vortex breakdown, as was the case in 2005.~~

565 ~~The most significant poleward aerosol transport by streamers observed in the last twenty years occurred in March 2017, when about 1 kt of sulfuric acid aerosols were transported towards the middle and high latitudes. This almost doubled the aerosol mass in high latitudes for the altitude range of 24.5–38.5 km of the streamer. On a local scale, the aerosol extinction coefficient increased by a factor of 27.~~ provide new insights into the poleward transport and persistence of tropical air in the middle stratosphere at the end of the polar vortex season.

570 *Data availability.* The Level 2 aerosol extinction coefficient data products of SCIAMACHY V3.0 and OMPS-LP V2.1 from the Institute of Environmental Physics at the University of Bremen are available at <https://www.iup.uni-bremen.de/DataRequest> (Rozanov, 2024, login required, retrieved: 13 June 2025). The SAGE III/ISS data used within this study are available on NASA's Atmospheric Science Data Center at https://doi.org/10.5067/ISS/SAGEIII/SOLAR_BINARY_L2-V6.0 (NASA/LARC/SD/ASDC, 2025, retrieved: 13 August 2025). The ERA5 data used are available from the ECMWF at <https://doi.org/10.24381/cds.bd0915c6> (Copernicus Climate Change Service, 2018, retrieved: 575 26 February 2025).

Author contributions. CL handled the different datasets and wrote the initial text for the paper. CL, RE and CvS discussed the observations and interpreted the results. AR and CP provided the SCIAMACHY and OMPS-LP data. All authors contributed to the scientific discussion and improvement of the initial paper.

Competing interests. The authors declare that they have no conflict of interest.

580 *Acknowledgements.* This work was funded by the Deutsche Forschungsgemeinschaft (research grant O-MSP-LI, project no. 516357253). ~~We want to acknowledge support by the University of Greifswald and thank the Earth Observation Data Group at the University of Oxford for providing the IDL Mie routines used in this study.~~ The University of Bremen team was funded in part by the European Space Agency (ESA) via the CREST project, by the German Research Foundation (DFG) via the research unit VolImpact (grant no. FOR2820), and by the University of Bremen and state of Bremen. The authors gratefully acknowledge the computing time granted by the Resource Allocation 585 Board and provided for the supercomputers Lise and Emmy at NHR@ZIB and NHR@Göttingen as part of the NHR infrastructure. The calculations for this research were conducted using computing resources under project no. hbk00098. We want to acknowledge support by the University of Greifswald and thank the Earth Observation Data Group at the University of Oxford for providing the IDL Mie routines used in this study. Thanks to Professor Don Grainger for his 'Some Useful Formulae for Particle Size Distributions and Optical Properties' available at <https://eodg.atm.ox.ac.uk/user/grainger/research/aerosols.pdf>, last accessed: 25 September 2025.

- Allen, D. R., Douglass, A. R., Manney, G. L., Strahan, S. E., Krossschell, J. C., Trueblood, J. V., Nielsen, J. E., Pawson, S., and Zhu, Z.: Modeling the Frozen-In Anticyclone in the 2005 Arctic Summer Stratosphere, *Atmospheric Chemistry and Physics*, 11, 4557–4576, <https://doi.org/10.5194/acp-11-4557-2011>, 2011.
- Allen, D. R., Douglass, A. R., and Strahan, S. E.: The large-scale frozen-in anticyclone in the 2011 Arctic summer stratosphere, *Journal of Geophysical Research: Atmospheres*, 118, 2656–2672, <https://doi.org/10.1002/jgrd.50256>, 2013.
- Andrews, D. G., Holton, J. R., and Leovy, C. B.: Middle atmosphere dynamics, no. v. 40 in *International geophysics series*, Academic Press, Orlando, ISBN 978-0-12-058575-5 978-0-12-058576-2, 1987.
- Appenzeller, C. and Davies, H. C.: Structure of stratospheric intrusions into the troposphere, *Nature*, 358, 570–572, <https://doi.org/10.1038/358570a0>, 1992.
- Appenzeller, C. and Holton, J. R.: Tracer lamination in the stratosphere: A global climatology, *Journal of Geophysical Research: Atmospheres*, 102, 13 555–13 569, <https://doi.org/10.1029/97JD00066>, 1997.
- Baldwin, M. P. and Dunkerton, T. J.: Stratospheric Harbingers of Anomalous Weather Regimes, *Science*, 294, 581–584, <https://doi.org/10.1126/science.1063315>, 2001.
- Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., Domeisen, D. I. V., Garfinkel, C. I., Garny, H., Gerber, E. P., Hegglin, M. I., Langematz, U., and Pedatella, N. M.: Sudden Stratospheric Warmings, *Reviews of Geophysics*, 59, e2020RG000 708, <https://doi.org/10.1029/2020RG000708>, 2021.
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission Objectives and Measurement Modes, *Journal of the Atmospheric Sciences*, 56, 127–150, [https://doi.org/10.1175/1520-0469\(1999\)056<0127:SMOAMM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2), 1999.
- Butler, A. H. and Domeisen, D. I. V.: The wave geometry of final stratospheric warming events, *Weather Clim. Dyn.*, 2, 453–474, <https://doi.org/10.5194/wcd-2-453-2021>, 2021.
- Butler, A. H. and Gerber, E. P.: Optimizing the Definition of a Sudden Stratospheric Warming, *Journal of Climate*, 31, 2337–2344, <https://doi.org/10.1175/JCLI-D-17-0648.1>, 2018.
- Butler, A. H., Seidel, D. J., Hardiman, S. C., Butchart, N., Birner, T., and Match, A.: Defining Sudden Stratospheric Warmings, *Bulletin of the American Meteorological Society*, 96, 1913–1928, <https://doi.org/10.1175/BAMS-D-13-00173.1>, 2015.
- Carn, S.: Multi-Satellite Volcanic Sulfur Dioxide L4 Long-Term Global Database V4, <https://doi.org/10.5067/MEASURES/SO2/DATA405>, 2025.
- Charlton, A. J. and Polvani, L. M.: A New Look at Stratospheric Sudden Warmings. Part I: Climatology and Modeling Benchmarks, *Journal of Climate*, 20, 449–469, <https://doi.org/10.1175/JCLI3996.1>, 2007.
- Chen, Z., Bhartia, P. K., Torres, O., Jaross, G., Loughman, R., DeLand, M., Colarco, P., Damadeo, R., and Taha, G.: Evaluation of the OMPS/LP stratospheric aerosol extinction product using SAGE III/ISS observations, *Atmospheric Measurement Techniques*, 13, 3471–3485, <https://doi.org/10.5194/amt-13-3471-2020>, 2020.
- Cisewski, M., Zawodny, J., Gasbarre, J., Eckman, R., Topiwala, N., Rodriguez-Alvarez, O., Cheek, D., and Hall, S.: The Stratospheric Aerosol and Gas Experiment (SAGE III) on the International Space Station (ISS) Mission, in: *SPIE Remote Sensing*, edited by Meynart, R., Neeck, S. P., and Shimoda, H., p. 924107, Amsterdam, Netherlands, <https://doi.org/10.1117/12.2073131>, 2014.

- Copernicus Climate Change Service: ERA5 hourly data on pressure levels from 1940 to present, <https://doi.org/10.24381/CDS.BD0915C6>, last access: 26 February 2025, 2018.
- Diallo, M., Ern, M., and Ploeger, F.: The advective Brewer–Dobson circulation in the ERA5 reanalysis: climatology, variability, and trends, *Atmospheric Chemistry and Physics*, 21, 7515–7544, <https://doi.org/10.5194/acp-21-7515-2021>, 2021.
- 630 Dobson, G. M. B.: The laminated structure of the ozone in the atmosphere, *Quarterly Journal of the Royal Meteorological Society*, 99, 599–607, <https://doi.org/10.1002/qj.49709942202>, 1973.
- Duchamp, C., Wrana, F., Legras, B., Sellitto, P., Belhadji, R., and von Savigny, C.: Observation of the Aerosol Plume From the 2022 Hunga Tonga—Hunga Ha’apai Eruption With SAGE III/ISS, *Geophysical Research Letters*, 50, e2023GL105076, <https://doi.org/10.1029/2023GL105076>, 2023.
- 635 Eswaraiah, S., Kumar, K. N., Kim, Y. H., Chalapathi, G. V., Lee, W., Jiang, G., Yan, C., Yang, G., Ratnam, M. V., Prasanth, P. V., Rao, S., and Thyagarajan, K.: Low-latitude mesospheric signatures observed during the 2017 sudden stratospheric warming using the fuke meteor radar and ERA-5, *Journal of Atmospheric and Solar-Terrestrial Physics*, 207, 105352, <https://doi.org/10.1016/j.jastp.2020.105352>, 2020.
- Eyring, V., Dameris, M., Grewe, V., Langbein, I., and Kouker, W.: Climatologies of subtropical mixing derived from 3D models, *Atmospheric Chemistry and Physics*, 3, 1007–1021, <https://doi.org/10.5194/acp-3-1007-2003>, 2003.
- 640 Gogoi, J., Bhuyan, K., Sharma, S. K., Kalita, B. R., and Vaishnav, R.: A comprehensive investigation of Sudden Stratospheric Warming (SSW) events and upper atmospheric signatures associated with them, *Advances in Space Research*, 71, 3357–3372, <https://doi.org/10.1016/j.asr.2022.12.003>, 2023.
- Gorkavyi, N., Rault, D. F., Newman, P. A., Da Silva, A. M., and Dudorov, A. E.: New stratospheric dust belt due to the Chelyabinsk bolide, *Geophysical Research Letters*, 40, 4728–4733, <https://doi.org/10.1002/grl.50788>, 2013.
- 645 Grainger, R. G.: Some Useful Formulae for Particle Size Distributions and Optical Properties, <http://eodg.atm.ox.ac.uk/user/grainger/research/aerosols.pdf>, last access: 10 October 2025, 2023.
- Gray, L. J., Anstey, J. A., Kawatani, Y., Lu, H., Osprey, S., and Schenzinger, V.: Surface impacts of the Quasi Biennial Oscillation, *Atmospheric Chemistry and Physics*, 18, 8227–8247, <https://doi.org/10.5194/acp-18-8227-2018>, 2018.
- Hamill, P., Toon, O. B., and Kiang, C. S.: Microphysical Processes Affecting Stratospheric Aerosol Particles, *Journal of the Atmospheric Sciences*, 34, 1104–1119, [https://doi.org/10.1175/1520-0469\(1977\)034<1104:MPASAP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1104:MPASAP>2.0.CO;2), 1977.
- 650 Harvey, V. L., Hitchman, M. H., Pierce, R. B., and Fairlie, T. D.: Tropical aerosol in the Aleutian High, *Journal of Geophysical Research: Atmospheres*, 104, 6281–6290, <https://doi.org/10.1029/1998JD200094>, 1999.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., De Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 655 Hocke, K., Schranz, F., Maillard Barras, E., Moreira, L., and Kämpfer, N.: An Atlantic streamer in stratospheric ozone observations and SD-WACCM simulation data, *Atmospheric Chemistry and Physics*, 17, 3445–3452, <https://doi.org/10.5194/acp-17-3445-2017>, 2017.
- Jaross, G., Bhartia, P. K., Chen, G., Kowitt, M., Haken, M., Chen, Z., Xu, P., Warner, J., and Kelly, T.: OMPS Limb Profiler instrument performance assessment, *Journal of Geophysical Research: Atmospheres*, 119, 4399–4412, <https://doi.org/10.1002/2013JD020482>, 2014.

- Khosrawi, F.: Intercomparison between Lagrangian and Eulerian simulations of the development of mid-latitude streamers as observed by CRISTA, *Atmos. Chem. Phys.*, 2005.
- 665 Knepp, T. N., Kovilakam, M., Thomason, L., and Miller, S. J.: Characterization of stratospheric particle size distribution uncertainties using SAGE II and SAGE III/ISS extinction spectra, *Atmospheric Measurement Techniques*, 17, 2025–2054, <https://doi.org/10.5194/amt-17-2025-2024>, 2024.
- Kouker, W., Offermann, D., Küll, V., Reddmann, T., Ruhnke, R., and Franzen, A.: Streamers observed by the CRISTA experiment and simulated in the KASIMA model, *Journal of Geophysical Research: Atmospheres*, 104, 16 405–16 418, <https://doi.org/10.1029/1999JD900177>,
670 1999.
- Kovilakam, M., Thomason, L. W., Verkerk, M., Aubry, T., and Knepp, T. N.: OMPS-LP aerosol extinction coefficients and their applicability in GloSSAC, *Atmospheric Chemistry and Physics*, 25, 535–553, <https://doi.org/10.5194/acp-25-535-2025>, 2025.
- Krüger, K., Langematz, U., Grenfell, J. L., and Labitzke, K.: Climatological features of stratospheric streamers in the FUB-CMAM with increased horizontal resolution, *Atmospheric Chemistry and Physics*, 5, 547–562, <https://doi.org/10.5194/acp-5-547-2005>, 2005.
- 675 Labitzke, K.: Stratospheric-mesospheric midwinter disturbances: A summary of observed characteristics, *Journal of Geophysical Research: Oceans*, 86, 9665–9678, <https://doi.org/10.1029/JC086iC10p09665>, 1981.
- Lahoz, W. A., Geer, A. J., and Orsolini, Y. J.: Northern Hemisphere stratospheric summer from MIPAS observations, *Quarterly Journal of the Royal Meteorological Society*, 133, 197–211, <https://doi.org/10.1002/qj.24>, 2007.
- Leovy, C., Sun, C.-R., Hitchman, M., Remsberg, E., Russell, J., Gordley, L., Gille, J., and Lyjak, L.: Transport of Ozone in the Middle
680 Stratosphere: Evidence for Planetary Wave Breaking, *Journal of the Atmospheric Sciences*, 42, 230–244, [https://doi.org/10.1175/1520-0469\(1985\)042<0230:TOOITM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1985)042<0230:TOOITM>2.0.CO;2), 1985.
- Loughman, R. P., Griffioen, E., Oikarinen, L., Postlyakov, O. V., Rozanov, A., Flittner, D. E., and Rault, D. F.: Comparison of radiative transfer models for limb-viewing scattered sunlight measurements, *Journal of Geophysical Research: Atmospheres*, 109, 2003JD003 854, <https://doi.org/10.1029/2003JD003854>, 2004.
- 685 Manney, G. L., Livesey, N. J., Jimenez, C. J., Pumphrey, H. C., Santee, M. L., MacKenzie, I. A., and Waters, J. W.: EOS Microwave Limb Sounder observations of “frozen-in” anticyclonic air in Arctic summer, *Geophysical Research Letters*, 33, 2005GL025 418, <https://doi.org/10.1029/2005GL025418>, 2006.
- Matsuno, T.: A Dynamical Model of the Stratospheric Sudden Warming, *Journal of the Atmospheric Sciences*, 28, 1479–1494, [https://doi.org/10.1175/1520-0469\(1971\)028<1479:ADMOTS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<1479:ADMOTS>2.0.CO;2), 1971.
- 690 McCormick, M. P.: Aerosol measurements from earth orbiting spacecraft, *Advances in Space Research*, 2, 73–86, [https://doi.org/10.1016/0273-1177\(82\)90331-3](https://doi.org/10.1016/0273-1177(82)90331-3), 1982.
- McCormick, M. P. and Veiga, R. E.: SAGE II measurements of early Pinatubo aerosols, *Geophysical Research Letters*, 19, 155–158, <https://doi.org/10.1029/91GL02790>, 1992.
- McInturff, R. M.: Stratospheric warmings: Synoptic, dynamic and general-circulation Aspects, NASA Reference Publ. NASA-RP-1017, p.
695 174, <https://ntrs.nasa.gov/citations/19780010687>, last access: 10 October 2025, 1978.
- McIntyre, M. E. and Palmer, T. N.: The ‘surf zone’ in the stratosphere, *Journal of Atmospheric and Terrestrial Physics*, 46, 825–849, [https://doi.org/10.1016/0021-9169\(84\)90063-1](https://doi.org/10.1016/0021-9169(84)90063-1), 1984.
- Mote, P. W., Rosenlof, K. H., McIntyre, M. E., Carr, E. S., Gille, J. C., Holton, J. R., Kinnarsley, J. S., Pumphrey, H. C., Russell, J. M., and
700 Waters, J. W.: An atmospheric tape recorder: The imprint of tropical tropopause temperatures on stratospheric water vapor, *Journal of Geophysical Research: Atmospheres*, 101, 3989–4006, <https://doi.org/10.1029/95JD03422>, 1996.

- Myhre, C. E. L., Nielsen, C. J., and Saastad, O. W.: Density and Surface Tension of Aqueous H₂ SO₄ at Low Temperature, *Journal of Chemical & Engineering Data*, 43, 617–622, <https://doi.org/10.1021/jc980013g>, 1998.
- NASA/LARC/SD/ASDC: SAGE III/ISS L2 Solar Event Species Profiles (Native) V006, https://doi.org/10.5067/ISS/SAGEIII/SOLAR_BINARY_L2-V6.0, last access: 13 August 2025, 2025.
- 705 Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective determination of the polar vortex using Ertel’s potential vorticity, *Journal of Geophysical Research: Atmospheres*, 101, 9471–9478, <https://doi.org/10.1029/96JD00066>, 1996.
- Offermann, D., Grossmann, K.-U., Barthol, P., Knieling, P., Riese, M., and Trant, R.: Cryogenic Infrared Spectrometers and Telescopes for the Atmosphere (CRISTA) experiment and middle atmosphere variability, *Journal of Geophysical Research: Atmospheres*, 104, 16 311–16 325, <https://doi.org/10.1029/1998JD100047>, 1999.
- 710 Ploeger, F., Diallo, M., Charlesworth, E., Konopka, P., Legras, B., Laube, J. C., Grooß, J.-U., Günther, G., Engel, A., and Riese, M.: The stratospheric Brewer–Dobson circulation inferred from age of air in the ERA5 reanalysis, *Atmospheric Chemistry and Physics*, 21, 8393–8412, <https://doi.org/10.5194/acp-21-8393-2021>, 2021.
- Plumb, R. A.: A “tropical pipe” model of stratospheric transport, *Journal of Geophysical Research: Atmospheres*, 101, 3957–3972, <https://doi.org/10.1029/95JD03002>, 1996.
- 715 Plumb, R. A.: Stratospheric Transport, *Journal of the Meteorological Society of Japan. Ser. II*, 80, 793–809, <https://doi.org/10.2151/jmsj.80.793>, 2002.
- Pohl, C., Wrana, F., Rozanov, A., Deshler, T., Malinina, E., von Savigny, C., Rieger, L. A., Bourassa, A. E., and Burrows, J. P.: Stratospheric aerosol characteristics from SCIAMACHY limb observations: two-parameter retrieval, *Atmospheric Measurement Techniques*, 17, 4153–4181, <https://doi.org/10.5194/amt-17-4153-2024>, 2024.
- 720 Randel, W. J., Gille, J. C., Roche, A. E., Kumer, J. B., Mergenthaler, J. L., Waters, J. W., Fishbein, E. F., and Lahoz, W. A.: Stratospheric transport from the tropics to middle latitudes by planetary-wave mixing, *Nature*, 365, 533–535, <https://doi.org/10.1038/365533a0>, 1993.
- Reid, S. J. and Vaughan, G.: Lamination in ozone profiles in the lower stratosphere, *Quarterly Journal of the Royal Meteorological Society*, 117, 825–844, <https://doi.org/10.1002/qj.49711750009>, 1991.
- Reid, S. J., Vaughan, G., and Kyro, E.: Occurrence of ozone laminae near the boundary of the stratospheric polar vortex, *Journal of Geophysical Research: Atmospheres*, 98, 8883–8890, <https://doi.org/10.1029/93JD00230>, 1993.
- 725 Rieger, L. A., Bourassa, A. E., and Degenstein, D. A.: Odin–OSIRIS detection of the Chelyabinsk meteor, *Atmospheric Measurement Techniques*, 7, 777–780, <https://doi.org/10.5194/amt-7-777-2014>, 2014.
- Rieger, L. A., Malinina, E. P., Rozanov, A. V., Burrows, J. P., Bourassa, A. E., and Degenstein, D. A.: A study of the approaches used to retrieve aerosol extinction, as applied to limb observations made by OSIRIS and SCIAMACHY, *Atmospheric Measurement Techniques*, 11, 3433–3445, <https://doi.org/10.5194/amt-11-3433-2018>, 2018.
- 730 Rozanov, A.: Satellite data download, <https://www.iup.uni-bremen.de/DataRequest>, last access: 13 June 2025, 2024.
- Rozanov, A., Pohl, C., Arosio, C., Bourassa, A., Bramstedt, K., Malinina, E., Rieger, L., and Burrows, J. P.: Retrieval of stratospheric aerosol extinction coefficients from sun-normalized Ozone Mapper and Profiler Suite Limb Profiler (OMPS-LP) measurements, *Atmospheric Measurement Techniques*, 17, 6677–6695, <https://doi.org/10.5194/amt-17-6677-2024>, 2024.
- 735 Scherhag, R.: Die Explosionsartige Stratosphärenenerwärmung des Spätwinters 1951/52, *Berichte des Deutschen Wetterdienstes in der US-Zone*, 6, 51–63, 1952.
- Schoeberl, M. and Newman, P.: MIDDLE ATMOSPHERE I Polar Vortex, in: *Encyclopedia of Atmospheric Sciences*, pp. 12–17, Elsevier, ISBN 978-0-12-382225-3, <https://doi.org/10.1016/B978-0-12-382225-3.00228-0>, 2015.

- Schoeberl, M. R. and Hartmann, D. L.: The Dynamics of the Stratospheric Polar Vortex and Its Relation to Springtime Ozone Depletions, *Science*, 251, 46–52, <https://doi.org/10.1126/science.251.4989.46>, 1991.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, New York Academy of Sciences Series, John Wiley & Sons, Incorporated, Newark, 1st edn., ISBN 978-1-118-94740-1 978-1-119-22116-6, 2016.
- Sofieva, V. F., Rozanov, A., Szélag, M., Burrows, J. P., Retscher, C., Damadeo, R., Degenstein, D., Rieger, L. A., and Bourassa, A.: CREST: a Climate Data Record of Stratospheric Aerosols, *Earth System Science Data*, 16, 5227–5241, <https://doi.org/10.5194/essd-16-5227-2024>, 2024.
- Sonkaew, T., von Savigny, C., Eichmann, K.-U., Weber, M., Rozanov, A., Bovensmann, H., Burrows, J. P., and Groö, J.-U.: Chemical ozone losses in Arctic and Antarctic polar winter/spring season derived from SCIAMACHY limb measurements 2002–2009, *Atmospheric Chemistry and Physics*, 13, 1809–1835, <https://doi.org/10.5194/acp-13-1809-2013>, 2013.
- Taha, G.: OMPS-NPP L2 LP Aerosol Extinction Vertical Profile swath daily 3slit V2, <https://doi.org/10.5067/CX2B9NW6FI27>, 2020.
- Thiéblemont, R., Huret, N., Orsolini, Y. J., Hauchecorne, A., and Drouin, M.-A.: Frozen-in anticyclones occurring in polar Northern Hemisphere during springtime: Characterization, occurrence and link with quasi-biennial oscillation, *Journal of Geophysical Research*, 116, D20 110, <https://doi.org/10.1029/2011JD016042>, 2011.
- Thiéblemont, R., Matthes, K., Orsolini, Y. J., Hauchecorne, A., and Huret, N.: Poleward transport variability in the Northern Hemisphere during final stratospheric warmings simulated by CESM(WACCM), *Journal of Geophysical Research: Atmospheres*, 121, <https://doi.org/10.1002/2016JD025358>, 2016.
- Thomason, L. W., Poole, L. R., and Randall, C. E.: SAGE III aerosol extinction validation in the Arctic winter: comparisons with SAGE II and POAM III, *Atmos. Chem. Phys.*, 2007.
- Thomason, L. W., Moore, J. R., Pitts, M. C., Zawodny, J. M., and Chiou, E. W.: An evaluation of the SAGE III version 4 aerosol extinction coefficient and water vapor data products, *Atmos. Chem. Phys.*, 2010.
- Trepte, C. R., Veiga, R. E., and McCormick, M. P.: The poleward dispersal of Mount Pinatubo volcanic aerosol, *Journal of Geophysical Research: Atmospheres*, 98, 18 563–18 573, <https://doi.org/10.1029/93JD01362>, 1993.
- van Loon, H., Jenne, R. L., and Labitzke, K.: Zonal harmonic standing waves, *Journal of Geophysical Research*, 78, 4463–4471, <https://doi.org/10.1029/JC078i021p04463>, 1973.
- Vogel, B., Volk, C. M., Wintel, J., Lauther, V., Clemens, J., Groö, J.-U., Günther, G., Hoffmann, L., Laube, J. C., Müller, R., Ploeger, F., and Stroh, F.: Evaluation of vertical transport in ERA5 and ERA-Interim reanalysis using high-altitude aircraft measurements in the Asian summer monsoon 2017, *Atmospheric Chemistry and Physics*, 24, 317–343, <https://doi.org/10.5194/acp-24-317-2024>, 2024.
- von Savigny, C. and Hoffmann, C. G.: Issues related to the retrieval of stratospheric-aerosol particle size information based on optical measurements, *Atmospheric Measurement Techniques*, 13, 1909–1920, <https://doi.org/10.5194/amt-13-1909-2020>, 2020.
- von Savigny, C., Ernst, F., Rozanov, A., Hommel, R., Eichmann, K.-U., Rozanov, V., Burrows, J. P., and Thomason, L. W.: Improved stratospheric aerosol extinction profiles from SCIAMACHY: validation and sample results, *Atmospheric Measurement Techniques*, 8, 5223–5235, <https://doi.org/10.5194/amt-8-5223-2015>, 2015.
- Waugh, D. W.: Seasonal variation of isentropic transport out of the tropical stratosphere, *Journal of Geophysical Research: Atmospheres*, 101, 4007–4023, <https://doi.org/10.1029/95JD03160>, 1996.
- Waugh, D. W. and Polvani, L. M.: Stratospheric polar vortices, in: *Geophysical Monograph Series*, edited by Polvani, L. M., Sobel, A. H., and Waugh, D. W., vol. 190, pp. 43–57, American Geophysical Union, Washington, D. C., ISBN 978-0-87590-479-5, <https://doi.org/10.1029/2009GM000887>, 2010.

- Waugh, D. W., Sobel, A. H., and Polvani, L. M.: What Is the Polar Vortex and How Does It Influence Weather?, *Bulletin of the American Meteorological Society*, 98, 37–44, <https://doi.org/10.1175/BAMS-D-15-00212.1>, 2017.
- Wernli, H. and Sprenger, M.: Identification and ERA-15 Climatology of Potential Vorticity Streamers and Cutoffs near the Extratropical Tropopause, *Journal of the Atmospheric Sciences*, 64, 1569–1586, <https://doi.org/10.1175/JAS3912.1>, 2007.
- Wrana, F., von Savigny, C., Zalach, J., and Thomason, L. W.: Retrieval of stratospheric aerosol size distribution parameters using satellite solar occultation measurements at three wavelengths, *Atmospheric Measurement Techniques*, 14, 2345–2357, <https://doi.org/10.5194/amt-14-2345-2021>, 2021.
- Wüst, S., Küchelbacher, L., Trinkl, F., and Bittner, M.: Gravity waves above the northern Atlantic and Europe during streamer events using Aeolus, *Atmospheric Measurement Techniques*, 18, 1591–1607, <https://doi.org/10.5194/amt-18-1591-2025>, 2025.




Figure 4. Polar stereographic maps from 30–90° N, with 0° E as the central longitude line from bottom to centre, for various weekly averaged parameters (rows) at 10 hPa for a period of 4 weeks in March and April 2017 (columns), indicated above the first row. First row contains the aerosol extinction coefficient at 869 nm from OMPS-UBr as in Fig. 3. Rows 2 to 6 show maps of ERA5 reanalysis data, including wind speed (e–h), potential vorticity in potential vorticity units and geopotential height in decametres (dam) (i–l), specific humidity (m–p), ozone mass mixing ratio (q–t) and temperature (u–x). Wind arrows are plotted at wind speeds of 10 m s⁻¹ or higher. The contour lines of the geopotential height are delineated at intervals of 0.2 km. The highest and lowest geopotential heights are denoted with 'H' and 'L', respectively, along with the corresponding height in dam.




Figure 5. Weekly averaged profiles of the aerosol extinction coefficient in middle to high latitude (55–70° N, 20–45° W) for background conditions before the streamer event (05–11 March) and inside the anticyclone during the streamer event (19–25 March), and in the tropics to subtropics (0–30° N, 0–45° E) before the start of the streamer event (26 February to 04 March).




Figure 6. Mean vertical profiles of aerosol extinction at 869 nm for two streamers in mid-latitudes from 26 March to 01 April 2017 (45–55° N, 25–50° W) and from 04–10 March 2018 (45–55° N, 10° W to 15° E) as retrieved from OMPS-LP and SAGE III/ISS measurements.




Figure 7. Polar stereographic maps of the weekly averaged aerosol extinction coefficient at 750 nm from SCIAMACHY interpolated at 30.5 km in from March to August 2005 in bins of 5° latitude and 15° longitude from $30\text{--}90^\circ$ N.




Figure 8. Polar stereographic maps from 30–90° N for ~~various~~ weekly averaged parameters (rows) for a period of 4 weeks in March and April 2005 (columns), indicated above the first row. First row contains the aerosol extinction coefficient at 750 nm from SCIAMACHY as in Fig. 7 at an altitude of 30.5 km. Rows 2 to 6 show maps of ERA5 reanalysis data at 10 hPa, including wind speed (e–h), potential vorticity in potential vorticity units and geopotential height in dam (i–l), specific humidity (m–p), ozone mass mixing ratio (q–t) and temperature (u–x). Wind arrows are plotted at wind speeds of 10 m s^{-1} or higher. The contour lines of the geopotential height are delineated at intervals of 0.2 km. The highest and lowest geopotential heights are denoted with 'H' and 'L', respectively, along with the corresponding height in dam.