

High-Resolution Paleo-Storm Reconstructions from Eastern Canada Aligns with Late Holocene Northwestern Atlantic Hurricane Records

Deleted: -

Antoine Lachance¹, Matthew Peros^{2,3}, Jeannine-Marie St-Jacques^{1,3}, Pierre Francus^{4,3}, Nicole K. Sanderson^{5,3}

¹ Department of Geography, Environment and Planning, Concordia University, Montreal, H3G 1M8, Canada

² Department of Environmental, Agriculture, and Geography, Bishop's University, Sherbrooke, J1M 1Z7, Canada

³ GEOTOP – Research Center in Earth System Dynamics, Montreal, H3C 3P8, Canada

⁴ INRS – Institut National de la Recherche Scientifique, Centre Eau, Terre, Environnement, Québec, G1K 9A9, Canada

⁵ Département de Géographie, UQAM – Université du Québec à Montréal, Montreal, H2X 3R9, Canada

10 Correspondence to: Antoine Lachance (antoine.lachance@mail.concordia.ca)

Abstract

Atlantic Canada experiences frequent major storms, particularly tropical cyclones transitioning into post-tropical storms. Events such as Hurricane Fiona (2022), Dorian (2019), and Juan (2003) have caused significant damage, loss of life, and coastal erosion, exacerbated by sea level rise and warming waters. Despite this, centennial- to millennial-scale storm records in the region remain scarce. Existing studies in North America focus primarily on marine and coastal overwash records, with limited use of aeolian mineral inputs in ombrotrophic peatlands as storm proxies. Here, we address these gaps by analysing grain-size and geochemical data from two peatlands in Quebec, Canada's Magdalen Islands.

Formatted: Indent: First line: 1.27 cm

Our 4000-year peat-based storm reconstructions reveal consistent storm signals with three key intervals of increased activity: 800–550 BCE, 500–750 CE, and 1300–1700 CE. These records align with marine overwash archives from eastern Canada, the US, and the Bahamas, supporting reduced storminess during the Medieval Climate Anomaly and heightened activity during the Little Ice Age. This regional coherence suggests storm variability at higher latitudes is strongly influenced by local climatic drivers, such as sea surface temperature gradients and storm intensification mechanisms, rather than tropical cyclone formation alone.

Deleted: Our two peat records reveal consistent storm signals over the past 4000 years, with three key periods of heightened activity: 800–550 BCE, 600–800 BCE, and 1300–1700 CE. These signals align with marine and overwash records spanning the past 2000 years across eastern Canada, the US, and the Bahamas, indicating low storm activity during the Medieval Climate Anomaly, followed by increased activity during the Little Ice Age. Our findings suggest that storm records in these regions are influenced by local climatic factors. Negative phases of the Atlantic Multidecadal Variability, which typically suppress hurricane activity in the North Atlantic, are associated with conducive hurricane formation and intensification north of the Bahamas. Additionally, the position of the Bermuda High seems to play a more significant role in directing storm tracks during different climatic phases. Our findings highlight the potential antiphase relationship in storm activity between regions north of the Bahamas and those in the Gulf of Mexico, suggesting broader climatic mechanisms that warrant further investigations.

Formatted: Indent: First line: 1.27 cm

Despite broad similarities, notable discrepancies in geochemical and mineralogical profiles between our two peatlands highlight the influence of site-specific factors, including proximity to sediment sources, bog size, and local geomorphology, on sediment deposition and storm signal strength. These findings underscore the importance of site selection and local context when interpreting peat-based storm records. Furthermore, challenges remain in calibrating peat proxies to historical storm events, limiting direct event attribution over the past 150 years.

Deleted: Despite the similarities between our two sites, discrepancies in geochemistry and mineralogical profiles highlight the importance of site-specific conditions in interpreting the storm record from peatlands, namely the distance of the sites to the coast and source of aeolian sediment, as well as peatland size. Challenges also remain in calibrating peat-based proxies with historical storm records, as identifying specific events from the past 150 years remains difficult.

Our study demonstrates the complementary value of peatland archives alongside marine overwash records in reconstructing paleo-storm activity, enriching understanding of storm dynamics and expanding potential reconstruction sites, particularly in mid-latitude coastal regions. Continued research is needed to refine calibration methods and clarify climatic mechanisms driving storm variability, which is essential for projecting future storm impacts in Atlantic Canada.

1 Introduction

60 Atlantic Canada is frequently impacted by major storms, which have resulted in considerable damage and occasional death, especially from storms of tropical origins that have transitioned into powerful post-tropical cyclones (Baker et al., 2021; Public Safety Canada, 2024). Most recently, in September 2022, Hurricane Fiona made landfall in Nova Scotia with winds as strong as a category 2 event before tracking northwards through the Gulf of St. Lawrence and striking southwestern Newfoundland, becoming the costliest extreme event in Atlantic Canada [on record](#) (Insurance Bureau of Canada, 2022; Pasch et al., 2023). In September 2019, Hurricane Dorian passed over central Nova Scotia, eastern Prince Edward Island, and the Magdalen Islands, and resulted in large-scale wind-induced damage to trees and infrastructure and severe storm surges up to [1.5 meters](#) above normal levels (Avila et al., 2020). In September 2003, Hurricane Juan made landfall east of Halifax, Nova Scotia, as a category 1 event, causing eight deaths (Avila, 2003). Such events, coupled with ongoing sea level rise, are also contributing to coastal erosion in areas of the Gulf of St. Lawrence such as the Magdalen Islands and Prince Edward Island, 70 while the waters surrounding these islands are warming at a rate faster than almost any other region in the world (Barnett et al., 2017; Rémillard et al., 2017; Barnett et al., 2019; Wu et al., 2022). Consequently, how the frequency and intensity of high magnitude storm events will change in the future are important questions to consider, especially given record-setting sea surface temperatures in the North Atlantic during the past several years (Lapointe et al., 2020; Kuhlbrodt et al., 2024).

75 [A long-term perspective can help answer these questions](#), but there has been very little research examining centennial- to millennial-scale storm records in Atlantic Canada. One exception is a study from Robinson Lake, Nova Scotia, which used sediment grain-size and geochemistry to infer a period of enhanced impacts interpreted as being caused by storms of tropical origins during the Little Ice Age (LIA) (Oliva et al., 2017), the results of which are comparable to similar studies from New England (Van De Plassche et al., 2006; Besonen et al., 2008; Boldt et al., 2010; Donnelly et al., 2015). In New Brunswick, an inland lake record revealed a high concentration of storm impacts during the early LIA (before *ca.* 1650 CE) and from 1895 80 CE to the present (Patterson et al., 2022). Moreover, evidence for a powerful (possibly category 4) post-tropical cyclone strike on Cape Breton Island, Nova Scotia, was also found to have occurred during the period from 1756–1763 (Dickie and Wach, 2024). In southern Newfoundland, sedimentary records from several coastal lagoons have been used to infer a series of marine flood events occurring over the last several hundred years, which were attributed to both storms and tsunamis (Pleskot et al., 2023). Finally, a 6.5 ka record of storms was developed from a marine core collected from the Scotian Shelf, offshore Nova 85 Scotia, using sand content and elemental ratios such as Zr/Fe and Zr/Ti (Yang et al., 2020). The results show that storm frequency was sometimes synchronized between low and middle latitudes, particularly during 4.5–2.5 ka and since 0.5 ka, while at other times it was asynchronous.

90 Throughout the western North Atlantic, paleo-storm reconstructions have largely relied on marine overwash signals in nearshore or littoral sedimentary basins, which have provided considerable insights into late Holocene storm dynamics and driving mechanisms (Liu and Fearn, 1993, 2000; Lane et al., 2011; Braun et al., 2017; Bregy et al., 2018; Oliva et al., 2018; Wallace et al., 2021b; Wang et al., 2024). At many locations in the Bahamian Archipelago, coastal karst basins, including blue

Deleted: ever

Deleted: 5 feet

Deleted: Answering these questions is aided

Deleted: by a

holes, have provided high-resolution, decadal to centennial sedimentary records of intense tropical cyclone strikes, revealing periods of enhanced activity, particularly during the Little Ice Age (Van Hengstum et al., 2014; Wallace et al., 2019; Wallace et al., 2021b; Winkler et al., 2023). Similarly, in the Gulf of Mexico, sedimentary records from lagoons, coastal lakes, and karst basins across western Florida, Texas, and Belize have provided comparable high-resolution, decadal to centennial data.

100 (Lane et al., 2011; Brandon et al., 2013; Denommee et al., 2014; Rodysill et al., 2020; Monica et al., 2024; Wang et al., 2024). However, these records indicate a reduction in tropical cyclone activity during the Little Ice Age. As records of marine incursions, such studies provide information largely about storm surges, and the degree to which these correlate with storm size is unclear (Irish et al., 2008). Moreover, marine incursions can also occasionally be confused with tsunami strikes, although such events are relatively rare in eastern North America (Ten Brink et al., 2014).

105 In contrast, paleo-storm studies in the European context have often focused on the use of allochthonous mineral input into ombrotrophic coastal peatlands located nearby beaches or dunes (Björck and Clemmensen, 2004; Goslin et al., 2019; Vandel et al., 2019; Kylander et al., 2020; Sjöström et al., 2024). The premise is that storm events transport fine sand to such peatlands, whose mineral input is predominately dependent on atmospheric input (Shotyk, 1988; Weiss et al., 2002). Such data, measured using grain-size and geochemical indicators, provide a record of aeolian sediment influx, and ultimately storm-

110 wind activity, which has been argued to reflect “storminess” — the frequency and intensity of storms over a given period (Björck and Clemmensen, 2004; Vandel et al., 2019). For example, a recent study from Sweden used mid-infrared (MIR) spectral data from a large coastal ombrotrophic bog and showed a series of six periods of enhanced activity over the last 5000 years (Kylander et al., 2023). Moreover, at a site in Denmark, mineral grain size was used to infer wind intensity, which was seen to be high during negative North Atlantic Oscillation (NAO) phases (Goslin et al., 2019). In addition to yielding 115 information on wind characteristics, peatlands also often have long-term morphometric stability, limited bioturbation, and abundant macrofossils for radiometric dating, facilitating the interpretation of such records. A critical question remains whether aeolian paleo-storm reconstructions can be directly compared with marine and overwash-based paleo-storm records from nearby coastal sites.

120 We are unaware of any published research on paleo-storms in eastern North America using aeolian mineral input into ombrotrophic peatlands. The purpose of our paper is thus threefold: 1) to add new, long-term data on the frequency of storm events from an understudied region, namely Atlantic Canada; 2) to develop wind-based paleo-storm reconstructions from 125 North American peatlands; and 3) to explore whether findings from these reconstructions align with marine records from the nearby region, the North American eastern seaboard. We focus on two previously unstudied ombrotrophic peatlands located on the Magdalen Islands, Quebec, Canada. Our detection of storms is based on a combination of sediment grain-size and high-resolution micro-XRF core scanning elemental data from both sites.

Deleted: been increasingly used to

Deleted: . These records have

Deleted: ed

Deleted: .

Deleted: However, these records indicate a reduction in tropical cyclone activity during the Little Ice Age

Deleted: the first

2 Context

2.1 Geological setting and sea-level history

135 The Magdalen Islands (47°26'54"N, 61°45'08"W) are an archipelago of eight rocky islands situated on the shallow
(< 100 m) Magdalen Shelf in the Gulf of the St. Lawrence (Fig. 1a) (Owens and McCann, 1980). Six of these islands are
connected by a system of barrier beaches, tombolos, and lagoons (Bernatchez et al., 2012b). The islands are organized around
basaltic hills surrounded by karstic foothills and sandstone platforms (Brisebois, 1981; Rémiillard et al., 2013; Héu et al.,
2020). Previous work on the Late Quaternary history of the archipelago has provided insights into the pre- and post-glacial
140 coastal environments of the area (Prest et al., 1976; Dredge et al., 1992). Deglaciation occurred approximately 20 ka before
present (BP), leaving the islands largely submerged until about 15 ka BP. Subsequent glacial isostatic rebound caused the
islands to emerge, resulting in a relative sea level (RSL) low-stand at 9–10 ka BP (Rémiillard et al., 2016). This was followed
by eustatic sea-level rise, which brought RSL up to about 20 m below present 6 ka BP. Since then, RSL has risen at a
decelerating rate, eventually reaching modern levels (Vacchi et al., 2018; Barnett et al., 2019). Foraminifera, testate amoeba,
145 and plant macrofossils from a salt marsh at Havre-Aubert Island show that RSL has risen at a rate of 1.3–2.0 mm/yr for the
last two millennia, a rate substantially slower than the > 4.0 mm/yr observed during the 20th century (Barnett et al., 2017). This
Late Holocene sea-level stabilization provided conditions favourable for the development of modern coastal landforms,
including dunes, tombolos, spits, and beaches.

2.2 Coastal geomorphology

150 The coastal geomorphology of the Magdalen Islands reflects the combined influence of marine transgression and
coastal erosion. Today, approximately 28% of the coastline consists of rocky cliffs (Bernatchez et al., 2012b). In the southern
islands, including Havre-Aubert Island where our research sites are located, these cliffs are composed mainly of relic coastal
sands, loose unconsolidated sediments (colluvium), and a thick, porous, and highly erodible layer of red sandstone, known as
the *Drift des Demoiselles*, all of which are predominantly composed of quartz (Brisebois, 1981; Dredge et al., 1992; Rémiillard
155 et al., 2013). Due to the friable nature of these materials, cliff retreat occurs at an average rate of 0.3 m/yr, supplying a steady
stream of sediment to adjacent depositional systems (Bernatchez et al., 2012b). These sediments are redistributed alongshore
by prevailing currents through longshore drift, contributing material to the intertidal zone and beaches (Morin, 2001). This
sediment supply supports a dynamic coastal dune system, whose morphology varies with accumulation rates. In areas of high
sand supply, dunes evolved into large-scale coastal features such as tombolos and sand spits, some of which have continued
160 to prograde despite ongoing sea level rise (Rémiillard et al., 2015). This has led to the formation of extensive beach ridge
systems, highlighting the sustained and abundant sediment inputs from the surrounding cliffs. In contrast, most beaches are
bordered by a single foredune that remains less stable due to more limited sand availability (Morin, 2001). These foredunes
are frequently eroded by storm waves, forming dune scarp that expose sand to wind transport. Aeolian processes often carry
this sand inland, where it accumulates in back-dune environments, sometimes resulting in landward dune migration.

- Deleted:** Location of study
- Deleted:** Located in,
- Deleted:** t
- Deleted:** (GSL)
- Deleted:** are an archipelago (Brisebois, 1981; Héu et al., 2020)
- Deleted:**
- Deleted:** (Héu et al., 2020)
- Deleted:** Magdalen Islands
- Deleted:** The Magdalen Islands were d
- Deleted:** ed
- Deleted:** by
- Deleted:** and were
- Deleted:** by marine water
- Deleted:** Following this,
- Deleted:** -
- Deleted:** resulted
- Deleted:** in
- Deleted:** nce
- Deleted:** of the islands and
- Deleted:** fell, resulting in a
- Deleted:** Next, due to
- Deleted:** processes
- Deleted:** submergence occurred, so that by
- Deleted:**
- Deleted:** RSL was approximately 20 m below present level, and then rose at a decelerating rate to its present position
- Deleted:** At Havre-Aubert Island, a recent study using f
- Deleted:** to reconstruct late Holocene sea levels showed
- Deleted:** been rising
- Deleted:** ,
- Deleted:** compared to the more
- Deleted:** experienced
- Deleted:** based on tide gauge data
- Formatted:** Heading 2, Tab stops: 6.03 cm, Left
- Deleted:**
- Formatted:** Font: Italic
- Deleted:** (Morin, 2001)
- Formatted:** Font color: Auto

200 The beach ridges located landward of tombolos and spits provide valuable insights into Late Holocene shoreline dynamics. Optically stimulated luminescence (OLS) dating indicates that many of these features developed over the last 2000–3000 years (Rémillard et al., 2015). The oldest known beach ridge system, located on the Havre-aux-Maisons Island, was dated to at least 2600 years ago. On Havre-Aubert Island, near our study area, a beach ridge complex in the north of the island formed between approximately 550 and 400 years ago.▼

205 **2.3 Local climate**
The region is characterized by a strongly maritime climate, with cool summers and mild winters (Hétu et al., 2020). Daily maximum temperature is highest in July, reaching 21°C for the period 1981–2010, and is lowest in February at 4.3°C for the same period (Environment and Climate Change Canada, 2024). Monthly precipitation is relatively stable throughout the year, although is highest in November at 109 mm and lowest in late winter (February, March, April) and July at around 75 mm (Environment and Climate Change Canada, 2024). Sea ice cover in the Gulf of St. Lawrence normally begins in December and extends until April each winter (Canadian Ice Service, 2022), although over the past two decades sea ice around the Magdalen Islands has been diminishing (Derksen et al., 2019).▼

210 **Prevailing winds in the Gulf of St. Lawrence and around the Magdalen Islands are predominantly westerlies, with a distinct seasonality.** Winds are primarily from the south southwest in spring and summer (40–50 km/hr for 15% of the time), and from the northwest in the fall and winter, which are occasionally much stronger (70–132 km/hr for 15% of the time) (Appendix A, Fig. A1). **These westerlies contribute to more pronounced coastal erosion and narrower beaches on the islands' western shores (narrower beaches), while the eastern coasts remain relatively sediment-rich** (Morin, 2001; Forbes et al., 2004).▼

Deleted: (Morin, 2001)(Dredge et al., 1992; Rémillard et al., 2013)(Bernatchez et al., 2012b)

Formatted: Heading 2, Indent: First line: 0 cm

Deleted: consisting of basaltic hills surrounded by karstic foothills and sandstone platforms (Hétu et al., 2020), with the sandstone primarily composed of quartz and being the main sand source for the island's beaches (Adomat and Gischler, 2017). These hills are connected by a system of barrier beaches, tombolos, and lagoons that formed for the most part after 600 BCE (Rémillard et al., 2015; Barnett et al., 2017). The Magdalen Islands'

Deleted: climate is

Deleted: ,

Deleted: ,

Deleted: , and steady winds from all directions

Deleted:

Formatted: Indent: First line: 1.27 cm

Deleted: The wind direction shows a

Deleted: ,

Deleted: with w

Formatted: Not Highlight




Figure 1. Map and context of the study area. (a) The Magdalen Islands, located in the Gulf of St. Lawrence between Newfoundland and Prince Edward Island, in Atlantic Canada. The extent of map (b) is indicated by the red ~~rectangle~~. (b) The study site is on Havre-Aubert Island, the southernmost island of the archipelago. The extent of map (c) is indicated by the red ~~rectangle~~. (c) The two coring sites, Tourbière-du-Lac-Maucôque (TLM) and Tourbière-de-l'Anse-à-la-Cabane (TAC), are marked by blue squares, surrounded by multiple beaches, indicated by orange circles. (d, e) The TLM core site is 2150 m from the nearest beach (du Cap Beach), while the TAC core site is 250 m from south-facing sandstone cliffs. Imagery panel (b): © OpenStreetMap contributors, licensed under ODbL. Imagery panel (c–e): © Google Earth Pro 7.3.6.9796 (2024), Magdalen Islands, Quebec, Canada. Image date 22 April 2023, © 2025 Airbus.

240

2.4 Study sites

We studied two peatbogs, *Tourbière-du-Lac-Maucôque* (TLM) ($47^{\circ}14' 30.9''$ N, $61^{\circ}55' 24.7''$ W, ~14 m ASL) and *Tourbière-de-l'Anse-à-la-Cabane* (TAC) ($47^{\circ}12' 54.0''$ N, $61^{\circ}57' 48.4''$ W, ~20 m ASL), located on Havre-Aubert Island, the southernmost island of the Magdalen Islands archipelago (Fig. 1a–c). TLM, a circular peatland with an area of 40.2 ha and an elevation 14 m above mean sea level, is situated approximately 2 km from coastal beaches on the south and northeast side of Havre-Aubert Island (Fig. 1d) (Ministère De L'environnement Et De La Lutte Contre Les Changements Climatiques, 2021b). TAC lies parallel to the coast, covers 9.0 ha, has an elevation between 20 m above mean sea level, and has its centre 250 m from the nearest south-facing red sandstone cliffs (Fig. 1d) (Ministère De L'environnement Et De La Lutte Contre Les

Deleted: (Morin, 2001) Recent work focused on the dating of beach dunes using optically stimulated luminescence has shown that many of these features developed over the last 2000–3000 years during this period of RSL rise, as sea water eroded sandstone cliffs, providing a source of sediment for local beaches and dunes (Rémillard et al., 2015). Indeed, some low-lying areas, including Havre-aux-Basques (a beach near one of the peatlands in our study) are still prograding, providing a constant source of sediment.[†]

Deleted: square

Deleted: square

Deleted: 2

Deleted: 4

Deleted: 1

Deleted: 0

Deleted: 18–21

Formatted: Not Highlight

Field Code Changed

Changements Climatiques, 2021a). Both sites are designated as protected areas by the *Gouvernement du Québec* due to their endangered populations of northern dwarf huckleberry (*Gaylussacia bigeloviana* [Fernald] Sorrie & Weakley), a small coastal shrub present at both locations. The dominant vegetation in the peatlands is *Sphagnum* spp. moss (Lindb.) and shrubs from the *Ericaceae* family, with black spruce (*Picea mariana* [Mill.] Britton, Sterns & Poggenburg) growing around the perimeter.

3 Methods

3.1 Core sampling

In August 2020, two peat cores were collected from each of TLM and TAC using a Russian peat corer in 50 cm x 5 cm drives, near the centre of each peat dome to target the deepest and oldest peat (Fig. 1d–e). Coring was conducted until dense bottom layers were reached that no longer permitted penetration of the corer. The two sequences at each site were less than 1 m apart and vertically offset by 25 cm. Due to the low peat density at the surface, the top 0–50 cm consisting of monoliths were extracted using a shovel. Cores and monoliths were stored at 4°C at the Climate and Environmental Change Laboratory at Bishop's University. Continuous composite cores for each site were assembled by integrating core sequences with the surface monoliths (details in Appendix B).

3.2 Chronological controls

Twenty-five radiocarbon (¹⁴C) dates (9 for TLM and 16 for TAC) were obtained from the A.E. Lalonde AMS Laboratory at the University of Ottawa, following the method described in Crann et al. (2017), and calibrated with the IntCal20 curve (Reimer et al., 2020). Most samples submitted for AMS dating were terrestrial macrofossils, with some bulk peat samples included. In addition, total lead-210 (²¹⁰Pb) activity was measured from 15 samples each taken at regular intervals from the surface monoliths of TLM and TAC at the GEOTOP Radiochronology Laboratory at the *Université du Québec à Montréal* (UQAM), using an acid-digestion sequence and alpha spectrometry, adapted from De Vleeschouwer et al. (2010).

Age-depth models were constructed using the Bayesian Plum algorithm from the *rplum* package version 0.5.1 in R (Aquino-López et al., 2018; Blaauw et al., 2024), which integrates ¹⁴C and ²¹⁰Pb data. While the ²¹⁰Pb profile from TAC reached the supported level, the profile for TLM did not extend deep enough to clearly identify it. However, because Plum uses prior information to estimate supported ²¹⁰Pb activity, we applied the supported level measured in TAC (supported mean = 17.5 Bq kg⁻¹) as a prior for TLM (Aquino-López et al., 2018; Chartrand et al., 2023; Cwanek et al., 2025). This approach is appropriate given the similar sedimentary and environmental conditions at the two nearby sites and allowed us to derive a reliable ²¹⁰Pb-based chronology for TLM despite the data limitation.

3.3 Peat identification

Peat type was identified using the Troels-Smith method (Troels-Smith, 1955). Samples of 1 cm³ were extracted every 8 cm (TLM) and 4 cm (TAC) and examined under a stereoscopic microscope (at 4–40x magnification) to estimate the relative

Deleted: surface

Deleted: in split PVC tubes and wrapped in cellophane, and monoliths were wrapped in cellophane, with all samples stored

Deleted: Peros Lab

Deleted: The surface monoliths were compacted during transport and storage, requiring a correction factor of 1.79 (TLM) and 1.61 (TAC). This factor was calculated by dividing the original monolith length of 50 cm at the time of collection by the monolith length at the time of analysis.

Formatted: Indent: First line: 0 cm

Deleted: (Fig. 2 & Appendix C)

Deleted: to

Deleted: Additionally

Deleted: (Aquino-López et al., 2018; Cwanek et al., 2025)

Deleted: The

Deleted: a

Deleted: for TLM and TAC

Deleted: model

Deleted: integrating

Deleted: the

Deleted: es

Deleted: As ²¹⁰Pb levels in the TLM monolith did not reach supported levels, the prior supported level from the TAC core was applied ...

Formatted: Not Highlight

Field Code Changed

abundance of each peat type. Four main peat types were identified: bryophytic peat (dominated by *Sphagnum* spp. moss remains), herbaceous peat (dominated by sedge remains), ligneous peat (dominated by twigs and bark fragments), and humous peat (highly degraded peat with indistinguishable components). Mineral-rich layers and some macrofossils (seeds, leaves, needles) were also noted using the Continental Paleoecology Laboratory collection from GEOTOP at UQAM as reference

325 (Garneau, 1995).

3.4 Loss-on-ignition

Organic and mineral content in TLM and TAC were measured using loss-on-ignition (LOI) (Dean, 1974; Bengtsson and Enell, 1986; Heiri et al., 2001). Wet sediment samples (5 cm^3) were extracted every 2 cm (TLM) and every 1 cm (TAC), dried during 15 to 17 hours at $105\text{ }^\circ\text{C}$ (dry weight, DW_{105}) to calculate dry bulk density, then combusted at $550\text{ }^\circ\text{C}$ (LOI_{550}) in crucibles with lids, with weights recorded at each step. Mineral content (%) was calculated as the ratio of LOI_{550} to DW_{105} .

Deleted: and aeolian sand influx

330

3.5 Aeolian Sand Influx

The Aeolian Sand Influx (ASI) index was used to estimate inputs of coarse minerals transported by wind into the peatlands. ASI represents the mass accumulation rate of the sand-sized mineral fraction ($\text{g m}^{-2} \text{ yr}^{-1}$) and has been used as a storm proxy in several peat-based paleoclimate studies (Björck and Clemmensen, 2004; Orme et al., 2016a; Kylander et al., 2020; Vaasma et al., 2025). To isolate this fraction, subsamples previously analysed for LOI_{550} were treated with 10% hydrochloric acid (HCl) to remove carbonate and recalcitrant organic matter, and 10% potassium hydroxide (KOH) to deflocculate sediments (Vaasma, 2008). The samples were then wet-sieved at TLM using a $63\text{ }\mu\text{m}$ sieve, and at TAC using a $125\text{ }\mu\text{m}$ sieve. In both cases, we retained the material that did not pass through the sieve, i.e., particles larger than $63\text{ }\mu\text{m}$ at TLM and larger than $125\text{ }\mu\text{m}$ at TAC. These thresholds correspond to the sand-silt boundary ($> 63\text{ }\mu\text{m}$) and fine/very fine sand boundary ($> 125\text{ }\mu\text{m}$), respectively, based on the Wentworth (1922) classification. Grain-size thresholds were selected based on established aeolian transport theory and the geomorphological context. The initiation of sand transport by wind depends on grain size, surface moisture, vegetation, and local topography (Pye and Tsoar, 2008; Goslin et al., 2019). At TAC, the $> 125\text{ }\mu\text{m}$ fraction would require minimum wind speeds of $\sim 22.5\text{ m s}^{-1}$ ($\sim 80\text{ km h}^{-1}$) for entrainment and transport in saltation or suspension, while the $> 63\text{ }\mu\text{m}$ fraction used at TLM has a lower transport threshold. Although the TLM site is farther from the coastline, justifying our choice of a lower mineral threshold, both peatlands are adjacent to unconsolidated sediment sources (e.g., sandstone cliffs, beaches), and the sparse vegetation likely facilitate aeolian input under storm conditions. ASI was calculated by multiplying the dry bulk density of the sieved sand fraction (g cm^{-3}) by the peat accumulation rate (cm yr^{-1}), yielding a sand mass accumulation rate ($\text{g m}^{-2} \text{ yr}^{-1}$).

Deleted: overnight

Deleted: PEAT

Deleted: ASH

Deleted: ASH

Deleted: PEAT

Formatted: Heading 2

Formatted: Indent: First line: 1.27 cm

335

340

345

3.6 Micro X-ray fluorescence (μ -XRF) core scanning

350 We measured the inorganic geochemical compositions of TLM and TAC using an ITRAX Core Scanner (ITRAX-CS) from the *Laboratoire de géochimie, imagerie et radiographie des sediments* (GIRAS) at the *Institut national de recherche*

Deleted: The ASI₅₅₀ samples were treated with 10% hydrochloric acid (HCl) and 10% potassium hydroxide (KOH) to remove carbonate matter and deflocculate sediments (Vaasma, 2008), then sieved using a $63\text{ }\mu\text{m}$ sieve for TLM and a $125\text{ }\mu\text{m}$ sieve for TAC to establish the Aeolian Sand Influx (ASI) profiles. These size boundaries correspond to the fine/very fine sand ($< 125\text{ }\mu\text{m}$) for TAC and sand-silt ($< 63\text{ }\mu\text{m}$) for TLM (Wentworth, 1922), with a finer sieve for TLM due to its lower mineral content. ASI was calculated by multiplying sand density (g cm^{-3}) by the peat accumulation rate (cm yr^{-1}), and reported in $\text{g m}^{-2} \text{ yr}^{-1}$, as described by Björck and Clemmensen (2004).⁴

Deleted: 5

Deleted: →

scientifique (INRS) in Quebec City. Typically applied to marine and lacustrine sediments (Longman et al., 2019), the use of μ -XRF has recently been expanded to peat cores (Kylander et al., 2011; Orme et al., 2016b; Kern et al., 2019). Scans were acquired at 2 mm (TLM) and 1 mm (TAC) intervals using a Cr tube set to 40 kV, 10 mA and an acquisition time of 20 s. Prior to scanning, the core surfaces were smoothed to minimize surface roughness and reduce measurement variability. We applied quality control procedures on the resulting dataset based on count rate (CPS) and mean squared error (MSE) to exclude outliers affected by poor data quality. For each element, measurements with CPS values exceeding ± 3 standard deviations from that element's mean, or exhibiting anomalously high MSE, were removed prior to analysis (Bishops, 2025). Our analysis used a subset of well-measured elements linked to terrigenous mineral inputs: K, Ti, Si, Fe, Mn, Ca, and S. Downcore variations are expressed as the proportion of each element relative to the total of these lithogenic elements (e.g., $Ti\% = 100 * Ti / [Ti + K + Si + Fe + Mn + Ca + S]$). This normalization reduces the influence of variations in sample density, water content, and organic matter on geochemical trends (Jansen et al., 1998; Bertrand et al., 2023).

3.7 Multivariate statistical analysis

3.7.1 Data processing

The LOI and μ -XRF datasets are compositional data, which require log-ratio transformations for multivariate analyses (Aitchison, 1982; Van Den Boogaart and Tolosana-Delgado, 2013; Bertrand et al., 2023). Raw μ -XRF data were first aggregated to 1-cm resolution by averaging raw counts within each depth interval to ensure consistency with the lower resolution LOI dataset. For μ -XRF elemental intensities, we computed centred-log-ratios (CLR) using the `clr` function in the `compositions` R package version 2.0-8 (Van Den Boogaart and Tolosana-Delgado, 2008; Van Den Boogaart et al., 2024). Any zero counts were replaced using a centred 10-point moving average. For LOI variables, which contain fewer than five components and do not meet CLR assumptions, we applied a log-ratio transformation of each component relative to the remainder (e.g., $\log(\text{Minerals} [\%] / (100 - \text{Minerals} [\%]))$) (Bertrand et al., 2023). The ASI was log-transformed to meet the assumption of normality.

3.7.2 Statistical methods

Core zonation of TLM and TAC were determined using stratigraphically constrained cluster analysis (CONISS) from the *rioja* package version 1.0-7 in R (Juggins, 2024) on LOI and μ -XRF variables. Correlation analyses were performed using Pearson's correlation coefficient to evaluate relationships among mineral content from LOI and μ -XRF variables within each zone identified by CONISS. Principal Components Analysis (PCA) was conducted on our subset of seven CLR-transformed elements (K, Ti, Si, Fe, Mn, Ca, S), with mineral content included as passive variable. This analysis, performed with the *factoMineR* package version 2.11 in R (Husson et al., 2024), used z-score-transformed variables and a correlation matrix to compute the eigenvalues. Measurements from TLM and TAC's basal sand layers were excluded from all statistical analyses.

3.8 Paleo-storm identification

475 Two independent paleo-storm reconstructions were performed, one for each of TLM and TAC. Storm reconstructions were based on the ombrotrophic portions of the peat cores, excluding the early developmental stages and sections where ombrotrophic conditions could not be confidently established. These zones likely reflect mixed sedimentary processes, including fluvial or groundwater transport, rather than purely aeolian input (Sjöström et al., 2020). We selected a paleo-storm proxy by identifying μ -XRF elemental signals that co-varied with ASI in ombrotrophic peat. To do this, we conducted a PCA on the ombrotrophic sections of the TLM and TAC cores using the subset of seven elements, with ASI included as a passive variable.

480 While μ -XRF data were CLR-transformed prior to performing the PCA, the paleo-storm identification method employed a μ -XRF storm proxy normalized to the sum of terrigenous elements, or $\text{Proxy}_{\text{normalized}} = \text{Proxy} / [\text{Ti} + \text{K} + \text{Si} + \text{Fe} + \text{Mn} + \text{Ca} + \text{S}]$. This approach mitigates matrix effects while preserving the natural variability and peak values essential for detecting extreme storm events, features that CLR transformation tends to smooth out.

485 To identify intervals where the μ -XRF storm proxy exceeded background levels, we used an approach adapted from Donnelly et al. (2015) and Lane et al. (2011). First, to account for variations attributed to long-term lithogenic changes, we detrended the storm proxy using a running average:

$$\text{Proxy}_{\text{detrended}}(z) = \text{Proxy}_{\text{normalized}}(z) - \text{RA}_{\text{proxy}}(z, w), \quad (3)$$

490 where $\text{RA}_{\text{proxy}}(z, w)$ is the running average of the storm proxy normalized at depth z , calculated over a window length w of 30 years. The threshold for identifying outlier proxy values was defined with the Tukey Rule of outlier detection (Tukey, 1977), following Winkler et al. (2022).

$$\text{Proxy}_{\text{threshold}} = Q_{0.75}(\text{Proxy}_{\text{detrended}} > 0) + x \cdot \text{IQR}(\text{Proxy}_{\text{detrended}} > 0), \quad (4)$$

495 where $Q_{0.75}(\text{Proxy}_{\text{detrended}} > 0)$ is the third quartile of detrended storm proxy values above zero, and $x \cdot \text{IQR}(\text{Proxy}_{\text{detrended}} > 0)$ is the interquartile range of detrended storm proxy values above zero, multiplied by a factor x of 0.75 for TLM and 1.5 for TAC, respectively, to account for differences in proxy values between the two cores. Detrended proxy values exceeding the threshold were considered storm events. To avoid overestimating the number of events due to consecutive storm proxy measurements associated with a single storm proxy peak, consecutive measurements were grouped together, and only the maximum value within each group was retained in the final paleo-storm record. Finally, the number of 500 events per century was calculated using a 100-year running-sum.

Deleted: 7 Paleo-storm identification

Deleted: (i.e., fen-dominated sections)

Deleted: ied, performing Ti and K. Ti was selected as the primary geochemical indicator of aeolian sand input. This decision is based on the potential for K, as an essential plant nutrient, to accumulate in plant tissues—especially near the surface—potentially complicating its interpretation as an aeolian indicator (Steinmann and Shotyk, 1997; Shotyk, 1988).

Formatted: English (UK)

Deleted: 

Deleted: Ti

Deleted: Ti

Deleted: using Eq. (3)

Deleted: Ti

Deleted: Ti

Deleted: raw

Deleted: Ti

Deleted: \rightarrow

Deleted: Ti

Deleted: raw

Deleted: Ti

Deleted: :

Deleted: 10 years for the period 1851–2019 CE and 30 years 

Deleted: \rightarrow

Deleted: Ti

Deleted: (Tukey, 1977)

Deleted: Winkler et al. (2020)

Deleted: Ti

Deleted: Ti

Deleted: Ti

Deleted: \rightarrow

Deleted: Ti

Deleted: normalized

Deleted: Ti

Deleted: Ti

Deleted: normalized

Deleted: Ti

Deleted: The lower x value for TLM accounts for its reduced 

Deleted: Ti


Deleted: Ti

Deleted: Ti

4.1 Core description

4.1.1 Chronostratigraphy

The TLM composite core measured 700 cm, with a basal sand layer between 693–700 cm. The TAC composite core was 325 cm long, with a basal sand layer between 317–325 cm (see Appendix C for core photographs). The age-depth models indicated that organic accumulation began above the basal sand layers around 2960 BCE for TLM and 2900 BCE for TAC, giving an approximate chronology of 5000 years of peat accumulation for both cores (Fig. 2, Table 1). The average peat accumulation rate (PAR) was 0.17 cm yr⁻¹ for TLM and 0.12 cm yr⁻¹ for TAC. At TAC, a marked decline in PAR to 0.02 cm yr⁻¹ occurred between 135–159 cm (870 CE–870 BCE), suggesting a possible hiatus in peat accumulation. This trend was not observed in TLM. In the upper 50 cm of both cores, which includes the acrotelm (the layer containing living plants) and was dated using ²¹⁰Pb, and the PAR increased significantly to 0.77 cm yr⁻¹ for TLM and 0.29 cm yr⁻¹ for TAC. Further details about the chronostratigraphy can be found in Appendix D.



580 Figure 2. *Rplum*-derived age-depth models for TLM (a) and TAC (b). Radiocarbon (¹⁴C) dates are shown in grey along the mean age model (dotted red). The rate of supported ²¹⁰Pb (right axis) is indicated by blue boxes on the left side (the blue boxes are the measured ²¹⁰Pb values, while the blue shadings represent the modelled ²¹⁰Pb values). A complete version with the upper graphs and blown-up version of the ²¹⁰Pb diagrams is available in Fig. C1.

585 Table 1. Accelerator mass spectrometry (AMS) radiocarbon (¹⁴C) results for TLM and TAC. Dated material includes bulk peat sediment (bulk) and wood fragments (wood). Uncalibrated ¹⁴C dates are shown as years before 1950 (BP). The ¹⁴C dates were processed using the Bayesian *rplum* algorithm in R and calibrated with the IntCal20 curve. Calibrated dates are presented as both calibrated years before 1950 (cal. BP) and as Common Era/Before Common Era (CE/BCE).

Sample depth (cm)	Lab ID	Material	AMS ¹⁴ C age BP (uncal.)	Calibrated 2 sigma range		Mean modelled age (uncertainty) (CE/BCE)
				Cal. BP	CE/BCE	

Deleted: 3.8 Historical storm record and compilation
 → We used the *International Best Track Archive for Climate Stewardship* (IBTrACTS) dataset (Knapp et al., 2010; Gahtan et al., 2024) to determine the number of cyclones with a tropical origin passing within 100 km of the Magdalen Islands from 1851 to 2019 CE. To assess the impact of these storms on the Magdalen Islands, we cross-referenced the IBTrACTS data with historical shipwrecks records and local newspaper archives. We searched the online press database of the *Bibliothèque et Archive National du Québec* (*Bibliothèque Et Archive Nationale Du Québec*, 2024) using French keywords such as *ouragan* and *Îles-de-la-Madeleine*, and notes were taken on storm-related damage in the Magdalen Islands. Additionally, we manually reviewed shipwreck records from the vicinity of the Magdalen Islands (Landry, 1989, 1994) for incidents attributed to storms identified in the IBTrACTS dataset. Storms mentioned in the newspaper articles or shipwreck records that were not present in the IBTrACTS dataset, but that had significant impacts on the Magdalen Islands, were added to the final historical storm dataset.

Deleted: Before consolidating the core drives into a composite core, we validated their alignment using cross-correlation analysis of overlapping raw μ -XRF measurements (Appendix B).

Deleted: D

Deleted: n abrupt decrease

Deleted: pronounced

Deleted: reduction in peat accumulation or a hiatus

Deleted:

Deleted: C

Deleted: r

Formatted: Caption

Deleted: For each core, the upper left panel shows the Markov Chain Monte Carlo (MCMC) iterations: followed by the prior (green curve) and posterior (grey histogram) distribution for the accumulation rate (second panel), memory (third panel), ²¹⁰Pb influx in Bq/m²·yr (fourth panel) and supported ²¹⁰Pb in Bq/kg (fifth panel). Since ²¹⁰Pb levels in TLM did not reach supported levels, the prior supported level from the TAC (s. mean = 17.5 Bq kg⁻¹) core was used.

TLM						
59-60	UOC-18434	Bulk	174 ± 21	289 – 73	1661 – 1916 CE	1825 (1750 – 1890) CE
122-123	UOC-14324	Wood	417 ± 36	526 – 327	1425 – 1624 CE	1455 (1350 – 1510) CE
243-244	UOC-15799	Wood	1582 ± 35	1532 – 1390	418 – 561 CE	495 (415 – 585) CE
390-391	UOC-15800	Wood	2491 ± 26	2723 – 2469	774 – 520 BCE	605 (745 – 470) BCE
484-485	UOC-15801	Wood	3014 ± 28	3335 – 3078	1386 – 1129 BCE	1270 (1380 – 1150) BCE
577-578	UOC-14325	Wood	3572 ± 37	3978 – 3724	2029 – 1775 BCE	1930 (2030 – 1805) BCE
634-635	UOC-15802	Wood	3813 ± 26	4346 – 4093	2397 – 2144 BCE	2365 (2490 – 2230) BCE
669-670*	UOC-18285	Wood	7923 ± 19	8979 – 8604	7030 – 6655 BCE	2730 (3015 – 2515) BCE
674-675*	UOC-14326	Wood	7650 ± 32	8536 – 8382	6581 – 6433 BCE	2780 (3080 – 2250) BCE

TAC						
31-32	UOC-18435	Bulk	6 ± 21	249 – 41	1701 – 1909 CE	1940 (1930 – 1950) CE
57-58	UOC-15803	Wood	162 ± 26	287 – 43	1664 – 1908 CE	1730 (1670 – 1785) CE
76-77	UOC-18286	Wood	341 ± 13	469 – 316	1482 – 1635 CE	1565 (1495 – 1625) CE
102-103	UOC-18436	Bulk	613 ± 20	649 – 551	1301 – 1400 CE	1340 (1300 – 1395) CE
122-123	UOC-15804	Wood	869 ± 26	900 – 693	1050 – 1257 CE	1150 (1060 – 1210) CE
128-129	UOC-18287	Wood	964 ± 14	921 – 797	1030 – 1154 CE	1070 (970 – 1140) CE
133-134	UOC-18437	Bulk	992 ± 21	957 – 798	994 – 1153 CE	945 (775 – 1040) CE
147-148	UOC-15805	Wood	2469 ± 27	2712 – 2371	763 – 422 BCE	475 (630 – 395) BCE
162-163	UOC-18288	Wood	2928 ± 14	3159 – 3001	1210 – 1052 BCE	970 (1080 – 840) BCE
185-186	UOC-18438	Bulk	3052 ± 21	3350 – 3178	1401 – 1229 BCE	1260 (1350 – 1145) BCE
199-200	UOC-15806	Wood	3178 ± 27	3452 – 3361	1503 – 1412 BCE	1400 (1470 – 1305) BCE
226-227	UOC-15807	Wood	3300 ± 28	3571 – 3455	1622 – 1506 BCE	1600 (1690 – 1525) BCE
258-259	UOC-15808	Wood	3508 ± 29	3870 – 3694	1921 – 1745 BCE	1885 (2005 – 1800) BCE
283-284	UOC-15809	Wood	3597 ± 28	3980 – 3834	2031 – 1885 BCE	2155 (2295 – 2035) BCE
296-297	UOC-15810	Wood	3771 ± 30	4240 – 3996	2291 – 2047 BCE	2370 (2490 – 2240) BCE
319-320	UOC-18289	wood	4376 ± 17	5025 – 4865	3076 – 2916 BCE	2960 (3065 – 2915) BCE

* The AMS ^{14}C dates at these depths were inverted and were considered outliers by the *plum* R algorithm. Therefore, the modelled mean dates associated with these depths do not reflect the AMS ^{14}C date.

625 4.1.2 Downcore variations in lithology

The TLM core was divided into three zones (Fig. 3a). Zone 1 (692–565 cm, 2950–1840 BCE) consisted mainly of ligneous peat with a high mineral content of 13 %. This zone included seeds from *Menyanthes trifoliata* L. (bog bean). A detritus peat layer between 657–652 cm (2590–2535 BCE) contained a peak mineral content of 34 %. Zone 2 (565–452 cm, 1840–1035 BCE) transitioned from ligneous to herbaceous peat with a sharp decrease in mineral content to an average of 2.3

630 %. There was an abrupt transition to bryophytic peat at 486 cm (1280 BCE). Zone 3 (450–0 cm, 1035 BCE–2019 CE) showed clear ombrotrophic conditions (Fig. 3) and alternated between well-preserved and degraded bryophytic peat, with a low mineral content of 1.5 %. The upper 50 cm showed a higher mineral content (2.5 %).

The TAC core was divided into 4 zones (Fig. 3b). Zone 1 (316–274 cm, 2890–2050 BCE) consisted of herbaceous peat interspersed with small ligneous fragments. It had a moderate mineral content of 6 %. A sand-enriched layer (28 % mineral content) between 309–305 cm (2700–2590 BCE) coincided chronologically with a similar layer in TLM. *Sphagnum* macrofossils first appeared at 284 cm (~ 2160 BCE). Zone 2 (274–159 cm, 2050–880 BCE) showed ombrotrophic conditions and alternated between well-preserved and degraded bryophytic peat, with the lowest mineral content (3 %). Zone 3 (159–135

Formatted: Font: Italic

Deleted: ~~1~~

Formatted: Superscript

Deleted: and a mean ASI of $3.4 \text{ g m}^{-2} \text{ yr}^{-1}$

Deleted: ~~1~~, a common shallow-water plant (Magnan et al., 2014)

Deleted: and an ASI of $0.06 \text{ g m}^{-2} \text{ yr}^{-1}$

Deleted: and a mean ASI of $0.1 \text{ g m}^{-2} \text{ yr}^{-1}$

Deleted: and ASI $(0.6 \text{ g m}^{-2} \text{ yr}^{-1})$

Deleted: and a mean ASI of $0.15 \text{ g m}^{-2} \text{ yr}^{-1}$

Deleted: and ASI $(0.08 \text{ g m}^{-2} \text{ yr}^{-1})$

cm, 880 BCE–870 CE) was a dense, mineral-rich zone (minerals content 27%) dominated by detritus peat with ligneous layers.

~~Two layers at 151 cm (~ 600 BCE) and 136 cm (~ 810 CE) had high mineral contents of 32 % and 55 %, respectively. This zone likely represents a hiatus in peat accumulation. The interruption appears to begin around 600 BCE, coinciding with the lower mineral layer at 151 cm, where the previously steady age–depth relationship begins to deviate. Peat accumulation resumes at a rate comparable to that of Zone 2 immediately above the upper mineral-rich layer at 136 cm. Due to a high degree of peat degradation, ombrotrophic conditions could not be confirmed for Zone 3. However, ombrotrophic conditions were re-established in Zone 4 (135–0 cm, 870–2019 CE), which consisted of well-preserved bryophytic peat with a mineral content of 21%.~~

Deleted: and ASI $8.1 \text{ g m}^{-2} \text{ yr}^{-1}$

Deleted: It contained two fine-grained sand


Deleted: layers

Deleted: , with

Deleted:

Deleted: and the highest ASI at $3.6 \text{ g m}^{-2} \text{ yr}^{-1}$.

Formatted: English (UK)



Deleted: 4
<object>

Figure 3. Lithological and geochemical profiles of the TLM (top) and TAC (bottom) composite cores. The peat type (bryophytic, herbaceous, ligneous, detritus, mineral) is color-coded, along with peatland classification (bog versus fen). Micro-XRF variables are displayed as percent ratios over the total counts from the presented elements (%). Grey lines indicate the original μ -XRF resolution (2 mm for TLM and 1 mm for TAC), while the black lines represent values averaged to 1 cm intervals. Principal components (PC 1 and PC 2) derived from seven CLR-transformed μ -XRF elements are shown, illustrating broad geochemical trends across the stratigraphy. Dark blue/red indicate positive loadings; pale blue/red indicate negative loadings.

Formatted: Caption

Deleted: Lithological and geochemical profiles of the TLM (a) and TAC (b) composite cores. The peat type (bryophytic, herbaceous, ligneous, detritus, mineral) is color-coded, along with peatland classification (bog versus fen). Grey shading in the LOI variables correspond to the \log_{10} of the mineral content and ASI to enhance visualization of small variations. Micro-XRF variables are displayed as CLRs. Grey lines indicate the μ -XRF original resolution (2 mm for TLM and 1 mm for TAC), while the black lines represent data averaged to 1 cm intervals. The first two principal components (PCA 1 and PCA 2) performed on seven CLR-transformed μ -XRF elements are presented. Dark blue/red indicate positive loading, while pale blue/red indicate negative loading.

4.1.3 Principal component analysis and geochemical profiles

685 The first principal components (PC1), derived from the PCA of seven elements of interest (K, Ti, Si, Fe, Mn, Ca, S) and mineral content added as a passive variable, explained 49 % and 52 % of the total data variance at TLM and TAC, respectively (Fig. 3). At both sites, PC1 was strongly negatively correlated with mineral content (TLM: $r = -0.94$; TAC: $r = -0.93$), indicating that positive PC1 values corresponded to low mineral/high organic content, and negative PC1 values indicated high mineral/low organic content. At TLM, PC1 showed the strongest negative correlations with Fe ($r = -0.84$), S ($r = -0.71$), 690 and Ti ($r = -0.48$), linking these elements to mineral content. In contrast, at TAC, PC1 was most strongly negatively correlated with Ti ($r = -0.90$) and K ($r = -0.75$), highlighting differences in the main geochemical composition between the two sites. However, PC1 at both locations was positively correlated with Mn and Ca, reflecting the consistent association of those elements with low mineral content. The second principal components (PC2) explained 21 % and 20% of the variance in TLM 695 and TAC, respectively, and showed no strong correlation with mineral content. At TLM, PC2 was positively correlated with S ($r = 0.54$), Ca ($r = 0.45$), and Fe ($r = 0.29$), and negatively correlated with Ti ($r = -0.71$) and K ($r = -0.60$). At TAC, PC2 was positively correlated with Si ($r = 0.55$), Fe ($r = 0.52$), and K ($r = 0.35$), and negatively correlated with S ($r = -0.81$). In both cases, PC2 likely reflected variations in the geochemical composition associated with changes in sources of mineral input.

700 Zone 1 of TLM (692–565 cm, 2950–1840 BCE) saw a shift in chemical composition indicated by PC2: Ti and K dominated the older (lower) part, while Fe, S, and Ca, commonly measured in higher proportion in fens (Shotyk, 1988), increased in the upper section. This shift corresponded to the detritus layer at 657–652 cm (2590–2535 BCE). In Zone 2 (565–452 cm, 1840–1035 BCE), PC2 remained positive, reflecting higher proportions of Fe and S, which are positively correlated with mineral content in that zone (Appendix Table E1). Zone 3 (450–0 cm, 1035 BCE–2019 CE) showed alternating PC2, indicating variable contributions from Ti, K, Fe, S, and Ca, though only Ti, K, and Fe were significantly positively associated to mineral content in this zone (Appendix Table E1). In the upper 50 cm, a negative shift in PC2 highlighted a dominance of 705 Ti and K in modern sediment inputs.

710 At TAC, Zone 1 (316–274 cm, 2890–2050 BCE) saw a shift in chemical composition indicated by PC1, with Ti and K dominating the older (lower) part, while Fe, Mn and Ca became more prominent in the upper part. As with TLM, this shift was associated to a mineral-enriched layer. A negative PC2 indicated a high S content, though S showed no significant correlation with mineral content (Appendix Table E1). In Zone 2 (274–159 cm, 2050–880 BCE), only Ti showed a positive correlation with the mineral content (Appendix Table E1). Zone 3 (159–135 cm, 880 BCE–870 CE) contained fine-grained layers enriched in Ti, K, and Si, as indicated by strong positive correlations with mineral content, and a negative PC2 suggested the presence of S, though it was not associated to mineral content (Table E1). In Zone 4 (135–0 cm, 870–2019 CE), LOI variables were positively correlated with Ti, K, and Fe, while in the upper 50 cm, a positive PC2 reflected increased input of Fe and Si.

Deleted: Only statistically significant results are described.

Deleted: two

Deleted: s (mineral content and ASI)

Deleted: 5

Deleted: -

Deleted: 2

Deleted: and ASI (TLM: $r = -0.92$, TAC: $r = -0.85$),

Deleted: and

Deleted: and ASI

Deleted: 19

Deleted: or ASI


Deleted: 4

Deleted: 2

Deleted: 7

Deleted: and ASI


Deleted: sand



PCA axis number	1	2
% variance	49.3	21.3
correlations r		
Ti	-0.48	-0.71
K	0.43	-0.60
Si	0.82	0.11
Fe	-0.84	0.29
Mn	0.82	0.18
Ca	0.69	0.45
S	-0.71	0.54
Minerals	-0.94	0

PCA axis number	1	2
% variance	52.3	19.4
correlations r		
Ti	-0.90	-0.16
K	-0.75	0.35
Si	-0.06	0.55
Fe	0.57	0.52
Mn	0.91	0.17
Ca	0.96	0.07
S	0.45	-0.81
Minerals	-0.93	0.14

Figure 4. Principal components analyses (PCA) of the seven μ -XRF elements for TLM (a) and TAC (b). Each CONISSL zone is depicted with a distinct coloured symbol. Solid black arrows represent the μ -XRF variables used in the PCA, while dashed blue arrows represent the supplementary variable Minerals. The Pearson correlation coefficients r between the principal components and all variables are displayed below the respective biplots, with significant correlations $r \geq 0.2$ highlighted in bold.



4.2 Paleo-storms

Deleted: 4

755 4.2.1 Ombrrophic zones

The paleo-storm reconstruction for TLM is based on Zone 3 (1035 BCE–2019 CE), while the TAC analysis is based on Zone 2 (2050–880 BCE) and Zone 4 (870–2019 CE). Zones 1–2 of TLM and Zone 1 of TAC were excluded, as they represent early stages of peat development, when sediment delivery was dominated by non-aeolian processes and thus unsuitable for storm reconstruction (see Discussion). TAC Zone 3 was also excluded; this section likely includes a hiatus and 760 may represent the time-integrated accumulation of up to 2000 years of mineral inputs and other processes. Moreover, the extremely low peat accumulation rate during this interval limits temporal resolution and makes it incompatible with our event detection methodology, which is based on a 30-year moving average.

4.2.2 Aeolian sand influx and geochemical proxies

Formatted: Not Highlight

765 While ASI was calculated across the full length of both TLM and TAC cores, the following analysis focuses on the identified ombrrophic sections.

In TLM, Zone 3 (450–0 cm; 1035 BCE–2019 CE) has a mean ASI of $0.1 \text{ g m}^{-2} \text{ yr}^{-1}$ (Fig. 5a). Log-transformed ASI values indicate a period of stability from 1035 BCE to around 1000 CE, with near-zero values interrupted by occasional small peaks. This is followed by a phase of increased variability between approximately 1000 and 1600 CE. The upper 50 cm of TLM show notably higher ASI values at $0.6 \text{ g m}^{-2} \text{ yr}^{-1}$. In TAC, Zone 2 (274–159 cm; 2050–880 BCE) has a mean ASI of 770 $0.08 \text{ g m}^{-2} \text{ yr}^{-1}$, whereas Zone 4 (135–0 cm; 870–2019 CE) exhibits a substantially higher mean ASI of $3.6 \text{ g m}^{-2} \text{ yr}^{-1}$ (Fig. 5b). As in TLM, the log-transformed ASI at TAC shows a period of variability in Zone 2 that matches similar trends in Ti.

775 A PCA was conducted on seven CLR-transformed elements, with ASI included as a passive variable, using data from the ombrrophic sections only. The first principal component (PC1) shows strong loadings for ASI, Ti, and K in both TLM and TAC (Fig. 5). In TLM, ASI, Ti, and K negatively correlate with PC1 with r values of -0.78, -0.68, and -0.73, respectively; in TAC, these negative correlations are even stronger, at -0.93, -0.87, and -0.86. These results reflect a strong correspondence between downcore variations in ASI and those of Ti and K, particularly visible in the log-transformed ASI.

780 PCA results identifies both Ti and K as strong candidates for tracking ASI and storm-related inputs. However, Ti exhibits greater amplitude between background and peak levels, particularly in low-ASI sections, making it a more sensitive indicator of aeolian sand influx. Although K appears to track ASI more closely than Ti in the upper 50 cm of both the TLM and TAC cores, its interpretation is complicated by its biological role. As an essential plant nutrient, K can accumulate in living peat near the surface, potentially obscuring its signal as an aeolian proxy (Steinmann and Shotyk, 1997; Shotyk, 1988). In contrast, Ti is a conservative lithogenic element that resists biogenic alteration and post-depositional remobilization, making it a more reliable tracer of mineral input and paleo-storm activity (Croudace and Rothwell, 2015).

Deleted: significantly




Figure 5. Aeolian Sand Influx (ASI) and elemental proxy analysis for (a) TLM and (b) TAC. Downcore profiles of ASI, log-transformed ASI, Ti, and K are shown by depth, with calibrated age provided as a secondary axis. Ombrotrophic intervals (Zone 3 for TLM; Zones 2 and 4 for TAC; see Fig. 3) are highlighted in black, while minerotrophic or uncertain intervals are shown in grey. Principal Component Analysis (PCA) was conducted on seven CLR-transformed elements, with ASI included as a passive variable, using only the ombrotrophic sections of each core.

Deleted: While both Ti and K correlate well with ASI and appear to be reliable proxies, Ti was selected as the primary geochemical indicator of aeolian sand input. This decision is based on the potential for K, as an essential plant nutrient, to accumulate in plant tissues—especially near the surface—potentially complicating its interpretation as an aeolian indicator (Steinmann and Shotyk, 1997; Shotyk, 1988).

Formatted: Caption

4.2.2 Paleo-storm reconstructions

800

We conducted separate paleo-storm reconstructions for TLM and TAC based on Ti measurements. These Ti-based event reconstructions are presented alongside the log-transformed ASI curves to emphasize their correspondence (Fig. 6).

At TLM, storm events were defined using a threshold set at 2.7% above the Ti moving average. Since 1035 BCE, the detrended Ti record exceeded this threshold 65 times (Fig. 6a), corresponding to an average of 2.3 events per century. Three multi-centennial periods show above-average storm frequency, with a peak of 6 events per century: 865–535 BCE, 500–750 CE, and 1510–1785 CE, while the recent period from 1945–2019 CE also shows above average event per century.

At TAC, the event threshold was set at 7.3 % above the Ti moving average. Excluding the non-ombrotrophic interval, the detrended Ti record surpassed this threshold 64 times since 2050 BCE (Fig. 6b), yielding an average of 3.2 events per century. Two prominent periods exceed the long-term average: 1320–1715 CE and 1815–2019 CE, with a maximum of 14 events per century during the early 20th century.

Deleted:

→ We did two separate paleo-storm reconstructions based on Ti measurements, one for each site. The TLM paleo-storm reconstruction (Fig. 5a) was based on its Zone 3 (1035 BCE–2019 CE; ~3050 years), while the TAC paleo-storm reconstruction (Fig. 5b) was based on its Zones 2–4 (2050 BCE–2019 CE; ~4070 years). Zones 1–2 of TLM and Zone 1 of TAC were excluded, as they each corresponded to early peat development stages when wetlands and/or fens conditions probably prevailed. Sediment delivery during these early periods was likely driven by non-aeolian processes, making them unsuitable for storm reconstruction. Additionally, the TAC core contained two fine-grained sand layers at 151 cm (600 BCE) and 136 cm (810 CE) (Zone 3, Fig. 3b) that corresponded to a very slow peat accumulation period or a hiatus (Fig. 2b). While similar sand deposits were not visible in TLM, the Ti data for this section of TAC (Fig. 5b) closely mirrored TLM's trend of events per century during the same interval (Fig. 5a), with heightened values between 800–600 CE and 600–900 BCE. Therefore, although this section of TAC lacked clear ombrotrophic conditions, we included it in our paleo-storm analysis due to its close similarity with the TLM data.

Formatted: Indent: First line: 1.27 cm

Formatted: Font: Not Bold

Formatted: Superscript

Formatted: English (CAN)




Figure 6. ASI (log), Ti (%), and storm events per century for TLM (a; in orange) and TAC (b; in purple). Solid black lines in (a) and (b) show ASI and Ti measurements during the periods of storm reconstruction, corresponding to the ombrotrophic portions of the core. Grey lines indicate the non-ombrotrophic portions of the cores, excluded from the reconstruction. Dashed red lines indicate the moving thresholds of event detection, calculated using the Tukey rule over a moving average (see Methods for details). Events are marked by orange (TLM) and purple (TAC) dots, with horizontal bars representing age uncertainty. Event frequencies are shown as a 100-year moving sum, with the average number of events per century shown as horizontal dashed lines.

Deleted: In TLM, the event thresholds were 63.6 Ti cps (1851–2019 CE) and 41.6 cps (pre-1851 CE) above the Ti moving average. Since 1035 BCE, the detrended Ti record exceeded the event threshold 66 times (Fig. 5a), yielding an average frequency of 2.3 events per century. Five periods exceeded the record's average events per century: 770–535 BCE, 605–800 CE, 1220–1320 CE, 1390–1785 CE, and 1955–2019 CE, with a maximum of ~6 events per century.⁴ For TAC, the event thresholds were 917.3 cps (1851–2019 CE) and 283.6 cps (pre-1851 CE) above the moving average. Since 2050 BCE, the detrended Ti exceeded the event threshold 70 times (Fig. 5b), corresponding to an average of 2.3 events per century. Four periods exceeded the record's average frequency: 835–685 BCE, 1315–1710 CE, 1750–1840 CE and 1880–2002 CE. Notably, TAC recorded a peak storm activity period centred around 1600 CE, with nearly 14 events per century, while the other periods exhibited 3–6 events per century. Importantly, the very low peat accumulation rate in TAC Zone 3 (135–159 cm, 880 BCE–870 CE) meant that the events per century during this time was almost certainly underestimated.⁴

Overall, the events per century reconstructions for both sites were remarkably similar, indicating enhanced mineral input to both peatlands, and hence storminess, during three main periods: 800–550 BCE, 600–800 CE, and 1300–1700 CE.⁴

... [8]

- Deleted:** 5
- Deleted:** Titanium (
- Deleted:**) counts per second (cps)
- Deleted:** vertical
- Deleted:** raw
- Deleted:** μ -XRF
- Deleted:** (except for the period corresponding to Zone 3 in TAC).
- Deleted:** (c) The annually-resolved Atlantic Multidecadal Variability (AMV) reconstruction based on titanium measurements from a varved lake record from Ellesmere Island, in Arctic Canada. An instrumental AMV record (1854–2023) is shown in dark blue (Huang et al., 2017).

900 5. Discussion

5.1 Peat development

Age-depth models indicate that peat initiation began around 5,000 years ago (Fig. 2). The basal sands in both cores are interpreted as marking the onset of peat accumulation, suggesting that the full organic sequences were recovered. These layers likely formed in a marine-influenced environment during a period of submergence before ca. 10.7 ka PB (Rémillard et al., 2017), though further data would be needed to confirm the depositional setting. The overlying organic-rich sediments are characteristic of early peatland development. At TLM, the presence of *Menyanthes trifoliata* seeds within mineral-rich lignous peat suggests that the site initially formed through the terrestrialization of a shallow water body (Hewett, 1964; Kuhry and Turunen, 2006). In contrast, the basal peat at TAC, composed of mostly herbaceous remains and lacking *Menyanthes trifoliata* seeds, suggests that the peatland formed directly on a mineral substrate in a well-drained environment, likely through primary paludification — a direct shift from bare land to peatland, without an intermediary wetland step (Kuhry and Turunen, 2006).

The fen-to-bog transition in TLM and TAC was characterized by the increasing dominance of *Sphagnum* mosses and a positive correlation between mineral content and lithogenic elements such as Ti and K. However, the timing of this transition differed substantially, with TAC transitioning around 2050 BCE and TLM following over a millennium later, around 1000 BCE. This temporal discrepancy may reflect differences in the initial conditions of peat formation at each site. Additionally, the smaller size of TAC, which currently extends over 9 ha compared to TLM's 40 ha, likely rendered it more sensitive to climate and hydrological changes, such as the sharp increase in mineral content around 2600 BCE, as has been found in a study from Sweden (Sjöström et al., 2020). This interpretation is supported by TAC's lower peat accumulation rate, shallower ^{210}Pb profile, and lower ^{210}Pb influx compared to TLM, suggesting that TAC is more exposed, has reduced snow accumulation, and, consequently, has more unstable hydrological conditions (Perrier et al., 2022).

A notable aspect of peat development at TAC are the two distinct mineral-rich layers in Zone 3, which are absent in the TLM record. The timing of these mineral intervals should nevertheless be interpreted with caution due to the very low resolution of the TAC age model in this zone. While the older mineral-rich layer is fairly well constrained by two radiocarbon dates spanning from 970–475 BCE, the younger layer is only constrained by one upper radiocarbon date at 945 CE (Table 1). Although their origin remains uncertain, they are unlikely to represent single depositional events. TAC is situated ~20 m above sea level and, based on sea-level indicators, was even higher during the late Holocene (Rémillard et al., 2016; Barnett et al., 2017), making marine overwash an improbable explanation. Likewise, an aeolian origin seems unlikely, as it would presumably have left a similar signal at TLM, which is not observed.

Although these layers are not reflected in TLM's mineral content or ASI records, their timing coincides with increased Ti extremes in TLM's storm reconstruction (Fig. 6a), suggesting that mineral inputs at TAC and Ti extremes at TLM may be linked to broader climatic variability. Indeed, the onset of slow peat accumulation at TAC coincides with the 2.8 ka event, a centennial-scale period generally characterized by cool and often dry conditions in the circum-North Atlantic (Van Geel et al., 1996) that was recorded in nearby peatlands from Prince Edward Island (Peros et al., 2016), Anticosti Island (Perrier et al.,

Deleted: 1

Deleted: Key similarities and differences between TLM and TAC

TLM and TAC showed storm signals that were generally consistent, suggesting that peatbogs can serve as reliable archives for paleo-storm reconstructions. Our reconstructions revealed three main periods of enhanced storm activity shared by both sites: 1300–1700 CE, 600–800 CE, and 800–550 BCE. However, notable differences in mineralogy and geochemistry between the two cores deserve further considerations to improve the interpretation of storm records. 1

5.1.1 Peat development

→ The development of peat at TLM and TAC reflects contrasting formation processes, hydrological sensitivities, and sedimentary influences, highlighting differences in their initial formation, later transitions, and responses to climatic and local environmental changes.

Deleted: (Rémillard et al., 2016; Barnett et al., 2017)

Deleted: (Perrier et al., 2022)(Peros et al., 2016)(Rémillard et al., 2017)

Formatted: Indent: First line: 1.27 cm

Deleted: .

Deleted: further

Deleted: Another notable difference

Deleted: in peat development was

Deleted: the presence of

Deleted: sand

Deleted: at TAC at depths corresponding approximately to 600 BCE (151 cm) and 810 CE (136 cm)

Formatted: Indent: First line: 1.27 cm

Deleted: , which are absent at TLM. The lack of abrupt contacts between the sand and the underlying and overlying peat (Appendix D) supports the interpretation that these units represent gradual accumulations of sand over time rather than short-lived deposition events. The presence of such prominent sand-layers may be due in part to the proximity of TAC to the coast and a sediment source (Fig. 1e). However, a more likely explanation is that these units represent an *in-situ* concentration of mineral matter due to a period of enhanced peat decomposition, possibly related to a change in hydrology that affected only TAC. Its relatively small size and its orientation perpendicular to a downward-sloping elevation gradient may have rendered this site more sensitive to changes in groundwater flow compared to TLM.

Deleted: T

Deleted: beginning

Deleted: of this period

Deleted: d

Deleted: , which was a

Deleted: . Just to the south, Baltic Bog, at Prince Edward Island, experienced a period of reduced peat accumulation or a hiatus around this time

2022), and the north shore of the Gulf of St. Lawrence (Magnan, 2014; Primeau and Garneau, 2021). The more pronounced response at TAC compared to TLM may reflect greater sensitivity to climate-driven hydrological shifts at this site, potentially due to TAC's smaller size and orientation perpendicular to a downslope gradient, which may have made it more vulnerable to groundwater fluctuations.

Deleted: and a similar event is recorded at a peatland in the Netherlands (Van Geel et al., 2014). While the specific mechanism behind the formation of these sand layers remains unclear, their timing and context suggest a complex interplay between local hydrology and broader climatic changes, necessitating further investigation. ...

Formatted: Font: Font color: Text 1, English (CAN)

Deleted: 1.

Deleted: Aeolian sediment sources

Formatted: Font: Not Bold

5.2 Modern aeolian sediment sources

TLM and TAC exhibit distinct patterns of aeolian sediment input (Fig. 5-6), reflecting differences in sediment availability and local geomorphology. In the modern period (post-1851 CE), ASI values at TLM are an order of magnitude lower than those at TAC (Fig. 4), even after adjusting the grain-size threshold (63 μm for TLM vs. 125 μm for TAC). This disparity likely reflects TLM's greater distance from active sediment sources: TLM lies \sim 2 km from the nearest beach, while TAC is situated only \sim 210 m from eroding south-facing sandstone cliffs (Figs. 1d–e). Although the precise sources cannot be confirmed without sedimentological or geochemical fingerprinting, proximity strongly suggests that the closest sources contribute most of the aeolian material to each site.

At TAC, the nearby sandstone cliffs contain a highly erodible unit known as the *Drift des Demoiselles*, a friable red sandstone rich in hematite (Fe_2O_3)-coated quartz (SiO_2), K-feldspar (KAlSi_3O_8), and Ti-bearing minerals derived from underlying basaltic rocks (Brisebois, 1981; Dredge et al., 1992; Rémillard et al., 2013). These cliffs retreat at an average rate of 0.3 m/yr (Bernatchez et al., 2012b), supplying a steady stream of sediment to coastal systems, which is then redistributed by longshore drift to local beaches (Morin, 2001). Importantly, the hematite coatings are typically removed during beach transport, meaning that Fe-rich grains are more likely to represent relatively non-weathered, locally sourced sediment from cliffs rather than beach-derived material.

While both cores show a strong association among ASI, Ti and K (Fig. 5), elements common in sandstone and beach-derived sands, the geochemical signals diverge in the modern period, suggesting different sediment sources. At TAC, positive correlations among ASI, Si, and Fe (Appendix Table E1) support a dominant input from nearby sandstone cliffs. Silicon is enriched in coarser grain-size fractions (Liu et al., 2019), which are more likely to be transported over short distances during storm events. The Fe enrichment further points to the presence of hematite-coated grains, characteristic of cliff-derived sediment.

In contrast, the TLM record suggests a primary source from local beaches, which are depleted in hematite due to longer transport pathways and reworking. This is consistent with TLM's greater distance from any major cliff exposures. However, beaches can still serve as significant aeolian sediment sources under storm conditions. Local observations show that under normal conditions, sand from exposed dune scarps is often transported inland and deposited in back-dune environments, contributing to landward dune migration (Morin, 2001). This supports the interpretation that, during storms, aeolian transport carries beach sand even farther inland and reaches the TLM bog, especially given its proximity to extensive beach ridge systems to the east (Fig. 1).

5.3 Ti and ASI as proxies for aeolian input and storms

5.3.1 2050 BCE – 1851 CE

Our paleo-storm reconstructions integrate ASI variability but rely primarily on identifying Ti extremes as a high-resolution proxy for aeolian input (Fig. 6).

← Formatted: Heading 3, Indent: First line: 0 cm

1025 During the early ombrotrophic phase at TAC (ca. 2050–880 BCE), both ASI and Ti values remain low, suggesting limited or distant sediment supply (Fig. 6b). This interpretation aligns with the timing of beach ridge development on Havre-Aubert Island, which began around 550 years ago (Rémillard et al., 2015). Similarly, the cliffs south of TAC, now eroding at ~0.3 m/yr (Bermatchez et al., 2012b), may have been located further seaward and thus would have been a less accessible 1030 sediment source at the time. The below-average storm frequency inferred for this period should therefore be interpreted with caution, as it may partly reflect limited sediment availability rather than reduced storm activity alone.

Deleted: (Kylander et al., 2011; Bertrand et al., 2023) →

1035 At TLM, ASI remained close to zero between 1035 BCE and 1000 CE (Fig. 6a), suggesting limited sediment availability. During this same interval, TAC shows similarly low ASI values. While other sedimentary processes may have influenced deposition at TAC during this period, the consistently low ASI at both sites suggests that local sediment sources were either inactive or insufficient. Nevertheless, the Ti record at TLM during this time shows numerous peaks, the largest of 1040 which correspond to ASI highs (Fig. 6a), but many smaller peaks are unmatched in the ASI records. This discrepancy may highlight a limitation of ASI: the grain-size threshold used to define aeolian input may be too coarse to capture the full range of storm-related sediment delivery. For examples, Vandel et al. (2019) found that ASI based on coarse sand with low overall grain amounts poorly reflected modern storm activity in Baltic bogs, while including finer grains improved the correlation. At TLM, ASI likely captures only the most intense aeolian events, whereas Ti reflects a broader range of storm-related inputs.

1045 After ca. 1000 CE, both sites show a slight increase in ASI variability, with a marked rise after ca. 1200 CE (Fig. 6). This shift may reflect intensified storm activity, drier climatic conditions enhancing aeolian transport, and/or the activation of new sediment sources, such as the beach ridges and sand spits east of Havre-Aubert (Fig. 1). The continued retreat of the southern sandstone cliffs may also have increased sediment availability, particularly at TAC. While ASI and Ti trends remain well-aligned at TAC, they diverge at TLM between 1200 and 1500 CE, when several ASI peaks are not mirrored in the Ti record (Fig. 6a). Given the strong ASI–Ti correlation at TAC, and the overall consistency of Ti trends at both sites, we place 1050 confidence in the Ti signal at TLM during this interval.▼

Deleted: TLM and TAC exhibited distinct patterns in aeolian sediment input, reflecting differences in their proximity to sediment sources and local geological settings.

ASI values decline between ca. 1600 CE and ca. 1750 CE at TAC, and until ca. 1880 CE at TLM, despite coinciding with the onset of beach ridge development in the region. Given the proximity of active sediment sources by this time, particularly beaches near TLM and eroding cliffs near TAC, this dip more likely reflects climatic controls, such as wetter conditions or reduced storm frequency, than sediment supply limitations. At TAC, Ti and ASI remain closely aligned, while at TLM, Ti suggests higher average storm frequency between 1510 and 1785 CE than ASI would indicate, reinforcing the idea that ASI may underestimate mineralogenic input when absolute values are very low.

Deleted: (Vandel et al., 2019)

5.3.2 1851–2019 CE

1060 From the late 18th century at TAC and post-1880 at TLM, ASI increases markedly (Fig. 6). While this may reflect continued storminess and increased sediment availability, it may also be influenced by human-driven landscape changes. Permanent settlement on the Magdalen Islands began in the mid-18th century and expanded after the mid-19th century (Fortin and Larocque, 2003), likely increasing sediment exposure through land clearance, grazing, and infrastructure development. For instance, the ASI between between 1915 and 1970 CE at TAC, not seen at TLM (Fig. 7a–b), may relate to 20th-century coastal gravel road construction near TAC (Fortin and Larocque, 2003).

1065 From 1851 CE onward, ASI and Ti records can be compared with storm tracks and intensities from the International Best Track Archive for Climate Stewardship (IBTrACS) dataset (Knapp et al., 2010; Gahtan et al., 2024) (Figs. 7c and 8; Appendix F). Since 1851 CE, 38 tropical cyclones have passed within 100 km of the archipelago, with wind speeds often reaching 75–125 km/h. These cluster around 1870–1900 CE, 1920–1950 CE, and ~ 2000 CE. The ASI and Ti records at TAC broadly reflect these trends, with 17 events marked by Ti extremes (Fig. 7b). In contrast, TLM shows only five Ti extremes, with four occurring between ca. 1980 and 2005 CE (Fig. 7a). This suggests that storm signal strength depends heavily on sediment availability and proximity, as seen at TAC.

1070 Attributing individual storms to specific peaks remains difficult. Six of the IBTrACS tropical cyclones had significant local impacts as reported in the newspaper and ship logs (Fig. 7c; Appendix F). For example, Hurricane Dorian (2019 CE), among the region's strongest recent storms (Jardine et al., 2021), is weakly expressed at TLM (possibly the surface ASI peak) and absent at TAC. Hurricane Blanche in 1975 CE (Public Safety Canada, 2024) is not detected at either sites. A major Ti peak around 1888 CE at TAC and 1886 CE at TLM may reflect the August Gale of 1873 CE, one of the most destructive storms on record for the Magdalen Islands (Landry, 1989, 1994), although increasing chronological uncertainty before 1930 CE limits confident attribution. Additionally, some Ti peaks may reflect unlisted events in IBTrACS, such as strong nor'easters and winter storms. For instance, the Ti peak at TLM around 1995 CE may correspond to a nor'easter on 7 November 1994, while peaks in both ASI and Ti around 1990 CE could relate to winter storms on 4 December 1989 and 16 December 1990 (Bernatchez et al., 2012a).

1075 Chronological uncertainty may arise from the downward movement of sand and mineral particles within the peat column due to bioturbation, water flow, or settling. For example, aeolian sands from Hurricane Dorian could have been displaced downward, potentially contributing to the prominent peak around 2010 CE. Such post-depositional movement could misdate an event or smear it across multiple depths. However, vertical migration appears limited in ombrotrophic peat. Although direct evidence is scarce, Allen et al. (2021) found that microplastic records in Pyrenean peat closely follow historical production, suggesting minimal post-depositional movement of similarly sized particles. Additionally, high modern accumulation rates at TLM (1.5–2 cm/yr) and TAC (0.45 cm/yr) reduce the likelihood of significant disturbance. As peat compacts with depth and becomes more isolated from surface processes, this risk further diminishes, though it cannot be entirely excluded.

Deleted: (Fortin and Larocque, 2003) TLM consistently displayed lower mineral content than TAC, particularly during ombrotrophic periods. For instance, in the modern period (post-1851 CE), ASI values at TLM were an order of magnitude smaller than those at TAC, even after adjusting the grain-size threshold to 63 µm for TLM compared to 125 µm for TAC (Fig. 3). This disparity likely reflects TLM's greater distance from sediment sources: TLM is situated 2 km from the nearest beach, while TAC is only 210 m from south-facing sandstone cliffs (Figs. 1d–e). The surface geology of the Magdalen Islands suggests that aeolian sediments in this region primarily originate from local beaches and sandstone cliffs, which are rich in quartz (SiO_2), k-feldspar (KAlSi_3O_8), and Ti-bearing minerals from basaltic rocks (Brisebois, 1981; Sabina, 2003; Rémillard et al., 2016). While both cores showed a strong association between ASI and both Ti and K – elements common in sandstone and beach-derived sediments — the geochemical signature diverged during the modern period. At TAC, the positive correlation of Si and Fe with ASI and mineral content (Table E1) supports the interpretation that sandstone cliffs serve as the primary sediment source. Silicon is enriched in coarser grain-size fractions (Liu et al., 2019), which are more likely to be transported over short distances during storms. Furthermore, the sandstone cliffs are rich in hematite- (Fe_2O_3) -coated quartz (Sabina, 2003), which would explain the elevated Fe content at TAC. In contrast, TLM's sediment appeared to be derived primarily from sand beach sands, which are depleted in hematite, and is consistent with the site's greater distance from the sandstone cliffs. Ti emerged as the common aeolian sand indicator in both TLM and TAC. While Ti and K consistently correlated with ASI in both cores, K, as an essential plant nutrient, can accumulate in plant tissues, potentially

... [91]

Deleted: 2

Deleted: Modern storm attribution (

Deleted: -

Deleted:)

Deleted: → To validate the relationship among ASI, Ti, and storm events, ASI and Ti data were compared with storm tracks an

... [10]

Formatted: Indent: First line: 1.27 cm

Deleted: The strongest storms affecting the Magdalen Islands are tropical cyclones, which typically track towards the north an

... [11]

Moved up [1]: The larger size of TLM and its greater distance to mineral sources (Fig. 1) likely explains this discrepancy, though

Moved (insertion) [1]

Deleted: The ASI and Ti record for 1851–2019 CE (Figs. 7a–b) show notable correspondence between the two sites, particul

... [12]

Deleted: →

Deleted: The relationship among ASI, Ti, and IBTrACS data (Fig. 7a–e) is more complex. For instance, the most recent major

... [13]

Deleted: .

Deleted: At TAC, one of these storms may correspond to the Ti and ASI peaks around 1990. Earlier, the mineral increase and

... [14]

Formatted: Indent: First line: 1.27 cm

Deleted: 1

... [15]

Deleted: Additionally, it seems that not all major events are captured; for instance, Hurricane Blanche in 1975 (Public Sa

... [16]





Figure 7. (a–b) Modern (1851–2019 CE) storm proxies for TLM (a; in orange) and TAC (b; in purple): the Aeolian Sand Index (ASI; $\text{g m}^{-2} \text{yr}^{-1}$) (grey silhouettes) and Ti measurements (CPS) (black lines). Dot-dashed red lines indicate the moving thresholds of event detection, calculated using the Tukey rule (see Methods for details). Events are marked by orange (TLM) and purple (TAC) dots, with horizontal bars representing age uncertainty. [The age in CE is based on our age-depth models](#). (c) Bar chart showing historical wind data from the IBTrACS dataset (Knapp et al., 2010; Gahtan et al., 2024) showing storms passing within 100 km of the Magdalen Islands and their peak winds (km h^{-1}). Storms with documented impacts in local newspapers and shipwreck records are highlighted in red. The Great August Gale of 1873, highlighted in green, was not in the IBTrACS dataset, but had known impacts on the Magdalen Islands. (d) Instrumental time-series of the Atlantic Meridional Mode (AMM; turquoise) from 1948–2023 (Chiang and Vimont, 2004) and of the Atlantic Multidecadal Variability (AMV; in dark blue) from 1854–2023 (Huang et al., 2017). [The dates in CE are calendar dates](#).

Deleted: <object>

Figure 6. (a) Map showing the location of the Magdalen Islands (pink star — this study) and tracks of storms passing within 100 km, based on the IBTrACS dataset from 1851–2019 CE (Knapp et al., 2010; Gahtan et al., 2024). Storms known to have impacted the Magdalen Islands are highlighted in red. Some paleo-storm records referenced in this paper are marked on the map: A (green circle) — Robinson Lake, New Brunswick, Canada (Patterson et al., 2022); B (orange circle) — Robinson Lake, Nova Scotia, Canada (Oliva et al., 2017); C and D (blue circles) — Mattapoisett Marsh (Boldt et al., 2010) and Salt Pond, Massachusetts (Donnelly et al., 2015); E–I (purple circles) — the studies compiled in the Bahamas compilation: Grand Bahamas (Winkler et al., 2023), Thatchpoint Blue Hole, Abaco Island (Winkler et al., 2020), South Andros (Wallace et al., 2019), Long Island (Wallace et al., 2021b), and Middle Caicos (Wallace et al., 2021a). (b) Enlarged view of the Magdalen Islands region, showing storm tracks with documented local impacts. (c) Bar chart showing peak winds (km h^{-1}) of the storms indicated in panels (a) and (b) as they passed within 100 km of the Magdalen Islands. The Great August Gale of 1873, highlighted in green, was not in the IBTrACS dataset, but had known impacts on the Magdalen Islands. Imagery panel (b): © Google Earth Pro 7.3.6.9796 (2024), Gulf of St. Lawrence, Canada. Image date 31 December 2023, © Landsat/Copernicus. [\[17\]](#)

... [17]

Deleted: 7



1335 **Figure 8.** (a) Map showing the location of the Magdalen Islands (pink star — this study) and tracks of storms passing within 100 km, based on the IBTrACKS dataset from 1851–2019 CE (Knapp et al., 2010; Gahtan et al., 2024). Storms known to have impacted the Magdalen Islands are highlighted in red. Some paleo-storm records referenced in this paper are marked on the map: A (black circle) — Robinson Lake, New Brunswick, Canada (Patterson et al., 2022); B (blue circle) — Robinson Lake, Nova Scotia, Canada (Oliva et al., 2017); C and D (green circle) — Mattapoisett Marsh (Boldt et al., 2010) and Salt Pond, Massachusetts (Donnelly et al., 2015); E–I (purple circles) — the studies compiled in the Bahamas compilation: Grand Bahamas (E) (Winkler et al., 2023), Thatchpoint Blue Hole (F), South Andros (G) (Wallace et al., 2019), Long Island (H) (Wallace et al., 2021b), and Middle Caicos (I) (Wallace et al., 2021a). (b) Enlarged view of the Magdalen Islands region, showing storm tracks with documented local impacts. (c) Bar chart showing peak winds (km h^{-1}) of the storms indicated in panels (a) and (b) as they passed within 100 km of the Magdalen Islands. The Great August Gale of 1873, highlighted in green, was not in the IBTrACS dataset, but had known impacts on the Magdalen Islands. Imagery panel (b): © Google Earth Pro 7.3.6.9796 (2024), Gulf of St. Lawrence, Canada. Image date 31 December 2023, © Landsat/Copernicus.

5.6 Western North Atlantic Basin comparison

The TLM and TAC records together offer a 4000-year chronology of storm activity for the Magdalen Islands, adding critical insights to the limited records available from the northern extent of the North Atlantic hurricane track. Despite being 1350 aeolian reconstructions, the Magdalen Islands records show notable similarity with marine overwash records from lakes and blue holes spanning the past 2000 years from eastern Canada, the US, and the Bahamas (Fig. 9). This north-south transect across the western North Atlantic reflects modern storm tracks from the Magdalen Islands (Fig. 8a) and enable broader regional comparisons.

During the instrumental period (1851 CE–present), the storm records from the Bahamas to the Magdalen Islands 1355 show significant heterogeneity (Fig. 9). The only temporally overlapping record from eastern Canada, at Harvey Lake, New Brunswick, also indicates elevated storms activity after ~1895 CE (Patterson et al., 2022). While the Harvey Lake study does not calculate events per century and is thus excluded from Fig. 9, it corroborates trends in TLM and TAC. In contrast, the

Deleted: 3

Deleted: 8

Deleted: 6

Deleted: 8

Deleted: TLM and TAC documented increased storm activity, particularly in the 20th century (Figs. 8a–b).

Deleted: 8

1365 record from Salt Pond, Massachusetts, does not show a comparable increase (Donnelly et al., 2015) (Fig. 9d), though a nearby record from Mattapoisett Marsh, just 18 km south of Salt Pond, documents numerous 20th century hurricane strikes (Boldt et al., 2010). Donnelly et al. (2015) attribute this discrepancy to Salt Pond's avoidance of several modern hurricanes. Similarly, the Bahamian compilation, aggregating five blue hole storm records across the archipelago, shows a quiet 20th century (Fig. 9e). This trend reflects mostly inactivity at the southern sites (South Andros, Long Island, and Middle Caicos), while northern 1370 sites (Grand Bahamas and Abaco) exhibit increased storm activity (Wallace et al., 2021a; Winkler et al., 2023).

Deleted: 8

The Little Ice Age (LIA, ~1300–1850 CE) (Magnan and Garneau, 2014) stands out as an active hurricane period across the western North Atlantic. At TLM, increased storm activity persisted between 1510–1785 CE (Fig. 9a). Similarly, TAC shows heightened activity 1320–1715 CE, with a calm period between 1400–1435 CE, and a peak between 1500–1600 CE (Fig. 9b). These trends align with Robinson Lake, Nova-Scotia, which shows increased storm activity between 1475–1670 1375 CE (Oliva et al., 2018) (Fig. 9c), and Harvey Lake, New Brunswick, where coarse silt beds were deposited between 1630–1640 CE (Patterson et al., 2022). Salt Pond also experienced elevated storm activity with 35 events between 1500–1600 CE (Donnelly et al., 2015) (Fig. 9d). In the Bahamas, storm activity peaked between 1570–1670 CE (Fig. 9e), matching patterns observed in the Magdalen Islands and nearby sites (Winkler et al., 2023). Notably, none of these records document storm activity during the early LIA (~1200–1400 CE), as seen in the Magdalen Islands.

Deleted: 8

1380 During the Medieval Climate Anomaly (MCA), which lasted between ~ 900–1300 CE in the Gulf of St. Lawrence (Magnan and Garneau, 2014), storm activity decreased across the Magdalen Islands, consistent with calmer periods at Salt Pond and in the Bahamas (Fig. 9d–e). However, brief upticks between 1000–1100 CE at Salt Pond and 980–1030 CE in the Bahamas contrast with calm conditions in the Magdalen Islands. In the Bahamas, the increase is attributed to southern sites like Southern Andros (Wallace et al., 2019) and Middle Caicos (Wallace et al., 2021a), while northern sites, such as the Grand 1385 Bahamas, recorded a decrease in storm frequency from 550–1090 CE (Winkler et al., 2023) (see Fig. 8a for location). Between 500–750 CE, storm frequency at TLM aligns with increased storm activity in the Bahamas (720–820 CE), while Salt Pond documented an earlier active period between 300–600 CE, partly overlapping with the TLM record (Fig. 9c). During this period, sediment delivery at TAC was also elevated (Fig. 6b). Unfortunately, the lack of pre-LIA eastern Canadian records limits our understanding of storm dynamics earlier in the record, while the shorter duration of the Salt Pond and Bahamas 1390 reconstructions (<2000 years) restricts longer-term comparison with the Magdalen Islands data. Overall, the similarities between our Magdalen Islands reconstruction and records from the eastern seaboard and the Bahamas lend additional confidence to our interpretations of ASI and Ti as storm proxies.

Deleted:), which lasted between

Deleted:

Deleted: in the Gulf of St. Lawrence

Deleted: 1220–1785 CE

Deleted: , interrupted only by a calm interval between 1320–1390 CE ...

Deleted: 8

Deleted: 15

Deleted: 0

Deleted: 7

Deleted: 8

Deleted: 6

Deleted: 600

Deleted: 800

Deleted: sediment delivery a

Deleted: t

Deleted: and TAC

Deleted: elevated sediment delivery at TAC

Deleted: 8




Figure 9. Events per century (Event/cent) from the Magdalen Islands and nearby regions reconstructions. (a–b) 100-year moving sum of storm events at TLM (orange line) and TAC (purple line). (c–e) Comparisons with previously published paleo-storm reconstructions. (c) 100-year moving sum of storm overwash events from Robinson Lake, Nova Scotia, between 1210–1925 CE (Oliva et al., 2017). (d) 100-year moving sum of storm overwash events from Salt Pond, Massachusetts, spanning the past 2000 years

(Donnelly et al., 2015). (e) A smoothed 1500-year compilation of storm records from five Bahamian blue hole records, with the shaded area representing the confidence interval (Wallace et al., 2019; Winkler et al., 2020; Wallace et al., 2021a; Wallace et al., 2021b; Winkler et al., 2023). The location of the studies in a–e is shown in Fig. 6a. (f) Smoothed 1500-year compilation of storm records from 12 regions spanning from Belize to New England from Yang et al. (2024). (g) The annually-resolved **Atlantic Multidecadal Variability (AMV) reconstruction based on titanium measurements from a varved lake record from Ellesmere Island, in Arctic Canada. An instrumental AMV record (1854–2023) is shown in dark blue (Huang et al., 2017).**

1435

5.7 Regional insights into long-term drivers of storm activity

Storm records from the Magdalen Islands, eastern Canada, New England, and the Bahamas reveal consistent patterns suggesting that our Magdalen Islands records capture storms originating in the Main Development Region (MDR), located in 1440 the tropical Atlantic off West Africa. These storms typically track through the Bahamas before recurring towards the US east coast and eastern Canada (Kossin et al., 2010), a pattern consistent with the IBTrACS storm tracks observed in the Magdalen Islands since 1851 CE (Fig. 8a).

Hurricane activity in the MDR is strongly influenced by the Atlantic Meridional Mode (AMM), a coupled ocean-atmosphere system driven by variations in tropical Atlantic sea surface temperatures (SSTs) and vertical wind shear (VWS) 1445 (Kossin and Vimont, 2007; Vimont and Kossin, 2007; Klotzbach and Gray, 2008; Kossin, 2017; Ting et al., 2019). Positive AMM phases, characterized by elevated SSTs and reduced VWS, creates favourable conditions for storm formation and intensification (Kossin, 2017). **Although long-term AMM reconstructions are unavailable, the AMM is positively correlated with the Atlantic Multidecadal Variability (AMV) (Fig. 7d), and variations in the AMV are thought to indirectly modulate long-term hurricane activity by influencing AMM trends** (Kossin and Vimont, 2007; Vimont and Kossin, 2007). The annually 1450 resolved AMV reconstruction from Lapointe et al. (2020) identifies a strong negative AMV during the LIA (not conducive to hurricane formation) and a more positive phase during the MCA (conducive to hurricane formation) (Fig. 9a). This trend is corroborated by other coarser-resolution AMV reconstructions (Mann et al., 2009b; Wang et al., 2017).

Nevertheless, storm records from eastern Canada (including **that** from the Magdalen Islands), New England, and the 1455 Bahamas show **the opposite trend: increased storminess during the LIA and reduced activity during the MCA**. For instance, Mann et al. (2009a), in a statistical reconstruction of paleo-hurricane frequency across the North Atlantic, estimated low hurricane counts during the LIA and higher counts during the MCA. **A recent synthesis of proxy-based paleo-storm reconstructions spanning from Belize to New England (Yang et al., 2024) supports this, showing elevated storm frequency from 900–1200 CE, following by a calm period from 1250–1700 CE, interrupted only by a peak in activity between 1375– 1475 CE (Fig. 9f). This contrasts with widespread evidence from eastern Canada, New England, and the Bahamas showing frequent storms throughout the LIA. This persistent antiphase with the tropical Atlantic suggests that regional factors, such as storm track shifts or local intensification, may dominate at higher latitudes during periods of low tropical storm activity.**

Several studies support this view. Patterson et al. (2022) and Oliva et al. (2018) proposed that elevated storm activity 1460 during the early LIA might reflect local warmer SSTs in the Gulf of Maine and the Sargasso Sea, despite cooler temperatures in the MDR. Dickie and Wach (2024) provide a concise summary of these mechanisms, highlighting that while the LIA experienced colder winter and lower annual mean temperatures, warmer summers may have been sufficient to fuel local storm

Deleted: Compilation methods are detailed in Wallace et al. (2021a).

Deleted: 4

Deleted: L

Deleted: climate drivers

Deleted: 5

Deleted: While

Deleted: long-term AMM reconstructions are currently unavailable,

Deleted: its correlation to the Atlantic Multidecadal Variability (AMV), a basin-wide pattern of decadal Atlantic SSTs fluctuation, provides an indirect mean of assessing long-term hurricane trends. The instrumental AMV record correlates positively with instrumental AMM data (Fig. 7d), and the AMV has been extensively reconstructed using proxies (e.g., Fig. 5c). Variations in AMV are thought to modulate long-term hurricane activity indirectly by exciting trends in AMM

Deleted: , based on a varved lake in the Canadian Arctic,

Deleted: 5

Deleted: c

Deleted: our own

Deleted: widespread agreement regarding heightened activity during the LIA, and moderate consensus on reduced activity during the MCA, which is contrary to what would be expected based on the AMV record.

Deleted: Additionally, a

Deleted: 12 regions

Deleted: e results from Mann et al. (2009a),

Deleted: 8

Deleted: 1

Deleted: Interestingly, the Yang et al. (2024) compilation contrasts markedly with our records, as well as others from eastern Canada, New England, and the Bahamas. This persistent antiphase pattern, alongside a negative correlation to the AMV (which typically promotes hurricane formation), suggests that during periods of suppressed hurricane activity in the North Atlantic – such as the LIA – local factors steered the fewer storms that formed towards the northern Bahamas, the US east coast, and eastern Canada. Our findings indicate that storm frequency in these regions is influenced less by factors driving storm genesis in the tropical Atlantic (e.g., AMM and AMV), and more by processes that either intensify storms locally or shift storm tracks towards higher latitudes. For example,

Formatted: Indent: First line: 1.27 cm

intensification in mid-latitude North America. Ting et al. (2019) documented a dipole pattern over the past 150 years, where positive AMV conditions in the MDR coincide with unfavourable conditions along the U.S. east coast—and vice versa—possibly shifting storm activity northward during negative AMV phases.

Deleted: s

Deleted: ¶

→ Supporting this hypothesis,

1510 The Magdalen Islands records add significant value to western North Atlantic storm reconstructions. They extend evidences suggesting that mid- to high-latitude storm activity is governed less by conditions in the MDR and more by regional controls such as SST gradients, storm steering, and local intensification. As a long, uninterrupted record from a previously underrepresented region, our data help bridge the spatial gap between well-studied tropical sites and the North American coast, 1515 offering new insights into the climatic drivers of storm variability over centennial to millennial scales.

6. Conclusion

1520 The Magdalen Islands, located near the centre of the Gulf of St. Lawrence in Atlantic Canada, have been impacted by post-tropical cyclones and other storm events over the last century. Our new 4000-year storm reconstruction, based on aeolian inputs in ombrotrophic peat, provides one of the longest and most northerly continuous paleo-storm records in the western North Atlantic, and in doing so offers critical insights into storm dynamics at the northern limit of the North Atlantic hurricane track.

1525 Across the western North Atlantic basin, our records align with regional reconstructions from eastern Canada, New England, and the Bahamas, confirming increased storm activity during the Little Ice Age (1300–1850 CE) and reduced activity during the Medieval Climate Anomaly (~900–1300 CE). Notably, these storm patterns show an antiphase relationship with tropical Atlantic variability, such as the AMM and the AMV, indicating that regional factors, including local sea surface temperature gradients, and storm intensification mechanisms, may play a dominant role in modulating storminess at higher latitudes, rather than tropical cyclone formation alone.

1530 The similarity between our work and marine overwash records from eastern North America demonstrates the complementary value of aeolian and peatland archives. These records enrich the paleo-storm literature by providing additional perspectives on past storm activity, including information on wind strength and direction. Moreover, they expand the range of potential sites for paleo-storm reconstructions, particularly in mid-latitude regions where data remain scarce.

1535 Our results also highlight important site-specific challenges in interpreting peat-based storm records. Despite their geographic proximity, the two study sites, TLM and TAC, exhibited critical differences in sediment input and signal strength. These differences stem largely from variations in bog size, proximity to sediment sources, and hydrological sensitivity. TAC, being smaller and closer to eroding sandstone cliffs, displayed a stronger and more variable storm signal sensitive to long-term environmental changes. In contrast, TLM's record showed a more muted but consistent signal over time, likely due to its greater distance from sediment sources. This underscores the need to carefully consider site selection and local geomorphological context in future research.

Deleted: identified a dipole pattern over the last 150 years, where favourable hurricane formation conditions in the MDR (positive AMV: higher SSTs, reduced VWS) coincide with unfavourable conditions along the US east coast (cooler SSTs, higher VWS), and vice versa. If this dipole operated during the LIA, regional conditions along the US east coast may have intensified hurricanes locally, while during the MCA, higher overall hurricane activity likely directed more storms towards the Gulf of Mexico and Caribbean. Indeed, a 1700 year-long western Florida compilation shows frequent storms between 750–1300 CE and lower activity between 1300–1800 CE (Wallace et al., 2021a; Winkler et al., 2023). This antiphase pattern seems to persist over the long-term with the Magdalen Islands records: a 5500-year record from Mullet Pond, in the Florida panhandle, indicates active periods between 150–350 CE, 150–1 BCE, 550–400 BCE and 950–800 BCE (Wang et al., 2024), which all correspond to quiet intervals at the Magdalen Islands (Fig. 5).

Deleted: ¶

→ Finally, another key factor influencing hurricane tracks is the position and strength of the North Atlantic subtropical high, also known as the Bermuda High. Modern observations show that a strengthened and westward-expanding Bermuda High drives hurricanes towards the Gulf of Mexico, while a weaker, eastward-shifted Bermuda High favours recurring tracks towards the US east coast (Kossin et al., 2010; Colbert and Soden, 2012). However, the absence of long-term reconstructions of Bermuda High dynamics limits our ability to analyse its influence over centennial to millennial timescales, as highlighted in several studies (Wallace et al., 2021b; Breyg et al., 2022). ¶

Formatted: English (CAN)

1570 Future research should also focus on elucidating the climatic mechanisms driving the storm variability recorded here, which is vital for projecting future storm impacts in eastern Canada. Further work is also needed to better understand depositional processes of sand in peat environments and to develop improved modern calibration techniques. Despite these challenges, paleo-storm studies such as ours remain critical for extending the temporal scope of storm records and enhancing understanding of storm-climate interactions in sensitive coastal regions.

1575 Data availability

All raw data can be provided by the corresponding author upon request. Paleo-storm reconstruction data are available on the *Borealis* data repository (<https://borealisdata.ca/>).

Author contributions

1580 AL, MP, and JSJ planned the study fieldwork campaign and conceptualization. AL designed the methodology, conducted the investigation, analysed the data, and wrote and prepared the original draft (writing and visualization). PF supervised μ -XRF data acquisition and supported data analysis. NKS supervised ^{210}Pb dating data acquisition and supported data analysis and age-depth modelling. AL, MP, JSJ, PF, and NKS reviewed and edited the draft. MP and JSJ supervised all steps of the research.

Competing interests

1585 The authors have the following competing interests: At least one of the (co-)authors is a member of the editorial board of Climate of the Past. Dr. Pierre Francus is a member of the editorial board of Climate of the Past.

Acknowledgements

We thank Sydney Moser and Tania O'Reilly for their assistance during the 2020 fieldwork campaign on the Magdalen Islands; Michelle Garneau and Leonie Perrier (GEOTOP – UQAM) for their valuable guidance in peat stratigraphy analysis; 1590 Arnaud De Coninck (INRS) for conducting the μ -XRF analysis and providing analytical support; and Steve Pratte (Zhejiang University) and Pascale Biron (Concordia University) for their insightful reviews of the master's thesis chapter upon which this paper is based. A special thank you to Luna for her quiet but constant support throughout the writing process.

Deleted:

→ The Magdalen Islands, located near the centre of the Gulf of St. Lawrence, have been impacted by post-tropical cyclones and other storm events over the last century, and our results provide important new data on paleo-storm history from one of the most northern sites studied for these events in the western North Atlantic, and one which is highly sensitive to ongoing climate warming.

Our 4000-year aeolian-based storm chronology from the Magdalen Islands demonstrates the viability of peatlands for such studies and offers critical insights into storm dynamics at the northern limit of the North Atlantic hurricane track. Furthermore, the comparability of our findings with existing North American paleo-storm records — primarily derived from marine and coastal overwash deposits — demonstrates the complementary value of aeolian and peatland records. These archives can enrich the paleo-storm literature in North America and provide a broader perspective on past storm activity.

→ One challenging task that emerged from our results is a proper calibration of peat-based proxies to the historical storm records. While our data show some correspondence with IBTRACS records over the past 150 years, identifying specific storms remains difficult. Future studies should prioritize the development of high-resolution age-models from undisturbed surface peat cores to refine calibration methods and better understand contemporary relationships between storm and sediment deposition in peatlands, supporting the interpretation of peat-based storm records.

→ Our results also highlight the specific challenges of working with peat records. Despite their proximity, our two study sites exhibited critical differences that were essential to address for an accurate storm record interpretation. For instance, differences in the amount and composition of sediments between the TLM and TAC storm records were linked to variations in bog size and distance from sediment sources, which influenced the strength of the storm signals. TAC displayed a stronger signal, sensitive to long-term environmental and climatic changes, whereas TLM's signal was more muted but exhibited greater consistency over time. These findings underscore the importance of considering such factors when selecting sites for future research.

→ Finally, our data reinforce the existence of an antiphase pattern in storms records between the Gulf of Mexico and the Caribbeans versus those north of the Bahamas. This finding extends the hypothesis that storm records north of the Bahamas may be in antiphase with basin-wide records from the North Atlantic. Future studies should investigate climatic mechanisms driving this antiphase pattern, as understanding these processes will prove important to estimate future storm impacts in eastern Canada.

Deleted: will also be


Formatted: Indent: First line: 1.27 cm

Deleted: upon publication

Formatted: Indent: First line: 1.27 cm

Formatted: Indent: First line: 1.27 cm

Figure A1. Seasonal wind roses displaying the 12 directions of incoming winds based on daily peak wind gusts data from 1984 to 2018 on the Magdalen Islands. The number of days (expressed as %) for any category of speed of peak wind gusts (in km h^{-1}) are indicated for each wind direction.



Deleted: 0
 Formatted: Font: Aptos, 12 pt
 Formatted: Font: Aptos
 Formatted: Font: Aptos, 12 pt
 Formatted: Font: Aptos
 Formatted: Font: Aptos, 12 pt
 Formatted: Font: Aptos
 Formatted: Font: Aptos
 Formatted: Font: Aptos, 12 pt
 Formatted: Font: Aptos
 Formatted: Font: Aptos, 12 pt
 Formatted: Font: Aptos
 Formatted: Font: Aptos

Appendix B: Supplementary information on core sampling and construction of the two composite cores

Deleted: C

Formatted: Heading 1, Indent: First line: 0 cm

1645 B1. Compression correction factors

The surface monoliths experienced compaction during transport and storage: the TLM monolith compressed from 50 cm to 28 cm, and the TAC monolith from 50 cm to 31 cm. To correct for this, a compression correction factor was calculated by dividing the original length (50 cm) by the compressed length at the time of analysis, resulting in factors of 1.79 (TLM) and 1.61 (TAC). Depths within each monolith were multiplied by the respective factor to restore their original, uncompressed position. These corrections applied only to the surface monoliths, not the deeper cores.

Formatted: Heading 2

1650 B2. Construction of composite cores

Table B1. Main cores drives from the TLM and TAC sites. The TLM composite core consists of the first 25 cm of TLM-monolith (0–25 cm), the first 75 cm of TLM-2 (25–100 cm) and the entire TLM-1 drive (100–700 cm). The TAC composite core consists of the first 25 cm of TAC-monolith (0–25 cm), the entire TAC-2 drive (25–275 cm), and a 50 cm segment from TAC-3 (275–325 cm). The remaining portion of TLM-2 and the entire TAC-1 core were archived for possible future analyses.

Core drive name	Actual depth (cm)	Portion used in composite core (cm)
<i>TLM</i>		
TLM-monolith	0–50	0–25
TLM-1	100–700	100–700
TLM-2	25–675	25–100
<i>TAC</i>		
TAC-monolith	0–50	0–50
TAC-1	0–315	None
TAC-2	25–275	50–275
TAC-3	275–325	275–325





Figure B1. Depth diagrams of core drives for TLM (a) and TAC (b). Note that the y-axis scales differ between the two panels. Solid orange rectangles represent core drives and segments used to construct the composite cores. Hatched orange rectangles indicate core drives and segments reserved for preliminary analyses, other research, or reference purposes. (a) The TLM composite core includes: the segment from the surface to 25 cm (TLM-monolith); 25 cm to 100 cm (TLM-2); and the entire TLM-1 core. (b) The TAC composite core includes: the full 50 cm length of TAC-monolith; 50 cm to 275 cm (TAC-2); and the full 50 cm length of TAC-3.



1665

Figure B2. Core drive alignment confirmed through cross-correlation analysis (CCA) of overlapping Fe μ -XRF measurements. Iron was used due to its abundance and consistent patterns across core drives. Black dots in the right-hand side CCA diagrams indicate significance ($p < 0.05$). (a) In the TLM composite core, cross-correlations between TLM-1 and TLM-2 (100–125 cm) and between TLM-2 and TLM-monolith (25–49 cm) showed strong positive correlation coefficient ($r = 0.8$ and $r = 0.6$, $p < 0.05$, at lag = 0), confirming alignment. (b) In the TAC composite core, TAC-2 and TAC-monolith's cross-correlation between 25–49 cm showed a strong positive correlation coefficient at lag = 0 ($r = 0.9$, $p < 0.05$), confirming alignment.

1670

Appendix G. Core pictures

(a) TLM



1680 (b) TAC

TAC-2	TAC-2	TAC-2	TAC-2	TAC-2	TAC-3
25-75	75-125	125-175	175-225	225-275	275-325



Figure G1. Photographs of the core segments taken at the TLM (a) and TAC (b) sites after extraction from the peatlands with depth indicated in cm. Only sections included in the TLM and TAC composite core are shown, except for the surface monoliths. (a) The basal sand unit is visible in TLM-1 650–700 between 693–700 cm. (b) The basal sand unit is visible in TAC-3 275–325 cm between 317–325 cm.

Deleted: Appendix C: Chronostratigraphy

Table C1. Accelerator mass spectrometry (AMS) radiocarbon (^{14}C) results for TLM and TAC. Dated material includes bulk peat sediment (bulk) and wood fragments (wood). Uncalibrated ^{14}C dates are shown as years before 1950 (BP). The ^{14}C dates were processed using the Bayesian *rlplum* algorithm in R and calibrated with the IntCal20 curve. Calibrated dates are presented as both calibrated years before 1950 (cal. BP) and as Common Era/Before Common Era (CE/BCE).

Sample depth (cm)

... [18]

Deleted: D


Formatted: Caption

Deleted: D

Appendix D: Chronostratigraphy

Table D1. Total ^{210}Pb activity results for TLM and TAC measured at the GEOTOP Radiochronology Laboratory at the *Université du Québec à Montréal* (UQAM). The background ^{210}Pb level for TLM was taken from the background ^{210}Pb level at TAC (17.5 Bq kg^{-1})

Sample depth (cm)	Lab ID	Dry Density (g cm^{-3})	^{210}Pb (Bq kg^{-1})	Mean modelled age (uncertainty) (CE)
TLM				
3.6	11366	0.04	300.67 \pm 22.61	2018 (2017 – 2018)
7.1	11367	0.04	331.2 \pm 24.12	2016 (2015 – 2017)
10.7	11368	0.04	335.54 \pm 24.4	2014 (2013 – 2015)
14.3	11369	0.04	261.04 \pm 19.6	2012 (2011 – 2013)
17.9	11370	0.05	494.38 \pm 35.6	2009 (2008 – 2010)
21.4	11371	0.06	405.92 \pm 28.81	2004 (2002 – 2006)
25	11372	0.06	325.84 \pm 24.3	1999 (1997 – 2001)
28.6	11373	0.05	274.68 \pm 20.79	1995 (1992 – 1999)
32.1	11374	0.06	310.76 \pm 22.45	1989 (1986 – 1993)
35.7	11375	0.06	383.36 \pm 26.56	1980 (1976 – 1984)
39.3	11376	0.06	313.56 \pm 21.67	1968 (1962 – 1976)
42.8	11377	0.07	234.26 \pm 16.49	1956 (1948 – 1967)
46.4	11378	0.07	188.63 \pm 13.49	1939 (1925 – 1956)
50	11379	0.07	190.79 \pm 14.44	1907 (1867 – 1942)
TAC				
3.3	11342	0.09	468.19 \pm 32.56	2009 (2003 – 2014)
6.7	11343	0.08	387.3 \pm 26.51	2002 (1997 – 2007)
10	11344	0.06	268.28 \pm 19.39	1997 (1991 – 2002)
13.3	11345	0.06	269.6 \pm 19.4	1991 (1985 – 1997)
16.7	11346	0.08	219.05 \pm 15.61	1985 (1979 – 1991)
20	11347	0.13	200.85 \pm 13.88	1975 (1968 – 1982)
23.3	11348	0.07	171.74 \pm 12.1	1967 (1960 – 1973)
26.7	11349	0.08	151.57 \pm 11.62	1957 (1949 – 1965)
30	11350	0.09	124.78 \pm 8.56	1946 (1937 – 1956)
33.3	11351	0.11	91.24 \pm 6.33	1932 (1922 – 1942)
36.7	11392	0.15	72.63 \pm 4.2	1905 (1887 – 1922)
40	11352	0.14	41.44 \pm 3.03	1868 (1827 – 1904)
43.3	11393	0.12	21.5 \pm 2.5	1839 (1789 – 1884)
46.7	11394	0.14	18.22 \pm 1.98	1812 (1751 – 1862)
50	11353	0.09	17.44 \pm 1.4	1785 (1720 – 1840)



705 **Figure D1.** *rplum*-derived age-depth models for TLM (left) and TAC (right). Radiocarbon (^{14}C) dates are shown in grey
 710 along the mean age model (dotted red). The rate of supported ^{210}Pb (right axis) is indicated by blue boxes on the left
 side (the blue boxes are the measured ^{210}Pb values, while the blue shadings represent the modelled ^{210}Pb values). For
 each core, the upper left panel shows the Markov Chain Monte Carlo (MCMC) iterations; followed by the prior (green
 curve) and posterior (grey histogram) distributions for the accumulation rate (second panel), memory (third panel),
 ^{210}Pb influx in $\text{Bq m}^{-2} \text{ yr}^{-1}$ (fourth panel) and supported ^{210}Pb in Bq kg^{-1} (fifth panel). A homing-in on the first 50 cm of
 the age-depth models is shown in the bottom panels.

Appendix E: Correlation results

Table E1. Pearson correlation coefficients among μ -XRF variables (CLR-transformed) and both ASI (log[ASI]) and mineral content (log[minerals/100-minerals]), presented for the core stratigraphic zones and upper 50 cm for TLM (a–b) and TAC (c–e; next page). Micro-XRF variables were averaged at 1 cm intervals to match the mineral content and ASI resolution.

(a)

	Ti	K	Ca	Mn	Fe	S	Si	ASI	Min
Ti		0.77***	-0.40***	-0.89***	-0.38***	-0.78***	-0.42***	0.65***	0.75***
K	0.00		-0.56***	-0.91***	-0.34***	-0.65***	-0.46***	0.54***	0.74***
Ca	-0.33***	0.22*		0.47***	-0.35***	0.14	0.55***	-0.44***	-0.48***
Mn	-0.42***	0.34***	0.77***		0.35***	0.71***	0.43***	-0.66***	-0.83***
Fe	-0.22*	-0.62***	0.03	-0.19*		0.69***	-0.53***	0.15	-0.06
S	-0.06	-0.64***	-0.62***	-0.77***	0.43***		-0.13	-0.41***	-0.62***
Si	-0.42***	0.24**	0.16	0.52**	-0.51***	-0.49***		-0.60***	-0.52***
ASI	0.21	-0.05	0.04	-0.30*	0.37**	0.21	-0.62***		0.85***
Minerals	0.14	-0.28*	-0.02	-0.49***	0.45***	0.45***	-0.63***	0.65***	

TLM (Zone 2 452–565 cm)

TLM (Zone 1 565–692 cm)

(b)

	Ti	K	Ca	Mn	Fe	S	Si	ASI	Min
Ti		0.12*	-0.65***	-0.40***	-0.15**	-0.40***	-0.52***	0.38***	0.47***
K	0.17		-0.02	-0.18***	0.13**	-0.43***	-0.75***	0.56***	0.43***
Ca	-0.41**	0.70***		0.32***	-0.13**	0.05	0.28***	-0.32***	-0.50***
Mn	-0.58***	-0.75***	-0.16		0.07	-0.31***	0.17***	-0.13*	-0.33***
Fe	0.19	-0.60***	-0.70***	0.37**		-0.18***	-0.25***	0.18**	0.24***
S	-0.51***	-0.49***	-0.15	0.41**	-0.00		0.39***	-0.30***	-0.17*
Si	-0.38**	-0.86***	-0.56***	0.66***	0.32*	0.38**		-0.65***	-0.58***
ASI	0.55**	0.67***	0.21	-0.77***	-0.34	-0.34	-0.62**		0.70***
Minerals	0.80***	0.17	-0.42	-0.58**	0.11	-0.22	-0.24	0.65**	

TLM (Upper 0–50 cm)

TLM (Zone 3 0–450 cm)

*** p < 0.001; ** p < 0.01; * p < 0.05

(c)

	Ti	K	Ca	Mn	Fe	S	Si	ASI	Min
Ti		0.82***	-0.93***	-0.94***	-0.87***	-0.46**	0.64***	0.49***	0.66***
K	-0.42***		-0.85***	-0.83***	-0.86***	-0.64***	0.71***	0.59***	0.67***
Ca	-0.90***	0.44***		0.97***	0.91***	0.39*	-0.81***	-0.57***	-0.72***
Mn	-0.88***	0.50***	0.96***		0.89***	0.37*	-0.78***	-0.58***	-0.72***
Fe	-0.53***	0.09	0.52***	0.48***		0.42**	-0.79***	-0.52***	-0.61***
S	-0.12	-0.27**	-0.12	-0.18	-0.31***		-0.57***	-0.42**	-0.48**
Si	-0.49***	-0.02	0.37***	0.35***	0.32***	-0.31***		0.61***	0.69***
ASI	0.06	0.02	-0.05	-0.02	0.04	-0.04	-0.10		0.88***
Minerals	0.40***	-0.09	-0.48***	-0.44***	-0.10	0.04	-0.27**	0.57***	

TAC (Zone 2 159–274 cm)

TAC (Zone 1 274–316 cm)

D

	Ti	K	Ca	Mn	Fe	S	Si	ASI	Min
Ti		0.95***	-0.96***	-0.92**	-0.79***	-0.93***	0.95***	0.29	0.61**
K	0.87***		-0.92***	-0.91***	-0.70***	-0.98***	0.96***	0.42*	0.69***
Ca	-0.89***	-0.78***		0.94***	0.77***	0.87***	-0.93***	-0.26	-0.53**
Mn	-0.79***	-0.87***	0.77***		0.73***	0.84***	-0.93***	-0.38	-0.61**
Fe	0.19*	0.25**	-0.18*	-0.14		0.62**	-0.76***	-0.14	-0.47*
S	-0.61***	-0.74***	0.45***	0.47**	-0.67***		-0.95***	-0.43*	-0.70***
Si	0.59***	0.75***	-0.60***	-0.74***	0.20*	-0.69***		0.41*	0.67***
ASI	0.68***	0.75***	-0.65***	-0.65***	0.36***	-0.64***	0.60***		0.83***
Minerals	0.83***	0.78***	-0.76***	-0.63***	0.17	-0.58***	0.53***	0.78***	

TAC (Zone 4 0–135 cm)

TAC (Zone 3 135–159 cm)

E

	Ti	K	Ca	Mn	Fe	S	Si	ASI	Min
Ti		0.76***	-0.87***	-0.49***	0.46***	-0.75***	0.63***	0.74***	0.81***
K			-0.52***	-0.82***	0.37**	-0.81***	0.85***	0.88***	0.69***
Ca				0.30*	-0.41**	0.50***	-0.43**	-0.54***	-0.72***
Mn					-0.05	0.48***	-0.87***	-0.71***	-0.40**
Fe						-0.77***	0.27	0.47**	0.49***
S							-0.73***	-0.82***	-0.73***
Si								0.80***	0.58***
ASI									0.83***
Minerals									

TAC (upper 0–50cm)

Appendix F: Historical hurricanes

We used the *International Best Track Archive for Climate Stewardship* (IBTrACTS) dataset (Knapp et al., 2010; Gahtan et al., 2024) to determine the number of cyclones with a tropical origin passing within 100 km of the Magdalen Islands from 1851

1725 to 2019 CE. To assess the impact of these storms on the Magdalen Islands, we cross-referenced the IBTrACTS data with historical shipwrecks records and local newspaper archives. We searched the online press database of the *Bibliothèque et Archive National du Québec* (Bibliothèque Et Archive Nationale Du Québec, 2024) using French keywords such as *ouragan* and *Îles-de-la-Madeleine*, and notes were taken on storm-related damage in the Magdalen Islands. Additionally, we manually reviewed shipwreck records from the vicinity of the Magdalen Islands (Landry, 1989, 1994) for incidents attributed to storms identified in the IBTrACTS dataset. Storms mentioned in the newspaper articles or shipwreck records that were not present in the IBTrACTS dataset, but that had significant impacts on the Magdalen Islands, were added to the final historical storm dataset.

Formatted: Not Highlight

1730 1735 Table F1. List of hurricanes that passed within 100 km or closer to the Magdalen Islands, Québec, with recorded wind of the hurricane at that time and whether or not there were known impacts.

Formatted: Normal

#	Date	Hurricane	Wind speed (km/h)	Known impacts	Listed in
1	2019-09-07	Dorian	139	Yes (Avila et al., 2020)	IBTrACTS
2	2014-07-06	Arthur	83	No	IBTrACTS
3	2010-09-04	Earl	101	No	IBTrACTS
4	2001-10-15	Karen	74	No	IBTrACTS
5	1999-09-18	Floyd	65	No	IBTrACTS
6	1996-10-10	Josephine	83	No	IBTrACTS
7	1996-07-14	Bertha	93	No	IBTrACTS
8	1988-08-08	Alberto	65	No	IBTrACTS
9	1979-09-07	David	93	No	IBTrACTS
10	1975-07-28	Blanche	111	Yes (Le Droit, July 29 1975; Le Soleil, July 29 1975)	IBTrACTS
11	1962-07-04	Not named	65	No	IBTrACTS
12	1959-07-12	Cindy	65	No	IBTrACTS
13	1950-08-21	Able	74	No	IBTrACTS
14	1944-09-15	Not named	83	No	IBTrACTS
15	1940-09-17	Not named	111	No	IBTrACTS
16	1937-09-26	Not named	111	No	IBTrACTS
17	1936-09-25	Not named	102	No	IBTrACTS
18	1934-09-10	Not named	46	No	IBTrACTS
19	1934-06-21	Not named	74	No	IBTrACTS

20	1933-10-30	Not named	93	No	IBTrACTS
21	1933-09-18	Not named	111	No	IBTrACTS
22	1927-08-25	1927 August Gale [†]	157	Yes (Landry, 1989, 1994)	IBTrACTS
23	1924-08-27	Not named	111	No	IBTrACTS
24	1917-08-11	Not named	83	No	IBTrACTS
25	1900-09-13	The Galveston Hurricane [†]	120	Yes (Landry, 1989, 1994)	IBTrACTS
26	1899-11-02	Not named	74	No	IBTrACTS
27	1898-10-06	Not named	46	No	IBTrACTS
28	1896-10-15	Not named	65	No	IBTrACTS
29	1891-10-15	Not named	65	No	IBTrACTS
30	1889-10-08	Not named	74	No	IBTrACTS
31	1888-08-23	Not named	93	No	IBTrACTS
32	1879-11-21	Not named	120	Yes (Landry, 1989, 1994)	IBTrACTS
33	1879-08-20	Not named	111	No	IBTrACTS
34	1877-11-30	Not named	74	No	IBTrACTS
35	1875-10-16	Not named	74	No	IBTrACTS
36	1873-08-25	1873 Great August Gale ^{†‡}	160	Yes (Landry, 1989, 1994) Shipwreck record(Landry, 1989, 1994)	IBTrACTS
37	1871-10-13	Not named	111	No	IBTrACTS
38	1860-08-27	Not named	65	No	IBTrACTS
39	1851-10-14	Not named	56	No	IBTrACTS

[†] Non-official but common names found in the literature to describe these events. [‡] The 1873 Great August Gale passed approximately 500 km south of the Magdalen Islands, near Sable Island in the North Atlantic Ocean. Nevertheless, it still was one of the most destructive storms in the Magdalen Islands and is well-documented.

1740 References

Aitchison, J.: The statistical analysis of compositional data, *J. Roy. Stat. Soc. B. Met.*, 44, 139-160, <https://doi.org/10.1111/j.2517-6161.1982.tb01195.x>, 1982.

Allen, D., Allen, S., Le Roux, G., Simonneau, A., Galop, D., and Phoenix, V. R.: Temporal Archive of Atmospheric Microplastic Deposition Presented in Ombrotrophic Peat, *Environ. Sci. Technol. Lett.*, 8, 954-960, 1745 <https://doi.org/10.1021/acs.estlett.1c00697>, 2021.

Aquino-López, M. A., Blaauw, M., Christen, J. A., and Sanderson, N. K.: Bayesian Analysis of 210-Pb Dating, 23, 317-333, 2018.

Avila, L.: National hurricane center tropical cyclone report - Hurricane Juan, NOAA National Hurricane Center, 11pp., 2003.

Avila, L., Stewart, S., Berg, R., and Hagen, A.: National hurricane center tropical cyclone report - Hurricane Dorian, NOAA 1750 National Hurricane Center, 74pp., 2020.

Baker, A. J., Hodges, K. I., Schiemann, R. K., and Vidale, P. L.: Historical variability and lifecycles of North Atlantic midlatitude cyclones originating in the tropics, *J. Geophys. Res.-Atmos.*, 126, e2020JD033924, <https://doi.org/10.1029/2020JD033924>, 2021.

Barnett, R., Bernatchez, P., Garneau, M., and Juneau, M. N.: Reconstructing late Holocene relative sea-level changes at the 1755 Magdalen Islands (Gulf of St. Lawrence, Canada) using multi-proxy analyses, *J. Quaternary Sci.*, 32, 380-395, <https://doi.org/10.1002/jqs.2931>, 2017.

Barnett, R., Bernatchez, P., Garneau, M., Brain, M., Charman, D., Stephenson, D., Haley, S., and Sanderson, N.: Late Holocene sea-level changes in eastern Québec and potential drivers, *Quaternary Sci. Rev.*, 203, 151-169, <https://doi.org/10.1016/j.quascirev.2018.10.039>, 2019.

Bengtsson, L. and Enell, M.: Chemical analysis, in: *Handbook of Holocene Palaeoecology and Palaeohydrology*, edited by: Berglund BE, Wiley Interscience, John Wiley & Sons Ltd., Chichester, <https://doi.org/10.1002/jqs.3390010111>, 1986.

Bernatchez, P., Boucher-Brossard, G., and Sigouin-Cantin, M.: Contribution des archives à l'étude des événements météorologiques et géomorphologiques causant des dommages aux côtes du Québec maritime et analyse des tendances, des fréquences et des temps de retour des conditions météo-marines extrêmes, Chaire de recherche en géoscience côtière, 1765 Laboratoire de dynamique et de gestion intégrée des zones côtières, Université du Québec à Rimouski, 140pp., 2012a.

Bernatchez, P., Drejza, S., and Dugas, S.: Marges de sécurité en érosion côtière: évolution historique et future du littoral des îles de la Madeleine, Chaire de recherche en géoscience côtière, Laboratoire de dynamique et de gestion intégrée des zones côtières, Université du Québec à Rimouski, 51pp., 2012b.

Bertrand, S., Tjallingii, R., Kylander, M. E., Wilhelm, B., Roberts, S. J., Arnaud, F., Brown, E., and Bindler, R.: Inorganic 1770 geochemistry of lake sediments: A review of analytical techniques and guidelines for data interpretation, *Eart-Sci. Rev.*, 104639, <https://doi.org/10.1016/j.earscirev.2023.104639>, 2023.

Besonen, M. R., Bradley, R. S., Mudelsee, M., Abbott, M. B., and Francus, P.: A 1,000-year, annually-resolved record of hurricane activity from Boston, Massachusetts, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL033950>, 2008.

Bibliothèque et Archive Nationale du Québec: BAnQ archive numérique, <https://numerique.banq.qc.ca/>, last access: 2024.

Bishops, T.: Using Itrax Data in R, <https://tombishop1.github.io/itrxBook/>, last access: July 12, 2025.

Björck, S. and Clemmensen, L. B.: Aeolian sediment in raised bog deposits, Halland, SW Sweden: a new proxy record of Holocene winter storminess variation in southern Scandinavia?, *Holocene*, 14, 677-688, 1775 <https://doi.org/10.1191/095963604hl746rp>, 2004.

Blaauw, M., Christen, J., and Aquino-Lopez, M.: rplum: Bayesian Age-Depth Modelling of Cores Dated by Pb-210 (R package version 0.5.1), CRAN [code], <http://doi.org/10.32614/CRAN.package.rplum>, 2024.

Boldt, K. V., Lane, P., Woodruff, J. D., and Donnelly, J. P.: Calibrating a sedimentary record of overwash from Southeastern New England using modeled historic hurricane surges, *Mar. Geol.*, 275, 127-139, <https://doi.org/10.1016/j.margeo.2010.05.002>, 2010.

1785 Brandon, C. M., Woodruff, J. D., Lane, D. P., and Donnelly, J. P.: Tropical cyclone wind speed constraints from resultant storm surge deposition: A 2500 year reconstruction of hurricane activity from St. Marks, FL, *Geochem. Geophys. Geosy.*, 14, 2993-3008, <https://doi.org/10.1029/ggpe.20217>, 2013.

Braun, E., Meyer, B., Deocampo, D., and Kiage, L. M.: A 3000 yr paleostorm record from St. Catherines Island, Georgia, *Estuar., Coast., Shelf S.*, 196, 360-372, <https://doi.org/10.1016/j.ecss.2017.05.021>, 2017.

1790 Bregy, J. C., Wallace, D. J., Minzoni, R. T., and Cruz, V. J.: 2500-year paleotempestological record of intense storms for the northern Gulf of Mexico, United States, *Mar. Geol.*, 396, 26-42, <https://doi.org/10.1016/j.margeo.2017.09.009>, 2018.

Brisebois, D.: Lithostratigraphie des strates permo-carbonifères, de l'archipel des Iles de la Madeleine, Direction générale des énergies conventionnelles, DPV-796, 48pp., 1981.

Canadian Ice Service: Seasonal Summary, Eastern Canada, Winter 2021-2022, Environment and Climate Change Canada, 29pp., 2022.

1795 Chartrand, P. G., Sonnentag, O., Sanderson, N. K., and Garneau, M.: Recent peat and carbon accumulation on changing permafrost landforms along the Mackenzie River valley, Northwest Territories, Canada, *Environ. Res. Lett.*, 18, 095002, <https://doi.org/10.1088/1748-9326/ace9ed>, 2023.

Chiang, J. C. H. and Vimont, D. J.: Analogous Pacific and Atlantic Meridional Modes of Tropical Atmosphere–Ocean Variability, *J. Climate*, 17, 4143-4158, <https://doi.org/10.1175/JCLI4953.1>, 2004.

1800 Crann, C. A., Murseli, S., St-Jean, G., Zhao, X., Clark, I. D., and Kieser, W. E.: First status report on radiocarbon sample preparation techniques at the AE Lalonde AMS Laboratory (Ottawa, Canada), *Radiocarbon*, 59, 695-704, <https://doi.org/10.1017/RDC.2016.55>, 2017.

Croudace, I. W. and Rothwell, R. G.: Micro-XRF Studies of Sediment Cores: Applications of a non-destructive tool for the environmental sciences, *Developments in Paleoenvironmental Research*, 17, Springer, Dordrecht, Netherlands, 656 pp., <https://doi.org/10.1007/978-94-017-9849-5>, 2015.

1805 Cwanek, A., Aquino-Lopez, M. A., Kołaczek, P., Lamentowicz, M., Fialkiewicz-Koziel, B., Galka, M., Słowiński, M., Łuców, D., Marcisz, K., Obremska, M., Czerwiński, S., and Łokas, E.: Strengthening potential of recent peat dating, 282, 107594, <https://doi.org/10.1016/j.jenrad.2024.107594>, 2025.

De Vleeschouwer, F., Sikorski, J., and Fagel, N.: Development of lead-210 measurement in peat using polonium extraction. A procedural comparison, *Geochronometria*, 36, 1-8, <https://doi.org/10.2478/v10003-010-0013-5>, 2010.

1810 Dean, W. E.: Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition; comparison with other methods, *J. Sediment Res.*, 44, 242-248, <https://doi.org/10.1306/74D729D2-2B21-11D7-8648000102C1865D>, 1974.

Denommee, K., Bentley, S., and Droxler, A.: Climatic controls on hurricane patterns: a 1200-y near-annual record from Lighthouse Reef, Belize, *Sci. Rep.*, 4, 3876, <https://doi.org/10.1038/srep03876>, 2014.

1815 Derksen, C., Burgess, D., Duguay, C., Howell, S., Mudryk, L., Smith, S., Thackeray, C., and Kirchmeier-Young, M.: Changes in snow, ice, and permafrost across Canada, in: Canada's changing climate report, edited by: Bush, E., and Lemmen, D. S., Government of Canada, 194-260, <https://doi.org/10.4095/308279>, 2019.

Dickie, J. and Wach, G.: A major midlatitude hurricane in the Little Ice Age, *Clim. Past.*, 20, 1141-1160, <https://doi.org/10.5194/cp-20-1141-2024>, 2024.

1820

Donnelly, J. P., Hawkes, A. D., Lane, P., MacDonald, D., Shuman, B. N., Toomey, M. R., van Hengstum, P. J., and Woodruff, J. D.: Climate forcing of unprecedented intense-hurricane activity in the last 2000 years, *Earth's Future*, 3, 49-65, <https://doi.org/10.1002/2014EF000274>, 2015.

1825 Dredge, L. A., Mott, R. J., and Grant, D. R.: Quaternary stratigraphy, paleoecology, and glacial geology, Îles de la Madeleine, Quebec, *Can. J. Earth Sci.*, 29, 1981-1996, <https://doi.org/10.1139/e92-154>, 1992.

Environment and Climate Change Canada: Canadian Climate Normals 1981-2010 Station Data - ILES DE LA MADELEINE A, https://climate.weather.gc.ca/climate_normals/results_1981_2010_e.html?stnID=5748&autofwd=1, last access: 30 July 2024.

1830 Forbes, D. L., Parkes, G. S., Manson, G. K., and Ketch, L. A.: Storms and shoreline retreat in the southern Gulf of St. Lawrence, *Mar. Geol.*, 210, 169-204, <https://doi.org/10.1016/j.margeo.2004.05.009>, 2004.

Fortin, J.-C. and Larocque, P.: *Histoire des Iles-de-la-Madeleine*, Institut québécois de recherche sur la culture, Les Éditions de l'IQRC et Les Presses de l'Université Laval, Sainte-Foy, 400 pp., <https://doi.org/10.7202/014675ar>, 2003.

1835 Gahtan, J., K. R. Knapp, C. J. Schreck, H. J. Diamond, J. P. Kossin, and Kruk, M. C.: International Best Track Archive for Climate Stewardship (IBTrACS) Project (Version 4r01) [North Atlantic Ocean], NOAA National Centers for Environmental Information [dataset], <https://doi.org/10.25921/82ty-9e16>, 2024.

Garneau, M.: Collection de référence de graines et autres macrofossiles végétaux de taxons provenant du Québec méridional et boréal et de l'arctique canadien, *Geol. Surv. of Can.*, Ottawa, Ont., Dossier publique 3048, 38 pp., 1995.

1840 Goslin, J., Galka, M., Sander, L., Fruergaard, M., Mokenbusch, J., Thibault, N., and Clemmensen, L. B.: Decadal variability of north-eastern Atlantic storminess at the mid-Holocene: New inferences from a record of wind-blown sand, western Denmark, *Global Planet. Change*, 180, 16-32, <https://doi.org/10.1016/j.gloplacha.2019.05.010>, 2019.

Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results, *J. Paleolimnol.*, 25, 101-110, <https://doi.org/10.1023/A:1008119611481>, 2001.

Hétu, B., Rémillard, A. M., Bernatchez, P., and St-Onge, G.: Landforms and Landscapes of the Magdalen Islands: The Role of Geology and Climate, in: *Landscapes and Landforms of Eastern Canada. World Geomorphological Landscapes*, edited by: Slaymaker, O., and Catto, N., Springer, Cham, 431-451, https://doi.org/10.1007/978-3-030-35137-3_19, 2020.

Hewett, D.: *Menyanthes trifoliata* L, *J. Ecol.*, 52, 723-735, <https://doi.org/10.2307/2257858>, 1964.

Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., Menne, M. J., Smith, T. M., Vose, R. S., and Zhang, H.-M.: Extended Reconstructed Sea Surface Temperature, Version 5 (ERSSTv5): Upgrades, Validations, and Intercomparisons, *J. Climate*, 30, 8179-8205, <https://doi.org/10.1175/JCLI-D-16-0836.1>, 2017.

Husson, F., Josse, J., Le, S., and Mazet, J.: FactoMineR: Multivariate Exploratory Data Analysis and Data Mining (R package version 2.11), CRAN [code], <http://doi.org/10.32614/CRAN.package.FactoMineR>, 2024.

Insurance Bureau of Canada: Hurricane Fiona causes \$660 million in insured damage, <http://www.ibc.ca/ns/resources/media-centre/media-releases/hurricane-fiona-causes-660-million-in-insured-damage>, last access: 31 October 2022.

1855 Irish, J. L., Resio, D. T., and Ratcliff, J. J.: The influence of storm size on hurricane surge, *J. Phys. Oceanograph.*, 38, 2003-2013, <https://doi.org/10.1175/2008JPO3727.1>, 2008.

Jansen, J. H. F., Van der Gaast, S. J., Koster, B., and Vaars, A. J.: CORTEX, a shipboard XRF-scanner for element analyses in split sediment cores, 151, 143-153, [https://doi.org/10.1016/S0025-3227\(98\)00074-7](https://doi.org/10.1016/S0025-3227(98)00074-7), 1998.

Jardine, D. E., Wang, X., and Fenech, A. L.: Highwater mark collection after post tropical storm Dorian and implications for Prince Edward Island, Canada, *Water*, 13, 3201, <https://doi.org/10.3390/w13223201>, 2021.

Juggins, S.: *rioja*: Analysis of Quaternary Science Data (R package version 1.0-7), CRAN [code], <http://doi.org/10.32614/CRAN.package.rioja>, 2024.

1865 Kern, O. A., Koutsodendris, A., Maechtle, B., Christianis, K., Schukraft, G., Scholz, C., Kotthoff, U., and Pross, J.: XRF core scanning yields reliable semiquantitative data on the elemental composition of highly organic-rich sediments: Evidence from the Fürmoos peat bog (Southern Germany), *Sci. Total Environ.*, 697, 134110, <https://doi.org/10.1016/j.scitotenv.2019.134110>, 2019.

1870 Klotzbach, P. J. and Gray, W. M.: Multidecadal variability in North Atlantic tropical cyclone activity, *J. Climate*, 21, 3929-3935, <https://doi.org/10.1175/2008JCLI2162.1>, 2008.

1875 Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., and Neumann, C. J.: The international best track archive for climate stewardship (IBTrACS) unifying tropical cyclone data, *B. Am. Meteorol. Soc.*, 91, 363-376, <https://doi.org/10.1175/2009BAMS2755.1>, 2010.

Kossin, J. P.: Hurricane intensification along United States coast suppressed during active hurricane periods, *Nature*, 541, 390-393, <https://doi.org/10.1038/nature20783>, 2017.

1880 Kossin, J. P. and Vimont, D. J.: A more general framework for understanding Atlantic hurricane variability and trends, *B. Am. Meteorol. Soc.*, 88, 1767-1782, <https://doi.org/10.1175/BAMS-88-11-1767>, 2007.

Kossin, J. P., Camargo, S. J., and Sitkowski, M.: Climate modulation of North Atlantic hurricane tracks, *J. Climate*, 23, 3057-3076, <https://doi.org/10.1175/2010JCLI3497.1>, 2010.

1885 Kuhlbrodt, T., Swaminathan, R., Ceppi, P., and Wilder, T.: A Glimpse into the Future: The 2023 Ocean Temperature and Sea Ice Extremes in the Context of Longer-Term Climate Change, *Bull. Am. Meteorol. Soc.*, 105, E474-E485, <https://doi.org/10.1175/BAMS-D-23-0209.1>, 2024.

Kuhry, P. and Turunen, J.: The postglacial development of boreal and subarctic peatlands, in: Boreal peatland ecosystems. Ecological Studies, vol. 188, edited by: Wieder, R. K., and Vitt, D. H., Springer, Berlin, Heidelberg, Germany, 25-46, https://doi.org/10.1007/978-3-540-31913-9_3, 2006.

Kylander, M. E., Ampel, L., Wohlfarth, B., and Veres, D.: High-resolution X-ray fluorescence core scanning analysis of Les Echets (France) sedimentary sequence: new insights from chemical proxies, *J. Quat. Sci.*, 26, 109-117, 2011.

1890 Kylander, M. E., Söderlindh, J., Schenk, F., Gyllencreutz, R., Rydberg, J., Bindler, R., Martínez Cortizas, A., and Skelton, A.: It's in your glass: a history of sea level and storminess from the Laphroaig bog, Islay (southwestern Scotland), *Boreas*, 49, 152-167, <https://doi.org/10.1111/bor.12409>, 2020.

Landry, F.: Derniere Course: Aventures maritimes dans le Golfe du Saint-Laurent, La Boussole, Havre-Aubert, Ile-de-la-Madeleine, 1989.

1895 Landry, F.: Piege de sable, La Boussole, Havre-Aubert, Iles-de-la-Madeleine, 1994.

Lane, P., Donnelly, J. P., Woodruff, J. D., and Hawkes, A. D.: A decadally-resolved paleohurricane record archived in the late Holocene sediments of a Florida sinkhole, *Mar. Geol.*, 287, 14-30, <https://doi.org/10.1016/j.margeo.2011.07.001>, 2011.

1900 Lapointe, F., Bradley, R. S., Francus, P., Balascio, N. L., Abbott, M. B., Stoner, J. S., St-Onge, G., De Coninck, A., and Labarre, T.: Annually resolved Atlantic sea surface temperature variability over the past 2,900 y, *P. Natl. Acad. Sci. USA*, 117, 27171-27178, <https://doi.org/10.1073/pnas.2014166117>, 2020.

Le Droit: Blanche a arraché de nombreux toits aux Iles-de-la-Madeleine, July 29 1975.

Le Soleil: Blanche isole les Iles-de-la-Madeleine du continent, 1, July 29 1975

Liu, D., Bertrand, S., and Weltje, G. J.: An empirical method to predict sediment grain size from inorganic geochemical measurements, *Geochem. Geophys. Geosy.*, 20, 3690-3704, <https://doi.org/10.1029/2018GC008154>, 2019.

1905 Liu, K.-b. and Fearn, M. L.: Lake-sediment record of late Holocene hurricane activities from coastal Alabama, *Geology*, 21, 793-796, [https://doi.org/10.1130/0091-7613\(1993\)021<0793:LSROLH>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<0793:LSROLH>2.3.CO;2), 1993.

1910 Liu, K.-b. and Fearn, M. L.: Reconstruction of prehistoric landfall frequencies of catastrophic hurricanes in northwestern Florida from lake sediment records, *Quaternary Res.*, 54, 238-245, <https://doi.org/10.1006/qres.2000.2166>, 2000.

Longman, J., Veres, D., and Wennrich, V.: Utilisation of XRF core scanning on peat and other highly organic sediments, *Quaternary Int.*, 514, 85-96, <https://doi.org/10.1016/j.quaint.2018.10.015>, 2019.

1915 Magnan, G. and Garneau, M.: Evaluating long-term regional climate variability in the maritime region of the St. Lawrence North Shore (eastern Canada) using a multi-site comparison of peat-based paleohydrological records, *J. Quaternary Sci.*, 29, 209-220, <https://doi.org/10.1002/jqs.2694>, 2014.

Mann, M. E., Woodruff, J. D., Donnelly, J. P., and Zhang, Z.: Atlantic hurricanes and climate over the past 1,500 years, *Nature*, 460, 880-883, <https://doi.org/10.1038/nature08219>, 2009a.

1920 Mann, M. E., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D., Ammann, C., Faluvegi, G., and Ni, F.: Global signatures and dynamical origins of the Little Ice Age and Medieval Climate Anomaly, *Science*, 326, 1256-1260, <http://doi.org/10.1126/science.1177303>, 2009b.

Ministère de l'Environnement et de la Lutte contre les changements climatiques: Habitat floristique de la Tourbière-de-L'Anse-à-la-Cabane, <https://www.environnement.gouv.qc.ca/biodiversite/habitats/tourbiere-anse-cabane/index.htm>, last access: 27 May 2021.

1925 Ministère de l'Environnement et de la Lutte contre les changements climatiques: Habitat floristique de la Tourbière-du-Lac-Maucôque, <https://www.environnement.gouv.qc.ca/biodiversite/habitats/tourbiere-lac-maucouque/index.htm>, last access: 27 May 2021

1930 Monica, S. B., Wallace, D. J., Wallace, E. J., Du, X., Dee, S. G., and Anderson, J. B.: 4500-year paleohurricane record from the Western Gulf of Mexico, *Coastal Central TX, USA, Mar. Geol.*, 473, 107303, <https://doi.org/10.1016/j.margeo.2024.107303>, 2024.

Morin, I.: Geomorphologie et evolution du système dunaire des îles de la Madeleine, Québec, Département de géographie, Université Laval, 2001.

1935 Oliva, F., Peros, M., and Viau, A.: A review of the spatial distribution of and analytical techniques used in paleotempestological studies in the western North Atlantic Basin, *Prog. Phys. Geog.*, 41, 171-190, <https://doi.org/10.1177/0309133316683899>, 2017.

Oliva, F., Peros, M., Viau, A., Reinhardt, E., Nixon, F., and Morin, A.: A multi-proxy reconstruction of tropical cyclone variability during the past 800 years from Robinson Lake, Nova Scotia, Canada, *Mar. Geol.*, 406, 84-97, <https://doi.org/10.1016/j.margeo.2018.09.012>, 2018.

1940 Orme, L. C., Reinhardt, L., Jones, R. T., Charman, D. J., Barkwith, A., and Ellis, M. A.: Aeolian sediment reconstructions from the Scottish Outer Hebrides: Late Holocene storminess and the role of the North Atlantic Oscillation, *Quat. Sci. Rev.*, 132, 15-25, <https://doi.org/10.1016/j.quascirev.2015.10.045>, 2016a.

Orme, L. C., Reinhardt, L., Jones, R. T., Charman, D. J., Croudace, I., Dawson, A., Ellis, M., and Barkwith, A.: Investigating the maximum resolution of μ XRF core scanners: A 1800 year storminess reconstruction from the Outer Hebrides, Scotland, UK, *Holocene*, 26, 235-247, <https://doi.org/10.1177/0959683615596819>, 2016b.

1945 Owens, E. and McCann, S.: The coastal geomorphology of the Magdalen Islands, Québec, 80-10, 1980.

Pasch, R. J., Penny, A. B., and Berg, R.: National hurricane center tropical cyclone report - Hurricane Fiona, NOAA National Hurricane Center, 2023.

Patterson, R. T., Nasser, N. A., Reinhardt, E. G., Patterson, C. W., Gregory, B. R., Mazzella, V., Roe, H. M., and Galloway, J. M.: End-Member Mixing Analysis as a Tool for the Detection of Major Storms in Lake Sediment Records, *Paleoceanogr. Paleoclimatol.*, 37, e2022PA004510, <https://doi.org/10.1029/2022PA004510>, 2022.

Peros, M., Chan, K., Magnan, G., Ponsford, L., Carroll, J., and McCloskey, T.: A 9600-year record of water table depth, vegetation and fire inferred from a raised peat bog, Prince Edward Island, Canadian Maritimes, *J. Quaternary Sci.*, 31, 512-525, <http://doi.org/10.1002/jqs.2875>, 2016.

Perrier, L., Garneau, M., Pratte, S., and Sanderson, N. K.: Climate-driven Holocene ecohydrological and carbon dynamics from maritime peatlands of the Gulf of St. Lawrence, eastern Canada, *Holocene*, 09596836221095978, <https://doi.org/10.1177/09596836221095978>, 2022.

Pleskot, K., Cwynar, L. C., Kowaleczyk, C., Kokociński, M., and Szczuciński, W.: Refining the history of extreme coastal events in southern Newfoundland, NW Atlantic, with lake sediment archives, *Quaternary Sci. Rev.*, 322, 108401, <https://doi.org/10.1016/j.quascirev.2023.108401>, 2023.

Prest, V., Terasme, J., Matthews, J., JV, and Lichti-Fedorovich, S.: Late-Quaternary History of Magdalen Islands, Quebec, *Atlantic Geol.*, 12, 39-60, <https://doi.org/10.4138/1835>, 1976.

Primeau, G. and Garneau, M.: Carbon accumulation in peatlands along a boreal to subarctic transect in eastern Canada, *Holocene*, 31, 858-869, <https://doi.org/10.1177/0959683620988031>, 2021.

Public Safety Canada: Canadian Disaster Database, https://bdc.securitepublique.gc.ca/prnt-eng.aspx?cultureCode=en_Ca&eventTypes=%27HU%27&normalizedCostYear=1, last access: 29 July 2024.

Pye, K. and Tsar, H.: Aeolian Sand and Sand Dunes, Springer Berlin, Heidelberg, 458 pp., <https://doi.org/10.1007/978-3-540-85910-9>, 2008.

Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP), *Radiocarbon*, 62, 725-757, <https://doi.org/10.1017/RDC.2020.41>, 2020.

Rémillard, A. M., Hétu, B., Bernatchez, P., and Bertran, P.: The Drift des Demoiselles on the Magdalen Islands (Québec, Canada): sedimentological and micromorphological evidence of a Late Wisconsinan glacial diamict, *Can. J. Earth Sci.*, 50, 545-563, <https://doi.org/10.1139/cjes-2011-0115>, 2013.

Rémillard, A. M., Buylaert, J.-P., Murray, A., St-Onge, G., Bernatchez, P., and Hétu, B.: Quartz OSL dating of late Holocene beach ridges from the Magdalen Islands (Québec, Canada), *Quat. Geochronol.*, 30, 264-269, <https://doi.org/10.1016/j.quageo.2015.03.013>, 2015.

Rémillard, A. M., St-Onge, G., Bernatchez, P., Hétu, B., Buylaert, J.-P., Murray, A. S., and Lajeunesse, P.: Relative sea-level changes and glacio-isostatic adjustment on the Magdalen Islands archipelago (Atlantic Canada) from MIS 5 to the late Holocene, *Quaternary Sci. Rev.*, 171, 216-233, <https://doi.org/10.1016/j.quascirev.2017.07.015>, 2017.

Rémillard, A. M., St-Onge, G., Bernatchez, P., Hétu, B., Buylaert, J. P., Murray, A. S., and Vigneault, B.: Chronology and stratigraphy of the Magdalen Islands archipelago from the last glaciation to the early Holocene: new insights into the glacial and sea-level history of eastern Canada, *Boreas*, 45, 604-628, <https://doi.org/10.1111/bor.12179>, 2016.

Rodysill, J. R., Donnelly, J. P., Sullivan, R., Lane, P. D., Toomey, M., Woodruff, J. D., Hawkes, A. D., MacDonald, D., d'Entremont, N., and McKeon, K.: Historically unprecedented Northern Gulf of Mexico hurricane activity from 650 to 1250 CE, *Sci. Rep.*, 10, 1-17, <https://doi.org/10.1038/s41598-020-75874-0>, 2020.

1990 Shotyk, W.: Review of the inorganic geochemistry of peats and peatland waters, *Earth-Sci. Rev.*, 25, 95-176, [https://doi.org/10.1016/0012-8252\(88\)90067-0](https://doi.org/10.1016/0012-8252(88)90067-0), 1988.

Sjöström, J. K., Cortizas, A. M., Hansson, S. V., Sánchez, N. S., Bindler, R., Rydberg, J., Mört, C.-M., Ryberg, E. E., and Kylander, M. E.: Paleodust deposition and peat accumulation rates—Bog size matters, *Chem. Geol.*, 554, 119795, <https://doi.org/10.1016/j.chemgeo.2020.119795>, 2020.

1995 Sjöström, J. K., Gyllencreutz, R., Cortizas, A. M., Nylund, A., Piilo, S. R., Schenk, F., McKeown, M., Ryberg, E. E., and Kylander, M. E.: Holocene storminess dynamics in northwestern Ireland: Shifts in storm duration and frequency between the mid- and late Holocene, *Quaternary Sci. Rev.*, 337, 108803, <https://doi.org/10.1016/j.quascirev.2024.108803>, 2024.

Steinmann, P. and Shotyk, W.: Geochemistry, mineralogy, and geochemical mass balance on major elements in two peat bog profiles (Jura Mountains, Switzerland), 138, 25-53, 10.1016/S0009-2541(96)00171-4, 1997.

2000 ten Brink, U. S., Chaytor, J., Geist, E. L., Brothers, D. S., and Andrews, B. D.: Assessment of tsunami hazard to the US Atlantic margin, *Mar. Geol.*, 353, 31-54, <https://doi.org/10.1016/j.margeo.2014.02.011>, 2014.

Ting, M., Kossin, J. P., Camargo, S. J., and Li, C.: Past and future hurricane intensity change along the US east coast, *Sci. Rep.*, 9, 1-8, <https://doi.org/10.1038/s41598-019-44252-w>, 2019.

2005 Troels-Smith, J.: Karakterisering af løse jordater (Characterization of unconsolidated sediments), *Danmarks Geologiske Undersøgelse IV. Række*, 3, 1-73, <https://doi.org/10.34194/raekke4.v3.6989>, 1955.

Tukey, J. W.: Exploratory data analysis, Springer1977.

Vaasma, T.: Grain-size analysis of lacustrine sediments: a comparison of pre-treatment methods, *Estonian J. Ecol.*, 57, <https://doi.org/10.3176/eco.2008.4.01>, 2008.

2010 Vaasma, T., Vandel, E., Sugita, S., Tönnisson, H., Suursaar, Ü., Kont, A., and Vilumaa, K.: Storminess reconstruction in the northeastern Baltic Sea region over the past 7600 years based on aeolian sand influx into coastal bogs, *Holocene*, 35, 61-74, <https://doi.org/10.1177/0959636241285783>, 2025.

Vacchi, M., Engelhart, S. E., Nikitina, D., Ashe, E. L., Peltier, W. R., Roy, K., Kopp, R. E., and Horton, B. P.: Postglacial relative sea-level histories along the eastern Canadian coastline, *Quaternary Sci. Rev.*, 201, 124-146, <https://doi.org/10.1016/j.quascirev.2018.09.043>, 2018.

2015 van de Plassche, O., Erkens, G., van Vliet, F., Brandsma, J., van der Borg, K., and de Jong, A. F.: Salt-marsh erosion associated with hurricane landfall in southern New England in the fifteenth and seventeenth centuries, *Geology*, 34, 829-832, <https://doi.org/10.1130/G22598.1>, 2006.

Van den Boogaart, K. G. and Tolosana-Delgado, R.: “Compositions”: a unified R package to analyze compositional data, *Comput. Geosci.*, 34, 320-338, <https://doi.org/10.1016/j.cageo.2006.11.017>, 2008.

2020 Van den Boogaart, K. G. and Tolosana-Delgado, R.: Analyzing compositional data with R, in: *Use R!*, edited by: Gentleman, R., Hornik, K., and Parmigiani, G. G., Springer, 258, <https://doi.org/10.1007/978-3-642-36809-7>, 2013.

Van den Boogaart, K. G., Tolosana-Delgado, R., and Bren, M.: compositions: Compositional Data Analysis (R package version 2.0-8), CRAN [code], 10.32614/CRAN.package.compositions, 2024.

2025 van Geel, B., Buurman, J., and Waterbolk, H. T.: Archaeological and palaeoecological indications of an abrupt climate change in The Netherlands, and evidence for climatological teleconnections around 2650 BP, *J. Quaternary Sci.*, 11, 451-460, [https://doi.org/10.1002/\(SICI\)1099-1417\(199611/12\)11:6<451::AID-JQS275>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1099-1417(199611/12)11:6<451::AID-JQS275>3.0.CO;2-9), 1996.

van Hengstum, P. J., Donnelly, J. P., Toomey, M. R., Albury, N. A., Lane, P., and Kakuk, B.: Heightened hurricane activity on the Little Bahama Bank from 1350 to 1650 AD, *Cont. Shelf Res.*, 86, 103-115, <https://doi.org/10.1016/j.csr.2013.04.032>, 2014.

2030 Vandel, E., Vaasma, T., Sugita, S., Tönnisson, H., Jaagus, J., Vilumaa, K., Anderson, A., and Kont, A.: Reconstruction of past storminess: Evaluation of an indicator approach using aeolian mineral grains buried in peat deposits, *Estonia, Quaternary Sci. Rev.*, 218, 215-227, <https://doi.org/10.1016/j.quascirev.2019.06.026>, 2019.

Vimont, D. J. and Kossin, J. P.: The Atlantic meridional mode and hurricane activity, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2007GL029683>, 2007.

2035 Wallace, E., Donnelly, J. P., van Hengstum, P. J., Wiman, C., Sullivan, R., Winkler, T., d'Entremont, N., Toomey, M., and Albury, N.: Intense hurricane activity over the past 1500 years at South Andros Island, The Bahamas, *Paleoceanogr. Paleoclimatol.*, 34, 1761-1783, <https://doi.org/10.1029/2019PA003665>, 2019.

Wallace, E., Donnelly, J., van Hengstum, P., Winkler, T., Dizon, C., LaBella, A., Lopez, I., d'Entremont, N., Sullivan, R., and Woodruff, J.: Regional shifts in paleohurricane activity over the last 1500 years derived from blue hole sediments offshore of 2040 Middle Caicos Island, *Quaternary Sci. Rev.*, 268, 107126, <https://doi.org/10.1016/j.quascirev.2021.107126>, 2021a.

Wallace, E., Donnelly, J., van Hengstum, P., Winkler, T., McKeon, K., MacDonald, D., d'Entremont, N., Sullivan, R., Woodruff, J., and Hawkes, A.: 1,050 years of hurricane strikes on Long Island in The Bahamas, *Paleoceanogr. Paleoclimatol.*, 36, e2020PA004156, <https://doi.org/10.1029/2020PA004156>, 2021b.

2045 Wang, J., Yang, B., Ljungqvist, F. C., Luterbacher, J., Osborn, T. J., Briffa, K. R., and Zorita, E.: Internal and external forcing of multidecadal Atlantic climate variability over the past 1,200 years, *Nat. Geosci.*, 10, 512-517, <https://doi.org/10.1038/ngeo2962>, 2017.

Wang, Y., Jahan, S., Burnett, W. C., Wu, Z., Elsner, J. B., Means, G. H., Liu, J., and Jiang, S.: Late Holocene tropical cyclones linked to climatic and solar variability, *Quaternary Sci. Rev.*, 334, 108710, <https://doi.org/10.1016/j.quascirev.2024.108710>, 2024.

2050 Weiss, D., Shotyk, W., Rieley, J., Page, S., Gloor, M., Reese, S., and Martinez-Cortizas, A.: The geochemistry of major and selected trace elements in a forested peat bog, Kalimantan, SE Asia, and its implications for past atmospheric dust deposition, *Geochim. Cosmochim. Ac.*, 66, 2307-2323, [https://doi.org/10.1016/S0016-7037\(02\)00834-7](https://doi.org/10.1016/S0016-7037(02)00834-7), 2002.

Wentworth, C. K.: A scale of grade and class terms for clastic sediments, *J. Geol.*, 30, 377-392, <https://doi.org/10.1086/622910>, 1922.

2055 Winkler, T., van Hengstum, P., Donnelly, J., Wallace, E., Albury, N., D'Entremont, N., Hawkes, A., Maio, C., Roberts, J., and Sullivan, R.: More frequent Hurricane passage across the Bahamian Archipelago during the little ice age, *Paleoceanogr. Paleoclimatol.*, 38, e2023PA004623, <https://doi.org/10.1029/2023PA004623>, 2023.

Winkler, T. S., van Hengstum, P. J., Donnelly, J. P., Wallace, E. J., Sullivan, R. M., MacDonald, D., and Albury, N. A.: Revising evidence of hurricane strikes on Abaco Island (The Bahamas) over the last 700 years, *Sci. Rep.*, 10, 16556, <https://doi.org/10.1038/s41598-020-73132-x>, 2020.

2060 Winkler, T. S., van Hengstum, P. J., Donnelly, J. P., Wallace, E. J., D'Entremont, N., Hawkes, A. D., Maio, C. V., Sullivan, R. M., and Woodruff, J. D.: Oceanic passage of hurricanes across Cay Sal Bank in The Bahamas over the last 530 years, *Mar. Geol.*, 443, 106653, <https://doi.org/10.1016/j.margeo.2021.106653>, 2022.

Wu, X., de Vernal, A., Fréchette, B., Moros, M., and Perner, K.: The signal of climate changes over the last two millennia in 2065 the Gulf of St. Lawrence, eastern Canada, *Quaternary Res.*, 106, 28-43, <https://doi.org/10.1017/qua.2021.56>, 2022.

Yang, W., Wallace, E., Vecchi, G. A., Donnelly, J. P., Emile-Geay, J., Hakim, G. J., Horowitz, L. W., Sullivan, R. M., Tardif, R., and van Hengstum, P. J.: Last millennium hurricane activity linked to endogenous climate variability, *Nat. Commun.*, 15, 816, <https://doi.org/10.1038/s41467-024-45112>, 2024.

2070 Yang, Y., Maselli, V., Normandeau, A., Piper, D. J., Li, M. Z., Campbell, D. C., Gregory, T., and Gao, S.: Latitudinal response
of storm activity to abrupt climate change during the last 6,500 years, *Geophys. Res. Lett.*, 47, e2020GL089859,
<https://doi.org/10.1029/2020GL089859>, 2020.