

1 **Evolution of nonstationary hydrological drought characteristics**
2 **in the UK under warming**

3 Srinidhi Jha^{1*}, Lucy J. Barker¹, Jamie Hannaford^{1,2}, Maliko Tanguy^{1,3}

4 ¹UK Centre for Ecology & Hydrology, Maclean Building, Crowmarsh Gifford, Oxfordshire,
5 OX108BB, United Kingdom

6 ²Irish Climate Analysis and Research UnitS (ICARUS), Maynooth University, Maynooth, Co.
7 Kildare, Ireland

8 ³European Centre for Medium Range Forecasts (ECMWF), Reading, United Kingdom

9 *Correspondence to: Srinidhi Jha (srijha@ceh.ac.uk)

10

11 **Abstract.** Although the United Kingdom (UK) is relatively wet, there is an increasing awareness of the
12 impacts of droughts, and an expectation that droughts will become worse in the future. This has
13 motivated studies that have developed projections of future UK drought characteristics. To date,
14 however, very few have addressed future changes in terms of probability of occurrence, and none
15 have quantified the evolution of rare nonstationary hydrological drought characteristics under
16 different warming conditions. This study investigates future changes in the hydrological drought
17 characteristics under varying ~~global~~ warming levels (1.5°C, 2°C, and 3°C), using nonstationary extreme
18 value analysis combined with a Bayesian uncertainty framework across 200 river catchments in the
19 UK. The analysis utilizes the enhanced future Flows and Groundwater (eFLaG) dataset, which is based
20 on the most recent UKCP18 climate projections, and incorporates outputs from four hydrological
21 models (G2G, PDM, GR4J, and GR6J). The findings indicate that rising temperatures will significantly
22 influence future drought duration, severity, and intensity across a majority of catchments, with rare
23 droughts (return period of 100-500 years) projected to be more severe in all seasons, particularly in
24 the southern UK. Further, relatively frequent summer droughts (return periods of 10 years) are
25 expected to become shorter but more severe and intense, particularly at higher warming. We observe
26 notable differences between stationary and nonstationary return periods across seasons, with the
27 change becoming more pronounced at longer return periods, particularly for drought severity.
28 Although the trends remain consistent across models under stationary and nonstationary conditions,
29 the results underscore the role of rarity, nonstationarity, and seasonal controls on the future evolution
30 of hydrological droughts in the region. Furthermore this framework could be used to support similar
31 analyses in other environments where analogous datasets of transient hydroclimate projections are
32 available

33

34

35 **1. Introduction**

36 With ongoing global climate change the United Kingdom (UK) is experiencing a pronounced
37 warming trend, with the most recent decade (2015-2024) averaging 1.24 °C above the 1961-
38 1990 baseline(Climate Change Committee, 2021; Kendon et al., 2024). Many notable drought
39 events have been recorded in the UK during the periods of 1975-76, 1988-89, 1990-92, 1995-
40 97, 2004-06, 2010-12, and 2022 Barker et al., 2024; Murphy et al., 2020; Turner et al., 2021).
41 Projections indicate that by 2050, several regions could face frequent water shortages, driven
42 by extended spells of hot and dry weather, which are expected to significantly affect river
43 flows and soil moisture levels (Bevan, 2019). In addition to the adverse impacts of climate
44 change, the increasing demand will pose water management challenges in the future, which
45 is particularly crucial for the south-eastern part of the UK, which is expected to experience
46 more significant changes in the long-term climate (Bevan, 2022). However, droughts are not
47 only expected to become more frequent, but also more spatially coherent, especially during
48 the summer season, which could further complicate drought management strategies(Tanguy
49 et al., 2023b). River-flow projections in the UK are known to be sensitive to seasonal variations
50 in precipitation and potential evapotranspiration, owing to their influence on the seasonal
51 wetting and drying cycles of the land surface (Parry et al., 2024a). Chan et al., (2024) further
52 highlighted that the likelihood of experiencing a summer month drier than the historically
53 driest recorded month is expected to rise with future warming in certain regions of UK. And
54 yet, deficits in the winter half-year have been a key driver of historical droughts, especially in
55 southeast England where faltering winter replenishment of groundwater resources also
56 impacts river flows. Hence it has been argued that it is important to consider hydrological
57 droughts in all seasons, and the interactions between them. Although these and other studies
58 highlight the importance of seasonal controls on UK droughts, a comprehensive probabilistic
59 analysis of drought return levels across characteristics and warming levels is still needed.

60

61 The growing awareness of drought as a major and increasing hazard and its impacts has
62 prompted a significant acceleration of research on changing drought risk in the UK, and
63 parallel changes in water resource management practices. In particular, the financial
64 regulators (OFWAT) and environmental regulators (Environment Agency) of the water
65 industry set out a 'duty of resilience' stipulates that water utilities must plan to ensure

66 security of supply to very extreme events (OFWAT, 2015; Environment Agency, 2023) in
67 practice, 1:500-year droughts. Understanding and preparing against these extreme
68 hydrological events is of most societal importance for the UK due to their disproportionate
69 impacts on water resources, agriculture, ecosystems, and public health. For instance, the cost
70 of relying on emergency drought measures in the UK is projected at £40 billion, whereas
71 proactively building water resilience would cost £21 billion over the same period (National
72 Infrastructure Commission, 2018). Furthermore, the annual cost to maintain resilience to
73 severe droughts is estimated at £60–600 million. For extreme droughts, this rises to £80–800
74 million per year (Climate Change Committee, 2019).

75 Given the relative brevity of most hydrological records, the need to ensure resilience to very
76 rare extremes has prompted the widespread adoption of stochastic simulation methods to
77 generate long time series from which we can sample such rare events. However, several lim-
78 itations and complexities arise from using such methods when understanding extreme event
79 evolution under anthropogenic climate change (Counsell and Durant, 2023; Environment
80 Agency, 2025), chief of which is the need to apply post-hoc climate change adjustments to
81 stochastic simulations based on the present day. There is therefore merit in directly analysing
82 climate change projections to assess the changing return levels of events of a given rarity,
83 including those very extreme events of the most importance for water resources planning. In
84 this study, return levels have been defined as the values of a variable (here duration, severity,
85 and intensity) expected to be exceeded on average once every T years, where T is the return
86 period. However, the complicated nature of the drought hazard and its relatively infrequent
87 occurrence, and the diverse and uncertain spatiotemporal patterns of hydrological droughts
88 make severity and rarity assessments complicated (Brunner et al., 2021). Further, under-
89 standing future changes in hydrological drought, in particular, remains limited for the UK, as
90 the majority of studies have primarily focused on analysing changes in drought magnitude
91 between current and future periods, using threshold-based metrics rather than exploring the
92 evolving nonstationary dynamics of various drought characteristics in the future (Barker et
93 al., 2019; Chan et al., 2022; Kay et al., 2021). More recently, Parry et al., (2024b) utilised a
94 newly developed nationally consistent, multi-model ensemble of hydrological projections en-
95 hanced future Flows and Groundwater (eFLaG) [dataset](#) (Hannaford et al., 2022a) [to quantify](#)
96 [future UK hydrological droughts which consists of transient time series \(continuous daily data](#)

97 ~~from 1980 to 2080), to explore changes in drought characteristics. These transient analyses~~
98 ~~capture how river flows evolve over time, rather than only comparing baseline and future~~
99 ~~time slices. However, they do not account for the probabilistic assessment of droughts or~~
100 ~~changes in their likelihood under future warming. datasetto quantify future UK hydrological~~
101 ~~droughts. The study conducts the analysis for baseline, and future periods as well as transient~~
102 ~~changes in low flows characteristics, but did not consider droughts in a probabilistic sense~~
103 ~~and could not therefore shed light on changing likelihood of very rare/extreme events.~~ Also,
104 there has been a lack of research focusing on understanding the evolution of hydrological
105 droughts in the UK under different warming conditions (1.5°C, 2°C, 3°C, and so on), which is
106 very important from a risk planning point of view(Tanguy et al., 2023a). ~~Global warming~~
107 ~~Warming~~ -level assessments can be used to support timely adaptation of drought manage-
108 ment strategies, inform policy decisions aligned with global targets, and ensure resilience un-
109 der plausible future warming scenarios.

110 The analysis in most of the previously mentioned research for the UK is based on the analyses
111 of extreme events relying on the assumption of stationarity, which assumes that the
112 probability distribution parameters of a drought characteristic remain constant over time (Wu
113 et al., 2024). However, it is well-accepted that rising temperatures introduce nonstationarity
114 into hydrological systems, challenging the conventional approaches to drought analysis. This
115 nonstationarity might lead to inaccuracies in estimating the return levels of extreme events
116 for any design return period under evolving climatic conditions. Coles, (2001) highlighted that
117 assuming stationarity can lead to an underestimation of extreme event probabilities.
118 Therefore, incorporating nonstationarity, particularly due to rising temperatures, is crucial for
119 accurately modelling future drought characteristics (Salas and Obeysekera, 2014). One of the
120 important aspects of probabilistic modelling of extreme hydroclimatic events is the
121 uncertainty in estimated parameters (Leng et al., 2024; Onyutha, 2017). Traditional methods,
122 such as L-moments (Parvizi et al., 2022), method of moments (Lück and Wolf, 2016) and
123 maximum likelihood estimation (Jha et al., 2022), typically rely on point estimates of
124 parameters, without adequately addressing this issue. However, Bayesian methods have
125 found their utility for addressing these challenges in parameter estimation processes (Baykal
126 et al., 2024; Liu et al., 2024). This approach allows for obtaining the posterior distribution of
127 parameters by integrating over the existing parameter space. Additionally, the introduction

128 of Markov Chain Monte Carlo (MCMC) methodology facilitates the approximation of integrals
129 by using a Markov chain with the posterior distribution (Chandra et al., 2015). This paper uses
130 a nonstationary extreme value analysis (EVA) framework with Bayesian uncertainty
131 assessment to analyse the evolution of future hydrological drought characteristics in the UK
132 with specifically including rare droughts (return period ≥ 100 years). Leveraging the benefits
133 of the eFLaG river flow datasets, which comprise four hydrological models' (GR4J, GR6J, PDM,
134 and G2G) outputs, this study analyses transient, in this case daily continuous, century-long
135 projections data over 200 catchments in the UK~~this study analyses transient, century-long~~
136 projections at a daily resolution over 200 catchments in the UK. It examines the evolution of
137 future hydrological drought characteristics under three different warming levels ~~Global~~
138 Warming Levels (GWs): 1.5°C, 2°C, and 3°C, with a particular focus on extreme droughts. By
139 focusing on a range of warming scenarios, we aim to capture the full spectrum of possible
140 future hydrological drought conditions under different climatic conditions. In doing so, this
141 study provides critical insights for policymakers and water resource managers to better
142 understand and prepare for future hydrological drought risks and their uncertainties under
143 the influence of climate change. In summary, the objectives of this study are: (i) to investigate
144 the projected changes in key hydrological drought characteristics (duration, severity, and
145 intensity) across 200 UK catchments under three future warming scenarios. (ii) to apply and
146 compare results from nonstationary and stationary EVA using a Bayesian framework to
147 quantify the role of nonstationarity in governing future hydrological drought risks. (iii) to
148 understand the future evolution of hydrological drought characteristics in UK, specifically for
149 rare events with robust estimation of uncertainty.

150

151 **2. Data and methods**

152 **2.1. eFLaG data set: hydrological models and future river flow projections**

153 This paper utilizes the eFLaG dataset which are nationally consistent and spatially coherent
154 hydrological river flow projections for the UK based on UKCP18 - the latest climate projections
155 from the UK Climate Projections programme (Hannaford et al., 2022a; Lowe et al., 2018;
156 Murphy et al., 2018). The eFLaG dataset are hydrological projections derived from a range of
157 hydrological models (Grid-to-Grid, PDM, GR4J and GR6J) and groundwater recharge model
158 ZOODRM (zooming object-oriented distributed-recharge model). However, in this paper we
159 have only focussed on the river flow projections for our analysis and did not consider the

160 ~~groundwater data. We considered the~~ The eFLaG dataset encompasses hydrological model
161 simulations of river flow ('simobs' and 'simrcm') for over 200 catchments in the UK. In this
162 context, 'simobs' refers to observation-driven simulations (1989-2018), while 'simrcm'
163 denotes outputs generated from hydrological modelling using 12km UKCP18 RCM (Regional
164 Climate Models) projections (up to 2080). The 'simrcm' projections consist of a 12-member
165 ensemble generated using perturbed-parameter runs of the Hadley Centre global climate
166 model (GCM, HadGEM3-GC3.05) and regional climate model (RCM, HadREM3-
167 GA705)(Murphy et al., 2018). Each ensemble member represents a plausible variation in
168 model parameters to capture uncertainty in the climate response, while all members share
169 the same underlying model framework and follow the high-emissions scenario (RCP8.5). The
170 12-member RCM perturbed-parameter ensemble is therefore valuable for representing
171 parameter uncertainty; however, because all members are based on the same model
172 structure and emissions scenario, they do not capture the full range of climate or scenario
173 uncertainties. The 'simrcm' projections comprise a 12-member ensemble generated through
174 perturbed-parameter runs of Hadley Centre climate models (GCM, HadGEM3-GC3.05) and
175 RCM (HadREM3-GA705) should be noted that all 12 ensemble members originate from the
176 same model framework and are based on the high-emissions scenario (RCP8.5).
177 GR4J and GR6J, members of the 'airGR' family, are lumped catchment rainfall-runoff models
178 known for their simplicity and efficient calibration function (Kuana et al., 2024). The
179 Probability Distributed Model (PDM) offers configurable options for catchment rainfall-runoff
180 modelling, allowing for various permutations to be tested across catchments (Moore, 2007).
181 Grid-to-Grid (G2G) is a distributed hydrological model utilized for simulating natural river
182 flows across Great Britain at a 1km resolution, providing consistent national-scale flow
183 estimates (Bell et al., 2018). These models have been successfully applied in diverse
184 hydrological studies, and several publications detail their versatility and wide-ranging
185 applicability (Kuana et al., 2024; Ndiaye et al., 2024; Tanguy et al., 2023b). Detailed metadata
186 and site listings are stored and accessible through the Environmental Informatics Data Centre,
187 which can be referred for more information(Hannaford et al., 2022b). ~~In this study, we have~~
188 ~~utilised all 200 catchments for our analysis.~~ For the nonstationary modelling of drought
189 characteristics for each catchment, we utilised the ~~recently developed~~ CHES-SCAPE
190 temperature datasets, which are bias-corrected 1km resolution gridded data also derived
191 from UKCP18 projections (Robinson et al., 2022a) as a covariate. The CHES-SCAPE

192 [temperature records are derived from UKCP18 projections that have been downscaled to 1](#)
193 [km resolution using methods that account for local topographic effects and pattern scaling](#)
194 [properties for different scenarios](#)(Robinson et al., 2022a), [however, the eFLaG dataset is](#)
195 [based directly on the original UKCP18 projections.](#)

196

197 **2.2. Nonstationary analysis of future drought characteristics**

198 The impact of adverse climate change effects has prompted scrutiny of the stationary
199 assumption regarding hydroclimatic variables, leading to heightened interest in the concept
200 of nonstationarity within the research community. The concept is also pertinent to planners
201 using projections of hydrological information and data in their decision-making. In this study,
202 the drought characteristics were fitted with the generalized extreme value (GEV) distribution
203 with a cumulative distribution function given by Eq. (1) (Coles, 2001):

$$204 \quad G(x; \mu, \sigma, \xi) = \begin{cases} \exp \left\{ - \left[1 + \left(\frac{(x-\mu)\xi}{\sigma} \right)^{-\frac{1}{\xi}} \right]^{-\xi} \right\}, & \sigma > 0, \quad 1 + \left(\frac{(x-\mu)\xi}{\sigma} \right)^{-\xi} > 0, \quad \xi \neq 0 \\ \exp \left\{ - \exp \left[- \frac{x-\mu}{\sigma} \right] \right\}, & \sigma > 0, \quad \xi = 0 \end{cases} \quad (1)$$

205 Here, μ , σ and ξ are the location, scale, and shape parameters of the distribution. Daily
206 temperature anomaly (ΔT) from the CHES-SCAPE data (Robinson et al., 2022a) was selected
207 as the covariate to quantify the temperature-dependent signals for future river flow. [Here,](#)
208 [daily temperature anomaly for each period were calculated relative to the mean temperature](#)
209 [over the UK for the reference period \(1989-2018\). After identifying drought events, we](#)
210 [matched the timestamp of each drought characteristic with the corresponding temperature](#)
211 [time series and used the mean reference-period temperature to compute the anomalies,](#)
212 [which were then used as covariates. Please refer to Section 2.4 for further details on the](#)
213 [event-calculation methodology to understand how seasonality and continuation of events](#)
214 [have been considered.](#)

215 The incorporation of linear dependency in the location parameter is a common practice in
216 nonstationary modelling, and similar applications to the scale parameter have been
217 advocated by Yilmaz and Perera, (2014). However, Gilleland and Katz, (2016) argue against
218 introducing covariates solely to the scale parameter without corresponding variations in the
219 location parameter. Further, the estimation of the shape parameter under a time-varying
220 framework is challenging due to the uncertain tail behaviour of the distribution, especially in

221 limited data settings, and is therefore often kept constant (Ragulina and Reitan, 2017). In our
 222 study, only the location parameter for historical and future streamflow extremes was
 223 assumed to be a linear function of temperature. Hence, the parameter set takes the form of
 224 $\mu(t) = \mu_0 + \mu_1 c(\Delta T)$, $\sigma(t) = \sigma$ and $\xi(t) = \xi$. Parameter estimation was conducted utilizing
 225 the maximum likelihood function, chosen for its capability to incorporate nonstationarity into
 226 the distribution parameter (Strupczewski et al., 2001) as given by Eq. (2):

$$227 \quad L(\theta) = -n \log \sigma - (1 + \frac{1}{\xi}) \sum_{i=1}^n \log \left[1 + \xi \left(\frac{x_i - \mu}{\sigma} \right) \right] - \sum_{i=1}^n \left[1 + \xi \left(\frac{x_i - \mu}{\sigma} \right) \right]^{\left(\frac{1}{\xi} \right)}, 1 + \xi \left(\frac{x_i - \mu}{\sigma} \right) > 0 \quad (2)$$

228
 229 Here, $L(\theta)$ is the likelihood function of the parameter vector θ and n is the sample size. By
 230 minimizing the above function, the distributions of parameters for both stationary and
 231 nonstationary cases were formulated. The comparative statistical significance of stationary
 232 and nonstationary models was assessed by using the likelihood ratio test (L.R. test) (Posada
 233 and Buckley, 2004) which is derived using Eq. (3):

$$234 \quad 2[nllh_s - nllh_{(NS)}] > c_\alpha \quad (3)$$

235 Here, $nllh_s$ and $nllh_{(NS)}$ are the negative log-likelihood values of stationary and
 236 nonstationary models. Further, c_α represents the $(1 - \alpha)$ quantile of the Chi-square
 237 distribution. The difference between the stationary and nonstationary models is expected to
 238 conform to an approximate chi-squared distribution at a specific significance level α (5% in
 239 this case). The null hypothesis in this study assumes that drought characteristics extremes are
 240 stationary, meaning their statistical properties do not change over time or with temperature.
 241 Using the likelihood ratio test, this hypothesis is evaluated by comparing the fit of stationary
 242 and nonstationary GEV models. The null hypothesis is rejected when the p-value falls below
 243 0.05, indicating that including temperature as a covariate significantly improves the model.
 244 Such an approach is consistent with standard methods in extreme value analysis for
 245 hydrological data (Das and Umamahesh, 2017; Salas and Obeysekera, 2014). The percentage
 246 of catchments showing nonstationary characteristics for different combinations of seasons,
 247 metrics, models and warming levels are mentioned in Table S1 in the supplementary
 248 information. The null hypothesis of stationarity is rejected when the p-value exceeds 0.05.

250 2.3. Bayesian framework for parameter uncertainty

251 As discussed above, parameters for both stationary and nonstationary methods are derived
 252 using the maximum likelihood approach, which only provides point estimates without
 253 accounting for uncertainty. Bayesian analysis aims at updating parameter uncertainty
 254 through a prior distribution using Bayes' theorem (Sarhadi et al., 2016). This approach
 255 combines the prior distribution and the data's likelihood function to form the posterior
 256 distribution, incorporating additional information to enhance predictive modelling. The
 257 posterior distribution is obtained by multiplying the likelihood function by the prior
 258 distribution of the parameter (Eq. 4):

$$259 \quad p(\theta | y) \propto p(y|\theta) p(\theta) \quad (4)$$

260 Here, $p(\theta | y)$ denotes the posterior distribution of the parameter vector $\theta = (\mu, \sigma, \xi)$, $p(\theta)$
 261 represents the prior distribution, and $p(y|\theta)$ denotes the likelihood function corresponding
 262 to the GEV distribution evaluated at $y_{i:n}$ where n is the number of observations. We utilised
 263 a non-informative prior distribution for location parameter modelling. Given the complexity
 264 of solving Eq. (4) analytically, numerical methods like MCMC sampling are utilized to produce
 265 numerous realizations from the posterior distribution (Reis and Stedinger, 2005). Further, we
 266 can estimate desired return levels for a given probability of occurrence (p) by employing Eq.
 267 (5):

$$268 \quad Z_p(\hat{\mu}, \hat{\sigma}, \hat{\xi}) = \hat{\mu} - \frac{\hat{\sigma}}{\hat{\xi}} \left\{ 1 - [-\log(1-p)]^{-\hat{\xi}} \right\} \quad \text{for } \xi \neq 0 \quad (5)$$

$$269 \quad Z_p(\hat{\mu}, \hat{\sigma}) = \hat{\mu} - \hat{\sigma} \log[-\log(1-p)] \quad \text{for } \xi = 0 \quad \text{---}$$

270 The Metropolis-Hastings algorithm is used to sample the parameter vector using the specified
 271 prior and likelihood function. It is crucial to monitor the convergence of the MCMC chain to
 272 ensure it accurately represents the posterior distribution. In this study, Heidelberger and
 273 Welch's convergence diagnostic is used to determine the necessary length of each simulation
 274 (Sharma and Mujumdar, 2022).

275

276 **2.4. Analysis of future drought return levels**

277 The whole analysis is set up to calculate the percentage changes in the return level of the
 278 hydrological drought characteristics in the warming level period as compared to the reference
 279 period. The 30-year reference period was 1989-2018, i.e., the available historical period in
 280 the eFLaG dataset. Relative to this reference period, three warming level periods (also 30-
 281 year) were calculated based on the recently developed CHESS-SCAPE temperature data

282 projections for the UK (Robinson et al., 2022a). In alignment with the objectives and directives
283 of the Paris Agreement about limiting global warming, a +1.5°C and +2°C rise in temperature
284 was considered (Jha et al., 2023). Moreover, a warming level of +3°C was also considered,
285 corresponding to the projected warming expected to be attained by the year 2100 under
286 existing nationally determined mitigation goals (Seneviratne and Hauser, 2020). The starting
287 year of each warming level period is defined as the initial year of the 30-year interval wherein
288 the mean warming exceeds the respective warming level. We considered the last 30-year time
289 period, in case, the +3°C warming period exceeded the end of the century. For example, in
290 cases where the warming period is identified as 2080-2110, we instead use the 2070-2100
291 window to remain within the 21st-century bounds. -The warming levels in this analysis should
292 be interpreted as regional UK warming levels rather than global warming levels, since CHES-
293 SCAPE provides only UKCP18 climate projections over the UK. While the CHES-SCAPE
294 framework does use global mean air temperature from UKCP18 GCMs and uses time shifting
295 and pattern scaling, the downscaled dataset contains only UK specific surface variables.
296 However, these warming levels are broadly aligned with global warming levels as UKCP18
297 assumes seasonal UK climate anomalies scale linearly with global mean temperature, and it
298 is known that UK temperature changes generally track global land-surface warming (Kendon
299 et al., 2024).

300 To identify hydrological drought events, we used a variable threshold-based approach that
301 has been widely applied for drought identification (Sarailidis et al., 2019)(Sarailidis et al.,
302 2019). For each of the 12 ensemble members of each hydrological model, we first calculated
303 the daily mean flow values for every day of the reference period using the eFLaG dataset. We
304 then applied a 30-day rolling window centred on each day of the year. For example, for 15
305 January, the window includes flows from 15 days before to 15 days after. This smoothing
306 method helps capture natural variability in daily flows and prevents the resulting statistics
307 from being overly influenced by short-lived extreme events. Using these rolling-window
308 values, we derived 365 Q90 thresholds, one for each day of the year, representing the 90th
309 percentile exceedance flow for the reference period. These thresholds were then used as the
310 baseline against which projected flow levels at different warming levels were compared.
311 Specifically, we calculated the difference between projected flows and the corresponding
312 daily Q90 threshold to identify high-flow anomalies or deficits relevant for drought analysis.
313 The resulting drought characteristics for each warming level were subsequently pooled across

314 all 12 ensemble members, and this pooled dataset was used to fit GEV distributions to assess
315 changes in extremes under future climate conditions.

316 -We selected the 90th percentile (Q90) threshold to ensure that the analysis captures
317 instances characterised by extremely low historical flows. This choice allows us to focus on
318 severe low-flow anomalies that are hydrologically meaningful, rather than relatively normal
319 variations in streamflow. The Q90 threshold has also been widely used in previous
320 hydrological drought assessments, providing both consistency and comparability with earlier
321 studies (Hasan et al., 2020; Janicka-Kubiak, 2025; Prudhomme et al., 2014)(Hasan et al., 2020;
322 Janicka-Kubiak, 2025; Prudhomme et al., 2014). Furthermore, Q90 is sufficiently stringent to
323 minimise the influence of short-term fluctuations, ensuring that the identified drought events
324 represent genuine low-flow conditions rather than transient anomalies. An additional
325 motivation for adopting the Q90 threshold is our emphasis on addressing uncertainties
326 associated with estimating rare drought characteristics. Using a high-percentile threshold
327 such as Q90 demonstrates that the methodology is robust for detecting extremely low-
328 occurrence drought events, thereby supporting the reliability of our drought characterisation
329 approach. Further, we have demonstrated the drought characteristics distribution for one
330 model (G2G) and one warming level 3°C using both Q90 and Q80 thresholds in Figure S1a-c
331 in the supplementary information.-A catchment was considered to be in drought on any given
332 day when the flow dropped below the baseline Q90 threshold for that day. A pooling
333 procedure across drought events was also applied, where two distinct events separated by a
334 single day were combined into a single drought event, provided the magnitude above the
335 threshold did not exceed the accumulated deficit before this single day similar to the
336 methodology used by Van Loon and Van Lanen, (2012) and Parry et al., (2024a)Van Loon and
337 Van Lanen, (2012) and Parry et al., (2024a). To reduce uncertainty arising from very short,
338 potentially non-significant drought events caused by daily variability in the threshold, we
339 excluded events with a duration of less than 30 days. Given that we focus on Q90 to derive
340 these events, even after applying precautionary measures such as a 30-day moving window
341 and a 12-member ensemble pool to ensure smoother and larger sample sizes, extreme value
342 analysis remains challenging, particularly for rare, small drought events. We acknowledge
343 that this threshold effectively imposes a hard lower bound on drought duration and may also
344 exclude smaller events such as flash droughts. Nevertheless, we chose 30 days which has

345 widely been used in similar analyses by Anderson et al., (2025) and Brunner and Chartier-
346 Rescan, (2024)Anderson et al., (2025) and Brunner and Chartier-Rescan, (2024), as
347 compromise to balance robustness of event statistics with capturing meaningful hydrological
348 droughts. To avoid uncertainty arising due to non significant drought events, we excluded
349 those with a standard duration of less than 30 days.

350 Figure 1 schematically represents the derivation of drought characteristics using the variable
351 threshold method and a flow chart of the methodology used. We chose the variable threshold
352 method as a more suitable and increasingly popular approach compared to the constant
353 (fixed) threshold method for defining hydrological droughts (Anderson et al., 2025; Brunner
354 and Chartier-Rescan, 2024)(Anderson et al., 2025; Brunner and Chartier-Rescan, 2024). This
355 method allows for smooth intra-annual variability and identifies drought events when flows
356 fall below the historically expected level on a given day, which would be overlooked by a
357 constant threshold. This is important when we consider drought is relative phenomenon, and
358 especially as we are looking at hydrological deficits in all seasons as argued in the
359 introduction. A variable approach -allows the identification of multi-season and indeed multi-
360 year droughts, whereas in strongly seasonal regimes a fixed threshold typically only identifies
361 'absolute' droughts in the 'low flow' period (in the summer half-year in the case of the UK),
362 and these naturally terminate in the autumn/winter simply given the fact flows in these
363 seasons are always higher than summer - even if they are in fact low for the season in question
364 relative to historical norms, and potentially part of multi-annual ongoing droughts. Wider
365 discussion on the use of both fixed and variable approaches is provided elsewhere (see e.g.
366 (Stahl et al., 2020; Tallaksen and Van Lanen, 2023)(Stahl et al., 2020; Tallaksen and Van Lanen,
367 2023) and quantitative comparisons have been made to highlight the impacts of such
368 decisions e.g. for the US, (Hammond et al., 2022)(Hammond et al., 2022).

369 Having identified individual events, three event characteristics were computed for each
370 season (i.e. winter: December-February, spring: March-May, summer: June-August and
371 autumn: September-November) which are duration(number of days)- the number of days
372 over which a drought occurs, severity - the accumulated flow deficit across all days(cumecs),
373 and intensity (cumecs per day)- the ratio of drought severity and duration of a drought event.
374 It should be noted that event detection is performed on the full continuous time series in
375 reference period and warming level periods, not within seasons. Seasonal metrics are

376 calculated only after drought events and their onset are identified, so physical continuity is
 377 preserved, and duration or severity are not artificially capped by seasonal or yearly
 378 boundaries except in the last year of the period. We have calculated the drought
 379 characteristics based on the starting and end points of the event and assigned the season
 380 based on starting month.

381




Figure 1. a) Variable threshold methodology used to identify and characterise drought events, b) Methodological framework utilized in the analysis.

382

383 **3. Results and discussion**

384 **3.1. Nonstationary properties and Bayesian parameter estimates**

385 Once the drought characteristics for all four models across all four seasons were calculated,
 386 the nonstationarity was assessed using the likelihood ratio test. Figure 2 and [Table S1 in the](#)
 387 [supplementary information](#) represent the percentage of nonstationary catchments for each
 388 drought characteristic across three warming levels and seasons. It shows that the
 389 nonstationary properties of catchments depend on the combination of the drought event
 390 characteristics, warming levels, and seasons. Future hydrological drought duration is found

391 to be nonstationary in most catchments across warming levels and seasons. This is most
392 noticeable at 3°C warming, where almost all catchments across seasons are depicting
393 nonstationarity in future hydrological drought duration. Interestingly, future drought
394 intensity at lower warming levels appears to be stationary. Only during the winter season
395 does drought intensity exhibit a trend of rising nonstationarity as the warming increases.
396 Further, at least half of the catchments display nonstationary hydrological drought severity
397 characteristics across warming levels, except during the summer season at lower warming
398 levels. The fluctuations in the nonstationarity properties of catchments specifically, the
399 number of nonstationary catchments declining from 1.5°C to 2°C warming but then increasing
400 at 3°C highlight the limitations of the pattern scaling assumption. This is central to CHES-
401 SCAPE and UKCP18 data considered, which is based on the assumption that local or regional
402 climate responses scale linearly with global mean temperature (Robinson et al.,
403 2022a)(Robinson et al., 2022a). The observed variations suggest that this assumption may
404 break down for certain warming levels or in specific regions, as illustrated in Figure 2, 3.
405 Examining the spatial distribution of nonstationarity across the UK provides insight into where
406 pattern scaling might hold and where caution is needed, highlighting regions dominated by
407 nonlinear responses. Therefore, changes in nonstationary properties, their dependence on
408 warming levels, catchment characteristics, and seasonal variability must be considered with
409 full caution when modelling the evolution of future hydrological droughts. Finally, ~~the~~
410 across models remains overall similar, and no noticeable difference in the ability to capture
411 nonstationarity was observed. ~~However, the changes in nonstationary properties, their~~
412 ~~dependence on warming conditions, characteristics, and seasons need consideration while~~
413 ~~modelling the evolution of future hydrological droughts.~~




Figure 2. Percentage of nonstationary catchments for each event characteristics, hydrological models and warming levels.

414
 415 Once the nonstationarity was assessed, we derived the parameter distribution for calculating
 416 the return levels of future and historical drought duration, intensity, and severity. Figure 4
 417 demonstrates the mean and standard deviation of the posterior distribution of parameters
 418 obtained using the Bayesian framework for the GR4J model during the summer season at
 419 +3°C. The spatial distribution of parameter means and standard deviation, particularly for
 420 duration, suggests that there is relatively higher uncertainty in the location parameter in the
 421 south-eastern catchments. The south-east not only experiences a higher magnitude of mean
 422 location parameter but also higher uncertainty which is in agreement with previous studies
 423 depicting more significant changes in future drought conditions in this region (Kay et al.,
 424 2021)(Kay et al., 2021). The variation of the location parameter across catchments for drought
 425 intensity and severity exhibits more or less similar behaviour. It can also be observed that
 426 catchments with a higher magnitude of the location parameter exhibit a higher standard
 427 deviation. This is crucial and calls for more caution as it denotes, for e.g., a catchment with a
 428 higher duration of drought might show higher uncertainty in the estimates. We also
 429 demonstrate the robustness of the employed method by comparing the curves of posterior
 430 distributions of location parameters for a sample catchment (Dee in Scotland, NRFA ID:

431 67018) for the reference period and +3°C warming (Figure 3). The location parameter for
 432 future drought duration shows a lower value, whereas intensity and severity are generally
 433 higher. This pattern is consistent with the findings from the return level analysis, which are
 434 presented in the next sections. Figure 3 also shows that the possible spread of location
 435 parameters for future drought characteristics is well constrained. This is critical as it ensures
 436 that the model provides robust estimates of parameters, especially for understanding future
 437 changes in drought characteristics under projected warming.




Figure 3. Posterior distribution of parameters for different drought characteristics for a sample (Dee in Scotland, NRFA ID: 67018) catchment in reference period and at 3°C warming level.

438




Figure 4. Mean and standard deviation of duration - number of days, severity - cumecs, intensity - cumecs/day parameter samples for GR4J model during summer season at 3°C warming level.

439

440 3.2. Return levels of different drought characteristics

441 Next, we calculated the return levels of drought duration, severity and intensity at different
 442 return periods (10, 100, and 500 years) using parameter samples from the posterior
 443 distribution obtained through Bayesian analysis. The return levels were calculated for both
 444 the reference period and the warming level periods, considering the stationary case as well
 445 as nonstationary case. As discussed, parameter uncertainty is a key aspect of the
 446 nonstationary hydrological drought risk assessment. To illustrate that the results are
 447 consistent, we computed results using four different summaries of the parameter
 448 distribution: the 25th percentile (Q25), the 75th percentile (Q75), the mean, and the median.
 449 The estimates of return level changes, as well as the differences between nonstationary and
 450 stationary return levels across these four summaries, demonstrate consistency and
 451 robustness throughout the analysis, as shown in Figures S2 and Figure S3. For the sake of
 452 brevity the results presented in the main text of this paper focus exclusively on the mean
 453 return levels, ~~however, different return levels corresponding to median, 75th, and 25th~~

454 ~~quantiles of the posterior parameter distribution were also calculated and can be referred to~~
455 ~~in the supplementary information (Figure a-c) for more insights about uncertainty in the~~
456 ~~estimates.~~

457 Figure 5,6 shows the model average percentage change in mean nonstationary return levels
458 for 10-year (frequent droughts) and 500-year (rare droughts) return levels, respectively. The
459 return level is dependent on the rarity of the drought, as changes in return levels are more
460 pronounced for a 500-year drought compared to a 10-year drought, with the former
461 exhibiting more distinct spatial characterisation. The overall distribution of percentage
462 changes in the mean 100-year return level is shown in the supplementary information
463 (Figure S2b, S3b, S4a-f). For drought duration, the overall return levels are expected to be
464 higher for 500-year droughts during the autumn and winter seasons, whereas they are
465 expected to be lower for 10-year droughts in the same seasons. This increase in the risk of
466 prolonged extreme droughts in autumn and winter is concerning, given that the winter half-
467 year is the critical time for replenishment of aquifers (in the south-east) and reservoirs (Barker
468 et al., 2019; Environment Agency, 2011) (Barker et al., 2019; Environment Agency, 2011). The
469 shorter duration of 10-year droughts may slightly ease water stress during more frequent
470 droughts in these seasons however, any potential benefits could be offset by increased
471 drought intensity, making the overall water management plan in the country still challenging.
472 In Fig. 6, which shows longer drought durations, regions in the north and west, which rely
473 almost entirely on surface water and lack the buffering capacity of groundwater, might be
474 significantly affected, whereas areas in the south-east dominated by groundwater-fed
475 systems might experience delayed drought impacts, offering a degree of resilience during
476 prolonged dry periods. Previous studies have also shown significant variability in
477 hydrometeorological drought characteristics, both in the current period and in future
478 projections, specifically in the southern part of the country (Barker et al., 2019; Di Nunno and
479 Granata, 2024; Reyniers et al., 2022) (Barker et al., 2019; Di Nunno and Granata, 2024;
480 Reyniers et al., 2022). Compared to intensity, duration return levels have more distinct
481 regional attributes for rare droughts - particularly in the spring and summer season where
482 some of the catchments show abrupt negative changes in return levels. Studies suggest that
483 the UK is likely to experience warmer and wetter winters alongside hotter and drier summers
484 in the future (Lowe et al., 2018) (Lowe et al., 2018).

485

Percentage Change in Mean Nonstationary 10 Year Return Levels (Model Avg.)




Figure 5. Percentage change in mean nonstationary 10-year return levels for different drought characteristics across all warming levels and seasons.

Percentage Change in Mean Nonstationary 500 Year Return Levels (Model Avg.)




Figure 6. Model average percentage change in mean nonstationary 500-year return levels for different drought characteristics across all warming levels and seasons.

488 Additionally, most projections indicate an overall increase in potential evapotranspiration,
489 with seasonal variations in the rate of change, but a consistent upward trend on an annual
490 basis (Robinson et al., 2022b)(Robinson et al., 2022b). This could be one of the possible drivers
491 of longer future drought durations for frequent droughts or higher severity of rarer droughts,
492 particularly in the summer season (Kay et al., 2020; Murphy et al., 2018)(Kay et al., 2020;
493 Murphy et al., 2018). Future severity is observed to be increasing for both frequent and rare
494 droughts in most catchments, except during the winter season for frequent droughts at lower
495 warming levels. Season-wise, the increasing changes in the severity of rare droughts in the
496 spring are highest, followed by summer, winter, and autumn. This increase is more substantial
497 at higher warming levels, which indicates that both rare and frequent droughts are, in general,
498 expected to be more severe in the future under the influence of rising temperature (Parry et
499 al., 2024b)(Parry et al., 2024b). Further, the intensity of droughts with a 10-year recurrence
500 interval is projected to increase during the autumn and summer seasons. Conversely, the
501 intensity of droughts with a 500-year return period is found to be decreasing in most seasons
502 across all warming levels. It should be noted that we have considered the mean intensity,
503 which is a function of both duration and severity, and highly intense frequent droughts in the
504 future, particularly in autumn and summer seasons, could be due to highly severe droughts
505 over a smaller duration (Figure 5).

506

507 **3.3. Difference between stationary and nonstationary return levels**

508 To understand the role of temperature in governing changes in future drought characteristics,
509 we compared the stationary return levels with the nonstationary return levels. Figure 7a,b
510 shows the distribution of model-average percentage change in nonstationary and the
511 stationary return levels for seasons and warming levels. The difference in percentage change
512 in hydrological drought intensity return levels for the stationary and nonstationary cases is
513 negative, particularly for higher return periods and warming levels across seasons. This might
514 be because most catchments for drought intensity exhibit stationary characteristics (Figure 2)
515 and show similar spatial patterns for stationary return levels as well (Figure S3a-c). For
516 drought severity, the changes in return levels tend to show a decreasing trend with increased
517 rarity. However, this is exclusive to the autumn season as drought severity in other seasons
518 exhibits higher return levels with higher return periods of droughts. Similar results were
519 observed for the stationary return levels; however, while the overall trend remains

520 consistent, there is a significant difference in the magnitude of the stationary and
 521 nonstationary return levels. Figure S3a-c in the supplementary information shows the spatial
 522 patterns of stationary return levels.




Figure 7 a, b. Spatial average percentage change in mean c) nonstationary and d) stationary return levels (10,100 and 500 years) for different drought characteristics across all warming levels and seasons.—

524 The incorporation of 100-year return levels also confirms the trends in the results, showing
525 that as droughts become less frequent, the changes in return levels become more
526 pronounced. ~~It can also be concluded that rarer droughts are not only accompanied by larger-~~
527 ~~scale changes in return levels but also by larger variability.~~It can also be concluded from Figure
528 7,8 that rarer droughts, which are inherently associated with larger uncertainty contributed
529 by factors such as event identification, estimation of distribution parameters, or an
530 interaction of these factors, are not only associated with larger changes but also with greater
531 overall spatial variability across catchments. This heightened variability underscores the need
532 for robust modelling approaches to better understand the impacts of rare hydrological
533 droughts in the UK under climate change. Most previous studies in the UK have considered
534 different climate model outputs or hydrological models but did not take into account the
535 variability induced due to warming on different drought events on the seasonal scale (Parry
536 et al., 2024b; Rudd et al., 2019)(Parry et al., 2024b; Rudd et al., 2019). Therefore, the results
537 of this analysis provide more comprehensive insights into the varying uncertainty of future
538 return levels.

539 **3.4. Inter-model differences in return levels**

540 In Further, Figure 9 shows the magnitude of the difference between the percentage changes
541 in nonstationary and stationary return levels for 3°C warming level. Results are shown for
542 each model to demonstrate the variability among models. The difference between the
543 nonstationary and stationary return levels is smaller for drought intensity compared to
544 drought duration and severity. This outcome was expected due to the relatively lower level
545 of nonstationarity detected in the drought intensity projections (Figure 2) and a higher
546 severity and lower duration compared to the reference period (Figure 5). This suggests that
547 the mean flow deficit relative to the historical drought threshold on any given day in the
548 future is less likely to be related to temperature change than for duration and severity.
549 However, the number of days over which drought might occur and the total accumulated flow
550 deficit across all days of a drought are more likely to be affected by these factors at higher
551 warming levels. Moreover, the duration of more frequent droughts being less affected by
552 rising temperatures is also confirmed by minimal difference between stationary and
553 nonstationary return levels across seasons, which changes significantly when higher return
554 levels are considered (Figure 9).

555 Overall, the results indicate that failing to incorporate temperature effects in modelling
 556 duration for longer return period droughts can lead to significant uncertainty regarding their
 557 future return levels. This underestimation and variability are most amplified for future
 558 drought severity, where it is evident that temperature influences across models, seasons, and
 559 warming levels might lead to more severe droughts. To further confirm this, we analysed the
 560 distribution of the 25th, 75th quantiles, and the median return levels for different warming
 561 levels (Figure S4a-f), which shows a similar trend. Further, assessing model performance for
 562 future periods compared to a baseline period is challenging because different hydrological
 563 models capture processes and uncertainties based on their individual structure and
 564 operational specifications. Therefore, it is important to incorporate multiple models for more
 565 confident estimates of future changes in drought characteristics (Hannaford et al., 2023; Lane
 566 et al., 2022)(Hannaford et al., 2023; Lane et al., 2022). In this setting, with four hydrological
 567 model outputs assessed, for each drought characteristic, the return levels across the UK are
 568 primarily driven by the rarity of the event in different seasons rather than the model itself.




Figure 8. Mean Percentage change in nonstationary return levels for duration, severity and intensity across different models, seasons and return periods.

569




Figure 9. Difference in spatial average percentage change in return levels for mean nonstationary and stationary return levels for different drought characteristics across all seasons and 3°C warming levels.

570

571 Although the results from this analysis are consistent across the hydrological models, a more
 572 detailed uncertainty partition analysis could be conducted in the future to gain a deeper
 573 understanding of the inter-model differences in the projected characteristics of future
 574 droughts. Further studies could also incorporate catchment hydrometeorological
 575 characteristics in the nonstationary modelling set-up to understand the role of changing
 576 catchment conditions in governing the drought characteristics. In this study, we have looked
 577 at the drought characteristics independently, however, the dependence of drought
 578 characteristics over time, as well as their evolution in a compound setting could give more
 579 useful insights about their interrelation in the future. Despite this, the findings from this
 580 analysis give crucial insights about the changing future hydrological drought characteristics in
 581 the UK under climate change. The results not only quantify the changes in the return level of

582 drought duration, severity, and intensity but also provide explicit estimates of uncertainty in
583 the GEV distribution parameters and associated return levels centred on the methodological
584 framework adopted in this study. The Bayesian approach allows full posterior distribution of
585 the GEV parameters to be explored, enabling return level estimates to be assessed across a
586 wide range of parameter values. This is further supported by using MCMC simulations whose
587 convergence is diagnosed with the Heidelberger-Welch test, which helps to ensure that the
588 posterior distributions are stable and reliable. These elements along with moving window
589 approach and pooling procedure to identify drought events ensure that thorough attention
590 has been given from the initial drought identification through to the estimation of return
591 levels, resulting in reliable and transparently quantified estimates of return level across
592 temporal scales, models, seasons and warming levels.~~Despite this, the findings from this~~
593 ~~analysis give crucial insights about the changing future hydrological drought characteristics in~~
594 ~~the UK under climate change. The results not only point out changing magnitudes of drought~~
595 ~~duration, severity, and intensity but also provide robust estimates of uncertainty on different~~
596 ~~spatial and temporal scales, which can be considered while designing more targeted and~~
597 ~~localized strategies against drought-related challenges in the future.~~

599 **4. Conclusions**

600 This study attempts to understand the evolution of future hydrological droughts in the UK
601 under different warming conditions, utilising nonstationary extreme value analysis with a
602 Bayesian framework for parameter uncertainty. We used the recently developed eFLaG
603 projections to investigate changes in drought characteristics in terms of return levels. The
604 findings indicate that future temperature changes contribute significantly and uniquely to
605 hydrological droughts' characteristics - duration, severity, and intensity. Results demonstrate
606 that the future changes in these characteristics are highly dependent on the season and the
607 rarity of droughts. Drought severity in most cases, irrespective of rarity and season, appears
608 to be increasing in the future at higher warming levels. However, future drought duration and
609 intensity are showing both increasing and decreasing trends depending on the season and
610 return period of droughts. This also underscores the varying degrees of nonstationarity
611 exhibited by different drought characteristics, which should be carefully considered while
612 planning measures against future drought risks in the UK. The projected return levels,
613 particularly for rare and high-impact events, also show a higher level of uncertainty in their

614 magnitude as compared to more frequent events, which can be critical for risk management
615 and adaptation strategies. Overall, this research underlines the importance of considering the
616 influence of temperature-induced nonstationarity in modelling future changes in hydrological
617 drought characteristics. Results from both stationary and nonstationary cases across different
618 seasons, rarities, and warming levels provide comprehensive insights that can be utilised by
619 policymakers and water managers to develop effective strategies against future risks.

620 We conclude that the most critical policy considerations for future hydrological droughts will
621 revolve around adapting to projected nonstationary changes in the nature of risk. As
622 mentioned, the finding that drought severity consistently increases across the majority of
623 catchments under higher warming adds to previous assessments of decreasing future water
624 availability, reaffirming that that policy reviews of water resource infrastructure and
625 management plans are necessary to create buffers against larger future deficits, as well as to
626 mitigate impacts of worsening hydrological droughts on the environment. However our
627 analysis provides greater granularity in terms of providing fine-detail spatial appraisals as well
628 as a multi-seasonal viewpoint as well as considering multiple characteristics of drought
629 (duration, severity and intensity) which is important given the widely varying nature and
630 timing of droughts which catchments, water resource systems and ecosystems alike are
631 vulnerable to around the UK (e.g. Barker et al. 2016; Counsell and Durant, 2023; Stubbington
632 et al. 2024).

633
634 As noted in our introduction, various reviews of the current frameworks for water resources
635 management have highlighted some of the limitations of current stochastic-based planning
636 approaches (Counsell & Durant, 2023; Environment Agency, 2025). Durant and Counsell
637 (2024) argued that ‘the future is transient’ and that more efforts should be directed towards
638 the use of continuous, transient projections like eFLaG, rather than focusing on change point
639 analyses based on time-slices. Here we provide a test-case further highlighting the added
640 value of such transient projections, although we acknowledge that our emphasis is on
641 hydrological droughts and further work is needed to look at the onward impacts on complex
642 water supply systems. Furthermore, our observation that changes in drought duration and
643 intensity are highly dependent on the season points toward a required shift from uniform,
644 year-round planning to seasonally specific risk management strategies. This bolsters the
645 argument (e.g. Environment Agency, 2025) for further investigation of ‘bottom-up’ storyline

646 [approaches to stress tests systems according to the types of drought they are vulnerable to,](#)
647 [in terms of seasonality and duration \(e.g. Chan et al. 2022\).](#) Finally, the higher uncertainty
648 [observed, particularly for rare high-impact droughts, such as the 1:200 and 1:500 year events](#)
649 [that are a cornerstone of planning indicates that future policy must explicitly integrate the](#)
650 [possibility of extreme outcomes beyond currently accepted limits of uncertainty, requiring](#)
651 [robust, nonstationary modelling in all risk management and adaptation strategies.](#)

652

653 **Code and data availability**

654 The eFLaG river flow projections analysed in this study are stored at the UKCEH's
655 Environmental Information Data Centre and can be freely accessed as DOI datasets. Please
656 ensure these data are cited in full when used in any application:

657 <https://catalogue.ceh.ac.uk/documents/1bb90673-ad37-4679-90b9-0126109639a9>. The

658 CHESS-SCAPE dataset can be downloaded from the NERC Environmental Data Service (EDS)
659 Centre for Environmental Data Analysis (CEDA) via the following link:

660 <https://doi.org/10.5285/8194b416cbee482b89e0dfbe17c5786c>. The R scripts used for
661 analysis were developed using publicly available packages, such as 'extRemes', 'evir', 'coda',
662 'foreach', and 'doparallel', which support extreme value analysis, Markov Chain Monte Carlo
663 diagnostics in a parallel environment.

664

665 **Author contribution**

666 Conceptualization was done by SJ, JH, MT, and LB. Methodology development and analysis
667 were carried out by SJ. The original draft was written by SJ and JH. Reviewing and editing of
668 the manuscript were performed by LB, JH, and MT. Supervision of the work was provided by
669 JH, LB, and MT.

670

671 **Competing interests statement**

672 The authors declare that they have no conflicts of interests.

673

674 **Acknowledgements**

675 We also acknowledge the use of the JASMIN high-performance computing facility for the
676 Bayesian analysis conducted in this study. JASMIN facility is operated by the Science and
677 Technology Facilities Council on behalf of the Natural Environment Research Council.

678

679 **Financial support**

680 This work has been financially supported by the Hydro-JULES Programme (NE/S017380/1)

681 funded by Natural Environment Research Council.

682 **Reference**

683 [Anderson, B. J., Muñoz-Castro, E., Tallaksen, L. M., Matano, A., Götte, J., Armitage, R., Magee, E., and](#)
684 [Brunner, M. I.: What is a drought-to-flood transition? Pitfalls and recommendations for defining](#)
685 [consecutive hydrological extreme events, *Hydrol. Earth Syst. Sci.*, 29, 6069–6092,](#)
686 <https://doi.org/https://doi.org/10.5194/egusphere-2025-1391>, 2025.

Formatted: Font colour: Black

687 [Barker, L. J., Hannaford, J., Parry, S., Smith, K. A., Tanguy, M., and Prudhomme, C.: Historic](#)
688 [hydrological droughts 1891–2015: systematic characterisation for a diverse set of catchments across](#)
689 [the UK, *Hydrol. Earth Syst. Sci.*, 23, 4583–4602, https://doi.org/10.5194/hess-23-4583-2019](#), 2019.

Formatted: Font: 11 pt, Font colour: Black

Formatted: Font colour: Black

690 [Barker, L. J., Hannaford, J., Magee, E., Turner, S., Sefton, C., Parry, S., Evans, J., Szczykulska, M., and](#)
691 [Haxton, T.: An appraisal of the severity of the 2022 drought and its impacts, *Weather*, 79, 208–219,](#)
692 <https://doi.org/10.1002/wea.4531>, 2024.

693 [Baykal, T., Taylan, E. D., Eriskin, E., and Terzi, Ö.: Predicting Hydrological Droughts of Long-Narrow](#)
694 [Type Drainage Basin Using Monte Carlo Technique, *J. Hydrol. Eng.*, 29, 04024013,](#)
695 <https://doi.org/10.1061/jhyeff.heeng-6144>, 2024.

696 [Bell, V. A., Kay, A. L., Rudd, A. C., and Davies, H. N.: The MaRIUS-G2G datasets: Grid-to-Grid model](#)
697 [estimates of flow and soil moisture for Great Britain using observed and climate model driving data,](#)
698 [Geosci. Data J., 5, 63–72, https://doi.org/10.1002/gdj3.55](#), 2018.

699 [Bevan, J.: Escaping the jaws of death: ensuring enough water in 2050, GOV. UK. Retrieved](#)
700 [November, 12, 2020, 2019.](#)

701 [Bevan, J.: Drought risk in the Anthropocene: from the jaws of death to the waters of life,](#)
702 [Philosophical Transactions of the Royal Society A, 380, 20220003,](#)
703 <https://doi.org/10.1098/rsta.2022.0003>, 2022.

704 [Brunner, M. I. and Chartier-Rescan, C.: Drought spatial extent and dependence increase during](#)
705 [drought propagation from the atmosphere to the hydrosphere, *Geophys. Res. Lett.*, 51,](#)
706 [e2023GL107918, https://doi.org/https://doi.org/10.1029/2023GL107918](https://doi.org/https://doi.org/10.1029/2023GL107918), 2024.

Formatted: Font colour: Black, German (Germany)

707 [Brunner, M. I., Slater, L., Tallaksen, L. M., and Clark, M.: Challenges in modeling and predicting floods](#)
708 [and droughts: A review, *Wiley Interdisciplinary Reviews: Water*, 8, e1520,](#)
709 <https://doi.org/10.1002/wat2.1520>, 2021.

Formatted: Font colour: Black

710 [Chan, W. C. H., Shepherd, T. G., Facer-Childs, K., Darch, G., and Arnell, N. W.: Storylines of UK](#)
711 [drought based on the 2010–2012 event, *Hydrol. Earth Syst. Sci.*, 26, 1755–1777,](#)
712 <https://doi.org/10.5194/hess-26-1755-2022>, 2022.

713 [Chan, W. C. H., Arnell, N. W., Darch, G., Facer-Childs, K., Shepherd, T. G., and Tanguy, M.: Added](#)
714 [value of seasonal hindcasts to create UK hydrological drought storylines, *Natural Hazards and Earth*](#)
715 [System Sciences, 24, 1065–1078, 2024.](#)

716 [Chandra, R., Saha, U., and Mujumdar, P. P.: Model and parameter uncertainty in IDF relationships](#)
717 [under climate change, *Adv. Water Resour.*, 79, 127–139,](#)
718 <https://doi.org/10.1016/j.advwatres.2015.02.011>, 2015.

719 [Climate Change Committee: National Infrastructure Commission: Preparing for a drier future-](#)
720 [England's water infrastructure needs, https://www.theccc.org.uk/wp-](#)
721 [content/uploads/2019/07/Outcomes-Water-stress-case-study.pdf](#), 2018.

722 [Climate Change Committee: Water Stress and Resilience, https://www.theccc.org.uk/wp-](#)
723 [content/uploads/2019/07/Outcomes-Water-stress-case-study.pdf](#), 2019.

724 [Climate Change Committee: Independent assessment of UK climate risk,](#)
725 <https://www.theccc.org.uk/wp-content/uploads/2021/07/Independent-Assessment-of-UK-Climate->
726 [Risk-Advice-to-Govt-for-CCRA3-CCC.pdf, 2021.](#)

727 [Coles, S.: An introduction to statistical modeling of extreme values, Springer, London,](#)
728 [Counsell, C. and Durant, M.: Water supply – observed and projected. In: Environment Agency,](https://doi.org/10.1007/978-1-4471-3675-0, 2001.</p><p>729 <a href=)
730 <https://www.gov.uk/government/publications/national-framework-for-water-resources-2025->
731 [water-for-growth-nature-and-a-resilient-future/forewords-executive-summary-and-structure-](#)
732 [national-framework-for-water-resources-2025, 2023.](#)

733 [Das, J. and Umamahesh, N. V.: Uncertainty and Nonstationarity in Streamflow Extremes under](#)
734 [Climate Change Scenarios over a River Basin, J. Hydrol. Eng., 22, 04017042,](#)
735 [Environment Agency: Impact of long droughts on water resources, Horizon House, Deanery Road,](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001571, 2017.</p><p>736 <a href=)
737 [Bristol, BS1 5AH, SC070079/R5-ISBN: 978-1-84911-273-4 pp., 2011.](#)

738 [Environment Agency: Review of stochastic and other approaches in water resources planning,](#)
739 [SC220039/R1, 2025.](#)

740 [Gilleland, E. and Katz, R. W.: extRemes 2.0: An Extreme Value Analysis Package in R, J. Stat. Softw.,](#)
741 [72, https://doi.org/10.18637/jss.v072.i08, 2016.](#)

742 [Hammond, J. C., Simeone, C., Hecht, J. S., Hodgkins, G. A., Lombard, M., McCabe, G., Wolock, D.,](#)
743 [Wieczorek, M., Olson, C., and Caldwell, T.: Going beyond low flows: Streamflow drought deficit and](#)
744 [duration illuminate distinct spatiotemporal drought patterns and trends in the US during the last](#)
745 [century, Water Resour. Res., 58, e2022WR031930, 2022.](#)

746 [Hannaford, J., Mackay, J., Ascott, M., Bell, V., Chitson, T., Cole, S., Counsell, C., Durant, M., Jackson,](#)
747 [C. R., and Kay, A.: eFLaG: enhanced future FLoWs and Groundwater. A national dataset of](#)
748 [hydrological projections based on UKCP18, Earth System Science Data Discussions, 2022, 1–40,](#)
749 [Hannaford, J., Mackay, J., Ascot, M., Bell, V., Chitson, T., Cole, S., Counsell, C., Durant, M., Facer-](https://doi.org/https://doi.org/10.5194/essd-15-2391-2023, 2022a.</p><p>750 <a href=)
751 [Childs, K., and Jackson, C.: Hydrological projections for the UK, based on UK Climate Projections 2018](#)
752 [\(UKCP18\) data, from the Enhanced Future Flows and Groundwater \(eFLaG\) project, Environmental](#)
753 [Information Data Centre. doi, 10, https://doi.org/10.5285/1bb90673-ad37-4679-90b9-](#)
754 [0126109639a9, 2022b.](#)

755 [Hannaford, J., Mackay, J. D., Ascott, M., Bell, V. A., Chitson, T., Cole, S., Counsell, C., Durant, M.,](#)
756 [Jackson, C. R., Kay, A. L., Lane, R. A., Mansour, M., Moore, R., Parry, S., Rudd, A. C., Simpson, M.,](#)
757 [Facer-Childs, K., Turner, S., Wallbank, J. R., Wells, S., and Wilcox, A.: The enhanced future Flows and](#)
758 [Groundwater dataset: Development and evaluation of nationally consistent hydrological projections](#)
759 [based on UKCP18, Earth Syst. Sci. Data, 15, 2391–2415, https://doi.org/10.5194/essd-15-2391-2023,](#)
760 [2023.](#)

761 [Hasan, H. H., Mohd Razali, S. F., Muhammad, N. S., Mohamed, Z. S., and Mohamad Hamzah, F.:](#)
762 [Assessment of probability distributions and minimum storage draft-rate analysis in the equatorial](#)
763 [region, Natural Hazards and Earth System Sciences Discussions, 2020, 1–29,](#)
764 [31](https://doi.org/https://doi.org/10.5194/nhess-21-1-2021, 2020.</p></div><div data-bbox=)

765 [Janicka-Kubiak, E.: Hydrological drought trends and seasonality in selected Polish catchments](#)
766 [between 1993 and 2022 using a threshold based approach, *Sci. Rep.*, 15, 40454,](#)
767 <https://doi.org/https://doi.org/10.1038/s41598-025-24133-1>, 2025.

768 [Jha, S., Goyal, M. K., Gupta, B. B., Hsu, C., Gilleland, E., and Das, J.: A methodological framework for](#)
769 [extreme climate risk assessment integrating satellite and location based data sets in intelligent](#)
770 [systems, *International Journal of Intelligent Systems*, 37, 10268–10288,](#)
771 <https://doi.org/10.1002/int.22475>, 2022.

772 [Jha, S., Gudmundsson, L., and Seneviratne, S. I.: Partitioning the uncertainties in compound hot and](#)
773 [dry precipitation, soil moisture, and runoff extremes projections in CMIP6, *Earths Future*, 11,](#)
774 [e2022EF003315](https://doi.org/10.1029/2022EF003315), <https://doi.org/10.1029/2022EF003315>, 2023.

775 [Kay, A. L., Griffin, A., Rudd, A. C., Chapman, R. M., Bell, V. A., and Arnell, N. W.: Climate change](#)
776 [effects on indicators of high and low river flow across Great Britain, *Adv. Water Resour.*, 151,](#)
777 [103909](https://doi.org/10.1016/j.advwatres.2021.103909), <https://doi.org/10.1016/j.advwatres.2021.103909>, 2021.

778 [Kay, G., Dunstone, N., Smith, D., Dunbar, T., Eade, R., and Scaife, A.: Current likelihood and dynamics](#)
779 [of hot summers in the UK, *Environmental Research Letters*, 15, 094099,](#)
780 <https://doi.org/10.1088/1748-9326/abab32>, 2020.

781 [Kendon, M., Doherty, A., Hollis, D., Carlisle, E., Packman, S., McCarthy, M., Jevrejeva, S., Matthews,](#)
782 [A., Williams, J., and Garforth, J.: State of the UK Climate 2023, *International Journal of Climatology*,](#)
783 [44, 1–117](https://doi.org/https://doi.org/10.1002/joc.8553), <https://doi.org/https://doi.org/10.1002/joc.8553>, 2024.

784 [Kuana, L. A., Almeida, A. S., Mercuri, E. G. F., and Noe, S. M.: Regionalization of GR4J model](#)
785 [parameters for river flow prediction in Paraná, Brazil, *Hydrol. Earth Syst. Sci.*, 28, 3367–3390,](#)
786 <https://doi.org/10.5194/hess-28-3367-2024>, 2024.

787 [Lane, R. A., Coxon, G., Freer, J., Seibert, J., and Wagener, T.: A large-sample investigation into](#)
788 [uncertain climate change impacts on high flows across Great Britain, *Hydrol. Earth Syst. Sci.*, 26,](#)
789 [5535–5554](https://doi.org/https://doi.org/10.5194/hess-26-5535-2022), <https://doi.org/https://doi.org/10.5194/hess-26-5535-2022>, 2022.

790 [Leng, J., Ma, K., Gu, S., Zhang, K., and He, D.: A non-stationary impactquant framework for assessing](#)
791 [the human activity impacts on hydrological drought in the Upper Red River, *Atmos. Res.*, 304,](#)
792 [107419](https://doi.org/10.1016/j.atmosres.2024.107419), <https://doi.org/10.1016/j.atmosres.2024.107419>, 2024.

793 [Liu, R., Yin, J., Slater, L., Kang, S., Yang, Y., Liu, P., Guo, J., Gu, X., Zhang, X., and Volchak, A.: Machine-](#)
794 [learning-constrained projection of bivariate hydrological drought magnitudes and socioeconomic](#)
795 [risks over China, *Hydrol. Earth Syst. Sci.*, 28, 3305–3326, https://doi.org/10.5194/hess-28-3305-](#)
796 [2024](https://doi.org/10.5194/hess-28-3305-2024), 2024.

797 [Van Loon, A. F. and Van Lanen, H.: A process-based typology of hydrological drought, *Hydrology and*](#)
798 [Earth System Sciences](#), 16, 1915–1946, 2012.

799 [Lowe, J. A., Bernie, D., Bett, P., Bricheno, L., Brown, S., Calvert, D., Clark, R., Eagle, K., Edwards, T.,](#)
800 [and Fosser, G.: UKCP18 science overview report, Met Office Hadley Centre: Exeter, UK, 1–73, 2018.](#)

801 [Lück, A. and Wolf, V.: Generalized method of moments for estimating parameters of stochastic](#)
802 [reaction networks, *BMC Syst. Biol.*, 10, 98, https://doi.org/10.1186/s12918-016-0342-8](#), 2016.

803 [Moore, R. J.: The PDM rainfall-runoff model, *Hydrol. Earth Syst. Sci.*, 11, 483–499,](#)
804 <https://doi.org/10.5194/hess-11-483-2007>, 2007.

805 [Murphy, C., Wilby, R. L., Matthews, T., Horvath, C., Crampsie, A., Ludlow, F., Noone, S., Brannigan, J.,](#)
806 [Hannaford, J., and McLeman, R.: The forgotten drought of 1765–1768: Reconstructing and re-](#)

807 [evaluating historical droughts in the British and Irish Isles, *International Journal of Climatology*, 40,](#)
808 [5329, <https://doi.org/10.1002/joc.6521>, 2020.](#)

809 [Murphy, J. M., Harris, G. R., Sexton, D. M. H., Kendon, E. J., Bett, P. E., Clark, R. T., Eagle, K. E., Fosser,](#)
810 [G., Fung, F., and Lowe, J. A.: UKCP18 land projections: science report, 2018.](#)

811 [Ndiaye, P. M., Bodian, A., Dezetter, A., Ogilvie, A., and Goudiaby, O.: Sensitivity of global hydrological](#)
812 [models to potential evapotranspiration estimation methods in the Senegal River Basin \(West Africa\),](#)
813 [J. Hydrol. Reg. Stud., 53, 101823, <https://doi.org/10.1016/j.ejrh.2024.101823>, 2024.](#)

814 [Di Nunno, F. and Granata, F.: Analysis of trends and abrupt changes in groundwater and](#)
815 [meteorological droughts in the United Kingdom, *J. Hydrol. \(Amst.\)*, 131430,](#)
816 [https://doi.org/10.1016/j.jhydrol.2024.131430, 2024.](#)

817 [OFWAT: Towards resilience: how we will embed resilience in our work,](#)
818 [https://www.ofwat.gov.uk/wp-](https://www.ofwat.gov.uk/wp-content/uploads/2015/07/pap_pos20151210towardsresiliencerev.pdf)
819 [content/uploads/2015/07/pap_pos20151210towardsresiliencerev.pdf, 2015.](#)

820 [Onyutha, C.: On rigorous drought assessment using daily time scale: Non-stationary frequency](#)
821 [analyses, revisited concepts, and a new method to yield non-parametric indices, *Hydrology*, 4, 48,](#)
822 [https://doi.org/10.3390/hydrology4040048, 2017.](#)

823 [Parry, S., Mackay, J. D., Chitson, T., Hannaford, J., Bell, V. A., Facer-Childs, K., Kay, A., Lane, R.,](#)
824 [Moore, R. J., Turner, S., and Wallbank, J.: Divergent future drought projections in UK river flows and](#)
825 [groundwater levels, <https://doi.org/10.5194/hess-28-417-2024>, 28 February 2024a.](#)

826 [Parry, S., Mackay, J. D., Chitson, T., Hannaford, J., Magee, E., Tanguy, M., Bell, V. A., Facer-Childs, K.,](#)
827 [Kay, A., and Lane, R.: Divergent future drought projections in UK river flows and groundwater levels,](#)
828 [Hydrol. Earth Syst. Sci., 28, 417–440, <https://doi.org/10.5194/hess-28-417-2024>, 2024b.](#)

829 [Parvizi, S., Eslamian, S., Gheysari, M., Gohari, A., and Kopai, S. S.: Regional frequency analysis of](#)
830 [drought severity and duration in Karkheh River Basin, Iran using univariate L-moments method,](#)
831 [Environ. Monit. Assess., 194, 336, <https://doi.org/10.1007/s10661-022-09977-8>, 2022.](#)

832 [Posada, D. and Buckley, T.: Model Selection and Model Averaging in Phylogenetics: Advantages of](#)
833 [Akaike Information Criterion and Bayesian Approaches Over Likelihood Ratio Tests, *Syst. Biol.*, 53,](#)
834 [793–808, <https://doi.org/10.1080/10635150490522304>, 2004.](#)

835 [Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., Fekete, B. M.,](#)
836 [Franssen, W., Gerten, D., and Gosling, S. N.: Hydrological droughts in the 21st century, hotspots and](#)
837 [uncertainties from a global multimodel ensemble experiment, *Proceedings of the National Academy*](#)
838 [of Sciences, 111, 3262–3267, <https://doi.org/https://doi.org/10.1073/pnas.1222473110>, 2014.](#)

839 [Ragulina, G. and Reitan, T.: Generalized extreme value shape parameter and its nature for extreme](#)
840 [precipitation using long time series and the Bayesian approach, *Hydrological sciences journal*, 62,](#)
841 [863–879, <https://doi.org/http://dx.doi.org/10.1080/02626667.2016.1260134>, 2017.](#)

842 [Reis, D. S. and Stedinger, J. R.: Bayesian MCMC flood frequency analysis with historical information,](#)
843 [J. Hydrol. \(Amst.\), 313, 97–116, <https://doi.org/10.1016/j.jhydrol.2005.02.028>, 2005.](#)

844 [Reyniers, N., Osborn, T. J., Addor, N., and Darch, G.: Projected changes in droughts and extreme](#)
845 [droughts in Great Britain strongly influenced by the choice of drought index, *Hydrology and Earth*](#)
846 [System Sciences Discussions, 2022, 1–36, <https://doi.org/10.5194/hess-27-1151-2023>, 2022.](#)

847 [Robinson, E. L., Huntingford, C., Shamsudheen, S., and Bullock, J.: CHES-SCAPE: Future projections](#)
848 [of meteorological variables at 1 km resolution for the United Kingdom 1980–2080 derived from UK](#)

849 [Climate Projections 2018, NERC EDS Centre for Environmental Data Analysis,](#)
850 <https://doi.org/http://dx.doi.org/10.5285/8194b416cbee482b89e0dfbe17c5786c>, 2022a.

851 [Robinson, E. L., Brown, M. J., Kay, A. L., Lane, R. A., Chapman, R., Bell, V. A., and Blyth, E. M.: Hydro-](#)
852 [PE: gridded datasets of historical and future Penman-Monteith potential evaporation for the United](#)
853 [Kingdom, Earth System Science Data Discussions, 2022, 1–44, 2022b.](#)

854 [Rudd, A. C., Kay, A. L., and Bell, V. A.: National-scale analysis of future river flow and soil moisture](#)
855 [droughts: potential changes in drought characteristics, Clim. Change, 156, 323–340,](#)
856 <https://doi.org/10.1007/s10584-019-02528-0>, 2019.

857 [Salas, J. D. and Obeysekera, J.: Revisiting the Concepts of Return Period and Risk for Nonstationary](#)
858 [Hydrologic Extreme Events, J. Hydrol. Eng., 19, 554–568, https://doi.org/10.1061/\(ASCE\)HE.1943-](#)
859 [5584.0000820](#), 2014.

860 [Sarailidis, G., Vasiliades, L., and Loukas, A.: Analysis of streamflow droughts using fixed and variable](#)
861 [thresholds, Hydrol. Process., 33, 414–431, 2019.](#)

862 [Sarhadi, A., Burn, D. H., Concepción Ausín, M., and Wiper, M. P.: Time-varying nonstationary](#)
863 [multivariate risk analysis using a dynamic Bayesian copula, Water Resour. Res., 52, 2327–2349, 2016.](#)

864 [Seneviratne, S. and Hauser, M.: Regional climate sensitivity of climate extremes in CMIP6 versus](#)
865 [CMIP5 multimodel ensembles, Earths Future, 8, e2019EF001474, 2020.](#)

866 [Sharma, S. and Mujumdar, P. P.: Modeling concurrent hydroclimatic extremes with parametric](#)
867 [multivariate extreme value models, Water Resour. Res., 58, e2021WR031519, 2022.](#)

868 [Stahl, K., Vidal, J.-P., Hannaford, J., Tjeldeman, E., Laaha, G., Gauster, T., and Tallaksen, L. M.: The](#)
869 [challenges of hydrological drought definition, quantification and communication: an interdisciplinary](#)
870 [perspective, Proceedings of the International Association of Hydrological Sciences, 383, 291–295,](#)
871 [2020.](#)

872 [Strupczewski, W. G., Singh, V. P., and Feluch, W.: Non-stationary approach to at-site flood frequency](#)
873 [modelling I. Maximum likelihood estimation, J. Hydrol. \(Amst\)., 248, 123–142, 2001.](#)

874 [Tallaksen, L. M. and Van Lanen, H. A. J.: Hydrological drought: processes and estimation methods for](#)
875 [streamflow and groundwater, 2023.](#)

876 [Tanguy, M., Magee, E., Hannaford, J., Bell, V., Rameshwaran, P., Baron, H., Keller, V., and Barker, L.:](#)
877 [CS-NOW-D2 Task 5: Analysis of future scenarios,](#)
878 [water-resources-output-analysis.pdf">https://assets.publishing.service.gov.uk/media/672b4fbbabb279b2de1e8c34/cs-now-d2-future-](https://assets.publishing.service.gov.uk/media/672b4fbbabb279b2de1e8c34/cs-now-d2-future-
879 <a href=)
[water-resources-output-analysis.pdf](#), 2023a.

880 [Tanguy, M., Chevuturi, A., Marchant, B. P., Mackay, J. D., Parry, S., and Hannaford, J.: How will](#)
881 [climate change affect the spatial coherence of streamflow and groundwater droughts in Great](#)
882 [Britain?, Environmental Research Letters, 18, 064048, 2023b.](#)

883 [Turner, S., Barker, L. J., Hannaford, J., Muchan, K., Parry, S., and Sefton, C.: The 2018/2019 drought in](#)
884 [the UK: a hydrological appraisal, Weather, 76, 248–253, 2021.](#)

885 [Wu, D., Chen, J., Xiong, L., Lee, J.-H., Kim, J.-S., and Moon, H.-T.: Assessing global drought conditions](#)
886 [under climate change: A comparison of stationary and non-stationary approaches and identification](#)
887 [of hotspot regions, J. Hydrol. \(Amst\)., 131663, 2024.](#)

888 [Yilmaz, A. G. and Perera, B. J. C.: Extreme Rainfall Nonstationarity Investigation and Intensity –](#)
889 [Frequency – Duration Relationship, J. Hydrol. Eng., 19, 1160–1172,](#)
890 [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000878.](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000878.), 2014.

891 [Anderson, B. J., Muñoz-Castro, E., Tallaksen, L. M., Matano, A., Götte, J., Armitage, R., Magee, E.,](#)
892 [and Brunner, M. I.: What is a drought-to-flood transition? Pitfalls and recommendations for defining](#)
893 [consecutive hydrological extreme events, Hydrol. Earth Syst. Sci., 29, 6069–6092,](#)
894 <https://doi.org/https://doi.org/10.5194/egusphere-2025-1391>, 2025.

895 [Barker, L. J., Hannaford, J., Parry, S., Smith, K. A., Tanguy, M., and Prudhomme, C.: Historic](#)
896 [hydrological droughts 1891–2015: systematic characterisation for a diverse set of catchments across](#)
897 [the UK, Hydrol. Earth Syst. Sci., 23, 4583–4602, https://doi.org/10.5194/hess-23-4583-2019, 2019.](#)

898 [Barker, L. J., Hannaford, J., Magee, E., Turner, S., Sefton, C., Parry, S., Evans, J., Szczykulska, M., and](#)
899 [Haxton, T.: An appraisal of the severity of the 2022 drought and its impacts, Weather, 79, 208–219,](#)
900 <https://doi.org/10.1002/wea.4531>, 2024.

901 [Baykal, T., Taylan, E. D., Eriskin, E., and Terzi, Ö.: Predicting Hydrological Droughts of Long-Narrow](#)
902 [Type Drainage Basin Using Monte-Carlo Technique, J. Hydrol. Eng., 29, 04024013,](#)
903 <https://doi.org/10.1061/jhyeff.heeng.6144>, 2024.

904 [Bell, V. A., Kay, A. L., Rudd, A. C., and Davies, H. N.: The MaRIUS-G2G datasets: Grid-to-Grid model](#)
905 [estimates of flow and soil moisture for Great Britain using observed and climate model driving data,](#)
906 [Geosci. Data J., 5, 63–72, https://doi.org/10.1002/gdj3.55, 2018.](#)

907 [Bevan, J.: Escaping the jaws of death: ensuring enough water in 2050, GOV. UK. Retrieved](#)
908 [November, 12, 2020, 2019.](#)

909 [Bevan, J.: Drought risk in the Anthropocene: from the jaws of death to the waters of life,](#)
910 [Philosophical Transactions of the Royal Society A, 380, 20220003,](#)
911 <https://doi.org/10.1098/rsta.2022.0003>, 2022.

912 [Brunner, M. I. and Chartier-Rescan, C.: Drought spatial extent and dependence increase during](#)
913 [drought propagation from the atmosphere to the hydrosphere, Geophys. Res. Lett., 51,](#)
914 [e2023GL107918, https://doi.org/https://doi.org/10.1029/2023GL107918](https://doi.org/https://doi.org/10.1029/2023GL107918), 2024.

915 [Brunner, M. I., Slater, L., Tallaksen, L. M., and Clark, M.: Challenges in modeling and predicting floods](#)
916 [and droughts: A review, Wiley Interdisciplinary Reviews: Water, 8, e1520,](#)
917 <https://doi.org/10.1002/wat2.1520>, 2021.

918 [Chan, W. C. H., Shepherd, T. G., Facer-Childs, K., Darch, G., and Arnell, N. W.: Storylines of UK](#)
919 [drought based on the 2010–2012 event, Hydrol. Earth Syst. Sci., 26, 1755–1777,](#)
920 <https://doi.org/10.5194/hess-26-1755-2022>, 2022.

921 [Chan, W. C. H., Arnell, N. W., Darch, G., Facer-Childs, K., Shepherd, T. G., and Tanguy, M.: Added](#)
922 [value of seasonal hindcasts to create UK hydrological drought storylines, Natural Hazards and Earth](#)
923 [System Sciences, 24, 1065–1078, 2024.](#)

924 [Chandra, R., Saha, U., and Mujumdar, P. P.: Model and parameter uncertainty in IDF relationships](#)
925 [under climate change, Adv. Water Resour., 79, 127–139,](#)
926 <https://doi.org/10.1016/j.advwatres.2015.02.011>, 2015.

927 [Climate Change Committee: National Infrastructure Commission: Preparing for a drier future-](#)
928 [England's water infrastructure needs, https://www.theccc.org.uk/wp-](#)
929 [content/uploads/2019/07/Outcomes-Water-stress-case-study.pdf, 2018.](#)

930 Climate Change Committee: Water Stress and Resilience, [https://www.theccc.org.uk/wp-](https://www.theccc.org.uk/wp-content/uploads/2019/07/Outcomes-Water-stress-case-study.pdf)
931 [content/uploads/2019/07/Outcomes-Water-stress-case-study.pdf](https://www.theccc.org.uk/wp-content/uploads/2019/07/Outcomes-Water-stress-case-study.pdf), 2019.

932 Climate Change Committee: Independent assessment of UK climate risk,
933 [https://www.theccc.org.uk/wp-content/uploads/2021/07/Independent-Assessment-of-UK-Climate-](https://www.theccc.org.uk/wp-content/uploads/2021/07/Independent-Assessment-of-UK-Climate-Risk-Advice-to-Govt-for-CCRA3-CCC.pdf)
934 [Risk-Advice-to-Govt-for-CCRA3-CCC.pdf](https://www.theccc.org.uk/wp-content/uploads/2021/07/Independent-Assessment-of-UK-Climate-Risk-Advice-to-Govt-for-CCRA3-CCC.pdf), 2021.

935 Coles, S.: An introduction to statistical modeling of extreme values, Springer, London,
936 <https://doi.org/10.1007/978-1-4471-3675-0>, 2001.

937 Counsell, C. and Durant, M.: Water supply—observed and projected. In: Environment Agency,
938 [https://www.gov.uk/government/publications/national-framework-for-water-resources-2025-](https://www.gov.uk/government/publications/national-framework-for-water-resources-2025-water-for-growth-nature-and-a-resilient-future/forewords-executive-summary-and-structure-national-framework-for-water-resources-2025)
939 [water-for-growth-nature-and-a-resilient-future/forewords-executive-summary-and-structure-](https://www.gov.uk/government/publications/national-framework-for-water-resources-2025-water-for-growth-nature-and-a-resilient-future/forewords-executive-summary-and-structure-national-framework-for-water-resources-2025)
940 [national-framework-for-water-resources-2025](https://www.gov.uk/government/publications/national-framework-for-water-resources-2025-water-for-growth-nature-and-a-resilient-future/forewords-executive-summary-and-structure-national-framework-for-water-resources-2025), 2023.

941 Das, J. and Umamahesh, N. V.: Uncertainty and Nonstationarity in Streamflow Extremes under
942 Climate Change Scenarios over a River Basin, *J. Hydrol. Eng.*, 22, 04017042,
943 [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0001571](https://doi.org/10.1061/(ASCE)HE.1943-5584.0001571), 2017.

944 Environment Agency: Impact of long droughts on water resources, Horizon House, Deanery Road,
945 Bristol, BS1 5AH, SC070079/R5 ISBN: 978-1-84911-273-4 pp., 2011.

946 Environment Agency: Review of stochastic and other approaches in water resources planning,
947 SC220039/R1, 2025.

948 Gilleland, E. and Katz, R. W.: extRemes 2.0: An Extreme Value Analysis Package in R, *J. Stat. Softw.*,
949 72, <https://doi.org/10.18637/jss.v072.i08>, 2016.

950 Hammond, J. C., Simeone, C., Hecht, J. S., Hodgkins, G. A., Lombard, M., McCabe, G., Wolock, D.,
951 Wiczorek, M., Olson, C., and Caldwell, T.: Going beyond low flows: Streamflow drought deficit and
952 duration illuminate distinct spatiotemporal drought patterns and trends in the US during the last
953 century, *Water Resour. Res.*, 58, e2022WR031930, 2022.

954 Hannaford, J., Mackay, J., Ascott, M., Bell, V., Chitson, T., Cole, S., Counsell, C., Durant, M., Jackson,
955 C. R., and Kay, A.: eFLaG: enhanced future Flows and Groundwater. A national dataset of
956 hydrological projections based on UKCP18, *Earth System Science Data Discussions*, 2022, 1–40,
957 <https://doi.org/https://doi.org/10.5194/essd-15-2391-2023>, 2022a.

958 Hannaford, J., Mackay, J., Ascot, M., Bell, V., Chitson, T., Cole, S., Counsell, C., Durant, M., Facer-
959 Childs, K., and Jackson, C.: Hydrological projections for the UK, based on UK Climate Projections 2018
960 (UKCP18) data, from the Enhanced Future Flows and Groundwater (eFLaG) project, *Environmental*
961 *Information Data Centre*. doi, 10, [https://doi.org/10.5285/1bb90673-ad37-4679-90b9-](https://doi.org/10.5285/1bb90673-ad37-4679-90b9-0126109639a9)
962 [0126109639a9](https://doi.org/10.5285/1bb90673-ad37-4679-90b9-0126109639a9), 2022b.

963 Hannaford, J., Mackay, J. D., Ascott, M., Bell, V. A., Chitson, T., Cole, S., Counsell, C., Durant, M.,
964 Jackson, C. R., Kay, A. L., Lane, R. A., Mansour, M., Moore, R., Parry, S., Rudd, A. C., Simpson, M.,
965 Facer-Childs, K., Turner, S., Wallbank, J. R., Wells, S., and Wilcox, A.: The enhanced future Flows and
966 Groundwater dataset: Development and evaluation of nationally consistent hydrological projections
967 based on UKCP18, *Earth Syst. Sci. Data*, 15, 2391–2415, <https://doi.org/10.5194/essd-15-2391-2023>,
968 2023.

969 Hasan, H. H., Mohd Razali, S. F., Muhammad, N. S., Mohamed, Z. S., and Mohamad Hamzah, F.:
970 Assessment of probability distributions and minimum storage draft rate analysis in the equatorial
971 region, *Natural Hazards and Earth System Sciences Discussions*, 2020, 1–29,
972 <https://doi.org/https://doi.org/10.5194/nhess-21-1-2021>, 2020.

973 Janicka-Kubiak, E.: Hydrological drought trends and seasonality in selected Polish catchments
974 between 1993 and 2022 using a threshold-based approach, *Sci. Rep.*, **15**, 40454,
975 <https://doi.org/https://doi.org/10.1038/s41598-025-24133-1>, 2025.

976 Jha, S., Goyal, M. K., Gupta, B. B., Hsu, C., Gilleland, E., and Das, J.: A methodological framework for
977 extreme climate risk assessment integrating satellite and location-based data sets in intelligent
978 systems, *International Journal of Intelligent Systems*, **37**, 10268–10288,
979 <https://doi.org/10.1002/int.22475>, 2022.

980 Jha, S., Gudmundsson, L., and Seneviratne, S. I.: Partitioning the uncertainties in compound hot and
981 dry precipitation, soil moisture, and runoff extremes projections in CMIP6, *Earths Future*, **11**,
982 e2022EF003315, <https://doi.org/10.1029/2022EF003315>, 2023.

983 Kay, A. L., Griffin, A., Rudd, A. C., Chapman, R. M., Bell, V. A., and Arnell, N. W.: Climate change
984 effects on indicators of high and low river flow across Great Britain, *Adv. Water Resour.*, **151**,
985 103909, <https://doi.org/10.1016/j.advwatres.2021.103909>, 2021.

986 Kay, G., Dunstone, N., Smith, D., Dunbar, T., Eade, R., and Scaife, A.: Current likelihood and dynamics
987 of hot summers in the UK, *Environmental Research Letters*, **15**, 094099,
988 <https://doi.org/10.1088/1748-9326/abab32>, 2020.

989 Kendon, M., Doherty, A., Hollis, D., Carlisle, E., Packman, S., McCarthy, M., Jevrejeva, S., Matthews,
990 A., Williams, J., and Garforth, J.: State of the UK Climate 2023, *International Journal of Climatology*,
991 **44**, 1–117, <https://doi.org/https://doi.org/10.1002/joc.8553>, 2024.

992 Kuana, L. A., Almeida, A. S., Mercuri, E. G. F., and Noc, S. M.: Regionalization of GR4J model
993 parameters for river flow prediction in Paraná, Brazil, *Hydrol. Earth Syst. Sci.*, **28**, 3367–3390,
994 <https://doi.org/10.5194/hess-28-3367-2024>, 2024.

995 Lane, R. A., Coxon, G., Freer, J., Seibert, J., and Wagener, T.: A large sample investigation into
996 uncertain climate change impacts on high flows across Great Britain, *Hydrol. Earth Syst. Sci.*, **26**,
997 5535–5554, <https://doi.org/https://doi.org/10.5194/hess-26-5535-2022>, 2022.

998 Leng, J., Ma, K., Gu, S., Zhang, K., and He, D.: A non-stationary impact quant framework for assessing
999 the human activity impacts on hydrological drought in the Upper Red River, *Atmos. Res.*, **304**,
1000 107419, <https://doi.org/10.1016/j.atmosres.2024.107419>, 2024.

1001 Liu, R., Yin, J., Slater, L., Kang, S., Yang, Y., Liu, P., Guo, J., Gu, X., Zhang, X., and Volchak, A.: Machine-
1002 learning-constrained projection of bivariate hydrological drought magnitudes and socioeconomic
1003 risks over China, *Hydrol. Earth Syst. Sci.*, **28**, 3305–3326, [https://doi.org/10.5194/hess-28-3305-](https://doi.org/10.5194/hess-28-3305-2024)
1004 2024, 2024.

1005 Van Loon, A. F. and Van Lanen, H.: A process-based typology of hydrological drought, *Hydrology and
1006 Earth System Sciences*, **16**, 1915–1946, 2012.

1007 Lowe, J. A., Bernie, D., Bett, P., Bricheno, L., Brown, S., Calvert, D., Clark, R., Eagle, K., Edwards, T.,
1008 and Fosser, G.: UKCP18 science overview report, Met Office Hadley Centre: Exeter, UK, 1–73, 2018.

1009 Lück, A. and Wolf, V.: Generalized method of moments for estimating parameters of stochastic
1010 reaction networks, *BMC Syst. Biol.*, **10**, 98, <https://doi.org/10.1186/s12918-016-0342-8>, 2016.

1011 Moore, R. J.: The PDM rainfall-runoff model, *Hydrol. Earth Syst. Sci.*, **11**, 483–499,
1012 <https://doi.org/10.5194/hess-11-483-2007>, 2007.

1013 Murphy, C., Wilby, R. L., Matthews, T., Horvath, C., Crampsie, A., Ludlow, F., Noone, S., Brannigan, J.,
1014 Hannaford, J., and McLeman, R.: The forgotten drought of 1765–1768: Reconstructing and re-

1015 evaluating historical droughts in the British and Irish Isles, *International Journal of Climatology*, 40,
1016 5329, <https://doi.org/10.1002/joc.6521>, 2020.

1017 Murphy, J. M., Harris, G. R., Sexton, D. M. H., Kendon, E. J., Bett, P. E., Clark, R. T., Eagle, K. E., Fosser,
1018 G., Fung, F., and Lowe, J. A.: UKCP18 land projections: science report, 2018.

1019 Ndiaye, P. M., Bodian, A., Dezetter, A., Ogilvie, A., and Goudiaby, O.: Sensitivity of global hydrological
1020 models to potential evapotranspiration estimation methods in the Senegal River Basin (West Africa),
1021 *J. Hydrol. Reg. Stud.*, 53, 101823, <https://doi.org/10.1016/j.ejrh.2024.101823>, 2024.

1022 Di Nunno, F. and Granata, F.: Analysis of trends and abrupt changes in groundwater and
1023 meteorological droughts in the United Kingdom, *J. Hydrol. (Amst.)*, 131430,
1024 <https://doi.org/10.1016/j.jhydrol.2024.131430>, 2024.

1025 OFWAT: Towards resilience: how we will embed resilience in our work,
1026 [https://www.ofwat.gov.uk/wp-](https://www.ofwat.gov.uk/wp-content/uploads/2015/07/pap_pos20151210towardsresiliencerev.pdf)
1027 [content/uploads/2015/07/pap_pos20151210towardsresiliencerev.pdf](https://www.ofwat.gov.uk/wp-content/uploads/2015/07/pap_pos20151210towardsresiliencerev.pdf), 2015.

1028 Onyutha, C.: On rigorous drought assessment using daily time scale: Non-stationary frequency
1029 analyses, revisited concepts, and a new method to yield non-parametric indices, *Hydrology*, 4, 48,
1030 <https://doi.org/10.3390/hydrology4040048>, 2017.

1031 Parry, S., Mackay, J. D., Chitson, T., Hannaford, J., Bell, V. A., Facer Childs, K., Kay, A., Lane, R.,
1032 Moore, R. J., Turner, S., and Wallbank, J.: Divergent future drought projections in UK river flows and
1033 groundwater levels, <https://doi.org/10.5194/hess-28-417-2024>, 28 February 2024a.

1034 Parry, S., Mackay, J. D., Chitson, T., Hannaford, J., Magee, E., Tanguy, M., Bell, V. A., Facer Childs, K.,
1035 Kay, A., and Lane, R.: Divergent future drought projections in UK river flows and groundwater levels,
1036 *Hydrol. Earth Syst. Sci.*, 28, 417–440, <https://doi.org/10.5194/hess-28-417-2024>, 2024b.

1037 Parvizi, S., Eslamian, S., Gheysari, M., Gohari, A., and Kopai, S. S.: Regional frequency analysis of
1038 drought severity and duration in Karkheh River Basin, Iran using univariate L moments method,
1039 *Environ. Monit. Assess.*, 194, 336, <https://doi.org/10.1007/s10661-022-09977-8>, 2022.

1040 Posada, D. and Buckley, T.: Model Selection and Model Averaging in Phylogenetics: Advantages of
1041 Akaike Information Criterion and Bayesian Approaches Over Likelihood Ratio Tests, *Syst. Biol.*, 53,
1042 793–808, <https://doi.org/10.1080/10635150490522304>, 2004.

1043 Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., Fekete, B. M.,
1044 Franssen, W., Gerten, D., and Gosling, S. N.: Hydrological droughts in the 21st century, hotspots and
1045 uncertainties from a global multimodel ensemble experiment, *Proceedings of the National Academy*
1046 *of Sciences*, 111, 3262–3267, <https://doi.org/https://doi.org/10.1073/pnas.1222473110>, 2014.

1047 Ragulina, G. and Reitan, T.: Generalized extreme value shape parameter and its nature for extreme
1048 precipitation using long time series and the Bayesian approach, *Hydrological sciences journal*, 62,
1049 863–879, <https://doi.org/http://dx.doi.org/10.1080/02626667.2016.1260134>, 2017.

1050 Reis, D. S. and Stedinger, J. R.: Bayesian MCMC flood frequency analysis with historical information,
1051 *J. Hydrol. (Amst.)*, 313, 97–116, <https://doi.org/10.1016/j.jhydrol.2005.02.028>, 2005.

1052 Reyniers, N., Osborn, T. J., Addor, N., and Darch, G.: Projected changes in droughts and extreme
1053 droughts in Great Britain strongly influenced by the choice of drought index, *Hydrology and Earth*
1054 *System Sciences Discussions*, 2022, 1–36, <https://doi.org/10.5194/hess-27-1151-2023>, 2022.

1055 Robinson, E. L., Huntingford, C., Shamsudheen, S., and Bullock, J.: CHES-SCAPE: Future projections
1056 of meteorological variables at 1 km resolution for the United Kingdom 1980–2080 derived from UK

1057 Climate Projections 2018, NERC EDS Centre for Environmental Data Analysis,
1058 <https://doi.org/http://dx.doi.org/10.5285/8194b416cbee482b89e0dfbe17c5786c>, 2022a.

1059 Robinson, E. L., Brown, M. J., Kay, A. L., Lane, R. A., Chapman, R., Bell, V. A., and Blyth, E. M.: Hydro-
1060 PE: gridded datasets of historical and future Penman-Monteith potential evaporation for the United
1061 Kingdom, *Earth System Science Data Discussions*, 2022, 1–44, 2022b.

1062 Rudd, A. C., Kay, A. L., and Bell, V. A.: National scale analysis of future river flow and soil moisture
1063 droughts: potential changes in drought characteristics, *Clim. Change*, 156, 323–340,
1064 <https://doi.org/10.1007/s10584-019-02528-0>, 2019.

1065 Salas, J. D. and Obeysekera, J.: Revisiting the Concepts of Return Period and Risk for Nonstationary
1066 Hydrologic Extreme Events, *J. Hydrol. Eng.*, 19, 554–568, [https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000820](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000820), 2014.

1068 Saraiidis, G., Vasiliades, L., and Loukas, A.: Analysis of streamflow droughts using fixed and variable
1069 thresholds, *Hydrol. Process.*, 33, 414–431, 2019.

1070 Sarhadi, A., Burn, D. H., Concepción Ausín, M., and Wiper, M. P.: Time-varying nonstationary
1071 multivariate risk analysis using a dynamic Bayesian copula, *Water Resour. Res.*, 52, 2327–2349, 2016.

1072 Seneviratne, S. and Hauser, M.: Regional climate sensitivity of climate extremes in CMIP6 versus
1073 CMIP5 multimodel ensembles, *Earths Future*, 8, e2019EF001474, 2020.

1074 Sharma, S. and Mujumdar, P. P.: Modeling concurrent hydroclimatic extremes with parametric
1075 multivariate extreme value models, *Water Resour. Res.*, 58, e2021WR031519, 2022.

1076 Stahl, K., Vidal, J. P., Hannaford, J., Tjeldeman, E., Laaha, G., Gauster, T., and Tallaksen, L. M.: The
1077 challenges of hydrological drought definition, quantification and communication: an interdisciplinary
1078 perspective, *Proceedings of the International Association of Hydrological Sciences*, 383, 291–295,
1079 2020.

1080 Strupczewski, W. G., Singh, V. P., and Feluch, W.: Non-stationary approach to at-site flood frequency
1081 modelling I. Maximum likelihood estimation, *J. Hydrol. (Amst.)*, 248, 123–142, 2001.

1082 Tallaksen, L. M. and Van Lanen, H. A. J.: Hydrological drought: processes and estimation methods for
1083 streamflow and groundwater, 2023.

1084 Tanguy, M., Magee, E., Hannaford, J., Bell, V., Rameshwaran, P., Baron, H., Keller, V., and Barker, L.:
1085 CS-NOW-D2 Task 5: Analysis of future scenarios,
1086 <https://assets.publishing.service.gov.uk/media/672b4fbbabb279b2de1e8c34/cs-now-d2-future-water-resources-output-analysis.pdf>, 2023a.

1088 Tanguy, M., Chevuturi, A., Marchant, B. P., Mackay, J. D., Parry, S., and Hannaford, J.: How will
1089 climate change affect the spatial coherence of streamflow and groundwater droughts in Great
1090 Britain?, *Environmental Research Letters*, 18, 064048, 2023b.

1091 Turner, S., Barker, L. J., Hannaford, J., Muchan, K., Parry, S., and Sefton, C.: The 2018/2019 drought in
1092 the UK: a hydrological appraisal, *Weather*, 76, 248–253, 2021.

1093 Wu, D., Chen, J., Xiong, L., Lee, J. H., Kim, J. S., and Moon, H. T.: Assessing global drought conditions
1094 under climate change: A comparison of stationary and non-stationary approaches and identification
1095 of hotspot regions, *J. Hydrol. (Amst.)*, 131663, 2024.

1096 ~~Yilmaz, A. G. and Perera, B. J. C.: Extreme Rainfall Nonstationarity Investigation and Intensity-~~
1097 ~~Frequency-Duration Relationship, J. Hydrol. Eng., 19, 1160-1172,~~
1098 ~~[https://doi.org/10.1061/\(ASCE\)HE.1943-5584.0000878](https://doi.org/10.1061/(ASCE)HE.1943-5584.0000878), 2014.~~
1099 -