

Dynamic Thinning and Grounding Line Retreat in Porpoise Bay, Wilkes Land, East Antarctica

Matilda Weatherley¹, Chris R. Stokes¹, Stewart S. R. Jamieson¹, Sindhu Ramanath², Alessandro Silvano³

¹Department of Geography, Durham University, Durham, DH1 3LE, UK

5 ²Remote Sensing Technology Institute, German Aerospace Centre, 82234 Weßling, Germany

³School of Ocean and Earth Science, University of Southampton, Southampton, SO14 3ZH, UK

Correspondence to: Matilda Weatherley (matilda.weatherley@durham.ac.uk)

Abstract. The East Antarctic Ice Sheet (EAIS) is often considered less vulnerable to climate change than the West Antarctic or Greenland ice sheets, but some regions of the EAIS have been losing mass over recent decades. In particular, mass loss in
10 Wilkes Land, which overlies the Aurora Subglacial Basin, may have accelerated over the past two decades. However, whilst several large outlet glaciers drain this region, few have been studied in detail. Here, we present new data on the recent ice dynamics of four outlet glaciers that drain into Porpoise Bay, Wilkes Land, which includes Holmes East, Holmes West, and Frost glaciers. We use optical satellite imagery, differential synthetic aperture radar interferometry, and a range of previously published datasets to describe changes in the ice-shelf front, grounding line position, ice surface velocity, and ice surface
15 elevation over the last three decades. Our results reveal evidence that is consistent with dynamic changes in the region, characterised by thinning of grounded ice and grounding line retreat, albeit with large uncertainties. We find indications of Circumpolar Deep Water proximal to the continental shelf break that could access the glaciers through deep cross-shelf troughs, which is consistent with previous estimates of high rates of basal melting beneath their floating tongues/ice shelves. In addition, our results support previous observations of near-synchronous ice-shelf calving across Porpoise Bay's ice
20 shelves/tongues, following the break-out of multi-year sea ice, together with a previously unidentified calving event in 2021/22. Taken together these findings highlight the potential vulnerability of this region of East Antarctica to ongoing and future changes in ocean and sea-ice conditions.

1 Introduction

The Antarctic Ice Sheet is the largest ice sheet worldwide, containing 57.9 m of global mean sea level equivalent (SLE)
25 (Morlighem et al., 2020). It comprises three main regions: the East Antarctic Ice Sheet (EAIS), the West Antarctic Ice Sheet (WAIS), and the Antarctic Peninsula. Historically, most work has focussed on changes in the Antarctic Peninsula (van den Broeke, 2005; Berthier et al., 2012; Cook et al., 2014, 2016; Leeson et al., 2020) and pronounced mass loss in the WAIS (Pritchard et al., 2009; Mouginot et al., 2014; Rignot et al., 2014; McMillan et al., 2015; Feldmann and Levermann, 2015; Pattyn and Morlighem, 2020), which contain an estimated 0.3 m and 5.3 m SLE, respectively (Morlighem et al., 2020). In

30 contrast, much less work has focused on the stability of the EAIS, containing 52.2 m SLE, an order of magnitude larger than for the WAIS (Morlighem et al., 2020; Stokes et al., 2022).

Recent mass balance studies show accelerating mass loss from the Antarctic Ice Sheet over recent decades, with the most recent reconciled estimate reporting 2671 ± 530 Gt of ice lost between 1992 and 2020 and equating to 7.4 ± 1.5 mm of sea-level rise (Otosaka et al., 2023). This mass loss is dominated by the WAIS (82 ± 9 Gt yr^{-1}) and the Antarctic Peninsula
35 (13 ± 5 Gt yr^{-1}), with the EAIS remaining close to balance (3 ± 25 Gt yr^{-1}), albeit with much larger uncertainties than for the WAIS or Antarctic Peninsula (Stokes et al., 2022; Otosaka et al., 2023). Mass loss from the WAIS is largely concentrated in the Amundsen Sea Embayment, primarily from Pine Island and Thwaites glaciers (Mouginot et al., 2014; Gardner et al., 2018; Shepherd et al., 2019). Here, mass loss has been linked to ice flow acceleration, thinning, and grounding line retreat, which is driven by ice-shelf debuttressing due to warm ocean water melting the base of ice shelves causing thinning (Mouginot et al.,
40 2014; Rignot et al., 2014; Fürst et al., 2016; Pattyn and Morlighem, 2020), in addition to iceberg calving losses (Greene et al., 2022; Davidson et al., 2023).

Comparatively, the EAIS has often been considered less vulnerable to recent climate change, with the latest IMBIE mass balance estimates revealing it is largely in balance or slightly increasing in mass (Otosaka et al., 2023). It has been noted that mass gain is generally concentrated in Dronning Maud Land and mass loss in Wilkes Land, with other areas largely close
45 to balance (Gardner et al., 2018; Smith et al., 2020a; Otosaka et al., 2023). Of note is that there is emerging evidence of a longer-term trend of regional mass loss from Wilkes Land (Rignot et al., 2019; Smith et al., 2020a; Wang et al., 2021; Stokes et al., 2022), which is often referred to as the ‘weak underbelly’ of the EAIS (Miles et al., 2016; Pelle et al., 2020). Recent work has also documented the collapse of the Conger-Glenzer Ice Shelf in March 2022, although the wider significance of this event is unclear (Walker et al., 2024).

50 Wilkes Land is drained by 39 outlet glaciers and has received growing attention in the past three decades due to the emerging signal of mass loss (Gardner et al., 2018; Rignot et al., 2019; Schröder et al., 2019a; Shepherd et al., 2019; Smith et al., 2020a; Nilsson et al., 2022; Thompson et al., 2023; Stokes et al., 2022, 2025). Similar to the WAIS, mass loss has been linked to the intrusion of warm modified Circumpolar Deep Water (mCDW) in sub-ice cavities through deep cross-shelf troughs, driving melt rates that are sometimes comparable to the Amundsen Sea Embayment (Rintoul et al., 2016; Silvano et al., 2016, Miles et al., 2016; Rignot et al., 2019). Indeed, observations have confirmed mCDW proximal to three key glaciers
55 in Wilkes Land: Vanderford, Totten, and Moscow University glaciers (Greenbaum et al., 2015; Rintoul et al., 2016; Silvano et al., 2017; Picton et al., 2023).

In a study of the cumulative mass loss from each drainage basin in East Antarctica between 1979 and 2017, Rignot et al. (2019) estimated that Totten (236 Gt) and Denman (191 Gt) glaciers showed the highest rates of mass loss, but that this
60 was followed by Frost (159 Gt) and Holmes (152 Gt) glaciers, which both drain into Porpoise Bay, Wilkes Land. However, despite Frost and Holmes glaciers losing mass, there have been few studies of these major outlet glaciers. Furthermore, Wilkes Land is underlain by a bed that lies below sea level and deepens inland to the Aurora Subglacial Basin, containing several deep

troughs (Morlighem, 2022; Pritchard et al., 2024), and making it potentially vulnerable to marine ice sheet instability (MISI), which may have occurred during past warm periods (Cook et al., 2013; Stokes et al., 2022).

65 1.1 Porpoise Bay

Porpoise Bay (76° S, 128° E) is a 150 km wide bay that lies east of Moscow University Glacier (~300 km) and Totten Glacier (~550 km) (Fig. 1a). It overlies the eastern Aurora Subglacial Basin and is drained by several marine-terminating outlet glaciers, of which the largest are Frost (0.84 m SLE) and Holmes (0.11 m SLE) (Rignot et al., 2019). Holmes Glacier has two distinct outlets that feed the same ice shelf (Fig. 1b). Following Miles et al. (2017), and to enable comparison with that study, we refer to these two outlets as Holmes East and Holmes West glaciers, although Holmes East is sometimes referred to as De Haven glacier (Rignot et al., 2026). A further enhanced area of flow lies between Frost and Holmes East glaciers and is referred to here as Glacier 1 (Fig. 1b).

Recent studies suggest a mass imbalance from some of the marine-based glaciers entering Porpoise Bay, including a potential combined sea level contribution of 0.8 mm from Holmes and Frost catchments between 1979 and 2017 (Rignot et al., 2019). In addition, there are reports of surface elevation changes between -0.07 m yr^{-1} and -0.87 m yr^{-1} (2003–2019) (Smith et al., 2020a), ice-shelf retreat linked to sea-ice break-out events (Miles et al., 2017), ice-shelf thinning (Smith et al., 2020a; Miles and Bingham, 2024), grounding line retreat (Konrad et al., 2018; Rignot et al., 2019; 2026), and ice flow speed up (Rignot et al., 2022). Despite these ice sheet-wide studies suggesting that dynamic change may be occurring, Porpoise Bay's glaciers have not been studied in detail.

This paper aims to enhance our understanding of the recent changes in the ice dynamics of the outlet glaciers in Porpoise Bay, East Antarctica. Following similar work undertaken in Vincennes Bay (Picton et al., 2023), also in Wilkes Land, we assessed four key parameters of the outlet glaciers: (1) ice-shelf frontal position, delineated from satellite imagery, (2) ice surface elevation, extracted from datasets from Schröder et al. (2019b), Smith et al. (2020b), and Nilsson et al. (2023), (3) ice surface velocity, extracted from ITS_LIVE (Gardner et al., 2025), (4) grounding line position, assessed using differential satellite synthetic aperture radar interferometry (DInSAR) and other published datasets (Bindenschadler and Choi, 2011; Rignot et al., 2016; Haran et al., 2018, 2021a, 2021b). The primary aim of the study is to examine the potential evidence for dynamic changes and then compare any such evidence to secondary datasets on ocean temperature data from EN4 (Good et al., 2013), MEOP casts (Treasure et al., 2017), Argo floats (Wong et al., 2020), and sea-ice concentrations from NSIDC (Fetterer et al., 2017), alongside bathymetry from Bedmap3 (Pritchard et al., 2024), BedMachine3 (Morlighem, 2022) and ANTGG (Rignot et al., 2024). A secondary aim is to examine the potential influence of sea-ice concentrations on major calving events, which have been known to occur in Porpoise Bay and which have been linked to major sea-ice break-out events here (Miles et al., 2017) and elsewhere in East Antarctica (e.g., Greene et al., 2018; Gomez-Fell et al., 2022).




Figure 1: (a) Ice sheet bed elevation and bathymetry of Porpoise Bay and the surrounding Wilkes land from Bedmap3 (Pritchard et al., 2024). (b) Ice velocity map of Porpoise Bay extracted from the 2021 ITS_LIVE ice velocity mosaic (Gardner et al., 2025). Central flowlines and sampling boxes are shown across the inland (IN), grounding line (GL), and floating ice tongue (FT) at each of the studied glaciers: Frost (FR), Glacier 1 (G1), Holmes East (HE, also known as De Haven Glacier), and Holmes West (HW). Black lines show the 1996 MEaSUREs grounding line (Rignot et al., 2016).

2 Methodology

The recent ice dynamics of Porpoise Bay's outlet glaciers were examined by combining a range of remote sensing approaches to measure four key glacier parameters (ice-shelf frontal position, ice surface elevation, ice surface velocity, and grounding line position). This depicts the scale of change and enables assessment of the potential driving mechanisms.





2.1 Ice-shelf frontal position

The Google Earth Engine Digitisation Tool (GEEDiT) was used to manually map the calving front of each outlet glacier following the criteria outlined in Table 1. GEEDiT uses imagery from Landsat 4, 5, 7, 8, and 9, ASTER, and Sentinel 1 and 2,

spanning from 1982 to the present day (Lea, 2018). In addition, we used co-registered and orthorectified Landsat 1 (Miles and Bingham, 2023) and declassified ARGON (Kim et al., 2007) imagery to extend our measurements back to 1963. Images were preferentially selected from austral summertime (December to February) to minimise complications with digitising sea ice. Cloud cover thresholds were not applied to avoid unnecessarily omitting images with visible ice-shelf edges. Instead, the images were manually checked to ensure the calving front was visible. The failure of the Scan Line Corrector onboard the Landsat 7 satellite resulted in striped data losses (Paul et al., 2017), but parts of the glaciers were still visible. In such images, the ice-shelf frontal position was, where possible, digitised across gaps, using an adjacent image with a differently striped pattern of data losses to inform digitisation (Black and Joughin, 2022). The spatial resolution of imagery ranged from 140 m (Landsat 1) to 10 m (Sentinel 2).

115

Table 1: Ice-shelf features and identification criteria adapted from Holt et al. (2013) and Arthur et al. (2021). Examples of ice-shelf features from Sentinel 2 are provided.

Feature	Example	Structure	Identification on satellite imagery
Ice calving front (ice-free conditions)			Abrupt transition from ice-shelf calving margin (bright feature) to open ocean (dark feature).
Ice calving front (mélange presence)			Transition from ice shelf (smooth feature) to mélange (tightly packed calved ice blocks with other calved materials); transition often marked by shadow.
Ice calving front (sea ice presence)			Sharp transition from ice shelf ('thick' feature) to sea ice (darker than glacial ice, 'flat' feature); transition often marked by shadow.
Calved ice blocks/icebergs			Distinct blocks detached from ice-shelf edge and surrounded by mélange or open ocean
Mélange			Amalgamation of sea ice, marine ice, firn, and densely packed calved icebergs

120

The outcomes of our mapping were exported to ArcGIS. The ice-shelf frontal position change was quantified using the box method outlined by Moon and Joughin (2008), which calculates the mean frontal change using the difference in ice-shelf geometry as measured by the area within an open-ended box across the central region of ice flow. This method accounts for asymmetric changes across glacier termini. We also calculated the percentage area loss relative to the mean ice-shelf area for the key calving events.

125 The availability of cloud-free digital satellite imagery limits this method as clouds conceal the ice-shelf edge. Between
1963 and 2000, only four observations were available, but we acquired approximately one image per year post-2000. When
mapping the ice-shelf edge, digitisation errors were minimised by using the highest-resolution imagery available and applying
contrast stretching filters to improve the shelf-ice-mélange and ice-water boundary visibility; the resultant digitisation
uncertainty was estimated to be similar to Miles et al. (2021) at 1 pixel (Table S1). A particular challenge was identifying the
130 mélange from the calving front, due to the blocky nature of the mélange. To minimise the subjectivity of our mapping this
boundary, we used the feature identification criteria outlined in Table 1.

2.2 Ice surface elevation

Monthly surface elevation changes from 1992 to 2020 were extracted from the following published ice surface elevation
change datasets: Schröder et al. (2019b), Smith et al. (2020b) and Nilsson et al. (2023). The error associated with each annual
135 velocity mosaic was provided at pixel scale, representing the standard deviation of the elevation values and quantified below.
Schröder et al. (2019b) calculated the monthly surface elevation changes between 1978 and 2017, relative to September 2010,
at a horizontal resolution of 10 km. The associated monthly uncertainties range from ± 0.3 to ± 9.7 m yr⁻¹ (Schröder et al.,
2019b). Schröder et al. (2019b) used altimetry data from Seasat, Geosat, ERS-1, ERS-2, Envisat, ICESat, and CryoSat-2
satellite missions. Ice surface elevation measurements include changes in surface mass balance and firn compactions rates
140 (Schröder et al., 2019b)

Nilsson et al. (2023) calculated monthly surface elevation changes between 1985 and 2020, relative to December
2013, at a horizontal resolution of 1920 m. The monthly error ranges from 0.06 to 3.8 m yr⁻¹ (Nilsson et al., 2023). Nilsson et
al. (2023) included additional data from the ICESat-2 mission, which uses a novel six-laser arrangement and is meshed with
previous data to extend the measurements to 2020.

145 The mean rates of elevation change from Schröder et al. (2019b) and Nilsson et al. (2023) were extracted from two
10 km² sampling boxes that were drawn to capture changes at key points along flow: at the grounding line (GL) and 10 km
inland (IN) (Fig. 1b). Since the boxes spanned multiple grid cells, we calculated a mean value for each box. Sampling from
two boxes was designed to facilitate the identification of any spatial patterns of dynamic thinning (e.g., propagation of thinning
from the glacier front to the inland portion of the glacier). The measurements were calculated relative to the earliest shared
150 time period across both datasets (September 1992), to enable direct comparison between them. Similar to Picton et al. (2022),
we calculated the elevation anomalies relative to the 1992–2017 mean and produced a 24-month moving average for both
datasets. Note that the gridded dataset from Schröder et al. (2019b) do not cover the GL box at Holmes West Glacier.

Smith et al. (2020b) use altimetry data from ICESat and ICESat-2 satellite missions, derived from the ATL 15 gridded
product (corrected with firn-models and tides) to calculate the rate of surface elevation change between 2003 and 2019, at a
155 horizontal resolution of 5 km. The dataset quantifies the root mean square error in each grid cell, which stems from instrument
precision, ground finding errors (errors in measuring the exact distance between the satellite and the ground), and interpolation
errors; the error from the sampling boxes ranges from 0.004 to 0.08 m. To enable comparison across the three datasets, mean

rates of elevation change were calculated for Nilsson et al. (2023) between 2003 and 2019 and for Schröder et al. (2019b) between 2003 and 2017 to match the temporal range in Smith et al. (2020b).

160 2.3 Ice surface velocities

The Making Earth Science Data Records for Use in Research Environments (MEaSUREs) ITS_LIVE product was used to extract annual ice surface velocity between 2000 and 2022 (Gardner et al., 2025). The ITS_LIVE mosaics were derived from Landsat 4, 5, 7, and 8 imagery using autonomous Repeat Image Feature Tracking (auto-RIFT) algorithms, an open-source Python module that analyses pixel displacement between two images to calculate ice velocity (Gardner et al., 2018).

165 The mean annual velocities were extracted from the GL and IN sampling boxes alongside a box positioned on the floating ice tongue (FT) (Fig. 1b). Sampling from multiple boxes depicts the spatial variability in ice velocity over time to reveal any pattern or propagation of dynamic change up-ice. For each outlet glacier, the central flowline was extracted from the 2021 annual velocity mosaic and used to position the sampling boxes. The 2021 mosaic was selected due to a relatively lower indicated error (Table S2).

170 The error associated with each annual velocity mosaic was provided at pixel scale, representing the standard deviation of the difference between the image-pair component velocities and the mean annual component velocities (Gardner et al., 2018). The mean annual velocity error was extracted from each FT, GL, and IN box; the error ranged from 0.7 to 412.8 m yr⁻¹ (see Table S2). Following the method in Picton et al. (2022), the annual velocity data were disregarded for mean errors over 50 % of the velocity magnitude. Most of the errors were below 5–10 % but this resulted in the loss of around 1 % of available
175 data.

Data scarcity is a limiting factor of the early ITS_LIVE product as the auto-RIFT processing chain is limited by the availability of image pairs across a given year. Incomplete coverage was seen across Holmes East and West before the launch of Landsat 8 in 2013. The annual velocity values were disregarded if less than 25 % data coverage was observed across the sampling box, resulting in the removal of 8 % of values.

180 2.4 Grounding line position

Grounding line positions were derived using DInSAR interferograms from Sentinel 1 imagery collected between 2019 and 2021 (Table 2) to enable comparison with older, previously published datasets (described below). DInSAR interferograms were generated by differencing two interferograms, each produced from three or more consecutive repeat pass SAR acquisitions. Assuming constant velocity over the observed time period, the phase change from tidal flexure at the ice sheet-
185 ice-shelf boundary can be detected from the resulting differential interferograms. These ice-shelf flexure zones are characterised by a dense fringe in the DInSAR phase.

To automatically delineate the grounding line, a Holistically-Nested Edge Detection (HED) neural network was applied to the DInSAR interferograms (Ramanath et al., 2025). The neural network was trained on the real and imaginary interferometric features, achieving a median and mean offset of 265 m and 421 m, respectively, from manual grounding line

delineations, and a predictive uncertainty of 401 m (Ramanath et al., 2025). The uncertainty was derived from calculating the pixel-wise standard deviations of each DInSAR interferogram of an ensemble of five HED neural networks, configured with the same hyperparameters, but differing in the initialization and trained with randomly shuffled training samples (Ramanath et al., 2025). After calculating the standard deviation of the network predictions, we aggregated those pixels that were within one standard deviation to get a contiguous area and converted it into a polygon/buffer geometry, i.e., 10 km is the maximum uncertainty because that is the width of the polygon around the 2020 grounding line at Holmes East. We identified two grounding lines at Holmes East Glacier but did not extract any coherent interferograms over Glacier 1 or Frost glaciers because the ice flow was too rapid to get coherence within the 6-day repeat-pass interval of the Sentinel 1 constellation. At Holmes West Glacier, we identified a 2020 grounding line, but it looked implausible relative to the other datasets and generated a very wide uncertainty buffer from a highly de-correlated interferogram. Thus, we disregarded the line. We also note that a recent study by Rignot et al. (2026) highlighted the challenges of identifying grounding line positions in Porpoise Bay due to the intense surface weathering, short floating sections, pinning points, and the steep transition between grounded and floating ice.

The grounding line positions were also extracted from several secondary datasets in the published literature. The MEaSURES grounding line product was generated using similar DInSAR techniques applied to ERS-1 and ERS-2 imagery collected in 1996 (Rignot et al., 2016). This dataset has a standard error of ± 100 m (Rignot et al., 2016). The Antarctic Surface Accumulation and Ice Discharge (ASAID) grounding line dataset was produced by manually delineating the most seaward continuous break in slope using Landsat 7 images between 1999 and 2003 and surface elevation data from ICESat satellite mission (Bindenschadler et al., 2011). This dataset has an estimated positional error of ± 502 m for outlet glacier boundaries (Bindenschadler and Choi, 2011). The Mosaic Of Antarctica (MOA) grounding lines were also produced using manual delineation of the most seaward break in slope, observed in 2004, 2009, and 2014 (Haran et al., 2018; 2021a; 2021b). The MOA grounding line products have associated errors of ± 250 m. The change in grounding line position was measured along the central flowline from the most seaward 1996 MEaSURES position.

The DInSAR results and MEaSURES product represent the hinge line, an approximation of the grounding line position (Rignot et al., 2016). Contrastingly, the ASAID and MOA datasets represent the break-in-slope; the change in ice surface at the transition from grounding and floating ice (Bindenschadler et al., 2011). The hinge line and break-in-slope are different components of the grounding zone (the break-in-slope is observed seaward of the true grounding line, and the hinge line is landward of the true grounding line). Hence, caution must be exercised when interpreting changes in grounding line position acquired from the different methods.

Table 2: Imagery used in the DInSAR grounding line extraction.

Satellite	T1	T2	T3
Sentinel 1	03/11/2019	09/11/2019	15/11/2019
Sentinel 1	21/11/2020	27/11/2020	03/12/2020

220 **2.5 Bed and ice surface topography**

To interpret grounding line migration in the context of the wider subglacial topography, we extracted bed and ice surface elevation profiles for the Porpoise Bay outlet glaciers. Bedrock elevation profiles were derived from BedMachine3 (Morlighem, 2022) and Bedmap3 (Pritchard et al., 2024) along the digitised flowlines. Bed elevation and the associated error were sampled at 500 m intervals (the horizontal resolution of the dataset). Although there are 47,492 line-kilometres of airborne radar profiles over the Aurora Subglacial Basin (Young et al., 2011), including over Porpoise Bay, the roughness and shape of the bed are not well characterised. Although the bed topography inland of the grounding line does deepen over distances of tens of kilometres, the presence of inland pinning points that may restrict future grounding line retreat are unknown due to a lack of detailed survey. The bed elevation error associated with Bedmap3 ranges from ± 7 to ± 276 m for the grounded bed and ± 60 m to ± 306 m for the interpolated seabed (Pritchard et al., 2024). For BedMachine3, the typical uncertainties are ± 36 m in well-constrained regions (driven by radar vertical resolution); uncertainties can exceed ± 200 m where data are sparse due to reliance on interpolations and can exceed ± 500 m beneath floating ice where sub-ice bathymetry data are lacking (Morlighem, 2022).

Bed elevation profiles along the marine area in front of the glaciers were digitised from the AntGG2021 dataset which is based on a 3D inversion of a circumpolar compilation of gravity anomalies constrained by measurement from BedMachine3, the International Bathymetric Chart of the Southern Ocean, and discrete seafloor measurements from seismic and ocean robotic probes (Rignot et al., 2024; Charrassin et al., 2025). This dataset provides insight into the seafloor elevation of ice-shelf cavities and continental shelf. The potential error of the estimated bathymetry for the Sabrina region, where Porpoise Bay is situated, is 185 m (Charrassin et al., 2025).

Surface topography profiles were extracted along each central flowline from the Reference Elevation Model of Antarctica (REMA) mosaic (Howat et al., 2022). The surface elevation product from REMA is provided at 100 m resolution with errors of less than 1 m (Howat et al., 2019). These were sampled at 500 m intervals to match the bedrock elevation. The ice-shelf base elevation was calculated by subtracting the Bedmap3 ice thickness from the Bedmap3 ice surface elevation (Pritchard et al., 2024). The uncertainty of the Bedmap3 ice thickness dataset ranges between ± 0 m for rock cells, ± 10 m for ice shelves, ± 10 – 20 m in surveyed cells, and ± 73 – 272 m in interpolated or poorly constrained cells (Pritchard et al., 2025). The Bedmap3 ice surface elevation error is, on average, ± 7 m (Pritchard et al., 2025).

2.6 Ocean and sea-ice conditions

We analyse ocean temperature and sea-ice conditions to understand their potential influence on any observed changes in ice dynamics. Due to an absence of direct, observational ocean temperature data across the continental shelf in the Wilkes Land, we used EN4.2.2 subsurface ocean temperature objective analysis data (Good et al., 2013), accessed from the UK Met Office. EN4 uses profile measurements of ocean temperature and salinity over time at point locations (e.g., from Argo programme, World Ocean Database, seal-based CTDs), applying quality control and bias corrections, to produce monthly gridded fields

($1^\circ \times 1^\circ$, 42 depth levels) of ocean temperature and salinity (Good et al., 2013; Sohail et al., 2026). We extracted the mean monthly temperature from four cells between 65° and 66°S , 127° and 128°E , located on the continental shelf proximal to Porpoise Bay (Fig. 1a), for each month between January 1990 and March 2025 at 24 depths. Due to the paucity of direct observation data from our study area, we acknowledge that the EN4 dataset has large uncertainties, but these data provide the only consistently derived indication of ocean temperature through time.

To scrutinise the observational data in more detail we also plotted empirical observations from the region. Most of the data are from marine mammals between 2004 and 2021, available from the Marine Mammal Exploring the Oceans Pole to Pole (MEOP) Consortium (Treasure et al., 2017). Each mammal is equipped with a Conductivity Temperature Depth Satellite Relay Data Logger (CTD-SRDL) and deployed for 72 to 136 days (Roquet et al., 2014; Treasure et al., 2017). Data are collected throughout a dive and once the mammal surfaces the data are telemetered through the Argo satellite system. The geolocation of each dive is extracted by the Argo satellite triangulation, and collected with an accuracy of about 4–5 km. The data are not uniform in space and time (i.e., there are numerous casts near the edge of the continental shelf but none within 130 km of the Porpoise Bay ice margin). This is supplemented by information from the Argo Program (Wong et al., 2020) that uses autonomous profiling floats that collect ocean temperature and salinity measurements across the upper 2 km of the ocean.

Sea-ice conditions in Porpoise Bay were analysed using the NSIDC Sea-ice Index, Version 3 (Fetterer et al., 2017). The data are available at 25 km spatial resolution. We extracted the monthly sea-ice concentrations from a $12,500 \text{ km}^2$ polygon between November 1979 and April 2025 (Fig. 1a). We recognise this region may include ice mélange, shelf ice, and grounded icebergs, so is likely to capture mélange and ice-shelf breakup as well as sea-ice concentration changes. The data are obtained from satellite passive micro-wave derived datasets (the Near-Real-Time DMSP SSMIS Daily Polar Gridded Sea-ice Concentrations and the Sea-ice Concentrations from Nimbus-7 SMMR and DMSP SMM/I-SSMIS Passive Microwave Data). The accuracy of the sea-ice concentration is usually cited within $\pm 5\%$ in winter and $\pm 15\%$ in summer (Fetterer et al., 2017). The error is reduced for thicker and higher sea-ice concentrations due to the reduced influence of open water or high cloud liquid content on the measured brightness temperature (Cavalieri, 1995; Spreen et al., 2008; Fetterer et al., 2017).

275 **3 Results**

3.1 Ice-shelf frontal position

The results show no overall trend of advance or retreat but there are several major calving events, followed by advance, (Fig. 2 and 3) at Frost, Glacier 1, Holmes East, and Holmes West, which fluctuated between 5.7 km and 11.3 km relative to the 1963 position, over the measurement period (1963–2025). All four glaciers advanced between 1963 and 1973, at a mean rate of 281 m yr^{-1} , and retreated between 1973 and 1991, at a mean rate of 386 m yr^{-1} , with Glacier 1 and Holmes East Glacier retreating by over $\sim 9 \text{ km}$. Contrastingly, 1991 to 2001 saw an advance from Frost Glacier (177 m yr^{-1}), Glacier 1 (329 m yr^{-1}), and Holmes East Glacier (545 m yr^{-1}) and a retreat from and Holmes West Glacier (52 m yr^{-1}). However, the lack of imagery before 2000 is likely to mask smaller-scale periods of advance and retreat, so these trends are evaluated with caution. We

observed the ice shelves advancing to a similar maximum position within 2 km of the previous advance limit (Fig. 2), before rapidly calving and beginning to advance once more.

The ice shelves in Porpoise Bay underwent almost synchronous cycles of advance and retreat between 2006 and 2025 (Fig. 3). Initially, the glaciers advanced from a post-calving position (having undergone a Bay-wide calving event between December 2005 and January 2007): extending 5.4 km at Frost Glacier (2008–2016), 6.2 km at Glacier 1 (2007–2015), 2.7 km at Holmes East Glacier (2006–2016) (underwent another calving event between 2010 and 2011), and 6.8 km at Holmes West Glacier (2006–2016). Between February 2016 and March 2017, the outlet glaciers underwent a calving event, moving −2.8 km at Frost Glacier, −8 km at Glacier 1, −0.9 km at Holmes East Glacier, and −4.5 km at Holmes West Glacier. Subsequently, Frost, Glacier 1, Holmes East, and West glacier gradually advanced 0.6 km, 3.7 km, 3.1 km, and 4.2 km, respectively, between 2017 and 2021. The ice shelves underwent a subsequent calving event between January 2021 and March 2022, with Glacier 1, Holmes East, and Holmes West glaciers moving −2.7 km, −7 km, and −5.4 km, respectively, with Frost Glacier moving −3.6 km between 2022 and 2023. Finally, the glaciers advanced 2.4 km (Frost), 1.2 km (Glacier 1), 3.7 km (Holmes East), and 0.6 km (Holmes West) by 2025.




Figure 2: (a) Changes in ice front position across Porpoise Bay. Panels b-e show a selection of minimum and maximum annual ice shelf frontal positions for each glacier: (b) Holmes East Glacier, (c) Frost Glacier, (d) Glacier 1, and (e) Holmes West Glacier. The background satellite imagery is a Landsat 9 image from 01/02/2024 and was downloaded from USGS EarthExplorer.




Figure 3: Width-averaged ice-shelf calving frontal position changes relative to the first measurement in 1963 at (a) Frost Glacier, (b) Glacier 1, (c) Holmes East Glacier, and (d) Holmes West Glacier. Frontal positions before 2000 are joined by dashed lines due to large gaps in satellite imagery availability. Vertical dashed lines display the 2007 and 2016 sea-ice break-out event from Miles et al. (2017). Grey lines show the timing of major calving events.

3.2 Ice surface velocity

3.2.1 Temporal variability

Moderate variations in ice surface velocity were observed between 2000 and 2022 (Fig. 4). All three glaciers saw an increase in ice flow velocity over the study period across parts of the glacier flowlines. Observations showed the greatest speed up at Holmes East Glacier (Fig. 4c), with ice surface velocity at the IN box increasing 32 % overall from 382.3 ± 37 in 2013 to 504 ± 121 m yr⁻¹ in 2022. However, the absolute value of velocity increase (122 m yr⁻¹) is similar to the associated error (see Table S2). We find significant ice surface velocity variations in the GL and IN boxes at Holmes East and West glaciers, with a period of slowdown measured between 2017 and 2020, followed by a subsequent speed-up. Interestingly, this variation in ice surface velocity was not translated to the floating tongue for either glacier. Glacier 1 increased to peak velocity in 2010 before slowing down to a lower constant velocity (Fig. 4b). The ice surface velocity at Frost Glacier consistently increased by 7–10 % over the study period (Fig. 4a).

3.2.1 Velocity and ice-shelf frontal position

Some of the outlet glaciers in Porpoise Bay displayed coincidental timing between changes in ice-shelf frontal position and velocity (Fig. 4, grey). Frost Ice Shelf underwent a calving event between 2007 and 2008, losing over 50 % ice-shelf area, after which the velocity increased by 2–6 % over the following two years across the glacier. Subsequently, the ice flow velocity increased following the small retreat between 2013 and 2014 and the ice-shelf calving between 2016 and 2019, increasing 3 % (IN), 10 % (GL), and 9 % (FT) over 8 years (Fig. 4a). The velocity across Glacier 1 increased 18 % (IN), 13 % (GL), and 9 % (FT) in the two years after minor calving in 2007, followed by a velocity change of -24 % (IN), -7 % (GL), and -10 % (FT) over the three years after 2010, as the ice shelf regrew (Fig. 4b). However, we do not record a velocity response to the latter two major calving events from Glacier 1 (Table S2).

Following the 2016 calving event, Holmes East Glacier displayed an increase in velocity of 13 % (IN), 10 % (GL), and 5 % (FT) between 2016 and 2017 (Fig. 4c). The glacier showed limited velocity response to the 2021 calving event; the velocity continued its upward trend. Holmes West Ice Shelf underwent a calving event between 2007 and 2008 that reduce the ice shelf area by over 50 %, during which a 6 % increase in velocity was recorded across the grounding line (Fig. 4d). The subsequent 2016–2017 calving event is coincidental with a 3 % increase in velocity across the floating tongue over 1 year, while the grounding line and inland sections decrease in velocity to a minimum in 2020 (Table S2). Conversely, the subsequent 2021 calving event, resulting significant ice-shelf area loss, was followed by a 5 % reduction in ice velocity across the floating tongue and further inland. Overall, the outlet glaciers display some changes in velocity that correspond to changes at the ice-shelf edge, but we do not record a consistent response to every calving event.




Figure 4: Mean annual velocity extracted from the inland (IN), grounding line (GL), and floating ice tongue (FT) boxes across (a) Frost Glacier, (b) Glacier 1, (c) Holmes East Glacier, and (d) Holmes West Glacier. Velocity data extracted from the ITS_LIVE velocity mosaic between 2000 and 2022 (Gardner et al., 2025). Dashed lines show the linear trend. Grey lines show the timing of major calving events at the glacier ice shelf; grey dashed lines depict a partial/minor calving event. The y-axis scales are different across Figure 4. For further information on velocity errors see Table S2.

3.3 Ice surface elevation change

There was general agreement between the two ice surface elevation datasets provided by Nilsson et al. (2023) and Schröder et al. (2019b) across the GL and IN boxes, especially after *ca.* 2003, when the uncertainties are lower. Indeed, each data point is associated with high levels of uncertainty, especially in the 1990s and early 2000s (see Section 2.2) and there is a greater spread of data points in this earlier period. Across the IN boxes, a clear and consistent trend of thinning was observed at Holmes East and West across the spread of individual data points (Fig. 5). Between 2003 and 2017, Schröder et al. (2019b) and Nilsson et al. (2023) observed elevation change at an average rate of -0.41 and -0.35 m yr^{-1} for Holmes East Glacier and -0.04 and -0.34 m yr^{-1} for Holmes West Glacier (values relative to 09/1992). Contrastingly, Frost Glacier and Glacier 1 displayed very little change in ice surface elevation and any minor changes are well inside the uncertainty indicated for this dataset (see Section 2.2).

Mirroring the elevation patterns inland, observations of surface elevation change across the GL boxes show moderate thinning at Holmes East Glacier and more rapid thinning at Holmes West Glacier (Fig. 6). Schröder et al. (2019b) and Nilsson et al. (2023) observed elevation change at an average rate of -0.41 and -0.40 m yr^{-1} for Holmes East Glacier and Nilsson et al. (2023) observed average elevation change at a rate of -1.22 m yr^{-1} for Holmes West Glacier between 2003 and 2017. Lower rates of surface elevation loss were similarly observed at Frost Glacier and Glacier 1.




Figure 5: Monthly surface elevation anomalies observed in each IN box at (a) Frost Glacier, (b) Glacier 1, (c) Holmes East Glacier, and (d) Holmes West Glacier between 1992 and 2020. Elevation anomalies are calculated relative to the 1992–2017 mean. Bold lines display the 24-month rolling mean.




Figure 6: Monthly surface elevation anomalies observed in each GL box at (a) Frost Glacier, (b) Glacier 1, (c) Holmes East Glacier, and (d) Holmes West Glacier between 1992 and 2020. Elevation anomalies are calculated relative to the 1992–2017 mean. Bold lines display the 24-month rolling mean.

Similar trends are observed by Smith et al. (2020b) using ICESat- and ICESat-2-derived data (Fig. 7). Porpoise Bay showed an overall decrease in ice surface elevation between 2003 and 2019, with a hotspot of thinning extending across the ice shelf and grounded ice around Holmes East and West glaciers. We observed a maximum surface elevation change of -1.11 m yr^{-1} on the grounded ice and -3.07 m yr^{-1} on the floating ice shelf.




Figure 7: Rate of ice-surface elevation change observed inland of Porpoise Bay between 2003 and 2019, calculated by Smith et al. (2020b). Dashed black lines depict the glacier flowlines.

3.4 Grounding line position and bathymetry

Notwithstanding some very large uncertainties (see Section 4.1), the highest grounding line retreat was observed at Frost and Holmes East glaciers (Fig. 8 and 9). Frost Glacier retreated 17.17 km between 1996 and 2014, at an average rate of -0.95 km yr^{-1} (Fig. 8a). This grounding line retreat occurred largely between 1996 and 2004, at an average rate of -2.14 km yr^{-1} , after which the grounding line was observed at a constant position. The grounding line retreat occurred across a largely retrograde slope, dropping in elevation from -615 m (Bedmap3)/ -587 m (BedMachine3)/ -758 m (ANTGG2021) to -982 m (Bedmap3)/ -875 m (BedMachine3), before becoming stabilised on a bedrock pinning point (Fig. 9a).

The grounding line position at Holmes East was observed to retreat 15.24 km between 1996 and 2001, at an average pace of -3.05 km yr^{-1} (Fig. 8c), where it stabilised for the following 13 years, with ASaID, and MOA datasets digitised at nearly identical locations. However, comparing across the DInSAR-derived grounding line positions, we found 3.42 km of retreat from 1996 (MEaSURES) to 2019, at an average rate of -0.15 km yr^{-1} . We digitised the 2019 position within a 10.0 km wide buffer zone, indicating retreat within the range of -9.69 to 0.34 km , relative to 1996. The retreat continues to 2020; our findings show 5.27 km of retreat from the 1996 position, calculated at -0.22 km yr^{-1} , with a maximum of 14.65 km and minimum of 2.75 km retreat within the 11.9 km buffer zone, at a mean rate of -0.61 km yr^{-1} to -0.11 km yr^{-1} . The grounding line retreated initially down a retrograde slope before stepping up a prograde slope (Fig. 9c). We observed the grounding line to retreat between 1996 and 2001, remain stable between 2001 and 2014, and then advance to 2019. The difference in grounding line position between the DInSAR and the ASaID and MOA products in Figure 8c highlights the uncertainty associated with comparing products delineating the hinge line from DInSAR and break-in-slope from manual delineation.




Figure 8: Changes in grounding line position measured relative to the minimum 1996 position observed at (a) Frost Glacier, (b) Glacier 1, (c) Holmes East Glacier, and (d) Holmes West Glacier. The 1996 position from MEaSUREs (Rignot et al., 2016), 2001 from ASAID (Bindshadler & Choi, 2011), 2004, 2009, and 2014 from MOA (Haran et al., 2018; 2021a; 2021b), 2019 and 2020 from this paper.

In contrast, the grounding line position of Holmes West Glacier was observed to be relatively stable between 1996 and 2014; the grounding lines were digitised at very similar positions across products (Fig. 8d). We recorded a -1.28 km change between 1996 and 2014, recorded at an average rate of -0.07 km yr $^{-1}$. The grounding line retreated down a very shallow slope but is positioned at the edge of a significant steepening retrograde bed that continues down a reverse slope for 20 km, with the grounding line position falling almost 700 m along only 2 km of retreat along the flowline (Fig. 9d).

The grounding line position at Glacier 1 showed retreat then advance (Fig. 8b). Between 1996 and 2004, the grounding line position moved -6.01 km, calculated at an average rate of -0.75 km yr $^{-1}$. The grounding line was subsequently observed to advance 4.18 km by 2014, at an average rate of 0.42 km yr $^{-1}$. Observations show an overall 1.83 km retreat across a largely flat bedrock elevation (Fig. 9b).




Figure 9: Ice surface and bed elevation profiles extracted along the central flowlines of (a) Frost Glacier, (b) Glacier 1, (c) Holmes East Glacier, and (d) Holmes West Glacier. Bed elevation: Bedmap3 (Pritchard et al., 2024), BedMachine3 (Morlighem, 2022), and ANTGG (Rignot et al., 2024), ice surface elevation: REMA (Howat et al., 2022), ice shelf base: Bedmap3 (Pritchard et al., 2024). The red lines depict the oldest (1996 DInSAR) and newest (2014 MOA or 2020 DInSAR) grounding line (dashed red in (c) shows the maximum uncertainty). Measurements are shown in the along-flow direction, in which 0 km represents the inland start point.

3.5 Variability in ocean and sea-ice conditions

Although there is substantial variability in EN4 ocean temperature over time and no obvious long-term trend (Fig. 10), these reanalysis data show the presence of warm ocean water on the continental shelf in Porpoise Bay over the last three decades or so. At the surface (shallower than 100 m depth), we record temperatures largely below -1°C , calculated at a mean temperature of -1.4°C between 1990 and 2025 (Fig. 11). There is annual or biennial seasonal warming to $>0^{\circ}\text{C}$ at depths above 50 m that last for a few summer months (Fig. 10). At intermediate depths (100 to 350 m depth), we calculate a mean ocean temperature of -0.7°C between 1990 and 2025 (Fig. 11), with seasonal cycles displaying approximately -0.5°C to -1°C in winter and 0°C to 0.5°C in summer (Fig. 10). At depth below 450 m, the ocean is largely warmer than 0°C with a seasonal cycle between -0.5°C (winter) and 0.5°C or warmer (summer). We calculate a mean temperature of 0.1°C for depths below 447 m between 1990 and 2025. We record cooler periods, during which temperatures warmer than 0°C are not present at depths greater than 500 m across 1997–1999, 2004–2005, 2007, 2011–2015, and 2019–2021 (Fig. 10). As noted above, we observe no long-term trends in ocean temperature over the 1990–2025 period across various depths (Fig. 11).






Figure 10: EN4 subsurface ocean temperature depth profile for 1990 to 2025 from 4 grid cells between 65° and 66°S , 127° and 128°E on the continental shelf, Porpoise Bay (Fetterer et al., 2017).



430 **Figure 11: Annual EN4 subsurface ocean temperature across three depth ranges (surface: above 98 m, intermediate: between 110 m and 373 m, deep: below 447 m). Dashed lines show mean temperature at each depth.**

18 MEOP casts recorded between 2004 to 2021 reveal warm, salty water between 200 m and 1000 m depth at the continental shelf break, close to Porpoise Bay (Fig. 12). Further offshore, warm, salty waters are recorded over the continental slope, ranging from 0°C to 2°C (Fig. 12a) with salinity of $\sim 34.8 \text{ g kg}^{-1}$ (Fig. 12b) between 200 m and 1000 m depth (Fig. 12c). Over the continental shelf there are relatively few MEOP casts, with none within 130 km of the ice sheet margin, possibly due to the persistence of sea ice in the region. However, *in situ* ocean temperatures on the outer continental shelf and shelf break, north-west of Porpoise Bay, are recorded above freezing at depth, with temperatures ranging from 0.19°C (at 1000 m depth) to 0.13°C (at 918 m depth) to 0°C (at 690 m depth) (Fig. 12a and 12c). Closer to the surface, we record cooler temperatures -1.77°C (at 152 m depth) to -1.68°C (at 324 m depth). The casts record salinity up to 34.7 g kg^{-1} on the continental shelf, with many of the values above 34.5 g kg^{-1} (Fig. 12b). mCDW in East Antarctica is characterised by high salinity ($>34.5 \text{ g kg}^{-1}$) and temperature above freezing (Schodlok et al., 2016). Our findings suggest the presence of mCDW around the continental shelf break near Porpoise Bay at depths at, or shallower than, the continental shelf edge.



445 Figure 12: Oceanic properties derived from the MEOP casts and Argo floats between 2004 and 2021 near Porpoise Bay: (a)
 maximum conservative temperature (°C) observed below 150 m depth at each seal dive, (b) absolute salinity (g kg^{-1}) at the depth of
 (a), (c) depth of observed maximum temperature (a), (d) bathymetry of the continental shelf from ANTGG (Rignot et al., 2024).
 Dashed line shows continental shelf break from Amblas (2018) at approximately 600–800 m, solid line shows 2007–2009 ice sheet
 and shelf edges from MEaSURES (Mouginot et al., 2017).

450

Our analysis of monthly sea-ice concentration from 1979 to 2025 (Fig. 13a) reveals lower-than-average sea-ice concentrations in March 2006, 2007, 2010, 2016, 2021, and 2022, with very low sea-ice concentrations in 2007 (38 %), 2010 (50 %), and 2021 (41 %). These abnormally low March sea-ice conditions coincide with the beginning of calving events recorded at the ice shelves (Fig. 3). Furthermore, the results show lower-than-average mean sea-ice concentration in the winter-spring (June to November) of 2006, 2015, and 2020, and the summer-autumn (Dec–May) of 2007, 2016, and 2021 (Fig. 13). The February–March mean sea-ice concentration is similarly low in 2007, 2016, and 2021 (Fig. 13d). These low seasonal sea-ice conditions coincide or directly precede calving events recorded at Frost, Glacier 1, and Holmes West ice shelves (Fig. 3).




Figure 13: Sea-ice concentration extracted from a 100 km × 125 km box extending 130 km offshore from Porpoise Bay (see Fig. 1a). (a) Monthly sea-ice concentration with March mean in dashed blue; pale grey depicts when one glacier underwent a calving event, dark grey denotes when three or more glaciers calved (see Fig. 3). Note that only four images are available prior to the early 2000s (Supplementary Table 1), which explains why some of the earlier sea-ice minima are not associated with calving events and why we only plot the mean change in ice shelf position (red line) from the early 2000s. (b) Winter–Spring mean sea-ice concentration (June to November). (c) Summer–Autumn (December to May). (d) February–March. Asterisks in (b)–(d) show when calving events occur at three or more ice-shelf fronts (Fig. 3). Dashed lines in (b)–(d) show the mean seasonal sea-ice concentration.

3.6 Summary of observations

Our key observations show that changes at Frost, Holmes East, and Holmes West glacier are consistent with dynamic changes and mass loss observed elsewhere in Antarctica (Table 3), albeit with a lower magnitude than some of the more dramatic changes observed in West Antarctica. Over the study period, we record grounded ice surface lowering, grounding line retreat, moderate ice surface velocity increases, and ice-shelf retreat of varying degrees across Frost, Holmes East, and Holmes West glaciers. The highest surface elevation change is seen at Holmes West (-1.22 m yr^{-1}), where the grounding line retreat is modest (70 m yr^{-1}), but where it sits on a deep bed with a retrograde bed-slope further inland. The surface elevation change at Frost and Holmes East is more moderate (-0.13 m yr^{-1} and -0.4 m yr^{-1} , respectively) but the grounding line retreat of both glaciers is high ($>500\text{ m yr}^{-1}$) compared to most other outlet glaciers in East Antarctica (Konrad et al., 2018; Stokes et al., 2022; Picton et al., 2023; Zhu et al., 2025; Rignot et al., 2026). The long-term trend of ice velocity is also increasing across all three of these glaciers, albeit very slowly, with their ice front position also retreating. In contrast, the much smaller Glacier 1 shows limited evidence of dynamic change (Table 3).

Table 3: Summary characteristics of each outlet glaciers. (a) Mean annual surface elevation change across the GL between 2003 and 2017 from Nilsson et al. (2023). (b) Mean annual grounding line position change between 1996 and 2014 or 2020 (maximum position within the buffer). (c) Mean annual percentage change in ice surface velocity between 2000 and 2022 (2014–2022 for HE and HW) (see Table S2). (d) Mean annual ice-shelf calving front change between 1963 and 2025. (e) Change in bed elevation between the 2014/2020 grounding line position and 10 km inland along the flowline from Bedmap3. Formatting indicates the degree to which each statistic reflects dynamic change: values in *bold italics* show the strongest evidence of dynamic change, values in *italics* show moderate evidence, and values in plain text show minimal or inconsistent evidence.

	Frost Glacier	Glacier 1	Holmes East Glacier	Holmes West Glacier
a. surface elevation (2003–2017)	<i>− 0.13 m yr^{−1}</i>	0 m yr ^{−1}	<i>− 0.4 m yr^{−1}</i>	<i>− 1.22 m yr^{−1}</i>
b. grounding line position (1996–2014/2020)	<i>− 950 m yr^{−1}</i>	<i>− 100 m yr^{−1}</i>	<i>− 610 m yr^{−1}</i>	<i>− 70 m yr^{−1}</i>
c. ice surface velocity (2000–2022)	+ 0.3 %	− 0.5 %	+ <i>1.7 %</i>	+ 0.1 %
d. ice-shelf frontal position (1963–2025)	<i>− 110 m yr^{−1}</i>	<i>− 17 m yr^{−1}</i>	− 3 m yr ^{−1}	<i>− 29 m yr^{−1}</i>
e. bed elevation	− 4 m	+ 18 m	+ 460 m	<i>− 783 m</i>

4 Discussion

4.1 Grounding line retreat and ice surface thinning

490 Porpoise Bay displays a pattern of localised grounding line retreat. If correct, our results would make Frost one of the fastest retreating glaciers in East Antarctica on decadal timescales at an average rate of -0.95 km yr^{-1} between 1996 and 2014 (Fig. 8a and 9a), with a more rapid rate of -2.15 km yr^{-1} between 1996 and 2001 (Fig. 8a). Neighbouring Holmes East Glacier underwent grounding line retreat at a maximum rate of -0.61 km yr^{-1} between 1996 and 2020 (Fig. 8c), observed at the maximum width of the grounding line uncertainty buffer zone. For context, only Vanderford Glacier in East Antarctica
495 sustained similarly high rates of grounding line retreat at -0.78 km yr^{-1} (1996–2020) (Picton et al., 2023). In the WAIS, Thwaites and Pine Island glaciers in the Amundsen Sea Embayment experienced similarly rapid rates of -0.8 km yr^{-1} and -0.95 km yr^{-1} between 1992 and 2011 (Park et al., 2013; Milillo et al., 2019).

We acknowledge that our measurements of grounding line retreat at Frost Glacier are subject to uncertainties due to a comparison across various products that not only delineate different parts of the grounding zone (i.e., hinge line vs break-in-slope), but also use different methods (i.e., DInSAR vs manual delineation) with varying levels of uncertainty. Furthermore,
500 the difference between the DInSAR and the ASaID and MOA positions at Holmes East Glacier (Fig. 8c) highlight the uncertainty between the ASaID and MOA grounding line position results at Frost Glacier (Fig. 8a), suggesting retreat could be less rapid than we record. However, even if the rates of retreat are not as high as we report in Figure 8a (due to uncertainties), our findings are broadly consistent with the grounding line retreat observed by Konrad et al. (2018), measured at -0.2 km yr^{-1}
505 between 2010 and 2016 using satellite altimetry and ice geometry, suggesting that grounding line retreat is almost certainly occurring. In addition, we note that a recent study by Rignot et al. (2026) also identified grounding line retreat at Frost Glacier (0.22 km yr^{-1}), Holmes West Glacier (0.37 km yr^{-1}), and Holmes East Glacier (0.29 km yr^{-1}) from 1996–2023, which they linked to bathymetry that channels warm mCDW towards grounding lines.

Grounded ice surface elevation lowering is also observed at Holmes East and West glaciers, propagating inland across
510 datasets (Schröder et al., 2019b; Smith et al., 2020b; Nilsson et al., 2023), indicating a potential signal of dynamic thinning. Holmes West Glacier displayed elevation change up to -1.22 m yr^{-1} at the grounding line (Fig. 6d), propagating 10 km inland at a rate of -0.34 m yr^{-1} (Fig. 5d). This rapid rate places Holmes West Glacier as one of the most rapidly thinning glaciers in East Antarctica over decadal timescales (Stokes et al., 2022); only at Totten Glacier are there observations of such extreme thinning, measured up to $-1.9 \pm 0.07 \text{ m yr}^{-1}$ (2003–2007) (Pritchard et al., 2009), $-1.22 \pm 0.1 \text{ m yr}^{-1}$ (2003–2019) (Smith et al., 2020b) and $-0.72 \pm 0.02 \text{ m yr}^{-1}$ (2010–2019) (Li et al., 2023). Since there have been no reported long-term changes in the precipitation or firn densification in the region (Schröder et al., 2019b), we attribute this elevation change to dynamic thinning. Our results also suggest that Frost, Holmes East, and West glaciers thinned whilst ice surface velocity increased modestly (Fig. 4), suggesting a connection between flow regime and ice thickness that is consistent with dynamic thinning (Pritchard et al., 2009).

520 Whilst we recorded moderate long-term thinning rates, closer inspection reveals that enhanced thinning took place over the same period that grounding line retreat was recorded. At Holmes East and West glaciers, the most rapid grounding line retreat took place between 1996 and 2001 (Fig. 8: 15.24 km and 1.81 km), during which we recorded enhanced thinning rates (Fig. 6: 0.63 m yr⁻¹ and 2.29 m yr⁻¹). This correlates with a pan-ice sheet study by Konrad et al. (2018), who found an approximately proportional relationship between ice thickness change and grounding line migration. Overall, the localised
525 grounding line retreat and the dynamic thinning, alongside some indication of ice surface velocity speed-up at Holmes East Glacier, are consistent with a sustained period of change in Porpoise Bay that may be indicative of a dynamic response to external forcing.

4.2 Role of ocean forcing in Porpoise Bay glacier dynamics

High rates of grounding line retreat and ice surface lowering inland have been widely attributed to the intrusion of warm
530 mCDW across the continental shelf towards Antarctic sub-ice cavities (Thoma et al., 2008; Paolo et al., 2015; Scambos et al., 2017; Rignot et al., 2019; Rignot et al., 2026). The rates of grounding line retreat observed at Frost Glacier and Holmes East Glacier are consistent with warm mCDW incursion beneath Frost Ice Shelf and Holmes East Ice Shelf. Although our data show no evidence of any long-term changes in ocean temperature (Fig. 11), recent observations of mid-depth CDW along the continental slope off East Antarctica (80–160° E), and including offshore Porpoise Bay, show warming since the 1990s
535 (Herraiz-Borreguero and Garabato, 2022). This warming is attributed to the southwards shift in the Antarctica Circumpolar Current (ACC), potentially driven by a poleward shift in the westerlies over the Southern Ocean that is linked to a positive trend in the Southern Annular Mode (Yamazaki et al., 2021; Herraiz-Borreguero and Garabato, 2022).

Furthermore, we recorded 18 *in situ* temperature and salinity measurements from MEOP casts that indicate mCDW presence on the outer continental shelf and shelf break. MEOP and Argo casts further offshore suggest CDW on the continental
540 slope between 2004 and 2021 (Fig. 12). In the absence of ocean temperature observations within 130 km of the ice-shelf margin, we use EN4 subsurface ocean temperatures to gain potential insight into the existence of mCDW across the continental shelf between 1990 and 2025 (Fig. 10). Though we acknowledge the uncertainties, EN4 data suggest the presence of mCDW at depth across the continental shelf since the 1990s, with an average ocean temperature of 0.7°C below 447 m, and a seasonal cycle of 0–0.5°C at intermediate depth (100–350 m) between 1990 and 2025. As noted above, mCDW in East Antarctica is
545 characterised by high salinity (>34.5 g kg⁻¹) and temperature above freezing (Schodlok et al., 2016) and the MEOP casts record salinity values up to 34.7 g kg⁻¹ on the continental shelf, with many of the values above 34.5 g kg⁻¹ (Fig. 12b). To summarise, our data suggests that CDW has existed periodically in the bay since at least the 1990s, although our data is unable to show whether its frequency has increased in recent years. The intrusion of mCDW has the potential to cause ice-shelf melting and thinning, delivering freshwater that could inhibit the formation of dense shelf water in Porpoise Bay, strengthening water
550 column stratification and enabling the enhanced intrusion of warm mCDW at depth (Ribeiro et al., 2021).

At Holmes Ice Shelf (East and West), high rates of ice-shelf basal melt have been estimated, which is also consistent with intrusions of mCDW. Using satellite radar altimetry with satellite-derived ice velocities and a model of firn-layer

evolution, Adusumilli et al. (2020) calculated an average basal melt rate of $13.3 \pm 2.9 \text{ m yr}^{-1}$ beneath the Holmes Ice Shelf between 1994 and 2018, the highest average basal melt rate of any ice shelf in East Antarctica, and fourth in Antarctica behind the ice shelves of Thwaites ($26.7 \pm 2.4 \text{ m yr}^{-1}$), Land ($20.4 \pm 2.7 \text{ m yr}^{-1}$), and Pine Island ($14.0 \pm 1.6 \text{ m yr}^{-1}$) glaciers. Adusumilli et al. (2020) reported that Holmes Ice Shelf had a higher average basal melt rate than Totten Ice Shelf ($11.5 \pm 2.0 \text{ m yr}^{-1}$), where hydrographic observations show warm mCDW at the ice front (Rintoul et al., 2016).

Studies also show that Holmes Ice Shelf has been thinning. Using ICESat- and ICESat-2-derived data, Smith et al. (2020b) recorded a maximum ice surface change rate of -3.07 m yr^{-1} (2003–2019) on the floating ice shelf. A recent study extended this thinning record using the surface expression of pinning points as a proxy for ice-shelf thickness; Miles and Bingham (2024) found that Holmes Ice Shelf began to thin at least 50 years ago, alongside Moscow University and Totten, evinced by a reduction in pinning points across three study epochs: 1973–1989, 1989–2000, and 2000–2022. Miles and Bingham (2024) concluded by noting that Holmes Ice Shelf was showing signs of acceleration and that the continuation of pinning-point loss would likely reduce buttressing and cause acceleration of ice discharge and mass loss.

Porpoise Bay’s pattern of localised rapid grounding line retreat and thinning is consistent with a dynamic response of glaciers flowing into thinning ice shelves due to warm mCDW intrusion. To enable mCDW to access the grounding line, sufficiently deep cross-shelf troughs must exist in front of Porpoise Bay to transfer warm water from the continental shelf break to the grounding line. Although bathymetric data are limited across Porpoise Bay due to the persistence of sea ice in the bay, a recent study using seal data found several dives at the edge of the continental shelf, offshore from Porpoise Bay, that were up to 500 m deeper than the reported bathymetry (McMahon et al., 2023). Indeed, a trough may exist that connects the open ocean and grounding line via a deeper corridor of topography across the continental shelf that could transport mCDW to the grounding line, causing enhanced melting (Fig. 12d). In addition, the ice-shelf cavity must be sufficiently deep for mCDW to access the grounding line. Using 3D inversion, seismic, and MBES/SBES data, Rignot et al. (2024) found the bed at the ice margin of all four outlet glaciers was deeper than reported by BedMachine3 or Bedmap3, by over 200 m for several tens of kilometres along the flowlines (Fig. 9 and 12d). Visual inspection of the dataset (Rignot et al., 2024) reveals deep sub-ice-shelf cavities below Frost, Holmes East, and Holmes West glaciers, between 650 m to 750 m depth, that deepen inland (Fig. 9). These cavities are sufficiently large to plausibly provide a pathway for mCDW, which the EN4 and MEOP datasets suggest may exist below approximately 450 m depth (Fig. 10), to circulate under the ice shelves and melt the base of the ice shelves. This is consistent with satellite-derived high basal melt rates (Adusumilli et al., 2020), ice-shelf thinning (Smith et al., 2020b; Miles and Bingham, 2024) and grounding line retreat (Rignot et al., 2026) reported elsewhere and likely causing the dynamic changes observed in Porpoise Bay.

The bed topography beneath the outlet glaciers may differentially enhance the impact of ocean forcing in Porpoise Bay. We record sufficiently low bed elevation across the observed grounding line positions at Frost (Fig. 9a: below 600 m in 1996 and 900 m in 2014) and Holmes East (Fig. 9c: below 800 m in 1996 and 900 m in 2020) for mCDW to plausibly access the grounding line, driving grounding line retreat. Another key detail revealed by the ANTGG2021 dataset is that Holmes West Glacier has been grounded on a bedrock high between 1996 and 2014 (Rignot et al., 2024). The bed seaward of the

grounding zone ranges from 600 m to 740 m depth but rises to 500 m to 485 m depth at the grounding zone (Fig. 9d). This bathymetry may limit warm mCDW from accessing the grounding line, which may explain the lower rate of grounding line retreat at Holmes West Glacier, compared to Frost and Holmes East glaciers (Fig. 8), alongside the rapid decrease in surface elevation observed (Fig. 6d), since Holmes West Glacier may flow into a rapidly thinning ice shelf, potentially caused by basal melting. Furthermore, the bed at Glacier 1 is entirely above 500 m depth, which may limit potential exposure to mCDW at the grounding line, explaining the lack of observed dynamic change at Glacier 1 (Table 3).

4.3 Role of sea ice in driving ice shelf and glacier flow changes

Variations in sea-ice conditions likely influenced the behaviour of calving events, ice-shelf area, and some moderate changes in ice flow velocity in Porpoise Bay. The ice shelves underwent near-simultaneous calving events over the study period; observations show calving events in 2006–2008, 2016–2017, and 2021–2022. Recent studies around Antarctica demonstrate that the presence of sea ice and mélange may exert a resistive backstress on the ice-shelf calving front, and calving events have been linked to the break-up of sea ice and the removal of mélange from the ice front in various settings, including East Antarctica (Amundson et al., 2010; Miles et al., 2017; Gomez-Fell et al., 2022; Kondo and Sugiyama, 2023; Parsons et al., 2024; Teder et al., 2025; Deakin et al., submitted), and the nearby Voyeykov Ice Shelf (Arthur et al., 2021). Resultant ice-shelf calving can reduce their buttressing potential (Dupont and Alley, 2005; Fürst et al., 2016), which can trigger dynamic thinning and ice flow acceleration across the glacier further upstream (Reese et al., 2018; Gudmundsson et al., 2019).

Our findings show that all three major calving events appear to coincide with lower-than-average winter-spring sea-ice concentrations the preceding year and lower-than-average summer-autumn and February–March sea-ice conditions at the beginning of each major calving event (2006–2008, 2016–2017, and 2021–2022) (Fig. 13). Furthermore, the monthly sea-ice concentration was anomalously low in March 2007 and March 2021 (Fig. 13a). This is supported by our observations from Landsat 7 and Landsat 8 satellite imagery that shows the break-up and removal of the mélange and sea ice adjacent to Holmes Ice Shelf in 2007, 2017 and 2021. Therefore, it is likely that the break-up of sea ice and the removal of iceberg mélange reduced the stabilising backstress on the ice-shelf fronts in Porpoise Bay, culminating in major, near-synchronous calving events. This is consistent with Miles et al. (2017), who attributed the 2007 and 2016 calving events to the break-up of multi-year sea ice and suggested that atmospheric circulation anomalies caused the sea-ice break-up. We also observed abnormally low sea-ice conditions in 2002 (Fig. 13), when we also recorded a calving event from Holmes East Ice Shelf (there are no frontal position record for the other ice shelves due to a lack of cloud-free imagery). Although we hypothesise that the sea-ice break out events may have triggered enhanced calving through the removal of the back-stress on the glacier tongues (e.g., Gomez-Fell et al., 2022; Greene et al., 2025), we acknowledge that correlation does not necessarily imply causation and that further work is required to examine these processes in more detail around East Antarctica (Deakin et al., submitted). For example, it has been shown that other factors might influence calving after sea-ice break-out, such as the role of wind stress and ocean swell (Surawy-Stepney et al., 2024; Teder et al., 2025)

Porpoise Bay's ice shelves advanced and retreated cyclically, each time advancing to a similar (± 1 km) maximum length prior to calving (Fig. 2). In a previous study of the ice shelves in Porpoise Bay, Miles et al. (2017) observed Holmes West Ice Shelf calving at the same frontal position in each cycle and found that this took place as the glacier advanced and pushed the sea ice (attached to the ice-shelf calving front) out of the bay towards the open ocean and deeper water, causing it to disintegrate. This mechanism may have operated at all four glaciers, in part facilitated by the winter-spring sea-ice conditions from the preceding years.

Our findings suggest that ice-shelf calving in Porpoise Bay, alongside the intermittent loss of stabilising sea ice and mélange, had some corresponding impact on ice flow velocity for the outlet glaciers, suggesting that some of the ice shelves buttressed ice flow. This is consistent with Fürst et al. (2016), who used an ice flow model to calculate that only 18.3 % of the ice-shelf area in Porpoise Bay was passive (i.e., has no dynamic influence on ice discharge), meaning that once calving exceeds this 18.3 % of ice-shelf area, further ice shelf calving will have a dynamic impact inland. The glaciers lost over 18.3 % of the ice shelf area in most major calving events. The most consistent velocity response followed the 2007 calving event, during which Frost, Glacier 1, and Holmes West glaciers lost over 25% ice-shelf area (Fig. 3) and displayed an increased velocity response over the following few years (Fig. 4). Given that the calving event coincided with low sea-ice conditions (Fig. 13), these findings suggest a sea-ice-driven reduction in buttressing force, indicating that the ice shelves calved beyond the passive-ice area (Fürst et al., 2016). The debuttressing velocity response to the 2016 calving event was more varied, with an increase in velocity observed between 2016 and 2017 at Holmes East and West glaciers (Fig. 4c and 4d). Given the ice-shelf retreat was relatively small at Holmes East Glacier, this debuttressing is likely caused by the sea-ice break-up and low sea-ice concentration leading up to the calving, rather than the loss of buttressing ice-shelf area. For Frost Glacier, the onset of flow speed-up coincided with a small calving event in 2013, after which the ice shelf broadly retreated, suggesting that the loss of back-stress exerted by the ice shelf is a potential driver of glacier speed-up. During high sea-ice concentrations between 2017 and 2020 (Fig. 13c: summer-autumn mean calculated 10 % above average) and the readvance of the ice shelves (Fig. 3), we record a decrease in ice flow velocity, with Holmes East and West slowing 32 % and 5 % across the grounding line and 20 % and 28 % across the inland ice (Fig. 4). Although we might have expected a greater decrease in speed across the grounding line than further up-ice, it suggests that, overall, glacier flow speeds were reduced by the accumulation of sea ice and ice mélange near the glacier front. Subsequently, between 2020 and 2021, we record autumn-winter and winter-spring sea-ice concentration below average, ice-shelf retreat, and ice flow velocity increase across all sampling boxes in Holmes East and West glaciers, suggesting that the low sea-ice conditions impacted glacier dynamics before the 2021/22 ice-shelf calving event. This mirrors the pattern found at nearby Totten Glacier, where Greene et al. (2018) attributed the onset of ice shelf acceleration each spring to the loss of buttressing from the break-up of seasonal landfast sea ice. It is difficult to assess the impact of the 2021/22 calving event on glacier flow dynamics and the impact of sea-ice-driven debuttressing that occurred due to the lack of ice flow velocity data beyond 2022.

In the future, we may see an increase in low sea-ice states that could impact the future ice-shelf stability in Porpoise Bay, as low sea-ice conditions continue to destabilise ice shelves/ice tongues, and drive ice flow velocity increase (Gomez-

Fell et al., 2022). The past decade has seen extreme Antarctic-wide sea-ice minima in 2017, 2022, 2023 and 2024, and a sea-ice decline in 2014–2016. There is a growing concern that there is a structural change taking place in the Antarctic atmosphere-ocean-sea-ice system (Purich and Dodderidge, 2023; Hobbs et al., 2024), revealed by analysis of reconstructed and satellite-observed sea-ice extent that shows increasing persistence in sea-ice extent anomalies, a pattern especially clear in East Antarctica (Raphael et al., 2025).

4.4 Future unpinning at Holmes West Glacier

The current grounding line position at Holmes West Glacier raises the possibility that it may be vulnerable to MISI in the near future. Our findings show the grounding line was pinned on a local bedrock high between 1996 and 2014, which might plausibly have inhibited mCDW from accessing the grounding line (Fig. 9d). The 2014 grounding line position is at the very edge of a steep retrograde bed and, over the next 2 km inland along the flowline, the bed elevation drops by 650–700 m (Morlighem, 2022, Pritchard et al., 2024). However, we note that the reverse-sloping bed topography is subject to some uncertainty. The BedMachine3 bathymetry was produced using mass conservation in the absence of radar data (Morlighem et al., 2020) and suggests the bed is reverse sloping for only 3–4 km and then stabilises. In contrast, the Bedmap3 reverse slope continues inland for the next 25 km along the Holmes West Glacier flowline, which was extracted/interpolated from radar flight paths that intersect the trough, albeit sparsely spaced compared to Totten Glacier or Moscow University Glacier troughs (Pritchard et al., 2025). More concerning, the ANTGG bathymetry suggests the reverse slope continues for over 35 km along the flowline (Charrassin et al., 2025). Regardless, given its current grounding line retreat and thinning rate of 1.22 m yr^{-1} at the grounding line, Holmes West Glacier is likely to unground from the local bedrock high and retreat to become grounded on a bed below 1000 m depth, which may be sufficient to initiate MISI at Holmes West Glacier. This is concerning given that the Holmes catchment stores 11 cm of SLE (Rignot et al., 2019), but there is an urgent need to resolve some of the bed elevation uncertainties to aid prognostic modelling of this system.

In addition, there is an adjacent 4700 km² ice rise directly west along the coastline from Holmes West Glacier (Matsuoka et al., 2015) that may currently exert a buttressing force on the ice sheet, retarding the flow of grounded ice to the ocean and contributing to grounding line stability (Favier and Pattyn, 2015; Matsuoka et al., 2015). If the current rates of grounding line retreat (70 m yr^{-1}) continue, Holmes West Glacier could unpin from this bedrock high in the next decade, losing contact with the stabilising ice rise, potentially resulting in threshold-like behaviour (Matsuoka et al., 2015).

Although we observed greater grounding line retreat at Holmes East and Frost glaciers, the likelihood of irreversible grounding line retreat or MISI is less concerning given the elevation of the glacier troughs along the flowlines. The bed inland of the Frost Glacier grounding line is relatively flat and the bed is prograding inland of Holmes East Glacier grounding line (Fig. 9). It is generally understood that retrograde slopes favour more extensive grounding line retreat for a given basal melt rate (Milillo et al., 2019; Millan et al., 2022).

5 Conclusion

685 To investigate recent glacier dynamics in Porpoise Bay, Wilkes Land, East Antarctica, we measured changes in ice-shelf calving position, ice surface velocity, grounded ice surface elevation, and grounding line position of four outlet glaciers over the last three decades. The observed glacier dynamics were compared to sea-ice conditions and ocean temperatures and salinity to assess the potential forcing of any observed changes. Despite uncertainties, our results suggest Frost and Holmes East glaciers have undergone substantial grounding line retreat, with more modest rates of grounding line retreat observed elsewhere
690 in Porpoise Bay. Observations of surface elevation change depict considerable thinning at Frost, Holmes East, and Holmes West glaciers between 2003 and 2019, which propagated inland, with Holmes West Glacier displaying potentially one of the highest rates of thinning in East Antarctica. In addition, some of the outlet glaciers underwent relatively moderate increases in velocity over the study period. Taken together, these findings are consistent with dynamic changes in this region, similar to previous work suggesting that it is losing mass and contributing to sea level rise (Rignot et al., 2019; Adusumilli et al., 2020; Smith et al., 2020a; Rignot et al., 2022; 2026). In addition, we report on a previously unidentified calving event in 2021/22,
695 that extends observations of near-synchronous ice-shelf calving events in 2007/08 and 2016/17. As in previous work (Miles et al., 2017), we show that the calving events appear to be linked to a reduction in sea ice in the preceding winter and summer and the collapse of a supportive mélange.

Our findings of dynamic change are consistent with observations of mCDW at the shelf break seaward of Porpoise
700 Bay, with EN4 ocean temperature data suggesting the presence of mCDW at depths below 450 m on the continental shelf. We find that bathymetric pathways beneath Frost, Holmes East and Holmes West glaciers could transport mCDW to the sub-ice cavities, driving high rates of basal melting that have been reported in previous work (Adusumilli et al., 2020). Furthermore, we find Frost and Holmes East glaciers are grounded sufficiently deep for mCDW to directly access the grounding line. In contrast, Holmes West is grounded on a local bedrock high that could inhibit year-round access of mCDW to the grounded
705 ice, which is also consistent with its relatively low grounding line retreat rate. Of concern is that Holmes West Glacier is currently grounded at the edge of a steep retrograde bed that would see the grounding line lose up to 700 m elevation in less than 2 km retreat along the flowline, providing access for mCDW to melt the grounded ice, which could initiate rapid and irreversible grounding line retreat down a retrograde bed that continues on a reverse slope. This slope may persist for up to 35 km inland (Charrassin et al., 2025), but there is an urgent need for more detailed and accurate bed topography in this region
710 in order to assess the potential for MISI, especially as the Holmes drainage basin contains around 11 cm of equivalent sea level rise.

Data availability. Ice shelf frontal positions, central flowlines, sampling boxes, and grounding lines positions are available to download from Zenodo (<https://doi.org/10.5281/zenodo.15847590>; Weatherley et al., 2025). The monthly ice surface elevation change dataset from Schröder et al. (2019b) is available from PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.897390>) (1978–2017), from Nilsson et al. (2023) is available from NSIDC (<https://doi.org/10.5067/L3LSVDZS15ZV>) (1985–2020), and from Smith et al. (2020b) is available from the Research Works Archive (<https://doi.org/10.5067/L3LSVDZS15ZV>). The 2000–2022 ITS_LIVE annual velocity mosaics (Gardner et al., 2022) are available from NSIDC (<https://nsidc.org/data/nsidc-0776/versions/1>). The MEaSUREs 1996 grounding line positions (Rignot et al., 2016) are available from NSIDC (<https://doi.org/10.5067/IKBWW4RYHF1Q>). The ASAD 2001 grounding line positions (Bindshadler and Choi, 2011) are available from the US Antarctic Program Centre (<https://doi.org/10.7265/N56T0JK2>). The MOA grounding line positions from 2004 (Haran et al., 2021a), 2009 (Haran et al., 2021b), and 2014 (Haran et al., 2018) are available from NSIDC (<https://doi.org/10.5067/68TBT0CGJSOJ>), (<https://doi.org/10.5067/4ZL43A4619AF>), and (<https://doi.org/10.5067/RNF17BP824UM>). The 1979–2025 monthly sea ice concentration dataset (Fetterer et al., 2017) is available from NSIDC (<https://doi.org/10.7265/N5K072F8>). The MEOP data (Treasure et al., 2017) is available from <https://meop.net/database/>. The Argo data (Wong et al., 2020) is available from the International Argo Program (<https://argo.ucsd.edu/>). The EN4 subsurface ocean temperature objective analyses (Good et al., 2013) are available from the Met Office Hadley Centre (<https://www.metoffice.gov.uk/hadobs/en4/>). The Bedmap3 bed elevation, ice surface elevation, and ice thickness datasets (Pritchard et al., 2024) are available from NERC EDS UK Polar Data Centre (<https://doi.org/10.5285/2d0e4791-8e20-46a3-80e4-f5f6716025d2>). The BedMachine3 bed elevation (Morlighem, 2022) is available from NSIDC (<https://doi.org/10.5067/FPSU0V1MWUB6>). The ANTGG bed elevation (Rignot et al., 2024) is available from Dryad (<https://doi.org/10.5061/dryad.rbnzs7hkc>). The REMA elevation dataset (Howat et al., 2022) is available from Harvard Dataverse (<https://doi.org/10.7910/dvn/x7ndny>).

Author contributions. MW, CRS, and SSRJ contributed to the design of the initial project. MW undertook the data collection and led the analysis, with guidance from CRS and SSRJ. SR completed the DInSAR grounding line mapping within Porpoise Bay, with MW writing up the method, results, and figures and additional figure contributions from AS. MW led the manuscript writing, with all authors editing the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. SSRJ and CRS are supported by Natural Environment Research Council (NERC) UK research grant NE/Y00115X/1. AS also acknowledges funding from NERC (NE/V014285/1). We are grateful to Hannah Picton and Bertie Miles for their guidance. The authors would also like to thank Bert Wouters and the three anonymous reviewers for their constructive comments on this paper.

References

- Adusumilli, S., Fricker, H.A., Medley, B., Padman, L., and Siegfried, M.R.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nat. Geosci.*, 13, 616–620, <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- Amundson, J., Fahnestock, M., Truffer, M., Brown, J., Lüthi, M., and Motyka, R.: Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland, *J. Geophys. Res.*, 115, F01005, <https://doi.org/10.1029/2009JF001405>, 2010.
- Amblas, D.: Antarctic continental shelf break (shapefile), PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.890863>, 2018.
- Arthur, J.F., Stokes, C.R., Jamieson, S.S.R., Miles, B.W.J., Carr, J.R. and Leeson, A.A.: The triggers of the disaggregation of Vöyeykov Ice Shelf (2007), Wilkes Land, East Antarctica, and its subsequent evolution. *J. Glaciol.*, 67, 933–951, <https://doi.org/10.1017/jog.2021.45>, 2021.
- Berthier, E., Scambos, T., Shuman, C.: Mass loss of Larsen B tributary glaciers (Antarctic Peninsula) unabated since 2002. *Geophys. Res. Lett.*, 39, 1–6, <https://doi.org/10.1029/2012GL051755>, 2012.
- Bindschadler, R. and Choi, H.: High-resolution Image-derived Grounding and Hydrostatic Lines for the Antarctic Ice Sheet. U.S. Antarctic Program (USAP) Data Center [data set] <https://doi.org/10.7265/N56T0JK2>, 2011.
- Bindschadler, R., Choi, H., Wichlacz, A., Bingham, R., Bohlander, J., Brunt, K., Corr, H., Drews, R., Fricker, H., Hall, M., Hindmarsh, R., Kohler, J., Padman, L., Rack, W., Rotschky, G., Urbini, S., Vornberger, P., and Young, N.: Getting around Antarctica: new high-resolution mappings of the grounded and freely-floating boundaries of the Antarctic ice sheet created for the International Polar Year, *The Cryosphere*, 5, 569–588, <https://doi.org/10.5194/tc-5-569-2011>, 2011.
- Black, T. E. and Joughin, I.: Multi-decadal retreat of marine-terminating outlet glaciers in northwest and central-west Greenland, *The Cryosphere*, 16, 807–824, <https://doi.org/10.5194/tc16-807-2022>, 2022.
- Cavalieri, D.J., St. Germain, K.M. and Swift, C.T.: Reduction of weather effects in the calculation of sea-ice concentration with the DMSP SSM/I, *J. Glaciol.*, 41, 455–464, <https://doi.org/10.3189/s0022143000034791>, 1995.
- Charrassin, R., Millan, R., Rignot, E. and Scheinert, M.: Bathymetry of the Antarctic continental shelf and ice shelf cavities from circumpolar gravity anomalies and other data, *Sci. Rep.*, 15, <https://doi.org/10.1038/s41598-024-81599-1>, 2025.
- Cook, C.P., van de Flierdt, T., Williams, T., Hemming, S.R., Iwai, M., Kobayashi, M., Jimenez-Espejo, F.J., Escutia, C., González, J.J., Khim, B.-K., McKay, R.M., Passchier, S., Bohaty, S.M., Riesselman, C.R., Tauxe, L., Sugisaki, S., Galindo, A.L., Patterson, M.O., Sangiorgi, F. and Pierce, E.L.: Dynamic behaviour of the East Antarctic ice sheet during Pliocene warmth, *Nat. Geosci.*, 765–769, <https://doi.org/10.1038/ngeo1889>, 2013.
- Cook, A., Vaughan, D., Luckman, A., Murray, T.: A new Antarctic Peninsula glacier basin inventory and observed area changes since the 1940s, *Antarct. Sci.*, 26, 614–624, <https://doi.org/10.1017/S0954102014000200>, 2014.

- 780 Cook, A., Holland, P., Meredith, M., Murray, T., Luckman, A., Vaughan, D.: Ocean forcing of glacier retreat in the western Antarctic Peninsula, *Science*, 353, <https://doi.org/10.1126/science.aae0017>, 283–286, 2016.
- Davison, B. J., Hogg, A. E., Gourmelen, N., Jakob, L., Wuite, J., Nagler, T., Greene, C. A., Andreasen, J., and Engdahl, M. E.: Annual mass budget of Antarctic ice shelves from 1997 to 2021, *Sci. Adv.*, 9, eadi0186, <https://doi.org/10.1126/sciadv.adi0186>, 2023.
- 785 Deakin, K.A., Stokes, C.R., Jamieson, S.S.R., and Carr, J.R.: The influence of sea ice on outlet glacier dynamics: recent progress and future challenges. *Rev. Geophys.*, <https://doi.org/10.22541/essoar.176350662.27264139/v1>, submitted.
- Dupont, T.K. and Alley, R.B.: Assessment of the importance of ice-shelf buttressing to ice-sheet flow, *Geophys. Res. Lett.*, 32, L04503, <https://doi.org/10.1029/2004gl022024>, 2005.
- Favier, L., and Pattyn, F.: Antarctic ice rise formation, evolution, and stability. *Geophys. Res. Lett.*, 42, 4456–4463. 790 <https://doi.org/10.1002/2015GL064195>, 2015.
- Feldmann, J. and Levermann, A.: Collapse of the West Antarctic Ice Sheet after local destabilization of the Amundsen Basin, *P. Natl. Acad. Sci. USA*, 112, 14191–14196, <https://doi.org/10.1073/pnas.1512482112>, 2015.
- Fetterer, F., Knowles, K., Meier, W. N., Savoie, M. and Windnagel, A. K.: Sea Ice Index. (G02135, Version 3), National Snow and Ice Data Center [data set], <https://doi.org/10.7265/N5K072F8>, 2017.
- 795 Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavad, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nat. Clim. Change*, 6, 479–482, <https://doi.org/10.1038/nclimate2912>, 2016.
- Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., van den Broeke, M., and Nilsson, J.: Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years, *The Cryosphere*, 12, 521–547, <https://doi.org/10.5194/tc-12-521-2018>, 2018.
- 800 Gardner, A., Fahnestock, M., and Scambos, T.: ITS_LIVE Regional Glacier and Ice Sheet Surface Velocities: Version 1, National Snow and Ice Data Center [data set], <https://doi.org/10.5067/6II6VW8LLWJ7>, 2025.
- Gomez-Fell, R., Rack, W., Purdie, H., and Marsh, O.: Parker Ice Tongue collapse, Antarctica, triggered by loss of stabilizing land-fast sea ice, *Geophys. Res. Lett.*, 49, e2021GL096156, <https://doi.org/10.1029/2021GL096156>, 2022.
- Good, S.A., Martin, M.J. and Rayner, N.A.: EN4: Quality Controlled Ocean Temperature and Salinity Profiles and Monthly 805 Objective Analyses with Uncertainty Estimates, *J. Geophys. Res. Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.
- Greenbaum, J. S., Blankenship, D. D., Young, D. A., Richter, T. G., Roberts, J. L., Aitken, A. R. A., Legresy, B., Schroeder, D. M., Warner, R. C., van Ommen, T. D., and Siegert, M. J.: Ocean access to a cavity beneath Totten Glacier in East Antarctica, *Nat. Geosci.*, 8, 294–298, <https://doi.org/10.1038/ngeo2388>, 2015.
- 810 Greene, C. A., Young, D. A., Gwyther, D. E., Galton-Fenzi, B. K., and Blankenship, D. D.: Seasonal dynamics of Totten Ice Shelf controlled by sea ice buttressing, *The Cryosphere*, 12, 2869–2882, <https://doi.org/10.5194/tc-12-2869-2018>, 2018.

- Greene, C.A., Gardner, A.S., Schlegel, N.J., and Fraser, A.S.: Antarctic calving loss rivals ice-shelf thinning, *Nature*, 609, 948–953, <https://doi.org/10.1038/s41586-022-05037-w>, 2022.
- 815 Gudmundsson, G.H., Paolo, F.S., Adusumilli, S. and Fricker, H.A.: Instantaneous Antarctic ice sheet mass loss driven by thinning ice shelves. *Geophysical Research Letters*, 46, 3903–13909, <https://doi.org/10.1029/2019gl085027>, 2019.
- Haran, T., Klinger, M., Bohlander, J., Fahnestock, M., Painter, T., and Scambos, T.: MEaSUREs MODIS Mosaic of Antarctica 2013–2014 (MOA2014) Image Map, Version 1. NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/RNF17BP824UM>, 2018.
- 820 Haran, T., Bohlander, J., Scambos, T., Painter, T., and Fahnestock, M.: MODIS Mosaic of Antarctica 2003–2004 (MOA2004) Image Map, Version 2. NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/68TBT0CGJSOJ>, 2021a.
- Haran, T., Bohlander, J., Scambos, T., Painter, T., and Fahnestock, M.: MODIS Mosaic of Antarctica 2008–2009 (MOA2009) Image Map, Version 2. NASA National Snow and Ice Data Center Distributed Active Archive Center [data set],
825 <https://doi.org/10.5067/4ZL43A4619AF>, 2021b.
- Herraiz-Borreguero, L. and Garabato, A. C. N.: Poleward shift of Circumpolar Deep Water threatens the East Antarctic Ice Sheet, *Nat. Clim. Change*, 12, 728–734, <https://doi.org/10.1038/s41558-022-01424-3>, 2022.
- Hobbs, W., Spence, P., Meyer, A., Schroeter, S., Fraser, A.D., Reid, P., Tian, T.R., Wang, Z., Liniger, G., Doddridge, E.W., and Boyd, P.W.: Observational evidence for a regime shift in summer Antarctic sea ice, *J. Clim.*, 37, 2263–2275,
830 <https://doi.org/10.1175/jcli-d-23-0479.1>, 2024.
- Holt, T. O., Glasser, N. F., Quincey, D. J., and Siegfried, M. R.: Speedup and fracturing of George VI Ice Shelf, Antarctic Peninsula, *The Cryosphere*, 7, 797–816, <https://doi.org/10.5194/tc-7-797-2013>, 2013.
- Howat, I. M., Porter, C., Smith, B. E., Noh, M.-J., and Morin, P.: The Reference Elevation Model of Antarctica, *The Cryosphere*, 13, 665–674, <https://doi.org/10.5194/tc-13-665-2019>, 2019.
- 835 Howat, I., Porter, C., Noh, M.-J., Husby, E., Khuvis, S., Danish, E., Tomko, K., Gardiner, J., Negrete, A., Yadav, B., Klassen, J., Kelleher, C., Cloutier, M., Bakker, J., Enos, J., Arnold, G., Bauer, G. and Morin, P.: The Reference Elevation Model of Antarctica - Mosaics, Version 2, Harvard Dataverse [data set], <https://doi.org/10.7910/dvn/x7ndny>, 2022.
- Kim, K., Jezek, K. C., and Liu, H.: Orthorectified image mosaic of Antarctica from 1963 Argon satellite photography: image processing and glaciological applications, *Int. J. Remote Sens.*, 28, 5357–5373,
840 <https://doi.org/10.1080/01431160601105850>, 2007.
- Kondo, K. and Sugiyama, S.: Calving, ice flow, and thickness of outlet glaciers controlled by land-fast sea ice in Lützow-Holm Bay, East Antarctica. *J. Glaciol.*, 69, 1–13, <https://doi.org/10.1017/jog.2023.59>, 2023.
- Konrad, H., Shepherd, A., Gilbert, L., Hogg, A., McMillan, M., Muir, A., and Slater, T.: Net retreat of Antarctic glacier grounding lines, *Nat. Geosci.*, 11, 258–262, <https://doi.org/10.1038/s41561-018-0082-z>, 2018.

- 845 Lea, J.M.: The Google Earth Engine Digitisation Tool (GEEDiT) and the Margin change Quantification Tool (MaQiT) – simple tools for the rapid mapping and quantification of changing Earth surface margins, *Earth Surf. Dynam.*, 6, 551–561. <https://doi.org/10.5194/esurf-6-551-2018>, 2018.
- Leeson, A., Forster, E., Rice, A., Gourmelen, N., van Wessem, J.: Evolution of Supraglacial Lakes on the Larsen B Ice Shelf in the Decades Before it Collapsed, *Geophys. Res. Lett.*, 47, 1–9, <https://doi.org/10.1029/2019GL085591>, 2020.
- 850 Li, T., Dawson, G.J., Chuter, S.J. and Bamber, J.L.: Grounding line retreat and tide-modulated ocean channels at Moscow University and Totten Glacier ice shelves, East Antarctica, *The Cryosphere*, 17, 1003–1022, <https://doi.org/10.5194/tc-17-1003-2023>, 2023.
- Matsuoka, K., Hindmarsh, R., Moholdt, G., Bentley, M.J., Pritchard, H.D., Brown, J.S., Conway, H., Drews, R., Durand, G., Goldberg, D.E., Hattermann, T., Kingslake, J., Lenaerts, J., Martin, C., Mulvaney, R., Nicholls, K.W., Pattyn, F.,
- 855 Ross, N., Scambos, T. and Whitehouse, P.L.: Antarctic ice rises and rumples: Their properties and significance for ice-sheet dynamics and evolution, *Earth-Sci. Rev.*, 150, 724–745. <https://doi.org/10.1016/j.earscirev.2015.09.004>, 2015.
- McMahon, C.R., Hindell, M.A., Charrassin, J.B., Coleman, R., Guinet, Christophe., Harcourt, R., Labrousse, S., Raymond, B., Sumner, M. and Ribeiro, N.: Southern Ocean pinnipeds provide bathymetric insights on the East Antarctic
- 860 continental shelf, *Commun. Earth Environ.*, 4, <https://doi.org/10.1038/s43247-023-00928-w>, 2023.
- McMillan, M., Shepherd, A., Sundal, A., Briggs, K., Muir, A., Ridout, A., Hogg, A., and Wingham, D.: Increased ice losses from Antarctica detected by CryoSat-2, *Geophys. Res. Lett.*, 41, 3899–3905, <https://doi.org/10.1002/2014GL060111>, 2015.
- Miles, B. W. J., Stokes, C. R., and Jamieson, S. S. R.: Pan-ice sheet glacier terminus change in East Antarctica reveals
- 865 sensitivity of Wilkes Land to sea-ice changes, *Sci. Adv.*, 2, 1–7, <https://doi.org/10.1126/sciadv.1501350>, 2016.
- Miles, B. W. J., Stokes, C. R., and Jamieson, S. S. R.: Simultaneous disintegration of outlet glaciers in Porpoise Bay (Wilkes Land), East Antarctica, driven by sea ice break-up, *The Cryosphere*, 11, 427–442, <https://doi.org/10.5194/tc-11-427-2017>, 2017.
- Miles, B. W. J., Jordan, J. R., Stokes, C. R., Jamieson, S. S. R., Gudmundsson, G. H., and Jenkins, A.: Recent acceleration of
- 870 Denman Glacier (1972–2017), East Antarctica, driven by grounding line retreat and changes in ice tongue configuration, *The Cryosphere*, 15, 663–676, <https://doi.org/10.5194/tc-15-663-2021>, 2021.
- Miles, B.W.J. and Bingham, R.G.: Landsat mosaics of Antarctic Ice Shelves from 1973 and 1989, Edinburgh DataShare [data set], <https://doi.org/10.7488/ds/3810>, 2023.
- Miles, B.W.J. and Bingham, R.G.: Progressive unanchoring of Antarctic ice shelves since 1973, *Nat.*, 626, 785–791,
- 875 <https://doi.org/10.1038/s41586-024-07049-0>, 2024.
- Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J., and Prats-Iraola, P.: Heterogeneous retreat and ice melt of Thwaites Glacier, West Antarctica, *Sci. Adv.*, 5, eaau3433, <https://doi.org/10.1126/sciadv.aau3433>, 2019.

- 880 Millan, R., Mouginot, J., Derkacheva, A., Rignot, E., Milillo, P., Ciraci, E., Dini, L., and Bjørk, A.: Ongoing grounding line retreat and fracturing initiated at the Petermann Glacier ice shelf, Greenland, after 2016, *The Cryosphere*, 16, 3021–3031, <https://doi.org/10.5194/tc-16-3021-2022>, 2022.
- Moon, T. and Joughin, I.: Changes in ice front position on Greenland's outlet glaciers from 1992 to 2007, *J. Geophys. Res.*, 113, F02022, <https://doi.org/10.1029/2007JF000927>, 2008.
- 885 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. van den, Ommen, T. D., van Wessem, M., and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nat. Geosci.*, 13, 132–137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- 890 Morlighem, M.: MEaSUREs BedMachine Antarctica. (NSIDC-0756, Version 3), NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/FPSU0V1MWUB6>, 2022.
- Mouginot, J., Rignot, E., and Scheuchl, B.: Sustained increase in ice discharge from the Amundsen Sea Embayment, West Antarctica, from 1973 to 2013, *Geophys. Res. Lett.*, 41, 1576–1584, <https://doi.org/10.1002/2013GL059069>, 2014.
- Mouginot, J., Scheuchl, B. & Rignot, E.: MEaSUREs Antarctic Boundaries for IPY 2007–2009 from Satellite Radar (NSIDC-0709, Version 2), NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/AXE4121732AD>, 2017.
- 895 Nilsson, J., Gardner, A. S., and Paolo, F. S.: Elevation change of the Antarctic Ice Sheet: 1985 to 2020, *Earth Syst. Sci. Data*, 14, 3573–3598, <https://doi.org/10.5194/essd-14-3573-2022>, 2022.
- Nilsson, J., Gardner, A. S. and Paolo, F.: MEaSUREs ITS_LIVE Antarctic Grounded Ice Sheet Elevation Change (NSIDC-0782, Version 1), NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/L3LSVDZS15ZV>, 2023.
- 900 Otosaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N.-J., Amory, C., van den Broeke, M. R., Horwath, M., Joughin, I., King, M. D., Krinner, G., Nowicki, S., Payne, A. J., Rignot, E., Scambos, T., Simon, K. M., Smith, B. E., Sørensen, L. S., Velicogna, I., Whitehouse, P. L., A. G., Agosta, C., Ahlstrøm, A. P., Blazquez, A., Colgan, W., Engdahl, M. E., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B. C., Harig, C., Helm, V., Khan, S. A., Kittel, C., Konrad, H., Langen, P. L., Lecavalier, B. S., Liang, C.-C., Loomis, B. D., McMillan, M., Melini, D., Mernild, S. H., Mottram, R., Mouginot, J., Nilsson, J., Noël, B., Pattle, M. E., Peltier, W. R., Pie, N., Roca, M., Sasgen, I., Save, H. V., Seo, K.-W., Scheuchl, B., Schrama, E. J. O., Schröder, L., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T. C., Vishwakarma, B. D., van Wessem, J. M., Wiese, D., van der Wal, W., and Wouters, B.: Mass Balance of the Greenland and Antarctic Ice Sheets from 1992 to 2020, *Earth Syst. Sci. Data*, 15, 1–33, <https://doi.org/10.5194/essd-15-1597-2023>, 2023.
- 910

- Paolo, F. S., Fricker, H. A., and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348, 327–331, <https://doi.org/10.1126/science.aaa0940>, 2015.
- Park, J., Gourmelen, N., Shepherd, A., Kim, S., Vaughan, D., and Wingham, D.: Sustained retreat of the Pine Island Glacier, *Geophys. Res. Lett.*, 40, 2137–2142, <https://doi.org/10.1002/grl.50379>, 2013.
- 915 Parsons, R., Sun, S., Gudmundsson, G.H., Wuite, J. and Nagler, T.: Quantifying the buttressing contribution of landfast sea ice and mélange to Crane Glacier, Antarctic Peninsula, *The Cryosphere*, 18, 5789–5801, <https://doi.org/10.5194/tc-18-5789-2024>, 2024.
- Pattyn, F. and Morlighem, M.: The uncertain future of the Antarctic Ice Sheet, *Science*, 367, 1331–1335, <https://doi.org/10.1126/science.aaz5487>, 2020.
- 920 Paul, F., Bolch, T., Briggs, K., Kääb, A., McMillan, M., McNabb, R., Nagler, T., Nuth, C., Rastner, P., Strozzi, T., and Wuite, J.: Error sources and guidelines for quality assessment of glacier area, elevation change, and velocity products derived from satellite data in the *Glaciers_cci* project, *Remote Sens. Environ.*, 203, 256–275, <https://doi.org/10.1016/j.rse.2017.08.038>, 2017.
- Pelle, T., Morlighem, M. and McCormack, F.S.: Aurora Basin, the Weak Underbelly of East Antarctica. *Geophys. Res. Lett.*, 925 47, 1–10, <https://doi.org/10.1029/2019GL086821>, 2020.
- Picton, H.J., Stokes, C.R., Jamieson, S.S.R., Floricioiu, D. and Krieger, L.: Extensive and anomalous grounding line retreat at Vanderford Glacier, Vincennes Bay, Wilkes Land, East Antarctica, *The Cryosphere*, 17, 3593–3616, <https://doi.org/10.5194/tc-17-3593-2023>, 2023.
- Pritchard, H. D., Arthern, R. J., Vaughan, D. G., and Edwards, L. A.: Extensive dynamic thinning on the margins of the 930 Greenland and Antarctic ice sheets, *Nature*, 461, 971–975, <https://doi.org/10.1038/nature08471>, 2009.
- Pritchard, H. D., Fretwell, P., Fremand, A. C., Bodart, J., Bamber, J. L., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Christianson, K., Conway, H., Corr, H. F. J., Cui, X., Damaske, D., Damm, V., Drews, R., Eagles, G., Eisen, O., Eisermann, H., Ferraccioli, F., Forsberg, R., Franke, S., Fujita, S., Gim, Y., Goel, V., Gogineni, P., Greenbaum, J., Hills, B., Hindmarsh, R. C. A., Holmlund, P., Holschuh, N., Holt, J. W., Humbert, A., 935 Jacobel, R. W., Jansen, D., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Langley, K. A., Lee, J., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., MacGregor, J., MacKie, E., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Paden, J. D., Pattyn, F., Popov, S. V., Kusk-Gillespie, M., Rignot, E., Rippin, D. M., Rivera, A., Ross, N., Ruppel, A., Schroeder, D. M., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tabacco, I., Tinto, B. K., Urbini, S., Vaughan, D. G., Welch, B. C., Wilson, D., Young, D. A., Zirizzotti, A., 940 Aitken, A., Roberts, R., Kirkham, J. D., Dorschel, B., Field, E., Hoffman, A. O., Horlings, A. N., and Moholdt, G.: BEDMAP3 – Ice thickness, bed and surface elevation for Antarctica – gridded products (Version 1.0), NERC EDC UK Polar Data Centre [data set], <https://doi.org/10.5285/2d0e4791-8e20-46a3-80e4-f5f6716025d2>, 2024
- Pritchard, H.D., Fretwell, P.T., Fremand, A.C., Bodart, J.A., Kirkham, J.D., Aitken, A., Bamber, J., Bell, R., Bianchi, C., Bingham, R.G., Blankenship, D.D., Casassa, G., Christianson, K., Conway, H., Corr, H.F.J., Cui, X., Damaske, D.,

- 945 Damm, V., Dorschel, B. and Drews, R.: Bedmap3 updated ice bed, surface and thickness gridded datasets for Antarctica. *Sci. Data*, 12, 1–20, <https://doi.org/10.1038/s41597-025-04672-y>, 2025.
- Purich, A. and Doddridge, E.W.: Record low Antarctic sea ice coverage indicates a new sea ice state, *Commun. Earth Environ.*, 4, 1–9, <https://doi.org/10.1038/s43247-023-00961-9>, 2023.
- Ramanath, S., Krieger, L., Floricioiu, D., Diaconu, C.-A., & Heidler, K.: Automatic grounding line delineation of DInSAR
950 interferograms using deep learning, *The Cryosphere*, 19, 2431–2455, <https://doi.org/10.5194/tc-19-2431-2025>, 2025.
- Raphael, M.N., Maierhofer, T.J., Fogt, R.L., Hobbs, W.R. and Handcock, M.S.: A twenty-first century structural change in Antarctica’s sea ice system, *Commun. Earth Environ.*, 6, 1–9, <https://doi.org/10.1038/s43247-025-02107-5>, 2025.
- Reese, R., Gudmundsson, G. H., Levermann, A., and Winkelmann, R.: The far reach of ice-shelf thinning in Antarctica, *Nat. Clim. Change*, 8, 53–57, <https://doi.org/10.1038/s41558-017-0020-x>, 2018.
- 955 Ribeiro, N., Herraiz-Borreguero, L., Rintoul, S. R., McMahon, C. R., Hindell, M., Harcourt, R., and Williams, G.: Warm Modified Circumpolar Deep Water Intrusions Drive Ice Shelf Melt and Inhibit Dense Shelf Water Formation in Vincennes Bay, East Antarctica, *J. Geophys. Res.-Oceans*, 126, 1–17, <https://doi.org/10.1029/2020JC016998>, 2021.
- Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H., and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophys. Res. Lett.*, 41, 3502–
960 3509, <https://doi.org/10.1002/2014GL060140>, 2014.
- Rignot, E., Mouginot, J., and Scheuchl, B.: MEaSUREs Antarctic Grounding Line from Differential Satellite Radar Interferometry, Version 2. NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/IKBWW4RYHF1Q>, 2016.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic
965 Ice Sheet mass balance from 1979–2017, *P. Natl. Acad. Sci. USA*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- Rignot, E., Mouginot, J., Scheuchl, B. and Jeong, S.: Changes in Antarctic ice sheet motion derived from satellite radar interferometry between 1995 and 2022, *Geophys. Res. Lett.*, 49, 1–10, <https://doi.org/10.1029/2022gl100141>, 2022.
- Rignot, E., Millan, R., Charrassin, R. and Scheinert, M.: Bathymetry of the Antarctic continental shelf and ice shelf cavities
970 from a 3D inversion of circumpolar gravity anomalies constrained by other data, *Dryad* [data set], <https://doi.org/10.5061/dryad.rbnzs7hkc>, 2024.
- Rignot, E., Scheuchl, B., Barre, J.B., Brancato, V., Charrier, L., Chan, H., Ciraci, E., Dinh, A., Herreid, S., Jeong, S., Li, X., Mitchell, T., Mohajerani, Y., Shamsian, S., Tolpeki, V., Velicogna, I. and Wollersheim, M. Thirty years of glacier grounding line retreat in Antarctica. *Proc. Natl. Acad. Sci. U.S.A.*, 123, e2524380123,
975 <https://doi.org/10.1073/pnas.2524380123>, 2026.
- Rintoul, S.R., Silvano, A., Pena-Molino, B., van Wijk, E., Rosenberg, M., Greenbaum, J.S. and Blankenship, D.D.: Ocean heat drives rapid basal melt of the Totten Ice Shelf, *Sci. Adv.*, 2(12), e1601610. <https://doi.org/10.1126/sciadv.1601610>, 2016.

- Roquet, F., Williams, G., Hindell, M.A., Harcourt, R., McMahon, C., Guinet, C., Charrassin, J.-B., Reverdin, G., Boehme, L.,
 980 Lovell, P. and Fedak, M.: A Southern Indian Ocean database of hydrographic profiles obtained with instrumented
 elephant seals, *Sci. Data*, 1, 140028, <https://doi.org/10.1038/sdata.2014.28>, 2014.
- Scambos, T. A., Bell, R. E., Alley, R. B., Anandakrishnan, S., Bromwich, D. H., Brunt, K., Christianson, K., Creyts, T., Das,
 S.B., DeConto, R., Dutrieux, P., Fricker, H. A., Holland, D., MacGregor, J., Medley, B., Nicolas, J. P., Pollard, D.,
 Siegfried, M. R., Smith, A. M., Steig, E. J., Trusel, L. D., Vaughan, D. G., and Yager, P. L.: How much, how fast?:
 985 A science review and outlook for research on the instability of Antarctica's Thwaites Glacier in the 21st century,
Global Planet. Change, 153, 16–34, <https://doi.org/10.1016/j.gloplacha.2017.04.008>, 2017.
- Schodlok, M. P., D. Menemenlis, and E. J. Rignot: Ice shelf basal melt rates around Antarctica from simulations and
 observations, *J. Geophys. Res. Oceans*, 121, 1085–1109, <https://doi.org/10.1002/2015JC011117>, 2016.
- Schröder, L., Horwath, M., Dietrich, R., Helm, V., van den Broeke, M. R., and Ligtenberg, S. R. M.: Four decades of Antarctic
 990 surface elevation changes from multi-mission satellite altimetry, *The Cryosphere*, 13, 427–449,
<https://doi.org/10.5194/tc-13-427-2019>, 2019a.
- Schröder, L., Horwath, M., Dietrich, R., Helm, V., van den Broeke, M. R., and Ligtenberg, S. R. M.: Gridded surface elevation
 changes from multi-mission satellite altimetry 1978-2017. PANGAEA [data set]
<https://doi.org/10.1594/PANGAEA.897390>, 2019b.
- 995 Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and
 Engdahl, M. E.: Trends in Antarctic Ice Sheet Elevation and Mass, *Geophys. Res. Lett.*, 46, 8174–8183,
<https://doi.org/10.1029/2019GL082182>, 2019.
- Silvano, A., Rintoul, S. and Herraiz-Borreguero, L.: Ocean-Ice Shelf Interaction in East Antarctica, *Oceanography*, 29, 130–
 143, <https://doi.org/10.5670/oceanog.2016.105>, 2016.
- 1000 Silvano, A., Rintoul, S.R., Peña-Molino, B., and Williams, G.D.: Distribution of water masses and meltwater on the continental
 shelf near the Totten and Moscow University ice shelves, *J. Geophys. Res.-Oceans*, 122, 2050–2068,
<https://doi.org/10.1002/2016jc012115>, 2017.
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho,
 B., Harbeck, K., Markus, T., Neumann, T., Siegfried, M. R., and Zwally, H. J.: Pervasive ice sheet mass loss reflects
 1005 competing ocean and atmosphere processes, *Science*, 368, 1239–1242, <https://doi.org/10.1126/science.aaz5845>,
 2020a.
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho,
 B., Harbeck, K., Markus, T., Neumann, T., Siegfried, M. R., and Zwally, H. J.: Ice-sheet height and thickness changes
 from ICESat to ICESat-2. ResearchWorks Archive [data set] <http://hdl.handle.net/1773/45388>, 2020b.
- 1010 Sohail, T., Zika, J. D., and Ehmen, T.: How accurate are salinity measurements around Antarctica? A machine learning based
 approach. *Machine Learning: Earth*, 2, 025001, <https://doi.org/10.1088/3049-4753/ae7113>, 2026.

- Spreen, G., Kaleschke, L. and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels. *Journal of Geophysical Research*, 113, 1–14, <https://doi.org/10.1029/2005jc003384>, 2008.
- 1015 Stokes, C. R., Abram, N. J., Bentley, M. J., Edwards, T. L., England, M. H., Foppert, A., Jamieson, S. S. R., Jones, R. S., King, M. A., Lenaerts, J. T. M., Medley, B., Miles, B. W. J., Paxman, G. J. G., Ritz, C., van de Flierdt, T., and Whitehouse, P. L.: Response of the East Antarctic Ice Sheet to past and future climate change, *Nature*, 608, 275–286, <https://doi.org/10.1038/s41586-022-04946-0>, 2022.
- Stokes, C.R., Bamber, J.L., Dutton, A., and DeConto, R.M.: Warming of +1.5 °C is too high for polar ice sheets. *Communications Earth and Environment*, 6, 351, <https://doi.org/10.1038/s43247-025-02299-w>, 2025.
- 1020 Surawy-Stepney, T., Hogg, A. E., Cornford, S. L., Wallis, B. J., Davison, B. J., Selley, H. L., Slater, R. A. W., Lie, E. K., Jakob, L., Ridout, A., Gourmelen, N., Freer, B. I. D., Wilson, S. F., and Shepherd, A.: The effect of landfast sea ice buttressing on ice dynamic speedup in the Larsen B embayment, Antarctica, *The Cryosphere*, 18, 977–993, <https://doi.org/10.5194/tc-18-977-2024>, 2024.
- Teder, N.J., Bennetts, L.G., Reid, P.A., Massom, R.A., Pitt, J.P., Scambos, T.A., and Fraser, A.D.: Large-scale ice-shelf calving events follow prolonged amplifications in flexure. *Nat. Geosci.*, 18, 599–606, <https://doi.org/10.1038/s41561-025-01713-4>, 2025.
- 1025 Thoma, M., Jenkins, A., Holland, D., and Jacobs, S.: Modelling Circumpolar Deep Water intrusions on the Amundsen Sea continental shelf, Antarctica, *Geophys. Res. Lett.*, 35, L18602, <https://doi.org/10.1029/2008GL034939>, 2008.
- Thompson, S. S., Kulesa, B., Luckman, A., Halpin, J. A., Greenbaum, J. S., Pelle, T., Habbal, F., Guo, J., Jong, L. M., Roberts, J. L., Sun, B., and Blankenship, D. D.: Glaciological history and structural evolution of the Shackleton Ice Shelf system, East Antarctica, over the past 60 years, *The Cryosphere*, 17, 157–174, <https://doi.org/10.5194/tc-17-157-2023>, 2023.
- 1030 Treasure, A., Roquet, F., Ansorge, I., Bester, M., Boehme, L., Bornemann, H., Charrassin, J.-B., Chevallier, D., Costa, D., Fedak, M., Guinet, C., Hammill, M., Harcourt, R., Hindell, M., Kovacs, K., Lea, M.-A., Lovell, P., Lowther, A., Lydersen, C. and McIntyre, T.: Marine Mammals Exploring the Oceans Pole to Pole: A Review of the MEOP Consortium, *Oceanography*, 30, 132–138, <https://doi.org/10.5670/oceanog.2017.234>, 2017.
- van den Broeke, M.: Strong surface melting preceded collapse of Antarctic Peninsula ice shelf, *Geophys. Res. Lett.*, 32, 1–4, <https://doi.org/10.1029/2005GL023247>, 2005.
- Wang, L., Davis, J. L., and Howat, I. M.: Complex Patterns of Antarctic Ice Sheet Mass Change Resolved by Time-Dependent Rate Modelling of GRACE and GRACE Follow-On Observations, *Geophys. Res. Lett.*, 48, e2020GL090961, <https://doi.org/10.1029/2020GL090961>, 2021.
- 1040 Walker, C.C., Millstein, J.D., Miles, B.W.J., Cook, S., Fraser, A.D., Colliander, A., Misra, S., Trusel, D., Adusumilli, S., Roberts, C., and Fricker, H.: Multi-decadal collapse of East Antarctica’s Conger–Glenzer Ice Shelf, *Nat. Geosci.*, 17, 1240–1248, 2024.

- 1045 Wong, A. P. S., Wijffels, S. E., Riser, S. C., Pouliquen, S., Hosoda, S., Roemmich, D., Gilson, J., Johnson, G. C., Martini, K.,
Murphy, D. J., Scanderbeg, M., Bhaskar, T. V. S. U., Buck, J. J. H., Merceur, F., Carval, T., Maze, G., Cabanes, C.,
André, X., Poffa, N., Yashayaev, I., Barker, P. M., Guinehut, S., Belbéoch, M., Ignaszewski, M., Baringer, M. O.,
Schmid, C., Lyman, J. M., McTaggart, K. E., Purkey, S. G., Zilberman, N., Alkire, M. B., Swift, D., Owens, W. B.,
Jayne, S. R., Hersh, C., Robbins, P., West-Mack, D., Bahr, F., Yoshida, S., Sutton, P. J. H., Cancouët, R., Coatanoan,
1050 C., Dobbler, D., Juan, A. G., Gournion, J., Kolodziejczyk, N., Bernard, V., Bourlès, B., Claustre, H., D’Ortenzio, F.,
Le Reste, S., Le Traon, P.-Y., Rannou, J.-P., Saout-Grit, C., Speich, S., Thierry, V., Verbrugge, N., Angel-Benavides,
I. M., Klein, B., Notarstefano, G., Poulain, P.-M., Vélez-Belchí, P., Suga, T., Ando, K., Iwasaka, N., Kobayashi, T.,
Masuda, S., Oka, E., Sato, K., Nakamura, T., Sato, K., Takatsuki, Y., Yoshida, T., Cowley, R., Lovell, J. L., Oke, P.
R., van Wijk, E. M., Carse, F., Donnelly, M., Gould, W. J., Gowers, K., King, B. A., Loch, S. G., Mowat, M., Turton,
1055 J., Rama Rao, E. P., Ravichandran, M., Freeland, H. J., Gaboury, I., Gilbert, D., Greenan, B. J. W., Ouellet, M., Ross,
T., Tran, A., Dong, M., Liu, Z., Xu, J., Kang, K., Jo, H., Kim, S.-D., and Park, H.-M.: Argo data 1999–2019: two
million temperature-salinity profiles and subsurface velocity observations from a global array of profiling floats,
Front. Mar. Sci., 7, 700, <https://doi.org/10.3389/fmars.2020.00700>, 2020.
- Weatherley, M., Stokes, C., Jamieson, S., Ramanath Tarekere, S., and Silvano, A.: Porpoise Bay Glacier Change - GIS files
1060 (Version 1.0), Zenodo [data set], <https://doi.org/10.5281/zenodo.15847590>, 2025.
- Yamazaki, K., Aoki, S., Katsumata, K., Hirano, D., and Nakayama, Y.: Multidecadal poleward shift of the southern boundary
of the Antarctic Circumpolar Current off East Antarctica, Sci. Adv., 7, eabf8755,
<https://doi.org/10.1126/sciadv.abf8755>, 2021.
- Young, D. A., Wright, A. P., Roberts, J. L., Warner, R. C., Young, N. W., Greenbaum, J. S., Schroeder, D. M., Holt, J. W.,
1065 Sugden, D. E., Blankenship, D. D., van Ommen, T. D., and Siegert, M. J.: A dynamic early East Antarctic Ice Sheet
suggested by ice-covered fjord landscapes, Nature, 474, 72–75, <https://doi.org/10.1038/nature10114>, 2011.
- Zhu, Y., Hogg, A. E., Hooper, A., and Wallis, B. J.: Short and Long-term Grounding Zone Dynamics of Amery Ice Shelf, East
Antarctica, The Cryosphere, 19, 3971–3989, <https://doi.org/10.5194/tc-19-3971-2025>, 2025.