

On the non-linear response of Antarctic ice shelf surface melt to warming

Marte Gé Hofsteenge¹, Willem Jan van de Berg¹, Christiaan van Dalum^{1,2}, Kristiina Verro^{1,3}, Maurice van Tiggelen¹, and Michiel van den Broeke¹

¹Institute for Marine and Atmospheric Research (IMAU), Utrecht University, Utrecht, the Netherlands

²Royal Netherlands Meteorological Institute, De Bilt, the Netherlands

³National Centre for Climate Research, Danish Meteorological Institute, Copenhagen, Denmark

Correspondence: Marte Gé Hofsteenge (m.g.hofsteenge@uu.nl)

Abstract. Surface meltwater can saturate firn, form melt ponds, and trigger hydrofracturing of Antarctic ice shelves, ultimately accelerating grounded ice flow and contributing to sea level rise. Although the response of surface melt to atmospheric warming (expressed by near-surface air temperature) is known to be non-linear, the mechanisms driving this non-linearity remain poorly understood. In this study we explain the non-linear temperature-melt relationship from an energy balance perspective and assess its spatial variability across Antarctic ice shelves. We use the regional climate model RACMO2.4p1, forced by ERA5 re-analysis and two global earth system models under the SSP3-7.0 high emission scenario, to simulate contemporary and future Antarctic climate and surface mass balance until 2100. We find that the temperature dependence of net shortwave radiation is the primary driver of the non-linearity. ~~On relatively cold ice shelves, warming on ice shelves in relatively dry climates. Warming~~ increases cloud cover and snowfall, ~~raising which both raise~~ albedo, reducing net shortwave radiation. ~~In contrast, on warmer ice shelves~~ When summer air temperatures approach -12°C, the snowmelt-albedo feedback dominates the response: warming leads to melt that reduces albedo, enhancing shortwave radiation absorption. The temperature-melt relationship also varies spatially: ice shelves in drier regions experience more melt at the same average summer temperatures than those in wetter regions, highlighting the role of snowfall in suppressing the albedo feedback. In wetter regions, elevated humidity and cloudiness enhance the sensitivity of melt to warming through changes in net longwave radiation, rather than albedo changes. When mean summer air temperatures reach or exceed the melting point (0 °C), ice shelves ~~becomes~~ become even more sensitive to warming. Surface temperatures can not rise above 0°C while the atmosphere can, allowing the sensible heat and net longwave radiation to increase. At the same time, snowfall transitions to rain, amplifying the albedo feedback. Our results suggest that currently colder, drier and stable ice shelves could experience rapid increases in melt under future warming, with implications for their long-term stability.

20 1 Introduction

Surface melt plays a critical role in the stability of Antarctic ice shelves, which act as barriers slowing the flow of inland ice into the ocean (Fürst et al., 2016). Episodes of intense or prolonged surface melting can lead to the formation of melt ponds on the surface (Kingslake et al., 2017), which may trigger hydrofracture and rapid ice shelf collapse (Kuipers Munneke et al., 2014; Lai et al., 2020). Such collapses reduce ice shelf buttressing, accelerate flow and mass loss of grounded ice and result in sea level rise (Bell et al., 2018). Whether melt ponds might form on the ice shelves depends on the balance between snow accumulation and snow melt: regions with low snowfall accumulation have limited capacity to store meltwater before saturation and ponding occurs (van Wessem et al., 2023; Kuipers Munneke et al., 2014). Surface melt and runoff from Antarctic ice shelves ~~is~~ are projected to increase with warming (Gilbert and Kittel, 2021; Kittel et al., 2021; Ligtenberg et al., 2013). However, the ~~relationship between temperature and melt~~ response of melt to warming is highly non-linear (Abram et al., 2013), resulting in a large spread of predicted melt rates at the end of the 21st century between different future scenarios (Trusel et al., 2015; Seroussi et al., 2020). The processes that drive this non-linearity are not yet fully understood.

Current surface melt rates over Antarctic ice shelves can be estimated from satellite products (Di Biase et al., 2026; Zheng et al., 2025) and regional climate models. Polar-adapted regional climate models (RCMs), such as the polar (p) version of the Regional Atmospheric Climate Model (RACMO), are currently among the most effective tools for quantifying current and predicting future Antarctic surface melt. These models include snowpack physics and surface energy balance (SEB) schemes to simulate key surface processes driving melt, and are typically run at much higher spatial resolutions (1-20 km) than global Earth system models (~ 100 km). However, due to the computational costs, RCMs are not typically used to estimate melt in large ensembles of future climate scenarios, such as those from CMIP, or coupled to ice sheet models to link the surface processes and ice dynamics. Instead, studies propose simplified empirical relationships or melt potential indices based solely on near-surface air temperature (Orr et al., 2023; Trusel et al., 2015; Vaughan, 2006; Zheng et al., 2023). These approaches rely on the fact that many SEB components are themselves dependent on temperature, such as incoming longwave radiation and sensible heat flux (Ambach, 1988; Braithwaite and Olesen, 1990; Ohmura, 2001). While widely used (Coulon et al., 2024; DeConto et al., 2021; Golledge et al., 2019), empirical relationships between temperature and melt such as positive degree day (PDD) models or temperature-melt index models, are often applied uniformly across space. This is despite the known spatial variability in melt sensitivity to temperature, which arises from differences in local surface conditions, cloudiness, and SEB regimes across Antarctic ice shelves (van den Broeke et al., 2023; Zheng et al., 2023).

The ~~strong~~ relationship between summer near-surface air temperature and surface melt on Antarctic ice shelves ~~can be characterized by an exponential relationship~~ has an exponential shape (Trusel et al., 2015; van Wessem et al., 2023). This means that for ice shelves already experiencing melt, even a small temperature increase can lead to a disproportionately large increase in melt, potentially reaching levels of melt at which ice shelves have collapsed in the past (Scambos et al., 2000; Trusel et al., 2015; van den Broeke, 2005). ~~The strong~~ One of the possible explanations for this non-linearity ~~in this~~

relationship has been linked to the snowmelt–albedo feedback (Jakobs et al., 2019), but the extent to which this mechanism, or other temperature-sensitive SEB components contribute to is the snowmelt-albedo feedback (Jakobs et al., 2019, 2021) in which melt and subsequent refreezing lowers the albedo of snow, increasing absorption of solar radiation and melt. The potential of this feedback to enhance surface melt is modulated by the frequency and timing of snowfall events in summer. But as noted by Jakobs et al. (2019), under warmer conditions such as those currently observed on the Antarctic Peninsula, the observed non-linearity remains unclear: snowmelt-albedo feedback becomes less important in enhancing melt, and other processes such as exposure of bare ice or turbulent fluxes also play a role. Other surface energy balance terms such as longwave radiation and the latent heat flux may also respond non-linearly to air temperature through changes in atmospheric moisture and cloud conditions that are associated with the atmospheric warming.

In this study we use output from RACMO to investigate the physical processes driving the non-linearity of the relationship between summer air temperature and surface melt across Antarctic ice shelves. We analyse both historical and future simulations to 1) assess the spatial variability in the temperature-melt relationship and 2) identify the dominant SEB components contributing to the non-linearity. Our findings will provide new insights into the physical controls on melt sensitivity and provide guidance on when and where temperature-based approaches to estimate surface melt are appropriate, and where they lack reliability due to more complex local SEB conditions.

2 Methods

2.1 The regional climate model RACMO

The Regional Atmospheric Climate Model (RACMO) is a hydrostatic regional climate model developed at the Royal Netherlands Meteorological Institute (KNMI). It incorporates atmospheric dynamics from HIRLAM (Undén et al., 2002), which uses a semi-Lagrangian approach with semi-implicit time stepping, and physical parameterizations from the ECMWF Integrated Forecasting System (IFS) (ECMWF, 2020). We use the polar version of RACMO which is developed and maintained at the Institute for Marine and Atmospheric Research Utrecht (IMAU). This version is specifically adapted for simulating the climate and surface processes of the polar regions and includes a multi-layer snow/firn model applied to glaciated surface tiles. The snow model simulates key processes to represent the surface energy and mass balance of snow, such as metamorphism, compaction, melt and refreezing of snow. In this study, we used the latest RACMO version (2.4p1) (van Dalum et al., 2024), in this paper referred to as RACMO. The main differences between RACMO2.4p1 and the previous operational version, RACMO2.3p2, are (van Dalum et al., 2024):

1. Update of IFS physics parameterizations from ECMWF cycle 33r1 to 47r1 (ECMWF, 2020), which includes improvements of cloud and precipitation physics such as a better representation of mixed phase clouds, separate prognostics variables for cloud water/ice, rain and snow that allows for advection of precipitation, the radiation scheme is replaced by ecRad,

aerosols prescription with Copernicus Atmospheric Monitoring Service (CAMS), and improved cloud optical properties from the Suite Of Community RAdiative Transfer codes based on Edwards and Slingo (SOCRATES). Additional processes that are included in the change to IFS cycle 47r1 are described in van Dalum et al. (2024);

2. Fractional ice cover, based on the BedMachine Antarctica version 3 ice mask (Morlighem et al., 2020), where ~~part of~~ grid cells can be partially glaciated and therefore ~~better-representing~~ provide a better representation of areas such as the McMurdo Dry Valleys;
3. A narrowband snow albedo model, which was introduced in the non-operational RACMO version 2.3p3 (van Dalum et al., 2022). This new albedo model explicitly resolves radiation penetration and subsurface heating in snow and ice. Because albedo is calculated wavelength-dependent, it captures the spectral effect of clouds on albedo. Cloud cover scatters much of the near-infrared radiation, which has a low albedo, leaving mostly visible light, for which the spectral albedo is higher (van Dalum et al., 2020).;
4. An updated snowdrift model (Gadde and van de Berg, 2024)

2.2 Surface energy balance

Within the updated snow model, shortwave radiation can penetrate ~~in~~ the snowpack and be absorbed in the subsurface. The total net absorbed shortwave radiation (SW_{net}) is thus split up in surface absorption ($SW_{net, surf}$) and internal absorption (SW_{pen}). Internal shortwave absorption leads to warming of the snowpack and internal melt when the snow reaches the melting point. As a result, melt can occur both at the surface and in the snowpack. The SEB of the skin layer is defined as:

$$Q_{M, surf} = LW_{net} + SW_{net, surf} + SH + LH + Q_G \quad (1)$$

where $Q_{M, surf}$ is the energy ~~used~~ available for surface melt, LW_{net} is net longwave radiation, SH and LH represent the turbulent fluxes of sensible and latent heat and Q_G is the subsurface conductive heat flux. All fluxes are defined as positive towards the surface and have as unit $W m^{-2}$.

Since we are interested in understanding the relationship between temperature and total melt; i.e. occurring both at the surface as well as in the subsurface, we use a pseudo-SEB for the analysis which is not strictly valid for a skin layer as in Eq. (1), but for the near-surface. This definition is comparable to previous SEB formulations, in which penetration of solar radiation was not included, and is as follows:

$$Q_M = LW_{net} + SW_{net} + SH + LH + Q_{G, rec}, \quad (2)$$

Where Q_M is the energy ~~used~~ available for melt (surface + internal melt) and $Q_{G, rec}$ is a reconstructed conductive heat flux defined as $Q_{G, rec} = Q_G + Q_{m, internal} - SW_{pen}$, with $Q_{m, internal}$ energy ~~used~~ available for internal melt. This formulation assumes

115 that, on seasonal timescales, the effects of short-term heat storage and energy redistribution in the near-surface are negligible, such that total SWnet (surface + penetrated) can be used as the radiative driver of total melt. In Appendix A we show that this pseudo-SEB produces comparable SEB terms to the classical formulation without shortwave penetration, supporting its use for the analysis in this study.

120 The resulting melt rate (kg m^{-2}) is then obtained by converting the available melt energy into a meltwater mass flux using the latent heat of fusion, such that the energy remaining after warming the snow to the melting point produces melt. Melt contributes to mass loss when the resulting meltwater is not refrozen or retained in the snow and instead leads to runoff (RU). The surface mass balance (SMB) is defined as the difference describes the balance between accumulation and ablation in the near-surface firn or ice:

$$125 \text{ SMB} = P_{tot} - \text{SU}_s - \text{SU}_{ds} - \text{RU} - \text{ER} \quad (3)$$

with P_{tot} total precipitation, SU_s is surface sublimation and SU_{ds} sublimation of blowing snow, RU runoff and ER drifting snow erosion which can be both ablative (erosion, $\text{ER} > 0$) and accumulative (deposition). ~~RU is simulated when the snow layers become saturated with meltwater or rain.~~, $\text{ER} < 0$.

2.3 Historical and future forcing

130 We use RACMO2.4p1 with a domain covering Antarctica and the southern tip of South America with a horizontal resolution of 11 km, forced with both ERA5 reanalysis and Earth-System-Model (ESM) datasets. Using ERA5 provides present-day conditions, while the ESM-forced simulations extend the analysis into future climates, allowing the temperature–melt relationship to be examined across a wider range of melt intensities and summer temperatures than occur today. The simulations for this domain ~~have been done~~ were performed as part of the PolarRES project, an EU Horizon 2020 funded project that uses RCMs
135 to simulate the current and future climate of the polar regions. ~~We use the RACMO simulation forced by ERA5 reanalysis data (1979–2023), previously evaluated with observations by van Dalum et al. (2025), as reference simulation for the historical period, which is hereafter referred to as RACMO(ERA5). Future projection simulations ((Gilbert et al., 2025b). Projections spanning 2015 –to 2099) were performed~~ were forced using boundary conditions from two CMIP6 ESMs: the Community Earth System Model 2 (CESM2) and the Max-Planck Institute Earth System model (MPI-ESM), under the high emission
140 SSP3-7.0 scenario.

~~As a second evaluation step, historical simulations (1985–2014) with these ESMs are compared to RACMO(ERA5). The RACMO simulations with ESM forcing are hereafter referred to as RACMO(SSP3-7.0 was chosen within the PolarRES framework because it is considered a more plausible high-emission pathway than SSP5-8.5, CESM2) and RACMO(MPI-ESM). CESM2 and MPI-ESM were selected within the PolarRES project MPI-ESM were then selected~~ from the CMIP6 ESMs using a
145 storyline approach to represent two contrasting but plausible Antarctic climate futures; CESM2 reflects a future with extensive sea ice loss and an earlier summertime stratospheric polar vortex breakdown, while MPI-ESM captures a scenario with limited

sea ice loss and a delayed polar vortex breakdown (Williams et al., 2024).

We use the RACMO simulation forced by ERA5 reanalysis data (Hersbach et al., 2017), previously evaluated with observations by van Dalum et al. (2025), as reference simulation for the historical period (1979–2023), hereafter referred to as RACMO(ERA5). At the lateral boundaries of the modelling domain, windspeed, temperature, pressure and humidity are specified using multi-level data from either ERA5 or ESM output. Sea surface temperature and sea-ice cover are defined at the ocean surface boundary and prescribed from ERA5 or ESM output. Sea ice temperature is calculated by a using the four-layer sea ice slab model with a fixed from the ECMWF IFS model, which assumes a fixed maximum thickness of 1.5 m. Furthermore, in the free atmosphere the temperature, wind field and humidity are gently relaxed to either the ERA5 or ESM output, following van de Berg and Medley (2016).

3 Results

2.1 Evaluation of RACMO historical simulations with ESM forcings

RACMO(ERA5) has been extensively evaluated against weather station and mass balance observations in van Dalum et al. (2025). Using observations from automatic weather stations on the Antarctic Peninsula and in Dronning Maud Land, this study shows that version 2.4p1 performs well in simulating Antarctica's near-surface air temperature (bias of -1.40 and an RMSE of 4.38 $^{\circ}\text{C}$) and shortwave radiation (bias of 8.5 and -8.8 W m^{-2} for downward and upward shortwave radiation, respectively), but has larger differences with observations for longwave radiative fluxes (bias of -20.4 and 11.7 W m^{-2} for downward and upward longwave radiation, respectively). Turbulent fluxes have small bias (-0.3 and 1.5 W m^{-2} for latent and sensible heat flux, respectively), but large spread (RMSE of 5.0 and 14.2 W m^{-2} , respectively). As the biases in longwave and shortwave radiation partially offset one another, the resulting melt rates are less affected. This is reflected in the good agreement between simulated meltwater presence in the snow and satellite-based estimates (van Dalum et al., 2025).

Before using RACMO for future scenario simulations, we need confidence in the model performance when forced by ESMs. To assess this, we compare the average climate in the historical simulation As a second evaluation step, historical simulations (1985–2014) with that in forced with ESMs are compared to RACMO(ERA5), since direct comparison with in situ observations is not possible for ESM output unconstrained by data assimilation. in Section 3.1. We evaluate whether the mean difference between the ESM-forced simulations and RACMO(ERA5) has been extensively evaluated against weather station and mass balance observations (?). The updated version 2.4p1 performs well in simulating Antarctica exceed the inter-annual standard deviation of RACMO(ERA5). The RACMO simulations forced by CESM2 and MPI-ESM are hereafter referred to as RACMO(CESM2) and RACMO(MPI-ESM), respectively.

2.1 Assessing temperature-dependency of SEB

180 To study the drivers of the non-linear response of melt to temperature, we assess how the individual SEB terms respond to temperature. Because this response may itself be non-linear, each SEB-temperature slope is evaluated across discrete temperature bins. As albedo and cloud feedbacks are expected to play a dominant role, we first assess their temperature sensitivity. The influence of clouds on albedo is examined by comparing clear-sky and all-sky albedo, making use of the fact that RACMO separately calculates radiative fluxes under clear-sky and cloudy-sky conditions and subsequently combines them
185 into total-sky fluxes based on cloud fraction. Based on this analysis, the following SEB analysis is split between ice shelves in relatively dry and wet climates, based on a threshold in annual snowfall. Following van Wessem et al. (2023), we use the median annual snowfall of $\sim 500 \text{ mm yr}^{-1}$ as the threshold separating ‘dry’ s-surface mass-balance, temperature, albedo, and shortwave radiation, whereas longwave radiation and turbulent fluxes still require improvement. Figure ?? ‘wet’ ice shelves, which results in two equally sized groups. A map with overview of the ice shelves that are classified in each category is given
190 in Fig. B1.

To systematically assess the temperature dependence of SEB terms and explain the non-linear temperature–melt relationship, we use the following approach. For each ice shelf grid cell that experiences an average summer melt of at least 1 mm during the historical period, we fit either a linear or exponential function relating the summer average SEB component to air temperature.
195 The function with the higher R^2 value is selected, and grid cells where the optimal fit parameters could not be reliably estimated (average $R^2 < 0.3$) are excluded from the analysis. We then calculate the average slope of the selected fits across all grid cells, grouped into temperature bins of 2°C . If the selected fit is exponential, the slope at the middle of the bin is taken. The resulting average slopes vary across temperature bins because each bin averages fits from different locations. We calculate uncertainty bands based on the standard deviation between the fits (reflecting variability between locations) weighted with the average R^2
200 value (reflecting the strength of the fit).

3 Results

First, the performance of RACMO forced by the ESMs is evaluated to gain confidence in the simulations before using them in the analysis. We then assess the temperature and melt trends in the future projection runs, which provide the basis for studying the temperature–melt relationship. Finally, we analyse the spatial variability in the relationship between temperature and melt, the role of albedo feedbacks in the non-linearity and systematically assess how all SEB terms depend on temperature and contribute to the melt response.

3.1 Evaluation of RACMO historical simulations with ESM forcings

Because ESM-forced simulations are not constrained by data assimilation, we evaluate the ESM-forced simulations by comparing the mean and variability of near-surface climate variables over 1985–2014 with those from RACMO(ERA5) for the same period. Figure 1 shows the differences in annual-DJF means of several key variables between simulations forced by ESMs and forced by ERA5. In the figure, hatching indicates areas where the mean difference between the ESM-forced simulations and RACMO(ERA5) is larger than the ERA5 interannual standard deviation over the historical period. RACMO(MPI-ESM) is warmer compared to RACMO(ERA5), especially over the sea ice zone and in the Ross Sea and Wedell Sea sectors (Fig. 1a). This coincides with the lower sea ice extent in RACMO(MPI-ESM) compared to RACMO(ERA5) (Fig. 1b). While the warm bias is smaller over the ice, the influence of the lower sea ice extent in RACMO(MPI-ESM) on air temperature becomes even clearer when examining the annual mean fields (Fig. C1). Over the Antarctic continent, RACMO(MPI-ESM) is still significantly warmer than temperatures differ little from RACMO(ERA5) in relative to the year-to-year variability. Exceptions are Dronning Maud Land, the coast of Marie Byrd Land and other coastal regions in East Antarctica which is warmer, and the high interior plateau in East Antarctica, where temperatures are lower than RACMO(ERA5) by more than the inter-annual standard deviation. The temperatures and sea ice conditions in RACMO(CESM2) are more realistic, but the simulations have show a cold bias over West Antarctica and large parts of East-Antarctica, where the 500 hPa geopotential height is lower compared to RACMO(ERA5) (contours in Fig. 1g).

RACMO(CESM2) and RACMO(MPI-ESM) have similar SMB-anomaly-precipitation difference patterns, with more positive SMB-precipitation in Dronning Maud Land and a more negative SMB in West less in other parts of coastal East Antarctica compared to RACMO(ERA5) (Fig. 1c,g), driven by differences in precipitation (Fig. 1d,h). The negative SMB anomaly in East-Antarctica in RACMO(CESM2), which is due to an underestimation of precipitation, is not present in RACMO(MPI-ESM). Over the Antarctic Peninsula, RACMO(MPI-ESM) simulates less precipitation on the western side and more on the eastern side compared to RACMO(ERA5). This pattern is associated with a weaker Amundsen Sea Low, reflected in a positive geopotential height anomaly-difference over the Amundsen-Ross Sea in Fig. 1e. In contrast, RACMO(CESM2) shows lower geopotential heights over the Bellingshausen Sea relative to RACMO(ERA5), which enhances westerly winds across the Antarctic Peninsula and orographic precipitation on the western slopes.




Figure 1. Difference in annual-DJF mean near-surface air temperature, sea ice concentration, surface mass balance, and precipitation and surface melt between RACMO simulations forced by MPI (a-d) or CESM (e-h) and those forced by ERA5 in the period 1985-2014. Hatched areas show where differences are smaller/larger than the standard deviation over the historical period in ERA5. Contour lines Contour lines in c and and g indicate corresponding difference in 500hPa geopotential height with RACMO(ERA5).

In this study we focus on melt, which occurs almost exclusively during the summer (DJF) season, so from here onwards we focus on the simulations in summer. Table C1 compares average summer near-surface air temperature, energy and mass balance terms over the Antarctic ice sheet. The simulations show good agreement in summer air temperature and most SEB terms. The underestimation of sea ice and warm bias over the sea ice zone and coastal regions is much smaller in summer (Appendix Fig. ??) In RACMO(MPI-ESM), melt rates are higher on the western side of the Antarctic Peninsula and lower on the eastern side compared to RACMO(ERA5); however, these differences do not exceed the inter-annual standard deviation. They likely reflect the impact of precipitation differences and albedo changes on SW_{net} (Fig. 2a). The opposite pattern holds for RACMO(CESM2), where melt rates are mostly underestimated on the western Antarctic Peninsula ice shelves, consistent with lower air temperatures and increased precipitation that reduces SW_{net} (Fig. 2e). Outside the Antarctic Peninsula, melt differences in RACMO(MPI-ESM) and RACMO(CESM2) are generally small compared to the annual-mean that was shown in Figure ??.

RACMO2.4p1 simulated summer averages (1985-2014) forced by ERA5, CESM2 and MPI-ESM. The table shows 30-year mean values and standard deviations for ice-sheet average near-surface air temperature, surface energy balance $W m^{-2}$ and ice-sheet integrated average summer totals of mass balance components $Gt summer^{-1}$. RACMO(ERA5) RACMO(CESM2) RACMO(MPI-ESM) $T_{2m}^{\circ}C$ 23.2 inter-annual variability in RACMO(ERA5). The Antarctic-integrated melt in RACMO(MPI-ESM)




Figure 2. As in Fig. 1 but for net shortwave radiation (SW_{net}), net longwave radiation (LW_{net}), sensible and latent heat flux (SH and LH).

is similar to that in RACMO(ERA5) (120 ± 0.8 – 24.0 ± 0.6 – 23.8 ± 0.6 SW_{in} $W m^{-2}$ 349.7 ± 1.9 350.6 ± 1.4 348.7 ± 2.0
 250 LW_{in} $W m^{-2}$ 155.0 ± 2.5 151.3 ± 1.9 154.3 ± 2.0 SW_{net} $W m^{-2}$ 58.4 ± 0.9 57.9 ± 1.3 56.6 ± 1.0 LW_{net} $W m^{-2}$ 63.7 ± 0.9
 – 64.0 ± 0.8 – 62.3 ± 0.9 SH $W m^{-2}$ 9.6 ± 0.5 10.2 ± 0.6 9.9 ± 0.3 LH $W m^{-2}$ 1.9 ± 0.1 1.8 ± 0.1 1.9 ± 0.2 Precipitation
 Gt summer 41 Gt yr^{-1} 565 ± 63 556 ± 68 608 ± 45 Surface sublimation Gt summer $^{-1}$ 70.8 ± 5.5 67.8 ± 5.6 72.1 ± 6.1
 Snowdrift sublimation Gt summer $^{-1}$ 24.8 ± 3.8 28.3 ± 4.2 27.6 ± 2.5 Melt Gt summer $^{-1}$ versus 115 ± 38 Gt yr^{-1} , see
 Appendix Table C1). In RACMO(CESM2) the Antarctic-integrated melt is lower (71 ± 25 120 ± 41 Refreezing Gt summer Gt
 255 yr^{-1} 110 compared to 115 ± 37 67 ± 25 116 ± 39 Runoff Gt summer 38 Gt yr^{-1} 4.3 ± 3.5 4.7 ± 4.0 5.0 ± 3.1 SMB Gt
 summer $^{-1}$ 461 ± 62 455 ± 66 503 ± 47 –

Overall, the dynamically downscaled SEB and SMB in summer from RACMO(CESM2) and RACMO(MPI-ESM) match well with that of RACMO(in ERA5). The largest differences in SMB terms–

260 The differences in SEB components between the simulations lie in melt and precipitation. Melt is comparable between also
 reflect atmospheric circulation-related differences between the ESMs and ERA5 (Fig. 2). In both ESM-forced simulations,
 enhanced moisture transport to Dronning Maud Land reduces SW_{net} and increases LW_{net} compared to RACMO(ERA5) and
 RACMO(MPI-ESM), but is underestimated in RACMO(CESM2), likely due to a cold bias especially in West-Antarctica.
 Precipitation in RACMO(MPI-ESM) is higher than in RACMO(ERA5) but the difference is still smaller than typical variability
 265 in summer precipitation. The difference in precipitation is dominated by positive anomalies on the western side of the Antarctic

Peninsula and over the coastal region of Dronning Maud Land (Appendix Fig. ??) through albedo and cloud effects. Away from Dronning Maud Land, RACMO(CESM2) is drier along much of coastal East-Antarctica compared to RACMO(ERA5), leading to higher SW_{net} and lower LW_{net} . Differences in the turbulent heat fluxes are generally small, except in regions that are influenced by the lower pressure over the Amundsen Sea in RACMO(CESM2). This pressure difference strengthens westerly winds across the Antarctic Peninsula, increasing sensible heating and sublimation through foehn winds on the eastern side, and also enhances the warm and dry downslope flow over the Ross Ice Shelf. Overall, the ESM-forced simulations reproduce Antarctic near-surface climate, SEB and melt patterns well over the ice shelves, with differences relative to RACMO(ERA5) generally within the inter-annual standard deviation, except in a few localized regions such as Wilkins and George VI ice shelves. This close agreement supports the reliability of these simulations for investigating the temperature-melt relationship over the 21st century.

3.2 Historical and future trends in Antarctic warming and surface melt

In our simulations under the SSP3-7.0 scenario, warming of the near-surface atmosphere over the 21st century is strongest in autumn and winter, especially over the sea ice zone (Fig. 3). We find large differences between RACMO(CESM2) and RACMO(MPI-ESM): the former consistently shows stronger warming across all seasons, with the most pronounced differences in autumn and winter that have substantial sea ice decline. Part of the different warming rates stem from the different starting conditions in the historical climate states, with MPI-ESM simulating significantly lower sea ice concentrations and higher air temperatures (Fig. ??b). Another reason for the difference in warming is likely the contrasting atmospheric circulation response of the ESMs. MPI-ESM has a stronger intensification and poleward shift of the jet stream compared to CESM2, which limits the advection of warm air from lower latitudes and therefore suppresses warming, especially in winter (Williams et al., 2024).

The warming over the Southern Ocean is weakest in summer, when sea ice concentration, and thus trends in sea ice concentration, are smallest. The large thermal inertia of the ocean inhibits a fast surface warming and subsequently a quick rise of the near-surface air temperature. Nevertheless, both RACMO(CESM2) and RACMO(MPI-ESM) show warming over the entire Antarctic continent during summer. ~~When focusing on near-surface air temperature over ice shelves, we find good agreement between the historical simulations of near-surface temperature, both in terms of the mean and inter-annual variability (Fig. 4). The largest differences are found on the ice shelves of West Antarctica (Fig. 4g), with RACMO(CESM2) being colder and RACMO(MPI-ESM) warmer compared to RACMO(ERA5).~~

We now focus on Antarctic ice shelves and assess their summer warming and surface melt evolution, which provides the context for the following analyses of the temperature-melt relationship and its physical drivers. For all regions, ice shelves are warming significantly over the 21st century ~~and consistently stronger in~~ (Fig. 4). ~~However, output from RACMO(CESM2) compared to consistently shows a greater rate of warming than~~ RACMO(MPI-ESM). Interestingly, for most regions melt in RACMO(MPI-ESM) and RACMO(CESM2) is more similar in the historical period and diverges in the future scenario, apart




Figure 3. Average near-surface air temperature change between the current climate (1985–2014) and future climate (2070–2099) as modeled by RACMO. Panels (a–d) are forced by MPI-ESM, while panels (e–h) are forced by CESM2. Hatched areas show where changes in temperature are smaller larger than the standard deviation over the historical period in ERA5.

300 from ice shelves in West Antarctica (Fig. 4g) that start with a large difference but converge in the future.

No significant trends in melt are found over the historical period, except for a decrease in melt on the ice shelves of the Antarctic Peninsula in RACMO(ERA5) (Fig. 4d). This agrees with previous studies that report a significant regional cooling since the late 1990s driven by changes in atmospheric circulation and increased sea ice advection (Turner et al., 2016; van
 305 Wessem et al., 2016), which changed in the mid-2010s together with changes in the large-scale climate modes (Carrasco et al., 2021). Melt starts to increase in the future simulations, showing significant increases over all regions. On the Filchner-Ronne and Ross Ice Shelves, the interannual variability relative to the overall melt trend is largest, particularly after 2070. Melt increases more rapidly in RACMO(CESM2) compared to RACMO(MPI-ESM) for all regions, consistent with the stronger temperature increase. As a result, for most ice shelves RACMO(CESM2) starts with lower temperatures and melt, but ends
 310 higher than RACMO(MPI-ESM).

3.3 Spatial variability in temperature-melt relationship

Given the strong link between temperature and melt suggested by previous research (e.g. van Wessem et al. (2023); Trusel et al. (2015)) (Trusel et al., 2015) and the future trends (Fig. 4), we now explore the temperature–melt relationship in more detail, showing
 315 that it is highly non-linear and varies significantly across regions (Fig. 5). Even though the simulations have different warming




Figure 4. Timeseries of summer average near-surface air temperature and melt over all ice shelves and subgroups of ice shelves, where AP represents the Antarctic Peninsula, EAIS is East Antarctica, WAIS is West Antarctica, and COLD includes the Filchner-Ronne and Ross Ice Shelves selected ice-shelf regions. Error bars indicate the mean and standard deviation over the historical period (1985–2014). Trendlines are shown where a significant trend (p -value < 0.05) is detected in either the historical (1985–2014) or SSP3-7.0 (2015–2099) simulations.

rates, the relationship between temperature and melt is consistent across the ERA5 and ESM model forcings (scatter plots in Fig. 5). The Filchner-Ronne and Ross ice shelves are coldest and while they experience significant warming, their melt rates remain small compared to other ice shelves, with shelves on the Antarctic Peninsula experiencing most melt. Summer melt is exponentially related to summer air temperature (Kittel et al., 2021; van Wessem et al., 2023), although the shape and steepness of this relationship vary across different ice shelves (Fig. 5).

Ice shelves in drier climates tend to experience more melt at the same temperatures compared to those in wetter climates. The boxplots in Fig. 5 illustrate the relative dryness of the highlighted ice shelves. For example, based on the exponential fits, a




Figure 5. Spatial variability in the temperature–melt relationship over Antarctic ice shelves. Scatter plots show the relationship between summer melt and temperature from RACMO simulations, forced by ERA5 (1979–2023, purple), CESM2 (1985–2099, pink), and MPI-ESM (1985–2099, yellow), with grey background scatter representing all ice shelves and all RACMO simulations. Black line shows exponential fit between temperature and melt for the considered ice shelf, with dashed line showing the 97.5th percentile confidence interval. Boxplots of summer snowfall rates are shown for each simulation. The boxplots show the distribution of summer snowfall for each simulation, with the box representing the middle 50% and whiskers extending to values within 1.5× the interquartile range. The map displays the average surface melt for 2070–2099 in RACMO(CESM2) (shading), overlaid on a grey elevation map of Antarctica.

summer-average air temperature of -4°C would lead to around 390 mm melt at the Amery Ice Shelf, that receives little snowfall, compared to 90 mm at the Amery ice shelf, which receives less than 100 mm snowfall during summer (boxplot in Fig. 5), has a temperature-melt curve where melt starts increasing at much lower temperatures compared to other ice shelves (grey scatter Fig. 5). In contrast, the Nickerson Ice Shelf where snowfall rates are higher. This shows that a given melt rate is reached at different air temperatures across the, located in a much wetter climate on the Marie Byrd Land coast, shows melt rates increasing only at substantially higher summer air temperatures. These examples show that the average summer temperature

330 required to reach a certain melt rate varies between ice shelves. For instance, reaching

To further quantify this, we use the fitted exponential temperature-melt relationships and determine for each ice shelf the summer air temperature at which 200 mm melt occurs between -6°C and -2.5°C , depending on the ice shelf mm of melt would occur. This temperature is lower for ice shelves with plotted against summer snowfall rates in Fig 6. Although the exact relationship differs between simulations, all show a clear pattern: ice shelves in drier climates (Appendix Fig. ??) reach 200 mm of melt at summer air temperatures several degrees lower than those in wetter climates.

335




Figure 6. Average DJF snowfall rates over the historical period vs. DJF near-surface air temperature required to produce 200 mm of surface melt according to fitted exponential relationships between temperature and melt per ice shelf, across the three RACMO simulations (columns). Each point represents one ice shelf, colored by the R^2 of an exponential fit. The red solid line shows a power law fit through the points, illustrating the relationship between snowfall and the temperature sensitivity of surface melt.

3.4 Variable albedo feedbacks

To better understand the relationship between temperature, surface melt, and the role of snowfall, we examine one of the primary drivers of Antarctic surface melt: the absorption of shortwave radiation (Elvidge et al., 2020; Gilbert et al., 2022; Hofsteenge et al., 2023; Jakobs et al., 2020). Figure 7 illustrates the temperature dependency of summer albedo across the major Antarctic ice shelves (ice shelf area $> 800 \text{ km}^2$). On cold ice shelves (where When summer air temperatures remain below approximately -10°C), such as on the Ronne-Filchner and Ross ice shelves, albedo is higher during warmer summers compared to colder ones. This can be attributed to increased atmospheric moisture content during warm summers, leading primarily to increased cloudiness and snowfall, with a smaller contribution from increased snowfall (not shown). Both fresh snowfall and increased cloudiness increase surface albedo. In RACMO2.4p1, albedo is wavelength-dependent, allowing the

345

model to capture the spectral effect of clouds on albedo. Cloud cover scatters much of the (near)-infrared radiation, which has a low albedo, leaving mostly visible light, for which the spectral albedo is higher (van Dalum et al., 2020). This effect is illustrated in Fig. ?? is illustrated in the second row of Fig. 7, which shows the difference in albedo between modelled (all-sky) and hypothetical clear-sky conditions across temperature bins. The figure demonstrates that, across all temperature ranges, warming is associated with increased cloud cover, which increases albedo. The cloud effect on albedo increases for lower albedos, as snow metamorphism primarily lowers the spectral albedo for red and near-infrared light, enhancing the cloud effect (Gardner and Sharp, 2010).




Figure 7. Temperature dependence of surface albedo and its relationship with snowfall rates in RACMO simulations forced by ERA5 (1979–2023) (a), CESM2 (1985–2099) (b), and MPI-ESM (1985–2099) (c). The row below shows the same analysis but for the effect of clouds on albedo, with DJF cloud cover indicated by color shading (d–f). Each scatter point represents the average over one summer season (DJF) over one major Antarctic ice shelf (ice shelves with area > 800 km²). Linear fits are shown per ice shelf where the R^2 value exceeds 0.3, with the color of each line indicating the average snowfall rate for that ice shelf.

355 As in Fig. 7 but for the effect of clouds on albedo, with DJF cloud cover in color shading.

On many of the warmer ice shelves ($T_{air, DJF}$ ~~At higher summer near-surface air temperatures ($T_{air, DJF} > -10^{\circ}$ – 10° C),~~
cloudiness increases with temperature (not shown), similar to cold regions. However, its impact on albedo warming still leads
to increased cloud cover that tends to enhance albedo (Fig. 7d-f), but this effect is outweighed by the effect of albedo reductions
associated with snow melt and snow metamorphism on albedo. While Fig. ?? shows 7d-f show that increased cloud cover can
360 raise albedo by up to 0.05, the net albedo change on the warmer ice shelves in Fig. 7 is a decrease of larger magnitude (Fig.
7a-c). Antarctic snow albedo is largely dependent on the snow grain size, and the coarsening of grains, snow metamorphism,
happens at a rate that increases with temperature (Picard et al., 2012; Taillandier et al., 2007). Additionally, when snow melts
and refreezes, the grain size increases, which enhances light absorption and further lowers albedo, a process known as the
snowmelt–albedo feedback (Jakobs et al., 2019, 2021). Whether melt or dry metamorphism plays a greater role in lowering
365 albedo depends on the the timing of snowfall relative to melt events. Besides, the The magnitude of the albedo decline depends
on snowfall rates (Fig. 7), as frequent snowfall brings small-grained, highly reflective snow to the surface. This is evident in
Fig. 7a-c, where the strongest decreases in albedo with increasing temperature occur at ice shelves with low snowfall rates
(e.g. Nansen and Publications ice shelves), whereas some ice shelves with high snowfall rates show little to no decline, or even
an increase in albedo (e.g. Swinburne ice shelf in RACMO(ERA5)). This helps explain our previous findings that ice shelves
370 that receive more snowfall have a lower melt sensitivity to near-surface air temperature compared to drier ice shelves. Frequent
fresh snowfall dampens metamorphism and the snowmelt-albedo feedback, consistent with Jakobs et al. (2021).

3.5 Temperature dependency of SEB components




Figure 8. Temperature-dependent average fitted slope between summer surface energy balance (SEB) components and summer air temperature over dry ice shelves (annual precipitation $< \sim 500$ mm). Each point represents the average slope of the SEB term with respect to temperature within a 2°C bin, calculated over all dry ice-shelf grid cells that experience melt. Shaded areas indicate the weighted standard deviation, representing the error due to spatial variability across grid cells and strength of the fit. Columns represent the different RACMO simulations and the number of grid cells used to compute the average slope per bin shown in g, h, i.

Next, we systematically assess how each SEB component responds to temperature and contributes to the non-linear relationship between temperature and melt. For each ice shelf grid cell that experiences an average summer melt of at least 1 mm during the historical period, we fit either a linear or exponential function relating the summer average SEB component to air temperature. The function with the higher R^2 value is selected, and grid cells where the optimal fit parameters could not be reliably estimated (average $R^2 < 0.3$) are excluded from the analysis. We then calculate the average slope of the selected fits across all grid cells, grouped into temperature bins. If the selected fit is exponential, the slope at the middle of the bin is taken. Figure 8 shows these average slopes for ‘dry’ ice shelves, defined as ice shelves with annual snowfall $< 500\text{ mm}$

following van Wessem et al. (2023). The resulting slopes vary across temperature bins because each bin averages multiple fits with different temperature sensitivities, and the temperature range covered by fits at each location represents only a subset of the full temperature range. Uncertainty bands are shown based on the standard deviation between the fits (reflecting variability between locations) weighted with the average R^2 value (reflecting the strength of the fit) the average SEB-temperature slopes

385 for dry ice shelves (see Section 2.1 how dry and wet ice shelves are defined). R_{net} represents net radiation ($SW_{net} + LW_{net}$).

Temperature-dependent average fitted slope between summer surface energy balance (SEB) components and summer air temperature over dry ice shelves (annual precipitation < 500 mm). Each point represents the average slope of the SEB term with respect to temperature within a 2°C bin, calculated over all dry ice shelf grid cells that experience melt. Shaded areas indicate the weighted standard deviation, representing the error due to spatial variability across grid cells and strength of the

390 fit. Columns represent the different RACMO simulations and the number of grid cells used to compute the average slope per bin is shown in g, h, i.

Net The relationship between net shortwave radiation (SW_{net}) responds differently to changes in temperature at low versus high temperatures and temperature is not constant across temperature bins (Fig. 8a,b,e). At a-c). In the lower temperature bins, the slope of SW_{net} decreases due to a decrease in is negative, indicating that SW_{net} decreases with warming. This decrease

395 occurs because higher atmospheric moisture and increased cloud cover reduce incoming shortwave radiation (SW_{in}) with rising temperature, caused by increased atmospheric moisture and cloud cover that reflect by reflecting more sunlight. This decrease in SW_{in} is larger than the reduction in outgoing shortwave radiation (SW_{out}), because surface albedo increases with temperature in this range and $\partial SW_{net} / \partial T$ remains negative (Fig. 7). However, at higher temperature bins, this pattern reverses

and the slope of SW_{out} becomes positive: SW_{out} begins to decrease faster than SW_{in} with warming as albedo drops due to snow

400 metamorphism and melt, leading to $\partial SW_{net} / \partial T > 0$. Thus, on dry ice shelves net shortwave radiation (SW_{net}) first declines with warming and then increases with further warming, producing a distinct crossover point visible across all three model combinations. The exact temperature at which this crossover occurs varies between -13°C and -8°C across the simulations, but it broadly aligns with the -11°C threshold where the strength of the snowmelt-albedo feedback increases, peaking in

Jakobs et al. (2021). The slope between SW_{net} and temperature continues to increase beyond this point in all simulations,

405 initially driven by snowmelt refreezing and dry snow metamorphism. At the highest temperature bins, albedo can decrease further due to the refreezing of rainfall and the increasing exposure of bare ice (not shown). In the RACMO simulations, rainfall begins to occur at summer mean air temperatures around -8°C as quantified by Jakobs et al. (2021) 7.5°C , increases to up to 50 mm per summer by -2.5°C , and then rises sharply above -2.5°C , following an exponential trend (not shown).

410 Longwave radiation components exhibit strong and consistent temperature response. Outgoing longwave radiation (LW_{out}) increases with near-surface air temperature due to its dependence on surface temperature, as governed by the Stefan-Boltzmann law. Meanwhile, incoming longwave radiation (LW_{in}) increases too due to a warmer, moister atmosphere. This increase in LW_{in} is generally stronger than the additional LW_{out} from a warmer surface, resulting in a net gain in longwave radiation. In the simulation forced by ERA5, however, LW_{out} begins to increase more rapidly than LW_{in} at -10°C (Fig. 8a), likely due to a

415 weakening in sensitivity of LW_{in} and difference in cloud response to the other model forcings. This decline in the sensitivity

of LW_{in} is weaker and occurs at higher temperatures in the ESM-forced simulations (Fig. 8b, c). Together, changes in net shortwave and net longwave radiation nearly compensate for each other at lower temperatures, yielding a small effect on net radiation (R_{net} in Fig. 8d, e, fd-f). But at higher temperatures, the ~~snowmelt-albedo-feedback~~ strong increase in temperature sensitivity of SW_{net} drives a strong increase in net radiation, amplifying melt.

420

The relative contributions of sensible and latent heat fluxes to the temperature response of the SEB are small compared to that of the radiative components, as the turbulent fluxes are generally smaller components in the SEB during summer (e.g. Fig. A1). ~~At higher temperatures, the~~ The sensible heat flux is on average positive in summer, as the surface temperature is colder than air temperature. The SH-temperature slope is very small or negative, especially for higher temperature bins in
 425 RACMO(ERA5). This indicates the sensible heat flux decreases despite rising air temperatures, because the darkening of the snow decreases the air-surface gradient due to increasing surface temperatures. The latent heat flux is on average negative in summer (not shown), indicating net sublimation. This flux becomes increasingly negative with rising air temperature, meaning that more energy is used for sublimation. Warmer air can hold more moisture and since the overlying air is typically dry due to its inland/katabatic origin, the humidity gradient between surface and atmosphere increases. Additionally, the saturation
 430 specific humidity at the ice surface increases exponentially with surface temperature, further increasing the vapor pressure gradient and sublimation. The energy losses through turbulent fluxes compensate for energy gains from net radiation, but this only holds at lower temperatures. At these low temperatures, most of the days included are non-melting days, and Q_M changes little for increasing temperature.

435 When considering wet ice shelves (defined by annual snowfall $> \sim 500$ mm, ~~following van Wessem et al. (2023)~~ see Section 2.1), which are generally warmer, the picture looks different than for dry ice shelves (Fig. 9). For these wet ice shelves, fewer grid cells fall into the lowest temperature bins, while a larger number of points contribute to the ~~warmer~~ highest bins. The response of ~~net shortwave radiation~~ SW_{net} to warming is weaker and less consistent across the model forcings, which may result from differences in cloud and precipitation patterns among the models in the coastal regions of the Amundsen Sea
 440 (Fig. ??1), where many of these wet ice shelves are located ~~-(Appendix Fig. B1)~~. The lower temperature sensitivity of ~~net shortwave radiation~~ SW_{net} over wet ice shelves is consistent with our findings in Fig. 7, which show that snowfall dampens the snowmelt-albedo feedback. The slope of SW_{net} is relatively constant across the temperature bins, except for the bin around 0 °C, where SW_{net} increases more rapidly with warming. In this temperature range, a fraction of the precipitation falls as rain, which removes the damping effect of fresh snowfall and reduces albedo through refreezing of rainwater.

445

A notable difference ~~from the dry-between dry and wet~~ ice shelves is the response of net longwave radiation to warming. ~~At higher temperature bins, On dry ice shelves, the increase in net longwave radiation change with warming remains positive on wet ice shelves, whereas it stabilizes or becomes negative on dry ice shelves. with temperature weakens or reverses at higher temperature bins, whereas on wet ice shelves it remains positive and even strengthens (Fig. 9a-c)~~. This shows the influence of
 450 increased humidity and cloudiness that ~~further~~ increase incoming longwave radiation and contribute to melt on wet ice shelves.

The increase in net longwave radiation contributes more to the increase in melt energy on wet ice shelves than net shortwave radiation, whereas the opposite is true at dry ice shelves.

~~For both dry and~~ Both sensible and latent heat fluxes generally decrease with warming on wet ice shelves, ~~several factors~~
455 ~~intensify melt rates when approaching summer air temperatures of~~ except as summer mean air temperatures approach 0°C :
~~first,~~ While the turbulent fluxes partly offset the increase in net radiation at lower temperatures, they intensify melt rates near
 0°C . First, air temperatures can exceed 0°C while the surface remains at the melting point, allowing sensible heat flux to
become positive and provide ~~extra heat,~~ additional energy to the surface (reflected in the shift to a positive slope in highest
temperature bin for SH in Fig. 9d-f). Second, energy loss through sublimation levels off ~~, since~~ because the saturation vapor
460 pressure over ice no longer increases when the surface temperature halts at 0°C , suppressing further sublimation while the
air in the boundary layer can become more humid. ~~Third, snowfall starts to transition to rain, therefore precipitation events~~
~~can no longer raise the surface albedo and refreezing of rainwater can lower the albedo, leading to further increases in net~~
~~shortwave radiation and net radiation. Rainfall begins to occur in the RACMO simulations at summer average air temperatures~~
~~around -7.5°C and increases to about 50 mm per summer by -2.5°C . Above -2.5°C , it starts to rise sharply, following an~~
465 ~~exponential trend (not shown). These combined processes contribute to~~ This transition to an increasing contribution from the
turbulent fluxes, together with a rapid increase in ~~melt on shelves near summer average air temperatures of 0°C~~ R_{net} , explains
the non-linear increase in energy available for melt.




Figure 9. As in Fig. 8 but for ice shelves in wet climates (annual precipitation $> \sim 500$ mm)

4 Discussion and conclusions

Our ~~This~~ study examines the spatial variability in the non-linear relationship between summer near-surface air temperature and surface melt across Antarctic ice shelves, and aims to identify the key physical processes driving this non-linearity. To investigate this, we used the regional climate model RACMO to dynamically downscale ERA5 reanalysis data for historical simulations and two historical and future climate scenarios from ESM simulations. We find that the temperature sensitivity of the surface energy balance components are similar across all simulations, indicating that the ESM-forced simulations reliably reproduce the relevant physical processes and can therefore be used to extend the temperature and melt range beyond that of the historical RACMO(ERA5) simulation. Building on Trusel et al. (2015), who related summer near-surface air temperature to surface melt using an exponential fit, we extended the analysis by examining its spatial variability and physical drivers. We find that there is no single relation between temperature and melt, but that there are differences in the exponential relationship between ice shelves in dry versus wet climates. For ~~a given the same~~ summer average air temperature, dry ice shelves tend to

480 experience more surface melt compared to wetter ones. While most ice shelves in drier climates are currently still cold and relatively stable with little melt, these ice shelves could experience rapid increases in melt under future warming. An example of a dry ice shelf that already experiences high melt rates is Amery Ice Shelf, where the largest amount of surface meltwater is observed from ~~satellites~~ satellites in East Antarctica (Tuckett et al., 2025).

485 Several studies have proposed possible explanations for the non-linearity in melt sensitivity to temperature (Trusel et al., 2015; van Wessem et al., 2023). However, this is the first study to systematically assess the ~~relationship between summer air temperature, surface energy balance (SEB) components, and their influence on surface melt in~~ temperature sensitivity of SEB components and their contribution to surface melt across Antarctica. We find that the non-linear relationship is strongly related to the temperature dependency of net shortwave radiation. At lower temperatures, warming ~~leads to increased~~ increases cloudiness and snowfall, ~~which reduce~~ reducing net shortwave radiation. ~~However, at higher temperatures~~ As summer average air temperatures approach -12 °C, the metamorphism-albedo and snowmelt-albedo feedback become ~~dominant, increasing~~ increasingly important, enhancing net shortwave radiation, particularly on dry ice shelves. This ~~dominant role of net shortwave radiation~~ is consistent with ~~findings from Franco et al. (2013) on the Greenland Ice Sheet, where albedo changes were found to be the most temperature-sensitive factor driving melt. However, similar to Antarctica, the relative contribution of individual~~ SEB components in Greenland also varies spatially. At the margins of the Greenland Ice Sheet, the sensible heat flux exhibits strong temperature sensitivity and plays a major role in driving melt anomalies (Braithwaite and Olesen, 1990; Franco et al., 2013; van den ~~-. In contrast, over Antarctic ice shelves we find that the sensible heat flux is generally less sensitive to temperature, though it increases significantly as summer air temperatures approach the melting point. This difference reflects the colder conditions over Antarctic ice shelves compared to the Greenland margins where air temperatures are more frequently positive.~~

495 ~~Once mean summer air temperatures reach the melting point, the processes governing surface melt begin to shift. Not only the sensible heat flux becomes more important near the melting point when surface temperatures become more often constrained at 0 °C (Jakobs et al. (2021), who found that the snowmelt-albedo feedback begins to strengthen around -12 °C, also the atmospheric temperatures and moisture content can continue to rise. As a result, energy losses through sublimation decrease and net longwave radiation transfers extra energy to the surface for melt. At the same time, the occurrence of rainfall increases. This~~ and peaks between -9 and -7 °C. At higher temperatures, additional albedo-lowering processes become important, including the transition from snowfall to rainfall ~~reduces surface albedo and further accelerates melt through a positive feedback. In these areas, rainfall and increasing exposure of bare ice. Rainfall~~ can precondition the ~~snow surface~~ snowpack for enhanced melt through refreezing (Nicolas et al., 2017; Vignon et al., 2021), something already observed on the Greenland Ice Sheet (Doyle et al., 2015). In our simulations, ~~annual rainfall averaged over the mean annual rainfall over~~ ice shelves increases by

500 ~~4-7 mm over the 21st century, with the largest increase over the Antarctic Peninsula (15 - 40 mm increase) and ice shelves in West Antarctica. West Antarctic ice shelves~~ (15-20 mm increase). However, ~~there is a large spread in~~ projected rainfall changes vary widely across CMIP6 models (Vignon et al., 2021). ~~Rainfall, highlighting the need for improved~~ observations and high-resolution ~~regional climate modeling are needed to improve rainfall simulation and to~~ modelling to assess the role of rainfall

510

in enhancing melt and affecting ice shelf stability (Gilbert et al., 2025a).

515

~~Surface melt is strongly related to temperature because the SEB components themselves are temperature dependent.~~ On wet ice shelves, changes in net longwave radiation with temperature drive melt more strongly than changes in net shortwave radiation, reflecting the effects of humidity and cloudiness. By contrast, on dry ice shelves, the temperature sensitivity of net longwave radiation is weak or even negative at higher temperatures, so it contributes little to enhanced melt. For average summer air temperatures below 0°C, when some melt is already occurring, turbulent heat fluxes respond weakly or decrease slightly with warming, partially offsetting increases in net radiation. As temperatures approach 0°C, both turbulent fluxes and longwave radiation increase strongly, providing additional energy for melt and amplifying the non-linear rise in melt with temperature. This happens because the surface temperature is constrained at the melting point, while air temperatures continue to rise. Such near-0°C conditions are still relatively rare over Antarctic ice shelves, so the strong contribution of turbulent fluxes to melt is limited. By contrast, at the margins of the Greenland Ice Sheet, where average summer air temperatures are often positive, sensible heat flux exhibits strong temperature sensitivity and plays a major role in driving melt anomalies (Braithwaite and Olesen, 1990; Franco et al., 2013; van den Broeke et al., 2008; Wang et al., 2021). In many other regions of Greenland however, net shortwave radiation and albedo changes are also the dominant factors controlling melt sensitivity (Franco et al., 2013).

530

We show that melt sensitivity to temperature varies spatially across Antarctic ice shelves, due to regional differences in snowfall, surface albedo and cloud cover, which influence the dominant SEB components and their sensitivity to temperature. These findings suggest that using a uniform, temperature-only approach, such as a fixed degree-day factor in positive degree day modeling, oversimplifies melt prediction and misses important regional differences and feedbacks. For example, ice shelves in dry climates tend to experience more melt at a given air temperature than ice shelves in wetter climates, partly due to lower snowfall rates, which limits fresh snow accumulation, and enhances the snowmelt–albedo feedback. Therefore, spatially varying degree-day factors are needed to more accurately represent melt sensitivity in different climate regimes (Zheng et al., 2023). Alternatively, melt parameterizations could be improved by incorporating additional variables such as precipitation or albedo (Garbe et al., 2023), or by shifting toward fully resolved SEB formulations.

540

Our findings have important implications for understanding the vulnerability of Antarctic ice shelves to climate change. The higher melt rates for the same temperature on dry ice shelves suggests that these regions may be more sensitive to future warming than previously recognized, as earlier estimates did not account for regional differences in melt sensitivity. Not only their higher melt sensitivity, but also their lower snowfall rates mean they are more likely to reach the melt-over-accumulation (MOA) threshold of around 0.7 at which meltwater ponding and hydrofracture become possible (van Wessem et al., 2023; Donat-Magnin et al., 2021). A recent study based on satellite observations confirmed that these drier ice shelves in East Antarctica are more favorable for meltwater ponding compared to ice shelves in West Antarctica (Tuckett et al., 2025). Our findings support and extend on the study by van Wessem et al. (2023) and Veldhuijsen et al. (2024), emphasizing an

underestimated risk of ice shelves in dry climates to contribute to Antarctic mass loss.

550

. Monthly SMB components, SEB components and near-surface variables (e.g. temperature, windspeed, pressure) from RACMO2.4p1 simulations forced by ERA5, CESM2 and MPI-ESM are available at: <https://doi.org/10.5281/zenodo.16902106>

. MGH led the conceptualization and analysis with guidance from WJvdB and MvdB. CvD and WJvdB developed this version of RACMO and, together with KV and MVT, performed the simulations and postprocessing. MGH prepared the manuscript with contributions from all
555 co-authors.

. One of the (co-)authors is a member of the editorial board of The Cryosphere. The authors have no other competing interests to declare.

. This publication was supported by PolarRES. The PolarRES project received funding from the European Union's Horizon 2020 research and innovation programme call H2020-LC-CLA-2018-2019-2020 under grant agreement number 101003590. This research was also supported by Ocean Cryosphere Exchanges in ANtarctica: Impacts on Climate and the Earth system, OCEAN ICE, which is funded by the European
560 Union, Horizon Europe Funding Programme for research and innovation under grant agreement Nr. 101060452, 10.3030/101060452. OCEAN ICE contribution number 35. [MvdB is supported by EMBRACER \(Summit grant SUMMIT.1.034\) financed by the Netherlands Organization for Scientific Research \(NWO\)](#). We acknowledge the ECMWF for storage facilities and computational time on their supercomputer

Appendix A: Demonstration of the pseudo-SEB approach

To illustrate that radiation penetration considerably alters the SEB, Figure A1a compares the SEB over the Larsen C Ice Shelf as simulated by RACMO2.3p2, which does not account for solar radiation penetration, with the SEB from RACMO2.4p1 (Eq. (1)). Because of penetration of solar radiation in RACMO2.4p1, the net shortwave radiation at the skin layer is smaller, and part of the radiation absorbed in the subsurface is resupplied as energy to the surface through Q_G . Because a large fraction of the melt occurs now as internal melt, $Q_{M, surf}$ is smaller in RACMO2.4p1 compared to RACMO2.3p2. Figure A1b shows the pseudo-SEB of RACMO2.4p1 (Eq. (2)), which is comparable to the SEB in RACMO2.3p2. Q_M is here the energy used for both surface and internal melt and agrees better with Q_M in RACMO2.3p2, which demonstrates that the pseudo-SEB approach is valid. Differences in the energy fluxes between the model versions also arise from differences in simulated near-surface climate, albedo (between Dec-Mar) as well as considerable year round changes in cloud cover leading to less downwelling longwave radiation (van Dalum et al., 2025), subsequently altering e.g. LW_{net} and SH.




Figure A1. Monthly mean surface energy balance (SEB) components from RACMO2.4p1 and RACMO2.3p2 for 2001–2016, spatially averaged over the Larsen C Ice Shelf. In a) the SEB from RACMO2.4p1 (solid lines) is given as in Eq. (1) and in b) the pseudo-SEB for RACMO2.4p1 (solid lines) is given as in Eq. (2). In both panels, RACMO2.3p2 estimates (dashed lines) are shown for comparison. The data are taken from van Dalum et al. (2024), as these simulations are on the exact same grid.




Figure B1. Maps of Antarctic ice shelves classified into dry (left) and wet (right) climates, separated by the median snowfall threshold of 500 mm yr^{-1} . Labels indicate the ice shelves within each category.

Appendix C: Evaluation of RACMO historical simulations with ESM forcings**for summer**

Exponential relationship temperature-melt and snowfall Average DJF snowfall rates over the historical period vs. DJF near-surface air temperature required to produce 200 mm of surface melt according to fitted exponential relationships (Fig. 5) for major Antarctic ice shelves, across the three RACMO simulations (columns). Each point represents one ice shelf, colored by the R^2 of an exponential fit between melt and temperature. Two linear regression lines are shown: a black dashed line fit to all data points, and a red solid line fit only to shelves where the exponential fit had a $R^2 \geq 0.5$. These lines illustrate the relationship between snowfall and the temperature sensitivity of surface melt.

Table C1. RACMO2.4p1 simulated summer averages (1985-2014) forced by ERA5, CESM2 and MPI-ESM. The table shows 30-year mean values and standard deviations for ice sheet average near-surface air temperature, surface energy balance [W m^{-2}] and ice sheet integrated average summer totals of mass balance components [Gt summer^{-1}].

	<u>RACMO(ERA5)</u>	<u>RACMO(CESM2)</u>	<u>RACMO(MPI-ESM)</u>
<u>T_{2m} [$^{\circ}\text{C}$]</u>	<u>-23.2 ± 0.8</u>	<u>-24.0 ± 0.6</u>	<u>-23.8 ± 0.6</u>
<u>SW_{in} [W m^{-2}]</u>	<u>349.7 ± 1.9</u>	<u>350.6 ± 1.4</u>	<u>348.7 ± 2.0</u>
<u>LW_{in} [W m^{-2}]</u>	<u>155.0 ± 2.5</u>	<u>151.3 ± 1.9</u>	<u>154.3 ± 2.0</u>
<u>SW_{net} [W m^{-2}]</u>	<u>58.4 ± 0.9</u>	<u>57.9 ± 1.3</u>	<u>56.6 ± 1.0</u>
<u>LW_{net} [W m^{-2}]</u>	<u>-63.7 ± 0.9</u>	<u>-64.0 ± 0.8</u>	<u>-62.3 ± 0.9</u>
<u>SH [W m^{-2}]</u>	<u>9.6 ± 0.5</u>	<u>10.2 ± 0.6</u>	<u>9.9 ± 0.3</u>
<u>LH [W m^{-2}]</u>	<u>-1.9 ± 0.1</u>	<u>-1.8 ± 0.1</u>	<u>-1.9 ± 0.2</u>
<u>Precipitation [Gt summer^{-1}]</u>	<u>565 ± 63</u>	<u>556 ± 68</u>	<u>608 ± 45</u>
<u>Surface sublimation [Gt summer^{-1}]</u>	<u>70.8 ± 5.5</u>	<u>67.8 ± 5.6</u>	<u>72.1 ± 6.1</u>
<u>Snowdrift sublimation [Gt summer^{-1}]</u>	<u>24.8 ± 3.8</u>	<u>28.3 ± 4.2</u>	<u>27.6 ± 2.5</u>
<u>Melt [Gt summer^{-1}]</u>	<u>115 ± 38</u>	<u>71 ± 25</u>	<u>120 ± 41</u>
<u>Refreezing [Gt summer^{-1}]</u>	<u>110 ± 37</u>	<u>67 ± 25</u>	<u>116 ± 39</u>
<u>Runoff [Gt summer^{-1}]</u>	<u>4.3 ± 3.5</u>	<u>4.7 ± 4.0</u>	<u>5.0 ± 3.1</u>
<u>SMB [Gt summer^{-1}]</u>	<u>461 ± 62</u>	<u>455 ± 66</u>	<u>503 ± 47</u>

References

Abram, N. J., Mulvaney, R., Wolff, E. W., Triest, J., Kipfstuhl, S., Trusel, L. D., Vimeux, F., Fleet, L., and Arrowsmith, C.: Acceleration
585 of snow melt in an Antarctic Peninsula ice core during the twentieth century, *Nature Geoscience*, 6, 404–411, <https://www.nature.com/articles/ngeo1787>, 2013.

Ambach, W.: Interpretation of the Positive-Degree-Days Factor by Heat Balance Characteristics-West Greenland, *Nordic Hydrology*, 19, 217–224, <http://iwaponline.com/hr/article-pdf/19/4/217/2574/217.pdf>, 1988.

Bell, R. E., Banwell, A. F., Trusel, L. D., and Kingslake, J.: Antarctic surface hydrology and impacts on ice-sheet mass balance, *Nature*
590 *Climate Change* 2018 8:12, 8, 1044–1052, <https://doi.org/10.1038/s41558-018-0326-3>, 2018.

Braithwaite, R. J. and Olesen, O. B.: Response of the Energy Balance on the Margin of the Greenland Ice Sheet to Temperature Changes, *Journal of Glaciology*, 36, 217–221, <https://doi.org/10.3189/s0022143000009461>, 1990.

Carrasco, J. F., Bozkurt, D., and Cordero, R. R.: A review of the observed air temperature in the Antarctic Peninsula. Did the warming trend
come back after the early 21st hiatus?, *Polar Science*, 28, 100 653, <https://doi.org/10.1016/J.POLAR.2021.100653>, 2021.

Coulon, V., Klose, A. K., Kittel, C., Edwards, T., Turner, F., Winkelmann, R., and Pattyn, F.: Disentangling the drivers of future Antarctic ice
loss with a historically calibrated ice-sheet model, *Cryosphere*, 18, 653–681, <https://doi.org/10.5194/TC-18-653-2024>, 2024.

DeConto, R. M., Pollard, D., Alley, R. B., Velicogna, I., Gasson, E., Gomez, N., Sadai, S., Condron, A., Gilford, D. M., Ashe, E. L., Kopp, R. E., Li, D., and Dutton, A.: The Paris Climate Agreement and future sea-level rise from Antarctica, *Nature* 2021 593:7857, 593, 83–89, <https://doi.org/10.1038/s41586-021-03427-0>, 2021.




Figure C1. As in Fig. 22-1 but for DJF annual means.

600 Di Biase, V., Kuipers Munneke, P., Wouters, B., van den Broeke, M. R., and van Tiggelen, M.: Estimating Antarctic surface melt rates using passive microwave data calibrated with weather station observations, *The Cryosphere*, 20, 87–96, <https://doi.org/10.5194/TC-20-87-2026>, 2026.

Donat-Magnin, M., Jourdain, N. C., Kittel, C., Agosta, C., Amory, C., Gallée, H., Krinner, G., and Chekki, M.: Future surface mass balance and surface melt in the Amundsen sector of the West Antarctic Ice Sheet, *Cryosphere*, 15, 571–593, <https://doi.org/10.5194/TC-15-571-2021>, 2021.

605 Doyle, S. H., Hubbard, A., van de Wal, R. S., Box, J. E., van As, D., Scharrer, K., Meierbachtol, T. W., Smeets, P. C., Harper, J. T., Johansson, E., Mottram, R. H., Mikkelsen, A. B., Wilhelms, F., Patton, H., Christoffersen, P., and Hubbard, B.: Amplified melt and flow of the Greenland ice sheet driven by late-summer cyclonic rainfall, *Nature Geoscience*, 8, 647–653, <https://doi.org/10.1038/NGEO2482>, 2015.

610 ECMWF: IFS Documentation CY47R1 - Part IV: Physical Processes, <https://www.ecmwf.int/en/elibrary/81189-ifs-documentation-cy47r1-part-iv-physical-processes>, 2020.

Elvidge, A. D., Kuipers Munneke, P., King, J. C., Renfrew, I. A., and Gilbert, E.: Atmospheric Drivers of Melt on Larsen C Ice Shelf: Surface Energy Budget Regimes and the Impact of Foehn, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2020JD032463>, 2020.

615 Franco, B., Fettweis, X., and Erpicum, M.: Future projections of the Greenland ice sheet energy balance driving the surface melt, *The Cryosphere*, 7, 1–18, <https://doi.org/10.5194/tc-7-1-2013>, 2013.

Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nature Climate Change*, 6, 479–482, <https://www.nature.com/articles/nclimate2912>, 2016.

- Gadde, S. and van de Berg, W. J.: Contribution of blowing-snow sublimation to the surface mass balance of Antarctica, *Cryosphere*, 18, 4933–4953, <https://doi.org/10.5194/TC-18-4933-2024>, 2024.
- Garbe, J., Zeitz, M., Krebs-Kanzow, U., and Winkelmann, R.: The evolution of future Antarctic surface melt using PISM-dEBM-simple, *Cryosphere*, 17, 4571–4599, <https://doi.org/10.5194/TC-17-4571-2023>, 2023.
- Gardner, A. S. and Sharp, M. J.: A review of snow and ice albedo and the development of a new physically based broadband albedo parameterization, *Journal of Geophysical Research: Earth Surface*, 115, 1009, <https://doi.org/10.1029/2009JF001444>, 2010.
- Gilbert, E. and Kittel, C.: Surface Melt and Runoff on Antarctic Ice Shelves at 1.5°C, 2°C, and 4°C of Future Warming, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL091733>, 2021.
- Gilbert, E., Orr, A., Renfrew, I. A., King, J. C., and Lachlan-Cope, T.: A 20-Year Study of Melt Processes Over Larsen C Ice Shelf Using a High-Resolution Regional Atmospheric Model: 2. Drivers of Surface Melting, *Journal of Geophysical Research: Atmospheres*, 127, <https://doi.org/10.1029/2021JD036012>, 2022.
- Gilbert, E., Pishniak, D., Torres, J. A., Orr, A., MacLennan, M., Wever, N., and Verro, K.: Extreme precipitation associated with atmospheric rivers over West Antarctic ice shelves: insights from kilometre-scale regional climate modelling, *Cryosphere*, 19, 597–618, <https://doi.org/10.5194/TC-19-597-2025>, 2025a.
- Gilbert, E., Torres-Alavez, J. A., Hofsteenge, M. G., van de Berg, W. J., Boberg, F., Christensen, O. B., van Dalum, C. T., Fettweis, X., Gumber, S., Hansen, N., Kittel, C., Lambin, C., Maure, D., Mottram, R., Olesen, M., Orr, A., Phillips, T., van Tiggelen, M., Verro, K., and Mooney, P. A.: The PolarRES dataset: a state-of-the-art regional climate model ensemble for understanding Antarctic climate, *EGU sphere*, pp. 1–35, <https://doi.org/10.5194/EGUSPHERE-2025-4214>, 2025b.
- Golledge, N. R., Keller, E. D., Gomez, N., Naughten, K. A., Bernales, J., Trusel, L. D., and Edwards, T. L.: Global environmental consequences of twenty-first-century ice-sheet melt, *Nature* 2019 566:7742, 566, 65–72, <https://doi.org/10.1038/s41586-019-0889-9>, 2019.
- Hersbach, H., Bell, B., Simmons, A., Berrisford, P., Dahlgren, P., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Radu, R., Schepers, D., Soci, C., Villaume, S., Bidlot, J. R., Haimberger, L., Woollen, J., Buontempo, C., and Thépaut, J. N.: Complete ERA5 from 1940: Fifth generation of ECMWF atmospheric reanalyses of the global climate, Copernicus Climate Change Service (C3S) Data Store (CDS), <https://doi.org/10.24381/cds.143582cf>, 2017.
- Hofsteenge, M. G., Cullen, N. J., Conway, J. P., Reijmer, C. H., Van Den Broeke, M. R., and Katurji, M.: Meteorological drivers of melt at two nearby glaciers in the McMurdo Dry Valleys of Antarctica, *Journal of Glaciology*, <https://doi.org/10.1017/jog.2023.98>, 2023.
- Jakobs, C. L., Reijmer, C. H., Kuipers Munneke, P., König-Langlo, G., and van den Broeke, M. R.: Quantifying the snowmelt-albedo feedback at Neumayer Station, East Antarctica, *The Cryosphere*, 13, 1473–1485, <https://doi.org/10.5194/tc-13-1473-2019>, 2019.
- Jakobs, C. L., Reijmer, C. H., Smeets, C. J., Trusel, L. D., Van De Berg, W. J., Van Den Broeke, M. R., and Van Wessem, J. M.: A benchmark dataset of in situ Antarctic surface melt rates and energy balance, *Journal of Glaciology*, 66, 291–302, <https://doi.org/10.1017/JOG.2020.6>, 2020.
- Jakobs, C. L., Reijmer, C. H., van den Broeke, M. R., van de Berg, W. J., and van Wessem, J. M.: Spatial Variability of the Snowmelt-Albedo Feedback in Antarctica, *Journal of Geophysical Research: Earth Surface*, 126, <https://doi.org/10.1029/2020JF005696>, 2021.
- Kingslake, J., Ely, J. C., Das, I., and Bell, R. E.: Widespread movement of meltwater onto and across Antarctic ice shelves, *Nature*, 544, 349–352, <https://doi.org/10.1038/nature22049>, 2017.

- 655 Kittel, C., Amory, C., Agosta, C., Jourdain, N. C., Hofer, S., Delhasse, A., Doutreloup, S., Huot, P. V., Lang, C., Fichet, T., and Fettweis, X.: Diverging future surface mass balance between the Antarctic ice shelves and grounded ice sheet, *Cryosphere*, 15, 1215–1236, <https://doi.org/10.5194/TC-15-1215-2021>, 2021.
- Kuipers Munneke, P., Ligtenberg, S. R., van den Broeke, M. R., and Vaughan, D. G.: Firn air depletion as a precursor of Antarctic ice-shelf collapse, *Journal of Glaciology*, 60, 205–214, <https://doi.org/10.3189/2014JoG13J183>, 2014.
- 660 Lai, C. Y., Kingslake, J., Wearing, M. G., Chen, P. H. C., Gentine, P., Li, H., Spergel, J. J., and van Wessem, J. M.: Vulnerability of Antarctica’s ice shelves to meltwater-driven fracture, *Nature*, 584, 574–578, <https://www.nature.com/articles/s41586-020-2627-8>, 2020.
- Ligtenberg, S. R., van de Berg, W. J., van den Broeke, M. R., Rae, J. G., and van Meijgaard, E.: Future surface mass balance of the Antarctic ice sheet and its influence on sea level change, simulated by a regional atmospheric climate model, *Climate Dynamics*, 41, 867–884, <https://doi.org/10.1007/S00382-013-1749-1>, 2013.
- 665 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat, W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel, A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. d., Ommen, T. D., Wessem, M. v., and Young, D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132–137, <https://doi.org/10.1038/S41561-019-0510-8>, 2020.
- 670 Nicolas, J. P., Vogelmann, A. M., Scott, R. C., Wilson, A. B., Cadetdu, M. P., Bromwich, D. H., Verlinde, J., Lubin, D., Russell, L. M., Jenkinson, C., Powers, H. H., Ryzek, M., Stone, G., and Wille, J. D.: January 2016 extensive summer melt in West Antarctica favoured by strong El Niño, *Nature Communications*, 8, 1–10, <https://doi.org/10.1038/NCOMMS15799>, 2017.
- Ohmura, A.: Physical Basis for the Temperature-Based Melt-Index Method, Tech. rep., 2001.
- Orr, A., Deb, P., Clem, K. R., Gilbert, E., Bromwich, D. H., Boberg, F., Colwell, S., Hansen, N., Lazzara, M. A., Mooney, P. A., Mottram, R., Niwano, M., Phillips, T., Pishniak, D., Reijmer, C. H., van de Berg, W. J., Webster, S., and Zou, X.: Characteristics of Surface “Melt Potential” over Antarctic Ice Shelves based on Regional Atmospheric Model Simulations of Summer Air Temperature Extremes from 1979/80 to 2018/19, *Journal of Climate*, 36, 3357–3383, <https://doi.org/10.1175/JCLI-D-22-0386.1>, 2023.
- 675 Picard, G., Domine, F., Krinner, G., Arnaud, L., and Lefebvre, E.: Inhibition of the positive snow-albedo feedback by precipitation in interior Antarctica, *Nature Climate Change*, 2, 795–798, <https://doi.org/10.1038/NCLIMATE1590>, 2012.
- 680 Scambos, T. A., Hulbe, C., Fahnestock, M., and Bohlander, J.: The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, *Journal of Glaciology*, 46, 516–530, <https://doi.org/10.3189/172756500781833043>, 2000.
- Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R., Dumas, C., Galton-Fenzi, B. K., Gladstone, R., Golledge, N. R., Gregory, J. M., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E., Leguy, G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N.-J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Breedam, J. V., Van De Wal, R. S. W., Winkelmann, R., Zhao, C., Zhang, T., and Zwinger, T.: ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *Matthew J. Hoffman*, 14, 3033–3070, <https://doi.org/10.5194/tc-14-3033-2020>, 2020.
- 685 Taillandier, A. S., Domine, F., Simpson, W. R., Sturm, M., and Douglas, T. A.: Rate of decrease of the specific surface area of dry snow: Isothermal and temperature gradient conditions, *Journal of Geophysical Research: Earth Surface*, 112, 3003, <https://doi.org/10.1029/2006JF000514>, 2007.
- 690 Trusel, L. D., Frey, K. E., Das, S. B., Karnauskas, K. B., Kuipers Munneke, P., Van Meijgaard, E., and Van Den Broeke, M. R.: Divergent trajectories of Antarctic surface melt under two twenty-first-century climate scenarios, <https://doi.org/10.1038/ngeo2563>, 2015.

- Tuckett, P. A., Sole, A. J., Livingstone, S. J., Jones, J. M., Lea, J. M., and Gilbert, E.: Continent-wide mapping shows increasing sensitivity of East Antarctica to meltwater ponding, *Nature Climate Change*, <https://doi.org/10.1038/s41558-025-02363-5>, 2025.
- 695 Turner, J., Lu, H., White, I., King, J. C., Phillips, T., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J., Mulvaney, R., and Deb, P.: Absence of 21st century warming on Antarctic Peninsula consistent with natural variability, *Nature* 2016 535:7612, 535, 411–415, <https://doi.org/10.1038/nature18645>, 2016.
- Undén, P., Rontu, L., Jarvinen, H., Lynch, P., Calvo Sánchez, F., Cats, G., Cuxart, J., Eerola, K., Fortelius, C., García-Moya, J., and Jones, C.: HIRLAM-5 scientific documentation, Tech. rep., <https://www.researchgate.net/publication/278962772>, 2002.
- 700 van Dalum, C. T., van de Berg, W. J., Lhermitte, S., and van den Broeke, M. R.: Evaluation of a new snow albedo scheme for the greenland ice sheet in the regional atmospheric climate model (racmo2), *Cryosphere*, 14, 3645–3662, <https://doi.org/10.5194/TC-14-3645-2020>, 2020.
- van Dalum, C. T., van de Berg, W. J., and van den Broeke, M. R.: Sensitivity of Antarctic surface climate to a new spectral snow albedo and radiative transfer scheme in RACMO2.3p3, *Cryosphere*, 16, 1071–1089, <https://doi.org/10.5194/TC-16-1071-2022>, 2022.
- 705 van Dalum, C. T., van de Berg, W. J., Gadde, S. N., van Tiggelen, M., van der Drift, T., van Meijgaard, E., van Uft, L. H., and van den Broeke, M. R.: First results of the polar regional climate model RACMO2.4, *The Cryosphere*, 18, 4065–4088, <https://doi.org/10.5194/tc-18-4065-2024>, 2024.
- van Dalum, C. T., van de Berg, W. J., van den Broeke, M. R., and van Tiggelen, M.: The surface mass balance and near-surface climate of the Antarctic ice sheet in RACMO2.4p1, *Cryosphere*, 19, 4061–4090, <https://doi.org/10.5194/TC-19-4061-2025>, 2025.
- 710 van de Berg, W. J. and Medley, B.: Brief Communication: Upper-air relaxation in RACMO2 significantly improves modelled interannual surface mass balance variability in Antarctica, *Cryosphere*, 10, 459–463, <https://doi.org/10.5194/TC-10-459-2016>, 2016.
- van den Broeke, M.: Strong surface melting preceded collapse of Antarctic Peninsula ice shelf, *Geophysical Research Letters*, 32, 1–4, <https://doi.org/10.1029/2005GL023247>, 2005.
- van den Broeke, M., Smeets, P., Ettema, J., van der Veen, C., van de Wal, R., and Oerlemans, J.: Partitioning of melt energy and meltwater fluxes in the ablation zone of the west Greenland ice sheet, *The Cryosphere*, 2, 179–189, <https://doi.org/10.5194/tc-2-179-2008>, 2008.
- 715 van den Broeke, M. R., Kuipers Munneke, P., Noël, B., Reijmer, C., Smeets, P., van de Berg, W. J., and van Wessem, J. M.: Contrasting current and future surface melt rates on the ice sheets of Greenland and Antarctica: Lessons from in situ observations and climate models, *PLOS Climate*, 2, <https://doi.org/10.1371/journal.pclm.0000203>, 2023.
- 720 van Wessem, J. M., Ligtenberg, S. R. M., Reijmer, C. H., van de Berg, W. J., van den Broeke, M. R., Barrand, N. E., Thomas, E. R., Turner, J., Wuite, J., Scambos, T. A., and van Meijgaard, E.: The modelled surface mass balance of the Antarctic Peninsula at 5.5 km horizontal resolution, *The Cryosphere*, 10, 271–285, <https://doi.org/10.5194/tc-10-271-2016>, 2016.
- van Wessem, J. M., van den Broeke, M. R., Wouters, B., and Lhermitte, S.: Variable temperature thresholds of melt pond formation on Antarctic ice shelves, *Nature Climate Change*, 13, 161–166, <https://doi.org/10.1038/s41558-022-01577-1>, 2023.
- 725 Vaughan, D. G.: Recent Trends in Melting Conditions on the Antarctic Peninsula and Their Implications for Ice-sheet Mass Balance and Sea Level, *Arctic, Antarctic, and Alpine Research*, 38, 147–152, [https://doi.org/10.1657/1523-0430\(2006\)038\[0147:RTIMCO\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2006)038[0147:RTIMCO]2.0.CO;2), 2006.
- Veldhuijsen, S. B., van de Berg, W. J., Kuipers Munneke, P., and van den Broeke, M. R.: Firn air content changes on Antarctic ice shelves under three future warming scenarios, *Cryosphere*, 18, 1983–1999, <https://doi.org/10.5194/TC-18-1983-2024>, 2024.

- 730 Vignon, Roussel, M. L., Gorodetskaya, I. V., Genthon, C., and Berne, A.: Present and Future of Rainfall in Antarctica, *Geophysical Research Letters*, 48, e2020GL092281, <https://doi.org/10.1029/2020GL092281>, 2021.
- Wang, W., Zender, C. S., van As, D., Fausto, R. S., and Laffin, M. K.: Greenland Surface Melt Dominated by Solar and Sensible Heating, *Geophysical Research Letters*, 48, e2020GL090653, <https://doi.org/10.1029/2020GL090653>, 2021.
- Williams, R. S., Marshall, G. J., Levine, X., Graff, L. S., Handorf, D., Johnston, N. M., Karpechko, A. Y., Andrew, O. R., van de Berg, W. J.,
 735 Wijngaard, R. R., and Mooney, P. A.: Future Antarctic Climate: Storylines of Midlatitude Jet Strengthening and Shift Emergent from CMIP6, *Journal of Climate*, 37, 2157–2178, <https://doi.org/10.1175/JCLI-D-23-0122.1>, 2024.
- Zheng, L., Shang, X., van den Broeke, M. R., Noël, B., Li, X., Fettweis, X., Liang, Q., Wang, K., Liu, J., and Cheng, X.: Rapid increases in satellite-observed ice sheet surface meltwater production, *Nature Climate Change* 2025 15:7, 15, 769–774, <https://doi.org/10.1038/s41558-025-02364-4>, 2025.
- 740 Zheng, Y., Golledge, N. R., Gossart, A., Picard, G., and Leduc-Leballeur, M.: Statistically parameterizing and evaluating a positive degree-day model to estimate surface melt in Antarctica from 1979 to 2022, *Cryosphere*, 17, 3667–3694, <https://doi.org/10.5194/tc-17-3667-2023>, 2023.