



The Biogeochemical Model for Hypoxic and Benthic Influenced areas: BAMHBI v1.0

Marilaure Grégoire¹, Luc Vandenbulcke¹, Séverine Chevalier¹, Mathurin Choblet¹, Ilya Drozd¹, Jean-François Grailet¹, Evgeny Ivanov¹, Loïc Macé¹, Polina Verezhenskaya¹, Haolin Yu¹, Lauranne Alaerts¹, Ny Riana Randresihaja¹, Victor Mangeleer¹, Guillaume Maertens de Noordhout¹, Arthur Capet^{2, 3}, Catherine Meulders¹, Anne Mouchet¹, Guy Munhoven⁴, and Karline Soetaert⁵

¹MAST-FOCUS, Astrophysics, Geophysics and Oceanography Department, University of Liège, B5 Sart-Tilman, 4000 Liège, Belgium

²ECOMOD, Royal Belgian Institute of Natural Sciences, Brussels, Belgium

³B-Geosys, Université Libre de Bruxelles, Brussels, Belgium

⁴LPAP–SPHERES Astrophysics, Geophysics and Oceanography Department, University of Liège, 4000 Liège, Belgium

⁵Estuarine and Delta System, NIOZ, Yerseke, The Netherlands

Correspondence: Marilaure Grégoire (mgregoire@uliege.be)

Abstract. This paper describes the ocean Biogeochemical Model for Hypoxic and Benthic Influenced areas (BAMHBI). BAMHBI is a moderate complexity marine biogeochemical model that describes the cycling of carbon, nitrogen, phosphorus, silicon and oxygen through the marine foodweb. It involves 22 state variables, extends from bacteria up to mesozooplankton and includes three phytoplankton functional types (PFTs), two zooplankton size-classes, a microbial loop with several classes of detritic materials. Five optional modules are available allowing to extend the model with the explicit modelling of Chlorophyll *a* (Chl*a*) in each PFT, benthic degradation, gelatinous dynamics, particles aggregation and the carbonate system. BAMHBI describes the degradation of organic matter according to oxygenation conditions using an approach similar to that used in the sediment to simulate early diagenesis. The model is particularly appropriate for modelling low oxygen environments and the generation of sulfidic waters. An optional benthic module solves the degradation of sedimentary organic matter and the benthic-pelagic fluxes of solutes using an efficient formulation based on meta-modelling.

This paper describes in details model formulations, implementation and coupling with the physics. BAMHBI's code is written in Fortran and can be coupled with many hydrodynamical models. Two case studies of application of BAMHBI in the Black Sea are described. One describes the application of BAMHBI to simulate the biogeochemical dynamics of the northwestern shelf during the eutrophication period. In particular, the ability of BAMHBI to simulate the oxygen dynamics at seasonal and interannual scales is assessed with a focus on the simulation of bottom hypoxia. We highlight the results of the benthic modelling module and its ability to represent benthic-pelagic fluxes. The second case study compares the BAMHBI simulated Chl*a* and oxygen dynamics in the deep sea with respect to biogeochemical Argo.



20 1 Introduction

Ocean biogeochemical models depict the cycles of essential chemical elements, mostly carbon (C), nitrogen (N), phosphorus (P) and oxygen (O₂), through the marine foodweb, what is often called the “green ocean.” The organic (living and dead) and inorganic forms of ocean materials and their transformation by biogeochemical processes are described with a level of details that differs according to the objectives of the study and the data at hand. Marine biogeochemical models mostly detail the foodweb from bacteria up to mesozooplankton and do not describe the species level. Instead plankton diversity is divided into groups (usually from two up to four) sharing common functional characteristics, the so-called Plankton Functional Types (PFTs). Each PFT aggregates a large number of species into one compartment. During the last three decades, we have had a considerable development of biogeochemical modelling to understand the mechanisms of environmental changes and to improve the quality of prediction. The increase in computing resources and the new knowledge acquired from observations have allowed the development of more complex biogeochemical models that incorporate new compartments, the description of the (un)coupled cycles of several chemical elements, the spectral light propagation, etc. Some of these biogeochemical models are updated regularly, well documented, openly available and used in operational production. This is the case for instance of ERSEM (Butenschön et al., 2016), BFM (Cossarini et al., 2017), PISCES (Aumont et al., 2015) and ERGOM (Daewel et al., 2019).

Here we describe the Biogeochemical Model for Hypoxic and Benthic Influenced areas (BAMHBI) that has been developed to simulate oxygen deficient environments and the generation of euxinic waters in the water column and on the bottom. BAMHBI is an intermediate complexity model that represents the cycles of C, N, Si, O₂, P through the pelagic and benthic compartments. BAMHBI is coded to be flexible in terms of model structure by (currently) including five optional modules that explicitly model 1) the Chlorophyll *a* (Chl*a*) content of each phytoplankton group, 2) the omnivorous and carnivorous gelatinous zooplankton, 3) the carbonate system, 4) the aggregation of organic matter and 5) the benthic compartment. This allows the user to obtain a (simplified) version of the model appropriate for the targeted application. BAMHBI is a stand-alone model that can be coupled with any hydrodynamical model.

BAMHBI has been coupled with GOTM and applied in the Black Sea to simulate the eutrophication period (Grégoire et al., 2008) and biogeochemical cycles (Grégoire and Soetaert, 2010). It has been coupled with the GHER hydrodynamical model to predict the occurrence of seasonal hypoxia over the northwestern shelf (Capet et al., 2013) and benthic-pelagic coupling (Capet et al., 2016). BAMHBI has also been applied to simulate the Ligurian sea ecosystem (Raick et al., 2005, 2006). BAMHBI is online coupled with the hydrodynamical model NEMO in the frame of the Copernicus Marine Service for forecasting (daily production of a 10-day forecast) and hindcasting (back to 1950) the Black Sea’s biogeochemistry (Ciliberti et al., 2021).

This paper describes in detail the BAMHBI conceptual and mathematical model. Two applications describe the coupling of BAMHBI with the NEMO hydrodynamical model in a three-dimensional framework.



2 The BAMHBI Model

BAMHBI models trophic interactions across the food web, spanning from bacteria to mesozooplankton throughout both the pelagic and benthic system. The number of state variables varies from 22 in the simplest version of BAMHBI up to 35 in the full version that includes all the optional modules (Figure 1, Table 1).




Figure 1. Schematic representation of the BAMHBI biogeochemical model showing the model state variables (in rectangular boxes) and interactions between them in the pelagic and benthic region differentiating the inorganic, living and non-living organic parts which are made of carbon, nitrogen and phosphorus, and also silicon. It also shows the oxygen (O_2) and oxygen demand unit (ODU) fluxes. Optional modules are framed in black: carbonate, benthic, gelatinous, aggregation modules. An additional optional module explicitly computes the Chla content of each Plankton Functional Type.

55 The model state variables are listed in Table 1 and their evolution is governed by the following general equation:

$$\frac{\partial C}{\partial t} + \mathbf{v} \cdot \nabla C + \frac{\partial(v_z C)}{\partial z} = \frac{\partial C}{\partial t} \Big|_{\text{Bio}} + \nabla_{\mathbf{H}}(K_D \nabla_{\mathbf{H}} C) + \frac{\partial}{\partial z} \lambda^C \frac{\partial C}{\partial z} \quad (1)$$

Equation 1 expresses that the rate of change of each concentration of biogeochemical component C , expressed in mmol m^{-3} , results from the three-dimensional transport by advection ($\mathbf{v} \cdot \nabla C$) by the main current with a velocity \mathbf{v} , sinking ($\frac{\partial(v_z C)}{\partial z}$)



with a sinking velocity $v_{z,C}$, diffusion $(\nabla_{\mathbf{H}}(K_D \nabla_{\mathbf{H}} C) + \frac{\partial}{\partial z} \lambda^C \frac{\partial C}{\partial z})$ with a vertical and horizontal diffusion coefficients re-
 spectively of λ^C and K_D and biogeochemical interactions $(\frac{\partial C}{\partial t}|_{\text{Bio}})$. Equations 2 until 25 give the biogeochemical interactions
 for each state variable listed in Table 1. In these equations, the subscript i refers to the three phytoplankton functional groups
 ($i = 1, 2, 3$) while and j refers to the two zooplankton (gelatinous) groups ($j = 1, 2$).

$$\frac{\partial CPHY_i}{\partial t} \Big|_{\text{Bio}} = \text{Photosynthesis}PHY_i - \text{Respiration}PHY_i - \text{Leakage}PHY_{C,i} - \text{Mort}PHY_{C,i} - \text{Graz}CPHY_i \quad (i = 1, 2, 3) \quad (2)$$

$$\frac{\partial NPHY_i}{\partial t} \Big|_{\text{Bio}} = (NOs_i^{\text{uptake}} + NHs_i^{\text{uptake}}) - \text{Leakage}PHY_{N,i} - \text{Mort}PHY_{N,i} - \text{Graz}NPHY_i \quad (i = 1, 2, 3) \quad (3)$$

$$65 \quad \frac{\partial CZOO_j}{\partial t} \Big|_{\text{Bio}} = \text{Growth}ZOO_{C,j} - \text{Mort}ZOO_{C,j} - \text{Graz}CZOO_j \quad (j = 1, 2) \quad (4)$$

$$\frac{\partial CBAC}{\partial t} \Big|_{\text{Bio}} = \text{Growth}BAC - \text{Mort}BAC_C - \text{Graz}CBAC \quad (5)$$

$$\frac{\partial NOs}{\partial t} \Big|_{\text{Bio}} = - \sum_{j=1}^3 NOs_j^{\text{uptake}} + NHs_{\text{oxid}O_2} NHs - \text{Resp}BAC_{\text{Nitrate}} \gamma_{NOs}^{POC} - NHs_{\text{oxid}NOs} NHs \gamma_{NOs}^{NHs} - ODU_{\text{oxid}NOs} ODU \gamma_{NOs}^{ODU} \quad (6)$$

$$\frac{\partial NHs}{\partial t} \Big|_{\text{Bio}} = \sum_{j=1}^2 (\text{Excr}ZOO_j) + \text{Excr}BAC - \sum_{j=1}^3 NHs_j^{\text{uptake}} - U_A - NHs_{\text{oxid}NOs} NHs - NHs_{\text{oxid}O_2} NHs \quad (7)$$

$$70 \quad \text{if Gelatinous} : + \sum_{j=1}^2 (\text{Adj} \text{Excr}GEL_j)$$

$$\frac{\partial POs}{\partial t} \Big|_{\text{Bio}} = \left[\sum_{j=1}^2 (\text{Excr}ZOO_j) + \text{Excr}BAC - U_A \right] (P : N) - \sum_{j=1}^3 POs_j^{\text{uptake}} - \text{Precipitation}POs \quad (8)$$

$$\text{if Gelatinous} : + \sum_{j=1}^2 (\text{Adj} \text{Excr}GEL_j)$$

$$\frac{\partial SiOs}{\partial t} \Big|_{\text{Bio}} = -SiOs^{\text{uptake}} + \text{Dissolution}SiDet \quad (9)$$

$$\frac{\partial DOCL}{\partial t} \Big|_{\text{Bio}} = \sum_{i=1}^3 \left(\text{Leakage}PHY_{C,i} + \gamma_1 \text{Extra} \text{Excr}PHY_i + (1 - \gamma_1) \delta_2 \text{Extra} \text{Excr}PHY_i \right) + \text{Hydrol}DOCSL - U_c \quad (10)$$

$$75 \quad + \delta_1 \left[\epsilon \sum_{i=1}^3 \text{Mort}PHY_{C,i} + \phi_i \sum_{j=1}^2 \text{Graz}ZOO_{C,j} + \text{Mort}BAC_C + \text{Hydrol}POC \right]$$

$$\frac{\partial DONL}{\partial t} \Big|_{\text{Bio}} = \sum_{i=1}^3 (\text{Leakage}PHY_{N,i}) \quad (11)$$

$$+ \delta_1 \left[\epsilon \sum_{i=1}^3 \text{Mort}PHY_{N,i} + \phi \sum_{j=1}^2 \text{Graz}ZOO_{N,j} + \frac{\text{Mort}BAC_C}{(C : N)_{BAC}} + \text{Hydrol}PON \right] + \text{Hydrol}DONSL - U_n$$

$$\frac{\partial DOCSL}{\partial t} \Big|_{\text{Bio}} = \sum_{i=1}^3 ((1 - \gamma_1)(1 - \delta_2) \text{Extra} \text{Excr}PHY_i) \quad (12)$$



Table 1. List of biogeochemical state variables, description, code declaration and units.

STATE VARIABLES	DESCRIPTION	Code Declaration	UNITS
<u>Pelagic module</u>			
NOs	Nitrate	NOs	mmol N m ⁻³
NHs	Ammonium	NHs	mmol N m ⁻³
POs	Phosphate	PHOs	mmol P m ⁻³
SiOs	Silicic acid	SiOs	mmol Si m ⁻³
O ₂	Dissolved oxygen concentration	DOX	mmol O ₂ m ⁻³
ODU	Oxygen demand unit concentration	ODU	mmol ODU m ⁻³
<i>NPHY₁, NPHY₂, NPHY₃</i>	Phototrophic small flagellates (nano and pico), large flagellates (micro), Diatoms in nitrogen	NEM, NFL, NDI	mmol N m ⁻³
<i>CPHY₁, CPHY₂, CPHY₃</i>	Phototrophic small flagellates (nano and pico), large flagellates (micro), Diatoms in carbon	CEM, CFL, CDI	mmol C m ⁻³
<i>CZOO1, CZOO2</i>	Micro- and meso- zooplankton, in carbon	MIC, MES	mmol C m ⁻³
<i>CBAC</i>	Bacteria in carbon	BAC	mmol C m ⁻³
<i>DOCL, DOCSL</i>	Labile and semi-labile dissolved organic carbon	DCL, DCS	mmol C m ⁻³
<i>DONL, DONSL</i>	Labile and semi-labile dissolved organic nitrogen	DNL, DNS	mmol N m ⁻³
<i>POC, PON</i>	Particulate organic carbon and nitrogen	POC, PON	mmol C m ⁻³ , mmol N m ⁻³
<i>SiDet</i>	Detrital silicate	SID	mmol Si m ⁻³
<u>Optional modules</u>			
<u>chlorophyll-Geider module</u>			
<i>ChlPHY1, ChlPHY2, ChlPHY3</i>	Chl _a content of phototrophic small flagellates (nano and pico), large flagellates (micro), Diatoms	CHE, CHF, CHD	mg Chl _a m ⁻³
<u>gelatinous module</u>			
<i>CGEL1, CGEL2</i>	Gelatinous omnivorous and carnivorous in carbon	NOC, GEL	mmol C m ⁻³
<u>carbonate module</u>			
DIC	Dissolved inorganic carbon	DIC	mmol C m ⁻³
Σ[-]	The excess negative charge	CHA	mmol m ⁻³
<u>aggregation module</u>			
<i>AggNum</i>	Aggregates number	AGG	m ⁻³
<u>benthic module</u>			
<i>S_C^s</i>	Vertically integrated sedimentary organic carbon (slow decay)	sCSED	mmol C m ⁻²
<i>S_C^f</i>	Vertically integrated sedimentary organic carbon (fast decay)	fCSED	mmol C m ⁻²
<i>S_S^s</i>	Vertically integrated sedimentary biogenic silica (slow dissolved)	sSSED	mmol Si m ⁻²
<i>S_S^f</i>	Vertically integrated sedimentary biogenic silica (fast dissolved)	fSSED	mmol Si m ⁻²
<i>S_{N:C}</i>	N:C ratio of sediment organic matter	NCrSED	mol N (mol C) ⁻¹



$$80 \quad \left. \frac{\partial DONSL}{\partial t} \right|_{\text{Bio}} = (1 - \delta_1) \left[\epsilon \sum_{i=1}^3 \text{MortPHY}_{C,i} + \phi \sum_{j=1}^2 \text{GrazZOO}_{C,j} + \text{HydrolPOC} + \text{MortBAC}_C \right] - \text{HydrolDOCSL} \\ + (1 - \delta_1) \left[\epsilon \sum_{i=1}^3 \text{MortPHY}_{N,i} + \phi \sum_{j=1}^2 \text{GrazZOO}_{N,j} + \text{HydrolPON} + \frac{\text{MortBAC}_C}{(C:N)_{BAC}} \right] - \text{HydrolDONSL} \quad (13)$$

$$\left. \frac{\partial POC}{\partial t} \right|_{\text{Bio}} = \sum_{i=1}^3 (1 - \epsilon) \text{MortPHY}_{C,i} + \sum_{j=1}^2 (\text{MortZOO}_{C,j} + \text{MortGEL}_{C,j}) + \sum_{j=1}^2 ((1 - \beta_C) I_{C,j}) - \text{HydrolPOC} - \text{Graz}_{POC} \quad (14)$$

$$\text{if Gelatinous : } + \sum_{j=1}^2 ((1 - \text{AssEffGEL}_j) \text{GrazGEL}_{C,j})$$

$$\left. \frac{\partial PON}{\partial t} \right|_{\text{Bio}} = \sum_{i=1}^3 (1 - \epsilon) \text{MortPHY}_{N,i} + \sum_{j=1}^2 \left(\frac{\text{MortZOO}_{C,j}}{(C:N)_{ZOO_j}} + \frac{\text{MortGEL}_{C,j}}{(C:N)_{GEL_j}} + (1 - \beta_N) I_{N,j} \right) - \text{HydrolPON} - \text{Graz}_{PON} \quad (15)$$

$$\text{if Gelatinous : } + \sum_{j=1}^2 ((1 - \text{AssEffGEL}_j) \text{GrazGEL}_{N,j})$$

$$85 \quad \left. \frac{\partial SiDet}{\partial t} \right|_{\text{Bio}} = -\text{DissolutionSiDet} + (\text{MortPHY}_{N,3} + \text{Graz}_{NPHY_3})(Si:N)_{PHY_3} \quad (16)$$

$$\left. \frac{\partial ODU}{\partial t} \right|_{\text{Bio}} = \gamma_{ODU}^{POC} \text{AnoxicRespBAC} - \text{ODU}_{oxidNOs} \text{ODU} - \text{ODU}_{oxidO_2} \text{ODU} - \text{SolidODU} \quad (17)$$

$$\left. \frac{\partial O_2}{\partial t} \right|_{\text{Bio}} = \sum_{i=1}^3 (\text{PhotosynthesisPHY}_i + \text{ExtraExcrPHY}_i) \gamma_{O_2}^C - \sum_{j=1}^2 (\text{RespZOO}_j) \gamma_{O_2}^C \\ - \text{RespBAC}_{Oxygen} \gamma_{O_2}^C - \text{NHs}_{oxidO_2} \text{NHs} \gamma_{O_2}^{NHs} - \text{ODU}_{oxidO_2} \text{ODU} \gamma_{O_2}^{ODU} \quad (18)$$

$$\text{if Gelatinous : } - \sum_{j=1}^2 (\text{RespGEL}_j + \text{AdjRespGEL}_j) \gamma_{O_2}^C$$

90 **Optional modules**

Chlorophyll-Geider module

$$\left. \frac{\partial \text{ChlPHY}_i}{\partial t} \right|_{\text{Bio}} = \text{SynthesisChlPHY}_i - \text{RespirationChlPHY}_i - \text{MortChlPHY}_i - \text{Graz}_{\text{ChlPHY}_i} \quad (i = 1, 2, 3) \quad (19)$$

Gelatinous module

$$95 \quad \left. \frac{\partial \text{CGEL}_j}{\partial t} \right|_{\text{Bio}} = \text{GrowthGEL}_{C,j} - \text{MortGEL}_{C,j} - \text{AdjRespGEL}_j \quad (j = 1, 2) \quad (20)$$

Carbonate module

$$\left. \frac{\partial \text{DIC}}{\partial t} \right|_{\text{Bio}} = \sum_{j=1}^2 (\text{RespZOO}_j) + \sum_{j=1}^3 (-\text{GrowthPHY}_j - \text{ExtraExcrPHY}_i) + \text{RespBAC} \quad (21)$$

$$\text{if Gelatinous : } + \sum_{j=1}^2 (\text{RespGEL}_j + \text{AdjRespGEL}_j)$$

$$\left. \frac{\partial \Sigma[-]}{\partial t} \right|_{\text{Bio}} = 0 \quad (22)$$

100 **Aggregation module**

$$\left. \frac{\partial \text{AggNum}}{\partial t} \right|_{\text{Bio}} = \left. \frac{\partial \text{PON}}{\partial t} \right|_{\text{Bio}} \frac{\text{AggNum}}{\text{PON}} - \text{Aggregation} \quad (23)$$



Benthic module

$$\frac{dS_i^j}{dt} = P_i^j - D_i^j - B_i^j \quad \text{with } i = C, Si \text{ and } j = f, s \text{ fast (f) and slowly (s) degraded} \quad (24)$$

$$\begin{aligned} \frac{dS_{N:C}}{dt} &= \frac{d}{dt} \left(\frac{S_N}{S_C} \right) = \frac{1}{S_C^2} \left(S_C \frac{dS_N}{dt} - S_N \frac{dS_C}{dt} \right), \quad \text{with } S_C = S_C^s + S_C^f \text{ and } S_N = S_N^s + S_N^f \\ 105 \quad &= \frac{1}{S_C} \left[(P_N - D_N - B_C \cdot S_{N:C}) - S_{N:C} (P_C - D_C - B_C) \right], \quad \text{with } D_C = D_C^s + D_C^f \text{ and } P_C = P_C^s + P_C^f \\ &= \frac{1}{S_C} \left[(P_N - S_{N:C} P_C) \right] \end{aligned} \quad (25)$$



Table 2. List of model ordinary variables.

VARIABLES	DESCRIPTION	UNITS
f^T	Temperature factor	–
<u>Optic</u>		
Q_{s0}	Solar Radiation at the sea surface	Wm^{-2}
PAR	Photosynthetically Active Radiation [400-700 nm]	Wm^{-2}
k_{short}	Attenuation coef. for short waves PAR ([400-580] nm)	m^{-1}
a_{short}	Total absorption coef. for short waves PAR ([400-580] nm)	m^{-1}
b_{short}	Total backscattering coef. for short waves PAR ([400-580] nm)	m^{-1}
k_{long}	Attenuation coef. for long waves PAR ([580-700] nm)	m^{-1}
a_{long}	Total absorption coef. for long waves PAR ([580-700] nm)	m^{-1}
b_{long}	Total backscattering coef. for long waves PAR ([580-700] nm)	m^{-1}
θ	Solar zenith angle above surface	–
<u>Phytoplankton</u>		
$(Chl : C)_{PHY_i}$	Phytoplankton i Chl a :C ratio	$g\ Chla\ (mol\ C)^{-1}$
$(Chl : N)_{PHY_i}$	Phytoplankton i Chl a :N ratio	$g\ Chla\ (mol\ N)^{-1}$
$(N : C)_{PHY_i}$	Phytoplankton i N:C ratio	$mol\ N\ (mol\ C)^{-1}$
$(N : Si)_{PHY_3}$	Phytoplankton 3 N:Si ratio	$mol\ N\ mol\ Si^{-1}$
$PhotosynthesisPHY_i$	Phytoplankton i photosynthesis rate	$mmol\ C\ m^{-3}\ d^{-1}$
$RespirationPHY_i$	Phytoplankton i respiration rate	$mmol\ C\ m^{-3}\ d^{-1}$
$RespirationChlPHY_i$	Phytoplankton i respiration in $Chla$	$mg\ Chla\ m^{-3},\ d^{-1}$
$LeakagePHY_{C,i}$	Phytoplankton i passive leakage in carbon	$mmol\ C\ m^{-3}\ d^{-1}$
$ExtraExcrPHY_i$	Phytoplankton i extra-excretion	$mmol\ C\ m^{-3}\ d^{-1}$
$NO_3^i\ uptake$	Phytoplankton i nitrate uptake rate	$mmol\ N\ m^{-3}\ d^{-1}$
$NH_4^i\ uptake$	Phytoplankton i ammonium uptake rate	$mmol\ N\ m^{-3}\ d^{-1}$
$SiO_3^i\ uptake$	Diatoms silicate uptake rate	$mmol\ Si\ m^{-3}\ d^{-1}$
v_{Phy_3}	Diatoms sinking rate	$m\ d^{-1}$
$DissolutionSiDet$	Dissolution rate of detrital particulate silicate	$mmol\ Si\ m^{-3}\ d^{-1}$
$LeakagePHY_{N,i}$	Phytoplankton i passive N leakage rate	$mmol\ N\ m^{-3}\ d^{-1}$
$MortPHY_{C,i}$	Phytoplankton i mortality C flux	$mmol\ C\ m^{-3}\ d^{-1}$
$MortPHY_{N,i}$	Phytoplankton i mortality N flux	$mmol\ N\ m^{-3}\ d^{-1}$
$MortPHY_{Chl,i}$	Phytoplankton i mortality Chl a flux	$mg\ Chla\ m^{-3}\ d^{-1}$
<u>Zooplankton</u>		
$GrazZOO_{C,j}$	Grazing C flux of zooplankton j	$mmol\ C\ m^{-3}\ d^{-1}$
$GrazZOO_{N,j}$	Grazing N flux of zooplankton j	$mmol\ N\ m^{-3}\ d^{-1}$

continued on next page ...



... continued from previous page

VARIABLES	DESCRIPTION	UNITS
$Graz_{Prey_i}$	Grazing flux of prey i by all its predators	$\text{mmol m}^{-3}\text{d}^{-1}$
$I_{C,j}$	Zooplankton j C intake	$\text{mmol C m}^{-3}\text{d}^{-1}$
$I_{N,j}$	Zooplankton j N intake	$\text{mmol N m}^{-3}\text{d}^{-1}$
$GrowthZOO_{C,j}$	Zooplankton j growth C flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
$GrowthZOO_{N,j}$	Zooplankton j growth N flux	$\text{mmol N m}^{-3}\text{d}^{-1}$
$ExcrZOO_j$	Zooplankton j excretion flux of ammonium	$\text{mmol N m}^{-3}\text{d}^{-1}$
$RespZOO_j$	Zooplankton j respiration flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
$MortZOO_{C,j}$	Zooplankton j mortality flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
<u>Bacteria</u>		
U_c	Uptake of DOC by bacteria	$\text{mmol C m}^{-3}\text{d}^{-1}$
U_n	Uptake of DON by bacteria	$\text{mmol N m}^{-3}\text{d}^{-1}$
U_A^*	Bacteria potential uptake of ammonium	$\text{mmol N m}^{-3}\text{d}^{-1}$
U_A	Bacteria uptake of ammonium	$\text{mmol N m}^{-3}\text{d}^{-1}$
$GrowthBAC$	Bacteria growth flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
$RespBAC$	Bacteria respiration flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
$ExcrBAC$	Bacteria excretion flux of ammonium	$\text{mmol N m}^{-3}\text{d}^{-1}$
$TestBAC$	intermediary variable	$\text{mmol N m}^{-3}\text{d}^{-1}$
$MortBAC_C$	Bacteria mortality C flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
<u>Detritus</u>		
$HydrolDOCSL$	Hydrolysis rate of Semi-labile DOC into Labile DOC	$\text{mmol C m}^{-3}\text{d}^{-1}$
$HydrolDONSL$	Hydrolysis rate of Semi-labile DON into Labile DON	$\text{mmol N m}^{-3}\text{d}^{-1}$
$HydrolPOC$	Hydrolysis rate of POC into Labile and Semi-Labile DOC	$\text{mmol C m}^{-3}\text{d}^{-1}$
$HydrolPON$	Hydrolysis rate of PON into Labile and Semi-Labile DON	$\text{mmol N m}^{-3}\text{d}^{-1}$
$RespBACOxygen$	Oxic bacterial respiration flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
$RespBACNitrate$	Bacterial respiration flux using nitrate as oxidant	$\text{mmol C m}^{-3}\text{d}^{-1}$
$AnoxicRespBAC$	Anoxic bacterial respiration flux	$\text{mmol C m}^{-3}\text{d}^{-1}$
$SolidODU$	the amount of ODU deposited as solids (metal sulfide formation)	$\text{mmol ODU m}^{-3}\text{d}^{-1}$
<u>Chemical reactions</u>		
$ODUoxid_{O_2}$	oxidation rate of ODU by oxygen	d^{-1}
$ODUoxid_{NO_3}$	oxidation rate of ODU by nitrate	d^{-1}
$NHs oxid_{O_2}$	oxidation rate of NHs by oxygen (Nitrification)	d^{-1}
$NHs oxid_{NO_3}$	oxidation rate of NHs by nitrate	d^{-1}
$Precipitation_{POs}$	Precipitation of POs by iron and manganese hydroxides.	mmol P d^{-1}

continued on next page ...



... continued from previous page

VARIABLES	DESCRIPTION	UNITS
<u>Chlorophyll module</u>		
$SynthesisChlPHY_i$	Phytoplankton i Chla synthesis rate	mg Chla $m^{-3}d^{-1}$
$V_{N,i}$	Phytoplankton carbon-specific nitrogen uptake rate	mol N mol $C^{-1}d^{-1}$
$\rho_{Chl,i}$	Chla synthesis regulation term	-
$RespirationChlPHY_i$	Phytoplankton i respiration	mg Chla $m^{-3}d^{-1}$
$MortChlPHY_i$	Phytoplankton i mortality	mg Chla $m^{-3}d^{-1}$
$Graz_{ChlPHY_i}$	Phytoplankton i grazing	mg Chla $m^{-3}d^{-1}$
<u>Gelatinous module</u>		
$GrazGEL_{C,j}$	Grazing C flux of gelatinous j $j = 1, 2$	mmol C m^{-3}
$GrazGEL_{N,j}$	Grazing N flux of gelatinous j	mmol N m^{-3}
$GrowthGEL_{C,j}$	Gelatinous j growth C flux	mmol C $m^{-3}d^{-1}$
$EgestGEL_{C,j}$	Gelatinous j egestion C flux	mmol C $m^{-3}d^{-1}$
$EgestGEL_{N,j}$	Gelatinous j egestion N flux	mmol N $m^{-3}d^{-1}$
$RespGEL_j$	Gelatinous j respiration C flux	mmol C $m^{-3}d^{-1}$
$AdjExcrGEL_j$	Potential excretion rate necessary to keep the N:C ratio of Gelatinous j constant	mmol N $m^{-3}d^{-1}$
$AdjRespGEL_j$	Potential additional respiration flux necessary to keep the N:C ratio of Gelatinous j constant	mmol C $m^{-3}d^{-1}$
$MortGEL_{C,j}$	Gelatinous j mortality rate	mmol C $m^{-3}d^{-1}$
<u>Carbonate module</u>		
pH	pH	
<u>Aggregation module</u>		
d	Particle size	m
ϵ	Exponent of the particles size distribution	-
$m(d)$	Nitrogen content of a particle of size d	mmol N m^{-3}
$w(d)$	Sinking rate of particles with a size $d < L$	$m d^{-1}$
$n(z, d)$	Particle size distribution at a certain depth z	nb. of partic.
m_l	Mass of a particle of size l (i.e. minimum mass of a single particle)	mmol N m^{-3}
m_L	Mass of a particle of size L	mmol N m^{-3}
w_l	minimum sinking rate of a particle	$m d^{-1}$
w_L	maximum sinking rate of a particle	$m d^{-1}$
$F_L(z)$	Fraction of particles with a size larger than L	-
$G_L(z)$	Fraction of mass of particles with a size larger than L	-

continued on next page ...



... continued from previous page

VARIABLES	DESCRIPTION	UNITS
S_L	$\frac{w_L}{w_i}$	–
$Cells_{PerAggregate}$	Number of Cells per Aggregate	–
Φ	Vertical flux of $AggNum$	nb. of partic. $m^{-2}d^{-1}$
Ψ	Vertical flux of PON	$mmol\ N\ m^{-2}d^{-1}$
w_{POM}	POM sinking rate	$m\ d^{-1}$
w_{AggNum}	Aggregates sinking rate	$m\ d^{-1}$
$Aggregation$	Aggregation rate of particles after collision due to differential settlement and shear	nb. of partic. $m^{-3}d^{-1}$
$AggrSettlement$	Aggregation rate of particles after collision due to differential settlement	nb. of partic. $m^{-3}d^{-1}$
$AggrSinking$	Aggregation rate of particles after collision due to differential shear	nb. of partic. $m^{-3}d^{-1}$
<u>Benthic module</u>		
P_i^j	Benthic-pelagic net exchange of the fraction j of the sedimentary component $i = C, N, Si; j = f, s.$	$mmol\ m^{-2}\ d^{-1}$
D_i^j	Degradation of the fraction j of the sedimentary component $i = C, N, Si;$ $j = f, s.$	$mmol\ m^{-2}\ d^{-1}$
B_i^j	Burial of the fraction j of the sedimentary component $i = C, N, Si; j = f, s.$	$mmol\ m^{-2}\ d^{-1}$
k	Mean first order degradation rate	d^{-1}
τ	Benthic shear stress	$N\ m^{-2}$
p_{nit}	Fraction of NHs oxidised to NOs	
p_{denit}	Fraction of organic matter mineralised by denitrification	
p_{anox}	raction of organic matter mineralised by anoxic degradation	
$p_{SolidDepo}$	Fraction of total production of anoxic substances that is permanently buried and escapes reoxidation	

2.1 The pelagic model

The pelagic part describes the foodweb from bacteria up to mesozooplankton and includes three groups of phytoplankton: diatoms, small and large phototrophic flagellates, two zooplankton groups: micro- and mesozooplankton, an explicit representation of the bacterial loop: bacteria, labile and semi-labile dissolved organic matter, particulate organic matter. Each organic variable is described by its carbon content, except the three groups of phytoplankton, dissolved and particulate detritus that are described by their carbon and nitrogen content. The N:P and, in the case of diatoms, the N:Si ratios are assumed constant. In addition, an optional module allows to explicitly model the Chl a content of each phytoplankton group.




Figure 2. Schematic representation of the trophic relationships in the BAMHBI biogeochemical model. It shows the model state variables (in rectangular boxes) and interactions between them differentiating the inorganic and organic flows made of carbon, nitrogen and phosphorus. The optional gelatinous module is framed in black.

2.1.1 The optical part

The Photosynthetically Active Radiation (PAR) is the fraction of radiation between 400 and 700 nm that can be used by autotrophs for photosynthesis. The sea surface PAR , PAR_0 , is obtained from the solar radiation reaching the sea surface, $Q_{s,0}$ after removing the reflected (*albedo*) and infrared (p_{IR}) fractions (Eq. (38)). The vertical penetration of PAR with depth is governed by the Inherent Optical Properties ($IOPs$) of the medium described here by the absorption (a) and (back)scattering (b) by the optically active components, i.e., here water, phytoplankton and particulate organic carbon (Eqs. (39), (42) and (43)). The absorption and scattering of light are constant for pure sea water (Smith and Baker, 1981), while for dead particles (Neukermans et al., 2012) and phytoplankton, they are represented as the product of a specific rate and the concentration of the optically active component. In the case of phytoplankton, the specific absorption coefficient varies according to a power law of



total chlorophyll concentration (Bricaud et al., 1995) while for the backscattering the effect of the three phytoplankton groups is differentiated (Vaillancourt et al., 2004). In addition, the effect of the Coloured Dissolved Organic Matter (*CDOM*) on light absorption is parameterized based on salinity (Kowalczyk et al., 2003; Mobley et al., 2004; Xu et al., 2005). The attenuation coefficients (k) are then computed using the formulations of Lee et al. (2005)(Eq. (40)). The model does not describe the spectral composition of the light field but rather differentiates two bands of wavelengths of *PAR*: a short [i.e., 400–540 nm] and a long [i.e., 540–700 nm] wavelengths bands. For each of these two bands, the *IOPs* are computed. The model has been tested and fitted with in-situ data.

2.1.2 The autotrophs

The primary producers are divided into three functional groups. The small flagellates lumping the pico-and nanophytoplankton constitute the first group. This group is characterized by high nutrients affinity due to their large surface to volume ratio, high photosynthetic efficiency, high growth and nutrient uptake rates. Large flagellates are lumped into the second group characterized by a high requirement for nutrients, low photosynthetic efficiency and growth rate. The phytoplankton silicifiers (e.g. diatoms) that convert dissolved silicate to particulate silicate and then contribute to the silicate pumping in surface waters constitute the third group. Each group is described by two state variables representing their N and C content. Phytoplankton N:C ratios vary around the Redfield ratio, between the limits $(N:C)_{PHY,min}$ and $(N:C)_{PHY,max}$. The phytoplankton N uptake and photosynthesis are modelled according to the model described in Tett (1998) and Smith and Tett (2000) which is based on the “cell–quota, threshold–limitation” algal growth model of Droop (1983) accounting for the potential control of growth by several nutrients and light. With this formulation, photosynthesis increases with the light availability, the N:C and Chl*a*:C ratios are zero when the N:C ratio reaches its minimum value or when there is no light (Eq. (44)).

Phytoplankton respiration consists of a basal and activity respiration (Eq. (45)). Phytoplankton excretion is the sum of two processes: leakage and active release (Eq. (46)). As in Anderson and Pondaven (2003), we assume that a constant fraction, (parameter γ_1), of all carbon fixed by photosynthesis is leaked from plant cells. In addition, metabolic instabilities in algae, for example, caused by shifts in environmental factors (light, nutrients) cause ‘extra’ carbon to be excreted by phytoplankton (Mague et al., 1980; Williams, 1990). This production of extra carbon is considered as an ‘overflow’ of photosynthesis under nutrient limitation and, thus, is calculated as a fraction (parameter γ_2) of the difference between the nutrient-limited (actual C:N ratio) and nutrient saturated (minimum C:N ratio) growth rate as in Anderson and Pondaven (2003) and Van den Meersche et al. (2004) (Eq. (46)). A fraction (parameter γ_1) of this carbon extra excretion is leaked from the cells and is exported towards the labile DOC (Eq. (10)), while the remainder $(1 - \gamma_1)$ is actively released and is partitioned between the labile (parameter δ_2) and semi-labile $(1 - \delta_2)$ DOC (Anderson and Pondaven, 2003) (Eqs. (10) and (12)).

Inorganic nitrogen can be taken in excess with respect to their immediate need for growth. The luxury uptake is stored and available for later use. For each phytoplankton group, the N:P and N:Si ratios are assumed constant following the Redfield hypothesis and the exchanges of N, P and Si are maintained in balance. The inorganic nutrient uptake increases at low phytoplankton *Nut*:C (*Nut* being N, P or Si) ratios and stops when the *Nut*:C ratio reaches a maximum value. It is not affected by the light intensity and can proceed in the dark (Laws and Wong, 1978). First, potential nitrogen, phosphorus and silicate



uptakes are estimated (Eq. (49)). Then, the realized uptake is governed by the most limiting element along the Liebig law of minimum (Eq. (50)). Ammonium inhibition on nitrate uptake is modelled using an hyperbolic equation. When the $Nut:C$ ratio exceeds the maximum value, an excretion of the nutrient is assumed (Eq. (51)).

The Chl_a of each phytoplankton is computed from the phytoplankton carbon content using a chlorophyll-to-carbon ratio (Chl:C) that is linked to the internal $N : C$ ratio according to a linear equation (Eqs. (47) and (48)). A fraction (ϵ) of phytoplankton mortality (Eq. (53)) contributes to DOM most easily available to bacteria, with the remainder becoming POM (Billen and Fontigny, 1987). The vertical sinking speed of diatoms varies with the internal C:N ratio. The higher this ratio (more nutrient limited), the higher the sinking speed (Eq. (54)).

2.1.3 The heterotrophs

The heterotroph community is described by two zooplankton size groups: the micro- and mesozooplankton, and by bacteria. All the heterotrophic groups are assumed to have a fixed stoichiometry in C, N and P.

2.1.4 Non-gelatinous zooplankton

The description of the growth of the micro- and mesozooplankton is based on the bioenergetic model developed by Anderson (1992). This model simulates the effect of food quality, estimated as the $N:C$ ratio of their food (Eq. (58)), on the growth rates, excretion and respiration (Eqs. (61)–(64)). The ingested food is used for growth and respiration, with a certain amount (parameters β_N and β_C) being undigested (non-assimilated). Messy feeding is considered and parameterized as a constant fraction (parameter ϕ) of grazed material (Eq. (59)). The internal composition of micro- and mesozooplankton is maintained constant through excretion and respiration; if the nitrogen content of the assimilated food is high compared to zooplankton requirements, then excess nitrogen is excreted. On the contrary, if the assimilated food is deficient in nitrogen, then the carbon surplus will be respired (set of Eqs. (61) to (64)).

By default, the mesozooplankton is the highest trophic level and does not have a predator. The mesozooplankton mortality term is one of the closure terms of the model taking into account the predation by fishes. As in Anderson and Pondaven (2003), a density-dependent mortality term is included for the micro- and mesozooplankton functional groups consisting of a Michaelis–Menten function of their biomass with a maximum mortality rate (parameter $mortZOO$) and a half-saturation constant (parameter $ksat_{mortZOO}$). In addition, a mortality term expressed as a function of the O_2 concentration is added representing the rapid mortality of zooplankton in anoxic conditions (Eq. (65)).

2.1.5 Bacteria

Labile DOC and DON are used primarily for bacteria growth. Bacteria consume all the available DOC using DON as a nitrogen source with NHs supplementing DON when the $C : N$ ratio of DOM is high, according to the stoichiometric model of Anderson (1992) (Eqs. (66)–(72)). According to this model, bacterial growth can be either limited by carbon (Eqs. (69)–(71)) or by nitrogen (Eq. (72)). In case of carbon limitation, when bacteria do not require to consume NHs for DOC consumption,



NHs is remineralized (Eqs. (71)) while when an uptake of NHs is needed, bacteria acts as a competitor of phytoplankton for NHs uptake (Eqs. (70)). In case of nitrogen limitation, the uptake of DOC is limited by nitrogen availability considering NHs and *DON* uptake. In that case, there is no excretion of NHs but a consumption (Eq. (72)). Bacterial mortality (Eq. (77)) contributes to the pool of DOM with a fraction (parameter δ_1) to the labile DOM while the remainder adds to the semi-labile DOM.




Figure 3. Schematic representation of the microbial loop in the BAMHBI biogeochemical model. The Figure shows the model state variables (in rectangular boxes) and interactions between them differentiating processes: mortality, hydrolysis, excretion, egestion, (messy) feeding and mortality. The gelatinous optional module is framed and the contribution of the gelatinous to the microbial loop is represented.

2.1.6 The dead organic pool

The dead organic matter is divided into the dissolved and particulate fractions. The dissolved pool is split into the labile (i.e., the only pool taken up directly by bacteria) and the semi-labile (i.e., the pool consisting of molecules whose eventual assimilation by the bacteria requires ectoenzyme hydrolysis to the labile pool) parts. Each dead organic pool is permitted a variable C:N ratio, giving rise to six state variables: DOCL, DONL, DOCSL, DONSL, POC and PON (see Table 1). The N:P ratio is assumed constant.

There are several sources of DOM in the model: phytoplankton excretion and lysis (Eqs. (46) and (53)), zooplankton "messy feeding", bacterial mortality (Eq. (77)) and detrital breakdown (Eqs. (80) and (81)). All these fluxes (except phytoplankton leakage which is assumed to produce only labile DOM) are partitioned between the labile and semi-labile pools with a single




Figure 4. Schematic representation of the flows of inorganic elements in the BAMHBI biogeochemical model showing the model state variables (in rectangular boxes) and interactions between them in anaerobic conditions.

parameter δ_1 as in Anderson and Pondaven (2003). The breakdown of the semi-labile DOM by bacteria into labile DOM is described by Michaelis-Menten kinetics (Eqs. (78) and (79)), with a maximum rate (parameters $MaxHydDOM$) and a half-saturation constant (parameter $k_{sat,hydDOM}$). The organic matter decomposition under anaerobic conditions is not as efficient as in aerobic conditions (Hedges et al., 1999). This lower efficiency is parameterized by multiplying the degradation rates by a
 210 limitation function of O_2 (Eqs. (80) and (81)).

For the dead particulate organic matter we consider a preferential decomposition of N (parameter $MaxHydPON$) relative to C (parameter $MaxHydPOC$) (Anderson and Pondaven, 2003). By default, POC and PON have a constant sinking speed $SinkingRatePOM_{constant}$. If the aggregation module is activated the sinking speed of POC and PON is computed using the aggregation model described in section 2.2.3

215 2.1.7 The inorganic pool

NOs , NHs , POs and for diatoms, $SiOs$, are taken up by phytoplankton according to Eqs. (50) and (51). NHs and POs are excreted by heterotrophs according to the Redfield N:P ratio. $SiOs$ is produced by the chemical dissolution of siliceous detritus (Eq. (82)).

In the oxygenated part of the water column, NHs is converted into NOs through nitrification. This process is described as
 220 first-order in NHs and limited by O_2 concentration (Eq. (83)). Bacterial respiration in C is computed according to Eqs. (69) and (72). The electron acceptor is successively O_2 , NOs and then other oxidants (i.e., MnO_2 , Fe_2O_3 , SO_4) that are not explicitly modelled because they are assumed not limiting. The consumption of O_2 and NOs by bacterial respiration is limited by









Figure 5. Schematic representation of the oxygen flows in the BAMHBI biogeochemical model showing the model state variables (in rectangular boxes) and oxygen flows between them. These oxygen flows include respiration, photosynthesis and oxidation reactions.

respectively the O_2 and NOs availability and additionally, NOs consumption is inhibited by the presence of O_2 (respectively Eqs. (73) and (74)). The approach used to describe the anaerobic degradation by MnO_2 , Fe_2O_3 and SO_4 as oxidants is based
 225 on the equations summarized in Table 3. These anaerobic respiration processes produce reduced substances (i.e., Mn^{2+} , Fe^{3+} and H_2S ; Table 3, Eqs. (29), (31) and (33)). Complete reoxidation of these reduced substances, formed through the oxidation of
 230 x moles of C, requires x moles of O_2 , regardless of the anoxic pathway considered (Table 3, Eqs. (30), (32) and (34)). Hence, as proposed by Soetaert et al. (1996) for the modelling of sediment diagenesis, instead of modeling each of these processes separately, they are lumped together into one process (Table 3, Eqs. (35) and 36), where degradation is only C limited (the quantity of oxidants is assumed non-limiting) and inhibited by O_2 and NOs (Tab. 3, Eq. (75)).

All the reduced substances formed by anoxic mineralisation are lumped into one state variable ODU (i.e. Oxygen Demand Unit) with $ODU = 0.5H_2S + 2Mn^{2+} + 4Fe^{2+}$. The terminal electron acceptors are not explicitly modeled; only the production of reduced substances is described. Anoxic mineralisation of 1 mole of C produces 1 mole of ODU and reoxidation of 1 mole of ODU requires 1 mole of O_2 (Table 1, Eqs. (35) and (36)). The ODU and NHs are directly oxidized by O_2 and NOs .
 235 The oxidation by O_2 is limited by the availability in O_2 while the oxidation by NOs is limited by NOs and inhibited by O_2 (Eqs. (83), (84), (85) and (86)). The denitrification process and the oxidation of NHs and ODU by NOs (respectively



Table 3. Idealized reactions representing the degradation of organic matter using different oxidants. The equations describe successively the aerobic organic matter mineralisation (Eq. (26)), nitrification (Eq. (27)), denitrification (Eq. (28)), and anoxic mineralisation via manganese, iron and sulfate reduction (Eqs. (29), (31) and (33), respectively). These three degradation pathways produce the same amount of CO_2 and the oxidation of the reduced substances they produced (respectively, 2Mn^{2+} , 4Fe^{2+} , 0.5S^{2-}) consume the same amount of O_2 (Eqs. (30), (32) and (34), resp.). They are lumped in the model into Eqs. (35) and (36) where ODU is the Oxygen Demand Unit ($\text{ODU} = 0.5\text{H}_2\text{S} + 2\text{Mn}^{2+} + 4\text{Fe}^{2+}$). y denotes the molar N:P ratio, x the molar C:P ratio in organic matter per mole of phosphorus (for Redfield Stoichiometry, $x = 106$, $y = 16$).



Eqs. (84) and (86)) produce dinitrogen gas and constitute a loss of fixed nitrogen for the system. A part of the formed ODU is permanently removed as metal sulfides where dissolved metals are available. This process is limited by iron availability according to a Michaelis–Menten function. A dissolved iron profile needs to be imposed to the model (Eq. (76)).

240 Phytoplankton

Light penetration

$$f^T = Q_{10}^{\frac{T-20}{10}} \quad (37)$$

$$PAR_0 = (1 - albedo) (1 - p_{IR}) Q_{s,0} \quad (38)$$

$$\frac{\partial PAR}{\partial z} = -PAR_0 \{p_{Long}(k_{Long}) + (1 - p_{Long})(k_{Short})\} \quad (39)$$

245 $k_\lambda = m_0 a_\lambda + m_1 (1 - m_2 \exp[m_3 a_\lambda]) b_\lambda \quad (40)$

$$m_0 = 1 + 0.005\theta \quad (41)$$

$$a_\lambda = a_{\lambda,sw} + a_{\lambda,chl}^{*} Chl^{\hat{a}_\lambda, Chl} + [a_{\lambda,CDOM}^{*} - a_{\lambda,CDOM}^{**} * Salinity] + a_{\lambda,POC}^{*} POC \quad (42)$$



$$b_{\lambda} = b_{\lambda,sw} + \sum_{i=1}^3 (b_{\lambda,PHY_i}^* CPHY_i) + b_{\lambda,POC}^* POC \quad (43)$$

Photosynthesis

$$250 \quad \text{Photosynthesis}PHY_i = P_{max,i} \lim_{light,i} CPHY_i \quad \text{for } (i=1, 2, 3) \quad (44)$$

$$\text{with } \begin{cases} \lim_{light,i} = 1 - \exp\left[\frac{-\alpha_i^{Chl} PAR}{\mu_{max,i}}\right] & \text{for } (i=1, 2, 3) \\ P_{max,i} = \mu_{max,i} f^T \lim_{nut,i} & \text{for } (i=1, 2, 3) \\ \lim_{nut,i} = \left(1 - \frac{(N:C)_{PHY_i,min}}{(N:C)_{PHY_i}}\right) & \text{for } (i=1, 2, 3) \end{cases}$$

$$\text{Respiration}PHY_i = \text{ActivityResp}PHY_i + \text{BasalResp}PHY_i \quad (45)$$

$$\text{with } \begin{cases} \text{ActivityResp}PHY_i = \text{ProdResp}_i \text{Photosynthesis}PHY_i \\ \text{BasalResp}PHY_i = \text{Resp}_i f^T CPHY_i \end{cases}$$

$$255 \quad \text{Excretion}CPHY_i = \text{Leakage}CPHY_i + \text{ExtraErr}CPHY_i \quad (46)$$

$$\text{with } \begin{cases} \text{Leakage}CPHY_i = \gamma_1 \text{Photosynthesis}PHY_i \\ \text{ExtraErr}CPHY_i = \gamma_2 \left[\text{Photosynthesis}PHY_i \left| \frac{(N:C)_{PHY_i}}{(N:C)_{PHY_i,max}} - \text{Photosynthesis}PHY_i \right| \right] \end{cases}$$

$$\text{Chlorophyll} = \sum_{i=1}^3 CPHY_i (Chl:C)_{PHY_i} \quad (47)$$

Chl:C

$$260 \quad \begin{cases} \text{for } (N:C)_{PHY_i,min} \leq (N:C)_{PHY_i} \leq (N:C)_{PHY_i,max} : \\ (Chl:C)_{PHY_i} = (N:C)_{PHY_i} \left\{ (Chl:N)_{PHY_i,min} + \left[(Chl:N)_{PHY_i,max} \dots \right. \right. \\ \left. \left. \dots - (Chl:N)_{PHY_i,min} \right] \times \frac{(N:C)_{PHY_i} - (N:C)_{PHY_i,min}}{(N:C)_{PHY_i,max} - (N:C)_{PHY_i,min}} \right\} \\ \text{for } (N:C)_{PHY_i} \leq (N:C)_{PHY_i,min} \\ (Chl:C)_{PHY_i} = (N:C)_{PHY_i,min} (Chl:N)_{PHY_i,min} \\ \text{for } (N:C)_{PHY_i} \geq (N:C)_{PHY_i,max} \\ (Chl:C)_{PHY_i} = (N:C)_{PHY_i,max} (Chl:N)_{PHY_i,max} \end{cases} \quad (48)$$

Nutrient Uptake : NOs_i^{uptake} , NHs_i^{uptake} , POs_i^{uptake} , $SiOs_3^{uptake}$

for $(Nut:C)_{PHY_i} \leq (Nut:C)_{PHY_i,max}$ with Nut=N, P, Si (for diatoms)

$$265 \quad \begin{cases} NOs_i^{uptake}|_{potential} = NO_{max,i} f^T \left(1 - \frac{(N:C)_{PHY_i}}{(N:C)_{PHY_i,max}} \right) \frac{NOs}{NOs+k_{NOs_i}} \frac{k_{in}}{k_{in}+NHs} CPHY_i \\ NHs_i^{uptake}|_{potential} = NH_{max,i} f^T \left(1 - \frac{(N:C)_{PHY_i}}{(N:C)_{PHY_i,max}} \right) \frac{NHs}{NHs+k_{NHs_i}} CPHY_i \\ POs_i^{uptake}|_{potential} = P_{max,i} f^T \left(1 - \frac{(P:C)_{PHY_i}}{(P:C)_{PHY_i,max}} \right) \frac{POs}{POs+k_{POs_i}} CPHY_i \\ SiOs_3^{uptake}|_{potential} = Si_{max,i} f^T \left(1 - \frac{(Si:C)_{PHY_i}}{(Si:C)_{PHY_i,max}} \right) \frac{SiOs}{SiOs+k_{SiOs_3}} CPHY_i \quad \text{for diatoms } i=3 \end{cases} \quad (49)$$



$$\left\{ \begin{array}{l} \text{Nitrogen}_{PHY_i}^{\text{uptake}} = \min \left(\text{NO}_{s_i}^{\text{uptake}}|_{\text{potential}} + \text{NH}_{s_i}^{\text{uptake}}|_{\text{potential}}, \frac{\text{PO}_{s_i}^{\text{uptake}}|_{\text{potential}}}{(P:N)_{PHY_i}}, \delta_{i,3} \frac{\text{SiO}_{s_3}^{\text{uptake}}|_{\text{potential}}}{(Si:N)_{PHY_3}} \right) \\ \text{NO}_{s_i}^{\text{uptake}} = \frac{\text{Nitrogen}_{PHY_i}^{\text{uptake}}}{\text{NO}_{s_i}^{\text{uptake}}|_{\text{potential}} + \text{NH}_{s_i}^{\text{uptake}}|_{\text{potential}}} \text{NO}_{s_i}^{\text{uptake}}|_{\text{potential}} \\ \text{NH}_{s_i}^{\text{uptake}} = \frac{\text{Nitrogen}_{PHY_i}^{\text{uptake}}}{\text{NO}_{s_i}^{\text{uptake}}|_{\text{potential}} + \text{NH}_{s_i}^{\text{uptake}}|_{\text{potential}}} \text{NH}_{s_i}^{\text{uptake}}|_{\text{potential}} \\ \text{PO}_{s_i}^{\text{uptake}} = \frac{\text{Nitrogen}_{PHY_i}^{\text{uptake}}}{(P:N)_{PHY_i}} \\ \text{SiO}_{s_3}^{\text{uptake}} = \left[\frac{\text{Nitrogen}_{PHY_3}^{\text{uptake}}}{(Si:N)_{PHY_3}} - \text{LeakageNPHY}_3 \right] (Si:N)_{PHY_3} \end{array} \right. \quad (50)$$

270

Excretion

for $(\text{Nut} : C)_{PHY_i} \geq (\text{Nut} : C)_{PHY_i, \text{max}}$ with Nut=N, P, Si (for diatoms)

We have a negative uptake that corresponds to an excretion

$$\left\{ \begin{array}{l} \text{NO}_{s_i}^{\text{uptake}} = 0 \\ \text{NH}_{s,3}^{\text{uptake}}|_{\text{potential}} = \text{NH}_{\text{max}_i} f^T \left(1 - \frac{(N:C)_{PHY_i}}{(N:C)_{PHY_i, \text{max}}} \right) C_{PHY_i} \\ \text{PO}_{s_i}^{\text{uptake}}|_{\text{potential}} = \text{P}_{\text{max}_i} f^T \left(1 - \frac{(P:C)_{PHY_i}}{(P:C)_{PHY_i, \text{max}}} \right) C_{PHY_i} \\ \text{SiO}_{s_3}^{\text{uptake}}|_{\text{potential}} = \text{SiO}_{\text{max}_3} f^T \left(1 - \frac{(Si:C)_{PHY_3}}{(Si:C)_{PHY_3, \text{max}}} \right) C_{PHY_3} \\ \text{NH}_{s_i}^{\text{uptake}} = \min \left(\text{NH}_{s_i}^{\text{uptake}}|_{\text{potential}}, \frac{\text{PO}_{s_i}^{\text{uptake}}|_{\text{potential}}}{(P:N)_{PHY_i}}, \delta_{i,3} \frac{\text{SiO}_{s_3}^{\text{uptake}}|_{\text{potential}}}{(Si:N)_{PHY_3}} \right) \\ \text{PO}_{s_i}^{\text{uptake}} = \frac{\text{NH}_{s_i}^{\text{uptake}}}{(P:N)_{PHY_i}}, \quad \text{SiO}_{s_3}^{\text{uptake}} = \frac{\text{NH}_{s_i}^{\text{uptake}}}{(Si:N)_{PHY_3}} \end{array} \right. \quad (51)$$

275

Leakage

$$\left\{ \begin{array}{l} \text{LeakageNPHY}_i = \gamma_1 \left(\text{NH}_{s,1}^{\text{uptake}} + \text{NO}_{s,1}^{\text{uptake}} \right) \\ \text{LeakageNPHY}_i = 0 \text{ when } (N:C)_{PHY_i} > (N:C)_{PHY_i, \text{max}} \end{array} \right. \quad (52)$$

Mortality

$$280 \quad \text{MortPHY}_{X,i} = \text{mort}_{PHY_i} f^T X_{PHY_i}, \quad X = C, N, \text{Chl} \quad (53)$$

Sedimentation of Diatoms

$$v_{PHY_3} = v_{PHY_3, \text{min}} + \left(v_{PHY_3, \text{max}} - v_{PHY_3, \text{min}} \right) \frac{\left((N:C)_{PHY_3, \text{max}} - (N:C)_{PHY_3} \right)}{\left((N:C)_{PHY_3, \text{max}} - (N:C)_{PHY_3, \text{min}} \right)} \quad (54)$$

Zooplankton

285

Feeding, ($i = 1, 2$)

$$\text{GrazZOO}_{C,i} = f^T \text{MaxGrazZOO}_i \frac{Ba_{C,ZOO_i}}{Ba_{C,ZOO_i} + k_{\text{sat},ZOO_i}} CZOO_i \quad (55)$$

$$\text{GrazZOO}_{N,i} = \text{GrazZOO}_{C,i} (N:C)_{\text{food},i} \quad (56)$$

$$Ba_{X,ZOO_i} = \sum_{\text{preys}} e_{\text{Prey},ZOO_i} X_{\text{Prey}} \quad X = C, N \quad (57)$$

$$290 \quad (N:C)_{\text{food},i} = Ba_{N,ZOO_i} / Ba_{C,ZOO_i} \quad (58)$$

$$I_{X,i} = (1 - \phi_i) \text{GrazZOO}_{X,i} \quad X = C, N \quad (59)$$



$$Graz_{Prey} = \sum_{i=1}^2 \left[\frac{GrazZOO_{C,i}}{Ba_{C,ZOO_i}} e_{Prey,ZOO_i} Prey \right] \quad Prey = CPHY_i, NPHY_i, ChlPHY_i, CBAC, POC, PON, CZOO_1 \quad (60)$$

Growth, ($i = 1, 2$)

$$(N : C)_i^* = (N : C)_{ZOO_i} \frac{k_{c,i} \beta_{C,i}}{\beta_{N,i}} \rightarrow \begin{cases} (N : C)_{food,i} < (N : C)_i^* & \Rightarrow \text{N limitation} \\ (N : C)_{food,i} > (N : C)_i^* & \Rightarrow \text{C limitation} \end{cases} \quad (61)$$

295

$$\text{If N limits: } \begin{cases} GrowthZOO_{N,i} = \beta_{N,i} I_{N,i} \\ GrowthZOO_{C,i} = GrowthZOO_{N,i} / (N : C)_{ZOO} \\ ExcrZOO_i = 0 \end{cases} \quad (62)$$

$$\text{If C limits: } \begin{cases} GrowthZOO_{C,i} = k_{c,i} \beta_{C,i} I_{C,i} \\ GrowthZOO_{N,i} = GrowthZOO_{C,i} (N : C)_{ZOO_j} \\ ExcrZOO_i = \beta_{N,i} I_{N,i} - GrowthZOO_{N,i} \end{cases} \quad (63)$$

$$RespZOO_i = \beta_{C,i} I_{C,i} - GrowthZOO_{C,i} \quad (64)$$

Mortality, ($i = 1, 2$)

300

$$MortZOO_{C,i} = f^T CZOO_i \left(mortZOO_i \frac{CZOO_i}{CZOO_i + ksat_{MortZOO_i}} + \left(1 - \frac{O_2}{O_2 + ksat_{AnoxMort}} \right) AnoxMort \right) \quad (65)$$

Microbial Loop

305

Bacteria Growth, respiration, excretion and mortality.

$$U_c = f^T \mu_{BAC} \frac{DOCL}{k_{BAC,DOC} + DOCL} CBAC \quad , \quad U_n = U_c \left(\frac{DONL}{DOCL} \right) \quad (66)$$

$$U_A^* = f^T \mu_{BAC} \frac{NHs}{k_{BAC,NHs} + NHs} CBAC (N : C)_{BAC} \quad (67)$$

$$TestBAC = U_n - \omega_B U_c (N : C)_{BAC} \quad (68)$$

$$\text{if } U_A^* > -TestBAC \rightarrow \text{Carbon limitation case:} \quad (69)$$

310

$$GrowthBAC = \omega_B U_c$$

$$RespBAC = (1 - \omega_B) U_c$$

$$\text{if } TestBAC < 0 \rightarrow \text{NHs uptake is necessary to consume all the DOCL} \quad (70)$$

$$U_A = -TestBAC$$

$$ExcrBAC = 0$$

315

$$\text{if } TestBAC > 0 \rightarrow \text{Bacteria act as remineralizers:} \quad (71)$$

$$U_A = 0$$

$$ExcrBAC = TestBAC$$

$$\text{if } U_A^* \leq -TestBAC \rightarrow \text{Nitrogen Limitation case:} \quad (72)$$

$$U_A = U_A^*$$

$$GrowthBAC = (U_n + U_A) / (N : C)_{BAC}$$

$$RespBAC = GrowthBAC (1 / \omega_B - 1)$$

$$ExcrBAC = 0$$

320



$$RespBAC_{Oxygen} = RespBAC \frac{O_2}{O_2 + k_{sat,O_2}^{OxicMin}} \quad (73)$$

$$RespBAC_{Nitrate} = RespBAC \frac{NOs}{NOs + k_{sat,NOs}^{Denit}} \left(1 - \frac{O_2}{O_2 + k_{in,O_2}^{Denit}} \right) \quad (74)$$

$$325 \quad AnoxicRespBAC = RespBAC \left(1 - \frac{NOs}{NOs + k_{in,NOs}^{AnoxMin}} \right) \left(1 - \frac{O_2}{O_2 + k_{in,O_2}^{AnoxMin}} \right) \quad (75)$$

$$SolidODU = R_{solid} IronLimitation \gamma_{POC}^{ODU} AnoxicRespBAC \quad (76)$$

with $IronLimitation = \frac{Iron}{Iron + k_{sat,iron}^{SolForm}}$

where Iron concentration is taken from the data

330

$$MortBAC = f^T mort_{BAC} CBAC \quad (77)$$

Labile and Semi-labile Dissolved Organic Matter

$$HydrolDOCSL = f^T MaxHydDOM \frac{DOCSL}{DOCSL + k_{sat,hydDOM}} CBAC \quad (78)$$

$$335 \quad HydrolDONSL = HydrolDOCSL \frac{DONSL}{DOCSL} \quad (79)$$

Particulate Detrital Matter

$$HydrolPOC = f^T MaxHydPOC \frac{O_2 + 1}{O_2 + 1 + k_{sat,OxicHydrol}} POC \quad (80)$$

$$HydrolPON = f^T MaxHydPON \frac{O_2 + 1}{O_2 + 1 + k_{sat,OxicHydrol}} PON \quad (81)$$

$$340 \quad DissolutionSidet = f^T diss_{SiDet} SiDet \quad (82)$$

Chemical submodel

$$NHs_{oxidO_2} = f^T \mu_{NHs}^{O_2} \frac{O_2}{O_2 + k_{sat,O_2}^{NHs_{oxid}}} \quad (83)$$

$$NHs_{oxidNOs} = f^T \mu_{NHs}^{NOs} \frac{NOs}{NOs + k_{sat,NOs}^{NHs_{oxidNOs}}} \left(1 - \frac{O_2}{O_2 + k_{in,O_2}^{NHs_{oxidNOs}}} \right) \quad (84)$$

$$345 \quad ODU_{oxidO_2} = f^T \mu_{ODU}^{O_2} \frac{O_2}{O_2 + k_{sat,O_2}^{ODU_{oxid}}} \quad (85)$$

$$ODU_{oxidNOs} = f^T \mu_{ODU}^{NOs} \frac{NOs}{NOs + k_{sat,NOs}^{ODU_{oxidNOs}}} \left(1 - \frac{O_2}{O_2 + k_{in,O_2}^{ODU_{oxidNOs}}} \right) \quad (86)$$

2.2 BAMHBI Modules

BAMHBI includes five optional modules that are summarized in Table 9 and described here below. The inclusion of such modules is explained in this section.



350 2.2.1 The chlorophyll module

If the chlorophyll module is activated, the Chla content of each phytoplankton group is dynamically computed using the formulation of Geider et al. (1998). Each phytoplankton group is then described by three state variables representing their N, C and Chla content. The dynamics of each phytoplankton group mechanistically represents the physiological acclimation process at cellular level that takes place in algae and consists in changing the cellular biochemical composition Chla:C and C:N
 355 of algae as a function of external environmental variables: light, inorganic nutrients and temperature. The synthesis of Chla is downregulated in conditions of inorganic nutrients limitations and is enhanced when light is limiting.

Chlorophyll

$$lim_{light,i} = 1 - \exp\left(\frac{-\alpha_i^{Chl}(Chl : C)_{PHY_i} PAR}{\mu_{max,i}}\right) \quad (87)$$

$$(Chl : C)_{max} = (Chl : N)_{max} (N : C)_{max} \quad (88)$$

$$360 (Chl : C)_{min} = (Chl : N)_{min} (N : C)_{min} \quad (89)$$

$$SynthesisChlPHY_i = \frac{\rho_{Chl,i} V_{N,i}}{(Chl : C)_{PHY_i}} ChlPHY_i \quad (90)$$

$$\text{If Nitrogen Uptake} > 0 \rightarrow V_{N,i} = \frac{Nitrogen_{PHY_i}^{uptake}}{CPHY_i} \quad (91)$$

$$\text{If Nitrogen Uptake} < 0 \rightarrow V_{N,i} = 0 \quad (92)$$

$$\text{If PAR} > 0 \rightarrow \rho_{Chl,i} = \frac{P_{max,i} lim_{light,i} (Chl : N)_{max}}{\alpha_i^{Chl} (Chl : C)_{PHY_i} PAR} \quad (93)$$

$$365 \text{ If PAR} = 0 \rightarrow \rho_{Chl,i} = 0 \quad (94)$$

$$RespirationChlPHY_i = Resp_i f^T ChlPHY_i \quad (95)$$

$$Chlorophyll = \sum_{i=1}^3 ChlPHY_i \quad (96)$$

Equations above describe the Geider module. The two mains differences with the other equation are

- 370 1. The chlorophyll content of each phytoplankton group are modelled as state variables
2. The photosynthesis - light curve depends now on the (Chl:C) ratio as shown by Eq. 87

In addition to these major changes, the (Chl:C) ratio is bounded between a minimal and maximal ratios computed by Eqs. 89 and 88. These bounds, also used in Aumont et al. (2015) and Butenschön et al. (2016), prevent the regulation term Eq. 93 and Eq. 90 from being to large or too small. The nitrogen to carbon ratio used for computing the carbon uptake Eq. 44 is bounded
 375 while it is not in the computation of nutrients uptake in Eq. 49. For diatoms, the (Si:C) computation based on the (N:C) ratio and the Redfield ratio do not use the bounded (N:C) ratio.



2.2.2 The gelatinous zooplankton module

When the `gelatinous` module is included, a carnivorous and an omnivorous group are added to the model. Both gelatinous groups are ‘top-predators’ and there is no trophic link between them. They have a grazing rate that linearly increases with prey concentrations without saturation and with no feeding when the food concentration is below a given threshold. The omnivorous gelatinous feeds on the three phytoplankton functional types, micro- and mesozooplankton and detritus while the carnivorous gelatinous can only feed on mesozooplankton. The internal composition of gelatinous organisms is held constant by adjusting the excretion and respiration terms (set of Eqs. (105), (106) and (107)). They are not consumed and represent trophic dead-ends. The mortality terms of the two gelatinous groups are represented by the sum of a first-order kinetics as in Lancelot et al. (2002) and an additional term traducing the rapid mortality of these organisms in anoxic conditions (Eq. (108)).

Gelatinous, ($j=1, 2$)

$$GrazGEL_{C,j} = f^T MaxGrazGEL_j Ba_{c,GEL_j} CGEL_j \quad (97)$$

$$GrazGEL_{N,j} = GrazGEL_{C,j} (N:C)_{food,j} \quad (98)$$

$$Ba_{x,GEL_j} = \sum_{preys} e_{Prey,GEL_j} X_{Prey}, \quad X = C, N \quad (99)$$

$$(N:C)_{food,GEL_j} = Ba_{n,GEL_j} / Ba_{c,GEL_j} \quad (100)$$

$$GrowthGEL_{C,j} = GrowthEffGEL_j AssEffGEL_j GrazGEL_{C,j} \quad (101)$$

$$EgestGEL_{C,j} = (1 - AssEffGEL_j) GrazGEL_{C,j} \quad (102)$$

$$EgestGEL_{N,j} = (1 - AssEffGEL_j) GrazGEL_{N,j} \quad (103)$$

$$RespGEL_j = (1 - GrowthEffGEL_j) AssEffGEL_j GrazGEL_{C,j} \quad (104)$$

$$(N:C)^* = \frac{GrazGEL_{C,j}(N:C)_{food,GEL_j} - EgestGEL_{N,j}}{GrazGEL_{C,j} - EgestGEL_{C,j} - RespGEL_j} \quad (105)$$

if $(N:C)_{GEL} < (N:C)^*$

$$\rightarrow \begin{cases} AdjRespGEL_j = 0 \\ AdjExcrGEL_j = (GrazGEL_{C,j}(N:C)_{food,GEL_j} - EgestGEL_{N,j}) \\ \quad - (N:C)_{GEL_i} (GrazGEL_{C,j} - EgestGEL_{C,j} - RespGEL_{C,j}) \end{cases} \quad (106)$$

400 if $(N:C)_{GEL} > (N:C)^*$

$$\rightarrow \begin{cases} AdjRespGEL_j = (GrazGEL_{C,j} - EgestGEL_{C,i} - RespGEL_j) \\ \quad - (GrazGEL_{C,j}(N:C)_{food,GEL_j} - EgestGEL_{N,j}) / (N:C)_{GEL_i} \\ AdjExcrGEL_j = 0 \end{cases} \quad (107)$$

$$MortGEL_{C,j} = f^T CGEL_j \left(mortGEL_j + \left(1 - \frac{O_2}{O_2 + ksat_{AnoxMort}} \right) AnoxMort \right) \quad (108)$$



$$Graz_{Prey} = \sum_{i=1}^2 \left[\frac{GrazZOO_{C,i}}{Ba_{C,ZOO_i}} e_{Prey,ZOO_i} Prey + \frac{GrazGEL_{C,i}}{Ba_{C,GEL_i}} e_{Prey,GEL_i} Prey \right] \quad (109)$$

405 with Prey = $CPHY_i, NPHY_i, ChlPHY_i, CBAC, POC, PON, CZOO_1, CZOO_2$

If the gelatinous module is not included, the effects of gelatinous on the trophic foodweb (grazing, excretion, mortality, respiration) are not taken into account without any compensation.

2.2.3 The aggregation module

When the aggregation module is included, BAMHBI simulates the formation of marine aggregates from detritus particles according to the aggregation model of Kriest (2002) (set of Eqs. (110) to (124)). Marine snow is described by two state variables: its mass (PON) and number of particles (i.e., state variable, $AggNum$). From that, it is assumed that the distribution of aggregate's size can be fully computed. The number of particles, $AggNum$, varies according to the production and destruction of PON and is reduced by aggregation into larger particles. Larger particles are formed by collision and adherence of smaller particles. There are two mechanisms that are responsible for collision: differential settlement and turbulent shear. Collisions due to turbulent shear are expected to be higher in the mixed layer where the shear is higher while collisions due to different sinking speeds is expected to dominate in the deeper layers (Kriest, 2002). As the number of particles varies their size distribution and sinking speed are also modified. The average particle size and sinking speed of PON and $AggNum$ are computed. It assumes that the number of particles $n(d_i)$ of size d_i (i.e. the size distribution spectrum), the mass of particles $m(d_i)$ (nitrogen content) and their sinking speeds $w(d_i)$ can be represented by a two-parameter power law of the following form: $y(d_i) = C d_i^{-x}$, where d_i is the diameter of a particle. The parameters C and x have been taken from Kriest (2002) and are derived from in-situ observations of marine snow; y stands for either m , n or w . The exponent describing the size distribution of particles depends on the average size of aggregates. ϵ , the exponent that describes the particles size distribution, is computed as a function of the number of cells per aggregate (Eq. (119)) according to Eq. (120).

Aggregates are assumed to be produced/removed at the same rate as detritus (Eq. (23)). The stickiness, (parameter $stick$), i.e., the probability that two particles stick together after contact may vary from 0 to almost 1. This parameter has been found to have a strong influence on the vertical export of organic matter and has been calibrated in order to reproduce sediment trap data.

Size distribution properties

$$m(d) = C d^\zeta \quad m_l = C l^\zeta; \quad m_L = C L^\zeta \quad (110)$$

$$w(d) = B d^\eta \quad w_l = B l^\eta; \quad w_L = B L^\eta; \quad S_L = \frac{w_L}{w_l} = \frac{B L^\eta}{B l^\eta} = \left(\frac{L}{l}\right)^\eta \quad (111)$$

$$n(d) = A d^{-\epsilon} \quad (112)$$

Estimation of ϵ and sinking velocity



$$AggNum = \int_l^\infty n(d) dd = \int_l^\infty d^{-\epsilon} dd = A \frac{l^{1-\epsilon}}{\epsilon-1} \text{ provided that } \epsilon > 1 \quad (113)$$

$$AggNum_{d>L} = \int_L^\infty n(d) dd = A \int_L^\infty d^{-\epsilon} dd = A \frac{L^{1-\epsilon}}{\epsilon-1} \text{ provided that } \epsilon > 1 \quad (114)$$

$$435 \quad F_L = \frac{A \frac{L^{1-\epsilon}}{\epsilon-1}}{A \frac{l^{1-\epsilon}}{\epsilon-1}} = \left(\frac{L}{l}\right)^{1-\epsilon} \quad (115)$$

$$PON = \int_l^\infty n(d) m(d) dd = AC \int_l^\infty d^{\zeta-\epsilon} dd = A m_l \frac{l^{1-\epsilon}}{\epsilon-1-\zeta} \text{ provided that } \epsilon > \zeta + 1 \quad (116)$$

$$PON = \int_L^\infty n(d) m(d) dd = AC \int_L^\infty d^{\zeta-\epsilon} dd = A m_l \frac{L^{1-\epsilon}}{\epsilon-1-\zeta} \text{ provided that } \epsilon > \zeta + 1 \quad (117)$$

$$G_L = \frac{A m_l \frac{L^{1-\epsilon}}{\epsilon-1-\zeta}}{A m_l \frac{l^{1-\epsilon}}{\epsilon-1-\zeta}} = \left(\frac{L}{l}\right)^{1+\zeta-\epsilon} \quad (118)$$

$$\text{if } AggNum > 0 \quad Cells_{PerAggregate} = \frac{PON}{AggNum m_l} \quad \text{else} \quad Cells_{PerAggregate} = 1 \quad (119)$$

$$440 \quad \text{if } Cells_{PerAggregate} > 1 \quad \epsilon = \frac{Cells_{PerAggregate} (1 + \zeta) - 1}{Cells_{PerAggregate} - 1} \quad \text{else} \quad \epsilon = 100 \quad (120)$$

$$\Psi = \left[\int_l^L n(d) w(d) dd + \int_L^\infty n(d) w_L dd \right] = AggNum \left[\frac{(\epsilon-1)v_l}{(\epsilon-\eta-1)} - \frac{F_L \eta v_L}{(\epsilon-\eta-1)} \right] \text{ assuming that for } d < L, w \neq w_L \quad (121)$$

$$\Phi = \left[\int_l^L n(d) w(d) m(d) dd + \int_L^\infty n(d) w_L m(d) dd \right] = PON \left[\frac{(\epsilon-\zeta-1)v_l}{(\epsilon-\eta-\zeta-1)} - \frac{G_L \eta v_L}{(\epsilon-\zeta-\eta-1)} \right] \quad (122)$$

$$v_{AggNum} = \frac{(\epsilon-1)v_l}{(\epsilon-\eta-1)} - \frac{F_L \eta v_L}{(\epsilon-\eta-1)}; \quad v_{POM} = \frac{(\epsilon-\zeta-1)v_l}{(\epsilon-\eta-\zeta-1)} - \frac{G_L \eta v_L}{(\epsilon-\zeta-\eta-1)} \quad (123)$$

Aggregation rates

$$445 \quad Aggregation = 0.5 stick (AggregSettlement + AggregShear) \quad (124)$$

where AggregSettlement and AggregShear are the number of particles colliding due to respectively differential settlement and shear rate as in Kriest (2002)

2.2.4 The carbonate module

When the carbonate module is activated, the carbonate system is solved, allowing a complete specification of *DIC* and the estimation of *pH*, Total alkalinity (TA) and of the air-sea exchange of CO_2 . The carbonate module is based on Soetaert et al. (2007). It solves two additional state variables (i.e. *DIC* and the "excess negative charge" Σ) and major acid-base reactions that affect the *pH* dynamics in the ocean. In the ocean, DIC is present in three forms: aqueous carbon dioxide $CO_{2(aq)}$, bicarbonate HCO_3^- , and carbonate ion CO_3^{2-} . A fourth form is H_2CO_3 (the true carbonic acid) but its concentration is much smaller than



that of $\text{CO}_{2(\text{aq})}$ (less than 0.3%). Therefore, the CO_2 notation usually refers to the sum of the two electrically neutral forms
 455 $\text{CO}_{2(\text{aq})}$ and H_2CO_3 .

Table 4 lists the acid–base reactions taken into account in BAMHBI. This list includes the dissociation and protonation of carbonates (Eqs. (125), (126)), water (Eq. (127)), borates (Eq. (128)), phosphates (Eqs. (129), (130) and (131)), silicates (Eqs. (132) and (133)), ammonium (Eq. (134)) and sulfide (Eqs. (136) and (137)). Nitrate is assumed to be fully dissociated (Eq. (135)) and in BAMHBI, the dissociation of nitrite, sulfates and fluoride is neglected.

460 Chemical reactions presented in Table 4 generally occur at rates which, compared with the slow biogeochemical reactions that consume or produce protons, are fast enough to consider that they operate at equilibrium. In many of these reversible reactions the effective equilibrium is reached within seconds to minutes (e.g. Wolf-Gladrow and Riebesell, 1997). The stoichiometric equilibrium speciation is expressed in terms of the total concentrations of species ($\sum X = \sum_i [H_i X^{i-a}]$) (Table 5). These lump sums of concentrations ($\sum X$) are (related to) model state variables. The equilibrium constants (K_*) are calculated as a
 465 function of temperature, salinity and pressure as given by Millero (1995) with typographical corrections according to Lewis and Wallace (1998); they are expressed on the total hydrogen ion pH scale. The borate concentrations are calculated as a function of salinity as described by Millero (1995).

Knowing the formulations for the equilibrium concentration of all the acid–base species, the dissociation constants and the lump sums of concentrations of species, the system of equations in Table 4 allows to solve the concentrations of all the ions and
 470 the proton concentration H^+ . As there are 16 unknowns, one more than the number of equations, an extra equation is required to find a unique solution. Usually, this additional equation is provided by TA:

$$TA = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] + [\text{HPO}_4^{2-}] + 2[\text{PO}_4^{3-}] + [\text{H}_3\text{SiO}_4^-] + 2[\text{H}_2\text{SiO}_4^{2-}] + [\text{NH}_3] + [\text{HS}^-] + 2[\text{S}^{2-}] - [\text{H}^+] - [\text{H}_3\text{PO}_4] - [\text{HSO}_4^-] - [\text{HF}] - [\text{HNO}_3] - [\text{HNO}_2] - 2[\text{H}_2\text{SO}_4] \quad (138)$$

In BAMHBI, the set of equations is completed by an additional equation for the 'excess negative charge' (denoted $\Sigma[-]$) as
 475 described in Soetaert et al. (2007). This quantity expresses the excess of negative charge as the difference in concentrations of negative charges over positive charges (both multiplied by their charge) of the acid-base system given in Table 4:

$$\Sigma[-] = [\text{HCO}_3^-] + 2[\text{CO}_3^{2-}] + [\text{B}(\text{OH})_4^-] + [\text{OH}^-] + [\text{H}_2\text{PO}_4^-] + 2[\text{HPO}_4^{2-}] + 3[\text{PO}_4^{3-}] + [\text{H}_3\text{SiO}_4^-] + 2[\text{H}_2\text{SiO}_4^{2-}] + [\text{NO}_3^-] + [\text{NO}_2^-] + [\text{F}^-] + [\text{HS}^-] + 2[\text{S}^{2-}] + [\text{HSO}_4^-] + 2[\text{SO}_4^{2-}] - [\text{NH}_4^+] - [\text{H}^+] \quad (139)$$

If we assume that uptake of ions is compensated by uptake or release of protons (electroneutrality), $\Sigma[-]$ is not impacted by
 480 changes in the concentrations of nitrate, nitrite, phosphate, ammonia/ammonium which is not the case of TA. It is straightforward to convert the $\Sigma[-]$ to TA:

$$TA = \Sigma[-] + \Sigma\text{NH}_3 - \Sigma\text{NO}_3 - \Sigma\text{NO}_2 - \Sigma\text{PO}_4 - 2\Sigma\text{SO}_4 - \Sigma\text{F} \quad (140)$$

This equation needs to be adapted according to the protocol used to measure TA. BAMHBI does not solve for ΣSO_4 , ΣF and ΣNO_2 . $\Sigma[-]$ is initialized with TA data using the conversion equation (140). Similarly, BAMHBI outputs TA as an ordinary
 485 variable that can be compared with observation.



Table 4. Main chemical reactions taken into account in the model that affect pH and the equilibrium concentrations (Soetaert et al., 2007). In the speciation relationships, H denotes $[H^+]$.













Table 5. Lump sums of concentrations.

ΣX	how it is calculated	model equivalent
ΣCO_2	$= [CO_2] + [HCO_3^-] + [CO_3^{2-}]$	Model state variable DIC
$\Sigma B(OH)_4$	$= [B(OH)_3] + [B(OH)_4^-]$	as a function of salinity: $\Sigma B(OH)_4 = 410.6S/35$
ΣPO_4	$= [H_3PO_4] + [H_2PO_4^-] + [HPO_4^{2-}] + [PO_4^{3-}]$	Model state variable <i>POs</i>
ΣSiO_4	$= [H_4SiO_4] + [H_3SiO_4^-] + [H_2SiO_4^{2-}]$	Model state variable <i>SiOs</i>
ΣNH_3	$= [NH_3] + [NH_4^+]$	Model state variable <i>NHs</i>
ΣNO_3	$= [NO_3^-] + [HNO_3]$	Model state variable <i>NOs</i>



The complete system of equations can be solved using a non-linear root-finding procedure, such as the Newton-Raphson technique for the H^+ concentration and pH. Once pH is known, the equilibrium proportions between the different acids and bases can be calculated. For additional details, an extensive description of the carbonate model is provided by Soetaert et al. (2007) and Hofmann et al. (2008).

490 2.2.5 The Benthic module

If the `Benthic` module is included, BAMHBI solves the benthic compartment dynamics. Otherwise, the benthic compartment is considered as impermeable (i.e. when it reaches the bottom, the sinking material accumulates and is progressively degraded as in the water column). The benthic compartment is not vertically resolved but is described by a vertically integrated dynamic sediment model using the Level-3 approach proposed by Soetaert et al. (2001). Only the solid part is explicitly simulated by
495 5 state variables that include the vertically integrated content of a fast and slow reacting pools of organic C (resp. S_C^f and S_C^s) and biogenic silica (resp. S_S^f and S_S^s). The fifth state variable is the N : C ratio of the sedimentary organic matter ($S_{N:C}$) that is used to compute the sedimentary organic nitrogen (Table 1). The labile and refractory parts have the same stoichiometry. The sedimentary content of organic phosphorus is derived from the nitrogen content using the same N : P ratio as for the pelagic part.

500 The sedimentary content of particulate organic carbon and biogenic silica (i.e., S_C^f , S_C^s , S_S^f and S_S^s) increases due to sinking and deposition and decreases due to degradation, erosion/resuspension and burial (Eq. (24)). P_i^j denotes the net exchange between the sediment and the overlying water column of the degradable fraction j ($j = f, s$) of the particulate materials i ($i = C, Si$). P_i^j is the difference between the potential deposition and erosion rates and is computed according to the equations presented in Table 6. Based on the value of the bottom shear stress (i.e. τ) compared to thresholds values for deposition (i.e.
505 τ_{dep}) and erosion (i.e. τ_{ero}), three situations are possible. For bottom shear stress below the threshold for deposition (i.e. τ_{dep}) only deposition occurs. The deposited materials is partitioned into the fast and slow degrading stock. For bottom shear stress in between the deposition and erosion threshold, there is neither deposition nor erosion. At high shear stress, larger than the erosion threshold, there is erosion of the stock. Because the slowly decaying stock is located deeper in the sediment and is more aggregated (Middelburg and Meysman, 2007), it is less easily resuspended than the fast degrading stock. Hence the slow
510 decaying stock is resuspended when all the fast decaying stock is resuspended and the erosion threshold of the fast decaying stock (i.e. τ_{ero}^f) is smaller than that of the slowly decaying fraction (i.e. τ_{ero}^s) (Table 6).

The bottom shear stress combines the stress due to the current and waves (Soulsby, 1997) and is computed at each time step according to Eqs. (147) to (153). The remineralization and dissolution rates are first order of the stock (Eqs. (142)–(142)) and a part of the slowly decaying stock is buried (Eq. (143)). The benthic carbon mineralization rate (D_C ; Eq. (142)) is derived
515 from the sum of the degradation of the two biodegradable carbon fractions. The model estimated N : C ratio is used to derive the benthic nitrogen mineralization (D_N ; Eq. (142)).

Porewater solutes are not explicitly simulated. The effect of benthic processes on pelagic dissolved inorganic constituents (i.e., NO₃, NH₄, O₂, PO₄, SiO₄, DIC) is parameterized based on mass budget considerations and assuming that there is no storage in the sediment. The main reactions that affect the O₂ cycle in the sediment are 1) oxic mineralization (Eq. (26)),



Table 6. Computation of the net flux P_i^j of particulate materials i ($i=C, Si$) of degradable fraction j ($j=f, s$) with $[X]_i = [POC], [CDI], [PON, NDI], [SID], [SDI]$ for respectively $i = C, N, Si$.

Deposition	No Depos./No eros.		Resuspension	
	$\tau_{dep} < \tau < \tau_{ero}^f$	$P_i^j = 0$	$\tau > \tau_{ero}^f$	$\tau > \tau_{ero}^s$
$\alpha_{dep} = (1 - \frac{\tau}{\tau_{dep}})$			$\alpha_{ero}^f = (\frac{\tau}{\tau_{ero}^f} - 1)$	$\alpha_{ero}^s = (\frac{\tau}{\tau_{ero}^s} - 1)$
$P_i = \alpha_{dep} \cdot \sum_{X \in X_i} w_X [X]$			$Me_i^{f'} = \frac{S_i^f}{\Delta t} (\frac{1}{\frac{\tau}{\tau_{ero}^f} - 1})$	
$P_i^f = P_i \cdot f_i$ (for $i=C, Si$ only)			$Me_i^{f*} = \min(Me_i^f, Me_i^{f'})$	$Me_i^{s*} = Me_i^s (1 - \frac{Me_i^{f*}}{Me_i^f})$
$P_i^s = P_i \cdot (1 - f_i)$ (for $i=C, Si$ only)			$P_i^f = \alpha_{ero}^f Me_i^{f*}$	$P_i^s = Me_i^{s*} (\frac{\tau}{\tau_{ero}^s} - 1)$
			$P_N = (P_C^f + P_C^s) S_{N:C}$	

520 2) nitrification (Eq. (27)) and 3) possible reoxydation of reduced substances produced during anoxic mineralization (Eq. (35)).
 The main reactions affecting the sedimentary nitrogen cycle are 1) production of NHs in the degradation process and possible
 re-oxidation into NOs, 2) consumption of NOs during the denitrification and production of N_2 .

The exchanges of solutes at the sediment-water interface are formulated as a function of variables p_{Nit} , p_{Denit} , p_{Anox} and
 $p_{SolidDepo}$ defined in Table 2. The efflux of NHs equals the total organic nitrogen mineralisation (D_N) minus the amount of
 525 NHs oxidised by nitrification (i.e., $p_{Nit} D_N$; Eq. (166) in Table 8). The NOs efflux equals the production of NOs by nitrification
 minus its consumption by denitrification (i.e., $0.8 D_C p_{Denit}$ assuming a consumption of 0.8 mole of NOs per mole of organic
 carbon remineralized; see Eq. (167)). The NOs efflux can be negative (i.e., NOs is consumed by the sediment) when the
 amount of NOs consumed by denitrification exceeds that produces by nitrification. The NHs efflux can also be negative when
 $p_{Nit} > 1$ which occurs when the sediment mineralization is low and the NHs concentrations are high (Eq. (166)). The O_2 flux
 530 to the sediment results from its consumption for the remineralisation and nitrification (respectively the first and second terms of
 Eq. (168) in Table 8). The consumed O_2 can be estimated from the amount of remineralized carbon (i.e., D_C) reduced by the
 amount of carbon degraded via denitrification and anoxic processes. O_2 is also consumed for the re-oxidation of the reduced
 substances that are produced by anoxic processes and are not deposited (i.e., $p_{Anox} (1 - p_{SolidDepo})$). We assume that the
 efflux of SiOs equals the dissolution rate of particulate silicate, that of DIC is the carbon mineralization (D_C) and that of POs
 535 is computed from the nitrogen mineralization rate (D_N) assuming a constant Redfield P:N ratio (Eqs. (163), (164) and (165),
 resp.). The adsorption of NHs and POs is neglected.

The parameters, p_{Nit} , p_{Oxic} and p_{Anox} are estimated by means of meta-modelling, while $p_{SolidDepo}$ is assumed to be
 constant (Table 7). These parameters are estimated using a regression model so that the vertically integrated model reproduces
 the results of a vertically resolved diagenetic model, OMEXDIA, implemented in the Black Sea by Wijsman et al. (2002).
 540 Monte Carlo simulations (~ 1000 runs) with the full diagenetic model are performed over a range of environmental conditions
 (e.g. bottom water inorganic nutrient concentrations, mineralization rate, carbon flux to the bottom, organic matter reactivity,
 bioturbation and bio-irrigation rates, water depth) that covers the expected variability of the system. The full model-estimated

nitrification and denitrification rates (resp. $p_{Nit} D_N$ and $p_{Denit} D_N$) and p_{Anox} are regressed as a function of environmental variables (Eqs. (153)–(155)). The values of the p_{Denit} presents a peak at intermediate mineralization rate while p_{Anox} increases
 545 with the mineralization rate. The benthic model and its validation are described in details in Capet et al. (2016).

Remineralization-Dissolution

$$D_i = \sum_{j=f,s} k_j f_i^j(T) S_i^j \quad \text{with} \quad f_i^j(T) = Q^{\frac{T-20}{10}} \quad (141)$$

with $i = C, Si$ and $j = f, s$

$$D_N = D_C \cdot S_{N:C} \quad (142)$$

550

Burial

$$B_i^s = b_i \cdot S_i^s \quad \text{with} \quad i = C, Si \quad (143)$$

$$B_i^f = 0 \quad (144)$$

The mean first-order degradation rate

$$k = \frac{\sum_{j=f,s} k_j f_i^j(T) S_i^j}{\sum_{j=f,s} S_i^j} \quad \text{with} \quad i = C, Si \quad \text{and} \quad j = f, s \quad (145)$$

555 Shear stress due to currents and waves

$$\tau_{\text{mean}} = \tau_{\text{cur}} \left[1 + 1.2 \left(\frac{\tau_{\text{wave}}}{\tau_{\text{wave}} + \tau_{\text{cur}}} \right)^{3.2} \right] \quad (146)$$

$$\tau = \sqrt{(\tau_{\text{mean}} + \tau_{\text{wave}} \cos(\Phi))^2 + (\tau_{\text{wave}} \sin(\Phi))^2} \quad (147)$$

where Φ is the angle between waves and currents

$$\tau_{\text{currents}} = \rho c_{db} U^2 \quad \text{with} \quad c_{db} = \left[\frac{1}{\kappa} \left(1 + \log \frac{z_0}{\Delta z_{\text{bottom}}} \right) \right]^{-2} \quad (148)$$

560

or set to constant value as in NEMO (ln_non_lin namelist option)

$$\tau_{\text{waves}} = 0.5 \rho f_w U_w^2 \quad (149)$$

with U_w is the maximum horizontal orbital wave velocity

$$f_w = \max(f_{wr}, f_{ws}) \quad (150)$$

The bottom friction is the maximum of rough and smooth bed friction

565

$$f_{wr} = 0.237 \left(\frac{A}{k_s} \right)^{-0.52} \quad \text{with} \quad A = \frac{U_w T}{2\pi} \quad \text{and} \quad k_s = 2.5 d_{50} \quad (151)$$

$$f_{ws} = BR_w^{-N} \quad \text{where} \quad R_w \text{ is the Reynolds number} \quad (152)$$

if $R_w < 5 \cdot 10^{-5}$ the flow is laminar and B=2, N=0.5

if $R_w > 5 \cdot 10^{-5}$ the flow is turbulent and B=0.0521, N=0.187

Bulk Parameterization

570

$$\ln(p_{Nit} \cdot D_N) = -6.280 - 0.286 (\ln D_C)^2 + 0.127 (\ln D_C)^3 + 0.006 (\ln D_C)^4 \quad (153)$$



575

$$\begin{aligned}
 &+ 0.098 (\ln D_C) \cdot \ln([O_2]) + 0.202 (\ln D_C)^2 \cdot \ln([O_2]) \\
 &- 0.021 (\ln D_C)^3 \cdot \ln([O_2]) - 0.031 (\ln D_C)^2 \cdot (\ln([O_2]))^2 \\
 &- 0.066 \ln(D_C) \cdot \ln([NHs]) + 0.009 (\ln D_C)^2 \cdot \ln([NOs]) \\
 &+ 0.823 \ln([O_2]) - 0.028 (\ln([O_2]))^2 - 0.012 \ln([O_2]) \cdot \ln([NHs]) \\
 &+ 0.119 \ln([NHs]) + 0.015 (\ln([NHs]))^2
 \end{aligned}$$

$$\ln(p_{Denit} \cdot D_C) = -5.475 - 0.786 (\ln D_C)^2 + 0.662 (\ln D_C)^3 + 0.042 (\ln D_C)^4 \quad (154)$$

580

$$\begin{aligned}
 &+ 0.064 (\ln D_C) \cdot (\ln([O_2])) + 0.794 (\ln D_C)^2 \cdot (\ln([O_2])) \\
 &- 0.082 (\ln D_C)^3 \cdot (\ln([O_2])) - 0.122 (\ln D_C)^2 \cdot (\ln([O_2]))^2 \\
 &+ 0.077 (\ln([O_2]))^2 - 0.155 (\ln([O_2])) \cdot (\ln([NOs])) \\
 &+ 0.875 \ln([NOs]) + 0.046 (\ln([NOs]))^2
 \end{aligned}$$

$$p_{anox} = 1.056 - 0.132 (\ln D_C) + 0.005 (\ln D_C)^4 \quad (155)$$

$$\begin{aligned}
 &+ 0.057 (\ln D_C) \cdot (\ln([O_2])) - 0.017 (\ln D_C)^2 \cdot (\ln([O_2])) \\
 &- 0.008 (\ln([O_2]))^2
 \end{aligned}$$



Table 7. Parameters values for biological processes.

PAR.	UNITS	VALUE			DESCRIPTION	REF.
Q_{10}	–	2			temperature coefficient	(1)
$Q_{10_{diatoms}}$	–	1.8			temperature coefficient for diatoms	(c)
p_{IR}	–	0.54			Part of the sea surface solar radiation $Q_{s,0}$ in the infra-red wave length (i.e. > 700 nm	(2)
p_{Long}	–	0.63			Part of PAR with a long wave length ([540 – 700] nm)	(c)
m_1	–	4.18				(3)
m_2	–	0.52				(3)
m_3	–	10.8				(3)
<u>Specific IOPs</u>		<u>Short</u>	<u>Long</u>			
$a_{\lambda,sw}$	m^{-1}	0.0196	0.24		(4)	
$a_{\lambda,Chl}^*$	-	0.029	0.007		(5)	
$a_{\lambda,Chl}$	-	0.6	0.1		(5)	
$a_{\lambda,POC}^*$	$m^2 mmolC^{-1}$	0.6	0.234		(6)	
$a_{\lambda,CDOM}^*$	m^{-1}	0.2522	0.2155		(c)	
$a_{\lambda,CDOM}^{**}$	$m^{-1} p.s.u.^{-1}$	0.0122	0.0113		(c)	
$b_{\lambda,sw}$	m^{-1}	0.0015	0.0007		(4)	
b_{λ,PHY_1}^*	$m^2 mmolC^{-1}$	$12.1 \cdot 10^{-5}$	$9.07 \cdot 10^{-5}$		(7)	
b_{λ,PHY_2}^*	$m^2 mmolC^{-1}$	$9.8 \cdot 10^{-5}$	$7.34 \cdot 10^{-5}$		(7)	
b_{λ,PHY_3}^*	$m^2 mmolC^{-1}$	$6.96 \cdot 10^{-5}$	$5.23 \cdot 10^{-5}$		(7)	
$b_{\lambda,POC}^*$	$m^2 mmolC^{-1}$	0.0055	0.005		(6, 8)	
<u>Phytoplankton</u>		<u>PHY₁</u>	<u>PHY₂</u>	<u>PHY₃</u>		
(N:C) _{PHY,min}	mol N (mol C) ⁻¹	0.05	0.05	0.05	minimum phytoplankton N:C ratio	(1)
(N:C) _{PHY,max}	mol N (mol C) ⁻¹	0.2	0.2	0.2	maximum phytoplankton N:C ratio	(1)
(N:Si)	mol N (mol Si) ⁻¹	–	–	5/6	phytoplankton N:Si ratio	(Redfield)
(N:P)	mol N (mol P) ⁻¹	16	16	16	phytoplankton N:P ratio	(Redfield)
(Chl:N) _{min}	g Chla (mol N) ⁻¹	1	1	1	minimum phytoplankton Chl:N ratio	(1)
(Chl:N) _{max}	g Chla (mol N) ⁻¹	2	2	2	maximum phytoplankton Chl:N ratio	(1)
α^{Chl}	($\mu\text{mol phot m}^{-2}$) ⁻¹ d ⁻¹	0.03	0.03	0.06	Chl a-specific initial slope of the photosynthesis-light curve	(c)

continued on next page ...



... continued from previous page

PAR.	UNITS	VALUE			DESCRIPTION	REF.
Resp	d^{-1}	0.01	0.01	0.01	Basal Respiration rate	(9)
ProdResp	–	0.1	0.1	0.1	Part of Primary Production used for respiration	(9)
μ_{max}	d^{-1}	3	1.5	3.5	Max. spec. growth rate	(c)
NOumax	$mol\ N\ (mol\ C)^{-1}\ d^{-1}$	1.5	0.5	1	Max. NOs uptake rate	(c)
NHumax	$mol\ N\ (mol\ C)^{-1}\ d^{-1}$	1.5	0.5	1	Max. NHs uptake rate	(c)
SiOsumax	$mol\ Si\ (mol\ C)^{-1}\ d^{-1}$	0	0	1	Max. SiOs uptake rate	(c)
Pumax	$mol\ P\ (mol\ C)^{-1}\ d^{-1}$	0	0	1	Max. POs uptake rate	(c)
k_{NOs}	$mmol\ N\ m^{-3}$	0.05	3	1	Half-Sat. Cst for NOs uptake	(c)
k_{NHs}	$mmol\ N\ m^{-3}$	0.05	3	1	Half-Sat. Cst for NHs uptake	(c)
k_{in}	$mmol\ N\ m^{-3}$	0.5	0.5	0.5	inhibition coefficient for NOs uptake by the presence of NHs	(1)
k_{SiOs}	$mmol\ Si\ m^{-3}$	–	–	2	Half-Sat. Cst for SiOs uptake	(c)
γ_1	–	0.02	0.02	0.02	Phyto. leakage fraction	(9)
γ_2	–	0.05	0.05	0.05	Ratio of extra DOC produced by N lim.	(9)
$mort_{PHY}$	d^{-1}	0.03	0.03	0.03	mortality rate	(1)
ϵ	–	0.34	0.34	0.34	DOM fraction of phytoplankton mortality	(10)
$v_{PHY3,max}$	$m\ d^{-1}$	0	0	1	max. diatoms sinking rate	(c)
$v_{PHY3,min}$	$m\ d^{-1}$	0	0	0.1	min. diatoms sinking rate	(c)
<u>Zooplankton</u>		<u>ZOO₁</u>	<u>ZOO₂</u>			
(C:N) _{ZOO}	$mol\ C\ (mol\ N)^{-1}$	5.5	4.67		internal ratio	(10)
MaxGrazZOO	d^{-1}	3.6	1		max. grazing rate	(c)
$k_{sat,ZOO}$	$mmol\ C\ m^{-3}$	5	5		half-sat. cst	(c)
β_N	–	0.77	0.77		Assimilation N effic.	(10)
β_C	–	0.64	0.64		Assimilation C effic.	(10)
k_c	–	0.8	0.8		net growth effic.	(1)
ϕ	–	0.23	0.23		messy feeding frac.	(10)
$mort_{ZOO}$	d^{-1}	0.3	0.3		max mortality rate	(9)
$ksat_{MortZOO}$	$mmol\ C\ m^{-3}$	1	1		half-sat. for mortality	(9)
AnoxMort	d^{-1}	0.25	0.25		mortality rate in anoxia	(c)
$ksat_{AnoxMort}$	$mmol\ O_2\ m^{-3}$	0.8	0.8		half-sat. for mortality in anoxia	(c)

continued on next page ...

585



... continued from previous page								PAR.	UNITS	VALUE	DESCRIPTION	REF.
Prey:	Phy1	Phy2	Phy3	Zoo1	Zoo2	Bac	POM					
$e_{\text{Prey,ZOO}_1}$	1	0	0	0	0	0.7	0			Capture efficiencies	(c)	
$e_{\text{Prey,ZOO}_2}$	0.4	0.4	1	1	0	0	0.8					
<i>Bacteria and dissolved organic matter</i>												
(C:N) _{BAC}								mol C (mol N) ⁻¹	5.1	bacteria internal ratio	(10)	
μ_{BAC}								d ⁻¹	13.3	Max. labile DOC or NHS uptake	(10)	
$k_{\text{BAC,DOC}}$								mmol C m ⁻³	25	half-sat. for labile DOC uptake	(10)	
$k_{\text{BAC,NH}_5}$								mmol N m ⁻³	0.5	half-sat. for NH ₄ uptake	(10)	
ω_{B}								–	0.17	bacterial gross growth efficiency	(10)	
m_{ortBAC}								d ⁻¹	0.04	mortality rate	(10)	
MaxHydDOM								d ⁻¹	4	max. semi-labile DOC hydrolysis	(10)	
$k_{\text{sat,hydDOM}}$								mmol C m ⁻³	417	Half-Sat. for semi-labile DOC uptake	(10)	
δ_1									0.7	lab. frac. phyto and non-phyto produced DOM	(9)	
δ_2									0.65	lab. frac. of phyto extra-excretion	(9)	
<i>Detritus</i>												
MaxHydPOC								d ⁻¹	0.04	Hydrolysis rate of POC	(10)	
MaxHydPON								d ⁻¹	0.055	Hydrolysis rate of PON	(10)	
$k_{\text{sat,OxicHydro}}$								mmol O ₂ m ⁻³	2.7	Half-Sat. for oxic hydrolysis	(c)	
diSSiDet								d ⁻¹	0.02	diss. rate of SiDet	(c)	
v_{SiDet}								m d ⁻¹	2	sinking rate of SiDet	(c)	
$\text{SinkingRatePOM}_{\text{constant}}$								m d ⁻¹	2	sinking rate of POM	(c)	
<i>Chemical Model</i>												
$\mu_{\text{NH}_5}^{\text{O}_2}$								d ⁻¹	0.03	max. NH ₅ oxidation rate by O ₂	(10)	
$\mu_{\text{NH}_5}^{\text{NO}_5}$								d ⁻¹	0.1	max. NH ₅ oxidation rate by NO ₅	(Assumed)	
$\mu_{\text{ODU}}^{\text{O}_2}$								d ⁻¹	0.1	max. ODU oxidation rate by O ₂	(Assumed)	
$\mu_{\text{ODU}}^{\text{NO}_5}$								d ⁻¹	0.1	max. ODU oxidation rate by NO ₅	(Assumed)	
$k_{\text{sat,O}_2}^{\text{NHsoxid}}$								mmol O ₂ m ⁻³	1	Half-sat. const. for O ₂ limit. in NH ₅ oxid. by O ₂	(11)	
$k_{\text{in,O}_2}^{\text{NHsoxidNO}_5}$								mmol O ₂ m ⁻³	30	Half Sat. const. for O ₂ inhib. in NH ₅ oxid. by NO ₅	(c)	
$k_{\text{sat,NO}_5}^{\text{NHsoxidNO}_5}$								mmol N m ⁻³	0.3	Half Sat. const. for NO ₅ limit. in NH ₅ oxid. by NO ₅	(c)	
$k_{\text{sat,O}_2}^{\text{ODUoxid}}$								mmol O ₂ m ⁻³	1	Half Sat. const. for O ₂ limit. in ODU oxid. by O ₂	(11)	



... continued from previous page

PAR.	UNITS	VALUE	DESCRIPTION	REF.
$k_{in,O_2}^{ODUoxidNOs}$	$\text{mmol O}_2 \text{ m}^{-3}$	30	Half Sat. const. for O_2 inhib. in ODU oxid. by NOs	(c)
$k_{sat,NOs}^{ODUoxidNOs}$	mmol N m^{-3}	0.3	Half Sat. const. for NOs limit. in ODU oxid. by NOs	(c)
$k_{sat,O_2}^{OxicMin}$	$\text{mmol O}_2 \text{ m}^{-3}$	3	Half Sat. const. for O_2 limit. in oxic respiration	(11)
k_{in,O_2}^{Denit}	$\text{mmol O}_2 \text{ m}^{-3}$	30	Half Sat. const. for O_2 inhib. in denitrification	(c)
$k_{sat,NOs}^{Denit}$	mmol N m^{-3}	0.3	Half Sat. const. for NOs limit. in denitrification	(c)
$k_{in,O_2}^{AnoxMin}$	$\text{mmol O}_2 \text{ m}^{-3}$	0.005	Half Sat. const. for O_2 inhib. in anoxic respiration	(c)
$k_{in,NOs}^{AnoxMin}$	mmol N m^{-3}	0.005	Half Sat. const. for NOs inhib. in anoxic respiration	(c)
R_{solid}		0.2	Perc. of ODU production that is deposited as a solid	(c)
$k_{sat,iron}^{SolForm}$	$\mu\text{mol Fe m}^{-3}$	100	Half Sat. const. for Iron limit. in solid ODU Formation	(c)
$\gamma_{O_2}^C$	$\text{mol O}_2 (\text{mol C})^{-1}$	1	Mol O_2 used per mol C in oxic respiration	(see equ. in Table 1)
$\gamma_{O_2}^{NHs}$	$\text{mol O}_2 (\text{mol NHs})^{-1}$	2	Mol O_2 needed to oxidize one mol of NHs in nitrification	(see equ. in Table 1)
$\gamma_{O_2}^{ODU}$	$\text{mol O}_2 (\text{mol ODU})^{-1}$	1	Mol O_2 needed to oxidize one mol of ODU	(see equ. in Table 1)
γ_{ODU}^{POC}	$\text{mol ODU} (\text{mol C})^{-1}$	1	Mol ODU formed per mol C in anoxic respiration	(see equ. in Table 1)
γ_{NOs}^{POC}	$\text{mol NOS} (\text{mol C})^{-1}$	0.8	Mol NOs used per mol C in denitrification	(see equ. in Table 1)
γ_{NOs}^{NHs}	$\text{mol NOS} (\text{mol NHs})^{-1}$	0.6	Mol NOs needed to oxidize one mol of NHs	(see equ. in Table 1)
γ_{NOs}^{ODU}	$\text{mol NOS} (\text{mol ODU})^{-1}$	0.8	Mol NOs needed to oxidize one mol of ODU	(see equ. in Table 1)
$burial_C$		0.28	Percentage of POC buried in the Sediments	(12)



... continued from previous page

PAR.	UNITS	VALUE				DESCRIPTION	REF.		
<i>burial_N</i>		0.28				Percentage of PON buried in the Sediments	(12)		
<u>Chlorophyll-Geider Module</u>		<u>PHY₁</u>	<u>PHY₂</u>	<u>PHY₃</u>					
α^{Chl}	mol C (g Chl _a) ⁻¹ ($\mu\text{mol phot m}^{-2}$) ⁻¹ d ⁻¹	0.062	0.026	0.052	Chl a-specific initial slope of the photosynthesis-light curve	(c)			
NHSNoUptake	mol N (mol C) ⁻¹ d ⁻¹	0	0	0	Minimal NHs uptake rate below which the uptake is set to zero	(c)			
PO4NoUptake	mol P (mol C) ⁻¹ d ⁻¹	8.64	8.64	8.64	Minimal POs uptake rate below which the uptake is set to zero	(c)			
<u>Gelatinous Module</u>		<u>GEL₁</u>	<u>GEL₂</u>						
(C:N) _{GEL}	molC molN ⁻¹	4.76	4		internal ratio	(13,14)			
<i>Q₁₀gelatinous</i>		3.5	3.5		<i>Q₁₀</i> factor	(15)			
MaxGrazGEL	d ⁻¹	0.06	0.3		max. grazing rate	(c)			
AssEffGEL	-	0.75	0.75		Assimilation efficiency	(13)			
GrowthEffGEL	-	0.2	0.2		Growth efficiency	(13)			
mortGEL	d ⁻¹	0.06	0.006		max mortality rate	(c)			
AnoxMort	d ⁻¹	0.25	0.25		mortality rate in anoxia	(c)			
<i>ksat_{AnoxMort}</i>	mmolO ₂ m ⁻³	8	8		half-sat for mortality in anoxia	(c)			
Prey:	Phy1	Phy2	Phy3	Zoo1	Zoo2	Bac	POM		
<i>e_{Prey,GEL1}</i>	1	0.5	1.0	1	0	0	1	Capture efficiencies	(c)
<i>e_{Prey,GEL2}</i>	0	0	0	0	1	0	0		
<u>Aggregation module</u>									
η	-					0.62		sinking exponent	(16)
B	m ^{-η} d ⁻¹					2294		sinking factor	(16)
stick	-					0.5		stickiness	(c)
ℓ	m					2 · 10 ⁻⁵		minimal cell size	(16)
<i>L</i>	m					0.01		maximal cell size	(16)
ζ	-					1.62		N content exponent	(16)
C	mmolN m ^{-ζ}					0.4744		N content coefficient	(16)
ϵ_{ini}						3.5		epsilon initial value	(16)
<u>Benthic module</u>									
<i>f_C</i>	-					0.29		Part of the particulate C deposition flux that goes to S_C^f	(17)



... continued from previous page				
PAR.	UNITS	VALUE	DESCRIPTION	REF.
f_{Si}	-	0.5	Part of the particulate Si deposition flux that goes to S_{Si}^f	(18)
k_C^f	d^{-1}	0.0753	Degradation rate of S_C^f at 20°C	(12)
k_C^s	d^{-1}	0.003	Degradation rate of S_C^s at 20°C	(17)
k_{Si}^f	d^{-1}	72×10^{-4}	Dissolution rate of S_{Si}^f at 20°C	(18)
k_{Si}^s	d^{-1}	72×10^{-5}	Dissolution rate of S_{Si}^s at 20°C	(18)
Q	--	2	Factor for Temperature effect on mineralization	(17, 19)
b_C, b_S	d^{-1}	2.16×10^{-4}	Burial rate for the slow degrading stocks	(c)
$p_{SolidDepo}$	-	0.11	Part of precipitating ODU	(17)
OC_{deg}	$molO (molN)^{-1}$	1	Oxygen consumed for oxic degradation	(12)
ON_{nit}	$molO (molN)^{-1}$	2	Oxygen consumed for nitrification	(12)
Me_C^s	$mmolC m^{-2} d^{-1}$	0.864	Erosion constant for S_C^s	(20)
Me_C^f	$mmolC m^{-2} d^{-1}$	0.864	Erosion constant for S_C^f	(20)
Me_{Si}^s	$mmolSi m^{-2} d^{-1}$	0.864	Erosion constant for S_{Si}^s	(21)
Me_{Si}^f	$mmolSi m^{-2} d^{-1}$	0.864	Erosion constant for S_{Si}^f	(21)
τ_{dep}	$N m^{-2}$	0.02	Bottom stress threshold for deposition	(21)
τ_{ero}^f	$N m^{-2}$	0.02	Bottom stress threshold for erosion of S_C^f, S_{Si}^f	(21)
τ_{ero}^s	$N m^{-2}$	0.04	Bottom stress threshold for erosion of S_C^s, S_{Si}^s	(21)
κ	--	0.4	Von Karman constant	
Δz_{bottom}	m		size of the bottom grid cell	-
d_{50}	m	250×10^{-6}	Median grain size	-
z_0	m	2.083×10^{-5}	Rugosity length scale for a grain size of 250×10^{-6}	(22)
ν	$m^2 s^{-1}$	12×10^{-6}	Kinematic viscosity	
$w_{POC} = w_{PON}$	$m d^{-1}$	$-[4 - 1.5]$	Particulate organic matter sinking velocity	(16)
$w_{CDI} = w_{NDI}$	$m d^{-1}$	-0.9	Diatoms sinking velocity	(23)
w_{SiDet}	$m d^{-1}$	-2	Biogenic silica sinking velocity	(23)



590 (c) after calibration. References: (1) Soetaert et al. (2001), (2) Mobley et al. (2004), (3) Lee et al. (2005), (4) Smith and Baker (1981), (5) Bricaud et al. (1995), (6) Neukermans et al. (2012), (7) Vaillancourt et al. (2004), (8) Doerffer and Schiller (2007), (9) Van den Meersche et al. (2004), (10) Anderson and Pondaven (2003), (11) Soetaert et al. (1996), (12) Muramoto et al. (1991), (13) Lancelot et al. (2002), (14) Nakamura (1998), (15) Kremer (1977), (16) Kriest (2002), (17) Wijsman et al. (2002), (18) Khalil et al. (2007), (19) Van Cappellen et al. (2002), (20) Stanev and Kandilarov (2012), (21) Capet et al. (2016), (22) Gayer et al. (2006), (23) Grégoire et al. (2008).

595 3 Horizontal interfaces

3.1 Air-sea interface

Gases, CO₂ and O₂, are exchanged at the air-sea interface. Net CO₂ flux at the air-sea interface (*AirSeaCO₂Flux*) depends on the difference in partial pressure of CO₂ (*pCO₂*) between air (*pCO_{2,air}*) and the surface water (*pCO_{2,water}*), the solubility coefficient *K₀(S,T)* and the gas transfer velocity *k(wind)*. *k(wind)* is parameterized as a function of the wind speed with the
600 formulation of Wanninkhof (1992):

$$AirSeaCO_2Flux = K_0(S,T)k(wind)(pCO_{2,air} - pCO_{2,water}) \quad (156)$$

$$k(wind) = 0.31wind^2 \frac{1}{\sqrt{S_c/660}} \quad (157)$$

$$\text{with } S_c = 2073.1 - 125.62T + 3.6276T^2 - 0.043219T^3$$

where *S_c* is the Schmidt number.

605 The net O₂ flux at the air-sea interface (*AirSeaO₂Flux*) depends on the difference in the O₂ saturation concentration (*O_{2,sat}*) and the surface water O₂ concentration (*O_{2,water}*), the solubility coefficient *K₀(S,T)* and the gas transfer velocity *k(wind)* which is parameterized as a function of the wind speed with the formulation of Ho et al. (2006):

$$AirSeaO_2Flux = k(wind)(O_{2,sat} - O_{2,water}) \quad (158)$$

$$k(wind) = 0.266wind^2 \frac{1}{\sqrt{S_c/660}} \quad (159)$$

610 with *S_c* = 1953.4 - 128.00*T* + 3.9918*T*² - 0.050091*T*³

where *S_c* is the Schmidt number.

Additionally the model includes the possibility to consider the deposition of dissolved inorganic nitrogen and phosphorus. None of the other biogeochemical state variables is exchanged across the air-sea interface.

3.2 Water-sediment interface

615 By default, the bottom is considered as frozen. The sedimenting particles accumulate on the bottom and are degraded in the bottom layer of the pelagic model. When the `benthic` module is activated, a sedimentary box is considered as explained in section 2.2.5. Table 8 lists the pelagic variables that have a non-zero flux imposed at the sediment-water interface. These variables are all the sedimenting and dissolved inorganic modelled components. When the flux of solutes can be directed towards the sediments (which is always the case for O₂ and may occur for NHs and NOs) it may happen that the computed



Table 8. Benthic-pelagic boundary conditions.

$$F_{\text{POC}} = P_C \quad (160)$$

$$F_{\text{PON}} = P_N \quad (161)$$

$$F_{\text{SiD}} = P_{\text{Si}} \quad (162)$$

$$F_{\text{SiOs}} = D_{\text{Si}} \quad (163)$$

$$F_{\text{DIC}} = D_C \quad (164)$$

$$F_{\text{POs}} = D_N r_{P:N} \quad (165)$$

$$F_{\text{NHs}} = D_N (1 - p_{\text{nit}}) \quad (166)$$

$$F_{\text{NOs}} = D_N p_{\text{nit}} - D_C p_{\text{denit}} 0.8 \quad (167)$$

$$F_{\text{O}_2}^* = -D_C (1 - p_{\text{denit}} - p_{\text{anox}} p_{\text{solid}}) OC_{\text{deg}} - D_N p_{\text{nit}} ON_{\text{nit}} \quad (168)$$

$$F_{\text{O}_2} = \max(F_{\text{O}_2}^*, -[\text{O}_2] \cdot \frac{\Delta z}{\Delta t}) \quad (169)$$

$$F_{\text{ODU}} = \max(0, F_{\text{O}_2} - F_{\text{O}_2}^*) \quad (170)$$

620 influx exceeds the availability of solutes in the last box of the pelagic model. This is because the meta-modelling formulation
 does not directly depend on the availability of the solute in the last box of the pelagic model. In that case, the modelled influx
 is adjusted to consume the quantity of solute in the bottom box (i.e. $[Solute] \frac{\Delta z}{\Delta t}$). Then, for O_2 the excess consumption is
 converted into a release of ODU (Eq. (169)) while for NHs and NOs, the parameters (respectively p_{Nit} , p_{Denit}) are adjusted
 for not exceeding the available amount and all the fluxes are recomputed to preserve mass conservation. It should be noted that
 625 these adjustments are rare.

4 Implementation

The BAMHBI model comes as an independent library. The source and sink terms of the pelagic variables (and benthic vari-
 ables if the `benthic` module is activated) are computed by BAMHBI in the `bamhbi.F90` module written in Fortran 90.
 BAMHBI parameters are declared in `bamhbi_params.F90`. Each optional module has its own namelist, variables declara-
 630 tion file (`bamhbi_optionalmodule_vars.F90`, e.g. `bamhbi_alkalinity_vars.F90` and FORTRAN code
 (`bamhbi_optionalmodule.F90`, e.g. `bamhbi_alkalinity.F90`). All of these files include the `BAMHBI.h90`
 header file containing pre-processor directives enabling or disabling specific parts of the code such as described in section
 4.3.

The time integration is done externally. BAMHBI needs to be forced by temperature, salinity, mixing coefficients and, in a



635 three-dimensional setting, currents. Most of the time these forcings are provided by an ocean physical model. BAMHBI has
already been coupled in 1D with GOTM (Grégoire et al., 2008), in 3D with the GHER model (Capet et al., 2013), in 1D and
3D with the "Nucleus for European Modelling of the Ocean" (NEMO) and in 1D in the R software environment. As much as
possible, BAMHBI is written to be easily coupled with hydrodynamical models without substantial adaptations. The notation
(e.g. names of loop indices and physical variables) respects the NEMO conventions. For each physical model, an interface
640 module has to "translate" the variable names between the respective physical model and BAMHBI. Here we describe the time
integration of BAMHBI in NEMO and in R. The interface module written for the coupling of BAMHBI with R can serve as
example for coupling to other models.

4.1 Coupling with NEMO

The definition of the variables in the BAMHBI code follows the NEMO nomenclature (e.g. *jpi* for the dummy loop index in the
645 longitudinal direction, *e3t* for the height of the grid cells, *tr* for the biogeochemical variables). The interface module between
NEMO and BAMHBI, called `bamhbi_driver_nemo.F90` is thus relatively short. The complete BAMHBI source code is
stored in the NEMO MY_SRC sub-directory and through `bamhbi_driver_nemo.F90`, it uses all the required variables
(e.g. grids, masks, physical variables (active tracers), biogeochemical pelagic and benthic variables (passive tracers), surface
wind stress, surface radiation, bottom wave stress).

650 The calls to the BAMHBI subroutines are done in the NEMO `trc_sms_my_trc` subroutine. At each time step, some prelimi-
nary operations are done (such as resetting all source and sinks to zero), then for each biogeochemical variable the sources and
sinks are computed, and from that the source minus sinks term (i.e. trends in NEMO's terminology).

4.2 Coupling with R

BAMHBI can be run in one dimension in the R environment. BAMHBI is integrated using the deSolve package. deSolve is an
655 R package that solves 1-D, 2-D or 3-D ordinary/partial/algebraic differential equations (Soetaert et al., 2010). The R/deSolve
combination allows the use of functions from external libraries (written in C, or Fortran like `BAMHBI.F90`) and thus provides
an efficient and user-friendly way of running BAMHBI in a 1-D context. Indeed, compared to a similar offline (i.e. the physical
variables are read from files rather than computed) or to the one-dimensional version of NEMO, the R/deSolve version uses
just 1 water column (as opposed to a 3x3 grid of columns in NEMO) and an adaptive time-step integration scheme.

660 Practically, an R script reads the model grid characteristics (both pelagic and benthic), model parameters, initial conditions
and external forcings. The forcing includes the physical fields that are needed to run the biogeochemical simulations (i.e. tem-
perature, salinity and vertical diffusion coefficients), the atmospheric conditions, the bottom shear stress and the vertical shear
of horizontal velocities (if the aggregation module is active). These forcing and initial fields are usually provided by another
model in netcdf files and are then assembled inside R.

665 The `bamhbi.R` script performs all the steps described above: it compiles the BAMHBI model (i.e. `BAMHBI.F90`) and depen-
dent modules into a library, loads the corresponding library into memory, the model grid, variables, parameters and forcings,
and calls the deSolve routines used to integrate the model in time. In particular, `ode.1D` is called using the `lsodar` method.



Table 9. List of available options: name and definition.

OPTIONS	DESCRIPTION	Code Declaration	NAMELISTS
Modelling modules			
chlorophyll module	Chlorophyll computed according to Geider	chlorophyll-Geider	namtrc-bamhbi-geider
gelatinous zooplankton module	Adding two groups of gelatinous (omnivorous, carnivorous)	gelatinous	namtrc-bamhbi-gelatinous
aggregation module	Sinking speed of detritus computed with an aggregation model	aggregation	namtrc-bamhbi-aggregation
carbonate module	Solving DIC, pH, alkalinity, pCO_2 and air-sea fluxes of CO_2	carbonate	namtrc-bamhbi-carbonates
benthic module	Solving the vertically integrated labile and refractory detrital materials C, Si and solutes fluxes	benthic	namtrc-bamhbi-benthic
Diagnostics modules			
Diagnostic Rates	Output the model rates	biodiagrates	
Diagnostic Trophic flows	Output the trophic flows	biodiagtrophic	
Conservation			
Diagnostic conservation	Compute the sum of sources and sinks terms in a closed version of the model	testcons	
Diagnostic budgets	Output the budgets of biogeochemical elements	testcons2	

The time-step is variable depending on the stiffness of the differential equations. deSolve extracts the forcing fields at the right instant from the time-series, and passes it on to the BAMHBI model. The latter returns the source and sinks to deSolve, which
 670 uses them to integrate the state variables in time.

`bamhbi.R` also provides supplementary routines, for example if no vertical diffusion coefficient is available in the forcing files, they are estimated from the temperature and salinity forcing.

R calls routines from `BAMHBI.F90` using two modules: `bamhbi_interface` and `bamhbi_driver_R`. `bamhbi_interface` passes from deSolve to BAMHBI the physical variables and other forcing fields, and the updated biogeochemical model state, and
 675 conversely returns to deSolve the `sms` (sources minus sinks) computed by `BAMHBI.F90`; it thus fulfills the same role as `trc_sms_my_trc` when BAMHBI is coupled to NEMO. `bamhbi_driver_R` translates the name of the physical variables provided by R/deSolve in the BAMHBI nomenclature, fulfilling the same role as `bamhbi_driver_nemo`. It also offers supplementary routines, such the possibility to impose a lateral transport over a predefined depth to mimic the effect of lateral advection to compensate for vertical loss due to sinking.

680



4.3 Compilation Options

The `BAMHBI.h90` header file contains pre-processor directives enabling or disabling specific parts of the code. Available options include the modelling modules described in details in section 2.2 and the estimation of model diagnostics described here below. All the compiling options are listed in Table 9.

685 4.3.1 Mass conservation

690 Compiling BAMHBI with option "testcons" allows to check that the model is conservative in terms of C, N, Si, P and O₂. This option cancels the calls to routines that are the net sources and sinks for the model (i.e. denitrification, ANAMMOX, nitrate reduction by ODU are losses of nitrogen since N₂ is not explicitly modelled). It then checks that at each time step the sum of the biogeochemical sources and sinks (i.e. $\frac{\partial y}{\partial t}|_{Bio}$) equals zero. With the option "testcons2" the inventory over the modelled domain of each chemical currency (i.e. C, N, Si, P and O₂) is computed closing the system (i.e. removing its interactions with the outside world). It remains the user responsibility to cancel fluxes at open boundaries, rivers, and atmosphere. For example, when BAMHBI is coupled with NEMO, these fluxes are specified in NEMO's `namelist_top_cfg` file respectively in the `BDY`, `SBC` and `CBC` parts.

4.3.2 Clipping

695 When integrating the biogeochemical variables, it may happen that negative concentrations are generated, when for instance, the sink term of a given biogeochemical variable exceeds its stock. These negative concentrations are not realistic and may create instabilities. Then, before computing the sources and sinks terms, negative state variables are set to a small value specified in the BAMHBI parameters (e.g. 10^{-12}) and the added material is computed over the whole domain to check its order of magnitude compared to the total mass. When BAMHBI is coupled to NEMO, the BAMHBI clipping procedure can be replaced with NEMO's `trc_rad` routine. This clips the biogeochemical variables after integration of the total trend, obtained as the sum of the source and sinks (computed by BAMHBI) and the transport (advection and diffusion, computed by BAMHBI).

4.3.3 Diagnostics

A list of diagnostics are proposed to estimate model rates and trophic interactions. These sets of diagnostics can be computed by activating respectively the options "biodiagrates" and "biodiagtrophic".

705

5 BAMHBI Applications

This section describes two applications of BAMHBI to simulate the Black Sea biogeochemistry. One of them describes the use of the model to estimate benthic-pelagic fluxes and bottom oxygen over the northwestern shelf. In the second application, the *Chla* and oxygen dynamics are simulated and compared with ARGO in the deep sea.



710 In these two applications BAMHBI is online coupled with NEMO. At each NEMO 3D time step biogeochemical variables are
integrated and transported (by advection and diffusion) in the NEMO framework. At the air-sea interface, the coupled model is
forced by ERA-5 data (Hersbach et al., 2020) that are used to estimate the heat and energy exchanges using bulk formulations.
Precipitations and nutrients fluxes (i.e. NOs, NHs, POs) are also imposed and the exchanges of O₂ and CO₂ are computed
as described in section 3. For the other biogeochemical variables, a zero flux condition is imposed. At the water-sediment
715 interface, the shear stress due to current and waves is imposed. The waves are imposed as an external forcing from Stanev
and Kandilarov (2012). For biogeochemical variables boundary conditions are described in section 3. The physics is initialized
from rest with typical vertical profiles of temperature and salinity. A spin-up of 15 years is performed forced by climatological
atmospheric forcings. The initial conditions of inorganic components like NOs, NHs, POs, SiOs, O₂, ODU, DIC, TA, and pH
are expressed on a density scale using typical profiles. For the other biogeochemical variables, homogeneous profiles with very
720 low values are used. In the sediment, the slowly and fast reacting components are initialized with a 5-year averaged deposition
rate divided by the first-order averaged decay rates. A new spin-up of 15 years is then performed with the coupled model to
adjust the initial biogeochemical conditions to the dynamics.

5.1 Case Study: shelf hypoxia and benthic–pelagic coupling

Since the early 80s, in response to eutrophication, the Black Sea northwestern shelf has been affected by bottom hypoxia (e.g.,
725 Zaitsev, 1993; Mee et al., 2005; Capet et al., 2013). Bottom hypoxia has affected benthic communities and the coupling of
the benthic and pelagic compartments. In this case study, the capability of BAMHBI to generate bottom shelf hypoxia and
benthic–pelagic fluxes is demonstrated. Figures 6 and 7 compare the simulated and observed bottom oxygen concentrations
over the northwestern shelf. Interannual (over 1980–2002) and seasonal (average cycle over 1980–2002) variations are shown.
The model is able to simulate the typical seasonal cycle of bottom oxygen concentrations with higher values in winter when
730 the whole water column is mixed and saturated in oxygen and minimum values at the end of summer–early fall when the water
column is stratified and prevents the mixing of bottom water. During this period bottom hypoxia occurs in regions of intense
degradation of organic matter such as the northern shelf. From July to November, 1–2 % of the shelf bottom is hypoxic in the
model and the observations. The median of the model bias is usually close to zero and always less than 20 μmol L⁻¹. On an
interannual scale, model and observation consistently show that the percentage of shelf area covered by hypoxia is the largest
735 at the end of the 80s and early 90s with a peak in 1991 and a disappearance after 1995. However, as shown in Capet et al.
(2013), the recovery from hypoxia after 1995 results from a sampling bias with data collected where and when there is no
hypoxia. The median of the model bias is still very small with usually a positive bias lower than 10 μmol L⁻¹ (except in 1998)
though in 1991 and 1997 the model underestimates the oxygen concentrations with a bias lower than 25 μmol L⁻¹.

Figure 8 shows the average seasonal cycle of the benthic fluxes of *NHs*, *NOs*, *O₂*, *POs*, *SiOs* over the northwestern
740 shelf for three regions defined on the base of their bathymetry. The shallowest region, Region 1, is located along the western
and northern coast of the shelf with a bathymetry shallower than 40 m. This is the region that receives the largest inputs of
organic matter. Region 2, 40–80 m depth, corresponds to a band of water offshore the coastal Region 1 and just before the
shelf break. Region 2 receives moderate inputs of organic matter with a peak in summer when the shelf circulation becomes



anticyclonic and the Danube's discharges are transported over the whole shelf. Region 3, 80–200 m depth, involved bottom
745 waters located over the shelf break including anoxic and even sulfidic bottom waters. It is in Region 1 that the benthic fluxes are
the highest and present the largest variability. The seasonal cycle is well-marked with maximum intensity in summer and early
fall when bottom temperatures are the highest and the organic matter produced during the winter-spring bloom accumulates on
the bottom. In the three regions, the model simulates an outflux of NH_s , PO_s and SiO_s except in Region 3 where the 2.5th
percentile gives negative fluxes (sediment consumption) of NH_s where the concentrations of NH_s are high while the quantity
750 of sedimentary organic material is low. The O_2 fluxes are always negative, O_2 is consumed by the sediments, with the largest
consumption in summer and in region where and when seasonal hypoxia occurs. In Region 3, the 75th percentile is close to
zero because there are areas (depth > 100 m) where there is no oxygen on the bottom. The model simulates an outflux of NO_s
except in Region 1 in early summer where NO_s is consumed by the sediment via denitrification. The observations are usually
contained within the spread of model results except that observations show highest denitrification rates and a consumption
755 of PO_s by the sediment in spring at two stations located in regions 1 and 3. In the model, the phosphorus diagenesis is
oversimplified and ignores the formation of chemical complexes involving PO_s . In agreement with observations, the benthic
flux of PO_s is always towards the water column and is estimated from the sedimentary nitrogen degradation rates assuming
Redfield stoichiometry.

5.2 Case Study: The deep sea biogeochemistry

760 The central part of the Black Sea has a depth of $\approx 2200m$ and is permanently stratified. The intrusion of the saline Mediter-
ranean waters through the Bosphorus Strait creates a permanent halocline that prevents water ventilation below 100-150 m. Waters
below that depth are then anaerobic and even sulfidic. Figures 9 and 10 compare over 2010-2023 the simulated and observed
(from biogeochemical Argos) O_2 and Chl a . Figure 9 shows that the model overestimates over the different years the oxygen
concentration with a median bias between 25-50 $\mu mol l^{-1}$. This difference is explained by an overestimation of the halocline
765 depth and then of all the gradients in biogeochemical quantities. The oxycline is too deep and oxygen penetrates too much at
depths. The deep chlorophyll maximum (DCM), which is strongly correlated with the nitracline depth, is also too deep for the
same reason (Ricour et al., 2021).

6 Conclusions

770 The Biogeochemical Model for Hypoxic and Benthic Influenced areas (BAMHBI) is an intermediate complexity biogeochem-
ical model that describes the cycling of N, C, O_2 , P and SiOs through the marine foodweb from bacteria up to mesozooplank-
ton. BAMHBI is a stand-alone biogeochemical model that can be coupled to any hydrodynamical model. BAMHBI has been
conceived to simulate anaerobic and euxinic environments. It describes the degradation of organic matter using a succession
of oxidants. Bacteria preferentially use oxygen and, in anaerobic conditions, first nitrate and then other oxidants (not explicitly
described). BAMHBI involves five optional modules that explicitly model 1) the Chl a content of each PFT, 2) two zooplankton
775 gelatinous groups, 3) the number of aggregates, 4) the carbonates, 5) the benthic compartment.




Figure 6. Seasonal cycle, average over 1980-2002, of the percentage of shelf area affected by bottom hypoxia (i.e. $O_2 < 63 \mu\text{mol L}^{-1}$) (top panel), bottom O_2 concentration (median, P_{25} and P_{75} values) (middle panel) and median of the bias between the model (cyan) and observations (red) estimates. Observations have been extracted from the World Ocean Database (<https://doi.org/10.25923/z885-h264>). Their number is indicated in the middle figure. Model results are interpolated at the data points.

BAMHBI is an open source model available on a Gitlab server. BAMHBI updates are regularly made and the group is developing a CD/CI pipeline for allowing automatic testing of the updates. BAMHBI is currently extended with several optional modules: 1) a spectral radiative transfer model based on RADTRANS (Dutkiewicz et al., 2015; Mace et al., 2024), 2) a model of suspended particulate matter, 3) a refinement of benthic diagenesis, 4) a module of fish modelling. BAMHBI is currently
780 implemented in the FABM framework. All of these new developments will be described in separate papers. A containerized version of BAHMBI is also under development for allowing its portability on different machines.




Figure 7. Interannual cycle of the percentage of shelf area affected by bottom hypoxia (i.e. $O_2 < 63 \mu\text{mol L}^{-1}$) (top panel), bottom O_2 concentration (median, P_{25} and P_{75} values) (middle panel) and median of the bias between the model (cyan) and observations (red) estimates. Observations have been extracted from the World Ocean Database (<https://doi.org/10.25923/z885-h264>). Their number is indicated in the middle figure. Model results are interpolated at the data points.

Code and data availability.

785 The BAMHBI model is distributed as open source software under the MIT licence through a public GitLab repository hosted by the University of Liège (Vandenbulcke and Grailet, 2025). A copy of version 1.0 can also be downloaded on Zenodo at <https://doi.org/10.5281/zenodo.16612928> (Marilaure and Vandenbulcke, 2025). The BAMHBI module, its dependencies, are available online at <https://gitlab.uliege.be/especes/mast/r-bamhbi>. Code to reproduce Figure 8 and the in-situ data behind it can be found in this github repository: https://github.com/mchoblet/benthic_model_assessment



Average annual cycle of benthic fluxes on shelf




Figure 8. Average (1990-1999) seasonal cycle of the benthic fluxes of (from the first to the last row) NH_4 , NO_3 , O_2 , PO_4 , SiO_4 observed (symbols) for regions of different bathymetry ranges and simulated by BAMHBI (continuous lines with the 25th and 75th (resp. 2.5th and 97.5th) percentiles superimposed in dark (resp. light) color). The distribution of the different regions as well as the data collection sites are shown in the top of the figure (first row). The sampling sites of Friedl et al. (1998) and Friedrich et al. (2002) are similar but are collected respectively in August 1995 (stars) and May 1997 (circles).



Figure 9. Average (2010–2023) monthly cycle of (top figure) the concentration of O_2 observed by Argos (red) and simulated by BAMHBI (cyan) median values with the 25th and 75th (resp. 2.5th and 97.5th) percentiles limiting the color bars (resp. dotted lines) and (bottom figure) the median value of the bias.

790

Author contributions.

MG wrote the first version of the BAMHBI code and the whole manuscript. LV implemented BAMHBI in NEMO and run the simulations. LV, MC, ID, LM, PV run the coupled NEMO-BAMHBI and fine-tuned its parameterization. MC, SC, LM and HY developed new BAMHBI modules. SC, MC and LM prepared model figures. JFG validated BAMHBI for the test cases and prepared figures. EI implemented BAMHBI in FABM. AC, CM, AM, GM, KS supported the development of BAMHBI. All the co-authors read and reviewed the manuscript.

795




Figure 10. Average (2010-2023) seasonal cycle of (top figure) the concentration of Chl a observed (red) and simulated by BAMHBI (cyan) median values with the 25th and 75th (resp. 2.5th and 97.5th) percentiles limiting the color bars (resp. dotted lines) and (bottom figure) the median value of the bias.

Competing interests.

The authors declare that they have no conflict of interest.

800 *Acknowledgements.* MG, LV, ID, EI, LM, PM, HY, AM, CM acknowledge the support from the Copernicus Marine Environmental and Monitoring service CMEMS, the Horizon Europe NECCTON project under grant agreement no. 101081273, EU H2020 BRIDGE-BS project under grant agreement no. 101000240. SC, MC and LA are supported by the Fonds pour la formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA/FNRS). EI and RN have been supported by the project FULLCONTINUUM funded by the Fonds de la Recherche



805 Scientifique–FNRS (T010221F), CM and EI were funded by the CE2COAST project funded by ANR (FR), BELSPO (BE), FCT (PT), IZM (LV), MI (IE), MIUR (IT), Rannis (IS), IRP MAST (Multiscale Adaptive Strategies) and RCN (NO) through the 2019 ‘JointTransnational Call on Next Generation Climate Science in Europe for Oceans’ initiated by JPI Climate and JPI Oceans. VM is supported by the ESA MITHO project funded under the Contract No. 4000142100/23/I-DT. GM is a Research Associate with the Fonds de la Recherche Scientifique–FNRS. Computational resources have been provided by the Consortium des Équipements de Calcul Intensif en Fédération Wallonie Bruxelles (CECI) funded by the Fonds de la Recherche Scientifique (F.R.S.-FNRS) under convention 2.5020.11.



810 References

- Anderson, T. R.: Modelling the influence of food C:N ratio, and respiration on growth and nitrogen excretion in marine zooplankton and bacteria, *Journal of Plankton Research*, 14, 1645–1671, 1992.
- Anderson, T. R. and Pondaven, P.: Non-Redfield carbon and nitrogen cycling in the Sargasso Sea: pelagic imbalances and export flux, *Deep-Sea Research I*, 50, 573–591, 2003.
- 815 Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., and Gehlen, M.: PISCES-v2: an ocean biogeochemical model for carbon ecosystem studies, *Geoscientific Model Development*, 8, 2465–2513, 2015.
- Billen, G. and Fontigny, A.: Dynamics of a phaeocystis-dominated spring bloom in belgian coastal waters. II. Bacterio-plankton dynamics, *Marine Ecology Progress Series*, 37, 249–257, 1987.
- Bricaud, A., Babin, M., Morel, A., and Claustre, H.: Variability in the chlorophyll specific absorption coefficients of natural phytoplankton:
820 Analysis and parameterization, *Journal of Geophysical Research*, 100, 13 321–13 332, 1995.
- Butenschön, M., Clark, J., Aldridge, J., Allen, J., Artioli, Y., Blackford, J., Bruggeman, J., Cazenave, P., Ciavatta, S., Kay, S., Lessin, G., van Leeuwen, S., van der Molen, J., de Mora, L., Polimene, L., Saille, S., Stephens, N., and Torres, R.: ERSEM 15.06: a generic model for marine biogeochemistry and the ecosystem dynamics of the lower trophic levels., *Geoscientific Model Development*, 9, 1293–1339, 2016.
- Capet, A., Beckers, J.-M., and Grégoire, M.: Drivers, mechanisms and long-term variability of seasonal hypoxia on the Black Sea northwest-
825 ern shelf -is there any recovery after eutrophication?, *Biogeosciences*, 10, 3943–3962, <https://doi.org/10.5194/bg-10-3943-2013>, 2013.
- Capet, A., Meysman, F. J. R., Akoumianaki, I., Soetaert, K., and Grégoire, M.: Integrating sediment biogeochemistry into 3D oceanic models : A study of benthic-pelagic coupling in the Black Sea, *Ocean Modelling*, 101, 83–100, <https://doi.org/10.1016/j.ocemod.2016.03.006>, 2016.
- Ciliberti, S., Grégoire, M., Staneva, J., Palazov, A., Coppini, G., Lecci, R., Peneva, E., Matreata, M., Marinova, V., Masina, S., Pinardi, N.,
830 Jansen, E., Lima, L., Aydoğdu, A., Creti, S., Stefanizzi, L., Azevedo, D., Causio, S., Vandenbulcke, L., Capet, A., Meulders, C., Ivanov, E., Behrens, A., Ricker, M., Gayer, G., Palermo, F., Ilicak, M., Gunduz, M., Valcheva, N., and Agostini, P.: Monitoring and Forecasting the Ocean State and Biogeochemical Processes in the Black Sea: Recent Developments in the Copernicus Marine Service, *Journal of Marine Science and Engineering*, 9, 1146, <https://doi.org/10.3390/jmse9101146>, 2021.
- Cossarini, G., Querin, S., Solidoro, C., Sannino, G., Lazzari, P., Di Biagio, V., and Bolzon, G.: Development of BFMCOUPLER (v1.0), the
835 coupling scheme that links the MITgcm and BFM models for ocean biogeochemistry simulations, *Geoscientific Model Development*, 10, 1423–1445, <https://doi.org/10.5194/gmd-10-1423-2017>, 2017.
- Daewel, U., Schrum, C., and Macdonald, J. I.: Towards end-to-end (E2E) modelling in a consistent NPZD-F modelling framework (ECOSMO E2E_v1.0): application to the North Sea and Baltic Sea, *Geoscientific Model Development*, 12, 1765–1789, <https://doi.org/10.5194/gmd-12-1765-2019>, 2019.
- 840 Doerffer, R. and Schiller, H.: The MERIS case 2 water algorithm, *International Journal of Remote Sensing*, 28, 517–535, 2007.
- Droop, M.: 25 years of algal growth kinetics - personal view, *Botanica Marina*, 26, 99–112, 1983.
- Dutkiewicz, S., Hickman, A., Jahn, O., Gregg, W., Mouw, C., and Follows, M.: Capturing optically important constituents and properties in a marine biogeochemical and ecosystem model, *Biogeosciences*, 12, 4447–4481, <https://doi.org/10.5194/egusphere-2024-3682>, 2015.
- 845 Friedl, G., Dinkel, C., and Wehrli, B.: Benthic fluxes of nutrients in the northwestern Black Sea., *Marine Chemistry*, 62, 77–88, 1998.



- Friedrich, J., Dinkel, C., Friedl, G., Pimenov, N. and Wijsman, J., Gomoiu, M.-T., Cociasu, A., Popa, L., and Wehrli, B.: Benthic Nutrient Cycling and Diagenetic Pathways in the North-western Black Sea, *Estuarine, Coastal and Shelf Science*, 54, 369–383, 2002.
- Gayer, G., Dick, S., Pleskachevsky, A., and Rosenthal, W.: Numerical modeling of suspended matter transport in the North Sea., *Ocean Dynamics*, 56, 62–77, <https://doi.org/10.1007/s10236-006-0070-5>, 2006.
- 850 Geider, R., H., M., and Kana, T.: A dynamic regulatory model of phytoplankton acclimation to light, nutrients and temperature, *Limnology and Oceanography*, 43, 679–694, [https://doi.org/10.1016/S0012-8252\(00\)00004-0](https://doi.org/10.1016/S0012-8252(00)00004-0), 1998.
- Grégoire, M. and Soetaert, K.: Carbon, nitrogen, oxygen and sulfide budgets in the Black Sea: A biogeochemical model of the whole water column coupling the oxic and anoxic parts, *Ecological Modelling*, 221, 2287–2301, <https://doi.org/10.1016/j.ecolmodel.2010.06.007>, 2010.
- 855 Grégoire, M., Raick, C., and Soetaert, K.: Numerical modeling of the central Black Sea ecosystem functioning during the eutrophication phase, *Progress in Oceanography*, 76, 286–333, <https://doi.org/10.1016/j.pocean.2008.01.002>, 2008.
- Hedges, J., Hu, F., Devol, A., Hartnett, H., Tsamakidis, E., and Keil, R.: Sedimentary organic matter preservation: A test for selective degradation under oxic conditions, *Am. J. Sci.*, 299, 529–555, 1999.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, <https://doi.org/10.1002/qj.3803>, 2020.
- 860 Ho, D. T., Law, C. S., Smith, M. J., Schlosser, P., Harvey, M., and Hill, P.: Measurements of air-sea gas exchange at high wind 820 speeds in the Southern Ocean: implications for global parameterizations, *Geophys. Res. Lett.*, 33, 16 611, 2006.
- Hofmann, A., Meysman, F., Soetaert, K. E., and Middelburg, J. J.: A step-by-step procedure for pH model construction in aquatic systems, *Biogeosciences*, 5, 227–251, 2008.
- 865 Khalil, K., Rabouille, C., Gallinari, M., Soetaert, K., DeMaster, D., and Ragueneau, O.: Constraining biogenic silica dissolution in marine sediments: A comparison between diagenetic models and experimental dissolution rates, *Marine Chemistry*, 106, 223–238, 2007.
- Kowalczyk, P., Cooper, W. J., Whitehead, R. F., Durako, M. J., and Sheldon, W.: Characterization of CDOM in an organic-rich river and surrounding coastal ocean in the South Atlantic Bight, *Aquatic Sciences*, 65, 384–401, 2003.
- 870 Kremer, P.: Respiration and excretion by the ctenophore *Mnemiopsis leidyi*, *Marine Biology*, 44, 20–43, 1977.
- Kriest, I.: Different parameterizations of marine snow in a 1D–model and their influence on representation of marine snow, nitrogen budget and sedimentation, *Deep–Sea Research I*, 49, 2133–2162, 2002.
- Lancelot, C., Staneva, J., Van Eeckhout, D., Beckers, J., and Stanev, E.: Modelling the Danube-influenced North-western Continental Shelf of the Black Sea. II: Ecosystem Response to Changes in Nutrient Delivery by Danube River after its damming in 1972, *Estuarine, Coastal and Shelf Science*, 54, 473–499, 2002.
- 875 Laws, E. and Wong, D.: Studies on carbon and nitrogen metabolism by three marine phytoplankton species in nitrate-limited continuous culture, *Journal of Phycology*, 14, 406–416, 1978.
- Lee, Z.-P., Du, K.-P., and Arnone, R.: A model for the diffuse attenuation coefficient of downwelling irradiance, *Journal of Geophysical Research*, 110, C02016, 2005.
- 880 Lewis, E. and Wallace, D. W. R.: Program Developed for CO_2 System Calculations, ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee., 1998.
- Mace, L., Vandenbulcke, L., Brankart, J.-M., Brasseur, P., and Grégoire, M.: Characterisation of uncertainties in an ocean radiative transfer model for the Black Sea through ensemble simulations, *Egusphere*, 2024, 1–36, <https://doi.org/10.5194/egusphere-2024-3682>, 2024.



- Mague, T., Friberg, E., Hughes, D., and Morris, I.: Extracellular release of carbon by marine phytoplankton: a physiological approach.,
885 *Limnology and Oceanography*, 25, 262–279, 1980.
- Marilaure, G. and Vandenbulcke, L.: Biogeochemical Model for Hypoxic and Benthic Influenced areas (BAMHBI) v1.0.1,
<https://doi.org/10.5281/zenodo.16964655>, 2025.
- Mee, L., Friedrich, J., and Gomoiu, M.-T.: Restoring the Black Sea in times of uncertainty, *Oceanography*, 18, 32–43, 2005.
- Middelburg, J. and Meysman, F.: Burial at Sea, *Science*, 18, 1775–1810, <https://doi.org/10.1126/science.1144001>, 2007.
- 890 Millero, F. J.: Thermodynamics of the carbon dioxide system in the oceans, *Geochimica et Cosmochimica Acta*, 59, 661–677, 1995.
- Mobley, C., Stramski, D., Bissett, W. P., and Boss, E.: Optical modeling of ocean waters: Is the case 1 - case 2 classification still useful?,
Oceanography, 17, 60–67, 2004.
- Muramoto, J., Honjo, S., Fry, B., Hay, B., Howarth, R., and Cisne, J.: Sulfur, iron and organic carbon fluxes in the Black Sea : sulfur isotopic
evidence for origin of sulfur fluxes, *Deep Sea Research II*, 38, 1151–1188, 1991.
- 895 Nakamura, Y.: Growth and grazing of a large heterotrophic dinoflagellate, *Noctiluca scintillans*, in laboratory cultures, *Journal of Plankton
Research*, 20, 1711–1720, 1998.
- Neukermans, G., Loisel, H., Mériaux, X., Astoreca, R., and McKee, D.: In situ variability of mass-specific beam attenuation and backscat-
tering of marine particles with respect to particle size, density, and composition, *Limnol. Oceanogr.*, 57, 124–144, 2012.
- Raick, C., Delhez, E., Soetaert, K., and Grégoire, M.: Study of the seasonal cycle of the biogeochemical processes in the Ligurian Sea using
900 an 1D interdisciplinary model, *Journal of Marine Systems*, 55, 177–203, <https://doi.org/10.1016/j.jmarsys.2004.09.005>, 2005.
- Raick, C., Soetaert, K., and Grégoire, M.: Model complexity and performance: how far can we simplify?, *Progress in Oceanography*, 70,
27–57, <https://doi.org/10.1016/j.pocean.2006.03.001>, 2006.
- Ricour, F., Capet, A., D’Ortenzio, F., Delille, B., and Grégoire, M.: Dynamics of the deep chlorophyll maximum in the Black Sea as depicted
by BGC-Argo floats, *Biogeosciences*, 18, 755–774, <https://doi.org/10.5194/bg-18-755-2021>, 2021.
- 905 Smith, C. and Tett, P.: A depth-resolving numerical model of physically forced microbiology at the European shelf edge, *Journal of Marine
Systems*, 26, 1–36, 2000.
- Smith, R. C. and Baker, K. S.: Optical properties of the clearest natural waters (200–800 nm), *Appl. Opt.*, 20, 177–184, 1981.
- Soetaert, K., Herman, P., and J., M.: A model of early diagenetic processes from the shelf to abyssal depths, *Geochimica et Cosmochimica
Acta*, 60, 1019–1040, 1996.
- 910 Soetaert, K., Herman, P. M. J., Middelburg, J. J., Heip, C., Smith, C. L., Tett, P., and Wild-Allen, K.: Numerical modelling of the shelf break
ecosystem: reproducing benthic and pelagic measurements, *Deep Sea Research Part II: Topical Studies in Oceanography*, 48, 3141–3177,
[https://doi.org/10.1016/S0967-0645\(01\)00035-2](https://doi.org/10.1016/S0967-0645(01)00035-2), 2001.
- Soetaert, K., Middelburg, J. J., Meysmann, F. J. R., and Greenwood, J.: The effect of biogeochemical processes on pH, *Marine Chemistry*,
105, 30–51, 2007.
- 915 Soetaert, K., Petzoldt, T., and Setzer, R.: Solving Differential Equations in R: Package deSolve, *Journal of Statistical Software*, 33, 1–25,
<https://doi.org/10.18637/jss.v033.i09>, 2010.
- Soulsby, R.: Dynamics of marine sands: a manual for practical applications, *Oceanographic Literature Review*, 9, 947, 1997.
- Stanev, E. and Kandilarov, R.: Sediment dynamics in the Black Sea: numerical modelling and remote sensing observations, *Ocean dynamics*,
62, 533–553, <https://doi.org/10.1007/s10236-012-0520-1>, 2012.
- 920 Tett, P.: Parameterising a microplankton model, Napier University, 1998.



- Vaillancourt, R. D., Brown, C. W., Guillard, R. R. L., and Balch, W. M.: Light backscattering properties of marine phytoplankton: relationships to cell size, chemical composition and taxonomy, *Journal of Plankton Research*, 26, 191—212, 2004.
- Van Cappellen, P., Dixit, S., and van Beusekom, J.: Biogenic silica dissolution in the oceans: reconciling experimental and field-based dissolution rates, *Global Biogeochemical cycles*, 16, 1075, <https://doi.org/10.1029/2001GB001431>, 2002.
- 925 Van den Meersche, K., Middelburg, J., Soetaert, K., van Rijswijk P., H., B., and Heip, C.: Carbon-nitrogen coupling and algal-bacterial interactions during an experimental bloom: Modeling a ¹³C tracer experiment, *Limnol. Oceanogr.*, 49, 862–878, 2004.
- Vandenbulcke, L. and Grailet, J.-F.: BAMHBI stable, <https://gitlab.uliege.be/ESPECES/MAST/bamhbi-stable>, GitLab repository, 2025.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *Journal of Geophysical Research*, 97, 7373–7382, 1992.
- 930 Wijsman, J., Herman, P., Middelburg, J. J., and Soetaert, K. E.: A model for early diagenetic processes in sediments of the continental shelf of the Black Sea, *Estuarine Coastal and Shelf Science*, 54, 403–421, <https://doi.org/10.1006/ecss.2000.0655>, 2002.
- Williams, P.: The importance of losses duringn microbial growth: commentary on the physiology, measurement and ecology of the release of dissolved organic material, *Marine Microbial Food Webs*, 4, 175–206, 1990.
- Wolf-Gladrow, D. and Riebesell, U.: Diffusion and reactions in the vicinity of plankton: a refined model for inorganic carbon transport, *Marine Chemistry*, 59, 17–34, [https://doi.org/10.1016/S0304-4203\(97\)00069-8](https://doi.org/10.1016/S0304-4203(97)00069-8), 1997.
- 935 Xu, J., Hood, R. R., and Chao, S.-Y.: A simple empirical optical model for simulating light attenuation variability in a partially mixed estuary, *Estuaries*, 28, 572–580, 2005.
- Zaitsev, Y.: Impact of eutrophication on the Black Sea fauna, vol. 64 of *Tech.Rep.*, General Fisheries Council for the Mediterranean, FAO, Rome, 1993.