

Monsoons, plumes, and blooms: intraseasonal variability of subsurface primary productivity in the Bay of Bengal

Tamara L Schlosser¹, Andrew J Lucas^{1,2}, Melissa Omand³, and J. Thomas Farrar⁴

¹Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA, USA.

²Dept. of Mechanical and Aerospace Engineering, University of California, San Diego, La Jolla, CA, USA.

³Graduate School of Oceanography, University of Rhode Island, Narragansett, RI 02882, USA.

⁴Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA.

Correspondence: Tamara L Schlosser (tamara.schlosser@utas.edu.au)

Abstract. During the southwest monsoon, seasonal storms bring torrential rainfall to the South Asian subcontinent and the northern Indian Ocean. Dense cloud cover limits the amount of sunlight that reaches the ocean surface, and sediment-laden river runoff limits the depths to which light can penetrate. Changing light availability should affect phytoplankton primary productivity and its dependent biogeochemical processes, yet little is known about how subtropical weather is linked to ecosystem processes below the ocean's surface. Here, using novel physical and bio-optical measurements from an array of free-drifting, autonomous systems in the Bay of Bengal, we show that the onset of cloudy conditions associated with 'active' monsoon conditions led to >50% reduction in gross chlorophyll productivity (GCP) near the subsurface chlorophyll maximum (SCM) relative to sunny 'break' conditions. Optical backscatter measurements confirm chlorophyll fluorescence fluctuations correspond to biomass variability of a similar scale. Simultaneous bioacoustic measurements collected onboard the autonomous platforms suggest this intraseasonal variability in SCM chlorophyll and biomass generated a response in higher trophic levels. Long-term measurements from biogeochemical (BGC) Argo floats in the bay confirm the presence of intraseasonal oscillations in chlorophyll-*a* concentration with days-to-weeks variability in magnitude similar to the regional annual cycle in the region. Our findings demonstrate that intraseasonal subtropical air-sea variability modulates important regional biogeochemical ocean processes in the Northern Indian Ocean with implications for the Indian Ocean carbon cycle.

1 Introduction

The ocean contributes roughly 50% of total global autotrophic carbon fixation, with more than 40% of ocean primary productivity occurring in vast but relatively low biomass oligotrophic seas (Antoine et al., 1996). Our current understanding is that net primary productivity in the subtropical oligotrophic ocean is limited by nutrient supply and not light availability (Mignot et al., 2014), but recently developed autonomous long-duration measurements of subsurface processes allow a more comprehensive analysis of the subsurface variability than has previously been possible (Johnson and Bif, 2021). Less is known about how subtropical primary productivity varies on small time- and space-scales, or its response to the rapidly changing subtropical weather patterns.

Current generation coupled atmosphere-ocean numerical models lack fidelity on 10-90 day timescales in the tropics and subtropics (Gadgil et al., 2005; Mandke et al., 2020; Shroyer et al., 2021). This so-called intraseasonal variability is a major component of global weather and is fundamental to predicting climate phenomena like the South Asian monsoon (Shroyer et al., 2021). How intraseasonal fluctuations impact biogeochemical processes in the subtropical ocean is poorly understood (Hansell, 2009). Since coupled ocean-atmosphere models struggle to accurately represent the magnitude or the timing of intraseasonal fluctuations (Gadgil et al., 2005; Mandke et al., 2020; Shroyer et al., 2021), these fluctuations must also be inaccurately represented in coupled biogeochemical models. Intraseasonal oscillations like the Madden-Julian Oscillation or the Monsoon Intraseasonal Oscillation impact much of the subtropical ocean, so the potential impact on biogeochemical processes may be significant.

Satellite observations reveal the decrease in surface productivity during the southwest monsoon (approximately June to September) in the Bay of Bengal due to the impact of monsoon weather on light availability at monthly scales (Kumar et al., 2010a, b). Light availability at the surface decreases via a reduction in shortwave radiation during the southwest monsoon (Weller et al., 2016), which oscillates at 10-90 day timescales, categorized as the sunny ‘break’ period and rainy and cloudy ‘active’ periods of the monsoon (e.g., Shroyer et al., 2021). Subsurface light availability additionally decreases due to the increasing diffuse attenuation following river run-off, where monsoon rain events inject high concentrations of sediments into the bay, even $\mathcal{O}(100\text{ km})$ from the coastline (Kumar et al., 2010a). The weather systems that reduce shortwave radiation also impact ocean color satellite measurements, so satellite measurements cannot observe biogeochemical processes over the entire intraseasonal oscillation. The subsurface impacts of intraseasonal varying light availability are generally unknown, contributing to the uncertainty in our understanding of the Indian Ocean carbon cycle. In particular, stratification, vertical salinity gradients and barrier layers have a dominating influence on subsurface biomass in the Bay of Bengal (Prasanth et al., 2023).

Autonomous technology to measure biogeochemical variability in the ocean – such as the BGC-Argo float program (Argo, 2000) – has provided new opportunities to quantify the variability of the ocean carbon system (Mignot et al., 2014; Johnson and Bif, 2021). These platforms provide information about biogeochemical transformations below the ocean’s surface that are largely invisible to satellite remote sensing (Mis, 2005). Although only recently deployed at scale, autonomous biogeochemical measurements have demonstrated that our capacity to make future predictions of the state of the climate depends sensitively on constraining biogeochemical ocean processes (Gray et al., 2018; Johnson and Bif, 2021).

The BGC-Argo float program typically samples at weekly time scales, which is sufficient to describe the intraseasonal biogeochemical variability but not diel processes, such as nonphotochemical quenching (NPQ) and the diel cycle in phytoplankton biomass. Near the surface, NPQ results in a non-biomass-related diel reduction in chlorophyll-*a* (Chl) fluorescence under high light conditions (Müller et al., 2001). In contrast, the diel cycle is distinct from NPQ in that concentrations of Chl (Marra, 1997; Prasanth et al., 2023), dissolved oxygen (Nicholson et al., 2015; Barone et al., 2019), and particulate carbon (White et al., 2017) increase with increasing irradiance. The diel cycle results from the competing processes of day-time gross primary production (GPP) and day- and night-time respiration and grazing losses (Marra, 1997). The tight coupling between the diel cycle and the solar cycle (i.e., sunrise and sunset) has allowed the development of robust methodologies for estimating GPP and loss rate for a range of biological variables (Marra, 1997; Nicholson et al., 2015). Although the diel cycle directly results

from the diel variability in subsurface irradiance, the sensitivity of the diel cycle to the short-term cloud- or turbidity-driven changes in irradiance remains under-explored.

60 An improved understanding of the Indian Ocean carbon cycle and its sensitivity to intraseasonal oscillations not only requires a better understanding of primary productivity but also the effect on higher trophic levels. Currently, most autonomous platforms like BGC-Argo deployed long-term do not observe grazers due to the difficulty in making accurate observations, either directly or indirectly. Acoustic backscatter measurements at high frequencies (i.e., 1 MHz) are sensitive to smaller zooplankton species, including grazers, as compared to lower frequency measurements of acoustic backscatter traditionally used to identify 65 fish and larger marine organisms (Lavery et al., 2007; Ohman et al., 2019; Gastauer et al., 2022). However, without confirming net or video observations (e.g., Lavery et al., 2007), these bioacoustic or high-frequency acoustic backscatter observations (as we present here) remain a relatively uncertain representation of grazer variability.

Here we report on physical, bio-optical, and bio-acoustic measurements gathered in the Bay of Bengal during the 2019 monsoon season, described in the next section. We then quantify the impact of ocean clarity and passing monsoon storms on 70 subsurface irradiance, and we statistically relate fluctuations in gross production to the subsurface irradiance. We contextualized our observations with an analysis of longer-term BGC-Argo measurements. Finally, we conclude by discussing some of the wider implications of our results, including regional patterns in subsurface productivity we may infer from our results, and evidence of higher trophic level variability. Taken together, our results demonstrate the variability in biogeochemical quantities on intraseasonal time scales, and show how increased autonomous observational capacity can contribute to our understanding 75 of the processes that shape the ocean climate system.

2 Materials and Methods

2.1 Field campaign

As part of a campaign to study intraseasonal oscillations in monsoon weather, we deployed an array of three densely instrumented buoys and ocean-wave-powered Wirewalker profiling vehicles (Pinkel et al., 2011) that gathered physical, bio-optical, 80 and irradiance measurements in July 2019 in the Bay of Bengal. Each Drogued Buoy Air-Sea Interaction System (DBASIS) was deployed for ~ 19 days (Fig. 1): M1 on July 9, M2 on July 10, and M3 on July 11, generally following the mesoscale flow. Each profiler had a metocean buoy on the surface, a vertically profiling Wirewalker over the upper 100 m (Pinkel et al., 2011), and six X-wings (1 m^2 drag elements) over 200 m to 210 m depth so that the system drifted with the subsurface currents at 200 m. They collected approximately 200 profiles per day spanning the upper 100 m of the water column with sub-meter 85 vertical resolution.

The buoy's meteorological package measured wind (2.9 m above mean sea level; MSL), air temperature (2.6 m above MSL), humidity (2.6 m above MSL), precipitation (2.9 m above MSL), barometric pressure (2.3 m above MSL), sea surface temperature (0.3 m and 1.5 m below MSL), and downwelling solar and infrared radiation (3.2 m above MSL). The metocean buoys were equipped with Kipp and Zonen SMP21 shortwave radiation sensors that measure downwelling radiation over wavelengths

90 of 285 nm to 2800 nm (50% points). These measurements were used to estimate the surface photosynthetically available radiation (PAR).

We equipped the Wirewalker profilers with RBR Inc. conductivity, temperature, and depth (CTD) sensors augmented by SBE Wetlabs Ecopucks measuring Chl fluorescence (ChlF, a proxy for Chl biomass), optical backscatter at 532 nm, and chromophoric dissolved organic material (CDOM; not used in the analysis presented here). We measured subsurface downwelling
95 irradiance onboard the Wirewalkers using a Satlantic OCR-504 multi-spectral radiometer at four bands, 380 nm, 412 nm, 490 nm and 532 nm. All parameters were collected continuously at 6 Hz and telemetered at that resolution in real-time via RUDICS Iridium modems on the surface buoy. Using the factory calibration, we convert the optical backscatter to Nephelometric Turbidity Unit, τ (NTU). The median time between profiles was 10 min at M2 and M3 and 20 min at M1. For all quantities, we only use measurements during the smooth upward ascent of the profiler.

100 To estimate the diffuse attenuation averaged by depth, K_d (m^{-1}), the raw measurements of downward irradiance were grouped in time and depth into 3 h and 4 m bins, respectively, and then least-squares fitted to an exponential curve. Irradiance was then interpolated back to the original $\mathcal{O}(10 \text{ min})$ time resolution (see supplementary materials for further details).

We also equipped the profilers with a downward-looking Nortek Signature with a frequency of 1 MHz, from which we show only the acoustic backscatter (Zheng et al., 2022). For each profiler’s upcast, we averaged the acoustic amplitudes of each
105 of the four beams over the first 20 bins (2.5 m depth range) closest to the transducer. We then interpolated range-averaged beam amplitudes onto a 0.25 m uniform depth grid before averaging all four beams. The 1 MHz backscatter was similar to the R/V Sally Ride acoustic backscatter that had a frequency of 150 kHz, with differences between frequencies as expected from literature (Lavery et al., 2007; Ohman et al., 2019; Gastauer et al., 2022).

All three DBASIS floats had the same bio-optical instrumentation and generally collected comparable measurements. We
110 present time series from only M3 here, which collected continuous (unlike M1) and high resolution (10 min) data, and show M1 and M2 observations in the supplementary materials. Our key results and findings were similar for all floats.

2.2 Chlorophyll-a fluorescence

The primary objective of the field campaign was to investigate air-sea interactions and the underlying physical variability in the BoB. As such, while biogeochemical observations were collected autonomously, we did not sample water in order to field
115 calibrate the chlorophyll sensors. Instead, using laboratory calibration carried out just prior to the experiment, we converted the observed ChlF in relative fluorescence units (RFU) to real units ($\mu\text{g L}^{-1}$) via $\text{ChlF} = 0.012 \times (\text{RFU} - 50)$. This conversion has an unknown uncertainty that linearly scales for all values (i.e., percentage error). For context, the observed surface ChlF from the drifting systems and MODIS-Aqua surface Chl from July 2019 were $0.17 \mu\text{g L}^{-1}$ and $0.33 \mu\text{g L}^{-1}$, respectively, with these points located an average of 54 km apart. The focus of our work is not on the absolute quantities of ChlF but rather on
120 their response to surface and penetrative solar radiation, so we focus on the robust trends in co-variability rather than absolute values.

Chlorophyll pigment concentration, ChlF, and phytoplankton biomass (i.e., carbon content) are highly variable under changing light conditions as phytoplankton adapt to changing light environments by adjusting their light-harvesting pigments (Cullen,

1982, 2015). Due to this, estimates of biomass from backscatter observations are typically a more robust estimate of carbon biomass than ChlF. However, in the situation encountered here, non-algal sources of backscatter at-times confounded measure of algal concentrations, for example in the high-sediment coastal waters we observed at M3 before July 15 (Fig. 2d). Focusing only on times when the backscatter signal was due to algal diel variability, ChlF and backscatter variability was similar. This analysis showed that ChlF was a good proxy for phytoplankton biomass at the depth of the subsurface chlorophyll maximum (SCM), which we discuss throughout the manuscript.

130 Near the surface, non-photochemical quenching (NPQ) results in a non-biomass-related diel reduction in fluorescence under high light conditions (Müller et al., 2001). To correct for NPQ in the ChlF measurements, we follow methods established for other rapidly profiling platforms (Davis et al., 2008; Todd et al., 2009; Schlosser et al., 2022). We least-squares fit irradiance at 490 nm to ChlF over the upper 20 m over one-day windows, stepping in time every 0.1 day and averaging overlapping time steps. We set a maximum correlation coefficient of -0.1 , ensuring we only modify ChlF if ChlF decreased when irradiance increased. Although we used only ChlF over the upper 20 m for the least-squares fitting, we corrected ChlF for all depths with non-zero irradiance. The NPQ correction to ChlF was small, with a 95th percentile of $0.016 \mu\text{g L}^{-1}$. Further details are available in the supplementary materials.

At all profilers, the depth of maximum ChlF was identified within a narrow density range. Excluding the initial 2 days observed at M1 (see supplementary materials), the SCM and the 1021.5 kg m^{-3} isopycnal were significantly correlated ($R^2 = 0.85$, $p < 0.001$) with a mean absolute error of 1.65 m. By defining the SCM as the depth of the 1021.5 kg m^{-3} isopycnal $\pm 1 \text{ m}$, the SCM can be matched to an along-isopycnal reference frame. Given the internal wave fluctuations ($\sim 15 \text{ m}$) observed during the campaign, employing an along-isopycnal reference frame was convenient for isolating the impact of changing surface irradiance and diffuse attenuation on ChlF at the SCM (denoted ChlF_{SCM}). The 1021.5 kg m^{-3} isopycnal had an average depth of 39 m at M3 (Fig. 2) that deepened to 50 m at M1.

145 2.3 Diel cycle method

ChlF and integrated ChlF reflects the standing stock of chlorophyll, and therefore integrates pigment concentration over days to weeks. To understand the rate at which ChlF is produced and lost due to the local conditions at the SCM, we applied the diel cycle method (Nicholson et al., 2015; Barone et al., 2019). We identify the rate of gross ChlF production (GCP_{SCM}) and ChlF loss (code available at <https://github.com/duebi/dielFit>) to the NPQ corrected ChlF_{SCM} . This loss term describes all processes resulting in a reduction of ChlF_{SCM} , including grazing, mortality, and downward export.

The diel cycle method linearly fits three terms via ordinary least-squares fitting to ChlF_{SCM} . This allows us to directly evaluate the response of SCM production to intraseasonally modulated PAR, rather than relying on changes in the accumulated concentration. We note that when deciding the depth region to vertically average ChlF_{SCM} over (2 m), averaging a larger depth range decreased the effectiveness of the diel cycle fit (i.e., the correlation coefficient decreased and mean absolute error increased). The three fitted terms in the diel cycle method included a constant, a constant loss rate for both day and night, and a variable in time GCP_{SCM} rate similar to the clear-skies PAR, with day-time growth maximum at noon and a zero rate of change at night.

Since we collected radiation measurements, we did not need to simulate the solar cycle. We thus modified the code to optionally include the observed PAR at the SCM (denoted PAR_{SCM}) to model growth (henceforward referred to as the PAR_{SCM} method). The PAR_{SCM} method is similar to the sinusoidal growth model but accounts for the impact of cloudiness, changes in diffuse attenuation, or perturbations of the SCM depth on growth. Before fitting to ChlF_{SCM} , we integrated each growth scheme in time. For all three profilers and days sampled, we fit each 26-h period (starting at 11 pm local), overlapping each day by one hour. We additionally modified the diel cycle methodology so that the fitted GCP_{SCM} and loss is always positive.

The performance of the fit was assessed by estimating the correlation coefficient and the mean absolute error, $\text{MAE} = 1/n \sum ||\text{ChlF}_{\text{SCM}} - \text{ChlF}_{\text{fit}}||$, where ChlF_{SCM} is the observed ChlF at the SCM, ChlF_{fit} is the diel cycle fit, and n is the number of samples. Of the tested linear, sinusoidal, and PAR_{SCM} schemes for GCP_{SCM} , the PAR_{SCM} method consistently returned the smallest MAE and better predicted the timing of the observed ChlF_{SCM} maxima by 45 min on average across all floats (Table 1). The sinusoid method also returned a good fit, with a larger MAE, but the linear method was relatively poor in describing the observed ChlF_{SCM} variability. The fitted mean absolute error doubled when fitting ChlF at a constant depth rather than along-isopycnal due to the vertical heaving of the SCM by passing internal waves (i.e., ChlF_{SCM} ; Table 1). The fitting routine was also effective at M1 and M2. At M1, the timing of peak afternoon ChlF_{SCM} was more variable, particularly when large-amplitude internal waves deepened the SCM during the day (see supplementary materials). We also applied the above diel cycle fitting methodology to turbidity to estimate gross turbidity production (GTP_{SCM}) and turbidity loss. We computed the confidence intervals of the model by bootstrapping the residuals with 200 iterations (Nicholson et al., 2015; Barone et al., 2019).

2.4 BGC-Argo analysis

To contextualize our observations and investigate the frequency of light-limited productivity at the SCM in the bay, we analyzed 12 BGC-Argo floats deployed in the BoB, obtained via the OneArgo-Mat routine (Frenzel et al., 2020). These floats were equipped with a CTD and bio-optical sensors (WET Labs ECO-FLBB AP2 or MCOMS, see Johnson et al., 2017) measuring temperature, salinity, pressure, chlorophyll fluorescence (ChlF), optical backscatter coefficient at 700 nm (at 124°), as well as other sensors not used here. The data were quality controlled using standard Argo protocols (Wong et al., 2022) and processed following Uchida et al. (2019). Quality-controlled data was used when available (quality flags 1, 2, 5 or 8), else visual inspection and simple outlier removal was performed (quality flags 0 or 3).

The raw signals were converted to ChlF ($\mu\text{g L}^{-1}$) and particulate backscattering coefficient (m^{-1}) following BGC-Argo procedures (Schmechtig et al., 2018). To account for variations between sensors, we standardized the minimum ChlF and backscatter value by removing the median ‘dark’ or background value at pressure >600 dbar for each float (Uchida et al., 2019). BGC-Argo floats vary in their temporal and vertical sampling frequencies, so following Uchida et al. (2019) we interpolated the quality-controlled float data onto uniform temporal grids with time steps equal to the minimum temporal sampling rate, which is 10 days for most floats. We standardized the vertical grid from 4 to 1000 m. It had 2 m resolution in the upper 300 m and 10 m resolution below that. This interpolation was performed using a hermite polynomial scheme (pchip in Mat-

lab), but depths outside the observed range were excluded (i.e., no extrapolation). We applied a 3-point moving-median in the vertical to remove measurement noise and aggregates.

The BGC-Argo subsurface ChlF was also maximum near a consistent isopycnal, like the DBASIS observations, except ChlF was maximum near the 1022 kg m^{-3} isopycnal, so we defined this isopycnal as the SCM depth. Individual months could have enhanced ChlF near a slightly different isopycnal, like the 1021.5 kg m^{-3} isopycnal used when analyzing the DBASIS floats, but we simply use the same isopycnal year-round. To estimate the ChlF_{SCM} , we averaged ChlF over the 10 m above and below this depth. Since the floats profile once every 10 days, and there are too few floats to apply more advanced statistical techniques (Johnson and Bif, 2021), we cannot estimate GCP_{SCM} from diel cycles for the BGC-Argo data. Instead, we examined the floats for any evidence of a correlation between ChlF_{SCM} and PAR at the surface and SCM (described below). We note the diel cycle in ChlF_{SCM} is prevalent in the Bay of Bengal (Lucas et al., 2016) and has the potential to alias multi-day trends in ChlF if measurements of ChlF are taken at different local times, as is the case for BGC-Argo measurements. To avoid this aliasing, in regions with diel cycles, ChlF and other biological or bio-optical variables should be measured at similar times of the day.

2.5 Satellite and reanalysis data

To understand the co-variance of light and BGC-Argo observed ChlF, we considered both MODIS Aqua PAR (NASA Goddard Space Flight Center, 2018b) and the atmospheric reanalysis ECMWF Reanalysis v5 (ERA5) product (Hersbach et al., 2018). On average, the PAR estimated at M3 was less than the satellite derived estimate but similar to the ERA5 estimate. We thereby estimated surface PAR from the ERA5 shortwave radiation, by dividing shortwave radiation by 2.114 (Britton and Dodd, 1976). Then, to estimate PAR at the SCM, we utilized the 490 nm diffuse attenuation (K_{d490}) from the monthly averaged MODIS-Aqua (Goddard Space Flight Center, 2018a) and converted to a PAR equivalent using the relation in Morel et al. (2007). If this data was missing due to cloud cover, we took the monthly climatology value instead. For a short-term estimate of 490 nm diffuse attenuation (K_{d490}) at the start of our DBASIS time series, we analyzed the Ocean Color CCI (OC-CCI) merged satellite product, which had a spatial resolution of 4 km and was averaged over the 5-day period of July 10 to 15, 2019 (Sathyendranath et al., 2019, 2023).

3 Results

3.1 Southwest monsoon campaign

The 2019 field campaign enabled a detailed investigation of the co-variability of subsurface physical and bio-optical properties on an hour-by-hour basis over a nearly 3-week deployment (Figs. 1 and 2). Conditions at deployment were consistent with the Southwest monsoon ‘break’ period with weak winds ($\sim 7.5 \text{ m s}^{-1}$), large net heat fluxes into the ocean (Q_N , daily average Q_N of 180 W m^{-2}), and no precipitation (Fig. 2).

The DBASIS array was deployed at the intersection of two eddies, as inferred from sea surface height (SSH, Fig. 1). Low-SSH ($\text{SSH} < 0.5 \text{ m}$) and high-SSH ($\text{SSH} > 0.5 \text{ m}$) eddies were present to the northwest and southwest, respectively. Between

these two eddies, a plume of elevated surface chlorophyll was advected more than 400 km from the coast toward the DBASIS array (Fig. 1a). This coastal plume exhibited elevated turbidity (τ) and surface chlorophyll-*a* fluorescence (ChlF), both of which reduced light penetration into the upper ocean (Fig. 2). Similar coastal plumes have been observed elsewhere at the intersection of cyclonic and anticyclonic eddies (i.e., eddy dipole, Malan et al., 2020).

Over the 19-day deployment, the array drifted southeastward and sampled two distinct regimes— one influenced by a coastal plume and one outside its influence. Surface turbidity and ChlF decreased after 5 days, following the profiler exiting the coastal plume. Below the surface, the pattern was reversed. ChlF at the subsurface chlorophyll maximum (SCM, ChlF_{SCM}) was initially relatively low and then approximately doubled after the drifting platforms left the plume (July 14 to 20). Meanwhile, turbidity near the SCM (τ_{SCM}) initially decreased and subsequently increased as ChlF_{SCM} increased (Fig. 2d-e). Outside the plume, turbidity is expected to primarily reflect algal biomass. The observed turbidity outside the plume, especially with its co-variability with ChlF near the SCM, indicates that variability in ChlF reflects changes in biomass and not just pigment concentrations.

Diffuse attenuation was strongly influenced by turbidity and was enhanced within the plume, limiting PAR_{SCM} (± 1 m of 1021.5 kg m⁻³ isopycnal, Fig. 3b). After the drifting buoys left the plume, however, turbidity and diffuse attenuation decreased, leading to a doubling of maximum day-time PAR_{SCM}, even though surface PAR was relatively steady. This increase in PAR_{SCM} was coincident with increasing ChlF_{SCM} (Fig. 3a). From July 18 to 23, wind speeds decreased to an average of 4.4 m s⁻¹, net heat fluxes were consistently large at a daily average of 139 W m⁻², and the total rainfall measured 4.4 mm (Fig. 2). Subsequently, the atmospheric conditions transitioned from mostly sunny conditions to mostly cloudy conditions around July 23 with the onset of the ‘active’ phase of the Southwest Monsoon (Fig. 2a). From July 23 to 28, surface-buoy-measured shortwave radiation and net heat fluxes decreased, with a daily average Q_N of -39 W m⁻², winds increased to an average of 8.8 m s⁻¹, and rainfall remained low, totaling 28 mm (Fig. 2). During this time, PAR_{SCM} dropped significantly and ChlF_{SCM} decreased, even though near-surface turbidity and K_d were at a minimum.

3.2 Gross ChlF production from diel cycles

The rapid vertical profiling of the Wirewalkers allowed the assessment of bio-optical variability from a time-varying, along-isopycnal frame of reference (Fig. 2d vs. 3a). We use the observed density to convert ChlF to an along-isopycnal reference frame. This approach moderates the effect of passing internal waves that confound measurements taken at discrete depths and highlighted the variability of ChlF at diel timescales (Fig. 3). Due to high variability in surface densities, we omit the upper 20 m in the along-isopycnal figure (Fig. 3a). We also note that due to the relatively dense surface densities from July 12 to 18, the along-isopycnal figure has missing data (black regions) to up to 30 m depth (≤ 1021.3 kg m⁻³ isopycnal), as no ChlF observations were made at these densities. The diel cycle at the SCM showed peak concentrations around 3 hours after local noon and minimum concentrations at dawn (Fig. 3a,c). The noon maximum in PAR_{SCM} (Fig. 3b teal) coincided with a rapid increase in ChlF_{SCM}, as found previously in the bay (Lucas et al., 2016). All records were corrected for NPQ, which had a fairly small effect on ChlF variability.

255 The consistent timing of irradiance and the diel cycle (e.g., maximum in ChlF rate of change at noon, not shown) has been used to estimate gross production and loss from dissolved oxygen and other biological variables (Marra, 1997; Nicholson et al., 2015; Barone et al., 2019; Johnson and Bif, 2021). Here we adapt these diel models to estimate the gross ChlF production (GCP) and the rate of ChlF loss from the on-isopycnal variability of ChlF and co-located PAR. We estimate at the SCM (GCP_{SCM}) and loss. By modeling growth from the co-located PAR (PAR_{SCM}), the impact of variable cloud cover, changes in ocean clarity, 260 or passing internal waves on growth can be accounted for, improving the performance of the model.

The observed diel periodicity and the multi-day trend in $ChlF_{SCM}$ were effectively emulated by the model for each DBASIS platform (Fig. 3c and supplementary materials). The fitted rate of $ChlF_{SCM}$ growth was closely coupled to $ChlF_{SCM}$ loss (Fig. 3d), reflecting strong recycling in this oligotrophic setting. When the rate of $ChlF_{SCM}$ loss exceeded GCP_{SCM} , daily-averaged $ChlF_{SCM}$ decreased even when GCP_{SCM} and the diel variability in $ChlF_{SCM}$ remained large (from July 21, Fig. 3c-d).

265 The observed GCP_{SCM} increased and decreased following respective changes in PAR_{SCM} . PAR_{SCM} was a function of short-wave radiation at the surface (largest effect, e.g., July 22 onwards, Fig. 3), diffuse attenuation (K_d) in the water column (e.g., July 12 to 15), and the SCM depth (e.g., July 20 to 21 at M1, see supplementary materials). Over the deployment, GCP_{SCM} linearly scaled with the daily averaged shortwave radiation and PAR_{SCM} (Fig. 4a,c), with a significant positive correlation ($r^2 \geq 0.34$, $p < 0.001$, and $n = 40$). From this linear regression, as shortwave radiation decreased by 80% following the tran- 270 sition from the break to active monsoon conditions (July 21 to 26), GCP_{SCM} decreased by 82%. Similarly, as the drifting systems left the coastal plume, PAR_{SCM} and the regression-estimated GCP_{SCM} increased by 58% (July 12 to 21). The fit with in situ PAR_{SCM} was superior to the fit with shortwave radiation due to the influence of the time-variable diffuse attenuation reflected in that measurement, with days with $K_d^{-1} \leq 15$ m decreasing the performance of the shortwave radiation fit. These findings suggest that spatial variability in diffuse attenuation, due to ocean clarity, and surface irradiance, due to intraseasonal 275 oscillations in monsoon weather, act together to influence subsurface productivity in the northern Bay of Bengal.

We also note that due to the SCM depth being approximately 10 m deeper at M1 compared to the other profilers, PAR_{SCM} was lowest, on average, at this profiler. We can slightly improve the fit (MAE decreased by 12%) of GCP_{SCM} to PAR_{SCM} (Fig. 4b) by accounting for phytoplankton light adaption by relating GCP_{SCM} to $PAR_{SCM}/\overline{PAR_{SCM}}$ instead of PAR_{SCM} , where $\overline{PAR_{SCM}}$ is the along-isopycnal average of the PAR_{SCM} at each profiler. This suggests phytoplankton were adapted to a lower 280 irradiance (i.e., isolume) at M1 than the other profilers, but regardless, GCP_{SCM} similarly increased for larger PAR_{SCM} .

Despite the occasional non-algal sources of high turbidity, especially at M3, we estimated a good diel cycle fit ($r^2 \geq 0.4$) in turbidity for 28 days in total across the three profilers. The estimated gross turbidity production at the SCM (GTP_{SCM}) linearly varied with PAR at the surface and SCM, similar to GCP_{SCM} variability (Fig. 4b,d). This indicates the co-variability of $ChlF_{SCM}$ and light availability is not limited to pigment concentration (e.g., photo-acclimation) but also corresponds to changes 285 in biomass.

Our findings of light-limited ChlF could additionally be found by analyzing the instantaneous $ChlF_{SCM}$ and depth-integrated ChlF, but the relationships were much weaker. We compared depth-integrated ChlF over the upper 80 m to depth-integrated PAR, and $ChlF_{SCM}$ to PAR_{SCM} . Both linear regressions returned weak correlations ($r^2 \leq 0.13$) that remained significant ($p < 0.001$, not shown). The instantaneous and depth-integrated ChlF was additionally influenced by ChlF loss and represents the

290 accumulated biomass rather than just growth. Despite this, the strong influence of light availability on ChlF variability remained evident in both instantaneous ChlF_{SCM} and GCP_{SCM}.

3.3 Barrier layer stability

In the southern BoB, SCM productivity was linked to barrier layer thickness, defined as the depth-region between the mixed layer depth (MLD) base and the isothermal layer (ITL, Prasanth et al., 2023). Here, we define the ITL depth as the shallowest
295 depth where the temperature was at least 1 °C cooler than the sea surface temperature ($\Delta T \leq 1$ °C), and the MLD as the shallowest depth where the density increase from the surface value corresponds to a temperature decrease of 1 °C. The ITL was on-average within 1.5 m of the SCM depth at M3 and the barrier layer thickness was highly dependent on salinity variations (Fig. 5a-c).

While M3 was within the coastal plume (before July 15), the MLD approximately equaled the SCM depth, which continued
300 til July 17 (Fig. 5a-b). While within the plume, water temperature and salinity at the surface and MLD were different, indicating a very thin barrier layer was present, but ChlF was the same (Fig. 5d-f). A passing storm resulted in a shallow rain-layer forming around July 17, the rapid shoaling of the MLD, diverging surface and MLD temperature and salinity (see Kerhalkar et al., 2025), and a rapid decrease in ChlF at the MLD base (Fig. 5). Over the remaining observation period, the MLD deepened until it reached the SCM around July 25. ChlF at the MLD and SCM then converged, and surface ChlF increased despite the
305 transition to active monsoon conditions and decrease in light availability. The thick barrier layer that formed from the rain event should theoretically isolate upper ocean mixing from the SCM. The observed increase in ChlF_{SCM} and GCP_{SCM} from July 18 may therefore be due to both the sustained sunny break conditions and a decrease in SCM mixing from the barrier layer formation (Fig. 3d).

3.4 Regional and seasonal context

310 To determine whether our observations of co-varying PAR_{SCM} and GCP_{SCM} in the central Bay of Bengal were representative of the region and season, we analyzed the BGC-Argo subsurface ChlF, ERA5 estimated PAR and MODIS-Aqua diffuse attenuation climatology (see Sections 2.4 and 2.5). To illustrate, we show one example of positively correlated PAR_{SCM} and ChlF_{SCM} over 12 months (WMO ID: 2902193; $r^2 = 0.58$, $p < 0.001$, and $n = 70$; Fig. 6). At the seasonal time scale, trends in PAR_{SCM} were more similar to SCM depth than the incident surface PAR, where PAR exponentially decays with depth and so
315 SCM depth can largely control PAR_{SCM}. In contrast, during the monsoon months of June to September, temporal variability was comparable between surface PAR and ChlF_{SCM}. This intraseasonal signal was sufficiently strong such that during the southwest monsoon period, surface PAR and subsurface ChlF (averaged over the upper 140 m) were significantly positively correlated ($r^2 = 0.26$, $p < 0.01$, and $n = 27$), even without accounting for changes in SCM depth.

The BGC-Argo float observed the largest change in ChlF_{SCM} during the southwest monsoon between the end of July and
320 start of August. The minimum and maximum PAR_{SCM} around this time were on July 23 and August 6, with observed ChlF_{SCM} on these days of 0.38 $\mu\text{g} \cdot \text{L}^{-1}$ and 1.65 $\mu\text{g} \cdot \text{L}^{-1}$, respectively (Fig. 6). The time of observation will also impact ChlF_{SCM} in this region of strong diel cycles. We note the time of sampling of the minimum and maximum PAR_{SCM} was at 4.12 pm and 3.08 am

local, respectively, where the DBASIS float observed daily maximum ChlF_{SCM} at 3 pm and minimum at 7 am, on average (Fig. 3c). The over 300% increase in ChlF_{SCM} observed by BGC-Argo was not only due to the transition from active to break monsoon conditions (Fig. 6), but also the shoaling of the SCM that further increased light availability.

If we contrast to the DBASIS platform observations, at M3 ChlF_{SCM} decreased from $0.96 \mu\text{g.L}^{-1}$ to $0.58 \mu\text{g.L}^{-1}$ on July 21 to 25, and at the same local time as the minimum and maximum BGC-Argo sample (Fig. 3c). This equates to a 40% difference at M3, compared to the 77% difference in the BGC-Argo float. We note the change in gross ChlF production at the SCM at the DBASIS platform was double the ChlF_{SCM} change, at an $\geq 80\%$ decrease, where this number is independent of sampling time (Fig. 3e). This may suggest changes in gross ChlF production due to intraseasonal oscillations at the BGC-Argo exceeded what we observed at M3, given variability in ChlF_{SCM} was larger at the BGC-Argo float than at M3.

4 Discussion

4.1 Primary productivity during the southwest monsoon

During the 2019 monsoon in the central Bay of Bengal and away from the coastal plume, ChlF and turbidity had a subsurface maximum, denoted the subsurface chlorophyll maximum (SCM, Fig. 2), and the SCM almost always coincided with the 1021.5 kg m^{-3} isopycnal. Both ChlF and turbidity also had a diel periodicity at the SCM, further suggesting that variations in ChlF at the SCM represent similar changes in biomass, as has been previously shown in the Indian Ocean waters (Cornec et al., 2021; Prasanth et al., 2023). The diel periodicity in ChlF has been observed elsewhere in the BoB (Lucas et al., 2016; Prasanth et al., 2023), and highlights a tight coupling between phytoplankton growth and loss in these oligotrophic waters. At the SCM, the gross production derived from both ChlF and turbidity varied with PAR_{SCM} (Fig. 4). Estimated gross ChlF production at the SCM (GCP_{SCM}) responded to variations in light from passing monsoon storms and changing ocean clarity (Fig. 4b and supplementary materials). GCP_{SCM} , estimated from the diel cycle in ChlF_{SCM} , linearly scaled with surface shortwave radiation at each DBASIS system, but the fit was sensitive to changing ocean color and turbidity. By accounting for variations in diffuse attenuation by linearly scaling GCP_{SCM} with PAR_{SCM} rather than PAR at the surface (PAR_0), we improved the linear fit (Fig. 4). We conclude we observed light limited SCM growth, contrary to prior studies showing the subtropical oligotrophic waters were typically nutrient limited (e.g., Mignot et al., 2014).

The estimated GCP_{SCM} rapidly decreased following the transition from sunny ‘break’ to cloudy ‘active’ monsoon conditions by $\geq 80\%$ in the northern BoB (Fig. 3d-e). This intraseasonal signal in ChlF gross production was also evident in BGC-Argo ChlF measurements from central BoB, with a similar difference in ChlF_{SCM} levels during the active vs. break periods as that observed by our measurements (e.g., July 24 to August 8 in Fig. 6a). Due to the southwest monsoon, surface PAR fluctuates on timescales of days to months over the entire bay (Gadgil et al., 2005; Mandke et al., 2020; Shroyer et al., 2021). Average surface PAR increased from 20°N to 8°N , while the monthly standard deviation in surface PAR decreases from north to south (Fig. 7). This indicates that the southern BoB waters have higher and more consistent surface PAR, and towards the north, intraseasonal oscillations have a larger impact on surface PAR. We hence predict our observed $\geq 80\%$ difference in SCM productivity due to intraseasonal oscillations was representative of northern and perhaps central BoB variability during the southwest monsoon,

confirmed by the BGC-Argo observations from central BoB (Fig. 6). From the surface PAR fluctuations, we expect weaker but still prevalent oscillations in productivity to occur in more southern waters.

Observations from the southern BoB have highlighted the importance of the barrier layer in supporting SCM productivity by impeding the mixing of the SCM into the mixed layer (Prasanth et al., 2023). Our observations deviated from the southern BoB observation but still confirmed the importance of the barrier layer. The formation of a shallow rain-layer rapidly shoaled the mixed layer, increasing the barrier layer thickness, and isolating the SCM from being eroded by upper ocean mixing (Fig. 5). Had the sunny break conditions not followed after a storm event, we expect observed intraseasonal oscillations in ChlF_{SCM} and GCP_{SCM} would have decreased. At M3, the erosion of the barrier layer co-occurred with minimum ChlF_{SCM} rather than when ChlF_{SCM} was enhanced, like observations by Prasanth et al. (2023). The erosion of the barrier hence had minimal impact on the already decreased SCM productivity, but likely contributed to the observed decrease in ChlF_{SCM} . We also note that at M3, the bulk of the SCM productivity was below the barrier layer, while Prasanth et al. (2023) observed enhanced ChlF within the barrier layer when it was present. As the SCM is typically located at the deepest depth where PAR is sufficient for growth and the shallowest depth where nutrients are available, our observations may suggest a relatively deeper nutricline than in the southern BoB. The barrier layer still separated the SCM from upper ocean mixing, and hence will influence SCM productivity, as found in the southern BoB (Prasanth et al., 2023).

Both the DBASIS and BGC-Argo observations indicate the intraseasonal variability in surface light project onto the SCM primary productivity. These observations suggest propagating coupled ocean-atmosphere intraseasonal weather patterns may leave a biological footprint of time- and space-variable primary productivity within the Bay of Bengal. Barrier layer formation is additionally impacted by intraseasonal oscillations (Thadathil et al., 2007; Girishkumar et al., 2011; Prasanth et al., 2023), and will impact ChlF vertical distributions (Prasanth et al., 2023) and hence SCM productivity. Other biogeochemical properties may also be dependent on intraseasonal weather patterns, since primary productivity modulates dissolved oxygen, particulate organic carbon, and dissolved inorganic carbon concentrations (Sarma et al., 2012). Similar patterns could also be evident in other tropical and subtropical oceans subject to intraseasonal atmospheric variability.

4.2 Higher trophic levels

Although we focus on the light limitation on primary productivity in this manuscript, grazing by higher trophic levels can also constrain phytoplankton variability. To provide some information on this component of the food web, we used acoustic backscatter measurements from 1 MHz Acoustic Doppler Current Profilers onboard the Wirewalkers (Fig. 8, Ohman et al., 2019). The backscatter measurements showed enhanced values at night in shallow waters with depths less than 50 m and density of 1022 kg m^{-3} or less (Fig. 8). Then during the day, backscatter measurements were enhanced in deeper waters at around 60 m depth. These observations are consistent with a shallow diel migration of zooplankton between around 60 m at daytime to near the SCM at nighttime. Although we do not have observations to confirm the size and species present, as highlighted in the ‘Introduction’, previous analysis with a 1 MHz acoustic backscatter confirms its sensitivity to smaller zooplankton species, including grazers (Lavery et al., 2007; Ohman et al., 2019; Gastauer et al., 2022). These smaller species

tend to have slower and shallower vertical migrations than larger species (Gastauer et al., 2022), so the shallow diel migration in Fig. 8b is consistent with a small zooplankton species.

The acoustic backscatter strength near the SCM increased over time and reached a maximum on July 24 (Fig. 8b). During this period of increasing acoustic backscatter, the rate of ChlF_{SCM} loss began to exceed that of GCP_{SCM} (Fig. 3d), leading to a decrease in ChlF_{SCM} from July 21. This suggests that grazing could have contributed to the increased loss rate and might provide a link between intraseasonal variability in primary production and higher trophic levels. Prasanth et al. (2023) highlighted that a better understanding of barrier layer formation under variable phytoplankton growth rates and zooplankton grazing is required. Here, we reveal important linkages between these processes and additionally highlight the importance of the intraseasonal oscillations in imparting periodicity to phytoplankton growth at the SCM.

5 Conclusions

The coupled bio-optical and physical observations collected from the DBASIS drifting buoy-profiler systems allowed concurrent measurements of the subsurface irradiance and the diel cycles of subsurface chlorophyll fluorescence during the southwest monsoon (Fig. 3a). These observations showed that changes in subsurface irradiance in the northern Bay of Bengal were coupled with intraseasonal variability in monsoon weather and spatial variability in the turbidity of surface waters, and led to major changes in subsurface primary productivity.

Coupled ocean-atmosphere global forecasting models are challenged to accurately represent intraseasonal variability in the tropical and subtropical ocean. Here we have shown that intraseasonal variability extends to the subsurface biogeochemical properties of the Northern Indian Ocean. Representing this variability correctly in climate predictions is necessary to reduce forecast uncertainty since it impacts both the ocean carbon system and the optical characteristics of the upper ocean, which modulate ocean heat content. Continued advances in autonomous measurement techniques, and widespread deployment of those platforms, is necessary to improve our understanding of the contribution of intraseasonal biogeochemical variability to the ocean ecosystem.

Code and data availability. Data underlying the presented figures and tables are freely available. Biogeochemical-Argo data are freely available through one of the two Global Data Assembly Centers (GDAC), using the WMO number of the float, which is its specific identifier. Argo data were collected and made freely available by the International Argo Program and the national programs that contribute to it. (<http://www.argo.ucsd.edu>, <http://argo.jcommops.org>). The Argo Program is part of the Global Ocean Observing System.

Ocean color satellite measurements are freely available through the NASA Goddard Space Flight Center, Ocean Ecology Laboratory, Ocean Biology Processing Group (2018). doi: 10.5067/AQUA/MODIS/L3M/KD490/2018, 10.5067/AQUA/MODIS/L3M/PAR/2018, and 10.5067/AQUA/MODIS/L3M/CHL/2018. Accessed August 2020 to October 2021. The VGPM NPP measurements were provided by the Ocean Productivity team at Oregon State University (<http://sites.science.oregonstate.edu/ocean.productivity/index.php>). The altimeter products were produced and distributed by the EU Copernicus Marine Service Information (doi: 10.48670/moi-00021). The modified diel cycle

420 code will be publicly available as a version update of the currently available code (<https://github.com/duebi/dielFit>). All other code used is
available upon request.

Author contributions. The manuscript contains observations from the Office of Naval Research MISO-BOB 2019 field campaign. AJL and JTF led the conception, planning, and execution of the field experiment and TLS assisted in the execution. TLS conducted the majority of the data analysis and MO contributed the biogeochemical Argo analysis. TLS and AJL drafted the initial manuscript. All authors approved
425 the final manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank the crew, volunteers, and scientists who aided in the field data collection aboard the R/V Sally Ride. We would like to thank Professor Debasis Sengupta of the Indian Institute of Science, Bangalore for his contributions to our understanding of intraseasonal variability in the Northern Indian Ocean. This study has been conducted using E.U. Copernicus Marine Service Information;
430 doi: 10.48670/moi-00021 and OC-CCI data (Sathyendranath et al., 2019, 2023). TLS, AJL, and JTF were supported by the Office of Naval Research “Monsoon Intra-Seasonal Oscillations Bay Of Bengal” (MISO-BOB) program. TLS and AJL were supported by ONR N00014-17-1-2391 and AJL was supported by ONR N00014-17-1-2987. MO was supported by AWD05945 and NSF 2048491.

References

- Characterizing the vertical diffuse attenuation coefficient for downwelling irradiance in coastal waters: Implications for wa-
 435 ter penetration by high resolution satellite data, *ISPRS Journal of Photogrammetry and Remote Sensing*, 60, 48–64,
<https://doi.org/10.1016/j.isprsjprs.2005.09.003>, 2005.
- Antoine, D., André, J.-M., and Morel, A.: Oceanic primary production 2. Estimation at global scale from satellite (coastal zone color scanner)
 chlorophyll, *Global Biogeochemical Cycles*, 10, 57–60, <https://doi.org/10.1029/2003GL016889>, 1996.
- Argo: Argo float data and metadata from Global Data Assembly Centre (Argo GDAC), SEANOE, <https://doi.org/10.17882/42182>, accessed
 440 on 09/11/2020, 2000.
- Barone, B., Nicholson, D., Ferrón, S., Firing, E., and Karl, D.: The estimation of gross oxygen production and community respira-
 tion from autonomous time-series measurements in the oligotrophic ocean, *Limnology and Oceanography: Methods*, 17, 650–664,
<https://doi.org/10.1002/lom3.10340>, 2019.
- Britton, C. and Dodd, J.: Relationships of photosynthetically active radiation and shortwave irradiance, *Agricultural Meteorology*, 17, 1–7,
 445 [https://doi.org/10.1016/0002-1571\(76\)90080-7](https://doi.org/10.1016/0002-1571(76)90080-7), 1976.
- Cornec, M., Claustre, H., Mignot, A., Guidi, L., Lacour, L., Poteau, A., D’Ortenzio, F., Gentili, B., and Schmechtig, C.: Deep
 Chlorophyll Maxima in the Global Ocean: Occurrences, Drivers and Characteristics, *Global Biogeochemical Cycles*, 35, 1–30,
<https://doi.org/10.1029/2020GB006759>, 2021.
- Cullen, J. J.: The Deep Chlorophyll Maximum: Comparing Vertical Profiles of Chlorophyll *<i>a</i>*, *Canadian Journal of Fisheries and*
 450 *Aquatic Sciences*, 39, 791–803, <https://doi.org/10.1139/f82-108>, 1982.
- Cullen, J. J.: Subsurface chlorophyll maximum layers: Enduring enigma or mystery solved?, *Annual Review of Marine Science*, 7, 207–239,
<https://doi.org/10.1146/annurev-marine-010213-135111>, 2015.
- Davis, R. E., Ohman, M. D., Rudnick, D. L., Sherman, J. T., and Hodges, B.: Glider surveillance of physics and biology in the southern
 California Current System, *Limnology and Oceanography*, 53, 2151–2168, https://doi.org/10.4319/lo.2008.53.5_part_2.2151, 2008.
- 455 Frenzel, H., Sharp, J., Fassbender, A., and Buzby, N.: OneArgo-Mat: A MATLAB toolbox for accessing and visualizing Argo data., Zenodo,
<https://doi.org/10.5281/zenodo.6588042>, 2020.
- Gadgil, S., Rajeevan, M., and Nanjundiah, R.: Monsoon prediction - Why yet another failure?, *Current Science*, 88, 1389–1400, 2005.
- Gastauer, S., Nickels, C. F., and Ohman, M. D.: Body size- and season-dependent diel vertical migration of mesozooplankton resolved
 acoustically in the San Diego Trough, *Limnology and Oceanography*, 67, 300–313, <https://doi.org/10.1002/lno.11993>, 2022.
- 460 Girishkumar, M. S., Ravichandran, M., McPhaden, M. J., and Rao, R. R.: Intraseasonal variability in barrier layer thickness in the south
 central Bay of Bengal, *Journal of Geophysical Research: Oceans*, 116, <https://doi.org/10.1029/2010JC006657>, 2011.
- Goddard Space Flight Center, Ocean Ecology Laboratory, O. B. P. G.: Moderate-resolution Imaging Spectroradiometer (MODIS)
 Aqua Downwelling Diffuse Attenuation Coefficient Data; 2018 Reprocessing, NASA OB.DAAC, Greenbelt, MD, USA,
<https://doi.org/10.5067/AQUA/MODIS/L3M/KD/2018>, accessed on 10/20/2021, 2018a.
- 465 Gray, A. R., Johnson, K. S., Bushinsky, S. M., Riser, S. C., Russell, J. L., Talley, L. D., Wanninkhof, R., Williams, N. L., and Sarmiento, J. L.:
 Autonomous Biogeochemical Floats Detect Significant Carbon Dioxide Outgassing in the High-Latitude Southern Ocean, *Geophysical*
Research Letters, 45, 9049–9057, <https://doi.org/10.1029/2018GL078013>, 2018.
- Hansell, D. A.: Dissolved organic carbon in the carbon cycle of the Indian Ocean, *Geophysical Monograph Series*, 185, 217–230,
<https://doi.org/10.1029/2007GM000684>, 2009.

- 470 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Sabater, J. M., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1940 to present, <https://doi.org/10.24381/cds.adbb2d47>, 2018.
- Johnson, K. S. and Bif, M. B.: Constraint on net primary productivity of the global ocean by Argo oxygen measurements, *Nature Geoscience*, 14, 769–774, <https://doi.org/10.1038/s41561-021-00807-z>, 2021.
- 475 Johnson, K. S., Plant, J. N., Coletti, L. J., Jannasch, H. W., Sakamoto, C. M., Riser, S. C., Swift, D. D., Williams, N. L., Boss, E., Haëntjens, N., Talley, L. D., and Sarmiento, J. L.: Biogeochemical sensor performance in the SOCCOM profiling float array, *Journal of Geophysical Research: Oceans*, 122, 6416–6436, <https://doi.org/10.1002/2017JC012838>, 2017.
- Kerhalkar, S., Tandon, A., Farrar, J., MacKinnon, J. A., Schlosser, T. L., Johnson, L., Lucas, A. J., Hormann, V., and Centurioni, L. R.: Modulation of Diurnal SST and Diurnal Warm Layer Variability by Salinity-Driven Stratification in the Bay of Bengal, *Journal of Physical*
 480 *Oceanography*, <https://doi.org/10.1175/jpo-d-25-0134.1>, 2025.
- Kumar, S. P., Narvekar, J., Nuncio, M., Kumar, A., Ramaiah, N., Sardesai, S., Gauns, M., Fernandes, V., and Paul, J.: Is the biological productivity in the Bay of Bengal light limited?, *Current Science*, 98, 1331–1339, 2010a.
- Kumar, S. P., Nuncio, M., Narvekar, J., Ramaiah, N., Sardesai, S., Gauns, M., Fernandes, V., Paul, J. T., Jyothibabu, R., and Jayaraj, K. A.: Seasonal cycle of physical forcing and biological response in the Bay of Bengal, *Indian Journal of Marine Sciences*, 39, 388–405, 2010b.
- 485 Lavery, A. C., Wiebe, P. H., Stanton, T. K., Lawson, G. L., Benfield, M. C., and Copley, N.: Determining dominant scatterers of sound in mixed zooplankton populations, *The Journal of the Acoustical Society of America*, 122, 3304–3326, <https://doi.org/10.1121/1.2793613>, 2007.
- Lucas, A., Nash, J., Pinkel, R., MacKinnon, J., Tandon, A., Mahadevan, A., Omand, M., Freilich, M., Sengupta, D., Ravichandran, M., and Boyer, A. L.: Adrift Upon a Salinity-Stratified Sea: A View of Upper-Ocean Processes in the Bay of Bengal During the Southwest
 490 Monsoon, *Oceanography*, 29, 134–145, <https://doi.org/10.5670/oceanog.2016.46>, 2016.
- Malan, N., Archer, M., Roughan, M., Cetina-Heredia, P., Hemming, M., Rocha, C., Schaeffer, A., Suthers, I., and Queiroz, E.: Eddy-Driven Cross-Shelf Transport in the East Australian Current Separation Zone, *Journal of Geophysical Research: Oceans*, 125, 1–15, <https://doi.org/10.1029/2019JC015613>, 2020.
- Mandke, S. K., Pillai, P. A., and Sahai, A. K.: Simulation of monsoon intra-seasonal oscillations in Geophysical Fluid Dynamics Laboratory
 495 models from Atmospheric Model Intercomparison Project integrations of Coupled Model Intercomparison Project phase 5, *International Journal of Climatology*, 40, 5574–5589, <https://doi.org/10.1002/joc.6536>, 2020.
- Marra, J.: Analysis of diel variability in chlorophyll fluorescence, *Journal of Marine Research*, 55, 767–784, <https://doi.org/10.1357/0022240973224274>, 1997.
- Mignot, A., Claustre, H., Uitz, J., Poteau, A., D’Ortenzio, F., and Xing, X.: Understanding the seasonal dynamics of phytoplankton biomass
 500 and the deep chlorophyll maximum in oligotrophic environments: A Bio-Argo float investigation, *Global Biogeochemical Cycles*, 28, 856–876, <https://doi.org/10.1002/2013GB004781>, 2014.
- Morel, A., Huot, Y., Gentili, B., Werdell, P. J., Hooker, S. B., and Franz, B. A.: Examining the consistency of products derived from various ocean color sensors in open ocean (Case 1) waters in the perspective of a multi-sensor approach, *Remote Sensing of Environment*, 111, 69–88, <https://doi.org/10.1016/j.rse.2007.03.012>, 2007.
- 505 Müller, P., Li, X. P., and Niyogi, K. K.: Non-photochemical quenching. A response to excess light energy, *Plant Physiology*, 125, 1558–1566, <https://doi.org/10.1104/pp.125.4.1558>, 2001.

- NASA Goddard Space Flight Center, Ocean Ecology Laboratory, O. B. P. G.: Moderate-resolution Imaging Spectroradiometer (MODIS) Aqua Photosynthetically Available Radiation Data; 2018 Reprocessing, NASA OB.DAAC, Greenbelt, MD, USA, <https://doi.org/10.5067/AQUA/MODIS/L3M/PAR/2018>, accessed on 10/20/2021, 2018b.
- 510 Nicholson, D. P., Wilson, S. T., Doney, S. C., and Karl, D. M.: Quantifying subtropical North Pacific gyre mixed layer primary productivity from Seaglider observations of diel oxygen cycles, *Geophysical Research Letters*, 42, 4032–4039, <https://doi.org/10.1002/2015GL063065>, 2015.
- Ohman, M. D., Davis, R. E., Sherman, J. T., Grindley, K. R., Whitmore, B. M., Nickels, C. F., and Ellen, J. S.: Zooglider: An autonomous vehicle for optical and acoustic sensing of zooplankton, *Limnology and Oceanography: Methods*, 17, 69–86, <https://doi.org/10.1002/lom3.10301>, 2019.
- 515 Pinkel, R., Goldin, M. A., Smith, J. A., Sun, O. M., Aja, A. A., Bui, M. N., and Huguen, T.: The Wirewalker: A vertically profiling instrument carrier powered by ocean waves, *Journal of Atmospheric and Oceanic Technology*, 28, 426–435, <https://doi.org/10.1175/2010JTECHO805.1>, 2011.
- Prasanth, R., Vijith, V., and Vinayachandran, P. N.: Formation, maintenance and diurnal variability of subsurface chlorophyll maximum during the summer monsoon in the southern Bay of Bengal, *Progress in Oceanography*, 212, 102974, <https://doi.org/10.1016/j.pocean.2023.102974>, 2023.
- 520 Sarma, V. V., Krishna, M. S., Rao, V. D., Viswanadham, R., Kumar, N. A., Kumari, T. R., Gawade, L., Ghatkar, S., and Tari, A.: Sources and sinks of CO₂ in the west coast of Bay of Bengal, *Tellus, Series B: Chemical and Physical Meteorology*, 64, 10961, <https://doi.org/10.3402/tellusb.v64i0.10961>, primary productivity influences distribution of both pCO₂ and DO, 2012.
- 525 Sathyendranath, S., Brewin, R., Brockmann, C., Brotas, V., Calton, B., Chuprin, A., Cipollini, P., Couto, A., Dingle, J., Doerffer, R., Donlon, C., Dowell, M., Farman, A., Grant, M., Groom, S., Horseman, A., Jackson, T., Krasemann, H., Lavender, S., Martinez-Vicente, V., Mazeran, C., Mélin, F., Moore, T., Müller, D., Regner, P., Roy, S., Steele, C., Steinmetz, F., Swinton, J., Taberner, M., Thompson, A., Valente, A., Zühlke, M., Brando, V., Feng, H., Feldman, G., Franz, B., Frouin, R., Gould, R., Hooker, S., Kahru, M., Kratzer, S., Mitchell, B., Muller-Karger, F., Sosik, H., Voss, K., Werdell, J., and Platt, T.: An Ocean-Colour Time Series for Use in Climate Studies: The Experience of the Ocean-Colour Climate Change Initiative (OC-CCI), *Sensors*, 19, 4285, <https://doi.org/10.3390/s19194285>, 2019.
- 530 Sathyendranath, S., Jackson, T., Brockmann, C., Brotas, V., Calton, B., Chuprin, A., Clements, O., Cipollini, P., Danne, O., Dingle, J., Donlon, C., Donlon, C., Groom, S., Krasemann, H., Lavender, S., Mazeran, C., Mélin, F., Müller, D., Steinmetz, F., Valente, A., Zühlke, M., Feldman, G., Franz, B., Frouin, R., Werdell, J., and Platt, T.: ESA Ocean Colour Climate Change Initiative (Ocean-Colour-cci): Global attenuation coefficient for downwelling irradiance (Kd490) gridded on a geographic projection at 4km resolution, Version 6.0, <https://catalogue.ceda.ac.uk/uuid/8ecae26f390b4938b67a97cbce3ecd8b/>, 2023.
- 535 Schlosser, T. L., Lucas, A. J., Jones, N. L., Nash, J. D., and Ivey, G. N.: Local winds and encroaching currents drive summertime subsurface blooms over a narrow shelf, *Limnology and Oceanography*, pp. 1–15, <https://doi.org/10.1002/lno.12043>, 2022.
- Schmechtig, C., Poteau, A., Claustre, H., D’Ortenzio, F., Dall’Olmo, G., and Boss, E.: Processing BGC-Argo particle backscattering at the DAC level. Version 1.4, 07 March 2018., Tech. rep., IFREMER for Argo Data Management, <https://doi.org/10.13155/39459>, 2018.
- 540 Shroyer, E., Tandon, A., Sengupta, D., Fernando, H. J., Lucas, A. J., Farrar, J. T., Chattopadhyay, R., de Szoeka, S., Flatau, M., Rydbeck, A., Wijesekera, H., McPhaden, M., Seo, H., Subramanian, A., Venkatesan, R., Joseph, J., Ramsundaram, S., Gordon, A. L., Bohman, S. M., Pérez, J., Simoes-Sousa, I. T., Jayne, S. R., Todd, R. E., Bhat, G., Lankhorst, M., Schlosser, T., Adams, K., Jinadasa, S., Mathur, M., Mohapatra, M., Rao, E. P. R., Sahai, A. K., Sharma, R., Lee, C., Rainville, L., Cherian, D., Cullen, K., Centurioni, L. R., Hormann, V., MacKinnon, J., Send, U., Anutaliya, A., Waterhouse, A., Black, G. S., Dehart, J. A., Woods, K. M., Creegan, E., Levy, G., Kantha, L. H.,

- 545 and Subrahmanyam, B.: Bay of Bengal Intraseasonal Oscillations and the 2018 Monsoon Onset, *Bulletin of the American Meteorological Society*, pp. 1–44, <https://doi.org/10.1175/bams-d-20-0113.1>, 2021.
- Thadathil, P., Muraleedharan, P. M., Rao, R. R., Somayajulu, Y. K., Reddy, G. V., and Revichandran, C.: Observed seasonal variability of barrier layer in the Bay of Bengal, *Journal of Geophysical Research: Oceans*, 112, <https://doi.org/10.1029/2006JC003651>, 2007.
- Todd, R. E., Rudnick, D. L., and Davis, R. E.: Monitoring the greater San Pedro Bay region using autonomous underwater gliders during fall
550 of 2006, *Journal of Geophysical Research: Oceans*, 114, 1–13, <https://doi.org/10.1029/2008JC005086>, 2009.
- Uchida, T., Balwada, D., Abernathey, R., Prend, C. J., Boss, E., and Gille, S. T.: Southern Ocean Phytoplankton Blooms Observed by Biogeochemical Floats, *Journal of Geophysical Research: Oceans*, 124, 7328–7343, <https://doi.org/10.1029/2019JC015355>, 2019.
- Weller, R., Farrar, J. T., Buckley, J., Matthew, S., Venkatesan, R., Lekha, J. S., Chaudhuri, D., Kumar, N. S., and Kuman, B. P.: Air-Sea Interaction in the Bay of Bengal, *Oceanography*, 29, 28–37, <https://doi.org/10.5670/oceanog.2016.36>, 2016.
- 555 White, A. E., Barone, B., Letelier, R. M., and Karl, D. M.: Productivity diagnosed from the diel cycle of particulate carbon in the North Pacific Subtropical Gyre, *Geophysical Research Letters*, 44, 3752–3760, <https://doi.org/10.1002/2016GL071607>, 2017.
- Wong, A., Keeley, R., and Carval, T.: Argo Quality Control Manual for CTD and Trajectory Data, Tech. rep., USA, France, <https://doi.org/10.13155/33951>, 2022.
- Zheng, B., Lucas, A. J., Pinkel, R., and Boyer, A. L.: Fine-Scale Velocity Measurement on the Wirewalker Wave-Powered Profiler, *Journal*
560 *of Atmospheric and Oceanic Technology*, 39, 133–147, <https://doi.org/10.1175/JTECH-D-21-0048.1>, 2022.




Figure 1. Overview of observations. a) Bay of Bengal with the study region (red square), location of analyzed biogeochemical (BGC) Argo profilers deployed from 2013 to 2018 (yellow, WMO ID: 2902086, 2902087, 2902114, 2902158, 2902160, 2902161, 2902189, 2902196, 2902217, 2902264), with a further analyzed 2016 profile highlighted (pink, WMO ID: 2902193; Argo, 2000), location of DBASIS profilers (purple), 5-day averaged (10-15 July, 2019) satellite chlorophyll (Chl, background color; Sathyendranath et al., 2019, 2023), and gray contours of the EU Copernicus Marine Service global ocean $1/12^\circ$ sea surface height (SSH, 4th July, 2019; <https://doi.org/10.48670/moi-00021>). b) Position and date for the three high-resolution DBASIS floats: M1 (dark blue), M2 (light blue), and M3 (green), with the depth-maximum chlorophyll-*a* fluorescence (ChlF) shown for M3 at noon each day. Gray contours show SSH (m). DBASIS floats were located in depths ~ 2600 m deep and ~ 400 km southeast of the nearest coastline.




Figure 2. Observations at M3. a) the net heat flux (Q_N), b) wind speed (blue left y-axis, W_s) and precipitation rate (orange right y-axis, P), c) Photosynthetically available radiation (PAR), d) ChlF, and e) turbidity (τ). In c) to e), two isopycnals (1021.5 kg m^{-3} and 1023 kg m^{-3}) are contoured (black), and in d) the depth of the isothermal layer (ITL, pink) and mixed layer depth (MLD, yellow) are shown.




Figure 3. Co-varying light and chlorophyll. At M3, a) on isopycnal ChlF, with depth representing the 48-h low-pass filtered isopycnal depth (white contour, right y-axis) and the SCM depth labeled (black contour). b) PAR at the surface (purple, PAR_0) and subsurface chlorophyll maximum (SCM, teal, PAR_{SCM}) multiplied by 10. c) ChlF_{SCM} (purple) and the diel cycle fit (black, see ‘Methods’). d) Rate of gross ChlF production (GCP_{SCM} , green) and loss (grey) at the SCM, with the 95% confidence interval labeled (error bars). e) Similar to panel b) but daily averaged (denoted by the over-bar). Note 0^+ and 0^- m indicate surface measurements in air and water, respectively, and we denote a conversion from shortwave radiation to PAR with $(\cdot)^*$.




Figure 4. Light-limited productivity. Daily averaged PAR_0 (y-axis, converted from shortwave radiation) vs. a) GCP_{SCM} and b) gross turbidity production (GTP_{SCM}) for all profilers and sample days with a good diel cycle fit ($r^2 \geq 0.4$), colored corresponding to the inverse diffuse attenuation (K_d^{-1}). b,d) Similar to the above panel, but with PAR_{SCM} . We least-squares linear fit each panels gross production and light variable, labeling the resulting linear regression (black), r^2 , p -value, and mean absolute error (MAE).




Figure 5. Barrier layer vs. the SCM. At M3, a,d) temperature, b,e) salinity, and c,f) ChlF, over the upper 70 m (top panels) and at specific depths (bottom panels). a-c) Depth of the isothermal layer (ITL, pink), SCM (black), and mixed layer depth (MLD, yellow). d-f) Variability at near-surface (black), and mixed layer depth (purple). f) Variability at the SCM (green).




Figure 6. Co-varying light and ChlF from a 2016 BGC-Argo deployment (WMO ID: 2902193, Figure 1a). ChlF concentrations a) depth-averaged over the 20 m centered on the 1022 kg m^{-3} isopycnal we designate as the SCM (green, left y-axis) and b) contoured over depth. Also on panel a), for each BGC-Argo profile we find the closest ERA5 surface PAR (purple, right y-axis; Hersbach et al., 2018) and MODIS-Aqua 490 nm diffuse attenuation (Goddard Space Flight Center, 2018a), converted to a PAR equivalent (Morel et al., 2007), to estimate PAR_{SCM} (black, right y-axis). Yellow shading in panel a) indicates the southwest monsoon period. In panel b), isopycnals are contoured in gray every 1 kg m^{-3} , with the SCM plotted in blue.




Figure 7. ERA5 derived monthly climatology (2012-2022) of surface PAR (PAR_0). ERA5 PAR_0 at fixed locations around the bay (color), a) monthly averaged and b) monthly standard deviation. Gray shading indicates the southwest summer monsoon period.




Figure 8. Diel vertical migration to the SCM. a) PAR_0 (converted from shortwave radiation), and b) on isopycnal acoustic backscatter at 1 MHz (depth as Figure 3a).

Table 1. The performance of the diel cycle fit averaged over all three profilers and days fitted. We tested three methods of estimating the gross chlorophyll production (GCP) along an isopycnal (1021.5 kg m^{-3}) estimated as the subsurface chlorophyll maximum (SCM): linear, sinusoidal and from the observed PAR_{SCM} . We also applied this method at a fixed depth corresponding to the average depth of the SCM at each profiler, labeled $\overline{\text{PAR}_{\text{SCM}}}$ where $(\bar{\cdot})$ indicates an average in time. We show the estimated correlation coefficient (r^2) and mean absolute error (MAE) from fitting ChlF, and we estimate the MAE in the timing of maximum ChlF (MAE_t).

Method	r^2 (p -value)	MAE ($\mu\text{g L}^{-1}$)	MAE_t (h)
Linear	0.69 (3.7×10^{-4})	0.042	3.5
Sinusoidal	0.78 (6.7×10^{-4})	0.032	2.1
PAR_{SCM}	0.81 (6.7×10^{-6})	0.029	1.4
$\overline{\text{PAR}_{\text{SCM}}}$	0.49 (1.0×10^{-2})	0.060	2.2