



Spatially contrasted response of Devonian anoxia to astronomical forcing

Justin Gérard¹, Alexandre Pohl², Loïc Sablon¹, Jarno J.C. Huygh³, Anne-Christine Da Silva³, and Michel Crucifix¹

¹Université catholique de Louvain (UCLouvain), Earth and Life Institute (ELI), 1348 Louvain-la-Neuve, Belgium

²Université Bourgogne Europe, CNRS, Biogéosciences UMR 6282, 21000 Dijon, France

³University of Liège, SediCClim Department of Geology, Sart Tilman, 4000 Liège, Belgium

Correspondence: Justin Gérard (justin.gerard@uclouvain.be)

Abstract. The Devonian period, spanning from 419 to 359 million years ago, was marked by a warmer-than-present climate and recurring ocean anoxic events, with evidence increasingly suggesting a link between these events and astronomical forcing. Here, we explore how astronomical forcing influences ocean oxygenation by modulating the continental weathering flux of phosphate within a Late Devonian climate framework. To investigate this, we performed transient simulations spanning 1.1 Myr, crossing a 2.4 Myr eccentricity node using the cGENIE Earth system model. These simulations were driven by spatially resolved fluxes of reactive phosphorus from continents, computed using the emulator developed by Sablon et al. (2025), trained on GEOCLIM and HadSM3 outputs. Our results provide new evidence supporting eccentricity maxima as a driver of Late Devonian anoxic events. Additionally, global analysis reveals that obliquity variations can imprint a distinct signal on global ocean oxygen levels via their influence on biological productivity, offering a plausible mechanism for obliquity-driven anoxia under greenhouse conditions. Regional analysis revealed pronounced spatial heterogeneity in the biogeochemical response to astronomical forcing. Local ocean circulation emerged as a critical factor in shaping these patterns. The simulations indicate that astronomical forcing can, through its impact on continental weathering fluxes, exert a dominant influence on ocean oxygenation, with regional oxygen concentrations varying by up to 35% and driving changes in regional anoxic volume of up to 19%. Finally, these findings help explain why proxy records from different locations may show divergent expressions of astronomical signals, potentially leading to contrasting interpretations of their role in driving ocean anoxia.

1 Introduction

Throughout the Devonian Period, multiple oceanic anoxic events (OAEs) occurred (Becker et al., 2020), with estimated durations ranging from 10^4 to 10^6 years (Kabanov et al., 2023). The Late Devonian, approximately 379–360 million years ago (Ma), was particularly marked by pronounced environmental crises and ocean anoxia, such as the well-known Kellwasser and Hangenberg events (Kaiser et al., 2016; Carmichael et al., 2019). There is still no consensus on the mechanisms responsible for the initiation and the duration of these Devonian OAEs (Haddad et al., 2018; Reershemius and Planavsky, 2021; Smart et al., 2023). Existing hypotheses include extensive volcanic activity (Ma et al., 2016; Paschall et al., 2019; Racki, 2020; Kabanov et al., 2023), the proliferation and expansion of terrestrial plants (Algeo and Scheckler, 1998; De Vleeschouwer et al., 2017;



Smart et al., 2023), climate cooling (Joachimski and Buggisch, 2002; Song et al., 2017; Huang et al., 2018; Pier et al., 2021), shifts in sea level and oceanic circulation dynamics (Wilde and Berry, 1984; Dopieralska, 2009; Chen et al., 2013), and changes in the position of the continents (Brugger et al., 2019; Gérard et al., 2025). There is also growing evidence that astronomical forcing could modulate at least some of these events (De Vleeschouwer et al., 2014; Wichern et al., 2024; Huygh et al., 2025, under revision). Specifically, studies conducted in Poland by De Vleeschouwer et al. (2013) identified that the deposition of anoxic black shales associated with the *Annulata*, Dasberg, and Hangenberg events is separated by approximately 2.4 million years (Myr). This suggests a role of eccentricity in the timing of these events, as the variance spectrum of eccentricity often contains such a cycle (Laskar et al., 2004). However, substantial uncertainties persist regarding the phase relationship of the astronomical forcing to the onset and development of Devonian OAEs. For instance, some cyclostratigraphic interpretations have reported the occurrence of the upper Kellwasser anoxic event, after a 2.4 Myr eccentricity minima (De Vleeschouwer et al., 2017; Da Silva et al., 2020; Wichern et al., 2024), while other studies reported strong obliquity and eccentricity, resulting in marine eutrophication leading to the upper Kellwasser Ma et al. (2022) and Huygh et al., (2025, under revision). Hence, these contrasting findings underscore the growing importance of better quantifying the contribution of astronomical forcing to Devonian oceanic oxygen levels and the pathways by which astronomical forcing may have modulated ocean oxygenation at that time.

Our objective is to assess the impact of astronomical forcing on ocean anoxia, specifically through the modulation of continental nutrient fluxes, within a Late Devonian climate context. To this end, we performed transient simulations using the Earth system model of intermediate complexity cGENIE (Ridgwell et al., 2007), coupled offline with a continental weathering flux of nutrients. This flux is computed using an emulator trained on outputs from the fully coupled general circulation model (GCM) HadSM3 (Valdes et al., 2017) and the geochemical model GEOCLIM (Maffre et al., 2024), as developed in Sablon et al. (2025). By incorporating a GCM, we account for the changes in precipitation and evaporation caused by astronomical forcing more accurately than with the standard version of cGENIE. In addition to improving the representation of hydrological responses, this approach further enhances the simulation of nutrient fluxes to the ocean, limited in the standard cGENIE model version due to its non-dimensional treatment of continental weathering (Colbourn et al., 2013). This modelling framework allows, on the one hand, for a mechanistic assessment of how variations in orbital parameters (namely eccentricity, precession, and obliquity) affect terrestrial weathering rates and nutrient fluxes to the ocean, thereby influencing the development and spatial extent of anoxic conditions. On the other hand, the influence of astronomical forcing on oxygen levels through changes in ocean circulation is captured by cGENIE. A detailed description of the model configuration and experimental design is presented in Section 2. In Section 3.1, we examine the global characteristics of the transient coupled simulations, followed by a regional analysis in Section 3.2.



2 Methods

2.1 Models description

cGENIE is an Earth system model of intermediate complexity (Claussen et al., 2002) designed to simulate long-term interactions between climate, biogeochemical cycles, and ocean–atmosphere processes (Ridgwell et al., 2007). The ocean circulation component (3D) uses a frictional-geostrophic model to simulate the transport of heat, salinity, and biogeochemical tracers through advection, convection, and parametrised mixing. The sea ice module (2D) uses a simplified thermodynamic–dynamic scheme to compute ice thickness and fractional coverage based on thermal balances and surface flux exchanges. The atmospheric module (2D) is an energy–moisture balance layer that transports heat and humidity through advection and diffusion, with precipitation triggered by a humidity threshold. These three components form the C-GOLDSTEIN framework, fully described in Edwards and Marsh (2005) and Marsh et al. (2011). The BIOGEM module simulates ocean biogeochemistry, including nutrient-limited biological productivity and tracer cycling, and handles air–sea gas exchange processes. The ATCHEM module tracks atmospheric chemical species and computes their partial pressures to represent changes in atmospheric composition (Ridgwell et al., 2007). For this study, we used an open-system configuration version of cGENIE with regard to carbon and phosphorus (P), which includes the SEDGEM and ROKGEM modules. SEDGEM provides a representation of sedimentary stratigraphy and the preservation of biogenic carbonates and organic carbon (C_{org}) deposited on the seafloor (Ridgwell and Hargreaves, 2007; Hülse and Ridgwell, 2024). ROKGEM is a spatially explicit weathering module that simulates temperature- and runoff-dependent weathering rates, allowing dynamic feedbacks between climate, weathering, and the carbon cycle over multimillion-year timescales (Colbourn et al., 2013).

In the standard configuration of cGENIE, continental weathering is simply a function of the global mean continental temperature. This is inadequate to capture the effects of the astronomical forcing, as the latter primarily affects climate through changes in the spatial and seasonal distribution of incoming solar radiation (insolation). To address this, we replaced this simplified scheme with a state-of-the-art continental weathering module, which accounts for the coupled contributions of chemical weathering and erosion. Specifically, we used the framework described in Sablon et al. (2025) to couple cGENIE with a climate–weathering emulator. This approach computes weathering maps for each steady-state climate condition simulated by the GCM HadSM3 (Valdes et al., 2017) under varying boundary conditions (orbital parameters and pCO_2 levels), assuming a regolith dynamic at equilibrium. We then build an emulator of these weathering maps using the same statistical methods developed for the runoff variable in Sablon et al. (2025), effectively bypassing GEOCLIM during the coupling procedure. The resulting weathering fluxes are instantaneously routed from continental source cells to the closest ocean coastal grid cell via a nearest-neighbour scheme, which remains suitable given the similar grid layouts, both adapted from Scotese and Wright (2018), and the limited knowledge of the Late Devonian hydrological network. A detailed overview of GEOCLIM’s continental weathering module is presented in Maffre et al. (2024), while a complete description of the emulator’s architecture and training process is documented in Sablon et al. (2025).

The coupling between the emulator and cGENIE is deliberately minimal and targeted, replacing the continental nutrient flux of dissolved phosphate (PO_4) provided by cGENIE with values computed by the emulator. This replacement is performed



offline every 1000 years, a frequency that is appropriate for capturing the long-term variability caused by the astronomical forcing. All other components and processes within the model remain unchanged, as our cGENIE setup uses a simplified, single-nutrient (PO_4) control on biological productivity (Ridgwell et al., 2007). An increase in surface PO_4 availability stimulates biological productivity, thereby enhancing microbial oxygen consumption through remineralization in the water column. Moreover, limiting the coupling to a single flux ensures that other variables computed by ROKGEM remain unaffected, maintaining consistency and simplicity in the overall setup. As a result, cGENIE now responds to spatially-resolved, astronomically-driven variations in continental PO_4 flux, thereby allowing a more robust investigation of the influence of orbital forcing on ocean anoxia.

In our simulations, the ocean is discretised on a 36×36 horizontal equal-area grid with 16 vertical levels that are logarithmically spaced (from 60 to 560 meters). We use the continental configuration on which the emulator from Sablon et al. (2025) was trained, namely that of Scotese and Wright (2018), which represents the palaeogeography at 370 Ma (see Fig. 1). Because the modelled climate system is older than the mid-Mesozoic advent of open-ocean planktic calcifiers (approximately 200 Ma), the representation of calcium carbonate (CaCO_3) production is adapted accordingly (Ridgwell, 2005). Specifically, CaCO_3 production is restricted to shallow-marine carbonate platforms, spatially constrained based on the reconstructions of Kiessling et al. (2003) (see Fig. 2). The surface albedo and zonally averaged wind stress profiles are prescribed from the HadCM3 simulation (Sablon et al., 2025), conducted with the same pCO_2 as in Valdes et al. (2020) and an orbital configuration characterized by an eccentricity of 0, a longitude of perihelion of 0° , and an obliquity of 23° . These fields are incorporated into cGENIE using the open-source “muffingen” suite v0.9.24 (Ridgwell, 2023), and are held constant throughout the transient simulation. The solar constant is set to 1325 W.m^{-2} based on Gough (1981), less than today because this younger Sun was also weaker. The atmospheric oxygen concentration is set to 80% of its present-day value to reflect the lower levels characteristic of the Late Devonian (Krause et al., 2018; Mills et al., 2023). Ocean and atmosphere parameter values follow the standard cGENIE setup of Cao et al. (2009) and are identical to those in Pohl et al. (2022) and in Gérard et al. (2025). In our setup, the empirical equation proposed by Dunne et al. (2007) represents C_{org} burial in marine sediments, while the empirical parameterization of Wallmann (2010) accounts for redox-dependent regeneration of P from sediments. Following Dunne et al. (2007), C_{org} burial scales with the flux of organic matter reaching the seafloor and is therefore a function of surface ocean productivity. Following Wallmann (2010), a significant portion of P initially bound in organic matter is degraded in surface sediments and released into the ocean as dissolved PO_4 . Furthermore, PO_4 release from sediments is enhanced under low-oxygen conditions in line with observations (Ingall and Jahnke, 1994). A detailed description of these schemes and their implementation in cGENIE is provided in Hülse and Ridgwell (2024).

2.2 Experimental setup

The astronomical forcing for the Devonian is not known precisely, due to the chaotic nature of the Solar System and the subtle, yet cumulative, effects of tidal dissipation on axial precession (Laskar et al., 2011). A series of plausible solutions is now available (Zeebe and Lantink, 2024), but our experiment design was finalised before that publication appeared. We therefore adopted the same hybrid approach as Sablon et al. (2025). It relies, for planetary motion, on the solution by Laskar et al. (2011)



(La11a), from which we extracted the interval between 123.3 Ma and 122.2 Ma. This interval features a resonance between the 2.4 Myr eccentricity period (associated with the $g4-g3$ terms) and the 1.2 Myr obliquity modulation (associated with the $s4-s3$ terms), as in most of the Cenozoic. This resonance is characterised by an increase in obliquity variance when eccentricity reaches a 2.4 Myr node, at which point its variance is lowest and the 100 thousand years (kyr) component of the eccentricity signal disappears. The node occurs approximately midway through the selected interval.

We coupled this solution with the precession model of Sharaf and Boudnikova (1967), following the procedure outlined in Berger and Loutre (1991). This involves first performing a harmonic decomposition of the planetary solution (using R code based on Šidlichovský and Nesvorný, 1997), and then specifying a reference obliquity and general precession rate. These reference values are given by Farhat et al. (2022) for the period around 370 Ma. With this construction, the solution yields obliquity frequencies of 30.555 and 37.335 years, and climatic precession frequencies of 16.470 and 19.386 years, which are considered plausible for the Late Devonian. The R code used to generate these time series is available in Crucifix (2025). We verified that the time series of obliquity and longitude of perihelion are nearly identical to those obtained by integrating, forward in time, the precession model of Laskar and Robutel (1993) with an appropriate Earth–Moon distance and dynamical ellipticity. The latter approach is admittedly more straightforward, but was not integrated into our workflow at the time the experiment was prepared.

The length and choice of this specific astronomical solution reflect a compromise between representativeness and computational feasibility, given that a 1.1 Myr simulation with cGENIE requires two months of continuous computation on our systems. In particular, we cannot know whether the ($g4-g3$) versus ($s4-s3$) resonance was active during the Devonian, but previous cyclostratigraphic interpretations assumed a well-expressed 2.4 Myr eccentricity node and pronounced obliquity cycles when crossing the node (De Vleeschouwer et al., 2017). The chosen solution satisfies these assumptions. Moreover, it conveniently includes within a single sequence a transition from well-expressed 100 kyr eccentricity cycles modulated by a 405 kyr cycle, to the 2.4 Myr node (during which the 100 kyr cycles disappear), and then back to well-expressed 100 kyr eccentricity cycles. For reference, we use the heliocentric convention for expressing the longitude of perihelion, which is therefore 0° when perihelion occurs in September (as in Sablon et al., 2025 and Gérard et al., 2025).

cGENIE reached equilibrium through a two-step spin-up process similar to Ridgwell and Hargreaves (2007), Hülse and Ridgwell (2024) and Vervoort et al. (2024). In the initial spin-up phase (SPIN1), ocean dynamics and biogeochemical cycling equilibrate with fixed atmospheric $p\text{CO}_2$ and $\delta^{13}\text{C}$ under a closed system. Burial of CaCO_3 , C_{org} , and P are balanced by global weathering inputs, and the resulting data are used to determine global equilibrium burial rates. The second spin-up phase (SPIN2) uses an open-system setup, with weathering fluxes set to the equilibrium with burial rates from SPIN1 and allowed to respond to temperature. A complete description of this two-step procedure in cGENIE, including C_{org} and P feedbacks, is detailed in Hülse and Ridgwell (2024). We found that during the SPIN2, the model exhibited significant drift, most notably in $p\text{CO}_2$, with the system requiring up to 500 kyr to reach equilibrium. To address this, we heuristically adjusted the initial $p\text{CO}_2$ concentration in SPIN1 to guide the model toward a geochemically consistent Late Devonian state. By the end of SPIN2, the system reached an atmospheric $p\text{CO}_2$ concentration of approximately 550 ppm, consistent with the estimates of Chen et al. (2021) and Dahl et al. (2022), and an average surface PO_4 concentration of $0.47 \mu\text{mol.kg}^{-1}$, in agreement with Sharoni and



Halevy (2023). Furthermore, the system achieved modern-like burial fluxes (similar to Hülse and Ridgwell, 2024), including $0.144 \text{ PgC.yr}^{-1}$ of CaCO_3 (reef only) (Frankignoulle and Gattuso, 1993), $0.075 \text{ PgC.yr}^{-1}$ of C_{org} (Dunne et al., 2007; Bradley et al., 2022), and $0.062 \text{ Tmol.yr}^{-1}$ of P (Benitez-Nelson, 2000; Ruttenberg, 2003). The cause of this drift remains unclear, but our testing indicates that drift is affected by the continental configuration and model resolution. A full investigation of its origin lies beyond the scope of this study, as it would require substantial computational time, with a 500 kyr simulation requiring roughly one month of runtime.

Starting from the equilibrium state (SPIN2), the continental PO_4 nutrient flux provided by cGENIE is replaced with values computed by the emulator. The model is then run for an additional 100 kyr with astronomical parameters fixed at their average value across the adopted astronomical solution, allowing the system to adjust to the new PO_4 flux boundary condition. This average astronomical configuration is characterised by an eccentricity of 0.027, an obliquity of 0.388 rad (22.23°) (Farhat et al., 2022) and longitude of perihelion set to 90° ($\sin \varpi = 1$). To ensure continuity, the global average flux of PO_4 computed by the emulator is rescaled (decreased by approximately 10%) to perfectly align with the global average flux of PO_4 computed by the built-in weathering module of cGENIE at the end of SPIN2. Following this adaptation period, a series of 3 transient simulations spanning the full 1.1 Myr astronomical solution were conducted. These include: (1) a run with all astronomical parameters evolving (Full); (2) a sensitivity experiment with only obliquity varying while eccentricity and precession were held constant at their average values (Obli); (3) a run in which only eccentricity and precession vary, with obliquity fixed at its average value (Ecc-prec). Additionally, a control simulation was performed in which all astronomical parameters were held constant at their mean values.

3 Results

3.1 Moderate global astronomical influence

The global time series of atmospheric pCO_2 , oceanic PO_4 , and dissolved O_2 concentrations ($[\text{PO}_4]$ and $[\text{O}_2]$ respectively) for each transient simulation conducted across the astronomical solution are shown in Fig. 3. Throughout the reference simulation with all astronomical parameters evolving (black line in Fig. 3), atmospheric pCO_2 fluctuates by up to 40 ppm around a mean value of 555 ppm. In the ocean, globally-averaged $[\text{PO}_4]$ varies by approximately $0.1 \mu\text{mol.kg}^{-1}$ around a mean of $2.67 \mu\text{mol.kg}^{-1}$, while global mean $[\text{O}_2]$ remains within a range of $5 \mu\text{mol.kg}^{-1}$ around an average of $54 \mu\text{mol.kg}^{-1}$. The temporal variability of both atmospheric pCO_2 and oceanic $[\text{PO}_4]$ is almost exclusively driven by the combined forcings of eccentricity and precession, with obliquity playing a very minor role (see Fig. 3a and b). Overall, variations in global mean concentrations caused by astronomical forcing remain of moderate amplitude. This is consistent with the results of Sablon et al. (2025) and unsurprising considering that global averages can mask spatially heterogeneous responses, including contrasting and potentially compensating regional signals.

Atmospheric pCO_2 and oceanic $[\text{PO}_4]$ evolve in anti-phase throughout the simulation (see black and red curves in Fig. 3a and b). This inverse relationship occurs because intensified weathering simultaneously enhances CO_2 consumption and increases the continental flux of PO_4 to the ocean, leading to lower atmospheric pCO_2 and higher oceanic $[\text{PO}_4]$. The dominant control



190 of eccentricity and precession on atmospheric $p\text{CO}_2$ and oceanic $[\text{PO}_4]$ arises from their pronounced effect on continental weathering fluxes. Specifically, these two orbital parameters can drive changes in PO_4 delivery to the ocean of up to 9.5%, compared to only about 1.5% under obliquity-driven forcing alone. When all three orbital parameters evolve simultaneously, this modulation can reach 10.5%. Eccentricity is positively correlated with PO_4 weathering, such that periods of high eccentricity are associated with enhanced oceanic $[\text{PO}_4]$. This outcome arises from the physical processes encoded in the emulator
 195 developed by Sablon et al. (2025), where the same feature was previously identified.

A global low-oxygen state characterises the ocean in these simulations (see Fig. 3c) compared to the modern climate (Schmidtke et al., 2017; Garcia et al., 2024), consistent with the findings of Gérard et al. (2025) under similar boundary conditions and the estimations made by Tostevin and Mills (2020) based on long-term geochemical proxy data. While all three orbital parameters affect global-mean $[\text{O}_2]$, they operate through distinct mechanisms. Eccentricity and precession primarily
 200 influence global $[\text{O}_2]$ by modulating the oceanic PO_4 inventory. The relationship between globally-averaged $[\text{PO}_4]$ and $[\text{O}_2]$ is evidenced by a strong correlation ($R^2 = 0.82$ and $p\text{-value} = 0.016$)¹ between the two red curves in Fig. 3b and c. These $[\text{PO}_4]$ variations arise from changes in the continental weathering flux of PO_4 , with high eccentricity associated with reduced global $[\text{O}_2]$. In contrast, obliquity exerts only a very weak effect on global $[\text{PO}_4]$ that is insufficient to explain the simulated $[\text{O}_2]$ variations. This is reflected by the almost absent correlation ($R^2 = 0.06$ and $p\text{-value} = 0.467$) between the two blue curves in
 205 Fig. 3b and c. Instead, obliquity primarily influences global $[\text{O}_2]$ through its strong modulation of high-latitude insolation, as also noted by Gérard et al. (2025). Because modelled biological productivity explicitly depends on local insolation (Ridgwell et al., 2007), obliquity substantially alters oxygen consumption in these regions. These variations lead to changes in subsurface (around 60–130 m depth) $[\text{O}_2]$ of up to $4 \mu\text{mol.kg}^{-1}$. As these $[\text{O}_2]$ anomalies emerge in deep-water formation zones, they are efficiently distributed throughout the global ocean via large-scale overturning circulation, resulting in consistent oxygen
 210 variations of approximately $4 \mu\text{mol.kg}^{-1}$ across the benthic ocean (not shown). Consequently, times of high obliquity are associated with decreased global $[\text{O}_2]$, driven by increased biological productivity in high-latitude regions. There, the productivity anomalies driven by obliquity are more than three times larger than those caused by eccentricity and precession combined. Hence, the combination of high eccentricity and high obliquity results in the most severe depletion of simulated global $[\text{O}_2]$ (see Fig. 3c and d).

215 3.2 High regional variability

To assess regional expressions of $[\text{O}_2]$ variability driven by astronomical forcing, we defined distinct regions, as shown together with their associated watersheds in Fig. 4a and b. They are named after their approximate paleogeographic positions: region SA (South America), region WL (western Laurussia), region Si (Siberia), region NP (northern pole), and region LG (Laurussia–Gondwana). These regions were initially constrained by the depth distribution of Devonian anoxia proxy data,
 220 which predominantly originates from shallow-marine settings. Because the ocean floor from the Devonian period has been lost to subduction and plate tectonic processes, proxies for deep-water properties are scarce (Granot, 2016; Scotese and Wright,

¹ All presented statistics account for the temporal autocorrelation inherent in our setup, using an effective sample size correction based on the N_{eff} method, following the approach of Bretherton et al. (1999).



2018). Consequently, grid cells deeper than 1000 meters were excluded, except for those associated with deep-water formation, which occurs in regions SA and NP. Within this depth-limited framework, we further refined the regions by grouping model grid points based on the temporal variance of $[PO_4]$ and $[O_2]$ (see Appendix A), with the final result shown in Fig. 4a. Hence, this approach enables us to assess whether astronomical forcing triggers distinct biogeochemical responses across various regions.

Fig. 4c displays the time series of regionally-averaged $[O_2]$ and anoxic volume anomalies, for each region (as defined in Fig. 4a), in the reference simulation with all astronomical parameters evolving concurrently (Full). The results highlight a marked spatial heterogeneity in the biogeochemical response to astronomical forcing. On the one hand, regions SA and LG exhibit strong sensitivity to astronomical variability. In these regions, $[O_2]$ fluctuates by 45 and $19.5 \mu\text{mol.kg}^{-1}$, respectively, equivalent to a 35% deviation from their regional means. Simulated $[O_2]$ reach their lowest values during periods of maximum eccentricity. These pronounced oxygen variations translate into substantial changes in anoxic volume, reaching approximately 19% in region SA and 13.5% in region LG. On the other hand, region Si is mostly unaffected by astronomical forcing. Temporal variability in both $[O_2]$ and anoxic volume is limited to approximately 5% and 2.5%, respectively, with the anoxic volume remaining stable throughout most of the simulation. Although the signal is subtle in magnitude, region Si exhibits an opposite precession phasing compared to regions SA and LG, a feature resulting from their locations in different hemispheres. Remarkably, region NP exhibits no anoxic conditions over the period, because of its consistently high $[O_2]$ and the limited magnitude of $[O_2]$ variations, which are insufficient for any grid cell in the region to cross the anoxic threshold.

The relative influence of individual astronomical parameters is clarified by the power spectrum analysis of the $[O_2]$ time series (Fig. 5). Precession emerges as the dominant driver of $[O_2]$ variability in regions SA, WL, Si, and LG, while $[O_2]$ in region NP is primarily shaped by obliquity. Although eccentricity produces no distinct spectral peak, it remains essential through its modulation of precession amplitude (see Fig. 4c). Notably, periods of high eccentricity tend to coincide with decreased oxygen levels across all regions, underscoring its consistent impact.

By definition, variations in anoxic volume result from changes in regional $[O_2]$. These $[O_2]$ fluctuations are closely linked to alterations in $[PO_4]$, where higher $[PO_4]$ leads to decreased $[O_2]$ through the stimulation of biological productivity, hence remineralization. In regions SA, WL, and LG, variations in $[PO_4]$ drive almost single-handedly the changes in $[O_2]$ (regional $R^2 \geq 0.88$ and $p\text{-values} < 10^{-3}$; results not shown). Region Si exhibits a more complex relationship between $[PO_4]$ and $[O_2]$. Although $[PO_4]$ remains the primary driver of $[O_2]$ variability ($R^2 = 0.62$ and $p\text{-value} < 10^{-3}$), a substantial secondary contribution is apparent, likely arising from insolation-driven changes in biological productivity. The limited amplitude of $[PO_4]$ variations in this region enhances the relative influence of these insolation-related effects, allowing them to leave a more discernible imprint on $[O_2]$ dynamics. In region NP, the influence of $[PO_4]$ on $[O_2]$ is minimal ($R^2 = 0.18$ and $p\text{-value} = 0.07$). As in region Si, limited $[PO_4]$ variability makes insolation-driven productivity changes more apparent, and they even become dominant here due to the strong effect of obliquity at high latitudes (as already noted in Section 3.1). Overall, when regional $[O_2]$ variability is high, it is almost entirely driven by $[PO_4]$. However, when $[PO_4]$ fluctuations are limited, other mechanisms, such as insolation-driven changes in biological productivity, become more apparent.



With the link between $[\text{PO}_4]$ and $[\text{O}_2]$ established, we now turn to the drivers of $[\text{PO}_4]$ variability. Figure 6 shows the time series of average $[\text{PO}_4]$ and PO_4 weathering, in addition to the linear regression between these two quantities for each region. Region NP has been excluded because there is no direct regional weathering flux of PO_4 coming from the continents. For correlation analysis, the $[\text{PO}_4]$ time series for each region (except region Si) was slightly shifted to best align with the weathering time series. This adjustment accounts for the natural time lag between the two quantities, as $[\text{PO}_4]$ typically increases when the weathering flux of PO_4 is high. Namely, even as continental weathering begins to decrease, $[\text{PO}_4]$ continues to rise as long as weathering remains high compared to its mean value, producing a lag of several thousand years between the weathering signal and the $[\text{PO}_4]$ response (see Fig. 6). To ensure the mechanism remains consistent with an astronomical-forcing interpretation, this lag was chosen to maximize the correlation but was limited to a maximum of 5 kyr (approximately one-fourth of a precession cycle). When a negative lag is applied to $[\text{PO}_4]$ in regions SA, WL, and LG, it becomes clear that regional weathering accounts for most of the variations in $[\text{PO}_4]$. Hence, in these regions, variations in $[\text{O}_2]$ and anoxic extent are predominantly driven by regional $[\text{PO}_4]$, hence PO_4 weathering, which in turn is mainly modulated by eccentricity and precession (see Fig. 6).

In contrast, region Si exhibits a markedly different relationship between $[\text{PO}_4]$ and PO_4 weathering. Specifically, the $[\text{PO}_4]$ time series slightly precedes that of the weathering flux, leading to a negative correlation when a negative lag is applied to PO_4 as was previously done for other regions (not shown; Fig. 6 hence uses no lag correction for region Si). This observation contradicts the mechanism described above, suggesting that regional weathering cannot account for the variations in $[\text{PO}_4]$ observed in this region. Local weathering fails to leave a strong imprint on $[\text{PO}_4]$ variations due to its relative weakness, about half that of region LG (see Fig. 6), and the physical oceanographic conditions in region Si, where strong currents cause important advection of weathered PO_4 , preventing its local accumulation (see Fig. 7). As a result, the regional $[\text{PO}_4]$ signature is strongly influenced by continental weathering fluxes originating outside of region Si, predominantly from the Northern Hemisphere. Consequently, the temporal evolution of $[\text{PO}_4]$ in region Si closely mirrors the broader-scale dynamics of the entire Northern hemisphere, rather than reflecting a local weathering-driven signal. This is supported by a strong correlation between the $[\text{PO}_4]$ time series of the Northern Hemisphere and region Si ($R^2 = 0.98$ and $p\text{-value} < 10^{-3}$). In this context, the variations in $[\text{PO}_4]$ are predominantly shaped by physical transport processes within the ocean, highlighting the central role of circulation and mixing in constraining the variation of $[\text{PO}_4]$, hence anoxia.

4 Discussion

4.1 Implications

We used cGENIE with an upgraded representation of weathering taken from Sablon et al. (2025), to investigate the influence of astronomical forcing on Late Devonian ocean oxygenation. We found that astronomical forcing leaves a discernible imprint on the global time series of atmospheric pCO_2 , oceanic $[\text{PO}_4]$, and dissolved $[\text{O}_2]$. Notably, the evolution of atmospheric pCO_2 under astronomical forcing closely mirrors the results reported by Sablon et al. (2025), exhibiting a similar amplitude of approximately 40 ppm, with higher atmospheric pCO_2 concentrations corresponding to periods of minimum eccentricity.



A similar $p\text{CO}_2$ range was also obtained by Vervoort et al. (2024) when running cGENIE with astronomical forcing, using a
 290 very different experimental setup featuring an idealized continental configuration and no representation of C_{org} burial. In our
 simulations, $[\text{O}_2]$ variations primarily occur through the influence of eccentricity and precession on continental weathering
 fluxes, as well as obliquity on high-latitude biological productivity. Periods of combined high eccentricity and high obliquity
 produce the most pronounced deoxygenation, although global mean $[\text{O}_2]$ variations remain within a modest magnitude range
 of $5 \mu\text{mol.kg}^{-1}$ throughout the simulations. Because the ocean exhibits substantial spatial heterogeneity in benthic $[\text{O}_2]$, with
 295 differences reaching $100 \mu\text{mol.kg}^{-1}$, it is unlikely for the simulated $[\text{O}_2]$ variations to drive global-scale benthic areas below
 the anoxia threshold. Based on our modelling, it is thus highly implausible that the astronomical forcing alone could induce an
 anoxic event over the entire global ocean (i.e., including the benthic ocean).

Building on the global-scale analysis, we performed a more localized investigation by defining specific regions to capture
 potentially distinct biogeochemical dynamics induced by astronomical forcing. The results reveal that eccentricity maxima
 300 commonly coincide with the most pronounced anoxic extent, supporting the proposal by Ma et al. (2022) and Huygh et al.
 (2025) that the Upper Kellwasser coincide with an eccentricity maxima configuration. However, both the magnitude and the
 dominant astronomical parameter driving this response vary markedly across regions, underscoring the spatial heterogeneity
 captured in our simulations. While certain regions exhibit strong sensitivity to astronomical forcing, with $[\text{O}_2]$ varying by up
 to 35% and the anoxic volume by up to 19%, others remain largely unaffected. Regions near continental margins (i.e., more
 305 directly influenced by weathering fluxes) exhibit a strong precessional control, with the largest variations in $[\text{O}_2]$ and anoxic
 volume occurring in these weathering-dominated settings. This influence is not limited to low-latitude areas; for instance,
 region SA exhibits a notable precession signal despite its relatively high latitude, whereas precession is generally thought to
 primarily affect low latitudes due to its strong control on precipitation patterns (De Vleeschouwer et al., 2012). In contrast,
 high-latitude regions far from continental margins (i.e., less directly influenced by weathering fluxes) are more sensitive to
 310 obliquity-driven changes, particularly through their influence on biological productivity. These findings underscore the pivotal
 role of PO_4 weathering and biological productivity in shaping regional biogeochemical responses, aligning with the 'top-down'
 hypothesis proposed in Carmichael et al. (2019). However, physical processes such as ocean circulation and tracer advection
 can override this control, as exemplified in region Si, where local $[\text{PO}_4]$ dynamics are decoupled from regional weathering
 due to strong transport. Our results further demonstrate how regions SA, WL and LG, characterized by relatively weak surface
 315 circulation, provide ideal conditions for astronomically driven weathering variations to leave a substantial imprint on local
 $[\text{O}_2]$.

Altogether, these results demonstrate that astronomical forcing can exert pronounced control on ocean biogeochemistry,
 with its impact shaped by local physical conditions. Furthermore, the regional analysis of the shallow ocean suggests that
 an astronomical signal may appear globally distributed across the observational record. However, it also indicates that proxy
 320 records from distinct locations may reflect fundamentally different expressions of an astronomical signal, potentially leading
 to contrasting interpretations of its role in driving ocean anoxia.

Despite the important contribution that astronomical forcing can exert on $[\text{O}_2]$, the biogeochemical response of the system
 remains relatively simple when compared to the forcing. Specifically, the regions often exhibit a quasi-linear response to the



forcing, lacking clear signs of long-term memory effects or strong non-linearities. This is particularly surprising given that
 325 redox-dependent P regeneration is represented in our model configuration (using the scheme from Wallmann, 2010), a feature
 explicitly proposed in Percival et al. (2020) to explain the sustainability of Late Devonian anoxia based on the investigation
 of sedimentary concentrations. Hence, the absence of prolonged anoxia in our simulations indicates that the PO_4 variations
 induced by astronomical forcing may be either too gradual or too limited in amplitude to trigger or maintain an anoxic state
 over extended timescales in our experimental setup (e.g., 10^5 years). Our results also suggest that anoxic events could manifest
 330 as a succession of smaller, transient episodes (see Fig. 4c and supplementary video) rather than a single, prolonged global
 event, similar to the scenario proposed by Hedhli et al. (2023). These findings motivate future studies centred on additional
 processes that may influence the system response, including vegetation–climate interactions driven by orbital forcing and the
 effects of large, episodic nutrient inputs. In particular, assessing the persistence and biogeochemical consequences of such
 perturbations could help constrain the conditions under which sustained anoxia might emerge.

335 4.2 Limitations

Because cGENIE is a relatively simple model, it suffers from well-known intrinsic limitations. Among these are its coarse spa-
 tial resolution, resulting in a simplified representation of continental geometry and shallow-water environments, the absence
 of explicitly resolved atmospheric dynamics, a basic hydrological scheme, and the lack of ice-sheet dynamics. Nevertheless,
 cGENIE has been widely used in previous studies and has demonstrated its capacity to reproduce large-scale ocean circulation
 340 features (Rahmstorf et al., 2005; Lunt et al., 2006; Cao et al., 2009; Marsh et al., 2013; Pohl et al., 2022; Gérard and Crucifix,
 2024; Laridon et al., 2025) and to accurately simulate key aspects of ocean biogeochemistry (Ridgwell et al., 2007; Ridgwell
 and Hargreaves, 2007; Colbourn et al., 2013; Ward et al., 2018; Adloff et al., 2020; Reinhard et al., 2020; Crichton et al., 2021;
 Van De Velde et al., 2021; Naidoo-Bagwell et al., 2024; Stappard et al., 2025). Furthermore, some of these limitations are
 addressed by our specific experimental design. The prescribed wind stress field, although not responsive to climate variability,
 345 was derived from HadCM3 simulations conducted by Sablon et al. (2025). Additionally, the emulator accounts for changes
 in precipitation associated with variations in astronomical configuration and atmospheric pCO_2 , thereby improving the repre-
 sentation of hydrological controls on PO_4 weathering. Together, these features enable a more realistic representation of two
 central climate components in our study: PO_4 weathering and ocean circulation.

Still, several important limitations of our approach carry implications for interpreting the results. Our results could depend
 350 on the selected paleogeographic reconstruction, given the substantial uncertainties that persist in deep-time topography and,
 by extension, in slope, erosion, and weathering patterns. Moreover, a simple nearest-neighbour scheme was used to route
 continental weathering products into the ocean; alternative watershed configurations could yield significantly different regional
 responses to astronomical forcing. The current setup also does not capture extreme events that may substantially influence
 sedimentary fluxes, and relies on a simplified organic matter sedimentation scheme that lacks oxygen dependence, highlighting
 355 the potential benefit of integrating more advanced models of organic matter diagenesis like OMEN-SED (Hülse et al., 2018) in
 future work. Additionally, the surface wind stress field is held constant throughout the simulations, preventing realistic changes
 in circulation patterns in response to evolving astronomical configurations. The exclusion of ice sheets and sea-level variations



removes critical mechanisms through which obliquity can strongly impact $[O_2]$, although the accurate representation of sea-level changes would remain challenging at the cGENIE grid resolution. Several studies have suggested that these processes may have contributed to, or exacerbated, Late Devonian anoxic events (Bond and Wignall, 2008; Chen et al., 2013; Huygh et al., 2025). As a result, the influence of obliquity identified in this study may be underestimated, though it still leaves a discernible imprint on the global ocean. For all these reasons, we emphasize the importance of further modelling efforts to better connect orbital forcing with ocean anoxia. In this context, Earth System Models of Intermediate Complexity (EMICs) offer a promising path forward, as they allow for long-timescale simulations while explicitly resolving many key physical and biogeochemical processes.

5 Conclusions

We have explored the role of astronomical forcing in driving ocean anoxia under a Late Devonian climate configuration. Our experimental setup relies on an offline coupling between cGENIE and a fast climate–weathering emulator, based on a GCM, developed by Sablon et al. (2025). It allowed us to capture the influence of astronomical forcing on oceanic $[O_2]$ via its modulation of continental PO_4 weathering fluxes. At the global scale, astronomical forcing exerts a moderate impact on ocean $[O_2]$, with the most pronounced deoxygenation occurring during intervals of combined high eccentricity and high obliquity. While eccentricity and precession primarily affect $[O_2]$ through their control on PO_4 weathering fluxes, obliquity exerts its influence by modulating high-latitude biological productivity, highlighting that even in a greenhouse world, obliquity can leave a discernible imprint on ocean biogeochemistry, hence sedimentary records. At the regional scale, eccentricity maxima also correspond to the most extensive anoxic conditions, providing new evidence supporting a strong impact of eccentricity maxima on the development of Late Devonian anoxic events. However, the biogeochemical response remains highly variable across regions. Our results show that astronomical forcing can act as a dominant control on ocean $[O_2]$, driving regional changes in anoxic volume of up to 19%. An unanticipated result is that local ocean circulation can represent a central driver of the spatial heterogeneity in $[O_2]$ responses to astronomical forcing. As a result, sedimentary records from different locations may reflect contrasting imprints of the same astronomical signal, potentially leading to contrasting interpretations of its role in pacing Late Devonian anoxia. Finally, while this study provides insights into how astronomical forcing affects Late Devonian ocean $[O_2]$, it also stresses the difficulty of explaining the duration of Devonian anoxic events and the uncertainties in their spatial extent, underscoring that much modelling and data integration work remains to fully understand the drivers and dynamics of Late Devonian anoxia.

Appendix A: Defining regions

This section clarifies how we refined the definition of the different regions based on the temporal variance of $[O_2]$ and $[PO_4]$. Figure A1 shows the normalized temporal variance of these two tracers, averaged over the upper 1000 m, consistent with the depth range applied in our regional analysis (see Section 3.2). First, a clear distinction between regions SA and WL



emerges from both $[O_2]$ and $[PO_4]$ variance fields, particularly around $30^\circ S$ and $60^\circ W$, providing strong justification for
390 separating them. Second, region LG was initially split into two sub-regions based on the spatial structure of the variance fields.
However, the temporal evolution and variability of $[O_2]$, anoxic volume, and $[PO_4]$ were found to be highly consistent across
these sub-regions. Consequently, we merged them into a single, unified region, which we now refer to as region LG. These
refinements provide a clear and consistent basis for the regional definitions presented in Fig. 4a. Although regional boundaries
are somewhat arbitrary, they follow coherent patterns of variability and are broad enough to avoid overinterpreting hyper-
395 localised phenomena (at the single grid cell level). The resulting regions are not intended as universal classifications, but rather
as a practical framework for exploring the spatial heterogeneity of climate–biogeochemical interactions.

Code availability. The code for the version of the ‘muffin’ release of the cGENIE Earth system model used in this paper, is tagged as v0.9.50,
and is assigned a DOI: *to be added*. Configuration files for the specific experiments presented in the paper can be found in the directory: *to*
be added. Details of the experiments, plus the command line needed to run each one, are given in the ‘readme.txt’ file in that directory. All
400 other configuration files and boundary conditions are provided as part of the code release. A manual detailing code installation, basic model
configuration, tutorials covering various aspects of model configuration, experimental design, and output, plus the processing of results, is
assigned the following DOI: <https://doi.org/10.5281/zenodo.13377225>. *Full directory and experiment information will be provided if the*
paper is accepted.

Video supplement. Spatial $[O_2]$ anomaly (bottom), averaged from the second vertical layer down to approximately 1000 m depth, evolving
405 through the 1.1 Myr astronomical solution (top). Dashed lines indicate the percentage of the water column (up to 1000 m) below the anoxic
threshold of $6.5 \mu\text{mol.kg}^{-1}$. The red dot represents climatic precession, the red cross eccentricity, and the blue dot obliquity. Periods of
reduced $[O_2]$ appear as distinct pulses rather than a single prolonged event. For instance, severe deoxygenation is particularly evident at 0.15
and 0.98 Myr.

Author contributions. JG conducted the cGENIE experiments, developed the code for the analysis, performed all computations, and drafted
410 the manuscript. AP contributed to the development of the open setup configuration of cGENIE. MC and ACDS contributed to the study
design. All authors contributed to the discussion of results, writing the manuscript and approved the final version.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Acknowledgements. This research is funded by the Belgian National Fund of Scientific Research (FNRS), project WarmAnoxia, PDR grant
no. T.0037.22. Computational resources have been provided by the supercomputing facilities of the Université catholique de Louvain (CIS-



- 415 M/UCL) and the Consortium des Équipements de Calcul Intensif en Fédération Wallonie Bruxelles (CÉCI). AP acknowledges the support of the French *Agence Nationale de la Recherche* (ANR) under reference ANR-23-CE01-0003 (project CYCLO-SED).



References

- Adloff, M., Ridgwell, A., Monteiro, F. M., Parkinson, I. J., Dickson, A., Pogge von Strandmann, P. A., Fantle, M. S., and Greene, S. E.: Inclusion of a suite of weathering tracers in the cGENIE Earth System Model—muffin release v. 0.9. 10, *Geoscientific Model Development Discussions*, 2020, 1–48, <https://doi.org/https://doi.org/10.5194/gmd-14-4187-2021>, 2020.
- Algeo, T. J. and Scheckler, S. E.: Terrestrial-marine teleconnections in the Devonian: links between the evolution of land plants, weathering processes, and marine anoxic events, *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 353, 113–130, <https://doi.org/10.1098/rstb.1998.0195>, 1998.
- Becker, R., Marshall, J., Da Silva, A.-C., Agterberg, F., Gradstein, F., and Ogg, J.: The Devonian Period, in: *Geologic Time Scale 2020*, pp. 733–810, Elsevier, ISBN 978-0-12-824360-2, <https://doi.org/10.1016/B978-0-12-824360-2.00022-X>, 2020.
- Benitez-Nelson, C. R.: The biogeochemical cycling of phosphorus in marine systems, *Earth-Science Reviews*, 51, 109–135, [https://doi.org/https://doi.org/10.1016/S0012-8252\(00\)00018-0](https://doi.org/https://doi.org/10.1016/S0012-8252(00)00018-0), 2000.
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years, *Quaternary Science Reviews*, 10, 297 – 317, [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- Bond, D. P. and Wignall, P. B.: The role of sea-level change and marine anoxia in the Frasnian–Famennian (Late Devonian) mass extinction, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 263, 107–118, <https://doi.org/https://doi.org/10.1016/j.palaeo.2008.02.015>, 2008.
- Bradley, J. A., Hülse, D., LaRowe, D. E., and Arndt, S.: Transfer efficiency of organic carbon in marine sediments, *Nature communications*, 13, 7297, <https://doi.org/https://doi.org/10.1038/s41467-022-35112-9>, 2022.
- Bretherton, C. S., Widmann, M., Dymnikov, V. P., Wallace, J. M., and Bladé, I.: The effective number of spatial degrees of freedom of a time-varying field, *Journal of climate*, 12, 1990–2009, 1999.
- Brugger, J., Hofmann, M., Petri, S., and Feulner, G.: On the Sensitivity of the Devonian Climate to Continental Configuration, Vegetation Cover, Orbital Configuration, CO₂ Concentration, and Insolation, *Paleoceanography and Paleoclimatology*, 34, 1375–1398, <https://doi.org/10.1029/2019PA003562>, 2019.
- Cao, L., Eby, M., Ridgwell, A., Caldeira, K., Archer, D., Ishida, A., Joos, F., Matsumoto, K., Mikolajewicz, U., Mouchet, A., Orr, J. C., and Yool, A.: The role of ocean transport in the uptake of anthropogenic CO₂, *Biogeosciences*, 6, 375–390, <https://doi.org/10.5194/bg-6-375-2009>, 2009.
- Carmichael, S. K., Waters, J. A., Königshof, P., Suttner, T. J., and Kido, E.: Paleogeography and paleoenvironments of the Late Devonian Kellwasser event: A review of its sedimentological and geochemical expression, *Global and Planetary Change*, 183, 102984, <https://doi.org/10.1016/j.gloplacha.2019.102984>, 2019.
- Chen, B., Ma, X., Mills, B. J., Qie, W., Joachimski, M. M., Shen, S., Wang, C., Xu, H., and Wang, X.: Devonian paleoclimate and its drivers: A reassessment based on a new conodont $\delta^{18}\text{O}$ record from South China, *Earth-Science Reviews*, 222, 103814, <https://doi.org/https://doi.org/10.1016/j.earscirev.2021.103814>, 2021.
- Chen, D., Wang, J., Racki, G., Li, H., Wang, C., Ma, X., and Whalen, M. T.: Large sulphur isotopic perturbations and oceanic changes during the Frasnian–Famennian transition of the Late Devonian, *Journal of the Geological Society*, 170, 465–476, <https://doi.org/10.1144/jgs2012-037>, 2013.
- Claussen, M., Mysak, L., Weaver, A., Crucifix, M., Fichet, T., Loutre, M.-F., Weber, S., Alcamo, J., Alexeev, V., Berger, A., Calov, R., Ganopolski, A., Goosse, H., Lohmann, G., Lunkeit, F., Mokhov, I., Petoukhov, V., Stone, P., and Wang, Z.: Earth system



- models of intermediate complexity: closing the gap in the spectrum of climate system models, *Climate Dynamics*, 18, 579–586, <https://doi.org/10.1007/s00382-001-0200-1>, 2002.
- 455 Colbourn, G., Ridgwell, A., and Lenton, T.: The rock geochemical model (RokGeM) v0. 9, *Geoscientific Model Development*, 6, 1543–1573, 2013.
- Crichton, K. A., Wilson, J. D., Ridgwell, A., and Pearson, P. N.: Calibration of temperature-dependent ocean microbial processes in the cGENIE.muffin (v0.9.13) Earth system model, *Geoscientific Model Development*, 14, 125–149, <https://doi.org/10.5194/gmd-14-125-2021>, 2021.
- 460 Crucifix, M.: Generation of sample climatic precession and obliquity solutions for running climate experiments over the Devonian, <https://doi.org/10.5281/zenodo.14894811>, 2025.
- Da Silva, A.-C., Sinnesael, M., Claeys, P., Davies, J. H. F. L., De Winter, N. J., Percival, L. M. E., Schaltegger, U., and De Vleeschouwer, D.: Anchoring the Late Devonian mass extinction in absolute time by integrating climatic controls and radio-isotopic dating, *Scientific Reports*, 10, 12 940, <https://doi.org/10.1038/s41598-020-69097-6>, 2020.
- 465 Dahl, T. W., Harding, M. A., Brugger, J., Feulner, G., Norrman, K., Lomax, B. H., and Junium, C. K.: Low atmospheric CO₂ levels before the rise of forested ecosystems, *Nature Communications*, 13, 7616, <https://doi.org/10.1038/s41467-022-35085-9>, 2022.
- De Vleeschouwer, D., Da Silva, A.-C., Boulvain, F., Crucifix, M., and Claeys, P.: Precessional and half-precessional climate forcing of Mid-Devonian monsoon-like dynamics, *Climate of the Past*, 8, 337–351, <https://doi.org/10.5194/cp-8-337-2012>, 2012.
- De Vleeschouwer, D., Rakociński, M., Racki, G., Bond, D. P., Sobieć, K., and Claeys, P.: The astronomical rhythm of Late-
 470 Devonian climate change (Kowala section, Holy Cross Mountains, Poland), *Earth and Planetary Science Letters*, 365, 25–37, <https://doi.org/10.1016/j.epsl.2013.01.016>, 2013.
- De Vleeschouwer, D., Crucifix, M., Bounceur, N., and Claeys, P.: The impact of astronomical forcing on the Late Devonian greenhouse climate, *Global and Planetary Change*, 120, 65–80, <https://doi.org/10.1016/j.gloplacha.2014.06.002>, 2014.
- De Vleeschouwer, D., Da Silva, A.-C., Sinnesael, M., Chen, D., Day, J. E., Whalen, M. T., Guo, Z., and Claeys, P.: Timing and
 475 pacing of the Late Devonian mass extinction event regulated by eccentricity and obliquity, *Nature Communications*, 8, 2268, <https://doi.org/10.1038/s41467-017-02407-1>, 2017.
- Dopieralska, J.: Reconstructing seawater circulation on the Moroccan shelf of Gondwana during the Late Devonian: Evidence from Nd isotope composition of conodonts, *Geochemistry, Geophysics, Geosystems*, 10, <https://doi.org/10.1029/2008GC002247>, 2009.
- Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the
 480 ocean interior and on the seafloor, *Global Biogeochemical Cycles*, 21, <https://doi.org/10.1029/2006GB002907>, 2007.
- Edwards, N. R. and Marsh, R.: Uncertainties due to transport-parameter sensitivity in an efficient 3-D ocean-climate model, *Climate Dynamics*, 24, 415–433, <https://doi.org/10.1007/s00382-004-0508-8>, 2005.
- Farhat, M., Auclair-Desrotour, P., Boué, G., and Laskar, J.: The resonant tidal evolution of the Earth-Moon distance, *Astronomy & Astrophysics*, 665, L1, <https://doi.org/10.1051/0004-6361/202243445>, publisher: EDP Sciences, 2022.
- 485 Frankignoulle, M. and Gattuso, J.-P.: Air-sea CO₂ exchange in coastal ecosystems, in: *Interactions of C, N, P and S biogeochemical cycles and global change*, pp. 233–248, Springer, https://doi.org/10.1007/978-3-642-76064-8_9, 1993.
- Garcia, H. E., Wang, Z., Bouchard, C., Cross, S. L., Paver, C. R., Reagan, J. R., Boyer, T. P., Locarnini, R. A., Mishonov, A. V., Baranova, O. K., et al.: *World Ocean Atlas 2023, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, Dissolved Oxygen Saturation and 30-year Climate Normal*, <https://doi.org/10.25923/rb67-ns53>, 2024.



- 490 Gough, D.: Solar interior structure and luminosity variations, in: *Physics of Solar Variations: Proceedings of the 14th ESLAB Symposium* held in Scheveningen, The Netherlands, 16–19 September, 1980, pp. 21–34, Springer, 1981.
- Granot, R.: Palaeozoic oceanic crust preserved beneath the eastern Mediterranean, *Nature Geoscience*, 9, 701–705, <https://doi.org/https://doi.org/10.1038/ngeo2784>, 2016.
- Gérard, J. and Crucifix, M.: Diagnosing the causes of AMOC slowdown in a coupled model: a cautionary tale, *Earth System Dynamics*, 15, 293–306, <https://doi.org/10.5194/esd-15-293-2024>, 2024.
- 495 Gérard, J., Sablon, L., Huygh, J. J. C., Da Silva, A.-C., Pohl, A., Vérard, C., and Crucifix, M.: Exploring the mechanisms of Devonian oceanic anoxia: impact of ocean dynamics, palaeogeography, and orbital forcing, *Climate of the Past*, 21, 239–260, <https://doi.org/10.5194/cp-21-239-2025>, 2025.
- Haddad, E. E., Boyer, D. L., Droser, M. L., Lee, B. K., Lyons, T. W., and Love, G. D.: Ichnofabrics and chemostratigraphy argue against persistent anoxia during the Upper Kellwasser Event in New York State, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 490, 178–190, <https://doi.org/10.1016/j.palaeo.2017.10.025>, 2018.
- 500 Hedhli, M., Grasby, S., Henderson, C., and Davis, B.: Multiple diachronous “Black Seas” mimic global ocean anoxia during the latest Devonian, *Geology*, 51, 973–977, <https://doi.org/10.1130/G51394.1>, 2023.
- Huang, C., Joachimski, M. M., and Gong, Y.: Did climate changes trigger the Late Devonian Kellwasser Crisis? Evidence from a high-resolution conodont $\delta^{18}\text{OPO4}$ record from South China, *Earth and Planetary Science Letters*, 495, 174–184, <https://doi.org/https://doi.org/10.1016/j.epsl.2018.05.016>, 2018.
- 505 Hülse, D. and Ridgwell, A.: Instability in the geological regulation of Earth’s climate, <https://doi.org/https://doi.org/10.31223/X5F13X>, 2024.
- Hülse, D., Arndt, S., Daines, S., Regnier, P., and Ridgwell, A.: OMEN-SED 1.0: a novel, numerically efficient organic matter sediment diagenesis module for coupling to Earth system models, *Geoscientific Model Development*, 11, 2649–2689, <https://doi.org/https://doi.org/10.5194/gmd-11-2649-2018>, 2018.
- 510 Huygh, J. J. C., Algeo, T. J., Sageman, B. B., Arts, M. C., Ver Straeten, C. A., Over, D. J., Gérard, J., Sablon, L., Crucifix, M., and Da Silva, A.-C.: Astronomical control on marine anoxia during the Kellwasser Crisis and Rhinestreet Event (Appalachian Basin, New York, USA), 2025.
- Ingall, E. and Jahnke, R.: Evidence for enhanced phosphorus regeneration from marine sediments overlain by oxygen depleted waters, *Geochimica et Cosmochimica Acta*, 58, 2571–2575, [https://doi.org/https://doi.org/10.1016/0016-7037\(94\)90033-7](https://doi.org/https://doi.org/10.1016/0016-7037(94)90033-7), 1994.
- 515 Joachimski, M. M. and Buggisch, W.: Conodont apatite $\delta^{18}\text{O}$ signatures indicate climatic cooling as a trigger of the Late Devonian mass extinction, *Geology*, 30, 711–714, [https://doi.org/https://doi.org/10.1130/0091-7613\(2002\)030<0711:CAOSIC>2.0.CO;2](https://doi.org/https://doi.org/10.1130/0091-7613(2002)030<0711:CAOSIC>2.0.CO;2), 2002.
- Kabanov, P., Hauck, T. E., Gouwy, S. A., Grasby, S. E., and Van Der Boon, A.: Oceanic anoxic events, photic-zone euxinia, and controversy of sea-level fluctuations during the Middle-Late Devonian, *Earth-Science Reviews*, 241, 104415, <https://doi.org/10.1016/j.earscirev.2023.104415>, 2023.
- 520 Kaiser, S. I., Aretz, M., and Becker, R. T.: The global Hangenberg Crisis (Devonian–Carboniferous transition): review of a first-order mass extinction, *Geological Society, London, Special Publications*, 423, 387–437, <https://doi.org/10.1144/SP423.9>, 2016.
- Kiessling, W., Flügel, E., and Golonka, J.: Patterns of Phanerozoic carbonate platform sedimentation, *Lethaia*, 36, 195–226, <https://doi.org/https://doi.org/10.1080/00241160310004648>, 2003.
- 525 Krause, A. J., Mills, B. J., Zhang, S., Planavsky, N. J., Lenton, T. M., and Poulton, S. W.: Stepwise oxygenation of the Paleozoic atmosphere, *Nature Communications*, 9, 4081, <https://doi.org/https://doi.org/10.1038/s41467-018-06383-y>, 2018.



- Laridon, A., Couplet, V., Gérard, J., Thiery, W., and Crucifix, M.: Connecting complex and simplified models of tipping elements: a nonlinear two-forcing emulator for the Atlantic meridional overturning circulation, *Open Research Europe*, 5, 87, 2025.
- Laskar, J. and Robutel, P.: The chaotic obliquity of the planets, *Nature*, 361, 608–612, <https://doi.org/10.1038/361608a0>, 1993.
- 530 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C., and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth, *Astronomy & Astrophysics*, 428, 261–285, <https://doi.org/https://doi.org/10.1051/0004-6361/20041335>, 2004.
- Laskar, J., Fienga, A., Gastineau, M., and Manche, H.: La2010: a new orbital solution for the long-term motion of the Earth, *Astronomy and Astrophysics*, 532, A89, <https://doi.org/10.1051/0004-6361/201116836>, 2011.
- Lunt, D., Williamson, M., Valdes, P., Lenton, T., and Marsh, R.: Comparing transient, accelerated, and equilibrium simulations of the last 30
 535 000 years with the GENIE-1 model, *Climate of the Past*, 2, 221–235, <https://doi.org/10.5194/cp-2-221-2006>, 2006.
- Ma, K., Hinnov, L., Zhang, X., and Gong, Y.: Astronomical climate changes trigger Late Devonian bio-and environmental events in South China, *Global and Planetary Change*, 215, 103 874, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2022.103874>, 2022.
- Ma, X., Gong, Y., Chen, D., Racki, G., Chen, X., and Liao, W.: The Late Devonian Frasnian–Famennian Event in South China—patterns and causes of extinctions, sea level changes, and isotope variations, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 448, 224–244,
 540 <https://doi.org/https://doi.org/10.1016/j.palaeo.2015.10.047>, 2016.
- Maffre, P., Goddérès, Y., Le Hir, G., Nardin, E., Sarr, A.-C., and Donnadieu, Y.: GEOCLIM7, an Earth System Model for multi-million years evolution of the geochemical cycles and climate, *Geoscientific Model Development Discussions*, 2024, 1–66, <https://doi.org/10.5194/gmd-2024-220>, 2024.
- Marsh, R., Müller, S. A., Yool, A., and Edwards, N. R.: Incorporation of the C-GOLDSTEIN efficient climate model into the GENIE frame-
 545 work: “eb_go_gs” configurations of GENIE, *Geoscientific Model Development*, 4, 957–992, <https://doi.org/10.5194/gmd-4-957-2011>, 2011.
- Marsh, R., Sobester, A., Hart, E. E., Oliver, K., Edwards, N., and Cox, S.: An optimally tuned ensemble of the “eb_go_gs” configuration of GENIE: parameter sensitivity and bifurcations in the Atlantic overturning circulation, *Geoscientific Model Development Discussions*, 6, 925–956, <https://doi.org/10.5194/gmd-6-1729-2013>, 2013.
- 550 Mills, B. J., Krause, A. J., Jarvis, I., and Cramer, B. D.: Evolution of atmospheric O₂ through the Phanerozoic, revisited, *Annual review of earth and planetary sciences*, 51, 253–276, 2023.
- Naidoo-Bagwell, A. A., Monteiro, F. M., Hendry, K. R., Burgan, S., Wilson, J. D., Ward, B. A., Ridgwell, A., and Conley, D. J.: A diatom extension to the cGENIE Earth system model–EcoGENIE 1.1, *Geoscientific Model Development*, 17, 1729–1748, <https://doi.org/https://doi.org/10.5194/gmd-17-1729-2024>, 2024.
- 555 Paschall, O., Carmichael, S. K., Königshof, P., Waters, J. A., Ta, P. H., Komatsu, T., and Dombrowski, A.: The Devonian–Carboniferous boundary in Vietnam: Sustained ocean anoxia with a volcanic trigger for the Hangenberg Crisis?, *Global and Planetary Change*, 175, 64–81, <https://doi.org/10.1016/j.gloplacha.2019.01.021>, 2019.
- Percival, L., Bond, D., Rakociński, M., Marynowski, L., Hood, A., Adatte, T., Spangenberg, J., and Föllmi, K.: Phosphorus-cycle disturbances during the Late Devonian anoxic events, *Global and Planetary Change*, 184, 103 070, <https://doi.org/10.1016/j.gloplacha.2019.103070>,
 560 2020.
- Pier, J. Q., Brisson, S. K., Beard, J. A., Hren, M. T., and Bush, A. M.: Accelerated mass extinction in an isolated biota during Late Devonian climate changes, *Scientific Reports*, 11, 24 366, <https://doi.org/https://doi.org/10.1038/s41598-021-03510-6>, 2021.
- Pohl, A., Ridgwell, A., Stockey, R. G., Thomazo, C., Keane, A., Vennin, E., and Scotese, C. R.: Continental configuration controls ocean oxygenation during the Phanerozoic, *Nature*, 608, 523–527, <https://doi.org/10.1038/s41586-022-05018-z>, 2022.



- 565 Racki, G.: A volcanic scenario for the Frasnian–Famennian major biotic crisis and other Late Devonian global changes: more answers than questions?, *Global and Planetary Change*, 189, 103–174, <https://doi.org/10.1016/j.gloplacha.2020.103174>, 2020.
- Rahmstorf, S., Crucifix, M., Ganopolski, A., Goosse, H., Kamenkovich, I., Knutti, R., Lohmann, G., Marsh, R., Mysak, L. A., Wang, Z., and Weaver, A. J.: Thermohaline circulation hysteresis: A model intercomparison, *Geophysical Research Letters*, 32, 2005GL023655, <https://doi.org/10.1029/2005GL023655>, 2005.
- 570 Reershemius, T. and Planavsky, N. J.: What controls the duration and intensity of ocean anoxic events in the Paleozoic and the Mesozoic?, *Earth-Science Reviews*, 221, 103–787, <https://doi.org/10.1016/j.earscirev.2021.103787>, 2021.
- Reinhard, C. T., Olson, S. L., Kirtland Turner, S., Pälike, C., Kanzaki, Y., and Ridgwell, A.: Oceanic and atmospheric methane cycling in the cGENIE Earth system model – release v0.9.14, *Geoscientific Model Development*, 13, 5687–5706, <https://doi.org/10.5194/gmd-13-5687-2020>, 2020.
- 575 Ridgwell, A.: A Mid Mesozoic Revolution in the regulation of ocean chemistry, *Marine Geology*, 217, 339–357, <https://doi.org/10.1016/j.margeo.2004.10.036>, 2005.
- Ridgwell, A.: derpycode/muffingen: v0.9.24, <https://doi.org/10.5281/zenodo.7545809>, 2023.
- Ridgwell, A. and Hargreaves, J.: Regulation of atmospheric CO₂ by deep-sea sediments in an Earth system model, *Global Biogeochemical Cycles*, 21, <https://doi.org/10.1029/2006GB002764>, 2007.
- 580 Ridgwell, A., Hargreaves, J., Edwards, N. R., Annan, J., Lenton, T. M., Marsh, R., Yool, A., and Watson, A.: Marine geochemical data assimilation in an efficient Earth System Model of global biogeochemical cycling, *Biogeosciences*, 4, 87–104, <https://doi.org/10.5194/bg-4-87-2007>, 2007.
- Ruttenberg, K.: The global phosphorus cycle, *Treatise on geochemistry*, 8, 585–643, <https://doi.org/10.1016/B0-08-043751-6/08153-6>, 2003.
- 585 Sablon, L., Maffre, P., Goddérès, Y., Valdes, P. J., Gérard, J., Huygh, J. J. C., Da Silva, A.-C., and Crucifix, M.: An Emulator-Based Modelling Framework for Studying Astronomical Controls on Ocean Anoxia with an Application on the Devonian, *EGUsphere*, 2025, 1–36, <https://doi.org/10.5194/egusphere-2025-1696>, 2025.
- Sarr, A., Donnadieu, Y., Laugié, M., Ladant, J., Suchéras-Marx, B., and Raison, F.: Ventilation Changes Drive Orbital-Scale Deoxygenation Trends in the Late Cretaceous Ocean, *Geophysical Research Letters*, 49, e2022GL099830, <https://doi.org/10.1029/2022GL099830>, 2022.
- 590 Schmidtko, S., Stramma, L., and Visbeck, M.: Decline in global oceanic oxygen content during the past five decades, *Nature*, 542, 335–339, <https://doi.org/10.1038/nature21399>, 2017.
- Scotese, C. R. and Wright, N.: PALEOMAP paleodigital elevation models (PaleoDEMS) for the Phanerozoic, Paleomap Proj, <https://doi.org/10.5281/zenodo.5348492>, 2018.
- Sharaf, S. G. and Boudnikova, N. A.: Secular variations of elements of the Earth’s orbit which influences the climates of the geological past, *Tr. Inst. Theor. Astron. Leningrad*, 11, 231–261, 1967.
- 595 Sharoni, S. and Halevy, I.: Rates of seafloor and continental weathering govern Phanerozoic marine phosphate levels, *Nature geoscience*, 16, 75–81, <https://doi.org/10.1038/s41561-022-01075-1>, 2023.
- Smart, M. S., Filippelli, G., Gilhooly, W. P., Ozaki, K., Reinhard, C. T., Marshall, J. E. A., and Whiteside, J. H.: The Expansion of Land Plants during the Late Devonian Contributed to the Marine Mass Extinction, *Communications Earth & Environment*, 4, 1–13, <https://doi.org/10.1038/s43247-023-01087-8>, 2023.
- 600



- Song, H., Song, H., Algeo, T. J., Tong, J., Romaniello, S. J., Zhu, Y., Chu, D., Gong, Y., and Anbar, A. D.: Uranium and carbon isotopes document global-ocean redox-productivity relationships linked to cooling during the Frasnian-Famennian mass extinction, *Geology*, 45, 887–890, <https://doi.org/10.1130/G39393.1>, 2017.
- Stappard, D. A., Wilson, J. D., Yool, A., and Tyrrell, T.: NutGENIE 1.0: nutrient cycle extensions to the cGENIE Earth system model to examine the long-term influence of nutrients on oceanic primary production, *EGUsphere*, 2025, 1–33, <https://doi.org/https://doi.org/10.5194/egusphere-2025-436>, 2025.
- Tostevin, R. and Mills, B. J.: Reconciling proxy records and models of Earth’s oxygenation during the Neoproterozoic and Palaeozoic, *Interface focus*, 10, 20190 137, <https://doi.org/https://doi.org/10.1098/rsfs.2019.0137>, 2020.
- Valdes, P. J., Armstrong, E., Badger, M. P. S., Bradshaw, C. D., Bragg, F., Crucifix, M., Davies-Barnard, T., Day, J. J., Farnsworth, A., Gordon, C., Hopcroft, P. O., Kennedy, A. T., Lord, N. S., Lunt, D. J., Marzocchi, A., Parry, L. M., Pope, V., Roberts, W. H. G., Stone, E. J., Tourte, G. J. L., and Williams, J. H. T.: The BRIDGE HadCM3 Family of Climate Models: HadCM3@Bristol v1.0, *Geoscientific Model Development*, 10, 3715–3743, <https://doi.org/10.5194/gmd-10-3715-2017>, 2017.
- Valdes, P. J., Scotese, C. R., and Lunt, D. J.: Deep ocean temperatures through time, *Climate of the Past Discussions*, 2020, 1–37, <https://doi.org/https://doi.org/10.5194/cp-17-1483-2021>, 2020.
- Van De Velde, S. J., Hülse, D., Reinhard, C. T., and Ridgwell, A.: Iron and sulfur cycling in the cGENIE.muffin Earth system model (v0.9.21), *Geoscientific Model Development*, 14, 2713–2745, <https://doi.org/10.5194/gmd-14-2713-2021>, 2021.
- Vervoort, P., Kirtland Turner, S., Rochholz, F., and Ridgwell, A.: Earth system model analysis of how astronomical forcing is imprinted onto the marine geological record: The role of the inorganic (carbonate) carbon cycle and feedbacks, *Paleoceanography and Paleoclimatology*, 39, e2023PA004 826, <https://doi.org/https://doi.org/10.1029/2023PA004826>, 2024.
- Wallmann, K.: Phosphorus imbalance in the global ocean?, *Global Biogeochemical Cycles*, 24, <https://doi.org/https://doi.org/10.1029/2009GB003643>, 2010.
- Ward, B. A., Wilson, J. D., Death, R. M., Monteiro, F. M., Yool, A., and Ridgwell, A.: EcoGENIE 1.0: plankton ecology in the cGENIE Earth system model, *Geoscientific Model Development*, 11, 4241–4267, <https://doi.org/10.5194/gmd-11-4241-2018>, 2018.
- Wichern, N. M. A., Bialik, O. M., Nohl, T., Percival, L. M. E., Becker, R. T., Kaskes, P., Claeys, P., and De Vleeschouwer, D.: Astronomically paced climate and carbon cycle feedbacks in the lead-up to the Late Devonian Kellwasser Crisis, *Climate of the Past*, 20, 415–448, <https://doi.org/10.5194/cp-20-415-2024>, 2024.
- Wilde, P. and Berry, W.: Destabilization of the oceanic density structure and its significance to marine “extinction” events, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 48, 143–162, [https://doi.org/https://doi.org/10.1016/0031-0182\(84\)90041-5](https://doi.org/https://doi.org/10.1016/0031-0182(84)90041-5), 1984.
- Zeebe, R. E. and Lantink, M. L.: A Secular Solar System Resonance that Disrupts the Dominant Cycle in Earth’s Orbital Eccentricity (g 2 ± g 5): Implications for Astrochronology, *The Astronomical Journal*, 167, 204, <https://doi.org/10.3847/1538-3881/ad32cf>, 2024.
- Šidlichovský, M. and Nesvorný, D.: Frequency Modified Fourier Transform and its Application to Asteroids, *The Dynamical Behaviour of our Planetary System*, p. 137–148, https://doi.org/10.1007/978-94-011-5510-6_9, 1997.




Figure 1. Late Devonian (370 Ma) bathymetry. Comparison between (a) the continental reconstruction from Scotese and Wright (2018), (b) the same reconstruction interpolated onto the HadCM3L model grid used in the emulator and (c) as per (b) on the cGENIE grid.




Figure 2. Carbonate platform location. Carbonate platform (designed as 'reef' in the cGENIE model) locations (in red) as represented in cGENIE based on Kiessling et al. (2003).




Figure 3. Global response to astronomical forcing. Global time series of (a) $p\text{CO}_2$, ocean (b) $[\text{PO}_4]$ and (c) $[\text{O}_2]$. The black curve represents the simulation in which all astronomical parameters evolve. The blue curve represents the simulation with only obliquity varying, while the red curve corresponds to the simulation where only eccentricity and precession evolve. (d) The astronomical solution used in the simulations, with eccentricity in dotted red, climatic precession in red and obliquity in blue. Here, e denotes the eccentricity, and ϖ represents the longitude of perihelion.




Figure 4. Regional variation of $[O_2]$ and anoxia. (a) Definition of the five regions. (b) Watersheds based on the nearest-neighbour continental runoff routing scheme. The colour of the continental grid cell indicates in which region the weathering flux of PO_4 computed by the emulator of Sablon et al. (2025) is routed. The hashed areas represent the regions as defined in cGENIE and are the same as in panel (a). The discrepancies between the watersheds and the regions arise from the different grid resolution of cGENIE and HadCM3. (c) Time series of average oceanic $[O_2]$ anomalies and anoxic volume anomalies (in %) for each region. These quantities were computed over the full extent of each region, from the second vertical layer down to approximately 1000 m depth, excluding the surface layer as it is in equilibrium with the atmosphere. The anoxic volume is shown in a darker colour, and the anomaly is expressed as a difference of %. The anoxic threshold is set at $6.5 \mu\text{mol.kg}^{-1}$ after Sarr et al. (2022). For both $[O_2]$ and anoxic volume, the y-axis scales are identical across all regional panels to allow direct comparison. As in Fig. 3, the bottom panel displays the astronomical forcing, but shown in black for clarity.




Figure 5. Normalized Fourier power spectrum of the regional $[O_2]$ time series. Vertical dashed lines indicate the characteristic periods of the astronomical parameters: 19 kyr for precession, 30 kyr for obliquity, and 100 kyr and 405 kyr for eccentricity.




Figure 6. Link between continental PO₄ weathering flux and regional [PO₄]. Displayed are correlations between regional [PO₄] and their corresponding regional weathering fluxes. A negative time lag is applied in the figures on the right, where the regression is computed, and is absent from the figures on the left. Regions SA, WL, and LG have all p-values smaller than 10^{-3} , while region Si has a p-value of 0.35. Region NP is excluded from the analysis due to the absence of direct continental PO₄ weathering.




Figure 7. Surface ocean circulation. Mean horizontal ocean velocity (m.s^{-1}) averaged over the upper 300 m. Arrow direction indicates flow orientation, while arrow length is proportional to velocity magnitude (also represented with background shading).




Figure A1. Temporal variance. Normalized temporal variance of (a) $[O_2]$ and (b) $[PO_4]$ averaged over the upper 1000 m. Dashed coloured lines indicate regional boundaries, as defined in Fig. 4.