

Technical note: How well do evapotranspiration partitioning approaches perform in moss-covered wetlands?

Yi Wang¹, Richard M. Petrone¹, Lei Zhang²

¹Hydrometeorology Research Group, University of Waterloo, Waterloo, Ontario, Canada

²School of Water Conservation, North China University of Water Resources and Electric Power, Zhengzhou, Henan, China

Correspondence to: Yi Wang (yi.wang1@uwaterloo.ca)

Abstract. Evapotranspiration (ET) is the dominant hydrologic flux in wetlands, and partitioning into transpiration (T) and evaporation (E) is essential for understanding water and carbon dynamics, guiding sustainable water management practices, and predicting responses to climate change in these systems. However, the presence of moss layers and the dynamic hydrological condition in many wetlands challenges the assumptions of commonly used partitioning methods. This study evaluates the performance of nine eddy covariance (EC)-based ET partitioning approaches. This study evaluates ten eddy covariance (EC)-based ET partitioning approaches representing three methodological groups, namely high-frequency EC-based, ecosystem carbon-water coupling, and machine learning approaches, across four moss-covered wetlands in the boreal region and the Canadian Rocky Mountains. ET partitioning estimates were evaluated against independent measurement-based T:ET estimates derived from flux chambers, micro-lysimeters, sap flow sensors, and EC measurements, and against the ensemble mean of all ET partitioning approaches, across multiple moss-covered wetland sites located in boreal and the Canadian Rocky Mountains. The partitioning results from each approach were compared against independent measurement-based estimates, which were obtained using flux chamber, micro-lysimeters, sap flow sensors, and EC systems. Results showed that none of the evaluated approaches consistently outperformed the others across sites or under both flooded and non-flooded conditions, and no methodological group demonstrated a clear overall advantage. ET partitioning performance is likely influenced by the combined effects of multiple environmental and methodological factors, including moss cover, flooding conditions, canopy structure, ecosystem heterogeneity, EC measurement configuration and data quality, and the assumptions underlying each approach. In addition, the evaluation results are also influenced by the choice of reference dataset, which highlights the uncertainty associated with benchmarking ET partitioning methods. These findings suggest that applying multiple ET partitioning approaches concurrently, whenever possible, can help identify sources of uncertainty and potential errors. The study also highlights the need to develop wetland-specific ET partitioning approaches that better represent moss-mediated carbon uptake, dynamic carbon-water coupling, and transitions between unsaturated and flooded conditions. none of the evaluated methods provided both accurate and precise estimates of ET partitioning (T:ET), and no single method emerged as the most suitable for studied ecosystems. Despite this, the general agreement between modelled and measured T:ET values indicates that many of these approaches still provide valuable insights. Applying multiple methods concurrently is recommended, where possible, to enhance confidence in partitioning results. For researchers with access to high frequency EC data, priority should be given to high frequency EC-based methods due to their more consistent performance across sites. The findings

also highlight the limitations of current partitioning approaches under evaporation-dominated conditions, and underscore the need to examine the mechanistic role of mosses, as well as to improve how optimal stomatal conductance theory is conceptualized and implemented in model formulations.

1. Introduction

Evapotranspiration (ET) represents the total water flux from terrestrial ecosystems to the atmosphere through vegetation transpiration (T) and direct evaporation (E) from canopy and ground surfaces. Distinguishing the contributions of T and E is important for understanding the terrestrial carbon cycle, as T is directly related to vegetation carbon assimilation processes (Jarvis and Mcnaughton, 1986; Farquhar et al., 1980; Farquhar and Sharkey, 1982). Despite its importance, there is a broad consensus within the biogeosciences community that accurately quantifying T and E, whether through direct measurements or computational methods, remains a challenge (Kool et al., 2014; Stoy et al., 2019; Nelson et al., 2020; Zahn et al., 2024). Briefly, directly quantifying T and E involves a combination of multiple field-based measurements, such as stable isotope analysis, lysimeters, porometers, flux chambers, sap flow measurements, and eddy covariance systems (Kool et al., 2014). These methods can be labor-intensive, require site-specific instrumentation, and are often constrained in both spatial (e.g., plot-scale) and temporal (e.g., depending on measurement frequency) resolutions (Albert-Saiz et al., 2025). In contrast, modelling approaches, especially the eddy covariance (EC)-based ET partitioning methods to partitioning ET into E and T at the ecosystem level, may be convenient and enable the exploration of temporal variability for estimating E and T at the ecosystem scale. However, they rely on assumptions that are ecosystem-dependent, require accurate parameterization, and model validation is not always available due to the lack of on-site measurements (Eichelmann et al., 2022; Zhou et al., 2016; Stoy et al., 2019). As reviewed by Stoy et al. (2019), these limitations highlight the need for ecosystem-scale experiments using multiple methods to measure E and T to understand their distinct responses to climate variability and change across diverse ecosystems and also to evaluate and refine the assumptions underlying ET partitioning approaches (Stoy et al., 2019).

Currently, research specifically addressing ET partitioning within wetland ecosystems remains notably scarce, even though wetlands, especially peatlands, are critical carbon sinks and play a critical role in regulating regional hydrology and global climate (Mitra et al., 2005; Leifeld and Menichetti, 2018; Strack et al., 2022; Frolking et al., 2006). Consequently, there is limited field data available to develop and evaluate reliable ET partitioning approaches for wetland ecosystems. Moreover, most of the existing studies have focused on hydroperiods where E mostly comes from standing water or saturated soils (Xu et al., 2011; Allen et al., 2017; Zhang et al., 2018; Hussey and Odum, 1992; Burba et al., 1999; Goulden et al., 2007; Kiniry et al., 2023; Bijoor et al., 2011; Eichelmann et al., 2022). However, it should be noted that many wetlands are not consistently in a saturated state (e.g., Jacobs et al. (2002), Wu et al. (2010), Kettridge et al. (2017), and Streich and Westbrook (2020)), which raises concerns about whether methods developed under flooded conditions can be applied reliably to non-flooded conditions or to estimate ET partitioning under changing environmental conditions. Furthermore, current studies report varying results regarding the dominance of T or E in wetland ET and their controlling environmental factors, which suggests that ET partitioning in wetland

ecosystems is regulated by both structural properties including vegetation composition, density, and areas of open water, and environmental conditions such as climatic variables and water table (Goulden et al., 2007; Kiniry et al., 2023; Warren et al., 2018; Allen et al., 2017; Eichelmann et al., 2022; Admiral and Lafleur, 2007). Given that many ET partitioning approaches rely on assumptions about T dominance or dominant environmental controls, it remains uncertain if there is a universally reliable ET partitioning method for wetland ecosystems.

Despite the limited research focused specifically on wetlands, ET partitioning approaches are relatively abundant, particularly those based on EC data, as EC systems have been widely deployed across diverse ecosystems. Several of these EC-based methods have been tested across a range of ecosystems, including forests, grasslands, woody savannas, and croplands, and have shown broad applicability, not restricted to water-limited ecosystems (Yu et al., 2022; Nelson et al., 2020; Zhou et al., 2016; Li et al., 2019; Zahn et al., 2024). However, most of these methods have not been evaluated in wetlands. In fact, simply not restricted to water-limited ecosystems does not guarantee that these methods are suitable for wetlands, as the unique characteristics of wetlands, particularly moss-covered ground surfaces in many peatlands, may not only influence the evaporative response of the wetlands (Speranskaya et al., 2024), but also violate key assumptions on which these ET partitioning approaches ~~these methods~~ primarily rely. Specifically, mosses can photosynthesize and contribute to ecosystem carbon sequestration (Pacheco-Cancino et al., 2024; Street et al., 2013; Badorek et al., 2011; Kokkonen et al., 2022), but they do not contribute to transpiration fluxes. This decoupling may violate the assumption in some methods that photosynthesis and transpiration share identical sources and sinks. Moreover, most mosses cannot actively regulate water transport and water fluxes; instead, they rely on passive capillary rise to transport water (Proctor, 1982). When free capillary rise is limited by low soil-water pressures, the water potential of moss cells can rapidly equilibrate with the surrounding air, reducing both evaporation and photosynthetic activity, even when atmospheric demand for evaporation remains high (Goetz and Price, 2015; Mccarter and Price, 2012; Proctor, 1982; Kettridge and Waddington, 2013; Ketcheson and Price, 2013). In addition, mosses can absorb water directly from their surroundings through their leaves and stems, and store water equivalent to 300 to 500% of their dry weight for extended periods (Rouse, 2000; Bayfield, 1973). This remarkable water storage capacity, combined with their ability to rapidly equilibrate their water content with ambient humidity, helps conserve moisture in deeper soil layers during dry periods (Ketcheson and Price, 2013; Kettridge and Waddington, 2013), and cools the soil surface (Chen et al., 2019). These unique behaviors of mosses may therefore violate assumptions about key environmental controls on T or E fluxes, as well as the assumed relationships between carbon uptakes and water exchange used by some methods to separate T and E fluxes.

Although the presence of mosses may violate some of the assumptions used in EC-based ET partitioning approaches, it is uncertain whether these violations render the methods completely unsuitable for wetland research, or whether these methods can still offer insights into the dynamics of T and E fluxes. Intuitively, the extent to which these violations affect the accuracy of EC-based ET partitioning approaches depends on the relative contribution of mosses to wetland gross primary production (GPP) compared to vascular plants at a given site. Although multi-site analyses have shown a positive correlation between peatland carbon assimilation and moss cover percentage (Strack et al., 2016; Pacheco-Cancino et al., 2024), quantitative data on moss contributions to ecosystem-level GPP remains

extremely scarce. Limited evidence suggests that mosses can play a considerable role: for instance, a field study in a raised mire complex in southern Finland reported moss contributions to cumulative community-level photosynthesis of a contribution of 1.5%, 35.2% and 41.2% in the undrained area in rich fen, poor fen, and bog, respectively (Kokkonen et al., 2022). Similarly, in a drained forested peatland dominated by moss on the forest floor, CO₂ assimilation by the forest floor represented 20 to 30% of the forest's total CO₂ uptake (Badorek et al., 2011). But ~~These findings~~ also indicate that while mosses can substantially contribute to ecosystem GPP, vascular plants generally dominate, accounting for over 50% of the ecosystem GPP. Consequently, the presence of mosses may not markedly compromise the performance of EC-based ET partitioning methods. Regardless of these considerations, there remains a need to evaluate the performance of ET partitioning approaches in moss-covered wetlands. ~~Currently, there has been no evaluation of these approaches in wetland ecosystems. Therefore, the objectives of this study are to assess the performance of existing EC-based ET partitioning methods in moss-covered wetlands.~~

The ET partitioning approaches discussed above, whose fundamental assumptions may be violated by the presence of moss cover, are all physically based methods. In contrast, machine learning (ML)-based ET partitioning methods have recently been developed for wetland ecosystems, and are based on assumptions that are, in principle, unaffected by the presence of mosses. For example, the approach proposed by Eichelmann et al. (2022) and subsequently extended by Stapleton et al. (2022) trained ML models to predict E using nighttime EC measurements under the assumption that nighttime T is negligible. T is then estimated as the difference between measured ET and the predicted E. These ML models performed well in four freshwater marshes in northern California. Their predicted evaporation agreed closely with EC-derived ET during flooding periods and winter, when T was also assumed to be negligible (Eichelmann et al., 2022)(Stapleton et al., 2022)(Stapleton et al., 2022; Eichelmann et al., 2022). In addition, Eichelmann et al. (2022) showed good agreement between model-estimated T and leaf-level T measurements of the two dominant plant species, tule and cattail.

Although the presence of mosses does not violate the assumption underlying these ML models, the assumption of negligible nighttime T may not be universally applicable across wetland ecosystems with different vegetation communities. Increasing evidence indicates that nighttime T occurs in a wide range of grasses, shrubs and trees (Eichelmann et al., 2022)(Daley and Phillips, 2006; Caird et al., 2007; Fisher et al., 2007), where it typically accounts for 10–30% of the total daily transpiration (Novick et al., 2009). Furthermore, nighttime EC measurements are generally subject to greater uncertainty because of weak atmospheric turbulence and reduced mixing within the nocturnal boundary layer (Baldocchi, 2003; Aubinet et al., 2012). Although nighttime T may indeed be small relative to E in many wetlands as suggested in Eichelmann et al. (2022), this assumption has not been directly validated. Moreover, these ML approaches have so far been evaluated only in four freshwater marshes, and their applicability to other wetland types remains uncertain. Despite these limitations, ML approaches have emerged as powerful tools in Earth system science because of their ability to identify complex, nonlinear relationships between environmental drivers and target variables without requiring explicit process-based formulations (Reichstein et al., 2019). Therefore, although this study primarily focuses on ET partitioning approaches whose underlying assumptions may be challenged

by the presence of moss cover, ML-based approaches are also included to assess their robustness and general applicability in wetland ecosystems.

The overall objective of this study is to evaluate the performance of existing EC-based ET partitioning methods in moss-covered wetlands. ~~Specifically, Currently, there has been no evaluation of these approaches in wetland ecosystems. Therefore, the objectives of this study are to assess the performance of existing EC-based ET partitioning methods in moss-covered wetlands.~~ we address the following research questions: (1) Is there any ET partitioning approach that consistently outperforms the others across different wetland sites? (2) Do these approaches perform well under both flooded and non-flooded conditions? (3) Do ML-based approaches outperform physically based approaches? And (4) What factors influence the performance of ET partitioning approaches in the studied wetlands?

2. Methods

2.1 Overview of EC data-based ET partitioning approaches considered

ET partitioning approaches selected for this study were chosen based on three criteria: (1) they rely primarily on EC-based CO₂ and H₂O fluxes, (2) focus on the ecosystem level with daily or finer temporal resolution, are not restricted to water-limited conditions, and (3) have publicly available formulations or programming codes. ~~Based on these criteria~~Consequently, this study evaluates ~~ten~~nine methods ~~were selected for evaluation~~ (Table 1). ~~Several existing approaches were excluded, including -and excludes the method of s-developed by~~ Wei et al. (2017), ~~which was not developed for daily or finer temporal scales; the methods of,~~ Scott and Biederman (2017) ~~and~~ Reich et al. (2024), ~~which were developed for water-limited ecosystems;~~ Rigden et al. (2018), ~~which primarily relies on weather data rather than EC-based measurements; and~~ Eichelmann et al. (2022), ~~for which no source code is available, -and Reich et al. (2024)~~Eichelmann et al. (2022)

~~To the best of the authors' knowledge, the ET partitioning methods evaluated in this study cover most of the existing EC-based approaches. Based on the type of EC data they utilize, either high-frequency measurements or fluxes aggregated to a half-hourly timestep, and the partitioning formulations they employed, either ecosystem carbon-water coupling or machine learning based, these approaches are grouped into three categories: (1) high-frequency EC data-based methods (Group 1), (2) ecosystem carbon-water coupling methods (Group 2), and (3) machine learning (ML) methods (Group 3). These approaches and their required data and limitations for application in moss-covered wetland ecosystems are summarized in Table 1. Note that the Transpiration Estimation Algorithm (TEA) is based on both the ecosystem carbon-water coupling and the Random Forest model which is an ML algorithm. Consequently, TEA is classified as both a Group 2 and Group 3 approach.~~

~~It should also be noted that -Note that~~Group 3 does not represent a collection of individual ET partitioning methods, but rather a framework developed by Stapleton et al. (2022) that comprises seven ML algorithms. The framework

Formatted: Font: (Default) Times New Roman, 10 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 10 pt, English (United States)

Formatted: English (United States)

identifies the best performing machine learning algorithm from the candidate models, determines the optimal combination of predictor variables, and ranks these variables according to their importance to predictive accuracy. It extends the ET partitioning approach proposed by Eichelmann et al. (2022), who applied an artificial neural network (ANNs) to partition ET. Although the ANN model and its source code are not publicly available, the underlying assumptions governing ET partitioning are consistent between the two studies.

The seven candidate ML algorithms included in the Stapleton et al. (2022) framework are Linear Regression, Ridge Regression, K-nearest Neighbors, Decision Trees, Gradient Boosting Decision Trees, LightGBM and XGBoost (Stapleton et al., 2022). Following the procedure described by Stapleton et al. (2022), only the ET partitioning predictions generated by the best-performing model identified by this framework were retained for evaluation in this study. Details of the performance of each candidate ML model and the rankings of the importance of the predictor variables are provided in Section S1 of the Supplementary Material.

Eichelmann et al. (2022) introduced an artificial neural networks (ANNs)-based ET partitioning methods for flooded ecosystems. Eichelmann et al. (2022), its code is not publicly available and was therefore not evaluated here. Stapleton et al. (2022). To the best of the authors' knowledge, the ET partitioning methods evaluated in this study cover most of the existing EC-based approaches. Based on the type of EC data they utilize, either high-frequency or values aggregated at a half-hourly timestep, these approaches are grouped into two categories: the high-frequency EC data-based methods (Group 1), and ecosystem carbon-water coupling methods (Group 2). These approaches and their required data and limitations for application in moss-covered wetland ecosystems are summarized in Table 1.

Table 1. Overview of data requirements and potential limitations of EC-based ET partitioning methods in moss-covered wetlands. Group 1 includes methods using high-frequency EC data, while Group 2 and Group 3 includes methods using ecosystem-level EC data aggregated at specific timesteps.

Method	Group	Required Data	Examples of Potential Issues When Applied to Moss-covered Wetlands	References
Flux Variance Similarity (FVS)	1	High-frequency c , q , u , v , w , and WUE_{canopy}	The assumption of identical source/sink for transpiration and photosynthesis is invalidated by moss photosynthesis.	Scanlon and Kustas (2010)
Conditional Eddy Accumulation method (CEA)	1	High-frequency c , q , u , v , and w	The uneven and open canopy often found in wetlands, together with moss photosynthesis, can blur ground and vegetation signals used for flux partitioning.	Businger and Oncley (1990) and Zahn et al. (2024)
Modified Relaxed-Eddy Accumulation method (MREA)	1	High-frequency c , q , u , v , and w	The presence of short or multi-layered canopies, often found in wetlands, along with moss photosynthesis and respiration in the understory, may violate the similarity of turbulent transport of non-stomatal components from the soil.	Thomas et al. (2008) and Zahn et al. (2024)
Conditional Eddy Covariance method (CEC)	1	High-frequency c , q , u , v , and w	Same as CEA	Zahn et al. (2022)
Underlying water-use efficiency method (uWUE)	2	GPP, ET, and VPD	The assumption that T dominates ET during periods of high GPP/ET may not hold in E-dominated wetlands; and carbon assimilation is assumed to occur only from leaf-level photosynthesis.	Zhou et al. (2016)
Pérez-Priego method (PP)	2	GPP, C_a , P_a , T_{air} , VPD, LE, H, U_{star} , WS, R_g , Q, and Z	The big leaf representation used to estimate optimal stomatal conductance from EC-based P_h does not account for moss photosynthesis; and the	Pérez-Priego et al. (2018)

			complexity of the optimization procedure can sometimes fail in reaching a numerical solution.	
Transpiration Estimation Algorithm (TEA)	2 and 3	GPP, RH, R _g , R _{g,pot} , T _{air} , VPD, Precip, WS, and year	The assumption that T:ET ~1 during periods with minimal surface moisture conditions may not hold in E-dominated wetlands.	Nelson et al. (2018)
Conductance Partitioning method (CP)	2	GPP, ET, VPD, and θ	The assumption that surface conductance is not sensitive to VPD may be invalidated by moss responses to ambient humidity.	Li et al. (2019)
Carbon-water relationship and stomatal conductance model (CWSC)	2	GPP, ET, VPD, C _a , P _a and g _i	Carbon assimilation is assumed to occur only from leaf-level photosynthesis; and publicly available g _i dataset by plant functional types (PFTs) are lacking for wetland ecosystems.	Yu et al. (2022)
Machine Learning Algorithms (ML)	3	All EC, micrometeorological, and environmental data measured at the study sites	The assumption of negligible nighttime T may introduce prediction biases, and nighttime EC measurements are generally less reliable.	Stapleton et al. (2022)

Formatted Table

Formatted: English (Canada)

Note. c, high-frequency CO₂ concentration; q, high-frequency H₂O concentration; u, high-frequency streamwise wind speed; v, high frequency cross-stream wind speed; w, high frequency vertical wind speed; WUE_{canopy}, canopy-level water-use efficiency; GPP, ecosystem gross primary production; ET, ecosystem evapotranspiration; VPD, vapor pressure deficit; C_a: mean atmospheric CO₂ concentration; P_a, atmospheric pressure; T_{air}, air temperature; LE, latent heat flux; H, sensible heat flux; U_{star}, friction velocity; WS, mean wind speed; R_g, incoming shortwave radiation; Q, photosynthetic active radiation; Z, altitude; RH: relative humidity; R_{g,pot}, daily potential radiation; Precip, precipitation; θ , soil moisture; g_i, a parameter in the optimal stomatal conductance model. Some required variables differ from those listed in the original publications and instead reflect the data inputs used in the publicly available codes provided by the authors of each method. Specifically, the FVS, CEA, MREA and CEC methods were applied using the ET partitioning code developed by Zahn (2024); the PP method by Pérez-Priego and Wutzler (2018); the uWUE and TEA methods by Nelson (2020); ~~and the CWSC method by Jiang and Yu (2021); and the ML method by~~ Stapleton (2022). Please refer to the corresponding references for further details.

2.2 Study sites

The best way to evaluate the ET partitioning approaches is by comparing their outputs to direct field measurements. However, to the best of the authors' knowledge, no publicly available datasets exist for wetland ET and its components. Therefore, we compiled data from four sites where field measurements of ET and its components had previously been conducted.

The first three sites, Sibbald, Burstall, and Bonsai, are located in the Canadian Rocky Mountains, where field measurements were conducted during the 2021 growing season. The fourth site, Poplar, is situated in the western boreal plain near Fort McMurray, Alberta, Canada, and its field measurements were conducted during the 2013 growing season. Among the four sites, only Sibbald and Poplar are peatlands. The study sites have been extensively described in detail in previous publications (Wang and Petrone, 2022; Wang et al., 2023; Wang, 2025; Gabrielli, 2016; Hathaway et al., 2022; Hrach et al., 2021; Streich and Westbrook, 2020; Morison et al., 2021); therefore, only the characteristics relevant to this study are briefly provided here. The geographic coordinates, wetland type, elevation, meteorological conditions during the study period, soil properties, water table depth, vegetation composition, and instrumentation used to measure evapotranspiration and its components are summarized in Table 2.

These sites are covered by mosses. At Sibbald, the moss layer is approximately 5 mm thick and includes species such as *Aulacomnium palustre* (Hedw.) Schwaegr., *Bryum pseudotriquetrum* (Hedw.) Gaertn. et. al., *Calliergon giganteum*

(Schimp.) Kindb., *Calliergon richardsonii* (Mitt.) Kindb. in Warnst., *Campylium stellatum* (Hedw.) C. Jens., *Hypnum lindbergii* (Mitt.), *Plagiomnium ellipticum* (Brid.) T. Kop., *Pohlia nutans* (Hedw.) Lindb., and *Warnstorfia fluitans* (Hedw.) Loeske (Lei, 2021). At Burstall, the moss layer, about 1 cm thick, is dominated by peat moss (*Sphagnum* spp.) and feather moss (*Eurhynchium* spp.). At Bonsai, the moss cover consists primarily of brown moss (*Tetradontium* spp.) with a thickness of up to 5 mm. At poplar, the moss layer is around 2.5 cm thick (Goetz and Price, 2015), dominated by *Tomenthypnum nitens* (Hedw.) Loeske and *Aulacomnium palustre* (Hedw.). It is important to clarify that the moss layer in this study refers mainly to the photosynthetically active (green) components of the mosses, including their shoots and capitula.

Although all sites are moss-covered wetlands, they differ considerably in meteorological conditions, water table levels, vegetation composition, and canopy structure (Table 2 and Fig. ~~ure~~ S3S4). During the study period, the growing season LAI was 4.56, 1.64, 1.33 and 2.82 for Sibbald, Burstall, Bonsai and Poplar, respectively, with the value at Poplar representing the tree canopy only. The average canopy height was 1.6 m for Sibbald and Burstall, 0.8 m for Bonsai and 4.2 m for Poplar. Notably, Sibbald and Bonsai remained under non-flooded conditions throughout the study period, whereas Burstall transitioned to flooded conditions after August 17, 2021. In contrast, Poplar remained flooded for almost entire study period. The water table dynamics and meteorological conditions for each study site during the respective study period are shown in Fig. S4.

Table 2. Summary of site characteristics, including geographic coordinates, wetland type, altitude, meteorological conditions during the study period, soil properties, water table depth, vegetation composition, and instrumentation used for measuring evapotranspiration and its components, and the study period.

Site	Wetland Class	Latitude	Longitude	Elevation (m)	Temp.* (°C)	Precip.* (mm)	Soil (0-15 cm)	WT depth* (m)	Vascular vegetation	Instruments	Study Period
Sibbald	Fen	51.06°N	114.87°W	1,490	14.0	137.7	Peat	0.5 to 1.24 to -0.56	The overstory is characterized by clusters of <i>Salix</i> spp., while the understory consists of dense <i>Carex</i> spp.	Micro-lysimeter, porometer and eddy covariance system	<u>July 1 to August 31, 2021</u>
Burstall	Marsh	50.78°N	115.34°W	1,900	11.6	32.2	Peat	-0.1 <u>5</u> to 0.1	The overstory is characterized by sparse <i>Salix</i> spp., while the understory consists of dense <i>Carex</i> spp.	Micro-lysimeter, porometer and eddy covariance system	<u>July 1 to August 31, 2021</u>

Formatted Table

Bonsai	Wet meadow	50.82°N	115.21°W	2,100	11.7	128.3	Silt and clay	2< ٠٠٠	The overstory consists of sparsely distributed species such as <i>Salix</i> spp., <i>Larix</i> spp., and <i>Abies</i> spp., while the understory is abundant with species including <i>Erigeron</i> spp., <i>parnassia</i> spp., and <i>Equisetum</i> spp.	Micro-lysimeter, porometer and eddy covariance system	July 27 to August 31, 2021
Poplar	Fen	56.56°N	111.33°W	322	18.4	320.8	Peat	-0.13 to 0.169	The overstory consists of a dense cover of <i>Larix laricina</i> , with sparse occurrences of <i>Picea mariana</i> , while the understory consists of a dense shrub layer including <i>Betula pumila</i> , <i>Equisetum fluviatile</i> , <i>Smilacena trifolia</i> , <i>Carex prairea</i> , and <i>Carex diandra</i>	Flux chamber, sap flow system, and eddy covariance system	May 18 to August 21, 2013

Formatted: Font: Bold

242 Note. Elevation is the elevation above sea level (in meters). Temp* represents the mean air temperature during the
243 measurement period, and Precip.* is total precipitation during the same period. Soil describes the dominant soil type
244 within the top 15 cm where micro-lysimeters were installed. WT depth* represents the mean water table position
245 relative to the ground surface during the study period, calculated from all available monitoring wells. Negative values
246 indicate that the water table was below the ground surface, whereas positive values indicate flooded conditions (i.e.,
247 the water table was above the ground surface)represents the position of the water table relative to the ground surface,
248 with positive values indicating that the water table is lower than the ground surface and negative values indicating
249 flooded conditions. Instruments summarize the main equipment used to measure or estimate E, T and ET at each site.
250 The study period for Bonsai begins on 27 July because the EC system malfunctioned before that date.

2.3 Field measurements of evaporation, transpiration and evapotranspiration

Ecosystem-scale water and carbon fluxes at all sites were measured by eddy covariance systems operating throughout the respective measurement periods. The ratio of EC measurement height to canopy height (z/h) is 1.88 at Sibbald, 1.25 at Burstall, 3.75 at Bonsai, and 1.55 at Poplar. EC data processing followed common FLUXNET protocols (Kaimal and Finnigan, 1994; Foken and Leclerc, 2004; Aubinet et al., 2012; Burba et al., 2012; Leuning and Judd, 1996; Webb et al., 1980; Pastorello et al., 2020). Latent heat flux (LE) was gap-filled using the MDS method (LE_F_MDS). Note that GPP is a required input for many ET partitioning approaches. In this study, it was estimated from EC-measured net ecosystem exchange (NEE_VUT_USTAR50) using the night-time flux partitioning method described by Reichstein et al. (2005).

In addition to EC measurements, evaporation at Sibbald, Burstall and Bonsai was measured using micro-lysimeters on six separate occasions during July and August in 2021, with each measurement period lasting mostly 2 to 4 days (Wang et al., 2023; Wang and Petrone, 2022). Stomatal conductance of the dominant vascular vegetation (Table 2) at these three sites were measured by leaf porometer (Wang, 2025). At the Poplar site, transparent flux chambers and sap flow systems were used to measure ground evaporation and canopy transpiration during the 2013 growing season (Gabrielli, 2016). These measurements were conducted within the EC footprint of each site. Detailed descriptions of site characteristics, field measurements, instrumentation, and the upscaling of plot-level data to estimate ecosystem-level daily transpiration-to-evapotranspiration ratios (T:ET) are provided in Wang (2025) and Gabrielli (2016). The errors and uncertainties of the measurements of E, T and ET are summarized in Table S4.

2.4 Reconstruction of daily ecosystem-scale measurement-based ET partitioning time series

~~, and are not repeated here.~~ For the Sibbald, Burstall, and Bonsai sites, the measurement-based T:ET time series are derived from a site-calibrated and validated Shuttleworth–Wallace (S–W) model rather than direct averaging of ground evaporation measurements. This approach was necessary because micro-lysimeters measure ground evaporation over multi-day intervals, whereas eddy covariance measurements of ET at the study sites are subject to intermittent data gaps. Using calibrated S–W model enabled the reconstruction of a continuous daily T:ET time series and characterization of its temporal variation.

The detailed calibration of the S–W model for each study site, together with the development of a surface resistance formulation that explicitly accounts for the effects of moss and litter on ground evaporation and its incorporation into the S–W model, are described in previous publications (Wang et al., 2023; Wang, 2025). The performance of the calibrated S–W model is further summarized in Section S3 of the Supplementary Material. Overall, the model accurately reproduced both latent heat flux (LE) and ground evaporation across the study sites. Because the S–W model is based on the surface energy balance, its predicted LE was evaluated against the energy balance-corrected EC measurements (LE_CORR). Across all sites, the relationship between modelled and measured LE was given by $LE_{\text{modelled}} = 0.97 \times LE_{\text{CORR}} + 2.23 \text{ W m}^{-2}$ ($R^2 = 0.83$). Agreement between modelled and measured ground evaporation (W m^{-2} , expressed as λE) was similarly strong, with $\lambda E_{\text{modelled}} = 0.91 \times \lambda E_{\text{measured}} - 4.46 \text{ W m}^{-2}$

Formatted: Level 2

($R^2 = 0.85$). These results support the use of the S–W model-reconstructed T:ET time series as the measurement-based, ecosystem-scale daily T:ET for the three study sites.

At the Poplar site, the measurement-based T:ET time series was constructed differently from the other three sites, as this site includes direct eddy-covariance ET measurements and upscaled sap flow measurements for dominant tree and understory species, informed by forest inventory data and LiDAR-based vegetation classification, as well as ground-based chamber flux measurements (Gabrielli, 2016). Ground evaporation was calculated as the residual (ET minus upscaled site transpiration (T)) and evaluated against chamber measurements dominated by moss. This comparison yielded a regression slope of 0.96, an intercept of 0.08 mm h^{-1} , and an R^2 of 0.26. Because both T and ET were measured at the same timestep, whereas chamber measurements are instantaneous and inherently variable, we used the sap-flow-derived transpiration together with eddy-covariance ET to construct the Poplar T:ET time series. We acknowledge the relatively low R^2 , which reflects the scatter in the chamber measurements, but the near-unity slope and small intercept indicate good overall agreement. On this basis, we consider the reconstructed T:ET time series to be a reasonable representation for this study site.

Formatted: Font: 10 pt, English (United States)

2.5.4 Data quality control procedure

Given the measurement techniques described above, it is important to clarify that evaporation measurements in this study represent only ground evaporation. However, during rainy periods, evaporation can occur from leaf surfaces. Excluding this canopy evaporation may introduce bias in estimating ecosystem-scale ET partitioning into E and T. To minimize this potential bias, all methods except for the CP method exclude rainy days and 1 to 2 days afterward, following the precipitation and potential evapotranspiration (PET)-based criteria described in Zhou et al. (2015) and Yu et al. (2022). Specifically, if $\text{PET} < \text{Precip} < 2\text{PET}$, 1 day after a rainy day was excluded, if $\text{Precip} > 2\text{PET}$, 2 days were excluded. PET was estimated using the Priestley-Taylor equation (Priestley and Taylor, 1972). For the CP method, data screening follows its original protocol, which removes data during rainy hours and the 6 hours following rainfall (Li et al., 2019).

In addition to excluding rainy periods, all methods were prescreened to include only data from the growing season, during daytime hours, and under conditions where net radiation, GPP, LE, H and VPD were greater than zero. Half-hourly EC fluxes data with poor quality (i.e., the quality flag (QC) > 1) were removed. For the CP method, additional filters were applied by excluding data with RH above 0.95, and by retaining only those periods where H exceeded 5 W m^{-2} and R_g exceeded 50 W m^{-2} (Li et al., 2019).

When aggregating half-hourly ET partitioning results to daily values, only days with more than 10 valid half-hourly values were included, following the procedure in Yu et al. (2022). Additionally, any half-hourly T:ET values that were negative or greater than 1 were excluded prior to daily aggregation.

2.6.5 Evaluation of ET partitioning approaches

Direct measurements of ET components offer an independent means to evaluate the performance of ET partitioning approaches. However, this study does not focus on comparing exact magnitudes of T:ET between measurement-based estimates and those derived from ET partitioning approaches. This is because most field measurements were conducted at the plot level, and upscaling to the ecosystem level introduces uncertainties, as a wetland is a mosaic of diverse habitats with varying extents of vegetation cover, open water, and exposed bare soil (Drexler et al., 2004), which makes spatial upscaling very challenging. In peatlands, such complexity is further increased by micro-topographical features, such as hummocks (local high points) and hollows (local low points), which have been shown to influence ground evaporation processes (Wang and Petrone, 2022).

In addition, the temporal resolution of measurements varies across methods. For example, flux chamber and porometer measurements provide instantaneous variables during daytime, whereas micro-lysimeters measurements in this study operated over multiple days, which yields averaged daily values. Reconstructing daily T:ET at the ecosystem level from these different measurements introduces additional biases. Moreover, all Group 2 ET partitioning approaches, strictly speaking, fundamentally rely on the coupling between carbon and water fluxes during the daytime period when photosynthesis takes place. Consequently, their ET partitioning results may not precisely align with measurements that include the nighttime period.

Furthermore, some required input variables were not directly measured and had to be approximated, which also introduces uncertainties. For example, the FVS method requires canopy-level water-use efficiency (WUE_{canopy}) as input, and its outputs are sensitive to the estimates of this parameter (Sulman et al., 2016). However, direct measurements of WUE_{canopy} and intercellular CO_2 concentration to estimate WUE_{canopy} were unavailable, as a result, the optimization method proposed by Scanlon et al. (2019) was used to estimate WUE_{canopy} . But this method is more appropriate for light-saturating conditions (Scanlon et al., 2019), which may not always occur. The FVS method that uses estimated WUE_{canopy} from this optimized method is hereafter referred to as FVSopt. The CWSC method requires the parameter g_l , which is not readily available for wetland ecosystems. Since most wetland plants are C3 species and mosses also follow a photosynthesis pathway similar to C3 plants (Aro and Gerbaud, 1984), g_l values for each site were derived from the reported values for C3 species based on a moisture index (MI) of each site, following Lin et al. (2015). Specifically, Sibbald and Burstall have MIs of 0.77 and 0.72, respectively, and were therefore both given a g_l value of $4.69 \text{ kPa}^{0.5}$. Bonsai, with an MI of 5.38, was assigned a g_l value of $4.02 \text{ kPa}^{0.5}$. Poplar, with an MI of 0.48, was given a g_l value of $3.77 \text{ kPa}^{0.5}$. However, these estimates may not accurately represent the actual g_l at the site, which potentially affects model performance.

Given those limitations, this study emphasizes evaluating whether T:ET values derived from partitioning approaches are broadly consistent with measurement-based estimates and exhibit similar temporal dynamics. Therefore, the coefficient of determination (R^2) was selected as the primary evaluation metric, following Eichelmann et al. (2022) and Stapleton et al. (2022), to quantify how well the predicted T:ET follows the overall pattern of the measurement-based estimates. Values of R^2 closer to 1 indicate better agreement in their temporal dynamics of T:ET. In addition,

Formatted: Superscript

the Nash–Sutcliffe model efficiency coefficient (NSE) was included to provide a complementary assessment of model performance. An NSE value of 1 indicates perfect agreement between predictions and measurements, a value of 0 suggests the model performs no better than the observed mean, and negative values imply the observed mean is a better predictor than the model.

Finally, because the measurement-based daily T:ET time series may be subject to measurement and upscaling uncertainties, model performance was also evaluated against the ensemble mean of all ET partitioning approaches, as ensemble averaging can reduce the influence of method-specific errors and thus provide a more stable and robust reference than any individual approach (Dietterich, 2000). For each day, the ensemble mean was calculated as the average of all available model predictions. In addition, the ensemble mean of each method group was evaluated against both the measurement-based T:ET time series and the overall ensemble mean to determine whether any group of ET partitioning approaches consistently outperformed the others.

It should be noted that evaporation measured by the microlysimeters was not compared with evaporation derived from the ensemble mean because the ensemble mean was derived from model predictions after applying the rainy-day exclusion procedure, whereas the microlysimeter measurements represented cumulative evaporation over multiple days, some of which included rainfall events (Table S3 and Fig. S4). Consequently, ensemble mean estimates were unavailable for portions of the microlysimeter observation periods. This mismatch in temporal coverage prevented a robust direct comparison. Since the measurement-based T:ET was independently validated against site measurements (Section S3 of the Supplementary Material), comparing the ensemble mean with the measurement-based T:ET provides an indirect assessment of how well the ensemble mean agrees with the observations.

3. Results and Discussions

3.1 Evaluation of the Group 1 approaches

The daily T:ET estimates derived from the Group 1 approaches, which are based on high-frequency EC measurements, are presented in Fig. 1. The measurement-based daily T:ET time series and the ensemble mean of all ET partitioning approaches are also shown for comparison. As shown in Fig. 1, the daily T:ET estimates produced by the Group 1 approaches do not always closely match either the measurement-based values or the overall ensemble mean. Likewise, the measurement-based values and the overall ensemble mean do not always agree, which reflects the uncertainty associated with both the reconstruction of the measurement-based T:ET time series and the ET partitioning approaches. Across four sites, the mean T:ET of the overall ensemble exceeds the mean of the measurement-based T:ET by 10.4%. The mean T:ET of the Group 1 ensemble is even higher, exceeding the mean measurement-based daily T:ET by 18.4%. Relative to the mean of all method ensemble, the mean of Group 1 ensemble is 5.3% higher.


Among the Group 1 approaches, MREA and CEC generally produce the highest daily T:ET estimates, whereas CEA consistently produce the lowest. Across four sites, the mean T:ET estimated by MREA exceeds the mean measurement-based by 26.4%, followed by CEC (18.3%). In contrast, the mean T:ET estimated by CEA is only 0.17%

Formatted: Font: 10 pt

Formatted: Font: (Default) +Body (Aptos), (Asian) +Body Asian (等线), 12 pt, Not Bold, Font color: Auto, Ligatures: Standard + Contextual

Formatted: Normal, Left, Level 2, Space Before: 0 pt, After: 0 pt

lower than the mean measurement-based values. FVSopt produces intermediate T:ET estimates among the four Group 1 approaches. The relatively lower T:ET estimates from FVSopt compared with MREA are consistent with the findings of Klosterhalfen et al. (2019b), who compared the FVS and MREA approaches across multiple ecosystems. They reported that FVS tends to overestimate the soil flux component (higher soil evaporation and thus lower T:ET), whereas MREA tends to underestimate the soil flux component. Klosterhalfen et al. (2019) The ET partitioning results derived from Group 1 methods, which are based on high-frequency EC data, are presented in Fig. 1. As anticipated, the T:ET estimates from these methods do not always align closely with the measurement-based values. Klosterhalfen et al. (2019) Klosterhalfen et al. (2019). Nevertheless, when using the measurement-based values as a reference, Group 1 methods appear to perform relatively better at Sibbald and Poplar than at Burstall and Bonsai. According to Zahn et al. (2024), Group 1 methods tend to be more reliable under specific conditions: no strong vertical stratification or convection, EC sampling height near the canopy top (i.e., $z/h \approx 1$), moderate or low LAI, and the canopy does not have gaps that are too wide. However, these site characteristics alone do not fully account for the better performance observed at Sibbald and Poplar. The only consistent difference between the better-performing sites (Sibbald and Poplar) and the others (Burstall and Bonsai) is the relatively higher T:ET values observed at Sibbald and Poplar. Although the methods varied in performance across sites and differed in the magnitude of T:ET estimates at the same site, they generally captured similar temporal patterns and fluctuations that correspond well with the measurement-based T:ET values. An exception is observed at Burstall after August 15, when the measurement-based T:ET exhibits a sharp decline, whereas the T:ET estimated by Group 1 methods shows only a slight decrease.



Formatted: Space After: 0 pt




Figure 1: Comparison of T:ET estimates from the high-frequency EC-based methods (Group 1) with the measurement-based values (black circles) and the ensemble mean of all ET partitioning approaches (open circles) for the across four study sites. Note that FVSopt did not produce any realistic T:ET values at Sibbald and was thus excluded. The light beige shading background (ML-measurement-periods) indicates highlights periods during which ground evaporation was measured using micro-lysimeters (ML) were used to measure ground evaporation at Sibbald, Burstall and Bonsai. The FVSopt method was excluded from the Sibbald analysis because it did not produce realistic T:ET values. Although FVSopt generated realistic T:ET estimates at Bonsai, no data remained after the exclusion of rainy periods. For periods outside of ML measurements, ET partitioning was obtained using the site-calibrated and validated Shuttleworth–Wallace (S–W) model (Wang, 2025).

The R^2 and NSE values for the four Group 1 approaches are presented in Table 3. Regardless of whether the approaches are evaluated against the measurement-based T:ET or the ensemble mean of all ET partitioning approaches, no single Group 1 approach consistently outperforms the others across all four study sites. Furthermore, most approaches perform no better than using the observed mean as the predictor, as indicated by the predominantly negative or near zero NSE values (Table 3). In terms of capturing the temporal variability of T:ET, as indicated by R^2 , CEA performs best at Sibbald when evaluated against the measurement-based T:ET, whereas FVSopt performs best at both Burstall and Poplar, and MREA performs best at Bonsai. The Group 1 ensemble mean does not outperform the best individual approach at any of the study sites. When evaluated against the ensemble mean of all ET partitioning approaches, MREA remains the best-performing approach at Sibbald, FVSopt continues to perform best at Burstall, and CEC

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, English (United Kingdom)

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, English (United Kingdom)

Formatted: Superscript

achieves the highest R^2 at Poplar. The Group 1 ensemble mean outperforms the individual Group 1 approaches only at Poplar.

of the four methods are provided in Table 3.

Table 3. Comparison of R^2 and NSE values for each Group 1 approach and the Group 1 ensemble mean (HF ensemble) relative to the measurement-based T:ET and the ensemble mean of all ET partitioning approaches.

Method	Sibbald		Burstall		Bonsai		Poplar		All sites	
	R^2	NSE	R^2	NSE	R^2	NSE	R^2	NSE	R^2	NSE
Comparison with the measurement-based T:ET:										
FVSopt	NA	NA	0.162	-4.101	NA	NA	0.336	-0.726	0.024	-2.200
MREA	0.174	-5.263	0.013	-6.366	NA	NA	0.031	-0.113	0.118	-1.383
CEA	0.552	-13.335	0.023	-0.727	NA	NA	0.039	-1.325	0.082	0.024
CEC	0.188	-4.836	0.062	-2.220	NA	NA	0.230	0.202	0.417	-0.151
HF ensemble	0.089	-4.469	0.025	-2.699	NA	NA	0.010	-0.162	0.289	-0.292
Comparison with the ensemble mean of all methods:										
FVSopt	NA	NA	0.234	-12.001	NA	NA	0.009	-1.591	0.002	-4.285
MREA	0.409	-4.098	0.270	-14.408	NA	NA	0.503	-2.678	0.437	-3.580
CEA	0.383	0.188	0.020	-7.532	NA	NA	0.465	-2.535	0.233	-3.083
CEC	0.189	-4.069	0.202	-2.421	NA	NA	0.550	-1.166	0.528	-1.111
HF ensemble	0.407	-2.587	0.450	-2.146	NA	NA	0.626	0.452	0.566	-0.274

Note. Values greater than 0.5 are shown in bold.

The performance of the Group 1 approaches has previously been shown to depend on site characteristics. For example, Klosterhalfen et al. (2019b) found that the performance of FVS decreases with increasing measurement height to canopy height ratio (z/h) and leaf area index to canopy height ratio (LAI/h), but improves with increasing canopy height (h) and measurement height (z). This finding may partly explain the poor performance of FVSopt at Sibbald and Bonsai, where both z/h and LAI/h were substantially higher than at the other two sites and few physically realistic T:ET estimates were obtained. For MREA, Klosterhalfen et al. (2019) reported that its performance is more strongly influenced by canopy structural characteristics, including h , LAI , and canopy gaps. More generally, they suggest that site characteristics increasing scalar dissimilarity, such as reduced turbulent mixing, greater source-sink separation, coherent turbulent structures, and ejection events, tend to improve the performance of both the FVS and MREA approaches (Klosterhalfen et al., 2019b).

To further examine how the performance of the Group 1 approaches is related to site characteristics, Spearman correlation analyses were conducted between site characteristics (canopy height, h ; measurement height, z ; LAI ; LAI/h ; z/h ; moss thickness; and mean water table depth, WTD) and the R^2 and NSE values of each Group 1 approach, as well as the Group 1 ensemble mean, relative to both the measurement-based T and the ensemble mean of all ET partitioning approaches. No significant correlations were identified between any site characteristic and the performance metrics.

However, this result should be interpreted with caution because only three sites were available for the correlation analysis (Table 3), which results in very limited statistical power.

Although no clear relationships between site characteristics and model performance were identified in this study, Zahn et al. (2024) extended the work of Klosterhalfen et al. (2019b) by evaluating all the Group 1 methods and suggested that the Group 1 methods tend to be more reliable under specific conditions: no strong vertical stratification or convection, EC sampling height near the canopy top (i.e., $z/h \approx 1$), moderate or low LAI, and the canopy does not have gaps that are too wide. Based on these criteria, the Group 1 methods would be expected to perform better at Burstall and Poplar than Sibbald and Bonsai. Nevertheless, considerable disagreement among the four approaches remains at both sites. At Burstall, the spread of daily T:ET estimates ranges from 0.10 to 0.42 across the study period, and at Poplar it ranges from 0.09 to 0.39, indicating substantial uncertainty in the Group 1 predictions even under conditions considered favorable for their application. Furthermore, Burstall transitions to flooded conditions after 17 August, during which the measurement-based T:ET exhibits a marked decline. In contrast, the T:ET estimates from the Group 1 approaches show only a slight decrease, suggesting that their predictive performance deteriorates under flooded conditions. However, a similar mismatch is not observed at Poplar, which remains flooded throughout the entire study period. At Poplar, the four Group 1 approaches produce relatively consistent T:ET estimates despite the persistent flooding. One possible explanation is the difference in vegetation structure between the two sites. Poplar is a densely forested wetland, whereas Burstall is dominated by shrubs with a lower and more open canopy. This observation suggests that the predictive performance of the Group 1 approaches may be more uncertain in waterlogged wetlands with sparse vegetation cover than in densely forested wetlands. Further evaluation across a broader range of wetland types is needed to confirm this hypothesis.

3.2 Evaluation of the Group 2 approaches

The daily T:ET estimates derived from the Group 2 approaches (uWUE, PP, TEA, CP, and CWSC) are presented in Fig. 2. Compared with the measurement-based T:ET and the ensemble mean of all ET partitioning approaches, the Group 2 approaches exhibit a considerably wider spread of daily T:ET estimates and generally larger temporal fluctuations. Nevertheless, the Group 2 ensemble mean shows closer agreement with both reference datasets than the Group 1 ensemble mean. Across all four sites, the mean T:ET of the Group 2 ensemble exceeds the mean measurement-based T:ET by only 0.32% and is 5.01% lower than the overall ensemble mean.

Although no single Group 2 approach consistently produces the highest or lowest T:ET across all four study sites, several systematic patterns emerge. PP consistently produces the lowest T:ET estimates, whereas CP generally produces T:ET values that are comparable to or higher than the measurement-based values. Among the five approaches, uWUE exhibits the largest temporal fluctuations, while TEA generally predicts higher T:ET than CWSC. Relative to the measurement-based T:ET, TEA overestimates T:ET by 17.44% across four sites, followed by CWSC (13.07%),

Formatted: Normal


Formatted: Font: (Default) +Body (Aptos), 12 pt, English (United Kingdom)

Formatted: Left, Level 2, Space After: 8 pt, Line spacing: Multiple 1.16 li

CP (11.45%), and uWUE (0.43%), whereas PP underestimates T:ET by 167.84%. Relative to the overall ensemble mean, CP overestimates T:ET by 10.39%, followed by TEA (3.84%), whereas uWUE, CWSC, and PP underestimate T:ET by 7.31%, 7.65%, and 67.71%, respectively.

The ET partitioning results from Group 2 methods, including uWUE, PP, TEA, CP and CWSC, are shown in Fig. 2. Compared to Group 1 methods, the performance of Group 2 approaches. Notably, TEA is the only method that captures the sharp decline in T:ET observed at Burstall after August 17, when the site became flooded. In contrast, PP substantially underestimates T:ET and exhibits contrasting temporal trends relative to the measurement-based values at Sibbald and Burstall. This observation is different from the findings in Nelson et al. (2020) who reported similar results between the PP and uWUE methods. The exact cause of this bias remains unclear. However, since other methods align more closely and PP is the only method in this group that requires photosynthetically active radiation (Q) as an additional input, the discrepancy is likely related to this variable. It is possible that Q measured at a single location may not adequately represent canopy-level conditions, potentially due to the uneven and multi-layered canopy structures in these study sites (Fig. S3). Another potential explanation is that the PP method's evaporation component may include not only ground evaporation but also canopy evaporation (e.g., dewfall), as suggested by Pérez-Priego et al. (2018). This inclusion of canopy evaporation could result in lower T:ET. However, this hypothesis could not be tested in the present study and requires further investigation.

is less consistent across sites. For example, the CP method performed well at Sibbald but failed at the other three sites; at Bonsai, all CP-estimated T:ET values exceeded 1 and were therefore excluded. Despite differences in magnitude among methods and across sites, most of Group 2 methods also captured the temporal patterns and fluctuations observed in the measurement-based T:ET values. Notably, TEA was the only method that captured the sharp decline in T:ET observed at Burstall after August 15. In contrast, the PP method substantially underestimated T:ET and exhibited contrasting temporal trends relative to the measurement-based values at Sibbald and Burstall. This observation is different from the findings in Nelson et al. (2020) who reported similar results between the PP and uWUE methods. The exact cause of this bias remains unclear. However, since other methods aligned more closely and PP is the only method in this group that requires photosynthetically active radiation (Q) as an additional input, the discrepancy is likely related to this variable. It is possible that Q measured at a single location did not adequately reflect canopy-level conditions, potentially due to the uneven and multi-layered canopy structures in these study sites (Fig. S1). Another potential explanation is that the PP method's evaporation component may include not only ground evaporation but also canopy evaporation (e.g., dewfall), as suggested by Pérez-Priego et al. (2018). This inclusion of canopy evaporation could result in lower T:ET. However, this hypothesis could not be tested in the present study and requires further investigation.






Figure 2: Comparison of T:ET estimates from the ecosystem carbon-water coupling methods (Group 2) with the measurement-based values (black circles) and the ensemble mean of all ET partitioning approaches (open circles) for the four study sites, against measurement-based values across four study sites. Note that the CP method did not produce any realistic T:ET values at Bonsai and thus was excluded. The PP method was not performed at Poplar due to missing photosynthetically active radiation data. The light beige shading indicates periods during which ground evaporation was measured using micro-lysimeters at Sibbald, Burstall and Bonsai. The light beige background (ML-measurement-periods) highlights periods during which micro-lysimeters (ML) were used to measure ground evaporation at Sibbald, Burstall and Bonsai.

The performance statistics for the Group 2 approaches are summarized in Table 4. Similar to the Group 1 approaches, no single Group 2 approach consistently outperforms the others across all four study sites, and most approaches perform no better than using the observed mean as the predictor. However, in terms of capturing the temporal variability of T:ET, as measured by R^2 , the Group 2 approaches generally perform better than the Group 1 approaches. For example, when evaluated against the measurement-based T:ET, TEA achieves the highest R^2 at Burstall, whereas uWUE and CWSC perform best at Bonsai. The Group 2 ensemble mean does not outperform the best individual approach at most sites, with the exception of Burstall. When evaluated against the overall ensemble mean, however, the Group 2 ensemble generally performs better than the individual Group 2 approaches at most sites. This result

should be interpreted with caution because Group 2 contains more approaches than either Group 1 or Group 3 and therefore likely exerts a stronger influence on the overall ensemble mean.

Figf

Table 4. Comparison of R² and NSE values for each Group 2 approach and the Group 2 ensemble mean (CW ensemble) relative to the measurement-based T:ET and the ensemble mean of all ET partitioning approaches.

Method	Sibbald		Burstall		Bonsai		Poplar		All sites	
	R ²	NSE	R ²	NSE	R ²	NSE	R ²	NSE	R ²	NSE
Comparison with the measurement-based T:ET:										
uWUE	0.331NA	-	0.14762	-	0.735NA	-	0.07933	-3.038-	0.0190.02	-2.043-
	41.666NA			2.7794.104		85.989NA	6	0.726	4	2.200
TEA	0.1780.174	-52.766-	0.5740.013	-2.608-	0.020NA	-	0.0100.0	-0.311-	0.2280.11	-0.886-
		5.263		6.366		62.917NA	31	0.113	8	1.383
CWSC	0.0160.552	-7.222-	0.1320.023	-3.581-	0.732NA	-	0.1050.0	-2.490-	0.0000.08	-
		13.335		0.727		32.363NA	39	1.325	2	0.9570.0
										24
PP	0.0000.188	-197.861-	0.2270.062	-32.868-	0.049NA	-	NA0.23	NA0.20	0.0440.41	-10.179-
		4.836		2.220		59.838NA	0	2	7	0.151
CP	0.2580.089	-1.121-	0.0440.025	-4.463-	NANA	NANA	0.1070.0	-0.589-	0.7110.28	0.450-
		4.469		2.699			10	0.162	9	0.292
CW ensemble	0.253	-36.742	0.664	0.636	0.141	-61.552	0.059	-0.783	0.192	-0.030
Comparison with the ensemble mean of all methods:										
uWUEFVSopt	0.284NA	-1.977NA	0.4200.234	-7.411-	0.433NA	-2.651NA	0.3170.0	-34.203-	0.3190.00	-2.686-
				12.004			09	1.594	2	4.285
TEAMREA	0.1920.409	-4.376-	0.5830.270	-16.882-	0.005NA	-1.001NA	0.2520.5	-0.575-	0.2860.43	-1.630-
		4.098		14.408			03	2.678	7	3.580
CWSCCEA	0.0310.383	-	0.5160.020	-3.033-	0.581NA	-0.479NA	0.7160.4	-8.217-	0.3870.23	-2.268-
		3.3530.18		7.532			65	2.535	3	3.083
		8								
PPCEG	0.0320.189	-214.522-	0.0180.202	-176.717-	0.830NA	-	NA0.55	NA-	0.7210.52	-27.959-
		4.069		2.421		22.169NA	0	1.166	8	1.111
CPHF ensemble	0.1170.407	-0.446-	0.2580.450	-1.113-	NANA	NANA	0.3800.6	-	0.3760.56	-0.284-
		2.587		2.146			26	0.7870.4	6	0.274
								52		
CW ensemble	0.760	0.234	0.428	-6.187	0.965	0.841	0.913	0.885	0.802	0.493

Note. Values greater than 0.5 are shown in bold.

Previous studies have shown that the performance of Group 2 approaches depends on both the quality of the input data, including EC-derived GPP and ET and environmental variables such as radiation, temperature, humidity, wind and precipitation, and the representation of ecosystem carbon-water coupling within each approach (Nelson et al., 2020; Li et al., 2019; Yu et al., 2022). Besides differences in the quality of the input variables used by the Group 2 approaches, differences in the formulation of ecosystem carbon-water coupling are likely to be a major source of

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Justified, Space After: 12 pt, Line spacing: 1.5 lines

Formatted Table

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman, 10 pt, English (Canada)

Formatted: Font: (Default) Times New Roman, 10 pt, English (Canada)

Formatted: Font: (Default) Times New Roman, 10 pt, English (Canada)

Formatted: Font: (Default) Times New Roman, 10 pt, English (Canada)

variation in the predicted T:ET among the approaches. All Group 2 methods are based, to varying extents, on the theory of optimal stomatal conductance. Among these, uWUE, CP, PP and CWSC are formulated using the unified stomatal conductance model proposed by Medlyn et al. (2011). The generally agreement between the measurement-based T:ET and the estimates from uWUE, CP, and CWSC (Fig. 3) supports previous conclusions that an optimal ecosystem response to VPD is a reasonable assumption (Stoy et al., 2019), even in some moss-covered wetlands such as those examined in this study.

However, as noted by Stoy et al. (2019) and Nelson et al. (2020), the way in which optimality is conceptualized and implemented in these models may need to be refined. Improvements could include incorporating effects of plant hydraulic architecture, soil moisture, temperature, atmospheric CO₂ concentration (Bernacchi et al., 2004; Medlyn et al., 2011; Lin et al., 2015; Nardini and Salleo, 2000), as well as considering the timescales over which assumptions about constant parameters in the unified stomatal conductance model can hold (Mäkelä et al., 1996), could improve model performance. This is illustrated by the comparison of results from uWUE, CP and CWSC in Fig. 3. Both CP and CWSC, which account for soil moisture and atmospheric CO₂ concentration effects respectively, produced a narrower spread of T:ET estimates than uWUE, which suggests that including these additional constraints improves model performance at sub-daily and daily scales.




Figure 3: T:ET estimates from each ET partitioning approach, pooled across the four study sites, with the measurement-based T:ET values (left two columns) and the ensemble mean of all ET partitioning approaches (right two columns). (Nelson et al., 2020) (Nelson et al., 2020)

3.3 Evaluation of the Group 3 approaches

The best-performing ML models were identified using the adjusted R^2 criterion defined by Stapleton et al. (2022), based on their performance on the night-time validation dataset. The flooding validation dataset was not used for model selection because all candidate models produced negative adjusted R^2 values under flooded conditions (Fig. S1). The best-performing ML algorithms are the LightGBM for Sibbald and Poplar and the XGBoost for Burstall and Bonsai, each with a different optimal predictor sets (Tables S1 and S2). This finding is consistent with Stapleton et al. (2022), who reported that no single ML algorithm or predictor set consistently performs best across all study sites. Instead, each site has its own best-performing algorithm, predictor combination, and variable importance ranking. Across the four study sites, sensible heat flux, air temperature, and net radiation consistently ranked among the five most important predictors of evaporation (Table S1).

The daily T:ET estimates produced by the best-performing ML model for each study site are presented in Fig. 4, and their performance statistics are summarized in Table 5. Similar to Groups 1 and 2 approaches, the ML models do not consistently outperform the other ET partitioning approaches across all four study sites and, in most cases, perform no better than using the observed mean as the predictor. In terms of reproducing the temporal variability of T:ET, as indicated by R^2 , the best-performing ML model achieves its highest agreement with the measurement-based T:ET at Poplar. When the overall ensemble mean is used as the reference, however, the highest R^2 is obtained at Bonsai. Additionally, the Group 3 ensemble is defined as the average of the predictions from the best-performing ML model identified for each study site and the TEA approach, calculated from all available predictions on each day. Across four sites, the mean T:ET of the Group 3 ensemble exceeds the mean measurement-based T:ET by 16.09% and is 6.14% higher than the overall ensemble mean.

Formatted: Normal, Level 2

Formatted: Not Highlight

Formatted: Not Highlight

Formatted: Space After: 10 pt




Figure 4. Comparison of T:ET estimates from the machine learning methods (Group 3) with the measurement-based values (black circles) and the ensemble mean of all ET partitioning approaches (open circles) for the four study sites. The light beige shading indicates periods during which ground evaporation was measured using micro-lysimeters at Sibbald, Burstall and Bonsai.

Table 5. Comparison of R^2 and NSE values for the Group 3 approach and the Group 3 ensemble mean (ML ensemble) relative to the measurement-based T:ET and the ensemble mean of all ET partitioning approaches.

Method	Sibbald		Burstall		Bonsai		Poplar		All sites	
	R^2	NSE	R^2	NSE	R^2	NSE	R^2	NSE	R^2	NSE
Comparison with the measurement-based T:ET:										
ML	0.107	-11.960	0.003	-3.884	0.115	-47.647	0.680	-0.417	0.023	-0.736
ML ensemble	0.128	-15.081	0.449	-2.368	0.153	-47.613	0.222	-0.304	0.181	-0.381
Comparison with the ensemble mean of all methods:										
ML	0.002	-1.393	0.116	-6.261	0.627	0.144	0.084	-2.446	0.179	-0.564
ML ensemble	0.007	-1.396	0.634	-5.488	0.442	0.412	0.253	-0.594	0.332	-0.187

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: Bold, Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: Bold, Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: Bold, Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: Bold, Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: Bold, Ligatures: None

Formatted: Font: (Default) Times New Roman, (Asian) Times New Roman, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Note. Values greater than 0.5 are shown in bold. The ML ensemble includes the best-performing ML model identified using the framework of Stapleton et al. (2022) and the TEA approach, which is also classified as a ML-based ET partitioning approach. Stapleton et al. (2022)

Unlike the results reported by Eichelmann et al. (2022) and Stapleton et al. (2022), the ML models developed under their framework (Eichelmann et al., 2022) fail to reproduce the sharp decline in the measurement-based T:ET when Burstall transitions from unsaturated to flooded conditions. Assuming that the measurement-based T:ET reasonably represents the actual temporal dynamics of the site, one possible explanation is that the dominant environmental controls governing ground evaporation and ET partitioning change substantially following flooding. Consequently, these ML models, which are trained under non-flooded conditions, rely on environmental relationships that are no longer representative once the site becomes inundated. In particular, the factors influencing WUE (i.e., the relationship between GPP and T) may change markedly following flooding. This interpretation is supported by the performance of the TEA approach, which is the only method that reproduces the observed decline in T:ET. Unlike the other carbon-water coupling approaches, TEA employs a random forest model to estimate a temporally varying WUE, which allows it to better adapt to changing environmental conditions. In contrast, the other carbon-water coupling approaches rely on presumed relationships between carbon and water fluxes and likewise fail to capture the transition in WUE following flooding. Although TEA demonstrates superior performance under these conditions, this does not necessarily imply that it is generally suitable for evaporation-dominated ecosystems, as Nelson et al. (2018) reported that TEA tends to overestimate transpiration at sites with a relatively constant evaporation component. Therefore, the improved performance of TEA at Burstall is more likely attributable to its ability to accommodate temporal changes in WUE during the transition from unsaturated to flooded conditions than to a general advantage in evaporation-dominated ecosystems. Together, these findings suggest that changes in WUE following inundation play an important role in determining ET partitioning.


Another important consideration is that, in both Eichelmann et al. (2022) and Stapleton et al. (2022), the ML models were validated by comparing predicted evaporation with EC-measured ET during flooded periods when vegetation within the flux footprint had not yet become established. Under these conditions, transpiration can reasonably be assumed to be negligible, such that nearly all measured ET originates from evaporation. In contrast, this study evaluated the ML models under a different flooding scenario. At our sites, flooding occurred after vegetation was already well established and transpiration remained an important component of ET. Consequently, these ML models may not perform equally well under all flooding scenarios, which may explain the negative adjusted R^2 values obtained when they were evaluated using the flooding validation dataset (Fig. S1). However, unlike at Burstall, the best-performing ML model at Poplar does not deviate substantially from the reference estimates (Fig. 4), suggesting that the framework may remain applicable to wetlands that are continuously flooded throughout the study period. Together, these findings suggest that, although the framework may perform well under conditions of initial flooding or persistent flooding, further evaluation is needed to determine its applicability to wetlands that transition between unsaturated and flooded conditions.

Formatted: Font: (Default) Times New Roman, 10 pt

The performance of these ML models Stapleton et al. (2022) depends on several factors, including the validity of the assumption that nighttime T is negligible, the quality of the nighttime EC data and associated gap-filling procedures, as well as the choice of ML architecture and its hyperparameters (Stapleton et al., 2022). Among these factors, the influence of data gaps and gap-filling procedures appears to be limited in the present study. Poplar and Sibbald have substantially larger proportions of missing EC data (approximately 60%) than Burstall and Bonsai (approximately 20%). Nevertheless, the best-performing ML model achieves its highest agreement with the measurement-based T:ET at Poplar, which indicates that the extent of data gaps and the associated gap-filling procedures are unlikely to be the primary factors controlling model performance in this study. The magnitude of nighttime transpiration at each study site and the choice of machine learning architecture and hyperparameter settings may also contribute to differences in model performance. However, evaluating these factors is beyond the scope of the present study. Overall, the results suggest that the performance of the ML models is likely governed by the combined effects of data quality, machine learning architecture, site-specific environmental conditions, and the validity of the underlying ET partitioning assumptions.

3.4 Agreement among ET Partitioning Approaches

The Spearman correlation analysis of the predicted T:ET among the ET partitioning methods evaluated in this study (Fig. 5) provides an additional perspective on the agreement among methods. Spearman correlation was preferred over Pearson correlation because it evaluates the monotonic relationship between methods without assuming a linear relationship or normally distributed data (Weaver et al., 2017). It is therefore more robust to outliers and better suited for comparing the temporal consistency of daily T:ET estimates produced by different ET partitioning approaches. Overall, the strength, direction, and statistical significance of the correlations vary considerably across the four study sites.



Formatted: Centered

Figure 5: Spearman correlation matrix of the predicted T:ET among the ET partitioning methods evaluated in this study. Blue indicates positive correlations, whereas orange indicates negative correlations. Statistical significance is denoted by * ($P < 0.05$), ** ($P < 0.01$), and *** ($P < 0.001$).

Notably, the measurement-based T:ET is not significantly correlated with the all-method ensemble at any of the study sites, and the correlation is negative at Poplar. Likewise, the measurement-based T:ET shows no significant correlation with any of the group ensemble means, except at Burstall, where it is positively correlated the Group 2 ensemble. In contrast, all-method ensemble exhibits positive and significant correlations with almost all three group. These contrasting correlation patterns indicate that the assessment of ET partitioning performance depends strongly on the choice of reference dataset. Although the present study does not provide sufficient evidence to determine whether the measurement-based T:ET or the all-method ensemble better represents the true site-scale T:ET dynamics, the discrepancy between these two references highlights the uncertainty associated with defining a benchmark for ET partitioning evaluation. Consequently, the choice of reference dataset should be carefully considered when comparing and interpreting the performance of different ET partitioning approaches.

Among the three methodological groups, the Group 1 ensemble is positively correlated with the Group 3 ensemble at all four study sites, although none of these correlations is statistically significant. In contrast, the correlation between

Formatted: Font: (Asian) Times New Roman, 9 pt, Bold, English (United Kingdom), Ligatures: None

Formatted: Font: (Asian) Times New Roman, 9 pt, Bold, English (United Kingdom), Ligatures: None

Formatted: Line spacing: single

the Group 1 and Group 2 ensembles is site dependent, with a significant positive correlation observed only at Poplar. The Group 2 ensemble is positively correlated with the Group 3 ensemble at all sites except Sibbald, and these correlations are significant at Burstall and Poplar. Overall, the Group 3 ensemble exhibits stronger and more consistent agreement with both the Group 1 and Group 2 ensembles than the agreement observed between Groups 1 and 2. This pattern suggests that the ML models capture temporal variability that is shared by both the high-frequency EC-based and the carbon-water coupling approaches, likely because the predictor variables used in the machine learning framework capture information related to both turbulence transport and ecosystem carbon-water coupling.

Regarding the individual methods, approaches within the same methodological group generally exhibit stronger positive correlations than approaches from different groups. For example, MREA agrees more with CEA and CEC than with uWUE, which reflects their shared theoretical assumptions and common input variables. However, notable contrasts from the within-group correlation patterns are observed for FVSopt at Burstall and Poplar, and for CWSC and PP at Sibbald, where these methods exhibit substantially weaker agreement with the other approaches in their respective groups. This may reflect the uncertainty associated with estimating leaf-level WUE, as FVSopt, CWSC, and PP all rely on prescribed relationships to estimate leaf-level WUE, and these relationships may not adequately represent the responses of leaf-level WUE to the highly dynamic environmental conditions in wetlands. For example, Klosterhalfen et al. (2019a) demonstrated that the FVS is vulnerable to errors in water use efficiency estimates and atmospheric conditions that violate the variance similarity assumption, like under conditions of weak turbulence, canopy heterogeneity, and poorly separated soil and canopy source-sink distributions.

The degree of agreement among approaches also differs across the four study sites. Poplar exhibits the strongest agreement among methods, whereas Sibbald shows the weakest. This difference may be related to the greater ecosystem heterogeneity at Sibbald. Compared with the relatively homogeneous canopy of the dense forested Poplar peatland, Sibbald is characterized by a multi-layered shrub canopy, large canopy gaps, and pronounced hummock-hollow microtopography (Fig. S3). Such structural heterogeneity is likely to increase the spatial variability of carbon and water fluxes within the EC footprint, and thereby introduces greater uncertainty into EC measurements and reduces the applicability of both scalar transport assumptions and carbon-water relationships used by the EC-based ET partitioning approaches examined in this study.

(Eichelmann et al., 2022)(Stapleton et al., 2022)(Eichelmann et al., 2022)(Stapleton et al., 2022)(Stapleton et al., 2022)
3.5 Effects of moss cover and flooding on the performance of ET partitioning

The primary objective of this study is to evaluate the performance of existing ET partitioning approaches in moss-covered wetlands. As discussed in Section 1 and 2.1, moss cover has the potential to violate several of the fundamental assumptions underlying these approaches. For the high-frequency EC-based approaches (Group 1), strong moss photosynthesis can produce a net negative surface CO₂ signal, thereby violating the assumption that the surface acts as a net CO₂ source. Evidence for this behavior was reported by Walker et al. (2017), whose chamber measurements on Sphagnum-peat columns showed frequent transitions between positive and negative hourly CO₂ fluxes (their Fig. 4), together with sustained daily net CO₂ uptake from mid-June to mid-July. These observations suggest that a moss

layer has the potential to compromise the applicability of high-frequency EC-based ET partitioning approaches that rely on a persistent positive surface CO₂ signal.(Walker et al., 2017)

In terms of our study sites, direct measurements of surface CO₂ exchange were unfortunately not available during the study period. Nevertheless, existing evidence indicates that negative surface CO₂ fluxes are plausible, particularly at Poplar. Following the 2016 wildfire, chamber measurements conducted in 2017 over moss-dominated plots at Poplar showed that the unburned forest floor remained a net CO₂ sink until early August (Van Beest (2019); Fig. 3-1). Although these measurements were not collected during the same study period, they provide independent evidence that strong moss photosynthesis can occur at this site and may influence the assumptions underlying Group 1 approaches.

As for the ecosystem carbon-water coupling approaches (Group 2), moss cover introduces a different source of uncertainty. These approaches estimate WUE under the assumption that GPP originates exclusively from vascular vegetation that also contributes to transpiration. However, mosses contribute to ecosystem GPP while making no contribution to transpiration. Consequently, moss photosynthesis may bias WUE estimates and propagate errors into ET partitioning. The methods that are more sensitive to GPP errors, such as PP and uWUE, as diagnosed in Nelson et al. (2020), are expected to exhibit greater uncertainty. The magnitude of this effect depends on the contribution of mosses to ecosystem GPP, which has been reported to account for approximately 20% to 30% of total forest CO₂ uptake in a drained forested peatland (Badorek et al., 2011). Therefore, this source of uncertainty may not be trivial.

Another potential source of uncertainty affecting ET partitioning performance is flooding. Under flooded conditions, EC measurements may become less reliable because of weak turbulence and enhanced advection effects (Dai et al., 2026). For the Group 1 approaches, flooding may also weaken or obscure surface CO₂ signals, further challenging the assumptions underlying these methods. To evaluate the influence of flooding, we recalculated the R² and NSE values for Burstall after excluding the flooded period (Table 6) and compared the resulting performance statistics with those obtained using the complete dataset (Tables 3, 4 and 5). When evaluated against the measurement-based T:ET, the R² values of CEA, MREA, PP, the best-performing ML model, and the Group 1 ensemble increased slightly after the flooded period was excluded. When the overall ensemble mean was used as the reference, only CEA, MREA and PP showed slight improvements in R². However, despite these modest increases in R², the NSE values decreased for all approaches except TEA, and all NSE values remained negative. These results suggest that the factors affecting ET partitioning performance are more complex and cannot be explained by flooding alone.

Table 6. Comparison of R² and NSE values relative to the measurement-based T:ET and the ensemble mean of all ET partitioning approaches for Burstall after excluding the flooding period.

Methods	Comparison with the measurement-based T:ET		Comparison with the ensemble mean of all methods	
	R ²	NSE	R ²	NSE
FVSopt	0.050	-13.991	0.017	-21.977
CEA	0.204 (+)	-2.835	0.040 (+)	-16.612

Formatted: Font: Bold

Formatted: Centered

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted Table

Formatted: Font: 8 pt, Superscript

Formatted: Font: 8 pt

Formatted: Font: 8 pt, Superscript

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

MREA	0.104 (+)	-19.574	0.310 (+)	-23.020
CEC	0.019	-7.533	0.095	-4.611
WUE	0.000	-10.159	0.286	-11.256
PP	0.236 (+)	-45.558	0.055 (+)	-237.481
TEA	0.015	-13.755	0.262	-15.987 (+)
CP	0.044	-4.463	0.193	-1.843
CWSC	0.005	-7.338	0.396	-6.875
ML	0.385 (+)	-12.533	0.110	-10.169
HF Ensemble	0.087 (+)	-8.624	0.381	-4.037
CW Ensemble	0.040	-0.383	0.399	-4.716
ML Ensemble	0.074	-11.887	0.436	-9.104

Note. (+) indicates that the corresponding performance metric improved after excluding the flooding period.

Overall, regardless of whether the measurement-based T:ET or the overall ensemble mean was used as the reference, none of the evaluated approaches consistently outperformed the observed mean as the predictor. This was true even for the ML approaches, whose underlying assumptions are not fundamentally violated by moss cover. Together, these findings suggest that the performance of ET partitioning approaches in wetlands is influenced by the combined effects of multiple factors, including moss cover, flooding conditions, canopy structure, canopy density, canopy openness, site heterogeneity, the measurement height relative to canopy height, the quality of EC and environmental measurements, and the validity of the assumptions underlying each ET partitioning approach. The overall performance of each method, pooled across all study sites, is presented in Fig. 3. Except for the PP method, most methods produced T:ET estimates that scatter around the 1:1 line relative to the measurement-based values. There was no clear pattern indicating how variations in water table levels near the ground affected model performance. Notably, as illustrated in Fig. 3, when T:ET exceeds 0.5, the estimates tend to align more closely with the 1:1 line. However, for T:ET values below 0.5, Group 1 methods generally exhibit substantial overestimation, while most Group 2 methods failed to provide realistic estimates and were therefore excluded. Surprisingly, the TEA method stands out as an exception among all methods when T:ET is lower than 0.5. However, this does not necessarily indicate its suitability for E-dominated ecosystems, as this method has been shown to overestimate T when sites have a relatively constant E component (Nelson et al., 2018). These findings highlight a limitation of these ET partitioning methods in evaporation-dominated ecosystems.

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Font: 8 pt


Formatted: Centered, Space After: 0 pt

Formatted: Font: 8 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: Not Italic

Formatted: Space After: 10 pt



Formatted: Justified, Space After: 10 pt, Line spacing: 1.5 lines

Figure 3: Performance of all ET partitioning methods pooled across sites, compared against measurement-based T:ET values.

Additionally, aside from MREA, CEA, and CEC, which are based on conditional sampling approaches, all other methods, to varying extents, utilize the theory of optimal stomatal conductance. Among these, uWUE, CP, and CWSC are formulated using the unified stomatal conductance model proposed by Medlyn et al. (2011). The general agreement between their T:ET estimates and the measurement-based values supports previous conclusions that an optimal ecosystem response to VPD is a reasonable assumption (Stoy et al., 2019), even in some moss-covered wetlands such as those examined in this study. However, as noted by (Nelson et al., 2018) Stoy et al. (2019) Nelson et al. (2020) (Nelson et al., 2018), the way in which optimality is conceptualized and implemented in these models may need to be refined. Improvements could include incorporating effects of plant hydraulic architecture, soil moisture, temperature, atmospheric CO₂ concentration (Bernacchi et al., 2004; Medlyn et al., 2011; Lin et al., 2015; Nardini and Sallee, 2000), as well as considering the timescales over which assumptions about constant parameters in the unified stomatal conductance model can hold (Mäkelä et al., 1996), could improve model performance. This is illustrated by the comparison of results from uWUE, CP and CWSC. Both CP and CWSC, which account for soil moisture and atmospheric CO₂ concentration effects respectively, produced a narrower spread of T:ET estimates than uWUE, which suggests that including these additional constraints improves model performance at sub-daily and daily scales.

In sum, despite the variability in T:ET estimates and the limitations discussed above, these methods collectively provide a plausible range within which the true T:ET likely falls and reasonably captures T:ET temporal dynamics. This indicates that, despite the presence of mosses, which may challenge some assumptions of ET partitioning approaches, these methods collectively still offer valuable insights into wetland ET partitioning, particularly its temporal dynamics. This enables investigation of some urgent research questions in moss-covered wetland ecosystems, like how transpiration and evaporation components respond to changing climate and environmental conditions.

4. Summary and closing thoughts

This study provides first comprehensive evaluations of ET partitioning approaches in moss-covered wetlands across contrasting vegetation structures and hydrological conditions. Overall, no single ET partitioning approach consistently outperformed the others across the four study sites or under both flooded and non-flooded conditions. Likewise, no methodological group, including the high-frequency EC-based approaches, ecosystem carbon-water coupling approaches, or machine learning approaches, demonstrated a clear overall advantage. The performance of ET partitioning approaches in wetlands is likely governed by the combined effects of multiple environmental and methodological factors. In addition to the assumptions underlying each approach, performance is influenced by EC measurement configuration, site characteristics such as canopy height and leaf area index, ecosystem heterogeneity, moss cover, and flooding conditions. These factors interact to influence both the quality of EC and environmental observations and the validity of the assumptions on which current ET partitioning approaches are based.

An important finding of this study is that the evaluation results depend strongly on the choice of reference dataset. The measurement-based T:ET and the ensemble mean of all ET partitioning approaches exhibit markedly different correlation patterns with the evaluated methods, highlighting the uncertainty associated with defining a reference for ET partitioning evaluation. This finding suggests that greater attention should be given to reference dataset selection and uncertainty quantification in future benchmarking studies.

Although several approaches perform reasonably well under specific site conditions, none adequately captures the full range of environmental conditions encountered in wetlands, particularly transitions between unsaturated and flooded states. Future work should therefore focus on developing wetland-specific ET partitioning approaches that explicitly account for dynamic hydrological conditions, improve the representation of carbon-water coupling, and incorporate the influence of mosses on ecosystem carbon exchange. Such developments will improve the reliability of ET partitioning in wetlands and advance our understanding of water and carbon cycling in these globally important ecosystems.

Since (The purpose of this study is to provide supporting evidence and recommendation for future research that plan to apply EC-based ET partitioning approaches in moss-covered wetlands, particularly in cases where direct validation through field measurements is not available. By comparing method-derived T:ET with direct field measurements of ET and its components at four different wetlands and by excluding rainy periods from analysis, this study found that all methods, despite having substantial differences in the magnitude of estimated T:ET, captured similar overall

temporal patterns of T:ET. This observation is consistent with the evaluation of three ET partitioning methods (TEA, uWUE and PP) conducted by Nelson et al. (2020). This study also found that no single approach consistently outperformed the others across all sites and time periods. Therefore several practical recommendations can be made. Based on our study, the ensemble of the high-frequency EC-based approaches and the machine learning approaches produced T:ET estimates of similar magnitude, and both tended to overestimate T:ET relative to both the measurement-based estimates and the overall ensemble mean. In contrast, the ensemble of the ecosystem carbon-water coupling approaches produced lower mean T:ET values but generally exhibited larger temporal fluctuations. Rather than relying on a single ET partitioning group, we recommend applying multiple approaches from all three methodological groups whenever possible. Comparing predictions across different approaches provides a more comprehensive assessment of uncertainty and helps identify potential sources of disagreement among methods. For example, our analysis suggests that the parameterization of WUE is one of the major factors contributing to disagreement among methods. This finding implies that the response of WUE to changing environmental conditions in wetland ecosystems may not yet be fully represented in current formulations. However, it should be noted that this recommendation does not imply that the ensemble mean or any single approach is consistently better than the others. Our results showed that the relative performance of individual approaches, as well as the ensemble mean, depended on the reference dataset used for evaluation. For researchers interested in ET partitioning in similar wetlands, applying multiple methods concurrently is recommended when possible.

Our recommendation is consistent with the suggestion provided by Zahn et al. (2024) in their evaluation of Group 1 methods. But building upon that recommendation, findings from this study further suggest that Group 2 methods exhibit greater variability and inconsistency across sites. This is likely due to the distinct dominant environmental controls on which each Group 2 method is based, as well as the uncertainties associated with the additional meteorological or soil variables required beyond EC data by these methods. While Group 1 methods may not yield the most accurate estimates, Group 2 methods may provide either the most accurate or the least accurate results, depending on site specific data quality. Without prior knowledge on the dominant environmental controls at a given wetland, and considering the greater potential for measurement uncertainties in Group 2 methods, Group 1 methods offer a more reliable starting point for estimating ET partitioning. However, high-frequency EC data are often accessible only to tower owners and are not widely shared. As such, this recommendation to prioritize Group 1 methods is intended for researchers with access to high-frequency data.

Finally, However, several caveats must be considered when interpreting the results of this study. First, the study did not examine the mechanisms by which mosses influence the results of ET partitioning approaches. This remains an important topic for future research. Secondly, the ET partitioning results presented here were filtered based on the data quality controls described in the Methods section. It is important to emphasize that rainy periods were excluded; if included, most Group 2 methods produced unrealistic T:ET values exceeding 1. Third, the evaporation component considered in this study refers exclusively to ground evaporation. Whether the findings can be extended to include canopy evaporation remains an open question. Lastly, the conclusions are most likely valid for sites with moss thickness similar to or less than that of those studied here. Uncertainty remains regarding their applicability to

ecosystems with thicker moss layers and fewer vascular plants. Nonetheless, the study sites represent common wetlands and are expected to reflect a broad range of wetland ecosystems. Therefore, the conclusions are highly likely applicable to many wetland ecosystems.

Data Availability Statement

The measurement-based T:ET estimates were previously published in Gabrielli (2016) and Wang (2025), and the raw measurement data will be made available upon reasonable request to the corresponding author. The FVS, CEA, MREA and CEC methods were applied using the ET partitioning code developed by Zahn (2024); the PP method by Pérez-Priego and Wutzler (2018); the uWUE and TEA methods by Nelson (2020); ~~and the CWSC method by Jiang and Yu (2021); and the machine learning framework by Stapleton (2022).~~ We sincerely thank these authors for generously sharing their codes.

Author contribution

YW and RP jointly developed the research concept. YW carried out data compilation, data analysis, and manuscript drafting. RP supervised the project, secured funding, and contributed to manuscript writing. LZ assisted with code adaptation, development, and manuscript editing. All authors approved the final manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

The authors would like to thank Lindsey Langs and Eric Murray for their assistance in data collection at Sibbald, Burstall and Bonsai, and Elise Gabrielli for data collection at Poplar. We thank Adam Green for eddy covariance data processing ~~and Dr. Myroslava Khomik for her suggestions on data analysis.~~ The University of Calgary must also be acknowledged for their hospitality and accommodations at the Biogeosciences Institute Barrier Lake Field Station during the field seasons. We also thank the Coldwater Laboratory in Canmore, Alberta, part of the University of Saskatchewan Global Water Future Project, for providing precipitation data for the Bonsai site. We are particularly grateful for ~~to the constructive feedback from~~ Dr. John Pomeroy (~~University of Saskatchewan~~), Dr. Cherie Westbrook, (~~University of Saskatchewan~~), and Dr. Rebecca Rooney (~~University of Waterloo~~), Dr. Elke Eichelmann, and the anonymous reviewers for their constructive comments, which greatly improved this manuscript.

854 **Declaration of Generative AI and AI-assisted technologies in the writing process**

855 During the preparation of this work the first author used ChatGPT-4.0 only for grammar checking and language
856 polishing. ChatGPT-4.0 was not used for content generation. After using this tool, the first author reviewed and edited
857 the content as needed and takes full responsibility for the content of the publication.

858 **Funding information**

859 We would like to thank the Mountain Water Futures project of the Global Water Futures programme (Canada First
860 Research Excellence Fund), Alberta Innovates – Energy and Environment Solutions, and the Canadian Natural Science
861 and Engineering Research Council (NSERC) Discovery (RGPN-04182-2017; Petrone) and CREATE (463960-2015;
862 Petrone) grants programs for the financial support.

863 **References**

864 Admiral, S. W. and Lafleur, P. M.: Partitioning of latent heat flux at a northern peatland, *Aquatic Botany*, 86, 107-
865 116, <https://doi.org/10.1016/j.aquabot.2006.09.006>, 2007.

866 Albert-Saiz, M., Stróżecki, M., Rastogi, A., and Juszczak, R.: A Multi-Model Gap-Filling Strategy Increases the
867 Accuracy of GPP Estimation from Periodic Chamber-Based Flux Measurements on Sphagnum-Dominated
868 Peatland, *Sustainability*, 17, 393, <https://doi.org/10.3390/su17020393>, 2025.

869 Allen, S. T., Reba, M. L., Edwards, B. L., and Keim, R. F.: Evaporation and the subcanopy energy environment in a
870 flooded forest, *Hydrological Processes*, 31, 2860-2871, <https://doi.org/10.1002/hyp.11227>, 2017.

871 Aro, E.-M. and Gerbaud, A.: Photosynthesis and Photorespiration in Mosses, in: *Advances in Photosynthesis*
872 *Research*, edited by: Sybesma, C., Springer Netherlands, Dordrecht, 867-870, [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-94-017-4973-2_198)
873 [94-017-4973-2_198](https://doi.org/10.1007/978-94-017-4973-2_198), 1984.

874 Aubinet, M., Vesala, T., and Papale, D.: *Eddy covariance: a practical guide to measurement and data analysis*,
875 Springer Science & Business Media, <https://doi.org/10.1007/978-94-007-2351-1>, 2012.

876 Badorek, T., Tuittila, E.-S., Ojanen, P., and Minkinen, K.: Forest floor photosynthesis and respiration in a drained
877 peatland forest in southern Finland, *Plant Ecology & Diversity*, 4, 227-241,
878 <https://doi.org/10.1080/17550874.2011.644344>, 2011.

879 Baldocchi, D. D.: Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of
880 ecosystems: past, present and future, *Global Change Biology*, 9, 479-492, [https://doi-](https://doi-org.proxy.lib.uwaterloo.ca/10.1046/j.1365-2486.2003.00629.x)
881 [org.proxy.lib.uwaterloo.ca/10.1046/j.1365-2486.2003.00629.x](https://doi-org.proxy.lib.uwaterloo.ca/10.1046/j.1365-2486.2003.00629.x), 2003.

882 Bayfield, N. G.: Notes on water relations of *Polytrichum commune* Hedw, *Journal of Bryology*, 7, 607-617,
883 <https://doi.org/10.1179/jbr.1973.7.4.607>, 1973.

884 Bernacchi, C. J., Singasaas, E. L., Pimentel, C., Portis Jr, A. R., and Long, S. P.: Improved temperature response
885 functions for models of Rubisco-limited photosynthesis, *Plant, Cell & Environment*, 24, 253-259,
886 <https://doi.org/10.1111/j.1365-3040.2001.00668.x>, 2004.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Indent Left: 0 cm, Hanging: 1.27 cm, Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

887 Bijoor, N. S., Pataki, D. E., Rocha, A. V., and Goulden, M. L.: The application of delta(1)(8)O and deltaD for
 888 understanding water pools and fluxes in a Typha marsh, Plant Cell Environ, 34, 1761-1775,
 889 <https://doi.org/10.1111/j.1365-3040.2011.02372.x>, 2011.

890 Burba, G., Schmidt, A., Scott, R. L., Nakai, T., Kathilankal, J., Fratini, G., Hanson, C., Law, B., McDermitt, D. K.,
 891 and Eckles, R.: Calculating CO₂ and H₂O eddy covariance fluxes from an enclosed gas analyzer using an
 892 instantaneous mixing ratio, Global Change Biology, 18, 385-399, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2011.02536.x)
 893 [2486.2011.02536.x](https://doi.org/10.1111/j.1365-2486.2011.02536.x), 2012.

894 Burba, G. G., Verma, S. B., and Kim, J.: Surface energy fluxes of Phragmites australis in a prairie wetland,
 895 Agricultural and forest meteorology, 94, 31-51, [https://doi.org/10.1016/S0168-1923\(99\)00007-6](https://doi.org/10.1016/S0168-1923(99)00007-6), 1999.

896 Businger, J. A. and Oncley, S. P.: Flux measurement with conditional sampling, Journal of Atmospheric and Oceanic
 897 Technology, 7, 349-352, [https://doi.org/10.1175/1520-0426\(1990\)007<0349:FMWCS>2.0.CO;2](https://doi.org/10.1175/1520-0426(1990)007<0349:FMWCS>2.0.CO;2), 1990.

898 Caird, M. A., Richards, J. H., and Donovan, L. A.: Nighttime Stomatal Conductance and Transpiration in C₃ and C₄
 899 Plants, Plant Physiology, 143, 4-10, <https://doi.org/10.1104/pp.106.092940>, 2007.

900 Chen, S., Yang, Z., Liu, X., Sun, J., Xu, C., Xiong, D., Lin, W., Li, Y., Guo, J., and Yang, Y.: Moss regulates soil
 901 evaporation leading to decoupling of soil and near-surface air temperatures, Journal of Soils and Sediments,
 902 19, 2903-2912, <https://doi.org/10.1007/s11368-019-02297-4>, 2019.

903 Dai, Y., Liu, B., Wei, J., Shi, Y., Li, Q., Liu, F., Luo, Y., and Cui, Y.: Impacts of surface heterogeneity on energy
 904 partitioning in paddy ecosystems: A dual eddy covariance study, Agricultural and Forest Meteorology, 384,
 905 111165, <https://doi.org/10.1016/j.agrformet.2026.111165>, 2026.

906 Daley, M. J. and Phillips, N. G.: Interspecific variation in nighttime transpiration and stomatal conductance in a
 907 mixed New England deciduous forest, Tree Physiology, 26, 411-419,
 908 <https://doi.org/10.1093/treephys/26.4.411>, 2006.

909 Dietterich, T. G.: Ensemble Methods in Machine Learning, Berlin, Heidelberg, 1-15, [https://doi.org/10.1007/3-540-](https://doi.org/10.1007/3-540-45014-9_1)
 910 [45014-9_1](https://doi.org/10.1007/3-540-45014-9_1).

911 Drexler, J. Z., Snyder, R. L., Spano, D., and Paw U, K. T.: A review of models and micrometeorological methods
 912 used to estimate wetland evapotranspiration, Hydrological processes, 18, 2071-2101,
 913 <https://doi.org/10.1002/hyp.1462>, 2004.

914 Eichelmann, E., Mantoani, M. C., Chamberlain, S. D., Hemes, K. S., Oikawa, P. Y., Szutu, D., Valach, A., Verfaillie,
 915 J., and Baldocchi, D. D.: A novel approach to partitioning evapotranspiration into evaporation and
 916 transpiration in flooded ecosystems, Global Change Biology, 28, 990-1007,
 917 <https://doi.org/10.1111/gcb.15974>, 2022.

918 Farquhar, G. D. and Sharkey, T. D.: Stomatal conductance and photosynthesis, Annual review of plant physiology,
 919 33, 317-345, <https://doi.org/10.1146/annurev.pp.33.060182.001533>, 1982.

920 Farquhar, G. D., von Caemmerer, S. v., and Berry, J. A.: A biochemical model of photosynthetic CO₂ assimilation in
 921 leaves of C₃ species, planta, 149, 78-90, <https://doi.org/10.1007/BF00386231>, 1980.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

922 Fisher, J. B., Baldocchi, D. D., Misson, L., Dawson, T. E., and Goldstein, A. H.: What the towers don't see at night:
 923 nocturnal sap flow in trees and shrubs at two AmeriFlux sites in California, *Tree Physiology*, 27, 597-610,
 924 10.1093/treephys/27.4.597, 2007.

925 Foken, T. and Leclerc, M. Y.: Methods and limitations in validation of footprint models, *Agricultural and Forest*
 926 *Meteorology*, 127, 223-234, <https://doi.org/10.1016/j.agrformet.2004.07.015>, 2004.

927 Frolking, S., Roulet, N., and Fuglestad, J.: How northern peatlands influence the Earth's radiative budget:
 928 Sustained methane emission versus sustained carbon sequestration, *Journal of Geophysical Research:*
 929 *Biogeosciences*, 111, <https://doi.org/10.1029/2005JG000091>, 2006.

930 Gabrielli, E. C.: Partitioning Evapotranspiration in Forested Peatlands within the Western Boreal Plain, Fort
 931 McMurray, Alberta, Canada, Master's thesis, Wilfrid Laurier University, 2016.

932 Goetz, J. D. and Price, J. S.: Role of morphological structure and layering of Sphagnum and Tomenthypnum mosses
 933 on moss productivity and evaporation rates, *Canadian Journal of Soil Science*, 95, 109-124,
 934 <https://doi.org/10.4141/cjss-2014-092>, 2015.

935 Goulden, M. L., Litvak, M., and Miller, S. D.: Factors that control Typha marsh evapotranspiration, *Aquatic Botany*,
 936 86, 97-106, <https://doi.org/10.1016/j.aquabot.2006.09.005>, 2007.

937 Hathaway, J. M., Westbrook, C. J., Rooney, R. C., Petrone, R. M., and Langa, L. E.: Quantifying relative
 938 contributions of source waters from a subalpine wetland to downstream water bodies, *Hydrological*
 939 *Processes*, 36, e14679, <https://doi.org/10.1002/hyp.14679>, 2022.

940 Hrach, D. M., Petrone, R. M., Van Huizen, B., Green, A., and Khomik, M.: The Impact of Variable Horizon Shade
 941 on the Growing Season Energy Budget of a Subalpine Headwater Wetland, *Atmosphere*, 12, 1473,
 942 <https://doi.org/10.3390/atmos12111473>, 2021.

943 Hussey, B. H. and Odum, W. E.: Evapotranspiration in tidal marshes, *Estuaries*, 15, 59-67,
 944 <https://doi.org/10.2307/1352710>, 1992.

945 Jacobs, J. M., Mergelsberg, S. L., Lopera, A. F., and Myers, D. A.: Evapotranspiration from a wet prairie wetland
 946 under drought conditions: Paynes Prairie Preserve, Florida, USA, *Wetlands*, 22, 374-385,
 947 [https://doi.org/10.1672/0277-5212\(2002\)022\[0374:EFAWPW\]2.0.CO;2](https://doi.org/10.1672/0277-5212(2002)022[0374:EFAWPW]2.0.CO;2), 2002.

948 Jarvis, P. G. and McNaughton, K.: Stomatal control of transpiration: scaling up from leaf to region, in: *Advances in*
 949 *ecological research*, Elsevier, 1-49, [https://doi.org/10.1016/S0065-2504\(08\)60119-1](https://doi.org/10.1016/S0065-2504(08)60119-1), 1986.

950 Jiang, K. and Yu, L.: A method for partitioning ecosystem evapotranspiration based on fluxnet data., *Zenodo* [code],
 951 <https://doi.org/10.5281/zenodo.4816690>, 2021.

952 Kaimal, J. C. and Finnigan, J. J.: Atmospheric boundary layer flows: their structure and measurement, Oxford
 953 university press, <https://doi.org/10.1093/oso/9780195062397.001.0001>, 1994.

954 Ketcheson, S. J. and Price, J. S.: Characterization of the fluxes and stores of water within newly formed Sphagnum
 955 moss cushions and their environment, *Ecohydrology*, 7, 771-782, <https://doi.org/10.1002/eco.1399>, 2013.

956 Kettridge, N. and Waddington, J. M.: Towards quantifying the negative feedback regulation of peatland evaporation
 957 to drought, *Hydrological Processes*, 28, 3728-3740, <https://doi.org/10.1002/hyp.9898>, 2013.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

958 Kettridge, N., Lukenbach, M. C., Hokanson, K. J., Hopkinson, C., Devito, K. J., Petrone, R. M., Mendoza, C. A.,
 959 and Waddington, J. M.: Low Evapotranspiration Enhances the Resilience of Peatland Carbon Stocks to
 960 Fire, *Geophysical Research Letters*, 44, 9341-9349, <https://doi.org/10.1002/2017gl074186>, 2017.
 961 Kiniry, J. R., Williams, A. S., Reisner, L. M., Hatfield, J. L., and Kim, S.: Effects of two categorically differing
 962 emergent wetland plants on evapotranspiration, *Agrosystems, Geosciences & Environment*, 6, e20331,
 963 <https://doi.org/10.1002/agg2.20331>, 2023.
 964 Klosterhalfen, A., Moene, A. F., Schmidt, M., Scanlon, T. M., Vereecken, H., and Graf, A.: Sensitivity analysis of a
 965 source partitioning method for H₂O and CO₂ fluxes based on high frequency eddy covariance data:
 966 Findings from field data and large eddy simulations, *Agricultural and Forest Meteorology*, 265, 152-170,
 967 <https://doi.org/10.1016/j.agrformet.2018.11.003>, 2019a.
 968 Klosterhalfen, A., Graf, A., Brüggemann, N., Drüe, C., Esser, O., González-Dugo, M. P., Heinemann, G., Jacobs, C.
 969 M. J., Mauder, M., Moene, A. F., Ney, P., Pütz, T., Rebmann, C., Ramos Rodríguez, M., Scanlon, T. M.,
 970 Schmidt, M., Steinbrecher, R., Thomas, C. K., Valler, V., Zeeman, M. J., and Vereecken, H.: Source
 971 partitioning of H₂O and CO₂ fluxes based on high-frequency eddy covariance data: a comparison between
 972 study sites, *Biogeosciences*, 16, 1111-1132, <https://doi.org/10.5194/bg-16-1111-2019>, 2019b.
 973 Kokkonen, N., Laine, A. M., Männistö, E., Mehtätalo, L., Korrensalo, A., and Tuittila, E.-S.: Two Mechanisms
 974 Drive Changes in Boreal Peatland Photosynthesis Following Long-Term Water Level Drawdown: Species
 975 Turnover and Altered Photosynthetic Capacity, *Ecosystems*, 25, 1601-1618,
 976 <https://doi.org/10.1007/s10021-021-00736-3>, 2022.
 977 Kool, D., Agam, N., Lazarovitch, N., Heitman, J. L., Sauer, T. J., and Ben-Gal, A.: A review of approaches for
 978 evapotranspiration partitioning, *Agricultural and Forest Meteorology*, 184, 56-70,
 979 <https://doi.org/10.1016/j.agrformet.2013.09.003>, 2014.
 980 Lei, C.: Vegetation diversity in mountain peatland systems, University of Waterloo, 2021.
 981 Leifeld, J. and Menichetti, L.: The underappreciated potential of peatlands in global climate change mitigation
 982 strategies, *Nat Commun*, 9, 1071, <https://doi.org/10.1038/s41467-018-03406-6>, 2018.
 983 Leuning, R. and Judd, M. J.: The relative merits of open- and closed-path analysers for measurement of eddy fluxes,
 984 *Global Change Biology*, 2, 241-253, <https://doi.org/10.1111/j.1365-2486.1996.tb00076.x>, 1996.
 985 Li, X., Gentine, P., Lin, C., Zhou, S., Sun, Z., Zheng, Y., Liu, J., and Zheng, C.: A simple and objective method to
 986 partition evapotranspiration into transpiration and evaporation at eddy-covariance sites, *Agricultural and*
 987 *Forest Meteorology*, 265, 171-182, <https://doi.org/10.1016/j.agrformet.2018.11.017>, 2019.
 988 Lin, Y.-S., Medlyn, B. E., Duursma, R. A., Prentice, I. C., Wang, H., Baig, S., Eamus, D., de Dios, Victor R.,
 989 Mitchell, P., Ellsworth, D. S., de Beeck, M. O., Wallin, G., Uddling, J., Tarvainen, L., Linderson, M.-L.,
 990 Cernusak, L. A., Nippert, J. B., Ocheltree, T. W., Tissue, D. T., Martin-StPaul, N. K., Rogers, A., Warren, J.
 991 M., De Angelis, P., Hikosaka, K., Han, Q., Onoda, Y., Gimeno, T. E., Barton, C. V. M., Bennie, J., Bonal,
 992 D., Bosc, A., Löw, M., Macinins-Ng, C., Rey, A., Rowland, L., Setterfield, S. A., Tausz-Posch, S.,
 993 Zaragoza-Castells, J., Broadmeadow, M. S. J., Drake, J. E., Freeman, M., Ghannoum, O., Hutley,
 994 Lindsay B., Kelly, J. W., Kikuzawa, K., Kolari, P., Koyama, K., Limousin, J.-M., Meir, P., Lola da Costa, A.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

995 C., Mikkelsen, T. N., Salinas, N., Sun, W., and Wingate, L.: Optimal stomatal behaviour around the world,
 996 Nature Climate Change, 5, 459-464, <https://doi.org/10.1038/nclimate2550>, 2015.

997 Mäkelä, A., Berninger, F., and Hari, P.: Optimal control of gas exchange during drought: theoretical analysis, Annals
 998 of botany, 77, 461-468, <https://doi.org/10.1006/anbo.1996.0056>, 1996.

999 McCarter, C. P. R. and Price, J. S.: Ecohydrology of Sphagnum moss hummocks: mechanisms of capitula water
 1000 supply and simulated effects of evaporation, Ecohydrology, 7, 33-44, <https://doi.org/10.1002/eco.1313>,
 1001 2012.

1002 Medlyn, B. E., Duursma, R. A., Eamus, D., Ellsworth, D. S., Prentice, I. C., Barton, C. V., Crous, K. Y., De Angelis,
 1003 P., Freeman, M., and Wingate, L.: Reconciling the optimal and empirical approaches to modelling stomatal
 1004 conductance, Global change biology, 17, 2134-2144, <https://doi.org/10.1111/j.1365-2486.2010.02375.x>,
 1005 2011.

1006 Mitra, S., Wassmann, R., and Vlek, P. L. G.: An appraisal of global wetland area and its organic carbon stock,
 1007 Current Science, 88, 25-35, <http://www.jstor.org/stable/24110090>, 2005.

1008 Morison, M., van Beest, C., Macrae, M., Nwaishi, F., and Petrone, R.: Deeper burning in a boreal fen peatland 1-
 1009 year post-wildfire accelerates recovery trajectory of carbon dioxide uptake, Ecohydrology, 14, e2277,
 1010 <https://doi.org/10.1002/eco.2277>, 2021.

1011 Nardini, A. and Salleo, S.: Limitation of stomatal conductance by hydraulic traits: sensing or preventing xylem
 1012 cavitation?, Trees, 15, 14-24, <https://doi.org/10.1007/s004680000071>, 2000.

1013 Nelson, J. A.: Code and examples of how to estimate transpiration from eddy covariance data, GitHub [code], 2020.

1014 Nelson, Jacob A., Carvalhais, N., Cuntz, M., Delpierre, N., Knauer, J., Ogée, J., Migliavacca, M., Reichstein, M.,
 1015 and Jung, M.: Coupling Water and Carbon Fluxes to Constrain Estimates of Transpiration: The TEA
 1016 Algorithm, Journal of Geophysical Research: Biogeosciences, 123, 3617-3632,
 1017 <https://doi.org/10.1029/2018jg004727>, 2018.

1018 Nelson, J. A., Perez-Priego, O., Zhou, S., Poyatos, R., Zhang, Y., Blanken, P. D., Gimeno, T. E., Wohlfahrt, G.,
 1019 Desai, A. R., Gioli, B., Limousin, J. M., Bonal, D., Paul-Limoges, E., Scott, R. L., Varlagin, A., Fuchs, K.,
 1020 Montagnani, L., Wolf, S., Delpierre, N., Berveiller, D., Gharun, M., Beletti Marchesini, L., Gianelle, D.,
 1021 Sigut, L., Mammarella, I., Siebicke, L., Andrew Black, T., Knohl, A., Hortnagl, L., Magliulo, V., Besnard,
 1022 S., Weber, U., Carvalhais, N., Migliavacca, M., Reichstein, M., and Jung, M.: Ecosystem transpiration and
 1023 evaporation: Insights from three water flux partitioning methods across FLUXNET sites, Glob Chang Biol,
 1024 26, 6916-6930, <https://doi.org/10.1111/gcb.15314>, 2020.

1025 Novick, K. A., Oren, R., Stoy, P. C., Siqueira, M. B. S., and Katul, G. G.: Nocturnal evapotranspiration in eddy-
 1026 covariance records from three co-located ecosystems in the Southeastern U.S.: Implications for annual
 1027 fluxes, Agricultural and Forest Meteorology, 149, 1491-1504,
 1028 <https://doi.org/10.1016/j.agrformet.2009.04.005>, 2009.

1029 Pacheco-Cancino, P. A., Carrillo-López, R. F., Sepulveda-Jauregui, A., and Somos-Valenzuela, M. A.: Sphagnum
 1030 mosses, the impact of disturbances and anthropogenic management actions on their ecological role in CO2

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

1031 fluxes generated in peatland ecosystems, *Global Change Biology*, 30, e16972,
1032 <https://doi.org/10.1111/gcb.16972>, 2024.

1033 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy,
1034 A., Humphrey, M., Isaac, P., Polidori, D., Reichstein, M., Ribeca, A., van Ingen, C., Vuichard, N., Zhang,
1035 L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M.,
1036 Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J.,
1037 Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G., Boike, J., Bolstad, P.
1038 V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brümmer, C., Buchmann, N.,
1039 Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T.
1040 R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P.
1041 S., D'Andrea, E., da Rocha, H., Dai, X., Davis, K. J., Cinti, B. D., Grandcourt, A. d., Ligne, A. D., De
1042 Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., Tommasi, P. d., Dolman, H., Domingo, F.,
1043 Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Dušek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A.
1044 M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R.,
1045 Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson,
1046 A., Goded, I., Goeckede, M., Goldstein, A. H., Gough, C. M., Goulden, M. L., Graf, A., Griebel, A.,
1047 Gruening, C., Grünwald, T., Hammerle, A., Han, S., Han, X., Hansen, B. U., Hanson, C., Hatakka, J., He,
1048 Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-
1049 Korczynski, M., Janouš, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox,
1050 S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J.,
1051 Kutsch, W. L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Y., Liddell, M., Limousin,
1052 J.-M., Lion, M., Liska, A. J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D.,
1053 Lucas-Moffat, A., Lüers, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G.,
1054 Marcolla, B., Margolis, H. A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J. H.,
1055 Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A. M. S., Merbold, L., Meyer, W., Meyers, T.,
1056 Miller, S. D., Minerbi, S., Moderow, U., Monson, R. K., Montagnani, L., Moore, C. E., Moors, E.,
1057 Moreaux, V., Moureaux, C., Munger, J. W., Nakai, T., Neiryneck, J., Nesic, Z., Nicolini, G., Noormets, A.,
1058 Northwood, M., Noretto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J. E., Ourcival, J.-M., Papuga,
1059 S. A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R. P., Pilegaard,
1060 K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S. M., Rambal, S., Rannik, Ü., Raz-Yaseef, N.,
1061 Rebmann, C., Reed, D., Dios, V. R. d., Restrepo-Coupe, N., Reverter, B. R., Roland, M., Sabbatini, S.,
1062 Sachs, T., Saleska, S. R., Sánchez-Cañete, E. P., Sanchez-Mejia, Z. M., Schmid, H. P., Schmidt, M.,
1063 Schneider, K., Schrader, F., Schroder, I., Scott, R. L., Sedlák, P., Serrano-Ortíz, P., Shao, C., Shi, P.,
1064 Shironya, I., Siebicke, L., Šigut, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R. M.,
1065 Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tomassucci, M.,
1066 Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K.,
1067 Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J. P., Walter-Shea, E.,

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

1068 Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y.,
 1069 Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., and Papale, D.: The
 1070 FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Scientific Data*, 7,
 1071 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.

1072 Pérez-Priego, O. and Wutzler, T.: Partitioning eddy covariance ET using optimal approaches, GitHub [code], 2018.

1073 Pérez-Priego, O., Katul, G., Reichstein, M., El-Madany, T. S., Ahrens, B., Carrara, A., Scanlon, T. M., and
 1074 Migliavacca, M.: Partitioning eddy covariance water flux components using physiological and
 1075 micrometeorological approaches, *Journal of Geophysical Research: Biogeosciences*, 123, 3353-3370,
 1076 <https://doi.org/10.1029/2018JG004637>, 2018.

1077 Priestley, C. H. B. and Taylor, R. J.: On the Assessment of Surface Heat Flux and Evaporation Using Large-Scale
 1078 Parameters, *Monthly Weather Review*, 100, 81-92, [https://doi.org/10.1175/1520-0493\(1972\)100<0081:OTAOSH>2.3.CO;2](https://doi.org/10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2), 1972.

1079 Proctor, M.: Physiological ecology: water relations, light and temperature responses, carbon balance, in: *Bryophyte
 1080 ecology*, Springer, 333-381, https://doi.org/10.1007/978-94-009-5891-3_10, 1982.

1081 Reich, E. G., Samuels-Crow, K., Bradford, J. B., Litvak, M., Schlaepfer, D. R., and Ogle, K.: A Semi-Mechanistic
 1082 Model for Partitioning Evapotranspiration Reveals Transpiration Dominates the Water Flux in Drylands,
 1083 *Journal of Geophysical Research: Biogeosciences*, 129, e2023JG007914,
 1084 <https://doi.org/10.1029/2023jg007914>, 2024.

1085 Reichstein, M., Camps-Valls, G., Stevens, B., Jung, M., Denzler, J., Carvalhais, N., and Prabhat: Deep learning and
 1086 process understanding for data-driven Earth system science, *Nature*, 566, 195-204,
 1087 <https://doi.org/10.1038/s41586-019-0912-1>, 2019.

1088 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer, C., Buchmann, N.,
 1089 Gilmanov, T., Granier, A., Grünwald, T., Havránek, K., Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T.,
 1090 Lohila, A., Loustau, D., Matteucci, G., Meyers, T., Miglietta, F., Ourcival, J. M., Pumpanen, J., Rambal, S.,
 1091 Rotenberg, E., Sanz, M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, D., and Valentini, R.: On
 1092 the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved
 1093 algorithm, *Global Change Biology*, 11, 1424-1439, <https://doi.org/10.1111/j.1365-2486.2005.001002.x>,
 1094 2005.

1095 Rigden, A. J., Salvucci, G. D., Entekhabi, D., and Short Gianotti, D. J.: Partitioning Evapotranspiration Over the
 1096 Continental United States Using Weather Station Data, *Geophysical Research Letters*, 45, 9605-9613,
 1097 <https://doi.org/10.1029/2018gl079121>, 2018.

1098 Rouse, W. R.: The energy and water balance of high-latitude wetlands: controls and extrapolation, *Glob Chang Biol*,
 1099 6, 59-68, <https://doi.org/10.1046/j.1365-2486.2000.06013.x>, 2000.

1100 Scanlon, T. M. and Kustas, W. P.: Partitioning carbon dioxide and water vapor fluxes using correlation analysis,
 1101 *Agricultural and Forest Meteorology*, 150, 89-99, <https://doi.org/10.1016/j.agrformet.2009.09.005>, 2010.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

1103 Scanlon, T. M., Schmidt, D. F., and Skaggs, T. H.: Correlation-based flux partitioning of water vapor and carbon
1104 dioxide fluxes: Method simplification and estimation of canopy water use efficiency, *Agricultural and*
1105 *Forest Meteorology*, 279, 107732, <https://doi.org/10.1016/j.agrformet.2019.107732>, 2019.

1106 Scott, R. L. and Biederman, J. A.: Partitioning evapotranspiration using long-term carbon dioxide and water vapor
1107 fluxes, *Geophysical Research Letters*, 44, 6833-6840, <https://doi.org/10.1002/2017GL074324>, 2017.

1108 Speranskaya, L., Campbell, D. I., Lafleur, P. M., and Humphreys, E. R.: Peatland evaporation across hemispheres:
1109 contrasting controls and sensitivity to climate warming driven by plant functional types, *Biogeosciences*,
1110 21, 1173-1190, <https://doi.org/10.5194/bg-21-1173-2024>, 2024.

1111 Stapleton, A.: ETPartitioning [code], <https://github.com/AdamStapleton/ETPartitioning>, 2022.

1112 Stapleton, A., Eichmann, E., and Roantree, M.: A framework for constructing machine learning models with
1113 feature set optimisation for evapotranspiration partitioning, *Applied Computing and Geosciences*, 16,
1114 100105, <https://doi.org/10.1016/j.acags.2022.100105>, 2022.

1115 Stoy, P. C., El-Madany, T. S., Fisher, J. B., Gentine, P., Gerken, T., Good, S. P., Klosterhalfen, A., Liu, S., Miralles,
1116 D. G., Perez-Priego, O., Rigden, A. J., Skaggs, T. H., Wohlfahrt, G., Anderson, R. G., Coenders-Gerrits, A.
1117 M. J., Jung, M., Maes, W. H., Mammarella, I., Mauder, M., Migliavacca, M., Nelson, J. A., Poyatos, R.,
1118 Reichstein, M., Scott, R. L., and Wolf, S.: Reviews and syntheses: Turning the challenges of partitioning
1119 ecosystem evaporation and transpiration into opportunities, *Biogeosciences*, 16, 3747-3775,
1120 <https://doi.org/10.5194/bg-16-3747-2019>, 2019.

1121 Strack, M., Davidson, S. J., Hirano, T., and Dunn, C.: The potential of peatlands as nature-based climate solutions,
1122 *Current Climate Change Reports*, 8, 71-82, <https://doi.org/10.1007/s40641-022-00183-9>, 2022.

1123 Strack, M., Cagampan, J., Fard, G. H., Keith, A., Nugent, K., Rankin, T., Robinson, C., Strachan, I., Waddington, J.,
1124 and Xu, B.: Controls on plot-scale growing season CO₂ and CH₄ fluxes in restored peatlands: Do they
1125 differ from unrestored and natural sites?, <https://doi.org/10.19189/MaP.2015.OMB.216>, 2016.

1126 Street, L. E., Subke, J. A., Sommerkorn, M., Sloan, V., Ducrottoy, H., Phoenix, G. K., and Williams, M.: The role of
1127 mosses in carbon uptake and partitioning in arctic vegetation, *New Phytol*, 199, 163-175,
1128 <https://doi.org/10.1111/nph.12285>, 2013.

1129 Streich, S. C. and Westbrook, C. J.: Hydrological function of a mountain fen at low elevation under dry conditions,
1130 *Hydrological Processes*, 34, 244-257, <https://doi.org/10.1002/hyp.13579>, 2020.

1131 Sulman, B. N., Roman, D. T., Scanlon, T. M., Wang, L., and Novick, K. A.: Comparing methods for partitioning a
1132 decade of carbon dioxide and water vapor fluxes in a temperate forest, *Agricultural and Forest*
1133 *Meteorology*, 226-227, 229-245, <https://doi.org/10.1016/j.agrformet.2016.06.002>, 2016.

1134 Thomas, C., Martin, J. G., Göckede, M., Siqueira, M., Foken, T., Law, B. E., Loescher, H., and Katul, G.: Estimating
1135 daytime subcanopy respiration from conditional sampling methods applied to multi-scalar high frequency
1136 turbulence time series, *agricultural and forest meteorology*, 148, 1210-1229,
1137 <https://doi.org/10.1016/j.agrformet.2008.03.002>, 2008.

1138 van Beest, C.: Deeper Burning Increases Available Phosphorus, Promotes Moss Growth, and Carbon Dioxide
1139 Uptake in a Fen Peatland One-Year Post-Wildfire in Fort McMurray, AB, University of Waterloo, 2019.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Walker, A. P., Carter, K. R., Gu, L., Hanson, P. J., Malhotra, A., Norby, R. J., Sebestyen, S. D., Wulschleger, S. D., and Weston, D. J.: Biophysical drivers of seasonal variability in Sphagnum gross primary production in a northern temperate bog, *Journal of Geophysical Research: Biogeosciences*, 122, 1078-1097, <https://doi.org/10.1002/2016JG003711>, 2017.

Wang, Y.: Uncovering the understudied role of microtopography and ground cover on evapotranspiration partitioning in high-elevation wetlands in the Canadian Rocky Mountains, Doctoral dissertation, University of Waterloo, 2025.

Wang, Y. and Petrone, R. M.: Effects of microforms on the evaporation of peat-bryophyte-litter column in a montane peatland in Canadian Rocky Mountain, *Ecohydrology*, 16, e2516, <https://doi.org/10.1002/eco.2516>, 2022.

Wang, Y., Petrone, R. M., and Van Huizen, B.: The dependence of evaporative efficiency of vegetated surfaces on ground cover mass fractions in vegetated soils in mesic ecosystems, *Hydrological Processes*, 37, e15036, <https://doi.org/10.1002/hyp.15036>, 2023.

Warren, R. K., Pappas, C., Helbig, M., Chasmer, L. E., Berg, A. A., Baltzer, J. L., Quinton, W. L., and Sonnentag, O.: Minor contribution of overstorey transpiration to landscape evapotranspiration in boreal permafrost peatlands, *Ecohydrology*, 11, e1975, <https://doi.org/10.1002/eco.1975>, 2018.

Weaver, K. F., Morales, V., Dunn, S. L., Godde, K., and Weaver, P. F.: Pearson's and Spearman's Correlation, in: *An Introduction to Statistical Analysis in Research*, 435-471, <https://doi.org/10.1002/9781119454205.ch10>, 2017.

Webb, E. K., Pearman, G. I., and Leuning, R.: Correction of flux measurements for density effects due to heat and water vapour transfer, *Quarterly Journal of the Royal Meteorological Society*, 106, 85-100, <https://doi.org/10.1002/qj.49710644707>, 1980.

Wei, Z., Yoshimura, K., Wang, L., Miralles, D. G., Jasechko, S., and Lee, X.: Revisiting the contribution of transpiration to global terrestrial evapotranspiration, *Geophysical Research Letters*, 44, 2792-2801, <https://doi.org/10.1002/2016GL072235>, 2017.

Wu, J., Kutzbach, L., Jager, D., Wille, C., and Wilmking, M.: Evapotranspiration dynamics in a boreal peatland and its impact on the water and energy balance, *Journal of Geophysical Research: Biogeosciences*, 115, <https://doi.org/10.1029/2009JG001075>, 2010.

Xu, S., Ma, T., and Liu, Y.: Application of a multi-cylinder evapotranspirometer method for evapotranspiration measurements in wetlands, *Aquatic Botany*, 95, 45-50, <https://doi.org/10.1016/j.aquabot.2011.03.009>, 2011.

Yu, L., Zhou, S., Zhao, X., Gao, X., Jiang, K., Zhang, B., Cheng, L., Song, X., and Siddique, K. H.: Evapotranspiration partitioning based on leaf and ecosystem water use efficiency, *Water Resources Research*, 58, e2021WR030629, <https://doi.org/10.1029/2021WR030629>, 2022.

Zahn, E.: Processing Eddy-Covariance Data: Five ET Flux Partitioning Methods (2.0.0), GitHub [code], 2024.

Zahn, E., Ghannam, K., Chamecki, M., Moene, A. F., Kustas, W. P., Good, S., and Bou-Zeid, E.: Numerical investigation of observational flux partitioning methods for water vapor and carbon dioxide, *Journal of*

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

1176 Geophysical Research: Biogeosciences, 129, e2024JG008025, <https://doi.org/10.1029/2024JG008025>,
1177 2024.

1178 Zahn, E., Bou-Zeid, E., Good, S. P., Katul, G. G., Thomas, C. K., Ghannam, K., Smith, J. A., Chamecki, M., Dias,
1179 N. L., and Fuentes, J. D.: Direct partitioning of eddy-covariance water and carbon dioxide fluxes into
1180 ground and plant components, Agricultural and Forest Meteorology, 315, 108790,
1181 <https://doi.org/10.1016/j.agrformet.2021.108790>, 2022.

1182 Zhang, S., Zhang, J., Liu, B., Zhang, W., Gong, C., Jiang, M., and Lv, X.: Evapotranspiration partitioning using a
1183 simple isotope-based model in a semiarid marsh wetland in northeastern China, Hydrological Processes,
1184 32, 493-506, <https://doi.org/10.1002/hyp.11430>, 2018.

1185 Zhou, S., Yu, B., Huang, Y., and Wang, G.: Daily underlying water use efficiency for AmeriFlux sites, Journal of
1186 Geophysical Research: Biogeosciences, 120, 887-902, <https://doi.org/10.1002/2015jg002947>, 2015.

1187 Zhou, S., Yu, B., Zhang, Y., Huang, Y., and Wang, G.: Partitioning evapotranspiration based on the concept of
1188 underlying water use efficiency, Water Resources Research, 52, 1160-1175,
1189 <https://doi.org/10.1002/2015wr017766>, 2016.

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Line spacing: 1.5 lines