

Observing the role of wind-driven processes in the evolution of warm marine cloud properties

Vishnu Nair¹, Edward Gryspeerdt¹, Antti Arola², Antti Lipponen², and Timo Virtanen²

¹Department of Physics, Imperial College, London, UK

²Finnish Meteorological Institute, Kuopio, Finland

Correspondence: Vishnu Nair (v.nair16@imperial.ac.uk)

Abstract. The cloud droplet effective radius is a key variable when evaluating the interactions between aerosols and clouds. The activation of fine-sized sea salt from the ocean results in the formation of more but smaller cloud droplets (reducing the effective radius) in marine stratocumulus. Coarse sea spray aerosols are generated for high surface wind speeds and act as giant cloud condensation nuclei, which activate to form larger droplets. This increases the effective radius and initiates precipitation.

5 These high wind speeds also lead to enhanced moisture fluxes from the ocean surface. Although the opposing impacts of wind-driven fine and coarse marine sea spray aerosols have been documented, their observations have been limited to instantaneous satellite images. In this work, a novel framework is introduced that uses short-timescale observations of the temporal evolution of clouds to identify, isolate, and extract the process fingerprints of marine sea-salt and surface fluxes on stratocumulus cloud properties. This method shows that changes in droplet size previously attributed to aerosol are actually due to increases in
10 evaporation from the ocean surface due to high surface wind speeds. However, when this is accounted for, a clear impact of giant cloud condensation nuclei is observed, reducing cloud droplet number concentrations by initiating precipitation in polluted clouds. By isolating the causal aerosol impact on clouds from confounding factors, this method provides a pathway to improved constraints on the human forcing of the climate, whilst also demonstrating how marine aerosols limit the effectiveness of anthropogenic aerosol perturbations.

15 1 Introduction

Aerosols affect the Earth's radiation budget directly by reflecting and absorbing incoming solar radiation, and indirectly by acting as nucleation sites on which cloud droplets form (Twomey, 1974; Bellouin et al., 2020). Indirect effects occur by changing existing or new cloud properties and can have a cooling effect on global surface temperatures, hence offsetting a large part of the greenhouse gas warming (Stocker et al., 2013). This is by modifying the cloud reflectivity, both by affecting droplet
20 size and by driving time-dependent 'adjustments' (Albrecht, 1989), modifying the evolution of cloud properties (Glassmeier et al., 2021; Fons et al., 2023; Gryspeerdt et al., 2022a). The effective radiative forcing from aerosol-cloud interactions (ACI) is the largest source of uncertainty in human forcing of the climate (Andreae et al., 2005; Bellouin et al., 2020). ACI contributions are mainly due to the instantaneous Twomey effect (Twomey, 1974) which affects the cloud microphysical properties, or due to adjustments to the cloud macrophysical properties.

25 The adjustments of low clouds, such as marine stratocumulus (MSC), to aerosol perturbations are crucial to the Earth's radiation budget (Slingo, 1990). There are significant changes in the budget for even a small change in MSC coverage and thickness, with even a 4% increase in the global area covered by low-level stratus clouds offsetting a 2-3K temperature increase from higher atmospheric CO₂ concentrations (Randall et al., 1984). Two key measures of the properties of clouds that affect its radiative properties and hence the effect of MSC on the ocean albedo are the liquid water path (LWP, a measure of the total liquid water in a cloud), and the cloud droplet number concentration (N_d , a measure of the number of droplets in a cloud).

The cloud droplet effective radius, r_e , is physically, the area weighted mean of the cloud droplet size distribution. For a constant LWP, an increase in aerosol concentration (or cloud condensation nuclei, CCN) leads to an increase in N_d and a decrease in r_e (Twomey, 1977). More numerous smaller cloud droplets with a larger total droplet surface area reflect more sunlight, leading to an increase in cloud albedo. The decrease in r_e can also modify the cloud macrostructure by suppressing precipitation (due to weakened collision-coalescence between droplets) which causes both the LWP and albedo to increase (Albrecht, 1989). On the other hand, lower r_e can also suppress in-cloud droplet sedimentation and enhance cloud-top evaporative cooling which causes an increase in turbulent entrainment of free tropospheric air. Depending on the humidity of the entrained air, this can lead to a decrease or increase in LWP (Ackerman et al., 2004; Bretherton et al., 2007).

As well as the aerosol impact on cloud, cloud processes (such as precipitation) can modify aerosol. These feedback loops of processes that occur simultaneously are difficult to unravel and are further dependent on different cloud and meteorological regimes, complicating the identification of causal aerosol impacts on cloud (Fons et al., 2023). There are differences in the estimates of the climate effects due to ACI from global climate models and observations. The accuracy of the representations of these separate adjustment processes in models is believed to be one of the reasons for this discrepancy (Mülmenstädt et al., 2024a, b). This creates a requirement for strong observational constraints on cloud processes, to ensure that models have accurate representations of ACI.

There are multiple processes that can modify the cloud r_e , either by changing the cloud N_d or by changing the LWP. A key process via the N_d pathway is the additional activation of cloud droplets on CCN from different sources. The marine aerosol population generally consists of aerosols generated from the ocean surface: sea-salt from the action of wind stresses at the ocean surface (primary aerosols) and the emission of biogenic sulfur gases (dimethyl sulfide) which oxidizes in the atmosphere to form sulfate aerosols (secondary aerosols), aerosols entrained from the free troposphere, and aerosols from anthropogenic sources (sulfates, black and organic carbon) and natural sources (dust) from the continent that are advected to the marine boundary layer. The role of sea spray aerosols is unique as the consequence of the ACI can vary depending on the size of sea salt generated. The cloud-top r_e retrieved from satellite observations has systematically higher values over the ocean than over land which has a higher fine anthropogenic aerosol (radii $< 1\mu\text{m}$) concentration (Bréon et al., 2002; Kaufman et al., 2005).

55 Fine (radii $< 1\mu\text{m}$) and coarse (radii $> 1\mu\text{m}$) sea salt, both mechanically generated by wind, coexist over the ocean and form a significant part of the respective (fine and coarse) aerosol modes. It is well established that their generation and subsequent concentrations and size distributions are strongly related to the sea surface (10m) wind speeds. Perturbing clouds with fine sea salt (FSS) would lead to a reduction in r_e thereby brightening clouds. However, coarse marine aerosols (CMA), which are generated for surface wind speeds greater than 4ms^{-1} (Lehahn et al., 2010), act as 'giant' CCNs and have been hypothesized

60 to enhance warm rain by accelerating the formation of larger cloud droplets (larger r_e) (Ludlam, 1951; Feingold et al., 1999; Lasher-Trapp et al., 2001; Rosenfeld et al., 2002; L'Ecuyer et al., 2009; Lehahn et al., 2011; Jung et al., 2015). By triggering rain and reducing the LWP, CMA can break up and hence reduce cloud reflectivity. However, other model studies have questioned this impact, showing that this depends on the aerosol concentration; CMA have a negligible impact on precipitation initiation in clean clouds (Dagan et al., 2015), or no impact at all (Blyth et al., 2013). More recent research (Liu et al., 2022; Liu et al., 65 2025) suggests that there is an optimal combination which can effectively brighten clouds due to reduced r_e and enhanced cloud cover. Although it is not an anthropogenic aerosol, sea-salt sets the background “unpolluted” state of the cloud, modifying the aerosol forcing. This makes it essential to correctly represent fine and coarse marine aerosol in any model used for future climate assessment.

The impact of different cloud adjustments to fine and coarse sea salt has potential implications for geoengineering through 70 marine cloud brightening (MCB). MCB deliberately injects clouds with aerosols (ideally seawater spray), to lower r_e and increase reflectivity. In addition to setting the cloud background condition, thus determining how effective MCB can be in some conditions, the size distribution of the seeded aerosol now becomes important (Hoffmann and Feingold, 2021). Due to technical limitations, the seawater sprayed often contains coarse particles as well which can cause cloud breakup by initiating precipitation, making it important to consider the combined and opposing effects of fine and coarse sea salt and the possible 75 consequences in MCB projects.

Cloud r_e modification occurs via changes in cloud LWP as well. An increase in cloud LWP leads to a vertically deeper cloud which results in a higher cloud top r_e under the adiabaticity assumption. At stronger surface wind speeds, there is more evaporation at the ocean surface and a consequent increase in surface moisture flux. This leads to a moist marine boundary layer, a lower cloud base, and the formation of thicker clouds with a larger LWP (Chen et al., 2011). Therefore, increased 80 low-level horizontal wind speeds can enhance the emission of fine sea salt and giant CCNs while at the same time evaporating and transporting more moisture into clouds. This makes the wind speed (w_s) a major driver of cloud change over the ocean through multiple pathways (figure 3). It is vital to distinguish between these two causal pathways ($w_s - N_d - r_e$ versus $w_s - \text{LWP} - r_e$) and extract the “fingerprints” of these different processes to ensure accurate assessments of the climate response to anthropogenic aerosol changes.

85 Current observational studies are based on instantaneous satellite imagery. Recent observational assessments of the combined effects of FSS and CMA on cloud r_e and warm rain (Liu et al., 2022), and cloud radiative effects (Liu et al., 2025) used instantaneous measures of the LWP, fixed to separate the combined effects of meteorological factors. However, studies including the evolution of nocturnal clouds over 12 hours (model studies, (Hoffmann et al., 2020; Glassmeier et al., 2021)) and short timescales (day time observations, (Gryspeerd et al., 2022a)) reveal that the LWP can evolve differently based on the 90 initial N_d perturbation. The impacts of N_d on LWP are not accurately captured by instantaneous measurements (Arola et al., 2022; Zhang et al., 2024). Temporal evolution of cloud properties provides a separate pathway to isolate aerosol impact on cloud (Meskhidze et al., 2009; Glassmeier et al., 2021; Gryspeerd et al., 2021, 2022a; Fons et al., 2023), removes any reliance on predetermined/instantaneous and possibly confounded (by meteorological and other cloud controlling factors) $N_d - \text{LWP}$ relationships, and ensures the accuracy of interpreted causal relationships (Zhang et al., 2024).

95 This work measures the impact of different processes on the evolution of r_e over short time scales using a variety of observational data sets. A new framework is introduced to identify the role of different cloud processes in observations of cloud evolution. By assessing the impact of different meteorological properties on cloud evolution over a three hour time period, this framework isolates the individual fingerprints of fine and coarse sea salt, and surface fluxes, on LWP and N_d changes. The analysis highlights the importance of the initial/background state of the cloud in the temporal evolution of its macrophysical and microphysical properties. Non-aerosol processes have a more effective role in controlling the production of larger cloud droplets whereas coarse marine aerosols limit the overall impact of anthropogenic aerosols on cloud properties in very polluted conditions.

2 Data and Methods

2.1 Observational and meteorological reanalysis data

105 The properties of the cloud field are calculated from the Moderate Resolution Imaging Spectroradiometer (MODIS) cloud retrieval (Platnick et al., 2017) onboard both the Aqua and Terra satellites over the 10-year period 2007-17 (inclusive). The Terra and Aqua satellites provide information about cloud properties three hours apart from the two daytime overpasses. The cloud N_d and LWP is derived from the values of r_e and the cloud optical depth τ from the $2.1 \mu\text{m}$ retrievals (Quaas et al., 2006; Grosvenor et al., 2018). The cloud N_d is calculated following the sampling criteria in Grosvenor et al. (2018), and sampling strategy G18 in Gryspeerdt et al. (2022b). Only single-layered, liquid-phase clouds are considered. This strategy also excludes pixels with solar zenith $> 65^\circ$ and satellite zenith angle $> 55^\circ$ to account for uncertainties at high viewing angles. An adiabatic cloud is assumed for all the selected pixels. The LWP is calculated using all available liquid pixels. These data are then aggregated to a $1^\circ \times 1^\circ$ latitude-longitude grid.

Following Gryspeerdt et al. (2021), boundary layer winds are used to account for advection between the observation times of the different instruments, restricting analysis to grid boxes where N_d and LWP are available for both instruments. The cloud field is advected using ERA5 reanalysis wind fields at 1000hPa. The advection is calculated on a $0.25^\circ \times 0.25^\circ$ resolution grid (as the movement of the cloud fields over 3h is expected to be less than 1°). Each grid box on the fine resolution grid is treated as a parcel trajectory and advected using the wind fields. The Aqua data are then sampled at the end points of these trajectories, and aggregated to a $1^\circ \times 1^\circ$ grid.

120 Surface wind speeds are obtained from ERA5, reanalysis product (on a $1^\circ \times 1^\circ$ grid) that offer an estimate of global atmospheric conditions that are collocated with the MODIS observations (Hersbach et al., 2023). Precipitating and not-precipitating cases are differentiated using the probability of precipitation (PoP) at each MODIS grid point, based on the proportion of liquid r_e retrievals greater than $15 \mu\text{m}$ (Rosenfeld et al., 2002). This is calculated from the MODIS level 3 daily gridded product (MOD08_D3) using the r_e histogram counts. The PoP is the ratio of small drops (sum of all histogram bins less than $15 \mu\text{m}$) to all drops in the distribution/histogram (sum of all bins). If $\text{PoP} > 70\%$, then the gridbox is considered to be precipitating. The results in sections 3.1 and 3.2 are from a region within the southeastern Atlantic stratocumulus deck between 40°S to 10°N and 30°W to 10°E .

2.2 Difference of rates (DoR)

The temporal evolution of the N_d and LWP (dN_d/dt and $dLWP/dt$ respectively), is obtained by calculating the difference in properties between the two daytime satellite overpasses, which are approximately three hours apart- at 10:30 and 13:30 local solar time for Terra and Aqua respectively. Joint histograms in the N_d - LWP space are then generated by binning the cloud data as a function of their initial N_d and LWP (Gryspeerdt et al., 2022a). These are then converted to relative rates of changes by dividing the differences first with the bin widths (N_d bins for dN_d , and the LWP bins for $dLWP$), and then with the time step of 3 hours (time difference between Aqua and Terra observations). The final result is expressed as a percentage with units of (%/hr) by multiplying with 100.

By strictly controlling the initial state of the cloud, we account for the confounding impact of existing meteorological variables which can change the properties of aerosol and the cloud simultaneously, and could introduce spurious correlations on the development of N_d -LWP relationships. If the clouds are advected across regions with a large gradient in meteorological properties, this would result in a large change in the cloud properties owing to how correlated the cloud is to a strong climatological change. Clouds with a high (low) initial value of LWP or N_d is likely to show a decrease (increase) in LWP or N_d , which is consistent with a 'regression to the mean' effect. This can happen as a statistical effect, where even when the cloud is remaining stationary, a positively biased first measurement (of r_e) is followed by a smaller second measurement. Since r_e is positively correlated with LWP and negatively correlated with N_d , this shows up as a highly negative $dLWP$ and a large positive dN_d . Previous studies (Eastman et al., 2016; Eastman and Wood, 2016) accounted for this by looking at anomalous changes across Lagrangian trajectories by removing seasonal means for day and night separately. On the other hand, it was shown by Gryspeerdt et al. (2021) (where $dLWP$ and dN_d were calculated in a similar way from MODIS Terra and Aqua) that the 'flowfields' (the rate of change of N_d and LWP) do not look the same when $dLWP$ and dN_d are binned by the final LWP and N_d . If there was indeed a regression to the mean effect, the flowfields should have looked the same when calculated from either direction. As stated in Gryspeerdt et al. (2021), while this does not completely rule out the impact of retrieval biases and the regression to the mean effect, it does rule out the possibility of the results being a statistical artefact caused by random biases.

Here, the difference-of-rates (DoR) method is introduced which accounts for these effects. The relative rates of change (in %/hr) are calculated by separating or stratifying the cloud population based on a cloud/meteorological variable (such as surface (10 m) wind speeds, w_s), and calculating the differences with respect to a reference data set. For example, in figure 2, the DoR (presented herewith with a symbol Δ) with respect to w_s is calculated as

$$\Delta_{w_s} dN_d = \Delta_{(w_s^+)} dN_d - \Delta_{(w_s^-)} dN_d \quad (1)$$

where $w_s^+ = w_s > 4 \text{ ms}^{-1}$ and $w_s^- = w_s < 4 \text{ ms}^{-1}$. Here, the cloud population with w_s^- is the reference data set. Multiple DoRs can be calculated by splitting w_s^+ into smaller bins/data ranges (different columns in figure 2).




Figure 1. The effect of precipitation on dN_d/dt (a,b) and $dLWP/dt$ (d,e). (a) dN_d/dt for precipitating clouds ($\text{PoP} > 70\%$), (b) dN_d/dt for not-precipitating clouds ($\text{PoP} < 70\%$), (c) $\Delta_{\text{PoP}} dN_d$, the difference between sub figures a and b, (d) $dLWP/dt$ for precipitating clouds ($\text{PoP} > 70\%$), (e) $dLWP/dt$ for not-precipitating clouds ($\text{PoP} < 70\%$), (f) $\Delta_{\text{PoP}} dLWP = d - e$.

3 Results and Discussion

3.1 The precipitation fingerprint

In addition to acting as a sink for the cloud N_d through the sedimentation of droplets, precipitation plays a key role in the scavenging of CCN (wet or below-cloud scavenging), which in turn can reduce N_d . The effects of precipitation are usually seen primarily in the (upper) left quadrant ($LWP > 50\text{gm}^{-2}$, $N_d < 50\text{cm}^{-3}$), i.e., for clouds with a high initial LWP and a low N_d . An overall positive change is seen in the dN_d field in this region for both precipitating and non-precipitating clouds (red region in figures 1a,b), which is possibly a regression to the mean effect (discussed in the next paragraph). In addition to precipitation, other processes such as the primary production of CCN from sea spray, and entrainment of aerosols from the free troposphere (especially closer to the coast) can possibly act as significant sources of N_d for clouds with an initially low N_d . However, precipitation rates as low as 1mm d^{-1} have been shown to be effective in reducing N_d by a factor of three over the SE Pacific (Wood et al., 2012). The DoRs between precipitating and not-precipitating clouds (figure 1c) reveal that precipitation acts as a sink for the cloud N_d , with a reduction of N_d observed in more strongly precipitating cases. Precipitation results in a smaller overall net increase in dN_d (i.e., the change in N_d over three hours) with figure 1a showing lighter reds and darker greens. There is a smaller decrease (larger increase) in dN_d for not-precipitating clouds in figure 1b (darker reds and lighter greens). Consequently, the corresponding DoR, $\Delta_{PoPd}N_d$ (difference between figures 1a and b, i.e., a- b), is negative (figure 1c).

The few positive values (red) in the DoR field in figure 1c are possibly not-precipitating clouds which were not filtered out using the r_e threshold. The positive and negative regions in figures 1a and b may be partly driven by the regression to the mean effect by using a filter in r_e , which is also used to calculate N_d and LWP. Similar patterns were obtained as in the figures 1a-c when an independent data source (Eastman et al., 2019) for precipitation was used (supplementary section A). The MODIS filters were subsequently chosen to identify precipitating clouds for the rest of the analysis. Using r_e as a measure of precipitation allows the precipitation data at the start of the time step to be included, such that the impact of precipitation on the cloud evolution is identified (rather than the impact of cloud evolution on precipitation, as is obtained using precipitation from the later overpass at the end of the timestep).

Droplet sedimentation at the cloud-top entrainment interfacial layer (EIL) depletes liquid water from this zone, leading to reduced entrainment and thicker clouds (high LWP) (Ackerman et al., 2004; Bretherton et al., 2007). In precipitating clouds the r_e is generally higher leading to droplets sedimenting out of the EIL. Conversely, in not-precipitating clouds, the r_e is smaller, and there are more, smaller cloud droplets in the entrainment interfacial layer at the cloud top. This leads to evaporative enhancement of entrainment of free tropospheric air, leading to a thinner cloud layer or a higher decrease (or a smaller increase) in LWP in not-precipitating clouds compared to precipitating ones. Consequently, the DoR for LWP is positive in figure 1f everywhere except in the region corresponding to the strongly negative region in figure 1c for N_d . Here, strongly precipitating clouds lose more liquid water (as rain) resulting in a more negative (less positive) change in LWP.




Figure 2. Effect of surface wind speed on N_d and LWP. All figures show DoRs, $\Delta dN_d = \Delta_{(w_s^+)} dN_d - \Delta_{(w_s^-)} dN_d$ and $\Delta dLWP = \Delta_{(w_s^+)} dLWP - \Delta_{(w_s^-)} dLWP$, where $w_s^- = w_s < 4\text{ms}^{-1}$ and, $w_s^+ =$ a) $4 < w_s < 7\text{ms}^{-1}$, b) $7 < w_s < 9\text{ms}^{-1}$, c) $9 < w_s < 12\text{ms}^{-1}$, d) $w_s > 12\text{ms}^{-1}$ for precipitating clouds. Figures (e) - (h) are ΔdN_d for not-precipitating clouds, figures (i) - (l) are $\Delta dLWP$ for precipitating clouds and figures (m)- (p) are $\Delta dLWP$ for not-precipitating clouds for similar w_s ranges.

3.2 Disentangling the impact of wind driven processes on cloud effective radius

With the new framework effectively extracting the precipitation fingerprint, we apply this technique to identify the different processes that modify the r_e . To investigate the simultaneous effects of FSS and CMA in altering r_e through changes in N_d , the DoRs are calculated by stratifying the data by surface wind speeds. Horizontal low-level (10m) winds have been shown to be strongly correlated with sea salt production (Lewis and Schwartz, 2004). In contrast to figure 1, all the plots in figure 2 are DoRs, $\Delta_{w_s} dN_d$ and $\Delta_{w_s} dLWP$, calculated for different wind speed ranges and with respect to a reference dataset with $w_s < 4\text{ms}^{-1}$ (as detailed in the methods section). Additionally, we treat initially precipitating and not-precipitating clouds separately as the impact of aerosol-induced precipitation suppression works differently for these two categories.

For initially precipitating clouds, precipitation suppression is expected to increase at higher w_s with the introduction of additional FSS. Therefore, dN_d would tend to be less (more) negative (positive) as w_s increases (due to reduced in-cloud scavenging). Consequently, $\Delta_{w_s} dN_d$ would be negative (blue) as seen in the left quadrant in figures 2a to d.

A positive fingerprint, consistent with FSS acting as CCN, strengthens with wind speed and dominates the precipitation effect at wind speeds greater than 12ms^{-1} in figure 2d. In addition to the increased aerosol burden, high w_s leads to stronger updrafts at the cloud base (Colón-Robles et al., 2006) increasing the activation of FSS and the formation of new droplets.

The effect of CMA is seen in not-precipitating clouds with a high initial N_d ($> 100\text{cm}^{-3}$) in figures 2e - h, with a negative trend in $\Delta_{w_s} dN_d$ consistent with the results from (Feingold et al., 1999; Yin et al., 2000). The CMA are usually the first to activate and form droplets at the cloud base. This depletes the supersaturation here, thereby inhibiting the activation of FSS into smaller droplets. This skews the droplet size distribution to larger sizes, enhancing the collision-coalescence rate (Rosenfeld et al., 2002). Both processes lead to a higher r_e which is a key precursor to precipitation initiation.

A negative $\Delta_{w_s} dN_d$ region is consistent with a giant CCN-induced reduction in dN_d . In addition to fewer new activated droplets, the smaller dN_d could also be due to giant CCN-induced precipitation (in initially not-precipitating clouds). Once giant CCNs are activated, condensational growth and collision-coalescence to raindrop sizes are expedited within this time scale (of three hours). Increasing w_s leads to the formation of more CMA which shows up as a stronger signal in the DoR. As expected, there is no perceptible impact of CMA on already raining clouds as drizzle is already active.

A relationship between CMA and r_e by altering N_d can clearly be seen. However, there is a strong positive signal in $\Delta dLWP$ across all initial N_d suggesting an alternate pathway to explain the $w_s - r_e$ correlation. Stronger surface winds over the ocean lead to an increase in surface fluxes through increased evaporation (Kazil et al., 2016; Eastman et al., 2022). This moistens the marine boundary layer and increases the moisture available at the cloud base. This can increase the cloud depth either by lowering the cloud base through condensation or increasing the cloud top height by entrainment (driven by buoyant production of kinetic energy in the updrafts (Kazil et al., 2016)). The current methodology does not allow us to distinguish between these two effects. In addition, higher wind speeds enhance wave formation - these waves can break and produce white caps and sea spray, which enhances the sea-to-air latent heat flux. This is an additional source to the cloud LWP leading to thicker clouds with larger droplets, and corresponding larger $dLWP$ over the observed period (figures 2i - l).




Figure 3. Variability in dN_d across the globe for different wind speed ranges shown here as the DoR $\Delta_{w_s} dN_d$ (in cm^{-3}) for initial $30 < N_d < 100 \text{ cm}^{-3}$. The boxes show the Sc decks/regions, data from which are used to generate the DoR joint histograms for the south east Atlantic (SEA), north Pacific (NP) and the south pacific (SP).

3.3 Geographical variability

225 The analysis presented thus far was restricted to the southeastern Atlantic (SEA) Sc clouds. In this section, we extend the analysis to the rest of the globe. DoR plots similar to figure 2 are generated for the stratocumulus decks over the north and south Pacific oceans (figures C1 and C2). In the north Pacific (NP) Sc deck, results are broadly consistent with those found over the SEA deck: for initially non-precipitating clouds, the influence of GCCN is evident, with a decrease in N_d at high initial N_d and $\text{LWP} > 30 \text{ gm}^{-2}$ (figures C1 e-h). For precipitating clouds, however, the impact of precipitation dominates the

230 N_d response, even at very high w_s (figures C1 a-d). Similar to the SEA deck, LWP in the NP deck increases with w_s (figure C1 i-p) for both not-precipitating and precipitating clouds. In contrast, for the south Pacific (SP) Sc clouds, the signals associated with the same processes are less clear when using the DoR method - the changes in N_d appear to show similar behavior as the NP clouds for both not-precipitating and precipitating clouds (figure C2 a-h), as does the change in LWP for precipitating




Figure 4. Variability in dN_d across the globe for different wind speed ranges ($\Delta_{w_s} dN_d$ in cm^{-3}) for $N_d > 100 \text{ cm}^{-3}$.

clouds. For not-precipitating clouds (figure C2 m-p), there is a strong negative and positive presence with no obvious pattern emerging.

To further investigate this behavior, global maps of the DoRs of dN_d ($\Delta_{w_s} dN_d$ in cm^{-3}) and dLWP ($\Delta_{w_s} \text{dLWP}$ in gm^{-2}) are analyzed for different w_s ranges. These are plotted separately for clouds with a mid-range ($30 < N_d < 100 \text{ cm}^{-3}$) and high ($N_d > 100 \text{ cm}^{-3}$) initial N_d . Figure 3, showing not-precipitating clouds with an initial N_d between 30 and 100 cm^{-3} , reveals $\Delta_{w_s} dN_d$ transitioning from positive to negative values in the northern hemisphere and regions off the shore of the African and south American continents with increasing w_s . The areas enclosed in the three boxes are the grid points used to make the joint histograms in figures 2, C1 and C2 for the SEA, NP and SP decks respectively.

It is possible that the negative DoRs are driven by competition for activation between sea salt and sulfates advected from the continent. Sea-salt aerosols are an important component of the so-called natural 'background aerosol/CCN' population. Introduction of more sea-salt aerosols in already polluted background states (over and offshore from the continent) results in competition between CCN particles for the liquid water/supersaturation required to activate to form cloud droplets. At higher wind speeds, sea salt becomes a dominant contributor to background aerosol concentrations, especially in the coarse mode.

The increased surface area enhances condensation of water vapor thereby reducing maximum supersaturation. This inhibits the activation of new sulfates or other small aerosols leading to a net reduction in N_d (Ghan et al., 1998; Fossum et al., 2020) or a smaller increase in dN_d compared to low w_s conditions. This could explain the negative values (blue regions) over the northern hemisphere and along the coast of the continents in the southern hemisphere. More pristine conditions exist further away from the continental coastlines in the southern hemisphere where the competition is less prominent, leading to almost constant changes (positive) as w_s increases in these regions.

The influence of GCCN in initiating precipitation is illustrated in figure 4, which shows $\Delta_{w_s} dN_d$ (in cm^{-3}) for different w_s for initially not-precipitating clouds with $N_d > 100 \text{ cm}^{-3}$. Outside the tropics, a robust decrease in dN_d is observed over the oceans in both hemispheres driven by precipitation.

Figure 5 shows the global distribution of DoRs of dLWP ($\Delta_{w_s} \text{LWP}$ in gm^{-2}) for not-precipitating clouds with an initial droplet number concentration $N_d > 30 \text{ cm}^{-3}$. A clear increase in dLWP with increasing w_s is observed. This signal is particularly pronounced over the subtropical and mid-latitude oceans in both hemispheres. Notably, a distinct increase in dLWP is evident for the SP deck, especially near the coast. This coastal signal is not clearly captured in the DoR joint histograms (figures C2 m-p), likely because those include clouds farther offshore, where stratocumulus transitions to cumulus regimes. For initially precipitating clouds, even though the change in LWP appears more positive (figure C3), no distinct signals can be observed in the maps even though the DoR histograms capture the increase in LWP. Coarsening the grid to a $5^\circ \times 5^\circ$ grid also did not reveal any clear patterns (not shown) in the maps. However, a combination of the DoR method along with global maps of the DoRs can provide a strong picture of the effects of wind-driven processes over the evolution of Sc decks across the globe along with any hemispherical differences.

4 Conclusions and Outlook

This article highlights the effectiveness of two different pathways through which surface wind-driven processes can modify the cloud droplet effective radius r_e . Using observations of the temporal evolution of clouds, process fingerprints of the effects of marine aerosols on the cloud properties were extracted. Giant CCNs were shown to reduce cloud N_d (figures 2e-h and 4) through two different pathways as illustrated in figure 6:

Pathway 1a: initiating precipitation in clouds with an initially high droplet number concentration.

Pathway 1b: depleting supersaturation by activating to form larger droplets. This increases competition for supersaturation among the remaining CCNs leading to the activation of fewer new droplets.

In addition to the role of fine and giant CCNs, we identify and highlight the role of wind-driven surface fluxes in the thickening of marine stratocumulus clouds and therefore in the increase of r_e (pathway 2 in figure 6). Similar modification in moisture in the STBL through stronger surface fluxes in response to higher surface wind speeds have been shown before (Kazil et al., 2016; Eastman et al., 2022).




Figure 5. Variability in dLWP across the globe for different wind speed ranges (Δ_{w_s} LWP in gm^{-2}) for all non-precipitating clouds with an initial $N_d > 30\text{cm}^{-3}$.

The percentage changes of r_e due to changes in N_d and LWP via the different pathways can be quantified by using equations B1 and B2, and using values for $d(\ln N_d)$ and $d(\ln \text{LWP})$ from figures 2f and j respectively. In figure 2f, for $N_d > 100\text{ cm}^{-3}$ and LWP across the range here, $d(\ln N_d)$ is approximately -2.5 % (light blue in the colorbar). Similarly, in figure 2n, $d(\ln \text{LWP})$ is approximately 5% for $N_d > 100$. This corresponds to an increase in $d(\ln r_e)$ of 7.5% and 30% respectively when used in equations B1 and B2 (details in appendix section B). This clearly identifies the second pathway - via increased surface fluxes - as the more dominant physical process that increases r_e , while simultaneously obscuring the impact of marine aerosol on clouds.

Investigating the N_d and LWP responses to changes in w_s across the globe yielded interesting results which makes it important to address the interactions of non-sea-salt aerosols with fine and coarse sea salt. Near-surface wind speeds have been shown to control the formation of sulfate aerosols through oceanic emissions of dimethyl sulfide (Huebert et al., 2004). Additionally, the presence of continents, and consequently a more polluted background state in the northern hemisphere causes different responses in cloud N_d and LWP compared to the more pristine conditions over the oceans in the southern hemisphere.




Figure 6. Different pathways to explain the $w_e - N_d/LWP - r_e$ correlation.

290 At low wind speeds, sulfate concentration exceeds that of sea salt, especially in the accumulation mode which provides the majority of the cloud CCN. However, as wind speed increases, the concentration of sea salt in the accumulation mode increases and they end up dominating the CCNs in cleaner clouds. For high (low) sulfate concentrations and weak (strong) updrafts, sea salt will reduce (increase) the maximum supersaturation and reduce the number concentration of activated sulfate particles (Ghan et al., 1998) as seen in figure 3. The wind-driven sea salt emissions, while acting as new CCN, can also suppress aerosol

295 nucleation by removing nucleating molecules.

However, sea salt is always preferentially activated as CCN at lower supersaturations as they are larger than sulfate particles. Also, majority of the marine aerosols in the coarse mode, ie, greater than 1μ , will be sea salt and hence we do not expect other aerosol sources to interfere with the GCCN results presented here.

It is important to acknowledge the possible role of surface winds on stratocumulus topped boundary layer (STBL) dynamics.

300 Specifically, the role of surface shear on cloud-top entrainment rate. Entrainment of free tropospheric (FT) aerosols across the cloud-top entrainment interfacial layer (EIL) can lead to the introduction of CCN. This possibly affects our results if a direct correlation exists between wind-generated surface shear and entrainment rates. However, even though there is considerable evidence of wind shear across the *inversion* enhancing the entrainment rate of dry, warm FT air and reducing the cloud fraction and LWP (Schulz and Mellado, 2018; Wang et al., 2008, 2012; Zamora Zapata et al., 2021), the effect of surface shear on

305 entrainment in STBL is still unclear. Increased low level vertical wind shear can contribute to a turbulent and well-mixed STBL. However, any increase to entrainment (a turbulent kinetic energy (TKE) consuming process) is dependent on the availability of the surface shear-generated TKE at the EIL. This surface shear generated TKE must be transferred through the entire depth of the STBL to the EIL as it goes through the energy cascade process of turbulence, before it can be used to drive entrainment. However, studies on the STBL specifically looking at the role of surface shear and cloud top shear failed to see conclusive
 310 evidence on an increase in entrainment rates due to *only* surface shear (Zamora Zapata et al., 2021). Studies on the interaction of a constant large-scale wind speed with the STBL showed that it is possible through buoyancy driven dynamics (rather than shear driven) for *geostrophic* wind to promote STBL growth and enhance entrainment throughout the diurnal cycle (Kazil et al., 2016). Higher surface moisture flux at increased wind speed boosts the latent heat release and buoyant production of TKE in cloud updrafts leading to increased entrainment. At the same time, they point out that features of boundary layer dynamics that
 315 determine entrainment exist, but require more in-depth study. This suggests that we cannot completely discount the possibility that wind driven entrainment of aerosols from the FT can affect the results in this work, especially in figure 2.

Additionally, depending on the humidity of the entrained FT air, there could be an increase or decrease in cloud LWP (Ackerman et al., 2004). Accounting for the contribution of surface winds to cloud-top entrainment and hence N_d and LWP would require controlling for both FT aerosol concentration and relative humidity (in addition to wind speeds) which complicates
 320 the analysis using DoRs. Since we expect the w_s - surface shear - entrainment correlation to be weak, and the w_s - sea salt correlation to be strong and dominate the N_d - LWP phase space, we choose to focus on the latter and reserve the former analysis for future work.

While retrieval biases can still affect the quantification of the initial state, the focus on time development reduces the impact of correlated errors in N_d and LWP that affected previous studies (Arola et al., 2022). Further studies are required to assess
 325 the impact of other factors on these fingerprints, particularly the diurnal cycle (Zhang et al., 2024). Including the effects of the diurnal cycle will account for variability in the N_d -LWP relationship over longer time scales. Observations from geostationary satellites are ideal for these analyses by evaluating cloud evolution through Lagrangian trajectories over longer time scales than those considered here. These will also have the advantage of using smaller time steps than the three hours considered in this work.

330 The impact of fine sea salt follows previous observational studies, increasing N_d , but coarse marine aerosol is shown to decrease N_d , particularly in initially not-precipitating cases with a high N_d . This shows that even small amounts of coarse aerosol can limit the efficacy of anthropogenic aerosol injections, providing an important constraint on the cloud response to aerosol and limiting the effectiveness of proposed marine cloud brightening programs.

Appendix A: Investigating regression to the mean effects using an independent data source for precipitation

335 It is possible that the positive and negative regions in figures 1a and b are partly driven by the regression to the mean effect. Whereby, a positively biased first measurement would likely be followed by a smaller second measurement. By applying a threshold on r_e (which is used to calculate both N_d and LWP) to distinguish between precipitating and not-precipitating clouds, there is a chance that similar regression to mean effects are inadvertently introduced in the DoRs in figure 1c.

We perform an alternate analysis using an independent data source for precipitation: warm rain rates inferred from AMSR/E and AMSR/2 89GHz passive microwave brightness temperatures trained using CloudSat rain rate observations (Eastman et al., 2019). The results for DoRs from the dataset provided by Eastman et al. (2019) in figures A1 and A2 suggest patterns similar to those in figures 1 and 2. This suggests that the patterns are indicative of precipitation effects rather than the regression to the mean (which might still be at play but less dominant). We refrain from using the data set from Eastman et al. (2019) to identify precipitating clouds in the main manuscript as these are collocated with data from Aqua, which is at the end of the time step in the context of this manuscript. By using the effective radius as a measure of precipitation we are using more information of the cloud microphysics and obtain data from the start of the time step. This allows us to identify the role of precipitation and other processes during the evolution of the cloud. Original CloudSat rain rate observations were also considered, but these are too sparse/patchy to provide reliable results.

Appendix B: Cloud r_e sensitivities to changes in LWP and N_d

350 The cloud N_d and LWP are calculated using

$$N_d = \frac{\sqrt{5}}{2\pi k \sqrt{\rho_l Q_{ext}}} \sqrt{f_{ad} \Gamma} \tau_c^{1/2} r_e^{-5/2},$$

$$\text{LWP} = 5/9 r_e \rho_l \tau_c.$$

where Q_{ext} is the extinction efficiency factor (unitless), f_{ad} is the adiabaticity fraction, τ_c is the cloud optical depth (unitless), $k = (r_v/r_e)^3$ where r_v is the volume-mean droplet radius, ρ_w is the density of liquid water (kg/m^3)

355 The change in r_e with a change in N_d can be represented as

$$\left. \frac{\partial \ln r_e}{\partial \ln N_d} \right|_{\text{LWP}} = -\frac{1}{3}, \tag{B1}$$

$$\left. \frac{\partial \ln r_e}{\partial \ln \text{LWP}} \right|_{N_d} = \frac{1}{6}. \tag{B2}$$

Appendix C: Geographical variability in Sc evolution




Figure A1. Calculation of DoRs as in figure 1 using AMSR data.




Figure A2. Effect of surface wind speed on N_d and LWP. As in figure 2 but with CloudSat data.




Figure C1. As in figure 2 but for the north pacific Sc deck.




Figure C2. As in figure 2 but for the south pacific Sc deck.




Figure C3. As in figure 5 but for precipitating clouds.

Data availability. The data supporting the conclusions, along with the code used for processing this data and generating the figures in this
360 study are available with the identifier <https://doi.org/10.5281/zenodo.16882487>.

Author contributions. EG and VN designed the study. VN performed the analysis and wrote the paper. All of the authors assisted in the interpretation of the results and commented on the paper.

Competing interests. The authors declare that no conflicting interests are present.

Acknowledgements. The authors acknowledge funding from Horizon Europe programme under Grant Agreement No 101137680 via project
365 CERTAINTY (Cloud-aERosol inTeractions & their impActs IN The earth sYstem).

References

- Ackerman, A. S., Kirkpatrick, M. P., Stevens, D. E., and Toon, O. B.: The impact of humidity above stratiform clouds on indirect aerosol climate forcing, *Nature*, <https://doi.org/10.1038/nature03174>, 2004.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <http://www.jstor.org/stable/1704234>, 1989.
- Andreae, M. O., Jones, C. D., and Cox, P. M.: Strong present-day aerosol cooling implies a hot future, *Nature*, 435, 1187–1190, <https://doi.org/10.1038/nature03671>, PMID: 15988515, 2005.
- Arola, A., Lipponen, A., Kolmonen, P., Virtanen, T. H., Bellouin, N., Grosvenor, D. P., Gryspeerdt, E., Quaas, J., and Kokkola, H.: Aerosol effects on clouds are concealed by natural cloud heterogeneity and satellite retrieval errors, *Nature Communications*, 13, 7357, <https://doi.org/10.1038/s41467-022-34948-5>, 2022.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Reviews of Geophysics*, 58, e2019RG000660, <https://doi.org/https://doi.org/10.1029/2019RG000660>, e2019RG000660 10.1029/2019RG000660, 2020.
- Blyth, A. M., Lowenstein, J. H., Huang, Y., Cui, Z., Davies, S., and Carslaw, K. S.: The production of warm rain in shallow maritime cumulus clouds, *Quarterly Journal of the Royal Meteorological Society*, 139, 20–31, <https://doi.org/https://doi.org/10.1002/qj.1972>, 2013.
- Bretherton, C. S., Blossey, P. N., and Uchida, J.: Cloud droplet sedimentation, entrainment efficiency, and subtropical stratocumulus albedo, *Geophysical Research Letters*, 34, <https://doi.org/https://doi.org/10.1029/2006GL027648>, 2007.
- Bréon, F.-M., Tanré, D., and Generoso, S.: Aerosol Effect on Cloud Droplet Size Monitored from Satellite, *Science*, 295, 834–838, <https://doi.org/10.1126/science.1066434>, 2002.
- Chen, Y.-C., Xue, L., Lebo, Z. J., Wang, H., Rasmussen, R. M., and Seinfeld, J. H.: A comprehensive numerical study of aerosol-cloud-precipitation interactions in marine stratocumulus, *Atmospheric Chemistry and Physics*, 11, 9749–9769, <https://doi.org/10.5194/acp-11-9749-2011>, 2011.
- Colón-Robles, M., Rauber, R. M., and Jensen, J. B.: Influence of low-level wind speed on droplet spectra near cloud base in trade wind cumulus, *Geophysical Research Letters*, 33, <https://doi.org/https://doi.org/10.1029/2006GL027487>, 2006.
- Dagan, G., Koren, I., and Altaratz, O.: Aerosol effects on the timing of warm rain processes, *Geophysical Research Letters*, 42, 4590–4598, <https://doi.org/https://doi.org/10.1002/2015GL063839>, 2015.
- Eastman, R. and Wood, R.: Factors controlling low cloud evolution over the eastern subtropical oceans: A Lagrangian perspective using the A-Train satellites, *J. Atmos. Sci.*, 73, 331–351, 2016.
- Eastman, R., Wood, R., and Bretherton, C. S.: Time Scales of Clouds and Cloud-Controlling Variables in Subtropical Stratocumulus from a Lagrangian Perspective, *Journal of the Atmospheric Sciences*, 73, 3079 – 3091, <https://doi.org/10.1175/JAS-D-16-0050.1>, 2016.
- Eastman, R., Lebsock, M., and Wood, R.: Warm Rain Rates from AMSR-E 89-GHz Brightness Temperatures Trained Using CloudSat Rain-Rate Observations, *Journal of Atmospheric and Oceanic Technology*, 36, 1033 – 1051, <https://doi.org/10.1175/JTECH-D-18-0185.1>, 2019.

- Eastman, R., McCoy, I. L., and Wood, R.: Wind, Rain, and the Closed to Open Cell Transition in Subtropical Marine Stratocumulus, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036795, <https://doi.org/https://doi.org/10.1029/2022JD036795>, e2022JD036795 2022JD036795, 2022.
- Feingold, G., Cotton, W. R., Kreidenweis, S. M., and Davis, J. T.: The Impact of Giant Cloud Condensation Nuclei on Drizzle
405 Formation in Stratocumulus: Implications for Cloud Radiative Properties, *Journal of the Atmospheric Sciences*, 56, 4100 – 4117, [https://doi.org/10.1175/1520-0469\(1999\)056<4100:TIOGCC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<4100:TIOGCC>2.0.CO;2), 1999.
- Fons, E., Runge, J., Neubauer, D., and Lohmann, U.: Stratocumulus adjustments to aerosol perturbations disentangled with a causal approach, *npj Climate and Atmospheric Science*, 6, <https://doi.org/10.1038/s41612-023-00452-w>, 2023.
- Fossum, K. N., Ovadnevaite, J., Ceburnis, D., Preißler, J., Snider, J. R., Huang, R.-J., Zuend, A., and O’Dowd, C.: Sea-spray regulates sulfate
410 cloud droplet activation over oceans, *npj Climate and Atmospheric Science*, 3, 14, <https://doi.org/10.1038/s41612-020-0116-2>, 2020.
- Ghan, S. J., Guzman, G., and Abdul-Razzak, H.: Competition between Sea Salt and Sulfate Particles as Cloud Condensation Nuclei, *Journal of the Atmospheric Sciences*, 55, 3340 – 3347, [https://doi.org/10.1175/1520-0469\(1998\)055<3340:CBSSAS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<3340:CBSSAS>2.0.CO;2), 1998.
- Glassmeier, F., Hoffmann, F., Johnson, J. S., Yamaguchi, T., Carslaw, K. S., and Feingold, G.: Aerosol-cloud-climate cooling overestimated by ship-track data, *Science*, 371, 485–489, <https://doi.org/10.1126/science.abd3980>, 2021.
- 415 Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., Boers, R., Cairns, B., Chiu, J. C., Christensen, M., Deneke, H., Diamond, M., Feingold, G., Fridlind, A., Hünerbein, A., Knist, C., Kollias, P., Marshak, A., McCoy, D., Merk, D., Painemal, D., Rausch, J., Rosenfeld, D., Russchenberg, H., Seifert, P., Sinclair, K., Stier, P., van Diedenhoven, B., Wendisch, M., Werner, F., Wood, R., Zhang, Z., and Quaas, J.: Remote Sensing of Droplet Number Concentration in Warm Clouds: A Review of the Current State of Knowledge and Perspectives, *Reviews of Geophysics*, 56, 409–453, <https://doi.org/https://doi.org/10.1029/2017RG000593>, 2018.
- 420 Gryspeerdt, E., Goren, T., and Smith, T. W. P.: Observing the timescales of aerosol–cloud interactions in snapshot satellite images, *Atmospheric Chemistry and Physics*, 21, 6093–6109, <https://doi.org/10.5194/acp-21-6093-2021>, 2021.
- Gryspeerdt, E., Glassmeier, F., Feingold, G., Hoffmann, F., and Murray-Watson, R. J.: Observing short-timescale cloud development to constrain aerosol–cloud interactions, *Atmospheric Chemistry and Physics*, 22, 11727–11738, <https://doi.org/10.5194/acp-22-11727-2022>, 2022a.
- 425 Gryspeerdt, E., McCoy, D. T., Crosbie, E., Moore, R. H., Nott, G. J., Painemal, D., Small-Griswold, J., Sorooshian, A., and Ziemba, L.: The impact of sampling strategy on the cloud droplet number concentration estimated from satellite data, *Atmospheric Measurement Techniques*, 15, 3875–3892, <https://doi.org/10.5194/amt-15-3875-2022>, 2022b.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Sabater, M., J., N., J., P., C., R., R., I., S., D., S., A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on pressure levels from 1940 to present., Copernicus Climate Change Service (C3S) Climate Data
430 Store (CDS), <https://doi.org/10.24381/cds.bd0915c6>, 2023.
- Hoffmann, F. and Feingold, G.: Cloud Microphysical Implications for Marine Cloud Brightening: The Importance of the Seeded Particle Size Distribution, *Journal of the Atmospheric Sciences*, 78, 3247 – 3262, <https://doi.org/10.1175/JAS-D-21-0077.1>, 2021.
- Hoffmann, F., Glassmeier, F., Yamaguchi, T., and Feingold, G.: Liquid Water Path Steady States in Stratocumulus: Insights from Process-Level Emulation and Mixed-Layer Theory, *Journal of the Atmospheric Sciences*, 77, 2203 – 2215, <https://doi.org/10.1175/JAS-D-19-0241.1>, 2020.
- 435 Huebert, B. J., Blomquist, B. W., Hare, J. E., Fairall, C. W., Johnson, J. E., and Bates, T. S.: Measurement of the sea-air DMS flux and transfer velocity using eddy correlation, *Geophysical Research Letters*, 31, <https://doi.org/https://doi.org/10.1029/2004GL021567>, 2004.

- Jung, E., Albrecht, B. A., Jonsson, H. H., Chen, Y.-C., Seinfeld, J. H., Sorooshian, A., Metcalf, A. R., Song, S., Fang, M., and Russell, L. M.: Precipitation effects of giant cloud condensation nuclei artificially introduced into stratocumulus clouds, *Atmospheric Chemistry and Physics*, 15, 5645–5658, <https://doi.org/10.5194/acp-15-5645-2015>, 2015.
- Kaufman, Y. J., Koren, I., Remer, L. A., Rosenfeld, D., and Rudich, Y.: The effect of smoke, dust, and pollution aerosol on shallow cloud development over the Atlantic Ocean, *Proceedings of the National Academy of Sciences*, 102, 11 207–11 212, <https://doi.org/10.1073/pnas.0505191102>, 2005.
- Kazil, J., Feingold, G., and Yamaguchi, T.: Wind speed response of marine non-precipitating stratocumulus clouds over a diurnal cycle in cloud-system resolving simulations, *Atmospheric Chemistry and Physics*, 16, 5811–5839, <https://doi.org/10.5194/acp-16-5811-2016>, 2016.
- Lasher-Trapp, S. G., Knight, C. A., and Straka, J. M.: Early Radar Echoes from Ultragiant Aerosol in a Cumulus Congestus: Modeling and Observations, *Journal of the Atmospheric Sciences*, 58, 3545 – 3562, [https://doi.org/10.1175/1520-0469\(2001\)058<3545:EREFUA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<3545:EREFUA>2.0.CO;2), 2001.
- L'Ecuyer, T. S., Berg, W., Haynes, J., Lebsock, M., and Takemura, T.: Global observations of aerosol impacts on precipitation occurrence in warm maritime clouds, *Journal of Geophysical Research: Atmospheres*, 114, <https://doi.org/https://doi.org/10.1029/2008JD011273>, 2009.
- Lehahn, Y., Koren, I., Boss, E., Ben-Ami, Y., and Altaratz, O.: Estimating the maritime component of aerosol optical depth and its dependency on surface wind speed using satellite data, *Atmospheric Chemistry and Physics*, 10, 6711–6720, <https://doi.org/10.5194/acp-10-6711-2010>, 2010.
- Lehahn, Y., Koren, I., Altaratz, O., and Kostinski, A. B.: Effect of coarse marine aerosols on stratocumulus clouds, *Geophysical Research Letters*, 38, <https://doi.org/https://doi.org/10.1029/2011GL048504>, 2011.
- Lewis, E. and Schwartz, S.: Sea Salt Aerosol Production: Mechanisms, Methods, Measurements and Models—A Critical Review, Washington DC American Geophysical Union Geophysical Monograph Series, 152, 3719–, <https://doi.org/10.1029/GM152>, 2004.
- Liu, F., Mao, F., Rosenfeld, D., Pan, Z., Zang, L., Zhu, Y., Yin, J., and Gong, W.: Opposing comparable large effects of fine aerosols and coarse sea spray on marine warm clouds, *Communications Earth and Environment*, 3, 232, <https://doi.org/10.1038/s43247-022-00562-y>, 2022.
- Liu, F., Mao, F., Rosenfeld, D., Pan, Z., Zang, L., Zhu, Y., and Gong, W.: Marine Stratocumulus Clouds With More Coarse Sea Spray Aerosols Are Brighter, *Geophysical Research Letters*, 52, e2024GL113 718, <https://doi.org/https://doi.org/10.1029/2024GL113718>, e2024GL113718 2024GL113718, 2025.
- Ludlam, F. H.: The production of showers by the coalescence of cloud droplets, *Quarterly Journal of the Royal Meteorological Society*, 77, 402–417, <https://doi.org/https://doi.org/10.1002/qj.49707733306>, 1951.
- Meskhidze, N., Remer, L. A., Platnick, S., Negrón Juárez, R., Lichtenberger, A. M., and Aiyyer, A. R.: Exploring the differences in cloud properties observed by the Terra and Aqua MODIS Sensors, *Atmospheric Chemistry and Physics*, 9, 3461–3475, <https://doi.org/10.5194/acp-9-3461-2009>, 2009.
- Mülmenstädt, J., Ackerman, A. S., Fridlind, A. M., Huang, M., Ma, P.-L., Mahfouz, N., Bauer, S. E., Burrows, S. M., Christensen, M. W., Dipu, S., Gettelman, A., Leung, L. R., Tornow, F., Quaas, J., Varble, A. C., Wang, H., Zhang, K., and Zheng, Y.: Can general circulation models (GCMs) represent cloud liquid water path adjustments to aerosol–cloud interactions?, *Atmospheric Chemistry and Physics*, 24, 13 633–13 652, <https://doi.org/10.5194/acp-24-13633-2024>, 2024a.
- Mülmenstädt, J., Gryspeerdt, E., Dipu, S., Quaas, J., Ackerman, A. S., Fridlind, A. M., Tornow, F., Bauer, S. E., Gettelman, A., Ming, Y., Zheng, Y., Ma, P.-L., Wang, H., Zhang, K., Christensen, M. W., Varble, A. C., Leung, L. R., Liu, X., Neubauer, D., Partridge, D. G., Stier,

- P., and Takemura, T.: General circulation models simulate negative liquid water path–droplet number correlations, but anthropogenic aerosols still increase simulated liquid water path, *Atmospheric Chemistry and Physics*, 24, 7331–7345, <https://doi.org/10.5194/acp-24-7331-2024>, 2024b.
- Platnick, S., Meyer, K. G., King, M. D., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Holz, R. E.,
 480 Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS Cloud Optical and Microphysical Products: Collection 6 Updates and Examples From Terra and Aqua, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.
- Quaas, J., Boucher, O., and Lohmann, U.: Constraining the total aerosol indirect effect in the LMDZ and ECHAM4 GCMs using MODIS satellite data, *Atmospheric Chemistry and Physics*, 6, 947–955, <https://doi.org/10.5194/acp-6-947-2006>, 2006.
- Randall, D. A., Coakley, J. A., Fairall, C. W., Kropfli, R. A., and Lenschow, D. H.: Outlook for Research on Subtropical Marine Stratiform Clouds, *Bulletin of the American Meteorological Society*, 65, 1290 – 1301, [https://doi.org/10.1175/1520-0477\(1984\)065<1290:OFROSM>2.0.CO;2](https://doi.org/10.1175/1520-0477(1984)065<1290:OFROSM>2.0.CO;2), 1984.
- 485 Rosenfeld, D., Lahav, R., Khain, A., and Pinsky, M.: The Role of Sea Spray in Cleansing Air Pollution over Ocean via Cloud Processes, *Science*, 297, 1667–1670, <https://doi.org/10.1126/science.1073869>, 2002.
- Schulz, B. and Mellado, J. P.: Wind Shear Effects on Radiatively and Evaporatively Driven Stratocumulus Tops, *Journal of the Atmospheric Sciences*, 75, 3245 – 3263, <https://doi.org/10.1175/JAS-D-18-0027.1>, 2018.
- 490 Slingo, A.: Sensitivity of the Earth’s radiation budget to changes in low clouds, , 343, 49–51, <https://doi.org/10.1038/343049a0>, 1990.
- Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., eds.: Climate Change 2013: The Physical Science Basis, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324>, contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental
 495 Panel on Climate Change, 2013.
- Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment*, 8, 1251–1256, [https://doi.org/https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of Atmospheric Sciences*, 34, 1149 – 1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOP>2.0.CO;2), 1977.
- 500 Wang, S., Golaz, J.-C., and Wang, Q.: Effect of intense wind shear across the inversion on stratocumulus clouds, *Geophysical Research Letters*, 35, <https://doi.org/https://doi.org/10.1029/2008GL033865>, 2008.
- Wang, S., Zheng, X., and Jiang, Q.: Strongly sheared stratocumulus convection: an observationally based large-eddy simulation study, *Atmospheric Chemistry and Physics*, 12, 5223–5235, <https://doi.org/10.5194/acp-12-5223-2012>, 2012.
- Wood, R., Leon, D., Lebsock, M., Snider, J., and Clarke, A. D.: Precipitation driving of droplet concentration variability in marine low
 505 clouds, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/https://doi.org/10.1029/2012JD018305>, 2012.
- Yin, Y., Levin, Z., Reisin, T. G., and Tzivion, S.: The effects of giant cloud condensation nuclei on the development of precipitation in convective clouds — a numerical study, *Atmospheric Research*, 53, 91–116, [https://doi.org/https://doi.org/10.1016/S0169-8095\(99\)00046-0](https://doi.org/https://doi.org/10.1016/S0169-8095(99)00046-0), 2000.
- Zamora Zapata, M., Heus, T., and Kleissl, J.: Effects of Surface and Top Wind Shear on the Spatial Organization
 510 of Marine Stratocumulus-Topped Boundary Layers, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034162, <https://doi.org/https://doi.org/10.1029/2020JD034162>, e2020JD034162 2020JD034162, 2021.

Zhang, J., Chen, Y.-S., Yamaguchi, T., and Feingold, G.: Cloud water adjustments to aerosol perturbations are buffered by solar heating in non-precipitating marine stratocumuli, *Atmospheric Chemistry and Physics*, 24, 10 425–10 440, <https://doi.org/10.5194/acp-24-10425-2024>, 2024, 2024.