

# 1 Recent intensification of extreme precipitation over Antarctica 2 driven by increases in greenhouse gases and stratospheric ozone

3  
4 Sai Prabala Swetha Chittella<sup>1,2</sup>, Andrew Orr<sup>2</sup>, Pranab Deb<sup>1</sup>, and Quentin Dalaiden<sup>3</sup>

5 <sup>1</sup>Centre for Ocean, River, Atmosphere and Land Sciences (CORAL), Indian Institute of Technology Khargapur, Khargapur,  
6 721302, India

7 <sup>2</sup>British Antarctic Survey, National Environmental Research Council, Cambridge, UK

8 <sup>3</sup>Nansen Environmental and Remote Sensing Center and Bjerknes Center for Climate Research, Bergen, Norway

9 *Correspondence to:* Sai Prabala Swetha Chittella (prabala.saiswetha@gmail.com)

10 **Abstract.** Extreme precipitation is a major contributor to the total precipitation over Antarctica, as well as its variability.  
11 However, it's still poorly understood whether any recent trends in extreme precipitation over Antarctica have occurred, and if  
12 so, whether they are anthropogenically driven. Here we address this knowledge gap by using ERA5 data from 1979 to 2023  
13 to identify six Antarctic drainage basins with significant positive trends in total and extreme precipitation. These basins  
14 include one in the Antarctic Peninsula, one in West Antarctica, and four in East Antarctica. We subsequently perform a  
15 detection and attribution analysis of these trends using precipitation outputs from CESM1 global climate model large  
16 ensembles based on “all-forcing” (ALL) simulations and “single-forcing” simulations that isolate the effects of greenhouse  
17 gases (GHG), anthropogenic aerosols (AER), and stratospheric ozone (O3). Our analysis indicates that a combination of both  
18 anthropogenic and natural forcings are responsible for these trends in five of the basins located in West and East Antarctica,  
19 specifically within the Filchner-Ronne basin, Dronning Maud Land, and Enderby Land. The results further show that the  
20 precipitation changes are robustly attributed to increases in greenhouse gases in four basins and stratospheric ozone depletion  
21 in three basins, and increases in anthropogenic aerosols in only one basin. Repeating the analysis using CESM2 ensembles  
22 for ALL, GHG, and AER also indicated that greenhouse gases are an important driver of the precipitation trends over these  
23 basins. However, these results suggest that increases in anthropogenic aerosol emissions can oppose / offset the increases in  
24 precipitation from increased greenhouse gases over our five basins, which contradicts the findings based on CESM1. Finally,  
25 we show that the precipitation trends are partly due to an increased occurrence of atmospheric rivers.

## 26 1 Introduction

27 Extreme precipitation events (EPEs, i.e., where the amount of precipitation at a location is substantially greater than normal)  
28 have increased in intensity and frequency globally in recent decades (Donat et al., 2013; Sun et al., 2021; Dunn et al., 2020;

Li et al., 2024), which has been attributed to anthropogenically-induced climate change (Min et al., 2011; Zhang et al., 2013; Paik et al., 2020; Kirchmeier-Young et al., 2020; Dong et al., 2020, 2021; Estrada et al., 2023; IPCC 2023). In Antarctica, EPEs account for over 40% of the continent's total precipitation and almost 70% of the variance in annual precipitation, and are often associated with intense atmosphere rivers (ARs) (Gorodetskaya et al., 2014; Turner et al., 2019; Adusumilli et al., 2021; Wille et al., 2021, 2024, 2025; Gehring et al., 2022; MacLennan et al., 2022, 2023; Simon et al., 2024). However, identifying positive trends in EPEs over recent decades in Antarctica is hampered by such events being rare and difficult to separate from the substantial interannual precipitation variability that characterises its climate, as well as limited observations (Connolley, 1997; Hosking et al., 2013; Yu et al., 2018, 2025). For example, Intergovernmental Panel on Climate Change (IPCC) assessments of observed extreme precipitation changes largely exclude Antarctica (Seneviratne et al., 2021). Nevertheless, positive trends in the number of EPEs occurring over recent decades have been identified over parts of East Antarctica (Yu et al., 2018; Simon et al., 2024).

Antarctic precipitation and EPEs are the major component of surface mass balance (Turner et al., 2019; Kittel et al., 2021; Mottram et al., 2021; Clem and Raphael, 2024; Davison et al., 2023; Wille et al., 2024) and thus play a crucial role in controlling the stability of the Antarctic ice sheet by offsetting dynamically driven ice-losses (Favier et al., 2017; Zwally et al., 2017; Paolo et al., 2018; Rignot et al., 2019; Kromer and Trusel, 2023; Otosaka et al., 2023; Wang et al., 2025). Furthermore, mass gained from precipitation and EPEs over Antarctic ice shelves can partially compensate for mass-losses caused by either basal or surface melting (Paolo et al., 2015; Pattyn et al., 2017; Gardner et al., 2018; Rignot et al., 2019; Nakayama et al., 2021; Johnson et al., 2022). This is especially critical as ice shelf thinning results in the acceleration of grounded ice toward the ocean, where the ice subsequently calves into the ocean as icebergs and causes increased sea-level rise (Rott et al. 1996; Pritchard et al. 2012; Rignot et al. 2019). Additionally, snowfall is an essential factor in minimizing the susceptibility of ice shelves to surface melt pond formation, which can initiate hydrofracturing of ice shelves, by counteracting any firm air depletion caused by surface melting (Scambos et al. 2000; Munneke et al., 2014; Banwell and MacAyeal 2015; Lai et al. 2020; Orr et al., 2023; Van Wessem et al., 2023).

Therefore, a better physical understanding of the trends in EPEs during the past decades over Antarctica is critical. Moreover, an enhanced understanding of the drivers of these trends, such as anthropogenically-induced changes (i.e., external climate forcing), is also of critical importance to better highlight how EPEs and associated impacts might change under future climate projections (Kittel et al., 2021; Vignon et al., 2021). For example, Dalaiden et al. (2022) identified that increased greenhouse gas emissions and stratospheric ozone depletion were the primary drivers of increased precipitation and temperature over West Antarctica since the middle of the twentieth century. These two drivers were also identified as causing recent changes in Southern Ocean temperature and salinity by Swart et al. (2018) and Hobbs et al. (2021). Additionally, increased anthropogenic aerosol emissions in Asia in recent decades have been suggested by Gu et al. (2025) to affect Antarctica's climate by Rossby-wave teleconnections. However, the strong internal variability prevailing in the Southern

61 Hemisphere high latitudes makes it challenging to detect the signal of anthropogenically-produced climatic changes from the  
62 observed changes during the past decades (e.g., Fyke et al., 2017), as well as when this signal will emerge from natural  
63 variability in the future (Trusel et al., 2015; Previdi and Polvani, 2016; Morioka et al., 2024).

64 This study addresses the knowledge gap around our understanding of the drivers of changes in EPEs over Antarctica by  
65 performing a detection and attribution (D&A) analysis to investigate the influence of external forcings and natural variability  
66 on annual trends in total and extreme precipitation across Antarctica from 1979 to the present-day. Our approach uses  
67 precipitation from: i) ERA5 atmospheric reanalysis to identify trends, and ii) large ensembles performed using the  
68 Community Earth System Model Versions 1 and 2 (CESM1 and CESM2), which are fully-coupled global climate models, to  
69 undertake D&A of the ERA5-based trends and determine the role of different external climate forcings versus internal  
70 climate variability.

## 71 **2 Data and Methods**

### 72 **2.1 Datasets**

73 In situ measurements of precipitation over Antarctica are extremely scarce and difficult to make due to the challenging  
74 remote environment and difficulties distinguishing between falling and wind-driven snow (Pritchard, 2021). Therefore, we  
75 use hourly precipitation (both rainfall and snowfall) data from the ERA5 reanalysis (Hersbach et al., 2020) at a horizontal  
76 resolution of  $0.25^\circ$  to identify present-day precipitation trends in Antarctica as this provides complete spatial and temporal  
77 coverage of this region for the period of interest. ERA5 precipitation has been used previously in studies examining daily  
78 precipitation over Antarctica (Vignon et al., 2021; Carter et al., 2022; Tewari et al., 2022) and for extreme event analysis (Yu  
79 et al., 2025). However, for this study we only use data from 1979 to 2023 (45 years), as prior to this period there were  
80 relatively few satellite observations over the Southern Ocean and Antarctica, which are necessary to constrain the reanalysis  
81 (Bromwich et al., 2024). Additionally, as ARs are an important contributor to EPEs (Gorodetskaya et al., 2014; MacLennan et  
82 al., 2022; Wille et al., 2025), we identify their occurrence using a 3-hourly AR detection catalog at a horizontal resolution of  
83  $0.5^\circ$ , derived from ERA5 reanalysis (Wille et al., 2021). Here, ARs are identified based on the meridional component of  
84 integrated vapor transport exceeding the 98<sup>th</sup> percentile of the monthly climatologies.

85 We also use daily precipitation (both rainfall and snowfall) output for the period 1979 to 2023 from the large ensemble  
86 simulations performed with the CESM1 and CESM2 models. These simulations were selected because they have a number  
87 of factors that are crucial for the D&A analysis, including: i) “all-forcing” (ALL) large ensembles that consider forcings from  
88 all anthropogenic and natural (such as solar variability and volcanic eruptions) sources, ii) “single-forcing” large ensembles  
89 that isolate the impact of specific external forcings resulting from anthropogenic activities, such as greenhouse gas (GHG)  
90 emissions, aerosol (AER) emissions, and stratospheric ozone depletion (O3) and iii) a relatively good representation of the

91 surface climate of Antarctica (Dunmire et al., 2022; Dalaiden et al., 2021, 2022; England et al., 2016; Landrum et al., 2017;  
 92 Lenaerts et al., 2016). Note that the atmosphere component of CESM1 is stratospheric-resolving and includes interactive  
 93 ozone chemistry, while CESM2 has a model top near 40 km and uses prescribed stratospheric ozone depletion (Kay et al.,  
 94 2015; Danabasoglu et al., 2020; Deser et al., 2020; Simpson et al., 2023).

95 CESM1 includes a 40-member ALL ensemble that spans the period from 1920 to 2100, based on CMIP5 (Coupled Model  
 96 Intercomparison Project Phase 5) historical radiative forcing from 1920 to 2005 and the RCP8.5 (Representative  
 97 Concentration Pathway 8.5, representing additional radiative forcing of 8.5 W m<sup>-2</sup> by 2100) scenario from 2006 to 2100 (Kay  
 98 et al., 2015). Its single-forcing large ensembles employ an “all-but-one” design (Deser et al., 2020): all external forcings  
 99 evolve in time as in the ALL large ensemble, except the forcing of interest, which is held fixed throughout the simulation. It  
 100 includes 20-member all-but-one forcing ensembles from 1920 to 2080 with greenhouse gases (xGHG) and anthropogenic  
 101 aerosols (xAER) fixed at 1920 levels. In the xAER ensemble, the aerosols included are from industrial, agricultural,  
 102 household, and transportation-related sources but exclude any anthropogenic biomass burning sources. Additionally, CESM1  
 103 also includes an 8-member all-but-one forcing ensemble from 1955 to 2005 with stratospheric ozone (xO3) fixed at 1955  
 104 levels (Landrum et al., 2017). Following Deser et al. (2020), the individual members of the single-forcing ensembles (GHG,  
 105 AER, O3) are calculated from the all-but-one forcing ensembles (xGHG, xAER, xO3) as follows:

$$106 \quad \text{GHG}_i = \text{xGHG}_i - 2\text{xGHG}_{\text{em}} + \text{ALL}_{\text{em}}, (1)$$

$$107 \quad \text{AER}_i = \text{xAER}_i - 2\text{xAER}_{\text{em}} + \text{ALL}_{\text{em}}, (2)$$

$$108 \quad \text{O3}_i = \text{xO3}_i - 2\text{xO3}_{\text{em}} + \text{ALL}_{\text{em}}, (3)$$

109 where the subscript “i” represents an individual ensemble member and “em” the ensemble mean. To assess internal  
 110 variability we use the CESM1 pre-industrial 2200-year long control simulation (with all external forcings fixed at 1850  
 111 values), but exclude the first 400 years of the simulation to avoid any artificial related to the initialization.

112 CESM2 includes a 50-member ALL ensemble that spans the period from 1850 to 2100 based on the CMIP6 (Coupled Model  
 113 Intercomparison Project Phase 6) historical forcings from 1850 to 2014 and the SSP370 (Shared Socioeconomic Pathways,  
 114 representing additional radiative forcing of 7 W m<sup>-2</sup> by 2100) scenario from 2015 to 2100. Its single-forcing large ensembles  
 115 employ an “only” design (Simpson et al., 2023): all external forcings are held fixed at 1850 values, except the forcing of  
 116 interest, which evolve in time throughout the simulation. It includes a 15-member GHG ensemble and a 20-member AER  
 117 ensemble from 1850 to 2050. In the AER ensemble, anthropogenic biomass burning sources are again excluded. To assess  
 118 internal variability we use the CESM2 pre-industrial 2000-year long control simulation (with external forcings fixed at 1850  
 119 values). Note that there are no CESM2 single-forcing simulations for stratospheric ozone depletion available (Simpson et al.,  
 120 2023).

## 121 2.2 Methods

122 Hourly ERA5 precipitation outputs are firstly re-gridded onto the CESM1 grid (resolution of 1°) using an area-weighted  
123 averaging method, and then summed over 24-hour intervals to produce daily values. We then compute trends in total and  
124 extreme precipitation for ERA5 and the CESM1 ensembles ALL, GHG, and AER for the period 1979 to 2023 and O3 for the  
125 period 1979 to 2005. Here, precipitation days are defined as days with precipitation exceeding a threshold of 0.02 mm day<sup>-1</sup>,  
126 and extreme precipitation as days with precipitation exceeding the 95<sup>th</sup> percentile of all precipitation days over the period  
127 1979 to 2023, following Swetha Chittella et al. (2022). These daily data are then annually aggregated to obtain an annual  
128 time-series of both total and extreme precipitation for ERA5 and each member of the ALL, GHG, and AER ensembles from  
129 1979 to 2023 and the O3 ensemble from 1979 to 2005. The ERA5 annual time-series are subsequently used to identify  
130 precipitation trends over Antarctica by calculating statistically significant (90% confidence interval): i) spatial distributions  
131 of linear trends of total and extreme precipitation , and ii) basin-averaged linear trends of total precipitation and extreme  
132 precipitation for Antarctica's 18 drainage basins (mapped using the Ice Sheet Mass Balance Inter-comparison (IMBIE)  
133 dataset shown in Fig. 1; Rignot et al., 2019).

134 Next we perform an initial comparison between the precipitation trends from ERA5 and the CESM1 ALL ensemble for the  
135 period 1979 to 2023. Here, if the basin-averaged linear trends in the total and extreme precipitation time-series from the ALL  
136 ensemble are significant at the 90% level and consistent with the trends estimated by ERA5, then the CESM1 ALL ensemble  
137 was judged to reproduce the ERA5-based trends for that basin. For the basins identified during this step, we subsequently  
138 conduct a preliminary attribution analysis by assessing the contribution of the trends in the total and extreme precipitation  
139 time-series from the GHG, AER and O3 ensembles to the trends in the ALL ensemble. This was primarily achieved by  
140 comparing probability density distributions for the precipitation trends for each of the basins, calculated using a kernel  
141 density estimation method (significant at the 90% confidence level).

142 We then apply a regression-based D&A technique for the selected drainage basins, based on a least-squares fitting statistical  
143 approach (Allen and Stott, 2003; Ribes et al., 2013; Dong et al., 2020; Hobbs et al., 2021; Dalaiden et al., 2022), to formally  
144 quantify to what extent the total and extreme precipitation trends from ERA5 is related to external forcing. This method  
145 calculates the scaling factors between observations (in this study, ERA5) and the estimated forced response from climate  
146 model ensembles (including all external forcing or only a subset), which is expressed as follows (Ribes et al., 2013; Dalaiden  
147 et al., 2022):

$$148 \quad y^* = \sum_{i=1}^n \beta_i x_i^*, (4)$$

149 where  $y^*$  is the “true” climate response to all external forcings,  $x_i^*$  the simulated climate response to the  $i^{\text{th}}$  specific external  
 150 forcing considered (up to a total of  $n$ ), and  $\beta_i$  the scaling factor (regression coefficient) for the  $i^{\text{th}}$  external forcing. Here the  
 151 actual observations  $y = y^* + \epsilon_0$ , with  $\epsilon_0$  corresponding to the noise term on  $y$ , i.e., internal climate variability. And  
 152  $x_i = x_i^* + \epsilon_i$  represents the ensemble mean of the simulated response to the  $i^{\text{th}}$  external forcing, with  $\epsilon_i$  the noise term on  
 153 forcing  $x_i$ , i.e., internal variability. This method also provides confidence intervals associated with the scaling factors. If the  
 154 scaling factor, including the confidence interval, is greater than zero, then the observed changes are outside the range of  
 155 internal variability. We therefore conclude that the observed changes can only be explained by external forcing (Dong et al.,  
 156 2020; Dalaiden et al., 2022). In the case the scaling factor, including the confidence interval, is greater (lower) than one, this  
 157 implies that the model significantly underestimates (overestimates) the observed trends. Finally, if the lower bound of the  
 158 confidence interval is smaller than or equal to zero, we cannot conclude a statistically impact of external forcing on the  
 159 observed changes.


160 Note that to improve the signal-to-noise ratio, D&A studies commonly apply a pre-whitening step to both observations and  
 161 model simulated responses to suppress the contribution from internal climate variability (e.g., Allen and Stott, 2003). For the  
 162 regression-based D&A technique, the ERA5 time series and the model responses are therefore pre-whitened using the  
 163 covariance matrix of internal variability (also referred to as the noise covariance matrix). We estimate this matrix using the  
 164 regularized optimal fingerprinting approach of Ribes et al. (2013). Additionally, as the ensemble mean smoothed out the  
 165 impact of internal variability, we scaled up the variance of the ensemble mean response by using the square root of ensemble  
 166 size (Allen and Stott, 2003; Ribes et al., 2013; Kirchmeier-Young et al., 2017).

167 Using the regression-based D&A technique, detection is performed by applying a one-signal analysis for CESM1, where the  
 168 basin-averaged total and extreme precipitation time-series from the ALL ensemble (along with internal variability) are  
 169 regressed onto the ERA5 time-series, resulting in scaling factors (and confidence intervals) for ALL (Dalaiden et al., 2022).  
 170 Following this, attribution is performed in CESM1 by applying a two-signal analysis, where the basin-averaged total and  
 171 extreme precipitation time-series from the ALL and all-but-one forcing ensembles are regressed onto the ERA5 time-series,  
 172 resulting in the scaling factors for a specific external forcing and for all other forcing except that specific forcing (Dalaiden et  
 173 al., 2022). In other words, regressing: i) ALL and xGHG against ERA5, results in the scaling factors for GHG and xGHG, ii)  
 174 ALL and xAER against ERA5, results in scaling factors for AER and xAER, and iii) ALL and xO3 against ERA5, results in  
 175 scaling factors for O3 and xO3 (Dalaiden et al., 2022). However, here we only focus on the scaling factors (and confidence  
 176 intervals) associated with GHG, AER, and O3. Both the one-signal and two-signal analysis involving the GHG and AER  
 177 ensembles were performed with the same number of members, which was therefore limited to 20, i.e., the number from the  
 178 GHG and AER ensembles, as this was less than the ALL (40) ensembles. However, the two-signal analysis involving the O3

ensemble was limited to 8 members. Note that the one-signal analysis results in CESM1 using all 40 members from the ALL ensemble were largely identical to those using 20 members (not shown). For CESM1, the noise covariance matrix is derived from pre-industrial control simulations. More details on the method are provided in Ribes et al. (2013) and Kirchmeier-Young et al. (2017).

The above analysis is then repeated using the CESM2 ensembles ALL, GHG, and AER to assess to what extent the ERA5 trends can be detected and attributed to specific external forcings using this model. For the regression-based D&A technique, we again apply a one-signal analysis to regress the ALL ensemble onto the ERA5 time-series. However, since the single-forcing ensembles for CESM2 were generated differently than for CESM1, attribution is also performed by applying a three-signal analysis, where the basin-averaged total and extreme precipitation time-series from the ALL, GHG and AER ensembles (along with internal variability) are regressed onto the ERA5 time-series (Dalaiden et al., 2022). This results in scaling factors (and confidence intervals) for GHG and AER, and also all forcings other than greenhouse gases and anthropogenic aerosols, which is labelled OTHERS (Gillett et al., 2021; Dalaiden et al., 2022). Similarly to CESM1, the noise covariance matrix is derived from the pre-industrial control simulation. Both one-signal and two-signal analysis were performed with a common number of members from each ensemble, which was therefore limited to 15, i.e., the number from the GHG ensemble, as this was less than both the ALL (50) and AER (20) ensembles. However, the one-signal analysis results in CESM2, using all 50 members from the ALL ensemble, was largely identical to using 15 members (not shown). Investigating results from two climate models is important as they enable an assessment of the robustness of the results. However, the investigation will predominately focus on analysis of the CESM1 results, as these include an O3 ensemble.

Finally, we examine trends in AR-associated total and extreme precipitation from 1980 to 2023 by combining the AR detection catalog with ERA5 precipitation. Here, daily values of AR-associated total and extreme precipitation are calculated by estimating the amount of ERA5 total and extreme precipitation occurring within each AR footprint. These daily data are then annually aggregated to obtain an annual time-series of both AR-associated total and extreme precipitation. Following the approach of MacLennan et al. (2022), this was subsequently used to investigate: i) the spatial distribution of statistically significant (90% confidence interval) trends in AR-associated total and extreme precipitation from 1980 to 2023, and ii) the relative contribution of AR-associated total and extreme precipitation trends to the ERA5-based total and extreme precipitation trends from 1980 to 2023 (which required regridding the total and extreme precipitation from ERA5 onto the AR detection catalog grid). Note that spuriously large percentages for the relative contributions of AR-associated total and extreme precipitation trends are produced if the ERA5-based total and extreme precipitation trends are close to zero, which is avoided by only calculating this for grid cells where the ERA5-based trends exceed a threshold of 0.05 mm year<sup>-1</sup>.



208

209 Figure 1: Map of Antarctica showing key regions of interest (names in blue) and the 18 major drainage basins (names in red), as  
 210 defined in the IMBIE dataset (Rignot et al., 2019).


## 211 3 Results

### 212 3.1 Identification of precipitation trends in ERA5

213 Figure 2 (a, e) shows annual trends in total and extreme precipitation from ERA5 from 1979 to 2023. This shows that broad  
 214 positive trends for both total and extreme precipitation (up to 2 mm year<sup>-1</sup>) are apparent from Ellsworth Land in West  
 215 Antarctica to the Amery ice shelf in East Antarctica, while negative trends (up to -2 mm year<sup>-1</sup>) are evident from Marie Byrd  
 216 Land in West Antarctica to Wilkes Land in East Antarctica (see Fig. 1 for locations). However, for some of these regions, the  
 217 trends are statistically insignificant, with only Ellsworth Land, the Filchner-Ronne basin, Dronning Maud Land, Enderby  
 218 Land, and the Amery Ice Shelf showing relatively large areas with significant positive trends. By contrast, the regions  
 219 showing significant negative trends are patchy and confined to Marie Byrd Land, Victoria Land, and Wilkes Land. Over the  
 220 Antarctic Peninsula, the ERA5 trends in total and extreme precipitation differ in direction, with total precipitation increasing  
 221 and extreme precipitation decreasing, although only the trends over the northern Antarctic Peninsula are significant.

222





223

224 **Figure 2: Annual trends in (a-e) total and (f-j) extreme precipitation over Antarctica from 1979 to 2023 ( $\text{mm year}^{-1}$ ) for (a, f)**  
 225 **ERA5 and the CESM1 ensemble mean for (b, g) ALL, (c, h) GHG, (d, i) AER and (e, j) O3. Stippling indicates regions where**  
 226 **trends are statistically significant at the 90% confidence level. Also shown in panel (a) are the locations of the I-Ipp drainage basin**  
 227 **in the Antarctic Peninsula, the J-Jpp basin in West Antarctica and the Jpp-K basin in East Antarctica (both part of**  
 228 **Filchner-Ronne basin), and the K-A, A-Ap (both part of Dronning Maud Land), and Ap-B (Enderby Land) drainage basins in**  
 229 **East Antarctica (outlined in black).**


230 Within the regions showing positive ERA5-based trends, we identified six drainage basins that showed significant  
 231 basin-averaged positive trends in both total and extreme precipitation (Figs. 2 and 3). These are basin I-Ipp in the Antarctic  
 232 Peninsula, basins J-Jpp in West Antarctica and Jpp-K in East Antarctica (both part of the Filchner-Ronne basin), basins K-A  
 233 and A-Ap in East Antarctica (both part of Dronning Maud Land), and basin Ap-B in East Antarctica (Enderby Land) (see  
 234 Figs. 1 and 2 for locations). The basins J-Jpp, A-Ap, and Ap-B all have similar basin-averaged ERA5 trends of about 1 mm  
 235  $\text{year}^{-1}$  for total precipitation and 0.5  $\text{mm year}^{-1}$  for extreme precipitation, whereas basins Jpp-K, K-A, and I-Ipp have trends of  
 236 about 0.1, 1.5, and 5  $\text{mm year}^{-1}$  for total precipitation respectively, and 0.1, 1, and 2.5  $\text{mm year}^{-1}$  for extreme precipitation  
 237 respectively (Fig. 3). Note that within the regions showing negative ERA5-based trends, none of the drainage basins showed  
 238 significant negative basin-averaged trends.

### 239 3.2 Detection of precipitation trends in CESM1

240 Figure 2 (b, g) demonstrates that the CESM1 ensemble mean for ALL also shows significant positive trends for both total  
 241 and extreme precipitation (up to 2  $\text{mm year}^{-1}$ ) from Ellsworth Land to the Amery ice shelf, which agree with ERA5. For

example, it largely captures the spatial pattern and magnitude of the significant positive trends from ERA5 for total and extreme precipitation over Ellsworth Land, the Filchner-Ronne basin, Dronning Maud Land, Enderby Land, and the Amery ice shelf. Additionally, the CESM1 ensemble mean for ALL also shows significant positive trends for total precipitation over the Antarctic Peninsula, which agree with the significant positive trends from ERA5. By contrast, the ensemble mean for ALL shows positive trends in total and extreme precipitation over Marie Byrd Land, Wilkes Land, and Victoria Land, and positive trends for extreme precipitation over the Antarctic Peninsula, which disagree with ERA5. However, some differences in trends between the ALL ensemble and ERA5 results are to be expected as the ALL results are based on an ensemble of model means, which would therefore artificially reduce the contribution from internal variability.


For the six drainage basins that are identified as having significant positive trends in both total and extreme precipitation from ERA5, Fig. 3 shows that for five of them (J-Jpp in West Antarctica, and Jpp-K, A-Ap, Ap-B, and K-A in East Antarctica) that: i) the basin-averaged trends from the CESM1 ensemble mean for ALL are close to the ERA5 trends (although this is less apparent for basin K-A), and ii) the range of the basin-averaged ALL trends from the different members includes the ERA5 trends. We therefore identify these five basins as having ALL trends that are in good agreement with ERA5 trends, and focus on these basins for the rest of our analysis. By contrast, the ERA5 trend for the remaining basin (I-Ipp in the Antarctic Peninsula) is outside the range of the ALL trends from the different members, and therefore is not considered for further analysis. Additionally, to further compare the basin-averaged trends between ERA5 and the ensemble mean for ALL, Fig. 4 shows their time-series of total and extreme precipitation from 1979 to 2023 for our five selected basins. Consistent with Fig. 3, this shows broadly similar positive trends for all five basins for both ERA5 and ALL. Additionally, for all basins the ensemble spread of ALL encompasses the ERA5-based precipitation amounts, indicating confidence in the ability of CESM1 to capture regional internal variability, as well as the trends.



262


263 Figure 3: Basin-averaged annual trends in (a) total and (b) extreme precipitation for drainage basins J-Jpp, Jpp-K (both part of  
 264 Filchner-Ronne basin), K-A, A-Ap (both part of Dronning Maud Land), and Ap-B (Enderby Land), and I-Ipp (Antarctic  
 265 Peninsula) from 1979 to 2023 ( $\text{mm year}^{-1}$ ) for ERA5 (thick red line) and the CESM1 ensemble mean for ALL (thick blue line).  
 266 Also shown are the basin-averaged trends from the different members (40) of the ALL ensemble (filled blue circles). The trends  
 267 are statistically significant with a 90% confidence level.

268 Figure 5 presents the one-signal D&A results for total and extreme precipitation across the five selected basins, based on  
 269 regression of the CESM1 ALL ensemble onto ERA5. This shows: i) scaling factors of around 0.9 to 1.1 for both total and  
 270 extreme precipitation, and ii) relatively small confidence intervals, with total precipitation showing narrower intervals than  
 271 extreme precipitation. Since the scaling factors are greater than zero and the confidence intervals are relatively small, and  
 272 because the confidence intervals include unity for basins K-A, A-Ap (both part of Dronning Maud Land) for total  
 273 precipitation and for basins K-A, A-Ap (both part of Dronning Maud Land) and Ap-B (Enderby Land) for extreme  
 274 precipitation, this indicates that the observed (i.e., ERA5-based) linear trends in total and extreme precipitation from 1979 to  
 275 2023 for these five basins is explained by external forcing (i.e., successful detection, and supporting the results from Figs. 2,  
 276 3 and 4). Additionally, since the scaling factors are slightly less than one for basins Jpp-K (part of Filchner-Ronne basin) and  
 277 Ap-B (Enderby Land) for total precipitation, and for basin J-Jpp (part of Filchner-Ronne basin) for extreme precipitation, this  
 278 suggests that for these basins that ALL tends to overestimate the associated ERA5-based trends. In contrast, the scaling  
 279 factors are slightly greater than one for basin J-Jpp (part of Filchner-Ronne basin) for both total and extreme precipitation,  
 280 suggesting that for this basin that ALL tends to underestimate the associated ERA5-based trends. Also, the slightly narrower  
 281 intervals for total precipitation indicates that the influence of internal variability is larger for extreme precipitation compared  
 282 to total precipitation.



283

284 Figure 4: Times series of basin-averaged annual (a-e) total and (f-j) extreme precipitation (mm) for drainage basins (a, f) J-Jpp,  
 285 (b, g) Jpp-K (both part of Filchner-Ronne basin), (c, h) K-A, (d, i) A-Ap (both part of Dronning Maud Land), and (e, j) Ap-B  
 286 (Enderby Land) from 1979 to 2023 from ERA5 (red) and the CESM1 ensemble mean for ALL (blue). Linear trends (statistically  
 287 significant at 90% confidence level) for ERA5 and ALL are represented by the red and blue lines, respectively. Blue shading  
 288 represents the maximum and minimum ensemble spread of ALL. Each panel also shows the linear trend and the standard error  
 289 (mm year<sup>-1</sup>).



290

291 Figure 5: One-signal estimates of basin-averaged scaling factors (solid blue circles) for annual (a) total and (b) extreme  
 292 precipitation for drainage basins J-Jpp, Jpp-K (both part of Filchner-Ronne basin), K-A, A-Ap (both part of Dronning Maud

293 Land), and Ap-B (Enderby Land) from 1979 to 2023 based on regressing the CESM1 ALL ensemble against ERA5. The error  
294 bars denote the 90% confidence intervals and the horizontal line indicates a scaling factor of one.

295 Table 1: Basin-averaged annual trends in total and extreme precipitation (mm year<sup>-1</sup>) for drainage basins J-Jpp, Jpp-K (both part  
296 of Filchner-Ronne basin), K-A, A-Ap (both part of Dronning Maud Land), and Ap-B (Enderby Land) from 1979 to 2023 (mm  
297 year<sup>-1</sup>) for the CESM1 ensemble mean for ALL, GHG, AER and O3. Trends that are statistically significant at the 90%  
298 confidence level are shown in bold.


|       | ALL                 |                       | GHG                 |                       | AER                 |                       | O3                  |                       |
|-------|---------------------|-----------------------|---------------------|-----------------------|---------------------|-----------------------|---------------------|-----------------------|
| Basin | Total precipitation | Extreme precipitation | Total precipitation | Extreme precipitation | Total precipitation | Extreme precipitation | Total precipitation | Extreme precipitation |
| J-Jpp | <b>0.60</b>         | <b>0.33</b>           | <b>0.28</b>         | <b>0.07</b>           | <b>0.36</b>         | <b>0.23</b>           | <b>0.44</b>         | <b>0.15</b>           |
| Jpp-K | <b>0.12</b>         | <b>0.08</b>           | <b>0.06</b>         | 0.01                  | <b>0.09</b>         | <b>0.07</b>           | <b>0.06</b>         | 0.04                  |
| K-A   | <b>0.54</b>         | <b>0.33</b>           | <b>0.31</b>         | <b>0.13</b>           | <b>0.35</b>         | <b>0.19</b>           | <b>0.53</b>         | <b>0.33</b>           |
| A-Ap  | <b>0.55</b>         | <b>0.35</b>           | <b>0.30</b>         | <b>0.07</b>           | <b>0.4</b>          | <b>0.23</b>           | <b>0.38</b>         | <b>0.20</b>           |
| Ap-B  | <b>0.86</b>         | <b>0.51</b>           | <b>0.42</b>         | <b>0.08</b>           | <b>0.6</b>          | <b>0.4</b>            | <b>0.65</b>         | <b>0.18</b>           |

### 299 3.3 Attribution of precipitation trends in CESM1

300 An initial assessment of the contribution from different external forcings to the statistically significant positive ensemble  
301 mean trends for CESM1 ALL for both total and extreme precipitation over Ellsworth Land, the Filchner-Ronne basin,  
302 Dronning Maud Land, Enderby Land, and the Amery ice shelf (i.e., the regions in agreement with ERA5) was made by  
303 comparing these trends with the CESM1 ensemble mean trends for GHG, AER and O3 (Fig. 2). These results show that the  
304 ensemble mean trends in total and extreme precipitation for GHG, AER, and O3 are mostly positive and significant over  
305 these regions, indicating that greenhouse gases, anthropogenic aerosols, and stratospheric ozone are all important drivers of  
306 the positive precipitation trends in the ALL response over these regions. However, for extreme precipitation, the trends for  
307 GHG and O3 are notably weaker, sometimes non-significant, and less spatially coherent than the AER trends, indicating that  
308 greenhouse gases and stratospheric ozone are less important for driving the positive extreme precipitation trends in the ALL  
309 response over these regions.

310 To assess the contribution from different external forcings to the positive ensemble mean trends for CESM1 ALL for both  
311 total and extreme precipitation for the five target basins (Figs. 3 and 4), Table 1 compares basin-averaged ensemble mean  
312 trends for ALL, GHG, AER and O3 for each of these sites. For total precipitation, the basin-averaged ensemble mean trends  
313 for ALL are significant and positive across all five basins, with values ranging from 0.12 to 0.86 mm year<sup>-1</sup>. By comparison,  
314 the trends for total precipitation for: i) GHG are also significant and positive across all basins, but with values around half  
315 that of the ALL trends, ranging from 0.06 to 0.42 mm year<sup>-1</sup>, and ii) significant and positive and largely similar for AER and  
316 O3 across all five basins, with values slightly larger than the GHG trends but lower than ALL, varying from 0.09 to 0.60 mm  
317 year<sup>-1</sup> for AER and from 0.06 to 0.65 mm year<sup>-1</sup> for O3. However, for basin K-A (Dronning Maud Land) the total

precipitation trend for O3 and ALL are 0.53 and 0.54 mm year<sup>-1</sup>, respectively, so comparable in size. For extreme precipitation, the ensemble mean trends for ALL are also significant and positive across all five basins, but with values considerably less than for the total precipitation trends, varying from 0.08 to 0.51 mm year<sup>-1</sup>. By comparison, the trends for extreme precipitation for: i) GHG are significant and positive for four of the five basins, with values ranging from 0.07 to 0.13 mm year<sup>-1</sup>, which are typically around a fifth of the ALL trends, ii) AER are significant and positive for all five basins, ranging from 0.07 to 0.4 mm year<sup>-1</sup>, which are much higher than the GHG trends and slightly less than the ALL trends (consistent with Fig. 2), and iii) O3 are significant and positive for four of the five basins, with values typically lower than AER but higher than GHG trends, ranging from 0.15 to 0.33 mm year<sup>-1</sup> (consistent with Fig. 2).




**Figure 6: Probability density functions of annual trends in (a-e) total and (f-j) extreme precipitation for drainage basins (a, f) J-Jpp, (b, g) Jpp-K (both part of Filchner-Ronne basin), (c, h) K-A, (d, i) A-Ap (both part of Dronning Maud Land), and (e, j) Ap-B (Enderby Land) from 1979 to 2023 (mm year<sup>-1</sup>) for all of the CESM1 ensembles for ALL (blue), GHG (green), AER (orange) and O3 (red). The results are based only on annual trends from grid points in the basin that are statistically significant at the 90% confidence level.**

The importance of greenhouse gases, anthropogenic aerosol and stratospheric ozone forcings for producing the positive trends in total and extreme precipitation over the five selected basins is also evident from comparison of the probability distribution functions of the trends for all of the CESM1 ensembles for ALL, GHG, AER and O3 (Fig. 6). These results show considerable similarities between the different distributions, which are generally shifted towards positive trends and largely overlap each other, although ALL is especially positively skewed and generally has a longer right-hand-side tail (consistent with Table 1). However, for total precipitation the results also show that the O3 distributions are especially positively skewed



338 and have a longer right-hand-side tail compared to GHG and AER (consistent with Table 1), and in several basins overlap  
339 with ALL (e.g., basin K-A, Dronning Maud Land). However, may partly reflect the shorter O3 simulation period  
340 (1979-2005) and the smaller ensemble size (8 members), both of which increase the sensitivity of the estimated trends to  
341 internal variability. For extreme precipitation, the results show that the GHG distributions are less positively skewed / weaker  
342 compared to O3 and AER (consistent with Table 1), suggesting again that greenhouse gases are less important for driving the  
343 positive extreme precipitation trends.

344 Figure 7 presents the two-signal D&A results for total and extreme precipitation across the five selected basins, showing  
345 scaling factors (and confidence intervals) for the CESM1 ensembles GHG, AER, and O3. The results indicate that the  
346 precipitation signal in four basins is formally related to increased greenhouse gas emissions. As for the stratospheric ozone  
347 depletion, the precipitation signal for three basins presents a statistically significant impact. Changes in anthropogenic  
348 aerosols are associated with precipitation changes in two basins. For GHG, its signal is found in basins J-Jpp  
349 (Filchner-Ronne basin, extreme precipitation only), Jpp-K (Filcher-Ronne basin, total and extreme precipitation), A-Ap  
350 (Dronning Maud Land, total and extreme precipitation), and Ap-B (Enderby Land, total and extreme precipitation). These  
351 were generally associated with relatively small confidence intervals, i.e., indicating high confidence that greenhouse gas  
352 forcing was responsible for the ERA5 precipitation changes in these basins. The one exception to this is extreme precipitation  
353 for basin Jpp-K, which had a relatively larger confidence interval, indicating lower confidence in the estimated scaling factor.  
354 For AER, its signal is found in basins J-Jpp (Filchner-Ronne basin, extreme precipitation only) and K-A (Dronning Maud  
355 Land, total and extreme precipitation). However, the result for extreme precipitation for basin J-Jpp was associated with a  
356 relatively small confidence interval (indicating higher confidence), while total precipitation for basin K-A was associated  
357 with a relatively large confidence interval (indicating lower confidence). For O3, its signal is found in basins J-Jpp  
358 (Filchner-Ronne basin, total and extreme precipitation), K-A (Dronning Maud Land, total precipitation only) and Ap-B  
359 (Enderby Land, total and extreme precipitation). The signals were again characterised by differences in confidence intervals,  
360 with the results for extreme precipitation associated with relatively small confidence intervals (indicating higher confidence),  
361 and total precipitation associated with relatively larger confidence intervals (indicating lower confidence). The remaining  
362 results were associated with confidence intervals that included zero, indicating that the precipitation signal cannot be  
363 formally associated with these external forcings.



364

365 Figure 7: As Fig. 5, but showing two-signal estimates of basin-averaged scaling factors for annual (a) total and (b) extreme  
 366 precipitation for drainage basins J-Jpp, Jpp-K (both part of Filchner-Ronne basin), K-A, A-Ap (both part of Dronning Maud  
 367 Land), and Ap-B (Enderby Land) from 1979 to 2023 for the CESM1 ensembles for GHG (solid green circles), AER (solid orange  
 368 circles), and O3 (solid red circles). The GHG scaling factors are based on regressing ALL and xGHG against ERA5, AER scaling  
 369 factors based on regressing ALL and xAER against ERA5, and O3 scaling factors based on regressing ALL and xO3 against  
 370 ERA5. The error bars denote the 90% confidence intervals and the horizontal line indicates a scaling factor of one.

### 371 3.4 Detection and attribution of precipitation trends using CESM2


372 Figures S1 to S6 in the Supplementary Material are analogous to Figs. 2 to 7, but based on results from the CESM2  
 373 ensembles for ALL, GHG, and AER. For the detection analysis, the results based on the CESM2 ALL ensemble are broadly  
 374 consistent with those based on the CESM1 ALL ensemble, showing significant positive annual trends in total and extreme  
 375 precipitation over the same regions, which agree with ERA5 (Fig. S1). Additionally, the results based on the CESM2 ALL  
 376 ensemble also identify the same five basins that are associated with significant positive trends in both total and extreme  
 377 precipitation, which agree with ERA5 (Fig. S2). Although one difference between the CESM1 and CESM2 results is that the  
 378 CESM2 ensemble mean for ALL shows considerably higher precipitation amounts than ERA5 over the basins at the start of  
 379 the period analysed (Fig. S3), reflecting a mean state bias in the CESM2 model (Dunmire et al., 2022). Additionally, the  
 380 one-signal D&A results for total and extreme precipitation across the five selected basins, based on regression of the CESM2  
 381 ALL ensemble onto ERA5, suggest with high confidence that the signal from the ALL ensemble is detectable in ERA5 (Fig.  
 382 S4), consistent with CESM1. Although, because the scaling factors are less than one, this suggests that the CESM2 ALL  
 383 ensemble tends to overestimate the trends apparent in ERA5.

384 For the attribution analysis, the results show that the ensemble mean trends in total and extreme precipitation from the  
 385 CESM2 GHG ensemble are in good agreement with those from the CESM2 ALL ensemble over much of Antarctica (Fig.



386 S1), i.e., indicating that greenhouse gas forcing is the primary driver of the positive trends in the CESM2 ALL ensemble,  
387 consistent with CESM1. For example, the CESM2 GHG ensemble captures the positive trends from Marie Byrd Land to the  
388 western section of Wilkes Land that are apparent in ALL (Fig. S1). By contrast, the trends of the ensemble mean of the  
389 CESM2 AER ensemble are mostly negative for total and extreme precipitation over Antarctica, although areas that show  
390 significant trends are rather patchy (Fig. S1), i.e., indicating that the effects of increased anthropogenic aerosols tend to offset  
391 the positive trends from increased greenhouse gases, which is inconsistent with CESM1 results. The importance of  
392 greenhouse gas forcing for producing the positive trends in total and extreme precipitation over the five selected basins is  
393 also evident from probability distribution functions of the trends from the CESM2 ensembles for ALL, GHG, and AER,  
394 which shows considerable similarities between the distributions for the GHG and ALL trends (Fig. S5). The distribution of  
395 the AER trends also confirms the offsetting effect of anthropogenic aerosols for the Jpp-K and K-A basins, where AAER  
396 distributions are skewed towards negative values. By contrast, for the remaining basins the AAER distributions are more  
397 mixed, with both positive and negative trends.

398 Finally, Fig. S6 presents the three-signal D&A results for total and extreme precipitation across the five selected basins,  
399 based on regression of the CESM2 ensembles for ALL, GHG, and AER against ERA5, which produces scaling factors for  
400 GHG, AER, and OTHERS (i.e., all forcings other than greenhouse gases and anthropogenic aerosols). Here, the scaling  
401 factors were all associated with large confidence intervals that included zero. Therefore, the signals from the CESM2  
402 ensembles for GHG and AER (and OTHERS) cannot be considered statistically significant for either total or extreme  
403 precipitation. One exception is basin Jpp-K, where the greenhouse gas forcing is found in the precipitation signal. However,  
404 this result still lacks consistency as the upper bound of the confidence interval is less than one, which suggests that GHG  
405 tends to overestimate the trends apparent in ERA5.



406

407 Figure 8: Annual trends in AR-associated (a) total and (b) extreme precipitation over Antarctica from 1980 to 2023 (mm year<sup>-1</sup>)  
 408 from the AR detection dataset. Also shown are the relative contribution of the (c) AR-associated total precipitation trend to the  
 409 ERA5-based total precipitation trend (%), and the (d) AR-associated extreme precipitation trend to the ERA5-based extreme  
 410 precipitation trend (%). Stippling indicates regions where trends are statistically significant at the 90% confidence level. Also  
 411 shown in panel (a) are the locations of drainage basins J-Jpp, Jpp-K (both part of Filchner-Ronne basin), K-A, A-Ap (both part of  
 412 Dronning Maud Land), and Ap-B (Enderby Land).

### 413 3.5 Role of atmospheric rivers in driving precipitation trends

414 Finally, to assess the role of ARs in driving the positive trends of total and extreme precipitation that are apparent in ERA5  
 415 over these five basins (Figs. 2 and 3), we also show the annual trends in AR-associated total and extreme precipitation from  
 416 1980 to 2023 (Fig. 8a, b), as well as the relative contribution of these trends to the total and extreme precipitation trends from  
 417 ERA5 (Fig. 8c, d). The results show broadly significant positive trends in AR-associated total and extreme precipitation over

the basins considered, with trends around  $0.5 \text{ mm year}^{-1}$  over basins J-Jpp and Jpp-K (both part of Filchner-Ronne basin) and up to  $2 \text{ mm year}^{-1}$  over basins K-A, A-Ap (both part of Dronning Maud Land), and Ap-B (Enderby Land). Additionally, the relative contribution of the AR trends to the ERA5-based total and extreme precipitation trends has increased across all five basins, with values of 10 to 40% over sections of basins J-Jpp, Jpp-K (both part of Filchner-Ronne basin), and K-A (Dronning Maud Land), and up to 100% over sections of basins A-Ap (Dronning Maud Land) and Ap-B (Enderby Land).

#### 4 Discussion and Conclusions

This study confirms a statistically significant increase in both total and extreme precipitation (up to  $2 \text{ mm year}^{-1}$ ) across six drainage basins in Antarctica from 1979 to 2023, based on ERA5 reanalysis (Fig. 2). These are basin I-Ipp in the Antarctic Peninsula, basin J-Jpp in West Antarctica and basin Jpp-K in East Antarctica (both part of Filchner-Ronne basin), basins K-A and A-Ap in East Antarctica (both part of Dronning Maud Land), and basin Ap-B in East Antarctica (Enderby Land). For five of these six basins (J-Jpp, Jpp-K, A-Ap, Ap-B, and K-A) the total and extreme precipitation trends from the CESM1 ensembles for ALL are in good agreement with those from ERA5 (Figs. 2, 3 and 4), i.e., suggesting that a combination of both anthropogenic and natural forcings are responsible for the present-day trends in total and extreme precipitation over these basins. Similar results were found using the CESM2 ensembles for ALL (Figs. S1, S2, and S3).

One of the main challenges in detecting a trend in total and extreme precipitation is separating the signal from forced variability and internal variability, which is exacerbated in Antarctica by the climate being particularly characterized by strong internal variability (Jones et al., 2016; Previdi and Polvani, 2016; Stenni et al., 2017). Thus, detection of the signal in ERA5 in the CESM1 ensembles for ALL using the one-signal regression-based D&A analysis is therefore only confined to five drainage basins (Fig. 5) where the forced signal emerges more clearly above internal variability, with four of the five basins located in East Antarctica and the other in West Antarctica. This agrees with Casado et al. (2023), who found weaker multi-decadal variability in East Antarctica compared to West Antarctica, i.e., the relatively weaker internal multi-decadal variability in East Antarctica likely increases the signal-to-noise ratio, and enables a more robust detection of the forced signal in this region. Similar results were found using the CESM2 ensembles for ALL (Fig. S4).

Our initial attribution analysis using the CESM1 ensembles of the contribution from different external forcings to the statistically significant positive trends for both total and extreme precipitation for our five target basins indicated that the evolution of greenhouse gases, anthropogenic aerosols, and stratospheric ozone since 1979 are all important drivers of the trends (Figs. 2 and 6, Table 1). By contrast, the two-signal regression-based D&A analysis using the CESM1 ensembles for ALL and GHG, ALL and AER, and ALL and O3 (resulting in scaling factors for GHG, AER, and O3) found robust attribution to greenhouse gases in four basins and stratospheric ozone in three basins, and for anthropogenic aerosols in only one basin (Fig. 7). Repeating this initial attribution analysis using the CESM2 ensembles also indicated that greenhouse gases are an important driver of the precipitation trends over these basins (Figs. S1 and S5). However, these results also suggest

that increases in anthropogenic aerosol emissions can oppose / offset the increases in precipitation from increased greenhouse gases over our five basins, which contradicts the findings based on CESM1. Moreover, the three-signal regression-based D&A analysis using the CESM2 ensembles ALL, GHG and AER ensembles (resulting in scaling factors for GHG, AER, and OTHERS) was unable to provide robust attribution for total or extreme precipitation to any specific single-forcing, including greenhouse gases (Fig. S6). These inconsistencies between the results demonstrate that detecting the fingerprint of anthropogenic forcing on recent Antarctic precipitation changes remains challenging (Previdi and Polvani, 2016; Dalaiden et al., 2022). Furthermore, the combined and potentially non-linear effects of different forcings is also likely important for representing precipitation, and especially for extreme precipitation where linear assumptions are least valid (Meehl et al., 2003; Pope et al., 2020). Nevertheless, the results demonstrate that the effects of greenhouse gases and stratospheric ozone are particularly important for driving the present-day increases in total and extreme precipitation over the five target basins, while the role of anthropogenic aerosols remains uncertain.

We also showed that the trends in total and extreme precipitation over the five target basins can be linked to increasing ARs (Fig. 8). This provides physical support for the analysis, highlighting the consistency between the identification of the trends and the underlying dynamical drivers. This result is also consistent with previous results that suggested that ARs contribute up to 20% of total precipitation in East Antarctica (Wille et al., 2021, 2025; MacLennan et al., 2022), and as much as 50 to 70% of extreme precipitation (Wille et al., 2021, 2025; Simon et al., 2024). The increased ARs are likely associated with anomalous northerly airflow, combined with increased atmospheric moisture from warmer oceans, which could be linked to recent regional climate changes such as sea-ice loss in the East Antarctic sector and a strengthening of the low-pressure system off the East Antarctic coast (Goosse et al., 2024). Moreover, additional anomalous transport from the midlatitudes through Rossby-wave teleconnections could also be a factor (Wille et al., 2025). In addition, future increases in greenhouse gas concentrations are expected to strengthen Antarctic total and extreme precipitation by causing warmer atmospheric conditions (leading to more moisture, via the Clausius-Clapeyron relation) and a reduction in sea-ice cover (Frieler et al., 2015; Lenaerts et al., 2016; Previdi and Polvani, 2016; Kittel et al., 2021; Vignon et al., 2021; Zhu et al., 2023; Nicola et al., 2023).

It's noteworthy that the effects of increased greenhouse gas emissions are generally considered to be a positive trend in the Southern Annular Mode (SAM) (Kushner et al., 2001; Arblaster and Meehl, 2006), which corresponds to stronger, poleward-shifted westerlies around Antarctica, resulting in reduced temperatures (and precipitation) over East Antarctica (Thompson and Solomon, 2002). Based on this, as greenhouse gases are one of the primary drivers of the present-day trends, a decrease in precipitation and extreme precipitation over our five target basins (four of which are in East Antarctica) would be expected, rather than an increase. However, one reason for this contradiction may be a reversal in the relationship between SAM and temperature anomalies across East Antarctica during the twenty-first century, which occurred in response to anomalous high pressure over East Antarctica (Marshall et al., 2013). This anomalous circulation would result in increased

northerly airflow into East Antarctica, i.e., consistent with our results showing increased precipitation and intensification of ARs. Additionally, increased anthropogenic aerosol emissions are generally considered to result in a negative trend in the SAM (Gillett et al., 2013; Pope et al., 2020), corresponding to weaker circumpolar westerlies, which would likely result in enhanced precipitation over Antarctica - consistent with the results based on CESM1 only. Also, the Antarctic ozone hole has been the primary driver of summertime atmospheric circulation changes in the Southern hemisphere from around 1980 onwards (Thompson and Solomon, 2002; Polvani et al., 2011; Orr et al., 2021), which Dalaiden et al. (2022) showed was partly responsible for recent increases in precipitation over West Antarctica.

As a future perspective, given the importance of ARs for Antarctic surface climate, applying the D&A framework to changes in their properties (e.g., intensity, duration, and spatial extent) would represent an important step toward identifying the contribution of external forcing versus internal variability in the AR changes, as well as their associated impacts (e.g., heatwave or surface melting).

## **Code and data availability**

The code used in this study can be obtained from the corresponding author upon reasonable request. Precipitation output from the ERA5 reanalysis dataset is publicly available through the Copernicus Climate Data Store: <https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels?tab=overview>.

Precipitation output from the CESM1 ensembles with all external forcings, single-forcings and pre-industrial control simulation are available at: <https://gdex.ucar.edu/datasets/d651027/>. Precipitation output from the CESM1 ensembles of stratospheric ozone are available at: <https://gdex.ucar.edu/datasets/d651053/>. Precipitation output from the CESM2 ensembles with all external forcings are available at: <https://rda.ucar.edu/datasets/d651056/>. Precipitation output from the CESM2 ensembles with single-forcings are available at: <https://rda.ucar.edu/datasets/d651055/dataaccess/#>. Precipitation output from the CESM2 pre-industrial control simulation are available at: [https://www.earthsystemgrid.org/dataset/ucar.cgd.cesm2.b.e21.B1850.f09\\_g17.CMIP6-piControl.001.atm.proc.daily\\_ave.PR\\_ECT.html](https://www.earthsystemgrid.org/dataset/ucar.cgd.cesm2.b.e21.B1850.f09_g17.CMIP6-piControl.001.atm.proc.daily_ave.PR_ECT.html). The AR detection dataset derived from ERA5 is available at: <https://zenodo.org/records/15830634>.

## **Author contributions**

SPSC designed the study and, together with PD and AO, conceptualized it. SPSC carried out the primary analysis with support from PD, AO, and QD. SPSC and AO were primarily responsible for writing the manuscript, with additions from PD and QD.

## 508 Competing interests

509 The corresponding author affirms that none of the authors have any financial or personal conflicts of interest to disclose.

## 510 Acknowledgement

511 The authors would like to thank the two reviewers for providing such thoughtful and expert comments, which greatly  
512 improved the scope and overall quality of the manuscript. We would also like to thank Jonathan Wille for his advice on using  
513 the AR detection dataset.

## 514 Financial support

515 AO received support from the European Union's Horizon 2020 funded project PolarRES (Polar Regions in the Earth System;  
516 Grant No. 101003590) and the Natural Environment Research Council funded project PICANTE (Processes, Impacts and  
517 Changes of ANTArctic Extreme weather; Grant No. NE/Z503356/1). SPSC and PD received funding from the Indian Institute  
518 of Technology Kharagpur and the Ministry of Education, Government of India. QD received support from European Union's  
519 Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 101149188.

## 520 References

- 521 Adusumilli, S., A Fish, M., Fricker, H. A., & Medley, B. (2021). Atmospheric river precipitation contributed to rapid  
522 increases in surface height of the west Antarctic ice sheet in 2019. *Geophysical Research Letters*, 48(5), e2020GL091076.  
523 <https://doi.org/10.1029/2020GL091076>
- 524 Arblaster, J. M., & Meehl, G. A. (2006). Contributions of external forcings to southern annular mode trends. *Journal of*  
525 *climate*, 19(12), 2896-2905. <https://doi.org/10.1175/JCLI3774.1>
- 526 Allen, M. R., & Stott, P. A. (2003). Estimating signal amplitudes in optimal fingerprinting, Part I: Theory. *Climate Dynamics*,  
527 21, 477-491. <https://doi.org/10.1007/s00382-003-0313-9>
- 528 Banwell, A. F., and D. R. MacAyeal, 2015: Ice-shelf fracture due to viscoelastic flexure stress induced by fill/drain cycles of  
529 supraglacial lakes. *Antarct. Sci.*, 27, 587–597, <https://doi.org/10.1017/S0954102015000292>.
- 530 Boening, C., Lebsock, M., Landerer, F., & Stephens, G. (2012). Snowfall-driven mass change on the East Antarctic ice sheet.  
531 *Geophysical Research Letters*, 39, L21501. <https://doi.org/10.1029/2012GL053316>

532 Bromwich, D. H., Ensign, A., Wang, S. H., & Zou, X. (2024). Major artifacts in ERA5 2-m air temperature trends over  
 533 Antarctica prior to and during the modern satellite era. *Geophysical Research Letters*, 51(21), e2024GL111907.  
 534 <https://doi.org/10.1029/2024GL111907>

535 Carter, J., A. Leeson, A. Orr, C. Kittel, J. Melchior van Wessem (2022), Variability in Antarctic surface climatology across  
 536 region climate models and reanalysis datasets, *The Cryosphere*, 16, 3815–3841, <https://doi.org/10.5194/tc-16-3815-2022>

537 Casado, M., Hébert, R., Faranda, D., & Landais, A. (2023). The quandary of detecting the signature of climate change in  
 538 Antarctica. *Nature Climate Change*, 13(10), 1082-1088. <https://doi.org/10.1038/s41558-023-01791-5>

539 Clem, K. R., & Raphael, M. N. (2024). STATE OF THE CLIMATE IN 2023. <https://doi.org/10.1175/BAMS-D-24-0099.1>

540 Connolley, W. M. (1997). Variability in annual mean circulation in southern high latitudes. *Climate Dynamics*, 13, 745-756.  
 541 <https://doi.org/10.1007/s003820050195>

542 Dalaiden, Q., Goosse, H., Rezsöhazi, J., & Thomas, E. R. (2021). Reconstructing atmospheric circulation and sea-ice extent  
 543 in the West Antarctic over the past 200 years using data assimilation. *Climate Dynamics*, 57(11), 3479-3503.  
 544 <https://doi.org/10.1007/s00382-021-05879-6>

545 Dalaiden, Q., Schurer, A. P., Kirchmeier-Young, M. C., Goosse, H., & Hegerl, G. C. (2022). West Antarctic surface climate  
 546 changes since the mid-20th century driven by anthropogenic forcing. *Geophysical Research Letters*, 49(16),  
 547 e2022GL099543. <https://doi.org/10.1029/2022GL099543>

548 Danabasoglu, G., Lamarque, J. F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., ... & Strand, W. G. (2020). The  
 549 community earth system model version 2 (CESM2). *Journal of Advances in Modeling Earth Systems*, 12(2),  
 550 e2019MS001916. <https://doi.org/10.1029/2019MS001916>

551 Davison, B. J., Hogg, A. E., Rigby, R., Veldhuijsen, S., van Wessem, J. M., van den Broeke, M. R., ... & Dutrieux, P. (2023).  
 552 Sea level rise from West Antarctic mass loss significantly modified by large snowfall anomalies. *Nature Communications*,  
 553 14(1), 1479. <https://doi.org/10.1038/s41467-023-36990-3>

554 Deb, P., Orr, A., Bromwich, D. H., Nicolas, J. P., Turner, J., & Hosking, J. S. (2018). Summer drivers of atmospheric  
 555 variability affecting ice shelf thinning in the Amundsen Sea Embayment, West Antarctica. *Geophysical Research Letters*,  
 556 45(9), 4124-4133. <https://doi.org/10.1029/2018GL077092>

557 Deser, C., Phillips, A. S., Simpson, I. R., Rosenbloom, N., Coleman, D., Lehner, F., ... & Stevenson, S. (2020). Isolating the  
 558 evolving contributions of anthropogenic aerosols and greenhouse gases: A new CESM1 large ensemble community resource.  
 559 *Journal of climate*, 33(18), 7835-7858. <https://doi.org/10.1175/JCLI-D-20-0123.1>



560 Dong, S., Sun, Y., & Li, C. (2020). Detection of human influence on precipitation extremes in Asia. *Journal of Climate*,  
561 33(12), 5293-5304. <https://doi.org/10.1175/JCLI-D-19-0371.1>

562 Dong, S., Sun, Y., Li, C., Zhang, X., Min, S. K., & Kim, Y. H. (2021). Attribution of extreme precipitation with updated  
563 observations and CMIP6 simulations. *Journal of Climate*, 34(3), 871-881. <https://doi.org/10.1175/JCLI-D-19-1017.1>

564 Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., Dunn, R. J., ... & Kitching, S. (2013). Updated analyses of  
565 temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset. *Journal of*  
566 *Geophysical Research: Atmospheres*, 118(5), 2098-2118. <https://doi.org/10.1002/jgrd.50150>

567 Dunmire, D., Lenaerts, J. T., Datta, R. T., & Gorte, T. (2022). Antarctic surface climate and surface mass balance in the  
568 Community Earth System Model version 2 during the satellite era and into the future (1979–2100). *The Cryosphere*, 16(10),  
569 4163-4184. <https://doi.org/10.5194/tc-16-4163-2022>

570 Dunn, R. J., Alexander, L. V., Donat, M. G., Zhang, X., Bador, M., Herold, N., ... & Bin Hj Yussof, M. N. A. (2020).  
571 Development of an updated global land in situ-based data set of temperature and precipitation extremes: HadEX3. *Journal of*  
572 *Geophysical Research: Atmospheres*, 125(16), e2019JD032263. <https://doi.org/10.1029/2019JD032263>

573 England, M. R., Polvani, L. M., Smith, K. L., Landrum, L., & Holland, M. M. (2016). Robust response of the Amundsen Sea  
574 Low to stratospheric ozone depletion. *Geophysical Research Letters*, 43(15), 8207–8213.  
575 <https://doi.org/10.1002/2016GL070055>

576 Estrada, F., Perron, P., & Yamamoto, Y. (2023). Anthropogenic influence on extremes and risk hotspots. *Scientific Reports*,  
577 13(1), 35. <https://doi.org/10.1038/s41598-022-27220-9>

578 Frieler, K., Clark, P. U., He, F., Buizert, C., Reese, R., Ligtenberg, S. R. M., et al. (2015). Consistent evidence of increasing  
579 Antarctic accumulation with warming. *Nature Climate Change*, 5(4), 348–352. <https://doi.org/10.1038/nclimate2574>

580 Favier, V., Krinner, G., Amory, C., Gallée, H., Beaumet, J., & Agosta, C. (2017). Antarctica-regional climate and surface  
581 mass budget. *Current Climate Change Reports*, 3(4), 303-315. <https://doi.org/10.1007/s40641-017-0072-z>

582 Fyke, J., Lenaerts, J. T. M., & Wang, H. (2017). Basin-scale heterogeneity in Antarctic precipitation and its impact on surface  
583 mass variability. *The Cryosphere*, 11(6), 2595–2609. <https://doi.org/10.5194/tc-11-2595-2017>

584 Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., Van Den Broeke, M., & Nilsson, J. (2018).  
585 Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years. *The Cryosphere*, 12(2), 521-547.  
586 <https://doi.org/10.5194/tc-12-521-2018>

587 Gehring, J., Vignon, É., Billault-Roux, A. C., Ferrone, A., Protat, A., Alexander, S. P., & Berne, A. (2022). Orographic flow  
588 influence on precipitation during an atmospheric river event at Davis, Antarctica. *Journal of Geophysical Research:*  
589 *Atmospheres*, 127(2), e2021JD035210. <https://doi.org/10.1029/2021JD035210>

590 Gillett, N. P., Stone, D. A., Stott, P. A., Nozawa, T., Karpechko, A. Y., Hegerl, G. C., ... & Jones, P. D. (2008). Attribution of  
591 polar warming to human influence. *Nature Geoscience*, 1(11), 750-754. <https://doi.org/10.1038/ngeo338>

592 Gillett, N. P., Fyfe, J. C., & Parker, D. E. (2013). Attribution of observed sea level pressure trends to greenhouse gas, aerosol,  
593 and ozone changes. *Geophysical Research Letters*, 40(10), 2302-2306. <https://doi.org/10.1002/grl.50500>

594 Gillett, N. P., Kirchmeier-Young, M., Ribes, A., Shiogama, H., Hegerl, G. C., Knutti, R., ... & Ziehn, T. (2021). Constraining  
595 human contributions to observed warming since the pre-industrial period. *Nature Climate Change*, 11(3), 207-212.  
596 <https://doi.org/10.1038/s41558-020-00965-9>

597 Gilbert, E., D. Pishniak, J. A. Torres, A. Orr, M. MacLennan, N. Wever, and K. Verro (2025), Extreme precipitation associated  
598 with atmospheric rivers over West Antarctic ice shelves: insights from kilometre-scale regional climate modelling, *The*  
599 *Cryosphere*, 19, 597–618, <https://doi.org/10.5194/tc-19-597-2025>.

600 Goosse, H., Dalaiden, Q., Feba, F., Mezzina, B., & Fogt, R. L. (2024). A drop in Antarctic sea ice extent at the end of the  
601 1970s. *Communications Earth & Environment*, 5(1), 628. <https://doi.org/10.1038/s43247-024-01793-x>

602 Gorodetskaya, I. V., Tsukernik, M., Claes, K., Ralph, M. F., Neff, W. D., & Van Lipzig, N. P. (2014). The role of atmospheric  
603 rivers in anomalous snow accumulation in East Antarctica. *Geophysical Research Letters*, 41(17), 6199-6206.  
604 Johnson <https://doi.org/10.1002/2014GL060881>

605 Gu, C., Luo, Y., Liu, F., Lu, J., & Chen, Z. (2025). Increased Asian aerosols contribute to historical climate change in  
606 Antarctica. *Geophysical Research Letters*, 52(13), e2025GL114888. <https://doi.org/10.1029/2025GL114888>

607 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... & Thépaut, J. N. (2020). The ERA5  
608 global reanalysis. *Quarterly journal of the royal meteorological society*, 146(730), 1999-2049.  
609 <https://doi.org/10.1002/qj.3803>

610 Hawkins, E., & Sutton, R. (2009). The potential to narrow uncertainty in regional climate predictions. *Bulletin of the*  
611 *American Meteorological Society*, 90(8), 1095-1108. <https://doi.org/10.1175/2009BAMS2607.1>

612 Hobbs, W. R., Roach, C., Roy, T., Sallée, J. B., & Bindoff, N. (2021). Anthropogenic temperature and salinity changes in the  
613 Southern Ocean. *Journal of Climate*, 34(1), 215-228. <https://doi.org/10.1175/JCLI-D-20-0454.1>

614 Hosking, J. S., Orr, A., Marshall, G. J., Turner, J., & Phillips, T. (2013). The influence of the Amundsen–Bellingshausen Seas  
 615 low on the climate of West Antarctica and its representation in coupled climate model simulations. *Journal of Climate*,  
 616 26(17), 6633–6648. <https://doi.org/10.1175/JCLI-D-12-00813.1>

617 IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment  
 618 Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva,  
 619 Switzerland, 184 pp. <https://doi.org/10.59327/IPCC/AR6-9789291691647>.

620 Johnson, A., R. Hock, and M. Fahnestock, 2022: Spatial variability and regional trends of Antarctic Ice Shelf surface melt  
 621 duration over 1979–2020 derived from passive microwave data. *J. Glaciol.*, 68, 533–546,  
 622 <https://doi.org/10.1017/jog.2021.112>.

623 Jones, J. M., Gille, S. T., Goosse, H., Abram, N. J., Canziani, P. O., Charman, D. J., ... & Vance, T. R. (2016). Assessing  
 624 recent trends in high-latitude Southern Hemisphere surface climate. *Nature Climate Change*, 6(10), 917–926.  
 625 <https://doi.org/10.1038/nclimate3103>

626 Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., et al. (2015). The community Earth system model (CESM)  
 627 large ensemble project: A community resource for studying climate change in the presence of internal climate variability.  
 628 *Bulletin of the American Meteorological Society*, 96(8), 1333–1349. <https://doi.org/10.1175/BAMS-D-13-00255.1>

629 Kirchmeier-Young, M. C., Zwiers, F. W., & Gillett, N. P. (2017). Attribution of extreme events in Arctic sea ice extent.  
 630 *Journal of Climate*, 30(2), 553–571. <https://doi.org/10.1175/JCLI-D-16-0412.1>

631 Kirchmeier-Young, M. C., & Zhang, X. (2020). Human influence has intensified extreme precipitation in North America.  
 632 *Proceedings of the National Academy of Sciences*, 117(24), 13308–13313. <https://doi.org/10.1073/pnas.1921628117>

633 Kittel, C., Amory, C., Agosta, C., Jourdain, N. C., Hofer, S., Delhasse, A., ... & Fettweis, X. (2021). Diverging future surface  
 634 mass balance between the Antarctic ice shelves and grounded ice sheet. *The Cryosphere*, 15(3), 1215–1236.  
 635 <https://doi.org/10.5194/tc-15-1215-2021>

636 Kromer, J. D., & Trusel, L. D. (2023). Identifying the impacts of sea ice variability on the climate and surface mass balance  
 637 of West Antarctica. *Geophysical Research Letters*, 50(18), e2023GL104436. <https://doi.org/10.1029/2023GL104436>

638 Kushner, P. J., Held, I. M., & Delworth, T. L. (2001). Southern Hemisphere atmospheric circulation response to global  
 639 warming. *Journal of Climate*, 14(10), 2238–2249. [https://doi.org/10.1175/1520-0442\(2001\)014<0001:SHACRT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<0001:SHACRT>2.0.CO;2)

640 Lai, C.-Y., J. Kingslake, M. G. Wearing, P.-H. Cameron Chen, P. Gentine, H. Li, J. J. Spergel, and J. M. van Wessem, 2020:  
 641 Vulnerability of Antarctica's ice shelves to meltwater-driven fracture. *Nature*, 584, 574–578,  
 642 <https://doi.org/10.1038/s41586-020-2627-8>.

643 Landrum, L. L., Holland, M. M., Raphael, M. N., & Polvani, L. M. (2017). Stratospheric ozone depletion: An unlikely driver  
 644 of the regional trends in Antarctic Sea ice in Austral fall in the late twentieth century. *Geophysical Research Letters*, 44(21),  
 645 11062–11070. <https://doi.org/10.1002/2017GL075618>

646 Lenaerts, J. T. M., Vizcaino, M., Fyke, J., van Kampenhout, L., & van den Broeke, M. R. (2016). Present-day and future  
 647 Antarctic ice sheet climate and surface mass balance in the Community Earth System Model. *Climate Dynamics*, 47(5),  
 648 1367–1381. <https://doi.org/10.1007/s00382-015-2907-4>

649 Lenaerts, J. T., Fyke, J., & Medley, B. (2018). The signature of ozone depletion in recent Antarctic precipitation change: A  
 650 study with the Community Earth System Model. *Geophysical Research Letters*, 45(23), 12–931.  
 651 <https://doi.org/10.1029/2018GL078608>

652 Li, S., Chen, Y., Wei, W., Fang, G., & Duan, W. (2024). The increase in extreme precipitation and its proportion over global  
 653 land. *Journal of Hydrology*, 628, 130456. <https://doi.org/10.1016/j.jhydrol.2023.130456>

654 MacLennan, M. L., Lenaerts, J. T., Shields, C., & Wille, J. D. (2022). Contribution of atmospheric rivers to Antarctic  
 655 precipitation. *Geophysical Research Letters*, 49(18), e2022GL100585. <https://doi.org/10.1029/2022GL100585>

656 MacLennan, M. L., Lenaerts, J., Shields, C. A., Hoffman, A. O., Wever, N., Thompson-Munson, M., ... & Wille, J. D. (2023).  
 657 Climatology and surface impacts of atmospheric rivers on West Antarctica. *The Cryosphere*, 17(2), 865–881.  
 658 <https://doi.org/10.5194/tc-17-865-2023>

659 Marshall, G. J. (2003). Trends in the Southern annular mode from observations and reanalyses. *Journal of Climate*, 16(24),  
 660 4134–4143. [https://doi.org/10.1175/1520-0442\(2003\)016<4134:TITSAM>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<4134:TITSAM>2.0.CO;2) Munneke

661 Marshall, G. J., Orr, A., & Turner, J. (2013). A predominant reversal in the relationship between the SAM and East Antarctic  
 662 temperatures during the twenty-first century. *Journal of Climate*, 26(14), 5196–5204.  
 663 <https://doi.org/10.1175/JCLI-D-12-00671.1>

664 Meehl, G. A., Washington, W. M., Wigley, T. M. L., Arblaster, J. M., & Dai, A. (2003). Solar and greenhouse gas forcing and  
 665 climate response in the twentieth century. *Journal of Climate*, 16(3), 426–444.  
 666 [https://doi.org/10.1175/1520-0442\(2003\)016<0426:SAGGFA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0426:SAGGFA>2.0.CO;2)

667 Mémin, A., Flament, T., Alizier, B., Watson, C., & Rémy, F. (2015). Interannual variation of the Antarctic Ice Sheet from a  
 668 combined analysis of satellite gravimetry and altimetry data. *Earth and Planetary Science Letters*, 422, 150–156.  
 669 <https://doi.org/https://doi.org/10.1016/j.epsl.2015.03.045>

670 Min, S. K., Zhang, X., Zwiers, F. W., & Hegerl, G. C. (2011). Human contribution to more-intense precipitation extremes.  
 671 *Nature*, 470(7334), 378–381. <https://www.nature.com/articles/nature09763>

672 Morioka, Y., Zhang, L., Cooke, W., Nonaka, M., Behera, S. K., & Manabe, S. (2024). Role of anthropogenic forcing in  
 673 Antarctic sea ice variability simulated in climate models. *Nature Communications*, 15(1), 10511.  
 674 <https://doi.org/10.1038/s41467-024-54485-7>

675 Mottram, R., Hansen, N., Kittel, C., Van Wessem, J. M., Agosta, C., Amory, C., ... & Souverijns, N. (2021). What is the  
 676 surface mass balance of Antarctica? An intercomparison of regional climate model estimates. *The Cryosphere*, 15(8),  
 677 3751–3784. <https://doi.org/10.5194/tc-15-3751-2021>

678 Munneke, P. K., Ligtenberg, S. R., Van Den Broeke, M. R., & Vaughan, D. G. (2014). Firm air depletion as a precursor of  
 679 Antarctic ice-shelf collapse. *Journal of Glaciology*, 60(220), 205–214. <https://doi.org/10.3189/2014JoG13J183>

680 Nakayama, Y., Greene, C. A., Paolo, F. S., Mensah, V., Zhang, H., Kashiwase, H., ... & Aoki, S. (2021). Antarctic slope  
 681 current modulates ocean heat intrusions towards Totten Glacier. *Geophysical Research Letters*, 48(17), e2021GL094149.  
 682 <https://doi.org/10.1029/2021GL094149>

683 Nicola, L., Notz, D., & Winkelmann, R. (2023). Revisiting temperature sensitivity: how does Antarctic precipitation change  
 684 with temperature?. *The Cryosphere*, 17(7), 2563–2583. <https://doi.org/10.5194/tc-17-2563-2023>

685 Orr, A., Lu, H., Martineau, P., Gerber, E. P., Marshall, G. J., & Bracegirdle, T. J. (2021). Is our dynamical understanding of  
 686 the circulation changes associated with the Antarctic ozone hole sensitive to the choice of reanalysis dataset?. *Atmospheric*  
 687 *Chemistry and Physics*, 21(10), 7451–7472. <https://doi.org/10.5194/acp-21-7451-2021>

688 Orr, A., Deb, P., Clem, K. R., Gilbert, E., Bromwich, D. H., Boberg, F., ... & Zou, X. (2023). Characteristics of surface “melt  
 689 potential” over Antarctic ice shelves based on regional atmospheric model simulations of summer air temperature extremes  
 690 from 1979/80 to 2018/19. *Journal of Climate*, 36(10), 3357–3383. <https://doi.org/10.1175/JCLI-D-22-0386.1>

691 Otosaka, I. N., Horwath, M., Mottram, R., & Nowicki, S. (2023). Mass balances of the Antarctic and Greenland ice sheets  
 692 monitored from space. *Surveys in Geophysics*, 44(5), 1615–1652. <https://doi.org/10.1007/s10712-023-09795-8>

693 Paik, S., Min, S. K., Zhang, X., Donat, M. G., King, A. D., & Sun, Q. (2020). Determining the anthropogenic greenhouse gas  
 694 contribution to the observed intensification of extreme precipitation. *Geophysical Research Letters*, 47(12), e2019GL086875.  
 695 <https://doi.org/10.1029/2019GL086875>

696 Paolo, F. S., Fricker, H. A., & Padman, L. (2015). Volume loss from Antarctic ice shelves is accelerating. *Science*,  
 697 348(6232), 327-331. DOI: [10.1126/science.aaa0940](https://doi.org/10.1126/science.aaa0940)

698 Paolo, F., Padman, L., Fricker, H., Adusumilli, S., Howard, S., & Siegfried, M. (2018). Response of Pacific-sector Antarctic  
 699 ice shelves to the el Niño/southern oscillation. *Nature Geoscience*, 11(2), 121–126.  
 700 <https://doi.org/10.1038/s41561-017-0033-0>

701 Pattyn, F. (2017). Sea-level response to melting of Antarctic ice shelves on multi-centennial timescales with the fast  
 702 Elementary Thermomechanical Ice Sheet model (f. ETISh v1. 0). *The Cryosphere*, 11(4), 1851-1878.  
 703 <https://doi.org/10.5194/tc-11-1851-2017>

704 Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy,  
 705 J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)). Cambridge University Press,  
 706 Cambridge, United Kingdom and New York, NY, USA, pp. 1513–1766, doi: [10.1017/9781009157896.013](https://doi.org/10.1017/9781009157896.013).

707 Pope, J. O., Orr, A., Marshall, G. J., & Abraham, N. L. (2020). Non-additive response of the high-latitude Southern  
 708 Hemisphere climate to aerosol forcing in a climate model with interactive chemistry. *Atmospheric Science Letters*, 21(12),  
 709 e1004. <https://doi.org/10.1002/asl.1004>

710 Previdi, M., & Polvani, L. M. (2016). Anthropogenic impact on Antarctic surface mass balance, currently masked by natural  
 711 variability, to emerge by mid-century. *Environmental Research Letters*, 11(9), 094001. DOI 10.1088/1748-9326/11/9/094001

712 Pritchard, H. D., S. R. M. Ligtenberg, H. A. Fricker, D. G. Vaughan, M. R. van den Broeke, and L. Padman, 2012: Antarctic  
 713 ice-sheet loss driven by basal melting of ice shelves. *Nature*, 484, 502–505, <https://doi.org/10.1038/nature10968>.

714 Pritchard, H. D. (2021), Global data gaps in our knowledge of the terrestrial cryosphere, *Front. Clim.*, 3,  
 715 <https://doi.org/10.3389/fclim.2021.689823>

716 Ribes, A., Planton, S., & Terray, L. (2013). Application of regularised optimal fingerprinting to attribution. Part I: Method,  
 717 properties and idealised analysis. *Climate dynamics*, 41, 2817-283 <https://doi.org/10.1007/s00382-013-1735-7>

718 Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., & Morlighem, M. (2019). Four decades of  
719 Antarctic ice sheet mass balance from 1979–2017. *Proceedings of the National Academy of Sciences - PNAS*, 116(4),  
720 1095–1103. <https://doi.org/10.1073/pnas.1812883116>

721 Rott, H., P. Skvarca, and T. Nagler, 1996: Rapid collapse of northern Larsen Ice Shelf, Antarctica. *Science*, 271, 788–792,  
722 <https://doi.org/10.1126/science.271.5250.788>.

723 Scambos, T. A., C. Hulbe, M. Fahnestock, and J. Bohlander, 2000: The link between climate warming and break-up of ice  
724 shelves in the Antarctic Peninsula. *J. Glaciol.*, 46, 516–530, <https://doi.org/10.3189/172756500781833043>.

725 Seneviratne, S.I., X. Zhang, M. Adnan, W. Badi, C. Dereczynski, A. Di Luca, S. Ghosh, I. Iskandar, J. Kossin, S. Lewis, F.  
726 Otto, I. Pinto, M. Satoh, S.M. Vicente-Serrano, M. Wehner, and B. Zhou, 2021: Weather and Climate Extreme Events in a  
727 Changing Climate. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth*  
728 *Assessment Report of the Intergovernmental Panel on Climate Change*[Masson-Delmotte, V., P. Zhai, A.

729 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M. R., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I.,  
730 Krinner, G., Nowicki, S., Payne, T., Scambos, T. A., Schlegel, N., A. G., Agosta, C., Ahlstrøm, A., Babonis, G., Barletta, V.,  
731 Blazquez, A., Bonin, J., Csatho, B., Cullather, R., Felikson, D., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A., Gilbert,  
732 L., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horvath, M., Khan, S., Kjeldsen, K. K., Konrad, H.,  
733 Langen, P., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P.,  
734 Mouginot, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B., Otosaka, I., Pattle, M. E., Peltier, W. R., Pie,  
735 N., Rietbroek, R., Rott, H., Sandberg-Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W.,  
736 Simonsen, S., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van Wessem,  
737 M., Vishwakarma, B. D., Wiese, D., Wouters, B., and team, T. I.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017,  
738 *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>

739 Smith, K. L., & Polvani, L. M. (2017). Spatial patterns of recent Antarctic surface temperature trends and the importance of  
740 natural variability: lessons from multiple reconstructions and the cmip5 models. *Climate Dynamics*, 48(7), 2653-2670.  
741 DOI: 10.1007/s00382-016-3230-4

742 Simon, S., Turner, J., Thamban, M., Wille, J. D., & Deb, P. (2024). Spatiotemporal variability of extreme precipitation events  
743 and associated atmospheric processes over Dronning Maud Land, East Antarctica. *Journal of Geophysical Research:*  
744 *Atmospheres*, 129(7), e2023JD038993. <https://doi.org/10.1029/2023JD038993>

745 Simpson, I. R., Rosenbloom, N., Danabasoglu, G., Deser, C., Yeager, S. G., McCluskey, C. S., ... & Rodgers, K. B. (2023).  
746 The CESM2 single-forcing large ensemble and comparison to CESM1: Implications for experimental design. *Journal of*  
747 *Climate*, 36(17), 5687-5711. <https://doi.org/10.1175/JCLI-D-22-0666.1>



748 Stenni, B., Curran, M. A., Abram, N. J., Orsi, A., Goursaud, S., Masson-Delmotte, V., ... & Frezzotti, M. (2017). Antarctic  
 749 climate variability on regional and continental scales over the last 2000 years. *Climate of the Past*, 13(11), 1609-1634.  
 750 <https://doi.org/10.5194/cp-13-1609-2017>

751 Sun, Q., Zhang, X., Zwiers, F., Westra, S., & Alexander, L. V. (2021). A global, continental, and regional analysis of changes  
 752 in extreme precipitation. *Journal of Climate*, 34(1), 243-258. <https://doi.org/10.1175/JCLI-D-19-0892.1>

753 Swart, N. C., Gille, S. T., Fyfe, J. C., & Gillett, N. P. (2018). Recent Southern Ocean warming and freshening driven by  
 754 greenhouse gas emissions and ozone depletion. *Nature Geoscience*, 11(11), 836-841.  
 755 <https://doi.org/10.1038/s41561-018-0226-1>

756 Swetha Chittella, S. P., Deb, P., & Melchior van Wessem, J. (2022). Relative contribution of atmospheric drivers to  
 757 “extreme” snowfall over the Amundsen Sea Embayment. *Geophysical Research Letters*, 49(16), e2022GL098661.  
 758 <https://doi.org/10.1029/2022GL098661>

759 Tewari, K., Mishra, S. K., Salunke, P., & Dewan, A. (2022). Future projections of temperature and precipitation for  
 760 Antarctica. *Environmental Research Letters*, 17(1), 014029. <https://doi.org/10.1088/1748-9326/ac43e2>

761 The Community Earth System Model Version 2 (CESM2), G. Danabasoglu et al., 2020,  
 762 <https://doi.org/10.1029/2022GL098661>

763 Thompson, D. W., & Solomon, S. (2002). Interpretation of recent Southern Hemisphere climate change. *Science*, 296(5569),  
 764 895-899. DOI: [10.1126/science.1069270](https://doi.org/10.1126/science.1069270)

765 Turner, J., Phillips, T., Thamban, M., Rahaman, W., Marshall, G. J., Wille, J. D., ... & Lachlan-Cope, T. (2019). The  
 766 dominant role of extreme precipitation events in Antarctic snowfall variability. *Geophysical Research Letters*, 46(6),  
 767 3502-3511. <https://doi.org/10.1029/2018gl081517>

768 Trusel, L. D., Frey, K. E., Das, S. B., Karnauskas, K. B., Kuipers Munneke, P., Van Meijgaard, E., & Van Den Broeke, M. R.  
 769 (2015). Divergent trajectories of Antarctic surface melt under two twenty-first-century climate scenarios. *Nature Geoscience*,  
 770 8(12), 927-932. <https://doi.org/10.1038/ngeo2563>

771 van Wessem, J. M., van den Broeke, M. R., Wouters, B., & Lhermitte, S. (2023). Variable temperature thresholds of melt  
 772 pond formation on Antarctic ice shelves. *Nature Climate Change*, 13(2), 161-166.  
 773 <https://doi.org/10.1038/s41558-022-01577-1>

774 Vignon, É., Roussel, M. L., Gorodetskaya, I. V., Genthon, C., & Berne, A. (2021). Present and future of rainfall in Antarctica.  
775 *Geophysical Research Letters*, 48(8), e2020GL092281. <https://doi.org/10.1029/2020GL092281>

776 Wang, S., Alexander, P. M., Alley, R. B., Huang, Z., Parizek, B. R., Willet, A. G., & Anandakrishnan, S. (2025). Recent  
777 variability in fracture characteristics and ice flow of Thwaites Ice Shelf, West Antarctica. *Journal of Geophysical Research:*  
778 *Earth Surface*, 130(5), e2024JF008118. <https://doi.org/10.1029/2024JF008118>

779 Wille, J. D., Favier, V., Gorodetskaya, I. V., Agosta, C., Kittel, C., Beeman, J. C., ... & Codron, F. (2021). Antarctic  
780 atmospheric river climatology and precipitation impacts. *Journal of Geophysical Research: Atmospheres*, 126(8),  
781 e2020JD033788. <https://doi.org/10.1029/2020JD033788>

782 Wille, J. D., Favier, V., Jourdain, N. C., Kittel, C., Turton, J. V., Agosta, C., ... & Berchet, A. (2022). Intense atmospheric  
783 rivers can weaken ice shelf stability at the Antarctic Peninsula. *Communications Earth & Environment*, 3(1), 90.  
784 <https://doi.org/10.1038/s43247-022-00422-9>

785 Wille, J. D., Alexander, S. P., Amory, C., Baiman, R., Barthélemy, L., Bergstrom, D. M., ... & Zou, X. (2024). The  
786 extraordinary March 2022 East Antarctica “heat” wave. Part II: Impacts on the Antarctic ice sheet. *Journal of Climate*, 37(3),  
787 779-799. <https://doi.org/10.1175/JCLI-D-23-0176.1>

788 Wille, J. D., Favier, V., Gorodetskaya, I. V., Agosta, C., Baiman, R., Barrett, J. E., ... & Zhang, Z. (2025). Atmospheric rivers  
789 in Antarctica. *Nature Reviews Earth & Environment*, 6(3), 178-192. <https://doi.org/10.1038/s43017-024-00638-7>

790 Yu, L., Yang, Q., Vihma, T., Jagovkina, S., Liu, J., Sun, Q., & Li, Y. (2018). Features of extreme precipitation at Progress  
791 Station, Antarctica. *Journal of Climate*, 31(22), 9087-9105. <https://doi.org/10.1175/JCLI-D-18-0128.1>

792 Yu, L., Zhong, S., Jagovkina, S., Sui, C., & Sun, B. (2025). Interannual Variability and Trends in Extreme Precipitation in  
793 Dronning Maud Land, East Antarctica. *Remote Sensing*, 17(2), 324. <https://doi.org/10.3390/rs17020324>

794 Zhang, X., Wan, H., Zwiers, F. W., Hegerl, G. C., & Min, S. K. (2013). Attributing intensification of precipitation extremes  
795 to human influence. *Geophysical research letters*, 40(19), 5252-5257. <https://doi.org/10.1002/grl.51010>

796 Zhu, Z., Liu, J., Song, M., & Hu, Y. (2023). Changes in extreme temperature and precipitation over the southern extratropical  
797 continents in response to Antarctic Sea ice loss. *Journal of Climate*, 36(14), 4755-4775.  
798 <https://doi.org/10.1175/JCLI-D-22-0577.1>

799 Zwally, H. J., Robbins, J. W., & Luthcke, S. B. (2017). Mass balance of east Antarctic ice sheet: Reconciling icesat altimetry  
800 with grace gravimetry and long-term ice history. In *Agu fall meeting abstracts* (Vol. 2017, p. C23D-08). DOI:  
801 <https://doi.org/10.1017/jog.2021.8>