

Measurement report: Three-year characteristics of sulfuric acid in urban Beijing and derivation of daytime sulfuric acid proxies applicable to inland sites

Yishuo Guo^{1,2}, Chao Yan^{2,3*}, Chang Li², Chenjuan Deng⁴, Ying Zhang^{2,3}, Ying Zhou², Haotian Zheng⁴, Yueqi Jiang⁴, Xin Chen², Wei Ma², Nina Sarnela⁵, Zhuohui Lin², Chenjie Hua², Xiaolong Fan⁶, Feixue Zheng², Zemin Feng², Zongcheng Wang², Yusheng Zhang², Jingkun Jiang⁴, Bin Zhao⁴, Markku Kulmala^{2,3,5}, Yongchun Liu^{2*}

Affiliations:

¹ College of Environmental and Chemical Engineering, Hebei Vocational University of Industry and Technology, Shijiazhuang, Hebei, China

² Aerosol and Haze Laboratory, Beijing Advanced Innovation Center for Soft Matter Science and Engineering, Beijing University of Chemical Technology, Beijing, China

³ Nanjing-Helsinki Institute in Atmospheric and Earth System Sciences, Nanjing University, Suzhou, China.

⁴ State Key Joint Laboratory of Environment Simulation and Pollution Control, State Environmental Protection Key Laboratory of Sources and Control of Air Pollution Complex, School of Environment, Tsinghua University, Beijing 100084, China

⁵ Institute for Atmospheric and Earth System Research / Physics, Faculty of Science, University of Helsinki, Finland

⁶ Center for Excellence in Regional Atmospheric Environment, Key Lab of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, China

*Correspondence to: Chao Yan (chaoyan@nju.edu.cn), Yongchun Liu (liuyc@buct.edu.cn)

Abstract Sulfuric acid (H_2SO_4) is a key precursor in atmospheric new particle formation and cluster early growth. However, long-term measurement of it is only available at a few sites. Although several proxies for estimating H_2SO_4 concentration have been proposed, they are always site-specific. Therefore, both reliable H_2SO_4 measurement and proxies with wider application are highly needed. Here, we conducted a long-term H_2SO_4 measurement in urban Beijing during 2019–2021, and derived three H_2SO_4 proxies based entirely on its formation and loss pathways. Results show that daytime H_2SO_4 concentration is $2.0\text{--}7.4 \times 10^6 \text{ molec cm}^{-3}$ and shows an overall decline with an average annual decrease of 14%. This decline is mainly due to the ongoing SO_2 emission controls. Daytime H_2SO_4 shows a clear seasonal variation that tracks UVB. Nighttime H_2SO_4 concentration is $1.6\text{--}6.3 \times 10^5 \text{ molec cm}^{-3}$, with higher levels in warmer seasons due to stronger sources and lower condensation sink (CS). The diurnal variations of H_2SO_4 across seasons follow those of photo-oxidation-related parameters, such as UVB, OH radical, and photolysis rate of NO_2 ($J(\text{NO}_2)$). All of the three proxies can reproduce H_2SO_4 concentration during 10:00–14:00. Importantly, they can estimate H_2SO_4 concentration at a boreal forest site in Hyytiälä, Finland, suggesting their applicability to sites with diverse environments. Furthermore, the parameters used in UVB- $\text{PM}_{2.5}$ based proxy are available at most observational sites. Further application of this proxy could provide H_2SO_4 concentrations covering many regions worldwide, which may further facilitate research on atmospheric nucleation and secondary aerosol growth of these sites.

1. Introduction

38 New particle formation (NPF) is a key contributor to the born of atmospheric aerosols (Merikanto et al.,
39 2009;Gordon et al., 2017), and thus can have a great influence on global climate and human health (Stocker,
40 2014;Lelieveld et al., 2015). Among all the precursors that drive atmospheric nucleation, the initial step of NPF,
41 sulfuric acid (H_2SO_4) has been shown to be the most important one from both laboratory experiments and field
42 observations (Kulmala et al., 2006;Riipinen et al., 2007;Paasonen et al., 2009;Erupe et al., 2010;Wang et al.,
43 2011;Kirkby et al., 2011;Yu et al., 2012;Almeida et al., 2013;Kürten et al., 2014;Riccobono et al., 2014;Lehtipalo
44 et al., 2018;Yao et al., 2018;Lee et al., 2019;Myllys et al., 2019;Yan et al., 2021). The clusters formed by sulfuric
45 acid and base molecules, such as ammonia and amines, provide the primary core for further condensation of other
46 low-volatility species, promoting aerosols growth to tens of nanometers, reaching the sizes of cloud condensation
47 nuclei (CNN) and ultrafine particles. Therefore, reliable measurement of sulfuric acid is of great importance.

48 Since the 1990s, sulfuric acid measurements have been conducted in various field campaigns that covered a
49 wide range of atmospheric environments, including urban (McMurry et al., 2005;Fiedler et al., 2005;Riipinen et al.,
50 2007;Mikkonen et al., 2011;Wang et al., 2011;Yao et al., 2018;Lu et al., 2019), rural (Berresheim et al.,
51 2000;Birmili et al., 2003;Paasonen et al., 2009;Erupe et al., 2010;Mikkonen et al., 2011;Kürten et al., 2016;Yang
52 et al., 2021a;Yang et al., 2023), mountainous (Weber et al., 1996;Weber et al., 1997;Boy et al., 2008;Mikkonen et
53 al., 2011), marine (Berresheim et al., 1993;Weber et al., 1995;Weber et al., 1996;Berresheim et al., 2002;O'Dowd
54 et al., 2002), forest environments (Fiedler et al., 2005;Riipinen et al., 2007;Petäjä et al., 2009;Nieminen et al.,
55 2009;Mikkonen et al., 2011;Jokinen et al., 2012), among others (Weber et al., 1998;Mauldin et al., 2001;Mauldin
56 et al., 2004;Sarnela et al., 2015;Jokinen et al., 2018). The locations of these sites, measurement periods, and
57 corresponding sulfuric acid concentrations are summarized in Table 1. In general, within the planetary boundary
58 layer, sulfuric acid concentration was around $0.2 - 15 \times 10^6 \text{ cm}^{-3}$, with the highest levels in urban areas, followed
59 by rural, mountainous, and marine regions, and the lowest in forest areas. This suggests that sulfuric acid levels
60 depend strongly on the intensity of human activity. In addition, most measurement campaigns lasted for less than
61 four months and concentrated mostly on warmer seasons (spring, summer and early autumn) when NPF usually
62 occurs (Dal Maso et al., 2005;Manninen et al., 2009;Dada et al., 2017;Nieminen et al., 2018;Chu et al., 2019;Qi et
63 al., 2015). Previous studies showed that NPF in Chinese megacities was also frequently observed in winter (Deng
64 et al., 2020;Chu et al., 2019), and thus sulfuric acid measurement in cold seasons is also crucial. To date, however,
65 long-term measurement of sulfuric acid worldwide is still lacking, which somewhat limits the investigation of NPF
66 processes.

67 To complement the limited sulfuric acid measurement, several sulfuric acid proxies were developed. The original
68 expression for estimating sulfuric acid concentration was derived from its production and loss pathways. Assuming
69 that sulfuric acid originates solely from OH-initiated oxidation of SO_2 and the only loss is the condensation sink
70 (CS) onto particle surfaces, the steady-state concentration of sulfuric acid can be expressed as $[\text{H}_2\text{SO}_4] =$
71 $k \cdot [\text{OH}][\text{SO}_2]/\text{CS}$, where k is the rate constant of $\text{OH} + \text{SO}_2$ reaction. As early as 1997, Weber et al. estimated sulfuric
72 acid concentrations at a marine site and a mountain site using this expression (Weber et al., 1996). Results showed

73 that the estimated daytime sulfuric acid for two selected days generally matched the measured one at both sites.
74 Later, Berresheim et al. also utilized this expression at a coastal site (Berresheim et al., 2002). However, when the
75 accommodation coefficient of CS calculation was chosen as 1, the estimated sulfuric acid concentration turned out
76 **Table 1.** Summary of atmospheric sites with sulfuric acid measurement.

Type of Site	Location	Measurement Period	H ₂ SO ₄ (× 10 ⁶ molec cm ⁻³)	References
Urban	Atlanta, Georgia, USA	2002/08	2.9	2005, McMurry et al. 2011, Mikkonen et al.
	Heidelberg, Germany	2004/03 - 04	3.0	2005, Fiedler, et al. 2007, Riipinen et al.
	Beijing, China	2008/07 - 09	1.0 - 9.0	2011, Wang et al.
	Shanghai, China	2014/03 - 2016/02	7.8	2018, Yao et al.
	Beijing, China	2018/02 - 03	4.9	2019, Lu, et al.
Rural	Hohenpeissenberg, Germany	151 days in 1998 186 days in 1999	0.1 - 10	2000, Berresheim et al.
		1998/04 - 2000/07	0.6	2003, Birmili et al. 2009, Paasonen et al. 2011, Mikkonen et al.
	Ohio, USA	2008/08 - 2009/11	winter: 0.6 spring: 5.2 summer: 2.9 autumn: 0.5	2010, Erupe et al.
	San Pietro Capofiume, Italy	2009/06 - 07	2.4	2011, Mikkonen et al.
	Melpitz, Germany	2008/05	2.9	2011, Mikkonen et al.
	Vielbrunn, Germany	2014/05 - 06	3.0	2016, Kürten et al.
	Nanjing, China	2017/12 - 2018/01	winter: 1.9	2021, Yang et al.
		2018/04	spring: 7.4	
		2018/07 - 08	summer: 4.5	
		2018/11	autumn: 9.0	
	Xiamen, China	2022/07 - 08	2.3	2023, Yang et al.
Mountain	Colorado, USA	1993/09	0.1 - 10	1996, Weber et al.
		2006/06 - 07	2.8	1997, Weber et al. 2008, Boy et al.
		2007/06 - 07	1.4	2011, Mikkonen et al.
Marine	Washington, USA	1991/04	0.03 - 32.0	1993, Berresheim et al.
	Hawaii, USA	1992/07	upslope: 1.2 downslope: 0.5	1995, Weber et al. 1996, Weber et al.
		1999/06	1.5	2002, Berresheim et al.
	Mace Head, Ireland	1998/09	2.0 - 15.0	2002, O'Wowd et al.
		1999/06		
Forest	Hyytiälä, Finland	2003/03 - 04	2.6	2005, Fiedler et al.
		2005/04 - 05	1.0 - 10	2007, Riipinen et al.
		2007/03 - 06	0.9 - 2.5	2009, Petäjä et al. 2009, Nieminen et al.
		2003/03 - 04	0.6	2011, Mikkonen et al.
		2007/03 - 06	0.2	2011, Mikkonen et al.
		2011/03 - 04	0.3 - 10	2012, Jokinen et al.
Others	Macquarie Island, Australia	11/27/1995	2.8	1998, Weber et al.
	Antarctica	1998/12	0.3	2001, Mauldin III et al.
		2000/11 - 12	0.3	2004, Mauldin III et al.
		2014/11 - 2015/01	0.2 - 10	2018, Jokinen et al.
	Kilpilahti, Finland	2012/06 - 07	oil refinery: 11.5 industrial: 4.4	2015, Sarnela et al.
			non-industrial: 1.3	

to be much lower than the measured one. The authors speculated that additional sulfuric acid sources might exist, likely the OH- or BrO-initiated oxidation of dimethyl disulfide or dimethyl sulfide, or the oxidation of SO₂ by non-OH oxidants. This also suggests that this proxy may not be suitable for coastal environments. In 2009, Petäjä et al. proposed the concept of sulfuric acid proxy clearly and derived three proxies based on its source-sink equilibrium (Petäjä et al., 2009). The first proxy, $P_1 = k_1 \cdot [\text{OH}][\text{SO}_2]/\text{CS}$, was very similar to that proposed by Weber et al., but the pre-factor k_1 was obtained by the fitting of measurement data. During daytime, OH radical mainly arises from photochemical reactions. Therefore, OH radical in P_1 could be replaced by UVB, yielding the second proxy of $P_2 = k_2 \cdot [\text{UVB}][\text{SO}_2]/\text{CS}$. Similarly, replacing OH radical with global radiation yielded the third proxy, $P_3 = k_3 \cdot [\text{Glob}][\text{SO}_2]/\text{CS}$. These three proxies showed good performance in estimating daytime sulfuric acid concentration. However, the authors noted that k_1 , k_2 and k_3 came from fits so that they are likely site-specific, which limits their transferability to other sites.

Later, Mikkonen et al. attempted to develop sulfuric acid proxies suitable for various environments. The datasets came from five sites, including one forest, one mountainous, two rural and one urban sites (Mikkonen et al., 2011). Five linear-fitting proxies, including one (Eq. 1) similar to the P_3 proxy proposed by Petäjä et al., were first built.

$$L1 = B \cdot k \cdot \text{Radiation} \cdot [\text{SO}_2] \cdot \text{CS}^{-1} \quad (1)$$

Results showed only minor differences among the five proxies, with L3 proxy (Eq. 2) generally performing the best.

$$L3 = B \cdot k \cdot \text{Radiation} \cdot [\text{SO}_2]^{0.5} \quad (2)$$

Based on this, the authors concluded that the pseudo-steady-state assumption for gaseous sulfuric acid could be somewhat unrealistic in atmospheric conditions, and then proposed five additional nonlinear-fitting sulfuric acid proxies. However, only correlation coefficients were used to evaluate the performance of five linear fitting proxies, and these correlation coefficients were close to each other. Thus, the above conclusion requires more data to be supported.

Based on these studies, Lu et al. developed seven nonlinear-fitting proxies to estimate daytime sulfuric acid concentration in urban Beijing (Lu et al., 2019). In proxies of N5–N7, O₃ and HONO were included to account for OH radical formation via photolysis of HONO. Results showed that seven proxies generally performed well in estimating daytime sulfuric acid concentration with similar correlation coefficients and relative errors. Nevertheless, the authors concluded that N7 proxy (Eq. 3) was the most suitable for estimating daytime sulfuric acid, as it took the CS loss pathway into account, had the lowest relative error and used the easily measured NO_x.

$$[\text{H}_2\text{SO}_4] = 0.0013 \cdot \text{UVB}^{0.13} \cdot [\text{SO}_2]^{0.40} \cdot \text{CS}^{-0.17} \cdot ([\text{O}_3]^{0.44} + [\text{NO}_x]^{0.41}) \quad (3)$$

Note that this proxy was developed for urban Beijing, and thus may not apply to other sites. A year later, Dada et al. constructed proxies based on the source-sink equilibrium of sulfuric acid at four different sites, including one boreal forest, one rural, one urban, and one megacity sites. The formation of sulfuric acid from the ozonolysis of alkenes was first considered. Results showed that P_1 (Eq. 4) and P_3 (Eq. 5) proxies with the alkene ozonolysis term could estimate both daytime and nighttime sulfuric acid well, while P_2 proxy (Eq. 6) without the alkene ozonolysis term could only estimate daytime sulfuric acid.

$$([H_2SO_4] = -\frac{CS}{2k_3} + \sqrt{\left(\frac{CS}{2k_3}\right)^2 + \frac{[SO_2]}{k_3}(k_1 \cdot GlobRad + k_2 \cdot [O_3][Alkenes])}) \quad (4)$$

$$[H_2SO_4] = \frac{k_1 \cdot GlobRad[SO_2] + k_2 \cdot [SO_2][O_3][Alkene]}{CS} \quad (5)$$

$$[H_2SO_4] = -\frac{CS}{2k_3} + \sqrt{\left(\frac{CS}{2k_3}\right)^2 + \frac{[SO_2]}{k_3} k_1 \cdot GlobRad} \quad (6)$$

Although the proxy equations were the same across sites, the parameters therein were different. Thus, these four proxies have limited application at other sites.

In this study, we characterize the interannual, seasonal, and diurnal variations of sulfuric acid in urban Beijing and derive proxies to estimate sulfuric acid concentration at various sites. Long-term measurement of sulfuric acid covering nearly three continuous years (from 1st January, 2019 to 11th November, 2021) was conducted in urban Beijing. First, the yearly and seasonal variation of sulfuric acid concentration, as well as its diurnal cycles were analyzed. Second, the performance of nine representative proxies, including seven steady-state based ones and two numerical regression ones, from previous studies at our site was investigated. Based on these analyses, three steady-state proxies were proposed according to the budget analysis of sulfuric acid, and their performance and limitations on estimating daytime sulfuric acid concentration were investigated in detail. These three proxies were then applied to estimate sulfuric acid concentration at a boreal forest site in Hyytiälä, Finland. Correlation coefficients and relative errors indicate that three proxies are able to reproduce daytime sulfuric acid well, suggesting that three proxies and parameters therein could be applicable at other atmospheric sites. Finally, a general suggestion on proxy selection with different available parameters was given.

2. Method

2.1 Measurement site

The measurements were conducted at the Aerosol and Haze Laboratory at the west campus of Beijing University of Chemical Technology (39.95° N, 116.31° E). It is a typical urban site surrounded by commercial and residential areas and three major roads (Liu et al., 2020; Yan et al., 2021; Guo et al., 2021; Yan et al., 2022). The datasets used in this study span nearly three continuous years from January 2019 to November 2022.

2.2 Measurement of sulfuric acid

Sulfuric acid was measured by a long time-of-flight chemical ionization mass specter (LTOF-CIMS, Aerodyne Research, Inc.) using nitric acid as reagent ions. The basic working principle of this instrument is described elsewhere (Jokinen et al., 2012), and the instrument configuration has been provided in our previous studies and has remained unchanged over the years. Briefly, air was drawn through a stainless-steel tube (1.6 m long, 3/4 inch in diameter). The inlet flow rate was maintained at 7.2 L min⁻¹. Additionally, a flush plate (Karsa Inc.) was installed to effectively remove water vapor in the sampled air.

Sulfuric acid concentration was quantified from the ratio of bisulfate ions (with counting rates unit in ions s⁻¹) to primary ions as follows:

$$\text{H}_2\text{SO}_4 = \frac{\text{HSO}_4^- + \text{H}_2\text{SO}_4\text{NO}_3^-}{\text{NO}_3^- + \text{HNO}_3\text{NO}_3^- + (\text{HNO}_3)_2\text{NO}_3^-} \times C \quad (7)$$

where C is the calibration coefficient, determined by direct calibration using known amounts of gaseous sulfuric acid injected into the instrument (Kürten et al., 2012). During the measurement period, the instrument ran stably. Calibration was performed every six months and after tuning. After correcting the diffusional wall loss of the sampling line (0.2129), the final calibration coefficients were $6.07 - 7.47 \times 10^9 \text{ molec cm}^{-3}$ over the three-year period.

2.3 Other ancillary measurements

Particle number concentration and size distribution was measured by a differential mobility particle sizer (DMPS, 6 – 840 nm) (Aalto et al., 2001) and a particle size distribution system (PSD, 3 nm – 10 μm) (Liu et al., 2016). The configuration of these two instruments have been described in our previous studies (Zhou et al., 2021, Yan et al., 2021). Based on these measurements, the condensation sink (CS) of sulfuric acid can be calculated from the following Eq. (8) (Kulmala et al., 2012):

$$CS = 4\pi D \int_0^{d_p^{max}} \beta_m(d'_p) d'_p N_{d'_p} dd'_p = 4\pi D \sum_{d'_p} \beta_{m,d'_p} d'_p N_{d'_p} \quad (8)$$

where D is the diffusion coefficient of sulfuric acid, d'_p is the particle diameter, $N_{d'_p}$ is the particle number concentration with diameter d'_p , and β_m represents the transition-regime correction. Size-resolved hygroscopic growth of aerosols was considered in the calculation of CS. CS values calculated from two instruments are shown in Figure S13. The datasets from PSD was chosen in priority as it measures wider size ranges. If PSD data was unavailable or not consecutive for more than 10–20 days, DMPS data was used. There were three periods during which measurements from both instruments were continuous and stable, and the CS comparison for these periods are shown in Figure S14. Compared with PSD CS values, DMPS CS values were on average $\sim 11.7\%$ lower.

Meteorological parameters were measured by a weather station (AWS310, Vaisala Inc.) located on the building rooftop. These parameters include ambient temperature, relative humidity (RH), pressure, visibility, UVB radiation, and horizontal wind speed and direction. Trace gases, including carbon monoxide (CO), sulfur dioxide (SO₂), nitrogen oxides (NO_x), and ozone (O₃), were monitored using four Thermo Environmental Instruments (models 48i, 43i-TLE, 42i, 49i, respectively). Calibrations of these instruments were performed every two weeks using standard gases of known concentrations. The mass concentration of PM_{2.5} and PM₁₀ were measured with a tapered element oscillating microbalance dichotomous ambient particulate monitor (TEOM 1405-DF, Thermo Fisher Scientific Inc, USA). The mass concentration of PM_{coarse} was obtained based on the difference between PM₁₀ and PM_{2.5}.

2.4 Modelling of OH radical, J(NO₂) and J(O¹D)

The Weather Research and Forecasting Model-Community Multiscale Air Quality (WRF-CMAQ) model was applied to simulate the concentration of OH radical, the photolysis rate of NO₂ ($J(\text{NO}_2)$) and the photolysis rate for producing excited atomic oxygen from O₃ ($J(\text{O}^1\text{D})$). Simulations covered the period from 1st January, 2019 to 19th February, 2020. The physical options in WRF (version 3.9.1) were the same as in Zheng et al. (2019a). The CMAQ

model (version 5.3.2) was coupled with the two-dimensional Volatility Basis Set (2D-VBS) (Zhao et al., 2016), where the SAPRC07 mechanism was adopted for gas-phase chemistry, and the AERO6 (Sarwar et al., 2011) was used for aerosol module. The modelling domain was the same as in Zheng et al. (2020), where the horizontal resolution was $27\text{ km} \times 27\text{ km}$ and the vertical grid had 14 layers. Default planetary boundary layer settings were used. To minimize the influence of initial conditions, simulations were spun up 5 days before the modelling period. This WRF-CMAQ model and the emission inventory have been widely applied and validated in previous studies using multiple lines of evidence, including ground-based monitoring networks and satellite retrievals (Zhao et al., 2018;Zheng et al., 2019a, b, 2023, 2024;Chang et al., 2023). Simulated concentrations of key pollutants agree well with observations in both magnitude and temporal variability (Tables S5–S7). These demonstrate that the modeling system reasonably reproduces the spatial and temporal variations of major air pollutants across China.

3. Results and discussion

3.1 Characteristics of measured sulfuric acid

3.1.1 Interannual and monthly variations of sulfuric acid

In the early morning, sulfuric acid concentration is influenced not only by the photochemical production and loss pathways, but also by additional sources, such as SO_2 oxidation on traffic-related black carbon (Yao et al., 2020). In addition, sulfuric acid from direct emission and the ozonolysis of alkenes cannot be ignored during daytime when far from noon (Yang et al., 2021a). Therefore, daytime window of 10:00–14:00 (local time) was chosen for proxy evaluation unless specified otherwise. This period also corresponds to the new particle formation time (Kulmala et al., 2007;Kulmala et al., 2013;Deng et al., 2020;Ma et al., 2021). The corresponding nighttime window was 22:00–02:00 next day (local time).

In urban Beijing, typical daytime sulfuric acid concentration ranges from 2.0×10^6 to $7.4 \times 10^6\text{ molec cm}^{-3}$ (monthly median concentration, Figure 1A and Table 2). Figure 1A and Figure S1E show that sulfuric acid concentration generally declines from 2019 to 2021, with an average annual decrease of 14%. During the three years, UVB intensity remains roughly constant (Figure 1B and Figure S1A), CS change is not significant (Figure 1D and Figure S1E), while SO_2 concentration decreases markedly (Figure 1C and Figure S1D). Thus, the yearly decline of sulfuric acid is mainly attributed to the decrease of SO_2 (by $\sim 25\%$ per year, Table S8). Figure 1A also reveals that sulfuric acid concentration has a clear seasonal variation, which is the highest in May ($4.2 - 7.4 \times 10^6\text{ molec cm}^{-3}$) and September ($5.0 - 7.2 \times 10^6\text{ molec cm}^{-3}$) and the lowest from November to February of the next year ($2.0 - 3.3 \times 10^6\text{ molec cm}^{-3}$). UVB shows the same monthly pattern as sulfuric acid, which reaches the highest from May to September and decreases to the lowest from November to February of next year, while SO_2 shows an opposite monthly trend to sulfuric acid. This indicates that the influence of UVB on sulfuric acid monthly variation outperforms that of SO_2 . Meanwhile, sulfuric acid concentration in July is much lower from May to September, likely driven by extremely low SO_2 despite a small decrease of UVB in that month. In Beijing, precipitation occurs more frequently in July and August than in May, June and September (Table S1). This reduces UVB and SO_2 in

these two months (Figure S2), and further leads to lower sulfuric acid concentration. Overall, UVB intensity and SO₂ concentration are the two key parameters determining sulfuric acid concentration.




Figure 1. Three-year (from 2019 January to 2021 November) monthly variations of (A) H₂SO₄ concentration, (B) UVB, (C) SO₂ and (D) condensation sink (CS) during daytime (10:00-14:00) when new particle formation mostly occurs. Blue triangles, green squares and orange circles represent data in 2019, 2020 and 2021, respectively. The up line, middle marker and bottom line stand for upper quartile, median and lower quartile values, respectively.

Table 2. Monthly concentration of H₂SO₄ (molec cm⁻³) during daytime (10:00-14:00) from 2019 to 2021. “NaN” means there is no data available.

Month	2019			2020			2021		
	Median	25th	75th	Median	25th	75th	Median	25th	75th
January	3.1×10 ⁶	2.6×10 ⁶	3.9×10 ⁶	3.3×10 ⁶	2.4×10 ⁶	4.6×10 ⁶	2.0×10⁶	1.4×10 ⁶	2.8×10 ⁶
February	3.1×10 ⁶	2.1×10 ⁶	4.1×10 ⁶	2.5×10 ⁶	1.1×10 ⁶	4.5×10 ⁶	2.5×10 ⁶	1.7×10 ⁶	3.4×10 ⁶
March	4.5×10 ⁶	3.5×10 ⁶	5.8×10 ⁶	NaN	NaN	NaN	2.1×10 ⁶	8.2×10 ⁵	4.5×10 ⁶
April	3.8×10 ⁶	2.4×10 ⁶	5.4×10 ⁶	4.9×10 ⁶	3.6×10 ⁶	8.1×10 ⁶	3.7×10 ⁶	2.4×10 ⁶	5.4×10 ⁶
May	4.2×10 ⁶	2.6×10 ⁶	6.1×10 ⁶	7.4×10⁶	4.7×10 ⁶	1.1×10 ⁷	5.5×10 ⁶	3.3×10 ⁶	7.6×10 ⁶
June	6.2×10 ⁶	3.7×10 ⁶	1.0×10 ⁷	5.9×10 ⁶	4.2×10 ⁶	9.4×10 ⁶	3.9×10 ⁶	2.1×10 ⁶	5.7×10 ⁶
July	4.1×10 ⁶	2.0×10 ⁶	6.6×10 ⁶	3.5×10 ⁶	1.7×10 ⁶	6.3×10 ⁶	2.6×10 ⁶	1.7×10 ⁶	4.0×10 ⁶
August	3.6×10 ⁶	1.9×10 ⁶	9.5×10 ⁶	5.3×10 ⁶	3.3×10 ⁶	8.1×10 ⁶	3.6×10 ⁶	2.2×10 ⁶	5.4×10 ⁶
September	7.2×10 ⁶	5.2×10 ⁶	9.8×10 ⁶	5.0×10 ⁶	2.6×10 ⁶	8.3×10 ⁶	NaN	NaN	NaN
October	4.5×10 ⁶	2.4×10 ⁶	6.8×10 ⁶	3.4×10 ⁶	2.4×10 ⁶	5.1×10 ⁶	3.0×10 ⁶	2.0×10 ⁶	4.5×10 ⁶
November	2.9×10 ⁶	2.1×10 ⁶	4.8×10 ⁶	2.5×10 ⁶	1.7×10 ⁶	3.6×10 ⁶	2.1×10 ⁶	6.3×10 ⁵	6.0×10 ⁶
December	2.8×10 ⁶	1.6×10 ⁶	4.6×10 ⁶	2.0×10⁶	1.6×10 ⁶	2.4×10 ⁶	NaN	NaN	NaN

Typical nighttime sulfuric acid concentration of urban Beijing ranges from 1.6×10^5 to 6.3×10^5 molec cm⁻³ (monthly median concentration, Figure S3 and Table S2), about one order of magnitude lower than that of daytime. Unlike daytime sulfuric acid, nighttime sulfuric acid concentration does not show a decreasing trend from 2019 to

2021. At night, under clean conditions, alkene ozonolysis is a major source of sulfuric acid (Guo et al., 2021); under more polluted conditions, primary emissions from vehicles or fresh plumes indicated by benzene also play an important role (Yang et al., 2021a). However, data of alkenes and benzene is unavailable in July–August 2019 and in 2020–2021, making it impossible to estimate the intensities of these nocturnal sulfuric acid sources. Thus, we are not able to give further explanation on the yearly variation of nighttime sulfuric acid. Figure S3A shows that nighttime sulfuric acid has a similar but weaker seasonal variation as that of daytime, i.e., concentration is generally higher from May to September than in other months, and concentration in July and August is significantly lower than in May, June and September. According to the data of 2019, the direct-emission source is higher from March to June (Figure S3B), and alkene-ozonolysis source is higher from March to September (Figure S3C), indicating that the sources of nighttime sulfuric acid are stronger in warmer seasons. Meanwhile, the CS level is lower from April to October (Figure S3D), resulting in lower losses of sulfuric acid. Together, these two factors lead to higher nighttime sulfuric acid concentrations during warmer seasons.

3.1.2 Diurnal variations of sulfuric acid and related parameters

The diurnal patterns of sulfuric acid across seasons are similar, starting increasing in the early morning (~ 04:00–06:00), peaking around noon (~ 11:00), and decreasing to a low level at nightfall (~19:00) (Figure 2A). The morning increase of sulfuric acid occurs earliest in summer, followed by spring, autumn and winter. Moreover, the peak width of sulfuric acid from the widest to narrowest follows the same seasonal trend. These diurnal patterns across the four seasons resemble those of UVB (Figure 2B), suggesting that radiation-driven photochemical reactions govern sulfuric acid formation. Moreover, the morning increase of sulfuric acid occurs earlier than UVB and global radiation but close to $J(\text{NO}_2)$ and OH radical (Figure S4D). This suggests that UVB and global radiation are not able to adequately represent the photochemical sources in early morning, whereas SO_2 oxidation by OH radical produced by NO_2 photolysis is a major source. HONO photolysis is another major formation pathway for OH radical (Tan et al., 2017; Tan et al., 2018; Ma et al., 2019; Yang et al., 2021b; Ma et al., 2022). HONO decreases in the morning at ~ 06:00–07:00, more than an hour after the morning increase of sulfuric acid and OH radical (Figure S4D). This suggests that sulfuric acid formation in the early morning is not likely caused by the oxidation of OH radical from HONO photolysis. The daytime peaking hour of sulfuric acid is close to $J(\text{NO}_2)$ and $J(\text{O}^1\text{D})$ (Figure S4B), indicating that sulfuric acid peaking hour is controlled by photochemical reactions related to $J(\text{NO}_2)$ and $J(\text{O}^1\text{D})$. The peak width of sulfuric acid is the widest, followed by $J(\text{NO}_2)$, global radiation, OH radical, $J(\text{O}^1\text{D})$ and UVB (Figure S4C). This implies that when using proxies with these parameters to estimate daytime sulfuric acid concentration, differences in peaking hours and peak widths may cause deviations from the measured concentration.

During the day, sulfuric acid concentration is the highest in summer, followed by spring and autumn, and then winter. This seasonal variation generally tracks UVB, except in autumn, when UVB and SO_2 are lower than spring, but daytime sulfuric acid concentration remains comparable to that in spring. In autumn, the frequency of sulfuric acid with high concentrations ($\geq 1.2 \times 10^7 \text{ molec cm}^{-3}$) is higher than spring (Figure S5), likely contributing to the overall higher level of sulfuric acid. At night, sulfuric acid concentrations are comparable across seasons, even

though SO₂ and CS levels vary. As aforementioned, additional sources such as benzene-related emissions (Yang et al., 2021a) are among the main nighttime sources of sulfuric acid, so nighttime sulfuric acid cannot be easily interpreted by proxies only including SO₂, CS and OH radical.




Figure 2. Three-year (from 2019 to 2021) diurnal variations of (A) H₂SO₄, (B) UVB, (C) SO₂ and (D) condensation sink (CS). Winter, spring, summer and autumn periods cover 15th November to 15th March of next year, 16th March to May, June to August, and September to 14th November, respectively. Lines are the mean values, and shaded areas denote the standard deviations of the data.

3.2 Performance of sulfuric acid proxies from previous study

In previous studies, several representative proxies for sulfuric acid that incorporate real physical and chemical considerations have been proposed. Before constructing the proxy for this study, we first evaluate the performance of existing proxies on estimating sulfuric acid concentration. The equations and internal parameters of these nine proxies are listed in Table 3 (Petäjä et al., 2009; Mikkonen et al., 2011; Lu et al., 2019; Dada et al., 2020). Figure 3 shows the measured sulfuric acid concentration and the proxy-estimated concentrations during daytime (10:00–14:00) in 2019. Table S3 summarizes the corresponding mean, standard deviation, median, lower quartile and upper quartile of sulfuric acid concentrations. To have a more accurate understanding on proxy performance, the correlation coefficients, power exponents, and slopes of the linear fittings between measured and estimated sulfuric acid, as well as the relative errors of estimated to measured sulfuric acid concentrations are further evaluated (Table 4). The relative error is calculated as follows (Lu et al., 2019):

$$RE = \frac{1}{n} \cdot \sum_{i=1}^n \frac{|[H_2SO_4]_{pro,i} - [H_2SO_4]_{mea,i}|}{[H_2SO_4]_{mea,i}} \quad (9)$$

Table 3. The equations and internal parameters of nine sulfuric acid proxies from literatures.

Proxy	Equation	Parameters	Reference
Proxy _{Petäjä OH-C}	$P_1 = \frac{k_1 \cdot [SO_2] \cdot [OH]}{CS}$	$k_1 = 2.2 \times 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$	(Petäjä et al., 2009)
Proxy _{Petäjä OH-F}		$k_1 = 8.6 \times 10^{-10} \times [OH]^{-0.48} \text{ cm}^3 \text{ s}^{-1}$	
Proxy _{Petäjä UVB-C}	$P_2 = \frac{k_2 \cdot [SO_2] \cdot UVB}{CS}$	$k_2 = 9.9 \times 10^{-7} \text{ m}^2 \text{ W}^{-1} \text{ s}^{-1}$	
Proxy _{Petäjä UVB-F}		$k_2 = 8.4 \times 10^{-7} \times UVB^{-0.68} \text{ m}^2 \text{ W}^{-1} \text{ s}^{-1}$	
Proxy _{Petäjä Glob-C}	$P_3 = \frac{k_3 \cdot [SO_2] \cdot Glob}{CS}$	$k_3 = 2.3 \times 10^{-9} \text{ m}^2 \text{ W}^{-1} \text{ s}^{-1}$	
Proxy _{Petäjä Glob-F}		$k_3 = 1.4 \times 10^{-7} \times Glob^{-0.70} \text{ m}^2 \text{ W}^{-1} \text{ s}^{-1}$	
Proxy _{Mikkonen et al.}	$a \cdot k \cdot \text{Radiation}^b \cdot [SO_2]^c \cdot (CS \cdot RH)^f$	$a = 8.21 \times 10^{-3}, b = 1, c = 0.62, f = -0.13$	(Mikkonen et al., 2011)
Proxy _{Lu et al.}	$k_0 \cdot UVB^a \cdot [SO_2]^b \cdot CS^c \cdot (O_3^d + NO_x^f)$	$k_0 = 0.0013, a = 0.13, b = 0.40, c = -0.17, d = 0.44, f = 0.41$	(Lu et al., 2019)
Proxy _{Lubna et al.}	$-\frac{CS}{2k_3} + \sqrt{\left(\frac{CS}{2k_3}\right)^2 + \frac{[SO_2]}{k_3}} \cdot k_1 \cdot Glob$	$k_1 = 1.0 \times 10^{-6}, k_3 = 1.6 \times 10^{-7}$	(Dada et al., 2020)

As shown in Figure 3 and Table 4, estimated sulfuric acid concentrations from Proxy_{Petäjä OH-C}, Proxy_{Petäjä OH-F} and Proxy_{Lu et al.} are closest to the measured one, with median deviations within twofold. This suggests that these three proxies provide the best estimates of sulfuric acid concentration. Estimated sulfuric acid concentrations from Proxy_{Petäjä Glob-C} and Proxy_{Dada et al.} are not too far away from the measured one, with median deviations within threefold. While estimated concentrations from other proxies differ substantially from the measured one, especially Proxy_{Petäjä UVB-F}, which underestimates sulfuric acid concentration markedly. The linear correlation coefficients of Proxy_{Petäjä OH-C}, Proxy_{Petäjä OH-F} and Proxy_{Petäjä UVB-C} proxies are closet to 1.0. Similarly, the power exponents of Proxy_{Petäjä OH-C}, Proxy_{Petäjä OH-F}, Proxy_{Petäjä UVB-C} and Proxy_{Petäjä Glob-C} are closet to 1.0. This indicates that the estimated sulfuric acid from former three proxies have the best linear correlation with the measurement. Only the slope of Proxy_{Petäjä OH-C} (0.85) is close to 1.0, suggesting that it performs the best in linear relationship. For Proxy_{Petäjä UVB-F}, Proxy_{Petäjä Glob-F}, Proxy_{Mikkonen et al.}, Proxy_{Lu et al.} and Proxy_{Dada et al.}, none of the linear correlation coefficients, power exponents or the slopes perform well, indicating that they fail to reproduce the linearity with measured sulfuric acid. The relative errors are within 50% for Proxy_{Petäjä OH-F}, Proxy_{Petäjä Glob-C} and Proxy_{Petäjä Glob-F}, and those for Proxy_{Petäjä OH-C}, Proxy_{Petäjä UVB-C}, Proxy_{Petäjä UVB-F} and Proxy_{Lu et al.} range from 57% to 91%.

Considering both linear correlation and concentration estimation accuracy, Proxy_{Petäjä OH-C} and Proxy_{Petäjä OH-F} are the two most suitable proxies for reproducing sulfuric acid concentration. Other four proxies from Petäjä et al., 2009 without the OH radical term underestimate the concentration of sulfuric acid. The reason might be that the scaling factors k_2 and k_3 were obtained by fitting measured sulfuric acid and other parameters rather than deriving from the direct relationships between UVB/global radiation and OH radical. Under this circumstance, scaling factors k_2 and k_3 are influenced by measured sulfuric acid, UVB and global radiation as well as calculated CS, which may introduce substantial uncertainties. Moreover, for both concentration estimation and linearity (R, exponent and linear slope), proxies with fitted scaling factors performed worse than those with constant scaling factors. This may

be due to the absence of linear relationships between proxies and photochemical terms (OH radical, UVB, or global radiation), since the fitted scaling factors k_1 , k_2 and k_3 all include the photochemical term (Table 3).




Figure 3. Sulfuric acid concentrations from measurement and estimated by proxies from literatures during the time window of 10:00-14:00 in 2019 (1st January to 31st December). Black value inside each box is the median concentration. “Proxy_{Petäjä OH-C}” “Proxy_{Petäjä OH-F}” “Proxy_{Petäjä UVB-C}” “Proxy_{Petäjä UVB-F}” “Proxy_{Petäjä Glob-C}” and “Proxy_{Petäjä Glob-F}” represent sulfuric acid proxies from the work of Petäjä et al., 2009, with P₁ proxy using OH radical with constant pre-factor k_1 , P₁ proxy using OH radical with fitted pre-factor k_1 , P₂ proxy using UVB with constant pre-factor k_2 , P₂ proxy using UVB with fitted pre-factor k_2 , P₃ proxy using global radiation with constant pre-factor k_3 , and P₃ proxy using global radiation with fitted pre-factor k_3 , respectively. “Proxy_{Mikonen et al.}” “Proxy_{Lu et al.}” and “Proxy_{Dada et al.}” are sulfuric acid proxies from the work of Mikkonen et al. 2011, Lu et al. 2019 and Dada et al. 2020, respectively. Gray area covers 50% to 200% of median concentration of measured sulfuric acid, and two gray lines cover 33.3% to 300% of median concentration of measured sulfuric acid.



Table 4. The correlation coefficients (R), power exponents (Exponent) and slopes (Linear Slope) of the linear fittings between measured sulfuric acid concentration and the estimated ones using proxies from literatures, the relative errors (RE) of the estimated sulfuric acid concentrations to the measured one, as well as the ratios of proxy concentrations to measured concentration using mean ($[\text{Proxy/Measured}]_{\text{mean}}$) and median ($[\text{Proxy/Measured}]_{\text{median}}$) values. All parameters are fitted with Bisquare fitting.

Proxy	R	Exponent	Linear Slope	RE (%)	$[\text{Proxy/Measured}]_{\text{mean}}$	$[\text{Proxy/Measured}]_{\text{median}}$	References
Proxy _{Petäjä OH-C}	0.97	1.17	0.85	74 %	2.09	1.18	(Petäjä et al., 2009)
Proxy _{Petäjä OH-F}	0.78	0.80	0.30	35 %	0.58	0.53	
Proxy _{Petäjä UVB-C}	0.84	1.08	0.09	57 %	0.16	0.14	
Proxy _{Petäjä UVB-F}	0.52	0.63	0.01	64 %	0.04	0.04	
Proxy _{Petäjä Glob-C}	0.59	0.86	0.16	40 %	0.43	0.43	
Proxy _{Petäjä Glob-F}	0.44	0.57	0.05	50 %	0.23	0.23	
Proxy _{Mikonen et al.}	0.46	0.49	2.40	565 %	7.98	8.70	(Mikkonen et al., 2011)
Proxy _{Lu et al.}	0.38	0.30	0.42	91 %	1.70	1.99	(Lu et al., 2019)
Proxy _{Dada et al.}	0.18	0.30	0.45	133 %	2.17	2.46	(Dada et al., 2020)

Note that the scaling factor k_1 of the OH-based proxy was obtained by replacing the left hand side of the equation with measured sulfuric acid concentration (Petäjä et al., 2009). Thus, k_1 was not derived from the chemical production pathways of sulfuric acid, and the best-fit value of k_1 may vary from site to site. This limitation restricts its applicability across a broader range of sites. Therefore, in this study, we aim to derive a proxy based entirely on the formation and loss pathways of sulfuric acid, where the parameters, related pre-factors and exponents all have chemical and physical meanings. Proxies of this kind should be applicable across different sites, since no site-dependent scaling factors or exponents are used.

3.3 Derivation of sulfuric acid proxies from its budget analysis

During daytime, the main formation pathway of sulfuric acid is the SO_2 oxidation by OH radical, followed by O_2 and H_2O addition (R1–R3) (Finlayson-Pitts and Pitts Jr., 2000):



As OH radical oxidation is the rate-limiting step, the production rate of sulfuric acid is nearly equivalent to that of HSO_3 and can be calculated as follows:

$$P_{[\text{H}_2\text{SO}_4]} = P_{[\text{HSO}_3]} = k_1 \cdot [\text{SO}_2] \cdot [\text{OH}] \quad (10)$$

Regarding sulfuric acid losses, the main loss pathway is its condensation sink onto particle surfaces (Dada et al., 2020; Guo et al., 2021; Yang et al., 2021a), which can be written as:

$$L_{[\text{H}_2\text{SO}_4]} = [\text{H}_2\text{SO}_4] \cdot \text{CS} \quad (11)$$

The production and loss rates of sulfuric acid are much faster than its net concentration change (Guo et al., 2021), so a pseudo-steady-state assumption can be applied:

$$k_1 \cdot [\text{SO}_2] \cdot [\text{OH}] \approx [\text{H}_2\text{SO}_4] \cdot \text{CS} \quad (12)$$

Then, the steady-state concentration of sulfuric acid can be estimated, which can be called as the OH-CS based proxy:

$$\text{Proxy}_{\text{OH,CS}} = [\text{H}_2\text{SO}_4] = \frac{k_{\text{SO}_2-\text{OH}} \cdot [\text{SO}_2] \cdot [\text{OH}]}{\text{CS}} \quad (13)$$

Here, $k_{\text{SO}_2-\text{OH}}$ is the rate constant of SO_2 oxidation by OH radical. It is taken as $1.3 \times 10^{-12} (\text{T}/300)^{-0.7} \text{ cm}^3 \text{ s}^{-1}$, where T is the temperature in Kelvin (Wine et al., 1984; Atkinson et al., 2004), $[\text{SO}_2]$ and $[\text{OH}]$ are concentrations of SO_2 and OH radical in molec cm^{-3} , and CS is condensation sink of sulfuric acid in s^{-1} . Compared with the proxy proposed by Petäjä et al. (2009), the pre-factor $k_{\text{SO}_2-\text{OH}}$ is not obtained by parameter fitting but is a verified reaction coefficient derived from experiments. Therefore, this proxy is chemically meaningful and has the potential to be used at various sites.

It is widely acknowledged that the OH radical is difficult to measure. Therefore, for most sites lacking OH radical measurements, the OH-CS based proxy cannot be applied. A major production pathway for OH radical is

the photolysis of NO₂ and O₃, along with radical recycling (Lu et al., 2012; Ma et al., 2022), all driven by solar radiation (Rohrer and Berresheim, 2006). Thus, UVB, a readily available parameter, can replace [OH] in Eq. (13) to derive the second proxy as follows:

$$\text{Proxy}_{\text{UVB,CS}} = \frac{k_{\text{UVB-CS}} \cdot [\text{SO}_2] \cdot \text{UVB}}{\text{CS}} \quad (14)$$

where $k_{\text{UVB-CS}}$ is the pre-factor, and [SO₂], UVB, and CS are in the units of molec cm⁻³, W m⁻² and s⁻¹, respectively. As shown in Figure S7A, OH radical and UVB has a linear correlation with R value of 0.86. The ratio of OH radical to UVB is 6.14×10^6 molec cm⁻³ W⁻¹ m². Accounting for this ratio yields $k_{\text{UVB-CS}}$ of 7.98×10^{-6} (T/300)^{-0.7} W⁻¹ m² s⁻¹. Replacing the left hand side of Eq. (14) with measured sulfuric acid concentration yields $k_{\text{UVB-CS}}$ of 7.5×10^{-6} (T/300)^{-0.7} W⁻¹ m² s⁻¹, which is close to the value derived from the OH-UVB relationship. This $k_{\text{UVB-CS}}$ is finally used as it brings less deviation between measured and estimated sulfuric acid concentrations.

Furthermore, calculating CS requires particle size distribution data, which is not always available. In this case, a surrogate parameter for CS is needed. The condensation sink of gaseous species onto particles is mainly determined by the aerosol surface area. PM_{2.5} measures the masses of particles. In principle, CS and PM_{2.5} should follow a power-law relationship with an exponent of 2/3. As expected, PM_{2.5}^{2/3} and CS are well linearly correlated (Figure S7B, R=0.92). Thus, replacing CS in Eq. (14) with PM_{2.5}^{2/3} yields the third proxy as follows:

$$\text{Proxy}_{\text{UVB,PM}_{2.5}} = \frac{k_{\text{UVB-PM}_{2.5}} \cdot [\text{SO}_2] \cdot \text{UVB}}{\text{PM}_{2.5}^{2/3}} \quad (15)$$

where $k_{\text{UVB-PM}_{2.5}}$ is the pre-factor, and [SO₂], UVB, and PM_{2.5} are in the units of molec cm⁻³, W m⁻² and µg m⁻³, respectively. The slope of CS to PM_{2.5}^{2/3} is 2.67×10^{-3} s⁻¹ µg^{-2/3} m². Then, substituting [OH] with UVB and CS with PM_{2.5}^{2/3} yields $k_{\text{UVB-PM}_{2.5}}$ of 2.99×10^{-3} µg^{2/3} W⁻¹. Replacing the left hand side of Eq. (15) with measured sulfuric acid concentration yields $k_{\text{UVB-PM}_{2.5}}$ of 2.8×10^{-3} µg^{2/3} W⁻¹, which is close to the value derived from the OH-UVB and CS-PM_{2.5} relationships and is finally used.

We summarize the three proxies incorporating the corresponding parameters as follows:

$$\text{Proxy}_{\text{OH,CS}} = (1.3 \times 10^{-12}) \times \left(\frac{T}{300}\right)^{-0.7} \times [\text{SO}_2] \times [\text{OH}] \div \text{CS} \quad (16)$$

$$\text{Proxy}_{\text{UVB,CS}} = (7.5 \times 10^{-6}) \times \left(\frac{T}{300}\right)^{-0.7} \times [\text{SO}_2] \times \text{UVB} \div \text{CS} \quad (17)$$

$$\text{Proxy}_{\text{UVB,PM}_{2.5}} = (2.8 \times 10^{-3}) \times [\text{SO}_2] \times \text{UVB} \div \text{PM}_{2.5}^{2/3} \quad (18)$$


The uncertainties of the OH-CS, UVB-CS, and UVB-PM_{2.5} based proxies, based on Eqns. (16–18), are estimated to be 41.7%, 96.1%, and 100.4%, respectively. Details are provided in Section S3.

3.4 Evaluation of different sulfuric acid proxies in this study

3.4.1 Performance of sulfuric acid proxies at Beijing Site

Figure 4 shows the overall concentrations of measured and estimated sulfuric acid from proxies. The estimated sulfuric acid concentrations from three proxies are generally in good agreement with the measured one, although the OH-CS-based proxy yields slightly lower concentration than measurement. Additionally, the concentration

386 ranges estimated by proxies are broader than the measured one. Detailed sulfuric acid concentrations, including
 387 mean, standard deviation, median, lower quartile and upper quartile values are summarized in Table S4.



388 **Figure 4.** Sulfuric acid concentrations from measurement and estimated by proxies in this study during daytime (10:00-14:00).
 389

390 The scatter plots of three proxies vs. measured sulfuric acid are shown in Figure 5. For all three proxies, the
 391 estimated sulfuric acid concentrations are well correlated with the measured one, with most data points falling on
 392 or near the 1:1 line. This suggests that the three steady-state based proxies generally perform well in estimating
 393 daytime sulfuric acid concentration. However, slight deviations between the least-square-fit lines and the 1:1 lines
 394 can be observed. To better understand these deviations, we summarize the correlation coefficients and power
 395 exponents of the fits between measured and estimated sulfuric acid concentrations, as well as the relative errors of
 396 the estimated concentrations (Table 5). The OH-CS-based proxy shows the best correlation ($R = 0.96$). The R values
 397 for the UVB-CS based proxy (0.83) and the UVB-PM_{2.5} based proxy (0.79) are also close to unity. The OH-CS
 398 based proxy has an exponent of 1.14, indicating that the relationship between proxy and measured sulfuric acid is
 399 not strictly linear, which could, to some extent, arise from the uncertainty in OH radical modelling. The exponents
 400 of UVB-CS based proxy (1.02) and UVB-PM_{2.5} based proxy (1.02) are very close to 1.0, suggesting excellent good
 401 linear relationships between proxies measured sulfuric acid. The relative errors of three proxies are all within 50%,
 402 which performs better than most proxies from previous studies (Table 4). Moreover, the ratios of proxy to measured
 403 concentrations give the same result that they are in the range of 0.72–1.22, much closer to 1.0 than most proxies
 404 from previous studies (Table 4).




Figure 5. Sulfuric acid concentrations estimated by proxies in this study vs. the measured concentration during daytime (10:00-14:00) for (A) OH-CS based proxy in 2019, (B) UVB-CS based proxy in 3 years, and (C) UVB-PM_{2.5} based proxy in 3 years. The black dashed lines are 1:1 lines, and the black lines are the distance weighted least square fits between proxy and measured sulfuric acid. Corresponding functions of the fits, correlation coefficients (R) and relative errors (RE) are shown in the legend. The triangle marker represents the binned data, where the up line, middle marker and bottom lines stand for upper quartile, median and lower quartile, respectively.


Table 5. The correlation coefficients (R) and power exponents (Exponent) of the linear fittings between measured sulfuric acid concentration and the estimated ones using proxies in this study, the relative errors (RE) of the estimated sulfuric acid concentrations to the measured one, as well as the ratios of proxy concentrations to measured concentration using mean ($[\text{Proxy}/\text{Measured}]_{\text{mean}}$) and median ($[\text{Proxy}/\text{Measured}]_{\text{median}}$) values.

Year	Parameters	R	Exponent	RE (%)	$[\text{Proxy}/\text{Measured}]_{\text{mean}}$	$[\text{Proxy}/\text{Measured}]_{\text{median}}$
2019	Proxy _{OH,CS}	0.96	1.14	46%	1.22	0.72
3-Year	Proxy _{UVB,CS}	0.83	1.02	32%	1.17	1.03
	Proxy _{UVB,PM_{2.5}}	0.79	1.02	40%	1.02	0.91

To have better understanding on the performance of sulfuric acid proxies at any given moment, the time variations of sulfuric acid concentrations from three proxies and measurement are shown in Figures S8 and S9. Generally, the OH-CS based proxy provides a good estimation on daytime sulfuric acid concentration (Figure S8). Specifically, in 2019, the concentration estimated by this proxy matches well with the measured one in January, February, March, April, August, and September. In other months of 2019, it underestimates or overestimates sulfuric acid concentration. This shows that although the OH-CS-based proxy generally performs well, sulfuric acid concentration at a given moment may deviate. Similarly, sulfuric acid concentrations estimated by UVB-CS based and UVB-PM_{2.5} based proxies generally match well with the measured one at most of the daytime, with deviations noticeable in several months over 3 years (Figures S8 and S9). These time variations are consistent with the findings in Section 3.1.2: the daily peak width of OH-CS based proxy is narrower than that of measured sulfuric acid, and the daily peak widths of UVB-CS based and UVB-PM_{2.5} based proxies are narrower than that of OH-CS based proxy. Furthermore, the OH-CS based proxy partially reproduces the formation of sulfuric acid at night and early morning, with evidence on most days of January 2019 and some days of February 2019. Although UVB-CS based and UVB-PM_{2.5} based proxies cannot estimate nighttime sulfuric acid, they provide a convenient, reliable and, more importantly, feasible way to trace the long-term daytime sulfuric acid concentration for sites without OH radicals.

Sulfuric acid concentration is estimated using OH radical, UVB, SO₂, CS and PM_{2.5}. We then use these parameters to assess how well the proxy-estimated concentrations match the measured values, and to determine the applicable parameter ranges of the proxies. Figure 6 shows that when $[\text{OH}]$ is lower than $4 \times 10^5 \text{ molec cm}^{-3}$, UVB

434 is lower than 0.10 W m^{-2} or SO_2 is lower than 0.5 ppb , all three steady-state based proxies underestimate sulfuric
 435 acid concentration. This suggests that when the OH radical, UVB, or SO_2 is low, other SO_2 oxidation pathways or
 436 additional sulfuric acid sources contribute more to sulfuric acid formation. As $[\text{OH}]$ increases, the ratio of proxy to
 437 measured sulfuric acid gradually rises above 1.0. These deviations of OH-CS based proxy may arise from
 438 uncertainties in OH radical modelling. As UVB and SO_2 increase, the ratios of proxies to measured sulfuric acid
 439 stabilize around 1.0. This suggests that although the OH-CS based proxy is derived entirely from sulfuric acid
 440 budget analysis, its long-term stability may not be as good as that of UVB-CS based or UVB- $\text{PM}_{2.5}$ based proxies,
 441 given the intrinsic uncertainty in OH modeling. The ratio of UVB-CS based proxy stays around 1.0 when CS is
 442 lower than 0.07 s^{-1} , accounting for $\sim 96.3\%$ of total data. Similarly, the ratio of UVB- $\text{PM}_{2.5}$ based proxy shows no
 443 clear dependence on $\text{PM}_{2.5}$ when it is lower than $200 \mu\text{g m}^{-3}$, accounting for $\sim 99.6\%$ of all datasets. This indicates
 444 that these two proxies can be applied across almost all CS and $\text{PM}_{2.5}$ ranges. For OH-CS based proxy, sulfuric acid
 445 concentration is underestimated when CS is lower than 0.015 s^{-1} ($\sim 32.2\%$) or higher than 0.07 s^{-1} ($\sim 3.6\%$). Higher
 446 CS is also associated with more polluted conditions when other sulfuric acid sources such as primary emissions
 447 may exist (Yang et al., 2021a). At lower CS, UVB-CS based proxy performs well, while OH-CS based proxy does
 448 not, suggesting that slightly poor performance of OH-CS based proxy may arise from OH radical modelling.
 449 Meanwhile, the performances of three steady-state based proxies show a clear dependence on RH. When RH is
 450 lower than 60% , the ratios of proxies to measured sulfuric acid stabilize around 1.0. When RH exceeds 60% ($\sim 13.6\%$
 451 of total data), these ratios increase with RH. Higher RH correlates with precipitation events with lower UVB and
 452 lower SO_2 , increasing the contribution of additional sulfuric acid sources. This may partly explain the
 453 underestimation of proxies at higher RH.






Figure 6. Left: The ratios of sulfuric acid concentrations estimated by proxies in this study to the measured one (Proxy/Measured) vs. concentration of OH radical ($[\text{OH}]$), UVB, SO_2 , CS, $\text{PM}_{2.5}$ and RH during daytime (10:00–14:00) of 2019. Different colored markers represent different proxies. The up line, middle marker and bottom line stand for upper quartile, median and lower quartile values respectively. Right: Frequency distributions of corresponding parameters classified by “NPF”, “Non-NPF”, “Undefined”, and “No Data” periods.

Sulfuric acid is a key precursor in NPF processes. Therefore, it is necessary to assess how well these proxies perform during NPF periods. As shown in Figures 6 and S10, about 30% of NPF cases fall outside the optimal range of SO_2 , while most NPF cases fall within the optimal ranges of OH radical, UVB, CS, $\text{PM}_{2.5}$, and RH. Consequently, during NPF periods, the performance of three proxies mainly depends on the SO_2 concentration at that time. As shown in Figure S11, restricting the analysis to data within the optimal parameter ranges reduces the number of data points that deviate from the 1:1 line and have extremely low estimated sulfuric acid concentrations. Meanwhile, the correlation coefficients between the estimated and measured sulfuric acid concentrations generally improved, while the relative errors increased, and the improvement in the slopes of linear fits was not significant. This suggests that data outside the optimal parameter ranges generally have little impact on the fitting results.

3.4.2 Performance of sulfuric acid proxies at Hyytiälä, Finland

Because the three proxies above are derived from the budget analysis of sulfuric acid, Eqns. (16–18) and their pre-factors should be applicable to other sites. To demonstrate this, we use datasets from a boreal forest site in Hyytiälä, Finland as test data. Figure 7 shows the scatter plots of UVB-CS based and UVB- $\text{PM}_{2.5}$ based proxies vs. measured sulfuric acid. For both proxies, most data points lie on or are near the 1:1 line, with R values close to 1.0, indicating good linear correlations between the estimated and measured sulfuric acid concentrations. The relative errors for UVB-CS based and UVB- $\text{PM}_{2.5}$ based proxies of Hyytiälä site are 97% and 80%, respectively, which are

only slightly larger than those of the Beijing site (Table 5) but still within an acceptable range. The above results suggest that both proxies perform well in estimating daytime sulfuric acid concentration at Hyytiälä.

For the UVB-CS based proxy, the pre-factor $k_{\text{UVB-CS}}$ in Eq. (17) was chosen the same as Beijing. This proxy estimates sulfuric acid concentrations well at both Beijing and Hyytiälä sites using the same $k_{\text{UVB-CS}}$ value, indicating that the OH-UVB relationships, or the k' values in $[\text{OH}] = k' \cdot \text{UVB}$, do not differ significantly between these two sites. This further suggests that k' values at other sites should not differ significantly, and that $k_{\text{UVB-CS}}$ values should be similar across sites.

For the UVB-PM_{2.5} based proxy, the pre-factors $k_{\text{UVB-PM}_{2.5}}$ in Eq. (18) are $2.8 \times 10^{-3} \mu\text{g}^{2/3} \text{W}^{-1}$ and $4.7 \times 10^{-3} \mu\text{g}^{2/3} \text{W}^{-1}$ for Beijing and Hyytiälä, respectively. This difference in $k_{\text{UVB-PM}_{2.5}}$ arises from the disparity of pre-factor in $\text{CS} = k \cdot \text{PM}_{2.5}^{2/3}$, where the values of k are $2.67 \times 10^{-3} \mu\text{g}^{-2/3} \text{m}^2$ and $1.59 \times 10^{-3} \mu\text{g}^{-2/3} \text{m}^2$ for Beijing (Figure S7B) and Hyytiälä (Figure S12A), respectively. Specifically, the ratios of 4.7×10^{-3} to 2.8×10^{-3} and of 2.67×10^{-3} to 1.59×10^{-3} are both 1.68. Therefore, considering the CS-PM_{2.5} relationships, Eq. (18) is also applicable to Hyytiälä. This tells us that when using the UVB-PM_{2.5} based proxy to estimate sulfuric acid concentration, the k value should be determined first to correct $k_{\text{UVB-PM}_{2.5}}$. Figures S8B-C show that the slope of CS to $\text{PM}_{2.5}^{2/3}$ is stable across years and seasons at a given site. Therefore, by conducting short-term synchronous measurement of PM_{2.5} and particle size distribution, a reliable k can be obtained. In summary, these steady-state based proxies are transferable proxies that can be widely used to estimate daytime sulfuric acid concentration at other atmospheric sites.




Figure 7. (A) UVB-CS based proxy ($\text{Proxy}_{\text{UVB,CS}}$) and (B) UVB-PM_{2.5} based proxy ($\text{Proxy}_{\text{UVB,PM}_{2.5}}$) vs. measured sulfuric acid of Hyytiälä, Finland during daytime (10:00-14:00) from 8th March to 13th Aug. 2018. The pre-factor of $\text{Proxy}_{\text{UVB,CS}}$ ($k_{\text{UVB,CS}}$) is $7.5 \times 10^{-6} (\text{T}/300)^{-0.7} \text{W}^{-1} \text{m}^2 \text{s}^{-1}$, which is the same as Beijing. The pre-factor of $\text{Proxy}_{\text{UVB,PM}_{2.5}}$ ($k_{\text{UVB,PM}_{2.5}}$) is $4.7 \times 10^{-3} \mu\text{g}^{2/3} \text{W}^{-1}$. In both two plots, the black dashed lines are 1:1 lines. Correlation coefficients (R) and the relative errors (RE) are shown in the legend.

4. Summary and Conclusions

505 In this study, long-term measurement of sulfuric acid from 2019 to 2021 was conducted in urban Beijing.
506 Daytime sulfuric acid concentration ranges from 2.0×10^6 to 7.4×10^6 molec cm^{-3} and shows a general declining trend,
507 with an average annual decrease of 14%, which is mainly due to SO_2 reduction. In addition, sulfuric acid
508 concentration shows a clear seasonal variation that tracks UVB, reaching the highest in May and September and
509 decreasing to the lowest from November to February of next year. In July and August, frequent precipitation lowers
510 UVB and SO_2 , resulting in lower sulfuric acid. Nighttime sulfuric acid concentration ranges from 1.6×10^5 to 6.3×10^5
511 molec cm^{-3} , about one order of magnitude lower than daytime. In warmer seasons, the sources of nighttime sulfuric
512 acid, such as benzene-related emissions and alkene ozonolysis, are stronger, and the losses are weaker, leading to
513 higher sulfuric acid level. The diurnal variations of photo-oxidation related parameters deviate slightly from sulfuric
514 acid. Sulfuric acid peaks earliest, followed by $\text{J}(\text{NO}_2)$, $\text{J}(\text{O}^1\text{D})$, UVB, global radiation, and OH radical. Meanwhile,
515 the peak width of sulfuric acid is the widest, followed by $\text{J}(\text{NO}_2)$, global radiation, OH radical, $\text{J}(\text{O}^1\text{D})$, and UVB.

516 The challenges in sulfuric acid measurement hinder its widespread observation. To obtain sulfuric acid proxies
517 applicable to most sites, we derive three sulfuric acid proxies directly from its steady-state budget analysis, named
518 as OH-CS based, UVB-CS based, and UVB- $\text{PM}_{2.5}$ based proxies. All three proxies perform well in estimating
519 sulfuric acid concentration during 10:00–14:00. We also evaluate the performance of nine sulfuric acid proxies
520 proposed in previous studies: seven based on formation and loss pathways (Petäjä et al., 2009; Dada et al., 2020)
521 and two derived from numerical regression (Mikkonen et al., 2011; Lu et al., 2019). Results show that $\text{Proxy}_{\text{Petäjä OH-C}}$
522 and $\text{Proxy}_{\text{Petäjä OH-F}}$ generally reproduce daytime sulfuric acid concentrations well, with estimated concentrations
523 closet to the measured one, correlation coefficients being 0.97 and 0.78, respectively, and relative errors being 74%
524 and 35%, respectively. However, the scaling factors therein are obtained by fitting the proxy equations. Thus, these
525 scaling factors are influenced by measurement reliability and have limited applicability at other sites. By contrast,
526 our proxies are derived directly from sulfuric acid budget analysis, and the parameters in the proxy equations are
527 transferable that can be used at a boreal forest site in Hyytiälä, Finland. Therefore, the three proxies developed in
528 this study have high potential for estimating daytime sulfuric acid concentrations at various sites.

529 It should be noted that the OH radical used in this study is not measured, but derived from a model simulation.
530 Under this circumstance, the OH-CS based proxy generally performs well, but has some deviations when OH radical
531 is in the range of $1.2\text{--}1.6 \times 10^7$ molec cm^{-3} and CS is lower than 0.015 s^{-1} . Although three steady-state-based proxies
532 generally perform well, they are not suitable under certain conditions. When OH radical, UVB and SO_2 are too low,
533 when CS and $\text{PM}_{2.5}$ are too high, or when RH exceeds 60%, estimated sulfuric acid concentration may deviate from
534 the actual concentration to a larger extent. Moreover, three proxies cannot fully reproduce sulfuric acid
535 concentration in early morning and at nightfall. This indicates that during these two periods, other sulfuric acid
536 sources, such as direct emission, alkenes ozonolysis and other formation pathways, are also important.

537 Here are some suggestions for the selection of three proxies. If one site has comprehensive measurement of OH
538 radical, particle size distribution and SO_2 , the OH-CS based proxy illustrated by Eq. (16) is preferred, since it
539 estimates daytime concentration well and partly captures diurnal variation and nighttime sulfuric acid. Moreover,

the pre-factor in Eq. (16) is the actual OH + SO₂ reaction rate, making it suitable to all atmospheric sites. Then, if OH radical is not directly measured, but UVB, SO₂, and particle size distribution are available, the UVB-CS based proxy illustrated by Eq. (17) is preferred. Although it cannot perfectly trace the diurnal variation of sulfuric acid, it estimates daytime concentration well. Moreover, because its pre-factor is transferable, it is convenient and straightforward to use. Finally, if neither OH radical nor particle size distribution is measured, but UVB, SO₂, and PM_{2.5} are available, the UVB-PM_{2.5} based proxy should be the right choice. These three parameters used are commonly measured, giving this proxy broad applicability. Noted that $k_{\text{UVB-PM}_{2.5}}$ in Eq. (18) varies across sites. For better accuracy, short-term synchronous measurement of particle size distribution and PM_{2.5} is suggested for obtaining the pre-factor (k) in $\text{CS} = k \cdot \text{PM}_{2.5}^{2/3}$ and then correcting $k_{\text{UVB-PM}_{2.5}}$.

The acquisition of fundamental sulfuric acid concentration datasets is of great significance for elucidating the global spatial distribution and long-term temporal trends of sulfuric acid. This may further promote researches on the mechanisms of atmospheric nucleation, cluster growth, secondary aerosol formation, and pollution event evolution at corresponding regions.

Data and materials availability: Datasets for this paper can be accessed at <https://zenodo.org/records/17216660> (Guo et al., 2025).

Author contributions:

YG, CY and YL designed the study and wrote the paper. CL, CD, YZhang, YZhou, XC, WM, NS, ZL, CH, XF, FZ, ZF, ZW, and YZ conducted the measurement and collected the data. HZ and YJ did the modelling. JJ, BZ and MK are acknowledged for valuable suggestions. And co-authors have read and commented on the paper.

Competing interests: The authors declare that they have no conflict of interests.

Acknowledgements and funds: This study is funded by the National Natural Science Foundation of China (NSFC) (grant No. 22327806), the Science and Technology Project of Hebei Education Department (grant No. QN2025049), and the Doctoral Fund of Hebei Vocational University of Industry and Technology (No. bz202402).

- 569 Aalto, P., Hämeri, K., Becker, E., Weber, R., Salm, J., Mäkelä, J. M., Hoell, C., O’ Dowd, C. D., Hansson, H.-C., Väkevä, M.,
570 Koponen, I. K., Buzorius, G., and Kulmala, M.: Physical characterization of aerosol particles during nucleation events, *Tellus*
571 *B: Chemical and Physical Meteorology*, 53, 344-358, 10.3402/tellusb.v53i4.17127, 2001.
- 572 Almeida, J., Schobesberger, S., Kuerten, A., Ortega, I. K., Kupiainen-Maatta, O., Praplan, A. P., Adamov, A., Amorim, A.,
573 Bianchi, F., Breitenlechner, M., David, A., Dommen, J., Donahue, N. M., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S.,
574 Flagan, R. C., Franchin, A., Guida, R., Hakala, J., Hansel, A., Heinritzi, M., Henschel, H., Jokinen, T., Junninen, H., Kajos,
575 M., Kangasluoma, J., Keskinen, H., Kupc, A., Kurten, T., Kvashin, A. N., Laaksonen, A., Lehtipalo, K., Leiminger, M., Leppa,
576 J., Loukonen, V., Makhmutov, V., Mathot, S., McGrath, M. J., Nieminen, T., Olenius, T., Onnela, A., Petaja, T., Riccobono,
577 F., Riipinen, I., Rissanen, M., Rondo, L., Ruuskanen, T., Santos, F. D., Sarnela, N., Schallhart, S., Schnitzhofer, R., Seinfeld,
578 J. H., Simon, M., Sipila, M., Stozhkov, Y., Stratmann, F., Tome, A., Troestl, J., Tsagkogeorgas, G., Vaattovaara, P., Viisanen,
579 Y., Virtanen, A., Vrtala, A., Wagner, P. E., Weingartner, E., Wex, H., Williamson, C., Wimmer, D., Ye, P., Yli-Juuti, T.,
580 Carslaw, K. S., Kulmala, M., Curtius, J., Baltensperger, U., Worsnop, D. R., Vehkamäki, H., and Kirkby, J.: Molecular
581 understanding of sulphuric acid-amine particle nucleation in the atmosphere, *Nature*, 502, 359-363, 10.1038/nature12663, 2013.
- 582 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe,
583 J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x, HO_x, NO_x and
584 SO_x species, *Atmospheric Chemistry and Physics*, 4, 1461-1738, 10.5194/acp-4-1461-2004, 2004.
- 585 Berresheim, H., Eisele, F. L., Tanner, D. J., McInnes, L. M., Ramsey-Bell, D. C., and Covert, D. S.: Atmospheric sulfur
586 chemistry and cloud condensation nuclei (CCN) concentrations over the northeastern Pacific Coast, *Journal of Geophysical*
587 *Research: Atmospheres*, 98, 12701-12711, <https://doi.org/10.1029/93JD00815>, 1993.
- 588 Berresheim, H., Elste, T., Plass-Dülmer, C., Eisele, F. L., and Tanner, D. J.: Chemical ionization mass spectrometer for long-
589 term measurements of atmospheric OH and H₂SO₄, *International Journal of Mass Spectrometry*, 202, 91-109,
590 [https://doi.org/10.1016/S1387-3806\(00\)00233-5](https://doi.org/10.1016/S1387-3806(00)00233-5), 2000.
- 591 Berresheim, H., Elste, T., Tremmel, H. G., Allen, A. G., Hansson, H. C., Rosman, K., Dal Maso, M., Mäkelä, J. M., Kulmala,
592 M., and O’Dowd, C. D.: Gas-aerosol relationships of H₂SO₄, MSA, and OH: Observations in the coastal marine boundary layer
593 at Mace Head, Ireland, *Journal of Geophysical Research: Atmospheres*, 107, PAR 5-1-PAR 5-12,
594 <https://doi.org/10.1029/2000JD000229>, 2002.
- 595 Birmili, W., Berresheim, H., Plass-Dulmer, C., Elste, T., Gilge, S., Wiedensohler, A., and Uhrner, U.: The Hohenpeissenberg
596 aerosol formation experiment (HAFEX): A long-term study including size-resolved aerosol, H₂SO₄, OH, and monoterpenes
597 measurements, *Atmospheric Chemistry and Physics*, 3, 361-376, 10.5194/acp-3-361-2003, 2003.
- 598 Boy, M., Karl, T., Turnipseed, A., Mauldin, R. L., Kosciuch, E., Greenberg, J., Rathbone, J., Smith, J., Held, A., Barsanti, K.,
599 Wehner, B., Bauer, S., Wiedensohler, A., Bonn, B., Kulmala, M., and Guenther, A.: New particle formation in the front range
600 of the colorado rocky mountains, *Atmospheric Chemistry and Physics*, 8, 1577-1590, 10.5194/acp-8-1577-2008, 2008.
- 601 Chang, X., Zheng, H., Zhao, B., Yan, C., Jiang, Y., Hu, R., Song, S., Dong, Z., Li, S., Li, Z., Zhu, Y., Shi, H., Jiang, Z., Xing,
602 J., and Wang, S.: Drivers of High Concentrations of Secondary Organic Aerosols in Northern China during the COVID-19
603 Lockdowns, *Environmental Science & Technology*, 57, 5521-5531, 10.1021/acs.est.2c06914, 2023.
- 604 Chu, B., Kerminen, V. M., Bianchi, F., Yan, C., Petäjä, T., and Kulmala, M.: Atmospheric new particle formation in China,
605 *Atmospheric Chemistry and Physics*, 19, 115-138, 10.5194/acp-19-115-2019, 2019.
- 606 Dada, L., Paasonen, P., Nieminen, T., Buenrostro Mazon, S., Kontkanen, J., Peräkylä, O., Lehtipalo, K., Hussein, T., Petäjä,
607 T., Kerminen, V. M., Bäck, J., and Kulmala, M.: Long-term analysis of clear-sky new particle formation events and nonevents
608 in Hyytiälä, *Atmospheric Chemistry and Physics*, 17, 6227-6241, 10.5194/acp-17-6227-2017, 2017.
- 609 Dada, L., Yliviikka, I., Baalbaki, R., Li, C., Guo, Y., Yan, C., Yao, L., Sarnela, N., Jokinen, T., Daellenbach, K. R., Yin, R.,
610 Deng, C., Chu, B., Nieminen, T., Kontkanen, J., Stolzenburg, D., Sipilä, M., Hussein, T., Paasonen, P., Bianchi, F., Salma, I.,
611 Weidinger, T., Pikridas, M., Sciare, J., Jiang, J., Liu, Y., Petäjä, T., Kerminen, V. M., and Kulmala, M.: Sources and sinks
612 driving sulphuric acid concentrations in contrasting environments: Implications on proxy calculations, *Atmospheric Chemistry*
613 *and Physics*, 20, 11747-11766, 10.5194/acp-20-11747-2020, 2020.
- 614 Dal Maso, M., Kulmala, M., Riipinen, I., and Wagner, R.: Formation and growth of fresh atmospheric aerosols: Eight years of
615 aerosol size distribution data from SMEAR II, Hyytiälä, Finland, *Boreal Environment Research*, 10, 323-336, 2005.

616 Deng, C., Fu, Y., Dada, L., Yan, C., Cai, R., Yang, D., Zhou, Y., Yin, R., Lu, Y., Li, X., Qiao, X., Fan, X., Nie, W., Kontkanen,
617 J., Kangasluoma, J., Chu, B., Ding, A., Kerminen, V.-M., Paasonen, P., Worsnop, D. R., Bianchi, F., Liu, Y., Zheng, J., Wang,
618 L., Kulmala, M., and Jiang, J.: Seasonal characteristics of new particle formation and growth in urban Beijing, *Environmental*
619 *Science & Technology*, 54, 8547-8557, 10.1021/acs.est.0c00808, 2020.

620 Erupe, M. E., Benson, D. R., Li, J., Young, L.-H., Verheggen, B., Al-Refai, M., Tahboub, O., Cunningham, V., Frimpong, F.,
621 Viggiano, A. A., and Lee, S.-H.: Correlation of aerosol nucleation rate with sulfuric acid and ammonia in Kent, Ohio: An
622 atmospheric observation, *Journal of Geophysical Research: Atmospheres*, 115, D23216, 10.1029/2010jd013942, 2010.

623 Fiedler, V., Dal Maso, M., Boy, M., Aufmhoff, H., Hoffmann, J., Schuck, T., Birmili, W., Hanke, M., Uecker, J., Arnold, F.,
624 and Kulmala, M.: The contribution of sulphuric acid to atmospheric particle formation and growth: A comparison between
625 boundary layers in Northern and Central Europe, *Atmospheric Chemistry and Physics*, 5, 1773-1785, 10.5194/acp-5-1773-
626 2005, 2005.

627 Finlayson-Pitts, B. J., and Pitts Jr., J. N.: *Chemistry of the Upper and Lower Atmosphere: Theory, Experiments, and*
628 *Applications*, Academic Press, San Diego, CA, USA, 298–299 pp., 2000.

629 Gordon, H., Kirkby, J., Baltensperger, U., Bianchi, F., Breitenlechner, M., Curtius, J., Dias, A., Dommen, J., Donahue, N. M.,
630 Dunne, E. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Frege, C., Fuchs, C., Hansel, A., Hoyle, C. R., Kulmala, M., Kürten, A.,
631 Lehtipalo, K., Makhmutov, V., Molteni, U., Rissanen, M. P., Stozhkov, Y., Tröstl, J., Tsagkogeorgas, G., Wagner, R.,
632 Williamson, C., Wimmer, D., Winkler, P. M., Yan, C., and Carslaw, K. S.: Causes and importance of new particle formation
633 in the present-day and preindustrial atmospheres, *Journal of Geophysical Research: Atmospheres*, 122, 8739-8760,
634 <https://doi.org/10.1002/2017JD026844>, 2017.

635 Guo, Y., Yan, C., Li, C., Ma, W., Feng, Z., Zhou, Y., Lin, Z., Dada, L., Stolzenburg, D., Yin, R., Kontkanen, J., Daellenbach,
636 K. R., Kangasluoma, J., Yao, L., Chu, B., Wang, Y., Cai, R., Bianchi, F., Liu, Y., and Kulmala, M.: Formation of nighttime
637 sulfuric acid from the ozonolysis of alkenes in Beijing, *Atmospheric Chemistry and Physics*, 21, 5499-5511, 10.5194/acp-21-
638 5499-2021, 2021.

639 Jokinen, T., Sipilä, M., Junninen, H., Ehn, M., Lönn, G., Hakala, J., Petäjä, T., Mauldin III, R. L., Kulmala, M., and Worsnop,
640 D. R.: Atmospheric sulphuric acid and neutral cluster measurements using CI-API-TOF, *Atmospheric Chemistry and Physics*,
641 12, 4117-4125, 10.5194/acp-12-4117-2012, 2012.

642 Jokinen, T., Sipilä, M., Kontkanen, J., Vakkari, V., Tisler, P., Duplissy, E. M., Junninen, H., Kangasluoma, J., Manninen, H.
643 E., Petaja, T., Kulmala, M., Worsnop, D. R., Kirkby, J., Virkkula, A., and Kerminen, V. M.: Ion-induced sulfuric acid-ammonia
644 nucleation drives particle formation in coastal Antarctica, *Science Advances*, 4, eaat9744, 10.1126/sciadv.aat9744, 2018.

645 Kirkby, J., Curtius, J., Almeida, J., Dunne, E., Duplissy, J., Ehrhart, S., Franchin, A., Gagne, S., Ickes, L., Kuerten, A., Kupc,
646 A., Metzger, A., Riccobono, F., Rondo, L., Schobesberger, S., Tsagkogeorgas, G., Wimmer, D., Amorim, A., Bianchi, F.,
647 Breitenlechner, M., David, A., Dommen, J., Downard, A., Ehn, M., Flagan, R. C., Haider, S., Hansel, A., Hauser, D., Jud, W.,
648 Junninen, H., Kreissl, F., Kvashin, A., Laaksonen, A., Lehtipalo, K., Lima, J., Lovejoy, E. R., Makhmutov, V., Mathot, S.,
649 Mikkilä, J., Minginette, P., Mogo, S., Nieminen, T., Onnela, A., Pereira, P., Petaja, T., Schnitzhofer, R., Seinfeld, J. H., Sipilä,
650 M., Stozhkov, Y., Stratmann, F., Tome, A., Vanhanen, J., Viisanen, Y., Vrtala, A., Wagner, P. E., Walther, H., Weingartner,
651 E., Wex, H., Winkler, P. M., Carslaw, K. S., Worsnop, D. R., Baltensperger, U., and Kulmala, M.: Role of sulphuric acid,
652 ammonia and galactic cosmic rays in atmospheric aerosol nucleation, *Nature*, 476, 429-433, 10.1038/nature10343, 2011.

653 Kulmala, M., Lehtinen, K. E. J., and Laaksonen, A.: Cluster activation theory as an explanation of the linear dependence
654 between formation rate of 3nm particles and sulphuric acid concentration, *Atmospheric Chemistry and Physics*, 6, 787-793,
655 10.5194/acp-6-787-2006, 2006.

656 Kulmala, M., Riipinen, I., Sipilä, M., Manninen, H. E., Petäjä, T., Junninen, H., Maso, M. D., Mordas, G., Mirme, A., Vana,
657 M., Hirsikko, A., Laakso, L., Harrison, R. M., Hanson, I., Leung, C., Lehtinen, K. E. J., and Kerminen, V.-M.: Toward Direct
658 Measurement of Atmospheric Nucleation, *Science*, 318, 89-92, 10.1126/science.1144124, 2007.

659 Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M., Aalto, P. P., Junninen, H.,
660 Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen, V.-M.: Measurement of the nucleation of
661 atmospheric aerosol particles, *Nature Protocols*, 7, 1651-1667, 10.1038/nprot.2012.091, 2012.

662 Kulmala, M., Kontkanen, J., Junninen, H., Lehtipalo, K., Manninen, H. E., Nieminen, T., Petäjä, T., Sipilä, M., Schobesberger,
663 S., Rantala, P., Franchin, A., Jokinen, T., Järvinen, E., Äijälä, M., Kangasluoma, J., Hakala, J., Aalto, P. P., Paasonen, P.,
664 Mikkilä, J., Vanhanen, J., Aalto, J., Hakola, H., Makkonen, U., Ruuskanen, T., Mauldin, R. L., Duplissy, J., Vehkamäki, H.,

665 Bäck, J., Kortelainen, A., Riipinen, I., Kurtén, T., Johnston, M. V., Smith, J. N., Ehn, M., Mentel, T. F., Lehtinen, K. E. J.,
 666 Laaksonen, A., Kerminen, V.-M., and Worsnop, D. R.: Direct observations of atmospheric aerosol nucleation, *Science*, 339,
 667 943-946, 10.1126/science.1227385, 2013.
 668 Kürten, A., Rondo, L., Ehrhart, S., and Curtius, J.: Calibration of a Chemical Ionization Mass Spectrometer for the measurement
 669 of gaseous sulfuric acid, *The Journal of Physical Chemistry A*, 116, 6375-6386, 10.1021/jp212123n, 2012.
 670 Kürten, A., Jokinen, T., Simon, M., Sipila, M., Sarnela, N., Junninen, H., Adamov, A., Almeida, J., Amorim, A., Bianchi, F.,
 671 Breitenlechner, M., Dommen, J., Donahue, N. M., Duplissy, J., Ehrhart, S., Flagan, R. C., Franchin, A., Hakala, J., Hansel, A.,
 672 Heinritzi, M., Hutterli, M., Kangasluoma, J., Kirkby, J., Laaksonen, A., Lehtipalo, K., Leiminger, M., Makhmutov, V., Mathot,
 673 S., Onnela, A., Petaja, T., Praplan, A. P., Riccobono, F., Rissanen, M. P., Rondo, L., Schobesberger, S., Seinfeld, J. H., Steiner,
 674 G., Tome, A., Troestl, J., Winkler, P. M., Williamson, C., Wimmer, D., Ye, P., Baltensperger, U., Carslaw, K. S., Kulmala, M.,
 675 Worsnop, D. R., and Curtius, J.: Neutral molecular cluster formation of sulfuric acid-dimethylamine observed in real time
 676 under atmospheric conditions, *Proceedings of the National Academy of Sciences*, 111, 15019-15024,
 677 10.1073/pnas.1404853111, 2014.
 678 Kürten, A., Bergen, A., Heinritzi, M., Leiminger, M., Lorenz, V., Piel, F., Simon, M., Sitals, R., Wagner, A. C., and Curtius,
 679 J.: Observation of new particle formation and measurement of sulfuric acid, ammonia, amines and highly oxidized organic
 680 molecules at a rural site in central Germany, *Atmospheric Chemistry and Physics*, 16, 12793-12813, 10.5194/acp-16-12793-
 681 2016, 2016.
 682 Lee, S. H., Gordon, H., Yu, H., Lehtipalo, K., Haley, R., Li, Y. X., and Zhang, R. Y.: New particle formation in the atmosphere:
 683 from molecular clusters to global climate, *Journal of Geophysical Research: Atmospheres*, 124, 7098-7146,
 684 10.1029/2018jd029356, 2019.
 685 Lehtipalo, K., Yan, C., Dada, L., Bianchi, F., Xiao, M., Wagner, R., Stolzenburg, D., Ahonen, L., Amorim, A., Baccarini, A.,
 686 Bauer, P., Baumgartner, B., Bergen, A., Bernhammer, A.-K., Breitenlechner, M., Brilke, S., Buchholz, A., Mazon, S., Chen,
 687 D., and Worsnop, D.: Multicomponent new particle formation from sulfuric acid, ammonia, and biogenic vapors, *Science*
 688 *Advances*, 4, eaau5363, 10.1126/sciadv.aau5363, 2018.
 689 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D., and Pozzer, A.: The contribution of outdoor air pollution sources to
 690 premature mortality on a global scale, *Nature*, 525, 367-371, 10.1038/nature15371, 2015.
 691 Liu, J., Jiang, J., Zhang, Q., Deng, J., and Hao, J.: A spectrometer for measuring particle size distributions in the range of 3 nm
 692 to 10 μ m, *Frontiers of Environmental Science & Engineering*, 10, 63-72, 10.1007/s11783-014-0754-x, 2016.
 693 Liu, Y., Yan, C., Feng, Z., Zheng, F., Fan, X., Zhang, Y., Li, C., Zhou, Y., Lin, Z., Guo, Y., Zhang, Y., Ma, L., Zhou, W., Liu,
 694 Z., Dada, L., Dällenbach, K., Kontkanen, J., Cai, R., Chan, T., Chu, B., Du, W., Yao, L., Wang, Y., Cai, J., Kangasluoma, J.,
 695 Kokkonen, T., Kujansuu, J., Rusanen, A., Deng, C., Fu, Y., Yin, R., Li, X., Lu, Y., Liu, Y., Lian, C., Yang, D., Wang, W., Ge,
 696 M., Wang, Y., Worsnop, D. R., Junninen, H., He, H., Kerminen, V.-M., Zheng, J., Wang, L., Jiang, J., Petäjä, T., Bianchi, F.,
 697 and Kulmala, M.: Continuous and comprehensive atmospheric observations in Beijing: A station to understand the complex
 698 urban atmospheric environment, *Big Earth Data*, 4, 295-321, 10.1080/20964471.2020.1798707, 2020.
 699 Lu, K. D., Rohrer, F., Holland, F., Fuchs, H., Bohn, B., Brauers, T., Chang, C. C., Häseler, R., Hu, M., Kita, K., Kondo, Y., Li,
 700 X., Lou, S. R., Nehr, S., Shao, M., Zeng, L. M., Wahner, A., Zhang, Y. H., and Hofzumahaus, A.: Observation and modelling
 701 of OH and HO₂ concentrations in the Pearl River Delta 2006: A missing OH source in a VOC rich atmosphere, *Atmospheric*
 702 *Chemistry and Physics*, 12, 1541-1569, 10.5194/acp-12-1541-2012, 2012.
 703 Lu, Y., Yan, C., Fu, Y., Chen, Y., Liu, Y., Yang, G., Wang, Y., Bianchi, F., Chu, B., Zhou, Y., Yin, R., Baalbaki, R., Garmash,
 704 O., Deng, C., Wang, W., Liu, Y., Petaja, T., Kerminen, V.-M., Jiang, J., Kulmala, M., and Wang, L.: A proxy for atmospheric
 705 daytime gaseous sulfuric acid concentration in urban Beijing, *Atmospheric Chemistry and Physics*, 19, 1971-1983,
 706 10.5194/acp-19-1971-2019, 2019.
 707 Ma, L., Zhu, Y., Zheng, M., Sun, Y., Huang, L., Liu, X., Gao, Y., Shen, Y., Gao, H., and Yao, X.: Investigating three patterns
 708 of new particles growing to the size of cloud condensation nuclei in Beijing's urban atmosphere, *Atmos. Chem. Phys.*, 21, 183-
 709 200, 10.5194/acp-21-183-2021, 2021.
 710 Ma, X., Tan, Z., Lu, K., Yang, X., Liu, Y., Li, S., Li, X., Chen, S., Novelli, A., Cho, C., Zeng, L., Wahner, A., and Zhang, Y.:
 711 Winter photochemistry in Beijing: Observation and model simulation of OH and HO₂ radicals at an urban site, *Science of The*
 712 *Total Environment*, 685, 85-95, <https://doi.org/10.1016/j.scitotenv.2019.05.329>, 2019.

Ma, X., Tan, Z., Lu, K., Yang, X., Chen, X., Wang, H., Chen, S., Fang, X., Li, S., Li, X., Liu, J., Liu, Y., Lou, S., Qiu, W., Wang, H., Zeng, L., and Zhang, Y.: OH and HO₂ radical chemistry at a suburban site during the EXPLORE-YRD campaign in 2018, *Atmospheric Chemistry and Physics*, 22, 7005-7028, 10.5194/acp-22-7005-2022, 2022.

Manninen, H. E., Nieminen, T., Riipinen, I., Yli-Juuti, T., Gagné, S., Asmi, E., Aalto, P. P., Petäjä, T., Kerminen, V. M., and Kulmala, M.: Charged and total particle formation and growth rates during EUCAARI 2007 campaign in Hyytiälä, *Atmospheric Chemistry and Physics*, 9, 4077-4089, 10.5194/acp-9-4077-2009, 2009.

Mauldin, R. L., Eisele, F. L., Tanner, D. J., Kosciuch, E., Shetter, R., Lefer, B., Hall, S. R., Nowak, J. B., Buhr, M., Chen, G., Wang, P., and Davis, D.: Measurements of OH, H₂SO₄, and MSA at the South Pole during ISCAT, *Geophysical Research Letters*, 28, 3629-3632, 10.1029/2000gl012711, 2001.

Mauldin, R. L., Kosciuch, E., Henry, B., Eisele, F. L., Shetter, R., Lefer, B., Chen, G., Davis, D., Huey, G., and Tanner, D.: Measurements of OH, HO₂+RO₂, H₂SO₄, and MSA at the South Pole during ISCAT 2000, *Atmospheric Environment*, 38, 5423-5437, <https://doi.org/10.1016/j.atmosenv.2004.06.031>, 2004.

McMurry, P. H., Fink, M., Sakurai, H., Stolzenburg, M. R., Mauldin, R. L., Smith, J., Eisele, F., Moore, K., Sjostedt, S., Tanner, D., Huey, L. G., Nowak, J. B., Edgerton, E., and Voisin, D.: A criterion for new particle formation in the sulfur-rich Atlanta atmosphere, *Journal of Geophysical Research: Atmospheres*, 110, D22S02, 10.1029/2005jd005901, 2005.

Merikanto, J., Spracklen, D. V., Mann, G. W., Pickering, S. J., and Carslaw, K. S.: Impact of nucleation on global CCN, *Atmospheric Chemistry and Physics*, 9, 8601-8616, 10.5194/acp-9-8601-2009, 2009.

Mikkonen, S., Romakkaniemi, S., Smith, J. N., Korhonen, H., Petäjä, T., Plass-Duelmer, C., Boy, M., McMurry, P. H., Lehtinen, K. E. J., Joutsensaari, J., Hamed, A., Mauldin, R. L., Birmili, W., Spindler, G., Arnold, F., Kulmala, M., and Laaksonen, A.: A statistical proxy for sulphuric acid concentration, *Atmospheric Chemistry and Physics*, 11, 11319-11334, 10.5194/acp-11-11319-2011, 2011.

Myllys, N., Chee, S., Olenius, T., Lawler, M., and Smith, J.: Molecular-level understanding of synergistic effects in sulfuric acid-amine-ammonia mixed clusters, *The Journal of Physical Chemistry A*, 123, 2420-2425, 10.1021/acs.jpca.9b00909, 2019.

Nieminen, T., Manninen, H. E., Sihto, S. L., Yli-Juuti, T., Mauldin, R. L., Petäjä, T., Riipinen, I., Kerminen, V. M., and Kulmala, M.: Connection of sulfuric acid to atmospheric nucleation in boreal forest, *Environmental Science & Technology*, 43, 4715-4721, 10.1021/es803152j, 2009.

Nieminen, T., Kerminen, V. M., Petäjä, T., Aalto, P. P., Arshinov, M., Asmi, E., Baltensperger, U., Beddows, D. C. S., Beukes, J. P., Collins, D., Ding, A., Harrison, R. M., Henzing, B., Hooda, R., Hu, M., Hörrak, U., Kivekäs, N., Komsaare, K., Krejci, R., Kristensson, A., Laakso, L., Laaksonen, A., Leaitch, W. R., Lihavainen, H., Mihalopoulos, N., Németh, Z., Nie, W., O'Dowd, C., Salma, I., Sellegri, K., Svenningsson, B., Swietlicki, E., Tunved, P., Ulevicius, V., Vakkari, V., Vana, M., Wiedensohler, A., Wu, Z., Virtanen, A., and Kulmala, M.: Global analysis of continental boundary layer new particle formation based on long-term measurements, *Atmospheric Chemistry and Physics*, 18, 14737-14756, 10.5194/acp-18-14737-2018, 2018.

O'Dowd, C. D., Aalto, P., Hmeri, K., Kulmala, M., and Hoffmann, T.: Atmospheric particles from organic vapours, *Nature*, 416, 497-498, 10.1038/416497a, 2002.

Paasonen, P., Sihto, S.-L., Nieminen, T., Vuollekoski, H., Riipinen, I., Plass-Dulmer, C., Berresheim, H., Birmili, W., and Kulmala, M.: Connection between new particle formation and sulphuric acid at Hohenpeissenberg (Germany) including the influence of organic compounds, *Boreal Environment Research*, 14, 616-629, 2009.

Petäjä, T., Mauldin, R. L., Kosciuch, E., McGrath, J., Nieminen, T., Paasonen, P., Boy, M., Adamov, A., Kotiaho, T., and Kulmala, M.: Sulfuric acid and OH concentrations in a boreal forest site, *Atmospheric Chemistry and Physics*, 9, 7435-7448, 10.5194/acp-9-7435-2009, 2009.

Qi, X. M., Ding, A. J., Nie, W., Petäjä, T., Kerminen, V. M., Herrmann, E., Xie, Y. N., Zheng, L. F., Manninen, H., Aalto, P., Sun, J. N., Xu, Z. N., Chi, X. G., Huang, X., Boy, M., Virkkula, A., Yang, X. Q., Fu, C. B., and Kulmala, M.: Aerosol size distribution and new particle formation in the western Yangtze River Delta of China: 2 years of measurements at the SORPES station, *Atmospheric Chemistry and Physics*, 15, 12445-12464, 10.5194/acp-15-12445-2015, 2015.

Riccobono, F., Schobesberger, S., Scott, C., Dommen, J., Ortega, I., Rondo, L., Almeida, J., Amorim, A., Bianchi, F., Breitenlechner, M., David, A., Downard, A., Dunne, E., Duplissy, J., Ehrhart, S., Flagan, R., Franchin, A., Hansel, A., Junninen, H., and Baltensperger, U.: Oxidation products of biogenic emissions contribute to nucleation of atmospheric particles, *Science*, 344, 717-721, 10.1126/science.1243527, 2014.

Riipinen, I., Sihto, S. L., Kulmala, M., Arnold, F., Dal Maso, M., Birmili, W., Saarnio, K., Teinila, K., Kerminen, V. M.,
 Laaksonen, A., and Lehtinen, K. E. J.: Connections between atmospheric sulphuric acid and new particle formation during
 QUEST III-IV campaigns in Heidelberg and Hyytiälä, *Atmospheric Chemistry and Physics*, 7, 1899-1914, 10.5194/acp-7-
 1899-2007, 2007.

Rohrer, F., and Berresheim, H.: Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet
 radiation, *Nature*, 442, 184-187, 10.1038/nature04924, 2006.

Sarnela, N., Jokinen, T., Nieminen, T., Lehtipalo, K., Junninen, H., Kangasluoma, J., Hakala, J., Taipale, R., Schobesberger,
 S., Sipila, M., Larnimaa, K., Westerholm, H., Heijari, J., Kerminen, V.-M., Petaja, T., and Kulmala, M.: Sulphuric acid and
 aerosol particle production in the vicinity of an oil refinery, *Atmospheric Environment*, 119, 156-166,
 10.1016/j.atmosenv.2015.08.033, 2015.

Sarwar, G., Appel, K. W., Carlton, A. G., Mathur, R., Schere, K., Zhang, R., and Majeed, M. A.: Impact of a new condensed
 toluene mechanism on air quality model predictions in the US, *Geosci. Model Dev.*, 4, 183-193, 10.5194/gmd-4-183-2011,
 2011.

Stocker, T.: Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment report of
 the Intergovernmental Panel on Climate Change, 2014.

Tan, Z., Fuchs, H., Lu, K., Hofzumahaus, A., Bohn, B., Broch, S., Dong, H., Gomm, S., Häseler, R., He, L., Holland, F., Li,
 X., Liu, Y., Lu, S., Rohrer, F., Shao, M., Wang, B., Wang, M., Wu, Y., Zeng, L., Zhang, Y., Wahner, A., and Zhang, Y.:
 Radical chemistry at a rural site (Wangdu) in the North China Plain: Observation and model calculations of OH, HO₂ and RO₂
 radicals, *Atmospheric Chemistry and Physics*, 17, 663-690, 10.5194/acp-17-663-2017, 2017.

Tan, Z., Rohrer, F., Lu, K., Ma, X., Bohn, B., Broch, S., Dong, H., Fuchs, H., Gkatzelis, G. I., Hofzumahaus, A., Holland, F.,
 Li, X., Liu, Y., Liu, Y., Novelli, A., Shao, M., Wang, H., Wu, Y., Zeng, L., Hu, M., Kiendler-Scharr, A., Wahner, A., and
 Zhang, Y.: Wintertime photochemistry in Beijing: Observations of RO_x radical concentrations in the North China Plain during
 the BEST-ONE campaign, *Atmospheric Chemistry and Physics*, 18, 12391-12411, 10.5194/acp-18-12391-2018, 2018.

Wang, Z. B., Hu, M., Yue, D. L., Zheng, J., Zhang, R. Y., Wiedensohler, A., Wu, Z. J., Nieminen, T., and Boy, M.: Evaluation
 on the role of sulfuric acid in the mechanisms of new particle formation for Beijing case, *Atmospheric Chemistry and Physics*,
 11, 12663-12671, 10.5194/acp-11-12663-2011, 2011.

Weber, R. J., McMurry, P., H. J. J. o. t. A. S.: Measurement of expected nucleation precursor species and 3-500-nm
 diameter particles at Mauna Loa, 1995.

Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measured atmospheric new particle
 formation rates: Implications for nucleation mechanisms, *Chemical Engineering Communications*, 151, 53-64, 1996.

Weber, R. J., Marti, J. J., McMurry, P. H., Eisele, F. L., Tanner, D. J., and Jefferson, A.: Measurements of new particle
 formation and ultrafine particle growth rates at a clean continental site, *Journal of Geophysical Research: Atmospheres*, 102,
 4375-4385, 10.1029/96jd03656, 1997.

Weber, R. J., McMurry, P. H., Mauldin, L., Tanner, D. J., Eisele, F. L., Brechtel, F. J., Kreidenweis, S. M., Kok, G. L.,
 Schillawski, R. D., and Baumgardner, D.: A study of new particle formation and growth involving biogenic and trace gas
 species measured during ACE 1, *Journal of Geophysical Research: Atmospheres*, 103, 16385-16396, 10.1029/97jd02465, 1998.

Wine, P. H., Thompson, R. J., Ravishankara, A. R., Semmes, D. H., Gump, C. A., Torabi, A., and Nicovich, J. M.: *Journal of*
Physical Chemistry, 88, 2104-2109, 1984.

Yan, C., Yin, R., Lu, Y., Dada, L., Yang, D., Fu, Y., Kontkanen, J., Deng, C., Garmash, O., Ruan, J., Baalbaki, R., Schervish,
 M., Cai, R., Bloss, M., Chan, T., Chen, T., Chen, Q., Chen, X., Chen, Y., Chu, B., Dällenbach, K., Foreback, B., He, X.,
 Heikkinen, L., Jokinen, T., Junninen, H., Kangasluoma, J., Kokkonen, T., Kurppa, M., Lehtipalo, K., Li, H., Li, H., Li, X., Liu,
 Y., Ma, Q., Paasonen, P., Rantala, P., Pileci, R. E., Rusanen, A., Sarnela, N., Simonen, P., Wang, S., Wang, W., Wang, Y.,
 Xue, M., Yang, G., Yao, L., Zhou, Y., Kujansuu, J., Petäjä, T., Nie, W., Ma, Y., Ge, M., He, H., Donahue, N. M., Worsnop, D.
 R., Veli-Matti, K., Wang, L., Liu, Y., Zheng, J., Kulmala, M., Jiang, J., and Bianchi, F.: The synergistic role of sulfuric acid,
 bases, and oxidized organics governing new-particle formation in Beijing, *Geophysical Research Letters*, 48, e2020GL091944,
<https://doi.org/10.1029/2020GL091944>, 2021.

Yan, C., Shen, Y., Stolzenburg, D., Dada, L., Qi, X., Hakala, S., Sundström, A. M., Guo, Y., Lipponen, A., Kokkonen, T.,
 Kontkanen, J., Cai, R., Cai, J., Chan, T., Chen, L., Chu, B., Deng, C., Du, W., Fan, X., He, X. C., Kangasluoma, J., Kujansuu,
 J., Kurppa, M., Li, C., Li, Y., Lin, Z., Liu, Y., Liu, Y., Lu, Y., Nie, W., Pulliainen, J., Qiao, X., Wang, Y., Wen, Y., Wu, Y.,

810 Yang, G., Yao, L., Yin, R., Zhang, G., Zhang, S., Zheng, F., Zhou, Y., Arola, A., Tamminen, J., Paasonen, P., Sun, Y., Wang,
 811 L., Donahue, N. M., Liu, Y., Bianchi, F., Daellenbach, K. R., Worsnop, D. R., Kerminen, V. M., Petäjä, T., Ding, A., Jiang, J.,
 812 and Kulmala, M.: The effect of COVID-19 restrictions on atmospheric new particle formation in Beijing, *Atmos. Chem. Phys.*
 813 *Discuss.*, 2022, 1-24, 10.5194/acp-2021-1079, 2022.
 814 Yang, C., Dong, H., Chen, Y., Wang, Y., Fan, X., Tham, Y. J., Chen, G., Xu, L., Lin, Z., Li, M., Hong, Y., and Chen, J.:
 815 Machine Learning Reveals the Parameters Affecting the Gaseous Sulfuric Acid Distribution in a Coastal City: Model
 816 Construction and Interpretation, *Environmental Science & Technology Letters*, 10, 1045-1051, 10.1021/acs.estlett.3c00170,
 817 2023.
 818 Yang, L., Nie, W., Liu, Y., Xu, Z., Xiao, M., Qi, X., Li, Y., Wang, R., Zou, J., Paasonen, P., Yan, C., Xu, Z., Wang, J., Zhou,
 819 C., Yuan, J., Sun, J., Chi, X., Kerminen, V.-M., Kulmala, M., and Ding, A.: Toward building a physical proxy for gas-phase
 820 sulfuric acid concentration based on its budget analysis in polluted Yangtze River Delta, East China, *Environmental Science*
 821 *& Technology*, 55, 6665-6676, 10.1021/acs.est.1c00738, 2021a.
 822 Yang, X., Lu, K., Ma, X., Liu, Y., Wang, H., Hu, R., Li, X., Lou, S., Chen, S., Dong, H., Wang, F., Wang, Y., Zhang, G., Li,
 823 S., Yang, S., Yang, Y., Kuang, C., Tan, Z., Chen, X., Qiu, P., Zeng, L., Xie, P., and Zhang, Y.: Observations and modeling of
 824 OH and HO₂ radicals in Chengdu, China in summer 2019, *Science of The Total Environment*, 772, 144829,
 825 <https://doi.org/10.1016/j.scitotenv.2020.144829>, 2021b.
 826 Yao, L., Garmash, O., Bianchi, F., Zheng, J., Yan, C., Kontkanen, J., Junninen, H., Mazon, S. B., Ehn, M., Paasonen, P., Sipila,
 827 M., Wang, M. Y., Wang, X. K., Xiao, S., Chen, H. F., Lu, Y. Q., Zhang, B. W., Wang, D. F., Fu, Q. Y., Geng, F. H., Li, L.,
 828 Wang, H. L., Qiao, L. P., Yang, X., Chen, J. M., Kerminen, V. M., Petaja, T., Worsnop, D. R., Kulmala, M., and Wang, L.:
 829 Atmospheric new particle formation from sulfuric acid and amines in a Chinese megacity, *Science*, 361, 278-281,
 830 10.1126/science.aao4839, 2018.
 831 Yao, L., Fan, X., Yan, C., Kurtén, T., Daellenbach, K. R., Li, C., Wang, Y., Guo, Y., Dada, L., Rissanen, M. P., Cai, J., Tham,
 832 Y. J., Zha, Q., Zhang, S., Du, W., Yu, M., Zheng, F., Zhou, Y., Kontkanen, J., Chan, T., Shen, J., Kujansuu, J. T., Kangasluoma,
 833 J., Jiang, J., Wang, L., Worsnop, D. R., Petäjä, T., Kerminen, V.-M., Liu, Y., Chu, B., He, H., Kulmala, M., and Bianchi, F.:
 834 Unprecedented ambient sulfur trioxide (SO₃) detection: Possible formation mechanism and atmospheric implications,
 835 *Environmental Science & Technology Letters*, 7, 809-818, 10.1021/acs.estlett.0c00615, 2020.
 836 Yu, H., McGraw, R., and Lee, S.-H.: Effects of amines on formation of sub-3 nm particles and their subsequent growth,
 837 *Geophysical Research Letters*, 39, 10.1029/2011gl050099, 2012.
 838 Zhao, B., Wang, S., Donahue, N. M., Jathar, S. H., Huang, X., Wu, W., Hao, J., and Robinson, A. L.: Quantifying the effect of
 839 organic aerosol aging and intermediate-volatility emissions on regional-scale aerosol pollution in China, *Scientific Reports*, 6,
 840 28815, 10.1038/srep28815, 2016.
 841 Zhao, B., Zheng, H., Wang, S., Smith, K. R., Lu, X., Aunan, K., Gu, Y., Wang, Y., Ding, D., Xing, J., Fu, X., Yang, X., Liou,
 842 K.-N., and Hao, J.: Change in household fuels dominates the decrease in PM_{2.5} exposure and premature mortality in China in
 843 2005–2015, *Proceedings of the National Academy of Sciences*, 115, 12401-12406, 10.1073/pnas.1812955115, 2018.
 844 Zheng, H., Cai, S., Wang, S., Zhao, B., Chang, X., and Hao, J.: Development of a unit-based industrial emission inventory in
 845 the Beijing–Tianjin–Hebei region and resulting improvement in air quality modeling, *Atmos. Chem. Phys.*, 19, 3447-3462,
 846 10.5194/acp-19-3447-2019, 2019a.
 847 Zheng, H., Zhao, B., Wang, S., Wang, T., Ding, D., Chang, X., Liu, K., Xing, J., Dong, Z., Aunan, K., Liu, T., Wu, X., Zhang,
 848 S., and Wu, Y.: Transition in source contributions of PM_{2.5} exposure and associated premature mortality in China during 2005–
 849 2015, *Environment International*, 132, 105111, <https://doi.org/10.1016/j.envint.2019.105111>, 2019b.
 850 Zheng, H., Song, S., Sarwar, G., Gen, M., Wang, S., Ding, D., Chang, X., Zhang, S., Xing, J., Sun, Y., Ji, D., Chan, C. K., Gao,
 851 J., and McElroy, M. B.: Contribution of Particulate Nitrate Photolysis to Heterogeneous Sulfate Formation for Winter Haze in
 852 China, *Environmental Science & Technology Letters*, 7, 632-638, 10.1021/acs.estlett.0c00368, 2020.
 853 Zheng, H., Chang, X., Wang, S., Li, S., Zhao, B., Dong, Z., Ding, D., Jiang, Y., Huang, G., Huang, C., An, J., Zhou, M., Qiao,
 854 L., and Xing, J.: Sources of Organic Aerosol in China from 2005 to 2019: A Modeling Analysis, *Environmental Science &*
 855 *Technology*, 57, 5957-5966, 10.1021/acs.est.2c08315, 2023.
 856 Zheng, H., Li, S., Jiang, Y., Dong, Z., Yin, D., Zhao, B., Wu, Q., Liu, K., Zhang, S., Wu, Y., Wen, Y., Xing, J., Henneman, L.
 857 R. F., Kinney, P. L., Wang, S., and Hao, J.: Unpacking the factors contributing to changes in PM_{2.5}-associated mortality in
 858 China from 2013 to 2019, *Environment International*, 184, 108470, <https://doi.org/10.1016/j.envint.2024.108470>, 2024.

