

1 Supplement of

2 **Operational chemical weather forecasting with the ECCC online** 3 **Regional Air Quality Deterministic Prediction System version 023** 4 **(RAQDPS023) - Part 1: System description**

5 Michael D. Moran^{1*}, Verica Savic-Jovicic¹, Craig A. Stroud¹, Sylvain Ménard², Wanmin Gong¹, Junhua
6 Zhang¹, Qiong Zheng¹, Jack Chen¹, Ayodeji Akingunola¹, Alexandru Lupu¹, Konstantinos Menelaou², and
7 Rodrigo Munoz-Alpizar²

8 ¹Air Quality Research Division, Environment and Climate Change Canada (ECCC), Toronto, Ontario, Canada

9 ²Canadian Centre for Meteorological and Environmental Prediction, ECCC, Montreal, Quebec, Canada

10 *Retired

13 **S1 GEM-related meteorological components and attributes of RAQDPS023**

14 **S1.1 Dynamical core**

15 The governing equations of atmospheric dynamics used by the GDPS 8.0.0, RDPS 8.0.0, and RAQDPS023 have been
16 described by Côté et al. (1998) and Girard et al. (2014). Prognostic variables included the horizontal wind components,
17 vertical velocity component in the hybrid ζ -coordinate (Sect. 2.1), virtual temperature, geopotential, specific humidity,
18 and physics-related atmospheric tracers such as cloud condensate mixing ratio and turbulent kinetic energy (TKE).
19 GEM was configured to be hydrostatic in all three forecast systems (e.g., CMC-RDPS-8.0.0, 2021b). Time integration
20 of this dynamical core was performed using a two-time-level, iterative-implicit, three-dimensional semi-Lagrangian
21 scheme (Côté et al., 1998; Girard et al., 2014; Husain and Girard, 2017). This semi-implicit, semi-Lagrangian
22 integration is unconditionally stable for any time step, thus allowing the choice of time step to be based solely on
23 accuracy requirements (Robert, 1981; Ritchie et al., 2022). A dynamics integration time step of 450 s was used by the
24 GDPS 8.0.0 and one of 300 s was used by both the RDPS 8.0.0 and RAQDPS023. The improved trajectory calculations
25 for semi-Lagrangian advection described by Husain and Girard (2017) were employed, which involve trapezoidal
26 averaging of wind components along air-parcel trajectories and cubic interpolation at the departure positions. These
27 calculations remove numerical inconsistencies in the discretization and interpolation used for the source terms between
28 the dynamical and trajectory equations.

29 Semi-implicit, semi-Lagrangian time integration schemes are highly efficient by virtue of circumventing the stiffness
30 of the dynamic equations, but semi-Lagrangian advection schemes are not inherently shape-preserving or mass-
31 conserving (e.g., Staniforth and Côté, 1991; Hansen et al., 2011; de Grandpré et al., 2016). Shape preservation for two
32 meteorological tracers (water vapour, cloud water) was implemented in all three forecast systems using a two-step,
33 iterative locally-mass-conserving (ILMC) monotonicity correction for tracer advection (Sørensen et al., 2013). The
34 ILMC scheme is an *a posteriori* filter that detects the lack of shape preservation of a forecasted solution, including

35 negative concentrations, and corrects it by redistributing the overshoot or undershoot mass to the surrounding cells
36 while preserving the mass of the forecasted solution. By imposing shape preservation, the creation of artificial local
37 minima or maxima by the advection scheme was avoided. The Bermejo and Conde (2002) global mass fixer was then
38 applied in the three forecast systems to impose mass conservation, where the LAM boundary-flux estimation was used
39 for the two regional systems (Aranami et al., 2015). It is worth mentioning in this context that Staniforth and Côté
40 (1991) observed that semi-Lagrangian advection schemes handle sharp discontinuities (e.g., clouds) better than
41 Eulerian advection schemes and dispersion errors are localized around the discontinuity, which suggests that mass-
42 fixing adjustments will tend to be localized as well. McTaggart-Cowan et al. (2019a) noted that the use of these two
43 schemes in GEM improved the conservation of liquid water static energy and total water mixing ratio, with the result
44 that total global precipitation in the GDPS was reduced by approximately 5%. A surface pressure adjustment was also
45 made for moist-air mass conservation (McTaggart-Cowan et al., 2019a; CMC-RDPS-8.0.0, 2021b).

46 While semi-Lagrangian advection schemes have some inherent numerical diffusion (e.g., Staniforth and Côté, 1991),
47 artificial horizontal diffusion was also implemented to damp spurious numerical noise at the shortest spatial scales
48 (e.g., Mailhot et al., 2006). In all three systems a Del-4 operator was applied to momentum variables (10%) and a
49 Del-6 operator (1%) was applied to potential temperature. As well, a Del-2 operator with augmented coefficients was
50 applied to temperature and momentum variables in a sponge layer below the model lid (top six levels) (CMC-GDPS-
51 8.0.0, 2021b; CMC-RDPS-8.0.0, 2021b).

52 **S1.2 Land-surface and water-surface schemes**

53 The ISBA (Interactions between Soil–Biosphere–Atmosphere) land-surface scheme was used in all three forecast
54 systems to parameterize land surface processes and atmosphere-surface exchanges of heat and moisture (Noilhan and
55 Planton, 1989; Bélair et al., 2003a,b). ISBA takes a mosaic approach that considers subgrid “tiles” of four major
56 surface types: land; glacier; water; and sea ice. A GEM surface grid cell may be composed of up to four tiles to account
57 for the presence of these different surface types. Ten prognostic variables are considered in the ISBA scheme for each
58 land surface tile, including two surface temperature variables, three soil moisture variables, one vegetation canopy
59 variable, and four variables describing snowpack characteristics. Each prognostic equation is solved based on an
60 energy or water budget formulation with a force-restore term to moderate diurnal variations. Initial values of land
61 surface temperature and moisture, deep-soil temperature and moisture, and snow depth, snow albedo and snow density
62 were derived from analyses and then predicted using the ISBA scheme. Emissivities of bare soil and snow were derived
63 from climatology and observations, respectively.

64 The treatment of glacier surface tiles closely followed that for land surfaces and combined a surface-energy-budget
65 model with a two-layer force-restore term to predict glacier temperature profiles (McTaggart-Cowan et al., 2019a).
66 Updated ice emissivities were based on observations.

67 The treatment of water surface tiles depends on the forecast system. The GDPS 8.0.0 was coupled to an ocean model
68 whereas the RDPS 8.0.0 and RAQDPS023 obtained initial sea-surface temperatures from surface analyses and then

69 employed the one-dimensional thermodynamic mixed-layer model of Zeng and Beljaars (2005) to predict the diurnal
70 evolution of the water skin temperature and underlying mixed-layer temperature. This two-layer scheme provides a
71 more realistic description of diurnal variations in water surface temperature and hence surface fluxes over water. Note
72 that the GDPS 8.0.0 also used the mixed-layer scheme over lakes as did the RDPS 8.0.0 and RAQDPS023 (McTaggart-
73 Cowan et al., 2019a; CMC-GDPS-8.0.0, 2021b; CMC-RDPS-8.0.0, 2021b). The surface roughness length for
74 momentum over water surfaces was based on the Charnock (1955) scheme while roughness lengths over water for heat
75 and moisture came from Deacu et al. (2012).

76 Sea-ice (and frozen lake) surfaces were parameterized using a one-dimensional heat diffusion model (Semtner, 1976)
77 with a simple, three-level vertical discretization (McTaggart-Cowan et al., 2019a). Initial sea-ice thickness values were
78 derived from climatology but then evolved in time, whereas sea-ice fractions were derived from surface analyses, were
79 fixed in time, and were checked for consistency with the sea-ice thickness. A 3% lead fraction in sea ice was assumed
80 over salt water (McTaggart-Cowan et al., 2019a).

81 Other geophysical variables needed by the ISBA land-surface scheme include surface roughness length (except over
82 water), surface albedo, vegetation characteristics (vegetation type and fraction, leaf area index,), soil texture and soil
83 thermal and hydraulic coefficients, and glacier fraction. For grid cells with heterogeneous land surface cover (e.g.,
84 some bare soil/some vegetation or some land/some water), some grid-scale geophysical fields such as roughness length,
85 albedo, and emissivity were determined as weighted averages of tile-specific geophysical fields.

86 **S1.3 Surface-layer scheme**

87 Monin-Obukhov surface-layer similarity theory was used in all three forecast systems to describe turbulent surface
88 fluxes of momentum, heat, and moisture and state variable profiles in the model surface layer, which extends from the
89 Earth's surface to the first prognostic model level (located at approximately 10 m AGL for temperature and moisture
90 and 20 m AGL for winds on the staggered Charney-Phillips vertical grid: Sect. 2.1). Separate profiles were calculated
91 over each of the four surface types (tiles) and were then aggregated as a weighted average to provide a grid-cell value
92 (McTaggart-Cowan et al., 2019a). Stability functions proposed by Delage and Girard (1992) and Beljaars and Holtslag
93 (1991) were used for unstable and stable stratifications, respectively (CMC-GDPS-8.0.0, 2021b; CMC-RDPS-8.0.0,
94 2021b).

95 For stable conditions the following minimum Obukhov lengths were imposed to prevent surface-atmosphere
96 decoupling (Foken, 2006; McTaggart-Cowan et al., 2019a): 20 m over soil surfaces; 5 m over glaciers; and 10 m over
97 water surfaces and sea ice (CMC-GDPS-8.0.0, 2021a; CMC-RDPS-8.0.0, 2021a). The higher value over land surfaces
98 is consistent with a dependence on surface roughness (van de Wiel et al., 2007; Zheng et al., 2017).

99 **S1.4 Boundary-layer scheme and vertical diffusion**

100 Vertical diffusion in the planetary boundary layer (PBL) and free troposphere was described in all three forecast
101 systems by the vertical diffusion coefficient K_z , which is parameterized using a 1.5-order turbulence closure scheme

102 based on a predictive equation for subgrid-scale (SGS) turbulent kinetic energy (Benoit et al., 1989; Bélair et al., 1999).
103 A key parameter of the scheme is the gradient Richardson number Ri . Two turbulence length scales are also needed
104 by this scheme. A turbulence-regime-specific mixing length λ , which reflects the significant eddy size, is needed to
105 calculate both TKE and K_z , and a dissipation length scale λ_e is also needed to calculate TKE. The required lower
106 boundary conditions are obtained from the surface-layer parameterization (see previous subsection). K_z from GEM
107 was also needed by MACH (Sect. 3.1).

108 Note that a number of enhancements have been made to the original formulation of Benoit et al. (1989). First, λ was
109 calculated in the GDPS 8.0.0, RDPS 8.0.0, and RAQDPS023 using a blended formulation (McTaggart-Cowan and
110 Zadra, 2015) based on a local scheme for λ (Blackadar, 1962) and a non-local scheme for λ (Bougeault and Lacarrère,
111 1989). A treatment for Richardson number hysteresis across the temporal transitions from stable-to-unstable conditions
112 and unstable-to-stable conditions that uses an Ri transition zone rather than a single critical value of Ri has been
113 implemented (McTaggart-Cowan and Zadra, 2015). A 900 s relaxation time scale was also imposed on λ to dampen
114 the effects of turbulent transitions, and an upper bound of 50 m was imposed on λ_e (Bélair et al., 1999; McTaggart-
115 Cowan et al., 2019a).

116 Lastly, the enhancement of turbulent diffusion near the top of the cloudy boundary layer due to updrafts, downdrafts,
117 and related entrainment associated with boundary-layer clouds, including stratus, stratocumulus, and shallow cumulus
118 clouds, was represented by an extension to the TKE scheme with statistical SGS cloudiness called the MoisTKE
119 scheme (Bechtold and Siebesma, 1998; Mailhot and Bélair, 2000; Bélair et al., 2005; Mailhot et al., 2006). Three key
120 parameters were considered: turbulent flux of total cloud water; SGS cloud fraction; and a nondimensional flux
121 enhancement factor (Bélair et al., 2005). These cloud effects were activated only if the surface buoyancy flux was
122 positive and the cloud layer resided below 1.5 times the PBL height (McTaggart-Cowan et al., 2019a).

123 PBL height, which is also an important quantity for air quality, was diagnosed every time step for use in the Kain-
124 Fritsch deep convection scheme (Sect. S1.7) as well as the MoisTKE scheme. A bulk Richardson number (Ri_b) method
125 was used in which PBL height was determined to be the lowest momentum vertical level at which Ri_b reaches a critical
126 value of 0.25 by scanning the Ri_b profile upward from the surface (Seidel et al., 2012; McTaggart-Cowan et al., 2019a).
127 PBL height from GEM was also needed by MACH (Sect. 3.13).

128 **S1.5 Radiation scheme**

129 Solar and infrared radiation were parameterized in all three forecast systems using a correlated-k distribution technique
130 (Li and Barker, 2005). Nine prescribed longwave bands and four shortwave bands were considered. The updated
131 version of this scheme that is now used by GEM included an improved treatment of the water vapour continuum and
132 solar radiation absorption by methane (McTaggart-Cowan et al., 2019a). A monthly climatological O_3 distribution
133 constructed from several data sets was used along with three-dimensional climatologies of other trace gases (CH_4 , NO ,
134 CFC-11, CFC-12) appropriate for the year 2015 (McTaggart-Cowan et al., 2019a). A simple parameterization of

135 methane oxidation by the hydroxyl radical (OH) was also considered as a source of water vapour in the middle
136 atmosphere (McTaggart-Cowan et al., 2019a).

137 The GEM v5.1 code includes the addition of O₃ as a new prognostic tracer, which made it possible for the GDPS 8.0.0
138 to assimilate satellite O₃ measurements (CMC-GDPS-8.0.0, 2021a). In order to represent O₃ as a standalone chemical
139 constituent (i.e., no explicit reactive chemistry), a linearized approach was used to parameterize the photochemical
140 production and loss terms (McLinden et al., 2000). The use of this parameterization scheme, called LINOZ, in GEM
141 has been shown to be both robust and very fast (de Grandpré et al., 2016).

142 **S1.6 Grid-scale cloud and precipitation scheme**

143 The Sundqvist grid-scale condensation scheme (Sundqvist et al., 1989; Pudykiewicz et al., 1992) was used in the GDPS
144 8.0.0, RDPS 8.0.0, and RAQDPS023 to represent grid-scale cloud and precipitation processes. Four water phases were
145 considered: water vapour; cloud water; rain water; and snow water. One modification to the original scheme was to
146 use an updated cloud-to-precipitable water autoconversion rate (McTaggart-Cowan et al., 2019a; CMC-GDPS-8.0.0,
147 2021b; CMC-RDPS-8.0.0, 2021b). Lastly, this grid-scale cloud scheme was coupled to the three SGS cloud schemes
148 described next via the liquid and solid condensate mass detrained from convective updrafts in these cumulus
149 parameterization schemes, which were added to the grid-scale condensate mass (McTaggart-Cowan et al., 2019a).

150 **S1.7 Cumulus parameterization schemes**

151 Three separate cumulus parameterization schemes were used in the GDPS 8.0.0, RDPS 8.0.0, and RAQDPS023 to
152 represent three different modes of SGS unstable moist vertical motions, namely deep convection, shallow convection,
153 and mid-level convection. The term “deep convection” denotes convective clouds connected to the PBL with depths
154 of at least 3 km; the term “shallow convection” refers to the subset of convective clouds whose bases are located in the
155 PBL but which have updraft depths between 0.5 and 3 km that penetrate the PBL capping inversion (thus distinguishing
156 them from the boundary-layer clouds parameterized in the MoisTKE scheme described in Sect. S1.4); and the term
157 “mid-level convection” (or elevated convection) refers to convective clouds whose bases are located above the PBL.
158 All three schemes are formulated to transport momentum as well as heat and moisture vertically. Each scheme is mass-
159 flux-based, and their implementations share five important structural elements: trigger; updraft; downdraft; solver; and
160 closure (McTaggart-Cowan et al., 2019a). Calculations were performed sequentially for these three schemes, starting
161 with the deep convection scheme and finishing with the mid-level convection scheme.

162 For SGS deep convection a modified version of the Kain–Fritsch deep-convection scheme (Kain and Fritsch, 1990,
163 1992) was used, which has been designed for horizontal grid spacing approaching the “gray zone” over which deep
164 convection is partially resolved (i.e., horizontal grid spacing on the order of 1–10 km). The modifications included a
165 Lagrangian treatment of convective initiation, object-based clouds, and a vertical transfer of momentum. See
166 McTaggart-Cowan et al. (2019a) for a summary and McTaggart-Cowan et al. (2019b) for a detailed description of
167 these modifications. Note that the trigger-function parameters are sensitive to horizontal grid spacing, so different

168 values were specified for the GDPS 8.0.0 versus the RDPS 8.0.0 and RAQDPS023 (see Table 2 of McTaggart-Cowan
169 et al., 2019a).

170 SGS shallow convection was represented with a mass-flux scheme based on Bechtold et al. (2001). Simple
171 representations of cloud entrainment (Siebesma et al., 2003) and constant fractional detrainment (Siebesma and
172 Cuijpers, 1995) were used in the GEM implementation along with a simple, non-iterative closure that assumes that the
173 shallow convective clouds in a grid cell are in a state of quasi-equilibrium with boundary-layer forcing (Bechtold et
174 al., 2014). Note that downdrafts and precipitation were not parameterized by this scheme.

175 Lastly, SGS mid-level convection was treated using components of the deep convection scheme if certain conditions
176 were met (McTaggart-Cowan et al., 2019a). One condition was that a minimum upward grid-scale mass flux was
177 present: the required threshold was higher for the GDPS 8.0.0 than the RDPS 8.0.0 / RAQDPS023 ($1 \times 10^7 \text{ kg s}^{-1}$ vs.
178 $7 \times 10^6 \text{ kg s}^{-1}$). Another condition was that there must be a layer at least 500 m AGL that can build a cloud at least 2 km
179 thick after the layer is raised to its lifting condensation level. Note that the deep and shallow convection schemes
180 would not be activated in a low convective-available-potential-energy (CAPE) environment, but the mid-level
181 convection scheme could be triggered even in low-CAPE environments.

182 **S1.8 Gravity wave drag and low-level orographic blocking schemes**

183 SGS orography is associated with two processes that must be parameterized: (i) vertical momentum transport due to
184 the breaking of mountain-induced gravity waves (“mountain wave drag”) and (ii) blocking of low-level upstream flows
185 by mountain barriers. Parameterizations due to McFarlane (1987) for mountain wave drag and Lott and Miller (1997)
186 for low-level blocking (with an enhanced drag coefficient: Wells et al., 2008; Vosper et al., 2009) were used by all
187 three forecast systems (McTaggart-Cowan et al., 2019a; CMC-GDPS-8.0.0, 2021b; CMC-RDPS-8.0.0, 2021b). In
188 addition, the Hines (1997a,b) Doppler spread parametrization for non-orographic gravity-wave drag due to imbalances
189 and geostrophic adjustment associated with a variety of transient mesoscale meteorological features was also used to
190 simulate the upper-level circulation accurately (Charron et al., 2002; McTaggart-Cowan et al., 2019a; CMC-GDPS-
191 8.0.0, 2021b; CMC-RDPS-8.0.0, 2021b)

192 The geophysical variables required by these schemes include gridded orography plus SGS orographic parameters for
193 gravity wave drag and low-level blocking. Note that it is a standard procedure to apply a smoothing operator to the
194 model orography field in order to minimize stationary noise generation due to small-scale resolved orographic features.
195 A numerical filter with a wavelength cutoff at $3\Delta x$ for the GDPS 8.0.0 and $5\Delta x$ for the RDPS 8.0.0 and RAQDPS023
196 was applied. SGS orographic parameters were computed using an analytically-extended reference orography spectrum
197 and a 5 km cut-off for horizontal terrain scales (Beljaars et al., 2004; McTaggart-Cowan et al., 2019a; CMC-GDPS-
198 8.0.0, 2021b; CMC-RDPS-8.0.0, 2021b).

199 **S2 Overview of anthropogenic national emissions inventories pre-processing and processing**

200 A number of adjustments had to be made to the projected national emissions inventories before they underwent the
201 processing required to create the SET4.0.0 emissions input files. For the projected 2020 Canadian Air Pollutant
202 Emissions Inventory (APEI), emissions from two emissions sectors, prescribed burning and agricultural-land wind
203 erosion, were removed from the inventory because of insufficient information to allocate these emissions to specific
204 days of the year or times of day. Projected 2020 aircraft in-flight cruise emissions were also removed because they
205 would not materially affect short-term surface pollutant concentrations due to the altitude (6-10 km above sea level) at
206 which they were emitted, whereas projected 2020 aircraft landing and takeoff emissions were kept. Projected 2020
207 residential wood combustion (RWC) emission values were replaced by 2010 APEI values due to concerns over some
208 aspects of the revised estimation methodology used to calculate RWC emissions for the 2015 APEI (ECCC, 2013,
209 2016). In addition, projected 2020 emission values for one power plant and one large industrial facility that had been
210 active in 2015 but that were known to have closed by 2020 were removed from the major-point-source emissions file
211 (Moran et al., 2021).

212 Adjustments were also made to Canadian projected 2020 fugitive dust emissions from paved and unpaved roads and
213 from construction activities. Fugitive dust emissions are the largest component of anthropogenic PM_{2.5} and PM₁₀
214 emissions, of which the main contributing emission sectors are vehicular traffic on paved and unpaved road surfaces
215 and agricultural and construction activities. Emissions estimation methodologies are always subject to re-evaluation
216 and improvement, and a new methodology to estimate fugitive dust emissions from paved and unpaved road surfaces
217 was introduced by ECCC in 2020 for the 2018 APEI (ECCC, 2020). In order to account for these methodological
218 changes in the projected 2020 APEI, whose base 2015 fugitive road dust emissions had been calculated using an older
219 methodology (ECCC, 2016), the paved-road fugitive dust emissions from the projected 2020 inventory for each
220 province and territory were scaled by a factor of 0.15 (i.e., decreased by 85%) and unpaved-road fugitive dust emissions
221 were scaled by a factor of 1.11 (i.e., increased by 11%).

222 One useful check on any Canadian CAC annual emissions inventory is to compare per capita Canadian and U.S.
223 emissions for the same emissions source sectors for the same year or closest year (since U.S. National Emissions
224 Inventories (NEIs) are only available every three years). Given that differences in population have been controlled for
225 by considering per capita values, one expects the per capita emissions values to be broadly similar given socioeconomic
226 and emission control similarities between the two countries. When this comparison was made for the scaled 2015
227 Canadian fugitive dust emissions from paved and unpaved road surfaces and from agricultural activities and residential
228 construction, the agreement with the closest U.S. CAC inventory, the 2014 U.S. NEI (<https://www.epa.gov/air-emissions-inventories/2014-national-emissions-inventory-nei-data>), was reasonable. For 2015 fugitive dust emissions
229 from heavy construction, however, per capita Canadian emissions were 40 times higher than 2014 per capita U.S.
230 emissions. Moreover, the estimation methodology used to estimate 2015 Canadian heavy construction emissions had
231 not changed since 2012 and heavy construction emissions in the APEI had been held constant since 2013, which
232 suggested a high level of uncertainty for these emissions. It was thus decided to scale Canadian projected 2020 fugitive
233

234 dust emissions for heavy construction by a factor of 0.05 so that 2015 per capita Canadian emissions from this sector
235 would be within a factor of two of the 2014 per capita U.S. emissions.

236 The Canadian projected 2020 APEI then underwent two final preparatory steps by ECCC before the emissions
237 processing step. First, the point source file was separated into two files, a major-point-source file and a minor-point-
238 source file, where a 20-m stack height threshold was applied to assign individual point sources to the major- or minor-
239 point-source files based on stack height. Similarly, the area source file was separated by emissions source type into
240 multiple files, including on-road mobile sources, off-road mobile sources, marine sources, agricultural sources, other
241 fugitive dust area sources, and other non-fugitive-dust area sources. This separation by source type made it possible
242 to apply special treatments such as meteorological modulation to an emissions subset (Sect. 3.12) and also to perform
243 quality-control checks more easily after the emissions processing step. The second final preparatory step was to convert
244 the Canadian projected 2020 APEI into a different format that would be readable by the emissions processing system.
245 This step included augmentation of the APEI to include some non-standard data such as point source stack
246 characteristics (Sect. 3.13) and facility operating schedules (e.g., Sassi et al., 2021).

247 Several minor adjustments were also required to the projected 2023 U.S. NEI for emissions from commercial marine
248 vessels. These emissions had been pre-processed by the U.S. EPA for three non-overlapping modelling grids used by
249 the EPA: (i) emissions within U.S. federal waters for the continental U.S. on a 12-km horizontal grid; (ii) emissions
250 for Alaska on a 9-km grid; and (iii) emissions for other Pacific and Atlantic Ocean areas on a 4-km grid. However,
251 there were spatial gaps among these three emissions regions, which were filled by inserting emissions from another
252 marine emissions data set that was used for Phase 4 of the Air Quality Model International Initiative (AQMEII-4)
253 project (Galmarini et al., 2021). The projected 2023 U.S. inventory also contained marine emissions over some
254 Canadian waters: these emissions were removed to avoid double counting with the Canadian inventory. Note that the
255 EPA had created an emissions-processing version of the projected 2023 U.S. NEI and had also separated the surface
256 emissions into multiple files by source type for easier processing and tracking (U.S. EPA, 2019).

257 **S2.1 Emissions processing**

258 Once the above inventory adjustments had been made, the three projected national inventories were processed using
259 version 4.7 of the SMOKE (Sparse Matrix Operation Kernel Estimation) emissions processing system software (see
260 <https://www.emascenter.org/smoke/>) to generate the SET4.0.0 set of major-point-source emissions files and gridded
261 surface emissions files. Two types of emissions files were needed because the separate major-point-source files
262 contained emissions for large individual sources such as power plants and smelters for which plume rise was calculated
263 by GEM-MACH (Sect. 3.13), whereas the surface emissions were the sum of area sources, on-road mobile sources,
264 off-road mobile sources, and minor point sources for which plume rise was not relevant. Note that there were also two
265 types of surface emissions identified in the SET4.0.0 set, one type identifying fugitive-dust area emissions and the
266 other type identifying all other surface emissions. This made it possible to apply meteorological modulation only to
267 fugitive-dust area emissions (cf. Sect. 3.12).

268 The SMOKE emissions processing system must perform four major processing tasks: (i) temporal allocation, where
269 annual/monthly inventory emissions were allocated to specific hours of specific days and months of the year based on
270 a set of source-sector-specific monthly, day-of-the-week (DOW), and diurnal temporal profiles; (ii) spatial allocation,
271 where emissions reported by jurisdiction such as a province or county were allocated to the group of model grid cells
272 that overlay that jurisdiction in as realistic a pattern as possible using source-sector-specific gridded spatial surrogate
273 fields; (iii) chemical speciation, where four of the CACs contained in the inventories (NO_x , VOC, $\text{PM}_{2.5}$, PM_{10}) were
274 separated into more detailed model species using source-sector-specific speciation profiles; and (iv) PM_x size
275 disaggregation ($x = 2.5$ or 10), where PM_x emissions from the inventory were allocated to different model PM size
276 sections or modes using source-sector-specific PM size distribution profiles (e.g., Dickson and Oliver, 1991; Houyoux
277 et al., 2000; Moran et al., 2013; Matthias et al., 2018; Zhang et al., 2018; Boutzis et al., 2020).

278 Each SET4.0.0 emissions file generated by SMOKE was specific to a particular day of the week and month of the year,
279 and each file contained 24 hours of gridded emission fields of model species (where NO_x emissions were split into
280 emissions of NO, NO_2 , and HONO, $\text{PM}_{2.5}$ and PM_{10} emissions were each split into emissions of six model PM chemical
281 components, and VOC emissions were split into emissions of 11 model lumped VOC species for the ADOM-2 gas-
282 phase chemistry mechanism) (Stroud et al., 2008; Zhang et al., 2018; see Table 3).

283 For the first emissions processing task, temporal allocation, SMOKE used 82 monthly temporal profiles, 22 DOW
284 profiles, and 29 diurnal profiles to process the Canadian emissions, 9,971 monthly profiles, 12 DOW profiles, and 95
285 diurnal profiles to process the U.S. emissions, and 161 monthly profiles, 15 DOW profiles, and 49 diurnal profiles to
286 process the Mexican emissions (CMC-RAQDPS-023, 2021). For the second task, spatial allocation, SMOKE
287 employed 55 source-sector-specific gridded spatial surrogate fields to allocate emissions for each Canadian province
288 or territory to one or more grid cells on the RAQDPS023 10-km continental Yin grid (Fig. 2), 42 gridded spatial
289 surrogate fields for each U.S. county, and 15 gridded spatial surrogate fields for each Mexican county (CMC-RAQDPS-
290 023, 2021). The gridded surface emissions from these different jurisdictions and source types were then summed for
291 each model grid cell. Major-point-source emissions, on the other hand, were assigned to a single model grid cell based
292 on their latitude-longitude coordinates while remaining in separate records for subsequent plume-rise calculations
293 (Sect. 3.13). For the third task, chemical speciation, SMOKE used 331 VOC and 85 PM speciation profiles from the
294 U.S. EPA's SPECIATE 4.5 database (<https://www.epa.gov/air-emissions-modeling/speciate-1>; Reff et al., 2009;
295 Simon et al., 2010). Only two speciation profiles were used to speciate NO_x emissions: one profile for on-road and
296 off-road sources, which included a small HONO component as well as NO and NO_2 components, and a generic profile
297 with only NO and NO_2 components for point and area sources. Lastly, the fourth task, PM size disaggregation, was
298 handled simply by assigning $\text{PM}_{2.5}$ emissions to the first RAQDPS023 PM size bin and assigning PM_{cf} emissions,
299 calculated as the difference between PM_{10} and $\text{PM}_{2.5}$ emissions, to the second RAQDPS023 PM size bin. Following
300 the chemical speciation and PM size disaggregation steps the SET4.0.0 output files produced by SMOKE contained
301 hourly gridded emission fields of 18 gases and 12 particle size bin/chemical components for a total of 30 emitted model

302 species: SO₂, (H₂)SO₄, NO, NO₂, HONO, CO, C₃H₈, ALKA, ETHE, ALKE, TOLU, AROM, HCHO, ALD₂, MEK,
303 CRES, ISOP, NH₃, SU₁, SU₂, NI₁, NI₂, AM₁, AM₂, EC₁, EC₂, PC₁, PC₂, CM₁, and CM₂.

304 It should be noted that two more adjustments were made to all fugitive dust emissions *after* SMOKE processing. One
305 adjustment was done immediately as a (static) post-processing step, where a transportable-fraction (TF) scaling factor
306 field was applied to account for “home-grid-cell” SGS particle settling and particle removal by interception and
307 impaction by vegetation (e.g., Pace, 2005). The TF scaling factor ranged between 0 and 1 and was dependent on the
308 fractional coverage of different land-use classes in each grid cell, where forest classes had a TF value of zero and
309 agricultural classes had a value of 0.75. As described in Sect. 3.12 the other (dynamic) adjustment was to apply
310 meteorological modulation at each chemistry time step during each RAQDPS023 simulation to account for the day-
311 specific dampening effects of precipitation and snow cover on fugitive dust emissions. In order to accommodate the
312 model calculations required for meteorological modulation and plume rise (Sects. 3.12 and 3.13), three sets of
313 emissions fields must be read by the RAQDPS023 each hour from two separate day-specific files: (i) major-point
314 source emissions; and (ii) surface emissions files consisting of fugitive dust emissions (type F) and other surface
315 emissions (type E). SET4.0.0 thus consisted of two input files per day of a typical week of each month, or 168 files in
316 total. More information related to the SMOKE processing required to create GEM-MACH emissions files may be
317 found in Zhang et al. (2018) and Sassi et al. (2021).

318 **S3 Description of Canadian Forest Fire Emissions Prediction System (CFFEPS) version 4.1**

319 CFFEPS v4.1 consists of a fire-growth module, a fire emissions module, and a thermodynamic-based module to
320 estimate the vertical penetration height of a smoke plume (Chen et al., 2019a,b). Outputs from the Canadian Wildland
321 Fire Information System (CWFIS), which is an operational product developed and managed by the Canadian Forest
322 Service (CFS), underpin the CFFEPS fire-growth module. The CWFIS is itself composed of three major subsystems:
323 (i) the Canadian Forest Fire Weather Index (FWI) subsystem, which accounts for the influence of four surface weather
324 elements (dry-bulb temperature, relative humidity, wind speed, and 24 hour accumulated precipitation) on the relative
325 risk for wildfire occurrence; (ii) the Fire Monitoring, Mapping, and Modeling (Fire M3) subsystem, which identifies
326 and locates active fires (“hotspots”) using NRT satellite imagery; and (iii) the Canadian Forest Fire Behaviour
327 Prediction (FBP) subsystem, which estimates fire fuel consumption and fire intensity
328 (<https://cwfis.cfs.nrcan.gc.ca/home>).

329 Input data needed by the FWI subsystem include observing station measurements of 2 m dry-bulb temperature and
330 relative humidity, 2 m dew point, 10 m wind speed, surface pressure, and past 24 hour rainfall from about 2500
331 Canadian and U.S. weather stations plus 72 hour ECCC weather forecasts for the same station locations. The six
332 primary outputs of the FWI subsystem are numerical moisture codes for the forest-floor fine-fuel layer (i.e., leaf litter,
333 needles, cured grass, small twigs; 1.2 cm deep), duff layer (layer of partially/fully decomposed organic materials
334 beneath the fine-fuel layer; 7 cm deep), and deep duff layer (beneath the duff layer; ~18 cm deep), an initial spread

335 index (a function of surface wind speed and fine-fuel moisture content), total subsurface fuel available for combustion,
336 and a combined FWI numerical value (Van Wagner, 1987; <https://cwfis.cfs.nrcan.gc.ca/background/summary/fwi>).

337 As of 2021 the fire hotspot data needed by the Fire M3 subsystem are obtained from the 4- μm and 11- μm infrared
338 channel imagery of two radiometer instruments carried on four U.S. polar-orbiting satellites: (i) the Moderate
339 Resolution Imaging Spectroradiometer (MODIS) on two NASA satellites; and (ii) the Visible Infrared Imaging
340 Radiometer Suite (VIIRS) on two more NASA satellites (<https://cwfis.cfs.nrcan.gc.ca/background/dsm/fm3>). One to
341 two retrievals covering all of Canada, including remote areas, are available from each satellite each day. However,
342 hotspot detection may not be possible due to cloud or thick smoke obscuration of the surface or if fire size or intensity
343 is small, so not every fire, even large fires, will be detected every day. In addition, internal data filtering and
344 measurements at other wavelengths are applied to remove false positives (e.g., industrial sources, roofs of large
345 buildings, cloud edges). To support ECCC the CFS has extended CWFIS capabilities to provide fire hotspots over the
346 U.S. as well as Canada throughout the year instead of just the May to September period, which historically was the
347 typical wildfire season in Canada. Note that the satellite measurements have a time lag or latency between one and
348 seven hours due to the time required for image acquisition and distribution, hence the NRT descriptor.

349 The third major CWFIS component is the FBP subsystem, which uses FWI system outputs together with fuel type and
350 topography to predict fire rate of spread (ROS; m min^{-1}), fire intensity (kW m^{-1}), crown fraction burned and fire type
351 (crown fire or surface fire), and total fuel consumption ($\text{kg dry biomass m}^{-2}$) at hotspot locations
352 (<https://cwfis.cfs.nrcan.gc.ca/background/summary/fbp>). Sixteen fuel types such as Boreal Spruce and Ponderosa
353 Pine–Douglas Fir are considered (e.g., <https://cwfis.cfs.nrcan.gc.ca/background/fueltypes/c2>).

354 **S3.1 Forest fire emission forecasts**

355 CFFEPS v4.1 includes the source code for the FWI and FBP subsystems, which allows meteorological forecast fields
356 from GEM to be considered for every hour and not just for noontime as in the CFS operational version of the CWFIS.
357 Hourly biomass burning (BB) emissions are then estimated in CFFEPS v4.1 using a bottom-up, process-based
358 approach, where emissions per fire (or hotspot) per day are the product of estimated burn area per hotspot, estimated
359 fuel consumption per unit area, and emission factors for different chemical species and combustion stages (Chen et al.,
360 2019a,b). Hotspot and associated fuel type information for the previous 24 hours is provided by the CWFIS Fire M3
361 subsystem. Initial daily burn area per hotspot is a prescribed value by province and fuel type that is updated periodically
362 by CFS based on their analysis of CWFIS historical averages of reported burn areas by eco-region and fuel type; it is
363 assumed to remain constant for each day of the forecast (i.e., persistence). Fuel consumption values are estimated by
364 fuel type via the FBP subsystem driven with forecast meteorology. The incorporation of the FBP subsystem allows
365 CFFEPS v4.1 to account for spatially variable biomass/fuel inputs using forest composition data from Canada's
366 National Forest Inventory (Beaudoin et al., 2014, 2018) (Chen and Menelaou, 2021).

367 To calculate fire emissions, CFFEPS v4.1 uses FBP fuel combustion estimates to account for emissions from three
368 combustion stages: flaming, smoldering, and residual combustion (Chen et al., 2019b). A look-up table is used to

369 allocate fuel consumption between these different stages (Table 2 of Chen et al., 2019b), which have different burn
370 durations: 15 minutes for flaming followed by several hours for smoldering and several hours for residual combustion,
371 depending on the thickness of the forest-floor organic duff layer (Chen et al., 2019b). Different emission factors, which
372 are constant parameters representing grams of emissions per kilogram of fuel consumed, are applied to estimate
373 emissions from a wildfire for each combustion stage for eight chemical species: CO, CH₄, NO_x, NH₃, SO₂, non-
374 methane hydrocarbons (NMHC), PM_{2.5}, and PM₁₀ (Table 3 of Chen et al., 2019b). Chemical speciation is further
375 applied for NMHC emissions (cf. Sect. S2) to calculate ADOM-2 VOC species emissions using the U.S. EPA
376 SPECIATEv4.5 speciation profiles for flaming (#95425) and smoldering and residual combustion (#95428)
377 (<https://www.epa.gov/air-emissions-modeling/speciate-1>; Chen et al., 2019a). Similarly, PM_{2.5} and PM₁₀ emissions
378 from CFFEPS are subdivided into six PM chemical components with the following EPA SPECIATEv4.5 wildfire PM
379 speciation profile: POM (78.5%), EC (9.5%), CM (9.7%), SU (1.3%), NI (0.1%), and AM (0.9%) (Chen et al., 2019a).

380 Fire emission forecasts were calculated by CFFEPS v4.1 prior to each RAQDPS-FW023 forecast simulation using the
381 latest hotspot information for the past 24 hours from the operational CFS version of CWFIS. To obtain 72 hour
382 forecasts of hourly BB emission values, current hotspots were assumed to persist for the next 72 hours and hourly
383 changes in biomass fuel consumption were parameterized for a 24-hour period based on burn duration for the three
384 combustion stages, fire ROS, and the average diurnal variation of the moisture content of the forest-floor litter layer,
385 which increases as moisture content decreases (Chen et al., 2019b).

386 In the previous version of CFFEPS, v2.06, the diurnal distribution of daily burn area, which is used to allocate daily to
387 hourly emissions, was estimated by a fixed weighted-normalization approach using a predefined diurnal profile, with
388 peak fire growth at 16:00 local standard time (LST) and minimal near constant growth between 0:00 to 8:00 LST (Chen
389 et al., 2019b). In CFFEPS v4.1, however, the diurnal distribution of daily burn area was determined based on a
390 weighted moving average of the ratio of hourly fire ROS as estimated by the FBP subsystem to the sum of hourly ROS
391 values for the previous 24 hours (Chen and Menelaou, 2021). This new scheme should be more realistic as it is based
392 on hourly forecast meteorology rather than a static temporal profile.

393 **S3.2 Plume rise**

394 Plume-rise schemes for smokestacks like the Briggs (1984) scheme (cf. Sect. 3.13) have also been used to estimate
395 wildfire plume rise, but wildland fires are much broader than smokestacks and they lack any initial vertical velocity;
396 instead, they generate their own buoyant plume. CFFEPS v4.1 uses an alternate thermodynamic approach described
397 by Anderson et al. (2011) to parameterize wildfire plume rise. Hourly plume injection height is estimated based on a
398 fire energy balance such that the buoyancy of a smoke plume is thermodynamically balanced with the forecast vertical
399 temperature profile of the atmospheric column above the fire while accounting for fuel moisture evaporation and
400 condensation.

401 In earlier versions of CFFEPS, the plume-rise scheme was based on lifting the smoke plume dry adiabatically at a
402 constant rate at hotspot locations and then equilibrating the buoyant plume with the local environmental lapse rate

403 derived from a simplified vertical temperature profile determined by temperatures on five pressure levels: surface, 850,
404 700, 500, and 250 hPa (Chen et al., 2019b). CFFEPS v4.1 instead uses the higher-resolution forecast vertical
405 temperature profiles that are available on the RAQDPS-FW023 84-layer vertical grid (Sect. 2.1). The thermodynamic
406 calculation of plume heat flux in CFFEPS v4.1 was also updated to account for the heat lost due to the latent heat of
407 evaporation of fuel moisture during combustion. At the same time, as the plume passes the atmospheric lifting
408 condensation level, the latent heat of condensation of plume water vapour is added to the energy of the plume aloft to
409 obtain the final injection height within the atmospheric column. These detailed physical processes are considered
410 hourly for each hotspot as the fire thermal energy and the atmospheric vertical temperature profile change, and the final
411 hourly plume injection height is used to prescribe the vertical distribution of emissions within the column below the
412 plume injection height (Chen et al., 2019b; Chen and Menelaou, 2021).

413 **S3.3 Outputs and file structure**

414 CFFEPS v4.1 outputs are saved in one file similar in format to the anthropogenic major-point-source emissions input
415 file. The outputs are estimated hourly plume injection height (ZPLM meters) and hourly speciated emissions (g s^{-1})
416 over the 72-hour forecast period at each hotspot location as specified by latitude/longitude. There are 21 ADOM-2
417 gas- and 12 particle-phase emitted species that are output (identified by a leading “E”): ECO, ECO2, ENO, ENO2,
418 ENH3, ESO2, ESO4, EA2, EA3, EALD, EARO, EC38, ECRE, EETH, EHCH, EISO, EMEK, ETOL, EOTH, EC1,
419 EC2, EAM1, EAM2, ECM1, ECM2, EEC1, EEC2, ENI1, ENI2, EPC1, EPC2, ESU1, and ESU2 (cf. Table 3). Note
420 that ECO2, EC1 (methane), and EC2 (ethane) are not needed by the ADOM-2 gas-phase chemistry mechanism but
421 may be used in other chemistry parameterizations. This BB emissions file was input directly to the RAQDPS-FW023,
422 which then allocated the emissions to model surface and elevated grid cells.

423 **S3.4 Limitations**

424 CFFEPS v4.1 has a number of limitations that should be kept in mind (Chen et al., 2019a). First, as already noted
425 above some fires may not be detected due to (i) their small size, (ii) their initiation after the last available satellite
426 overpass, or (iii) the presence of thick cloud or heavy smoke above the fire. Second, detected fires are assumed to
427 continue burning but to remain the same size over the duration of the forecast. Third, fire emissions are treated as point
428 sources and hence are assigned to a single grid cell. And fourth, wildfire combustion heat and emitted smoke are
429 assumed not to affect meteorological fields, which may not be a good assumption for large, intense fires.

430

431 **Supplement References**

432 Anderson, K., Pankratz, A., and Mooney, C.: A thermodynamic approach to estimating smoke plume heights, in: Proc. Ninth
433 Symposium on Fire and Forest Meteorology, Oct. 17-20, Palm Springs, California, 8 pp.,
434 <https://ams.confex.com/ams/9FIRE/webprogram/Manuscript/Paper192294/FFM9 - Smoke - final.pdf>, 2011.

435 Aranami, K., Davies, T., and Wood, N.: A mass restoration scheme for limited-area models with semi-Lagrangian advection, Q.
436 J. R. Meteorol. Soc., 141, 1795–1803, <https://doi.org/10.1002/qj.2482>, 2015.

- 437 Beaudoin, A., Bernier, P. Y., Guindon, L., Villemaire, P., Guo, X. J., Stinson, G., Bergeron, T., Magnussen, S., and Hall, R. J.:
 438 Mapping attributes of Canada's forests at moderate resolution through k NN and MODIS imagery, *Can. J. For. Res.*, 44, 521–532,
 439 <https://doi.org/10.1139/cjfr-2013-0401>, 2014.
- 440 Beaudoin, A., Bernier, P. Y., Villemaire, P., Guindon, L., and Guo, X. J.: Tracking forest attributes across Canada between 2001
 441 and 2011 using a k nearest neighbors mapping approach applied to MODIS imagery, *Can. J. For. Res.*, 48, 85–93,
 442 <https://doi.org/10.1139/cjfr-2017-0184>, 2018.
- 443 Bechtold, P. and Siebesma, P.: Organization and representation of boundary layer clouds, *J. Atmospheric Sci.*, 55, 888–895,
 444 [https://doi.org/10.1175/1520-0469\(1998\)055<0888:OAROBL>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<0888:OAROBL>2.0.CO;2), 1998.
- 445 Bechtold, P., Bazile, E., Guichard, F., Mascart, P., and Richard, E.: A mass-flux convection scheme for regional and global models,
 446 *Q. J. R. Meteorol. Soc.*, 127, 869–886, <https://doi.org/10.1002/qj.49712757309>, 2001.
- 447 Bechtold, P., Semane, N., Lopez, P., Chaboureau, J.-P., Beljaars, A., and Bormann, N.: Representing equilibrium and
 448 nonequilibrium convection in large-scale models, *J. Atmospheric Sci.*, 71, 734–753, <https://doi.org/10.1175/JAS-D-13-0163.1>,
 449 2014.
- 450 Bélair, S., Mailhot, J., Strapp, J. W., and MacPherson, J. I.: An examination of local versus nonlocal aspects of a TKE-based
 451 boundary layer scheme in clear convective conditions, *J. Appl. Meteorol.*, 38, 1499–1518, [https://doi.org/10.1175/1520-0450\(1999\)038<1499:AEOLVN>2.0.CO;2](https://doi.org/10.1175/1520-0450(1999)038<1499:AEOLVN>2.0.CO;2), 1999.
- 453 Bélair, S., Crevier, L.-P., Mailhot, J., Bilodeau, B., and Delage, Y.: Operational implementation of the ISBA land surface scheme
 454 in the Canadian regional weather forecast model. Part I: Warm season results, *J. Hydrometeorol.*, 4, 352–370,
 455 [https://doi.org/10.1175/1525-7541\(2003\)4<352:OIOTIL>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)4<352:OIOTIL>2.0.CO;2), 2003a.
- 456 Bélair, S., Brown, R., Mailhot, J., Bilodeau, B., and Crevier, L.-P.: Operational implementation of the ISBA land surface scheme
 457 in the Canadian regional weather forecast model. Part II: Cold season results, *J. Hydrometeorol.*, 4, 371–386,
 458 [https://doi.org/10.1175/1525-7541\(2003\)4<371:OIOTIL>2.0.CO;2](https://doi.org/10.1175/1525-7541(2003)4<371:OIOTIL>2.0.CO;2), 2003b.
- 459 Bélair, S., Mailhot, J., Girard, C., and Vaillancourt, P.: Boundary layer and shallow cumulus clouds in a medium-range forecast of
 460 a large-scale weather system, *Mon. Weather Rev.*, 133, 1938–1960, <https://doi.org/10.1175/MWR2958.1>, 2005.
- 461 Beljaars, A. C. M. and Holtslag, A. A. M.: Flux parameterization over land surfaces for atmospheric models, *J. Appl. Meteorol.*,
 462 30, 327–341, [https://doi.org/10.1175/1520-0450\(1991\)030<0327:FPOLSF>2.0.CO;2](https://doi.org/10.1175/1520-0450(1991)030<0327:FPOLSF>2.0.CO;2), 1991.
- 463 Beljaars, A. C. M., Brown, A. R., and Wood, N.: A new parametrization of turbulent orographic form drag, *Q. J. R. Meteorol.*
 464 *Soc.*, 130, 1327–1347, <https://doi.org/10.1256/qj.03.73>, 2004.
- 465 Benoit, R., Côté, J., and Mailhot, J.: Inclusion of a TKE boundary layer parameterization in the Canadian Regional Finite-Element
 466 Model, *Mon. Weather Rev.*, 117, 1726–1750, [https://doi.org/10.1175/1520-0493\(1989\)117<1726:IOATBL>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1726:IOATBL>2.0.CO;2), 1989.
- 467 Bermejo, R. and Conde, J.: A conservative quasi-monotone semi-Lagrangian scheme, *Mon. Weather Rev.*, 130, 423–430,
 468 [https://doi.org/10.1175/1520-0493\(2002\)130<0423:ACQMSL>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<0423:ACQMSL>2.0.CO;2), 2002.
- 469 Blackadar, A. K.: The vertical distribution of wind and turbulent exchange in a neutral atmosphere, *J. Geophys. Res.*, 67, 3095–
 470 3102, <https://doi.org/10.1029/JZ067i008p03095>, 1962.
- 471 Bougeault, P. and Lacarrere, P.: Parameterization of orography-induced turbulence in a mesobeta--scale model, *Mon. Weather*
 472 *Rev.*, 117, 1872–1890, [https://doi.org/10.1175/1520-0493\(1989\)117<1872:POOITI>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1872:POOITI>2.0.CO;2), 1989.
- 473 Boutzis, E. I., Zhang, J., and Moran, M. D.: Expansion of a size disaggregation profile library for particulate matter emissions
 474 processing from three generic profiles to 36 source-type-specific profiles, *J Air Waste Manag Assoc.*, 70, 1067–1100,
 475 <https://doi.org/10.1080/10962247.2020.1743794>, 2020.
- 476 Briggs, G. A.: Plume rise and buoyancy effects, in: *Atmospheric Sciences and Power Production*, DOE/TIC-27601 (DE84005177),
 477 edited by D. Randerson, Technical Information Center, U.S Department of Energy, Oak Ridge, Tennessee, 327–366, 1984.
- 478 Charnock, H.: Wind stress on a water surface, *Q. J. R. Meteorol. Soc.*, 81, 639–640, <https://doi.org/10.1002/qj.49708135027>, 1955.

479 Charron, M., Manzini, E., and Warner, C. D.: Intercomparison of gravity wave parameterizations: Hines Doppler-spread and
480 Warner and McIntyre ultra-simple schemes., *J. Meteorol. Soc. Jpn. Ser II*, 80, 335–345, <https://doi.org/10.2151/jmsj.80.335>, 2002.

481 Chen, J. and Menelaou, K.: Regional Air Quality Deterministic Prediction System with near-real-time wildfire emissions
482 (RAQDPSFW): Upgrade to version 023, Technical note, November, Canadian Centre for Meteorological and Environmental
483 Prediction, Montreal, 31 pp.,
484 https://collaboration.cmc.ec.gc.ca/cmc/CMOI/product_guide/docs/tech_notes/technote_raqdpsfw_e.pdf, 2021.

485 Chen, J., Moran, M. D., Pavlovic, R., and Munoz-Alpizar, R.: RAQDPS-FireWork v20.2 (RAQDPS020.2FW): Operational
486 version of the Regional Air Quality Deterministic Prediction System with near-real-time wildfire emissions from the Canadian
487 Forest Fire Emission Prediction System (CFFEPS), Technical note, April, Canadian Centre for Meteorological and Environmental
488 Prediction, Montreal, 14 pp.,
489 https://collaboration.cmc.ec.gc.ca/cmc/cmoe/product_guide/docs/tech_notes/technote_raqdps020.2fw_20190415_e.pdf, 2019a.

490 Chen, J., Anderson, K., Pavlovic, R., Moran, M. D., Englefield, P., Thompson, D. K., Munoz-Alpizar, R., and Landry, H.: The
491 FireWork v2.0 air quality forecast system with biomass burning emissions from the Canadian Forest Fire Emissions Prediction
492 System v2.03, *Geosci. Model Dev.*, 12, 3283–3310, <https://doi.org/10.5194/gmd-12-3283-2019>, 2019b.

493 CMC-GDPS-8.0.0: Changes to the Global Deterministic Prediction System from version 7.1.0 to version 8.0.0, Technical note,
494 December, Canadian Centre for Meteorological and Environmental Prediction, Montreal, 112 pp.,
495 http://collaboration.cmc.ec.gc.ca/cmc/cmoe/product_guide/docs/tech_notes/technote_gdps-800_e.pdf, 2021a.

496 CMC-GDPS-8.0.0: The Global Deterministic Prediction System (GDPS) version 8.0.0 of the Meteorological Service (MSC) of
497 Canada: Technical specifications document, December, Canadian Centre for Meteorological and Environmental Prediction,
498 Montreal, 14 pp.,
499 https://collaboration.cmc.ec.gc.ca/cmc/cmoe/product_guide/docs/tech_specifications/tech_specifications_GDPS_8.0.0_e.pdf,
500 2021b.

501 CMC-RAQDPS-023: The Regional Air Quality Deterministic Prediction System (RAQDPS) version 023 and the Regional Air
502 Quality Deterministic Prediction System with Near-Real-Time Wildfire Emissions (RAQDPSFW) version 023 of the
503 Meteorological Service of Canada (MSC): Technical Specifications Document, November, Canadian Centre for Meteorological
504 and Environmental Prediction, Montreal, 19 pp.,
505 https://collaboration.cmc.ec.gc.ca/cmc/cmoe/product_guide/docs/tech_specifications/tech_specifications_RAQDPS_023_e.pdf,
506 2021.

507 CMC-RDPS-8.0.0: Changes to the Regional Deterministic Prediction System (RDPS) from version 7.1.0 to version 8.0.0,
508 Technical note, December, Canadian Centre for Meteorological and Environmental Prediction, Montreal, 36 pp.,
509 https://collaboration.cmc.ec.gc.ca/cmc/cmoe/product_guide/docs/tech_notes/technote_rdps-800_e.pdf, 2021a.

510 CMC-RDPS-8.0.0: The Regional Deterministic Prediction System (RDPS) version 8.0.0 of the Meteorological Service of Canada
511 (MSC): Technical specifications document, December, Canadian Centre for Meteorological and Environmental Prediction,
512 Montreal, 10 pp.,
513 https://collaboration.cmc.ec.gc.ca/cmc/cmoe/product_guide/docs/tech_specifications/tech_specifications_RDPS_8.0.0_e.pdf,
514 2021b.

515 Côté, J., Gravel, S., Méthot, A., Patoine, A., Roch, M., and Staniforth, A.: The operational CMC–MRB Global Environmental
516 Multiscale (GEM) model. Part I: Design considerations and formulation, *Mon. Weather Rev.*, 126, 1373–1395,
517 [https://doi.org/10.1175/1520-0493\(1998\)126<1373:TOCMGE>2.0.CO;2](https://doi.org/10.1175/1520-0493(1998)126<1373:TOCMGE>2.0.CO;2), 1998.

518 Deacu, D., Fortin, V., Klyszejko, E., Spence, C., and Blanken, P. D.: Predicting the net basin supply to the Great Lakes with a
519 hydrometeorological model, *J. Hydrometeorol.*, 13, 1739–1759, <https://doi.org/10.1175/JHM-D-11-0151.1>, 2012.

520 Delage, Y. and Girard, C.: Stability functions correct at the free convection limit and consistent for both the surface and Ekman
521 layers, *Bound.-Layer Meteorol.*, 58, 19–31, <https://doi.org/10.1007/BF00120749>, 1992.

522 Dickson, R. J. and Oliver, W. R.: Emissions models for regional air quality studies, *Environ. Sci. Technol.*, 25, 1533–1535,
523 <https://doi.org/10.1021/es00021a003>, 1991.

524 ECCC: Manual for the compilation of Canada’s air pollutant emissions, Environment Canada, Gatineau, Québec, 41 pp.,
525 https://publications.gc.ca/collections/collection_2013/ec/En14-87-2013-eng.pdf, 2013.

- 526 ECCC: 1990–2015 Air Pollutant Emission Inventory Report, Environment and Climate Change Canada, Gatineau, Quebec, 97
527 pp., https://publications.gc.ca/collections/collection_2017/eccc/En81-26-2015-eng.pdf, 2016.
- 528 ECCC: Canada’s Air Pollutant Emissions Inventory Report 1990–2018, Environment and Climate Change Canada, Gatineau,
529 Quebec, 96 pp., https://publications.gc.ca/collections/collection_2020/eccc/En81-30-2018-eng.pdf, 2020.
- 530 Foken, T.: 50 Years of the Monin–Obukhov similarity theory, *Bound.-Layer Meteorol.*, 119, 431–447,
531 <https://doi.org/10.1007/s10546-006-9048-6>, 2006.
- 532 Galmarini, S., Makar, P., Clifton, O. E., Hogrefe, C., Bash, J. O., Bellasio, R., Bianconi, R., Bieser, J., Butler, T., Ducker, J.,
533 Flemming, J., Hodzic, A., Holmes, C. D., Kioutsioukis, I., Kranenburg, R., Lupascu, A., Perez-Camanyo, J. L., Pleim, J., Ryu, Y.-
534 H., San Jose, R., Schwede, D., Silva, S., and Wolke, R.: Technical note: AQMEII4 Activity 1: evaluation of wet and dry deposition
535 schemes as an integral part of regional-scale air quality models, *Atmospheric Chem. Phys.*, 21, 15663–15697,
536 <https://doi.org/10.5194/acp-21-15663-2021>, 2021.
- 537 Girard, C., Plante, A., Desgagné, M., McTaggart-Cowan, R., Côté, J., Charron, M., Gravel, S., Lee, V., Patoine, A., Qaddouri, A.,
538 Roch, M., Spacek, L., Tanguay, M., Vaillancourt, P. A., and Zadra, A.: Staggered vertical discretization of the Canadian
539 Environmental Multiscale (GEM) model using a coordinate of the log-hydrostatic-pressure type, *Mon. Weather Rev.*, 142, 1183–
540 1196, <https://doi.org/10.1175/MWR-D-13-00255.1>, 2014.
- 541 de Grandpré, J., Tanguay, M., Qaddouri, A., Zerroukat, M., and McLinden, C. A.: Semi-Lagrangian advection of stratospheric
542 ozone on a Yin–Yang grid system, *Mon. Weather Rev.*, 144, 1035–1050, <https://doi.org/10.1175/MWR-D-15-0142.1>, 2016.
- 543 Hansen, A. B., Brandt, J., Christensen, J. H., and Kaas, E.: Semi-Lagrangian methods in air pollution models, *Geosci. Model Dev.*,
544 4, 511–541, <https://doi.org/10.5194/gmd-4-511-2011>, 2011.
- 545 Hines, C. O.: Doppler-spread parameterization of gravity-wave momentum deposition in the middle atmosphere. Part 1: Basic
546 formulation, *J. Atmospheric Sol.-Terr. Phys.*, 59, 371–386, [https://doi.org/10.1016/S1364-6826\(96\)00079-X](https://doi.org/10.1016/S1364-6826(96)00079-X), 1997a.
- 547 Hines, C. O.: Doppler-spread parameterization of gravity-wave momentum deposition in the middle atmosphere. Part 2: Broad
548 and quasi monochromatic spectra, and implementation, *J. Atmospheric Sol.-Terr. Phys.*, 59, 387–400,
549 [https://doi.org/10.1016/S1364-6826\(96\)00080-6](https://doi.org/10.1016/S1364-6826(96)00080-6), 1997b.
- 550 Houyoux, M. R., Vukovich, J. M., Coats, C. J., Wheeler, N. J. M., and Kasibhatla, P. S.: Emission inventory development and
551 processing for the Seasonal Model for Regional Air Quality (SMRAQ) project, *J. Geophys. Res. Atmospheres*, 105, 9079–9090,
552 <https://doi.org/10.1029/1999JD900975>, 2000.
- 553 Husain, S. Z. and Girard, C.: Impact of consistent semi-Lagrangian trajectory calculations on numerical weather prediction
554 performance, *Mon. Weather Rev.*, 145, 4127–4150, <https://doi.org/10.1175/MWR-D-17-0138.1>, 2017.
- 555 Kain, J. S. and Fritsch, J. M.: A one-dimensional entraining/detraining plume model and its application in convective
556 parameterization, *J. Atmospheric Sci.*, 47, 2784–2802, [https://doi.org/10.1175/1520-0469\(1990\)047<2784:AODEPM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1990)047<2784:AODEPM>2.0.CO;2),
557 1990.
- 558 Kain, J. S. and Fritsch, J. M.: The role of the convective ‘trigger’ function in numerical forecasts of mesoscale convective systems,
559 *Meteorol. Atmospheric Phys.*, 49, 93–106, <https://doi.org/10.1007/BF01025402>, 1992.
- 560 Li, J. and Barker, H. W.: A radiation algorithm with correlated-k distribution. Part I: Local thermal equilibrium, *J. Atmospheric*
561 *Sci.*, 62, 286–309, <https://doi.org/10.1175/JAS-3396.1>, 2005.
- 562 Lott, F. and Miller, M. J.: A new subgrid-scale orographic drag parametrization: Its formulation and testing, *Q. J. R. Meteorol.*
563 *Soc.*, 123, 101–127, <https://doi.org/10.1002/qj.49712353704>, 1997.
- 564 Mailhot, J. and Bélair, S.: An examination of a unified cloudiness-turbulence scheme with various types of cloudy boundary layers,
565 in: *Proc. 14th Symposium on Boundary Layer and Turbulence*, 7-11 August, Aspen, Colorado, American Meteorological Society,
566 215–218, 2000.

567 Mailhot, J., Bélair, S., Lefavre, L., Bilodeau, B., Desgagné, M., Girard, C., Glazer, A., Leduc, A., Méthot, A., Patoine, A., Plante,
568 A., Rahill, A., Robinson, T., Talbot, D., Tremblay, A., Vaillancourt, P., Zadra, A., and Qaddouri, A.: The 15-km version of the
569 Canadian regional forecast system, *Atmosphere-Ocean*, 44, 133–149, <https://doi.org/10.3137/ao.440202>, 2006.

570 Matthias, V., Arndt, J. A., Aulinger, A., Bieser, J., Denier Van Der Gon, H., Kranenburg, R., Kuenen, J., Neumann, D., Pouliot,
571 G., and Quante, M.: Modeling emissions for three-dimensional atmospheric chemistry transport models, *J. Air Waste Manag.*
572 *Assoc.*, 68, 763–800, <https://doi.org/10.1080/10962247.2018.1424057>, 2018.

573 McFarlane, N. A.: The effect of orographically excited gravity wave drag on the general circulation of the lower stratosphere and
574 troposphere, *J. Atmospheric Sci.*, 44, 1775–1800, [https://doi.org/10.1175/1520-0469\(1987\)044<1775:TEOOEG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1987)044<1775:TEOOEG>2.0.CO;2), 1987.

575 McLinden, C. A., Olsen, S. C., Hannegan, B., Wild, O., Prather, M. J., and Sundet, J.: Stratospheric ozone in 3-D models: A simple
576 chemistry and the cross-tropopause flux, *J. Geophys. Res. Atmospheres*, 105, 14653–14665,
577 <https://doi.org/10.1029/2000JD900124>, 2000.

578 McTaggart-Cowan, R. and Zadra, A.: Representing Richardson Number hysteresis in the NWP boundary layer, *Mon. Weather*
579 *Rev.*, 143, 1232–1258, <https://doi.org/10.1175/MWR-D-14-00179.1>, 2015.

580 McTaggart-Cowan, R., Vaillancourt, P. A., Zadra, A., Chamberland, S., Charron, M., Corvec, S., Milbrandt, J. A., Paquin-Ricard,
581 D., Patoine, A., Roch, M., Separovic, L., and Yang, J.: Modernization of atmospheric physics parameterization in Canadian NWP,
582 *J. Adv. Model. Earth Syst.*, 11, 3593–3635, <https://doi.org/10.1029/2019MS001781>, 2019a.

583 McTaggart-Cowan, R., Vaillancourt, P. A., Zadra, A., Separovic, L., Corvec, S., and Kirshbaum, D.: A Lagrangian perspective on
584 parameterizing deep convection, *Mon. Weather Rev.*, 147, 4127–4149, <https://doi.org/10.1175/MWR-D-19-0164.1>, 2019b.

585 Moran, M. D., Dastoor, A., and Morneau, G.: Long-range transport of air pollutants and regional and global AQ modelling, in: *Air*
586 *Quality Management -- Canadian Perspectives on a Global Issue*, E. Taylor and A. McMillan, Editors, Springer, Dordrecht,
587 https://link.springer.com/content/pdf/10.1007%2F978-94-007-7557-2_4, 69–98, 2013.

588 Moran, M. D., Ménard, S., and Kornic, D.: Regional Air Quality Deterministic Prediction System (RAQDPS): Upgrade from
589 version 022 to version 023, Technical note, December, Canadian Centre for Meteorological and Environmental Prediction,
590 Montreal, 48 pp., https://collaboration.cmc.ec.gc.ca/cmc/cmoei/product_guide/docs/lib/technote_raqdps023_20211130_e.pdf,
591 2021.

592 Noilhan, J. and Planton, S.: A simple parameterization of land surface processes for meteorological models, *Mon. Weather Rev.*,
593 117, 536–549, [https://doi.org/10.1175/1520-0493\(1989\)117<0536:ASPOLS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<0536:ASPOLS>2.0.CO;2), 1989.

594 Pace, T. G.: Methodology to estimate the transportable fraction (TF) of fugitive dust emissions for regional and urban scale air
595 quality analyses (8/3/2005 Revision), U.S. Environmental Protection Agency, Research Triangle Park, North Carolina, 12 pp.,
596 <https://www.nrc.gov/docs/ML1321/ML13213A386.pdf>, 2005.

597 Pudykiewicz, J., Benoit, R., and Mailhot, J.: Inclusion and verification of a predictive cloud-water scheme in a regional numerical
598 weather prediction model, *Mon. Weather Rev.*, 120, 612–626, [https://doi.org/10.1175/1520-0493\(1992\)120<0612:IAVOAP>2.0.CO;2](https://doi.org/10.1175/1520-0493(1992)120<0612:IAVOAP>2.0.CO;2), 1992.

600 Reff, A., Bhawe, P. V., Simon, H., Pace, T. G., Pouliot, G. A., Mobley, J. D., and Houyoux, M.: Emissions inventory of PM_{2.5} trace
601 elements across the United States, *Environ. Sci. Technol.*, 43, 5790–5796, <https://doi.org/10.1021/es802930x>, 2009.

602 Ritchie, H., Bélair, S., Bernier, N. B., Buehner, M., Charron, M., Fortin, V., Garand, L., Houtekamer, P., Husain, S., Laroche, S.,
603 Lemieux, J.-F., Lin, H., McTaggart-Cowan, R., Milbrandt, J., Mitchell, H., Pellerin, P., Pudykiewicz, J., Separovic, L., Smith, G.
604 C., Tanguay, M., and Vaillancourt, P. A.: Recherche en Prévision Numérique contributions to numerical weather prediction,
605 *Atmosphere-Ocean*, 60, 35–64, <https://doi.org/10.1080/07055900.2022.2038071>, 2022.

606 Robert, A.: A stable numerical integration scheme for the primitive meteorological equations, *Atmosphere-Ocean*, 19, 35–46,
607 <https://doi.org/10.1080/07055900.1981.9649098>, 1981.

608 Sassi, M., Zhang, J., and Moran, M. D.: 2015 SMOKE-ready Canadian Air Pollutant Emission Inventory (APEI) package version
609 1 [Data set], <http://doi.org/10.5281/zenodo.4883639>, 2021.

610 Seidel, D. J., Zhang, Y., Beljaars, A., Golaz, J.-C., Jacobson, A. R., and Medeiros, B.: Climatology of the planetary boundary layer
611 over the continental United States and Europe, *J. Geophys. Res. Atmospheres*, 117, D17106,
612 <https://doi.org/10.1029/2012JD018143>, 2012.

613 Siebesma, A. P. and Cuijpers, J. W. M.: Evaluation of parametric assumptions for shallow cumulus convection, *J. Atmospheric*
614 *Sci.*, 52, 650–666, [https://doi.org/10.1175/1520-0469\(1995\)052<0650:EOPAFS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<0650:EOPAFS>2.0.CO;2), 1995.

615 Siebesma, A. P., Bretherton, C. S., Brown, A., Chlond, A., Cuxart, J., Duynkerke, P. G., Jiang, H., Khairoutdinov, M., Lewellen,
616 D., Moeng, C.-H., Sanchez, E., Stevens, B., and Stevens, D. E.: A large eddy simulation intercomparison study of shallow cumulus
617 convection, *J. Atmospheric Sci.*, 60, 1201–1219, [https://doi.org/10.1175/1520-0469\(2003\)60<1201:ALESIS>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)60<1201:ALESIS>2.0.CO;2), 2003.

618 Simon, H., Beck, L., Bhave, P. V., Divita, F., Hsu, Y., Luecken, D., Mobley, J. D., Pouliot, G. A., Reff, A., Sarwar, G., and Strum,
619 M.: The development and uses of EPA’s SPECIATE database, *Atmospheric Pollut. Res.*, 1, 196–206,
620 <https://doi.org/10.5094/APR.2010.026>, 2010.

621 Sørensen, B., Kaas, E., and Korsholm, U. S.: A mass-conserving and multi-tracer efficient transport scheme in the online integrated
622 Enviro-HIRLAM model, *Geosci. Model Dev.*, 6, 1029–1042, <https://doi.org/10.5194/gmd-6-1029-2013>, 2013.

623 Staniforth, A. and Côté, J.: Semi-Lagrangian integration schemes for atmospheric models—A review, *Mon. Weather Rev.*, 119,
624 2206–2223, [https://doi.org/10.1175/1520-0493\(1991\)119<2206:SLISFA>2.0.CO;2](https://doi.org/10.1175/1520-0493(1991)119<2206:SLISFA>2.0.CO;2), 1991.

625 Stroud, C. A., Morneau, G., Makar, P. A., Moran, M. D., Gong, W., Pabla, B., Zhang, J., Bouchet, V. S., Fox, D., Venkatesh, S.,
626 Wang, D., and Dann, T.: OH-reactivity of volatile organic compounds at urban and rural sites across Canada: Evaluation of air
627 quality model predictions using speciated VOC measurements, *Atmos. Environ.*, 42, 7746–7756,
628 <https://doi.org/10.1016/j.atmosenv.2008.05.054>, 2008.

629 Sundqvist, H., Berge, E., and Kristjánsson, J. E.: Condensation and cloud parameterization studies with a mesoscale numerical
630 weather prediction model, *Mon. Weather Rev.*, 117, 1641–1657, [https://doi.org/10.1175/1520-0493\(1989\)117<1641:CACPSW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1641:CACPSW>2.0.CO;2), 1989.

632 Van Wagner, C. E.: Development and structure of the Canadian Forest Fire Weather Index System, Forestry Technical Report 35,
633 Canadian Forestry Service, Ottawa, Ontario, 35 pp., <https://cfs.nrcan.gc.ca/publications?id=19927>, 1987.

634 Vosper, S. B., Wells, H., and Brown, A. R.: Accounting for non-uniform static stability in orographic drag parametrization: Non-
635 uniform stability in orographic drag, *Q. J. R. Meteorol. Soc.*, 135, 815–822, <https://doi.org/10.1002/qj.407>, 2009.

636 Wells, H., Vosper, S. B., Ross, A. N., Brown, A. R., and Webster, S.: Wind direction effects on orographic drag, *Q. J. R. Meteorol.*
637 *Soc.*, 134, 689–701, <https://doi.org/10.1002/qj.247>, 2008.

638 van de Wiel, B. J. H., Moene, A. F., Steeneveld, G. J., Hartogensis, O. K., and Holtslag, A. A. M.: Predicting the collapse of
639 turbulence in stably stratified boundary layers, *Flow Turbul. Combust.*, 79, 251–274, <https://doi.org/10.1007/s10494-007-9094-2>,
640 2007.

641 Zeng, X. and Beljaars, A.: A prognostic scheme of sea surface skin temperature for modeling and data assimilation: Sea surface
642 skin temperature scheme, *Geophys. Res. Lett.*, 32, L14605, 4 pp., <https://doi.org/10.1029/2005GL023030>, 2005.

643 Zhang, J., Moran, M. D., Zheng, Q., Makar, P. A., Baratzadeh, P., Marson, G., Liu, P., and Li, S.-M.: Emissions preparation and
644 analysis for multiscale air quality modeling over the Athabasca Oil Sands Region of Alberta, Canada, *Atmospheric Chem. Phys.*,
645 18, 10459–10481, <https://doi.org/10.5194/acp-18-10459-2018>, 2018.

646 Zheng, W., Ek, M., Mitchell, K., Wei, H., and Meng, J.: Improving the stable surface layer in the NCEP Global Forecast System,
647 *Mon. Weather Rev.*, 145, 3969–3987, <https://doi.org/10.1175/MWR-D-16-0438.1>, 2017.

648