

On the Mechanisms that Control the Rainy Season Transition Periods in the Equatorial Congo Basin

Sarah Worden^{1,2}, Rong Fu²

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, 91109, USA

5 ²Atmospheric and Oceanic Sciences Department, University of California Los Angeles, Los Angeles, 90095, USA

Correspondence to: Sarah Worden (sarah.r.worden@jpl.nasa.gov)

Abstract. The equatorial Congo Basin (2°S-2°N, within the watershed), experiences two rainy seasons (RSs) in boreal spring and fall. Previously, the mechanisms driving the transitions to these RSs have not been investigated systematically. We show that both RS transitions begin with increases in the low-level atmospheric moisture transport from the Atlantic Ocean into the region, ~1.5 months prior to the spring RS and ~2 months prior to the fall RS. Evapotranspiration contributes the most to atmospheric moisture but does not change significantly throughout both transition periods. Sharp precipitation jumps 10 days before the start of the spring and fall RSs result from increasing boundary layer moisture that is orographically uplifted to the lower troposphere by the East African Rift and Congo Basin Cell ascending branch. This destabilizes the lower free troposphere, lowering the level of free convection and decreasing convective inhibition energy. Meanwhile, the African Easterly Jet-North and westerly return flow of the Congo Basin Cell induce vertical shear for the spring and fall RSs, respectively. Mid-tropospheric convergence strengthens due to return flow at ~700 hPa from shallow meridional cells that direct low-level moisture towards the West African Heat Low prior to the spring RS, and to the Congo Air Boundary prior to the fall RS. Therefore, the RS onsets over the equatorial region are driven by seasonal changes in large-scale atmospheric circulation, contrasting with the pivotal role of increasing evapotranspiration in driving the transition to the RS onset over the southern Congo Basin.

© 2025. All rights reserved.

1 Introduction

The Congo Basin is a global water and carbon centre hosting vast rainforest, rivers, and precipitation (Brummett et al., 2009; Alsdorf et al., 2016; Xu et al., 2021). However, due to a lack of in situ data and limited infrastructure for research (Alsdorf et al., 2016), its meteorology and hydrological cycle have been less studied compared to the rest of Africa (Washington et al., 2013; Biasutti, 2019; Biasutti and Sobel, 2009; Lv et al., 2024; Seth et al., 2013; Yang et al., 2015; Thorncroft et al., 2011). Most studies examining process controls on seasonal rainfall have focused on specific months during the dry or rainy seasons (Nicholson, 2018; 2022; Cook and Vizy, 2022; Pokam Mba et al., 2022). As such, the processes that control the transition to the rainy season onsets are virtually unknown. To fundamentally understand observed rainfall variability (Jiang et al., 2019; Zhou et al., 2014; Hua et al., 2016; 2018) within the equatorial region (defined as the area between 2°S-2°N, within the Congo Basin watershed), which experiences two rainy seasons, in boreal spring and fall,

respectively (Fig. 1; Pokam Mba et al., 2022; Nicholson, 2022), necessitates a full understanding of the mechanisms controlling its seasonal rainfall. In addition, understanding the mechanisms controlling the transition periods to the rainy seasons can aid identification of errors in climate models for seasonal rainy season onset forecasting, which is important climate information for agricultural production (Zampieri et al., 2023; Tchinda et al., 2022).




Figure 1: (a) Above-ground biomass averaged between 2000-2019. Black outline denotes the Congo Basin watershed and the dashed lines at 2°S and 2°N represent the boundary of the equatorial region. (b) Precipitation climatology for the equatorial region. Shades are the standard error of the mean over of the different years.

The equatorial Congo Basin is unique in that its rainy seasons are not associated with strong near surface moisture flux convergence (Nicholson, 2018; Yang et al., 2015). Instead, near-surface moisture diverges away from the equatorial Congo basin, while mid-tropospheric moisture convergence appears to be central for rainfall formation (Nicholson, 2022). Atmospheric moisture for rainfall mainly comes from evapotranspiration (ET) both within Congo basin and from the adjacent land (e.g., Risi et al., 2016; Sori et al., 2022; van der Ent et al., 2010; Worden et al., 2021), though moisture transport from the surrounding oceans is important for its rainy seasons as well (e.g., Pokam et al., 2012). Several dynamic systems play important roles: a shallow zonal circulation referred to as the Congo Basin Cell (Longandjo and Rouault, 2020), the Congo Air Boundary (Howard and Washington, 2019) where warm, humid air from the Congo meets with relatively cool and dry air from the subtropical southern hemispheric Africa, the African Easterly Jet North and South (Nicholson, 2022; Chen 2005; Kuete et al., 2020), a deep Walker-like zonal circulation (Longandjo and Rouault, 2020), and shallow meridional overturning cell (Longandjo and Rouault, 2024). Additionally, the East African Rift helps to induce uplift of near surface air and hence precipitation on the Congo Basin's eastern boundary (Nicholson, 2022).

However, the moisture contributions from land and terrestrial sources, and the presence and intensity of these dynamical systems vary throughout the year, likely leading to differences in the mechanisms controlling the transitions to the spring and fall rainy seasons. For example, the annual presence of the zonal Congo Basin Cell is debated. While Longandjo and Rouault (2020) argue that the cell is present year-round, Neupane (2016) and Cook and Vizy (2016) indicate that it is

55 present only in the second half of the year. The Congo Air Boundary is strongest between August-October, and weakest between December-January (Howard and Washington, 2019). The AEJ-N is present year-round but moves latitudinally such that it is centred near the equator during the spring rainy season but centred outside of the Congo Basin around 10°N during the fall rainy season (Kouete et al., 2020; Nicholson and Grist, 2003). The AEJ-S only develops in the boreal fall in September (Kouete et al., 2020).

60 The influences of the heat lows over Sahara and Kalahari Desert, respectively, on the equatorial Congo region also change between the two rainy seasons, which is important for meridional moisture transport. During the fall rainy season, the southern African dry heat low is centred over the plateau region as the Angola heat low, but prior to the spring rainy season, a moist Angola thermal low migrates to the Kalahari region to become the Kalahari heat low (Howard and Washington, 2018; 2019; Attwood et al., 2024). Meanwhile, the northern hemisphere West African heat low is centred around the Sahelian region
65 during the spring rainy season (around $10^{\circ}\text{N} - 15^{\circ}\text{N}$). During the fall rainy season, it transitions back towards the northern Congo Basin (Lavaysse et al., 2009). The different position of the heat lows can affect meridional moisture flow in and out of the basin (Cook et al., 2020). The mechanisms controlling the transition periods to the rainy seasons are likely related to the seasonal presence or movement of these key features.

ET provides about 70% of the atmospheric moisture over the Congo basin (e.g., van de Ent et al. 2010; Risi et al.
70 2013; Worden et al. 2021), and as much as 83% of the atmospheric moisture in February and March, during the transition to spring rainy season. Worden and Fu (2025) further shows that, over the Southern Congo ($2^{\circ}\text{S} - 14^{\circ}\text{S}$, $15^{\circ}\text{E} - 30^{\circ}\text{E}$), the increase of ET driven by increasing surface solar radiation plays a pivotal role to increase moisture in the atmospheric boundary layer, which, in turn, increases deep convection and leads to rainy season onset in boreal fall. On the other hand, most previous studies emphasize increasing moisture transport as a potential key driver for the rainy season (e.g., Pokam et al. 2012;
75 Nicholson and Dezfuli, 2013; Balas et al., 2007). For the equatorial Congo basin, whether ET plays a similar role as over the Southern Congo basin has been unclear. In this study, we will investigate the roles of the aforementioned dynamic and thermodynamic processes in driving the transition to the spring and fall wet seasons, respectively, over the equatorial Congo basin.

2 Data and Methods

80 2.1 Data

Reanalysis datasets differ in terms of magnitude of seasonal rainfall and atmospheric circulation characteristics but generally agree on major features and seasonal evolution over Central Africa (Hua et al. 2019; Kenfack et al. 2023; Nicholson & Klotter 2021). The European Centre for Medium-Range Weather Forecasts 5th generation reanalysis, ERA (Hersbach et al. 2020), is commonly adopted as a primary reanalysis for Congo Basin studies (Cook & Vizy 2022; Kenfack et al. 2024;
85 Longandjo & Rouault 2020) and other African regions (Cook & Vizy 2022; Vizy & Cook 2019). Its relatively high resolution

helps capture topographic structure relevant to regional climate (Vizy & Cook 2019), and its precipitation is broadly consistent with recommended observational products for the basin (Nicholson et al. 2019; Cook & Vizy 2022).

90 We use satellite data and reanalysis products in our study. We resample all the daily data to 5-day pentads, except for the Tropospheric Emission Spectrometer (TES) data, in which we use daily HDO and H₂O measurements in this analysis instead of pentad measurements as the number of observations is limited compared to the other products. For all other datasets, we average our products over 2000-2018 or 2003-2018 depending on when the datasets are available.

95 The TES data requires quality control processing described here: the following quality flags were used when retrieving the data from the TES satellite: “Species Retrieval Quality = 1,” “Degrees of Freedom for Signal >1,” and “Average Cloud Optical Depth <0.4” to ensure good quality data as suggested by Worden et al. (2012). The accuracy of these data is ~6 per mil (6 parts per thousand relative to the reference value) with a precision of 20 per mil (Worden et al., 2012) for the vertical range used in this analysis (~900-420 hPa, or about 1-6 km above sea level). We use data between 2005-2011 when the quality of estimates is suitable for our analysis (Worden et al., 2021).

Table 1: Data used in this study

Source	Data	Resolution	Years	Reference
Tropical Rainfall Measuring Mission (TRMM) 3B42/3B43	Precipitation	0.25° × 0.25° Daily, Monthly	2000-2020	Huffman et al., 2007
Atmospheric Infrared Sounder (AIRS) L3 V7 Standard Physical Retrieval Ascending TqJoint	atmospheric-layer mass mixing ratio, surface mass mixing ratio, atmospheric-layer relative humidity, atmospheric-layer temperature	1° × 1° Daily	2003-2018	Aumann et al., 2019, Pagano et al., 2003, Irion et al., 2018, DeSouza-Machado et al., 2018
Clouds and the Earth's Radiant Energy System (CERES) Syn1deg Ed 4.1	cloud top temperature, cloud top height, cloud area fraction (total, high, mid-high, mid-low, and low)	1° × 1° Daily	2003-2018	Doelling, et al., 2013, Doelling et al., 2016, Su, et al., 2005, Rutan et al., 2015
Global Land Evaporation Amsterdam Model (GLEAM) v3.7b	Evapotranspiration	0.25 × 0.25° daily	2000-2018	Martens et al., 2017, Miralles et al., 2011
Ground plots, lidar, global modeling, and satellite data	Aboveground biomass	0.1 × 0.1° yearly	2000-2018	Xu et al., 2021
Tropospheric Emission Spectrometer version 6 (v006_Litev01.00)	Deuterium (<i>HDO</i>), water vapor (<i>H₂O</i>)	Level 2 daily	2005-2011	Worden et al., 2012, Worden et al., 2021
European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5	zonal and meridional (u and v) winds, vertically integrated moisture flux convergence, total column water vapor (CWV), vertical velocity, divergence, temperature, vertical integral of eastward total water vapour flux, vertical integral of northward total water vapour flux, and specific humidity	0.25° × 0.25° daily	2000-2018	Hersbach et al., 2020

2.2 Methods

100 2.2.1 Calculating the Onsets and Ends of the Rainy Seasons

Using TRMM precipitation data, we calculate the rainy season onsets and ends (RSOs and RSEs, respectively) in the equatorial Congo Basin similar to the approach used in Chakraborty et al. (2021), originally developed by Li and Fu (2004),

for RSOs and RSEs in the Amazon. This definition is designed to capture persistently high rainfall during the rainy season and persistent low rainfall in the dry season using the annual mean rainfall as the threshold. We first calculate the climatological pentad (5 day mean) rainfall for each year over our domain, then determine the RSOs and RSEs in the following steps.

For the spring rainy season, the onset is defined as the first pentad of each year that meets the following criterion: four out of the previous six pentads are less than the climatological annual mean, and four out of the following seven pentads are more than the climatological annual mean. Likewise, the end of the spring rainy season is defined as the first pentad of each year that meets the following criterion: four out of the previous six pentads are more than the climatological annual mean, and four out of the following six pentads are less than the climatological annual mean.




Figure 2: Rainy season onset (RSO) and rainy season end (RSE) for the spring and fall rainy seasons for each year in the equatorial Congo

For the fall rainy season, the onset is defined as the first pentad of each year that meets the following criterion: five out of the previous eight pentads are less than the climatological annual mean, and five out of the following eight pentads are more than the climatological annual mean. The fall RSE is defined as the first pentad of each year that meets the following

120 criterion: six out of the previous eight pentads are more than the climatological annual mean, and six out of the following eight pentads are less than the climatological annual mean. We note that this criterion for the fall rainy seasons differs from that of the spring rainy season because the latter is shorter and weaker than that of the fall rainy season (Fig. 2). Therefore, a shorter period and less persistent threshold was needed to best capture the starts and ends of persistently high and low rainfall for the spring RSO and RSE.

Figure 2 shows the RSOs and RSEs for the spring and fall rainy seasons between the years 2000-2020. We excluded the year 2000 when calculating the spring RSO because we needed to consider the ends of the prior years in our RSO calculations. Similarly, we excluded the year 2020 when calculating the fall RSE because we needed to consider the start of the next years in our RSE calculations. We additionally excluded from our following analyses any years during which a RSO or RSE was unable to be identified, or if the length of a rainy season was less than 5 pentads. The spring RSO ranges from the 6th to 14th pentad (with Days 1-5 defined as Pentad “0”), corresponding to early-February to mid-March. On average, the spring RSO occurs on 20 February-24 February. The spring RSE ranges from the 25th-32nd pentads, corresponding to mid-May to mid-June. On average, the spring RSE occurs on 21 May-25 May. The fall RSO ranges from the 40th-50th pentads, corresponding to mid-July to early-September. On average, the fall RSO occurs on 18 August-23 August. The fall RSE ranges from the 64th-71st pentads, corresponding to mid-November to late-December. On average, the fall RSE occurs on 2 December-6 December. We show 15 pentads prior to the spring RSO and 19 pentads prior to the fall RSO as these periods capture the key changes in processes that drive the dry to rainy season transition as in Li and Fu (2004) and Wright et al. (2017).

2.2.2 Computation of Key Variables

135 We calculate the area-mean equivalent potential temperature (θ_e) from AIRS gridded data at daily time resolution to examine atmospheric instability during and after the transition to the rainy season. We calculate θ_e using the following equation from Bolton (1980):

$$\theta_e = T_k \left(\frac{1000}{p} \right)^{0.2854(1-0.28 \times 10^{-3} * r)} \times \exp \left[\left(\frac{3.376}{T_L} - 0.00254 \right) \times r (1 + 0.81 \times 10^{-3} r) \right] \quad \text{Eq. 1}$$

140 Where T_k is the absolute temperature (K) at pressure level p , p is pressure (hPa), r is the mixing ratio (g/kg), T_L is the absolute temperature at the lifting condensation level (K) calculated by the following:

$$T_L = \frac{2840}{3.5 \ln T_{k0} - \ln e - 4.805} + 55 \quad \text{Eq. 2}$$

145 Where $e = \frac{p * r}{622 + r}$ is the water vapor pressure, and here T_{k0} is the absolute temperature at the surface, defined here as 925 hPa as the surface of the equatorial region is above sea level. We use the AIRS mass mixing ratio, temperature, and pressure in these calculations. We similarly calculate saturated equivalent potential temperature θ_{e_s} by letting $r = r_s$, where r_s is the saturated mixing ratio. We calculate r_s by the following:

We calculate e_s as in Bolton (1980):

$$e_s = 6.112 \exp\left(\frac{17.67T}{T+243.5}\right) \quad \text{Eq. 3}$$

Where T is the temperature in Celsius ($^{\circ}C$). We calculate saturated vapor pressure by:

$$r_s = \frac{\varepsilon e_s}{p - e_s} \quad \text{Eq. 4}$$

150 Where $\varepsilon = 0.622$ is ratio of molar masses of vapor and dry air, and e_s is the saturation vapor pressure, calculated as in Bolton (1980):

$$e_s(T) = 6.112 \exp\left(\frac{17.67T}{T+243.5}\right) \quad \text{Eq. 5}$$

We also calculate moist static energy (MSE) where MSE is represented by:

$$MSE = c_p T + Lq + gz. \quad \text{Eq. 6}$$

155 Where c_p is the specific heat capacity at constant pressure, L is the latent heat of evaporation and g is gravity.

Finally, we examine changes in moisture in the equatorial Congo with the following. We use vertically integrated moisture flux $MF = \int_{p_s}^{p_{TOA}} q \cdot \vec{V} dp / g$, where q is the specific humidity, and $\vec{V} = (\mathbf{u}, \mathbf{v})$, is the horizontal wind, with u the zonal, and v the meridional wind components, respectively. P_s denotes the surface pressure and P_{TOA} denotes the pressure of the top of the atmosphere. In particular, we calculate the net zonal and meridional MF as follows: We take the zonal vertically
 160 integrated $MF_{zonal} = \int_{p_s}^{p_{TOA}} q \cdot u dp / g$ across the western and eastern boundaries of the equatorial Congo basin, and the meridional vertically integrated $MF_{meridional} = \int_{p_s}^{p_{TOA}} q \cdot v dp / g$ across its northern and southern boundaries. For simplicity, we choose the northern boundary along $2^{\circ}N$ and between $14 - 30^{\circ}E$, and the southern boundary along $2^{\circ}S$ and between $14 - 30^{\circ}E$. The western boundary is along $14^{\circ}E$ and between $2^{\circ}S - 2^{\circ}N$ and the eastern boundary is along $30^{\circ}E$ and between $2^{\circ}S - 2^{\circ}N$. To be able to compare to vertically integrated MF convergence, we calculate the length between the
 165 western/eastern boundaries and the northern/southern boundaries, respectively, and integrate average MF_{zonal} and $MF_{meridional}$ across these boundaries as in Satyamurty et al. (2013):

$$Q_E = \int MF_{zonal,E} dy$$

$$Q_W = \int MF_{zonal,W} dy$$

$$Q_N = \int MF_{meridional,N} dx$$

$$170 \quad Q_S = \int MF_{meridional,S} dx$$

Where $MF_{zonal,E}$ is MF_{zonal} integrated along the eastern boundary, $MF_{zonal,W}$ is MF_{zonal} integrated along the western boundary, $MF_{meridional,N}$ is $MF_{meridional}$ integrated along the northern boundary, and $MF_{meridional,S}$ is $MF_{meridional}$ integrated along the southern boundary. y is the length of the northern/southern boundaries, and x is the length of the western/eastern boundaries. To calculate the net zonal vertically integrated MF, we calculate $Q_{zonal} = Q_W - Q_E$. To calculate

175 the net meridional vertically integrated MF, we calculate $Q_{meridional} = Q_S - Q_N$. This ensures that positive values of Q_{zonal} and $Q_{meridional}$ means net convergence, and negative means net divergence.

We calculate the first component of the moisture flux convergence term (named the moisture convergence term, $q * \nabla \cdot \vec{V}$) and moisture transport vectors ($q * \vec{V}$) averaged between different pressure levels to represent the lower and middle tropospheric moisture transport. We do not examine the evolution of moisture advection term ($\vec{V} * \nabla \cdot q$) as it is small compared
180 to that of the moisture convergence term (Cook and Vizu 2022; Fig. S1-S4).

2.2.3 Examining the ET Contribution to Atmospheric Moisture

Due to a lack of reliable ET measurements, we use the deuterium content of water (HDO) derived from satellite measurements and an isotope mixing model to estimate the fractional contribution of ET to atmospheric moisture. HDO is expressed as the relative ratio of the number of HDO molecules to the total number of H_2O molecules in parts per thousand
185 (%‰) relative to the isotopic composition of ocean water as shown below:

$$\delta D = 1000 \times \left(\frac{R - R_{std}}{R_{std}} \right) \quad \text{Eq. 8}$$

Where R is of the ratio of HDO molecules to the total number of H_2O molecules and R_{std} is the corresponding ratio in a reference standard, taken here to be the Vienna Standard Mean Ocean Water: $R_{std} = 3.11 \times 10^{-4}$ (e.g. Wright et al., 2017 and references therein). The isotopic composition of water vapor in the free troposphere is due to a mixture of air parcels
190 originating from different sources (Galewsky, 2018; Galewsky and Hurley, 2010). We can use the isotopic composition of an air mass to trace its source to either vegetation or ocean because δD values from ocean evaporation are distinctively different from those by rainforest ET. Atmospheric moisture coming from transpiration will generally be more isotopically enriched (or heavier) than evaporation because lighter isotopes preferentially evaporate (Risi et al., 2020; Tremoy et al., 2014; Worden et al., 2007). However, observed values of free-tropospheric deuterium content also depend on atmospheric processes, such as
195 the type of convection present, described further in Galewsky et al. (2016). This is because they vary by the first order with changes in specific humidity (Bailey et al., 2017).

Therefore, we compare the observed isotopic composition of water vapor in the free troposphere to two models, a mixing model and a Rayleigh model, to identify which air parcels are likely influenced by land (i.e., transpiration) versus ocean (i.e., evaporated water; Noone, 2012). A mixing model describes what happens to a mixture of two air masses with
200 different water vapor isotopic compositions. Meanwhile, under the Rayleigh distillation model, as an air mass moves upward (or towards cooler conditions), condensate enriched with heavy isotope is completely removed immediately after it forms under the assumption of pseudo adiabatic process (Galewsky et al., 2016; Wright et al., 2017). A further description of these models can be found in the Supplementary and in S. Worden et al. (2021).

2.2.4 Key Dynamic Features

205 We examine the seasonal evolution of key dynamic features that transport moisture and energy within and around the Congo Basin. They are discussed further in several review papers: Nicholson, (2022); Pokam et al. (2026).

- 1) Low Level Westerlies (LLWs): These winds transport moisture from the Atlantic Ocean into Central Africa, with varying seasonal intensity (Pokam et al., 2012; 2014; Vondou et al., 2010)
- 2) Congo Basin Cell: The LLWs are the lower branch of both shallow and deep zonal circulations found over the Congo Basin. The shallow zonal circulation is called the Congo Basin Cell, with ascent over the Congo Air Boundary (see below), return at mid-levels via an easterly jet, and subsidence over the Atlantic Ocean (Longandjo and Rouault 2020)
- 3) Shallow meridional overturning cell: Here, moisture moves meridionally from the Congo Basin to the Sahel heat low, with return leading to mid-tropospheric moisture convergence over the Congo Basin (Longandjo and Rouault 2024)
- 4) African Easterly Jet North and South (AEJ-N, AEJ-S): The AEJ-N is a year-round feature with seasonal meridional migration from 5°N-17°N, while the AEJ-S is present between September-November with its core at ~8°S (Nicholson and Grist 2003; Kuete et al., 2020)
- 5) Congo Air Boundary: An extreme gradient in temperature and specific humidity that denotes the southern limit of Congo Basin, present between August to November (Howard and Washington 2019).


3 Results

220 3.1 Transition from the Dry Season to the Spring RSO

3.1.1 Early Transition

The first part of the spring transition, i.e., the early-transition period, starts with a turning point from strengthening to weakening vertically integrated MF divergence (Fig. 3a) 10 pentads (50 days) before the spring RSO. To explain this weakening, we evaluate Q_{zonal} and $Q_{meridional}$ (Fig. 3c). Negative values indicate net moisture leaving the basin for both Q_{zonal} and $Q_{meridional}$. Variations in Q_{zonal} during the early transition period are driven by increases in low-level moisture transport from the Atlantic Ocean, by the lower branch of the Congo Basin Cell, which goes through the equatorial Congo, towards the West African Heat Low (Lavaysse et al., 2009) in the northern Congo (Fig. 4a, b; Fig. B1a, e). This is driven by increasing near-surface temperature gradients between the Atlantic Ocean and West African Heat Low (Fig. S5a, b; Fig. S6). These increases in low-level moisture transport, combined with increases in the second half of the transition period of mid-level moisture transport into the basin across its eastern boundary, acts against zonal moisture leaving the basin at mid-levels, across the western boundary (Fig. B1a, b), but is not enough to lead to Q_{zonal} convergence (Fig. 3c). Meridionally, mid-level moisture entering the basin across the northern boundary does not change significantly, although low-level moisture leaving the basin across the northern boundary increases in response to increased low-level moisture transport towards the West African Heat Low (Fig. 4a, b; Fig. B1c, f). Across the southern boundary, moisture transport entering the basin at low levels

235 increases, while moisture transport leaving the basin at mid-levels does not significantly change (Fig. B1d, f). Ultimately, it is changes in low-level meridional moisture transport that drive changes in $Q_{meridional}$ divergence (Fig. 3c).



240 **Figure 3: For the transition to the spring RSO: (a) Precipitation (P), vertically integrated MF convergence (MFC), and evapotranspiration (ET); (b) column water vapor (CWV); and (c) Net vertically integrated MF across the zonal and meridional boundaries (Q_{zonal} , $Q_{meridional}$; Methods, Units: kg/s*1e7). Positive means convergence, negative means divergence. All are relative to the RSO (denoted as ‘0’ in the graph). Time series are smoothed using a Savitsky-Golay filter (number of coefficients: 5, polynomial order 2). Shades represent the standard error of the mean of the different years.**

245 Meanwhile, the contribution of ET to atmospheric moisture increases somewhat compared to the pre-transition ($f_{mix_{pre}} = 0.611 \pm 0.093$; $f_{mix_{early}} = 0.772 \pm 0.12$), and is clearly the dominant contributor to moisture (Fig. 5a, b). This is consistent with S. Worden et al., (2021) for a similar domain. Therefore, ET is important for maintaining high background moisture while weakening in vertically integrated MF divergence initiates the transition to the spring RSO.

Furthermore, the atmosphere appears thermodynamically primed for deep convection, with large convective available potential energy (CAPE) and conditional instability as shown by $\frac{d\theta_e}{dp} < 0$ (Fig. 6c, d). However, weak ascent or even subsidence still exists above 650 hPa (Fig. 6e). Additionally, the level of free convection (LFC; Fig. 7a), and therefore convective inhibition energy (CIN, Fig. 6c) do not change significantly throughout the early transition period. This indicates that the atmospheric conditions, especially the dynamic conditions, are still not ready for large-scale increases of deep convection and subsequently, rainfall.




Figure 4: Moisture transport vectors ($q * v$) and moisture convergence term ($q * DIV$; shades, positive means convergence) for (a)-(c) average of 925-875 hPa for the “before” (15-11 pentads prior to the spring RSO), early, and late transition periods, respectively; (d)-(f) average of 850-600 hPa for the “before,” early, and late transition periods, respectively. Vectors are scaled up by a factor of 2 to better visualize the changes in magnitude and direction between periods. White masks in (a)-(c) represent areas where the elevation is higher than 875 hPa. Aqua contours in (a)-(c) represent sharp 925 hPa specific humidity gradients, which overlap with the area of convergence.

We explain this as following: with weakening atmospheric vertically integrated MF divergence, boundary layer specific humidity begins to increase (Fig. 7d), due to increases in moisture transport from the equatorial Atlantic Ocean into the equatorial Congo (Fig. 4a, b). The increase of boundary layer moisture, rather than increase of near-surface temperature, drives boundary layer increases in MSE (Fig. 7b-d). However, at the mid-levels, MSE initially decreases and then increases in mid-level specific humidity (Fig. 7b), which is driven by mid-level moisture divergence. Therefore, while the boundary layer MSE is favourable for deep convection, additional changes are needed to support deep convection in the mid-layers of the atmosphere.





Figure 5: Fraction of observed δD above a series of mixing (solid) and Rayleigh (dashed) models. Blue shades represent a density map of the TES δD observations. Green indicates land-based water vapor models, while black indicates ocean-based water vapor models. f_{mix} is the fraction of observed δD above the uppermost, land-based mixing

270 **model. For (a) the “before” period (15-11 pentads prior to the spring RSO); (b) the early-transition; and (c) the late-transition.**

3.1.2 Late Transition Period

The late-transition period starts with a rapid increase in atmospheric column water vapor (CWV; Fig. 3b), a key condition for deep convection (Bretherton et al., 2004; Schiro et al., 2018). This is accompanied by rapid increases in precipitation and vertically integrated MF convergence (Fig. 3a). ET does not change significantly ($f_{mixlate} = 0.778 \pm 0.139$; Fig. 5c), revealing the importance of large-scale circulation changes in altering atmospheric moisture over the equatorial Congo prior to the spring rainy season.



280 **Figure 6:** Prior to the spring RSO: (a) cloud area fraction for high, mid-high, mid-low, and low clouds; (b) cloud top temperature and cloud top height; (c) CAPE; and CIN; (d) equivalent potential temperature (θ_e ; shades) and saturated equivalent potential temperature ($\theta_{e,s}$); (e) vertical velocity; and (f) black contours: 650 hPa zonal wind (u_{650}) greater than $6.5 m s^{-1}$, denoting the AEJ-N. Shaded contours represent the wind shear ($u_{900} - u_{650}$). Shades of the time series in a)-c)

represent the standard error of the mean of the different years. All the time series have been smoothed using a Savitsky-Golay filter (number of coefficients: 5, polynomial order 2).

Q_{zonal} and $Q_{meridional}$ rapidly switch from divergence towards convergence (Fig. 3c). Zonally, further increases in
 285 near surface temperatures over the West African Heat Low drive stronger LLWs from the Atlantic and across the equatorial
 region along the way (Fig. B1a; Fig. S5c; Fig. S6; Fig. 4c). This, together with increases in mid-level (above 800 hPa) zonal
 moisture transport across the eastern boundary (Fig. B1b) compensate the increases in mid-level moisture transport leaving
 the basin across the western boundary (Fig. B1a). The latter is likely due to the formation of the AEJ-N (Fig. 6f). Therefore,
 Q_{zonal} becomes convergent, although weakly so (Fig. 3c). Meridionally, moisture transport entering the basin at low levels
 290 through the southern boundary increases (Fig. B1c), bringing with it moist air from the southern Congo (Fig. S7c), which is
 experiencing its rainy season during this time (Worden and Fu, 2025). However, moisture transport leaving the basin across
 its northern boundary towards the West African Heat Low increases as well (Fig. B1d), leading to net low level meridional
 moisture transport out of the basin (Fig. 4c). In contrast, moisture convergence at mid-levels increases (Fig. 4f) as mid-level
 moisture leaving the basin across the southern boundary decreases (Fig. B1d, f). Altogether, this leads to $Q_{meridional}$
 295 convergence (Fig. 3c).




Figure 7: Evolution of (a) the LFC; (b) MSE at 925 hPa and 700 hPa; (c) MSE component $C_p T$ 925 hPa and 700 hPa; and (d) MSE component $L_v q$ at 925 hPa and 700 hPa. All are relative to the spring RSO, averaged between $14^\circ E - 26^\circ E$ to avoid topography errors in the calculations, and smoothed using the Savitsky-Golay filter. Shades represent the standard error of the mean of the different years.

These changes in atmospheric moisture transport increase boundary layer moisture, as indicated by increasing relative humidity (Fig. 8a, b), but subsidence from the surface to ~ 850 hPa (Fig. 6e; Fig. 8a, b) between $10^{\circ}\text{E} - 26^{\circ}\text{E}$ (west of the East African Rift) likely prevents this moisture from being uplifted within large parts of the interior equatorial region. Instead increases in westerly wind within the interior of the region pushes this moisture towards the East African Rift (Fig. 8), where orographic lifting forces this moist air into the free troposphere above 800 hPa. Additionally, moisture is uplifted by the ascending branch of the Congo Basin Cell over the equatorial coastline (highlands of Gabon and Cameroon) at the western edge of the equatorial Congo (Fig. 8). Therefore, this results in increasing mid-level moisture and mid-level MSE while decreasing LFC and CIN (Fig. 6c; Fig. 7a, b, d). Upward vertical velocity increases in strength starting in the mid-levels and reaches 300 hPa for the first time (Fig. 6e). Meanwhile, shear increases with the increasing strength of the AEJ-N (Fig. 6f), thus creating conditions favourable for deep convection and the start of the spring rainy season. This is indicated by increases in high clouds, increases in cloud top height, and decreases in cloud top temperature (Fig. 6a, b).

310





Figure 8: Relative humidity (shades) and vectors of u and $\omega * 10^2$ for (a) 10-3 pentads prior to the spring RSO; (b) 2-0 pentads prior to the spring RSO. Zonal wind (u) for (c) 10-3 pentads prior to the spring RSO; (d) 2-0 pentads prior to the spring RSO. Grey shading in (a)-(b) and white shading in (c)-(d) denotes the topography of the region.

3.2.1 Early Transition Period



320 **Figure 9: For the transition to the fall RSO: (a) Precipitation (P), vertically integrated MF convergence (MFC), and evapotranspiration (ET); (b) column water vapor (CWV); (c) Q_{zonal} and $Q_{meridional}$ (Methods; Units: $\text{kg/s} \cdot 1e7$). Positive means convergence, negative means divergence. All are relative to the RSO (denoted as '0' in the graph). All have been smoothed using a Savitsky-Golay filter (number of coefficients: 5, polynomial order 2). Shades represent the standard error of the mean of the different years.**

As in the transition to the spring RSO, the early-transition period starts 14 pentads before the fall RSO with weakening in vertically integrated MF divergence over the equatorial Congo (Fig. 9a), indicated reduced moisture export out of the region. This is mainly due to decreases in Q_{zonal} divergence (Fig. 9c), which becomes convergent about halfway through the early transition period. This acts against the change of $Q_{meridional}$, which starts convergent and switches to divergence during the transition period (Fig. 9c).




Figure 10: Moisture transport vectors ($q * v$) and moisture convergence ($q * DIV$; shades, positive means convergence) for (a)-(c) average of 925-875 hPa. The westerly winds represent the lower branch inflow of the Congo Basin Cell; the aqua shades denote areas of sharp 925 hPa specific humidity gradients. The southern aqua gradients therefore note the rough location of the Congo Air Boundary. (d-f) average of 850-800 hPa; and (g-i) average of 775-600 hPa. The easterly winds represent the upper branch return flow of the Congo Basin Cell. Vectors have been multiplied by a scale factor of 2 in all panels. Area of convergence between $4^{\circ}S$ and $8^{\circ}S$ in a-c) represents the rough location of the Congo Air Boundary (CAB). Mainly zonal moisture transport between $6^{\circ}N - 15^{\circ}N$ represents the AEJ-N.

Changes in initial Q_{zonal} divergence are primarily due to increases in low-level moisture transport into the basin, across the western boundary (Fig. 10a, b; Fig. B2a), which combined with steady moisture transport between 800-600 hPa into the basin across the eastern boundary (Fig. B2b), acts against weakly decreasing moisture transport out of the western boundary above 800 hPa (Fig. B2a) via the return branch of the Congo Basin Cell (Longandjo et al., 2020). Increases in low-level atmospheric moisture across the western boundary are driven by the near-surface temperature gradient between the Atlantic Ocean and the equatorial and southern Congo (Fig. S8a, b; Fig. S9), which strengthens the lower branch of the Congo Basin Cell (Longandjo and Rouault, 2020).





Figure 11: Fraction of observed δD above a series of mixing (solid) and Rayleigh (dashed) models. Blue shades represent a density map of the TES δD observations. Green indicates land-based water vapor models, while black indicates ocean-based water vapor models. f_{mix} is the fraction of observed δD above the uppermost, land-based mixing

model. For (a) the “before” period (19-14 pentads prior to the fall RSO); (b) the early-transition; and (c) the late-transition.


350 $Q_{meridional}$ convergence initially weakens towards zero (Fig. 9c). Across the southern boundary, moisture transport leaving the basin at low levels increases (Fig. B2d, f), moving towards the Congo Air Boundary (aqua contours in Fig. 10b denote sharp surface humidity gradients that show the rough location of the Congo Air Boundary; Howard and Washington, 2019) and associated low-level convergence band (Fig. 10b). Additionally, moisture transport into the equatorial Congo across the southern boundary, above 850 hPa, decreases in strength (Fig. 10d, e). This acts against increases in low-level moisture entering the basin across the northern boundary (Fig. B2d). Together, this causes the $Q_{meridional}$ divergence by the end of the transition period.



355 **Figure 12: Prior to the fall RSO: (a) cloud area fraction for high, mid-high, mid-low, and low clouds; (b) cloud top temperature and cloud top height; (c) CAPE; and CIN; (d) equivalent potential temperature (θ_e ; shades) and saturated**

equivalent potential temperature (θ_{e_s}); (e) vertical velocity; and (f) black contours: 650 hPa zonal wind greater than 6.5 m/s, denoting the AEJ-N. Shaded contours represent the wind shear ($u_{900} - u_{650}$). Shades of the time series in (a)-(c) represent the standard error of the mean of the different years. All the time series have been smoothed using a Savitsky-Golay filter (number of coefficients: 5, polynomial order 2).

Meanwhile, the contribution of ET to atmospheric moisture does not change significantly: $f_{mix} = 0.612 \pm 0.082$ during the “before” period, while $f_{mix} = 0.665 \pm 0.168$ during the early transition (Fig. 11 a, b). Therefore, it is the increases of low-level moisture from the Atlantic Ocean that initiate the early transition period prior to the fall RSO.



365 **Figure 13: Evolution of (a) LFC, (b) MSE at 925 hPa and 700 hPa; (c) $C_p T$ at 925 hPa and 700 hPa; and (d) $L_v q$ at 925 hPa and 700 hPa. All are prior to the fall rainy season. All are averaged between $14E^\circ - 26E^\circ$ to account for effects of topography in the eastern part of the region. All have been smoothed using a Savitsky-Golay filter. Shades represents the standard error of the mean of the different years.**

Despite an overall unstable lapse rate ($\frac{d\theta_e}{dp} < 0$), the thermodynamic condition of tropospheric column does not yet support deep convection. In the first half of the early transition, decreases in low-level atmospheric moisture (Fig. 13d) weakens negative $\frac{d\theta_e}{dp}$, coinciding with rising CIN and LFC, as well as decreasing CAPE and low and mid-level MSE (Fig. 12c, d; Fig. 13b, c). Changes in MSE at both levels are mainly due to changes in $L_v q$ rather than $C_p T$ (Fig. 13b, c, d). In the second half of the early transition, low-level moisture begins to increase, corresponding to increases in θ_e in the low-levels and strengthening

negative (destabilizing) $\frac{d\theta_e}{dp}$ (Fig. 12c; Fig. 13d). This corresponds to the LFC stabilizing (no longer consistently increasing) between -7 to -2 pentads prior to the RSO (Fig. 13a) and decreases in CIN (Fig. 12c). Additionally, the vertical velocity is near zero above 600 hPa (Fig. 12e). Therefore, atmospheric dynamic conditions are not yet ready to facilitate increase of deep convection, consistent with relatively low and constant mid-high and high cloud covers (Fig. 12a), cloud top temperatures, and cloud top heights (Fig. 12b).




Figure 14: The zonal and vertical cross-section of relative humidity (shades) and vectors of u and ω averaged between $2^{\circ}S$ - $2^{\circ}N$ for (a) the early transition period, i.e., 14-3 pentads prior to the fall RSO; (b) the late transition period, i.e., 2-0 pentads prior to the fall RSO. The vertical wind, ω , is magnified by multiple 100. As in (a)-(b), but for zonal wind speed (u) for (c) 14-3 pentads prior to the fall RSO; and d) 2-0 pentads prior to the fall RSO. Grey mask in (a)-(b) and white mask in (c)-(d) denote the topography of the region.

3.2.2 Late Transition Period

The late transition period to the fall rainy season starts with sharp increases in precipitation, vertically integrated MF convergence, and CWV (Fig. 9a, b). Meanwhile, the contribution of ET to atmospheric moisture ($f = 0.613 \pm 0.194$) does not change significantly (Fig. 11c). Q_{zonal} convergence sharply increases, while $Q_{meridional}$ divergence weakens to near zero (Fig. 9c).

Zonally, the speed of low-level westerly wind that carry moisture from the Atlantic Ocean further increases (Fig. 10b,c). This is due to further increases in the zonal temperature gradient as the Atlantic cold tongue develops (Fig. S8b, c; Fig.

390 S9; Caniaux et al., 2011), which increases the strength of the lower branch of the Congo Basin Cell (Longandjo and Rouault, 2020). Meanwhile, the strength of the mid-level zonal moisture transport does not change as much compared to its low-level counterpart, indicating that the sharp increase in Q_{zonal} convergence is driven by increases low-level moisture transport (Fig. 9c; Fig. B2a, b, e).

Meridionally, in the low-levels, moisture transport out of the southern boundary into the southern Congo continues
395 to increase as the low-level winds move towards the Congo Air Boundary (Fig. 10b, c). However, this is compensated by moisture transport into the equatorial Congo across the southern boundary above 850 hPa (Fig. 10e, f, h, g; Fig. B2d). Across the northern boundary, low level moisture transported into the basin increases, acting against mid level moisture leaving the basin (Fig. B2c). Therefore, $Q_{meridional}$ divergence decreases (Fig. 9c).

At the low-levels, atmospheric moisture increases rapidly due to increases in westerly winds (Fig. 14) and further
400 destabilizes (strengthens negative $\frac{d\theta_e}{dp}$; Fig. 12d). The increased atmospheric moisture lifts up into the mid-troposphere above 800 hPa via the ascending branch of the Congo Basin Cell (located between $10^{\circ}E - 20^{\circ}E$) as well as orographically along the slope of the East African Rift (Fig. 14a, b), Therefore, mid-tropospheric relative humidity increases (Fig. 14b) despite subsidence in the low troposphere in the interior of the basin (Fig. 12e; Fig. 14b). As the mid-troposphere moistens, the thermodynamic atmospheric condition becomes favourable for deep convection rapidly, as shown by decreases in CIN (Fig.
405 12c) and LFC (Fig. 13a), and increases in CAPE (Fig. 12c).

Meanwhile, despite the absence of the AEJ-N or AEJ-S over the equatorial region (Fig. 12f), the return branch of the Congo Basin Cell provides the vertical wind shear (Fig. 12f, on average within the equatorial region $8.30 m s^{-1}$ during the late transition period prior to the fall RSO compared to $8.55 m s^{-1}$ during the late transition period prior to the spring RSO) needed for favourable dynamic conditions for mesoscale convective systems (MCSs). These conditions enable rapid increase
410 of P (Fig. 9a), and lead to the fall RSO as indicated by increases in the strength of vertical velocity from the lower atmosphere to 300 hPa (Fig. 12e), sharp increases (decreases) in high/mid-high (low) clouds, sharp increases in cloud top height, and sharp decreases in cloud top temperatures (Fig. 12a, b).

In summary, the transition from the boreal summer dry season to the fall RSO is mainly initiated by increasing westerly moisture transport from Atlantic Ocean into the equatorial Congo basin in the boundary layer via the lower branch of
415 the Congo Basin Cell (925-875 hPa). These changes start 14 pentads (around mid-June) before the fall RSO (around late August). This weakens the vertically integrated MF divergence, which, in turn, slows down the drying trend during the dry season. Low-level moisture increases due to increases in moisture transport from the Atlantic Ocean during the early transition (14-3 pentads before the fall RSO). Mid-level atmospheric moisture increases as a result of increased uplift via the ascending branch of the Congo Basin Cell and orographic uplift against the East African Rift during the late transition period. This
420 increases CWV, which subsequently reduces CIN and lows the LFC, while increasing CAPE, MSE and vertical wind shear. Subsequently, the fall rainy season begins.

4 Discussion

4.1 Comparing the Transition to the Spring vs Fall Rainy Seasons

Here, we compare how conditions evolve to trigger the spring and fall rainy seasons, clarifying the similarities and
425 differences in the controlling processes and their interactions that lead to each season's onset.

Prior to the spring rainy season, low-level moisture from the equatorial region is transported to the West African Heat
Low, mixes with warm air from the Sahelian region (Fig. 4c; Fig. S5), uplifts, and then returns to the equatorial Congo (Fig.
4f) at the mid-levels (850-600 hPa). Therefore, a shallow meridional overturning cell exists that moves moisture northward at
low levels, and southward at mid-levels. In contrast, prior to the fall rainy season, low-level moisture from the Atlantic Ocean
430 turns towards the Congo Air Boundary located in the southern Congo. This low-level moisture transport into the equatorial
Congo across the western boundary is greater in the fall rainy season compared to the spring rainy season, consistent with
Pokam et al. (2012) who found that the Atlantic Ocean plays an important role in low-level moisture flux between August-
November. As this low-level moisture reaches the Congo Air Boundary, it mixes with warm air from the southern Congo
Basin (Fig. S8e, f) and South Africa (Worden and Fu 2025), uplifts, and then returns to the equatorial Congo (Fig. 10d-i)
435 Therefore, a shallow meridional overturning cell exists that moves moisture southward at low levels, and northward at mid-
levels.

Furthermore, the Congo Basin Cell moves further into the interior of the basin prior to the fall RSO compared to the
spring RSO, with its ascending branch located around $20^{\circ}E$ prior to the fall and located around $15^{\circ}E$ prior to the spring rainy
season (Fig. 8a, b; Fig. 14a, b). Therefore, less moisture is brought into the region across its western boundary prior to the
440 spring (Fig. B1a; Fig. B2a) and prior to the fall. Finally, the cause of the lower to middle tropospheric wind shear differs
between the two rainy seasons. The zonal shear increases prior to the start of the rainy season is induced by the formation of
the AEJ-N, present in the equatorial region. Meanwhile, prior to the fall rainy season, shear exists due to the return branch of
the Congo Basin Cell, at similar pressure levels, but it is weaker compared to the wind shear prior to the spring RSO.

Additionally, the thermodynamic conditions are more favourable for convection prior to the fall RSO than the spring
445 RSO. CIN prior to the spring rainy season ranges from 226-308 J/Kg and is 226 J/kg 1 pentad prior to RSO, while CIN prior
to the fall rainy season ranges from 149-222 J/kg and is at 149 J/kg 1 pentad prior to the RSO. The LFC is also higher by the
beginning of the spring RSO compared to the fall RSO. Therefore, the cloud base is higher, and convective air must overcome
stronger CIN prior to the spring RSO.

Finally, both transition periods show increasing low-level atmospheric moisture before increases in mid-level
450 atmosphere, and the moistening of the mid-level during late transition seasons that is important for increasing deep convection
and initiating both RSOs. The increase of low-level zonal moisture transport is mainly responsible for increasing the lower-
level moisture, whereas the increase of mid-level is main due to lifting of the low-level moisture to the mid-level by ascending
branch of the Congo Basin Cell over the western equatorial Congo basin and by the East African rift over the eastern equatorial
Congo basin.

455 4.2 Comparing the Transition to the Fall Rainy Season in the Equatorial vs Southern Congo

We next compare the transition period to the fall rainy season in the equatorial versus southern Congo. The fall RSO is earlier over the equatorial Congo basin, beginning on average 18 August, than that over the southern Congo, beginning on average 9 October (Worden and Fu, 2025). This difference in timing is likely driven, in part, by the seasonal movement of the Congo Air Boundary (Longandjo and Rouault, 2020), which is centred closer to the equator prior to the equatorial fall RSO (Fig. 10b, c) and then moves towards the southern Congo prior to its fall RSO (Worden and Fu, 2025). This southward movement brings with it moisture from the equatorial region, supporting the southern Congo RSO in boreal fall. Furthermore, moisture contributions between the two regions differ greatly. ET is the dominant contributor to atmospheric moisture prior to the fall rainy season in the equator; however, it does not change significantly during the transition period. Meanwhile, for the southern region, ET increases significantly and contributes equally to atmospheric moisture as the oceanic moisture transport by the late transition period. Therefore, the increases of moisture prior to the RSOs in the equatorial region are primarily controlled by advected oceanic moisture, while over the southern Congo, moisture increases are controlled by increases of both moisture transport from the ocean and from ET over land.

Additionally, atmospheric instability differs strongly. In the equatorial region, CAPE is much higher than CIN for the entire transition period, helping initiate a large jump in precipitation once CIN decreases as there is plenty of energy for deep convection. However, in the southern region, CAPE is lower, and CIN is higher than in the equatorial Congo, where above 700 hPa is dominated by adiabatic warming due to subsidence until the late transition period. Therefore, the equatorial region has larger potential for storms and heavy rain to occur than in the southern Congo basin.

Finally, the AEJs play different roles between the equatorial region and the southern Congo. In the equatorial region, both the AEJ-S and AEJ-N do not play a direct role in determining vertical wind shear prior to the fall RSO. In contrast, the AEJ-S directly contributes to the vertical wind shear prior to the RSO in the southern Congo. Therefore, it is important to consider the sub-basin scale variability in the evolution of atmospheric conditions prior to the transition to the RSOs that will likely react differently to climate perturbations and hence the seasonal rainfall cycle.

5 Conclusion

Using a suite of satellite and reanalysis data, we examined the mechanisms controlling the transition periods to the
480 spring and fall rainy season in the equatorial Congo. We summarize these mechanisms below.




Figure 15: Summary of the key systems controlling moisture transport over the equatorial Congo Basin prior to (a) the spring RSO and (b) the fall RSO.

Spring rainy season onset:

- 485 (1) *Early-transition*: This period starts with a turning point from strengthening to weakening of the vertically integrated MF divergence. Changes in both Q_{zonal} and $Q_{meridional}$ lead to near zero vertically integrated MF divergence by the end of this period. This is due to (1) increases in low-level moisture entering the basin from the Atlantic Ocean; and (2) initial decreases and then increases in mid-level moisture entering the basin from the Indian Ocean and leaving the basin back towards the Atlantic Ocean. Changes in $Q_{meridional}$ are due to increases in low-level moisture leaving the basin across the northern boundary towards the West African heat low level moisture entering the basin across the southern boundary. ET contributes the most to atmospheric moisture but does not change significantly. Meanwhile, large CAPE, and negative $\frac{d\theta_e}{dp}$, but large CIN and high LFC, as well as the presence of subsidence in the upper troposphere, indicates that thermodynamic conditions are beginning to change in favour of deep convection.
- 490
- (2) *Late-transition*: This period starts with sharp increases in CWV, indicating atmospheric moistening, vertically integrated MF convergence, and precipitation. Increases in low-level moisture transport into the region from the Atlantic Ocean and mid-level transport into the region from the Indian Ocean act against moisture leaving the region at mid-levels back towards the Atlantic Ocean. Low-level moisture is lifted to the mid-troposphere by the ascending branch of the Congo Basin Cell on the western edge of the region, and by orographic uplift in the eastern part of the region as moisture is pushed up the East African Rift. Meridionally, more moisture transport enters the region across the northern boundary than leaves the basin across the southern boundary, supporting mid-level MF convergence (most important feature for predicting precipitation; Table 2). Atmospheric moisture increases, causing decreases in CIN (second-most important feature for predicting precipitation; Table 2) and lowering of the LFC. Additionally, the AEJ-N moves over the equatorial region, providing vertical wind shear. This all leads to sharp increases in precipitation. That ET contribution to atmospheric moisture does not change significantly indicates that it provides background moisture but does not contribute to
- 500
- 505 transitioning the atmospheric conditions to be conducive for deep convection.

Fall rainy season onset:

- (1) *Early Transition*: This period starts from a turning point from increasing to weakening vertically integrated MF divergence, which reaches convergence about halfway through the period. These changes are primarily due to large increases in Q_{zonal} convergence, as increases in low-level moisture transport from the Atlantic Ocean, due to the strengthening of the lower-branch of the Congo Basin Cell (Longandjo and Rouault, 2020), act against mid-level moisture transport out of the region back towards the Atlantic Ocean. Therefore, low-level moisture increases in the second half of the transition period, increasing CWV and strengthening the negative $\frac{d\theta_e}{dp}$ gradient. However, as mid-level atmospheric moisture decreases to a minimum, atmospheric thermodynamic conditions are not yet ready to support deep convection, as CIN is high and the LFC is elevated.
- 510
- (2) *Late Transition*: CWV, vertically integrated MF convergence, and precipitation increase sharply. ET remains the same, but it contributes an equal amount of atmospheric moisture for rainfall as in the early transition period. Q_{zonal} convergence increases as low-level moisture from the Atlantic Ocean increases. $Q_{meridional}$ divergence goes to near zero despite
- 515

520 increases in low-level moisture transport towards the Congo Air Boundary across the southern boundary. Low-level moisture is pushed upwards by the ascending branch of the Congo Basin Cell, located more centrally in the region compared to prior to the spring RSO, as well as up against the East African Rift. This, along with increases in mid-level meridional MF convergence (moisture transport entering the basin is greater than moisture transport leaving the basin above 850 hPa) supports mid-level MF convergence (most important feature for predicting precipitation; Table 2) and moistens the mid-troposphere, lowering CIN and the LFC. The westerly return branch of the Congo Basin Cell provides the vertical shear needed. Therefore, conditions are conducive for the start of the rainy season.

525 **Table 2: Summary of changes to key features controlling the rainy season transition periods for the spring and fall rainy seasons. A ridge regression was performed to show which features are most important to controlling P during the transition period. The R^2 represents the performance of the test dataset (30% of data withheld when training the model).**

Spring Early Transition Phase	Spring Late Transition Phase
$P_{sprRSO} = 0.45MF_{mid-level} - 0.27CIN + 0.16MF_{low-level} + 0.014Shear - 0.005MSE_{low-level} - 0.002CAPE + 3.51$ $R^2 = 0.58$	
$MF_{low-level}$ ($q.DIV$ averaged between 925-875 hPa): Convergence decreasing. Moisture transport from Atlantic Ocean, through equatorial Congo, towards West African Heat Low.	$MF_{low-level}$: Convergence decreasing. Moisture transport from Atlantic Ocean, through equatorial Congo, towards West African Heat Low.
$MF_{mid-level}$ ($q.DIV$ averaged between 850-600 hPa): Convergence increasing. Moisture transport from Indian Ocean, through equatorial Congo, towards Atlantic Ocean.	$MF_{mid-level}$: Convergence increasing. Zonal Moisture transport from Indian ocean, through equatorial Congo, towards Atlantic Ocean.
CIN: Steady	CIN: Decreasing
CAPE: Increasing	CAPE: Increasing
$MSE_{low-level}$ (averaged between 925-875 hPa): Increasing	$MSE_{low-level}$: Increasing
Shear: Steady	Shear: Increasing , from return branch of Congo Basin Cell
Precipitation: Steady	Precipitation: Increasing
Fall Early Transition Phase	Fall Late Transition Phase

$P_{fallRSO} = 0.24MF_{mid-level} + 0.13MSE_{low-level} + 0.10CAPE + 0.09MF_{low-level} - 0.06CIN - 0.014Shear + 3.49$	
$R^2 = 0.43$	
$MF_{low-level}$: Convergence increasing. Moisture transport switch from northeast direction to southeast direction: from Atlantic Ocean, through equatorial Congo, towards Congo Air Boundary	$MF_{low-level}$: Convergence increasing. Moisture transport from Atlantic Ocean, through equatorial Congo, towards Congo Air Boundary.
$MF_{mid-level}$: Convergence increasing. Moisture transport from Southern Congo and Indian Ocean, through equatorial Congo, towards Atlantic Ocean.	$MF_{mid-level}$: Convergence increasing. Moisture transport from Southern Congo and Indian ocean, through equatorial Congo, towards Atlantic Ocean.
CIN: Decreasing	CIN: Decreasing
CAPE: Decreasing	CAPE: Increasing
$MSE_{low-level}$: Decreasing	$MSE_{low-level}$: Increasing
Shear: Steady	Shear: Steady , from return branch of Congo Basin Cell
Precipitation: Steady	Precipitation: Increasing

The clarification of the atmospheric processes that drive the dry to rainy season transition provided by this study will lay the groundwork for understanding variability in the seasonal cycle of precipitation. This includes changes induced by ENSO, the Indian Ocean Dipole, and Madden-Julian Oscillations (e.g., Kebacho and Sarfo, 2023; Moihamette et al., 2022; Jury et al., 2009; Creese et al., 2019), variability induced by aerosols (Chakraborty et al., 2020), by the poleward movement of the northern and southern heat lows (Cook et al., 2020), and by the long-term AMJ drought (Jiang et al., 2019). Overall, the equatorial region seasonal rainfall is influenced by a combination of local (e.g., ET providing background moisture, and orographic uplift) and regional factors (e.g., moisture from the Atlantic Ocean, the West African Heat Low, and the Congo Air Boundary). Therefore, it is important to consider the influence of climate variability on the rainy season timing and intensity on both scales.

6 Appendices

6.1 Appendix A

The following text further explains the comparison of the observed δD measurements to the mixing and Rayleigh models, as well as the error calculations involved:

A mixing model is described as follows:

$$\delta_{mix} = q_0(\delta_0 - \delta_F) \frac{1}{q} + \delta_F \tag{Eq. A1}$$

Where q_0 and δ_0 are, respectively, the specific humidity and its δD value of the dry air mass in the upper troposphere, and q_F and δ_F are, respectively, the specific humidity and δD value of the air mass at the surface. $q = q_0 + q_F$ is the specific humidity of the mixed air mass between dry air mass in the upper troposphere and humid air mass from its surface source (Noone, 2012).

We also examine the observed δD values in relation to a Rayleigh model, which describes the change of δD with water vapor mixing rate as liquid water evaporates in equilibrium with temperature:

$$\delta_{ray} \simeq (\alpha - 1) \ln\left(\frac{q}{q_0}\right) + \delta_0 \quad \text{Eq. A2}$$

where α is set to equal the temperature-dependent equilibrium fractionation factor between liquid and water vapor (Majoube, 1971). Further discussion of δD and these models can be found in (Worden et al., 2021).

Following Worden et al. (2021), we compare the observed δD measurements to the mixing and Rayleigh models to identify the relative contribution of ET to atmospheric moisture. The initial values of modeled δD (δ_F) were -50% and -80% for land-based and ocean-based models, respectively, chosen based on representative values of δD from land and ocean. Any observed δD above the land-based mixing model is considered as transpired by the plants over land. Any δD value shown between the land-based and ocean-based mixing models likely comes from some mixture of the two sources. The fraction of atmospheric moisture coming from ET is represented by the fraction of observed δD above the land-based mixing model (f_{mix}). This fraction can be considered a lower bound on the amount of ET contributing to atmospheric moisture because rainfall processes as well as vapor originating from oceans or evaporation and mixing with vapor from ET will decrease δD . To calculate the error on this fraction, we assume a 12% error in the δD observations, and recalculate the fraction of δD above the land-based mixing model with 12% added and subtracted from the observations as in Worden et al. (2021). We then find the differences between the original fraction and the new fractions and average the two to calculate our final error.





Figure B1: (a) Zonal moisture transport ($q * u$) across $15^\circ E$ and between $2^\circ S - 2^\circ N$; (b) zonal moisture transport ($q * v$) across $30^\circ E$ and between $2^\circ S - 2^\circ N$. Note that contours below 875 hPa should not be considered as they are below the East African Rift, but all pressure levels are shown to emphasize the mid-level moisture transport. (c) meridional

570 moisture transport ($q * v$) across $2^{\circ}N$ and between $15^{\circ}E - 30^{\circ}E$; d) meridional moisture transport ($q * v$) across $2^{\circ}S$
and between $15^{\circ}E - 30^{\circ}E$; e) zonal moisture transport at 925 hPa (orange) and 700 hPa (purple) across the western
and eastern boundaries; and f) meridional moisture transport at 925 hPa (green) and 700 hPa (grey) across the
575 northern and southern boundaries. For (a) and (b), positive values equal eastward transport and negative equal
westward transport. For (c) and (d), positive equals northward transport and negative values equal southward
transport. Additionally, for longitude values greater than $26^{\circ}E$, meridional moisture transport values have been
masked out between 925-875 hPa to account for the topography of the East African Rift. Therefore, shades below 875
hPa represent meridional moisture transport between $15^{\circ}E - 26^{\circ}E$. Prior to the spring RSO.



580 **Figure B2:** a) zonal moisture transport ($q * u$) across $15^\circ E$ and between $2^\circ S - 2^\circ N$; b) zonal moisture transport ($q * v$) across $30^\circ E$ and between $2^\circ S - 2^\circ N$. Note that contours below 875 hPa should not be considered as they are below the East African Rift, but all pressure levels are shown to emphasize the mid-level moisture transport. c) meridional moisture transport ($q * v$) across $2^\circ N$ and between $15^\circ E - 30^\circ E$; and d) meridional moisture transport ($q * v$) across $2^\circ S$ and between $15^\circ E - 30^\circ E$; e) zonal moisture transport at 925 hPa (orange) and 700 hPa (purple) across the western and eastern boundaries; and f) meridional moisture transport at 925 hPa (green) and 700 hPa (grey) across the

585 northern and southern boundaries. For a) and b), positive values equal eastward transport and negative values equal westward transport. For c) and d), positive values equal northward transport and negative values equals southward transport. Additionally, for longitude values greater than $26^{\circ}E$, meridional moisture transport values have been masked out between 925-875 hPa to account for the topography of the East African Rift. Therefore, shades below 875 hPa represent meridional moisture transport between $15^{\circ}E - 26^{\circ}E$. Prior to the fall RSO.

7 Code Availability

590 Code can be made available by request to the corresponding author (Sarah Worden, sarah.r.worden@jpl.nasa.gov)

8 Data Availability

GLEAM evapotranspiration estimates can be downloaded via the GLEAM website: <https://www.gleam.eu/>. Version 3.7b can be made available by request to the corresponding author Sarah Worden (sarah.r.worden@jpl.nasa.gov). All AIRS TqJoint data and TRMM 3b42 daily precipitation can be downloaded using the NASA GES DISC: <https://disc.gsfc.nasa.gov>.
605 ERA5 products can be found at: <https://cds.climate.copernicus.eu/cdsapp#!/home>. TES HDO and H2O estimates can be found at: <https://catalog.data.gov/dataset/tes-aura-l2-hdo-limb-v006>. CERES cloud and products can be found at: <https://ceres-tool.larc.nasa.gov/ord-tool/jsp/SYN1degEd41Selection.jsp>

9 Author Contribution

All authors contributed to the study conception and design. S.W. performed experimental design, data collection, analysis, and manuscript writing. R.F. performed experimental design and manuscript writing. All authors read and approved the final manuscript.
600

10 Competing Interests

The authors declare that they have no conflict of interest.

11 Acknowledgments

605 Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004). © 2025. All rights reserved.

12 Financial Support

This work was supported by the NASA FINNEST grant #80NSSC20K1654 and NSF Grant Number 1917781.

13 References

- 610 Alsdorf, D., Beighley, E., Laraque, A., Lee, H., Tshimanga, R., O'Loughlin, F., et al. (2016). Opportunities for hydrologic research in the Congo Basin. *Reviews of Geophysics*, 54(2), 378–409. <https://doi.org/10.1002/2016RG000517>
- Attwood, K., Washington, R., & Munday, C. (2024). The Southern African Heat Low: Structure, Seasonal and Diurnal Variability, and Climatological Trends. *Journal of Climate*. <https://doi.org/10.1175/JCLI-D-23-0522.1>
- 615 Aumann, H. H., Broberg, S., Manning, E., & Pagano, T. (2019). Radiometric Stability Validation of 17 Years of AIRS Data Using Sea Surface Temperatures. *Geophysical Research Letters*, 46, 12504–12510. <https://doi.org/10.1029/2019GL085098>.
- Balas, N., Nicholson, S. E., & Klotter, D. (2007). The relationship of rainfall variability in West Central Africa to sea-surface temperature fluctuations. *International Journal of Climatology*, 27(10), 1335–1349. <https://doi.org/10.1002/joc.1456>
- Bolton, D. (1980). The computation of equivalent potential temperature. *Monthly weather review*, 108(7), 1046-1053. [https://doi.org/10.1175/1520-0493\(1980\)108<1046:TCOEPT>2.0.CO;2](https://doi.org/10.1175/1520-0493(1980)108<1046:TCOEPT>2.0.CO;2)
- 620 Bretherton, C. S., Peters, M. E., & Back, L. E. (2004). Relationships between water vapor path and precipitation over the tropical oceans. *Journal of Climate*, 17(7), 1517–1528. [https://doi.org/10.1175/1520-0442\(2004\)017<1517:RBWVPA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<1517:RBWVPA>2.0.CO;2)
- 625 Brummett, R.E., C. Tanania, A. Pandi, J. Ladel, Y. Munzimi, A. Russell, et al. (2009). Ressources en eau et biens et services liés à l'écosystème forestier. In: Wasseige, C., Devers, D., de Marcen, P., Eba'a Atyi, R., Nasi, R., & Mayaux, P. (Eds.) Les forêts du Bassin du Congo. Office des publications de l'Union européenne, Luxembourg. 10.2788/32456.
- Caniaux, G., Giordani, H., Redelsperger, J. L., Guichard, F., Key, E., & Wade, M. (2011). Coupling between the Atlantic cold tongue and the West African monsoon in boreal spring and summer. *Journal of Geophysical Research: Oceans*, 116(C4). <https://doi.org/10.1029/2010JC006570>
- 630 Chakraborty, S., Jiang, J. H., Su, H., & Fu, R. (2020). Deep convective evolution from shallow clouds over the Amazon and Congo rainforests. *Journal of Geophysical Research: Atmospheres*, 125(1), e2019JD030962. <https://doi.org/10.1029/2019JD030962>
- 635 Chakraborty, S., Jiang, J. H., Su, H., & Fu, R. (2021). On the role of aerosol radiative effect in the wet season onset timing over the Congo rainforest during boreal autumn. *Atmospheric Chemistry and Physics*, 21(17), 12855-12866. <https://doi.org/10.5194/acp-21-12855-2021>.
- Chen, T. C. (2005). Maintenance of the midtropospheric North African summer circulation: Saharan high and African easterly jet. *Journal of climate*, 18(15), 2943-2962. <https://doi.org/10.1175/JCLI3446.1>
- 640 Cook, K. H., Liu, Y., & Vizy, E. K. (2020). Congo Basin drying associated with poleward shifts of the African thermal lows. *Climate Dynamics*, 54, 863-883. <https://doi.org/10.1007/s00382-019-05033-3>
- Cook, K. H., & Vizy, E. K. (2016). The Congo Basin Walker circulation: Dynamics and connections to precipitation. *Climate Dynamics*, 47(3–4), 697–717. <https://doi.org/10.1007/s00382-015-2864-y>
- Cook, K. H., & Vizy, E. K. (2022). Hydrodynamics of regional and seasonal variations in Congo Basin precipitation. *Climate Dynamics*, 59(5-6), 1775-1797. <https://doi.org/10.1007/s00382-021-06066-3>

- 645 Creese, A., Washington, R., & Jones, R. (2019). Climate change in the Congo Basin: processes related to wetting in the December–February dry season. *Climate Dynamics*, 53, 3583-3602. <https://doi.org/10.1007/s00382-019-04728-x>
- Cook, K. H., Liu, Y., & Vizy, E. K. (2020). Congo Basin drying associated with poleward shifts of the African thermal lows. *Climate Dynamics*, 54(1), 863-883. <https://doi.org/10.1007/s00382-019-05033-3>
- 650 DeSouza-Machado, S., Strow, L. L., Tangborn, A., Huang, X., Chen, X., Liu, X., Wu, W., and Yang, Q. (2018). Single-footprint retrievals for AIRS using a fast TwoSlab cloud-representation model and the SARTA all-sky infrared radiative transfer algorithm. *Atmos. Meas. Tech.*, 11, 529–550. <https://doi.org/10.5194/amt-11-529-2018>
- Doelling, D. R., N. G. Loeb, D. F. Keyes, M. L. Nordeen, D. Morstad, C. Nguyen, B. A. Wielicki, D. F. Young, M. Sun. (2013). Geostationary Enhanced Temporal Interpolation for CERES Flux Products, *Journal of Atmospheric and Oceanic Technology*, 30(6), 1072-1090. doi: 10.1175/JTECH-D-12-00136.1.
- 655 Doelling, D. R., M. Sun, L. T. Nguyen, M. L. Nordeen, C. O. Haney, D. F. Keyes, P. E. Mlynchzak. (2016). Advances in Geostationary-Derived Longwave Fluxes for the CERES Synoptic (SYN1deg) Product, *Journal of Atmospheric and Oceanic Technology*, 33(3), 503-521. doi: 10.1175/JTECH-D-15-0147.1
- Galewsky, J. (2018). Using stable isotopes in water vapor to diagnose relationships between lower-tropospheric stability, mixing, and low-cloud cover near the island of Hawaii. *Geophysical Research Letters*, 45(1), 297–
- 660 305. <https://doi.org/10.1002/2017GL075770>
- Galewsky, J., & Hurley, J. V. (2010). An advection-condensation model for subtropical water vapor isotopic ratios. *Journal of Geophysical Research*, 115(D16), D16116. <https://doi.org/10.1029/2009JD013651>
- Hart, N. C., Washington, R., & Maidment, R. I. (2019). Deep convection over Africa: Annual cycle, ENSO, and trends in the hotspots. *Journal of Climate*, 32(24), 8791-8811. <https://doi.org/10.1175/JCLI-D-19-0274.1>
- 665 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., et al. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. <https://doi.org/10.1002/QJ.3803>
- Hua, W., Zhou, L., Chen, H., Nicholson, S. E., Raghavendra, A., & Jiang, Y. (2016). Possible causes of the Central Equatorial African long-term drought. *Environmental Research Letters*, 11(12). <https://doi.org/10.1088/1748-9326/11/12/124002>
- 670 Hosking, J. S., Russo, M. R., Braesicke, P., & Pyle, J. A. (2010). Modelling deep convection and its impacts on the tropical tropopause layer. *Atmospheric Chemistry and Physics*, 10(22), 11175-11188. <https://doi.org/10.5194/acp-10-11175-2010>
- Howard, E., & Washington, R. (2018). Characterizing the synoptic expression of the Angola low. *Journal of Climate*, 31(17), 7147-7165. <https://doi.org/10.1175/JCLI-D-18-0017.1>
- Howard, E., & Washington, R. (2019). Drylines in southern Africa: Rediscovering the Congo air boundary. *Journal of Climate*, 32(23), 8223-8242. <https://doi.org/10.1175/JCLI-D-19-0437.1>
- 675 Hua, W., Zhou, L., Nicholson, S. E., Chen, H., & Qin, M. (2019). Assessing reanalysis data for understanding rainfall climatology and variability over Central Equatorial Africa. *Climate Dynamics*, 53(1–2), 651–669. <https://doi.org/10.1007/s00382-018-04604-0>

- 680 Huffman, G. J., Bolvin, D. T., Nelkin, E. J., Wolff, D. B., Adler, R. F., Gu, G., et al. (2007). The TRMM multisatellite precipitation analysis (TMPA): Quasi-global, multiyear, combined-sensor precipitation estimates at fine scales. *Journal of Hydrometeorology*, 8(1), 38–55. <https://doi.org/10.1175/JHM560.1>
- Irion, F. W., Kahn, B. H., Schreier, M. M., Fetzer, E. J., Fishbein, E., Fu, D., Kalmus, P., Wilson, R. C., Wong, S., and Yue, Q. (2018). Single-footprint retrievals of temperature, water vapor and cloud properties from AIRS. *Atmos. Meas. Tech.*, 11, 971–995. <https://doi.org/10.5194/amt-11-971-2018>.
- 685 Jiang, Y., Zhou, L., Tucker, C. J., Raghavendra, A., Hua, W., Liu, Y. Y., & Joiner, J. (2019). Widespread increase of boreal summer dry season length over the Congo rainforest. *Nature Climate Change*, 9, 617–622. <https://doi.org/10.1038/s41558-019-0512-y>
- Jury, M. R., Matari, E., & Matitu, M. (2009). Equatorial African climate teleconnections. *Theoretical and applied climatology*, 95, 407-416. <https://doi.org/10.1007/s00704-008-0018-4>
- 690 Kebacho, L. L., & Sarfo, I. (2023). Why Eastern Africa was not dry during the 2020 short rainy season despite La Niña and a negative Indian Ocean Dipole: Interplay between the Madden-Julian Oscillation and La Niña in modulating short rain. *Theoretical and Applied Climatology*, 153(3), 1191-1201. <https://doi.org/10.1007/s00704-023-04519-9>
- Kuete, G., Pokam Mba, W., & Washington, R. (2020). African Easterly Jet South: control, maintenance mechanisms and link with Southern subtropical waves. *Climate Dynamics*, 54(3-4), 1539-1552. <https://doi.org/10.1007/s00382-019-05072-w>
- 695 Lacour, J.-L., Risi, C., Worden, J., Clerbaux, C., & Coheur, P.-F. (2018). Importance of depth and intensity of convection on the isotopic composition of water vapor as seen from IASI and TES δ D observations. *Earth and Planetary Science Letters*, 481, 387–394. <https://doi.org/10.1016/j.epsl.2017.10.048>
- Lavaysse, C., Flamant, C., Janicot, S., Parker, D. J., Lafore, J. P., Sultan, B., & Pelon, J. (2009). Seasonal evolution of the West African heat low: a climatological perspective. *Climate Dynamics*, 33, 313-330. <https://doi.org/10.1007/s00382-009-0553-4>.
- 700 Li, W., & Fu, R. (2004). Transition of the large-scale atmospheric and land surface conditions from the dry to the wet season over Amazonia as diagnosed by the ECMWF re-analysis. *Journal of Climate*, 17(13), 2637-2651. [https://doi.org/10.1175/1520-0442\(2004\)017<2637:TOTLAA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2637:TOTLAA>2.0.CO;2)
- 705 Longandjo, G. N. T., and Rouault, M. (2020). "On the Structure of the Regional-Scale Circulation over Central Africa: Seasonal Evolution, Variability, and Mechanisms". *Journal of Climate* 33.1, 145-162. <https://doi.org/10.1175/JCLI-D-19-0176.1>
- Longandjo, G. N. T., & Rouault, M. (2024). Revisiting the Seasonal Cycle of Rainfall over Central Africa. *Journal of Climate*. <https://doi.org/10.1175/JCLI-D-23-0281.1>
- Majoube, M. (1971). Fractionnement en oxygène 18 et en deutérium entre l'eau et sa vapeur. *Journal de Chimie Physique*, 68, 1423–1436. <https://doi.org/10.1051/jcp/1971681423>
- 710 Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., et al. (2017). GLEAM v3: satellite-based land evaporation and root-zone soil moisture. *Geosci. Model Dev.*, 10, 1903–1925. <https://doi.org/10.5194/gmd-10-1903-2017>.

- Mba, W.P., Vondou, D.A. & Kamsu-Tamo, P.H. (2022). Central African Climate. In Congo Basin Hydrology, Climate, and Biogeochemistry (eds R.M. Tshimanga, G.D.M. N'kaya and D. Alsdorf). <https://doi.org/10.1002/9781119657002.ch2>.
- 715 Moihamette, F., Pokam, W. M., Diallo, I., & Washington, R. (2022). Extreme Indian Ocean dipole and rainfall variability over Central Africa. *International Journal of Climatology*, 42(10), 5255-5272. <https://doi.org/10.1002/joc.7531>
- Neupane, N. (2016). The Congo Basin zonal overturning circulation. *Adv. Atmos. Sci.*, 33, 767–782, <https://doi.org/10.1007/s00376-015-5190-8>.
- Nicholson, S. E. (2018). The ITCZ and the seasonal cycle over equatorial Africa. *Bulletin of the American Meteorological Society*, 99(2), 337–348. <https://doi.org/10.1175/BAMS-D-16-0287.1>
- 720 Nicholson, S.E. (2022). The Rainfall and Convective Regime over Equatorial Africa, with Emphasis on the Congo Basin. In Congo Basin Hydrology, Climate, and Biogeochemistry (eds R.M. Tshimanga, G.D.M. N'kaya and D. Alsdorf). <https://doi.org/10.1002/9781119657002.ch3>.
- Nicholson, S. E., & Dezfuli, A. K. (2013). The relationship of rainfall variability in western equatorial Africa to the tropical oceans and atmospheric circulation. Part I: The boreal spring. *Journal of Climate*, 26(1), 45–65. <https://doi.org/10.1175/JCLI-D-11-00653.1>
- 725 Nicholson, S. E., & Grist, J. P. (2003). The seasonal evolution of the atmospheric circulation over West Africa and equatorial Africa. *Journal of climate*, 16(7), 1013-1030. [https://doi.org/10.1175/1520-0442\(2003\)016<1013:TSEOTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<1013:TSEOTA>2.0.CO;2)
- Noone, D. (2012). Pairing measurements of the water vapor isotope ratio with humidity to deduce atmospheric moistening and dehydration in the tropical midtroposphere. *Journal of Climate*, 25(13), 4476–4494. <https://doi.org/10.1175/JCLI-D-11-00582.1>
- 730 Pagano, T. S., Aumann, H. H., Hagan, D. E., & Overoye, K. (2003). Prelaunch and in-flight radiometric calibration of the Atmospheric Infrared Sounder (AIRS). *IEEE transactions on geoscience and remote sensing*, 41(2), 265-273. 10.1109/TGRS.2002.808324
- 735 Pokam, W. M., Bain, C. L., Chadwick, R. S., Graham, R., Sonwa, D. J., & Kamga, F. M. (2014). Identification of processes driving low-level westerlies in West Equatorial Africa. *Journal of Climate*, 27(11), 4245–4262. <https://doi.org/10.1175/JCLI-D-13-00490.1>
- Pokam, W. M., Djiotang, L. A. T., & Mkankam, F. K. (2012). Atmospheric water vapor transport and recycling in Equatorial Central Africa through NCEP/NCAR reanalysis data. *Climate Dynamics*, 38(9–10), 1715–1729. <https://doi.org/10.1007/s00382-011-1242-7>
- 740 Pokam, W., Nicholson, S., Vondou, D., Kuete, G., Longandjo, G. N., Guenang, M., ... & Fotso-Nguemo, T. (2026). Climate of the Congo Basin. In *Resilience and Sustainability in the Congo Basin: Retracing the Past, Looking to the Future* (pp. 1-32). Cham: Springer Nature Switzerland. https://doi.org/10.1007/978-3-032-02023-9_3-1
- Risi, C., Muller, C., & Blossey, P. (2020). What controls the water vapor isotopic composition near the surface of tropical oceans? Results from an analytical model constrained by large-eddy simulations. *Journal of Advances in Modeling Earth Systems*, 12(8). <https://doi.org/10.1029/2020MS002106>
- 745

- Risi, C., Noone, D., Frankenberg, C., & Worden, J. (2013). Role of continental recycling in intraseasonal variations of continental moisture as deduced from model simulations and water vapor isotopic measurements. *Water Resources Research*, 49(7), 4136–4156. <https://doi.org/10.1002/wrcr.20312>
- 750 Rutan, D. A., S. Kato, D. R. Doelling, F. G. Rose, L. T. Nguyen, T. E. Caldwell, and N. G. Loeb. (2015): CERES synoptic product: Methodology and validation of surface radiant flux. *J. Atmos. Oceanic Technol.*, 32, 1121–1143, doi: 10.1175/JTECH-D-14-00165.1.
- Satyamurty, P., da Costa, C. P. W., & Manzi, A. O. (2013). Moisture source for the Amazon Basin: a study of contrasting years. *Theoretical and Applied Climatology*, 111(1), 195-209. <https://doi.org/10.1007/s00704-012-0637-7>
- 755 Schiro, K. A., Ahmed, F., Giangrande, S. E., & David Neelin, J. (2018). GoAmazon2014/5 campaign points to deep-inflow approach to deep convection across scales. *Proceedings of the National Academy of Sciences of the USA*, 115(18), 4577–4582. <https://doi.org/10.1073/pnas.1719842115>
- Sohn, B. J., Choi, M. J., & Ryu, J. (2015). Explaining darker deep convective clouds over the western Pacific than over tropical continental convective regions. *Atmospheric Measurement Techniques*, 8(11), 4573-4585. [https://doi.org/10.5194/amt-8-](https://doi.org/10.5194/amt-8-4573-2015)
760 4573-2015
- Sorí, R., Stojanovic, M., Nieto, R., Liberato, M.L. & Gimeno, L. (2022). Spatiotemporal variability of droughts in the Congo River Basin: The role of atmospheric moisture transport. In *Congo Basin Hydrology, Climate, and Biogeochemistry* (eds R.M. Tshimanga, G.D.M. N'kaya and D. Alsdorf). 10.1002/9781119657002.ch11.
- Su, W., T. P. Charlock, and F. G. Rose. (2005): Deriving surface ultraviolet radiation from CERES surface and atmospheric radiation budget: Methodology. *J. Geophys. Res.*, 110(D14209), doi: 10.1029/2005JD005794.
765
- Tchinda, A. F., Tanessong, R. S., Mamadou, O., & Orou, J. B. C. (2022). Assessing precipitation seasonal forecasts in Central Africa using North American Multimodel Ensemble (NMME) Assessing precipitation seasonal forecasts in Central Africa using North American Multimodel Ensemble (NMME). *Theoretical and Applied Climatology*, 147(3), 1309-1325. <https://doi.org/10.1007/s00704-021-03915-3>
- 770 Thorncroft, C. D., Nguyen, H., Zhang, C., & Peyrillé, P. (2011). Annual cycle of the West African monsoon: regional circulations and associated water vapour transport. *Quarterly Journal of the Royal Meteorological Society*, 137(654), 129-147. <https://doi.org/10.1002/qj.728>
- Tremoy, G., Vimeux, F., Soumana, S., Souley, I., Risi, C., Favreau, G., & Oï, M. (2014). Clustering mesoscale convective systems with laser-based water vapor $\delta^{18}\text{O}$ monitoring in Niamey (Niger). *Journal of Geophysical Research*, 119(9), 5079–5103. <https://doi.org/10.1002/2013JD020968>
775
- van der Ent, R. J., Savenije, H. H. G., Schaeffli, B., & Steele-Dunne, S. C. (2010). Origin and fate of atmospheric moisture over continents. *Water Resources Research*, 46(9). <https://doi.org/10.1029/2010WR009127>
- Vondou, D. A., Nzeukou, A., Lenouo, A., & Mkankam Kamga, F. (2010). Seasonal variations in the diurnal patterns of convection in Cameroon–Nigeria and their neighboring areas. *Atmospheric Science Letters*, 11(4), 290-300.
780 <https://doi.org/10.1002/asl.297>
- Washington, R., James, R., Pearce, H., Pokam, W. M., & Moufouma-Okia, W. (2013). Congo basin rainfall climatology: Can we believe the climate models? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1625). <https://doi.org/10.1098/rstb.2012.0296>

- 785 Worden, J., Kulawik, S., Frankenberg, C., Payne, V., Bowman, K., Cady-Peirara, K., et al. (2012). Profiles of CH₄, HDO, H₂O, and N₂O with improved lower tropospheric vertical resolution from Aura TES radiances. *Atmospheric Measurement Techniques*, 5(2), 397–411. <https://doi.org/10.5194/amt-5-397-2012>
- Worden, J., Noone, D., Bowman, K., Beer, R., Eldering, A., Fisher, B., et al. (2007). Importance of rain evaporation and continental convection in the tropical water cycle. *Nature*, 445(7127), 528–532. <https://doi.org/10.1038/nature05508>
- 790 Worden, J., Saatchi, S., Keller, M., Bloom, A., Liu, J., Parazoo, N., et al. (2021). Satellite observations of the tropical terrestrial carbon balance and interactions with the water cycle during the 21st century. *Reviews of Geophysics*, 59, e2020RG000711. <https://doi.org/10.1029/2020RG000711>.
- Worden, S., Fu, R., Chakraborty, S., Liu, J., & Worden, J. (2021). Where Does Moisture Come From Over the Congo Basin?. *Journal of Geophysical Research: Biogeosciences*, 126(8), e2020JG006024. <https://doi.org/10.1029/2020JG006024>
- 795 Wright, J. S., Fu, R., Worden, J. R., Chakraborty, S., Clinton, N. E., Risi, C., et al. (2017). Rainforest-initiated wet season onset over the southern Amazon. *Proceedings of the National Academy of Sciences of the USA*, 114(32), 8481–8486. <https://doi.org/10.1073/pnas.1621516114>
- Xu, L., Saatchi, S., Yang, Y., Yu, Y., Pongratz, J., Bloom, A., et al. 2021. Changes in global terrestrial live biomass over the 21st century. *Science Advances*, 7(27): eabe9829. 18 p. <https://doi.org/10.1126/sciadv.abe9829>.
- 800 Yang, W., Seager, R., Cane, M. A., & Lyon, B. (2015). The annual cycle of East African precipitation. *Journal of Climate*, 28, 2385–2404. <https://doi.org/10.1175/JCLI-D-14-00484.1>
- Zampieri, M., Toreti, A., Meroni, M., Bojovic, D., Octenjak, S., Marcos-Matamoros, R., ... & Hoteit, I. (2023). Seasonal forecasts of the rainy season onset over Africa: Preliminary results from the FOCUS-Africa project. *Climate Services*, 32, 100417.
- 805 Zhang, Z., Xu, C. Y., El-Tahir, M. E. H., Cao, J., & Singh, V. P. (2012). Spatial and temporal variation of precipitation in Sudan and their possible causes during 1948–2005. *Stochastic environmental research and risk assessment*, 26, 429–441. <https://doi.org/10.1007/s00477-011-0512-6>
- Zhou, L., Tian, Y., Myneni, R. B., Ciais, P., Saatchi, S., Liu, Y. Y., et al. (2014). Widespread decline of Congo rainforest greenness in the past decade. *Nature*, 508(7498), 86–90. <https://doi.org/10.1038/nature13265>