

21st century change in precipitation on the Greenland Ice Sheet using high resolution regional climate models

Fredrik Boberg¹, Xavier Fettweis², Nicolaj Hansen¹, Ruth Mottram¹, Michiel R. van den Broeke³

- 5 ¹National Centre for Climate Research, Danish Meteorological Institute (DMI), Copenhagen, Denmark
²Laboratory of Climatology, SPHERES research unit, Geography Department, University of Liège, Liège, Belgium
³Institute for Marine and Atmospheric Research (IMAU), Utrecht University, Utrecht, The Netherlands

Correspondence to: Fredrik Boberg (fbo@dmu.dk)

10

Abstract. An ensemble of regional climate model simulations downscaling global climate models within Coupled Model Intercomparison Project Phase 6 (CMIP6) is used to estimate future precipitation changes for the Greenland ice sheet under a range of climate change pathways. The regional climate models are HIRHAM5, MAR3.12 and RACMO2.3p2 contributing a total of eleven simulations for the SSP5-8.5 scenario, five simulations for the SSP2-4.5 scenario and four simulations for the
15 SSP1-2.6 scenario. The Greenland ice sheet is divided into six drainage basins to evaluate the change in total (snow plus rain) precipitation in regions with different precipitation characteristics. Compared with the reference period 1971–2000, the ensemble median change in precipitation for the full Greenland ice sheet for the SSP5-8.5 scenario is estimated to be about +60 Gt (+8%) per year during the 2031–2060 period and +170 Gt (+24%) per year during the 2071–2100 period. We see similar ensemble median change for the 2031–2060 period for the SSP1-2.6 and SSP2-4.5 scenarios while the 2071–2100
20 change is +40 and +80 Gt (+6 and +11%) per year for SSP1-2.6 and SSP2-4.5, respectively. Using linear regression on the annual mean change in near-surface (2 m) air temperature and precipitation over the ice sheet, we estimate an increase of about 35 Gt per year in precipitation (equal to about 5%) for every degree of warming during the 21st century. The change in precipitation phase shows a non-linear increase in rainfall with temperature, particularly along the outer edge and the southern part of the ice sheet. The regional climate model output is compared with an ensemble of global climate models within CMIP6
25 showing similar patterns in precipitation change but with overall larger changes in the CMIP6 ensemble median compared with the regional climate model ensemble median.


1 Introduction

Solid precipitation (snowfall) is the largest positive contributor to the (surface) mass balance of the Greenland ice sheet. Runoff from the Greenland ice sheet is the largest negative surface mass balance component and currently a main contributor to sea
30 level rise (Box et al., 2022a). Early work (Huybrechts et al., 1991) suggested that the ice sheet would not contribute significantly to future sea level rise because of the projected increase in snowfall (Bailey and Hubbard, 2025), compensating

for increased runoff. However, the observed increase in surface melt/runoff and ice dynamic changes leading to increased solid ice discharge since the late 1990s shows conclusively that, in spite of an increase in observed precipitation (Box et al., 2022a), the ice sheet is net losing mass as increases in melt and runoff rates outpace increases in snowfall. The accumulation of snow is not only important for the ice sheet mass balance (the sum of surface mass balance and solid ice discharge) but also has process implications for ablation via surface albedo feedbacks and firn pore space and energy budgets that determine the retention and refreezing of melt water in the firn pack (Firn Symposium team, 2024; Glaude et al., 2024). If warming-enhanced precipitation occurs under future climate change scenarios, it is likewise important to determine the phase. Liquid precipitation percolates into near-surface snow/firn layers and thereby decreases the pore space in the firn. During the right conditions (heavy rain on low temperature firn), the higher temperature of rain compared to melt water can also transport energy deeper into the firn, warming it and reducing refreezing potential (Harper et al., 2023). Where fresh snow generally increases surface albedo, rain usually has a darkening effect on surface snow, particularly in near infrared wavelengths, and can therefore also enhance melt-albedo feedbacks.

Huai et al. (2025) used CMIP5 and CMIP6 ensembles and found a model mean increase in precipitation for the entire Greenland ice sheet during the 21st century with changes in excess of 200 mm yr⁻¹ along the outer edges for the SSP5-8.5 scenario. However, some individual models showed a decrease in the southeast. Bochow et al. (2024) used CMIP6 ensembles for four different emission scenarios and found an increase in precipitation in correlation with the increase in surface temperature for the 21st century. The aim of this study is to study changes (amount and phase) in the ice sheet precipitation for the 21st century using an ensemble of high resolution regional climate model simulations and compare these results with the forcing CMIP6 models.

2 Data



55 **Figure 1: Model domain and orography for the HIRHAM (a), MAR (b) and RACMO (c) RCM simulations. Sea points are given in blue, glacier free land points in green/brown and glacier points in white with added surface elevation contour lines in black. The location of seven observational stations (red dots) and six drainage basins (blue contour lines; Rignot and Mouginot, 2012) are shown in panel a. The stations in order 1–7 are Kangerlussuaq, Nuuk, Narsarsuaq, Tasiilaq, Ittoqqortoormiit, Danmarkshavn and Station Nord.**

60

2.1 Models

A total of eleven global climate model (GCM) – regional climate model (RCM) combinations are used in this study. The RCMs used are: HIRHAM5 (from now on: HIRHAM; one SSP1-2.6 and two SSP5-8.5 simulations), MAR3.12 (from now on: MAR; two SSP1-2.6, four SSP2-4.5 and eight SSP5-8.5 simulations) and RACMO2 (from now on: RACMO; one for each of the three SSP scenarios) (see Table 1). The reference period for this study is 1971–2000, taken from the historical run of the RCM simulations. Two future periods (2031–2060 and 2071–2100) are extracted from the three SSP scenarios. We use the period 1991–2022, using data for both the historical period (from 1991 to 2014) and the SSP5-8.5 period (from 2015 to 2022), for model evaluation against observations and reanalysis data.

70 The two HIRHAM (combining the dynamics of the HIRLAM model (Undén et al., 2002) and the physical parameterization schemes of the ECHAM model (Roeckner et al., 2003)) simulations are performed on a rotated latitude-longitude grid with a horizontal resolution of 0.05° (about 5.5 km) (Figure 1a; cf. Boberg et al. (2022) for a more detailed model setup description). The first HIRHAM simulation is forced with the CMIP6 GCM EC-Earth3 for the period 1970–2100 (where the first year is used for spin-up) using the SSP1-2.6 and SSP5-8.5 scenarios whereas the second HIRHAM simulation is forced by CMIP6 GCM CESM2-L for the SSP5-8.5 scenario and the same model set-up and period as for the HIRHAM/EC-Earth3 simulation.

75 The MAR (Modèle Atmosphérique Régional) simulations (Figure 1b; Hofer et al., 2020) are performed with a horizontal resolution of 15 km. The CMIP6 GCMs that are downscaled with MAR are CESM2, CESM2-L, CNRM-CM6, CNRM-ESM2,

80 IPSL-CM6A-LR, MPI-ESM1.2-HR, NorESM2 and UKESM1.0-LL. There are in total two SSP1-2.6 simulations, four SSP2-4.5 simulations and eight SSP5-8.5 simulations. The MAR data are regridded onto the HIRHAM grid to enable a model ensemble comparison (see Glaude et al., 2024). After regridding, we apply a common HIRHAM ice mask to all data since the ice sheet extent for the three RCMs are comparable in size (within $\pm 0.6\%$).

85 The RACMO simulations (Noël et al., 2020) are forced by CESM2-L and run at a horizontal resolution of 11 km (see Figure 1c) with one simulation per SSP scenario. The RACMO data are also regridded onto the HIRHAM grid. For a more detailed model description of RACMO, see Noël et al. (2018).

The RCM ensemble is compared with 34 CMIP6 GCM simulations (see Table 2) for the historical and the SSP5-8.5 scenario, selecting only one realization per GCM. The GCM outputs are regridded onto the EC-Earth3 grid with a horizontal resolution of about 80 km (Döscher et al., 2022).

Model	Res. [km]	P 1971–2000 [Gt]	P 2031–2060 [Gt]	P 2071–2100 [Gt]	ΔP 2031–2060		ΔP 2071–2100		RMSE 1991–2022 [mm]	Bias 1991–2022 [mm]
					[Gt]	[%]	[Gt]	[%]		
CARRA	2.5	842*	N/A	N/A	N/A	N/A	N/A	N/A	188	-88
HIRHAM/EC-Earth3	5.5	751	835	883	84	11	132	18	125	-10
			N/A	N/A	N/A	N/A	N/A	N/A		
HIRHAM/CESM2-L	5.5	762	N/A	N/A	N/A	N/A	N/A	N/A	199	106
			N/A	N/A	N/A	N/A	N/A	N/A		
RACMO/CESM2-L	11	700	765	931	3	0	170	22	352	109
			744	696	40	6	-4	-1		
			717	720	17	2	20	3		
MAR/CESM2	15	671	715	849	15	2	148	21	456	-290
			774	716	103	15	45	7		
			742	752	71	11	81	12		
MAR/CESM2-L	15	697	752	845	81	12	174	26	475	-312
			N/A	N/A	N/A	N/A	N/A	N/A		
			N/A	N/A	N/A	N/A	N/A	N/A		
MAR/CNRM-CM6	15	703	708	835	11	2	139	20	529	-386
			N/A	N/A	N/A	N/A	N/A	N/A		
			N/A	N/A	N/A	N/A	N/A	N/A		
MAR/CNRM-ESM2	15	696	754	896	51	7	193	27	514	-373
			N/A	N/A	N/A	N/A	N/A	N/A		
			N/A	N/A	N/A	N/A	N/A	N/A		
MAR/IPSL-CM6	15	703	720	815	24	3	119	17	598	-439
			N/A	N/A	N/A	N/A	N/A	N/A		
			N/A	N/A	N/A	N/A	N/A	N/A		
MAR/MPI-ESM1.2	15	690	799	957	97	14	255	36	548	-383
			726	720	36	5	30	4		
			721	766	31	4	76	11		
MAR/NorESM2	15	714	746	845	56	8	155	22	422	-233
			N/A	N/A	N/A	N/A	N/A	N/A		
			778	795	64	9	81	11		
			805	852	91	13	138	19		

MAR/UKESM1.0	15	684	N/A	N/A	N/A	N/A	N/A	N/A	494	-322
			824	954	140	20	270	39		
			855	1043	171	25	359	52		
Ensemble median	N/A	700	759	718	62	9	38	6	473	-307
			742	766	64	9	81	11		
			754	852	56	8	170	24		

* The CARRA value is for the period 1991–2022

Table 1: List of CARRA and RCM simulations used in this study and the annual mean total precipitation for the Greenland ice sheet for the reference period, the mid-century period and the end of century period. Also given are the absolute and relative changes for the two scenario periods. For each of the RCM and the ensemble median, the three lines refer to the SSP1-2.6, SSP2-4.5 and SSP5-8.5 scenarios, respectively. Note that the CARRA value is for the period 1991–2022. N/A refers to data not available, either missing scenario simulations (RCM) or time period not covered by the data (CARRA). Also given are the RMSEs and mean biases for 1991–2022 (using historical data and SSP5-8.5 data) relative to observations for the seven weather stations (cf. Figure 3).

GCM	P [Gt]	ΔP [Gt]	GCM	P [Gt]	ΔP [Gt]	GCM	P [Gt]	ΔP [Gt]
ACCESS-CM2	575	243	CNRM-CM6-1-HR	799	337	KACE-1-0-G	583	176
ACCESS-ESM1-5	590	287	CNRM-ESM2-1	882	156	MIROC6	771	180
AWI-CM-1-1-MR	655	242	E3SM-1-1	699	832	MIROC-ES2L	830	230
AWI-ESM-1-REcoM	663	253	EC-Earth3	702	320	MPI-ESM1-2-HR	634	159
BCC-CSM2-MR	718	237	EC-Earth3-Veg	720	322	MPI-ESM1-2-LR	668	206
CAMS-CSM1-0	574	75	FGOALS-f3-L	690	260	MRI-ESM2-0	887	218
CanESM5-1	606	540	FGOALS-g3	740	72	NESM3	581	340
CESM2	805	128	FIO-ESM-2-0	687	247	NorESM2-MM	641	127
CESM2-WACCM	780	165	GFDL-ESM4	665	119	TaiESM1	717	183
CMCC-CM2-SR5	723	344	INM-CM4-8	655	172	UKESM1-0-LL	548	333
CMCC-ESM2	709	281	INM-CM5-0	677	204	Ensemble		
CNRM-CM6-1	848	196	IPSL-CM6A-LR	853	393	median	694	233

Table 2: List of the 34 CMIP6 GCMs used in this study together with the annual mean total Greenland ice sheet precipitation for 1971–2000 and the change in total precipitation 2071–2100 relative to 1971–2000. Also given are the ensemble median values. The names in bold face are the GCMs downscaled by the RCMs. We have restricted the model selection and only using one ensemble member per GCM.

2.2 Observations and reanalysis

There are limited long-term observations of precipitation in Greenland and observed precipitation, particularly solid, is notoriously difficult to accurately measure (Allerup et al., 1997). Nevertheless, corrected datasets from automatic weather stations (AWS) operated by the Danish Meteorological Institute (DMI) have long time series that have been previously used to evaluate climate models (Huai et al., 2025; Boberg et al., 2018, Huai et al., 2021; Box et al., 2023; Niwano et al., 2021). On the ice sheet, the PROMICE (Fausto et al., 2021) and GC-Net (Steffen and Box, 2001) networks of AWS (Fausto et al., 2025) assess surface height changes, using two or three different techniques, but these are hard to compare with precipitation in a climate model as they measure a different variable affected by for example, melt, firm densification and drifting snow processes. We have therefore chosen to concentrate on using the observed precipitation from the DMI AWS. The station data are undercatch corrected (i.e. positive correction of the amount of water caught by the rain gauge due to winds) using the mean correction factors and mean rainfall fractions from Huai et al. (2021).

The Copernicus Arctic Regional Reanalysis (CARRA; Yang et al., 2020; Schyberg et al., 2020) is available from 1991 onward for two Arctic regions where one covers Greenland. CARRA data have a horizontal grid spacing of 2.5 km compared with the European Center for Medium-range Weather Forecasts fifth-generation reanalysis (ERA5) dataset at 31 km. Compared with ERA5, CARRA includes a number of improvements such as a larger set of local observations, a more realistic ice mask and coastline mask, a satellite derived glacier albedo dataset and high resolution sea surface temperatures and sea ice concentrations. The CARRA outperforms ERA5 when comparing the two datasets with in-situ observations of a range of variables including snowfall, snow depth, pressure, temperature and wind speeds across the Arctic (Køltzow et al., 2022; Box et al., 2023).

Eleven RCM simulations (historical 1991–2014 and SSP5-8.5 for 2015–2022) and CARRA are validated against observations for precipitation from seven stations with at least 10 000 daily measurements (out of a maximum of 11 688) for the 32-year period 1991–2022. The stations (see Figure 1) are located along the south-western and eastern coast of Greenland.

The six drainage basins (see Figure 1a) are taken from Rignot and Mouginot (2012), who used ice surface velocity maps from satellite radar interferometry data to divide the ice sheet into six subregions of similar size.

2.2 Model evaluation

Figure 2a shows the mean annual precipitation for the CARRA dataset for 1991–2022, with values exceeding 2000 mm year⁻¹ in the southeast and southwest whereas the northern part of the ice sheet is very dry with values below 200 mm year⁻¹ in large areas. Figure 2b to Figure 2l show the relative model difference for the eleven RCM simulations with respect to CARRA. We see similar patterns for all models with large areas with a negative difference of about -10 to -30% in the central parts of the ice sheet and smaller areas with a positive difference up to 50% along the Greenland coastline. The model with the largest positive difference (in regions NW and NO with values in the range 10-50%) over the ice sheet are MAR driven by CNRM-CM6, CNRM-ESM2 and MPI-ESM1.2 while the two HIRHAM simulations and the RACMO simulation have the largest positive differences outside the ice sheet with values in the range 30-70% with peaks in excess of 70%.




Figure 2: Mean annual precipitation for 1991–2022 by CARRA (a) and the model difference relative CARRA for HIRHAM/EC-EARTH (b), HIRHAM/CESM2-L (c), MAR/CESM2 (d), MAR/CESM2-L (e), MAR/CNRM-CM6 (f), MAR/CNRM-ESM2 (g), MAR/IPS�-CM6 (h), MAR/MPI-ESM1.2 (i), MAR/NorESM2 (j), MAR/UKESM1.0 (k) and RACMO/CESM2-L (l). For the RCM data, 1991–2014 is taken from the historical run and 2015–2022 is taken from the SSP5-8.5 scenario run. Black contour lines show the six drainage basins.

145


150

Figure 3 compares station observations of precipitation with CARRA and the eleven RCM models for the grid point closest to the weather stations (cf. Figure 1a). For station 1 all model simulations agree with observations except MAR, driven by IPSL, showing a dry bias while RACMO shows a clear wet bias of about 60% (130 mm). For station 2 all MAR simulations show a dry bias of about -25% (-250 mm) while the other simulations are closer to observations. For station 3, CARRA is close to observations while the two HIRHAM simulations have a wet bias in the range 35 to 70% (250 to 500 mm), RACMO show a

wet bias of about 110% (800 mm) and five of the MAR simulations have a dry bias of about -35% (-250 mm). The two HIRHAM simulations are close to observations for station 4 while CARRA and RACMO show a dry bias close to -20% (-350 mm) while all eight MAR simulations have a clear dry bias close to -65% (-1300 mm). For station 5, CARRA and one HIRHAM simulation are close to observations while the other HIRHAM simulation together with two MAR simulations and RACMO have a dry bias of about -20% (-180 mm). The other MAR simulations have a clear dry bias in the range -30 to -60% (-330 to -530 mm). A similar pattern is seen for station 6 with the exception for RACMO here showing a wet bias of about 25% (90 mm). Finally, for station 7 one HIRHAM simulation is close to observations while the other HIRHAM simulation together with RACMO have a wet bias of about 15% (80 mm) while CARRA has a dry bias of -15% (70 mm) and all eight MAR simulations have a dry bias in the range -15 to -45% (-60 to -210 mm).

The two rightmost columns of Table 1 give the RMSE and mean bias for CARRA and the eleven RCM simulations, showing low RMSE values and low mean bias for the two HIRHAM simulations and CARRA while the MAR simulations have a relatively high RMSE and negative bias.

CARRA models different climate variables well compared with ERA5 (Isaksen et al., 2022; Box et al., 2023; Køltzow et al., 2022). However, when comparing CARRA with station observations (see Figure 3 and Table 1) and RCM output, there are no clear evidence justifying a bias adjustment of the RCM data towards CARRA. Instead, we will use the raw model output in this study.



175 **Figure 3: Mean annual precipitation for 1991–2022 (dots) for seven weather station locations (cf. Figure 1), showing observations, CARRA, two HIRHAM simulations, eight MAR simulations and one RACMO simulation. For the RCM data, 1991–2014 is taken from the historical run and 2015–2022 is taken from the SSP5-8.5 scenario run. The vertical lines show the standard deviation of the 32 yearly precipitation sums. The precipitation for CARRA and the HIRHAM, MAR and RACMO simulations are taken from the land grid point closest to the AWS locations.**

3 Results

180 Figure 4 shows the model spread of the relative change in precipitation for a mid-century and an end of the century period relative to 1971–2000 using the RCM ensemble for all three SSP scenarios. For the mid-century median for all three scenarios (Figure 4b, 4h and 4n) we see a positive change between 10 and 50% for most of Greenland except region SE with no clear change. For the mid-century 25th percentile (Figure 4a, 4g and 4m) we mostly see a positive change (10 to 30%) for regions NE and NO and a negative change (-10 to -30%) in region SE. For the mid-century 75th percentile (Figure 4c, 4i and 4o) all of Greenland shows a positive change that is most pronounced in regions NE and NO with values in the range 30 to 70%.

185

Towards the end of the century, we see an overall amplification of the mid-century signal. The ensemble median shows a reduction in precipitation compared with the mid-century values for the SSP1-2.6 scenario (Figure 4e) while SSP2-4.5 (Figure 4k) and SSP5-8.5 (Figure 4q) show an increase (except region SE) with values pct reaching 100% for region NE for SSP5-8.5. For the end of century 25th percentile (Figure 4d, 4j and 4p), we see similar patterns except for region SE with a -10 to -30% change along the outer edge. For the end of century 75th percentile (Figure 4f, 4l and 4r) we see a positive change for most of Greenland with values reaching 30 to 70% in region NE for SSP2-4.5 and values reaching 70 to 110% for SSP5-8.5.




Figure 4: Relative change in Greenland precipitation using the RCM ensembles for SSP1-2.6 (panels a–f), SSP2-4.5 (panels g–l) and SSP5-8.5 (panels m–r). Panels a–c, g–i and m–o are for 2031–2060 relative 1971–2000 while panels d–f, j–l and p–r are for 2071–2100 relative 1971–2000. First and fourth columns represent the 25th percentile, second and fifth columns are the median and the third and sixth columns represent the 75th percentile.




Figure 5: Ensemble median relative change in precipitation using the subset of CMIP6 models used when downscaling the RCMs for 2031–2060 (a) and 2071–2100 (d) relative to 1971–2000 for the SSP5-8.5 scenario. The corresponding ensemble median relative change using the full set of 34 CMIP6 GCMs are shown in panels b and e for 2031–2060 and 2071–2100, respectively. The two panels c and f in the right column show number of CMIP6 models, using the full 34 model set, with a positive relative change in precipitation for 2031–2060 (c) and 2071–2100 (f) relative to 1971–2000 for the SSP5-8.5 scenario, respectively.

Figure 5a and 5d show the ensemble mean relative change, using the eight CMIP6 GCMs dynamically downscaled by the RCMs (cf. Table 2), for the SSP5-8.5 scenario for 2031–2060 and 2071–2100 relative 1971–2000, respectively. The mid-century relative change is between 10 and 30% for regions SW, NO and NE with areas reaching 50% for region NE. For the end of the century, all regions except SE show a positive change with areas reaching 90% in NO and NE. Figure 5b and 5e give the corresponding maps when using the full 34 CMIP6 GCM ensemble for the SSP5-8.5 scenario. Compared with the eight-model ensemble, the full model ensemble has larger changes for all six subregions but most pronounced for region SE where we now see a positive change in the range 10 to 30%. Figure 5c and 5f show the number of CMIP6 models that give positive changes in precipitation for the mid-century and end of century, pixel by pixel. The CMIP6 ensemble show good

agreement for all drainage basins except for region SE where about 6 of the 34 members show a negative change for the end of the century, in line with Huai et al. (2025).

Figure 6 shows the change from solid to liquid precipitation for the SSP5-8.5 scenario, represented by the rainfall fraction (rainfall amount divided by the total precipitation amount) where a value of 50% would mean that rainfall and snowfall contribute equal amounts to the total precipitation. This indicates that not only will there be an increase in precipitation in the future, but an increasing fraction of that will be rainfall, especially around the ice sheet margins. At the end of the century for the SSP5-8.5 scenario (Figure 6c), we see that a large fraction of the precipitation is rainfall, especially in the south (basins SW and SE; cf. Fig. 1a). Figure 6d shows how the rainfall fraction of total precipitation for the six drainage basins changes with time. For the historical period, the mean rainfall fraction is around 5% for basins SW and SE while the other basins show mean values below 2%. At the end of the century for the SSP5-8.5 scenario, we see that a large fraction of the precipitation is rainfall (total ice sheet: around 15%), especially for basins SW (above 25%) and SE (around 20%).






Figure 6: Ensemble median rainfall fraction of total precipitation in Greenland, using the eleven-model ensemble for the historical

230 period 1971–2000 (a) and the two SSP5-8.5 scenario periods 2031–2060 (b) and 2071–2100 (c). Also shown is a time series of ensemble median annual values of the rainfall fraction for all six drainage basins and the total ice sheet (d).



235 **Figure 7: Mean precipitation (in Gt per year) for the RCM simulations over the six drainage basins and the full ice sheet for the historical period 1971–2000, for CARRA (1991–2022) and the scenario periods 2031–2060 and 2071–2100. The horizontal lines denote the ensemble median, the boxes represent the 25th to 75th percentiles while the vertical bars show the full model spread. For the two scenario periods the boxes are split into three segments: the left for SSP1-2.6, the middle for SSP2-4.5 and the right for SSP5-8.5. The y axis to the right is used for the full ice sheet (black) while the values for the six individual drainage basins are given by the y axis to the left.**

240



245 **Figure 8: Annual sum of precipitation (a), snowfall (b) and rainfall (c) over the Greenland ice sheet as a function of annual mean near-surface temperature for the period 1971–2100, using the SSP5-8.5 scenario, with respect to the 1971–2000 mean. The data are separated for each RCM (MAR: blue, HIRHAM: red and RACMO: brown), including a least square fit (linear for a and b, second degree polynomial for c). Also shown is the least square fit to the 34 CMIP6 model ensemble mean values (black).**

250 Figure 7 shows the annual total precipitation for the six drainage basins (using the left y-axis) as well as for the total ice sheet (using the right y-axis) for CARRA and the RCM simulations. The vertical extent of each bar represents the model spread for each basin while the boxes show the 25th, 50th and the 75th percentiles of the ensemble. For the two scenario periods the bar is split into three segments to show the difference between the three SSPs. The largest uncertainties are, as expected, found in

difference in the ensemble median. The CARRA values match the higher end of the model spread for each basin, giving a total annual amount that is significantly higher than any of the other models for the reference period.

255 A warmer atmosphere has an exponentially higher saturation specific humidity (can hold more water vapour), termed the
Clausius-Clapeyron relationship with about a 7% increase per degree warming (Trenberth et al. 2003). Figure 8a shows the
relationship between annual temperature and annual precipitation over the ice sheet for the three RCMs using data for the
historical period and the SSP5-8.5 scenario only to have a more uniform distribution of the data points. Here we see that the
integrated precipitation over the ice sheet increases with 31.8, 35.5, and 37.3 Gt per degree change in temperature for RACMO,
260 MAR and HIRHAM respectively. Figure 8b and 8c shows that the increase in rainfall per degree of warming is larger than the
increase in snowfall per degree of warming. Snowfall is expected to increase by about 13 Gt per degree of warming while
rainfall is expected to increase with above 20 Gt per degree of warming during the 21st century but with a non-linear trend
indicating a much higher value at the end of the century.

265 In absolute terms (cf. Figure 9a), for the end of century SSP5-8.5 scenario the largest changes are expected, using the RCM
ensemble, in the south west of Greenland with values exceeding 200 mm yr⁻¹ and along the outer edge of the ice sheet with
values between 100 and 200 mm yr⁻¹ (except for the south east with values close to 0). A similar pattern is seen using the
subset of GCMs (cf. Figure 9b) but with even lower (and negative) values in the south east. For the full set of GCMs (cf. Figure
9c) we find increases for all of the ice sheet with the highest values in the south west and south east.

270




Figure 9: Ensemble median absolute change in precipitation for SSP5-8.5 for 2071–2100 relative 1971–2000 using the RCM ensemble (a), the subset of downscaled GCMs (b) and the full set of GCMs (c).

4 Discussion

275 4.1 Implications for mass budget

Using an ensemble of high resolution RCM simulations for the SSP1-2.6/SSP2-4.5/SSP5-8.5 scenarios, we predict an increase of precipitation on the ice sheet during the 21st century, though with a change in the relative distribution of precipitation as well as a change in the phase. By the end of the century, the ice sheet is expected to receive between 40 and 170 Gt more precipitation annually, depending on the climate scenario (cf. Table 1), an increase between 6 and 24% compared to the present
280 day. The largest relative differences are predicted in the north east of Greenland, where all models see a marked increase. In the southeast by contrast, the RCM ensemble median give relative changes close to zero (cf. Figure 4 and Figure 7). Results from the GCM ensemble from CMIP6 (cf. Figure 5) show that the north east signal is robust across most models while the south east signal is positive for 28 out of the 34 models for the end of the century (cf. Figure 5f). The RCMs follow the forcing GCM climate in this respect but their higher resolution shows more detail on the local scale.

285

The precipitation phase is important because rainfall serves as a crucial climate indicator in polar regions and plays a multifaceted role in cryospheric processes. When heavy rain falls on low temperature snow or ice surfaces, its infiltration not only contributes directly to surface melt, but also initiates a sequence of physical changes in the snowpack (Box et al., 2022a). The freezing of rainwater within the firn releases latent heat, which further enhances snowpack collapse by warming the
290 surrounding ice and snow (Harper et al., 2023). In addition, rainfall events lower the surface albedo making the surface more susceptible to future melting due to increased absorption of solar radiation (Noël et al., 2022; Firn Symposium team, 2024; Gilbert et al., 2025; Box et al., 2022b).

A reduction in Arctic sea ice cover, due to increases in atmospheric CO₂, can enhance transfer of heat and moisture between
295 the ocean and the atmosphere leading to an increase in temperature and precipitation over Greenland (Stroeve et al., 2017; Noël et al., 2014). This increase in temperature will increase the ice melt and thereby decreasing the surface albedo creating a positive feedback loop, resulting in a warmer atmosphere with the capacity to hold more water vapor and thereby intensifying the hydrological cycle (Zeitz et al., 2021).

4.2 Sources of spread in precipitation projections

300 Huai et al. (2025) used CMIP6 GCM outputs for the SSP5-8.5 scenario and found a northeastward shift of the Icelandic low leading to lower precipitation amounts in southeast Greenland. The future location of the Icelandic low has therefore an important influence on precipitation changes over southeast Greenland, as CMIP6 models have some important biases in the Iceland Low and representation of the North Atlantic Oscillation. Furthermore, Greenland blocking events are difficult to estimate by climate models (Davini et al. 2021) and can have a large impact on precipitation, especially during winter.

305

CARRA has shown clear advantages to other reanalysis products (Køltzow et al., 2022; Box et al., 2023) and CARRA performs well when compared with station observations outside the ice sheet (cf. Figure 3). Compared with our RCM ensemble (cf. Figure 7) over the ice sheet, CARRA overestimate precipitation for all basins except NO and NW. This may simply reflect the local conditions around the observation sites, which are typically located in or near topographically complex terrain, difficult to capture even at 2.5 km resolution. To accurately capture extreme weather events in Greenland, DMI use a sub-km resolution model for numerical weather prediction purposes (Yang, 2018; Yang, 2019).

Using the mean values of total precipitation, separated by RCM, in Table 1 for the period 1971–2000 together with the relationship between precipitation and temperature in Figure 8, we get 4.9%, 5.1% and 4.5% increase in precipitation per Kelvin for HIRHAM, MAR and RACMO, respectively. These values are lower than the app. 7% K⁻¹ predicted by the Clausius-Clapeyron relationship (Trenberth et al., 2003). However, the near-surface temperature is not an accurate measure of the vertically weighted atmospheric temperature where the moisture resides. Moreover, the total ice sheet precipitation is for an important part topographically forced, i.e. very sensitive to changes in the large-scale circulation. Bochow et al. (2024) used 32 GCMs within the CMIP6 project and found a mean value between 3 and 6% K⁻¹ following the scenario used for the ice sheet but with a strong spatial dependence. A 5% increase in precipitation per degree warming over the ice sheet were found by Gregory and Huybrechts (2006) and by Fettweis et al. (2013). There are several ways of determining the relationship between temperature and precipitation changes. Frieler et al. (2015) used a wide range of methods for Antarctica and derived a combined sensitivity of $5 \pm 1\% \text{ K}^{-1}$. The individual forcing models in this study give similar values. Box et al. (2013) found the theoretically expected 7% K⁻¹ using reconstructions of snow accumulation from ice cores together with the average near-surface air temperature in the Northern Hemisphere for the period 1880–1962 during which the temperature increase was relatively low (less than +1°C) and during which changes in general circulation or natural variability could have also impacted this precipitation increase vs temperature increase. However, values close to the theoretical 7% K⁻¹ relationship are only expected for heavy rain conditions (Trenberth et al., 2003).

Figure 6 shows that a significant fraction of future precipitation, particularly over the southern ice sheet, will fall as rain. On the broad scale, rain replacing snow implies enhanced ablation over the ice sheet due to surface energy budget and firn processes (Box et al., 2023). On a local to regional scale, increased rainfall, particularly close to the end of the melt season, has also been shown to affect outlet glacier dynamics (Doyle et al., 2014) with local short-lived increases in ice velocity. The CMIP6 GCMs are unlikely to capture this transition from snow to rain, and along with the higher resolution show how RCMs add value to the GCMs when considering future climate change. Zhang et al. (2024) used 21 CMIP6 models for the SSP5-8.5 scenario giving a projected annual increase in precipitation over the ice sheet of 280 Gt for 2071–2100 relative 1986–2005. The RCM ensemble median gives 170 Gt with individual model values ranging from 120 to 360 Gt.

340 A clear limitation of this study is that RCMs often exhibit large biases in simulating point-wise precipitation, while the observations also have large uncertainties, especially when undercatch is prominent. The large-scale circulation and sea surface temperatures from forcing GCMs often have errors that are subsequently inherited by the RCMs. Furthermore, small-scale convective processes are not (fully) resolved by the RCMs used in this study. Also, the lack of a gridded observational product for Greenland makes it difficult to perform bias adjustment of the modeled precipitation. These limitations are, to some degree, compensated when looking at relative changes and not absolute values and the use of an RCM ensemble instead of looking at
345 a single RCM.

5 Conclusions

Our work, consistent with theory, finds that the Greenland ice sheet faces a wetter future in all regions, with as a possible exception the southeast. The fraction of rain relative to the total precipitation increases exponentially with temperature during the 21st century. Our analysis clearly shows a linear relationship between annual precipitation amounts and annual mean
350 temperature as well as a dependency on shared socioeconomic pathway (SSP). We also see a dependency on driving model; when all three RCMs downscale the same GCM (CESM2-L) we obtain similar results on change in precipitation for both scenario periods (cf. Table 1). For the historical period, the different MAR simulations show similar results while HIRHAM, and to some extent RACMO, present different characteristics (cf. Figure 3). At the end of the century, HIRHAM, driven by EC-Earth3, and MAR, driven by IPSL-CM6 and UKESM1.0, give much higher changes in precipitation over the ice sheet
355 than the other RCM simulations, indicating a dependency on the choice of GCM. The more detailed topography from the higher resolution RCMs shows a complexity in the distribution of precipitation suggesting that using emulation techniques on RCM output to expand future ensembles of climate simulations is feasible.

Acknowledgments

Fredrik Boberg, Nicolaj Hansen and Ruth Mottram are supported by the Danish State through the National Centre for Climate
360 Research (NCKF) and also supported by the European Union's Horizon 2020 project PROTECT (grant no. 869304); furthermore, Nicolaj Hansen has been supported by the Novo Nordisk Foundation project PRECISE (grant no. NNF23OC0081251). Finally, we would like to thank Thomas Mölg for editorial handling and Jason Box and an anonymous reviewer for helpful and supportive comments that improved this paper.

Author contributions

365 FB designed the study with contributions from all co-authors. FB performed the HIRHAM simulations, the data collection and the analysis. XF and MvdB provided the MAR and RACMO data, respectively. FB wrote the paper with contributions from all co-authors.

Competing interests

370 Some authors are members of the editorial board of The Cryosphere. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Data availability

All GCM, RCM, CARRA and station observation data used in this study will be made available upon request. CARRA data can be downloaded from the Copernicus Arctic Regional Reanalysis webpage at <https://climate.copernicus.eu/copernicus-arctic-regional-reanalysis-service>. The drainage basin information can be downloaded from <http://imbie.org/imbie-2016/drainage-basins/>.

References

- Allerup, P., Madsen, H., and Vejen, F.: A comprehensive model for correcting point precipitation, *Nordic Hydrology*, 28, 1–20, 1997.
- 380 Bailey, H., and Hubbard, A.: Snow mass recharge of the Greenland ice sheet fueled by intense atmospheric river. *Geophysical Research Letters*, 52, e2024GL110121. <https://doi.org/10.1029/2024GL110121>, 2025.
- Boberg, F., Langen, P. L., Mottram, R. H., Christensen, J. H., and Olesen, M.: 21st-century climate change around Kangerlussuaq, west Greenland: From the ice sheet to the shores of Davis Strait, *Arctic, Antarctic, and Alpine Research*, 50(1), <https://doi.org/10.1080/15230430.2017.1420862>, 2018.

385

- Boberg, F., Mottram, R., Hansen, N., Yang, S., and Langen, P. L.: Uncertainties in projected surface mass balance over the polar ice sheets from dynamically downscaled EC-Earth models, *The Cryosphere*, 16, 17–33, <https://doi.org/10.5194/tc-16-17-2022>, 2022.
- 390 Bochow, N., Poltronieri, A., and Boers, N.: Projections of precipitation and temperatures in Greenland and the impact of spatially uniform anomalies on the evolution of the ice sheet, *The Cryosphere*, 18, 5825–5863, <https://doi.org/10.5194/tc-18-5825-2024>, 2024.
- Box, J.E., N. Cressie, D.H. Bromwich, J. Jung, M. van den Broeke, J.H. van Angelen, R.R. Forster, C. Miège, E. Mosley-
395 Thompson, B. Vinther, J.R. McConnell: Greenland ice sheet mass balance reconstruction. Part I: net snow accumulation (1600–2009), *J. Climate*, 26, 3919–3934, doi:10.1175/JCLI-D-12-00373.1, 2013.
- Box, J. E., Hubbard, A., Bahr, D.B. et al.: Greenland ice sheet climate disequilibrium and committed sea-level rise. *Nat. Clim. Chang.*, 12, 808–813, <https://doi.org/10.1038/s41558-022-01441-2>, 2022a.
- 400 Box, J. E., Wehrlé, A., vanAs, D., Fausto, R. S., Kjeldsen, K. K., Dachauer, A., et al.: Greenland ice sheet rainfall, heat and albedo feedback impacts from the mid-August 2021 atmospheric River. *Geophysical Research Letters*, 49, e2021GL097356. <https://doi.org/10.1029/2021GL097356>, 2022b.
- 405 Box, J. E., Nielsen, K. P., Yang, X., Niwano, M., Wehrlé, A., van As, D., et al.: Greenland ice sheet rain climatology, extremes and atmospheric river rapids, *Meteorological Applications*, 30(4), e2134, <https://doi.org/10.1002/met.2134>, 2023.
- Davini, P., Weisheimer, A., Balmaseda, M., Johnson, S. J., Molteni, F., Roberts, C. D., Senan, R., and Stockdale, T. N.: The representation of winter Northern Hemisphere atmospheric blocking in ECMWF seasonal prediction systems, *Q. J. Roy. Meteor. Soc.*, 147, 1344–1363, <https://doi.org/10.1002/qj.3974>, 2021.
- 410 Doyle, S. H., Hubbard, A., Fitzpatrick, A. A. W., van As, D., Mikkelsen, A. B., Pettersson, R., and Hubbard, B.: Persistent flow acceleration within the interior of the Greenland ice sheet, *Geophysical Research Letters*, 41(3), 899–905. <https://doi.org/10.1002/2013GL058933>, 2014.
- 415 Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arsouze, T., Bergman, T., Bernardello, R., Boussetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblas-Reyes, F. J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenigk, T., Makkonen, R., Massonnet, F., Ménégou, M., Miller, P. A., Moreno-Chamarro, E., Nieradzic, L., van Noije, T., Nolan, P.,

- 420 O'Donnell, D., Ollinaho, P., van den Oord, G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-Earth3 Earth system model for the Coupled Model Intercomparison Project 6, *Geosci. Model Dev.*, 15, 2973–3020, <https://doi.org/10.5194/gmd-15-2973-2022>, 2022.
- 425 Fausto, R. S., van As, D., Mankoff, K. D., Vandecrux, B., Citterio, M., Ahlstrøm, A. P., Andersen, S. B., Colgan, W., Karlsson, N. B., Kjeldsen, K. K., Korsgaard, N. J., Larsen, S. H., Nielsen, S., Pedersen, A. Ø., Shields, C. L., Solgaard, A. M., and Box, J. E.: Programme for Monitoring of the Greenland Ice Sheet (PROMICE) automatic weather station data, *Earth Syst. Sci. Data*, 13, 3819–3845, <https://doi.org/10.5194/essd-13-3819-2021>, 2021.
- 430 Fausto, R. S., How, P., Vandecrux, B., Lund, M. C., Box, J. E., Mankoff, K. D., Andersen, S. B., van As, D., Bahbah, R., Citterio, M., Colgan, W., Jakobsgaard, H. T., Karlsson, N. B., Kjeldsen, K. K., Larsen, S. H., Olsen, C., Oraschewski, F., Rutishauser, A., Shields, C. L., Solgaard, A. M., Stevens, I. T., Svendsen, S. H., Langley, K., Messerli, A., Bjørk, A. A., Andersen, J. K., Abermann, J., Steiner, J., Prinz, R., Hynek, B., Lea, J. M., Brough, S., and Ahlstrøm, A. P.: PROMICE | GC-
- 435 NET automatic weather station data, *Earth Syst. Sci. Data Discuss.* [preprint], <https://doi.org/10.5194/essd-2025-687>, in review, 2025.
- Fettweis, X., Franco, B., Tedesco, M., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., and Gallée, H.: Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *The Cryosphere*, 7, 469–489, <https://doi.org/10.5194/tc-7-469-2013>, 2013.
- Firn Symposium team, The, Firn on ice sheets, *Nat. Rev. Earth Environ.*, 5, 79–99, <https://doi.org/10.1038/s43017-023-00507-9>, 2024.
- 445 Frieler, K., Clark, P. U., He, F., Buizert, C., Reese, R., Ligtenberg, S. R. M., van den Broeke, M. R., Winkelmann, R. and Levermann, A.: Consistent evidence of increasing Antarctic accumulation with warming, *Nature Climate Change*, vol. 5, no. 4, pp. 348-352. <https://doi.org/10.1038/nclimate2574>, 2015.
- Gilbert, E., Pishniak, D., Torres, J. A., Orr, A., Maclennan, M., Wever, N., and Verro, K.: Extreme precipitation associated with atmospheric rivers over West Antarctic ice shelves: insights from kilometre-scale regional climate modelling. *The Cryosphere*, 19(2), 597-618, 2025.
- 450

- 455 Glaude, Q., Noel, B., Olesen, M., Van den Broeke, M., van de Berg, W. J., Mottram, R., et al.: A factor two difference in 21st-century Greenland ice sheet surface mass balance projections from three regional climate models under a strong warming scenario (SSP5-8.5), *Geophysical Research Letters*, 51, e2024GL111902, <https://doi.org/10.1029/2024GL111902>, 2024.
- Gregory J. M. and Huybrechts P.: Ice-sheet contributions to future sea-level change, *Phil. Trans. R. Soc. A.*, 364, 1709–1732, <http://doi.org/10.1098/rsta.2006.1796>, 2006.
- 460 Harper, J., Saito, J., and Humphrey, N.: Cold season rain event has impact on Greenland's firn layer comparable to entire summer melt season, *Geophysical Research Letters*, 50, e2023GL103654. <https://doi.org/10.1029/2023GL103654>, 2023.
- Hofer, S., Lang, C., Amory, C. et al.: Greater Greenland Ice Sheet contribution to global sea level rise in CMIP6. *Nat. Commun.*, 11, 6289, <https://doi.org/10.1038/s41467-020-20011-8>, 2020.
- 465 Huai, B., Van Den Broeke, M. R., Reijmer, C. H., and Cappellen, J.: Quantifying Rainfall in Greenland: A Combined Observational and Modeling Approach, *Journal of Applied Meteorology and Climatology*, 60(8), 1171–1188. <https://doi.org/10.1175/JAMC-D-20-0284.1>, 2021.
- 470 Huai, B., Ding, M., van den Broeke, M.R. et al.: Future large-scale atmospheric circulation changes and Greenland precipitation, *npj Clim. Atmos. Sci.*, 8, 10, <https://doi.org/10.1038/s41612-025-00899-z>, 2025.
- Huybrechts, P., Letreguilly, A., Reeh, N.: The Greenland ice sheet and greenhouse warming, *Paleogeogr. Paleoclim. Paleoecol.*, 89, 399–412, 1991.
- 475 Isaksen, K., Nordli, Ø., Ivanov, B. et al.: Exceptional warming over the Barents area, *Sci. Rep.* 12, 9371, <https://doi.org/10.1038/s41598-022-13568-5>, 2022.
- 480 Køltzow, M., Schyberg, H., Støylen, E., and Yang, X.: Value of the Copernicus Arctic Regional Reanalysis (CARRA) in representing near-surface temperature and wind speed in the north-east European Arctic, *Polar Research*, 41, <https://doi.org/10.33265/polar.v41.8002>, 2022.

Niwano, M., Box, J. E., Wehrlé, A., Vandecrux, B., Colgan, W. T., and Cappelen, J.: Rainfall on the Greenland ice sheet: Present-day climatology from a high-resolution non-hydrostatic polar regional climate model. *Geophysical Research Letters*, 48, e2021GL092942. <https://doi.org/10.1029/2021GL092942>, 2021.

485

Noël, B., Fettweis, X., van de Berg, W. J., van den Broeke, M. R., and Erpicum, M.: Sensitivity of Greenland Ice Sheet surface mass balance to perturbations in sea surface temperature and sea ice cover: a study with the regional climate model MAR, *The Cryosphere*, 8, 1871–1883, <https://doi.org/10.5194/tc-8-1871-2014>, 2014.

490 Noël, B., van de Berg, W. J., van Wessem, J. M., van Meijgaard, E., van As, D., Lenaerts, J. T. M., Lhermitte, S., Kuipers Munneke, P., Smeets, C. J. P. P., van Uft, L. H., van de Wal, R. S. W., and van den Broeke, M. R.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, 2018.

495 Noël, B., van Kampenhout, L., van de Berg, W. J., Lenaerts, J. T. M., Wouters, B., and van den Broeke, M. R.: Brief communication: CESM2 climate forcing (1950–2014) yields realistic Greenland ice sheet surface mass balance, *The Cryosphere*, 14, 1425–1435, <https://doi.org/10.5194/tc-14-1425-2020>, 2020.

Noël, B., Lenaerts, J. T., Lipscomb, W. H., Thayer-Calder, K., and van den Broeke, M. R.: Peak refreezing in the Greenland firn layer under future warming scenarios, *Nat. Commun.*, 13, 6870, <https://doi.org/10.1038/s41467-022-34524-x>, 2022.

500

Rignot, E. and Mouginot, J.: Ice flow in Greenland for the International Polar Year 2008-2009: ICE FLOW GREENLAND 2009, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051634>, 2012.

505 Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblueh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U. and Tompkins, A.: The atmospheric general circulation model ECHAM5. Part 1. Model description, Report no. 349, Max-Planck-Institut für Meteorologie (MPI-M), 2003.

510 Schyberg, H., Yang, X., Køltzow, M. A. Ø., Amstrup, B., Bakketun, Å., Bazile, E., Bojarova, J., e., B., Dahlgren, P., Hagelin, S., Homleid, M., Horányi, A., Hoyer, J., Johansson, Å., Killie, M. A., Körnich, H., LeMoigne, P., Lindskog, M., Manninen, T., Nielsen, Englyst, P., Nielsen, K. P., Olsson, E., Palmason, B., Peralta, Aros, C., Randriamampianina, R., Samuelsson, P., Stappers, R., Støylen, E., Thorsteinsson, S., Valkonen, T., and Wang, Z. Q.: Arctic regional reanalysis on single levels from 1991 to present, <https://doi.org/10.24381/cds.713858f6>, 2020.

515 Steffen, K. and Box, J.: Surface climatology of the Greenland Ice Sheet: Greenland Climate Network 1995–1999, *J. Geophys. Res.-Atmos.*, 106, 33951–33964, <https://doi.org/10.1029/2001jd900161>, 2001.

- Stroeve, J. C., Mioduszewski, J. R., Rennermalm, A., Boisvert, L. N., Tedesco, M., and Robinson, D.: Investigating the local scale influence of sea ice on Greenland surface melt, *The Cryosphere*, 11, 2363–2381, <https://doi.org/10.5194/tc-11-2363-2017>, 2017.
- 520
- Trenberth, K. E., Dai, A., Rasmussen, R. M., and Parsons, D. B.: The changing character of precipitation, *Bull. Am. Meteorol. Soc.*, 84, 1205–1217, 2003.
- Undén, P., Rontu, L., Järvinen, H., Lynch, P., Calvo, J., Cats, G., Cuxart, J., Eerola, K., Fortelius, C., Garcia-Moya, J. A., Jones, C., Lenderink, G., McDonald, A., McGrath, R., Navascues, B., Woetman Nielsen, N., Ødegaard, V., Rodrigues, E., Rummukainen, M., Room, R., Sattler, K., Hansen Sass, B., Savijärvi, H., Wichers Schreur, B., Sigg, R. The, H. and Tijm, A.: HIRLAM-5 Scientific Documentation. Scientific Report, 2002.
- 525
- Yang, X.: Sub-km HARMONIE and on-demand setup for storm forecast, *Joint ALADIN-HIRLAM Newsletter* 10, pp 35–39, Jan 2018, 2018.
- 530
- Yang, X.: TAS, an operational forecast model at hectometric scale. *Joint ALADIN-HIRLAM Newsletter* 12, pp 40–49, Jan 2019, 2019.
- 535
- Yang X., Schyberg H., Palmason B., Bojarova J., Box J., Pagh Nielsen K., Amstrup B., Peralta C., Høyer J., Nielsen Englyst P., Homleid M. Køltzow M.A.Ø., Randriamampianina R., Dahlgren P., Støylen E., Valkonen T., Thorsteinsson S., Kornich H., Lindskog M. and Mankoff K.: C3S Arctic regional reanalysis—full system documentation. Accessed on the internet at <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-carra-single-levels?tab=doc> on 1 November 2021, 2020.
- 540
- Zeitz, M., Reese, R., Beckmann, J., Krebs-Kanzow, U., and Winkelmann, R.: Impact of the melt–albedo feedback on the future evolution of the Greenland Ice Sheet with PISM-dEBM-simple, *The Cryosphere*, 15, 5739–5764, <https://doi.org/10.5194/tc-15-5739-2021>, 2021.
- 545
- Zhang, Q., Huai, B., Ding, M., Sun, W., Liu, W., Yan, J., Zhao, S., Wang, Y., Wang, Y., Wang, L., Che, J., Dou, J., Kang, L.: Projections of Greenland climate change from CMIP5 and CMIP6 *Global Planet. Change*, 232, Article 104340, [10.1016/j.gloplacha.2023.104340](https://doi.org/10.1016/j.gloplacha.2023.104340), 2024.