

Eddy kinetic energy and ~~baroclinic and barotropic~~ energy conversion rates along the Atlantic Water boundary current north of Svalbard

Kjersti Kalhagen^{1,2}, Ilker Fer^{2,3,1}, Till M. Baumann⁴, Jon Albretsen⁴, and Lukas Frank^{1,2*}

¹Department of Arctic Geophysics, University Centre in Svalbard (UNIS), Longyearbyen, Svalbard, Norway

²Geophysical Institute, University of Bergen, Bergen, Norway

³Bjerknes Centre for Climate Research, Bergen, Norway

⁴Institute of Marine Research, Bergen, Norway

*Now at SINTEF Ocean, Trondheim, Norway

Correspondence: Kjersti Kalhagen (kjerstik@unis.no)

Abstract. On the continental slope north of Svalbard, the boundary current carrying Atlantic Water (AW) loses heat as it flows eastward. This cooling cannot be fully attributed to atmospheric heat loss or turbulent mixing. Lateral exchange, potentially linked to mesoscale activity, has previously been proposed as a contributing factor, based on limited observations of eddies. Here, we analyse a year-long dataset of hydrography and velocity observations from two mooring arrays, supplemented by output from an eddy-resolving ocean model, to quantify the seasonal variability of eddy kinetic energy (EKE) and baroclinic and barotropic energy conversion rates over time-scales from days to months. Both EKE and conversion rates peak in autumn and winter, coinciding with the strongest boundary current and the warmest AW. Local EKE variability, however, is only weakly associated with conversion rates, suggesting advection from upstream generation sites or unresolved variability from limited measurements. Conversion is mainly baroclinic, ~~indicating through~~ boundary current instability~~and~~, providing conditions favourable for offshore propagation of warm-core eddies. The model ~~underscores the need for adequate spatial output~~ allows adequate volume and temporal averaging~~and reveals~~, and reveals spatial structure with substantial conversion rates on the offshore, deeper side of the boundary current, with comparable contributions from baroclinic and barotropic processes. Resulting mesoscale activity ~~likely~~ enhances lateral stirring and heat loss from the boundary current, particularly in winter and spring, contributing to the along-stream cooling of AW.

1 Introduction

Atlantic Water (AW) is the primary oceanic source of heat for the Arctic Ocean and plays a central role in shaping its changing physical environment (e.g., Carmack et al., 2015) and ecosystems (Ardyna and Arrigo, 2020). The largest volume of warm AW enters the Arctic Ocean with the West Spitsbergen Current (WSC), which flows through Fram Strait – the main deep gateway connecting the Arctic Ocean to the global oceans (Aagaard et al., 1987; Beszczynska-Möller et al., 2012). The AW flow splits into several poleward and recirculating branches as it approaches the Yermak Plateau (Fig. 1), and the inflow eventually merges downstream and contributes to the Arctic Circumpolar Boundary Current (Rudels et al., 1999). The region north of

Svalbard has been shown to have far-reaching signatures in the Arctic Ocean (Polyakov et al., 2017), such as recent changes reported hundreds of kilometres downstream in the Eurasian Basin, including warming of the AW core (Richards et al., 2022), weakening of the halocline and shoaling of the AW (Polyakov et al., 2020b, a), thus affecting the exchange with sea ice and the surface layer.

The continental slope north of Svalbard is an important area for the modification of AW in the boundary current. The drivers of water mass transformation include air-sea-ice interactions and mixing (Renner et al., 2018; Kolås et al., 2020; Pérez-Hernández et al., 2019; Koenig et al., 2022b), entrainment of surface waters (Richards et al., 2022) and shelf waters (Schauer et al., 1997), and mesoscale variability and eddies (Koenig et al., 2022b; Crews et al., 2018; Wekerle et al., 2020; Athanase et al., 2020; Pérez-Hernández et al., 2017). We define the mesoscale as low Rossby number flows ($Ro = U/fL < 1$, where U and L are the characteristic horizontal velocity and length-scales and f is the Coriolis frequency) that occur on time-scales of days to months and length-scales of 10 km to 100 km. In particular, mesoscale eddies [in this region](#) facilitate slope-basin exchange and lateral heat loss (Våge et al., 2016; Pérez-Hernández et al., 2017; Renner et al., 2018). In this study, we investigate the mesoscale eddy-variability and energy conversion rates that transfer energy from the mean flow and stratification into eddy energy, using moored observations and a high-resolution numerical ocean model north of Svalbard.

Mesoscale eddies can form through barotropic ~~and baroclinic~~ and [baroclinic](#) instabilities of the flow (~~Cushman-Roisin and Beckers, 2011~~ [\(Killworth, 1980\)](#)). Barotropic instability may arise from horizontal velocity shear, drawing kinetic energy from the mean flow and converting it into eddy kinetic energy (EKE) (e.g., Cushman-Roisin and Beckers, 2011; Teigen et al., 2010). For a boundary current over a steep slope, the shape of the current determines whether barotropic instability can form. Steep slopes stabilise the current (von Appen et al., 2016), requiring the current to be narrow and fast to become unstable (Håvik et al., 2017). Baroclinic instability, in contrast, may form in regions with a horizontal density gradient, hence strong thermal-wind shear, and converts available potential energy into eddy energy, which enforces flattening of the pycnocline. Bottom slopes also impact (suppress) baroclinic instability (Isachsen et al., 2024).

In the Arctic Ocean, eddies play an important role in variability ~~, transport of inflow waters, and of inflow water properties~~ [and transport and](#) ice–ocean interaction. High mesoscale variability is observed over continental slopes, in Fram Strait, and in the Arctic Circumpolar Boundary Current (von Appen et al., 2022; Wang et al., 2020). These regions, together with the Barents Sea, have relatively higher EKE and baroclinic conversion compared to the rest of the Arctic Ocean (Li et al., 2024). Eddies that shed from unstable fronts along the boundary of the Arctic Ocean transport inflow waters across the basin (Carmack et al., 2015). Eddies also mediate ice–ocean interaction. Under-ice cyclonic eddies can bring AW upward into the surface layer and towards the ice, increasing the heat fluxes and enhancing melting (Müller et al., 2024). Cyclonic eddies may trap and advect sea ice into warmer waters (Manucharyan and Thompson, 2017). Sea ice has been shown to dampen near-surface eddies (Meneghello et al., 2020). The ongoing decline of Arctic sea ice may reduce this effect, causing an increase in near-surface eddy activity and EKE (Armitage et al., 2020; Manucharyan and Thompson, 2022; Müller et al., 2024; Li et al., 2024).

In our study region, north of Svalbard, eddies have been registered both in situ and in models. Scarce observations show evidence for anticyclonic eddies carrying warm anomalies offshore from the shelf break (Våge et al., 2016). These eddies laterally stir and redistribute heat, and thereby contribute to the heat loss from the boundary current (Renner et al., 2018; Koenig




Figure 1. a) Bathymetry (shading and contours every 500 m; IBCAO version 4, Jakobsson et al. (2020)) and circulation of AW (red arrows) in Fram Strait and north of Svalbard. The black box outlines the map in (b), the red box shows the model subdomain. The inset in (a) shows the location in the Arctic Ocean (BS: Barents Sea, GR: Greenland). b) The study region showing the bathymetry and mooring positions.

et al., 2022b). Estimates of along-path heat loss of AW revealed that turbulent heat fluxes and heat loss to the atmosphere alone could not account for the observed cooling, suggesting a substantial contribution through lateral heat loss (Koenig et al., 2022b). High-resolution model results support this view, showing anticyclonic eddies carrying AW spawning from the boundary current (Crews et al., 2018), propagating offshore towards the deeper basin (Wekerle et al., 2020), facilitating slope–basin exchange and lateral transport of AW. However, the seasonal magnitude and variability of the associated energy conversion rates have not been previously reported from observations in this area.

In the Nansen LEGACY project, the slope north of Svalbard was chosen for targeted studies, including mooring arrays to quantify the properties of the boundary current, specifically its volume ~~and heat transport, along-slope~~ transport and along-path cooling (Koenig et al., 2022b), and the mesoscale variability (this study). These detailed observations are analysed here to quantify energy conversion rates and are supplemented using outputs from an eddy-resolving ocean model. The combined results provide insight into where and how the boundary current flow energises variability at mesoscale eddy scales that could contribute to the lateral heat loss from the boundary current.

2 Data and Methods

2.1 Moorings

Ocean temperature, salinity, and horizontal current data ~~obtained from~~ were measured at two mooring arrays across the continental slope north of Svalbard ~~are used in this study~~ (Fig. 1). ~~Measurements were collected,~~ for approximately one year from September 2018. The western array (18° E) was recovered in September 2019, and the eastern array (24° E) was recovered in

Table 1. Mooring positions, total depth, and temporal coverage of the deeper mooring pairs used in this study.

Mooring	Latitude	Longitude	Depth (m)	Deployed-Recovered Temporal coverage
W2	N 81°22.686'	E 18°23.789'	727	15.09.2018 1420 –21.09.2018 0800
W3	N 81°27.356'	E 18°23.730'	1202	20.09.2018 1810 –21.09.2018 1330
E2	N 81°30.813'	E 23°59.853'	706	16.09.2018 1120 –23.11.2019 1905
E3	N 81°35.453'	E 23°59.982'	1222	16.09.2018 1445 –23.11.2019 2150

November 2019. The data are available from Fer et al. (2022) and have been presented in Koenig et al. (2022b). Each mooring array consisted of one upper slope mooring (W1 and E1) near the 300 m – 400 m isobath, one middle slope mooring (W2 and E2) near the 700 m isobath, and one lower slope mooring (W3 and E3) near the 1200 m isobath, providing a good coverage of the boundary current (Fig. 1b). In addition, a mooring was deployed onshore of the western array (W0). In this study, we used data from the deeper ~~pair of moorings, mooring pairs~~ W2–W3 and E2–E3 (Table 1) ~~–as the distance within each pair – approximately 9 km – is small enough for calculating lateral gradients (see Section 2.3.3).~~

Mooring instrumentation coverage and setup, and details of data processing can be found in Koenig et al. (2022a). ~~The vertical coverage of instrumentation at the moorings for September–February and March–August is summarised in Fig. 2.~~ Temperature, salinity, and horizontal current records were checked for inter-consistency and compared and corrected against ship-based profiles taken at mooring deployment/recovery. All time series were hourly averaged (the sampling interval of instruments varied between 5 min and 1 hour) and linearly interpolated onto the same hourly time vector. At each hourly time step, instrument depths were obtained by linear interpolation between instruments equipped with a pressure sensor, to account for mooring knockdowns. The data were finally gridded onto a regular (t, z) grid with 1 h temporal and 10 m vertical resolution.

~~The analysis is based on data from the deeper mooring pairs which provided current profiles with a 5 m–10 m vertical resolution from the seafloor to approximately 300 m depth. Here, only three levels of density measurements at approximately 300 m, 500 m, and 700 m depth were available. Our analysis is therefore based on layer averages over 300 m to 700 m (Section 2.3.3), focusing on the levels close to the CTD sensors to limit the errors from interpolation. For context, we present seasonal mean sections of velocity and hydrography at both mooring arrays (Fig. 2). For computing mean structure of the current velocity and hydrography across the slope, a spline-Laplacian routine was used for interpolation, described in detail in Koenig et al. (2022b). While current measurements cover the water column at W1, there were no temperature and salinity measurements available, and the hydrographic structure over the upper slope was supplemented by historical hydrographic data. The details on the gridding and the associated uncertainties are described in the Appendix of Koenig et al. (2022b). The uncertainties in constructing the sections do not affect the analysis in this paper.~~

2.2 Regional Ocean Model

In order to better evaluate and interpret the analysis based on the mooring observations, we used the output from a high-resolution ocean model applying ROMS (Regional Ocean Modeling System; Shchepetkin and McWilliams (2005)) in a domain

100 covering the continental shelf and slope north of Svalbard ([Frank et al., 2025a](#))([Frank et al., 2025b](#)). The model used the same configuration as the Norkyst model system documented in Asplin et al. (2020) but was set up with a horizontal resolution of 500 m and 35 vertical [stretched](#) bathymetry-following levels. The model configuration also included a sea ice module (Budgell, 2005). The operational ocean forecasting model Barents2.5 (Röhrs et al., 2023) provided boundary conditions including tidal forcing, and the operational weather forecasting model AROME-Arctic (Müller et al., 2017) provided atmospheric input. The
105 model [was validated by Frank et al. \(2025b\), and a brief summary of this validation is provided in Appendix B.](#)

[The model](#) simulation was initialised in April 2019, and we used two-year daily output from October 2019 to align seasonally with the mooring observations. Hourly saved fields were daily averaged, and the daily fields were then detided for the fortnightly and monthly tidal constituents using UTide (Codiga, 2011). [The across-slope structure of the modelled boundary current and hydrography is shown in Fig. B1 and compared with the mooring-based sections in Appendix B.](#)

110 2.3 Methods for energy analysis

2.3.1 Wavelet analysis

In order to analyse the time-variability of energetic frequency bands, we used wavelet transforms with generalised Morse wavelets following Lilly and Olhede (2010, 2012) with the parameters $\gamma = 3$ for symmetric wavelets (Lilly and Olhede, 2009) and $\beta = 9$ for a reasonable time and frequency resolution over the relevant time-scales studied here. We denote the wavelet
115 transform of a time series x as W_x and its complex conjugate as W_x^* .

2.3.2 Eddy kinetic energy and energy conversion rates

The calculations of eddy energetics and conversion rates require definition of fluctuations, background conditions, and averaging. We use primed values (e.g., u', ρ') to denote fluctuations from the mean, and an overbar to indicate the background conditions and time averaging (e.g., $\bar{u}, \bar{\rho}$). For the moorings, fluctuations were obtained by band-pass filtering the hourly time
120 series using cut-off frequencies corresponding to 35 hours and two weeks, and the averaging was done over 30-day moving windows. For the background conditions, we used a low-pass filter with a cut-off frequency corresponding to 30 days. From the model output, primed values were obtained by Reynolds-decomposition, e.g., $u' = u - \bar{u}$, where $\bar{u}$ values are 30-day means centred on the 15th of each month and u' values are daily fluctuations from the mean.

Eddy kinetic energy density (EKE), the barotropic energy conversion (BT), and the baroclinic energy conversion (BC) rates
125 are defined as

$$\text{EKE} = \frac{1}{2} \left(\overline{u'^2} + \overline{v'^2} \right), \quad (1)$$

$$\text{BT} = -\rho_0 \left(\overline{u'u'} \frac{\partial \bar{u}}{\partial x} + \overline{u'v'} \left(\frac{\partial \bar{u}}{\partial y} + \frac{\partial \bar{v}}{\partial x} \right) + \overline{v'v'} \frac{\partial \bar{v}}{\partial y} \right), \quad (2)$$

and

$$BC = g \left(\frac{\partial \bar{\rho}}{\partial z} \right)^{-1} \left(\overline{u' \rho'} \frac{\partial \bar{\rho}}{\partial x} + \overline{v' \rho'} \frac{\partial \bar{\rho}}{\partial y} \right), \quad (3)$$

130 where $\rho_0 = 1027 \text{ kg m}^{-3}$ is a reference density, $\overline{u'u'}$, $\overline{u'v'}$, and $\overline{v'v'}$ are eddy momentum fluxes, g is the gravitational acceleration, $\overline{u'\rho'}$, $\overline{v'\rho'}$ are eddy density fluxes, $\partial/\partial y$ and $\partial/\partial x$ denote lateral gradients, and $\partial \bar{\rho}/\partial z$ is the mean vertical stratification.

In the model subdomain, EKE, BT, and BC were calculated on the model grid. Then, x and y denote the right-handed orthogonal coordinates in the native model grid, and u and v the velocity components in these directions. For the moorings, x and y denote the local along- and across-slope directions, and u and v the along- and across-slope current components. The
 135 coordinate system was rotated with 20° (i.e., counter-clockwise) at the western and -5° (i.e., clockwise) at the eastern array, respectively.

2.3.3 Simplified calculations for conversion rates

Conversion rate estimates from the mooring records were made using simplified forms of Equations 2 and 3, with the common assumptions that the across-isobath gradients dominate and that there is no along-isobath variability:

$$140 \quad BT = -\rho_0 \overline{u'v'} \frac{\partial \bar{u}}{\partial y}, \quad (4)$$

$$BC = g \overline{v' \rho'} \frac{dz}{dy}, \quad \frac{dz}{dy} = \frac{\partial \bar{\rho}}{\partial y} / \frac{\partial \bar{\rho}}{\partial z}. \quad (5)$$

In BT (Eq. 4), the lateral gradient was obtained by first differencing of ~~$\overline{u}$~~ that was layer-averaged between 300 m and 700 m (280 m and 680 m at the eastern array), and the eddy momentum flux was calculated at each mooring and then averaged to be representative of the flux between the moorings. For layer averages, standard error is calculated as $SE(t) = \sigma(t)/\sqrt{n(t)}$,
 145 where $n(t)$ is the number of estimations at time t , and $\sigma(t)$ is the standard deviation of the n estimations. In BC (Eq. 5), dz/dy is the mean isopycnal slope, estimated by the quotient of the across-slope gradient of the mean density $\partial \bar{\rho}/\partial y$ and the mean stratification $\partial \bar{\rho}/\partial z$. The potential density ρ was calculated from gridded T and S fields, at selected depths near the target depths of the sensors, to minimise error from linear interpolation across the halocline.

Calculations of BT and BC require adequate lateral gradient estimations. First, the separation between two moorings must be sufficiently small, limiting the analysis to the deeper moorings pairs. Second, simultaneous measurements at both moorings are needed, constraining the analysis to the 300 m to 700 m layer because of substantial data gaps higher in the water column at the deepest moorings. EKE and eddy momentum and density fluxes can be calculated over a broader depth range at all seven moorings (Fig. 1), capturing more of the energetic part of the water column where the data loss was greatest. Their vertical structure is shown in Appendix A.
 150

155 We conducted additional calculations of BT and BC from the model output using the simplified equations (Eqs. 4 and -5) applied to the time series of velocity and density extracted at ~~locations~~ grid points representative of the moorings (“virtual moorings”) as well as ~~70~~along 16 km and ~~50~~13 km transects (“segments”) with ~~101 and 71~~ 24 and 19 grid points across the

deeper part of the slope, covering the virtual moorings. Virtual mooring calculations mimic the data and the method used for the in situ moorings, which we compare to the full volume-averaged or segment-averaged conversion rates in order to assess their limitations.

2.3.4 Calculations along the boundary current pathway

We analysed the EKE and energy conversion rates, and eddy temperature fluxes $(\overline{u'T'}, \overline{v'T'})$ obtained from the model to describe the temporal and spatial variability along the pathway of the boundary current. Upon inspection of the model outputs, we identified the 1400 m isobath to be representative for the energetic part of the boundary current on the upper slope. The structure along the isobath is obtained by smoothing the monthly fields using a 2D Gaussian filter with a 20 km length-scale and interpolating the results onto the isobath. The smoothing length-scale ensures that we capture the boundary current structure between, typically, 800 m and 2000 m isobaths.

As an indication for conditions that might allow barotropic instability along the boundary current pathway, we calculate the cross-isobath gradient of potential vorticity, $q_y = \beta_{\text{topo}} - \partial^2 \bar{u} / \partial y^2$, where $\beta_{\text{topo}} = -(f/H) \partial H / \partial y$ is the topographic β , H is the water depth, f is the Coriolis parameter, u is the along-stream current velocity component, and y is the across-slope direction. A necessary, but not sufficient, condition for barotropic instability is that q_y must change sign within the domain (Vallis and Maltrud, 1993). Along the 1400 km isobath, we calculated q_y on 20 km lines oriented perpendicular to and centred on the isobath. To approximately visualise this instability condition, we show the product of the minimum and maximum values of q_y calculated along each line, i.e., $f(q_y) \equiv \min(q_y) \cdot \max(q_y)$, as a change in sign of q_y would result in negative values.

3 Results

3.1 Seasonal and mesoscale variability of the boundary current

Mean structure and seasonal variability of the boundary current observed in the mooring arrays have been reported in Koenig et al. (2022b). Here, we summarise the mean hydrography and current structure relevant to our analysis (Fig. 2) using the three-monthly averages from Koenig et al. (2022b) but grouped into September–February and March–August periods.

The boundary current was on average stronger from September to February and weaker from March to August (Fig. 2). In September–February, the velocity core was at around 100 m depth over the 800 m isobath, carrying AW warmer than 3 °C at a speed exceeding 20 cm s⁻¹ (Fig. 2a–b). In March–August, the velocity core was weaker (10 cm s⁻¹ to 20 cm s⁻¹), deeper, and relatively diffuse in both vertical and lateral directions (Fig. 2c–d). Water with the highest temperatures, exceeding 2.5 °C, was at 200 m depth seawards of the 500 m isobath, but did not coincide with the deeper velocity core.

The mesoscale activity was elevated variability is demonstrated using records from W2, and shows elevated activity in autumn and winter (Fig. 3), in the same period when the boundary current was strongest and carried the warmest water (Fig. 2a–b). The strongest velocity variability occurred on time-scales between 2 and 6 days and, like the elevated mesoscale




Figure 2. Average along-slope current velocity u (colours) on (a, c) the western and (b, d) eastern mooring arrays during (a, b) autumn and winter (September–February) and (c, d) spring and summer (March–August). **Gray** contours show velocity every 10 cm s^{-1} , white dashed isopycnals every 0.1 kg m^{-3} , and black isotherms every 1°C . Triangles at the top mark the mooring locations (except W0). **The two deeper instrumentation is shown at the mooring pairs used locations as indicated in this study are illustrated in the lower panels: vertical legend. Vertical bars show the current profiling coverage by the ADCPs (light gray: 50–80% and dark gray: 90–100% of the time in each season); large circles indicate CTD sensors, small circles temperature sensors, triangles upward-facing ADCPs, diamonds pairs of up- and down-facing ADCPs, and squares RCMs.** Wind roses (right) show the relative frequency of wind direction and speed during (a, b) September–February and (c, d) March–August in the region $18^\circ \text{ E}–24.5^\circ \text{ E}$, $81.2^\circ \text{ N}–81.7^\circ \text{ N}$, extracted from the Copernicus C3S Arctic Regional Reanalysis (CARRA, Schyberg et al., 2020).

activity, was observed in autumn and winter (Fig. 3a, c). From October 2018 to January 2019, the offshore velocity component v and the temperature T co-varied on a 3–10-day time-scale (Fig. 3d). At that time, the current core was located near the 800 m isobath, close to W2, and carried AW (Fig. 2a).

The offshore eddy temperature flux, $\overline{v'T'}$, at 250 m depth was positive in the autumn of 2018, peaking in mid-November, and became negative in December 2018 (Fig. 3b), consistent with mesoscale activity driving cross-isobath heat transport. Averaged over September–November 2018, $\overline{v'T'}$ at W2 was positive at depths 150 m to 350 m, spanning the temperature core. At W3, the eddy temperature flux in the upper water column could not be calculated in autumn due to mooring knockdown, but was positive at and below 400 m depth (not shown). The eddy temperature flux divergence between W3 and W2 was over four




Figure 3. Time series of a) along-slope and across-slope current velocity u , v (left axis; v is offset by -10 cm s^{-1}), and temperature T (right axis, purple) measured at 250 m depth at the western mooring W2 (a level in the core which consistently has both temperature and velocity measurements), low-pass filtered at 35 hours (thin curves) and 30 days (thick curves). Corresponding b) offshore eddy temperature flux $\overline{v'T'}$ (positive for warm anomalies towards deeper water and cold anomalies towards shallower water), c) wavelet power spectrum, $\frac{1}{2} (|W_u|^2 + |W_v|^2)$, and d) wavelet cross-spectrum $|W_v \times W_T^*|$ between v and T . The cone of influence (solid white curve) in (c, d) indicates the areas affected by boundary effects.

times higher in autumn and winter (September–February) than in spring and summer (March–August; [not shown](#)), suggesting stronger lateral heat loss from the boundary current in autumn and winter.

While the relatively higher eddy temperature flux and [divergence](#) in autumn and winter indicate increased mesoscale activity, we cannot directly quantify net lateral heat transport because the rotational and the divergent components of the flux could not be separated (Marshall and Shutts, 1981; Guo et al., 2014).

3.2 Energetics and conversion rates from mooring data

[Despite their limited spatial coverage and sampling volume, the moorings enable year-round estimates of eddy kinetic energy and associated energy conversion rates. EKE and BT, together with its contributing terms, are shown in Fig. 4. BC and its contributing terms are shown in Fig. 5. To provide context for the magnitudes of conversion rates and their effect on changes in EKE, note that a conversion rate of \$1 \times 10^{-5} \text{ W m}^{-3}\$ sustained over one day would produce \$1 \times 10^{-3} \text{ J kg}^{-1}\$ corresponding to an EKE increase of \$10 \text{ cm}^2 \text{ s}^{-2}\$.](#)




Figure 4. Time series of (a, b) EKE, (c, d) lateral shear of the mean along-slope current, (e, f) eddy momentum flux, and (g, h) BT at the (left) western and (right) eastern arrays. Blue curves are values at W2 and E2, red curves at W3 and E3, and black curves for the average between the pair of moorings. Shading shows standard error calculated over estimations at every hundred meters between 300 m and 700 m depth.

Several features are consistent at both mooring arrays. EKE was higher in autumn and winter than in spring and summer at both mooring arrays (Fig. 4a, b), and it was surface intensified with values at 300 m depth about four times higher approximately four times larger than at 700 m (not shown). In contrast, BT remained low-weak year-round at both sites relative to the conversion rate expected from the observed EKE variability (Fig. 4g, h). It was negligible in summer and varying otherwise fluctuated within $\pm 0.5 \times 10^{-5} \text{W m}^{-3}$ otherwise. The low BT resulted from a combination of averaging. The persistently low BT reflects both cancellation of oppositely signed momentum fluxes (Fig. 4e, f) and occasionally low velocity shears periods of weak velocity shear (Fig. 4c, d). BC was generally largest in autumn and winter at both sites, ranging from $-3 \times 10^{-5} \text{W m}^{-3}$ to $4 \times 10^{-5} \text{W m}^{-3}$, and decreasing to near-zero values in spring and summer (Fig. 5e, f). BC was generally strongest at the uppermost measurement level (310 m–320 m), where variability in density and cross-slope velocity was greatest, and decreased with depth (Fig. 5c, d). Despite these similarities, BC estimates showed substantial variability between depths and

moorings, particularly in autumn and winter, primarily due to differences in eddy density fluxes despite being measured only 100 m apart vertically, e.g. in October and November 2018 (Fig. 5c, d).

220 At the western array, the autumn- and wintertime EKE reached nearly $40 \text{ cm}^2 \text{ s}^{-2}$, ~~while the summertime EKE dropped and decreased to~~ below $20 \text{ cm}^2 \text{ s}^{-2}$ ~~in summer~~ (Fig. 4a). BT was ~~mostly low except for two events: generally weak, with two notable exceptions.~~ In January 2019, BT was positive (Fig. 4g) ~~when the values in the upper water column. Based on estimates at 100 m intervals (not shown), the values at the uppermost level (300 m) reached $1 \times 10^{-5} \text{ W m}^{-3}$ due to negative momentum flux and driven by negative momentum fluxes combined with~~ positive shear. ~~These decreased. This signal weakened~~ with depth, reducing the layer-average BT. In late March 2019, ~~simultaneous a second event occurred, when~~ strong positive velocity shear and positive momentum flux at W2 ~~resulted in produced~~ a similarly large but negative and ~~shorter-lasting shorter-lived~~ BT (Fig. 4g, c, e).

The eastern array had generally lower EKE than the western array, but with a ~~similar seasonal contrast: autumn–winter values reached or exceeded $20 \text{ cm}^2 \text{ s}^{-2}$, peaking at $35 \text{ cm}^2 \text{ s}^{-2}$ in February 2019 (Fig. 4b).~~ BT in the eastern array was quite 230 ~~similar to BT in the west, generally low but with two events. The eastern conversion events appeared comparable seasonal contrast. BT showed a similar overall behaviour to the western array. Typically low values increased in two events, but with opposite signs as the ones in the to the~~ western array. BT was negative in January 2019 due to modest momentum fluxes and shear, while from late March to early April 2019, BT was positive due to a period of negative momentum fluxes and an offshore ~~drift displacement~~ of the boundary current (Fig. 4h, d, f). ~~When EKE peaked in February, BT was During the February EKE~~ 235 ~~maximum, BT remained~~ negligible due to near-zero shear and ~~momentum~~ fluxes.

BC was generally largest in autumn and winter at both mooring arrays, ranging within $-3 \times 10^{-5} \text{ W m}^{-3}$ and $4 \times 10^{-5} \text{ W m}^{-3}$, decaying to negligible values through spring and summer (Fig. 5e, f). BC values were typically largest in the upper water column, due to higher fluctuations in density and cross-slope velocity, and decreased with depth (Fig. 5e, d). Notably, BC estimates differed substantially between depths and moorings, especially during autumn and winter (Fig. 5e, f), mainly due 240 ~~to differences in eddy density fluxes despite being measured only 100 m apart vertically, e.g. in October and November 2018 (Fig. 5e, d).~~

At the western array, the most notable BC event occurred at 320 m at W2 from mid-October through November 2018. BC increased from negligible values to $4 \times 10^{-5} \text{ W m}^{-3}$ and remained ~~high elevated~~ for nearly a month. This event ~~was associated coincided~~ with a strong mean current, high EKE (Fig. 4a), a modest and ~~steady persistent~~ negative isopycnal slope, 245 and an elevated eddy density flux ~~due to from~~ an increased variability and ~~co-variability covariance~~ of v and ρ ~~of periodicity, with dominant periods of 2–10 days (similar to Fig. 3d).~~ During and following this event, EKE at W2 ~~was high and peaked remained high, peaking~~ at $88 \text{ cm}^2 \text{ s}^{-2}$ at 300 m depth. However, such ~~coincident concurrent~~ increases in EKE and BC were not ~~observed frequently commonly observed~~. A similar but shorter-lasting event ~~was observed occurred~~ at 310 m depth at E2 from mid-November to early December 2018 (Fig. 5f). ~~Here, an eddy density flux of $-4 \times 10^{-4} \text{ kg m}^{-2} \text{ s}^{-1}$ resulted in BC~~ 250 ~~exceeding $4 \times 10^{-5} \text{ W m}^{-3}$. EKE at 280 m depth reached a local maximum of $55 \text{ cm}^2 \text{ s}^{-2}$ during this event (not shown).~~

In summary, ~~the~~ current strength, temperature variability, EKE, ~~the~~ eddy momentum and density fluxes, BC, and to ~~some degree a lesser extent~~ BT, were all generally stronger in autumn and winter ~~compared to than in~~ spring and summer. ~~Although~~




Figure 5. Time series of (a, b) isopycnal slope, (c, d) eddy density flux, and (e, f) BC at the (left) western and (right) eastern arrays. A positive isopycnal slope occurs when density decreases with offshore distance.

the measurements do not resolve the upper ocean and surface layer, the available vertical coverage indicates a tendency for enhanced EKE, BT, and BC were generally surface intensified at the uppermost measurement levels. While EKE was relatively enhanced during the months with often elevated during periods of higher fluxes and BC, there was no consistent or strong the relationship between them was neither consistent nor robust.

Advection of eddies past the moorings complicates the interpretation of our estimates in several ways. A fraction of the calculated EKE may be attributed to eddies advected by the mean flow, adding to the local conversion from mean to eddy energy, potentially leading to an overestimation of locally generated EKE. In addition, the use of a band-pass filter to estimate fluctuations may attenuate contributions from mesoscale features when the filter cutoff exceeds the eddy advection time scale. The advection time scale is 24–40 h, based on along-slope velocities of 10 cm s^{-1} to 15 cm s^{-1} and an eddy lateral length scale of twice the baroclinic Rossby radius of deformation ($\sim 7 \text{ km}$). This range overlaps with, and partly exceeds, the 35 h cutoff used in the EKE calculations. Alternative calculations using a shorter cutoff of $\approx 24 \text{ h}$, excluding the diurnal tidal band while retaining most mesoscale variability, show no significant differences (Appendix C; Fig. C1a, b).

3.3 Energetics and conversion rates from the ocean model data

We analyse the output from the high-resolution model to evaluate assess whether the conversion estimates using the simplified expressions for BT and BC from, based on limited depth levels and locations of the mooring records mooring locations, are

representative for a larger area. In order to assess this broader region. To do so, we calculate EKE, BT, and BC using all terms of the conversion equations the full conversion terms (Eqs. 2, 3) and averaging in the vertical vertically average between 100 m and 2000 m depth to obtain the to obtain their spatial structure (Fig. 6 and Section 3.3.1). We next then show the temporal evolution of the energetics averaged along the along the boundary current pathway by averaging over a 20-km diameter region centered on the 1400 m isobath (Fig. 7 and Section 3.3.2). Additionally, we use the simplified equations apply the simplified formulations (Eqs. 4, 5) applied to time series of to velocity and density extracted at the mooring locations from the model (“virtual moorings”) time series extracted at virtual moorings, using the same calculation method as for the moorings (Fig. 8) and Section 3.3.3). These results are compared with both the observational estimates and the spatially averaged values along the boundary current pathway.

3.3.1 Mean spatial distribution




Figure 6. Maps of the two-year mean (October 2019 to September 2021) model fields, averaged from 100 m to the seafloor or to 2000 m depth if deeper, for (a) EKE current speed, (b) EKE, (c) BT, and (d) BC. To highlight the AW boundary current in (a), velocity vectors are shown between the model subdomain averaged over 600 m and 2800 m isobaths, and the two years from October 2019 to September 2021. 100–500 m layer-averaged 2°C isotherm is shown with purple curves. The positions of the virtual moorings are marked with circles. Isobaths are drawn every 400 m between 200 m and 2600 m (gray), and the 1400 m isobath is highlighted in black. Black dots mark the distance along the 1400 m isobath every 5025 km starting at 0 km with the first dot. The colour bar ranges are set to the 99.5 percentiles of the time-averaged fields. In (a), the circle at 20° E of diameter 20 km shows the length-scale of the spatial filtering applied in Fig. 7.

The core of the AW boundary current is evident in the vertically and temporally averaged current speed, located over the upper continental slope between the 1000 and 1400 m isobaths, with peak velocities exceeding 24 cm s^{-1} (Fig. 6a). The mooring pairs capture this boundary current structure well. Waters warmer than 2°C within the 100–500 m depth range extend from the lower continental slope across the current core and onshore toward the shelf, where the mean flow gradually weakens. Applied 20-km diameter averaging along the 1400 m isobath covers the core of the boundary current carrying AW and captures

the energetic part of the slope. EKE values along the slope, averaged over 2 years, were typically $20\text{--}30\text{--}50\text{ cm}^2\text{ s}^{-2}$, with higher values, up to $45\text{--}60\text{ cm}^2\text{ s}^{-2}$, e.g. around and west of the western mooring array around and upstream of both mooring arrays (Fig. 6a). The largest BT and BC magnitudes were located b).

Time-averaged BT had strong spatial variability along the slope (Fig. 6c). The largest maxima were found on the steep slope offshore of the 1000 m isobath, with particularly elevated values in the region of complex bathymetry upstream of the western array, where local values reached $15\text{--}30 \times 10^{-5}\text{ W m}^{-3}$ (Fig. 6b, e). Areas where the two-year mean EKE, BT, and BC all exceeded high-percentile values were primarily located near between the 1000 m and 1800 m isobaths in the vicinity c, these values exceed the colorbar range). Other strong positive regions were found near the 1400 m isobath, approximately 20 km downstream of the western mooring array. Such areas tended to be more localised for BT and BC but parts of larger maxima for EKE array ($25 \times 10^{-5}\text{ W m}^{-3}$), and approximately 20 km upstream of the eastern array ($15 \times 10^{-5}\text{ W m}^{-3}$). The zones of BT maxima were anisotropic, elongated in the across-isobath direction. Aside from the strong positive maxima, time-averaged BT alternated in sign along the slope, characterized by transitions between local minima and maxima, often associated with variations in slope geometry and isobath curvature.

Along the slope, BT showed a pattern with positive values where isobaths veer leftward downstream, and negative values in areas where isobaths veer rightward.

BC had comparable spatial variability and magnitude to BT (Fig. 6b). These zones were persistent in time (not shown). An example of two such oppositely signed zones is d). The largest time-mean BC values were also found on the steep slope near the western mooring array where the values reached $\pm 7 \times 10^{-5}\text{ W m}^{-3}$ in the two-year average. A particularly strong positive BT region was. A pronounced BC maximum occurred near the 1400 m isobath approximately 20 km downstream from of the western array, where the average BT exceeded $20 \times 10^{-5}\text{ W m}^{-3}$ coincident with the strong BT maximum there, though with roughly half the magnitude.

BC had comparable spatial variability and magnitude to BT (Fig. 6c). A hot-spot upstream of the western array averaged to $24 \times 10^{-5}\text{ W m}^{-3}$, coincident with substantial EKE. The strong BT region 20 km downstream, also co-located with BT, was upstream of the western array also had large BC occurring near the 1400 m isobaths.

Several of the regions along the slope that were energetic on average were associated with maxima in time-mean conversion rates. In some energetic locations, BT and BC were co-located or overlapping, while in others, primarily one of the conversion types dominated.

3.3.2 Structure along the boundary current pathway

Along the 1400 m isobath, EKE, BT, and BC varied both spatially and temporally (Fig. 7). Time-mean EKE values were elevated near the two mooring arrays and approximately 40 km upstream, but was slightly elevated over broad regions near both mooring arrays, with the highest mean EKE occurred at the southwestern boundary of the domain value of $50\text{ cm}^2\text{ s}^{-2}$ occurring 30 km upstream of the western array (Fig. 6ab, Fig. 7a). Monthly averages over the span full 180 km extent of the isobath ranged from below $20\text{ cm}^2\text{ s}^{-2}$ to above 60 in month 20 (May) to $65\text{ cm}^2\text{ s}^{-2}$ in month 14 (November).

BT averaged to $0.34 \times 10^{-5} \text{ W m}^{-3}$. When averaged along the 1400 m isobath over the full period, its time-mean was mostly positive, ranging from -0.9×10^{-5} . BT was $0.47 \times 10^{-5} \text{ W m}^{-3}$, one order of magnitude larger than BC ($0.05 \times 10^{-5} \text{ W m}^{-3}$ to 2.2×10^{-5}). Both conversion terms displayed substantial along-isobath variability. Time-mean BT ranged from a negative value of $-1.7 \times 10^{-5} \text{ W m}^{-3}$, peaking about in the region between the mooring arrays to a maximum positive rate of $3.5 \times 10^{-5} \text{ W m}^{-3}$ located 20 km downstream of the western array, with a secondary maximum of 1.7×10^{-5} mooring array. A secondary BT maximum of $2.8 \times 10^{-5} \text{ W m}^{-3}$ located was located about 10 km upstream of the western array. Further east, two additional local maxima were found 15 km downstream-upstream of the eastern array ($2.5 \times 10^{-5} \text{ W m}^{-3}$) and at the array ($1.4 \times 10^{-5} \text{ W m}^{-3}$). When averaged Along the isobath, BT remained generally positive and reached its maximum value of 1.5×10^{-5} reached a maximum of $2.0 \times 10^{-5} \text{ W m}^{-3}$ in month 14, which coincided with the EKE maximum. No peak. The minimum value of $-0.3 \times 10^{-5} \text{ W m}^{-3}$ occurred in month 10 (July). Overall, no clear seasonality was observed.

The time-mean BC varied within $\pm 2 \times 10^{-5}$ ($-2.1, +2.1$) $\times 10^{-5} \text{ W m}^{-3}$. The largest positive BC was approximately 10 km upstream and in a wide zone 15 km to 45 km of the western array, coincident with a BT local maximum, and within a broad region extending 15–40 km downstream of the western array. Near the eastern array, where BC remained above $1.0 \times 10^{-5} \text{ W m}^{-3}$. In contrast, BC was negative on average near the eastern array. Averaged along the isobath, BC was typically within $\pm 0.5 \times 10^{-5}$ varied within $(-0.6, +0.9) \times 10^{-5} \text{ W m}^{-3}$, except for a couple of larger minima and a peak of $0.6 \times 10^{-5} \text{ W m}^{-3}$ in month 14, again coinciding with the EKE maximum.

Although there were occasional coincidences between peaks in EKE and conversion rates on the 1400 m isobath, the variability of EKE generally did not match the local conversion rates. This lack of a robust relation between EKE variability and conversion rates mirrors the findings in the mooring observations and suggests that increasing EKE in a region does not necessarily indicate that strong local energy conversion is taking place. While there is not a conclusive, direct correspondence between local conversion rates and EKE variability, several segments along the isobath exhibit statistically significant correlations (not shown; Pearson's correlation coefficient, r , ranging from 0.65 to 0.92). In particular, BT and EKE are significantly positively correlated in regions where BT is mainly positive (directly upstream and 20 km downstream of the western array, and across a broad region surrounding the eastern array (Fig. 7b). Near the western array, the predominantly positive BC also shows a positive but weaker correlation with EKE, (r from 0.51 to 0.67).

The cross-isobath gradient of the effective potential vorticity, q_y , suggests some regions along the boundary current may experience be susceptible to barotropic instability. At To diagnose this, we evaluate at each along-isobath point, the product of the minimum and maximum values of q_y across the 1400 km isobath, $f(q_y)$, which visualises whether q_y changes sign (Fig. 7d) and is also discussed further in Section 4.3. Some zones were primarily. Several zones show near-zero or negative, e.g., a narrow zone values of $f(q_y)$, implying sign changes in q_y . These include a narrow region 10 km upstream and a wider zone 40 of the western array and a broader zone 10 km downstream of the western moorings, indicating that, suggesting that the necessary condition for barotropic instability may occur locally with the current conditions in those two years. Other zones had both signs, be locally satisfied during the two-year period. In other zones, $f(q_y)$ alternates in sign, indicating that topographic β was low small enough to be outweighed by the curvature of the current at the 1400 m isobath during in some months. Some

~~positive BT was found even~~ Positive BT is also found in regions where the necessary condition for barotropic instability ~~was~~ is not met, indicating that eddy–mean flow interaction may occur in regions that are not locally unstable.

3.3.3 ~~Variability at the virtual moorings~~

The divergence of eddy temperature flux integrated along the 1400 m isobath (Fig. 7e) has pronounced temporal variability, ranging from $-1.8^{\circ}\text{C cm s}^{-1}$ to $2.3^{\circ}\text{C cm s}^{-1}$, with a mean value of $0.3^{\circ}\text{C cm s}^{-1}$. The five months with the largest integrated eddy temperature flux divergence occurred between November and February. Averaged in time, the cumulative integral of eddy temperature flux divergence increases starting from near the western mooring array to until 10 km downstream of the eastern array (Fig. 7e, upper panel, left axis). The largest contributions to the cumulative integral are from a broad zone extending downstream from the western mooring array (35 km to 70 km along the isobath) and a narrower zone centred on the eastern array (125 km to 140 km; Fig. 7e, upper sub-panel, right axis). The broad zone starting from the western array coincides with areas where both BT and BC are positive on average (Fig. 7b, c), while the more localized smaller maximum near the eastern array is approximately co-located with the local BT maximum there.

3.3.3 ~~Variability at the virtual moorings compared to in situ moorings and along-isobath estimates~~

The virtual moorings ~~reproduced, although slightly overestimated, the seasonal~~ tended to overestimate the magnitude and variability of EKE ~~seen both in the observations and in the~~ relative to the spatially averaged model estimates ~~at each mooring site along the isobath~~ along the 1400 m isobath sampled at the mooring sites (Fig. 8a, b). EKE was for example on average 1.8 higher, and at most 4 times higher, at the virtual mooring than in the spatially averaged along-isobath estimates. In contrast, EKE estimates from the virtual moorings match well those obtained from cross-slope segments spanning the vertical and lateral extent of the moorings, as well as from longer segments covering most of the water column (not shown), indicating that point-based sampling can capture the energetics of a broader cross-slope region. EKE in the model was generally larger than the observed EKE, especially during summer when observed values were reduced to approximately half of the autumn- and winter values.

BT estimated from the virtual moorings was generally larger ~~than that from the in situ moorings in~~ magnitude than both the along-isobath estimates and, in particular, the in situ mooring estimates (Fig. 8b, e–c, d). BC estimates from the virtual moorings showed variability ~~similar to those from in situ moorings and depended similarly on~~ comparable to the in situ observations and showed similarly sensitivity to sampling depth and the ~~specific mooring pair used~~ choice of mooring pair (Fig. 8e, f). ~~We note that although the virtual and in situ moorings were placed at approximately the same geographical and slope locations, they still might not be equivalently placed~~ As with BT, virtual-mooring BC estimates diverged from the spatially averaged values. While BT from the virtual-moorings broadly agreed with segment-based estimates, BC showed larger disagreement (not shown).

Differences between the model-based and observational estimates could be attributed to the different time periods covered, the representativeness of mooring locations relative to the boundary current, ~~and note also that the modelled and observed data sets cover different time periods.~~

The virtual moorings also agreed fairly well with the simplified estimates of BT made on cross-slope segments (not shown).
 385 However, isopycnal slopes and shear, and potential shortcomings of the model in representing the meso- and submesoscale
 processes. EKE and conversion rate estimates using the full BT equation equations for the conversion rates and averaging over
 the full depth and across the region, the resulting spatial averages differed substantially from the simplified virtual mooring and
 segment-based estimates, highlighting the effects of both the calculation method and the degree of averaging. Segment-based
 averages of BC were generally smaller than point-wise virtual mooring estimates (not shown), showing that spatial averaging
 390 tends to reduce variability. As with BT, spatially averaged BC values also diverged from the virtual mooring results. importance
 of adequate volume averaging.

4 Discussion

4.1 Enhanced EKE and conversion in autumn and winter rates

We observed higher EKE during autumn and winter from the mooring observations (Fig. 4a–b). This agrees with Crews et al.
 395 (2018) who found higher EKE between October and March in a high-resolution model and with Koenig et al. (2022b) who
 found increased geostrophic surface EKE in the autumn and winter months. Upstream at the 78° 50' N mooring array, the WSC
 is most energetic from December to May, with a peak in February (von Appen et al., 2016). In AW inflow regions, such as the
 Barents Sea Opening, Fram Strait, and the western Nansen Basin, wintertime EKE can be relatively higher. This is attributed
 to the typically lower sea ice cover in these areas, which permits stronger wind forcing (von Appen et al., 2022). In addition,
 400 the lower sea ice cover in the inflow zones and supports weaker stratification and higher instability (von Appen et al., 2022).

During the same periods of elevated EKE —in autumn and winter—, we also observed larger energy conversion, with BC as
 the dominant contribution (Figs. 4g–h, 5e–f). The important role of BC in autumn and winter seen in the mooring observations
 is supported by the model, lending confidence on to the simplified calculations from the moorings. Previous studies from up-
 stream regions in Fram Strait and west of Spitsbergen have similarly shown that the boundary current is baroclinically unstable
 405 (Teigen et al., 2011; von Appen et al., 2016; Wekerle et al., 2020) (Teigen et al., 2011; von Appen et al., 2016; Wekerle et al., 2020; Fer et
 . The high wintertime EKE in the WSC has been linked to baroclinic instability due to weak stratification and a stronger vertical
 shear (von Appen et al., 2016). Similarly, in the Eurasian Basin, Müller et al. (2024) reported a spatial agreement between EKE
 and BC, suggesting that baroclinic instability supplies eddy energy locally. Our finding that both EKE and BC are greater in
 autumn and winter indicates that baroclinic instability may similarly drive higher EKE north of Svalbard as well.

410 While BC was stronger in the mooring observations, the volume-averaged conversion rates in the model show similar
 contributions of BT and BC (Fig. 6c, d). Averaged along the boundary current path, BT was comparable to and larger than
 BC for most of the time (Fig. 7b, c). Considering the uncertainties and limitations of point-based estimates of conversion
 rates, highlighted by the limited ability of the virtual moorings to reproduce the volume-averaged conversion rates along
 the boundary current pathway (Section 3.3.3 and Fig. 8), we may assume there can be a substantial contribution to energy
 415 conversion from barotropic instability. Upstream of our mooring site, the boundary current west of Spitsbergen has been

shown to be barotropically unstable (Teigen et al., 2010), and the results of Wekerle et al. (2020) also indicate that barotropic conversion is non-negligible.

4.2 ~~Comparison to other regions~~

Conversion rates have been estimated in several other studies using mooring observations and models. Across the pathway of
420 the WSC in Fram Strait, conversion rates inferred from moored observations were highest near the shelf break during winter,
with a pronounced seasonal cycle, peaking in February and reaching its minimum in August (von Appen et al., 2016). Their
largest ~~BC value, 15×10^{-5}~~ monthly mean BT, 2.5×10^{-5} W m^{-3} at 75 m depth in February, is about ten times higher than
our ~~November maximum of $4 \times 10^{-5} \text{W m}^{-3}$ at 320 m~~ layer-averaged maximum during January 2019 at the western array. At
the same location, their ~~BT in February reached 2.5×10^{-5}~~ BC value reaches 15×10^{-5} W m^{-3} , which ~~is about ten times~~
425 higher than our layer-averaged maximum during January 2019 at the western array. However, in our data exceeds our largest
monthly mean, $2.9 \times 10^{-5} \text{W m}^{-3}$ at 320 m in November. In our data, both BT and BC were larger at shallower depths
between roughly 300 m and 700 m, so it is plausible that values at 75 m could be closer to those of von Appen et al. (2016).
It should be noted that we used slightly different passbands than von Appen et al. (2016) to obtain fluctuations. A comparison
between choices of passbands and filters is shown in Appendix C. If we use their passband of 48 hours to 30 days instead
430 of our 35 hours to 14 days, our November-mean BC increases by 25 %. Conversion rates in Fram Strait estimated using two
high-resolution models (Wekerle et al., 2020), reach magnitudes of $5 \times 10^{-5} \text{m}^3 \text{s}^{-3}$ along the slope. Depth-integrated values
in $\text{m}^3 \text{s}^{-3}$ are numerically equivalent to depth-averaged values with units W m^{-3} for a ~ 1000 m-thick layer of seawater with
density $\sim 1000 \text{kg m}^{-3}$.

~~Further downstream of WSC in Fram Strait, Fer et al. (2023) estimated conversion rates from mooring records on the On~~
435 ~~the southern flank of the Yermak Plateau at 80°N , further downstream of the mooring observations of von Appen et al. (2016)~~
~~, Fer et al. (2023) estimated conversion rates from mooring records.~~ Similar to our results, they observed that both BT and BC
were highly variable in depth and time, ~~with periods of positive and negative values. The conversion rates showed no~~ with no
consistent or strong correlation with EKE, ~~consistent with our findings from both mooring and model data.~~ Their observed
EKE levels exceeded what would be expected from the local conversion rates, and could be partly accounted for advection
440 from an energetic region upstream of the moorings reported in von Appen et al. (2016).

~~Conversion rates in Fram Strait have also been estimated using two high-resolution models (Wekerle et al., 2020), resulting~~
~~in values along the slope within $\pm 5 \times 10^{-5} \text{m}^3 \text{s}^{-3}$ (equivalent to $\pm 5 \times 10^{-5} \text{W m}^{-3}$ at 1000 m depth). BT had alternating~~
~~sign along the slope, indicating a strong influence of bathymetry, consistent with our findings, a point we return to in Section 4.3.~~

445 Negative conversion rates, observed in both our mooring records (lasting weeks) and ~~in~~ the model (lasting months in some
regions), have also been reported elsewhere in other regions (e.g., Spall et al., 2008; Håvik et al., 2017; Wekerle et al., 2020;
Fer et al., 2023). Theoretically, negative BT implies eddy energy feeding a transfer of energy from eddies to the mean flow,
while negative BC implies energy transfer that steepens isopycnals by lifting denser water onto toward the shelf. ~~In our mooring~~
~~data~~ Our observations provide some support for these interpretations. For example, a period of negative BC in January–February

2019 was followed by an increase in isopycnal slope (Fig. 5c, a). In the model, a wide zone between 80 km and 110 km along the isobath with predominantly negative BT (Fig. 7b), shows a significant negative correlation (r -0.47 to -0.72) with the along-slope current velocity (not shown). Here, decreasing BT is associated with increasing current strength, consistent with a transfer of energy from eddies and to the mean flow. Correspondingly, the mean velocity transitions from below- to above-average values downstream (Fig. 7a, right axis of upper panel), suggesting a local strengthening of the boundary current. However, this relation between negative BT and an increase in $\bar{u}$ is not found in all regions. Due to uncertainties in the estimates from the moorings, we cannot determine whether ~~this reflects a physical process or the observed isopycnal slope variability coincidentally aligned with the estimated negative BC~~ the observed change in the isopycnal slope is a direct consequence of negative BC or a coincidental alignment. However, the occurrence of negative conversion rates ~~seen in both our measurements and in both observations and~~ high-resolution ocean model studies support models supports the interpretation that ~~such processes may occur in the ocean, although further investigation is required to confirm this~~ energy can be transferred to the mean flow or into available potential energy.

4.2 Mesoscale activity and heat loss from the boundary current

Understanding ~~the heat loss of AW as it flows in from~~ AW along the boundary current is key to regional climate and oceanography north of Svalbard, with implications for the broader Arctic Ocean. Koenig et al. (2022b) estimated the seasonal evolution of along-stream heat loss of AW between the western and the eastern mooring arrays and showed that air-sea heat fluxes ~~explain much of the AW cooling between 18°E and 24°E during autumn, but are insufficient to alone cannot~~ account for the cooling rates observed ~~observed cooling rates~~ in winter, spring, and summer. This implies that additional mechanisms contribute to the along-stream heat loss of AW, such as lateral exchange proposed by Crews et al. (2018) and Kolås et al. (2020).

Several observations from the slope north of Svalbard indicate the presence of such lateral processes: ~~temperature.~~ Temperature variability exhibited an onshore-offshore mode with a 6–7 day periodicity that was correlated with across-slope velocity variability, and surface geostrophic EKE was elevated near the current core (Koenig et al., 2022b). Anticyclonic eddies are a likely agent for this exchange. Limited observations north of Svalbard showed anticyclones carrying warm anomalies offshore from the boundary current (Våge et al., 2016) and other eddy features linked to baroclinic instability (Pérez-Hernández et al., 2017); ~~which could contribute to lateral heat loss from the current.~~

Anticyclonic eddies carrying AW or warm anomalies have also been identified in high-resolution modelling studies both north of Svalbard (Crews et al., 2018) and upstream (Wekerle et al., 2020). ~~Model results reported~~ The model results presented here also indicate substantial conversion rates ~~, not only in the upper slope where moorings were placed but also farther both near the moorings sites and further~~ offshore (Fig. 6). ~~This offshore activity agrees with earlier observations of eddies over the deeper slope (Våge et al., 2016; Pérez-Hernández et al., 2017), as well as frequent offshore eddy propagation detected in a high-resolution model (Crews et al., 2018). Furthermore, from their mooring at the 800 m isobath, Renner et al. (2018) detected few eddies, suggesting that most warm-core eddies detach further offshore. Our observations and model results, together with previously published evidence, indicate mesoscale activity consistent with eddy-driven exchange contributing to AW heat loss from the boundary current.~~

Using the mooring arrays analysed here, Koenig et al. (2022b) reported that the

485 Seasonal contrasts further point to a role for mesoscale processes in the heat budget. In autumn, along-stream heat loss
of AW in autumn could be accounted for by the vertical heat loss to the atmosphere. During this period, can largely be
explained by direct atmospheric cooling, when the AW core is warm, shallow, and the study area was ice-free, and the AW
core was at its warmest and closest to the surface region is ice-free (Koenig et al., 2022b). In winter and spring, however, when
along-stream heat loss increases substantially while the AW core subducted and sea ice cover limited direct atmospheric
490 cooling, 75 W m^{-2} out of 302 W m^{-2} along-stream cooling in winter, and 35 W m^{-2} out of 60 W m^{-2} in spring, was
unaccounted for by heat loss subducts and becomes less exposed to the atmosphere. Our analysis of the measured EKE and
conversion rates suggests that the contribution of mesoscale activity can explain much of this missing. During this period,
EKE remains elevated, suggesting that lateral exchange associated with mesoscale activity becomes more important. In spring,
surface heat loss is strongly suppressed by sea ice, while mesoscale activity persists at reduced levels, again indicating a
495 contribution from eddy-driven lateral exchange. Consistent with this interpretation, eddy temperature flux divergence estimated
from the moorings (Section 3.1) indicates stronger lateral heat loss from the boundary current in spring and winter (autumn and
winter. Although these local estimates are uncertain, their seasonal structure is consistent with the periods when atmospheric
heat fluxes cannot account for the observed cooling. Model-based estimates of integrated eddy temperature flux divergence
along the boundary current show a similar seasonal signal, with consistently positive divergence during winter, and the largest
500 values occurring between November and February (Section 3.3.2, Fig. 4a, b7e).

During summer, however, when the surface gained heat from the atmosphere, Koenig et al. (2022b) estimated average
along-stream cooling of AW comparable to winter, suggesting that all the heat loss in summer must be lateral or downward.
EKE in this period was, however, at its lowest. In contrast, summer is characterized by weak mesoscale activity, small EKE
(Figs. 3c, 4a–b), and the calculated conversion rates were negligible, 7a), negligible observed (Figs. 4g–h, 5e–f). This implies
505 that mesoscale eddies were and below-average modelled (Fig. 7b, c) energy conversion rates, and small eddy temperature
fluxes and their divergence. Mesoscale eddies are therefore unlikely to be the dominant agents for heat loss in summer, and
other processes must play a role. However, Koenig et al. (2022b) estimated average along-stream cooling of AW in summer
comparable to winter, suggesting that all the heat loss in summer must be lateral or downward by turbulence at the base of the
AW layer. A likely candidate for further summer lateral heat loss is topographic vorticity-Rossby waves, which have previ-
510 ously been linked to increased heat loss from the WSC (Nilsen et al., 2006). An ongoing study (unpublished F. Nilsen, personal
communication) indicates that the topographic slope shape of the continental slope near the western mooring array, together
with the lower-reduced stratification and slower current observed in summer, could allow for vorticity support topographic
Rossby waves with diurnal periods, which are energized by the diurnal tide in the area near the western mooring array.

Our observations and model results, together with previously published evidence, indicate mesoscale activity and eddy-driven
515 exchange play an important role in the heat loss of AW from the boundary current.

4.3 Mechanisms behind the energy conversion

The pattern of alternating signs ~~of in the~~ energy conversion rates along the continental slope (Fig. 6bc) is consistent with findings from other regions such as Fram Strait (Wekerle et al., 2020) and the Lofoten escarpment (Fer, 2020). North of Svalbard, these ~~time-persistent-alternating~~ patterns were connected to ~~the-veering-of-the-veering~~ isobaths (Section ??3.3.1), indicating ~~the-a~~ close connection to bathymetry, ~~as also~~ noted by Wekerle et al. (2020). This connection is likely a manifestation of the necessary condition for barotropic instability, which depends on ~~the-variation-spatial variations~~ in bathymetry through ~~β_{topo}~~ the topographic β (defined in Section 2.3.4).

Our ~~quantification-of-metric for~~ the lateral sign change of the PV gradient (Fig. 6d) ~~indicated some-7d~~ identifies several areas along the 1400 m isobath where the necessary condition for barotropic instability could be satisfied. However, the spatial and temporal patterns of BT are not directly ~~explained by these areas aligned with these regions~~ (Fig. 6b)-7b). For example, BT was positive in several regions ~~—for example-such as~~ upstream of the eastern array—, where the condition for instability was ~~never met, and conversely, BT was low or negative-not met, while~~ in some areas where the condition was fulfilled, ~~BT was weak or negative~~. The latter ~~can happen since meeting is expected since~~ the necessary condition does not ensure instability will occur. Overall, the connection between the potential for barotropic instability and the observed barotropic conversion ~~was~~ is inconclusive. This can partly be explained by ~~the mean-current-adveeting-advection:~~ instabilities or eddies ~~from-upstream, where they were triggered in areas satisfying-generated upstream, in regions where~~ the instability condition ~~is met, may be transported into downstream areas with different local conditions~~. Similar observations were made by e.g., Håvik et al. (2017).

~~For baroclinic instability to occur, Baroclinic instability requires that~~ the cross-stream gradient of the effective potential vorticity ~~must change-changes~~ sign with depth (Charney and Stern, 1962; Spall and Pedlosky, 2008). With only two moorings at each array, we cannot reliably ~~determine-whether-this-condition-is-met~~assess this condition. However, small values of the geostrophic Richardson number, $Ri = N^2/S^2$, may indicate conditions favourable for baroclinic instability (von Appen et al., 2016; Crews et al., 2018), as ~~stratification-is-expected-to-suppressand-shear-to-favour~~strong stratification tends to suppress, ~~while strong shear promotes~~ instability. We use the background fields of stratification, $N^2 = -(g/\rho_0) \partial\rho/\partial z$, and shear, $S^2 = (\partial\bar{u}/\partial z)^2$, instead of the thermal-wind shear, which are representative of the geostrophic scales. Overall, Ri was significantly smaller during winter than in summer, consistent with ~~the~~ enhanced EKE and ~~dominant-stronger~~ BC in winter observed in the mooring records. Median ~~value-of-Ri-was-values of Ri in summer were~~ 2 to 20 times larger than in winter, depending on the mooring and the measurement level analysed. During winter, Ri was less than 50 ~~,21%-21%~~ of the time at W2 (~~30%-30%~~ at W3), ~~but only 5%-of-the-time-compared to only 5%~~ during summer. Strong vertical shear, connected to the horizontal density gradient through the thermal wind relation, was the main factor reducing Ri.

The mean flow advecting eddies can also explain differences between measured energy conversion and observed EKE. Eddies generated in regions of high instability can be transported downstream by the mean circulation – for instance, eddies observed in the eastern Eurasian Basin have been traced as far back upstream as the Yermak Plateau, the area north of Svalbard, and the slope between Svalbard and Franz Josef Land (Pnyushkov et al., 2018) – resulting in enhanced EKE away from the generation sites (von Appen et al., 2016; Wekerle et al., 2020; Fer et al., 2023). Furthermore, some EKE can originate from

550 meandering of the current, hence elevated EKE levels do not necessarily indicate local eddy occurrence (Wekerle et al., 2020; Crews et al., 2018).

In summary, several factors complicate the interpretation of instability diagnostics, conversion rate estimates, and local EKE: the necessary condition for barotropic instability identifies regions susceptible to instability, but may still not coincide with regions of high positive barotropic conversion; local conversion rates may not align with local EKE due to advection of
555 eddies from upstream; and EKE levels themselves may not reflect the eddy formation or occurrence.

4.4 Limitations of using mooring observations for conversion rate estimations

As noted by earlier studies (e.g., Fer et al., 2020, 2023), conversion rates based on mooring observations are subject to significant uncertainty and therefore should be interpreted with caution. A primary limitation is that moorings provide fixed-point measurements, which may not capture the full spatial and temporal variability of energy conversion. Model results show that
560 conversion rates are patchy and intermittent, as illustrated by differences between volume-averaged rates and estimates from virtual moorings or cross-slope segments. Sufficient averaging in time and space is therefore essential for reliable conversion rate estimates.

Another limitation is that mooring arrays do not allow estimation of all terms in the conversion rate equations. This limitation is illustrated using the estimates from the simplified formulas at virtual mooring and segments versus the full equations.
565 In addition, a comparison of the four terms in the BT formula (Eq. 2) showed that the two divergent terms were more often positive, while the rotational terms were occasionally negative. In a similar comparison for the two terms in the BC formula (Eq. 3), the signs often opposed. Hence, a negative value in one term estimated from moorings does not preclude a positive total conversion rate, or vice versa. Even under the assumption that point-based estimates from one term adequately represent regional conditions, uncertainties would still arise from methodological and instrumental limitations associated with the mooring data: for example a larger mooring spacing relative to the Rossby radius will underestimate the isopycnal slope and thus
570 BC (von Appen et al., 2016); and similarly, inadequate resolution of the lateral velocity shear will underestimate BT.

Our model results suggest that much of the conversion occurs deeper and further offshore than our moorings (Fig. 6). Nonetheless, a considerable fraction of the conversion occurs over the slope and at the depth sampled by the moorings, as supported by both analyses of virtual moorings (Fig. 8) and by comparing full-depth cross-slope segment estimates with those
575 at limited-depth segments with shorter lateral span. The virtual mooring results further demonstrate that the substantial spread observed with the BC rates at various depths (Fig. 5) is to be expected. As such, mooring-based estimates are better regarded as order-of-magnitude approximations indicating the possible conversion for that time in that region.

5 Summary and Conclusions

Motivated by the potential role of mesoscale eddies in transporting AW and heat offshore from the boundary current, we
580 investigated mesoscale variability and energy conversion rates on the continental slope north of Svalbard. Mooring observations revealed large differences between autumn–winter and spring–summer both in EKE and energy conversion rates (Figs. 3, 4, 5).

The relatively energetic autumn and winter periods coincided with the strongest boundary current and warmest AW (Fig. 2). The largest conversion rates were primarily baroclinic (Figs. 4, 5), indicating baroclinic instability of the boundary current. Our findings from year-long measurements provide observational support to previous limited observations from a cruise (Våge et al., 2016) and inferences from a model (Crews et al., 2018), which suggested baroclinic instability and associated warm-core eddies carrying AW offshore.

To put these findings obtained from limited mooring observations into perspective, and to evaluate whether the conversion calculations are representative for the larger slope area, we used a 500-m horizontal resolution ocean model. The model results revealed that the mooring arrays were placed in energetic zones along the continental slope where considerable energy conversion can occur (Fig. 6). They also revealed that substantial conversion occurs also deeper and offshore of the mooring arrays.

Although mooring data indicated that baroclinic conversion exceeded barotropic conversion, these estimates are subject to uncertainties, for example due to limited sampling in vertical and cross-slope extents as well as the assumption of along-slope homogeneity. Model results provided important context, showing that both barotropic and baroclinic conversion can occur at comparable magnitudes, with substantial variability in time and space, and that considerable energy conversion may take place further offshore than the mooring arrays. These findings highlight the need for broad spatial sampling and adequate averaging to obtain representative estimates of energy conversion on the slope. Furthermore, conversion rates were only weakly connected to local EKE variability in both mooring observations and model data, suggesting that advection of eddies from upstream unstable regions by the mean circulation may be important.

Our inferences from the mesoscale-band variability and the energetics help explain the missing along-stream heat loss from the boundary current in winter and spring that was reported in Koenig et al. (2022b) using the same moorings. Enhanced mesoscale activity in winter, and to a lesser extent in spring, facilitated mesoscale eddy stirring and lateral heat exchange, which become particularly important in periods when the AW has subducted and isolated from the atmosphere by sea ice cover.

Data availability. The Nansen Legacy mooring data (W1–W3, E1–E3, Fer et al., 2022) are available at the Norwegian Marine Data Centre: doi.org/10.21335/NMDC-1852831792. ROMS model output is stored at Institute of Marine Research data servers and can be made available upon request. CARRA (Schyberg et al., 2020) data are available from doi.org/10.24381/cds.713858f6.

Appendix A: EKE and eddy fluxes

In this section, we present EKE, eddy momentum and eddy density fluxes across all moorings and depth levels where sufficient data are available for their calculation. EKE was generally larger during autumn and winter than during spring and summer at all moorings (Fig. A1a,b). Between September and February, EKE at 300 m was typically 2–4 times higher than during the period from March to August. A similar seasonal contrast was observed at shallower depths, including at 100 m on the

upper slope (W1 and E1) and at 50 m on the shelf (W0). After recovery of the moorings W1–W3, W0 recorded a maximum EKE of $65 - 75 \text{ cm}^2 \text{ s}^{-2}$ in November 2019. The largest values, between $75 \text{ cm}^2 \text{ s}^{-2}$ and $145 \text{ cm}^2 \text{ s}^{-2}$, occurred around 200 m depth through November and December 2018 at W2. In general, EKE increased toward the uppermost measurement level at all moorings. This vertical structure was most pronounced at W2, where EKE at 300 m was at least twice that at 600 m for most of the year, and exceeded it by more than a factor of three during autumn 2018. A similar decrease with depth was observed at the other moorings, although the strength of the intensification toward the upper ocean varied. At the deepest moorings, W3 and E3, local EKE maxima occurred at 1000 m depth in November 2018, and EKE at depth remained slightly elevated throughout winter.

Similar to EKE, the eddy momentum flux was also larger in magnitude during autumn and winter (Fig. A1c, d), typically 2–4 times higher between September and February, with the largest values at W2. On the middle and lower slope, the flux was predominantly negative in the upper part of the water column and positive at depth. On the upper slope and shelf, it was generally weaker, except for a pronounced minimum of $-50 \text{ cm}^2 \text{ s}^{-2}$ at W0 in November 2019, coincident with the local EKE maximum. At the eastern array, the momentum flux magnitudes were similar but slightly smaller than at the western array. At E1, the momentum flux generally varied in both sign and magnitude across depth and time, but for about one month it showed a stronger, vertically coherent negative signal spanning the water column, reaching a minimum value of $-25 \text{ cm}^2 \text{ s}^{-2}$ at 100 m, more than half of the magnitude of the winter minimum at E2.

At the western array, the eddy density flux was largest in magnitude at W2 during September–February (Fig. A1e). During autumn 2018, values were negative between 150 m and 300 m, while from December through winter, positive values dominated throughout the water column. Magnitudes at W0 and W3 were lower. At the eastern array, the largest density fluxes were also observed at E2. There, the eddy density flux was largest above 300 m depth, with elevated positive values during spring and summer and maxima in March and August at 150 m depth. Large positive values were also observed at E3 above 300 m from December to July (note, however, this layer was not sampled outside this period). Deeper, flux magnitudes were generally smaller. At E1, the density fluxes were generally small, apart from a one-month maximum in March at 100 m depth, which coincided with, but was notably shorter-lived than, the peak observed at E2.

Appendix B: Model setup validation

B1 Validation against observations on the West Spitsbergen Shelf

The model setup used in this study was originally implemented by Frank et al. (2025b) and validated against observations from Isfjorden and the West Spitsbergen Shelf obtained through the ocean observation program at the University Centre in Svalbard (Skogseth et al., 2020). Hydrographic and current measurements selected along a section across the West Spitsbergen Shelf and the WSC during three autumns were contemporary with the model simulations. At this section, the model had a systematic temperature bias, with colder surface waters and warmer deep water over the slope. Salinity was biased low, both in the surface layer and the AW layer. These compensating biases in temperature and salinity resulted in good agreement in density. The model realistically reproduced the circulation patterns in the region.

Temporal variability in the model simulations was validated against contemporaneous mooring records from the mouth of Isfjorden on the west coast of Svalbard. Simulated temperatures were generally slightly lower than observed but reproduced the temporal variability well. Simulated salinity and temperature were both slightly lower during winter, while agreement was good during the rest of the year. Density time series were well captured by the model. The inflow into Isfjorden was also well represented, with simulated current speeds in good agreement with the observations, although peak velocities were slightly overestimated.

B2 Model–observation comparison north of Svalbard

Because the model simulations were originally designed for the West Spitsbergen Shelf study by Frank et al. (2025b), they do not overlap in time with the mooring observations used here. Nevertheless, we can compare the seasonal mean cross-slope structure of the boundary current and hydrography (Fig. B1).

The model bathymetry is based on IBCAO version 3 (Jakobsson et al., 2012), which is too shallow in the vicinity of the western mooring array. As a result, the steeper part of the slope is positioned farther offshore in the model relative to observations. To account for this, the virtual moorings were shifted offshore to align with comparable depths and slope characteristics as the in situ moorings (Sec. 3.3.3). Because the modelled bathymetry differs from the real bathymetry, direct comparisons of the full cross-slope sections are challenging. However, we can directly compare the hydrographic and current structure at the virtual moorings (Fig. B1) with those from the corresponding deeper mooring pairs (Fig. 2).

The simulated boundary current is located farther offshore and has a different seasonality than observed (Fig. B1). In September–February, the simulated current core, with 18 cm s^{-1} , is centred near the 1100 m isobath and extends from 100 m to 1000 m depth at the western array (Fig. B1a). The observed current core during the same season reaches 20 cm s^{-1} , is located near the 700 m isobath at 100 m depth, and is weakening more rapidly with increasing depth (Fig. 2a). Both the simulated and observed currents in September–February strengthen downstream along-path from the western to the eastern array, reaching 23 cm s^{-1} (Fig. B1b and Fig. 2b). In March–August, the simulated current is stronger, with maximum speeds increasing to 24 cm s^{-1} at the western array and almost 30 cm s^{-1} at the eastern array (Fig. B1c, d). This contrasts with the observations, which show a weakening of the current during this period (Fig. 2c, d).

The offshore displacement of the simulated velocity core relative to observations likely contributes to the lack of a clear seasonal signal in EKE and conversion rates in the model. It may also explain why the largest conversion rates occur offshore of the mooring arrays.

Both the observed and simulated temperature maxima were approximately 0.5°C higher in September–February than in March–August (Fig. B1 and Fig. B1). However, the observed temperature maxima were approximately 0.5°C higher than those in the model in both seasons. Despite being colder, the simulated water masses were also less dense. For example, the 1027.9 kg m^{-3} and 1028.0 kg m^{-3} isopycnals were located at depths of approximately 300–400 m and 600–650 m in the observations, while in the model they were found deeper, near 600–650 m and 1000 m respectively.

The virtual moorings and the 1400 m isobath selected for the analysis are positioned to capture both the simulated temperature core and the boundary current core.

Various studies have used slightly different cutoff frequencies for the band-pass filter in their analyses. To illustrate the sensitivity to passband choice, we compare the original 35 h–14 d passband with a wider 48 h–30 d band used by von Appen et al. (2016) (Fig. C1). We also examine the effect of lowering the high-frequency cutoff toward 24 h, motivated by advective time scales ranging from 24 to 40 h. To approach a 24 h cutoff while avoiding contamination from the diurnal tidal band, we use a finite impulse response Kaiser filter, which provides a sharper transition between passband and stopband. The original 35 h–14 d passband using a Butterworth filter yields similar results to the Kaiser filter, with some minor differences at the longer time scales.

The different filter choices are generally in good agreement, and our results are not strongly sensitive to the filter choices. The smallest difference is between the Butterworth and Kaiser filters when applied with the same passband. Time series with different passbands typically show similar magnitudes and variability, although the 48 h–30 d band occasionally deviates. The results are at times sensitive to the choice of high-frequency cutoff (27 h, 35 h, or 48 h). For example, in November 2018, BC shows the largest spread between passbands, as a result of differences in the fluctuations of v and ρ arising from the different cutoffs.

Author contributions. KK analysed the mooring data and performed calculations on the model output with support from IF and TMB. JA and LF set up and ran the ocean model. IF developed the research idea. IF and TMB provided guidance through the project. KK prepared the original draft with advice and support from IF. All authors discussed the results and finalized the paper.

Competing interests. Ilker Fer is a member of the editorial board of Ocean Science.

Acknowledgements. This work was funded by the Research Council of Norway through the project The Nansen Legacy (RCN # 276730). We thank the captains, officers, and crew on the research vessel *Kronprins Haakon* and our ~~scientific and technical colleagues on the cruises~~ colleagues for good cooperation during the deployment and recovery of the moorings. We are also grateful to ~~two colleagues~~ Eva Falck for reading the manuscript and providing valuable feedback. We thank Andrey Pnyushkov and Wilken-Jon von Appen for reviewing our manuscript. Their helpful and constructive feedback has contributed to a substantial improvement of our paper.

References

- Aagaard, K., Foldvik, A., and Hillman, S. R.: The West Spitsbergen Current: Disposition and Water Mass Transformation, *Journal of Geophysical Research: Oceans*, 92, 3778–3784, <https://doi.org/10.1029/JC092iC04p03778>, 1987.
- 705 Ardyna, M. and Arrigo, K. R.: Phytoplankton Dynamics in a Changing Arctic Ocean, *Nature Climate Change*, 10, 892–903, <https://doi.org/10.1038/s41558-020-0905-y>, 2020.
- Armitage, T. W. K., Manucharyan, G. E., Petty, A. A., Kwok, R., and Thompson, A. F.: Enhanced Eddy Activity in the Beaufort Gyre in Response to Sea Ice Loss, *Nature Communications*, 11, 761, <https://doi.org/10.1038/s41467-020-14449-z>, 2020.
- 710 Asplin, L., Albretsen, J., Johnsen, I. A., and Sandvik, A. D.: The Hydrodynamic Foundation for Salmon Lice Dispersion Modeling along the Norwegian Coast, *Ocean Dynamics*, 70, 1151–1167, <https://doi.org/10.1007/s10236-020-01378-0>, 2020.
- Athanase, M., Provost, C., Pérez-Hernández, M. D., Sennéchal, N., Bertosio, C., Artana, C., Garric, G., and Lellouche, J.-M.: Atlantic Water Modification North of Svalbard in the Mercator Physical System From 2007 to 2020, *Journal of Geophysical Research: Oceans*, 125, e2020JC016463, <https://doi.org/10.1029/2020JC016463>, 2020.
- 715 Beszczynska-Möller, A., Fahrbach, E., Schauer, U., and Hansen, E.: Variability in Atlantic Water Temperature and Transport at the Entrance to the Arctic Ocean, 1997–2010, *ICES Journal of Marine Science*, 69, 852–863, <https://doi.org/10.1093/icesjms/fss056>, 2012.
- Budgell, W. P.: Numerical Simulation of Ice-Ocean Variability in the Barents Sea Region: Towards Dynamical Downscaling, *Ocean Dynamics*, 55, 370–387, <https://doi.org/10.1007/s10236-005-0008-3>, 2005.
- Carmack, E., Polyakov, I., Padman, L., Fer, I., Hunke, E., Hutchings, J., Jackson, J., Kelley, D., Kwok, R., Layton, C., Melling, H., Perovich, D., Persson, O., Ruddick, B., Timmermans, M.-L., Toole, J., Ross, T., Vavrus, S., and Winsor, P.: Toward Quantifying the Increasing Role of Oceanic Heat in Sea Ice Loss in the New Arctic, *Bulletin of the American Meteorological Society*, 96, 2079–2105, <https://doi.org/10.1175/BAMS-D-13-00177.1>, 2015.
- 720 Charney, J. G. and Stern, M. E.: On the Stability of Internal Baroclinic Jets in a Rotating Atmosphere, *Journal of Atmospheric Sciences*, 19, 159–172, [https://doi.org/10.1175/1520-0469\(1962\)019<0159:OTSOIB>2.0.CO;2](https://doi.org/10.1175/1520-0469(1962)019<0159:OTSOIB>2.0.CO;2), 1962.
- 725 Codiga, D. L.: Unified Tidal Analysis and Prediction Using the UTide Matlab Functions, Tech. Rep. Technical Report 2011-01, Graduate School of Oceanography, University of Rhode Island, Narragansett, RI, USA, <https://doi.org/10.13140/RG.2.1.3761.2008>, 2011.
- Crews, L., Sundfjord, A., Albretsen, J., and Hattermann, T.: Mesoscale Eddy Activity and Transport in the Atlantic Water Inflow Region North of Svalbard, *Journal of Geophysical Research: Oceans*, 123, 201–215, <https://doi.org/10.1002/2017JC013198>, 2018.
- Cushman-Roisin, B. and Beckers, J.: Introduction to Geophysical Fluid Dynamics: Physical and Numerical Aspects, *International Geophysics*, Elsevier Science, ISBN 978-0-12-088759-0, 2011.
- 730 Fer, I.: Physical Oceanography Data from the Cruise KB 2018616 with R.V. Kristine Bonnevie., <https://doi.org/10.21335/NMDC-2047975397>, 2020.
- Fer, I., Bosse, A., and Dugstad, J.: Norwegian Atlantic Slope Current along the Lofoten Escarpment, *Ocean Science*, 16, 685–701, <https://doi.org/10.5194/os-16-685-2020>, 2020.
- 735 Fer, I., Koenig, Z., Kolås, E. H., Baumann, T. M., and Kalhagen, K.: Physical Oceanography Data from Moorings North of Svalbard, September 2018 - September 2019, <https://doi.org/10.21335/NMDC-1852831792>, 2022.
- Fer, I., Peterson, A. K., and Nilsen, F.: Atlantic Water Boundary Current Along the Southern Yermak Plateau, Arctic Ocean, *Journal of Geophysical Research: Oceans*, 128, e2023JC019645, <https://doi.org/10.1029/2023JC019645>, 2023.

- Frank, L., Albretsen, J., Skogseth, R., Nilsen, F., and Jonassen, M. O.: Warm-Water Intrusions onto the West Spitsbergen Shelf during Winter, <https://doi.org/10.5194/egusphere-2025-1721>, 2025a.
- Frank, L., Albretsen, J., Skogseth, R., Nilsen, F., and Jonassen, M. O.: Mechanisms of Warm-Water Intrusions onto the West Spitsbergen Shelf during Winter, *Ocean Science*, 21, 2419–2442, <https://doi.org/10.5194/os-21-2419-2025>, 2025b.
- Guo, C., Ilicak, M., Fer, I., Darelus, E., and Bentsen, M.: Baroclinic Instability of the Faroe Bank Channel Overflow, *Journal of Physical Oceanography*, 44, 2698–2717, <https://doi.org/10.1175/JPO-D-14-0080.1>, 2014.
- Håvik, L., Våge, K., Pickart, R. S., Harden, B., von Appen, W.-J., Jónsson, S., and Østerhus, S.: Structure and Variability of the Shelfbreak East Greenland Current North of Denmark Strait, *Journal of Physical Oceanography*, 47, 2631–2646, <https://doi.org/10.1175/JPO-D-17-0062.1>, 2017.
- Isachsen, P. E., Vogt-Vincent, N. S., Johnson, H. L., and Nilsson, J.: Instability and Mesoscale Eddy Fluxes in an Idealized 3-Layer Beaufort Gyre, *Journal of Geophysical Research: Oceans*, 129, e2023JC020 757, <https://doi.org/10.1029/2023JC020757>, 2024.
- Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J. A., Forbes, S., Fridman, B., Hodnesdal, H., Noormets, R., Pedersen, R., Rebesco, M., Schenke, H. W., Zarayskaya, Y., Accettella, D., Armstrong, A., Anderson, R. M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M., Gardner, J. V., Hall, J. K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad, R., Mosher, D., Nghiem, S. V., Pedrosa, M. T., Travaglini, P. G., and Weatherall, P.: The International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL052219>, 2012.
- Jakobsson, M., Mayer, L. A., Bringensparr, C., Castro, C. F., Mohammad, R., Johnson, P., Ketter, T., Accettella, D., Amblas, D., An, L., Arndt, J. E., Canals, M., Casamor, J. L., Chauché, N., Coakley, B., Danielson, S., Demarte, M., Dickson, M.-L., Dorschel, B., Dowdeswell, J. A., Dreutter, S., Fremand, A. C., Gallant, D., Hall, J. K., Hehemann, L., Hodnesdal, H., Hong, J., Ivaldi, R., Kane, E., Klauke, I., Krawczyk, D. W., Kristoffersen, Y., Kuipers, B. R., Millan, R., Masetti, G., Morlighem, M., Noormets, R., Prescott, M. M., Rebesco, M., Rignot, E., Semiletov, I., Tate, A. J., Travaglini, P., Velicogna, I., Weatherall, P., Weinrebe, W., Willis, J. K., Wood, M., Zarayskaya, Y., Zhang, T., Zimmermann, M., and Zinglersen, K. B.: The International Bathymetric Chart of the Arctic Ocean Version 4.0, *Scientific Data*, 7, 176, <https://doi.org/10.1038/s41597-020-0520-9>, 2020.
- Killworth, P. D.: Barotropic and Baroclinic Instability in Rotating Stratified Fluids, *Dynamics of Atmospheres and Oceans*, 4, 143–184, [https://doi.org/10.1016/0377-0265\(80\)90013-5](https://doi.org/10.1016/0377-0265(80)90013-5), 1980.
- Koenig, Z., Kalhagen, K., and Fer, I.: Ocean Current, Temperature and Salinity Measurements from Moorings North of Svalbard: September 2018 - November 2019, Tech. rep., University of Bergen, 2022a.
- Koenig, Z., Kalhagen, K., Kolås, E., Fer, I., Nilsen, F., and Cottier, F.: Atlantic Water Properties, Transport and Heat Loss From Mooring Observations North of Svalbard, *Journal of Geophysical Research: Oceans*, 127, e2022JC018 568, <https://doi.org/10.1029/2022JC018568>, 2022b.
- Kolås, E. H., Koenig, Z., Fer, I., Nilsen, F., and Marnela, M.: Structure and Transport of Atlantic Water North of Svalbard From Observations in Summer and Fall 2018, *Journal of Geophysical Research: Oceans*, 125, e2020JC016 174, <https://doi.org/10.1029/2020JC016174>, 2020.
- Li, X., Wang, Q., Danilov, S., Koldunov, N., Liu, C., Müller, V., Sidorenko, D., and Jung, T.: Eddy Activity in the Arctic Ocean Projected to Surge in a Warming World, *Nature Climate Change*, 14, 156–162, <https://doi.org/10.1038/s41558-023-01908-w>, 2024.
- Lilly, J. M. and Olhede, S. C.: Higher-Order Properties of Analytic Wavelets, *IEEE Transactions on Signal Processing*, 57, 146–160, <https://doi.org/10.1109/tsp.2008.2007607>, 2009.
- Lilly, J. M. and Olhede, S. C.: On the Analytic Wavelet Transform, *IEEE Transactions on Information Theory*, 56, 4135–4156, <https://doi.org/10.1109/TIT.2010.2050935>, 2010.

- Lilly, J. M. and Olhede, S. C.: Generalized Morse Wavelets as a Superfamily of Analytic Wavelets, *IEEE Transactions on Signal Processing*, 60, 6036–6041, <https://doi.org/10.1109/TSP.2012.2210890>, 2012.
- Manucharyan, G. E. and Thompson, A. F.: Submesoscale Sea Ice-Ocean Interactions in Marginal Ice Zones, *Journal of Geophysical Research: Oceans*, 122, 9455–9475, <https://doi.org/10.1002/2017JC012895>, 2017.
- Manucharyan, G. E. and Thompson, A. F.: Heavy Footprints of Upper-Ocean Eddies on Weakened Arctic Sea Ice in Marginal Ice Zones, *Nature Communications*, 13, 2147, <https://doi.org/10.1038/s41467-022-29663-0>, 2022.
- Marshall, J. and Shutts, G.: A Note on Rotational and Divergent Eddy Fluxes, *Journal of Physical Oceanography*, 11, 1677–1680, [https://doi.org/10.1175/1520-0485\(1981\)011<1677:ANORAD>2.0.CO;2](https://doi.org/10.1175/1520-0485(1981)011<1677:ANORAD>2.0.CO;2), 1981.
- Meneghello, G., Marshall, J., Lique, C., Isachsen, P. E., Doddridge, E., Campin, J.-M., Regan, H., and Talandier, C.: Genesis and Decay of Mesoscale Baroclinic Eddies in the Seasonally Ice-Covered Interior Arctic Ocean, *Journal of Physical Oceanography*, 51, 115–129, <https://doi.org/10.1175/JPO-D-20-0054.1>, 2020.
- Müller, M., Batrak, Y., Kristiansen, J., Køltzow, M. A. Ø., Noer, G., and Korosov, A.: Characteristics of a Convective-Scale Weather Forecasting System for the European Arctic, *Monthly Weather Review*, 145, 4771–4787, <https://doi.org/10.1175/MWR-D-17-0194.1>, 2017.
- Müller, V., Wang, Q., Koldunov, N., Danilov, S., Sidorenko, D., and Jung, T.: Variability of Eddy Kinetic Energy in the Eurasian Basin of the Arctic Ocean Inferred From a Model Simulation at 1-Km Resolution, *Journal of Geophysical Research: Oceans*, 129, e2023JC020139, <https://doi.org/10.1029/2023JC020139>, 2024.
- Nilsen, F., Gjevik, B., and Schauer, U.: Cooling of the West Spitsbergen Current: Isopycnal Diffusion by Topographic Vorticity Waves, *Journal of Geophysical Research: Oceans*, 111, <https://doi.org/10.1029/2005JC002991>, 2006.
- Pérez-Hernández, M. D., Pickart, R. S., Pavlov, V., Våge, K., Ingvaldsen, R., Sundfjord, A., Renner, A. H. H., Torres, D. J., and Erofeeva, S. Y.: The Atlantic Water Boundary Current North of Svalbard in Late Summer, *Journal of Geophysical Research: Oceans*, 122, 2269–2290, <https://doi.org/10.1002/2016JC012486>, 2017.
- Pérez-Hernández, M. D., Pickart, R. S., Torres, D. J., Bahr, F., Sundfjord, A., Ingvaldsen, R., Renner, A. H. H., Beszczynska-Möller, A., von Appen, W.-J., and Pavlov, V.: Structure, Transport, and Seasonality of the Atlantic Water Boundary Current North of Svalbard: Results From a Yearlong Mooring Array, *Journal of Geophysical Research: Oceans*, 124, 1679–1698, <https://doi.org/10.1029/2018JC014759>, 2019.
- Pnyushkov, A., Polyakov, I. V., Padman, L., and Nguyen, A. T.: Structure and Dynamics of Mesoscale Eddies over the Laptev Sea Continental Slope in the Arctic Ocean, *Ocean Science*, 14, 1329–1347, <https://doi.org/10.5194/os-14-1329-2018>, 2018.
- Polyakov, I. V., Pnyushkov, A. V., Alkire, M. B., Ashik, I. M., Baumann, T. M., Carmack, E. C., Goszczko, I., Guthrie, J., Ivanov, V. V., Kanzow, T., Krishfield, R., Kwok, R., Sundfjord, A., Morison, J., Rember, R., and Yulin, A.: Greater Role for Atlantic Inflows on Sea-Ice Loss in the Eurasian Basin of the Arctic Ocean, *Science*, 356, 285–291, <https://doi.org/10.1126/science.aai8204>, 2017.
- Polyakov, I. V., Alkire, M. B., Bluhm, B. A., Brown, K. A., Carmack, E. C., Chierici, M., Danielson, S. L., Ellingsen, I., Ershova, E. A., Gårdfeldt, K., Ingvaldsen, R. B., Pnyushkov, A. V., Slagstad, D., and Wassmann, P.: Borealization of the Arctic Ocean in Response to Anomalous Advection From Sub-Arctic Seas, *Frontiers in Marine Science*, 7, <https://doi.org/10.3389/fmars.2020.00491>, 2020a.
- Polyakov, I. V., Rippeth, T. P., Fer, I., Alkire, M. B., Baumann, T. M., Carmack, E. C., Ingvaldsen, R., Ivanov, V. V., Janout, M., Lind, S., Padman, L., Pnyushkov, A. V., and Rember, R.: Weakening of Cold Halocline Layer Exposes Sea Ice to Oceanic Heat in the Eastern Arctic Ocean, *Journal of Climate*, 33, 8107–8123, <https://doi.org/10.1175/JCLI-D-19-0976.1>, 2020b.

- Renner, A. H. H., Sundfjord, A., Janout, M. A., Ingvaldsen, R. B., Beszczynska-Möller, A., Pickart, R. S., and Pérez-Hernández, M. D.: Variability and Redistribution of Heat in the Atlantic Water Boundary Current North of Svalbard, *Journal of Geophysical Research: Oceans*, 123, 6373–6391, <https://doi.org/10.1029/2018JC013814>, 2018.
- Richards, A. E., Johnson, H. L., and Lique, C.: Spatial and Temporal Variability of Atlantic Water in the Arctic From 40 Years of Observations, *Journal of Geophysical Research: Oceans*, 127, e2021JC018358, <https://doi.org/10.1029/2021JC018358>, 2022.
- Röhrs, J., Gusdal, Y., Rikardsen, E. S. U., Durán Moro, M., Brændshøi, J., Kristensen, N. M., Fritzner, S., Wang, K., Sperrevik, A. K., Idžanović, M., Lavergne, T., Debernard, J. B., and Christensen, K. H.: Barents-2.5km v2.0: An Operational Data-Assimilative Coupled Ocean and Sea Ice Ensemble Prediction Model for the Barents Sea and Svalbard, *Geoscientific Model Development*, 16, 5401–5426, <https://doi.org/10.5194/gmd-16-5401-2023>, 2023.
- Rudels, B., J. Friedrich, H., and Quadfasel, D.: The Arctic Circumpolar Boundary Current, *Deep Sea Research Part II: Topical Studies in Oceanography*, 46, 1023–1062, [https://doi.org/10.1016/S0967-0645\(99\)00015-6](https://doi.org/10.1016/S0967-0645(99)00015-6), 1999.
- Schauer, U., Muench, R. D., Rudels, B., and Timokhov, L.: Impact of Eastern Arctic Shelf Waters on the Nansen Basin Intermediate Layers, *Journal of Geophysical Research: Oceans*, 102, 3371–3382, <https://doi.org/10.1029/96JC03366>, 1997.
- Schyberg, H., Yang, X., Køltzow, M. A. Ø., Amstrup, B., Bakketun, Å., Bazile, E., Bojarova, J., Box, J. E., Dahlgren, P., Hagelin, S., Homleid, M., Horányi, A., Høyer, J., Johansson, Å., Killie, M. A., Körnich, H., LeMoigne, P., Lindskog, M., Manninen, T., Nielsen-Englyst, P., Nielsen, K. P., Olsson, E., Palmason, B., PeraltaAros, C., Randriamampianina, R., Samuelsson, P., Stappers, R., Støylen, E., Thorsteinsson, S., Valkonen, T., and Wang, Z. Q.: Arctic Regional Reanalysis on Single Levels from 1991 to Present, <https://doi.org/10.24381/cds.713858f6>, 2020.
- Shchepetkin, A. F. and McWilliams, J. C.: The Regional Oceanic Modeling System (ROMS): A Split-Explicit, Free-Surface, Topography-Following-Coordinate Oceanic Model, *Ocean Modelling*, 9, 347–404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- Skogseth, R., Olivier, L. L. A., Nilsen, F., Falck, E., Fraser, N., Tverberg, V., Ledang, A. B., Vader, A., Jonassen, M. O., Søreide, J., Cottier, F., Berge, J., Ivanov, B. V., and Falk-Petersen, S.: Variability and Decadal Trends in the Isfjorden (Svalbard) Ocean Climate and Circulation – An Indicator for Climate Change in the European Arctic, *Progress in Oceanography*, 187, 102394, <https://doi.org/10.1016/j.pocean.2020.102394>, 2020.
- Spall, M. A. and Pedlosky, J.: Lateral Coupling in Baroclinically Unstable Flows, *Journal of Physical Oceanography*, 38, 1267–1277, <https://doi.org/10.1175/2007JPO3906.1>, 2008.
- Spall, M. A., Pickart, R. S., Fratantoni, P. S., and Plueddemann, A. J.: Western Arctic Shelfbreak Eddies: Formation and Transport, *Journal of Physical Oceanography*, 38, 1644–1668, <https://doi.org/10.1175/2007JPO3829.1>, 2008.
- Teigen, S. H., Nilsen, F., and Gjevik, B.: Barotropic Instability in the West Spitsbergen Current, *Journal of Geophysical Research: Oceans*, 115, <https://doi.org/10.1029/2009JC005996>, 2010.
- Teigen, S. H., Nilsen, F., Skogseth, R., Gjevik, B., and Beszczynska-Möller, A.: Baroclinic Instability in the West Spitsbergen Current, *Journal of Geophysical Research: Oceans*, 116, <https://doi.org/10.1029/2011JC006974>, 2011.
- Våge, K., Pickart, R. S., Pavlov, V., Lin, P., Torres, D. J., Ingvaldsen, R., Sundfjord, A., and Proshutinsky, A.: The Atlantic Water Boundary Current in the Nansen Basin: Transport and Mechanisms of Lateral Exchange, *Journal of Geophysical Research: Oceans*, 121, 6946–6960, <https://doi.org/10.1002/2016JC011715>, 2016.
- Vallis, G. K. and Maltrud, M. E.: Generation of Mean Flows and Jets on a Beta Plane and over Topography, *Journal of Physical Oceanography*, 23, 1346–1362, 1993.

- 850 von Appen, W.-J., Schauer, U., Hattermann, T., and Beszczynska-Möller, A.: Seasonal Cycle of Mesoscale Instability of the West Spitsbergen Current, *Journal of Physical Oceanography*, 46, <https://doi.org/10.1175/JPO-D-15-0184.1>, 2016.
- von Appen, W.-J., Baumann, T., Janout, M., Koldunov, N., Lenn, Y.-D., Pickart, R., Scott, R., and Wang, Q.: Eddies and the Distribution of Eddy Kinetic Energy in the Arctic Ocean, *Oceanography*, <https://doi.org/10.5670/oceanog.2022.122>, 2022.
- Wang, Q., Koldunov, N. V., Danilov, S., Sidorenko, D., Wekerle, C., Scholz, P., Bashmachnikov, I. L., and Jung, T.: Eddy Kinetic
855 Energy in the Arctic Ocean From a Global Simulation With a 1-Km Arctic, *Geophysical Research Letters*, 47, e2020GL088550, <https://doi.org/10.1029/2020GL088550>, 2020.
- Wekerle, C., Hattermann, T., Wang, Q., Crews, L., von Appen, W.-J., and Danilov, S.: Properties and Dynamics of Mesoscale Eddies in Fram Strait from a Comparison between Two High-Resolution Ocean–Sea Ice Models, *Ocean Science*, 16, 1225–1246, <https://doi.org/10.5194/os-16-1225-2020>, 2020.




Figure 7. Spatio-temporal evolution along the 1400 m isobath of modelled a) EKE, b) BT, c) BC, and d) $f(q_y) \equiv \min(q_y) \cdot \max(q_y)$ (negative values indicate a measure of cross-isobath sign-change change of sign of the lateral PV gradient $-q_y$. Temporal averaging is 1 month and the spatial smoothing is 20 km across the isobath) e) cumulative along-isobath integral of eddy temperature flux divergence, $\int_0^x \nabla \cdot \mathbf{q}_T dx$, where $\mathbf{q}_T = (\overline{u'T'}, \overline{v'T'})$. Each group of panels include the spatio-temporal field (colours), its time-average (top), and its along-isobath average (right), except in panel e) where the panel on the right shows the product of along-isobath integral evaluated at the minimum and maximum end of q_y calculated across the isobath, ca. 180 km. In addition, time-averaged $\bar{u}$ (negative values are indicative of a change in sign). Line plots in horizontal panels are time averages and those in vertical panels (e) are shown (green, top panel on the right are along-isobath averages axes). Temporal averaging is 1 month and the spatial smoothing is 20 km. October corresponds to months 1 and 13, April to months 7 and 19. Triangles mark approximate locations of the virtual moorings. Distance-axis tick labels intervals every 25 km correspond to the dots on the 1400 m isobath in Fig. 6. All variables are layer-averaged between 100 m and 2000 m or the seabed, except $\nabla \cdot \mathbf{q}_T$ which is between 100 m and 500 m.




Figure 8. Time series of (a, [eb](#)) EKE, ([bc](#), [ed](#)) BT, and ([ee](#), f) BC from in situ moorings (black pluses), from the model using virtual moorings (purple dots), and the average along the 1400 m isobath near moorings (green open circles). The [first-two](#) model [year-is-shown-in-dark](#), and [the-second-year-in-pale-colours](#) years were averaged together. In ([ee](#), f), BC are shown for the same levels as the moorings (Fig. 5), [and-pale-purple-dots](#). The in situ mooring time series are offset by five days for clarity [as in Figs. 4, 5](#) but monthly averaged.




Figure A1. Time series of (a, b) EKE, (c, d) eddy momentum flux, (e, f) and eddy density flux, at the (left) western and (right) eastern arrays. In each panel a–f, the moorings are ordered from the offshore (top) to onshore (bottom), e.g. W3, W2, W1, W0. The depth range at W0 was 30 m to 80 m, and the vertical axis is scaled by a factor of 2 for clarity.




Figure B1. Average simulated along-slope current velocity u (colours) at (a, c) the western and (b, d) eastern mooring arrays during (a, b) autumn and winter (September–February) and (c, d) spring and summer (March–August) averaged over the two-year period. Grey contours show velocity every 10 cm s^{-1} , white dashed isopycnals every 0.1 kg m^{-3} , and black isotherms every 1°C . Triangles at the top mark the virtual mooring locations (a, b) and vertical white dash-dotted lines mark the location of the 1400 m isobath used in Section 3.3.2.




Figure C1. Example time series of EKE, BT, and BC at 460 m depth at the western (left) and eastern (right) mooring arrays showing the sensitivity to the choice of passband and filter type used in the band-pass filtering for fluctuation calculations.