

Seasonal to long-term variability of natural and anthropogenic carbon concentrations and transports in the subpolar North Atlantic Ocean

Raphaël Bajon ^{1,2}, Lidia I. Carracedo ¹, Herlé Mercier ¹, Rémy Asselot ³, and Fiz F. Pérez ⁴

¹Laboratoire d'Océanographie Physique et Spatiale (UMR 6523 LOPS), Univ. Brest, CNRS, IFREMER, IRD, IUEM, IFREMER Centre de Bretagne, 29280 Plouzané, FRANCE

²Now at Department of Oceanography, School of Earth and Space Science and Technology, University of Hawai'i at Mānoa, Honolulu, HI, 96822, USA

³BRGM - French Geological Survey, F-33600 Pessac, France

⁴Oceanographic department, IIM, Street, Vigo, 10587, Pontevedra, SPAIN

Correspondence: Raphaël Bajon (raphael.bajon@ifremer.fr)

Abstract. The Atlantic Meridional Overturning Circulation (AMOC) is integral to the climate system, transporting heat and anthropogenic carbon across the North Atlantic (NA) from subtropical to subpolar latitudes. This physical mechanism promotes the uptake and sequestration of atmospheric CO₂ through surface cooling as warm water advances northward and consequently sinks through deep winter convection. Using ship-based observations, ocean reanalyses, neural networks, and a back-calculation approach, we present a 30-year monthly time series of contemporary carbon (natural, C_{nat} and anthropogenic, C_{ant}) concentrations and transports at the A25-OVIDE hydrographic section in the subpolar NA Ocean, and assess their variability from seasonal to long-term scales. We divided the section into essential layers, including the upper branch of the AMOC (uMOC) and the mixed layer (ML). Our findings indicate that the full-section-averaged C_{nat} concentration shows no significant trend over the 30-year period. In contrast, the full-section-averaged C_{ant} concentration increased by more than one third over the 30-year period, attributed to ~~anthropogenic influences and~~ the anthropogenic increase in atmospheric CO₂ ~~increase~~. Seasonal and interannual variability is more pronounced in the uMOC and in the ML, where deep convection and biological activity impact their concentration, than in the deeper ocean. The seasonal deepening of the ML in winter contributes two thirds and one half of its ML concentration for C_{nat} and C_{ant} , respectively, the rest being attributed to biology and solubility. The C_{ant} and C_{nat} transports are predominantly determined by the variability of volume transport, except for the decadal trend in C_{ant} transport which is primarily influenced by changes in C_{ant} concentration. The variability in tracer transport is the largest in the uMOC, which exhibits a seasonal peak-to-peak amplitude of approximately 25% of the annual mean tracer transport. These results offer new insights to refine model representations and improve our understanding of the subpolar NA carbon dynamics.

1 Introduction

20 The atmospheric CO₂ concentration has surged by 50% since the onset of the industrial revolution in the 1750s, ~~currently~~ surpassing 420 ppm in 2023 (Keeling et al., 2005). In response to this rapid increase, the ocean acts as a vital ~~natural reservoir~~ ~~(carbon sink)~~carbon sink, absorbing $2.9 \pm 0.4 \text{ PgC yr}^{-1}$ from the atmosphere and therefore compensating for approximately a quarter of total annual CO₂ emissions (~~Friedlingstein et al., 2023~~)(Friedlingstein et al., 2025). This oceanic uptake is facilitated by the massive carbon storage capacity of the ocean –its dissolved inorganic carbon (DIC) reservoir is approximately 50
25 times larger than the atmospheric reservoir (~~Friedlingstein et al., 2023~~). ~~Consequently, the ocean DIC pool comprises both~~ (Friedlingstein et al., 2025). The oceanic DIC pool can be divided into its natural (C_{nat}) and anthropogenic (C_{ant}) DIC fractions, such that $\text{DIC} = C_{\text{nat}} + C_{\text{ant}}$. C_{nat} includes preformed DIC as well as DIC resulting from the biological and carbonate pumps.

The ocean's ability to absorb atmospheric CO₂ and redistribute DIC within the water column is governed by the interplay of physical and biological processes — namely physical and biological carbon pumps (PCP and BCP), respectively — oper-
30 ating within or close to the ocean mixed layer (ML). ~~The ML represents the ocean surface layer in direct exchange with the atmosphere, characterized by vertically uniform properties such as density, salinity (S); oceanic mixed layer (ML) corresponds to the near-surface layer of the ocean where turbulent processes, primarily induced by wind forcing, buoyancy fluxes, and wave breaking, maintain quasi-homogeneous~~ temperature (T), salinity (S) and dissolved oxygen (O₂) ~~profiles. It represents the portion of the ocean directly interacting with the atmosphere, where weak vertical gradients may still persist. The depth~~
35 of the ML is generally defined from a threshold criterion based on potential density or temperature relative to surface values. Within the ML, the PCP operates through CO₂ solubility – enhanced in colder waters – and vertical mixing. Meanwhile, the BCP encompasses the photosynthetic fixation of DIC by phytoplankton, followed by sinking and remineralization of organic matter at depth (Dall'Olmo et al., 2016; Diaz et al., 2021; Lacour et al., 2019). By modulating carbon uptake and redistribution, the ML thus plays a pivotal role in climate regulation (Sallée et al., 2021).

40 The ocean's capacity to absorb and store CO₂ is not spatially homogeneous. The ~~NA~~North Atlantic (NA), characterized by the deepest ML in the world, complex physical dynamics, and strong biological activity, plays a significant role in the uptake and storage of global CO₂ uptake and storage (Dall'Olmo et al., 2016; Resplandy et al., 2018; Pérez et al., 2010, 2013, 2024). Despite covering only 15% of the global surface ocean, the NA accounts for approximately 25% of contemporary global CO₂ ocean uptake and a quarter of the global ocean C_{ant} inventory (DeVries, 2014; Friedlingstein et al., 2019; Gruber et al., 2019;
45 Khatiwala et al., 2013), which is the highest per unit area of the global ocean (Gruber et al., 2023; Sabine et al., 2004). The latter is partly related to the Atlantic Meridional Overturning Circulation (AMOC). The upper limb of the AMOC drives the poleward transport of C_{ant} from the subtropics to the subpolar NA (Brown et al., 2021; Pérez et al., 2013; Zunino et al., 2015, 2014), a regional convergence zone where the deepest penetration of C_{ant} occurs (Sabine et al., 2004; Mikaloff Fletcher et al., 2006). Previous studies have highlighted the sensitivity of C_{ant} transport to the strength of the AMOC (Boers, 2021;
50 Caesar et al., 2021, 2018; Jackson et al., 2022; Thornalley et al., 2018), with implications for carbon sequestration and air-sea fluxes (Pérez et al., 2013; Zunino et al., 2015; Brown et al., 2021) (Liu et al., 2023; Pérez et al., 2013).

Discrepancies persist between observational estimates and model simulations, particularly with regard to ocean C_{ant} and C_{nat} transports (Racapé et al., 2018; Tjiputra et al., 2010; Mikaloff Fletcher et al., 2006), and the sea surface partial pressure of CO_2 ($p\text{CO}_2$) driving air-sea fluxes (Rodgers et al., 2023). Although individual cruises provide indispensable reference estimates, they do not fully capture the temporal variability in the transport and concentration of C_{ant} and C_{nat} (Zunino et al., 2015, 2014; McCarthy et al., 2015; Mercier et al., 2024). The magnitude, variability and factors that govern the contribution of ocean circulation to regional storage of C_{ant} or C_{nat} , and consequently the resilience of the ocean carbon sink to global changes, thus remain largely unexplored (Hauck et al., 2020; Henson et al., 2022).

In particular for the SubPolar North Atlantic (SPNA), most studies have focused mainly on decadal changes, with biennial cruises measuring C_{nat} and C_{ant} transports (Zunino et al., 2014, 2015; Fontela et al., 2019), or on high-temporal-resolution volume transports (McCarthy et al., 2015; Caesar et al., 2021; Fu et al., 2023; Tooth et al., 2023; Mercier et al., 2024), but there is still a dearth of high-temporal-resolution tracer transport data. This study addresses this research gap by presenting the first 30-year observation-based monthly time series of surface-to-bottom C_{nat} and C_{ant} transport across the A25-OVIDE Greenland to Portugal section in the northern North Atlantic (Fig. 1) between 1993 and 2022. By combining ship-based data, ocean reanalyses, neural networks (NN), and a back-calculation (BC) approach for ocean C_{ant} estimation, this research aims to improve our understanding of seasonal to long-term variability in C_{nat} and C_{ant} concentrations (hereinafter marked as $[C_{nat}]$ and $[C_{ant}]$) and transports. It will contribute to improve the predictions of carbon uptake and storage in the NA by providing a novel and comprehensive assessment of the regional surface-to-bottom seasonal cycles and long-term trends. We decompose the net transport of C_{nat} and C_{ant} into vertical layers to differentiate various signals: we split the water column into the upper and lower limbs of the AMOC (uMOC and lMOC, respectively) and also distinguish between the ML and what lies beneath it (bML).

2 Data and methods

This study is based on the Greenland-to-Portugal OVIDE section, known as A25 by GO-SHIP (Sloyan et al., 2019). We used two types of data: (1) hydrographic data (T, S, $[\text{O}_2]$, nutrients, total alkalinity $[A_T]$, pH, velocities) from the 1997 FOUREX cruise and from the 2002–2018 A25 OVIDE biennial repeats (section 2.1), referred to as reference dataset; and (2) ocean reanalysis data (velocities, T, S monthly gridded fields) at the A25-OVIDE section (section 2.2). We applied neural networks (NN) algorithms to the reanalysis property data to generate monthly gridded fields of $[\text{O}_2]$, nutrients, $[A_T]$ and $[\text{DIC}]$ (section 2.4). Using hydro- and NN-based reanalysis property datasets, we used a back-calculation approach (Section 2.3) to calculate $[C_{ant}]$, $[C_{nat}]$ being the difference between $[\text{DIC}]$ and $[C_{ant}]$. The so-derived $[C_{ant}]$ and $[C_{nat}]$ fields were then combined with the corresponding velocity fields to compute the time series of cross-A25 section C_{ant} and C_{nat} transports (Section 2.5). C_{ant} and C_{nat} transports were divided into different vertical regions (Section 2.6) to assess their seasonal to interannual and long-term variability (Section 2.7). Finally, we evaluated the performance of our ocean reanalysis-NN-BC method (hereinafter referred to as OR-NN-BC method) and its uncertainty error in section 2.8.

2.1 GO-SHIP A25 OVIDE hydrographic section

85 The Portugal-to-Greenland GO-SHIP A25 OVIDE hydrographic section (Fig. 1), referred to here as A25, has been repeated biannually in summer since 2002 (Mercier et al., 2024; Sloyan et al., 2019). This study uses data from nine A25 cruise repeats that span 2002–2018. To extend the reference dataset further back in time, data from the FOUREX 1997 cruise (Álvarez et al., 2003; Lherminier et al., 2007), which differs slightly from the A25 positions, have also been included. The T and S of the CTD sensors are collocated with nutrients (nitrate, phosphate, and silicate), $[A_T]$, pH and $[O_2]$ from bottle samples. BGC
90 data consistency was ensured by applying the GLODAP recommended adjustments to the measured values of $[O_2]$, nitrate, phosphate, silicate, pH and $[A_T]$ (see "Recommended adjustment values" at <https://glodapv2.geomar.de>) (Olsen et al., 2019). We used these data to compute C_{nat} and C_{ant} (2.3), which are used here in conjunction with absolute velocities to compute property transports (2.5) (Daniault et al., 2016; Zunino et al., 2017, 2015, 2014; Pérez et al., 2013; Fontela et al., 2019; Lherminier et al., 2007, 2010; Gourcuff et al., 2011; Mercier et al., 2024). Geostrophic velocities were derived by integrating
95 geostrophic shears, calculated from T and S data obtained at hydrographic stations, using as reference simultaneous velocity measurements from a Ship-mounted Acoustic Doppler Current Profiler (S-ADCP). Ekman transport, estimated from NCEP data, was incorporated into the surface layer (0–30 m). An inverse model was applied to compute velocity corrections for each pair of hydrographic stations to ensure volume conservation (see Lherminier et al. (2007, 2010); Gourcuff et al. (2011); Mercier et al. (2024, 2015); Daniault et al. (2016); Zunino et al. (2014)). The A25 velocities v used here are normal to the
100 section and correspond to geostrophic velocities plus Ekman velocities (Lux et al., 2001; Lherminier et al., 2007).

2.2 Ocean products

The ocean reanalysis datasets used (Table 1) are GLOSEA5 (Scaife et al., 2014; MacLachlan et al., 2015), ECCO (Fenty and Wang, 2020), EN4 (Good et al., 2013) and CORA (Szekely et al., 2019). [Ocean reanalyses are data-driven products with varying levels of complexity, and each individual product has its own strengths and limitations. Using the mean concentration across products helps mitigate product-specific biases and errors, while the spread among them provides an estimate of the associated uncertainty.](#) All reanalyses provide gridded T and S at monthly resolution. The velocity fields (v) of CORA and EN4 are geostrophic velocities derived from T, S, surface altimetry, while the Ekman velocities are derived from NCEP (Mercier et al., 2024) (Table 1). Velocities for GLOSEA5 and ECCO are full general circulation model (GCM) dynamics (Mercier et al., 2024). EN4 and CORA were interpolated to the positions of the A25 section. For ECCO and GLOSEA5 GCM, the nearest native grid
110 points to the A25 section were used. The reader is referred to Mercier et al. (2024) for a detailed discussion of seasonal to long-term volume transport variability at A25 from GLOSEA5, ECCO, EN4, and CORA. To complement the ocean reanalysis datasets, we also considered the use of the GOBAI- O_2 gridded product (Sharp et al., 2022) ([version 1.1](#)). GOBAI- O_2 provides O_2 monthly fields computed by applying NN to ~~the monthly fields T, S gridded~~ [gridded monthly T and S fields](#) derived from Argo (Roemmich and Gilson, 2009). The original GOBAI- O_2 data ($1^\circ \times 1^\circ$ resolution on 58 depth levels) were interpolated
115 to the positions of the A25 section (Table 1). [The use of GOBAI- \$O_2\$ might introduce additional depth-dependent biases in](#)




Figure 1. North Atlantic subpolar region and the [GOSHIPGO-SHIP](#) A25 OVIDE hydrographic section (called A25 here). The arrows represent the main currents at the A25 section: the NAC, the Western Boundary Current (WBC) [including the East Greenland Current \(EGC\)](#), the Iceland–Scotland Overflow Water (ISOW) and the Denmark Strait Overflow Water (DSOW). Natural carbon [C_{nat}] and anthropogenic carbon [C_{ant}] calculated from the T, S fields from the GLOSEA5 reanalysis (see [Methods2.4](#)) are displayed at A25 for the last year of the study (2022) in the bottom and top panels, respectively (see Section 2). Winter and summer mean mixed layer depths from GLOSEA5 are shown as solid and dashed lines respectively for [C_{nat}]. The isopycnal σ_{MOC} , separating the upper and lower limbs of the AMOC, from GLOSEA5 is shown on [C_{ant}] for April (32.24 kg m^{-3} for a density anomaly referenced at 1000 db) and October (32.13 kg m^{-3}), which are the months with the strongest and weakest seasonal transports in the upper branch of the uMOC respectively. Bathymetry is from GEBCO.

O₂ associated to Argo-O₂ sensors (Bushinsky et al., 2025). Our error estimation method, based on the comparison to the A25 ground truth (see 2.8), ensures that any potential bias from the use of GOBAI-O₂ is reflected in the reported uncertainties.

2.3 Anthropogenic and natural carbon estimates

To determine the C_{ant} fraction from [DIC], we used the carbon-based BC φC_T^0 approach (Pérez et al., 2008; Vázquez-Rodríguez et al., 2009). This BC approach has been broadly applied to study the inventory of C_{ant} , its storage rates, its variability (Pérez et al., 2008; Vázquez-Rodríguez et al., 2009; Pérez et al., 2013; Fröb et al., 2018; Asselot et al., 2024) and the influence of C_{ant} on ocean acidification (Pérez et al., 2018). The input variables for this approach include date, geographical location, T, S, O₂, macronutrients (NO₃⁻, PO₄³⁻ and Si(OH)₄), [A_T] and [DIC]. To estimate the natural carbon **fraction concentration** [C_{nat}], [C_{ant}] (2.3) is subtracted from [DIC] ($[C_{nat}] = [DIC] - [C_{ant}]$). For reference data, T, S, [O₂], macronutrients, [A_T], and pH are measured at A25 bottles. [DIC] is derived from [A_T] and pH from the in situ bottle measurements using the **CO2SYS toolbox (Humphreys et al., 2022)** PyCO2SYS toolbox (Humphreys et al., 2022) (version 1.8). This toolbox also provides the Revelle factor shown in this study. NNs are used to derive the necessary parameters for ocean reanalysis to obtain [C_{ant}] (2.4).

2.4 Choice and application of neural networks

Two different NNs were sequentially applied to ocean reanalysis data to estimate [DIC] -and the evaluation of the error is detailed in section 2.8. Each NN has limitations, and the use of two different NNs was intended to build on the strengths of each NN. First, we applied ESPER NN (Carter et al. (2021), Eq. 8) to the T and S fields of the reanalysis (as well as date and position, i.e., longitude, latitude, and depth) as input to determine [O₂] and macronutrients. Second, we calculate [DIC] and [A_T] using CANYON-B NN - CONTENT (Bittig et al., 2018). CANYON-B uses the same input data as ESPER NN plus [O₂] (O₂ derived from ESPER_NN). The choice of ESPER NN and CANYON-B-CONTENT for the estimation of oxygen and macronutrients and carbon variables, respectively, is based on the performance of the corresponding NNs for each variable (i.e. lowest final **uncertainty error**) (Asselot et al., 2024). Using ESPER alone to calculate [DIC], we found that [DIC] values diverged from cruise-based estimates after 2010, whereas between CANYON-B-CONTENT and observations there was a better agreement (Fig. S1). Instead of a time-dependent prediction of [DIC] as in CANYON-B (Bittig et al., 2018), [DIC] is given for a reference year (2002) in ESPER. Within this NN, the anthropogenic component of DIC (C_{ant}) is calculated as an exponential increase (see Carter et al. (2021), their Eq. 1), assuming that C_{ant} is in a transient steady state (Tanhua et al., 2007) (that is, exponential increases in atmospheric anthropogenic CO₂ should result in the concentration of marine C_{ant} that increases at rates proportional to the concentration of atmospheric anthropogenic CO₂). The use of ESPER NN to estimate [O₂] and macronutrients and CANYON-B-CONTENT to estimate [DIC] and [A_T] is therefore a reliable compromise for applying NN to T and S fields in the subpolar gyre (Fig. S1). In the particular case of the GOBAI-O₂ dataset (Table 1), we applied ESPER NN only to retrieve macronutrients, then CANYON-B-CONTENT to retrieve [DIC] and [A_T] (Asselot et al., 2024). It is important to note that the A25 2002-2014 data used in this study to obtain the reference estimates are among the source data used for the training phase of both NNs: GLODAPv2.2019 (Olsen et al., 2019; Carter et al., 2021) and GLODAPv2 (Bittig

Table 1. Ocean products, with their time band and their depth range at the A25 section. All ocean products include monthly average values of temperature (T) and salinity (S). Ocean reanalyses also include velocity data and GOBAl-O₂, dissolved oxygen (O₂). (*) Note that for EN4, the T and S fields are provided for the entire water column, but the velocity data are restricted to the depth range 0–2000 m (see Mercier et al. (2024)) (Table 2). The reader is referred to Mercier et al. (2024) for more details on the ocean reanalysis data.

Ocean reanalysis	Variables	Time band	Spatial Resolution (horizontal x vertical) - depth range	Method for T, S	Description of v
GLOSEA5	T, S, v	1993–2022	253 x 75 - full	3D-var	full GCM dynamics
ECCO	T, S, v	1992–2017	67 x 50 - full	state estimate	full GCM dynamics
EN4	T, S, v	1993–2021	108 x 42 - full (*)	objective mapping	geostrophy (obtained with altimetry) + Ekman
CORA	T, S, v	1993–2020	108 x 152 - 0-2000m	objective mapping	geostrophy (obtained with altimetry) + Ekman
GOBAl-O ₂	T, S, O ₂	2004–2021	108 x 152 - 0-2000m	optimal interpolation	/

at the A25 section. All ocean products include monthly average values of temperature (T) and salinity (S). Ocean reanalyses also include velocity data and GOBAl-O₂, dissolved oxygen (O₂). (*) Note that for EN4, the T and S fields are provided for the entire water column, but the velocity data are restricted to the depth range 0–2000 m (see Mercier et al. (2024)) (Table 2). The reader is referred to Mercier et al. (2024) for more details on the ocean reanalysis data.

Ocean products, with their time band and their depth range

et al., 2018; Olsen et al., 2016). However, the reference estimates from the A25 data for 2016 and 2018 can be considered
 150 entirely independent of those obtained using the OR-NN-BC method.

2.5 Transport calculation

2.5.1 Definition

Transport refers to the cross-section volume transport at A25. Net transport at a given time t (both for the A25 hydrographic
 data and ocean reanalyses), $T(t)$ (Eq. 1) is expressed in Sverdrup ($10^6 \text{m}^3 \text{s}^{-1}$), with the integration performed over z from
 155 surface ($z_1 = 0$) to bottom ($z_2 = z_{max}(x)$) and over x from Portugal to Greenland along the A25 line.

$$T(t) = \frac{1}{10^6} \int_x \int_{z_1}^{z_2} v(x, z, t) dx dz \quad (1)$$

The velocity field $v(x, z, t)$ (Table 1) refers to the absolute velocities normal to the section. The concentration $c(x, z, t)$ of
 [C_{nat}] or [C_{ant}] (expressed in $\mu \text{mol kg}^{-1}$) and the velocities $v(x, z, t)$ obtained from the A25 cruise and ocean reanalysis are
 used to calculate the net transport of property $T_p(t)$, in kmol s^{-1} or PgC yr^{-1} . $T_p(t)$ is calculated by integrating the section
 160 and multiplying the property by the flow velocity and density (Eq. 2). The transport of C_{nat} ($T_{C_{nat}}$) and the transport of C_{ant}
 ($T_{C_{ant}}$) are thus determined as:

$$T_p(t) = \frac{1}{10^9} \int_x \int_{z_1}^{z_2} \rho(x, z, t) \cdot c(x, z, t) \cdot v(x, z, t) dx dz \quad (2)$$

For the different transport estimations, volume and tracer transports (Eq. 1, 2), positive (negative) transport means northward
 (southward) transport. The integration limits in z may vary according to the layer of the water depth column considered (see
 165 2.6 for details on the vertical layer separations).

2.5.2 Diapycnal and isopycnal decomposition

Following previous studies on heat (Mercier et al., 2015), fresh water (McDonagh et al., 2015) or property transport (Álvarez
 et al., 2003; Zunino et al., 2014), we decomposed net property transport into a diapycnal and isopycnal term (Eq. 3). The
 diapycnal term refers to the transport of property associated with the overturning circulation, which accounts for the conversion
 170 of light to dense water masses north of the section (Grist et al., 2014). The isopycnal term refers to the gyre circulation and is
 the area integration of the covariance of the volume transport and property anomalies at each longitude and density level along
 the A25 section. This term is called horizontal circulation when decomposition is performed in pressure coordinates (Böning
 and Herrmann, 1994). The net' transport is the net transport of property through the section related to the net northward volume
 transport of approximately 1 Sv associated with the Arctic mass balance (Lherminier et al., 2007). We decompose the property
 175 and velocity as $c(x, \sigma, t) = \langle c \rangle_x(\sigma, t) + c''(x, \sigma, t)$ (for C_{ant} and C_{nat}), $v(x, \sigma, t) = \langle v \rangle_{x, \sigma}(t) + \langle v \rangle_x(\sigma, t) + v'''(x, \sigma, t)$

, where given a quantity a and some spatial direction b we define $\langle a \rangle_b = \frac{\int_b a(b) db}{\int_b db}$. As explained in Zunino et al. (2015), Eq. 2 can be rewritten as:

$$T_p(t) = T_p^{net'}(t) + T_p^{diap}(t) + T_p^{isop}(t) \quad (3)$$

where $T_p^{net'}(t) = \langle v \rangle_{x,\sigma}(t) \cdot \int_x \int_\sigma \langle \rho \cdot c \rangle_x(\sigma, t) dx d\sigma$, $T_p^{diap}(t) = \int_x \int_\sigma \langle v \rangle_x(\sigma, t) \cdot \langle \rho \cdot c \rangle_x(\sigma, t) dx d\sigma$ and $T_p^{isop}(t) = \int_x \int_\sigma v'''(x, \sigma, t) \cdot \rho \cdot c''(x, \sigma, t) dx d\sigma$.

~~We also calculated the same estimator as in Zunino et al. (2014) to further describe C_{ant} diapycnal transport variability (Eq. ??). The estimator is a function of the C_{ant} difference between uMOC and IMOC (ΔC_{ant}) multiplied by the uMOC transport calculated in the density coordinates (Mercier et al., 2015), along with the water density (ρ).~~

$$T_{est} = \rho \cdot \Delta C_{ant} \cdot \text{uMOC}$$

185 2.6 Region separation

2.6.1 uMOC and IMOC

In the SPNA, the upper and lower parts of the MOC, noted uMOC, IMOC respectively, are determined in σ levels (Table 2, see Lherminier et al. (2007, 2010); Mercier et al. (2015); Lozier et al. (2019)). This relies on finding the density coordinate, σ_{MOC} , where the AMOC stream function ($\Psi(\sigma, t)$) is maximum. To do so, we compute the meridional overturning stream function by integrating the across-section transport in density referenced to 1000 dbar (σ_1) from the surface to σ_1 ($\Psi(\sigma_1, t) = \int_{surface}^{\sigma_1} \int_{Portugal}^{Greenland} v(x, \sigma_1, t) dx d\sigma$). The density at which the overturning stream function is maximum, called $\sigma_{1,MOC}$, is bounding uMOC and IMOC (Eq. 4).

$$\sigma_1(x, z, t) \in \{\text{uMOC if } \sigma_1(x, z, t) \leq \sigma_{1,MOC}(t), \text{IMOC if } \sigma_1(x, z, t) > \sigma_{1,MOC}(t)\} \quad (4)$$

Computing the MOC in density coordinates provides a better representation of the thermohaline circulation at the latitudes of the A25 section than separating in depth levels. This is because the northward flow of warm waters transported by the North Atlantic Current (NAC) and the southward flow of colder, denser waters carried by the ~~East Greenland Irminger Current~~ [EGC](#) occur at overlapping depths so that they partially cancel each other when using depth coordinates to find the MOC, as explained by Lherminier et al. (2007); Mercier et al. (2015); Lozier et al. (2019); Mercier et al. (2024). The maximum value of the stream function may vary over time (Mercier et al., 2015), as well as the associated value of $\sigma_{1,MOC}$. For each monthly time step t of the ocean reanalysis (and year of the A25 section), the transport of the uMOC was estimated by setting $z_1 = 0$ (sea surface) and $z_2 = \sigma_{1,MOC}(t)$ in Eq. 1, 2 while the transport of the IMOC corresponds to $z_1 = \sigma_{1,MOC}(t)$ and $z_2 = z_{max}(x)$ (bottom) in Eq. 1, 2.

2.6.2 Mixed-layer depth

Several methodologies have been developed to determine the ML depth using T, S, or density profiles (Brainerd and Gregg, 1995; Thomson and Fine, 2003; de Boyer Montégut, 2004; Holte and Talley, 2009; Holte et al., 2017). The ML depth z_{ML} is defined here as the depth at which the potential density, referenced to the ocean surface and denoted as σ_0 , exceeds the density of the water at a fixed depth of 10 m by a predefined threshold of 0.03 kg m^{-3} (Eq. 5).

$$\sigma_0(x, z = z_{ML}, t) = \sigma_0(x, z = 10, t) + 0.03 \quad (5)$$

The threshold of 0.03 kg m^{-3} has been shown to effectively identify the base of the ML in various oceanic regions around the world (de Boyer Montégut, 2004; Holte and Talley, 2009; Holte et al., 2017). The threshold criterion of 0.01 kg m^{-3} , previously used for Argo-profiling floats in the region (Piron et al., 2016), often resulted in inappropriate ML detections when applied to reanalysis products, particularly during deep convection periods. The use of monthly means (described in 2.2) made it unlikely to capture a profile of perfectly constant density in the ML; thus a more flexible criterion was necessary. In addition, freshwater flows in the Irminger Sea (IS) create a density front on the sea surface. This led us to set the reference density σ_0 at 10 meters instead of the usual 0 meters. The water above z_{ML} is considered part of the ML. The region below z_{ML} that exceeds the criteria σ_0 in Eq. 5 is noted as bML. ML transport is calculated by setting $z_1 = 0$ (sea surface) and $z_2 = z_{ML}(x, t)$ in Eq. 1, 2 while bML transport is calculated by setting $z_1 = z_{ML}(x, t)$ and $z_2 = z_{max}(x)$ (bottom) in the equations.

2.7 Seasonal cycle, interannual filtering~~and~~, trends and annual profile

From the monthly time series of concentration and transport, seasonal signal time series were obtained by applying a two-year high-pass filter over the time series. The new high-frequency time series were grouped by months and the mean for each month was calculated to derive the seasonal (also referred to as intraannual) signal.

Interannual time series were computed by subtracting the high-frequency time series (obtained using the high-pass filter) from the initial one. The Standard Deviation (STD) of this interannual time series will be used as a metric of interannual variability. In the following, the interannual signal will refer to this low-frequency signal. The standard error (STDn) of the average over the reanalyses is defined as the STD between reanalyses divided by the square root of the number of reanalysis used to compute the mean.

Trends are calculated as the slope coefficient of a linear fit. The uncertainty in trends is estimated using the Moving Block Bootstrap method (Kunsch, 1989).

To better quantify the effects of seasonal changes in ML and uMOC thickness on the seasonality of $[C_{nat}]$ and $[C_{ant}]$, we calculated an idealized seasonal anomaly for ML and uMOC using the annual mean vertical profile of the property ($[C_{annual}]$), rather than monthly varying concentrations, and the full variability of ML depth and uMOC thickness. This approach isolates the impact on ML concentrations of physically-driven seasonal changes in layer thickness from biologically-driven seasonal variations in concentration.

Table 2. Names of the vertical layers and ocean reanalyses used in this study. In the context of transport estimates, a layer denotes the transport across the layer (positive northward). Conversely, in terms of concentrations, it refers to the average concentration over the layer. CORA and GOBAI-O₂ cover the depth range 0–2000m (Table 1) and their use is therefore limited to the uMOC and mixed-layer (ML) layers. Only GLOSEA5 and ECCO contribute to the calculation of net, IMOC and bML transports. uMOC* is the average of the uMOC quantity (concentration or transport) over these two ocean reanalyses only.

Names	Region signification	Ocean product used to compute the quantity	
		Concentration	Transport
net, FSA	full section	GLOSEA5, ECCO, EN4	GLOSEA5, ECCO
uMOC	upper branch of the AMOC	GLOSEA5, ECCO, EN4, CORA, GOBAI-O ₂	GLOSEA5, ECCO, EN4, CORA
IMOC	lower branch of the AMOC	GLOSEA5, ECCO, EN4	GLOSEA5, ECCO
uMOC*	uMOC with only the ocean reanalyses that discretized the full water column	GLOSEA5, ECCO	GLOSEA5, ECCO
ML	Mixed Layer	GLOSEA5, ECCO, EN4, CORA, GOBAI-O ₂	GLOSEA5, ECCO, EN4, CORA
bML	below the Mixed Layer	GLOSEA5, ECCO, EN4	GLOSEA5, ECCO

Names of the vertical layers and ocean reanalyses

used in this study. In the context of transport estimates, a layer denotes the transport across the layer (positive northward). Conversely, in terms of concentrations, it refers to the average concentration over the layer. CORA and GOBAI-O₂ cover the depth range 0–2000m (Table 1) and their use is therefore limited to the uMOC and mixed-layer (ML) layers. Only GLOSEA5 and ECCO contribute to the calculation of net, IMOC and bML transports. uMOC* is the average of the uMOC quantity (concentration or transport) over these two ocean reanalyses only.

2.8 Error estimation

235 The error in the estimation of $[C_{nat}]$ and $[C_{ant}]$ is the sum of the errors associated with the combined use of ocean reanalyses, NN, and the BC approach. The error associated with the use of NNs is evaluated at the hydrographic section A25, using estimates directly obtained from seawater samples as a reference, first at the sampling points (2.8.1) and then for integrated variables such as averaged concentrations (2.8.2) and transports (2.8.3) in predefined layers. The overall errors resulting from both the use of NN and reanalyses have been estimated altogether for average regional concentrations and transports and
240 comparing them with hydrographic data. The resulting ~~RMSD~~ Root Mean Square Deviation (RMSD) give the final errors for integrated concentrations (2.8.2) and transports (2.8.3), where all errors have been taken into account. The error of the BC approach is fixed according to previous studies to $5.2 \mu\text{mol kg}^{-1}$ (Vázquez-Rodríguez et al., 2009; Pérez et al., 2013).

Using error propagation to calculate an error on an integral quantity (e.g., average concentration per layer or property transport by uMOC) would require postulating the error correlation between different variables. This correlation is unknown.
245 Our method of direct comparison with observational results avoids this issue, enabling us to compare final integrated values to reference hydrographic data.

2.8.1 Neural networks evaluation on A25 hydrographic data

The error resulting from the use of NN was quantified by applying NNs to the T and S A25 bottle data to estimate [DIC], and comparing these values to the original [DIC] estimated directly derived from observations (T, S, O_2 , nutrients, pH, and
250 A_T bottle data) (section 2.8). The ~~Root Mean Square Deviation (RMSD)~~ RMSD between both estimates was $9.7 \mu\text{mol kg}^{-1}$ (mean for all A25 cruises) for [DIC] (Table S1). The uncertainty of the reference [DIC] from the A25 data is equal to $5.8 \mu\text{mol kg}^{-1}$ (see Supplementary Text S1) and comes from the uncertainties on the measurements of A_T and pH in seawater samples. The RMSD is nearly equal to the median uncertainty of [DIC] of $9.1 \mu\text{mol kg}^{-1}$ for CANYON-B-CONTENT (Bittig et al. (2018), see their Table 2). The $[\text{O}_2]$ generated by ESPER has a RMSD of $7.8 \mu\text{mol kg}^{-1}$ with the A25 bottle (Table S1), which
255 is within the mean uncertainty of $9.1 \mu\text{mol kg}^{-1}$ provided in the North Atlantic region by ESPER NN (Carter et al., 2021). The mean uncertainty given by GOBAI- O_2 in $[\text{O}_2]$ is $6.3 \mu\text{mol kg}^{-1}$ in the SPNA (Sharp et al., 2022). ESPER also retrieves confident nutrient values. For example, nitrate shows an RMSD of $2.8 \mu\text{mol kg}^{-1}$ with the A25 data. Applying φC_T^0 on our NN-generated fields (from the initial T, S of the A25 bottle data), we find a RMSD of $5.1 \mu\text{mol kg}^{-1}$ for $[C_{ant}]$ (Table S1) and $11.3 \mu\text{mol kg}^{-1}$ for $[C_{nat}]$. Considering that the BC φC_T^0 approach error of $5.2 \mu\text{mol kg}^{-1}$ (Vázquez-Rodríguez et al., 2009; Pérez et al., 2013) is independent of that due to the use of NNs for $[C_{ant}]$, the final error of our $[C_{ant}]$ estimates is $7.3 \mu\text{mol kg}^{-1}$. As highlighted in Asselot et al. (2024), uncertainties in $[\text{O}_2]$ and [DIC] may cancel each other out, resulting in relatively low errors on the final $[C_{ant}]$ values, given uncertainties in $[\text{O}_2]$ and [DIC].

The use of $[\text{O}_2]$ as an input variable for ESPER NN reduces the uncertainty error of the predicted variables (Carter et al. (2021); see their Table 10, difference between Eq. 7, including $[\text{O}_2]$, and Eq. 8, without $[\text{O}_2]$). ~~We evaluated this using data from the~~ The use of ESPER to estimate $[\text{O}_2]$ was validated by comparing NN-derived estimates against A25 bottle .Although our results also pointed to a reduced uncertainty for the estimates of measurements. $[C_{ant}]$ [DIC] and RMSD increases only

marginally when using ESPER-estimated rather than measured $[C_{nat}O_2]$ when oxygen was used as predictor in ESPER-NN (9.7 vs. 9.5 $\mu\text{mol kg}^{-1}$, Table S1), the difference with the calculations based solely on T and S (as does $[O_2C_{ant}]$ being estimated by ESPER-NN) is small (RMSD (5.1 vs. 4.7 $\mu\text{mol kg}^{-1}$, Table S1), supporting the suitability of ESPER for $[O_2]$ estimation in this framework. Hence, the NN predictions at A25 based on T, S are considered sufficiently robust and coherent with respect to the final values of $[C_{nat}]$ and $[C_{ant}]$.

Our concentration errors are consistent with those derived from error propagation by Asselot et al. (2024), using 3 Argo- O_2 floats data and a similar NN-based approach. They reported [DIC] and $[C_{ant}]$ errors of 10.5 and 5.9 $\mu\text{mol kg}^{-1}$, respectively, compared to 9.7 and 5.1 $\mu\text{mol kg}^{-1}$ in the present study using reference hydrographic data.

2.8.2 Neural networks applied to ocean reanalyses

Using ocean reanalysis introduces additional errors stemming from errors in the T and S fields. These errors are included when comparing the section averaged $[C_{nat}]$ and $[C_{ant}]$ obtained from OR-NN-BC with the A25 bottle estimates. In this comparison, we compare the synoptic A25 hydrographic sections to the monthly average ocean reanalyses, including the set of errors. We find a RMSD for $[C_{nat}]$ averaged over reanalyses of 1.4, 2.4, and 1.9 $\mu\text{mol kg}^{-1}$ for the net, uMOC and IMOC layers, respectively (Table 3,S2). The RMSD for $[C_{ant}]$ is of the same order (1.2, 1.2, 0.7 $\mu\text{mol kg}^{-1}$, for the net, uMOC and IMOC layers, respectively) (Table 3,S2). Considering the spreading between ocean reanalysis (Table 4,S3), we find a RMSD for uMOC in the range 2.9–5.2 $\mu\text{mol kg}^{-1}$ for $[C_{nat}]$ and 1.4–2.0 $\mu\text{mol kg}^{-1}$ for $[C_{ant}]$.

The averaging of the variables reduces the NN errors compared to those calculated for the same variable at the sample points (Tables S1, 3, 4, S2, S3), suggesting that the errors in the concentrations calculated by NNs are mainly random. Examining the differences between the results of the reanalyses and the A25 data, the uMOC biases for $[C_{nat}]$ range from -3.1 to +4.3 $\mu\text{mol kg}^{-1}$ depending on the reanalysis with 2.5 to 3.9 $\mu\text{mol kg}^{-1}$ as STD. For $[C_{ant}]$ the biases vary from -0.52 to +0.1 $\mu\text{mol kg}^{-1}$ with 1.3 to 2 $\mu\text{mol kg}^{-1}$ as STD. Opposite signs in bias reduce RMSD while averaging between reanalyses, especially for $[C_{nat}]$. The STDs are of the same order of magnitude as the biases for $[C_{nat}]$ and greater than the biases for $[C_{ant}]$.

2.8.3 Tracer transport

For the hydrographic section A25, the uncertainties of the transport of the property are calculated with the inverse formalism as in Zunino et al. (2015). The average uncertainty in $T_{C_{nat}}$ is equal to 2.4, 1.1, 2.0 PgC yr^{-1} for the net, uMOC and IMOC layers, respectively, while we find 0.04, 0.02 and 0.03 PgC yr^{-1} for $T_{C_{ant}}$ for the net, uMOC and IMOC layers, respectively.

To evaluate the impact of our method on tracer transport estimates, we calculated RMSD between $T_{C_{nat}}$ and $T_{C_{ant}}$ calculated with A25 bottle data (2.8.1) and those derived from ocean reanalysis using OR-NN-BC (Tables 3,4,S4). $T_{C_{nat}}$ presents the RMSD for 2002–2018 of 1.7 PgC yr^{-1} for the uMOC layer averaged over all reanalyses, while $T_{C_{ant}}$ presents the RMSD for 2002–2018 of 0.05 PgC yr^{-1} for the same layer (Table 3,4,S4). Both property concentration and volume transport errors are included in this final error estimate, which considers the A25 hydrographic sections as a reference. When calculating RMSD for each ocean reanalysis within the uMOC layer, a range of 2.3–3.7 PgC yr^{-1} is found for $T_{C_{nat}}$, and 0.05–0.09 PgC yr^{-1} for $T_{C_{ant}}$ (Table 4,S5). Like for the property, taking the average transport between products also reduces the RMSD within

Table 3. RMSD (for each layer) between $[C_{nat}]$, $[C_{ant}]$, $T_{C_{nat}}$, $T_{C_{ant}}$ computed with the OR-NN-BC method (mean of all ocean reanalyses) and the estimations derived from the bottle measurements at A25 (2002–2018). The units are $\mu\text{mol kg}^{-1}$ for concentration and PgC yr^{-1} for transport of properties. The comparison is made in June for all years in which the A25 cruises were carried out. For the hydrographic section A25, the uncertainties of the transport of the property are calculated with the inverse formalism as in Zunino et al. (2015). The average uncertainty in $T_{C_{nat}}$ is equal to 2.4, 1.1, 2.0 PgC yr^{-1} for the net, uMOC and IMOC layers, respectively, while we find 0.04, 0.02 and 0.03 PgC yr^{-1} for $T_{C_{ant}}$.

A25 2002–2018	net	uMOC	IMOC
$[C_{nat}]$	1.4	2.4	1.9
$[C_{ant}]$	1.2	1.2	0.7
$T_{C_{nat}}$	0.8	1.7	2.6
$T_{C_{ant}}$	0.04	0.05	0.05

RMSD(for each layer) between C_{nat} , C_{ant} , $T_{C_{nat}}$, $T_{C_{ant}}$ computed with the OR-NN-BC

method (mean of all ocean reanalyses) and the estimations derived from the bottle measurements at A25 (2002–2018). The units are $\mu\text{mol kg}^{-1}$ for concentration and PgC yr^{-1} for transport of properties. The comparison is made in June for all years in which the A25 cruises were carried out.

Table 4. RMSD between $[C_{nat}]$, $[C_{ant}]$, T , $T_{C_{nat}}$, $T_{C_{ant}}$ for the uMOC computed with the OR-NN-BC method (by ocean reanalysis) and the estimations derived from sea bottle measurements at the A25 cruise (2002–2018). The units are $\mu\text{mol kg}^{-1}$ for concentration, Sv for transport and PgC yr^{-1} for transport of properties. The comparison is made in June same as above.

A25 2002–2018	GLOSEA5	ECCO	EN4	CORA	GOBAI-O ₂	Mean (Table 3)
uMOC $[C_{nat}]$	2.9	5.2	4.9	4.1	3.3	2.4
uMOC $[C_{ant}]$	1.5	1.3	2.0	1.9	1.4	1.2
uMOC T	4.3	3.2	2.8	3.1	/	2.0
uMOC $T_{C_{nat}}$	3.7	2.7	2.3	2.6	2.4	1.7
uMOC $T_{C_{ant}}$	0.09	0.06	0.05	0.06	0.06	0.05

RMSDbetween C_{nat} , C_{ant} , T , $T_{C_{nat}}$

$T_{C_{ant}}$ for the uMOC computed with the OR-NN-BC method (by ocean reanalysis) and the estimations derived from sea bottle measurements at the A25 cruise (2002–2018). The units are $\mu\text{mol kg}^{-1}$ for concentration, Sv for transport and PgC yr^{-1} for transport of properties. The comparison is made in June same as above.

300 the A25 observations. Using only GLOSEA5 and ECCO (see Table 2), $T_{C_{nat}}$ presents the RMSD for 2002–2018 of 0.8 and 2.6 PgC yr⁻¹ for the net and IMOC layers, respectively, while $T_{C_{ant}}$ presents the RMSD for 2002–2018 of 0.04 and 0.05 PgC yr⁻¹ for the same layers, respectively (Table 3).

3 Results

3.1 C_{ant} and C_{nat} concentrations

305 Along the A25 OVIDE section, the C_{nat} fraction represents, on average, 98% of the DIC, while the C_{ant} fraction accounts for only the remaining 2%. The $[C_{nat}]$ distribution shows a vertical gradient of generally lower to higher concentrations from surface to depth (Fig. 1), which is largely shaped by the biological carbon pump (Passow and Carlson, 2012). In contrast, the $[C_{ant}]$ distribution shows an opposite vertical gradient, with the highest concentrations in the upper layers due to direct contact with the atmosphere (Fig. 1,2). $[C_{ant}]$ decreases with depth and also from East to West, as the cold subpolar waters contain
310 less $[C_{ant}]$ than the warmer subtropical waters (Sabine et al., 2004). In particular, the highest $[C_{ant}]$ signature is found in and east of the North Atlantic Current (NAC) (Fig. 1) which transports C_{ant} -loaded subtropical waters to higher latitudes. This current represents the primary source of C_{ant} to the subpolar gyre from advection (Brown et al., 2021; Pérez et al., 2013).

The results discussed in this section are the averages over all reanalyses, unless otherwise indicated. We recall that typical errors of our method on integrated concentration range from 2.9 to 5.2 $\mu\text{mol kg}^{-1}$ for $[C_{nat}]$ and from 1.4 to 2.0 $\mu\text{mol kg}^{-1}$ for $[C_{ant}]$ (Section 2.8).
315

3.1.1 Seasonal

The amplitude of the $[C_{nat}]$ seasonal cycle is the highest for ML (Fig. 2,3b). We find a maximum positive anomaly of $32.1 \pm 1.4 \mu\text{mol kg}^{-1}$ in March and a maximum negative anomaly of $-37.1 \pm 1.9 \mu\text{mol kg}^{-1}$ in March and August, respectively August (Fig. 3b); with zero anomaly in May and November. The seasonal variability of $[C_{nat}]$ in ML (ML $[C_{nat}]$ hereafter) may be due
320 to the variation in depth of the ML seasonal variations of both $[C_{nat}]$ and the ML depth (as it deepens, the ML incorporates higher values of $[C_{nat}]$) and the seasonal variation of C_{nat} . To better understand the effect of varying the ML depth on the seasonal variability of ML C_{nat} , we calculated the effect that a change the seasonality in ML depth applied to an annual mean $[C_{nat}]$ profile would have (Fig. 3b), see 2.7), the difference between the two being related to the seasonality of the biological activity. We observe that seasonal variations in the ML depth account for approximately two-thirds of the seasonal ML $[C_{nat}]$
325 amplitude in March, and only up to one-third in August (Fig. 3b). All ocean reanalyses show a deepening of the ML in winter (Fig. S2) so that the deep layers with higher $[C_{nat}]$ have a greater contribution to ML $[C_{nat}]$. The deepening is more pronounced in the Irminger Sea (IS) than in the NAC (Iceland Basin) or east of the NAC (Iberian Basin) (Fig. S3) which, combined with the downward increase in $[C_{nat}]$, explains that the amplitude of the ML $[C_{nat}]$ seasonal cycle is larger in the IS ($70.6 \mu\text{mol kg}^{-1}$) than in the NAC ($60.2 \mu\text{mol kg}^{-1}$).




Figure 2. Area mean time series of (a) $[C_{nat}]$ and (b) $[C_{ant}]$ in all the layers of interest: the upper and lower branches of the Atlantic Meridional Overturning Circulation (uMOC in green, and lMOC in dark green, respectively), Mixed Layer (ML) in yellow green, and below the ML (bML) in dark blue. The full section averaged concentration (referred to as FSA) is shown in black. For the uMOC and ML layers, the concentration values were computed as the mean of the estimates from GLOSEA5, ECCO, CORA, EN4 and GOBAI-O₂ (Table 2). For the net, lMOC, and bML layers, the concentration values represent the average of the estimates derived from GLOSEA5, ECCO and EN4. The gray shading along the lines represents the standard deviation of all product estimates used in the monthly averaging, divided by the square root of the number of reanalysis. The dashed lines for C_{nat} (top) represent a low-pass filter time series (cutoff frequency of 24 months). For C_{ant} (bottom), dashed lines indicate the linear trend. The A25 cruise $[C_{nat}]$ and $[C_{ant}]$ estimates for uMOC, lMOC, and net are shown in red, with red vertical lines denoting the error bars (see methods). The grey shading at the end of the time series highlights the period for which only the GLOSEA5 reanalysis is available.

330 Looking at the sea surface, the C_{nat} averaged at 5-m depth along the section (Fig. 4a) is maximum in March ($2066.3 \mu\text{mol kg}^{-1}$) and minimum in August ($2009.7 \mu\text{mol kg}^{-1}$), which results in a seasonal amplitude of $57.2 \mu\text{mol kg}^{-1}$ (Fig. 4a). It shows the same pattern as the changes in seasonal concentration of ML and is a proxy for the variation of C_{nat} in ML, although a deviation of up to $16.0 \mu\text{mol kg}^{-1}$ between the two is observed in March, coming from the IS ($17.0 \mu\text{mol kg}^{-1}$) (Fig. 4a).

The seasonal cycle of uMOC [C_{nat}] varies in phase with the seasonal cycle of ML, but is six times lower (Fig. 3a,b). A maximum positive anomaly of $5.7 \pm 0.5 \mu\text{mol kg}^{-1}$ is observed in the uMOC in April, and a maximum negative anomaly of $-4.6 \pm 0.6 \mu\text{mol kg}^{-1}$ in November, with zero anomaly occurring in June (Fig. 3a). As for the ML, the seasonal cycle of uMOC [C_{nat}] appears to be related to changes in the surface area of uMOC and changes in the biological pump. Consistent with the downward increase in [C_{nat}], the maximum (minimum) thickness of the uMOC in April (fall) (Fig. S2) corresponds to the maximum (minimum) [C_{nat}] (Fig. 1,4c)(Fig. 1b). The Quantifying the proportion of seasonality related to biological activity can be quantified for the full section as the difference between the seasonality of uMOC C_{nat} and the seasonality calculated by taking an annual value of C_{nat} (Fig. 3a). A a maximum difference is observed in March–April ($1.7 \mu\text{mol kg}^{-1}$) and September ($1.9 \mu\text{mol kg}^{-1}$). Mercier et al. (2024) showed that the thickness of the uMOC varies seasonally in opposite directions in the IS compared to the rest of the section, decreasing (increasing) in winter (summer) in the IS, but increasing (decreasing) in the eastern part of the section. Fig. 4c shows that the amplitude of the seasonal cycle for uMOC [C_{nat}] in the IS is markedly different from the rest of the section. Although the maximum anomaly observed in April is in phase with the rest of the section, there is a pronounced minimum in August that we relate to maximum biological activity (see, e.g., Lacour et al. (2015)). This suggests that seasonal changes in the uMOC [C_{nat}] in the IS depend more than the rest of the section on biologically-driven biologically-driven changes.

Smaller seasonal variations are observed in the lMOC and bML layers (of 0.7 and $4.6 \mu\text{mol kg}^{-1}$, respectively), which can be interpreted as these layers being less affected by seasonal forcing due to their greater distance from the surface (Fig. 3). Despite their small magnitudes, the signals remain significant due to the small associated STD. The amplitude of the full section-average seasonal [C_{nat}] anomaly of $1.2 \mu\text{mol kg}^{-1}$ is largely determined by these low-amplitude signals of lMOC and bML, as these layers occupy a greater surface area.

As for [C_{nat}], the seasonality of [C_{ant}] is the highest in ML and the second highest in the uMOC (Fig. 3c,d). ~~However, the amplitude of the~~ The [C_{ant}] seasonality is an order of magnitude smaller for the ML, and about a factor of three smaller for the uMOC than the amplitude of the seasonal C_{nat} . The seasonal minimum in March–April for the upper layers in C_{ant} coincides with the seasonal maximum of C_{nat} at that time for the same layers. The C_{ant} values are minimal when [C_{nat}] is maximal and vice versa (Fig. 3,4).

~~A positive~~ A positive [C_{ant}] anomaly of $3.0 \pm 0.4 \mu\text{mol kg}^{-1}$ and a negative [C_{ant}] anomaly of $-2.5 \pm 0.2 \mu\text{mol kg}^{-1}$ are observed for the seasonal cycle of ML C_{ant} in August and April, respectively, for hence a $5.5 \mu\text{mol kg}^{-1}$ seasonal amplitude (Fig. 3). Following the same approach as for [C_{nat}], we note that the seasonal variation in thickness of the ML applied to an annual mean [C_{ant}] profile explains at most a third of the ML [C_{ant}] seasonal signal (Fig. 3d). The latter can be explained by the Revelle factor that creates seasonality in the surface [C_{ant}]. The Revelle factor decreases (increases) with increasing (decreasing) temperature, and the surface [C_{ant}] follows the seasonal cycle temperature temperature seasonal cycle with a

365 peak-to-peak amplitude of about $4 \mu\text{mol kg}^{-1}$ (Fig. S4), in phase with that caused by the ~~MLD-ML depth~~ seasonal cycle. In all regions, the cooling and winter deepening of ML decrease ML $[C_{ant}]$ due to the lower concentration at depth and larger Revelle factor (Fig. 4).

Looking at the sea surface, the surface C_{ant} is minimum in March ($54.7 \mu\text{mol kg}^{-1}$) and maximum in August ($59.5 \mu\text{mol kg}^{-1}$), which results in a seasonal amplitude of $4.8 \mu\text{mol kg}^{-1}$ (Fig. 4b). Higher values of C_{ant} are present in the NAC due to C_{ant} -laden water from subtropics, and the seasonal surface amplitude of the NAC is the highest ($4.9 \mu\text{mol kg}^{-1}$) (Fig. 4b). As for C_{nat} , the seasonal changes in ML C_{ant} mirror those of near-surface C_{ant} except in winter (Fig. 4b). However, the magnitude in the difference between surface and ML for C_{ant} is mostly constant for the rest of the year. We observe a discrepancy in winter (Fig. 4b). This winter difference is caused by a slight decrease (increase) in C_{ant} (C_{nat}) with depth in the winter ML. The largest seasonal amplitude in the ML for C_{ant} accounts for $8.3 \mu\text{mol kg}^{-1}$, and is observed in the middle of the IS, as for C_{nat} ($70.6 \mu\text{mol kg}^{-1}$) (Fig. 4a,b). The reader is referred to the point raised in the discussion section on the method for further information on the surface C_{ant} (4.5).

With a minimum in April ($-1.6 \pm 0.1 \mu\text{mol kg}^{-1}$) and a maximum in December ($+1.4 \pm 0.1 \mu\text{mol kg}^{-1}$), the peak-to-peak seasonal amplitude of uMOC $[C_{ant}]$ is $3.0 \mu\text{mol kg}^{-1}$ and approximately half that of ML $[C_{ant}]$ (Fig. 3c). We note that the variation in thickness of the uMOC applied to an annual mean $[C_{ant}]$ profile explains most of the seasonal $[C_{ant}]$ uMOC signal (Fig. 3c). Regionally, the deepening of the uMOC in winter in the NAC (see Mercier et al. (2024)) decreases the uMOC $[C_{ant}]$ (Fig. 4d). The reverse holds for summer, when the volume of uMOC decreases in the NAC. In the IS, a different mechanism prevails. The uMOC is confined to the surface layer and essentially belongs to the ML. The seasonal cycle of $[C_{ant}]$ is therefore similar to that of ML $[C_{ant}]$ and not the one we would expect, given that the thickness of the uMOC is less in winter than in summer (Mercier et al., 2024) (Fig. 4).

385 The bML and lMOC layers have reduced seasonal signals. The summer-to-winter difference ~~of the bML layer is is equal to~~ $1.9 \mu\text{mol kg}^{-1}$. The seasonal changes of C_{ant} in lMOC are equal to $0.8 \mu\text{mol kg}^{-1}$ ~~for the bML, lMOC, respectively~~. The seasonal amplitude of the full section averaged is negligible ($+0.2 \mu\text{mol kg}^{-1}$ between April and October).

3.1.2 Interannual to long-term

As evidenced by the time series in Figure 5a, the interannual variability in $[C_{nat}]$ is characterized by a 4–6 year periodic signal in the uMOC and, although a little less clear, in the ML (Fig. 5a). The peak-to-peak amplitude of the signal ranges from 8.5 to $11.2 \mu\text{mol kg}^{-1}$ in the uMOC and from 7.5 to $14.6 \mu\text{mol kg}^{-1}$ in the ML (Fig. 5a). Considering the STD of the mean ML and uMOC $[C_{nat}]$ values (\pm STD) for the whole period 1993–2022 (3.6 and $2.9 \mu\text{mol kg}^{-1}$, for a mean value of 2050.3 and $2105.9 \mu\text{mol kg}^{-1}$, respectively), along with the method uncertainty ($2.4 \mu\text{mol kg}^{-1}$ for the uMOC) (see Table 3), we conclude that the signal is statistically significant.

395 These quasiperiodic changes in $[C_{nat}]$ are ~~found to be~~ in phase with the changes in the average depth of the ML and the thickness of the uMOC (Fig. S2), ~~and not strongly correlated to the North Atlantic Oscillation (pearson coefficient of 0.32 for interannual uMOC $[C_{nat}]$ and 0.55 for interannual ML $[C_{nat}]$)~~. We have plotted in Fig. 5c,e the interannual variability of $[C_{nat}]$ due solely to variations in the depth of ML and the thickness of the uMOC, obtained using $[C_{nat}]$ averaged over the total




Figure 3. (a,b) Natural carbon $[C_{nat}]$ and (c,d) anthropogenic carbon $[C_{ant}]$ seasonal anomalies. Left (a,c) and right (b,d) panels show respectively uMOC/IMOC and ML/bML (Fig. 1, Table 2). Black lines represent the full section-average (referred to as FSA in the legend), light colors are used for the layers the closest to the sea surface (uMOC or ML), and dark colors account for the deeper layers (IMOC or bML). Each monthly value represents the mean value calculated from the ensemble of ocean reanalyses, and the shaded areas the standard errors. The dashed lines represent the idealized seasonal anomaly computed using annual mean property fields and only taking into account changes in area surfaces of layers (see 2.7 for details). The difference between the seasonal cycle of $[C_{nat}]$ in the ML and its idealized anomaly, attributed to biological activity, is shown in purple (b).




Figure 4. Seasonal cycles of surface (5-m depth) (a) natural carbon ML $[C_{nat}]$ (solid black lines) and (b) anthropogenic carbon ML $[C_{ant}]$ (black) for the full section (continuous dark color line), the middle-center of the Irminger Sea (between -42°E and -32°E , cross symbols line with x markers) and the NAC (between -27°E and -17°E , plus symbols). Seasonal ML (a) C_{nat} and (b) C_{ant} are displayed for the full section (continuous color line with + markers), the middle-, Seasonality of Irminger Sea (cross symbols colored line) and the NAC (plus symbols colored line). Seasonal uMOC (c) uMOC $[C_{nat}]$ and (d) uMOC $[C_{ant}]$ are also shown for the full section (continuous color line), the Irminger Sea (cross symbols colored line with x markers), the Iceland Basin-NAC region (IB-NAC) (left triangle symbols colored line with ◀ markers) and the East of the NAC (right triangle symbols colored line with ▶ markers). Here, The seasonal cycles are obtained by grouping the time series signal data by month and averaging over all ocean reanalyses. The time series are subsets of the complete series of Fig. 2, sampled from 2004 to 2017, so that each reanalysis has the same weight in order to have all reanalyses defined in the period construction of the seasonality (Table 1). Values represent the average of all ocean reanalyses.

duration of the time series. Variability due to changes in uMOC thickness and ML depth explain most of the interannual variability in $[C_{nat}]$, which confirms the previous visual inspection. This is further illustrated in Figs. S3 and S5 from GLOSEA5, which show that a maximum in the surface section area is associated with a maximum in $[C_{nat}]$. In the ML, the interannual signal in the section area is dominated by the varying maximum depth of the winter ML (Fig. S2), that is, the interannual variability in seasonal processes (such as the winter ML mixing) causes interannual concentrations to vary. For example, the deep ML observed in the IS in winter 2015 and 2016 appears to have had the most significant impact on interannual variability of ML $[C_{nat}]$ (Fig. S3). The interannual changes in the area of the uMOC layer are the greatest in the West European Basin (Fig. S5), where the interannual changes in σ_{MOC} density results in uMOC thickness variations (Mercier et al., 2024). The interannual, impacting interannual uMOC $[C_{nat}]$ variability in the uMOC follows most of these uMOC thickness changes over 1993–2022 (shown for GLOSEA5 in Fig. S5). Regarding long-term changes, the uMOC $[C_{nat}]$ does not show any significant tendency. In contrast, the ML shows a significant increase in its variability from 2008 onward. It is concomitant with the intermittent resumption of deep convection in the NASP and documented events in 2008, 2012 and 2015 (Piron et al., 2016, 2017) associated with maxima of $[C_{nat}]$ creating an apparent $[C_{nat}]$ (linear) increase of $0.22 \pm 0.01 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ over the period (Fig. 5a). IMOC and bML $[C_{nat}]$ did not show interannual variability and no trends. These layers have the largest $[C_{nat}]$ mean values ($2153.4 \pm 0.5 \mu\text{mol kg}^{-1}$ and $2146.5 \pm 0.6 \mu\text{mol kg}^{-1}$ for IMOC and bML, respectively) and the largest surface section areas. They contribute the most to the section average mean $[C_{nat}]$, which is mostly constant over 1993–2022, showing which shows no interannual variability and no trend, with a mean value (\pm STD) of $2143.5 \pm 0.8 \mu\text{mol kg}^{-1}$ (Fig. 5).

Unlike $[C_{nat}]$, $[C_{ant}]$ is dominated by the long-term signal across all layers considered (Fig. 5). The ML shows the highest mean $[C_{ant}]$, and the highest rate of $[C_{ant}]$ increase of $0.825 \pm 0.016 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ (Fig. 5b). It is followed by the uMOC, with a $[C_{ant}]$ increase rate of $0.625 \pm 0.027 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ (increase in $[C_{ant}]$ uMOC of $45.6 \pm 2.0 \%$ from 38.4 to $55.9 \mu\text{mol kg}^{-1}$ between 1993 and 2021). However, IMOC, not highly concentrated in $[C_{ant}]$, experiences the same relative increase in $[C_{ant}]$ of $40.2 \pm 0.7 \%$ (which increased from 17.0 to $23.9 \mu\text{mol kg}^{-1}$ between 1993 and 2021 at a linear rate of $0.245 \pm 0.005 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$) as bML and section average. In terms of the section average (net) $[C_{ant}]$, the increase rate is $0.301 \pm 0.004 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$, with a mean $[C_{ant}]$ value varying from $21.7 \mu\text{mol kg}^{-1}$ in 1994 to $30.4 \mu\text{mol kg}^{-1}$ in 2021. For $[C_{ant}]$, the interannual variability is mainly observed in ML and uMOC, similar to $[C_{nat}]$, although in opposite phases due to inverse vertical concentration gradients between C_{ant} and C_{nat} . The amplitude of the interannual signal ranges from 0.7 to $3.4 \mu\text{mol kg}^{-1}$ for the ML, and from 2.6 to $3.9 \mu\text{mol kg}^{-1}$ for the uMOC. These values lie above the method's error ($1.2 \mu\text{mol kg}^{-1}$ for uMOC $[C_{ant}]$, Table 3).

3.2 C_{ant} and C_{nat} transports

The time series of volume, C_{nat} and C_{ant} transports across A25 for each of the ocean reanalysis are presented in Figure 6, with independent cruise-based estimates shown in red.




Figure 5. Low pass filtered signal (continuous lines) with their linear trends (dashed lines, plotted when significant at the 90 percent significance level) of (a) $[C_{nat}]$ and (b) $[C_{ant}]$ in the layers of interest uMOC, IMOC, ML and bML based on time series reported in Fig. 2. Full section average (FSA) concentration is in black. For the uMOC (c,d) and (e,f) ML, the signal is plotted in an anomaly along with the corresponding signal for a mean concentration $[C]_{MEAN}$. For $[C_{ant}]$, the linear trends have been removed in the anomaly plots. Grey shading indicates when only GLOSEA5 reanalysis is available. STDns (\pm) are shown in grey shading. [Linear fit \(\$\pm\$ confidence interval at 90%\) on interannual \$\[C_{nat}\]\$: \$0.22 \pm 0.04 \mu\text{mol kg}^{-1} \text{yr}^{-1}\$ \(ML\).](#) [Linear fit values on interannual \$\[C_{ant}\]\$: \$0.8 \pm 0.008 \mu\text{mol kg}^{-1} \text{yr}^{-1}\$ \(ML\), \$0.3 \pm 0.002 \mu\text{mol kg}^{-1} \text{yr}^{-1}\$ \(bML\), \$0.6 \pm 0.012 \mu\text{mol kg}^{-1} \text{yr}^{-1}\$ \(uMOC\), \$0.3 \pm 0.002 \mu\text{mol kg}^{-1} \text{yr}^{-1}\$ \(IMOC\) and \$0.3 \pm 0.001 \mu\text{mol kg}^{-1} \text{yr}^{-1}\$ \(FSA\).](#)




Figure 6. (a) Volume, (b) C_{nat} and (c) C_{ant} transports. Each panel (a,b,c) shows the net transport value across A25 (average of ECCO and GLOSEA5), the uMOC transport (average of ECCO, GLOSEA5, EN4, CORA transports) and the IMOC transport (average of ECCO and GLOSEA5). The standard errors (STDns) for the mean values computed as the STDs divided by the square root of the number of reanalysis are shown as grey shading. The cruise estimates with their uncertainties indicated in red are from Danialt et al. (2016); Lherminier et al. (2007, 2010); Zunino et al. (2014, 2015, 2017); Mercier et al. (2024). Vertical grey shading indicates when only the GLOSEA5 reanalysis is available.

3.2.1 Seasonal

Maximum seasonal northward C_{nat} transport anomalies of 2.66 and 2.63 PgC yr^{-1} are observed in March within uMOC and ML, respectively (Fig. 7c,d). The seasonal anomaly of uMOC C_{nat} transport ~~becomes null in June and~~ reaches a minimum of $-1.80 \text{ PgC yr}^{-1}$ in September. ~~For the ML, the C_{nat} transport anomaly becomes null mid-April.~~ Between June and November, 435 the ML C_{nat} transport anomaly is negative with a slight minimum of $-1.19 \text{ PgC yr}^{-1}$ in June. ~~The~~ uMOC* C_{nat} transport seasonality is dampened compared to ~~that of the~~ uMOC while ML* C_{nat} transport seasonality is strengthened compared to that of ML. The seasonality of the transport of IMOC (bML) C_{nat} is the opposite to that of uMOC* (ML*) C_{nat} transport seasonality. ~~The~~ C_{nat} transport in the IMOC reaches a ~~maximum of~~ $-1.24 \text{ PgC yr}^{-1}$ maximum to the South in March and a ~~maximum seasonal anomaly to the North of~~ 1.15 PgC yr^{-1} maximum to the North in September and November (Fig. 7c) ~~while~~ 440 ~~the~~ C_{nat} transport in the bML reaches $-4.06 \text{ PgC yr}^{-1}$ to the South in March and a maximum seasonal anomaly to the North of 2.11 PgC yr^{-1} in September (Fig. 7d). The seasonal anomaly of ~~the~~ net C_{nat} transport is minimum in May ($-0.28 \text{ PgC yr}^{-1}$) and maximum in September (0.34 PgC yr^{-1}) (Fig. 7c,d). Its amplitude is an order of magnitude lower than the seasonal anomaly of the ~~other~~ uMOC, IMOC, ML and bML layers.

The seasonality of ~~the~~ C_{ant} transport is mainly in phase with the seasonality of C_{nat} transport (Fig. 7,c-e,d-f). In the uMOC, 445 there is a corresponding northward maximum of 0.06 PgC yr^{-1} in March and a reduced southward transport in September of $-0.04 \text{ PgC yr}^{-1}$ (Fig. 7e). The seasonality of uMOC* C_{ant} transport is dampened compared to that of the uMOC. The seasonality of the transport of ML C_{ant} is slightly higher than that of uMOC in the late winter, with 0.08 PgC yr^{-1} to the ~~north~~ North in March (Fig. 7f). There is a rapid drop in April in the transport of ML C_{ant} and a negative anomaly of $-0.04 \text{ PgC yr}^{-1}$ in July, corresponding to half of the maximum ML in late winter (Fig. 7f). The seasonality of ML* C_{ant} transport is 450 slightly larger than that of ML C_{ant} transport. The transport of bML C_{ant} is equal to $-0.09 \text{ PgC yr}^{-1}$ southward in March and shows a plateau between June and November of almost 0.05 PgC yr^{-1} (Fig. 7e). The net C_{ant} transport seasonality ~~shows two~~ ~~maximums in March and September, which corresponds to the seasonal maximum in the uMOC and the northward anomaly in the IMOC, respectively~~ is very small. Seasonal net C_{ant} transport is influenced by both the seasonal variation of net volume transport and full section averaged concentration (Fig. 7a,b and 3c,d).

455 The seasonal cycle of the C_{ant} and C_{nat} transports closely reflects the seasonality of the volume transport, as evidenced by the similarity between the seasonal cycles of the volume transport and the property transport (Fig. 7). The strong influence of ocean circulation on the seasonal transport of C_{ant} and C_{nat} is also supported by the fact that, for a given layer (ML or MOC branches), although $[C_{nat}]$ and $[C_{ant}]$ seasonal anomalies are opposite, C_{nat} transport and C_{ant} transport seasonal signals are synchronous (Fig. 7). The seasonal variations of volume transport of 4.88 and 2.95 Sv for uMOC and IMOC, respectively, 460 represent 26.1 % and 17.3 % of their annual mean values corresponding to seasonal changes of 4.46 and 2.40 PgC yr^{-1} for C_{nat} , and 0.10 and 0.04 PgC yr^{-1} for C_{ant} for uMOC and IMOC, respectively (Fig. 7). The seasonal amplitudes for the uMOC and the IMOC for the transport of C_{nat} (C_{ant}) represent 29.3 % and 17 % (29.8 % and 17.9 %) of their annual mean value (Fig. 6a,c and 7a,e). The sole observed effect of the seasonality of $[C_{ant}]$ is on the seasonality of the net C_{ant} transport.




Figure 7. Volume (a,b), C_{nat} (c,d), and C_{ant} (e,f) seasonal transport anomalies. Left (right) panels show uMOC/IMOC (ML/bML). Light colors represent the upper-ocean layers (either uMOC or ML), dark colors the deeper-ocean layers (either IMOC or bML), and the black lines the net transport (section integration). The seasonal anomalies presented here are computed using the average of all reanalyses available. uMOC* and ML* represent the average of only GLOSEA5 and ECCO, to align with the IMOC and bML estimates, which are based solely on these two datasets (Table 2). The shaded areas is the standard error for the mean (STDn) computed as the standard deviation for each month, divided by the square root of the number of reanalysis.




Figure 8. Low-pass filtered (a) volume, (b) C_{nat} and (c) C_{ant} transports time series for the uMOC (upper panels), IMOC (middle panels) and net section (bottom panels). These interannual time series were obtained by subtracting the high frequency time series obtained using a one-year high-pass filter to the original series, as reported in Figure 6 (see [Methods-2.7](#)). The dashed straight lines correspond to the linear trends on the low-pass filtered signals. Vertical light grey shading indicates where only GLOSEA5 reanalysis is available. The standard errors (STDns) are shown in dark grey shading for the mean values.

3.2.2 Interannual to long-term

465 The net volume transport along the section is close to 1 Sv for the 30-year time series, the uMOC and IMOC transports having opposite signs (Fig. 6). These opposite signs will be reflected in the signs of uMOC and IMOC C_{nat} (C_{ant}) transports (Figs. 6, 8).

The net C_{nat} transport averages to 0.40 ± 0.44 PgC yr⁻¹ northward over 1993–2021 (Fig. 8b). This transport did not show a tendency before 2010 and increased from 0.13 ± 0.15 to 1.04 ± 0.09 PgC yr⁻¹ between 2010 and 2021 (Fig. 8b). The STDn of the net C_{nat} transport increases over time, from 0.18 PgC yr⁻¹ for 1993–2010 to 0.33 PgC yr⁻¹ for 2011–2018. Between 1993 and 2021, the C_{nat} transports for the uMOC and lMOC average to 15.3 ± 1.1 and -14.1 ± 1.1 PgC yr⁻¹, respectively. There is noticeable variability in C_{nat} uMOC transport with a longer-term weakening period before 2010 and growth after this year. In 2021, uMOC C_{nat} transport reaches 17.78 ± 0.63 PgC yr⁻¹ (Fig. 8b). The C_{nat} uMOC and lMOC transports show the same variability as the uMOC and lMOC volume transport (Fig. 8a,b). The reduction of 1.15 (1.94) Sv in the volume transport of uMOC (lMOC) between the 1993–1997 pentad and the 2008–2012 pentad results in a decrease of 1.17 (1.58) PgC yr⁻¹ in the C_{nat} uMOC (lMOC) transport during the same period (Fig. 8a,b). This result indicates that the interannual variations of $[C_{nat}]$ are negligible compared to volume changes (Fig. 3b, 8a). Pearson coefficient correlation of 0.98 (0.99) is found between interannual volume and C_{nat} transport in the uMOC (lMOC), highlighting that changes in volume transport outweigh changes in concentration. The interannual variability of uMOC C_{nat} transport amounts to 0.86 PgC yr⁻¹ (uMOC STD in Fig. 8b). It is an order of ~~amplitude~~ magnitude smaller than the seasonal amplitude of transport of uMOC C_{nat} (4.46 PgC yr⁻¹). The dispersion between ocean reanalysis C_{nat} transport is measured by STDns. The uMOC STDns average at 1.06 PgC yr⁻¹ for 1993–2021. A larger dispersion in the reanalyses is observed within the C_{nat} lMOC transport after 2010, the STDns increasing from 1.15 PgC yr⁻¹ for 1993–2010 to 1.36 PgC yr⁻¹ for 2011–2018. Breaking down the net interannual C_{nat} transport time series following Eq. 3, we note that most of the variability is explained by the net' C_{nat} transport i.e. the product at each time step of the net volume transport by the average property. This is confirmed in Fig. 9 where 99–100 % of the C_{nat} transport variance (r^2) is explained by net' C_{nat} transport and thus volume transport variability for ECCO and GLOSEA5, respectively. The variability of C_{nat} transport in the SPNA at A25 is shown to be driven by volume transport both in the uMOC, lMOC and net sections at the interannual and long-term time scales (Fig. 6–8).

The net C_{ant} transport doubled in thirty years, from 0.07 in 1993 to 0.14 PgC yr⁻¹ in 2021 northward corresponding to an increase of 0.023 PgC per decade. Its evolution is closely approximated by a linear increase (Fig. 8c). C_{ant} transport in the uMOC increased from 0.29 ± 0.03 PgC yr⁻¹ to 0.45 ± 0.02 PgC yr⁻¹ between 1993 and 2021 (0.056 PgC per decade). ~~An increase in~~ In contrast, the lMOC C_{ant} southward transport shows no trend until 2008, after which it increases. This results in an overall increase between 1993 and 2021 from -0.19 ± 0.03 PgC yr⁻¹ to -0.32 ± 0.04 PgC yr⁻¹ ~~between 1993 and 2021~~ (0.047 PgC per decade, Fig. 8c) ~~is also observed~~. The southward transport of lMOC compensates in part for the northward transport of uMOC. The relative increase in $[C_{ant}]$ in uMOC is close in percentage (45.6 ± 2.0 %, 48.4 ± 2.6 % for uMOC*) to that in lMOC (37.4 ± 0.7 %) (Fig. 5). However, the net northward $[C_{ant}]$ transport increases due to higher $[C_{ant}]$ values in the upper layers and an increase in the difference in $[C_{ant}]$ between uMOC and lMOC from 1994 to 2021, uMOC $[C_{ant}]$ being 19% larger than lMOC $[C_{ant}]$ in 1993 but more than twice in 2021 (Fig. 5). The section averaged $[C_{ant}]$ shows an increase of $9.1 \mu\text{mol kg}^{-1}$ (42.5 ± 0.8 %) between 1993 ($21.4 \mu\text{mol kg}^{-1}$) and 2021 ($30.5 \mu\text{mol kg}^{-1}$) (Fig. 5b). For the net C_{ant} transport, there is a first period of small increase before 2010, which is resulting from an increase in $[C_{ant}]$ compensated for by a decrease in the water mass transport over the period (Fig. 5a and 8c) for both branches. After 2008, the increase is more rapid, with a rate of 0.038 PgC per decade compared to 0.013 PgC per decade previously, as both volume transport and concentration are intensified.

The interannual C_{ant} transport for uMOC and IMOC follows the interannual transport of volume (Fig. 8). Pearson correlation coefficients of 0.91 and 0.93 were obtained between the two interannual transports for uMOC and IMOC, respectively.

505 The large correlations between the diapycnal component (Eq. 3) and the net C_{ant} transport (r^2 values of 0.71, 0.81 for ECCO, GLOSEA5) (Fig. 9c,d) show that the diapycnal component T_{diap} drives the net C_{ant} transport. The strong positive T_{diap} for C_{ant} comes from the higher concentration of C_{ant} in uMOC than in IMOC (Fig. 5b). Separating the two components of the diapycnal transport (velocity and $[C_{ant}]$) into a time-mean term (overline) and a fluctuation term (prime) in Fig. 10 shows that the interannual changes in diapycnal transport come from the changes in volume and velocity multiplied by the average
510 concentration of $[C_{ant}]$, for both GLOSEA5 and ECCO (Fig. 10). ~~93% and 81% of the variability of the interannual diapycnal component are explained by the estimator T_{est} in GLOSEA5 and ECCO, respectively, as shown by the Pearson correlation coefficient.~~




Figure 9. (a,b) Natural carbon (C_{nat}) transport as a function of its net' term along with the (c,d) anthropogenic carbon (C_{ant}) transport as a function of its diapycnal term, for (a,c) GLOSEA5 and (b,d) ECCO. The data points are colored according to their date. The linear function with the r^2 coefficient is shown on the figures.




Figure 10. (a,b) Low-pass filtered signal anomaly of the diapycnal component of the net anthropogenic carbon (C_{ant}) transport along with its decomposition into a mean over time profile ($\langle c \rangle_t(\sigma)$, $\langle v \rangle_t(\sigma)$) and a perturbation ($c(\sigma, t)'$, $v(\sigma, t)'$) for the velocity and concentration terms in Eq. 3, for (a) GLOSEA5 and (b) ECCO.

4 Discussion

4.1 Time series evaluation

515 The $[C_{ant}]$ derived from ocean reanalysis temperature and salinity, using NNs and the BC φC_T^0 approach, collectively referred to here as the OR-NN-BC method, shows good agreement with the 2002–2018 cruise-based estimates presented here using the same BC approach (RMSD of $5.1 \mu\text{mol kg}^{-1}$ Table S1 and Fig. 2b), and with previous estimates at A25 reported by Zunino et al. (2014). The [agreement between reanalysis products and bottle data was evaluated and summarized in Tables 4, S3, S5, and discussed in Sections 2.8.2 and 2.8.3.](#) The $[C_{ant}]$ increase rates obtained from our $[C_{ant}]$ time series are also in good
520 agreement with the cruise-based rates. Here, we found $[C_{ant}]$ increase rates of 0.7 ± 0.03 , 0.2 ± 0.006 and $0.3 \pm 0.004 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ for the uMOC, IMOC and the section average, respectively, during the entire period (1993–2022) (Fig. 2b). These results closely match the $[C_{ant}]$ increase rate estimates by Zunino et al. (2014) of 0.6 ± 0.3 , 0.2 ± 0.3 and $0.3 \pm 0.2 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ for the same layers, based on A25 cruise data spanning 1997–2010. The agreement between both estimates, considering the two distinct periods used for the calculations, highlights the linear nature of the $[C_{ant}]$ increase over time for all the layers
525 under consideration. Looking at specific longitudinal regions of the A25 section, the $[C_{ant}]$ increase rate in the IS ($0.36 \pm 0.005 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$) is slightly higher than the increase rate for the section average (Fig. S3ij), likely due to the transfer of $[C_{ant}]$ to intermediate depths in the IS by deep convection (Pérez et al., 2018; Asselot et al., 2024). This increase rate is comparable to that observed in the North East Atlantic Deep Water in the Labrador Sea ($0.3 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ for 1986–2016), but substantially lower than the $0.8 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ rate found in Labrador Sea Water over the same period (Raimondi et al.,
530 2021). This difference reflects the enhanced $[C_{ant}]$ storage capacity of Labrador Sea compared to that of the IS, which results from its direct exposure to frequent deep convection events that efficiently transport $[C_{ant}]$ from the surface to depths up to 2000 m (Raimondi et al., 2021; Yashayaev, 2024).

A comparison of C_{ant} transports with previous studies at A25 also reveals good consistency of the results. The cruise-based estimate of $0.092 \pm 0.010 \text{ PgC yr}^{-1}$ by Pérez et al. (2013) for June 2004 is similar to our estimate of $0.09 \pm 0.02 \text{ PgC yr}^{-1}$ for
535 the same period (Fig. 7c). The 2002–2010 C_{ant} transport of $0.096 \pm 0.011 \text{ PgC yr}^{-1}$ ($254 \pm 29 \text{ kmol s}^{-1}$) reported by Zunino et al. (2014) is comparable to the $0.098 \pm 0.02 \text{ PgC yr}^{-1}$ that we find for the same period. Our $[C_{nat}]$ estimates are also in good agreement with the 2002–2018 cruise-based $[C_{nat}]$ values (Table S3), showing no long-term trend. The C_{nat} transport estimates derived from ocean reanalysis (net, uMOC and IMOC) are generally consistent with the A25 cruise-based estimates presented in this study, lying within their uncertainty range (Fig. 6, Table S7), except in 2006, when the discrepancy between
540 the two values exceeded the uncertainty (Fig. 6a,c). Taking into account what happens on an intraannual scale, the seasonality of ML $[C_{nat}]$ and the seasonality of surface $[C_{nat}]$ and [DIC] (Fig. 4a, S6) are consistent with seasonal amplitudes in the range of $42\text{--}60 \mu\text{mol kg}^{-1}$ observed for surface [DIC] at high latitudes (Keppler et al., 2020; Hagens and Middelburg, 2016) and $\approx 60 \mu\text{mol kg}^{-1}$ for surface [DIC] in the SPNA (Leseurre et al., 2020; Reverdin et al., 2018). Finally, we note the lack of previous research on the seasonality of C_{ant} and C_{nat} transports in the SPNA.

545 This study corroborates previous estimates (Zunino et al., 2015; Pérez et al., 2013), which relied on cruise observations, and provides a more detailed perspective on C_{ant} and C_{nat} concentration and transport variability at seasonal to interannual time scales.

4.2 Mechanisms involved in seasonality

The deepening of the ML in winter favors the enrichment of [C_{nat}] within the ML through the entrainment of DIC-rich
550 thermocline waters (Keppler et al., 2020; Takahashi et al., 1993), increasing the seawater partial pressure of CO_2 ($p\text{CO}_2$)
and thus leading to surface saturation or slight supersaturation. In this study, we quantify that this physical mechanism of the
mixed-layer pump accounts for approximately two-thirds of the winter increase in ML [C_{nat}] (Fig. 3). ~~The remaining one-third
can reasonably be attributed to the combined effects of solubility and lateral advection/mixing, as biological activity is minimal
during this period (Henson et al., 2006, 2013).~~ ML [C_{nat}] decreases in summer. During this period, changes in the depth of ML
555 account for only one-third of the ML [C_{nat}] variability, with the remaining two-thirds largely favored by extensive biological
carbon consumption (Keppler et al., 2020) resulting in surface undersaturation (Olsen et al., 2008; Lacour et al., 2015; Tjiputra
et al., 2012).

~~As inferred above, changes in DIC concentration in the ML are one of the main drivers of the $p\text{CO}_2$ seasonal variability
in the SPNA (Tjiputra et al., 2012; Landschützer et al., 2018; Rodgers et al., 2023; Pérez et al., 2024), and hence of the air-sea
560 CO_2 flux.~~

It is important to note that the OR-NN-BC method relies solely on T, S, date, and position as predictor variables to obtain
[DIC] from which the [C_{nat}] and [C_{ant}] components are derived (see [Methods 2.4](#)). Hence, by methodology, we expect that
the surface [C_{nat}] and [C_{ant}] correlate or anti-correlate to a certain extent with temperature (Fig. S7,S8). However, this does
not imply that the temperature itself (i.e. solubility) is the sole or dominant driving mechanism of the observed seasonal
565 surface [C_{nat}] and [C_{ant}] variability. For [C_{nat}], the seasonal cycle, which anti-correlates with temperature (Fig. S7a,b), still
remains when considering no seasonal temperature signal at all (Fig. S7c,d). For [C_{ant}], however, there is a positive correlation
with temperature (Fig. S8), and when the seasonal cycle of temperature is removed (Fig. S8c,d), the [C_{ant}] seasonal signal
largely disappears. This ~~does not imply that C_{ant} seasonal variability is driven by temperature-dependent solubility. Rather, it
effect~~ reflects seasonal changes in the ocean's CO_2 uptake capacity rather than temperature-driven solubility control: summer
570 conditions show minimum Revelle factor values (Fig. S4), which indicates enhanced seawater buffer capacity, thereby favoring
[C_{ant}] uptake. Together, summer changes in both [C_{nat}] and [C_{ant}] components are indicative of an enhanced uptake of
atmospheric CO_2 in the SPNA (Rodgers et al., 2023; Rustogi et al., 2023).

~~Seasonal variability in volume transport dominates the seasonal variability of uMOC and IMOC C_{nat} and C_{ant} transports
at A25. The seasonal variability of the volume transport of the uMOC and the IMOC can be attributed both to the volume
575 variations of the uMOC in the East Greenland Current and to the~~ At the intraannual scale, C_{nat} and C_{ant} transports are
dominated by seasonal variations in volume transport, driven by uMOC volume changes in the EGC and velocity variations in
the eastern boundary current (Mercier et al., 2024). At A25, this combined effect of uMOC thickness variations and velocity
changes on seasonal tracer transport variability differs from C_{ant} transport variability at 26.5°N in the subtropical gyre, which

was attributed primarily to velocity variability (Brown et al., 2021) (Fig. S9). Less seasonal variation in C_{ant} transport is
580 observed in the SPNA at the A25 section (amplitude of $0.015 \text{ PgC yr}^{-1}$) (Fig. 7e–f) than at 26.5°N (peak-to-trough amplitude
of 0.08 PgC yr^{-1}) (Brown et al., 2021). However, the large seasonal variations for uMOC and IMOC represent more than
25% of the annual average C_{ant} for volume and tracer transports at A25. Likewise, at 26.5°N between 2004 and 2012, the seasonal
variability in the transport of uMOC C_{ant} constituted 27% of its mean value (Brown et al., 2021).

4.3 Interannual to long-term mechanisms

585 The long-term $[C_{ant}]$ increase at A25 is linear (Fig. 8). This rise is primarily driven by the atmospheric increase in CO_2 and the
increase in air-sea CO_2 fluxes in the NA (Gruber et al., 2023), as suggested by the trend-less C_{ant} transport at 26.5°N between
2004 and 2012 (Brown et al., 2021). The decadal rate of increase in $[C_{ant}]$ in the water column depends on the distance of
the layer from the sea surface, with higher increase rates closer to the surface where atmospheric $[C_{ant}]$ enters the ocean via
air-sea exchange (Fig. 5). For example, a rate of increase of $6.0 \mu\text{mol kg}^{-1}$ per decade was found for the uMOC compared
590 to $2.4 \mu\text{mol kg}^{-1}$ per decade for the IMOC. However, the relative increase for each layer is the same. The atmospheric CO_2
growth was about 61.35 ppm for 1993–2022 (1993: 357.21 ppm , 2022: 418.56 ppm) (Lan et al., 2023). This rise resulted in an
almost double-doubling of $[C_{ant}]$ in ML (~~49.8~~ 49.8 %). $[C_{ant}]$ variability is dominated by long-term changes in concentration,
due to atmospheric CO_2 forcing, while $[C_{nat}]$ variability is dominated by seasonal and interannual changes.

The interannual variability in ML $[C_{nat}]$ and $[C_{ant}]$ is the strongest when particularly deep ML are found during deep
595 convection periods, before 1995 (for $[C_{nat}]$) and between 2008–2016 (for both) (Fig. 5,S3). The interannual variability of
seasonal deep convection events shapes the interannual $[C_{nat}]$ and $[C_{ant}]$ in the ML. The signal is the largest in the IS and in
agreement with previously reported events (Piron et al., 2016, 2017). In turn, the alternation of shallow and deep ML and the
persistence of the signal at depth cause interannual variability in concentrations.

At 24.5°N , the rates of increase in $[C_{ant}]$ between 0.25 and $0.88 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ were observed between 1992 and 2011
600 for deep waters and surface layers, respectively (Guallart et al., 2015). This is consistent with our increase rate for 1993–
2011 of $0.2 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ for deep waters and more than $0.8 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ in ML. Motivated by the study of Müller
et al. (2023) on GLODAP data spanning over 1994 (1989–1999), 2004 (2000–2009) and 2014 (2010–2020) decades, we com-
puted the accumulation rates obtained with our method for these three decades at A25 (the first decade for us is limited and
starts in 1993). Despite using a different methodology, we observe the reduction in $[C_{ant}]$ accumulation rate in the North
605 Atlantic ~~Steinfeldt et al. (2024)~~ (Steinfeldt et al., 2024) at A25, with a lower $[C_{ant}]$ increase rate above 3000 m in the 2014
decade ($3.4 \mu\text{mol kg}^{-1} \text{ dec}^{-1}$) than in the 1994 and 2004 decades ($4.0 \mu\text{mol kg}^{-1} \text{ dec}^{-1}$) (Fig. 11). This reduction is con-
comitant with the observation of deeper ML at the end of the 2011–2020 period (Fig. S2). The maximum $[C_{ant}]$ reduction is
located on the Reykjanes Ridge (Fig. 11b), suggesting that the NAC brought less $[C_{ant}]$ during the 2014 decade. The early
1994 decade and the 2014 decade correspond to periods when the subpolar gyre ~~is was~~ is was more intense and ~~has less influence~~
610 ~~from the waters of the NAC (Häkkinen and Rhines, 2004; Marzocchi et al., 2015; Zunino et al., 2020; Holliday et al., 2020)~~
less affected by NAC waters (Häkkinen and Rhines, 2004; Marzocchi et al., 2015; Zunino et al., 2020; Holliday et al., 2020).




Figure 11. (a) C_{ant} increase rate for the 0–3000 m depth range according to the estimates calculated from GLOSEA5 reanalysis. The decades are centered like in Müller et al. (2023). (b) $\Delta\Delta C_{ant}$: difference in increase rate between 2014–2004 and 2004–1994 at A25 section.

In addition, deeper ML (as in the 2014 decade) would lead to a minimum in $[C_{ant}]$, even if the $[C_{ant}]$ from the NAC was higher or unchanged.

Looking at volume transport, an interannual decrease is observed in the net and MOC transports until the early 2010s as in Jackson et al. (2022) (Fig. 8a) and is consistent with the weakest state of the AMOC in recent decades (Caesar et al., 2018, 2021; Boers, 2021; Jackson et al., 2022; Mercier et al., 2024) before increasing since then (Fig. 8a).

For C_{ant} transport, the decadal time scale that shows a linear trend is not driven by circulation but by the linear growth in $[C_{ant}]$, compared to C_{nat} , a result that refines previous results, obtained over a shorter time frame in the subtropical and subpolar gyres, which concluded that the variability of C_{ant} transport is driven by circulation (Brown et al., 2021; Pérez et al., 2013). Most of the net C_{ant} transport variability comes from its diapycnal component, which is set by $[C_{ant}]$ growth (Fig. 9, 10). There is a significant imbalance between the C_{ant} transports of the uMOC and the IMOC due to the $[C_{ant}]$ strong vertical gradient so that the uMOC is the main driver of the North Atlantic northward transport of C_{ant} (Brown et al., 2021; Pérez et al., 2013). The increase in $[C_{ant}]$ results in almost a doubling of the northward C_{ant} transport from 0.08 PgC yr^{-1} in 1992 to 0.15 PgC yr^{-1} in 2022. Over the shorter pre–2008 period, C_{ant} transport exhibited a slower increase rate ($0.013 \text{ PgC per decade}$ compared to $0.038 \text{ PgC per decade}$ after 2008), attributed to weakened MOC strength (uMOC decreased by $1.8 \text{ Sv per decade}$ before 2008) that attenuated the increase in $[C_{ant}]$.

The variability of the C_{nat} net transport is driven by the variability of volume transport from intraannual to long-term scales, in agreement with Zunino et al. (2014, 2015), and that $[C_{nat}]$ is mainly constant throughout the time range of this study. Relative changes in mean $[C_{nat}]$ are small (Fig. 2), so long-term C_{nat} transport changes are mainly determined by

630 volume transport changes. Since C_{nat} transport strongly correlates with net volume transport in the subpolar gyre, having the latter well constrained and solved is crucial. The interannual net C_{ant} transport variability is also circulation-driven because relative changes in $[C_{ant}]$ are not significant in successive years contrary to the relative changes in volume transport. The net volume transport of GLOSEA5 increases while ECCO one stays constant, which is reflected as well on net C_{nat} transports. The transport of uMOC C_{nat} also shows the periods of reduction and strengthening before and after 2008–2012, and is based
635 on the diversity in the physics of ocean reanalysis. Due to the strong volume transport-driven variability of net C_{nat} transport, the spread of circulation in ECCO and GLOSEA5 has a greater influence on net C_{nat} decadal transport than on decadal net C_{ant} decadal transport, being more influenced by $[C_{ant}]$ ([see Section 2.8.3 and Tables 4.S5 for a comparison of ocean analysis products](#)).

4.4 NA C_{ant} budget implications

640 The C_{ant} budget of an oceanic region is the result of the balance between lateral advection, air-sea fluxes, and storage. Net advective transport plays an important role in the NA budget, contributing to $65 \pm 13\%$ of the NA C_{ant} storage rate (reference to 2004, Pérez et al. (2013)). The observed increase in net northward C_{ant} transport over the 30-year period is evident in both subtropical and subpolar regions of the NA, as documented through decadal GO-SHIP section repeats (Caínzos et al., 2022). At [36°NA03](#), C_{ant} transport experienced an increase from 0.058 ± 0.036 to 0.104 ± 0.035 PgC yr^{-1} between 2000–2009
645 and 2010–2019, while at 26.5°N C_{ant} transport passed from 0.128 ± 0.032 to 0.222 ± 0.024 PgC yr^{-1} (Caínzos et al., 2022) (Fig. S9). During the same time periods (2000–2009 and 2010–2019), we found at A25 an increase in northward C_{ant} transport from 0.09 ± 0.02 PgC yr^{-1} to 0.11 ± 0.03 PgC yr^{-1} . At AR07, north of A25, similar values as at A25 are found: 0.088 ± 0.038 for 2000–2009 and 0.115 ± 0.042 PgC yr^{-1} for 2010–2019 (Caínzos et al., 2022) (Fig. S9). The 73–79% increase in C_{ant} transport in the subtropical region between 2000–2009 and 2010–2019 is greater than the 22–32% increase in the subpolar
650 region. However, stable C_{ant} transport is found at 26.5°N over a shorter period (2004–2012) at high resolution (Brown et al., 2021), suggesting an analysis over a longer period to better identify long-term trends and a better consensus between distinct methods. The 2004–2012 average C_{ant} transport convergence between A25 and 26.5°N is equal to 0.091 ± 0.033 PgC yr^{-1} : 0.191 ± 0.013 PgC yr^{-1} at 26.5°N (Brown et al., 2021) minus 0.10 ± 0.02 PgC yr^{-1} at A25. Therefore, the northward transport of C_{ant} at A25 is half the one observed at 26.5°N , similar to previous findings (Pérez et al., 2013). Due to the northward
655 increase in C_{ant} transport at A25 concomitant with its constant supply at 26.5°N , the convergence is greater (reduced) at the beginning (end) of the time period. We found that the lateral C_{ant} transport convergence between A25 and 26.5°N averages to 0.13 ± 0.07 PgC yr^{-1} in 2004 and 0.10 ± 0.05 PgC yr^{-1} in 2012. The documented increase in air-sea CO_2 fluxes between A25 and 26.5°N during this period (Gruber et al., 2023) may partially offset this reduced convergence; however, this compensation would imply a decrease in storage.

660 4.5 Limits of the OR-NN-BC method

This study represents the first quantitative assessment of $[C_{ant}]$ seasonality in the SPNA, and no comparable studies exist in the literature to validate or challenge these findings. The seasonality of $[C_{ant}]$ may depend on the approach used to calculate it,

and further investigations based on other $[C_{ant}]$ estimation approaches (TTD, TrOCA, C_{IPSL}^o , ΔC^*) may be of interest. The abiotic model, used as an assumption in the BC approach to determine the saturation of $[C_{ant}]$ in the first 25 meters based solely on the fraction of atmospheric CO_2 , is a good approximation to constrain $[C_{ant}]$ seasonality at the sea surface. We obtained a seasonal variation of ML $[C_{ant}]$ of ± 2.5 - $3.0 \mu\text{mol kg}^{-1}$, that is $1.1 \mu\text{mol kg}^{-1}$ higher on average compared to the seasonal amplitude of (on average) compared to seasonal $[C_{ant}]$ in the ML for an annual $[C_{ant}]$ profile (difference between ML and ML annual in Fig. 3d) (the maximum difference is $2.3 \mu\text{mol kg}^{-1}$ in summer). The natural variability of [DIC] coming from NN is an order of magnitude higher (Fig. 3b,d). We found a $\pm 12.5 \mu\text{mol kg}^{-1}$ amplitude increase on average in ML $[C_{nat}]$ compared to ML $[C_{nat}]$ applied to annual $[C_{nat}]$ profile (Fig. 3b). The natural variation of the CO_2 fraction in surface water, motivated by natural processes, and here incorporated in the OR-NN-BC method owing to NN, is then an order of magnitude higher than the variability caused by the increase of in the anthropogenic fraction of CO_2 in the atmosphere.

~~The large winter ML depth in the IS~~ Our results show that the seasonal variability of ML $[C_{nat}]$ and C_{ant} is generally well captured by surface C_{nat} and C_{ant} concentrations (Fig. S3) combined with a concentration gradient at depth creates discrepancies between the S10), as the difference between surface and ML values during the season: the observed ML mean concentrations remains nearly constant throughout most of the year (within $-8.6 \pm 4.9 \mu\text{mol kg}^{-1}$ for $[C_{nat}]$, and $1.8 \pm 0.4 \mu\text{mol kg}^{-1}$ for C_{ant}). Surface concentrations therefore provide a reasonable proxy for seasonal variability within the ML, except in winter. In March, deviations in $[C_{nat}]$ ($[C_{nat}]$) is below (over) saturated in winter in IS compared to the surface of up to 16.8 (2.7) $\mu\text{mol kg}^{-1}$ are observed between surface and ML values. These differences result from the combined effect of a slight vertical decrease (increase) in $[C_{ant}]$ ($[C_{nat}]$) within the winter ML, and a deeper winter ML in the IS relative to the rest of the section (Fig. 4a,b S3). The non-uniformity of ML $[C_{ant}]$ for winter deep ML has also been observed reported using a similar methodology on with Argo- O_2 profilers in the IS (Asselot et al., 2024). This point is raised by the method. Moreover, in In our case, this effect is further amplified by the NN-generated $[O_2]$ shows a, which exhibits a vertical gradient within the ML, which causes the gradient observed in. This pattern likely reflects a methodological artifact rather than a physical signal, as Argo- O_2 observations generally show a vertically homogeneous $[O_2]$ distribution within the winter ML in the IS (Piron et al., 2016, 2017; Wolf et al., 2018). Consequently, the NN-induced $[O_2]$ vertical gradient amplifies the vertical gradients in ML $[C_{ant}]$ and $[C_{nat}]$ in the ML. In general, a homogeneous, and such methodological limitation should be taken into account when applying the OR-NN-BC method.

As for the inter-product comparison, GLOSEA5 simulates deeper ML values in winter and shallower ML in summer than ECCO (Fig. S2), producing higher (lower) ML $[O_2 C_{nat}]$ across the ML is found in the IS using Argo- O_2 data (Piron et al., 2016, 2017; Wolf et al., 2018). Values in winter (summer), thereby amplifying the seasonal carbon signal. In contrast, the weaker ECCO ML depth seasonal amplitude dampens the seasonal variability of this property. The opposite behavior is observed for $[C_{ant}]$ owing to its opposed vertical gradient.

Furthermore, while studying interannual and long-term variability, a suitable choice of period for future studies (not ad hoc) is especially important, as When looking at the interannual-to-long-term signal, this study suggests that the results in C_{ant} increase rates depend strongly on the period chosen. Interannual variability is added to the linear trend, giving different results

depending on the chosen period. Continuing to address $[C_{ant}]$ variability at higher resolution on multiple time scales is also of interest.

700 ~~We~~ To summarize, and despite the methodological limitations discussed in this section, we are confident in our results ~~for~~
~~the main intraannual variability regarding the main interannual variability signals~~ and long-term trends in concentrations and
transport, as well as for their mean seasonal cycle ~~for the different layers of~~ $[C_{nat}]$, which are supported by other independent
studies. For $[C_{ant}]$, however, this study represents the first quantitative assessment of $[C_{ant}]$ seasonality in the SPNA, and no
comparable studies exist in the literature to validate or challenge these findings. Each reanalysis shows good agreement with
A25. However, year-round hydrographic data is not available for comparison with the seasonal transport estimates from the
705 reanalysis. This would be all the more interesting in winter, when seasonal maxima for most of the concentrations and transports
presented in this study are observed. In addition, the study of specific high-frequency processes will require additional data
sources to further discuss the results and explore each of the reanalysis rather than an average signal. Finally, our results will
depend even more on the reanalysis chosen if volume transport is predominant in the process, i.e. for $[C_{nat}]$ transport (seasonal
and interannual) and seasonal $[C_{ant}]$ transport.

710 5 Conclusions

This study establishes the first nearly 30-year time series of natural (C_{nat}) and anthropogenic (C_{ant}) carbon properties and transports in the SPNA at the A25 section. The novelty of this study lies in the combination of neural networks (NN) and ocean reanalysis temperature and salinity fields to estimate C_{nat} and C_{ant} (with errors), yielding monthly-resolution results consistent with biennial summer observations. The resulting time series enable the analysis of variability across time scales,
715 from seasonal to interannual and long-term, relative to hydrographic sections.

Large seasonal variability was evidenced in the mixed layer (ML) and the upper branch of the meridional overturning circulation (uMOC), the ML showing the strongest seasonal variations in tracer concentrations. For C_{nat} concentrations, seasonal variability dominates over interannual and long-term signals. We found that winter ML deepening (here referred to as mixed layer pump) accounts for two-thirds of the ML C_{nat} seasonal cycle. In summer, however, the changes in ML depth explain
720 only one-third of the ML C_{nat} seasonal cycle, with the remainder mainly attributed to biological activity, which peaks at this time of year. In contrast, for C_{ant} concentrations, the long-term signal dominates over seasonal and interannual variability. The long-term increase averages to $0.3 \pm 0.004 \mu\text{mol kg}^{-1} \text{yr}^{-1}$ for the entire section and reaches $0.825 \pm 0.016 \mu\text{mol kg}^{-1} \text{yr}^{-1}$ in the ML. Despite the relatively small seasonal amplitude of C_{ant} concentrations, ML C_{ant} also exhibited intraannual variability. In this case, the mixed layer pump accounted for about one-quarter of the mean ML C_{ant} variation in summer and half of the
725 seasonal signal in winter, the remainder likely linked to seasonal changes in the Revelle factor.

In terms of lateral tracer transport, we observed seasonal peak-to-peak amplitudes of about 25% of the annual average for the uMOC (iMOC). Therefore, significant differences in C_{nat} and C_{ant} uMOC transports can arise when comparing annual to synoptic data. Since observations of the A25 OVIDE cruise are summer dependent, further reference observations in winter might be a key element in better constraining the seasonal cycles of carbon and corresponding transport.

730 The variability of tracer transport is largely governed by volume transport for both C_{nat} and C_{ant} on all evaluated time scales, except long-term C_{ant} transport, which is driven by changes in $[C_{ant}]$ - ultimately linked to rising atmospheric CO_2 concentrations. We show the rapid increase in $[C_{ant}]$ at different rates in the water column and how it influences northward C_{ant} transport. In general, the uniform vertical relative increase in $[C_{ant}]$ combined with steep vertical gradients creates a disequilibrium in meridional $[C_{ant}]$ transport, and this imbalance intensifies as atmospheric concentrations continue to rise. We
735 ~~suspect-speculate~~ that the projected reduced AMOC, which may already have begun (e.g. Caesar et al. (2018)), will reinforce the $[C_{ant}]$ gradient between the upper and lower limbs of the MOC as the accumulation of $[C_{ant}]$ at depth will be less rapid and $[C_{ant}]$ will continue to grow.

Regarding interannual variability, the most pronounced anomaly in volume transport occurred around 2010, also affecting tracer transport due to circulation-driven variability. A significant C_{ant} transport anomaly in 2010, previously documented at
740 26.5°N, was attributed to a decrease in volume transport during that year. Here we found that a persistent long-term decrease in the uMOC volume transport was observed up to 2010. The two ocean reanalysis used to evaluate net transports (GLOSEA5 and ECCO) presented similar values before 2010, supporting the reliability of this anomaly, but diverged afterwards, leading to larger discrepancies in the tracer transport post 2010. This discrepancy in the datasets is not attributable to an inherent signal but rather to the methodologies employed in the construction of the reanalyses. The period of reanalysis divergence aligns with
745 the freshening of the eastern subpolar gyre that began in the late 2010s and peaked in 2016 (Fox et al., 2022). Together, these observations and our results indicate a transition period around 2010 with impacts on the carbon dynamics.

This study, based on ocean reanalysis, connects with studies using high-resolution Argo- O_2 Lagrangian profilers to analyze biogeochemical properties (Asselot et al., 2024) and helps to resolve natural and anthropogenic oceanic carbon cycles, providing results that can contribute to the evaluation of the impacts of climate change on marine ecosystems amid uncertain ocean
750 responses (Henson et al., 2022). Expanding the BGC-Argo float network (including Argo- O_2 profilers), combined with NN and multiple linear regression (MLR) techniques (Keppler et al., 2020; Müller et al., 2023; Brown et al., 2021), will improve understanding of biogeochemical cycles and parameterization of biogeochemical processes in Earth system models. NN and MLR techniques are effective in filling data gaps, but perform poorly in data-scarce regions due to limited training. Ultimately, enhancing the spatio-temporal coverage of ocean observations will refine climate projections and assessments of ocean carbon
755 storage. Increasing data from BGC-Argo will also support NN training to capture diverse oceanic states under climate change, particularly for carbon dynamics and acidification (Pérez et al., 2013; Gruber et al., 2023). GLODAP's extensive North Atlantic coverage (Olsen et al., 2019) aids in reliable reference data for NN training, ensuring minimal errors. Although in this study we did not find that the use of O_2 as input to NN improved the estimation of layer-averaged variables, O_2 plays a central role in biological processes such as photosynthesis/remineralization. Thereby, including O_2 data, for example, through Argo- O_2
760 profilers, will reduce errors in the estimation of carbon variables in discrete locations, especially close to the sea surface (Bittig et al., 2018; Carter et al., 2021; Asselot et al., 2024).

A logical next step following the present study would be to apply the method used here to 4D gridded fields to quantify $[C_{ant}]$ between several ocean sections and to identify the connections between advection and inventory changes in C_{ant} in

different ocean regions. Properly accounting for uncertainty in the C_{ant} inventory opens up many possibilities for future work
765 aimed at better understanding the processes governing ocean's anthropogenic and natural carbon budgets.

Data availability. The original reanalysis data can be downloaded from their different owners on the native grid but should not be interpolated to section A25 for volume transport conservation. The estimates of $[C_{nat}]$ and $[C_{ant}]$ and their transports needed to evaluate the conclusion of the article can be downloaded from Zenodo (Bajon, 2025). The GLODAPv2 database with A25 points is available free of charge in netcdf format along with its various versions <https://glodap.info/index.php/data-access/>.

770 *Author contributions.* R.B., L.I.C., and H.M. designed and developed the concept of the study. R.B. conducted the data analysis with inputs from L.I.C., H.M., R.A., and F.F.P. R.B. drafted the first version of the paper. All co-authors read and reviewed the paper, and all co-authors agreed on the final version of the paper.

Competing interests. The authors declare no competing interests.

775 *Acknowledgements.* R.B. has received funding to do his PhD from the Ifremer institute and from the Brittany Region with the "Allocations de Recherche Doctorale" (A.R.E.D.). L.I.C. acknowledge support from Ifremer. H.M. was supported by CNRS. F.F.P. was supported by the BOCATS2 (PID2019-104279GB-C21) project funded by MCIN/AEI/10.13039/501100011033. R.A. was supported by BRGM. This work is a contribution to CSIC's Thematic Interdisciplinary Platform PTI WATER:iOS and the OVIDE programme. We gratefully acknowledge support from the French Oceanographic Fleet to OVIDE (<https://doi.org/10.18142/140>; OVIDE Group, 2025). The data were collected and made freely available.

780 The authors would **also** like to thank Pascale Lherminier, for providing transport data from the A25 2002–2010 campaigns, Marcos Fontela for double-checking the C_{ant} A25 transport and Laura L. Jackson for making available the GLOSEA5 reanalysis at the A25 section. The authors would also like to thank Sébastien Hervé, who helped draw Fig. 1. We are grateful to all contributors to the xarray Python package (Hoyer and Hamman, 2017, v2022.3.0) that was used for the analyzes in this work.

785 Finally, we would like to thank the editor, Olivier Sulpis; the reviewer, Louise Delaigue; and the anonymous reviewer for helping us improve our paper.

References

- Asselot, R., Carracedo, L. I., Thierry, V., Mercier, H., Bajon, R., and Pérez, F. F.: Anthropogenic carbon pathways towards the North Atlantic interior revealed by Argo-O₂, neural networks and back-calculations, *Nature Communications*, 15, 1630, <https://doi.org/10.1038/s41467-024-46074-5>, 2024.
- 790 Bajon, R.: Monthly natural and anthropogenic carbon concentrations and transports in the subpolar North Atlantic Ocean over 1993-2021, <https://doi.org/10.5281/ZENODO.17091522>, 2025.
- Bittig, H. C., Steinhoff, T., Claustre, H., Fiedler, B., Williams, N. L., Sauzède, R., Körtzinger, A., and Gattuso, J.-P.: An Alternative to Static Climatologies: Robust Estimation of Open Ocean CO₂ Variables and Nutrient Concentrations From T, S, and O₂ Data Using Bayesian Neural Networks, *Frontiers in Marine Science*, 5, 328, <https://doi.org/10.3389/fmars.2018.00328>, 2018.
- 795 Boers, N.: Observation-based early-warning signals for a collapse of the Atlantic Meridional Overturning Circulation, *Nature Climate Change*, 11, 680–688, <https://doi.org/10.1038/s41558-021-01097-4>, 2021.
- Brainerd, K. E. and Gregg, M. C.: Surface mixed and mixing layer depths, *Deep Sea Research Part I: Oceanographic Research Papers*, 42, 1521–1543, [https://doi.org/10.1016/0967-0637\(95\)00068-H](https://doi.org/10.1016/0967-0637(95)00068-H), 1995.
- Brown, P. J., McDonagh, E. L., Sanders, R., Watson, A. J., Wanninkhof, R., King, B. A., Smeed, D. A., Baringer, M. O., Meinen, C. S.,
800 Schuster, U., Yool, A., and Messias, M.-J.: Circulation-driven variability of Atlantic anthropogenic carbon transports and uptake, *Nature Geoscience*, 14, 571–577, <https://doi.org/10.1038/s41561-021-00774-5>, 2021.
- Bushinsky, S. M., Nachod, Z., Fassbender, A. J., Tamsitt, V., Takeshita, Y., and Williams, N.: Offset Between Profiling Float and Shipboard Oxygen Observations at Depth Imparts Bias on Float pH and Derived *p* CO₂, *Global Biogeochemical Cycles*, 39, e2024GB008185, <https://doi.org/10.1029/2024GB008185>, 2025.
- 805 Böning, C. W. and Herrmann, P.: Annual Cycle of Poleward Heat Transport in the Ocean: Results from High-Resolution Modeling of the North and Equatorial Atlantic, *Journal of Physical Oceanography*, 24, 91–107, [https://doi.org/10.1175/1520-0485\(1994\)024<0091:ACOPHT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<0091:ACOPHT>2.0.CO;2), 1994.
- Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed fingerprint of a weakening Atlantic Ocean overturning circulation, *Nature*, 556, 191–196, <https://doi.org/10.1038/s41586-018-0006-5>, 2018.
- 810 Caesar, L., McCarthy, G. D., Thornalley, D. J. R., Cahill, N., and Rahmstorf, S.: Current Atlantic Meridional Overturning Circulation weakest in last millennium, *Nature Geoscience*, 14, 118–120, <https://doi.org/10.1038/s41561-021-00699-z>, 2021.
- Carter, B. R., Bittig, H. C., Fassbender, A. J., Sharp, J. D., Takeshita, Y., Xu, Y., Álvarez, M., Wanninkhof, R., Feely, R. A., and Barbero, L.: New and updated global empirical seawater property estimation routines, *Limnology and Oceanography: Methods*, 19, 785–809, <https://doi.org/10.1002/lom3.10461>, 2021.
- 815 Cañzos, V., Velo, A., Pérez, F. F., and Hernández-Guerra, A.: Anthropogenic Carbon Transport Variability in the Atlantic Ocean over Three Decades, *Global Biogeochemical Cycles*, <https://doi.org/10.1029/2022GB007475>, 2022.
- Dall’Olmo, G., Dingle, J., Polimene, L., Brewin, R. J. W., and Claustre, H.: Substantial energy input to the mesopelagic ecosystem from the seasonal mixed-layer pump, *Nature Geoscience*, 9, 820–823, <https://doi.org/10.1038/ngeo2818>, 2016.
- Daniault, N., Mercier, H., Lherminier, P., Sarafanov, A., Falina, A., Zunino, P., Pérez, F. F., Ríos, A. F., Ferron, B., Huck, T., Thierry, V., and
820 Gladyshev, S.: The northern North Atlantic Ocean mean circulation in the early 21st century, *Progress in Oceanography*, 146, 142–158, <https://doi.org/10.1016/j.pocean.2016.06.007>, 2016.

- de Boyer Montégut, C.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *Journal of Geophysical Research*, 109, C12 003, <https://doi.org/10.1029/2004JC002378>, 2004.
- DeVries, T.: The oceanic anthropogenic CO₂ sink: Storage, air-sea fluxes, and transports over the industrial era, *Global Biogeochemical Cycles*, 28, 631–647, <https://doi.org/10.1002/2013GB004739>, 2014.
- 825 Diaz, B. P., Knowles, B., Johns, C. T., Laber, C. P., Bondoc, K. G. V., Haramaty, L., Natale, F., Harvey, E. L., Kramer, S. J., Bolaños, L. M., Lowenstein, D. P., Fredricks, H. F., Graff, J., Westberry, T. K., Mojica, K. D. A., Haëntjens, N., Baetge, N., Gaube, P., Boss, E., Carlson, C. A., Behrenfeld, M. J., Van Mooy, B. A. S., and Bidle, K. D.: Seasonal mixed layer depth shapes phytoplankton physiology, viral production, and accumulation in the North Atlantic, *Nature Communications*, 12, 6634, <https://doi.org/10.1038/s41467-021-26836-1>,
830 2021.
- Fenty, I. and Wang, O.: ECCO Ocean Velocity - Monthly Mean 0.5 Degree (Version 4 release 4), <https://doi.org/10.5067/ECG5M-OVE44>, 2020.
- Fontela, M., Mercier, H., and Pérez, F. F.: Long-term integrated biogeochemical budget driven by circulation in the eastern subpolar North Atlantic, *Progress in Oceanography*, 173, 51–65, <https://doi.org/10.1016/j.pocean.2019.02.004>, 2019.
- 835 Fox, A. D., Handmann, P., Schmidt, C., Fraser, N., Rühls, S., Sanchez-Franks, A., Martin, T., Oltmanns, M., Johnson, C., Rath, W., Holliday, N. P., Biastoch, A., Cunningham, S. A., and Yashayaev, I.: Exceptional freshening and cooling in the eastern subpolar North Atlantic caused by reduced Labrador Sea surface heat loss, *Ocean Science*, 18, 1507–1533, <https://doi.org/10.5194/os-18-1507-2022>, 2022.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L.,
840 Buitenhuis, E., Chandra, N., Chevallier, F., Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S., Gruber, N., Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J. O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregón, A., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., Smith, N., Tans, P. P.,
845 Tian, H., Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2019, *Earth System Science Data*, 11, 1783–1838, <https://doi.org/10.5194/essd-11-1783-2019>, 2019.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra,
850 N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, O., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien,
855 K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., Van Der Werf, G. R., Van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth System Science Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.

- 860 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H., Luijckx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Arneeth, A., Arora, V., Bates, N. R., Becker, M., Bellouin, N., Berghoff, C. F., Bittig, H. C., Bopp, L., Cadule, P., Campbell, K., Chamberlain, M. A., Chandra, N., Chevallier, F., Chini, L. P., Colligan, T., Decayeux, J., Djeutchouang, L. M., Dou, X., Duran Rojas, C., Enyo, K., Evans, W., Fay, A. R., Feely, R. A., Ford, D. J., Foster, A., Gasser, T., Gehlen, M., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, O.,
- 865 Harris, I., Hefner, M., Heinke, J., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A. K., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Kato, E., Keeling, R. F., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Lan, X., Lauvset, S. K., Lefèvre, N., Liu, Z., Liu, J., Ma, L., Maksyutov, S., Marland, G., Mayot, N., McGuire, P. C., Metz, N., Monacci, N. M., Morgan, E. J., Nakaoka, S.-I., Neill, C., Niwa, Y., Nützel, T., Olivier, L., Ono, T., Palmer, P. I., Pierrot, D., Qin, Z., Resplandy, L., Roobaert, A., Rosan, T. M., Rödenbeck, C., Schwinger, J., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Steinhoff, T., Sun, Q., Sutton, A. J., Séférian, R., Takao, S., Tatebe, H., Tian,
- 870 H., Tilbrook, B., Torres, O., Tourigny, E., Tsujino, H., Tubiello, F., Van Der Werf, G., Wanninkhof, R., Wang, X., Yang, D., Yang, X., Yu, Z., Yuan, W., Yue, X., Zaehle, S., Zeng, N., and Zeng, J.: Global Carbon Budget 2024, *Earth System Science Data*, 17, 965–1039, <https://doi.org/10.5194/essd-17-965-2025>, 2025.
- Fröb, F., Olsen, A., Pérez, F. F., García-Ibáñez, M. I., Jeansson, E., Omar, A., and Lauvset, S. K.: Inorganic carbon and water masses in the Irminger Sea since 1991, *Biogeosciences*, 15, 51–72, <https://doi.org/10.5194/bg-15-51-2018>, 2018.
- 875 Fu, Y., Lozier, M. S., Biló, T. C., Bower, A. S., Cunningham, S. A., Cyr, F., De Jong, M. F., deYoung, B., Drysdale, L., Fraser, N., Fried, N., Furey, H. H., Han, G., Handmann, P., Holliday, N. P., Holte, J., Inall, M. E., Johns, W. E., Jones, S., Karstensen, J., Li, F., Pacini, A., Pickart, R. S., Rayner, D., Straneo, F., and Yashayaev, I.: Seasonality of the Meridional Overturning Circulation in the subpolar North Atlantic, *Communications Earth & Environment*, 4, 181, <https://doi.org/10.1038/s43247-023-00848-9>, 2023.
- Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.
- 880 Gourcuff, C., Lherminier, P., Mercier, H., and Le Traon, P. Y.: Altimetry Combined with Hydrography for Ocean Transport Estimation, *Journal of Atmospheric and Oceanic Technology*, 28, 1324–1337, <https://doi.org/10.1175/2011JTECHO818.1>, 2011.
- Grist, J. P., Josey, S. A., Marsh, R., Kwon, Y.-O., Bingham, R. J., and Blaker, A. T.: The Surface-Forced Overturning of the North Atlantic: Estimates from Modern Era Atmospheric Reanalysis Datasets, *Journal of Climate*, 27, 3596–3618, <https://doi.org/10.1175/JCLI-D-13-00070.1>, 2014.
- 885 Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T., and Wanninkhof, R.: The oceanic sink for anthropogenic CO₂ from 1994 to 2007, *Science*, 363, 1193–1199, <https://doi.org/10.1126/science.aau5153>, 2019.
- Gruber, N., Bakker, D. C. E., DeVries, T., Gregor, L., Hauck, J., Landschützer, P., McKinley, G. A., and Müller, J. D.: Trends and variability in the ocean carbon sink, *Nature Reviews Earth & Environment*, <https://doi.org/10.1038/s43017-022-00381-x>, 2023.
- 890 Guallart, E. F., Schuster, U., Fajar, N. M., Legge, O., Brown, P., Pelejero, C., Messias, M.-J., Calvo, E., Watson, A., Ríos, A. F., and Pérez, F. F.: Trends in anthropogenic CO₂ in water masses of the Subtropical North Atlantic Ocean, *Progress in Oceanography*, 131, 21–32, <https://doi.org/10.1016/j.pocean.2014.11.006>, 2015.
- Hagens, M. and Middelburg, J. J.: Attributing seasonal pH variability in surface ocean waters to governing factors, *Geophysical Research Letters*, 43, <https://doi.org/10.1002/2016GL071719>, 2016.
- 895

- Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C. E., Bopp, L., Chau, T. T. T., Gürses, O., Ilyina, T., Landschützer, P., Lenton, A., Resplandy, L., Rödenbeck, C., Schwinger, J., and Séférian, R.: Consistency and Challenges in the Ocean Carbon Sink Estimate for the Global Carbon Budget, *Frontiers in Marine Science*, 7, 571–720, <https://doi.org/10.3389/fmars.2020.571720>, 2020.
- 900 Henson, S. A., Robinson, I., Allen, J. T., and Waniek, J. J.: Effect of meteorological conditions on interannual variability in timing and magnitude of the spring bloom in the Irminger Basin, North Atlantic, *Deep Sea Research Part I: Oceanographic Research Papers*, 53, 1601–1615, <https://doi.org/10.1016/j.dsr.2006.07.009>, 2006.
- Henson, S. A., Painter, S. C., Penny Holliday, N., Stinchcombe, M. C., and Giering, S. L. C.: Unusual subpolar North Atlantic phytoplankton bloom in 2010: Volcanic fertilization or North Atlantic Oscillation?: Unusual North Atlantic Bloom, *Journal of Geophysical Research: Oceans*, 118, 4771–4780, <https://doi.org/10.1002/jgrc.20363>, 2013.
- 905 Henson, S. A., Laufkötter, C., Leung, S., Giering, S. L. C., Palevsky, H. I., and Cavan, E. L.: Uncertain response of ocean biological carbon export in a changing world, *Nature Geoscience*, 15, 248–254, <https://doi.org/10.1038/s41561-022-00927-0>, 2022.
- Holliday, N. P., Bersch, M., Berx, B., Chafik, L., Cunningham, S., Florindo-López, C., Hátún, H., Johns, W., Josey, S. A., Larsen, K. M. H., Mulet, S., Oltmanns, M., Reverdin, G., Rossby, T., Thierry, V., Valdimarsson, H., and Yashayaev, I.: Ocean circulation causes the largest freshening event for 120 years in eastern subpolar North Atlantic, *Nature Communications*, 11, 585, [https://doi.org/10.1038/s41467-020-](https://doi.org/10.1038/s41467-020-14474-y)
- 910 14474-y, 2020.
- Holte, J. and Talley, L.: A New Algorithm for Finding Mixed Layer Depths with Applications to Argo Data and Subantarctic Mode Water Formation*, *Journal of Atmospheric and Oceanic Technology*, 26, 1920–1939, <https://doi.org/10.1175/2009JTECHO543.1>, 2009.
- Holte, J., Talley, L. D., Gilson, J., and Roemmich, D.: An Argo mixed layer climatology and database, *Geophysical Research Letters*, 44, 5618–5626, <https://doi.org/10.1002/2017GL073426>, 2017.
- 915 Hoyer, S. and Hamman, J.: xarray: N-D labeled Arrays and Datasets in Python, *Journal of Open Research Software*, 5, 10, <https://doi.org/10.5334/jors.148>, 2017.
- Humphreys, M. P., Lewis, E. R., Sharp, J. D., and Pierrot, D.: PyCO2SYS v1.8: marine carbonate system calculations in Python, *Geoscientific Model Development*, 15, 15–43, <https://doi.org/10.5194/gmd-15-15-2022>, 2022.
- Häkkinen, S. and Rhines, P. B.: Decline of Subpolar North Atlantic Circulation During the 1990s, *Science*, 304, 555–559, <https://doi.org/10.1126/science.1094917>, 2004.
- 920 Jackson, L. C., Biastoch, A., Buckley, M. W., Desbruyères, D. G., Frajka-Williams, E., Moat, B., and Robson, J.: The evolution of the North Atlantic Meridional Overturning Circulation since 1980, *Nature Reviews Earth & Environment*, 3, 241–254, <https://doi.org/10.1038/s43017-022-00263-2>, 2022.
- Keeling, C. D., Piper, S. C., Bacastow, R. B., Wahlen, M., Whorf, T. P., Heimann, M., and Meijer, H. A.: Atmospheric CO₂ and ¹³CO₂ Exchange with the Terrestrial Biosphere and Oceans from 1978 to 2000: Observations and Carbon Cycle Implications, in: *A History of Atmospheric CO₂ and Its Effects on Plants, Animals, and Ecosystems*, edited by Baldwin, I., Caldwell, M., Heldmaier, G., Jackson, R. B., Lange, O., Mooney, H., Schulze, E.-D., Sommer, U., Ehleringer, J. R., Denise Dearing, M., and Cerling, T. E., vol. 177, pp. 83–113, Springer-Verlag, New York, ISBN 978-0-387-22069-7, https://doi.org/10.1007/0-387-27048-5_5, series Title: Ecological Studies, 2005.
- 925 Kepler, L., Landschützer, P., Gruber, N., Lauvset, S. K., and Stemmler, I.: Seasonal Carbon Dynamics in the Near-Global Ocean, *Global Biogeochemical Cycles*, 34, <https://doi.org/10.1029/2020GB006571>, 2020.
- 930 Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G. A., Murata, A., Ríos, A. F., and Sabine, C. L.: Global ocean storage of anthropogenic carbon, *Biogeosciences*, 10, 2169–2191, <https://doi.org/10.5194/bg-10-2169-2013>, 2013.

- Kunsch, H. R.: The Jackknife and the Bootstrap for General Stationary Observations, *The Annals of Statistics*, 17, 935 <https://doi.org/10.1214/aos/1176347265>, 1989.
- Lacour, L., Claustre, H., Prieur, L., and D'Ortenzio, F.: Phytoplankton biomass cycles in the North Atlantic subpolar gyre: A similar mechanism for two different blooms in the Labrador Sea: THE LABRADOR SEA BLOOMS, *Geophysical Research Letters*, 42, 5403–5410, <https://doi.org/10.1002/2015GL064540>, 2015.
- Lacour, L., Briggs, N., Claustre, H., Ardyna, M., and Dall'Olmo, G.: The Intraseasonal Dynamics of the Mixed Layer Pump 940 in the Subpolar North Atlantic Ocean: A Biogeochemical-Argo Float Approach, *Global Biogeochemical Cycles*, 33, 266–281, <https://doi.org/10.1029/2018GB005997>, 2019.
- Lan, X., Tans, P., Thoning, K., and NOAA Global Monitoring Laboratory: Trends in globally-averaged CO₂ determined from NOAA Global Monitoring Laboratory measurements., <https://doi.org/10.15138/9N0H-ZH07>, 2023.
- Landschützer, P., Gruber, N., Bakker, D. C. E., Stemmler, I., and Six, K. D.: Strengthening seasonal marine CO₂ variations due to increasing 945 atmospheric CO₂, *Nature Climate Change*, 8, 146–150, <https://doi.org/10.1038/s41558-017-0057-x>, 2018.
- Leseurre, C., Lo Monaco, C., Reverdin, G., Metzl, N., Fin, J., Olafsdottir, S., and Racapé, V.: Ocean carbonate system variability in the North Atlantic Subpolar surface water (1993–2017), *Biogeosciences*, 17, 2553–2577, <https://doi.org/10.5194/bg-17-2553-2020>, 2020.
- Lherminier, P., Mercier, H., Gourcuff, C., Alvarez, M., Bacon, S., and Kermabon, C.: Transports across the 2002 Greenland-Portugal Ovide section and comparison with 1997, *Journal of Geophysical Research*, 112, C07 003, <https://doi.org/10.1029/2006JC003716>, 2007.
- 950 Lherminier, P., Mercier, H., Huck, T., Gourcuff, C., Perez, F. F., Morin, P., Sarafanov, A., and Falina, A.: The Atlantic Meridional Overturning Circulation and the subpolar gyre observed at the A25-OVIDE section in June 2002 and 2004, *Deep Sea Research Part I: Oceanographic Research Papers*, 57, 1374–1391, <https://doi.org/10.1016/j.dsr.2010.07.009>, 2010.
- Liu, Y., Moore, J. K., Primeau, F., and Wang, W. L.: Reduced CO₂ uptake and growing nutrient sequestration from slowing overturning circulation, *Nature Climate Change*, 13, 83–90, <https://doi.org/10.1038/s41558-022-01555-7>, 2023.
- 955 Lozier, M. S., Li, F., Bacon, S., Bahr, F., Bower, A. S., Cunningham, S. A., De Jong, M. F., De Steur, L., deYoung, B., Fischer, J., Gary, S. F., Greenan, B. J. W., Holliday, N. P., Houk, A., Houpert, L., Inall, M. E., Johns, W. E., Johnson, H. L., Johnson, C., Karstensen, J., Koman, G., Le Bras, I. A., Lin, X., Mackay, N., Marshall, D. P., Mercier, H., Oltmanns, M., Pickart, R. S., Ramsey, A. L., Rayner, D., Straneo, F., Thierry, V., Torres, D. J., Williams, R. G., Wilson, C., Yang, J., Yashayaev, I., and Zhao, J.: A sea change in our view of overturning in the subpolar North Atlantic, *Science*, 363, 516–521, <https://doi.org/10.1126/science.aau6592>, 2019.
- 960 Lux, M., Mercier, H., and Arhan, M.: Interhemispheric exchanges of mass and heat in the Atlantic Ocean in January–March 1993, *Deep Sea Research Part I: Oceanographic Research Papers*, 48, 605–638, [https://doi.org/10.1016/S0967-0637\(00\)00033-9](https://doi.org/10.1016/S0967-0637(00)00033-9), 2001.
- MacLachlan, C., Arribas, A., Peterson, K. A., Maidens, A., Fereday, D., Scaife, A. A., Gordon, M., Vellinga, M., Williams, A., Comer, R. E., Camp, J., Xavier, P., and Madec, G.: Global Seasonal forecast system version 5 (GloSea5): a high-resolution seasonal forecast system: GloSea5: A High-Resolution Seasonal Forecast System, *Quarterly Journal of the Royal Meteorological Society*, 141, 1072–1084, 965 <https://doi.org/10.1002/qj.2396>, 2015.
- Marzocchi, A., Hirschi, J. J.-M., Holliday, N. P., Cunningham, S. A., Blaker, A. T., and Coward, A. C.: The North Atlantic subpolar circulation in an eddy-resolving global ocean model, *Journal of Marine Systems*, 142, 126–143, <https://doi.org/10.1016/j.jmarsys.2014.10.007>, 2015.
- McCarthy, G., Smeed, D., Johns, W., Frajka-Williams, E., Moat, B., Rayner, D., Baringer, M., Meinen, C., Collins, J., and Bryden, H.: Measuring the Atlantic Meridional Overturning Circulation at 26°N, *Progress in Oceanography*, 130, 91–111, 970 <https://doi.org/10.1016/j.pocean.2014.10.006>, 2015.

- McDonagh, E. L., King, B. A., Bryden, H. L., Courtois, P., Szuts, Z., Baringer, M., Cunningham, S. A., Atkinson, C., and McCarthy, G.: Continuous Estimate of Atlantic Oceanic Freshwater Flux at 26.5°N, *Journal of Climate*, 28, 8888–8906, <https://doi.org/10.1175/JCLI-D-14-00519.1>, 2015.
- 975 Mercier, H., Lherminier, P., Sarafanov, A., Gaillard, F., Daniault, N., Desbruyères, D., Falina, A., Ferron, B., Gourcuff, C., Huck, T., and Thierry, V.: Variability of the meridional overturning circulation at the Greenland–Portugal OVIDE section from 1993 to 2010, *Progress in Oceanography*, 132, 250–261, <https://doi.org/10.1016/j.pocean.2013.11.001>, 2015.
- Mercier, H., Desbruyères, D., Lherminier, P., Velo, A., Carracedo, L., Fontela, M., and Pérez, F. F.: New insights into the eastern subpolar North Atlantic meridional overturning circulation from OVIDE, *Ocean Science*, 20, 779–797, <https://doi.org/10.5194/os-20-779-2024>, 2024.
- 980 Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean, *Global Biogeochemical Cycles*, 20, 2005GB002530, <https://doi.org/10.1029/2005GB002530>, 2006.
- Müller, J. D., Gruber, N., Carter, B., Feely, R., Ishii, M., Lange, N., Lauvset, S. K., Murata, A., Olsen, A., Pérez, F. F., Sabine, C., Tanhua, T., Wanninkhof, R., and Zhu, D.: Decadal Trends in the Oceanic Storage of Anthropogenic Carbon From 1994 to 2014, *AGU Advances*, 985 4, e2023AV000875, <https://doi.org/10.1029/2023AV000875>, 2023.
- Olsen, A., Brown, K. R., Chierici, M., Johannessen, T., and Neill, C.: Sea-surface CO₂ fugacity in the subpolar North Atlantic, *Biogeosciences*, 5, 535–547, <https://doi.org/10.5194/bg-5-535-2008>, 2008.
- Olsen, A., Key, R. M., Van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The Global Ocean Data Analysis Project version 2 (GLODAPv2) – an internally consistent data product for the world ocean, *Earth System Science Data*, 8, 297–323, <https://doi.org/10.5194/essd-8-297-2016>, 990 2016.
- Olsen, A., Lange, N., Key, R. M., Tanhua, T., Álvarez, M., Becker, S., Bittig, H. C., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jones, S. D., Jutterström, S., Karlsen, M. K., Kozyr, A., Lauvset, S. K., Lo Monaco, C., Murata, A., Pérez, F. F., Pfeil, B., Schirnack, C., Steinfeldt, R., Suzuki, T., Telszewski, M., Tilbrook, B., Velo, A., and Wanninkhof, R.: 995 GLODAPv2.2019 – an update of GLODAPv2, *Earth System Science Data*, 11, 1437–1461, <https://doi.org/10.5194/essd-11-1437-2019>, 2019.
- Passow, U. and Carlson, C.: The biological pump in a high CO₂ world, *Marine Ecology Progress Series*, 470, 249–271, <https://doi.org/10.3354/meps09985>, 2012.
- Piron, A., Thierry, V., Mercier, H., and Caniaux, G.: Argo float observations of basin-scale deep convection in the Irminger sea during winter 1000 2011–2012, *Deep Sea Research Part I: Oceanographic Research Papers*, 109, 76–90, <https://doi.org/10.1016/j.dsr.2015.12.012>, 2016.
- Piron, A., Thierry, V., Mercier, H., and Caniaux, G.: Gyre-scale deep convection in the subpolar North Atlantic Ocean during winter 2014–2015, *Geophysical Research Letters*, 44, 1439–1447, <https://doi.org/10.1002/2016GL071895>, 2017.
- Pérez, F. F., Vázquez-Rodríguez, M., Louarn, E., Padín, X. A., Mercier, H., and Ríos, A. F.: Temporal variability of the anthropogenic CO₂ storage in the Irminger Sea, *Biogeosciences*, 5, 1669–1679, <https://doi.org/10.5194/bg-5-1669-2008>, 2008.
- 1005 Pérez, F. F., Vázquez-Rodríguez, M., Mercier, H., Velo, A., Lherminier, P., and Ríos, A. F.: Trends of anthropogenic CO₂ storage in North Atlantic water masses, *Biogeosciences*, 7, 1789–1807, <https://doi.org/10.5194/bg-7-1789-2010>, 2010.

- Pérez, F. F., Mercier, H., Vázquez-Rodríguez, M., Lherminier, P., Velo, A., Pardo, P. C., Rosón, G., and Ríos, A. F.: Atlantic Ocean CO₂ uptake reduced by weakening of the meridional overturning circulation, *Nature Geoscience*, 6, 146–152, <https://doi.org/10.1038/ngeo1680>, 2013.
- 1010 Pérez, F. F., Fontela, M., García-Ibáñez, M. I., Mercier, H., Velo, A., Lherminier, P., Zunino, P., de la Paz, M., Alonso-Pérez, F., Guallart, E. F., and Padin, X. A.: Meridional overturning circulation conveys fast acidification to the deep Atlantic Ocean, *Nature*, 554, 515–518, <https://doi.org/10.1038/nature25493>, 2018.
- Pérez, F. F., Becker, M., Goris, N., Gehlen, M., López-Mozos, M., Tjiputra, J., Olsen, A., Müller, J. D., Huertas, I. E., Chau, T. T. T., Cainzos, V., Velo, A., Benard, G., Hauck, J., Gruber, N., and Wanninkhof, R.: An Assessment of CO₂ Storage and Sea-Air Fluxes for the Atlantic Ocean and Mediterranean Sea Between 1985 and 2018, *Global Biogeochemical Cycles*, 38, e2023GB007862, <https://doi.org/10.1029/2023GB007862>, 2024.
- 1015 Racapé, V., Zunino, P., Mercier, H., Lherminier, P., Bopp, L., Pèrèz, F. F., and Gehlen, M.: Transport and storage of anthropogenic C in the North Atlantic Subpolar Ocean, *Biogeosciences*, 15, 4661–4682, <https://doi.org/10.5194/bg-15-4661-2018>, 2018.
- 1020 Raimondi, L., Tanhua, T., Azetsu-Scott, K., Yashayaev, I., and Wallace, D.: A 30 -Year Time Series of Transient Tracer-Based Estimates of Anthropogenic Carbon in the Central Labrador Sea, *Journal of Geophysical Research: Oceans*, 126, e2020JC017092, <https://doi.org/10.1029/2020JC017092>, 2021.
- Resplandy, L., Keeling, R. F., Rödenbeck, C., Stephens, B. B., Khatiwala, S., Rodgers, K. B., Long, M. C., Bopp, L., and Tans, P. P.: Revision of global carbon fluxes based on a reassessment of oceanic and riverine carbon transport, *Nature Geoscience*, 11, 504–509, <https://doi.org/10.1038/s41561-018-0151-3>, 2018.
- 1025 Reverdin, G., Metzl, N., Olafsdottir, S., Racapé, V., Takahashi, T., Benetti, M., Valdimarsson, H., Benoit-Cattin, A., Danielsen, M., Fin, J., Naamar, A., Pierrot, D., Sullivan, K., Bringas, F., and Goni, G.: SURATLANT: a 1993–2017 surface sampling in the central part of the North Atlantic subpolar gyre, *Earth System Science Data*, 10, 1901–1924, <https://doi.org/10.5194/essd-10-1901-2018>, 2018.
- Rodgers, K. B., Schwinger, J., Fassbender, A. J., Landschützer, P., Yamaguchi, R., Frenzel, H., Stein, K., Müller, J. D., Goris, N., Sharma, S., Bushinsky, S., Chau, T., Gehlen, M., Gallego, M. A., Gloege, L., Gregor, L., Gruber, N., Hauck, J., Iida, Y., Ishii, M., Keppler, L., Kim, J., Schlunegger, S., Tjiputra, J., Toyama, K., Vaittinada Ayar, P., and Velo, A.: Seasonal Variability of the Surface Ocean Carbon Cycle: A Synthesis, *Global Biogeochemical Cycles*, 37, e2023GB007798, <https://doi.org/10.1029/2023GB007798>, 2023.
- 1030 Roemmich, D. and Gilson, J.: The 2004–2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program, *Progress in Oceanography*, 82, 81–100, <https://doi.org/10.1016/j.pocean.2009.03.004>, 2009.
- 1035 Rustogi, P., Landschützer, P., Brune, S., and Baehr, J.: The impact of seasonality on the annual air-sea carbon flux and its interannual variability, *npj Climate and Atmospheric Science*, 6, 66, <https://doi.org/10.1038/s41612-023-00378-3>, 2023.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The Oceanic Sink for Anthropogenic CO₂, *Science*, 305, 367–371, <https://doi.org/10.1126/science.1097403>, 2004.
- 1040 Sallée, J.-B., Pellichero, V., Akhoudas, C., Pauthenet, E., Vignes, L., Schmidtko, S., Garabato, A. N., Sutherland, P., and Kuusela, M.: Summertime increases in upper-ocean stratification and mixed-layer depth, *Nature*, 591, 592–598, <https://doi.org/10.1038/s41586-021-03303-x>, 2021.
- Scaife, A. A., Arribas, A., Blockley, E., Brookshaw, A., Clark, R. T., Dunstone, N., Eade, R., Fereday, D., Folland, C. K., Gordon, M., Hermanson, L., Knight, J. R., Lea, D. J., MacLachlan, C., Maidens, A., Martin, M., Peterson, A. K., Smith, D., Vellinga, M., Wallace, E.,

- 1045 Waters, J., and Williams, A.: Skillful long-range prediction of European and North American winters, *Geophysical Research Letters*, 41, 2514–2519, <https://doi.org/10.1002/2014GL059637>, 2014.
- Sharp, J. D., Fassbender, A. J., Carter, B. R., Johnson, G. C., Schultz, C., and Dunne, J. P.: GOBAI-O<sub>2</sub>: temporally and spatially resolved fields of ocean interior dissolved oxygen over nearly two decades, preprint, ESSD – Ocean/Chemical oceanography, <https://doi.org/10.5194/essd-2022-308>, 2022.
- 1050 Sloyan, B. M., Wanninkhof, R., Kramp, M., Johnson, G. C., Talley, L. D., Tanhua, T., McDonagh, E., Cusack, C., O'Rourke, E., McGovern, E., Katsumata, K., Diggs, S., Hummon, J., Ishii, M., Azetsu-Scott, K., Boss, E., Ansorge, I., Perez, F. F., Mercier, H., Williams, M. J. M., Anderson, L., Lee, J. H., Murata, A., Kouketsu, S., Jeansson, E., Hoppema, M., and Campos, E.: The Global Ocean Ship-Based Hydrographic Investigations Program (GO-SHIP): A Platform for Integrated Multidisciplinary Ocean Science, *Frontiers in Marine Science*, 6, 445, <https://doi.org/10.3389/fmars.2019.00445>, 2019.
- 1055 Steinfeldt, R., Rhein, M., and Kieke, D.: Anthropogenic carbon storage and its decadal changes in the Atlantic between 1990–2020, *Biogeosciences*, 21, 3839–3867, <https://doi.org/10.5194/bg-21-3839-2024>, 2024.
- Szekely, T., Gourrion, J., Pouliquen, S., and Reverdin, G.: The CORA 5.2 dataset for global in situ temperature and salinity measurements: data description and validation, *Ocean Science*, 15, 1601–1614, <https://doi.org/10.5194/os-15-1601-2019>, 2019.
- Takahashi, T., Olafsson, J., Goddard, J. G., Chipman, D. W., and Sutherland, S. C.: Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: A comparative study, *Global Biogeochemical Cycles*, 7, 843–878, <https://doi.org/10.1029/93GB02263>, 1993.
- 1060 Tanhua, T., Körtzinger, A., Friis, K., Waugh, D. W., and Wallace, D. W. R.: An estimate of anthropogenic CO₂ inventory from decadal changes in oceanic carbon content, *Proceedings of the National Academy of Sciences*, 104, 3037–3042, <https://doi.org/10.1073/pnas.0606574104>, 2007.
- Thomson, R. E. and Fine, I. V.: Estimating Mixed Layer Depth from Oceanic Profile Data, *Journal of Atmospheric and Oceanic Technology*, 20, 319–329, [https://doi.org/10.1175/1520-0426\(2003\)020<0319:EMLDFO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2003)020<0319:EMLDFO>2.0.CO;2), 2003.
- 1065 Thornalley, D. J. R., Oppo, D. W., Ortega, P., Robson, J. I., Brierley, C. M., Davis, R., Hall, I. R., Moffa-Sanchez, P., Rose, N. L., Spooner, P. T., Yashayaev, I., and Keigwin, L. D.: Anomalously weak Labrador Sea convection and Atlantic overturning during the past 150 years, *Nature*, 556, 227–230, <https://doi.org/10.1038/s41586-018-0007-4>, 2018.
- Tjiputra, J. F., Assmann, K., and Heinze, C.: Anthropogenic carbon dynamics in the changing ocean, *Ocean Science*, 6, 605–614, <https://doi.org/10.5194/os-6-605-2010>, 2010.
- 1070 Tjiputra, J. F., Olsen, A., Assmann, K., Pfeil, B., and Heinze, C.: A model study of the seasonal and long-term North Atlantic surface <i>CO₂</i> variability, *Biogeosciences*, 9, 907–923, <https://doi.org/10.5194/bg-9-907-2012>, 2012.
- Tooth, O. J., Johnson, H. L., Wilson, C., and Evans, D. G.: Seasonal overturning variability in the eastern North Atlantic subpolar gyre: a Lagrangian perspective, *Ocean Science*, 19, 769–791, <https://doi.org/10.5194/os-19-769-2023>, 2023.
- 1075 Vázquez-Rodríguez, M., Touratier, F., Lo Monaco, C., Waugh, D. W., Padin, X. A., Bellerby, R. G. J., Goyet, C., Metzl, N., Ríos, A. F., and Pérez, F. F.: Anthropogenic carbon distributions in the Atlantic Ocean: data-based estimates from the Arctic to the Antarctic, *Biogeosciences*, 6, 439–451, <https://doi.org/10.5194/bg-6-439-2009>, 2009.
- Wolf, M. K., Hamme, R. C., Gilbert, D., Yashayaev, I., and Thierry, V.: Oxygen Saturation Surrounding Deep Water Formation Events in the Labrador Sea From Argo-O₂ Data, *Global Biogeochemical Cycles*, 32, 635–653, <https://doi.org/10.1002/2017GB005829>, 2018.
- 1080 Yashayaev, I.: Intensification and shutdown of deep convection in the Labrador Sea were caused by changes in atmospheric and freshwater dynamics, *Communications Earth & Environment*, 5, 156, <https://doi.org/10.1038/s43247-024-01296-9>, 2024.

- Zunino, P., Garcia-Ibañez, M. I., Lherminier, P., Mercier, H., Ríos, A. F., and Pérez, F. F.: Variability of the transport of anthropogenic CO₂ at the Greenland–Portugal OVIDE section: controlling mechanisms, *Biogeosciences*, 11, 2375–2389, <https://doi.org/10.5194/bg-11-2375-2014>, 2014.
- 1085 Zunino, P., Pérez, F. F., Fajar, N. M., Guallart, E. F., Ríos, A. F., Pelegrí, J. L., and Hernández-Guerra, A.: Transports and budgets of anthropogenic CO₂ in the tropical North Atlantic in 1992-1993 and 2010-2011: TRANSPORTS AND BUDGET OF C_{ANT} IN THE TNA, *Global Biogeochemical Cycles*, 29, 1075–1091, <https://doi.org/10.1002/2014GB005075>, 2015.
- Zunino, P., Lherminier, P., Mercier, H., Danault, N., García-Ibañez, M. I., and Pérez, F. F.: The GEOVIDE cruise in May–June 2014 reveals an intense Meridional Overturning Circulation over a cold and fresh subpolar North Atlantic, *Biogeosciences*, 14, 5323–5342, <https://doi.org/10.5194/bg-14-5323-2017>, 2017.
- 1090 Zunino, P., Mercier, H., and Thierry, V.: Why did deep convection persist over four consecutive winters (2015–2018) southeast of Cape Farewell?, *Ocean Science*, 16, 99–113, <https://doi.org/10.5194/os-16-99-2020>, 2020.
- Álvarez, M., Ríos, A. F., Pérez, F. F., Bryden, H. L., and Rosón, G.: Transports and budgets of total inorganic carbon in the subpolar and temperate North Atlantic, *Global Biogeochemical Cycles*, 17, <https://doi.org/10.1029/2002GB001881>, 2003.