

Spatial variability of greenhouse gas concentrations and fluxes in shallow coastal bays of the western Baltic Sea

Julika Zinke¹, Joakim P. Hansen¹, Martijn Hermans¹, Alexis Fonseca^{1,2}, Sofia A. Wikström¹, Linda Kumblad¹, Emil Rydin¹, Marc Geibel¹, Matthew E. Salter^{1,3}, and Christoph Humborg^{1,3}

¹Baltic Sea Centre, Stockholm University, Stockholm, Sweden

²Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

³Department of Environmental Science, Stockholm University, Stockholm, Sweden

Corresponding author: Julika Zinke (julika.zinke@su.se)

Abstract. Coastal ecosystems play a crucial role in greenhouse gas (GHG) dynamics but are less studied than open oceans or terrestrial systems. This study measured concentrations of carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) in six shallow bays of the wider Stockholm Archipelago during spring (April) and autumn (September–October) 2024 using cavity ring-down spectroscopy combined with a water equilibration system. We explored how GHG levels relate to bay physical characteristics (i.e. topographic openness, sediment properties, vegetation cover) and seawater properties (temperature, salinity, dissolved-oxygen saturation, chlorophyll-a, organic carbon, and nutrient concentrations), revealing significant seasonal variation of concentrations. Surface water pCO₂ ranged from 225–1372 ppm, CH₄ from 3.6–580 nmol L⁻¹, and N₂O from 8–20.8 nmol L⁻¹ with pCO₂ and CH₄ higher in autumn and N₂O higher in spring. CH₄ concentrations below 250 nmol L⁻¹ were negatively correlated with N₂O, while higher CH₄ levels showed a positive correlation, indicating a shift in biogeochemical processes suggesting differences in the dominant sedimentary microbial pathways. All bays except the two most open ones (which acted as net sinks in spring) served as GHG sources at the time of sampling. Most bays acted as net GHG sinks in April and sources in September, with only one bay showing net source behaviour in both seasons. with one anthropogenically degraded bay One bay that is subject to substantial human impacts (e.g. dredging, high nutrient loading, reduced vegetation cover) showed CO₂-equivalent CH₄ emissions that surpassed CO₂ uptake in this particular bay. CO₂-equivalent fluxes ranged from -76.1 -195.2 to 710.8 793.6 mg CO₂-eq m⁻² d⁻¹ (median: 56.9 131.5 mg CO₂-eq m⁻² d⁻¹). This study is distinctive in simultaneously measuring all three major GHGs across multiple bays in relation to diverse environmental controls, offering a uniquely integrated understanding of coastal GHG dynamics. These findings highlight the variability and complexity of coastal ecosystems and demonstrate the importance of high-resolution measurements for accurate up-scaling of fluxes from these dynamic environments.

1 Introduction

Coastal zones, particularly inshore habitats, are critical for understanding global GHG emissions since they are directly impacted by human activities at the land-ocean interface. Vegetated coastal ecosystems are highly productive and play an important role in carbon cycling (Al-Haj and Fulweiler, 2020) by capturing organic matter and taking up CO₂ from the atmosphere. However, this carbon sequestration is partly counterbalanced by the release of CH₄ and N₂O which have 100-year sustained global warming potentials 45 and 270 times greater than CO₂, respectively (Neubauer and Megonigal, 2015). Recent studies have shown that coastal habitats such as mangroves and salt marshes constitute significant sources of both CH₄ (Rosentreter et al., 2021a; Weber et al., 2019) and N₂O (Denman et al., 2007).

While mangroves, salt marshes, and seagrass ecosystems have been extensively studied, current estimates of coastal GHG emission budgets inadequately represent the diversity of coastal habitats, particularly shallow enclosed bays in brackish waters. Estimating GHG emissions in these diverse coastal environments is complex due to substantial spatial and temporal variability (Resplandy et al., 2024). Key influencing factors include vegetation type and density, sediment characteristics (organic content and porosity), salinity and corresponding sulfate availability, and eutrophication status (Rosentreter et al., 2021a; Al-Haj and Fulweiler, 2020). Additionally, GHG emissions show seasonal patterns driven by both biotic activity and abiotic factors such as oxygen availability, seawater temperature, wind speed and ice cover (e.g. Bange et al., 2024; Laine et al., 2024). This strong spatiotemporal variability makes scaling up GHG emissions from coastal areas using bottom-up approaches particularly challenging (Lundevall-Zara et al., 2021).

The biogeochemical processes underlying GHG production in coastal sediments are well understood (e.g. Bauer et al., 2013). CO₂ is produced through respiration and decomposition of organic matter and can be consumed by photosynthesis of phytoplankton and vegetation. N₂O is generated as a by-product of nitrification by ammonia-oxidizing bacteria (AOB) and archaea (AOA) or as an intermediate of denitrification under suboxic and hypoxic conditions (Elkins et al., 1978; Codispoti, 2010) and has been shown to be controlled by dissolved inorganic nitrogen (DIN) and oxygen availability (Murray et al., 2015). The relative importance of these pathways is regulated by dissolved inorganic nitrogen (DIN) availability and oxygen concentrations (Murray et al., 2015). Following the oxygen-based classification of Naqvi et al. (2010), these processes primarily occur under hypoxic to suboxic conditions, with nitrification becoming increasingly inhibited at O₂ concentrations below ~1–2 mL L⁻¹ and denitrification dominating at O₂ ≤ 0.1 mL L⁻¹ (Elkins et al., 1978; Codispoti, 2010). CH₄ is primarily produced via methanogenesis during organic matter degradation in anoxic sediments (Reeburgh, 2007; Amaral et al., 2021) and reaches the air-sea-interface through diffusive gas transfer and ebullition (e.g. McGinnis et al., 2006; Hermans et al., 2024; Bisander et al., 2025), though during upward diffusion through the water column, dissolved CH₄ may be aerobically oxidized by methanotrophic bacteria (e.g. Hanson and Hanson, 1996; Venetz et al., 2024) or consumed by anaerobic methanotrophic archaea (Knittel and Boetius, 2009), thereby limiting atmospheric flux.

Nevertheless, coastal eutrophication from increased nutrient input via river run-off and anthropogenic sources can destabilize alter the delicate equilibrium between methane CH₄ production by methanogens and oxidation by methanotrophs, such that net CH₄ emissions may increase or decrease depending on environmental conditions (Żygadłowska et al., 2023; Venetz

et al., 2024). Enhanced phytoplankton blooms and subsequent organic matter deposition on the seafloor lead to bottom-water oxygen depletion, which stimulates sediment ~~methane~~ CH₄ generation while reducing CH₄ oxidation efficiency by methanotrophic microorganisms (e.g. Broman et al., 2017; Egger et al., 2016). While extensive oxygen depletion typically occurs in deeper coastal waters below the photic zone, it can also develop in shallower wave-protected areas where slow water exchange promotes organic matter accumulation (Virtanen et al., 2019; Wikström et al., 2025). ~~As this material decomposes, microbial respiration consumes oxygen faster than it can be replenished, leading to hypoxic or anoxic conditions (Heip et al., 1995).~~ The extensive archipelago regions of Sweden and Finland exemplify this phenomenon, containing numerous shallow, sheltered bays that accumulate substantial organic matter and function as potential carbon sinks (Gubri et al., 2025; Wikström et al., 2025). ~~Shallow areas with water depths < 5 m comprise up to about 30,000 km², or roughly 7% of the Baltic Sea (Roth et al., 2022; Jakobsson et al., 2019), though the coverage of sheltered shallow bays, such as those investigated in this study, is likely smaller.~~ ~~These shallow, enclosed bays cover approximately 142 km² across the Stockholm and Uppsala archipelagos, Åland islands, and southwestern Finnish archipelago~~ Focusing only on the Stockholm and Uppsala archipelagos, Åland islands, and southwestern Finnish archipelago, these shallow, enclosed bays cover approximately 142 km² (Gubri et al., 2025). Similar archipelago morphology, characterized by numerous embayments, is also found further north and south along the Swedish and Finnish coasts. Despite the Baltic Sea's well-documented eutrophication (e.g. Żygadłowska et al., 2024) and its effects on coastal ecosystems, we currently lack sufficient knowledge to accurately upscale GHG emissions from these ecologically important shallow bay systems.

~~Recent~~ Advances in in situ measurement techniques, particularly cavity ring-down spectroscopy (CRDS), have enabled high-resolution, real-time monitoring of GHG concentrations in coastal waters (Rosentreter et al., 2021b; Roth et al., 2022). Using this technique, we conducted measurements of CH₄, CO₂, and N₂O in the surface waters of six shallow, sheltered, vegetated bays in the wider Stockholm Archipelago during two seasonal campaigns in April and September/ October 2024. ~~These sampling periods were selected to cover the pre-spring bloom period and the post-summer bloom period.~~ Our aim was to characterize the spatial ~~and temporal~~ variability of surface water GHG concentrations in these ~~underrepresented~~ understudied systems and to identify key environmental drivers. Our central hypothesis was that GHG concentrations and fluxes increase along a eutrophication gradient and are influenced by geomorphological and physical factors such as ~~topographic openness~~ affecting water retention time and sediment composition. We further expected that the three GHGs would show distinct spatial patterns, with hotspots emerging in different niches within a bay, highlighting the need for detailed mapping to better estimate their overall climate feedback. To this end, we examined how GHG concentrations relate to bay characteristics including topographic openness, water chemistry including eutrophication indicators, sediment properties and ~~seafloor~~ vegetation cover. These data provide critical insights into the functioning of shallow enclosed bays and contribute to more accurate scaling of coastal GHG emissions.

2 Methods

2.1 Study area

Continuous day-time measurements of CO₂, CH₄, and N₂O were conducted in the surface waters of six shallow bays in the wider Stockholm archipelago, Sweden (see Fig. 1). This region is characterized by a complex coastline with numerous shallow, sheltered bays that are variably separated from the surrounding open Baltic Sea. The six study bays were selected to represent gradients in topographic openness and trophic status observed across the region, based on previous investigations of more than 20 shallow bays (e.g. Wikström et al., 2025; Gubri et al., 2025). Bay openness was quantified using the topographic openness index (Ea), calculated as $Ea = 100 \cdot \frac{At}{a}$, where At is the cross-sectional area of the bay opening and a is the total bay area. ~~Three bays were classified as enclosed ($Ea \approx 0.01$), while three were semi-open ($Ea = 0.03 - 0.06$), and ranged between ~ 0.01 and 0.06 in the study bays (see Table 1). This metric is a reliable predictor of water retention time (Persson et al. 1994), sediment characteristics (Wikström et al., 2025) and biological communities (Munsterhjelm, 1997; Hansen et al., 2008; Snickars et al., 2009; Scheinin and Mattila, 2010), all of which can influence GHG cycling.~~ Bay openness strongly influences water retention time (Persson et al. 1994), sediment characteristics such as grain size, organic-matter content, and redox conditions (Wikström et al., 2025), as well as the composition of benthic and macrophyte communities (Munsterhjelm, 1997; Hansen et al., 2008; Snickars et al., 2009; Scheinin and Mattila, 2010). In enclosed bays, reduced water exchange promotes the accumulation of fine sediments and organic matter, creating conditions favourable for anaerobic decomposition and CH₄ production in the sediment. Conversely, open bays often are characterized by coarser, more oxygenated sediments that enhance aerobic respiration and CH₄ oxidation. Likewise, differences in macrophyte cover influence sediment oxygenation through root oxygen release and alter organic-matter deposition.

~~Within each openness category, we selected bays with varying eutrophication status based on historical total phosphorus and chlorophyll-a concentrations (see Table 1). Due to~~ Furthermore, longer water retention times in the more enclosed bays ~~naturally accumulated more~~ lead to accumulation of nutrients and had higher chlorophyll-a concentrations than in semi-open bays. It is noteworthy that Högklykeviken (HV) had significantly higher total phosphorus and chlorophyll-a concentrations than the other bays (see Table 1). Högklykeviken represents a system that has shifted from benthic vegetation dominance to plankton dominance after an extensive dredging of the opening area. The bays is subjected to a restoration measure since May 2024 (after our initial measurements in April). The restoration has consisted of an aluminum-based geoengineering treatment to decrease internal load of phosphate from the sediment (Rydin et al., 2025). All study bays were small (6 to 22 ha) and shallow (1.8–3.4 m, see Table 1), though ~~semi-~~ the more open bays were slightly deeper than enclosed ones.




Figure 1. Location of the sampling bays in the wider Stockholm Archipelago in the Western Baltic Sea. Basemap data: ©Esri, TomTom, Garmin, FAO, MET/NASA/NOAA, USGS.

Table 1. Characteristics of the six study bays, including **openness-category**; topographic openness index (Ea), physical dimensions, and **the eutrophication indicators** phosphorus and chlorophyll concentrations in seawater. Mean and maximum total phosphorus (TP) and chlorophyll a (Chl a) concentrations represent historical data from 4 to 7 sampling occasions per year during 2020-2024 (spring through autumn) and were used to characterize eutrophication status for bay selection. Bodviken (BV) was not sampled in 2020; Östra Lermaren (ÖL) not sampled in 2022-2023, the other bays were sampled all years.

Bay name	Bay category	Topographic openness (Ea)	Bay area (ha)	Maximum depth (m)	Mean (max) TP ($\mu\text{g L}^{-1}$)	Mean (max) Chl a ($\mu\text{g L}^{-1}$)
Östra Myttingeviken (ÖM)	Enclosed	0.003	6	2.5	39 (66)	8 (23)
Bodviken (BV)	Enclosed	0.006	9	1.8	42 (73)	10 (26)
Högklykeviken (HV)	Enclosed	0.010	12	2.4	67 (114)	19 (58)
Sandviken (SV)	Semi-open	0.026	22	3.4	41 (66)	8 (19)
Assöviken (AV)	Semi-open	0.032	17	3.2	37 (68)	7 (20)
Östra Lermaren (ÖL)	Semi-open	0.063	11	3.0	30 (49)	5 (10)

115 2.2 Continuous measurements of GHG concentrations

Measurements were conducted during midday in April and September/ October 2024 from a small boat. A cavity ring-down spectrometer (CRDS, model Picarro G2508, Picarro Inc., USA) coupled with a custom-built water equilibration gas analyzer system (WEGAS) was used to measure the concentrations of atmospheric and dissolved CO₂, CH₄ and N₂O. The instrument was factory-calibrated by the manufacturer in 2022, and the measurements presented here represent its first field deployment following calibration. According to the manufacturer's specifications, the precision of 1-minute averaged measurements is <300 ppb + 0.05% of reading for CO₂, <7 ppb + 0.05% for CH₄, and <10 ppb + 0.05% for N₂O. The CRDS technique is characterized by negligible instrumental drift which was confirmed by a post-campaign calibration (0.2% for CO₂ and -0.5% for CH₄ over three years since the last calibration). The offset determined by the post-campaign calibration was significantly smaller than the concentration ranges sampled in this study (-0.4 ppm for CO₂ and 0.03 ppm for CH₄ in April; -1.2 ppm CO₂ and 0.04 ppm CH₄ in September). Given that the study focuses on relative spatial and seasonal differences measured with the same instrument, such a small systematic offset would not affect the interpretation of the results. April measurements in Bodviken were conducted using a Picarro G2201-i instead of the G2508, which measured the concentrations of CO₂ and CH₄ but not N₂O.

2.2.1 The WEGAS system

130 The WEGAS system is described in detail in Humborg et al. (2019). Briefly, seawater from just below the surface (at approximately 30 cm depth) was continuously passed through a water handling system consisting of a thermosalinograph (SBE45 MicroTSG, Seabird Scientific, US) measuring seawater temperature, salinity, and conductivity; a flowmeter maintaining stable flow at $\sim 3 \text{ L min}^{-1}$; and a showerhead equilibrator (RAD-AQUA, Durrige, US). After the seawater was equilibrated with a flow of ambient air, the air stream was passed through a custom-built cryocooler that cooled the gas to a dew point of 4°C to reduce excess humidity before analysis by the CRDS. A gas handling system controlled airflow switching between ambient air measurements and equilibrator measurements. Each sampling cycle consisted of 5 minutes of ambient air followed by 40 minutes of equilibrator air, with cycles repeated until horizontal profiling of each bay was completed. Transition periods between ambient and equilibrator air were excluded from analysis. Sampling durations lasted between 60 and 90 minutes (typically ~ 75 minutes). Measurements were conducted both inside and outside bay areas. To distinguish between "inner bay" and "outer bay" sampling points, we delineated the bay boundary at the narrowest part of the inlet connecting each bay to the open Baltic Sea. This location represents the transition in water exchange, residence time, and mixing characteristics. For cross-bay comparisons, concentrations were first averaged within each bay, and summary statistics (e.g., median) were then calculated across bays using one value per bay, treating each bay as an independent unit rather than applying area-weighted averaging.

2.2.2 Gas concentration calculations

145 Mole fractions (ppm) of CO₂ were converted to partial pressures (μatm) using the Seacarb (v.3.3) x2pCO₂ function (Gattuso et al., 2021). Partial pressures Mole fractions (ppm) of CH₄ and N₂O were converted to molar concentrations using Henry's law (equation 1), assuming full equilibration in the equilibrator at ambient pressure:

$$C = p \times K_H \quad (1)$$

where C is concentration (mol L^{-1}), p is the partial pressure (~~atm=10⁶ ppm~~ 1 ppmv corresponds to 1 μatm at an ambient pressure of 1 atm), and K_H is the temperature-corrected Henry's law constant:

$$K_H = K_H^* \times \exp\left(\frac{-\Delta_{\text{sol}}H}{R} \times \left(\frac{1}{T_K} - \frac{1}{298.15 \text{ K}}\right)\right). \quad (2)$$

where K_H^* is the Henry's law constant at reference temperature (298.15 K), $\Delta_{\text{sol}}H$ is the enthalpy of dissolution, R is the gas constant and T_K is water temperature in Kelvin. Constants were obtained from Sander (2015).

Gas solubilities were calculated using the Bunsen solubility coefficient:

$$155 \quad \beta = \exp\left(A_1 + A_2\left(\frac{100}{T}\right) + A_3 \ln\left(\frac{T}{100}\right) + S\left(B_1 + B_2\left(\frac{T}{100}\right) + B_3\left(\frac{T}{100}\right)^2\right)\right) \quad (3)$$

where β is the dimensionless Bunsen coefficient, A_1 – A_3 and B_1 – B_3 are gas-specific constants from Wiesenburg and Guinasso Jr (1979), T is temperature (K), and S is salinity (g kg^{-1}). For N₂O, the solubility constant is given by $K_0 = \beta \cdot F$, whereas for CH₄ (-assuming ideal gas behaviour)-, the solubility constant is calculated as $K_0 = \beta(R \times 273.15 \text{ K})$.

2.2.3 Air-sea flux calculations

160 Air-sea fluxes of GHGs were estimated using:

$$F = k \times K_0 \times (pX_{\text{seawater}} - pX_{\text{air}}). \quad (4)$$

where F is flux, k is gas transfer velocity (m s^{-1}), K_0 is solubility, and pX represents partial pressures in seawater and air. The gas transfer velocity was calculated following Cole and Caraco (1998) which is representative for lake environments:

$$k = (2.07 + 0.215 \times U_{10}^{1.7}) \times \left(\frac{Sc}{600}\right)^{-0.5}. \quad (5)$$

165 where U_{10} is wind speed and Sc is the Schmidt number. Schmidt numbers for brackish Baltic Sea conditions were interpolated between freshwater and seawater values (Wanninkhof, 2014):

$$Sc = (Sc_{\text{seawater}} - Sc_{\text{freshwater}}) \times \frac{S}{35} + Sc_{\text{freshwater}} \cdot \quad (6)$$

170 ~~Wind speed was not measured during this study. Given the sheltered nature of all bays and absence of white-capping (typically occurring at $U \geq 4 \text{ m s}^{-1}$), we used a constant wind speed of 2 m s^{-1} for all flux calculations.~~ Wind speed at 10 m height was obtained from the ICON-EU numerical weather prediction model (Deutscher Wetterdienst, Germany). Model output at $\sim 7 \text{ km}$ horizontal resolution was accessed through the Ventusky online visualization platform (<https://www.ventusky.com>). We extracted 10-m wind values corresponding to the sampling dates and coordinates of each site. The derived wind speeds were 1.67 m s^{-1} (Sandviken), 7.0 m s^{-1} (Assöviken), 6.67 m s^{-1} (Högklykeviken), and 4.4 m s^{-1} (Bodviken) in April; 3.3 m s^{-1} (Sandviken), 3.9 m s^{-1} (Assöviken), 7.2 m s^{-1} (Högklykeviken), and 3.6 m s^{-1} (Bodviken) in September; and 2.5 m s^{-1} in both Östra Lermaren and Östra Myttingeviken in October.

2.2.4 CO₂-equivalent fluxes

To derive CO₂-equivalent fluxes, calculated fluxes ($\mu\text{mol m}^{-2}\text{day}^{-1}$) were converted to mass units ($\text{mg m}^{-2}\text{day}^{-1}$) using respective molar masses, then multiplied by 100-year sustained global warming potentials of 45 for CH₄ and 270 for N₂O (Neubauer and Megonigal, 2015).

180 2.3 Collection and analysis of seawater and sediment samples

2.3.1 Water sample collection and laboratory analysis

Surface water samples (0–1 m depth) were collected from the centre of each bay and kept cool until analysis at the certified Erken laboratory, Uppsala University (ISO/IEC 17025). Dissolved concentrations of nitrite-N and nitrate-N (NO₂-N + NO₃-N, Swedish Standards Institute (SIS), 1996), ammonium-N (NH₄-N, SIS, 2005) and phosphate-P (PO₄-P, SIS, 2004) were determined colorimetrically using an AutoAnalyzer 3 (SEAL Analytical, US) or a U-2910 analyser (Hitachi, Japan). Total nitrogen (TN, Swedish Standards Institute (SIS), 1996) and phosphorus (TP, SIS, 2004) concentrations were determined as NO₃-N and PO₄-P after persulfate digestion.

Chlorophyll a (Chl a) was determined spectrophotometrically after acetone extraction (SIS, 1980). Total organic carbon (TOC, SIS, 2024) was analyzed using a 680°C combustion catalytic oxidation method with a TOC-L analyser (Shimadzu, Japan). Organic content was estimated as loss on ignition (LOI) after combustion at 550°C (SIS, 1983). Turbidity was measured as Formazin Nephelometric Units (FNU, SIS, 2016) using a 2100 P ISO turbidity meter (Hach, CO, USA).

2.3.2 In-situ water measurements

Temperature, salinity and dissolved oxygen were measured using a WTW Multi 3420 probe (Xylem, US), and pH was measured with a YSI Pro10 pH meter (Xylem, US). All measurements were taken in the centre of each bay, adjacent to the water sampling location.

2.3.3 Vegetation surveys

Aquatic vegetation cover was ~~visually estimated~~ recorded in all basins by a free-diver in August, a few weeks prior to the September GHG measurements. For two bays (Högklykeviken and Bodviken), vegetation cover was also estimated in May, a few weeks after the April GHG measurements. For the other two bays with GHG measurements in May (Assöviken and Sandviken), we retrieved May vegetation data from a previously conducted survey (in 2022). ~~The interannual variation in total vegetation cover is rather small in these bays.~~ Survey ~~plots~~ sites consisting of 7–13 circular areas (5 m radius, $\sim 80 \text{ m}^2$) were distributed evenly from the bay opening to the innermost areas. ~~Plots~~ Survey sites were randomly allocated within subareas along a distance-from-opening gradient, excluding nearshore areas with $< 0.5 \text{ m}$ water depth. Within each ~~plot~~ survey site, percentage cover of individual taxa ~~was recorded~~ and total cover of all macroscopic autotrophs (including filamentous algae and cyanobacteria) was visually estimated. ~~The vegetation assessment method follows a standardized national protocol that has been widely applied in this region (e.g., Hansen et al., 2019).~~ For this study, we used two vegetation indicators: (1) total vegetation cover and (2) cumulative cover of all rooted vegetation (sum of all rooted taxa cover) ~~because they capture distinct functional aspects of benthic vegetation that are relevant for greenhouse-gas dynamics. Total vegetation cover provides an integrated measure of overall primary producer abundance, which can influence water-column oxygen dynamics and carbon cycling through photosynthesis and respiration. Rooted vegetation cover specifically reflects the presence of macrophytes capable of affecting sediment–water exchange processes through below-ground gas transport in addition to photosynthesis and respiration. These indicators therefore represent the most ecologically meaningful metrics for assessing vegetation-related controls on GHG concentrations in these shallow bays.~~

2.3.4 Sediment sampling and analysis

Sediment cores were collected using a gravity corer (63 mm inner diameter) and sectioned on-site immediately upon return to land. ~~Cores were sliced at the following depth intervals: 0–1 cm, 1–3 cm, 3–5 cm, 5–7 cm, 7–9 cm, 9–11 cm, 11–13 cm, 13–15 cm, 20–22 cm and 28–30 cm.~~ For this study, we used only data from the uppermost sediment layer (0–1 cm), which represents the sediment-water interface where redox-sensitive processes and exchanges directly influence surface-water GHG concentrations. We note that deeper sediment layers may be important for methane production and ebullition dynamics, but ~~were beyond the scope of the present study.~~ Sediment samples ~~from each interval~~ were homogenized in sterile containers and transferred to pre-weighed polypropylene vials for analysis. Samples were freeze-dried and pulverized to fine powder. Porosity was calculated from weight loss after freeze-drying, assuming a sediment dry density of 2.65 g cm^{-3} (Burdige, 2020). Sediment water content was determined after freeze-drying, and organic content was determined by loss on ignition (LOI) at 550°C for 2 h (U.S. Environmental Protection Agency, 1971). Organic carbon (C_{org}) and nitrogen (N_{org}) content were determined using an Elemental Combustion System (ECS 4010, Costech Analytical Technologies Inc, US). ~~For this study, we used only data from the uppermost sediment layer (0–1 cm) as it represents the most biogeochemically active zone and is most relevant to surface-water GHG dynamics.~~

3 Results and Discussion

3.1 Spatio-temporal variability of GHG across shallow bays

230 Surface water concentrations of CH₄, pCO₂, and N₂O exhibited substantial spatial and temporal variability across the six study bays, between seasons, and between areas inside and outside the bays (see Fig. 2 and Fig. A1-A6 in the appendix). Statistical analysis using Kruskal-Wallis tests (based on 10% of the data to avoid interdependence between neighbouring measurement points) confirmed significant differences in GHG concentrations between bays (see Table A1 in the appendix). Calculating post-hoc Bonferroni corrected p-values allowed us to discern which bays differed from each other (see Table A2). A Wilcoxon
235 rank-sum test further showed significant differences between inside and outside bay areas for all gases (see Table A3).

3.1.1 CO₂ concentrations and patterns

CO₂ was generally ~~slightly supersaturated~~ close to saturation in surface waters (see Table 2), with concentrations differing significantly between bays (see Fig. 2 and Tables 2 and A1). The highest concentrations were observed in Bodviken ~~during in~~ April (mean 1022±121.6 ppm) and in Östra Lermaren and Östra Myttingeviken ~~during in~~ October (mean 1108±117.7 ppm and
240 1033±83.1 ppm, respectively). These bays showed significantly higher CO₂ concentrations inside compared to outside areas (see Table A2). ~~In contrast, bays where CO₂ was near saturation showed no significant inside–outside differences.~~ Overall, no consistent patterns emerged ~~between across the bay openness categories gradient~~ or between inside versus outside areas across all bays and seasons, ~~indicating high spatial and temporal variability in CO₂ dynamics.~~

~~The bays with elevated CO₂ concentrations shared several characteristics: extensive vegetation cover, high sediment organic carbon content and lower eutrophication status compared to other study sites (see Tables 3 and 4). While it might seem counter-intuitive that the least eutrophied and most vegetated bays, act as hotspots for CO₂ emissions, this might be due to high autochthonous input from decaying plant matter. This pattern aligns with other coastal studies documenting seasonal CO₂ hotspots in areas with elevated organic matter input due to remineralization processes (Amaral et al., 2021; Asmala and Scheinin, 2024).~~

250 ~~The CO₂ concentrations observed in our study bays were at the high end of or exceeded those reported from more open archipelago areas in the Eastern Baltic Sea.~~ Previous studies from the Tvärminne archipelago in southwestern Finland reported values that were of a similar magnitude or exceeded the concentrations measured in our study: 750 ppm (Humborg et al., 2019), 4.5–13,100 ppm (Asmala and Scheinin, 2024), and ~~92–697 ppm (Brunberg et al., 2015)~~ 160–2521 ppm (Geilfus et al., 2025). Long-term measurements across the open Baltic Sea, that were conducted on the Finnmaid ferry between Travemünde
255 and Helsinki (Bittig et al., 2023), reported values ranging between 18–1238 μatm (mean 293±60 μatm) in April and 14–1198 μatm (mean 375±50 μatm) in September. Similar measurements by Schneider et al. (2014) yielded values of <200 μatm in summer and ~400 μatm in September. Notably, maximum values in our study compared well with the maximum values reported in Bittig et al. (2023) as well as the mean concentrations of 1288 ppm reported from Swedish lakes (Humborg et al., 2010). ~~This similarly suggests that these transitional environments between lakes and fully marine environments may share~~




Figure 2. Spatial variation of surface water concentrations of (a-b) CO_2 , (c-d) CH_4 and (e-f) N_2O across six shallow bays in April and September/October 2024. Box plots show median, quartiles, outliers and range for measurements inside (blue) and outside (red) each bay. N_2O data were not available for Bodviken, and no outside measurements were obtained for Östra Myttingeviken in October. Bays are arranged from left to right in increasing order of topographic openness.

260 ~~characteristics with lake systems regarding GHG emissions~~ The sheltered nature of the bays may resemble lake-like conditions with respect to air–water CO_2 exchange, but not necessarily other gases.

These findings ~~indicate suggest~~ that ~~while although~~ shallow bays often accumulate organic matter and are significant reservoirs of carbon and nutrients accumulated from surrounding areas (Gubri et al., 2025; Wikström et al., 2025), ~~they may simultaneously represent an overlooked coastal CO₂ source to the atmosphere, particularly during periods of low pelagic production~~ their role in atmospheric CO₂ exchange is not uniform. Instead, they may function either as CO₂ sources or sinks depending on seasonal conditions and bay-specific properties such as openness, vegetation cover, and eutrophication status. However, our measurements represent only snapshots from two seasons and capture transitional states rather than peak or minimum seasonal conditions. In temperate coastal environments, growth of phytoplankton and algae in spring reduces pCO₂ in the water column, while biomass decay in autumn results in elevated pCO₂. Recent studies by Honkanen et al. (2021) and Pönisch et al. (2025) reported diurnal variability in surface-water pCO₂ and CH₄ in the Baltic Sea that could be linked to biological and physical drivers such as solar radiation, temperature or biological activity. We acknowledge that our measurements, which were always conducted around noon, do not capture these diurnal fluctuations and thus likely introduce a small but systematic bias relative to true daily mean conditions. While such measurements remain valuable, more extensive, long-term monitoring is required to identify the environmental parameters that drive these systems to function as CO₂ sources or sinks across different temporal scales.

3.1.2 CH₄ concentrations and patterns

CH₄ was strongly supersaturated in all study bays (see Table 2) and significantly higher inside bays compared to open water (as see Fig. 2 and shown using a Wilcoxon rank-sum test, see Table A2 in the appendix). Concentrations were generally higher in autumn compared to spring (see Fig. 2c,d and Table 2), likely reflecting enhanced organic matter degradation and increased activity of methanogenic archaea in anoxic sediments (Conrad, 2009). The highest concentrations were recorded in Högklykeviken, reaching 181 nmol L⁻¹ in April and 580 nmol L⁻¹ in September. Östra Myttingeviken also showed elevated levels up to 494 nmol L⁻¹. Both are enclosed bays, with Högklykeviken representing a more degraded disturbed system that has shifted from benthic vegetation dominance to plankton dominance.

Methane CH₄ production occurs primarily through methanogenic archaea in oxygen-depleted sediments (Schubert and Wehrli, 2019). In enclosed bays with narrow openings, limited water exchange minimizes sediment disturbance by waves and currents, allowing organic matter to accumulate (Gubri et al., 2025) and creating conditions conducive to elevated CH₄ production (Egger et al., 2016). Recent studies have shown that such shallow, sheltered bays are significant organic carbon reservoirs, with higher accumulation correlated with vegetation cover, coastal morphology, and landscape characteristics (Wikström et al., 2025).

~~Our observations align with previous research showing that sediment porosity and organic content positively influence CH₄ concentrations (Lundevall-Zara et al., 2021; Roth et al., 2022), while rooted vegetation can negatively impact CH₄ concentrations, likely because plant roots transport oxygen to sediments, suppressing methanogenesis while enhancing methane oxidation by aerobic methanotrophs. In April, higher CH₄ concentrations were associated with warmer seawater temperature, consistent with previous studies (e.g. Lundevall-Zara et al., 2021; Roth et al., 2022). Elevated temperatures accelerate micro-~~

bial metabolic rates and enzymatic activities boosting, thereby increasing methanogenic rates (Yvon-Durocher et al., 2014; Reay et al., 2018). Warmer seawater temperatures also decrease CH_4 solubility and enhance stratification, reducing mixing and potentially increasing concentrations. Higher CH_4 concentrations were observed in bays with lower salinity during autumn, consistent with previous research (e.g. Lundevall-Zara et al., 2021; Roth et al., 2022). Salinity inhibits CH_4 production through multiple mechanisms, including increased sulfate availability that favours sulfate-reducing bacteria over methanogens (Camacho et al., 2017; Poffenbacher et al., 2011). Another factor that can contribute substantially to CH_4 emissions in shallow, organic rich sediments is ebullition (McGinnis et al., 2006; Hermans et al., 2024; Bisander et al., 2025). Recently, Bisander et al. (2025) showed that ebullition from sandy sediments can be substantial. The WEGAS system measures CH_4 from both benthic diffusion and bubble dissolution. ~~Without isotopic data, distinguishing between diffusive transport or ebullition events is not possible.~~ Consequently, the observed CH_4 concentrations represent the combined effect of these pathways, and without isotopic information we cannot distinguish between diffusive transport and ebullition. Although no visible bubbling was observed during sampling, we cannot exclude the possibility that episodic ebullition events might have impacted our measurements. This measurement limitation should be considered when interpreting the relationships between CH_4 and the environmental parameters described in section 3.1.4.

Our measured concentrations are comparable to previous studies in the Baltic Sea region. Studies in the southern Stockholm Archipelago around Askö reported 6–460 nmol L^{-1} (Roth et al., 2022) and 26–6596 nmol L^{-1} (Lundevall-Zara et al., 2021), while Tvärminne Archipelago studies from the southwestern coast of Finland reported ranges of ~~12.8–114 nmol L^{-1} (Brunberg et al.)~~ 19–469 nmol L^{-1} (Geilfus et al., 2025), 44 nmol L^{-1} (Humborg et al., 2019), 130–665 nmol L^{-1} (Myllykangas et al., 2020), and 0–6767 nmol L^{-1} (Asmala and Scheinin, 2024). The CH_4 concentrations measured in our study are significantly higher than values reported from long-term measurements in the open Baltic Sea, ranging between 3.5–6 nmol L^{-1} (Schneider et al., 2014), 2.8–18.6 nmol L^{-1} (Jacobs et al., 2020) and 3.2–22 nmol L^{-1} (Gülzow et al., 2013). The consistent observation of high spatial variability and local CH_4 hotspots across these studies underscores the need for high-resolution sampling to accurately characterize GHG dynamics in shallow coastal ecosystems.

3.1.3 N_2O concentrations and patterns

N_2O concentrations showed pronounced seasonal variation, with higher values in spring (13.7–20.8 nmol L^{-1}) than in autumn (8–11.75 nmol L^{-1}). Most bays were slightly subsaturated or close to saturation (see Table 2). ~~A clear spatial pattern emerged in April, with higher concentrations~~ In April, N_2O concentrations were higher in open bays compared to more enclosed bays (see Fig. 2e), while no such trend was apparent in the September data. In addition, N_2O concentrations were generally higher outside bays than inside, contrasting with the patterns observed for CH_4 .

~~Higher N_2O concentrations were associated with lower seawater temperatures in April. This relationship is likely driven by two key factors: (1) increased N_2O solubility at lower temperatures, and (2) the temperature sensitivity of denitrification enzymes. Specifically, cold temperatures inhibit the nosZ enzyme that encodes nitrous oxide reductase, a crucial enzyme in the~~

denitrification process responsible for reducing N_2O to N_2 . Contrary to findings reported by Murray et al. (2015), we could not observe a correlation between the concentrations of $\text{NO}_2 + \text{NO}_3$ and N_2O across the bays.

The consistently higher N_2O concentrations outside the bays can may be explained by hydrodynamic and sedimentological conditions that favour coupled nitrification-denitrification (Marchant et al., 2016). Higher flow-rates water currents enhance oxygen penetration into coarser sediments (sand, gravel, stones) that promote denitrification processes which promotes nitrification in the oxic surface layer and denitrification in underlying anoxic microzones (Murray et al., 2015). Lower concentrations inside bays are likely the result of reduced flow-rates water currents and the accumulation of fine organic matter, creating different biogeochemical conditions. These conditions promote weaker ventilation, stronger sediment-water coupling, and lower oxygen availability, which tend to suppress nitrification and favour complete denitrification to N_2 rather than N_2O , ultimately reducing dissolved N_2O concentrations. Additionally, higher N_2O concentrations outside the bays may partly reflect wind-induced mixing in the more exposed areas, where longer fetch and higher wind speeds enhance vertical exchange and stimulate nitrification-denitrification dynamics. In contrast, the sheltered bay interiors experience reduced wind forcing, limiting mixing and potentially suppressing N_2O production and release. N_2O production in these systems results from nitrification and denitrification processes mediated by distinct microbial communities. Ammonia-oxidizing bacteria and archaea (AOB and AOA) initiate nitrification, producing nitrite and nitrate, which are then reduced by denitrifying microbes under low oxygen conditions potentially producing N_2O as an intermediate product (Stein, 2020).

Few studies have simultaneously measured CO_2 , CH_4 and N_2O in shallow Baltic Sea bays. Our results are similar to those of Brunberg et al. (2025) Geilfus et al., (2025), who reported similar seasonal patterns in the Tvärminne Archipelago: higher N_2O concentrations up to 20.2 nmol L^{-1} in April compared to $8\text{--}12 \text{ nmol L}^{-1}$ in August and October concentrations of $9\text{--}25 \text{ nmol L}^{-1}$ in August/ September 2023. They suggested that spring N_2O peaks relate to sedimentation of the phytoplankton spring bloom and associated nitrification/denitrification processes. Similarly, studies from the southern Baltic Sea (Ma et al., 2019; Cheung et al., 2025) observed pronounced seasonal patterns with higher concentrations in winter and spring and low concentrations in autumn when anoxic and hypoxic conditions predominate. Seasonal patterns in dissolved N_2O observed in our shallow Baltic Sea bays, with relatively high concentrations in spring (April) and lower concentrations in autumn (September/October), are consistent with patterns reported from other Baltic coastal settings. Long term observations in the Kiel Bay (Boknis Eck time series station in Eckernförde Bay) likewise show elevated N_2O in winter and early spring followed by reduced concentrations in autumn, particularly under hypoxic or anoxic conditions (Ma et al., 2019). At that site, seasonal declines in dissolved oxygen and nutrient dynamics were closely coupled to N_2O variability, with lower autumn N_2O attributed to increased denitrification to N_2 under suboxic conditions that consume N_2O (Ma et al., 2019). Likewise, Cheung et al. (2025) identified pronounced seasonal N_2O variation in coastal Baltic waters and linked it to shifts in redox conditions and stratification that modulate microbial nitrification and denitrification pathways—processes that are both oxygen sensitive and seasonally dynamic. In shallow bays, spring mixing and higher oxygen availability may enhance nitrification and partial denitrification, leading to relatively elevated N_2O , whereas prolonged summer stratification and oxygen depletion in late summer and early autumn favour complete denitrification and N_2O consumption, resulting in lower observed N_2O concentrations. These sea-

sonally varying oxygen and nitrogen transformation dynamics offer a plausible mechanistic framework for the spring–autumn N_2O trend observed in our study..

Table 2. Mean concentrations (averaged over inside or outside bay area), ranges, and saturation percentages of CH₄, CO₂, and N₂O inside and outside of six bays in April and September.

Bay Name	Month	Location	CH ₄ range (mmol L ⁻¹)	CH ₄ mean (mmol L ⁻¹)	CH ₄ sat. (%)	pCO ₂ range (μ atm)	pCO ₂ mean (μ atm)	CO ₂ sat. (%)	N ₂ O range (mmol L ⁻¹)	N ₂ O mean (mmol L ⁻¹)	N ₂ O sat. (%)
Östra Mytingeviken	October	Inside	93.6–993.9	225.6±76.7	5604±1898.3	299.6–1221.3 788.7–1216.1	1033.6–833.1 1020.7±81.9	422.9–32.5 238.7±19.2	9.1–11.65	9.9±0.4	81.4±4.4
	April	Inside	26.9–64.8	52.1±10.9	564.9±118.5	899.7–1226.4 823.5–1216.9	402.2–124.6 1014.8±120.7	444.5–52.2 236.1±28.1	-	-	-
	September	Outside	6.3–20.7	11.1±4.5	113.3±47.6	230.3–532.2 228.4–527.9	285.9±72.8 72.2	176.1–31.2 66.0±16.8	-	-	-
		Inside	119.4–215.9	172.9±17.1	4304.4±448.0	347.2–296.4 342.2–785.1	469.7±109.4 462.3±107.8	184.4–43.9 108.5±25.3	8.3–11.5	9.3±0.6	86.4±4.3
Högklykeviken	September	Outside	15.1–112.4	43.5±24.2	1084.6±636.3	551.4–294.6 543.8–779.6	638.5±70.6 626.9±69.2	258.7–31.2 147.5±16.3	9.4–12.7	11.7±0.8	104.7±3.6
	April	Inside	49.9–180.9	133.8±35.5	1425.9±374	292.4–533.2 234.8–529.8	465.3±28.5 465.3±27.3	202.1–12.2 107.5±6.6	13.7–16.3	14.5±0.6	90±1.4
		Outside	40.6–128.4	66.7±27.6	674±270.1	218.8–394.5 219.5–406.3	352.1±53.5 259.3±60.3	111.2–22.9 38.2±12.4	14.6–17.3	16.3±0.6	95.7±1.8
	September	Inside	44.5–580.3	327.7±131.1	8267.7±3330.9	267.0–671.2 262.4–660.3	363.3±97.4 357.2±95.8	141.8–38.4 83.9±22.5	9.0–11.8	10.0±0.7	95.4±6.1
Sandvikén	April	Outside	14.7–52.5	23.3±8.6	572.6±214.4	346.5–456.4 341.1–449.5	413.4±31.2 407.0±30.8	163.1–12.5 95.5±7.2	10.9–12.1	11.6±0.2	105.8±1.6
	September	Inside	15.3–42.3	27.6±7.9	546.2±155.9	235.1–443.8 223.3–440.1	321.7±78.1 319.0±77.5	149.8–36.4 74.3±18.0	18.2–20.8	19.5±0.5	116.6±2.7
		Outside	17.9–21.4	19.2±0.97	380.1±19.3	423.6–449.1 420.1–447.0	432.7±86.6 435.2±8.8	203.8–3.0 101.1±2.0	18.8–20.9	19.9±0.4	119.2±2.15
	April	Inside	55.9–113.5	94.7±13.6	2367.9±345.6	366.7–762.7 380.0–749.5	483.6±86.6 478.9±46.4	189.2–48.3 111.7±11.2	8.9–9.9	9.4±0.17	89.9±1.7
Assöviken	September	Outside	9.8–90.3	38.2±25.3	1253.9±739.5	361.3–576.3 355.2–566.5	433.1±60.3 423.3±61.7	166.9–24.2 100.0±13.9	8.7–11.1	10.1±0.6	98.6±5.98
	April	Inside	3.7–135.9	78.6±45.0	850.9±489.1	227.9–440.4 226.5–412.8	321.7±78.1 267.0±40.4	118.2–21 63.0±11.3	15.7–17.6	16.7±0.4	106±1.8
		Outside	6.6–122.5	57.5±36.6	588.8±380.5	239.7–486.7 242.8–483.8	437.7±85.5 420.3±66.9	185.7–33.1 97.0±17.1	16.5–18.5	17.5±0.3	103.55±2.4
	September	Inside	3.9–171.5	85.9±99.8	2925.6±1245.2	413.6–763.8 393.7–750.6	483.6±86.6 538.4±106.0	311.8–44.2 125.7±23.9	8.1–11.0	9.3±0.9	86.5±5.3
Östra Lermaren	Outside	3.6–18.3	10.9±5.6	446.9±262.5	499.3–883.2 363.1–543.5	433.1±60.3 392.0±38.3	144.6–14.7 90.7±7.3	9.1–11.3	10.6±0.5	104.2±2.1	
	October	Inside	56.1–149.3	106.4±22.9	2540.2±553.7	861.9–1371.8 849.8–1352.2	1408.2±117.7 1092.4±116.1	443.6–47.2 255.9±27.2	8.7–10.3	9.4±0.3	82.1±2.4
Outside	9.2–33.7	15.2±6.8	360.2±159.6	523.8–902.3 514.5–886.6	38.1±3.9 603.7±84.8	245.8–34.4 141.7±19.9	10.3–12.2	11.2±0.45	97.9±4.9		

Table 3. Seawater properties in the different bays in April and September, including seawater temperature (T_{SW}), salinity (S), dissolved oxygen at seafloor, chlorophyll-a (Chl-a) concentration, turbidity, pH, total organic carbon (TOC), loss of ignition (LOI) as well as dissolved concentrations of total phosphorus (TP), phosphate (PO_4), total nitrogen (TN), nitrite and nitrate (NO_2+NO_3).

Bay abbrev.	Month	T_{SW} (°C)	S ($g \cdot kg^{-1}$)	Dissolved oxygen %	Chl-a ($\mu g L^{-1}$)	Turbidity (FNU)	pH	TOC ($\mu g L^{-1}$)	LOI ($mg L^{-1}$)	TP ($\mu g L^{-1}$)	PO_4 ($\mu g L^{-1}$)	TN ($\mu g L^{-1}$)	NO_2+NO_3 ($\mu g L^{-1}$)
ÖM	Oct.	9.6	4.5	59	4.1	0.8	7.4	7.0	2.1	26.1	2.7	495.2	0.46
BV	April	6.4	3.2	96	9.3	4.2	7.7	11.85	2.4	35.7	3.95	644	2.76
	Sept.	13.3	5.3	76	22.0	2.3	7.8	8.85	7	60.6	0.8	799.7	1
HV	April	3.3	4.1	104	6.9	1.7	7.8	7.4	3.4	34.9	0.85	464	1.85
	Sept.	14.9	5.4	91	13.6	3.2	7.8	10.8	7.5	58.6	0.4	944.15	1.98
SV	April	3.8	4.5	109	11.1	2.3	7.94	6.9	2.3	25.2	1.94	397	1.37
	Sept.	15	5.5	96	6.3	4.3	7.94	6.6	8.0	44.67	5.4	539.4	1.44
AV	April	3.6	4.8	102	9.5	1.2	7.7	6.1	1.6	29.8	4.3	417	27.2
	Sept.	15.1	5.6	85	13.5	3.1	7.7	7.7	4.2	46.7	0	719.6	1.8
ÖL	Oct.	11.4	5.6	NA	4.7	1.36	7.56	6.1	2.4	23.5	0	530.2	1.98

Table 4. Sediment and vegetation properties in the different bays in April and September including the total cover of aquatic vegetation, cumulative cover of rooted vegetation, organic carbon (OC_{sed}) and organic nitrogen (ON_{sed}) in the sediment and sediment porosity.

Bay abbrev.	Month	Total vegetation %	Rooted vegetation %	OC_{sed} (0-1 cm) (wt%)	ON_{sed} (0-1 cm) (wt%)	Porosity (0-1 cm)
ÖM	Oct.	65	46	28.6	4.0	0.98
BV	April	79	18	-	-	-
	Sept.	96	60	10.2	1.2	0.96
HV	April	29	3	-	-	-
	Sept.	32	20	13.3	1.7	0.97
SV	April	12	7	-	-	-
	Sept.	32	23	6.73	0.9	0.94
AV	April	30	4	-	-	-
	Sept.	54	27	8.75	1.07	0.93
ÖL	Oct.	77	63	22.3	2.4	0.98

365 3.1.4 Correlation of surface water GHG concentrations with environmental parameters across bays

To identify environmental factors associated with variability in surface-water GHG concentrations, we conducted a Spearman's rank correlation analysis using bay-averaged GHG concentrations and environmental parameters measured in the center of each bay (see Fig. 3 as well as Figures A7 and A8 in the appendix). To increase statistical power and assess general trends, data from April and September/October were pooled.

370 CO₂ concentrations were positively correlated with LOI ($r = 0.67$, $p = 0.04$) and showed negative correlation trends with chlorophyll-a ($r = -0.64$, $p = 0.05$) and pH ($r = -0.59$, $p = 0.07$), as well as a positive correlation trend with rooted vegetation cover ($r = 0.60$, $p = 0.06$). The negative relationship with chlorophyll-a and pH suggests that periods or locations of enhanced primary production are associated with CO₂ drawdown and elevated pH, whereas positive correlations with LOI and vegetation indicate that respiration and mineralization of organic matter—particularly from macrophyte-derived inputs—can offset photosynthetic uptake and elevate CO₂ concentrations in surface waters. This interpretation is supported by the observation that the bays with the highest CO₂ concentrations (Östra Lermaren, Östra Myttingeviken, and Bodviken) shared extensive rooted vegetation cover and elevated sediment organic carbon content. In Östra Lermaren and Östra Myttingeviken, which also exhibited the lowest eutrophication status as measured by TP and chlorophyll-a concentrations, high CO₂ concentrations may appear counter-intuitive but are likely driven by substantial autochthonous organic matter inputs from decaying vegetation, consistent with coastal studies documenting seasonal CO₂ hotspots linked to remineralization of organic-rich material (Amaral et al., 2021; Asmala and Scheinin, 2024). In contrast, Bodviken combined high CO₂ concentrations with comparatively higher eutrophication, suggesting that enhanced internal mineralization under nutrient-rich conditions may dominate CO₂ production in this system. Although the correlations with pH and rooted vegetation were slightly above the conventional 5% significance threshold, they are consistent with the expected coupling between primary production, organic matter mineralization, and CO₂ dynamics in shallow coastal systems. Given the limited number of bays, these trends should be interpreted as exploratory and warrant confirmation through studies with higher spatial and temporal resolution.

385 CH₄ concentrations showed a significant negative correlation with dissolved oxygen ($r = -0.75$, $p = 0.03$) and a positive correlation with LOI ($r = 0.67$, $p = 0.04$). These relationships are consistent with enhanced methanogenesis under low-oxygen conditions and increased availability of degradable organic substrates in the water column, which together promote CH₄ production and accumulation.

390 In contrast, N₂O concentrations exhibited significant negative correlations with temperature ($r = -0.82$, $p = 0.01$), TN ($r = -0.72$, $p = 0.04$), total vegetation cover ($r = -0.71$, $p = 0.04$), and rooted vegetation ($r = -0.78$, $p = 0.02$). The negative relationship of N₂O and temperature is likely driven by two key factors: (1) increased N₂O solubility at lower temperatures, and (2) the temperature sensitivity of denitrification enzymes. Under low-temperature conditions, enzymatic activity of N₂O reductase may be reduced, potentially slowing conversion of N₂O to N₂ and thereby increasing net N₂O emissions (Wang et al., 2014). In addition, vegetation-associated oxygenation of surface sediments may suppress N₂O-producing pathways while promoting N₂O reduction, contributing to lower observed N₂O concentrations. Contrary to findings reported by Murray et al. (2015), we could not observe a correlation between the concentrations of NO₂+NO₃ and N₂O across the bays. This decoupling

likely reflects the dominance of local-scale processes characteristic of shallow, sheltered bay environments. In particular, N₂O production may be spatially decoupled from water-column NO_x concentrations if it occurs primarily in sediments, where nitrate availability, redox gradients, and organic matter supply differ substantially from overlying waters. In organic-rich bay sediments, denitrification may proceed efficiently to N₂, thereby limiting N₂O accumulation despite elevated NO_x in the water column. In addition, rapid biological uptake of inorganic nitrogen by phytoplankton and benthic vegetation can reduce ambient NO₂⁻+NO₃⁻ concentrations without proportionally affecting N₂O production. Physical processes such as advection, sediment–water exchange, and episodic ebullition may further bypass water-column NO_x controls on dissolved N₂O. Finally, differences in spatial scale, environmental setting, and sampling strategy between this study and the global synthesis of Murray et al. (2015) likely contribute to the contrasting relationships observed.




Figure 3. Spearman correlation matrix between environmental parameters and CH₄, CO₂ and N₂O for data pooled from April and September/October. Blue indicates a negative correlation, red indicates a positive correlation. Significance (at the 95% confidence level) is indicated by p-values.

3.1.5 Correlation between N₂O and CH₄

Negative correlations between N₂O and CH₄ were observed across different bays and seasons (see Fig. 4). ~~Similar patterns were observed in measurements conducted around Askö in September 2020 and March 2021 (see Fig. 3a, unpublished data collected by Florian Roth, measured using the same sampling setup).~~ A similar negative correlation was reported by Woszczyk and Schubert (2021).

This negative correlation can be likely explained by the spatial distribution of the gases. N₂O concentrations were generally highest outside the bays and in the channels that connect to the open sea, where the flow water current velocities are higher and coarser substrates (sand, gravel, stones) dominate. ~~This aligns with previous studies (as in reviewed in Murray et al., 2015), who observed that high energy environments with turbulent regimes and coarser sediment types are N₂O hot spots.~~ This pattern is consistent with previous observations that N₂O hotspots often occur in hydrodynamically energetic settings, where strong

420 currents, turbulent mixing, and coarse substrates (sand and gravel) enhance oxygen penetration into sediments and stimulate nitrification (Murray et al., 2015). Such conditions also promote rapid porewater–water column exchange, facilitating the release of N_2O produced during coupled nitrification–denitrification. Our elevated N_2O concentrations in channels and outer-bay areas therefore align well with the mechanistic understanding established by earlier studies. Additionally, higher N_2O concentrations outside the bays may partly reflect wind-induced mixing in the more exposed areas, as already discussed earlier. In contrast, CH_4 was highest inside the bays, where sedimentary organic matter accumulates in fine muddy sediments. ~~This creates a gradient where decreasing flow velocity from bay openings toward inner areas corresponds with decreasing N_2O and increasing CH_4 .~~

425 However, in Högklykeviken and Östra Myttingeviken, the bays with the highest autumn CH_4 concentrations, we observed an interesting shift from negative correlations at CH_4 concentrations $<250 \text{ nmol L}^{-1}$ to positive correlations at CH_4 concentrations $>250 \text{ nmol L}^{-1}$. This threshold-like behaviour suggests that different biogeochemical processes dominate at high versus low CH_4 concentrations, which is indicative of the complex ~~redox carbon-nitrogen cycling~~ dynamics of these systems. ~~Importantly, in these two bays, high N_2O concentrations were also measured above shallow, muddy areas furthest into the bays, contradicting the general spatial pattern.~~

430 The different spatial distributions of CH_4 and N_2O ~~are most likely due to~~ may partly reflect their different optimal oxygen conditions: CH_4 production occurs mainly in anoxic regions, while N_2O production is maximal at suboxic levels ~~near oxygen minimum zones~~ where denitrification dominates (Naqvi et al., 2010; Ji et al., 2018; Barnes and Upstill-Goddard, 2018). ~~○ Although our measured dissolved oxygen levels measurements in the central bay locations indicate generally oxic conditions in both Högklykeviken ($\text{O}_{2,\text{dissolved}} = 8.3 \text{ mg L}^{-1} \approx 91\%$ saturation) and Östra Myttingeviken ($\text{O}_{2,\text{dissolved}} = 5.6 \text{ mg L}^{-1} \approx 59\%$ saturation), though minimum levels reached 71% and 23% saturation, respectively. Since measurements were taken in the center of the bays (Table 3 and also marked location in the maps, Fig. A1–A6) there might have been local hypoxia or anoxia where CH_4 peaks were observed. The local peaks in N_2O might be due to periodic O_2 entrainment events where suboxia occurs close enough to the sea surface to allow respiration (Codispoti, 2010).~~ we cannot resolve small-scale oxygen heterogeneity and therefore can only speculate that oxygen-reduced microenvironments may existed in areas of high CH_4 concentrations. Beyond oxygen availability, several additional mechanisms could explain the shift from a negative to a positive CH_4 - N_2O correlation. As mentioned earlier, increased inputs of labile organic matter can stimulate methanogenesis further inside the bays, while changes in the availability of alternative electron acceptors (e.g., nitrate, sulfate, iron) alter competition among metabolic pathways, which can suppress or enhance methanogenesis and modulate N_2O production or consumption. Coupled processes such as nitrate-dependent anaerobic methane oxidation can also link CH_4 and N cycling in non-linear ways. Ebullition would provide a pathway for CH_4 accumulation by bypassing water-column oxidation and decoupling CH_4 from dissolved N_2O dynamics. However, as mentioned previously, our measurement set-up does not allow us to discern between bubble-mediated and diffusive CH_4 . Changes in rooted vegetation and bioturbation may further modify sediment oxygen penetration and bubble release, influencing the relative dominance of CH_4 and N_2O -producing pathways. Finally, sediment disturbance from the research vessel in very shallow areas could explain these anomalous patterns. In order to resolve which of these factors operates in our bays would require targeted process data, limiting our discussion to speculations.

In Högklykeviken, an additional factor may have influenced these relationships. As part of a coastal restoration project, an aluminum-based geoengineering treatment was conducted on 13 May 2024 in the area where both CH₄ and N₂O exhibited high concentrations and positive correlations. This treatment involved injecting an aluminum solution into the sediment to increase the phosphorus retention and reduce eutrophication. Previous research has suggested that aluminium can decrease organic matter remineralization, possibly slowing methane CH₄ production (Reitzel et al., 2006; Zhou et al., 2018; Scalize et al., 2021), while Zhou et al. (2018) reported enhanced nitrogen fixation in the presence of aluminum. Whether this sediment disturbance altered microbial communities and affected GHG emissions requires further investigation that is beyond the scope of this study.




Figure 4. Correlations between N₂O and CH₄ across different bays, and seasons and studies, also including data from Askö (data collected by Florian Roth).

3.2 Flux estimates from shallow bays

To determine whether these bays acted as net sources or sinks of GHG, air-sea fluxes were calculated using the methods described in section 2.2. Individual gas flux densities had high variability between bays and seasons ~~-, similar to the complex spatial and temporal patterns observed in the surface water concentrations.~~

465 CO₂ flux densities were highly variable ranging from ~~-68 to 272~~ -661 to 1591 mg CO₂ m⁻² d⁻¹ in spring and ~~-29 to 303~~ -526 to 1218 mg CO₂ m⁻² d⁻¹ in autumn. The negative values indicate CO₂ uptake (sinks), while positive values are representative of emissions to the atmosphere (sources). Our estimated CO₂ flux densities are generally at the lower end of values reported from previous studies in the Baltic Sea, Swedish lakes, and global estimates for other coastal habitats (see table 5).

470 CH₄ flux densities were generally positive across all bays and seasons and ranged from ~~0.05 to 1.1~~ 0.001 to 15.6 mg CH₄ m⁻² d⁻¹, suggesting that all study sites acted as CH₄ sources. These estimates are similar to those reported for similar habitats by ~~Roth et al. (2022)~~ Ma et al. (2019) and fall within a similar range of ~~global estimates for tidal systems, lagoons, and fjords (Rosentreter et al., 2023)~~ as estimates from several other studies (see Table 5).

475 N₂O fluxes ~~showed strong seasonal patterns: predominantly positive in spring (0.003 to 0.02 mg N₂O m⁻² d⁻¹) and negative in autumn (-0.02 to -0.003 mg N₂O m⁻² d⁻¹). This seasonal switch from source to sink behaviour is consistent with findings by Brunberg et al. (2025) and Cheung et al. 2025, who reported similar patterns in Eastern and Southern coastal Baltic areas.~~ were small, ranging between -0.13 and 0.08 mg N₂O m⁻² d⁻¹ across bays and seasons.

480 Our ~~estimated flux densities are generally at the lower end of values reported from previous studies in the Baltic Sea, Swedish lakes, and global estimates for other coastal habitats (see table 5).~~ The more moderate fluxes observed in our study sites compared to other studies likely reflect the sheltered nature of these shallow bays and the relatively low wind speeds we encountered during our measurements. Furthermore, estimates of air-water GHG fluxes are highly sensitive to the choice of gas-transfer velocity parameterization. In this study, we applied the formulation by Cole and Caraco (1998), which was developed for shallow, sheltered, fetch-limited systems and allows for non-zero gas exchange under low wind speeds. This is particularly relevant for the studied bays, which are characterized by weak currents and limited wind-driven turbulence. Alternative parameterizations such as the open-ocean parameterization of Wanninkhof (2014) or the estuarine parameterization of Borges et al. (2004) produce significantly lower or higher estimates, respectively. These differences highlight that absolute flux values are strongly dependent on the assumed turbulence regime and caution against direct inter-study comparisons without careful consideration of the underlying gas-transfer assumptions.

Table 5. Range and median values (if available) of flux densities reported from different coastal habitats.

Study	F _{CO₂} (mg CO ₂ m ⁻² d ⁻¹)	F _{CH₄} (mg CH ₄ m ⁻² d ⁻¹)	F _{N₂O} (mg N ₂ O m ⁻² d ⁻¹)	Location	Flux model
Western Baltic Sea					
This study	-68-303 (-27.5) -691- 1591 (113)	0.05-1.1 (0.48) 0.001-15.60 (1.3)	-0.02-0.02 (-0.005) - 0.13-0.08 (0.03)	Stockholm Archipelago	Cole and Caraco (1998)
Bisander et al. (2025)	-937-3512	0.1-26 (diffusion), 232 (ebullition)	-	Stockholm Archipelago	Chamber measurements
Lundevall-Zara et al. (2021)	-	0.3-162	-	Askö	Wanninkhof (2014)
Roth et al. (2022)	-	0.05-0.69 (0.19)	-	Askö, mixed vegetated	Wanninkhof (2014)
	-	0.03-0.51 (0.16)	-	Askö, algae-dominated	
	-	0.03-0.465 (0.11)	-	Askö, bare sediments	
Eastern Baltic Sea					
Brunberg et al. Geilfus et al., 2025	-1798-1112(-72) - 1584.4-6601.6	-0.17-2.5 (-0.3) 2.2- 22.3	-0.04-0.48 (-0.08) - 0.09-1.67	Tvärminne Archipelago	Borges et al. (2004)
Asmala and Scheinin (2024)	-7000-108000 (180±4000)	-0.9-478 (31.0±50.0)	-	Tvärminne Archipelago	Raymond and Cole (2001)
Humborg et al. (2019)	3300-12000	-	-	Tvärminne Archipelago	Wanninkhof (2014)
Southern Baltic Sea					
Woszczyk and Schubert (2021)	12700	21.7	0.74	Coastal lakes, Poland	Crusius and Wanninkhof (2003)
Cheung et al. (2025)	-	-	-0.07-0.48 (0.19)	Curonian lagoon	Wanninkhof (2014)
	-	-	-0.04-0.25 (0.11)	Oder lagoon	
	-	-	-0.02-0.19 (0.09)	Vistula lagoon	
Bange et al. (1998)	-	0.82-5.9	0.02-0.31	Bodden waters	Wanninkhof (1992)
Bange et al. (2010)	-	0.1-0.23	-	Boknis Eck	Raymond and Cole (2001)
Ma et al. (2019)	-	-	-0.6-1.33	Boknis Eck	Nightingale et al. (2000)
Ma et al. (2020)	-	0.005-11.97	-	Boknis Eck	Nightingale et al. (2000)
Heyer and Berger (2000)	-	2.4-2496	-	Rügen	Chamber measurements
Pönisch et al. (2025)	2880±3840	12.2±13.4	-	Polder Drammendorf, rewetted peatland	Wanninkhof (2014)
Open Baltic					
Gülzow et al. (2013)	-	0.01-1.59	-	Gotland Basin, Mecklenburg Bight, Arkona Basin, Gulf of Finland; on board Finnmaid	Wanninkhof et al. (2009)
Schneider et al. (2014)	-301-241	-	-	On board Finnmaid	Wanninkhof et al. (2009)
Swedish lakes					
Bastviken et al. (2004)	-	0.19-4.2	-		Chamber measurements
Humborg et al. (2010)	320-883.6	-	-		Cole and Caraco (1998)
Global					
Rosentreter et al. (2023)	1020-1490 (1220)	0.67-0.85 (0.77)	0.20-0.29 (0.25)	Tidal systems	NA (Compilation)
	570-950 (710)	0.77-1.54 (1.32)	0.09-0.20 (0.15)	Lagoons	-
	-610-100 (-340)	0.03-0.06 (0.04)	0.23-0.22 (0.17)	Fjords	-
	-8310- -6780 (-7250)	4.64-7.51 (6.11)	0.05-0.23 (0.13)	Mangroves	-
	-2700- -2090 (-2130)	6.23-15.36 (10.57)	0.01-0.20 (0.11)	Salt marshes	-
	-2960- -500 (-1630)	1.20-1.63 (1.47)	-0.05- -0.3 (-0.04)	Sea grasses	-

3.3 CO₂-equivalent fluxes and net greenhouse gas balance

To assess the overall climate impact, individual gas fluxes were converted to CO₂-equivalent fluxes using 100-year sustained global warming potentials of 45 for CH₄ and 270 for N₂O. Total net CO₂-equivalent fluxes, varied significantly between bays and seasons, ranging from ~~-76.1~~ -195.2 mg CO₂-eq m⁻² d⁻¹ (net sink) in ~~Assöviken~~ Sandviken in Spring to ~~710.8~~ 793.6 mg CO₂-eq m⁻² d⁻¹ (net source) in Östra Lermaren in autumn, with a median of ~~56.9~~ 131.5 mg CO₂-eq m⁻² d⁻¹ across all measurements (not area-weighted).

~~Most bays acted as net greenhouse gas sources, with only Sandviken and Assöviken showing net sink behaviour in April~~ Most bays acted as net GHG sinks in April and net sources in September. However, in Högklykeviken and Bodviken, mean CO₂-equivalent fluxes were close to zero and associated with large variability in April, indicating a near-balanced system that alternated between weak sink and source behaviour. Bodviken showed a slightly positive mean flux in both seasons, but the large uncertainty in April suggests that this pattern should be interpreted cautiously. CO₂ fluxes generally dominated the greenhouse gas balance. However, in Högklykeviken, CH₄ emissions nearly balanced CO₂ uptake in spring and even exceeded CO₂ influx in autumn (see Fig. 5 and Table A4 in the appendix) highlighting the potential importance of CH₄ in degraded coastal systems. N₂O contributions were generally minor, except in Sandviken in April, where N₂O efflux accounted for 15% of the net flux. The large variability observed across bays and seasons underscores the challenge of scaling up fluxes from such heterogeneous environments. Nevertheless, to ~~provide estimates of~~ constrain potential regional contributions, we ~~multiplied~~ scaled our total CO₂-equivalent fluxes ~~by the estimated~~ using two area estimates: (1) the total area of shallow, enclosed bays in the archipelagos around Stockholm, Uppsala, Åland and southwestern Finland (142 km², Gubri et al., 2025) as a lower estimate and (2) the total area shallower than 5 m in the Baltic Sea (~ 30,000 km², Jakobsson et al., 2019; Roth et al., 2022) as an upper estimate. ~~This calculation results in~~ The resulting total carbon fluxes ~~ranging~~ ranged from ~~-2.95~~ -7.5 tC d⁻¹ to ~~27.5~~ 30.7 tC d⁻¹ ~~with a median of 2.2~~ (median 5.1 tC d⁻¹) for the lower limit and -1596 tC d⁻¹ to 6492 tC d⁻¹ (median 1076 tC d⁻¹) for the upper limit. These estimates highlight both the potential regional significance of these shallow bay systems and the enormous uncertainty when extrapolating from limited spatial and temporal measurements. ~~As such, these scaled fluxes provide a first-order indication of their potential regional relevance, but should not be interpreted as a closed regional budget due to spatial heterogeneity and limited spatial coverage.~~ The wide range emphasizes the need for more comprehensive monitoring to better constrain regional greenhouse gas budgets from coastal ecosystems.




Figure 5. CO₂ equivalent fluxes of CH₄, CO₂, N₂O and total fluxes from all bays in (a) April and (b) September/October estimated based on the parameterization by Cole and Caraco (1998). Bars represent mean values (averaged over bay area) and the error bars represent the standard deviation.

4 Conclusions

515 This study provides a **comprehensive spatially resolved** assessment of GHG emissions (CO₂, CH₄ and N₂O) from shallow coastal bays in the **Baltic-Sea wider Stockholm Archipelago**, and is one of the few investigations to simultaneously measure all three major GHGs across multiple bay environments. The results highlight the complex and highly variable nature of GHG dynamics in these systems. Our findings demonstrate that shallow Baltic Sea bays are significant but highly variable sources of GHGs, with net CO₂-equivalent fluxes ranging from **-76.1 -195.2** mg CO₂-eq m⁻² d⁻¹ in Spring to **710.8 793.6** mg CO₂-eq m⁻² d⁻¹ in autumn (median **56.9 131.5** mg CO₂-eq m⁻² d⁻¹ across all bays and seasons). Each GHG showed different behaviour with differing spatial and temporal variability: CO₂ has the highest variability and generally dominated **CO₂-equivalent** fluxes, CH₄ was routinely elevated inside the bays and increased from spring to autumn, while N₂O showed opposite seasonal trends with higher concentrations outside the bays **and lower concentrations in autumn than in spring**.

520

525 Interestingly, we observed a threshold behaviour in N₂O-CH₄ correlations. In the two bays with the highest concentrations of CH₄, ~~we observed a shift from negative correlations at CH₄ levels below 250 nmol L⁻¹ to positive correlations above this threshold. This is a pattern that has not previously been reported in coastal environments.~~ We observed a change in the relationship between CH₄ and N₂O, with negative correlations at CH₄ concentrations below 250 nmol L⁻¹ and positive correlations at higher concentrations. To our knowledge, such a pattern has rarely been reported for shallow coastal bay environments and highlights the complexity of coupled nitrogen and carbon cycling under variable redox and hydrodynamic conditions. This shift likely reflects a transition from conditions where nitrification and coupled nitrification–denitrification dominate to more reduced, microbially active regimes in which methanogenesis become more prevalent.

535 ~~By contextualizing GHG fluxes within a broader environmental framework, this study provides important observational insights into the potential environmental controls on coastal GHG dynamics. Bay characteristics, including openness, eutrophication status, sediment properties, and vegetation cover, influenced gas concentrations and fluxes, although the high spatial and temporal variability complicated statistical analysis with our limited number of study sites.~~ By placing GHG concentrations and fluxes in the context of measured environmental parameters, this study identifies observational relationships between bay characteristics and seawater properties with variability in coastal GHG dynamics. CO₂ was positively correlated with LOI and exhibited negative correlation trends with chlorophyll-a and pH as well as a positive correlation trend with rooted vegetation cover, while CH₄ was negatively correlated with dissolved oxygen and positively correlated with LOI. N₂O was negatively correlated with seawater temperature, TN, total vegetation and rooted vegetation. At the same time, the pronounced spatial and temporal heterogeneity across bays and seasons, together with the limited number of study sites, constrained our ability to quantitatively attribute individual drivers, underscoring the need for targeted process-based studies to resolve the mechanisms underlying these patterns.

545 The substantial variability observed between bays and seasons underscores both the complexity of these systems and the challenges in scaling up coastal GHG estimates. However, our findings suggest that shallow enclosed bays may represent an ~~overlooked understudied~~ but important component of coastal GHG budgets.

This study represents temporal and spatial snapshots that compare four to six bays in two seasons. Scaling up from such limited measurements risks substantial under- or overestimation of coastal ecosystems contributions to global GHG budgets. 550 Given this, future research should prioritize two key areas. Firstly, ~~long-term monitoring combining eddy covariance flux measurements with seawater monitoring would better capture annual and interannual variability while identifying the biological drivers of emissions, particularly methanogenic and methanotrophic communities~~ long-term monitoring that combines eddy-covariance flux measurements with high-resolution monitoring of seawater chemistry, oxygen levels, sediments, and microbial community composition. This combined approach would enable attribution of CH₄ flux variability to specific biogeochemical drivers, such as methanogenic production in sediments and methanotrophic consumption in the water column. Secondly, 555 research is needed to differentiate between ebullitive and diffusive CH₄ fluxes and to analyze factors that promote ebullition across seasonal timescales.

The increasing frequency of seasonal anoxia in coastal areas of the Baltic Sea, driven by eutrophication and climate change, will likely intensify GHG emissions from coastal areas. As such, understanding these dynamics is becoming increasingly important as coastal development and nutrient pollution continue to impact these systems.

Future research is needed to develop management frameworks that consider GHG emissions alongside traditional water quality concerns. Finally, this research provides important baseline data and methodological approaches for future investigations of GHG dynamics in shallow coastal ecosystems, and importantly, the results contribute to a more accurate scaling of coastal GHG emissions and highlight the importance of including these systems in regional and global GHG budgets.

Data availability. Data will be made available prior to publication.

Author contributions. The study was conceptualized jointly by all authors (JH, SW, LK, ER, CH, JZ, MH, AF, MS). JZ performed the water column GHG measurements (with help from CH and MS) and AF and MH collected and processed the sediment cores (with assistance from all other co-authors). JZ carried out the data analysis and visualization, and prepared the initial manuscript draft, with input from all co-authors. JH provided vegetation and seawater property data, and MH contributed the sediment data. MG developed the WEGAS system and trained JZ in its use.

Competing interests. The authors declare no competing interests.

Disclaimer.

Acknowledgements. We gratefully acknowledge the financial support provided by the BalticWaters foundation (project “Thriving Bays”). The study is conducted in collaboration with the Centre for Coastal Ecosystem and Climate Change Research (www.coastclim.org). We extend our thanks to Ulf Lindqvist and Maria Arvidsson (Naturvatten in Roslagen AB) for assistance in the field and Prof. Johan S. Eklöf (Department of Ecology, Environment and Plant Sciences, Stockholm University) for sharing vegetation data from one of the bays.

A1 Tables

Table A1. A Kruskal-Wallis test was conducted on 10% of the data in each bay to test whether the concentrations of GHGs inside the different bays were significantly different. A difference is significant if $p < 0.01$.

	April	September/ October
CO ₂	$p = 0.007$	$p = 3.3e^{-24}$
CH ₄	$p = 4.9e^{-29}$	$p = 1.78e^{-11}$
N ₂ O	$p = 1.3e^{-46}$	$p = 5.4e^{-5}$

Table A2. Bonferroni-adjusted p -values from post hoc pairwise comparisons between bays; values less than 0.05 indicate statistically significant differences. No data is available for Östra Lermaren (ÖL) and Östra Myttingeviken (ÖM) in April. Furthermore, no N₂O data is available for Bodenviken (BV) in April.

Group A	Group B	April			September/October		
		p_{CO_2}	p_{CH_4}	$p_{\text{N}_2\text{O}}$	p_{CO_2}	p_{CH_4}	$p_{\text{N}_2\text{O}}$
SV	AV	1	$4.8e^{-8}$	$2.9e^{-23}$	1	1	1
SV	HV	0.014	$1.1e^{-24}$	$4.1e^{-31}$	0.46	0.0025	0.003
SV	BV	0.58	1	-	1	0.217	1
SV	ÖL	-	-	-	$3.7e^{-8}$	1	1
SV	ÖM	-	-	-	$2.24e^{-9}$	$1.07e^{-7}$	0.915
AV	HV	0.046	0.001	0.35	0.197	$1.55e^{-4}$	$6.75e^{-5}$
AV	BV	0.62	$1.38e^{-5}$	-	1	0.035	0.24
AV	ÖL	-	-	-	$2.98e^{-8}$	1	1
AV	ÖM	-	-	-	$1.86e^{-9}$	$1.8e^{-9}$	0.15
HV	BV	1	$1.5e^{-15}$	-	0.02	1	0.49
HV	ÖL	-	-	-	$4.56e^{-15}$	0.033	0.007
HV	ÖM	-	-	-	$4.74e^{-16}$	0.27	1
BV	ÖL	-	-	-	$5.14e^{-6}$	1	1
BV	ÖM	-	-	-	$3.08e^{-7}$	0.004	1
ÖL	ÖM	-	-	-	1	$4.04e^{-6}$	1

Table A3. A Wilcoxon ranksum test was conducted to test whether the concentrations of GHGs inside and outside the bays were significantly different. A difference is significant if $h = 1$ and $p < 0.05$. No data is available outside Östra Myttingeviken and no N_2O data is available for Bodviken (BV).

	April				September/October				
	SV	AV	HV	BV	SV	AV	HV	BV	ÖL
CO ₂	$h = 1$ $p = 6e^{-183}$	$h = 1$ $p = 4.1e^{-57}$	$h = 1$ $p = 5.1e^{-68}$	$h = 1$ $p = 1.2e^{-58}$	$h = 1$ $p = 5.9e^{-20}$	$h = 1$ $p = 2.15e^{-37}$	$h = 1$ $p = 6.85e^{-16}$	$h = 1$ $p = 1e^{-38}$	$h = 1$ $p = 9.8e^{-29}$
CH ₄	$h = 1$ $p = 9.3e^{-70}$	$h = 1$ $p = 1.6e^{-8}$	$h = 1$ $p = 7.85e^{-47}$	$h = 1$ $p = 1.2e^{-58}$	$h = 1$ $p = 7e^{-61}$	$h = 1$ $p = 5.9e^{-34}$	$h = 1$ $p = 1.1e^{-46}$	$h = 1$ $p = 1.65e^{-52}$	$h = 1$ $p = 7.5e^{-29}$
N ₂ O	$h = 1$ $p = 8.7e^{-79}$	$h = 1$ $p = 1.7e^{-66}$	$h = 1$ $p = 3.4e^{-62}$	NA	$h = 1$ $p = 2.2e^{-36}$	$h = 1$ $p = 4.6e^{-36}$	$h = 1$ $p = 1.65e^{-43}$	$h = 1$ $p = 1.8e^{-52}$	$h = 1$ $p = 7.7e^{-29}$

Table A4. Percentage contribution to the total net flux calculated as $\frac{\text{abs}(\text{mean}(F_{X,CO_2-eq}))}{\text{abs}(\text{mean}(F_{CO_2})) + \text{abs}(\text{mean}(F_{CH_4,CO_2-eq})) + \text{abs}(\text{mean}(F_{N_2O,CO_2-eq}))}$. No N_2O data is available for Bodviken in April.

	April				September/October					
	SV	AV	HV	BV	SV	AV	HV	BV	ÖL	ÖM
CO ₂	80%	76%	53.2%	96.5%	55.4%	68.5%	33.5%	73.7%	93.8%	87.8%
CH ₄	5%	15.6%	46.2%	3.5%	37.2%	26.3%	65%	23.5%	4.5%	10.5%
N ₂ O	15%	8.4%	0.5%	-	7.5%	5.3%	1.5%	2.8%	1.7%	1.7%

A2 Figures




Figure A1. Surface water CH_4 concentrations in the different bays in April. Note the differences in scale between the different panels. The gray dots marks the sediment/water sampling locations during the GHG measurements, while red triangles mark long-term water monitoring stations. The dashed line marks the division between inside and outside bay area.




Figure A2. Surface water CH_4 concentrations in the different bays in September - October. Note the differences in scale between the different panels. The gray dots marks the sediment/water sampling locations during the GHG measurements, while red triangles mark long-term water monitoring stations. The dashed line marks the division between inside and outside bay area.




Figure A3. Surface water $p\text{CO}_2$ concentrations in the different bays in April. The gray dots marks the sediment/water sampling locations during the GHG measurements, while red triangles mark long-term water monitoring stations. The dashed line marks the division between inside and outside bay area.




Figure A4. Surface water $p\text{CO}_2$ concentrations in the different bays in September - October. The gray dots marks the sediment/water sampling locations during the GHG measurements, while red triangles mark long-term water monitoring stations. The dashed line marks the division between inside and outside bay area.




Figure A5. Surface water N_2O concentrations in the different bays in April. No N_2O data is available for Bodviken. The gray dots marks the sediment/water sampling locations during the GHG measurements, while red triangles mark long-term water monitoring stations. The dashed line marks the division between inside and outside bay area.




Figure A6. Surface water N_2O concentrations in the different bays in September - October. The gray dots marks the sediment/water sampling locations during the GHG measurements, while red triangles mark long-term water monitoring stations. The dashed line marks the division between inside and outside bay area.




Figure A7. Relationships between seawater properties and GHG concentrations. Correlation coefficient and significance level are given for the combined data from April and September/October.




Figure A8. Relationships between bay characteristics (openness, vegetation cover and sediment properties) and GHG concentrations. Correlation coefficient and significance level are given for the combined data from April and September/October.

References

- 580 Al-Haj, A. N. and Fulweiler, R. W.: A synthesis of methane emissions from shallow vegetated coastal ecosystems, *Global Change Biology*, 26, 2988–3005, 2020.
- Amaral, V., Ortega, T., Romera-Castillo, C., and Forja, J.: Linkages between greenhouse gases (CO₂, CH₄, and N₂O) and dissolved organic matter composition in a shallow estuary, *Science of the Total Environment*, 788, 147 863, 2021.
- Asmala, E. and Scheinin, M.: Persistent hot spots of CO₂ and CH₄ in coastal nearshore environments, *Limnology and Oceanography Letters*, 585 9, 119–127, 2024.
- Bange, H. W., Dahlke, S., Ramesh, R., Meyer-Reil, L.-A., Rapsomanikis, S., and Andreae, M.: Seasonal study of methane and nitrous oxide in the coastal waters of the southern Baltic Sea, *Estuarine, Coastal and Shelf Science*, 47, 807–817, 1998.
- Bange, H. W., Bergmann, K., Hansen, H. P., Kock, A., Koppe, R., Malien, F., and Ostrau, C.: Dissolved methane during hypoxic events at the Boknis Eck time series station (Eckernförde Bay, SW Baltic Sea), *Biogeosciences*, 7, 1279–1284, 2010.
- 590 Bange, H. W., Mongwe, P., Shutler, J. D., Arévalo-Martínez, D. L., Bianchi, D., Lauvset, S. K., Liu, C., Löscher, C. R., Martins, H., Rosentreter, J. A., et al.: Advances in understanding of air–sea exchange and cycling of greenhouse gases in the upper ocean, *Elementa: Science of the Anthropocene*, 12, 2024.
- Barnes, J. and Upstill-Goddard, R. C.: The denitrification paradox: The role of O₂ in sediment N₂O production, *Estuarine, Coastal and Shelf Science*, 200, 270–276, 2018.
- 595 Bastviken, D., Cole, J., Pace, M., and Tranvik, L.: Methane emissions from lakes: Dependence of lake characteristics, two regional assessments, and a global estimate, *Global biogeochemical cycles*, 18, 2004.
- Bauer, J. E., Cai, W.-J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. A.: The changing carbon cycle of the coastal ocean, *Nature*, 504, 61–70, 2013.
- Bisander, T., Prytherch, J., and Brüchert, V.: Methane ebullition as the dominant pathway for carbon sea-air exchange in coastal, shallow 600 water habitats of the Baltic Sea, *Biogeosciences*, 22, 4779–4796, 2025.
- Bittig, H. C., Jacobs, E., Neumann, T., and Rehder, G.: A regional pCO₂ climatology of the Baltic Sea, <https://doi.org/10.1594/PANGAEA.961119>, 2023.
- Borges, A. V., Delille, B., Schiettecatte, L.-S., Gazeau, F., Abril, G., and Frankignoulle, M.: Gas transfer velocities of CO₂ in three European estuaries (Randers Fjord, Scheldt, and Thames), *Limnology and Oceanography*, 49, 1630–1641, 2004.
- 605 Broman, E., Sjöstedt, J., Pinhassi, J., and Dopson, M.: Shifts in coastal sediment oxygenation cause pronounced changes in microbial community composition and associated metabolism, *Microbiome*, 5, 96, 2017.
- Burdige, D. J.: *Geochemistry of marine sediments*, 2020.
- Cheung, H. L., Zilius, M., Politi, T., Lorre, E., Vybernaite-Lubiene, I., Santos, I. R., and Bonaglia, S.: Nitrate-driven eutrophication supports high nitrous oxide production and emission in coastal lagoons, *Journal of Geophysical Research: Biogeosciences*, 130, e2024JG008 510, 610 2025.
- Cole, J. J. and Caraco, N. F.: Atmospheric exchange of carbon dioxide in a low-wind oligotrophic lake measured by the addition of SF₆, *Limnology and Oceanography*, 43, 647–656, 1998.
- Conrad, R.: The global methane cycle: recent advances in understanding the microbial processes involved, *Environmental microbiology reports*, 1, 285–292, 2009.

- 615 Crusius, J. and Wanninkhof, R.: Gas transfer velocities measured at low wind speed over a lake, *Limnology and Oceanography*, 48, 1010–1017, 2003.
- Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., et al.: Couplings between changes in the climate system and biogeochemistry, *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change The Physical*
- 620 *Science Basis*, pp. 499–587, 2007.
- Egger, M., Lenstra, W., Jong, D., Meysman, F. J., Sapart, C. J., Van der Veen, C., Röckmann, T., Gonzalez, S., and Slomp, C. P.: Rapid sediment accumulation results in high methane effluxes from coastal sediments, *PloS one*, 11, e0161609, 2016.
- Gattuso, J., Epitalon, J., Lavigne, H., Orr, J., Gentili, B., Hofmann, A., Proye, A., Soetaert, K., and Rae, J.: seacarb: seawater carbonate chemistry with R, R package version 3.2. 16, 2021.
- 625 Geilfus, N.-X., Delille, B., Villnäs, A., and Norkko, A.: Spatial heterogeneity of GHG dynamics across an estuarine ecosystem, *EGUphere*, 2025, 1–27, 2025.
- Gubri, B., Hansen, J. P., Wikström, S. A., Snickars, M., Dahl, M., Gullström, M., Rydin, E., Masqué, P., Garbaras, A., Björk, M., et al.: Shallow Coastal Bays as Sediment Carbon and Nutrient Reservoirs in the Baltic Sea, *Estuaries and Coasts*, 48, 1–16, 2025.
- Gülzow, W., Rehder, G., Schneider v Deimling, J., Seifert, T., and Tóth, Z.: One year of continuous measurements constraining methane
- 630 emissions from the Baltic Sea to the atmosphere using a ship of opportunity, *Biogeosciences*, 10, 81–99, 2013.
- Hansen, J. P., Sundblad, G., Bergström, U., Austin, Å. N., Donadi, S., Eriksson, B. K., and Eklöf, J. S.: Recreational boating degrades vegetation important for fish recruitment, *Ambio*, 48, 539–551, 2019.
- Hanson, R. S. and Hanson, T. E.: Methanotrophic bacteria, *Microbiological reviews*, 60, 439–471, 1996.
- Heip, C., Goosen, N., Herman, P., Kromkamp, J., Middelburg, J., and Soetaert, K.: Production and consumption of biological particles in
- 635 temperate tidal estuaries, *Oceanography and Marine Biology: an annual review*, 1995.
- Hermans, M., Stranne, C., Broman, E., Sokolov, A., Roth, F., Nascimento, F. J., Mörth, C.-M., Ten Hietbrink, S., Sun, X., Gustafsson, E., et al.: Ebullition dominates methane emissions in stratified coastal waters, *Science of the Total Environment*, 945, 174–183, 2024.
- Heyer, J. and Berger, U.: Methane emission from the coastal area in the southern Baltic Sea, *Estuarine, Coastal and Shelf Science*, 51, 13–30, 2000.
- 640 Honkanen, M., Müller, J. D., Seppälä, J., Rehder, G., Kielosto, S., Ylöstalo, P., Mäkelä, T., Hatakka, J., and Laakso, L.: The diurnal cycle of pCO₂ in the coastal region of the Baltic Sea, *Ocean Science*, 17, 1657–1675, 2021.
- Humborg, C., Mörth, C.-M., Sundbom, M., Borg, H., Blenckner, T., Giesler, R., and Ittekkot, V.: CO₂ supersaturation along the aquatic conduit in Swedish watersheds as constrained by terrestrial respiration, aquatic respiration and weathering, *Global Change Biology*, 16, 1966–1978, 2010.
- 645 Humborg, C., Geibel, M. C., Sun, X., McCrackin, M., Mörth, C.-M., Stranne, C., Jakobsson, M., Gustafsson, B., Sokolov, A., Norkko, A., et al.: High emissions of carbon dioxide and methane from the coastal Baltic Sea at the end of a summer heat wave, *Frontiers in Marine Science*, 6, 493, 2019.
- Jacobs, E., Bittig, H. C., Gräwe, U., Graves, C. A., Glockzin, M., Müller, J. D., Schneider, B., and Rehder, G.: Upwelling-induced trace gas dynamics in the Baltic Sea inferred from 8 years of autonomous measurements on a ship of opportunity, *Biogeosciences Discussions*,
- 650 2020, 1–38, 2020.
- Jakobsson, M., Stranne, C., O'Regan, M., Greenwood, S. L., Gustafsson, B., Humborg, C., and Weidner, E.: Bathymetric properties of the Baltic Sea, *Ocean Science*, 15, 905–924, 2019.

- Ji, Q., Buitenhuis, E., Suntharalingam, P., Sarmiento, J. L., and Ward, B. B.: Global nitrous oxide production determined by oxygen sensitivity of nitrification and denitrification, *Global Biogeochemical Cycles*, 32, 1790–1802, 2018.
- 655 Knittel, K. and Boetius, A.: Anaerobic oxidation of methane: progress with an unknown process, *Annual review of microbiology*, 63, 311–334, 2009.
- Lainela, S., Jacobs, E., Luik, S.-T., Rehder, G., and Lips, U.: Seasonal dynamics and regional distribution patterns of CO₂ and CH₄ in the north-eastern Baltic Sea, *Biogeosciences*, 21, 4495–4519, 2024.
- Lundevall-Zara, M., Lundevall-Zara, E., and Brüchert, V.: Sea-air exchange of methane in shallow inshore areas of the Baltic Sea, *Frontiers in Marine Science*, 8, 657459, 2021.
- 660 Ma, X., Lennartz, S. T., and Bange, H. W.: A multi-year observation of nitrous oxide at the Boknis Eck Time Series Station in the Eckernförde Bay (southwestern Baltic Sea), *Biogeosciences*, 16, 4097–4111, 2019.
- Ma, X., Sun, M., Lennartz, S. T., and Bange, H. W.: A decade of methane measurements at the Boknis Eck time series station in Eckernförde Bay (southwestern Baltic Sea), *Biogeosciences*, 17, 3427–3438, 2020.
- 665 Marchant, H., Holtappels, M., Lavik, G., Ahmerkamp, S., Winter, C., and Kuypers, M.: Coupled nitrification–denitrification leads to extensive N loss in subtidal permeable sediments, *Limnology and Oceanography*, 61, <https://doi.org/10.1002/lno.10271>, 2016.
- McGinnis, D., Greinert, J., Artemov, Y., Beaubien, S. E., and Wüest, A.: Fate of rising methane bubbles in stratified waters: How much methane reaches the atmosphere?, *Journal of Geophysical Research: Oceans*, 111, 2006.
- Murray, R. H., Erler, D. V., and Eyre, B. D.: Nitrous oxide fluxes in estuarine environments: response to global change, *Global change biology*, 21, 3219–3245, 2015.
- 670 Myllykangas, J.-P., Hietanen, S., and Jilbert, T.: Legacy effects of eutrophication on modern methane dynamics in a boreal estuary, *Estuaries and Coasts*, 43, 189–206, 2020.
- Naqvi, S., Bange, H. W., Farías, L., Monteiro, P., Scranton, M., and Zhang, J.: Marine hypoxia/anoxia as a source of CH₄ and N₂O, *Biogeosciences*, 7, 2159–2190, 2010.
- 675 Neubauer, S. C. and Megonigal, J. P.: Moving beyond global warming potentials to quantify the climatic role of ecosystems, *Ecosystems*, 18, 1000–1013, 2015.
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochemical Cycles*, 14, 373–387, 2000.
- 680 Pönisch, D. L., Bittig, H. C., Kolbe, M., Schuffenhauer, I., Otto, S., Holtermann, P., Premaratne, K., and Rehder, G.: Variability of CO₂ and CH₄ in a coastal peatland rewetted with brackish water from the Baltic Sea derived from autonomous high-resolution measurements, *Biogeosciences*, 22, 3583–3614, 2025.
- Raymond, P. A. and Cole, J. J.: Gas exchange in rivers and estuaries: Choosing a gas transfer velocity, *Estuaries*, 24, 312–317, 2001.
- Reeburgh, W. S.: Oceanic methane biogeochemistry, *Chemical reviews*, 107, 486–513, 2007.
- 685 Reitzel, K., Ahlgren, J., Gogoll, A., and Rydin, E.: Effects of aluminum treatment on phosphorus, carbon, and nitrogen distribution in lake sediment: a 31P NMR study, *Water research*, 40, 647–654, 2006.
- Resplandy, L., Hogikyan, A., Müller, J. D., Najjar, R. G., Bange, H. W., Bianchi, D., Weber, T., Cai, W.-J., Doney, S. C., Fennel, K., et al.: A synthesis of global coastal ocean greenhouse gas fluxes, *Global Biogeochemical Cycles*, 38, e2023GB007803, 2024.
- Rosentreter, J. A., Al-Haj, A. N., Fulweiler, R. W., and Williamson, P.: Methane and nitrous oxide emissions complicate coastal blue carbon 690 assessments, *Global Biogeochemical Cycles*, 35, e2020GB006858, 2021a.

- Rosentreter, J. A., Borges, A. V., Deemer, B. R., Holgerson, M. A., Liu, S., Song, C., Melack, J., Raymond, P. A., Duarte, C. M., Allen, G. H., et al.: Half of global methane emissions come from highly variable aquatic ecosystem sources, *Nature Geoscience*, 14, 225–230, 2021b.
- 695 Rosentreter, J. A., Laruelle, G. G., Bange, H. W., Bianchi, T. S., Busecke, J. J., Cai, W.-J., Eyre, B. D., Forbrich, I., Kwon, E. Y., Maavara, T., et al.: Coastal vegetation and estuaries are collectively a greenhouse gas sink, *Nature Climate Change*, 13, 579–587, 2023.
- Roth, F., Sun, X., Geibel, M. C., Prytherch, J., Brüchert, V., Bonaglia, S., Broman, E., Nascimento, F., Norkko, A., and Humborg, C.: High spatiotemporal variability of methane concentrations challenges estimates of emissions across vegetated coastal ecosystems, *Global change biology*, 28, 4308–4322, 2022.
- Rydin, E., Huser, B. J., Agstam-Norlin, O., and Kumblad, L.: Continuous phosphorus binding and accumulation in euxinic Baltic Sea sediment a decade after aluminium treatment, *Water Research*, 284, 123 945, 2025.
- 700 Sander, R.: Compilation of Henry’s law constants (version 4.0) for water as solvent, *Atmospheric Chemistry and Physics*, 15, 4399–4981, 2015.
- Scalize, P., Albuquerque, A., and Di Bernardo, L.: Impact of alum water treatment residues on the methanogenic activity in the digestion of primary domestic wastewater sludge, *Sustainability*, 13, 8783, 2021.
- 705 Schneider, B., Gülzow, W., Sadkowiak, B., and Rehder, G.: Detecting sinks and sources of CO₂ and CH₄ by ferrybox-based measurements in the Baltic Sea: Three case studies, *Journal of Marine Systems*, 140, 13–25, 2014.
- Schubert, C. J. and Wehrli, B.: Contribution of methane formation and methane oxidation to methane emission from freshwater systems, in: *Biogenesis of hydrocarbons*, pp. 401–430, Springer, 2019.
- SIS: Determination of chlorophyll in water – Extraction with acetone; Spectrophotometric method (SS 28146), <http://www.sis.se>, 1980.
- 710 SIS: Determination of suspended solids in waste water and their residue on ignition (SS 28112), <http://www.sis.se>, 1983.
- SIS: Water quality – Determination of phosphorus – Ammonium molybdate spectrometric method (ISO 6878:2004), <http://www.sis.se>, 2004.
- SIS: Water quality – Determination of ammonium nitrogen – Method by flow analysis (CFA and FIA) and spectrometric detection (ISO 11732:2005), <http://www.sis.se>, 2005.
- SIS: Water quality – Determination of turbidity – Part 1: Quantitative methods (ISO 7027-1:2016), <http://www.sis.se>, 2016.
- 715 SIS: Water analysis - Guidelines for the determination of total organic carbon (TOC) and dissolved organic carbon (DOC) (SS-EN 1484), <http://www.sis.se>, 2024.
- Swedish Standards Institute (SIS): Water quality – Determination of nitrite nitrogen and nitrate nitrogen and the sum of both by flow analysis (CFA and FIA) and spectrometric detection (ISO 13395:1996), <http://www.sis.se>, 1996.
- U.S. Environmental Protection Agency: Method 160.4: Residue, Volatile (Gravimetric, Ignition at 550°C) by Muffle Furnace, Epa analytical method, U.S. Environmental Protection Agency, 1971.
- 720 Venetz, J., Żygadłowska, O. M., Dotsios, N., Wallenius, A. J., van Helmond, N. A., Lenstra, W. K., Klomp, R., Slomp, C. P., Jetten, M. S., and Veraart, A. J.: Seasonal dynamics of the microbial methane filter in the water column of a eutrophic coastal basin, *FEMS Microbiology Ecology*, 100, fae007, 2024.
- Virtanen, E. A., Norkko, A., Nyström Sandman, A., and Viitasalo, M.: Identifying areas prone to coastal hypoxia—the role of topography, *Biogeosciences*, 16, 3183–3195, 2019.
- 725 Wang, X., Yang, X., Zhang, Z., Ye, X., Kao, C. M., and Chen, S.: Long-term effect of temperature on N₂O emission from the denitrifying activated sludge, *Journal of bioscience and bioengineering*, 117, 298–304, 2014.

- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *Journal of Geophysical Research: Oceans*, 97, 7373–7382, 1992.
- 730 Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean revisited, *Limnology and Oceanography: Methods*, 12, 351–362, 2014.
- Wanninkhof, R., Asher, W. E., Ho, D. T., Sweeney, C., and McGillis, W. R.: Advances in quantifying air-sea gas exchange and environmental forcing, *Annual review of marine science*, 1, 213–244, 2009.
- Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nature communications*, 735 10, 4584, 2019.
- Wiesenburg, D. A. and Guinasso Jr, N. L.: Equilibrium solubilities of methane, carbon monoxide, and hydrogen in water and sea water, *Journal of chemical and engineering data*, 24, 356–360, 1979.
- Wikström, S. A., Gubri, B., Asplund, M. E., Dahl, M., Gullström, M., Hansen, J. P., Kumblad, L., Rydén, E., Garbaras, A., and Björk, M.: Influence of landscape characteristics and submerged aquatic vegetation on sediment carbon and nitrogen storage in shallow brackish 740 water habitats, *Scientific Reports*, 15, 7808, 2025.
- Woszczyk, M. and Schubert, C. J.: Greenhouse gas emissions from Baltic coastal lakes, *Science of the Total Environment*, 755, 143 500, 2021.
- Zhou, L., Tan, Y., Huang, L., Fortin, C., and Campbell, P. G.: Aluminum effects on marine phytoplankton: implications for a revised iron hypothesis (iron–aluminum hypothesis), *Biogeochemistry*, 139, 123–137, 2018.
- 745 Żygadłowska, O. M., Venetz, J., Klomp, R., Lenstra, W. K., van Helmond, N. A., Röckmann, T., Wallenius, A. J., Martins, P. D., Veraart, A. J., Jetten, M. S., et al.: Pathways of methane removal in the sediment and water column of a seasonally anoxic eutrophic marine basin, *Frontiers in Marine Science*, 10, 1085 728, 2023.
- Żygadłowska, O. M., Roth, F., van Helmond, N. A., Lenstra, W. K., Venetz, J., Dotsios, N., Rockmann, T., Veraart, A. J., Stranne, C., Humborg, C., et al.: Eutrophication and deoxygenation drive high methane emissions from a Brackish Coastal system, *Environmental 750 Science & Technology*, 58, 10 582–10 590, 2024.