

Reviewer #3:**Comments and Suggestions for Authors:**

Study on the life cycle of ice crystal cloud over the Taklimakan desert using multi-source data" by Su et al. presents a case study of a cloud through four stages of existence as observed by stationary lidar, several satellite products, reanalysis data, and trajectory modeling. The paper excellently links signals in the data with meteorological phenomena at lead to the formation and dissipation of an ice cloud from dust. However, there is a lack of discussion regarding the advantages or disadvantages of this method compared to others. For example, could this multi-source method be used to demonstrate the life cycle of other ice clouds in the region? When comparing the case in the paper to others, are any of the data sources relatively unimportant? Because of this, I recommend this paper be accepted, though with potentially major revisions to address the questions above.

Response: We sincerely appreciate your positive evaluation of our data analysis and phenomenological descriptions. The three specific questions you raised are crucial for perfecting our methodological evaluation. Accordingly, we have supplemented the conclusion section of our revised manuscript.

1、 Advantages and limitations: The core advantage lies in the perfect complementation between "vertical microscopic dynamics" and "horizontal macroscopic thermodynamics/phase states". The CDWL provides extremely high vertical spatiotemporal resolution, which is completely unattainable for meteorological satellites. Conversely, satellite and reanalysis data compensate for the CDWL's inability to directly acquire large-scale cloud phase states and three-dimensional thermodynamic fields. The disadvantage is that CDWL, as a fixed-point Eulerian observation method, relies heavily on large-scale cloud systems passing horizontally over the observation station. Additionally, CDWL cannot directly obtain cloud phase states and heavily relies on cross-validation from satellite observations.

2、 Applicability to other ice clouds: This multi-source framework can definitely be used to demonstrate the life cycle of other ice clouds in the region. As described in Section 2.2, as long as the cloud system undergoes advection across the observation station, CDWL can record the spatial development characteristics of the cloud system at different times like a "slice". This method is particularly suitable for relatively enclosed topographies like the Taklamakan Desert where weather systems exhibit somewhat predictable movement patterns.

3、 Relative importance of data sources: In this case and future similar studies, no single data source is completely "unimportant", but their relative importance changes significantly at different stages of the ice cloud life cycle. For instance, in the pre-formation stage (Stage 1-2), HYSPLIT trajectories and MERRA-2 reanalysis data are vital for tracing the source and climbing dust aerosols. The following table lists the irreplaceable functions of various types of data.

Table 1: Table of irreplaceable functions of various data

| Data Source | Key Irreplaceable Functions |
|--------------------------------------|---|
| CDWL | Vertical structure, turbulence, cloud base height, ice virga detection |
| FY-4A/Himawari-8 | Cloud phase, horizontal extent, cloud top temperature |
| ERA5 | Temperature and humidity profiles, large-scale wind fields, ice water content |
| MERRA-2 | Dust source regions, transport pathways, deposition amount |
| HYSPLIT | Trajectory verification, source region confirmation |
| Ground-based meteorological stations | Near-surface validation, visibility |

Change: Line 290-298. To capture such complex interactions, this multi-source approach synergizes CDWL's high vertical resolution with satellites' broad horizontal coverage, capturing both microphysical dynamics and macro-scale thermodynamics. However, CDWL's fixed-point Eulerian nature requires

clouds to advect overhead and relies on auxiliary data for phase classification. While applicable to other Taklamakan ice clouds, the importance of specific datasets shifts dynamically: HYSPLIT and MERRA-2 are crucial for initial dust tracking, whereas CDWL becomes essential later for observing localized ice virga evaporation. In view of this, our future research plans to integrate polarization lidar and weather radar (Ma et al., 2015; Yin et al., 2021; Wu et al., 2015; Yuan et al., 2022b). This will enable precise classification of clouds and aerosols, thereby deepening our understanding of ice cloud formation mechanisms in desert environments.

Minor/technical comments:

1、As far as I am aware, the name of the desert is typically romanized as "Taklamakan". "Ice crystal cloud" can simply be written as "ice cloud"

Response: The term 'ice cloud' is a more rigorous meteorological classification. Additionally, although both 'Taklamakan' and 'Taklimakan' are frequently used, 'Taklamakan' remains the most universally accepted English spelling. We have carefully revised the manuscript based on your valuable comments.

Change: Line 1-2. **Study on the life cycle of an ice cloud system over the Taklamakan desert using multi-source data**

Line 13. on the southern edge of the Taklamakan Desert.

...

2、There are several cases, such as in the title and abstract, where the indefinite article "an" is missing from before "ice crystal cloud".

Response: We deeply appreciate your meticulous review. Based on your feedback, several sentences have been modified to improve clarity.

Change: Line 1-2. **Study on the life cycle of an ice cloud system over the Taklamakan desert using multi-source data**

Line 12. and decomposition of an ice cloud event

...

3、In lines 80-85, references are given both in-text and parenthetically in each sentence. Only one of the two is necessary.

Response: Thank you for pointing this out. We have corrected the redundant citations in lines 80-85 in the revised manuscript.

Change: Line 79-84. To ensure the reliability of the data, the CDWL data with a CNR less than -17 dB were filtered out. Banakh et al. described the calculation method and error analysis of turbulent kinetic energy dissipation rate (TKEDR) (Banakh et al., 2017). Li et al. introduced the inversion of backscattering coefficient (BSC) in detail (Li et al., 2023). Wei et al. analyzed the error of wind speed and wind direction (Wei et al., 2019). The planetary boundary layer height (PBLH) is calculated by using the TKEDR method (Wang et al., 2021). Su et al. provided a detailed explanation of the parameter inversion method in the article (Su et al., 2024a).

4、In section 3.1.2, it is unclear how mixed phase is differentiated from a combination of ice, mixed, and supercooled. See Fig. 3 at -20 to -15 C, for example.

Response: Thank you for raising this critical point. To clarify, we detail the classification mechanism of the "mixed phase" from two distinct perspectives: the pixel-level algorithmic retrieval and the statistical meaning of the figure:

1、Pixel-level retrieval mechanism: The FY-4A cloud phase product identifies phases based on brightness temperature differences across multi-channel infrared bands (specifically 8.5, 10.8, and 12.0 μm). For a single 4-km spatial resolution pixel: if it exhibits solely the radiative signatures of ice crystals

or supercooled water, it is classified as "Ice" or "Supercooled" phase, respectively. However, if the pixel contains the dual optical signals of both water and ice, the algorithm categorizes it as an independent "Mixed phase."

2、 Statistical meaning of figure 3 (-20°C to -15°C): Figure 3 displays the temporal statistical probability of all pixels passing over the observation site during the entire study period (February 5-6), rather than a snapshot of the internal cloud profile at any single moment. Consequently, the co-occurrence of three phases within this temperature range signifies that, throughout the entire observation period, pure supercooled water pixels, mixed-phase pixels, and pure ice-phase pixels were independently recorded under these specific temperature conditions.

Change: Line 147-148. Fig. 3 shows the statistical probability distribution of these pre-classified cloud phase pixels.

Line 149-150. within the -10 °C to -17.5 °C, individual pixels of supercooled, mixed, and ice phases all occur statistically, indicating that the ice cloud is formed under relatively warm temperature conditions.

5、 Line 175: The HYSPLIT results are in Fig. 7, not Fig. 6.

Response: Thank you for catching this error. We have corrected it in the revised manuscript.

Change: Line 181. Fig. 7 shows the 24 h HYSPLIT backward trajectory results of dust transport. Each stage of the ice cloud life cycle is analyzed as follows.

6、 Line 188: Change "proves" to "supports".

Response: Thank you for catching this error. We have corrected it in the revised manuscript.

Change: Line 194. which supports that there is a dust layer here.

7、 Line 200: The units of dust emission contain "ug" instead of "µg".

Response: Thank you for catching this error. We have corrected it in the revised manuscript.

Change: Line 207. Fig. S4k-1 is also greater than $0.5 \text{ ug m}^{-2} \text{ s}^{-1}$

8、 Lines 215 - 216, 239: How can you be sure about this assertion? Generally, ice clouds formed from heterogeneous nucleation, as is the case here, have fewer crystals relative to those produced by homogeneous freezing. Is it not possible that there are actually a smaller number of crystals that are growing rapidly by vapor deposition to the point of precipitating? Collisions and aggregation would not be necessary, especially considering that the part of the cloud has dendritic growth zone conditions. Could the aggregation hypothesis be supported by the elevated TKEDR in the cloud, or perhaps by the Doppler velocity of the virga is higher than expected for individual crystals?.

Response: We deeply appreciate your insightful comments regarding the microphysical processes of the ice cloud. Your assertion is highly accurate; since the ice crystals in this case formed via heterogeneous nucleation, their number concentration is indeed likely lower than that produced by homogeneous freezing. We completely agree that in such an environment, especially within the dendritic growth zone conditions (-10°C to -17.5°C)—rapid growth via vapor deposition plays a dominant role in developing precipitating crystals. However, we still believe that collision and aggregation occurred simultaneously, and this hypothesis is directly supported by your excellent suggestion regarding the Doppler velocity. According to our CDWL observations, the maximum falling velocity of the ice virga reached 1.38 m/s. Individual pristine ice crystals (like dendrites) typically have much lower terminal velocities (around 0.3 to 0.5 m/s). The observed high fall speed of 1.38 m/s strongly indicates the presence of larger aggregates formed by collisions. Thank you for guiding us to look deeper into the Doppler velocity data. We have revised the manuscript to emphasize the primary role of vapor deposition while using the fall speed to justify the occurrence of aggregation.

Change: Line 239-241. the gradual thickening of the cloud layer provides a favorable environment for

rapid crystal growth via vapor deposition, and simultaneously increases the probability of collision and aggregation of ice crystals, thus creating conditions for ice phase precipitation in Stage-4.

Line 394-402. It is obvious that the large-grained ice crystals, primarily grown by rapid vapor deposition and subsequent aggregation, rush down from 3 km to 1 km under the function of gravity conditions, forming an ice virga. The aggregation process is firmly supported by the maximum falling velocity of the ice virga (1.38 m/s), which significantly exceeds the typical terminal velocity of individual pristine ice crystals.

9、 Figure 6: Perhaps it would be more useful to show the relative humidity with respect to ice in panel (h). Also, the caption refers to the figure as number 5 twice in line 229.

Response: We sincerely thank the reviewer for suggesting the use of Relative Humidity with respect to ice (RH_i), which we agree is highly relevant for ice cloud microphysics. While we considered this conversion, ERA5 only provides a standard "Relative humidity" product. However, according to the ECMWF Integrated Forecasting System (IFS) documentation, the calculation of this standard ERA5 Relative humidity inherently accounts for the ice phase: the saturation vapor pressure is evaluated over liquid water for temperatures above 0 °C, over pure ice for temperatures below -23 °C, and uses a quadratic mixed-phase interpolation for temperatures between 0 °C and -23 °C. Since our studied cloud temperatures predominantly range from -10 °C to -35 °C (Fig. 6i), the native ERA5 product naturally reflects ice-phase thermodynamics. To maintain the dataset's thermodynamic consistency and avoid artifacts from manual empirical conversions, we retained the original product in Figure 6h.

Regarding the typographical error in the caption, we sincerely apologize for the oversight. We have corrected "Figs. 5a and 5b" to "Figs. 6a and 6b" as you correctly pointed out. Thank you again for enhancing the rigor of our manuscript.

Change:

Line 98-100. It should be noted that according to the ECMWF formulation, standard ERA5 relative humidity is calculated with respect to water above 0 °C, ice below -23 °C, and uses mixed-phase interpolation in between.

Line 217. In Figs. 6a and 6b, the dotted boxes A and B represent the existing dust layer and the upward transport process of dust aerosol, respectively

10、 Line 237: This is the first time virga is defined in the paper, but it was discussed earlier in the previous paragraph.

Response: Thank you for catching this. To ensure a better logical flow, we have removed the premature mention of 'virga' from the earlier paragraph. As a result, the term is now first introduced and immediately defined in Stage-4.

Change: Line 228. the laser beam does not penetrate the ice cloud (Wu and Yi, 2017; Cheng and Yi, 2020)

11、 The header for the Conclusions section should be numbered 4.

Response: Thank you for pointing this out. We have corrected the section number for the Conclusions to 4 in the revised manuscript.

Change: Line 276. **4 Conclusions**