

Direct Estimation of Wildfire Emissions at High Latitudes from Combined Polar Orbiter FRP and Sentinel-5P CO Data

William M. Maslanka^{1,2}, Martin J. Wooster^{1,2,3}, Zixia Liu^{1,2}, Jiangping He^{1,2}

¹Department of Geography, King's College London, London, WC2R 2ND, UK

²National Centre for Earth Observation (NCEO), London, UK

³Leverhulme Centre for Wildfires, Environment and Society, London, UK

Correspondence to: William M. Maslanka (william.maslanka@kcl.ac.uk)

Abstract. High Latitude (HL) landscape fires are an important source of greenhouse gases and aerosols, with growing significance under rapid anthropogenic climate change-induced warming. Current fire emission inventories are mostly ‘bottom-up’ in nature; combining, or relying on linear regressions between, satellite remote sensing data and process-based model outputs. However, these methods rely on uncertainties surrounding fuel load and combustion completeness. Here, we adapt the ‘top-down’ Fire Radiative Energy Emission (FREM) approach for HL fires (HLFREM), linking Fire Radiative Energy (FRE) directly to emissions via coefficients derived solely from satellite observations. We derive biome-specific emission coefficients by combining Fire Radiative Power (FRP) from GFAS v1.4 with TROPOMI Total Column Carbon Monoxide plume observations, for the HL’s four most fire-prone biomes; Deciduous and Evergreen Needleleaf Forests, Grasslands, and Shrublands. By applying these coefficients to daily GFAS v1.2 FRE totals (2003-2024), we estimate CO and total carbon emissions across the HL using HLFREM. HLFREM-derived CO emissions were found to be in good temporal and spatial agreement, but smaller in emission totals than generally agree with other widely used inventories (GFAS v1.2, ~~FEERv1.0 GFASv1.2~~, and GFEDv4.1s) in forested biomes, as well as with the MODIS-based FEER approach, with annual average differences of ~~-3254%~~ to ~~-4362%~~ smaller for Deciduous Needleleaf Forests, and ~~-2859%~~ to ~~-4370%~~ smaller for Evergreen Needleleaf forests. For Shrublands and Grassland biomes, HLFREM estimates are ~~3465-4375%~~ and ~~6171-8085%~~ lower respectively. Total carbon emissions, using Emission Factors, were found to show consistent patterns with CO across all biomes. ~~Our results represent the first HL fire emissions dataset based only on satellite data of a major carbon containing gas (CO) emitted by fires and the rate of fire radiative energy release.~~

1. Introduction

Landscape fire is Earth’s largest natural disturbance agent, burning, according to the latest satellite-derived datasets, an average of approximately 5.5% of Earth’s land surface annually (Chen et al., 2023). These landscape fires greatly affect ecology, land carbon stores, atmospheric composition, air quality, and human health. The true magnitude of these effects may even be higher than currently estimated, due to satellite-based burned area mapping often missing many of the highly-numerous smaller burns that together make up a significant fraction of total global burned area (Ramo et al., 2021), even when adjusted for some of this low bias (Chen et al., 2023).

Satellite data indicates that vegetation fires are extensive across every vegetated continent apart from Antarctica, including even High Latitude (HL, $\geq 60^\circ\text{N}$) areas (Jones et al., 2022) up to 75°N (Masrur et al., 2018). Such HL burns include so-called ‘Arctic-fires’ (i.e. fires taking place at latitudes above 66°N) which are of increasing interest, in part because evidence suggests an increase in both their frequency and magnitude; potentially driven by the rapid warming of Earth’s climate occurring in

37 more northerly regions (McCarty et al., 2021). These Arctic-fires, for example those in Siberia in 2020 and 2021 (Kharuk et
38 al., 2021; Liu et al., 2022) can be seen as an extension of the wider northern (i.e. Boreal) regions' fire activity (McCarty et al.,
39 2021), which is itself apparently increasing (Masrur et al., 2018). This is most commonly attributed to lengthening of the 'fire
40 season', the period of the year when vegetation, litter and/or organic soils are sufficiently warm and dry for significant
41 landscape fires to ignite and spread.

42 Frequent lightning, and the extensive uninterrupted fuel (primarily forest) cover, promotes fire in many HL areas; with
43 certain tree species having evolved to include fire as part of their reproductive cycle (Hodges et al., 2021; Hutto, 2008). Such
44 adaptations and their regionally varying occurrence in HL regions are apparently behind many of the spatial differences seen
45 in certain remotely sensed fire signatures (Rogers et al., 2015; Wooster and Zhang, 2004). Because many of the forests of the
46 HL regions ultimately require fire to thrive, wildfires are often left to burn without intervention if they pose little-to-no threat
47 to humans or property. However, there is a concern that the massive stores of ground carbon contained in the underlying peat
48 and organic soils of northern regions, some of which have remained frozen for millennia, may also be becoming more
49 accessible to fire (Davidson and Janssens, 2006; Turetsky et al., 2002). Burning of these carbon stores results in net greenhouse
50 gases (GHGs) emissions that are unreplaced by photosynthetic carbon reassimilation (i.e. vegetation regrowth) on the decades-
51 to-century time-scales typical of forest recovery (Friedlingstein et al., 2021). Quantifying HL fire activity and any trends in its
52 nature is therefore valuable for understanding fires contribution to net atmospheric greenhouse gas concentrations (Mekonnen
53 et al., 2022), as well as for issues such as air quality (Warneke et al., 2023).

54 Calculating the amount of land carbon (i.e. live vegetation, litter and, organic soil) consumed by fire, including in HL fires,
55 as well as emissions of Carbon Monoxide (CO) and other GHGs, reactive gases, and aerosols, typically involves a so-called
56 "bottom-up" calculation (Crutzen and Andreae, 1990), with the most modern implementations combining modelling and
57 satellite-derived datasets (Ichoku and Ellison, 2014; Kaiser et al., 2012; van der Werf et al., 2010, 2017). The most common
58 implementation, the Global Fire Emission Database (GFED, (Randerson et al., 2017)), is driven by satellite-derived measures
59 of burned area, but is still classed as an 'indirect' approach since it relies heavily on mathematical models and/or in-situ
60 assessments for the fuel load and combustion completeness parameters that remain subject to significant uncertainty at any
61 particular location (Reid et al., 2009). An alternative satellite-based 'direct' approach relies on Fire Radiative Power (FRP)
62 measures rather than burned area, with FRP representing the radiant energy released per unit time by the combustion process.
63 These FRP data can be time-integrated to estimate the Fire Radiative Energy (FRE) released by the burn, which then links to
64 total fuel consumption (Roberts et al., 2005; Wooster et al., 2005, 2015). FRP-based approaches avoid the need for assumed
65 or modelled fuel load and combustion completeness terms (Kasischke and Penner, 2004), but are not always fully independent
66 of the burned area-based methods since the necessary 'FRP-to-fuel consumption' conversion coefficients are themselves often
67 derived from correlations and regressions made between FRE datasets and the outputs of the more indirect burned area-based
68 approach; such as with the Global Fire Assimilation System (GFAS, (Kaiser et al., 2012)). Alternatively, such conversion
69 coefficients have been based on limited numbers of direct comparisons between FRE and fuel consumption, but in mostly
70 small-scale or laboratory burns which may not accurately represent the dynamics of real landscape scale fires, nor the full

71 characteristics of the FRP measurements made from Earth orbit (Freeborn et al., 2008; Nguyen and Wooster, 2020; Wooster
72 et al., 2005).

73 More recently, even more direct “top-down” FRP-based emissions assessment methodologies have emerged; now relating
74 smoke emissions directly to the observed FRP of landscape fires. Their uniqueness is that, in order to reduce the
75 aforementioned issues and embedded assumptions that arise when using satellite-derived FRE totals with ‘FRP-to-fuel
76 consumption’ conversion coefficients, these new approaches use direct satellite-based measurements of fire emissions and
77 FRP, via a set of ‘matchup fires’, to generate a set of landscape-fire ‘emissions coefficients’ that can be used to generate smoke
78 emissions estimates directly from all other FRP observations of the area.

79 These more direct assessments have included both airborne observations of landscape fire and their associated emitted
80 smoke species (Hayden et al., 2022; Owsley-Brown et al., 2024; Stockwell et al., 2022) as well as satellite-based observations
81 (Adams et al., 2019; Griffin et al., 2024; Mebust et al., 2011); calculating emission coefficients of observed landscape fires
82 categorised in terms of field campaign and land cover type (Griffin et al., 2024).

83 One of the most recent such approaches, termed the Fire Radiative Energy Emissions (FREM) method (Mota and Wooster,
84 2018; Nguyen and Wooster, 2020), derives these emissions coefficients directly from geostationary satellite FRE estimates
85 and matching smoke plume Earth observation data (Fisher et al., 2020; Mota and Wooster, 2018; Nguyen and Wooster, 2020).
86 Emissions coefficients derived in this way are appropriate to real landscape fires of all sizes, minimally reliant on any
87 assumptions, and appropriate to apply to further satellite FRE data to convert them into smoke emissions estimates (Nguyen
88 et al., 2023). Thus far, the FREM approach has relied on geostationary satellites to provide the high imaging frequency FRP
89 data with which to accurately estimate FRE via temporal integration. Observations made by equatorial orbit geostationary
90 satellites are, however, of mostly poor quality at high latitudes. However, whilst FRP-to-FRE calculations based on polar
91 orbiting satellite data can be challenging at lower latitude fires due to the limited temporal sampling of FRP provided (Mota
92 and Wooster, 2018), at higher latitudes orbital convergence generally provides far more polar-orbiting based estimates of FRP
93 per day (Fig. 1).




Figure 1: (a) Mean number of daily overpasses at any Earth location made by MODIS on AQUA and TERRA, and (b) zonal mean number of daily overpasses. Daily averages based on 16 days of data taken between 11/08/24 00:00 UTC and 27/08/24 00:00 UTC, gridded to a regular $0.5^\circ \times 0.5^\circ$ grid. At latitudes where HL fires occur ($60^\circ - 75^\circ\text{N}$, green line and above), there is a latitudinal observation frequency dependence that provides on average between 8 and 18 daily satellite observations from these two systems alone. Each cloud-free observation provides an FRP measure, and which can be used to estimate FRE via temporal integration.

Given that geostationary observations of FRP are unsuitable for fire-prone HL regions, ~~Here~~ we aim to take advantage of the higher latitude, higher temporal resolution sampling of FRP of polar orbiting FRP observations- to develop a HL version of the current FREM approach of Nguyen et al., (2023); generating ~~for the first time~~ a dataset of HL fire carbon and trace gas emissions completely independent of satellite-derived burned area measures and/or of the FRP-to-fuel consumption conversion coefficients influenced by them. These conversion coefficients are also not derived from small-scale experiments, but from Earth observation data of the landscape fires themselves. We compare our fire emissions assessments to those of existing state-of-the-art burned area and FRP based approaches, and assess the characteristics and benefits of our approach with respect to these alternatives. In Sect. 2, we provide the background to the FREM approach. In Sect. 3, we describe its adaptation for HL fires (hereby referring to the method as HLFREM) and using satellite derived polar orbiter FRP data and plume-integrated Total Column Carbon Monoxide (TCCO) observations we derive biome (b) specific emission coefficients for Carbon Monoxide (EC_{CO}^b) and Carbon (EC_C^b), as well as total CO and carbon emissions via their use with FRP time-series data. In

112 Sect. 4, we calculate the total CO and Carbon emissions, using EC_{CO}^b and EC_C^b , and compare them to the fire emissions
113 estimates present in other widely used inventories, before closing with a Summary and Conclusion in Sect. 5.

114 2. FREM Background

115 As detailed in the review of Wooster et al., (2021), both geostationary satellites (Wooster et al., 2015; Xu et al., 2010, 2017,
116 2021) and polar-orbiting satellites (Giglio et al., 2016; Schroeder et al., 2014; Wooster et al., 2012; Xu et al., 2020) offer the
117 capability to observe and quantify wildfire activity using Active Fire (AF) approaches involving FRP retrievals. Well-known
118 inventories using these methods include the Fire Inventory from NCAR (FINN, Wiedinmyer et al., (2011)), the Fire Energetic
119 and Emission Research (FEER) approach (Ichoku and Ellison, 2014)), the Quick Emission Dataset (QFED, Darmenov and da
120 Silva, (2015)), and GFAS (Kaiser et al., 2012). The most recently developed FREM method (Nguyen et al., 2023) directly
121 links satellite-derived FRP data to a fires CO emission rate via a set EC_{CO}^b derived from a ‘matchup fire’ subset of satellite-
122 derived FRP and CO observations. Specifically, estimates of EC_{CO}^b are calculated using FRE and plume-integrated Total
123 Column Carbon Monoxide (TCCO) data collected at a subset of ‘matchup fires’ where both datasets are ‘well-observed’ (e.g.
124 no gaps in the FRP record, and a clear fire emissions CO plume is seen). Once derived, application of EC_{CO}^b to the FRP
125 observations at all the regions fires enables the regions fire-related CO emission rates to be calculated without the need for any
126 other information (Reid et al., 2009). Hence the approach is extremely direct, and with the emissions coefficients derived from
127 the same types of satellite FRP data that they will ultimately be applied to, and thus which are impacted by the same factors
128 such as overstory impacts on FRP measures and minimum FRP detection limits (Nguyen and Wooster, 2020).

129 Each FREM iteration thus far has relied on geostationary FRP data, due to its frequent 10-to-30 min imaging frequency
130 enabling easy derivation of FRE (Wooster et al., 2021). Rather than CO however, the first FREM (v1) iteration focused on
131 deriving aerosol optical depth (AOD) related total particulate matter (TPM) emissions from FRP observations (Mota and
132 Wooster, 2018), an approach then enhanced by Nguyen and Wooster, (2020) and Fisher et al., (2020) through the use of
133 improved AOD datasets. FREMv2 (Nguyen et al., 2023) replaced use of AOD data with trace gas column amounts of CO
134 (from ~~the the~~ S5P (Sentinel-5 Precursor) satellites observations using the TROPOMI (TROPOspheric Monitoring Instrument)
135 instrumentsensor), directly estimating fluxes of CO rather than TPM. CO is the second largest fire-emitted compound after
136 CO₂ (Akagi et al., 2011) and CO emissions factors are significantly less variable than those of TPM (Andreae, 2019). This
137 makes any subsequent estimation of total fuel consumption using the FREM approach more appropriately conducted using
138 CO pathway rather than one based on TPM fire emissions data.

139 3. High Latitude FREM Adaptation

140 ~~The polar-orbiting based HL-FREM framework developed herein builds on the geostationary FREM methods by identifying~~
141 ~~a set of matchup fires for which both the total amount of CO contained in the fire plume is assessed, as well as the total amount~~

~~of FRE released over the time it took the plume to form. As mentioned in Section 2, previous FREM studies have utilised geostationary observations of FRP, due to its high temporal sampling resolution of 10-30 minutes. However, a disadvantage of using geostationary platforms for FRP analysis is that the geostationary pixel areas grow rapidly at locations very far from nadir (> 40° view zenith angle, (Nguyen and Wooster, 2020; Xu et al., 2021)), thus making geostationary platforms unsuitable for HL FRP studies.~~

The polar-orbiting based HLFREM framework developed herein builds on the geostationary FREM methods by identifying a set of matchup fires for which both the total amount of CO contained in the fire plume is assessed, as well as the total amount of FRE released over the time it took the plume to form (Section 3.4). Using these polar-orbiting matchups, a set of EC_{CO}^b are generated using ordinary least squares (OLS) linear best fits to the co-incident FRE and CO data of all matchup fires in that biome. These EC_{CO}^b can then be multiplied by the regionally complete FRP time-series, including potentially hundreds of thousands more fires than are included in the matchup datasets, generating a spatially and temporally comprehensive set of biomass burning CO emissions covering all fires. Dividing these CO emissions by the standard biome-specific Emissions Factors (EFs, (Akagi et al., 2011; Andreae, 2019)) provides estimates of dry biomass consumption and also of total carbon (assuming biomass is $50\% \pm 5\%$ carbon), or via use of EF ratios of any trace gas to CO also estimates of other trace gas emissions totals (Nguyen et al., 2023; Nguyen and Wooster, 2020).

A brief description of the data used for the HLFREM biome specific emission coefficient is shown in Sect. 3.1, with a description of the region of interest in Sect. 3.2. The full selection process for the fire matchups to which this procedure is applied is described in Sect. 3.23; and at each fire the relevant FRE was calculated as described in Sect. 3.34 using as the temporal integration period the time between the early morning fire activity minimum and the S-NPP VIIRS overpass made almost simultaneously to that of S5P (Sect. 3.45). Once sufficient matchup fires existed for each fire biome covered (see Sect. 3.56), we generated EC_{CO}^b for these biomes as detailed in Sect. 3.76. These emission coefficients were then multiplied by the full GFAS v1.2 FRP record (Jan 2003 – Dec 2024, Sect. 4.1), to generate a HLFREM CO emission timeseries. Additionally, via the application of the appropriate EFs, it was also possible to generate a HLFREM Carbon emission timeseries using GFAS v1.2 (Sect 4.2). For both the CO and Carbon emission timeseries, the HLFREM emissions will be compared to pre-existing emission databases; namely FEER, GFAS v1.2, and GFED v4.1s.

3.1. HLFREM Earth Observation Data Used

3.1.1. VIIRS and MODIS multispectral

Satellite observations from the Visible Infrared Imaging Radiometer Suite (VIIRS) onboard the Suomi National Polar-orbiting Partnership (SNPP) and the Moderate Resolution Imaging Spectroradiometer (MODIS) instruments onboard the Terra and Aqua platforms were used in order to manually identify matchup fire locations (Sect. 3.3).

The SNPP satellite was launched in 2011 and operates in a sun-synchronous orbit with equatorial crossing local times of approximately 01:30 and 13:30. The VIIRS instrument provides simultaneous, co-registered measurements at 375 and 750 m spatial resolution across 22 spectral bands spanning the visible to longwave infrared spectrum. These observations support the generation of a wide range of operational environmental products. In this study, fire detections were obtained from the VIIRS active fire product (VNP14, Schroeder et al., 2014), while aerosol loading was characterised using Aerosol Optical Depth (AOD) retrievals produced by the Enterprise Processing System (AOD EPS, NOAA, 2020).

The MODIS instruments onboard Terra and Aqua provide a longer observational record, with launches in 1999 and 2002, respectively. Both instruments operate in sun-synchronous orbits, with initial equatorial crossing local times of 10:30/22:30 for Terra and 01:30/13:30 for Aqua. However, gradual orbital drift has resulted in changes to these nominal observation times over the course of the missions. MODIS provides simultaneous, co-registered observations at 250, 500, and 1000 m spatial resolutions across 36 spectral bands covering the visible through longwave infrared wavelengths. Like with the VIIRS observations, The MODIS active fire products (MOD14 and MYD14, Giglio et al., 2016) were also used for matchup fire identification.

The regionally comprehensive FRP time series used herein is the Global Fire Assimilation System (GFAS v1.2) FRP dataset described in, generated as part of the Copernicus Atmospheric Monitoring Service (CAMS: <https://atmosphere.copernicus.eu/>) and widely used. The GFAS v1.2 record is based on per-pixel FRP retrievals contained within the Terra/Aqua MOD14/MDY14 active fire products, using these to generate daily mean cloud-adjusted FRP data gridded at 0.1° back to almost the start of the MODIS mission.

~~However, whilst the GFAS v1.2 FRP data dating back to 2003 are considered a suitably long FRP time series for generating a long-term CO emissions record, they are less well suited to the initial generation of the EC_{CO}^b coefficients linking FRP to rate of CO emissions since they provide only daily average FRP values. Specifically, the daily average FRP data contained in GFAS v1.2 cannot be used to estimate the amount of FRE emitted by a fire from the night-time FRP minimum up until the early afternoon TROPOMI overpass, since only daily average FRP is provided, and yet this sub-daily FRE value is required at the set of matchup fires to generate the initial EC_{CO}^b emissions coefficients. For this reason, these coefficients were instead derived using the alternative GFAS v1.4, which applies the fire diurnal cycle model of to the 0.1° gridded MODIS FRP used to generate GFAS v1.2 but now to obtain hourly gridded FRP estimates. At the location of each matchup fire, these hourly FRP data can then be integrated over the required time period from the night-time fire minimum up until the S5P (TROPOMI) satellite overpass time to generate the FRE released. Though GFAS v1.4 goes back only to 2019, this is perfectly sufficient for deriving the EC_{CO}^b coefficients.~~

3.1.2. TROPOMI TCCO

The TROPOMI sensor onboard the European Space Agency (ESA) S5P satellite orbits the globe with an equatorial ascending and descending node of approximately 13:30 and 01:30 respectively; very similar to that of the Visible Infrared Radiometer Suite (VIIRS) onboard Suomi National Polar-orbiting Partnership (S-NPP, Jing et al., 2022). Onboard TROPOMI are four spectrometers, covering the ultraviolet-visible, near infrared, and shortwave infrared (SWIR) spectral bands (Veefkind et al., 2012). Amongst other species, such as NO₂, CH₄, and O₃, TCCO is observed, at a spatial resolution of approximately

208 5.5 x 7.0 km (August 2019 to present, 7.0 x 7.0 km prior to August 2019, (Inness et al., 2022)), derived using SWIR
209 observations with the Shortwave Infrared CO Retrieval (SICOR, (Landgraf et al., 2016)) algorithm. The TROPOMI TCCO
210 observations have been validated against both ground-based remote sensing techniques (such as the Total Carbon Column
211 Observing Network, TCCON, (Borsdorff et al., 2018)) and through space-based remote sensing techniques (Martínez-Alonso
212 et al., 2020; Neyra-Nazarrett et al., 2025).

213 The S5P TCCO product includes a quality flag, indicating the potential presence of cloud in the resulting TCCO pixel. For
214 this study, we have used observations with a quality flag of ≥ 0.5 , aligning with the recommended quality threshold (Apituley
215 et al., 2018) and other TCCO studies (Griffin et al., 2024; Nguyen et al., 2023; Rowe et al., 2022). The impact of using only
216 pixels with a quality flag of ≥ 0.5 was small, given that the quality flags in the TCCO product were carried out in clear-sky
217 only conditions; a requirement of the FREMv2 methodology (Borsdorff et al., 2018; Nguyen et al., 2023).

218 **3.1.3. Global Fire Assimilation System (GFAS) FRP**

219 The regionally comprehensive FRP time-series used herein, replacing the geostationary FRP time-series used in FREMv2,
220 is the Global Fire Assimilation System (GFAS v1.2) FRP dataset described in (Kaiser et al., 2012), generated as part of the
221 Copernicus Atmospheric Monitoring Service (CAMS: <https://atmosphere.copernicus.eu/>) and widely used (e.g. Di Giuseppe
222 et al., 2018; Inness et al., 2019; Popovicheva et al., 2022) The GFAS v1.2 record is based on per-pixel FRP retrievals contained
223 within the Terra/Aqua MOD14/MDY14 active fire products (Giglio et al., 2016), using these to generate daily mean cloud-
224 adjusted FRP data gridded at 0.1° back to almost the start of the MODIS mission (Kaiser et al., 2012). At the high latitudes,
225 the GFAS FRP record will include more FRP observations from the polar-orbiting MODIS platform, due to orbital
226 convergence (Figure 1).

227 However, whilst the GFAS v1.2 FRP data dating back to 2003 are considered a suitably long FRP time-series for generating
228 a long-term CO emissions record, they are less well suited to the initial generation of the EC_{CO}^b coefficients linking FRP to rate
229 of CO emissions since they provide only daily average FRP values. Specifically, the daily average FRP data contained in
230 GFAS v1.2 cannot be used to estimate the amount of FRE emitted by a fire from the night-time FRP minimum up until the
231 early afternoon TROPOMI overpass, since only daily average FRP is provided, and yet this sub-daily FRE value is required
232 at the set of matchup fires to generate the initial EC_{CO}^b emissions coefficients. For this reason, these coefficients were instead
233 derived using the alternative GFAS v1.4 implementation (Kaiser et al., 2024), which applies the fire diurnal cycle model of
234 (Andela et al., 2015) to the 0.1° gridded MODIS FRP used to generate GFAS v1.2 but now to obtain hourly gridded FRP
235 estimates. A comparison of a GFASv1.2 and GFASv1.4 global FRE estimates for 2020 showed that they were in good
236 agreement with each other (Fig. 2), with a coefficient of determination (r^2) of 0.908, and a mean bias of 0.13 PJ, indicating
237 negligible systematic differences between the two datasets, and an RMSE of 3.53 PJ and a standard deviation of 3.53 PJ,
238 reflecting variability in the FRE estimates between the two datasets, due to the inclusion of a diurnal FRP model into
239 GFASv1.4.




Figure 23: (Randerson et al., 2017) (a) Daily Global FRE estimated from GFASv1.2 (black) and GFASv1.4 (red) throughout 2020. (b) Scatter of Daily global FRE estimates from GFASv1.2 and GFASv1.4.



At the location of each matchup fire, these hourly FRP data can then be integrated over the required time period from the night-time fire minimum up until the S5P (TROPOMI) satellite overpass time to generate the FRE released. Though GFAS v1.4 goes back only to 2019, this is perfectly sufficient for deriving the EC_{CO}^b coefficients.

~~The full selection process for the fire matchups to which this procedure is applied is described in Sect. 3.2; and at each fire the relevant FRE was calculated as described in Sect. 3.3 using as the temporal integration period the time between the early morning fire activity minimum and the S NPP VIIRS overpass made almost simultaneously to that of S5P (Sect. 3.4). Once sufficient matchup fires existed for each fire biome covered (see Sect. 3.5), we generated EC_{CO}^b for these biomes as detailed in Sect. 3.6. These emission coefficients were then multiplied by the full GFAS v1.2 FRP record (Jan 2003 – Dec 2024, Sect. 4.1), to generate a HLFREM CO emission timeseries. Additionally, via the application of the appropriate EF_s, it was also possible to generate a HLFREM Carbon emission timeseries using GFAS v1.2 (Sect 4.2). For both the CO and Carbon emission timeseries, the HLFREM emissions will be compared to pre-existing emission databases; namely FEER, GFAS v1.2, and GFED v4.1c.~~

3.21. Region of Interest Selection

HL fire is seasonal in nature, dominantly present in the Northern Hemispheric Summer months (June, July, August, Fig. 32a). Four HL ROIs and matching time periods were used from within this period to generate the set of matchup fire, and each of the ROI extractions focused on a period when fire activity was maximised in that ROI, as shown in Figs. 32b and 23c, and detailed in Table 1. Landcover data, natively at 300 m grid resolution, taken from the ESA Climate Change Initiative (ESA-CCI) 2018 land cover map (ESA, 2017) was used to classify biomes within the ROI's, based on 11 'fire-biome' classes aggregated from the original 37 landcovers (Table A1), as shown in Fig. 32b. The two ROIs in Boreal Asia (HL-BOAS-20

264 and HL-BOAS-21) were dominated by the Deciduous Needleleaf Forest fire biome, whereas those in Boreal North America
265 (HL-BONA-19 and HL-BONA-23) were dominated by Evergreen Needleleaf Forest fire biome.



268 **Figure 23: HL fire information. (a) Mean monthly FRE (2003 – 2024) for HL fires ($\geq 60^\circ\text{N}$) in the regions used by the Global Fire**
269 **Emissions Database (GFED; (Randerson et al., 2017), BONA: Boreal North America, EURO: Europe, BOAS: Boreal Asia), as derived**
270 **from GFAS v1.2. (b) HL regions used for the mean monthly FRE estimation in (a), as defined by GFED. (cb) ROI and aggregated**
271 **fire biome map, as derived from the CCI Land Cover 2018 map (ESA, 2017) and with biome aggregation detailed in Table A1. (ed)**
272 **The four ROIs used in this study within which the 833 fire-matchup (shown as crosses) were identified.**

273 **Table 1: Acronyms and Geographic bounding boxes of ROIs, as shown in Fig. 23c.**

ROI Full Name	ROI Acronym	Bounding Box
High Latitude Boreal North America 2019	HL-BONA-19	60° N - 70° N, 105° W - 165° W
High Latitude Boreal Asia 2020	HL-BOAS-20	60° N - 75° N, 100° E - 175° E
High Latitude Boreal Asia 2021	HL-BOAS-21	60° N - 70° N, 100° E - 140° E
High Latitude Boreal North America 2023	HL-BONA-23	60° N - 70° N, 100° W - 140° W

274 3.32. Data-Used-and-Plume Identification

275 Within each of the four ROIs shown in Figs. 2b and 2c, the potential cloud-free matchup fires were manually identified and
276 examined via visual inspection of multispectral and AF Products detailed in Table 2, based on the following criteria:

- 277 • Clear smoke emission visible in the VIIRS true colour (Red: M05, Green: M04, Blue: M03) imagery, and without
278 apparent mixing with other nearby smoke plumes.
- 279 • Clear visual indication of a developed smoke plume, spread by the wind, in the same VIIRS true colour imagery.
- 280 • Clear identification of a region of shortwave infrared thermal emission (hereby referred to as the “burning area”)
281 resulting from active combustion, as seen in a VIIRS false colour (Red: M11, Green: I02, Blue: I01) imagery.
- 282 • AF Pixels present within the “burning area” as deduced from the VIIRS VNP14 and MODIS MOD14/MYD14 AF
283 Products.


284 **Table 2: Satellite data products used during the manual inspection and digitization of matchup fire polygons.**

Parameter	Satellite	Product Code	Resolution (nadir)
VIIRS Moderate Resolution L1B Calibrated Radiances	S-NPP	VNP02MOD	750 m x 750 m
VIIRS Moderate Resolution L1B Terrain Corrected Geolocation	S-NPP	VNP03MOD	750 m x 750 m
VIIRS Imagery Resolution L1B Calibrated Radiances	S-NPP	VNP02IMG	375 m x 375 m
VIIRS Imagery Resolution L1B Terrain Corrected Geolocation	S-NPP	VNP03IMG	375 m x 375 m
VIIRS Thermal Anomalies/Fires	S-NPP	VNP14v002	750 m x 750 m
MODIS Thermal Anomalies and Fire	TERRA	MOD14v001	1.0 km x 1.0 km

MODIS Thermal Anomalies and Fire	AQUA	MYD14v001	1.0 km x 1.0 km
VIIRS Enterprise Processing System Aerosol Optical Depth	S-NPP	AOD EPS	750 m x 750 m
TROPOMI Level 2 Total Column Carbon Monoxide	Sentinel-5P	L2__CO__	7.0 km x 3.5 km

285

286 Each potential matchup fire then had its smoke plume and associated burning area extent digitized and examined; based on
287 manual interpretation of the VIIRS multispectral imagery, matching VIIRS AF and Aerosol optical depth (AOD) products
288 (NOAA, 2020), and TROPOMI TCCO products, as shown in Fig. 43.




289

290 **Figure 34:** Example of a matchup fire imaged in Siberia (62.39° N, 167.49° E) on 28/07/2020 at 01:42 UTC (13:42 PETT). The
291 digitised outline of the plume (cyan) and burning area (red) are shown superimposed on (a) RGB (M05-M04-M03) imagery via S-
292 NPP (VIIRS), (b) AOD (processed via EPS) from S-NPP (VIIRS), (c) Sentinel-5P TCCO data, (d) False Colour Composites (M11-
293 I02-I01) via S-NPP (VIIRS), (e) VIIRS Active Fire Product (VNP14) from S-NPP (VIIRS). Green: Non burning vegetation. Yellow:
294 Active Fire Hotspot, Blue: Water, White: Cloud, Black: No Data. (f) Geographic location and extent of the plots shown (red).

295 3.43. Fire Radiative Energy Estimation

296 To calculate the FRE for each matchup fire, each digitised burning area polygon was converted to a binary mask and used
297 to extract the FRP time-series from the hourly-timestep FRP record held within GFAS v1.4 (see example in Fig. 54) for all
298 GFAS v1.4 pixels that intersect with or are contained by the burning area polygon. HL orbital convergence ensured the
299 calculation was based on significantly more polar-orbiting observations than would be the case at lower latitudes (Fig. 1), and,

300 following (Nguyen et al., 2023), the FRP integration took place from the 06:00 hrs local time fire activity minima (Fig. A1)
 301 up until the time of the almost simultaneous early afternoon S5P and Suomi S-NPP overpasses. Mean FRP record length for
 302 all matchup fires was 7.3 hrs (minimum 4 and maximum 11 hours), with the variation due to the orbital convergence (Fig. 1)
 303 and this results in the match-up fires being observed by Sentinel-5P up to three times per day from different ascending
 304 overpasses.
 305



306
 307 **Figure 54:** Example of the hourly FRP time-series from GFAS v1.4, used to generate an FRE measure for a single matchup fire (i.e.
 308 the match-up fire in Fig. 43, as shown above). The hourly FRP timeseries data were generated from the set of individual FRP
 309 observations provided by MODIS on AQUA and TERRA, using the diurnal cycle model of Andela et al., (2015). Additionally, the
 310 VIIRS observations on S-NPP used for fire-matchup identification are also shown. Times of the individual satellite overpasses, as
 311 well as of the S5P overpass time used to provide the TCCO data for this fire-matchup, are indicated.

312 3.45. Excess CO Estimation

313 For each fire-matchup, the smoke plume polygon with a surrounding buffer of one pixel was also used to generate a binary
 314 mask employed to extract the relevant S5P TROPOMI TCCO data within the plume for a given S5P overpass, along with a
 315 measure of the ambient background TCCO ($TCCO_{BG}$, taken as the minimum TCCO within the buffered mask). Total
 316 excess plume CO (CO_{EX}) was then calculated for each fire-matchup, as the sum of all CO_{EX} from each S5P pixel associated
 317 with each fire-matchup, using Eq. (1).

$$318 \quad CO_{EX} = \sum (TCCO_M - TCCO_{BG}) \cdot A \cdot M_{CO} \quad (1)$$

319 where $TCCO_M$ is the TCCO within the buffered binary mask (units: mol m^{-2}), $TCCO_{BG}$ is the ambient background TCCO
 320 (units: mol m^{-2}), A is the S5P pixel size (units: m^2) calculated from the geographic coordinates of the pixel corners, and M_{CO}
 321 the molecular weight of CO (units: g mol^{-1}). Due to orbital convergence, a single fire can be observed multiple times by
 322 Sentinel-5P. Each individual overpass is treated as an independent observation. As a result, a single landscape fire event may
 323 correspond to multiple matchup fire events within the matchup dataset, each associated with a different estimate of excess CO
 324 (due to overpasses occurring approximately 100 minutes apart).


325 3.56. Fire Biome Estimation

326 All MODIS and S-NPP AF pixels within the fires burning area polygon and timed during the FRP temporal integration
 327 period had their fire-biome determined using the aggregated biome data shown in Fig. 32b. For a single fire if $\geq 50\%$ of the

328 AF pixels were from a single fire-biome then that was set as the fires “dominant” fire-biome. Fires having no such dominant
329 class were discarded, as their derived emission coefficients could not be properly allocated to a fire-biome.

330 **3.67. Biome Specific Emission Coefficient Generation**

331 In total, 833 individual matchup fires were initially identified across the four regions of interest (Table 1), in their respective
332 Northern Hemispheric months, with 125 of those being discarded as having no dominant fire-biome. Of the 708 fires
333 remaining, 468 were located in Deciduous Needleleaf Forest (DecNeedle), 186 in Evergreen Needleleaf Forests (EGNeedle),
334 26 in Grassland, 22 in Shrubland, and six in a fire-biome described as ‘Sparee’Sparse. The locations of the matchup fires for
335 the four most well-sampled fire-biomes identified (DecNeedle, EGNeedle, Grassland, and Shrubland) are mapped in Fig. 32c.
336 Analysis of the mean annual FRE totals from GFAS v1.2 (Fig. 65, Table 3) shows that these four biomes are responsible for
337 the vast majority (~93%) of the total FRE generated by HL fires.



338
339 **Figure 65: Mean annual HL FRE (a) totals (PJ), and (b) contribution (%) of each of the four most well sampled biomes identified**
340 **as part of the fire-matchups database, as well as from additional biomes, taken from GFAS v1.2 (2003-2024).**

341 **Table 3: Mean Annual FRE totals (PJ) and contribution (%) for each of the four most well sampled biomes, compared to all other**
342 **HL biomes, calculated using GFAS v1.2, between 2003 and 2024.**

HL Biome	Mean Annual FRE Total (PJ)	Standard Deviation (PJ)	Mean annual FRE Contribution (%)	Standard Deviation (%)
All	205.4	115.5	100	0.0
DecNeedle	100.2	106.9	43.0	24.7
EGNeedle	67.8	59.9	34.2	19.1
Grassland	17.9	12.4	11.6	8.4

Shrubland	6.7	6.0	4.2	4.1
All other biomes	12.8	9.9	7.0	3.3

Following Nguyen et al., (2023), the FRE and Excess CO data of each fire-biomes matchup fires were used to derive a set of EC_{CO}^b emissions coefficients via zero-intercept OLS linear regression (Fig. 76); with results reported in Table 4. As detailed in Nguyen and Wooster, (2020), “FEER-equivalent” EC_{CO}^b values are also reported therein, calculated from the grid cell (not biome) based FREP-based total particulate matter (TPM) emissions coefficients (EC_{TPM}^b) reported in Ichoku and Ellison, (2014) (available at www.feer.gsfc.nasa.gov/data/emissions/) and the relevant biomes TPM-to-CO EF ratios (Nguyen and Wooster, 2020). The HLFREM EC_{CO}^b values reported in Fig. 7 and Table 4 are similar in magnitude to the emission coefficients reported in Griffin et al., (2024), who developed global emission coefficients using a mass balance approach to determine a relationship between TROPOMI observations of CO and MODIS observations of FRP, according to different biome classifications. Whilst the GFEDv4 biome classification is not limited to HL regions only, the similarity in the forested HLFREM EC_{CO}^b values to the (Griffin et al., 2024) BOREF EC_{CO}^b is encouraging.

Table 4: Emissions coefficients for CO (EC_{CO}^b) as derived in Fig. 7, along with FEER-equivalent values for comparison and those for carbon (EC_C^b) calculated using EC_{CO}^b and EFs from (Andreae and Merlet, 2001) and (Akagi et al., 2011) - labelled -AM, and -AG respectively, in Sect. 4.2. Also shown are EC_{CO}^b for GFEDv4 biomes (in brackets), as derived by (Griffin et al., 2024).

HLFREM Biome	Sentinel-5P TCCO-derived EC_{CO}^b [g MJ ⁻¹]	FEER-equivalent EC_{CO}^b [g MJ ⁻¹]	Sentinel-5P TCCO-derived EC_{C-AM}^b [g MJ ⁻¹]	FEER-equivalent EC_{C-AM}^b [g MJ ⁻¹]	Sentinel-5P TCCO-derived EC_{C-AG}^b [g MJ ⁻¹]	Griffin et al., (2024) EC_{CO}^b [g MJ ⁻¹]
DecNeedle	81.54	198.3	367.51	893.7	298.55	91 ± 2 (BOREF)
EGNeedle	66.74	169.5	300.81	764.1	244.36	91 ± 2 (BOREF)
Grassland	53.78	198.3	242.39	893.7	196.91	60 ± 1 (SAVA)
Shrubland	53.14	151.7	239.51	683.8	194.56	60 ± 1 (SAVA)


Analysis comparing the spatial distribution of FRP across the entire Northern Hemispheric Summer (JJA) HL GFASv1.4 dataset with those relating to the matchup fires can be found in Fig. 8. It can be seen that the matchup fires are spatially well sampled across the JJA GFASv1.4 dataset (Fig. 8a, 8b, and 8d), with matchup fires being found across the fire-prone HL-BONA region, and the eastern half of the fire-prone HL-BOAS region ($>100^\circ\text{E}$). Additionally, it can be seen the matchup fires tended to be comprised of a greater proportion of higher FRP observations (Fig. 8e), which did not have a zonal or a longitudinal bias in them, as the zonal and longitudinal mean FRP of the matchup fires were consistently higher than those of the JJA GFASv1.4 dataset (Fig. 8c and Fig. 8e). A Mann-Whitney U test was performed on the matchup fire FRP values (containing $n = 73951$ pixels, median FRP: 76.5 MW) and remaining JJA GFASv1.4 FRP values (containing = 4414203 pixels, median FRP: 11.4 MW), showing that the distributions differed significantly ($U = 7.92 \times 10^{10}$, $p < 0.001$, rank-biserial correlation = 0.515), indicating preferable sampling of higher FRP values. This is to be expected, as the plume identification criteria (Sect. 3.3) requires a clear visual indication of a developed smoke plume with an associated area of active combustion; the likelihood of which increases with increasing FRP, when occurring under the same conditions. Smaller fires are likely not going to produce an optically visible smoke plume, and are thus unlikely to result in selection for digitization. Fires that occur

underneath pre-existing smoke plumes are also not going to be picked for plume digitization, as the CO enhancement could not be accredited to an individual smoke plume.

An OLS regression was selected because the uncertainties associated with the excess CO and FRE estimates for a given matchup fire data point included potentially unmeasured sources of variability. Although both variables contain measurement and retrieval uncertainties (Griffin et al., 2024; Rowe et al., 2022), these uncertainties do not represent all sources of variability affecting the relationship between excess CO and fire activity, with some contributions of uncertainty being unquantifiable (Nguyen et al., 2023). Due to this, a regression method that is driven by data point uncertainty (such as orthogonal distance regression) was deemed unsuitable for use.

Table 4: Emissions coefficients for CO (EC_{CO}^b) as derived in Fig. 7, along with FEER-equivalent values for comparison and those for carbon (EC_C^b) calculated using EC_{CO}^b and EFs from and — labelled — AM, and — AG respectively, in Sect. 4.2.

HLFREM Biome	Sentinel-5P TCCO-derived EC_{CO}^b [g MJ ⁻¹]	FEER-equivalent EC_{CO}^b [g MJ ⁻¹]	Sentinel-5P TCCO-derived EC_{CO}^b [g MJ ⁻¹]	FEER-equivalent EC_{CO}^b [g MJ ⁻¹]	Sentinel-5P TCCO-derived EC_{CO}^b [g MJ ⁻¹]
DecNeedle	122.88	198.3	553.84	893.7	449.90
EGNeedle	121.30	169.5	546.71	764.1	444.12
Grassland	69.02	198.3	311.08	893.7	252.71
Shrubland	105.20	151.7	474.2	683.8	385.17






Figure 76: HLFREM biome-specific CO emission coefficients (EC_{CO}^b , in g MJ^{-1}) derived from the set of matchup fires for (a) Deciduous Needleleaf Forests, (b) Evergreen Needleleaf Forests, (c) Grasslands, and (d) Shrubland fire-biomes. Each datapoint represents a single matchup fire that had its plume total Excess CO and total released FRE assessed for a matching time-period, and the EC_{CO}^b value for the fire-biome is derived from the slope of the OLS linear best fit (solid black line) to these data. Colour of the scatter points denotes the ROI containing the fire (see Fig. 32c). Error bars denote the standard deviation of FRP values and the uncertainty in the S5P TCCO produce respectively, as calculated following (Nguyen et al., 2023). The shaded area indicates the uncertainty on the slope of the linear best fit, taken to be the uncertainty on the derived EC_{CO}^b value.




Figure 8: Spatial analysis comparison of GFASv1.4 FRP values (2019-2024, JJA) and matchup fires selected from GFASv1.4. (a) Relative frequency of FRP observations plotted against matchup-fire pixel locations. (b) Zonal FRP relative frequency (1° bins) for All GFASv1.4 (JJA) pixels and matchup-fire pixels. (c) Zonal mean FRP value frequency (1° bins) for all GFASv1.4 (JJA) pixels and matchup-fire pixels. (d) Longitudinal FRP relative frequency (2.5° bins) for all GFASv1.4 (JJA) pixels and matchup-fire pixels. (e) Longitudinal mean FRP (2.5° bins) for all GFASv1.4 (JJA) pixels and matchup-fire pixels. (f) FRP relative frequency for all GFASv1.4 (JJA) pixels and matchup-fire pixels.

It should be noted that the emission coefficients shown in Fig. 7, as well as the matchup fire dataset they are derived, assume a single relationship between CO emissions and FRE for a given biome. In reality, this is unlikely to be the case, as combustion phase and the relative contributions of flaming and smouldering combustion can heavily influence the fire behaviour, and resulting smoke and emission characteristics (Freeborn et al., 2008; Reid et al., 2009). However, no current space-based EO approach for estimating landscape fire emissions explicitly accounts for variability in combustion phase (Owsley-Brown et al., 2024). Additionally, satellite-derived FRP represents an instantaneous measurement of radiative power and provides no direct information on the relative proportions of flaming and smouldering combustion occurring within the observed fire.

The uncertainty bars associated with the emission coefficients in Fig. 7 account for two primary sources of uncertainty. Firstly, the estimated uncertainty in the Sentinel-5P CO retrievals, including the approximate 10% bias relative to TCCON observations (Sha et al., 2021). Secondly, the spatial variability of both the excess CO estimations and the FRP within each fire polygon, represented by their standard deviations. In addition to these quantified sources of uncertainty, the spread of the individual matchup fires contributing to each biome-specific emission coefficient is also likely to reflect natural variability in the combustion phases, particularly differences in the proportion of flaming and smouldering activity between fires.

Additionally, the emission coefficients shown in Fig. 7, together with the matchup fire dataset they are derived from, do not explicitly account for peat or organic soil combustion, and therefore do not distinguish between regions of high or negligible

organic soil content, which may lead to an underestimation of CO emission from fires within peat-rich environment. Organic soil content was considered as a potential additional classification variable during the matchup fire dataset creation, via use of the global PEATMAP product (Xu et al., 2018). However, incorporating this information reduced the sample sizes of each individual classification, limiting the reliability of the respected OLS regressions. Therefore, as with the combustion phase, the emission coefficients implicitly assumes a uniform contribution of organic soil combustion across the entire matchup fire dataset. This assumption, like that with combustion phase, likely contributes the variability of the individual matchup fires within Figure 7, which is reflected in the uncertainty of the fitted OLS regressions.

In order to assess the impact of using the one-pixel buffer around the digitized plume polygon, the excess CO calculation was repeated, using an increasing number of pixels as the buffer (specifically, a one-, two-, three-, four-, five-, and six-pixel buffer) to capture $TCCO_{BG}$. EC_{CO}^b were recalculated, using the new excess CO values from the corresponding pixel buffer (Fig. A2). Table 5 details the calculated EC_{CO}^b for each biome and pixel buffer, along with the percentage change from the one-pixel buffer. It can be seen that for the forested (DecNeedle and EGNeedle) and shrubland biomes, the EC_{CO}^b increased with increasing pixel buffer, before approaching a plateau between three and four pixels. Beyond this point, further expansion of the buffer resulted in relatively small increases in EC_{CO}^b ($< 3\%$ for both forested biomes, and $< 1\%$ for shrubland biomes), suggesting that most of the excess CO associated with the plume had already been captured. In contrast, the grassland EC_{CO}^b exhibited a stronger dependence on plume buffer size, increasing approximately 35% between the one- and five-pixel buffers, indicating greater sensitivity to plume buffer size. However, the one-pixel buffer was adopted for all subsequent analyses, to maintain methodological consistency with previous FREMv2 studies. The sensitivity analysis demonstrated that this choice introduces relatively small differences for the forest and shrubland biomes (Coefficient of variation, CV = 3.1 – 6.8%), although a larger sensitivity was observed for grassland (CV = 9.8%).

Table 5: EC_{CO}^b values calculated using one- to six-pixel buffers to estimate $TCCO_{BG}$. Scatterplots from which EC_{CO}^b was calculated can be found in Fig. 7 (for one-pixel buffer) and A2 (for two- to six-pixel buffer).

Pixel Buffer	DecNeedle		EGNeedle		Grassland		Shrubland	
	EC_{CO}^b [g MJ ⁻¹]	% change from one-pixel	EC_{CO}^b [g MJ ⁻¹]	% change from one-pixel	EC_{CO}^b [g MJ ⁻¹]	% change from one-pixel	EC_{CO}^b [g MJ ⁻¹]	% change from one-pixel
One-	81.54	0.0	66.74	0.0	53.78	0.0	53.14	0.0
Two-	85.35	4.7	70.89	6.2	61.39	14.2	55.50	4.4
Three-	87.21	7.0	71.97	7.8	67.63	25.8	59.56	12.1
Four-	88.16	8.1	72.67	8.9	68.30	27.0	63.19	18.9
Five-	88.85	9.0	73.33	9.9	71.01	32.0	63.33	19.2
Six-	89.53	9.8	73.56	10.2	72.77	35.3	63.36	19.2
CV	3.1%	=	3.2%	=	9.8%	=	6.8%	=

437 4. Emission Inventory Comparisons

438 4.1. High Latitude Carbon Monoxide Emission Timeseries


439 To generate a long-term CO emissions timeseries for the HL biomes, we applied the derived EC_{CO}^b emissions coefficients to
440 the GFAS v1.2 FRP data available from the current time back to almost the start of the MODIS record (2003). The resulting
441 monthly and annual CO emissions timeseries are shown in Fig. 97. Also shown are the FEER-equivalent CO timeseries, the
442 CO emissions that come as part of the GFAS v1.2 dataset derived from the FRP values as per Kaiser et al., (2012), and the
443 GFED v4.1s CO emissions (2003 – 2024). The mean annual CO emissions for the four biomes using HLFREM, FEER, GFAS
444 v1.2, and GFED v4.1 can be found in Table 65, and note that of these four datasets, the same MODIS FRP data drive the
445 emissions estimates in HLFREM, GFAS v1.2 and FEER, whilst GFED v4.1s uses primarily MODIS burned area data (Giglio
446 et al., 2013). However, also note that the conversion coefficients linking daily mean FRP to biomass burned in GFED v1.2
447 as a prior step to its CO emissions estimation were generated using linear regression against burned biomass estimates of an
448 earlier version of GFED (Kaiser et al., 2012). HLFREM removes this link by directly relating the daily FRP measures to CO
449 fluxes via the conversion coefficients derived from FRP and Sentinel-5P CO data at the matchup fires.

450



451 Table 56: Mean annual (2003-2024) CO emissions for the inventory-biome pairs. Uncertainty values are the standard error of the
452 mean.

Inventory	DecNeedle	EGNeedle	Grassland	Shrubland
HLFREM	5.247.83 ± 1.2292 Tg	2.945.00 ± 0.574.03 Tg	0.6281 ± 0.103 Tg	0.2346 ± 0.0540 Tg
FEER	12.7364 ± 23.9640 Tg	7.466.99 ± 1.453 Tg	2.2833 ± 0.357 Tg	0.667 ± 0.145 Tg
GFAS v1.2	11.48 ± 2.66 Tg	7.18 ± 1.39 Tg	4.03 ± 0.80 Tg	0.80 ± 0.16 Tg
GFED v4.1s	13.63 ± 4.07 Tg	9.938.74 ± 3.882.23 Tg	2.132 ± 0.302 Tg	0.678 ± 0.156 Tg

453



454



455 **Figure 97: Timeseries of Monthly (a,c,e,g) and Annual (b,d,f,h) HL Wildfire emissions of CO for (a,b) DecNeedle, (c,d) EGNeedle,**
 456 **(e,f) Grassland, and (g,h) Shrubland biomes, calculated using the EC_{CO}^b from HLFREM (blue) and FEER (orangered) as detailed in**
 457 **Table 4 using daily FRE totals generated by GFAS v1.2. GFAS v1.2 (green) and GFED v4.1s (redorange) CO emissions for the same**
 458 **biomes are shown in green. OLS regression analysis can be found in Fig. A23.**

459 It can be seen from Fig. 78 that the temporal patterns of HLFREM CO emissions across all biomes are consistent with those
 460 from the other three inventories, with the temporal patterns being more pronounced in the two forested and Shrubland biomes.
 461 This is to be expected with respect to GFAS v1.2 and FEER-EQ, as HLFREM, GFAS, and FEER-EQ emissions all use the
 462 same FRP input (GFAS v1.2). The peaks also match independent reports of fire activity, for example with the DecNeedle
 463 biome showing CO peaks in both 2020 and 2021 consistent with the high fire activity reported across HL Russia (dominated
 464 by DecNeedle) in these years (Kharuk et al., 2021; Ponomarev et al., 2021). Similarly, the CO increase in 2023 in the EGNeedle
 465 biome is consistent with reported high fire activity across HL North America (Byrne et al., 2024; Dodd et al., 2018). It is clear
 466 from Fig. 97 and Table 56 that the CO emissions from HLFREM across the two forested biomes are consistently smaller than
 467 those of the other inventories (DecNeedle: ~~3859%~~, ~~3254%~~, and ~~4362%~~ smaller than FEER, GFAS v1.2, and GFED v4.1s
 468 respectively, EGNeedle: ~~2861%~~, ~~3059%~~ and ~~4370%~~ smaller than FEER, GFAS v1.2, and GFED v4.1s respectively). The
 469 HLFREM CO emissions for Shrublands are also smaller by the same magnitude (~~3465%~~, ~~4371%~~, and ~~3366%~~ smaller than
 470 FEER, GFAS v1.2, and GFED v4.1s respectively). For Grasslands, however, the HLFREM CO emissions are significantly
 471 smaller (~~6573%~~, ~~8085%~~, and ~~6271%~~ smaller than FEER, GFAS v1.2, and GFED v4.1s respectively), though the grassland
 472 EC_{CO}^b emission coefficient (Fig. 96c) has a smaller r^2 value, potentially making them more uncertain. Additionally, there appear
 473 to be some unusual patterns in the Grassland biome; the CO emissions from GFAS v1.2 exhibit a peak in June and July 2015
 474 and July 2021, peaks not seen in the HLFREM- nor FEER-derived versions. On investigation, these peaks are due to an above-
 475 average number of grassland fires occurring in regions defined within GFAS as having peat (Kaiser et al., 2012), as opposed
 476 to extratropical forests with organic soils (EFOS). The presence of peat dramatically increases the conversion factor linking
 477 FRP to dry matter combustion in the GFAS system via the linear regression performed against GFED biomass burned estimates
 478 (Kaiser et al., 2012). Therefore, in the GFAS v1.2 CO emissions dataset, the same FRE values for grassland fires in these
 479 ‘PEAT’ regions results in significantly more CO than identical FRE fires in the EFOS regions. The HLFREM and FEER-
 480 equivalent inventories do not make this distinction, and so their CO timeseries do not show such elevated CO emissions peaks
 481 related to differing Grassland fire locations. Figure 108 shows the fractional dominant fire type as denoted by GFAS v1.2 and
 482 GFED v3.1 for the four HL biomes used in this study. The Grassland biome has a larger fraction of PEAT (11.7%), compared
 483 to the other three biomes (DecNeedle: 2.8%, EGNeedle: 1.6%, Shrubland: 3.1%). Nevertheless, since that CO emissions from
 484 fires in Grassland only represent ~12% of the multi-biome total FRE (Table 3) the overall impact of this difference is limited
 485 on the total CO emissions values.






Figure 810: GFAS v1.2 land cover classes, based on dominant fire type in GFED v3.1 and organic soils and peat maps, taken from (Kaiser et al., 2012). (a) Land cover classes map, for Savannah (SA), Savannah with organic soils (SAOS), Agriculture (AG), Agriculture with organic soils (AGOS), Tropical Forest (TF), Peat (PEAT), Extratropical Forest (EF), and Extratropical Forests

491 with organic soils (EFOS). 60°N is denoted with a red line. (b-e) Fractional land cover for (b) HL DecNeedle, (c) HL EGNeedle, (d)
492 HL Grassland, and (e) HL Shrubland.

493 In terms of spatial patterns, Fig. ~~119~~ maps the annual total GFAS v1.2 FRE values for the a subregion of the Sakha Republic
494 in Russia in 2021 and the Northwest Territories in Canada 2023 respectively~~HL BONA 2023 and HL BOAS 2021 ROIs,~~
495 ~~detailed in Table 1, from which the HLFREM EC_{CO}^b were derived from.~~ The elevated annual FRE totals (Figs. ~~119~~a and ~~119~~b)
496 align with CO emissions across all four inventories (Figs. ~~911~~c-h), whose spatial pattern is very similar, and corresponded to
497 the extreme wildfire activity reported in these years and regions (Byrne et al., 2024; Kharuk et al., 2021). This is to be expected,
498 as HLFREM, FEER-equivalent, and GFASv1.2 emissions all stem from GFASv1.2 FRE totals.





500

501 **Figure 911:** Annual FRE totals from GFAS v1.2 (a, b) and CO emissions (c-j) for a subregion of the Northwest Territories, Canada
 502 in 2023 (the HL-BONA-23 (a, c, e, h, i) and a subregion of the Sakha Republic, Russia in 2021 (HL-BOAS-21 (b, d, f, h, j) ROIs from
 503 the HLFREM (c, d), FEER-equivalent (e, f), GFAS v1.2 (g, h), and GFED v4.1s (i, j), respectively. Grey areas denote regions that do
 504 not fall under one of the four HLFREM biomes, whilst blue denote waterbodies.

505 4.2. High Latitude Carbon Emission Timeseries

506 Following the approach of (Mota and Wooster, 2018; Nguyen et al., 2023; Nguyen and Wooster, 2020), emissions estimates
 507 of total carbon, as well as any other GHG or trace gas, can be derived from the CO emissions provided by the HLFREM
 508 approach, simply via use of standard ratios of the relevant gaseous emission factors. Numerous different summarised EF
 509 inventories exist, with for example GFAS v1.2 using (Andreae and Merlet, 2001) and GFEDv4.1s using (Akagi et al., 2011).
 510 HL wildfire emissions of Carbon, for example, can be calculated using its biome-specific emission coefficient (EC_C^b), derived
 511 using the EF ratio between Carbon and CO, and the HLFREM CO emission coefficient (EC_{CO}^b), via Eq. (2):

512

$$EC_C^b = \frac{EF_C^b}{EF_{CO}^b} EC_{CO}^b \quad (2)$$

513

514 Where the emission coefficients have units: g MJ⁻¹ and the emission factors units: g kg⁻¹. EF_{CO}^b values are taken from the
aforementioned emission inventories, and EF_C^b being calculated using Eq. (3):

515

$$EF_C^b = \frac{12}{44} EF_{CO_2}^b + \frac{12}{28} EF_{CO}^b + \frac{12}{16} EF_{CH_4}^b \quad (3)$$

516

517 Where EF_C^b , $EC_{CO_2}^b$, EC_{CO}^b , and $EC_{CH_4}^b$ are the biome specific wildfire EFs for burnt carbon, CO₂, CO, and CH₄ respectively,
518 assuming these make up more than 95% of burnt carbon emitted in gaseous form (Akagi et al., 2011) and that by comparison
519 carbon in aerosols is negligible by mass (Akagi et al., 2011; Andreae and Merlet, 2001). Multiplying these carbon emissions
by a factor of two then provides an estimate of the amount of biomass burned.


520

521 Similar to Fig. 7, Fig. 10 shows our multi-inventory Carbon emissions calculated across the four most dominant HL fire
522 affected biomes. As GFAS v1.2 uses EFs from Andreae and Merlet, (2001), we calculated EC_C^b for both HLFREM and FEER
523 using these same EFs, as shown in Table 4. However, as GFED v4.1s uses EFs from (Akagi et al., 2011), we also calculated
524 EC_C^b for HLFREM using these same EFs, also shown in Table 4. The HLFREM and FEER EC_C^b were applied to the GFAS
525 v1.2 daily FRE totals, and, similar to Table 6, mean annual HL Wildfire Carbon emissions totals were calculated, and are
shown in Table 7.

526

527 The HLFREM carbon emission timeseries exhibits a very similar behaviour as the CO emission timeseries from which it
528 was derived; with a strong temporal similarity with the carbon emissions of the other inventories. The difference between the
529 HLFREM carbon emissions and the other inventories for the two forested biomes (DecNeedle: 38%, 30%, and 42% smaller
than FEER, GFAS v1.2, and GFED v4.1s respectively, EGNeedle: 28%, 30% and 42% smaller than FEER, GFAS v1.2, and
530 GFED v4.1s respectively) and the Shrubland biomes (31%, 41%, and 32% smaller than FEER, GFAS v1.2, and GFED v4.1s
531 respectively), are very similar to those of the CO emissions, differing by $\leq 2\%$ from these. The HLFREM carbon emission
532 timeseries from Grassland also exhibits a similar behaviour to those of the Grassland HLFREM CO emission timeseries (65%,
533 74%, and 61% smaller than FEER, GFAS v1.2, and GFED v4.1s respectively); in that the emission estimations are significantly
534 smaller than those from the other inventories, although the difference from GFAS v1.2 (6%) is larger than those from the other
535 two inventories (2%).

536



537




Figure 102: Timeseries of HL Wildfire Emissions of Carbon for (a, b) Deciduous Needleleaf Forest, (c, d) Evergreen Needleleaf Forest, (e, f) Grasslands, and (g, h) Shrublands biomes, for (a, c, e, g) FEER, GFAS v1.2, and HLFREM (using EFs from (Andreae and Merlet, 2001)), and (b, d, f, h) GFEDv4.1s and HIFREM (using EFs from (Akagi et al., 2011)). Regression analysis on the timeseries can be found in Fig. A43.

Table 76: Mean annual (2003-2024) C emissions for the inventory-biome pairs. HLFREM_{AM} details the mean annual HL Carbon emissions, using EC_{C-AM}^b , whilst HLFREM_{AG} details the mean annual HL Carbon emission using EC_{C-AG}^b . HLFREM and FEER HL Carbon emissions are generated using GFAS v1.2 FRE totals. Uncertainty values are the standard error of the mean.

Inventory	DecNeedle	EGNeedle	Grassland	Shrubland
HLFREM _{AM}	23.4235.29 ± 58.7567 Tg	12.4022.53 ± 2.544.62 Tg	2.843.65 ± 0.450.57 Tg	1.052.08 ± 0.240.47 Tg
HLFREM _{AG}	19.0328.67 ± 4.677.04 Tg	10.0718.30 ± 2.073.76 Tg	2.31 ± 0.362.96 ± 0.47 Tg	0.851.69 ± 0.190.38 Tg
FEER	56.95 ± 13.99 Tg	31.49 ± 6.46 Tg	10.48 ± 1.65 Tg	3.00 ± 0.67 Tg
GFAS v1.2	50.72 ± 12.04 Tg	32.14 ± 6.38 Tg	14.02 ± 2.49 Tg	3.52 ± 0.74 Tg
GFED v4.1s	50.1549.60 ± 14.1482 Tg	35.621.40 ± 8.687.96 Tg	7.6357 ± 1.0813 Tg	2.427 ± 0.5457 Tg

5. Summary and Conclusions

Taking advantage of the increased number of polar orbiting satellite overpasses nearer the poles, we have extended the Fire Radiative Energy Emissions (FREM) approach to direct “top down” fire emissions estimation to High Latitudes (HL) fires, and to polar-orbiting FRP datasets. Previously the approach was limited to use on low-to-mid latitude geostationary FRP data (Nguyen et al., 2023). This highly direct approach to estimating fire emissions uses only satellite observations of fire radiative power and CO, removing the need for assuming the amount of biomass per unit area, pre-fire fuel loads and combustion completeness, or correlations between FRE and burned biomass developed in the laboratory, in small scale fires, or elsewhere. Negating the requirements for these parameters removes a key source of uncertainty associated with other emissions estimation techniques (Kasischke and Penner, 2004; Reid et al., 2009; Wooster et al., 2015), and provides a means of converting the FRE data from the Global Fire Assimilation system (GFAS) into fire emissions without use of factors derived via linear regressions against the GFED burned biomass totals, as is used currently within GFAS (Kaiser et al., 2012).

Via a set of 708 matchup fires across four regions of interest in the HL region above 60°N, we used S5P TROPMI Total Column Carbon Monoxide (TCCO) data to calculate total plume CO and total FRE up to the time of the S5P overpass from GFAS v1.4. From these data a set of biome specific CO emission coefficients (EC_{CO}^b) were generated using the approach of (Nguyen et al., 2023) that can then be used to convert further FRP data of fires in the four dominant HL biomes (Deciduous Needleleaf Forests, Evergreen Needleleaf Forests, Grasslands, and Shrublands) to emission rates of CO. Application of these emissions coefficients to the daily FRE totals from GFAS v1.2 enabled us to produce new CO emissions inventory for these four biomes in the HL region, unrelant on any additional parameters or datasets.

564 The CO emissions derived from this new approach were compared to three other widely used fire emission inventories,
 565 those produced and outputted by GFAS v1.2 itself, those from FEERv1.0-GFASv1.2 (calculated using GFAS FRE and a
 566 FEER-equivalent EC_{CO}^b ; (Nguyen et al., 2023)), and those from GFED v4.1s. Total carbon emissions estimates were also
 567 calculated based on emission factor ratios between CO and Carbon. The HLFREM CO and Carbon emissions timeseries and
 568 CO spatial patterns derived using the HLFREM approach were found to be in good temporal and spatial agreement with the
 569 FEER-equivalent, GFAS v1.2, and GFED v4.1s inventories, particularly for the two forested biomes and shrublands, producing
 570 relatively similar but generally smaller emissions totals. Mean annual HLFREM CO and carbon emission totals were found to
 571 be ~~2854-4371~~% smaller than totals produced by FEER, GFAS v1.2, and GFAS v4.1s for the two forested biomes and Shrubland
 572 biomes. HLFREM CO and carbon emission totals from Grassland were also in good temporal and spatial agreement with the
 573 other inventories, although the emissions were significantly lower (~~7161-8085~~%) than those from other inventories; and whilst
 574 this could be correct, it is also possibly a result of the relatively small number of fire-matchups found during the generation of
 575 the HLFREM $EC_{CO}^{Grassland}$ and perhaps a low bias. Compounding this, the largest difference (GFAS v1.2, 859% lower CO
 576 emissions, 8074% lower carbon emissions) was found over the Grassland biomes, which in certain of the other databases has
 577 some consideration of peat burning included that is not accounted for in the HLFREM methodology. ~~On the other hand, if the~~
 578 ~~matchup fires included areas having peat burning, the impact of this would already be in the derived emissions coefficients/~~
 579 Not accounting for peat burning, as well as not accounting for changes in combustion phase, by assuming a single
 580 relationship between CO emissions and FRE within each fire biome is a limitation of this HLFREM methodology. While
 581 combustion phase cannot be currently be resolved using space-based EO observations, organic soil content was evaluated as
 582 an additional classification variable but was not incorporated because it substantially reduced the sample size within each class.
 583 Future developments in advancing the HLFREM approach will include consideration of this organic soil burning, as well
 584 as expansion of both the biomes covered and the number of fire matchups in each. Our aim is to develop a long-term and
 585 geographically continuous HL fire emissions dataset that does not rely on parameters taken from modelling, from correlations
 586 between inventories, or from conversion coefficients taken from laboratory or small-scale fire experiments when converting
 587 the satellite observations into emissions estimates. Instead, the FREM approach uses information extracted from the satellite
 588 data itself to generate the necessary conversion coefficients, an approach that we feel has great potential to expand further as
 589 both active fire and trace gas remotely sensed datasets continue to advance.
 590 Whilst the HLFREM framework is readily transferable to fires globally, through a global matchup fire database using
 591 TROPOMI TCCO observations and a global GFAS v1.4 dataset, its application would likely benefit more from the use of
 592 geostationary FRP observations from other platforms, such as the National Oceanic and Atmospheric Administration (NOAA)
 593 Geostationary Operational Environment Satellite (GOES) or Japan Aerospace Exploration Agency (JAXA) Himawari
 594 platforms, where available. Compared to with the hourly GFAS FRP product, which relies on a prescribed diurnal cycle
 595 (Andela et al., 2015) to distribute fire activity between solar overpasses, geostationary observations provide much higher
 596 temporal resolution, and directly capture the natural diurnal variability of fire behaviour. This would allow the temporal
 597 evolution of emissions to be represented more realistically, particularly for fires with rapidly changing intensities.


598

599 **Appendix A**

600

Table A1: Aggregated CCI landcover classes (taken from ESA 2017) assigned to each HLFREM biome.

HLFREM Biome (≥60°N)	Full Biome Name	Assigned CCI Class Codes
Managed	Managed	10, 11, 12, 20, 30, 40
EGBroad	Evergreen Broadleaf Forests	50, 160, 170
DecBroad	Deciduous Broadleaf Forests	60, 61, 62, 90
EGNeedle	Evergreen Needleleaf Forests	70, 71, 72
DecNeedle	Deciduous Needleleaf Forests	80, 81, 82
Grassland	Grassland	100, 110, 130, 180, 190
Shrubland	Shrubland	120, 121, 122
Sparce	Sparce	140, 150, 151, 152, 153
Bare	Bare	200, 201, 202
Snow/Ice	Snow and Ice	220



601

602 **Figure A1: Median (blue) and mean (orange) hourly FRP of all plumes identified in the HLFREM study, during the day before**
603 **observation (negative hours relative to local midnight) and day of observation (positive hours relative to local midnight) from the**
604 **hourly FRP dataset. Local midnight and 06:00 local time is shown with the black and red line, respectively.**





Figure A2: HLFREM biome-specific CO emission coefficients (EC_{CO}^b , in $g\ MJ^{-1}$) derived for Deciduous Needleleaf Forests (DecNeedle, a, e, i, m, q, u), Evergreen Needleleaf Forests (EGNeedle, b, f, j, n, r, v), Grassland (Grassland, c, g, k, o, s, w), and Shrubland (Shrubland, d, h, l, p, t, x), using a one- (a, b, c, d), two- (e, f, g, h), three- (i, j, k, l), four- (m, n, o, p), five- (q, r, s, t), and six-pixel buffer (u, v, w, x) to determine the background TCCO values, used in Table 5. R^2 values are also shown.







Figure A32: OLS Regression analysis of the Annual HL CO Emissions, comparing the HLFREM emissions (calculated using GFAS v1.2 FRE totals) with (a-d) FEER, (e-h) GFAS v1.2, and (i-l) for Deciduous Needleleaf Forests (a,e,i), Evergreen Needleleaf Forests (b,f,j), Grassland (c,g,k), and Shrubland (d,h,l) biomes. OLS regression and associated errors are shown on each plot (displayed as the solid line and shading respectively), with the R² value being shown in the brackets. A 1:1 line (dashed line) is also shown.






Figure A43: OLS Regression analysis of the Annual HL C Emissions, comparing the HLFREM emissions (calculated using GFAS v1.2 FRE totals) with (a-d) FEER, (e-h) GFAS v1.2, and (i-l) for Deciduous Needleleaf Forests (a,e,i), Evergreen Needleleaf Forests (b,f,j), Grassland (c,g,k), and Shrubland (d,h,l) biomes. As the HLFREM C Emissions have been calculated via EF ratios with respect to Carbon Monoxide, different EF ratios have been used to compare FEER and GFAS v1.2 (using EFs from (Andreae and Merlet, 2001), referred to as HLFREM_{AM}), and GFEDv4.1s (using EFs from (Akagi et al., 2011), referred to as HLFREM_{AG}). OLS regression and associated errors are shown on each plot (displayed as the solid line and shading respectively), with the R² value being shown in the brackets. A 1:1 line (dashed line) is also shown.

7. Code Availability

Code is available upon request to William Maslanka (william.maslanka@kcl.ac.uk).

8. Data Availability

Sentinel-5P products are distributed freely by the Copernicus Data Space Ecosystem (<https://dataspace.copernicus.eu/>, last accessed 19 March 2025), as is the VIIRS and MODIS AF and Multispectral Products by the LAADS DAAC (<https://ladsweb.modaps.eosdis.nasa.gov/>, last accessed 19 March 2025), and the VIIRS AOD EPS product on the NOAA

631 Comprehensive Large Array-Data Stewardship System (CLASS, <https://www.aev.class.noaa.gov/>, last accessed 19 March
632 2025). GFAS v1.2 data is distributed freely through the ECMWF Atmosphere Data Store
633 (<https://ads.atmosphere.copernicus.eu/>, last accessed 19 March 2025). GFAS v1.4 data used in this study is available upon
634 request. GFED v4.1s data is distributed freely from the GFED web portal (<https://www.globalfiredata.org/>, last accessed 19
635 March 2025). FEER data is distributed freely on the Fire Energetics and Emission Research website
636 (<https://feer.gsfc.nasa.gov/index.php>, last accessed on 19 March 2025).

637 **9. Author Contributions**

638 W.M.: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Software,
639 Validation, Visualization, Writing - original draft, and Writing - review & editing.

640 M.W.: Conceptualization, Formal analysis, Funding acquisition, Methodology, Supervision, Writing - original draft, and
641 Writing - review & editing.

642 Z.L.: Data curation, Resources, Software, and Writing - review & editing.

643 J.H.: Resources, Software, and Writing - review & editing.

644 **10. Competing Interests**

645 The authors declare that they have no conflict of interest.

646 **11. Acknowledgements**

647 Neither the European Commission nor ECMWF is responsible for any use that may be made of the information it contains.
648 The authors would like to thank Mark Parrington and Enza Di Tomaso of the ECMWF for the provision of GFAS v1.4 data,
649 as well as members of the KCL Wildfire and Earth Observation research group, who have helped to improve the quality and
650 content of the research and of this paper. We would also like to thank the anonymous reviewers for their comments, which
651 have helped to improve the quality of this manuscript.

652 **12. Financial Support**

653 This work was supported by the Natural Environment Research Council (NERC) National Centre for Earth Observation
654 (NCEO) funding under grants NE/R016518/1 and NE/Y006216/1. This research is also an output of the Leverhulme Centre
655 for Wildfires, Environment and Society through the Leverhulme Trust, grant number RC-2018-023.

656 **References**

657 Adams, C., McLinden, C. A., Shephard, M. W., Dickson, N., Dammers, E., Chen, J., Makar, P., Cady-Pereira, K. E., Tam, N.,
658 Kharol, S. K., Lamsal, L. N., and Krotkov, N. A.: Satellite-derived emissions of carbon monoxide, ammonia, and nitrogen
659 dioxide from the 2016 Horse River wildfire in the Fort McMurray area, *Atmos. Chem. Phys.*, 19, 2577–2599,
660 <https://doi.org/10.5194/acp-19-2577-2019>, 2019.

661 Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.:
662 Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072,
663 <https://doi.org/10.5194/acp-11-4039-2011>, 2011.

664 Andela, N., Kaiser, J. W., van der Werf, G. R., and Wooster, M. J.: New fire diurnal cycle characterizations to improve fire
665 radiative energy assessments made from MODIS observations, *Atmos. Chem. Phys.*, 15, 8831–8846,
666 <https://doi.org/10.5194/acp-15-8831-2015>, 2015.

667 Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*,
668 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.

669 Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, 15,
670 955–966, <https://doi.org/10.1029/2000GB001382>, 2001.

671 Apituley, A., Pedergrana, M., Sneep, M., Veeffkind, J. P., Loyola, D., Landgraf, J., Borsdorff, T., and Mandal, S.: Sentinel-5
672 precursor/TROPOMI Level 2 Product User Manual Carbon Monoxide, 1–126 pp., 2018.

673 Borsdorff, T., van de Brugh, J., Hu, H., Hasekamp, O., Susmann, R., Rettinger, M., Hase, F., Gross, J., Schneider, M., Garcia,
674 O., Stremme, W., Grutter, M., Feist, D. G., Arnold, S. G., De Mazière, M., Kumar Sha, M., Pollard, D. F., Kiel, M., Roehl, C.,
675 Wennberg, P. O., Toon, G. C., and Landgraf, J.: Mapping carbon monoxide pollution from space down to city scales with
676 daily global coverage, *Atmos. Meas. Tech.*, 11, 5507–5518, <https://doi.org/10.5194/amt-11-5507-2018>, 2018.

677 Byrne, B., Liu, J., Bowman, K. W., Pascolini-Campbell, M., Chatterjee, A., Pandey, S., Miyazaki, K., van der Werf, G. R.,
678 Wunch, D., Wennberg, P. O., Roehl, C. M., and Sinha, S.: Carbon emissions from the 2023 Canadian wildfires, *Nature*, 633,
679 835–839, <https://doi.org/10.1038/s41586-024-07878-z>, 2024.

680 Chen, Y., Hall, J., van Wees, D., Andela, N., Hantson, S., Giglio, L., van der Werf, G. R., Morton, D. C., and Randerson, J.
681 T.: Multi-decadal trends and variability in burned area from the fifth version of the Global Fire Emissions Database (GFED5),
682 *Earth Syst. Sci. Data*, 15, 5227–5259, <https://doi.org/10.5194/essd-15-5227-2023>, 2023.

683 Crutzen, P. J. and Andreae, M. O.: Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical
684 Cycles, *Science* (1979)., 250, 1669–1678, 1990.

685 Darmenov, A. and da Silva, A.: The quick fire emissions dataset (QFED) – Documentation of versions 2.1, 2.2 and 2.4.
686 NASA/TM-2015-104606, Greenbelt, MD 20771, 1–212 pp., 2015.

687 Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change,
688 *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.

689 Dodd, W., Scott, P., Howard, C., Scott, C., Rose, C., Cunsolo, A., and Orbinski, J.: Lived experience of a record wildfire
690 season in the Northwest Territories, Canada, *Canadian Journal of Public Health*, 109, 327–337,
691 <https://doi.org/10.17269/s41997-018-0070-5>, 2018.

692 ESA: Land Cover CCI Product User Guide Version 2. Tech. Rep., 2017.

693 Fisher, D., Wooster, M. J., Xu, W., Thomas, G., and Lestari, P.: Top-down estimation of particulate matter emissions from
694 extreme tropical peatland fires using geostationary satellite fire radiative power observations, *Sensors* (Switzerland), 20, 1–
695 25, <https://doi.org/10.3390/s20247075>, 2020.

696 Freeborn, P. H., Wooster, M. J., Hao, W. M., Ryan, C. A., Nordgren, B. L., Baker, S. P., and Ichoku, C.: Relationships between
697 energy release, fuel mass loss, and trace gas and aerosol emissions during laboratory biomass fires, *Journal of Geophysical*
698 *Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008679>, 2008.

699 Friedlingstein, P., Lovenduski, N., Lewis, S., Schuur, E., Turetsky, M., and Smith, A.: Climate change weakens carbon sinks
700 and further amplifies climate change, *Critical Issues in Climate Change Science*, 1–4,
701 <https://doi.org/https://doi.org/10.5281/zenodo.5596739>., 2021.

702 Giglio, L., Randerson, J. T., and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-
703 generation global fire emissions database (GFED4), *J. Geophys. Res. Biogeosci.*, 118, 317–328,
704 <https://doi.org/10.1002/jgrg.20042>, 2013.

705 Giglio, L., Schroeder, W., and Justice, C. O.: The collection 6 MODIS active fire detection algorithm and fire products, *Remote*
706 *Sens. Environ.*, 178, 31–41, <https://doi.org/10.1016/j.rse.2016.02.054>, 2016.

707 Di Giuseppe, F., Rémy, S., Pappenberger, F., and Wetterhall, F.: Using the Fire Weather Index (FWI) to improve the estimation
708 of fire emissions from fire radiative power (FRP) observations, *Atmos. Chem. Phys.*, 18, 5359–5370,
709 <https://doi.org/10.5194/acp-18-5359-2018>, 2018.

710 Griffin, D., Chen, J., Anderson, K., Makar, P., McLinden, C. A., Dammers, E., and Fogal, A.: Biomass burning CO emissions:
711 exploring insights through TROPOMI-derived emissions and emission coefficients, *Atmos. Chem. Phys.*, 24, 10159–10186,
712 <https://doi.org/10.5194/acp-24-10159-2024>, 2024.

713 Hayden, K. L., Li, S.-M., Liggio, J., Wheeler, M. J., Wentzell, J. J. B., Leithead, A., Brickell, P., Mittermeier, R. L., Oldham,
714 Z., Mihele, C. M., Staebler, R. M., Moussa, S. G., Darlington, A., Wolde, M., Thompson, D., Chen, J., Griffin, D., Eckert, E.,
715 Ditto, J. C., He, M., and Gentner, D. R.: Reconciling the total carbon budget for boreal forest wildfire emissions using airborne
716 observations, *Atmos. Chem. Phys.*, 22, 12493–12523, <https://doi.org/10.5194/acp-22-12493-2022>, 2022.

717 Hodges, J. A., Price, J. N., Nicotra, A. B., Neeman, T., and Guja, L. K.: Smoke and heat accelerate and increase germination
 718 in fire-prone temperate grassy ecosystems, *Ecosphere*, 12, <https://doi.org/10.1002/ecs2.3851>, 2021.

719 Hutto, R. L.: The Ecological Importance of Severe Wildfires: Some Like It Hot, *Ecological Applications*, 18, 1827–1834,
 720 <https://doi.org/10.1890/08-0895.1>, 2008.

721 Ichoku, C. and Ellison, L.: Global top-down smoke-aerosol emissions estimation using satellite fire radiative power
 722 measurements, *Atmos. Chem. Phys.*, 14, 6643–6667, <https://doi.org/10.5194/acp-14-6643-2014>, 2014.

723 Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R.,
 724 Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy,
 725 S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, *Atmos. Chem. Phys.*, 19, 3515–3556,
 726 <https://doi.org/10.5194/acp-19-3515-2019>, 2019.

727 Inness, A., Aben, I., Ades, M., Borsdorff, T., Flemming, J., Jones, L., Landgraf, J., Langerock, B., Nedelec, P., Parrington,
 728 M., and Ribas, R.: Assimilation of S5P/TROPOMI carbon monoxide data with the global CAMS near-real-time system,
 729 *Atmos. Chem. Phys.*, 22, 14355–14376, <https://doi.org/10.5194/acp-22-14355-2022>, 2022.

730 Jing, X., Uprety, S., Liu, T.-C., Zhang, B., and Shao, X.: Evaluation of SNPP and NOAA-20 VIIRS Datasets Using RadCalNet
 731 and Landsat 8/OLI Data, *Remote Sens. (Basel)*, 14, 3913, <https://doi.org/10.3390/rs14163913>, 2022.

732 Jones, M. W., Abatzoglou, J. T., Veraverbeke, S., Andela, N., Lasslop, G., Forkel, M., Smith, A. J. P., Burton, C., Betts, R.
 733 A., van der Werf, G. R., Sitch, S., Canadell, J. G., Santin, C., Kolden, C., Doerr, S. H., and Le Quéré, C.: Global and Regional
 734 Trends and Drivers of Fire Under Climate Change, *Reviews of Geophysics*, 60, <https://doi.org/10.1029/2020RG000726>, 2022.

735 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M.
 736 G., Suttie, M., and Van Der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on
 737 observed fire radiative power, <https://doi.org/10.5194/bg-9-527-2012>, 2012.

738 Kaiser, K., Parrington, M., and Armenteras, D.: Biomass Burning in “State of the Climate 2023,” *Bull. Am. Meteorol. Soc.*,
 739 105, S107–S109, <https://doi.org/10.1175/BAMS-D-24-0116.1>, 2024.

740 Kasischke, E. S. and Penner, J. E.: Improving global estimates of atmospheric emissions from biomass burning, *Journal of*
 741 *Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004972>, 2004.

742 Kharuk, V. I., Ponomarev, E. I., Ivanova, G. A., Dvinskaya, M. L., Coogan, S. C. P., and Flannigan, M. D.: Wildfires in the
 743 Siberian taiga, *Ambio*, 50, 1953–1974, <https://doi.org/10.1007/s13280-020-01490-x>, 2021.

744 Landgraf, J., aan de Brugh, J., Scheepmaker, R., Borsdorff, T., Hu, H., Houweling, S., Butz, A., Aben, I., and Hasekamp, O.:
 745 Carbon monoxide total column retrievals from TROPOMI shortwave infrared measurements, *Atmos. Meas. Tech.*, 9, 4955–
 746 4975, <https://doi.org/10.5194/amt-9-4955-2016>, 2016.

747 Liu, Z., Eden, J. M., Dieppois, B., Drobyshev, I., Gallo, C., and Blackett, M.: Were Meteorological Conditions Related to the
 748 2020 Siberia Wildfires Made More Likely by Anthropogenic Climate Change?, *Bull. Am. Meteorol. Soc.*, 103, S44–S49,
 749 <https://doi.org/10.1175/BAMS-D-21-0168.1>, 2022.

750 Martínez-Alonso, S., Deeter, M., Worden, H., Borsdorff, T., Aben, I., Commane, R., Daube, B., Francis, G., George, M.,
 751 Landgraf, J., Mao, D., McKain, K., and Wofsy, S.: 1.5 years of TROPOMI CO measurements: comparisons to MOPITT and
 752 ATom, *Atmos. Meas. Tech.*, 13, 4841–4864, <https://doi.org/10.5194/amt-13-4841-2020>, 2020.
 753 Masrur, A., Petrov, A. N., and DeGroote, J.: Circumpolar spatio-temporal patterns and contributing climatic factors of wildfire
 754 activity in the Arctic tundra from 2001–2015, <https://doi.org/10.1088/1748-9326/aa9a76>, 1 January 2018.
 755 McCarty, J. L., Aalto, J., Paunu, V. V., Arnold, S. R., Eckhardt, S., Klimont, Z., Fain, J. J., Evangeliou, N., Venäläinen, A.,
 756 Tchebakova, N. M., Parfenova, E. I., Kupiainen, K., Soja, A. J., Huang, L., and Wilson, S.: Reviews and syntheses: Arctic fire
 757 regimes and emissions in the 21st century, <https://doi.org/10.5194/bg-18-5053-2021>, 15 September 2021.
 758 Mebust, A. K., Russell, A. R., Hudman, R. C., Valin, L. C., and Cohen, R. C.: Characterization of wildfire NO_x emissions
 759 using MODIS fire radiative power and OMI tropospheric NO₂ columns, *Atmos. Chem. Phys.*, 11, 5839–5851,
 760 <https://doi.org/10.5194/acp-11-5839-2011>, 2011.
 761 Mekonnen, Z. A., Riley, W. J., Randerson, J. T., Shirley, I. A., Bouskill, N. J., and Grant, R. F.: Wildfire exacerbates high-
 762 latitude soil carbon losses from climate warming, *Environmental Research Letters*, 17, 094037, <https://doi.org/10.1088/1748-9326/ac8be6>, 2022.
 763
 764 Mota, B. and Wooster, M. J.: A new top-down approach for directly estimating biomass burning emissions and fuel
 765 consumption rates and totals from geostationary satellite fire radiative power (FRP), *Remote Sens. Environ.*, 206, 45–62,
 766 <https://doi.org/10.1016/j.rse.2017.12.016>, 2018.
 767 Neyra-Nazarrett, O. A., Miyazaki, K., Bowman, K. W., and Saide, P. E.: An Assessment of TROPES CrIS and TROPOMI
 768 CO Retrievals and Their Synergies for the 2020 Western U.S. Wildfires, *Remote Sens. (Basel)*, 17, 1854,
 769 <https://doi.org/10.3390/rs17111854>, 2025.
 770 Nguyen, H. M. and Wooster, M. J.: Advances in the estimation of high Spatio-temporal resolution pan-African top-down
 771 biomass burning emissions made using geostationary fire radiative power (FRP) and MAIAC aerosol optical depth (AOD)
 772 data, *Remote Sens. Environ.*, 248, <https://doi.org/10.1016/j.rse.2020.111971>, 2020.
 773 Nguyen, H. M., He, J., and Wooster, M. J.: Biomass burning CO, PM and fuel consumption per unit burned area estimates
 774 derived across Africa using geostationary SEVIRI fire radiative power and Sentinel-5P CO data, *Atmos. Chem. Phys.*, 23,
 775 2089–2118, <https://doi.org/10.5194/acp-23-2089-2023>, 2023.
 776 NOAA: Visible Infrared Imaging Radiometer Suite (VIIRS) Enterprise Aerosol Optical Depth and Aerosol Particle Size
 777 Products User’s Guide. Version 1, 2020.
 778 Owsley-Brown, F., Wooster, M. J., Grosvenor, M. J., and Liu, Y.: Can the remote sensing of combustion phase improve
 779 estimates of landscape fire smoke emission rate and composition?, *Atmos. Meas. Tech.*, 17, 6247–6264,
 780 <https://doi.org/10.5194/amt-17-6247-2024>, 2024.
 781 Ponomarev, E., Yakimov, N., Ponomareva, T., Yakubailik, O., and Conard, S. G.: Current Trend of Carbon Emissions from
 782 Wildfires in Siberia, *Atmosphere (Basel)*, 12, 559, <https://doi.org/10.3390/atmos12050559>, 2021.

783 Popovicheva, O. B., Evangeliou, N., Kobelev, V. O., Chichayeva, M. A., Eleftheriadis, K., Gregorič, A., and Kasimov, N. S.:
 784 Siberian Arctic black carbon: gas flaring and wildfire impact, *Atmos. Chem. Phys.*, 22, 5983–6000,
 785 <https://doi.org/10.5194/acp-22-5983-2022>, 2022.

786 Ramo, R., Roteta, E., Bistinas, I., van Wees, D., Bastarrika, A., Chuvieco, E., and van der Werf, G. R.: African burned area
 787 and fire carbon emissions are strongly impacted by small fires undetected by coarse resolution satellite data, *Proceedings of*
 788 *the National Academy of Sciences*, 118, <https://doi.org/10.1073/pnas.2011160118>, 2021.

789 Randerson, J. T., Van Der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire Emissions Database,
 790 Version 4.1 (GFEDv4), <https://doi.org/10.3334/ORNLDAAAC/1293>, 2017.

791 Reid, J. S., Hyer, E. J., Prins, E. M., Westphal, D. L., Zhang, J., Wang, J., Christopher, S. A., Curtis, C. A., Schmidt, C. C.,
 792 Eleuterio, D. P., Richardson, K. A., and Hoffman, J. P.: Global Monitoring and Forecasting of Biomass-Burning Smoke:
 793 Description of and Lessons From the Fire Locating and Modeling of Burning Emissions (FLAMBE) Program, *IEEE J. Sel.*
 794 *Top. Appl. Earth Obs. Remote Sens.*, 2, 144–162, <https://doi.org/10.1109/JSTARS.2009.2027443>, 2009.

795 Roberts, G., Wooster, M. J., Perry, G. L. W., Drake, N., Rebelo, L. -M., and Dipotso, F.: Retrieval of biomass combustion
 796 rates and totals from fire radiative power observations: Application to southern Africa using geostationary SEVIRI imagery,
 797 *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2005JD006018>, 2005.

798 Rogers, B. M., Soja, A. J., Goulden, M. L., and Randerson, J. T.: Influence of tree species on continental differences in boreal
 799 fires and climate feedbacks, *Nat. Geosci.*, 8, 228–234, <https://doi.org/10.1038/ngeo2352>, 2015.

800 Rowe, J. P., Zarzana, K. J., Kille, N., Borsdorff, T., Goudar, M., Lee, C. F., Koenig, T. K., Romero-Alvarez, J., Campos, T.,
 801 Knote, C., Theys, N., Landgraf, J., and Volkamer, R.: Carbon Monoxide in Optically Thick Wildfire Smoke: Evaluating
 802 TROPOMI Using CU Airborne SOF Column Observations, *ACS Earth Space Chem.*, 6, 1799–1812,
 803 <https://doi.org/10.1021/acsearthspacechem.2c00048>, 2022.

804 Schroeder, W., Oliva, P., Giglio, L., and Csizsar, I. A.: The New VIIRS 375 m active fire detection data product: Algorithm
 805 description and initial assessment, *Remote Sens. Environ.*, 143, 85–96, <https://doi.org/10.1016/j.rse.2013.12.008>, 2014.

806 Sha, M. K., Langerock, B., Blavier, J.-F. L., Blumenstock, T., Borsdorff, T., Buschmann, M., Dehn, A., De Mazière, M.,
 807 Deutscher, N. M., Feist, D. G., García, O. E., Griffith, D. W. T., Grutter, M., Hannigan, J. W., Hase, F., Heikkinen, P., Hermans,
 808 C., Iraci, L. T., Jeseck, P., Jones, N., Kivi, R., Kumps, N., Landgraf, J., Lorente, A., Mahieu, E., Makarova, M. V., Mellqvist,
 809 J., Metzger, J.-M., Morino, I., Nagahama, T., Notholt, J., Ohyama, H., Ortega, I., Palm, M., Petri, C., Pollard, D. F., Rettinger,
 810 M., Robinson, J., Roche, S., Roehl, C. M., Röhling, A. N., Rousogonous, C., Schneider, M., Shiomi, K., Smale, D., Stremme,
 811 W., Strong, K., Sussmann, R., Té, Y., Uchino, O., Velazco, V. A., Vigouroux, C., Vrekoussis, M., Wang, P., Warneke, T.,
 812 Wizenberg, T., Wunch, D., Yamanouchi, S., Yang, Y., and Zhou, M.: Validation of methane and carbon monoxide from
 813 Sentinel-5 Precursor using TCCON and NDACC-IRWG stations, *Atmos. Meas. Tech.*, 14, 6249–6304,
 814 <https://doi.org/10.5194/amt-14-6249-2021>, 2021.

815 Stockwell, C. E., Bela, M. M., Coggon, M. M., Gkatzelis, G. I., Wiggins, E., Gargulinski, E. M., Shingler, T., Fenn, M., Griffin,
 816 D., Holmes, C. D., Ye, X., Saide, P. E., Bourgeois, I., Peischl, J., Womack, C. C., Washenfelder, R. A., Veres, P. R., Neuman,

817 J. A., Gilman, J. B., Lamplugh, A., Schwantes, R. H., McKeen, S. A., Wisthaler, A., Piel, F., Guo, H., Campuzano-Jost, P.,
 818 Jimenez, J. L., Fried, A., Hanisco, T. F., Huey, L. G., Perring, A., Katich, J. M., Diskin, G. S., Nowak, J. B., Bui, T. P.,
 819 Halliday, H. S., DiGangi, J. P., Pereira, G., James, E. P., Ahmadov, R., McLinden, C. A., Soja, A. J., Moore, R. H., Hair, J.
 820 W., and Warneke, C.: Airborne Emission Rate Measurements Validate Remote Sensing Observations and Emission Inventories
 821 of Western U.S. Wildfires, *Environ. Sci. Technol.*, 56, 7564–7577, <https://doi.org/10.1021/acs.est.1c07121>, 2022.
 822 Turetsky, M., Wieder, K., Halsey, L., and Vitt, D.: Current disturbance and the diminishing peatland carbon sink, *Geophys.*
 823 *Res. Lett.*, 29, <https://doi.org/10.1029/2001GL014000>, 2002.
 824 Veefkind, J. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J., de Haan, J. F., Kleipool, Q.,
 825 van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R.,
 826 Visser, H., and Levelt, P. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the
 827 atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120, 70–83,
 828 <https://doi.org/10.1016/j.rse.2011.09.027>, 2012.
 829 Warneke, C., Schwarz, J. P., Dibb, J., Kalashnikova, O., Frost, G., Al-Saad, J., Brown, S. S., Brewer, Wm. A., Soja, A., Seidel,
 830 F. C., Washenfelder, R. A., Wiggins, E. B., Moore, R. H., Anderson, B. E., Jordan, C., Yacovitch, T. I., Herndon, S. C., Liu,
 831 S., Kuwayama, T., Jaffe, D., Johnston, N., Selimovic, V., Yokelson, R., Giles, D. M., Holben, B. N., Goloub, P., Popovici, I.,
 832 Trainer, M., Kumar, A., Pierce, R. B., Fahey, D., Roberts, J., Gargulinski, E. M., Peterson, D. A., Ye, X., Thapa, L. H., Saide,
 833 P. E., Fite, C. H., Holmes, C. D., Wang, S., Coggon, M. M., Decker, Z. C. J., Stockwell, C. E., Xu, L., Gkatzelis, G., Aikin,
 834 K., Lefer, B., Kaspari, J., Griffin, D., Zeng, L., Weber, R., Hastings, M., Chai, J., Wolfe, G. M., Hanisco, T. F., Liao, J.,
 835 Campuzano Jost, P., Guo, H., Jimenez, J. L., and Crawford, J.: Fire Influence on Regional to Global Environments and Air
 836 Quality (FIREX-AQ), *Journal of Geophysical Research: Atmospheres*, 128, <https://doi.org/10.1029/2022JD037758>, 2023.
 837 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin,
 838 Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat
 839 fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, <https://doi.org/10.5194/acp-10-11707-2010>, 2010.
 840 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E.,
 841 Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth*
 842 *Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
 843 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire
 844 INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model*
 845 *Dev.*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.
 846 Wooster, M. J. and Zhang, Y. H.: Boreal forest fires burn less intensely in Russia than in North America, *Geophys. Res. Lett.*,
 847 31, <https://doi.org/10.1029/2004GL020805>, 2004.
 848 Wooster, M. J., Roberts, G., Perry, G. L. W., and Kaufman, Y. J.: Retrieval of biomass combustion rates and totals from fire
 849 radiative power observations: FRP derivation and calibration relationships between biomass consumption and fire radiative
 850 energy release, *Journal of Geophysical Research Atmospheres*, 110, 1–24, <https://doi.org/10.1029/2005JD006318>, 2005.

851 Wooster, M. J., Xu, W., and Nightingale, T.: Sentinel-3 SLSTR active fire detection and FRP product: Pre-launch algorithm
852 development and performance evaluation using MODIS and ASTER datasets, *Remote Sens. Environ.*, 120, 236–254,
853 <https://doi.org/10.1016/j.rse.2011.09.033>, 2012.

854 Wooster, M. J., Roberts, G., Freeborn, P. H., Xu, W., Govaerts, Y., Beeby, R., He, J., Lattanzio, A., Fisher, D., and Mullen,
855 R.: LSA SAF Meteosat FRP products – Part 1: Algorithms, product contents, and analysis, *Atmos. Chem. Phys.*, 15, 13217–
856 13239, <https://doi.org/10.5194/acp-15-13217-2015>, 2015.

857 Wooster, M. J., Roberts, G. J., Giglio, L., Roy, D., Freeborn, P., Boschetti, L., Justice, C., Ichoku, C., Schroeder, W., Davies,
858 D., Smith, A., Setzer, A., Csiszar, I., Strydom, T., Frost, P., Zhang, T., Xu, W., de Jong, M., Johnston, J., Ellison, L., Vadrevu,
859 K., McCarty, J., Tanpipat, V., Schmidt, C., and San-Miguel, J.: Satellite remote sensing of active fires: History and current
860 status, applications and future requirements, <https://doi.org/10.1016/j.rse.2021.112694>, 15 December 2021.

861 Xu, J., Morris, P. J., Liu, J., and Holden, J.: PEATMAP: Refining estimates of global peatland distribution based on a meta-
862 analysis, *Catena (Amst.)*, 160, 134–140, <https://doi.org/10.1016/j.catena.2017.09.010>, 2018.

863 Xu, W., Wooster, M. J., Roberts, G., and Freeborn, P.: New GOES imager algorithms for cloud and active fire detection and
864 fire radiative power assessment across North, South and Central America, *Remote Sens. Environ.*, 114, 1876–1895,
865 <https://doi.org/10.1016/j.rse.2010.03.012>, 2010.

866 Xu, W., Wooster, M. J., Kaneko, T., He, J., Zhang, T., and Fisher, D.: Major advances in geostationary fire radiative power
867 (FRP) retrieval over Asia and Australia stemming from use of Himarawi-8 AHI, *Remote Sens. Environ.*, 193, 138–149,
868 <https://doi.org/10.1016/j.rse.2017.02.024>, 2017.

869 Xu, W., Wooster, M. J., He, J., and Zhang, T.: First study of Sentinel-3 SLSTR active fire detection and FRP retrieval: Night-
870 time algorithm enhancements and global intercomparison to MODIS and VIIRS AF products, *Remote Sens. Environ.*, 248,
871 111947, <https://doi.org/10.1016/j.rse.2020.111947>, 2020.

872 Xu, W., Wooster, M. J., He, J., and Zhang, T.: Improvements in high-temporal resolution active fire detection and FRP retrieval
873 over the Americas using GOES-16 ABI with the geostationary Fire Thermal Anomaly (FTA) algorithm, *Science of Remote*
874 *Sensing*, 3, 100016, <https://doi.org/10.1016/j.srs.2021.100016>, 2021.

875