

1 Mitigating *Mazuku* Hazards: Implementation and Effectiveness of Local Dry-Gas 2 Degassing Measures in the Goma Area (Virunga Volcanic Province)

3 Blaise Mafuko-Nyandwi*

4 Ecole Supérieure de Volcanologie, de Gestion des Risques et des Catastrophes, Université de Goma
5 (Campus du Lac, Rue Eugène Serufuli 43, Quartier Katindo, Goma, North Kivu province, Congo)

6 *Corresponding authors: BMN, blaise.mafuko.nyandwi@unigom.ac.cd or tbmafuko@gmail.com

7 **Keyword:** Volcanic risk, risk mitigation, carbon dioxide, Nyiragongo

8 1. Abstract

9 Mitigation of carbon dioxide diffuse degassing hazards remains underexplored in comparison to
10 other volcanic hazards such as eruptions, despite their persistent and deadly impacts on
11 communities living in active volcanic regions. This study uses a mixed-methods approach—
12 combining quantitative surveys and qualitative interviews—to assess household perceptions of
13 the implementation and effectiveness of risk mitigation measures against *mazuku*, a locally
14 known hazard caused by emissions of carbon dioxide in the western part of Goma, Virunga
15 Volcanic Province. Data were collected across three sampling zones, capturing demographic
16 characteristics, eruption risk experiences, and perceptions regarding the implementation of
17 *mazuku* risk mitigation measures.

18 Findings reveal three locally recognised categories of mitigation measures: (1) emission-limiting
19 measures, such as blocking gas with waste materials; (2) adaptive measures, such as house
20 ventilation or living on upper floors; and (3) awareness measures based on orally transmitted
21 local knowledge such as avoiding *mazuku* zone early morning. Financial resources, gender and
22 prior risk experience—often linked to length of residence—emerged as significant positive
23 determinants of both motivation and perceived efficacy for the first two categories. Perceptions
24 of awareness measures showed no significant variation across zones even between demographic
25 profile groups. Spatial patterns in perceived implementation and perceived efficacy appear to
26 reflect collective community mitigation implementation rather than based on individual risk
27 mitigation assessment, with some measures perceived as effective despite limited physical
28 evidence of reduced gas concentration.

29 The study supports the importance of co-creating mitigation strategies with local communities,
30 adapting interventions to socio-economic realities and avoiding the importation of external
31 mitigation measures that may lack contextual relevance. It also calls for complementary research
32 measuring the actual effectiveness of these measures through physical monitoring of *mazuku*
33 concentrations. These insights, grounded in a Global South context—characterised by rapid
34 uncontrolled urbanisation, offer a valuable perspective for the development of inclusive and
35 effective strategies of carbon dioxide diffuse degassing risk management.

36 **2. Introduction**

37 Volcanic hazards are the surface manifestations of Earth's internal activity. They can be short-
38 lived, such as eruptions, or long-term, like carbon dioxide (CO₂) diffuse degassing and
39 hydrothermal activities(Loughlin et al., 2015). Despite the dangers posed by these hazards,
40 numerous societies have settled near active volcanoes (Brown & Jenkins, 2017), including in
41 areas with intense CO₂ diffuse degassing. Exposition to CO₂ diffuse degassing represents a
42 significant threat to human health and safety (Edmonds et al., 2017; Hansell & Oppenheimer,
43 2004). The CO₂, an odourless and colourless gas, acts as an inert asphyxiant and displace oxygen
44 in the air down to dangerously low levels. Lethal concentrations—exceeding 10 vol.%— cause
45 rapid loss of consciousness, asphyxiation, and death of human and nonhuman beings(Viveiros &
46 Silva, 2024).

47 The short-term exposure limit for CO₂ is set at 3 vol%, while the permissible limit for an 8-hour
48 exposure is 0.5 vol% (Hansell & Oppenheimer, 2004). When these thresholds are exceeded,
49 specific symptoms may appear depending on the concentration level and duration of exposure.
50 These include accelerated breathing and increased heart rate, followed by dyspnoea and
51 headaches, and in more severe cases sweating, dizziness, ringing in the ears, vertigo, vomiting,
52 and muscular weakness (Viveiros et al., 2016). Viveiros et al. (2024) note that although CO₂
53 diffuse degassing is often considered a neglected natural hazard, it has caused the deaths of more
54 than 2,000 people over the past decades. Considering the potential impact of CO₂ on human
55 health and its silent infiltration into buildings in diffuse degassing areas, studies on their
56 mitigation measures are crucial to inform disaster risk mitigation programs.

57 This paper examines the diffuse CO₂ degassing (called mazuku) in the Goma area in eastern
58 Democratic Republic of the Congo (DR Congo), located within the Virunga Volcanic Province.
59 Locally in Goma, the term mazuku is derived from Swahili and translates as “evil wind” (or “evil
60 wind that spreads and kills during the night”). It is used to refer both to the diffused CO₂-rich gas
61 and the areas where it is emitted. The hazardous gas accumulates in low-lying depressions where
62 they become concentrated due to their heavier-than-air nature (Wauthier et al., 2018). CO₂
63 concentrations in Mazuku can largely exceed the minimum exposure limits for humans or fauna,
64 reaching high concentration ranging from 45 to 80 vol%, with diurnal–nocturnal fluctuations of
65 up to 80% (Balagizi et al., 2018a). The rapid growth of Goma, as observed in many cities of the
66 Global South (Quesada-Román, 2022), has been driven largely by intense migration linked to
67 recurrent armed conflicts in the region and the search for professional opportunities (Pech et al.,
68 2018; Pech & Lakes, 2017). This expansion has extended the city westward into areas highly
69 concentrated in *mazuku*, thereby exposing a large population.

70 However, previous mazuku related studies in the region have focused primarily on hazard
71 assessments, including its formation, vent locations, and the geographical distribution of its
72 concentrations (M. M. Kasereka et al., 2017; Smets et al., 2010; Wauthier et al., 2018), or
73 evaluating the changes in its magnitude following a volcanic eruption (Vaselli et al., 2003). To
74 date, *mazuku* mitigation measures are poorly studied. In addition, it has been observed in the
75 region that while official awareness campaigns encourage people to avoid high-risk areas by
76 installing warning panels that call people to avoid settling in mazuku, these signs are frequently
77 removed. Residents continue to stay or others to come and settle in known hazardous zones and
78 subsequently they develop their own local mitigation strategies.

79 In this perspective, this study seeks to examine household representatives’ perceptions regarding
80 the implementation of local mazuku mitigation measures, focusing on their perceived efficacy,
81 associated costs, extent of implementation, and motivations for adoption. To achieve this, the
82 research employs a mixed-methods approach. Qualitative data were collected through 32
83 interviews and three focus group discussions, identifying, describing, and categorizing 12
84 principal local mitigation measures. Additionally, a large-scale survey of over 500 households
85 was conducted to evaluate quantitatively public perceptions regarding the implementation of
86 these measures.

87 This study provides a new perspective on volcanic disaster risk management, highlighting that in
88 a context of scarcity of risk information and mitigation strategies, exposed population developed
89 their own mitigation measures. This makes individual mitigation an imperative if there are no
90 other options for where to live. It means, for instance, when the necessity of settling in volcanic
91 areas with high concentrations of mazuku outweighs the risks of living in regions around Goma
92 affected by armed conflict, the local population seeks to work out practical strategies to mitigate
93 the mazuku hazard and therefore resettle or remain in these high-risk zones. Consequently,
94 hazard mitigation strategies that incorporate local practices prove more effective than those
95 imported from outside (Lutete Landu et al., 2023).

96 After this introduction, this article provides a detailed overview of mazuku within the human,
97 geological, and geographical context of the Goma region. Then it presents the used methodology
98 and the results followed by a discussion both on the challenges and successes in implementing
99 local mazuku mitigation measures. The paper concludes with key insights and recommendations
100 to strengthen volcanic risk mitigation measures among local communities, drawing on evidence
101 from this case study of Goma.


102 **3. Goma region: socio-environmental setting and *mazuku* formation**

103 Goma is located in eastern DR Congo on the northern shore of Lake Kivu (~1,460 m a.s.l.), at
104 the international border with Rwanda (Gisenyi), within the Virunga Volcanic Province (VVP) of
105 the western branch of the East African Rift System (Smets, 2016; Vlassenroot & Büscher, 2011).
106 The city is built entirely on the active lava field of Mount Nyiragongo (3,470 m a.s.l.), situated
107 about 18 km north of Goma inside Virunga National Park (Fig1). Nyiragongo is a rift-controlled
108 stratovolcano with a summit crater of about 1.2 km width and a semi-persistent lava lake
109 (Barrière et al., 2022). Fractures and eruptive fissures on its southern flank strongly control lava
110 pathways (Kervyn et al., 2024). Historical flank eruptions (1977, 2002 and 2021) produced
111 highly fluid lava flows directed towards Goma and the Lake Kivu basin (Kervyn et al., 2024;
112 Smets, 2016). In addition to lava flows, the region is exposed to other volcanic hazards. These
113 include persistent SO₂-rich plumes that can degrade air quality and promote acid deposition
114 (Smets et al., 2017), diffuse CO₂ degassing (Smets et al., 2010), and recurrent seismicity (Subira
115 Muhindo, 2024) and ground deformation linked to magma intrusions along rift structures

116 (Geirsson et al., 2017; Smittarello et al., 2022). Furthermore, lake Kivu contains large amounts
117 of dissolved carbon dioxide (CO₂) and methane (CH₄), which represent a potential limnic gas-
118 release hazard in case of lake destabilisation (Hirslund & Morkel, 2020).

119 Despite its exposure to multiple natural hazards, Goma is a rapidly expanding border city whose
120 urban development has been strongly shaped by ongoing conflict, the presence of the
121 humanitarian sector, and cross-border trade—including natural-resource economics (Pech et al.,
122 2018; Vlassenroot & Büscher, 2009). Populations repeatedly seek refuge mostly from insecurity
123 affecting surrounding areas of eastern DRC. These dynamics have contributed to intense and
124 largely informal urbanisation (Pech et al., 2018). Between 2002 and 2020, the city's population
125 doubled from around half a million to more than one million inhabitants, increasing settlement
126 density and demand for housing and services (Adalbert et al., 2021). At the same time, the city's
127 physical expansion is strongly constrained by Lake Kivu to the south, Virunga National Park to
128 the northwest and the Rwanda border to the east, forcing growth northwards towards
129 Nyiragongo's flanks and recent lava fields even in mazuku-affected area (Fig.1),

130 The Mazuku-affected area under study, which is now inhabited (Fig.1), was unoccupied three
131 decades ago (Pech et al., 2018). At that time, the region was covered by an open woodland
132 typical of the area. According to testimonies gathered from local elders, people used to cross it at
133 dusk to reach the lake Kivu for fishing, or early in the morning when returning with their catch.
134 It also served as a hunting ground for Gambian rats and as pastureland for livestock before it was
135 settled. These activities mostly took place in the evening or early morning when the mazuku
136 concentration is high. Therefore, many people, as well as livestock, lost their lives asphyxiated,
137 which was then regarded as an evil wind—one that had neither a smell nor a visible form
138 (Vaselli et al., 2003). Today, the area is inhabited by new residents and Internally Displaced
139 Persons (IDP) with a more urban lifestyle than the earlier inhabitants and the term was kept.



140

Figure 1: Study area of the mazuku zone within the Nyiragongo lava flows (Goma, eastern DR Congo). The map shows the extent of the 1977, 2002 and 2021 lava flows, eruptive fissures, urban areas, and the sampling zone investigated in this study. Part of shapefiles credit: GeoRiskA, (<https://georiska.africamuseum.be/>)

141 Mazuku denotes depressions into which dense CO₂—heavier than air—emanates and
 142 accumulates (Fig.2). Such phenomena also occur in other volcanic areas around the globe,
 143 including Mammoth Mountain (USA), Royat (France), and the Siena Graben (Italy); however,
 144 they differ in terms of both gas origin mechanisms and patterns of human occupancy(Edmonds et
 145 al., 2017; Hansell & Oppenheimer, 2004). Despite their long-standing recognition, the formation
 146 mechanisms of these gases remain poorly understood and widely debated (Williams-Jones &
 147 Rymer, 2015).



148

Figure 2: Picture of warning panel. The Mazuku emission zone is outlined with red dotted lines, and in the middle stands a warning panel located within an IDP camp. The inscriptions on the warning panel are in French, with a Swahili translation, reading: “High-risk zone. Beware of gas. Watch over and prevent children from playing near gas areas.” Nearby, we can also observe (a) public latrines with uncemented septic pits, and (b) small tent shelters occupied by the IDPs (Photo credit Blaise Mafuko).

149


150 In the VPP they are common in the vicinity of Goma—particularly between Lake Kivu and the
 151 west part of lava flow fields of Nyiragongo and Nyamuragira (Smets et al., 2010). Wauthier et
 152 al., (2018) explain that these occur where a deep magmatic CO₂ source connects to the surface
 153 via a network of fractures, and where topographical depressions enable the gas to settle. The
 154 expansion of Goma has led to the occupation of lakeshore areas in the west of the city, along
 155 Lake Kivu (Büscher & Vlassenroot, 2010; Pech et al., 2018), where these mazuku are highly

156 concentrated. The official mitigation strategy involves mapping gas-emission zones and
157 installing warning panels. Nonetheless, these *mazuku* continue to cause fatalities over extended
158 periods, and livestock asphyxiation remains a frequent occurrence.

159 **4. Methodology**

160 **4.1. Data collection**

161 This study adopted a sequential mixed-methods design integrating qualitative and quantitative
162 approaches (Fig. 3). We first conducted semi-structured interviews in neighbourhoods previously
163 identified by the Goma Volcanological Observatory (GVO) as *mazuku* emission zones to
164 document local experiences and identify existing mitigation measures. We then organised three
165 focus group discussions (with community representatives, street leaders, and manual septic-pit
166 diggers) to validate, clarify, and categorise the measures and refine contextual understanding of
167 their implementation. Finally, insights from these qualitative steps informed the construction of a
168 questionnaire survey, which was administered at scale to quantitatively assess household
169 perceptions of the identified measures, particularly regarding perceived effectiveness, costs,
170 levels of implementation, and motivations for adoption. In this study, street leaders refer to
171 locally recognised neighbourhood-level representatives who coordinate 10 households and
172 facilitate communication between residents and local authorities.



173

Figure 3: Sequential mixed-methods design showing the intersections between qualitative interviews, focus group discussions, and the questionnaire survey used to identify, validate, categorise, and quantitatively assess local mazuku mitigation measures. Icons credit: Shutterstock (<https://www.shutterstock.com/>).

174

175 4.1.1. The interviews

176 The interviews were conducted between 1 October and 10 October 2024. We interviewed 32
 177 individuals—17 women and 15 men—focusing exclusively on adult household heads.
 178 Participants were selected at random, with an aim of interviewing three people per main street:
 179 one at the beginning, one in the middle, and one at the end. The entire area identified by the
 180 Goma Volcanological Observatory as a high-risk mazuku zone was covered. Verbal consent was
 181 obtained from all participants prior to the interviews. The interviews were structured and
 182 addressed the following themes: (1) the respondent's experience of volcanic risk in Goma; (2)
 183 their knowledge of the existence and formation of mazuku; (3) indicators used to identify areas
 184 with high mazuku concentrations; (4) impacts recorded as a result of mazuku exposure; and (5)
 185 mitigation measures against mazuku-related risks.


186 4.1.2. Focus groups

187 In addition to the interviews, we organised three focus group discussions (FGD) towards the end
 188 of October 2024. The FGDs covered the same themes as the interviews but adopted a debate-

189 based approach among participants to identify the spatial and daily temporal variations in the
190 occurrence of mazuku. The first FGD brought together 10 participants, including 5IDPs and
191 local residents. The aim was to capture differences in perception between the various social
192 groups living in the same area. The second FGD comprised 8 men who manually dig septic pits.
193 They work in the area extracting stones for sale as well as digging toilet septic pits. They are
194 familiar with the history of land occupation and are well aware of the areas with high gas
195 concentrations, although without any scientific assessment of the levels. This discussion enabled
196 the oral history of the area's occupation to be reconstructed.

197 Finally, we brought together 9 street leaders to discuss the same themes, with a stronger focus on
198 local mechanisms for managing this risk. The FGDs concluded with a walk-through in the area
199 for observations involving 4 street leaders, 3 diggers, and 3 community members who were
200 available. This exercise allowed us to distinguish 3 types of land occupation according to the
201 nature of the houses and the period of settlement (Fig.4): a highly urbanised area occupied by
202 high-income residents; a transitional area undergoing urbanisation with sporadic permanent
203 constructions; and a rural area mainly inhabited by indigenous populations and IDPs.

204



205

Figure 4: **Maps of the Study Area:** Map (A) shows the location of the city of Goma, and the lava flows from the last three eruptions of Nyiragongo. Map (B) indicates the three sampling zones and the pattern of housing structures, derived from Google Earth.

206

207 4.1.3. Questionnaire survey

208 The data gathered from qualitative evaluations enabled us to describe and classify 12 risk
 209 mitigation measures (Mafuko Nyandwi, 2025). Subsequently, we conducted a large-scale
 210 survey—carried out by trained enumerators—to assess population perception regarding the
 211 implementation of these measures.

212 The questionnaire focused on:

- 213 1. **Demographic profile:** including participants' age, gender, experience with risk, household
 214 size, monthly household income, number of rooms in the house, duration of residence, and
 215 residential status.

216 2. **Perceptions of measure implementation:** covering respondents' individual motivation to
217 implement each mazuku mitigation measure over the next six months; the perceived
218 efficacy of each measure in reducing risk across within their neighbourhood; the perceived
219 cost of implementation; and finally, how they perceived the current level of
220 implementation of each measure within their neighbourhood.

221 The sample size was determined based on the population of the Goma targeted neighbourhoods
222 (Kyeshero and Lac Verts). With an estimated population of approximately 100,000—according
223 to data collected from the respective neighbourhoods offices during our survey—our sample of
224 573 individuals at a 95 % confidence level far exceeded the minimum required for statistical
225 representativeness (Morgan, 1970).

226 We randomly distributed around 600 sampling points over a Landsat image from Google Earth,
227 across the identified high-risk mazuku zone, maintaining an approximately equidistant spacing of
228 40 m between points. Enumerators were instructed to survey the household closest to each
229 sampling point, following a previous developed protocol (Mafuko Nyandwi, Kervyn,
230 Habiyaemye, et al., 2023). We targeted only adult household heads as respondents.

231 **4.2. Data analysis**

232 The qualitative data were analysed using content analysis to list all mazuku mitigation measures,
233 followed by thematic analysis to identify recurring patterns and key themes related to their
234 implementation and categorisation. We then employed descriptive statistics to characterise the
235 measures by evaluating the proportion of the population at each level of perception. Cronbach's
236 alpha was used to measure internal consistency across the three categories of mitigation
237 measures, enabling us to aggregate motivation and perceived efficacy within each group.
238 Aggregation was performed using the mean when the coefficient of variation (CV) was less than
239 25 %, and the median when the CV was 25 % or higher — the CV, being the ratio of standard
240 deviation to the mean, provides a standardised measure of variability.

241 Non-parametric tests were applied to assess how motivation for implementation varied across
242 demographic variables. Statistically significant variations were represented on boxplots. Pairwise
243 Spearman's rank-order correlations were calculated to evaluate the strength and direction of
244 monotonic relationships between ranked variables—motivation, perceived efficacy, and

245 perceived cost—and the results were visualised using bar charts that display the correlation
246 coefficient for each pair. Finally, chi-square tests were conducted to evaluate spatial variations in
247 aggregated efficacy and the level of implementation of each measure across the three sampling
248 zones.

249 **5. Results**

250 **5.1. Demographic profile of participants**

251 Our survey targeted only adult household heads (Table 1). The majority of these heads were
252 under 45 years of age (77.31%), with the majority of respondents being women (61.78%).
253 Households are large. Over 80% have between 4 and 10 members. Despite this large household
254 size, the average monthly income per household remains very low, with 58.12% of households
255 living on around USD 150 per month and 28.97% on an income of between USD 151 and USD
256 300. This situation is even more pronounced in zone 3, where almost all households (91.5%) live
257 on less than USD 150 per month. Zone 3 is more unusual in that it is home to more displaced
258 people from the wars than the other zones. Zone 1, which is located further east, i.e. on the city
259 centre side, has the lowest proportion of war-displaced people (8.9%). Generally speaking, the
260 western part of Goma that we surveyed had a high rate of new arrivals. 62.13% had lived there
261 for less than 5 years and 22.16% for between 6 and 11 years.

262

Table 1: Demographic characteristics of respondents²⁶³

Age (years)	18-30	31-45	46-55	56-65	Over 65
Zone 1	46 (24%)	93 (48.4%)	40 (20.8%)	8 (4.2%)	5 (2.6%)
Zone 2	39 (20.3%)	109 (56.8%)	33 (17.2%)	9 (4.7%)	2 (1%)
Zonz 3	82 (43.4%)	74 (39.2%)	18 (9.5%)	11 (5.8%)	4 (2.1%)
Total per age group	167 (29.14%)	276 (48.17%)	91 (15.88%)	28 (4.89%)	11 (1.92%)

Income (USD)	0-151	151-300	301-450	451-600	Over 600
Zone 1	60 (31.2%)	73 (38%)	38 (19.8%)	17 (8.9%)	4 (2.1%)
Zone 2	100 (52.1%)	79 (41.1%)	11 (5.7%)	2 (1%)	0 (0%)
Zonz 3	173 (91.5%)	14 (7.4%)	2 (1.1%)	0 (0%)	0 (0%)
Total per income range	333 (58.12%)	166 (28.97%)	51 (8.90%)	19 (3.32%)	4 (0.70%)

Household size (persons)	1-3	4-6	7-10	11-13	Over 15
Zone 1	17 (8.9%)	63 (32.8%)	97 (50.5%)	12 (6.2%)	3 (1.6%)
Zone 2	17 (8.9%)	106 (55.2%)	64 (33.3%)	5 (2.6%)	0 (0%)
Zonz 3	17 (9%)	74 (39.2%)	88 (46.6%)	10 (5.3%)	0 (0%)
Total per size range	51 (8.90%)	243 (42.41%)	249 (43.46%)	27 (4.71%)	3 (0.52%)

Eruption experience	No	2021	2002	2002&2021	1977 2002&2021
Zone 1	20 (10.4%)	81 (42.2%)	2 (1%)	85 (44.3%)	4 (2.1%)
Zone 2	45 (23.4%)	52 (27.1%)	7 (3.6%)	80 (41.7%)	8 (4.2%)
Zonz 3	85 (45%)	65 (34.4%)	0 (0%)	38 (20.1%)	1 (0.5%)
Total per experience group	150 (26.18%)	198 (34.55%)	9 (1.57%)	203 (35.43%)	13 (2.27%)

Duration of residence	0 to 5 yrs	6 to 11 yrs	12 to 16 yrs	17 to 21 yrs	26 yrs and more
Zone 1	108 (56.2%)	47 (24.5%)	20 (10.4%)	11 (5.7%)	6 (3.1%)
Zone 2	115 (59.9%)	51 (26.6%)	20 (10.4%)	2 (1%)	4 (2.1%)
Zonz 3	133 (70.4%)	29 (15.3%)	11 (5.8%)	4 (2.1%)	12 (6.3%)
Total per duration	356 (62.13%)	127 (22.16%)	51 (8.90%)	17 (2.97%)	22 (3.48%)

Residence status	IDP	Inhabitant
Zone 1	17 (8.9%)	175 (91.1%)
Zone 2	64 (33.3%)	128 (66.7%)
Zonz 3	84 (44.4%)	105 (55.6%)
Total per status	165 (28.80%)	408 (71.20)

Gender	Female	Male
Zone 1	132 (68.8%)	60 (31.2%)
Zone 2	81 (42.2%)	111 (57.8%)
Zonz 3	141 (74.6%)	48 (25.4%)
Total per gender group	354 (61.78%)	219 (38.22%)

264

265 5.2. Description of mitigation measures

266 Through the analysis of interview discourse, we identified 12 key local strategies for mazuku
 267 risk mitigation. Additionally, follow-up focus group discussions, held in a participatory manner,

268 enabled the classification of these measures into three categories based on whether they aim to
269 prevent mazuku, reduce its impact, or inform the population of its occurrences.

270 For preventing mazuku emission, on the one hand, local residents explained that they use
271 household waste mixed with mud to cover areas emitting mazuku (measure 1), hoping to reduce
272 gas emission. On the other hand, households with sufficient financial means tend to cement all
273 potential emission points within their plots with concrete, such as house floors (measure 2),
274 courtyards (measure 3), and septic systems (measure 4).

275 *“We use household waste mixed with mud to cover the mazuku areas, hoping to reduce the*
276 *emissions, especially when the mazuku is located in a public area ... These zones are already*
277 *known to us, so we organise regularly community works to prevent or reduce the mazuku*
278 *emissions.”*

279 (Elderly man, street leader, 16 years living in a mazuku zone)

280 *“Some houses have uncemented floors, so mazuku emissions can occur in bedrooms or living*
281 *rooms... When households have the financial means, they cement all potential emission sources*
282 *like septic tanks or backyards. But for public spaces, we mostly use household waste.”*

283 (27-year-old woman, born, raised, and now married in the same mazuku area)


284 When it is not possible to prevent the emission of mazuku, local communities have developed
285 adaptive strategies and or convey local knowledge—passed down orally from generation to
286 generation and also between long-time residents to newcomers—to help avoid high-risk areas
287 within neighbourhoods or in public areas.

288 To cope with high concentrations of mazuku within their homes, residents elevate beds (measure
289 1), live on upper floors when available (measure 2), or improve ventilation by enlarging
290 windows and keeping them open during the day or sometimes at night (measure 3). In cold
291 conditions, certain households reported heating courtyards or indoor areas to facilitate the
292 dispersion of mazuku (measure 4). In addition, to keep the wider community informed about
293 mazuku occurrences, residents raise awareness about avoiding known mazuku zones. They are
294 encouraged to work together with officials from the volcanic observatory and civil protection to
295 install warning signs and follow their instructions (measure 1). Furthermore, based on

296 experience, it is best to avoid mazuku particularly in the early morning (measure 2) or after
297 rainfall (measure 3). For those raising livestock or poultry, it is recommended that animals be
298 kept in very well-ventilated areas (measure 4). Descriptive statistics further characterise these
299 measures by examining individual perceived motivation, response efficacy, associated costs, and
300 levels of implementation.

301 **5.2.1. Level of perception of blocking gas emission measures**

302 Measures aimed at blocking *mazuku* emissions that require greater financial resources—such as
303 cementing different parts of the household environment—were evaluated similarly by the
304 population (Fig.5). The majority perceive these measures as costly, although nearly all agree that
305 they are effective or very effective. Their perceived high cost may explain the mixed views when
306 it comes to households to evaluate their motivation for their implementation. Among this group
307 of measures, the highest proportion (53%) of respondents reporting a high or very high
308 motivation to implement relates to the use of household waste—a measure which, as expected, is
309 perceived by the majority (68%) as having low or very low cost and perceived to be largely
310 implemented in the zone



311

Figure 5: (A) Level of perceptions of different indicators for blocking gas mitigation measure. The percentages on the left indicate the proportion who perceived this likelihood as low or very low, while the middle percentages represent those with a moderate perception of likelihood. (B) The level of implementation


312

313 5.2.2. Level of perception of adaptive mitigation measures

314 Opinions are divided when it comes to evaluating the motivation, perceived efficacy, and even
 315 the cost associated with measures such as raising the bed level or improving house ventilation
 316 (Fig.6). Yet, among these adaptation strategies, these two are the most widely implemented in
 317 the region. The least implemented are heating the courtyard or living upstairs. An elder from the
 318 neighbourhood offers insight into why:

319 *“We burn dry grass or sometimes cardboard boxes from nearby shops—especially when the cold*
 320 *persists for over 24 hours—to help evaporate the mazuku. Living upstairs is certainly better, but*
 321 *not everyone can afford it. My neighbour, who has an upper-floor dwelling, told me that all the*

322 bedrooms are upstairs to avoid being caught unawares at night by a high mazuku concentration.
 323 On particularly cold days, he said that his family decide outright not to stay on the ground floor
 324 at all.”



325

Figure 6: (A) Level of perceptions of different indicators for adaptative mitigation measure (B) The level of perceived implementation within the neighbourhood

326

327 **5.2.3. Level of perception of community based awareness measures**

328 Knowing which areas have high concentrations of mazuku—so as to avoid them in the early
 329 morning, during rain, or simply when temperatures drop during the day—is among the most
 330 widely implemented measures (Fig.7). Approximately 85 % of the population report that these
 331 two measures are effective and they are motivated to implement them. As might be expected,

332 nearly everyone surveyed—around 90 %—perceive their implementation cost to be very low,
333 which may explain why they are so frequently adopted.

334 *Mazuku incidents tend to be more concentrated in the evening or early morning, and when the*
335 *temperature is low especially during the rainy seasons. You cannot see the mazuku or detect any*
336 *odour, but sometimes, on a path, you suddenly feel suffocated as though someone were pressing*
337 *on your chest, and you cannot breathe. At that moment you must act quickly and leave the area*
338 *while you still have strength....*


339 *Just after the dry season — at the beginning of September when children return to school — the*
340 *first critical period begins and lasts until December. It is followed by a second critical period*
341 *during the second rainy season, from February to May every year. These periods are*
342 *particularly hazardous because they coincide with the school term, when children have to leave*
343 *home early for classes.*

344 *In this context, we do our best to inform our children, newcomers or everyone in the*
345 *neighbourhood, about the locations of these mazuku zones: we encourage them to identify them*
346 *and to stay well away from them, especially when it is cold.*

347 (A mother of 4 children at primary school, 13 years of residence in the area)

348 A significant proportion of respondents (75 %) believe that installing panels is effective in
349 reducing the risk of mazuku exposure; however, opinions remain divided when it comes to
350 motivation to implement or the cost of installation. Similarly, views are mixed regarding the
351 measure of keeping livestock or poultry in well-ventilated spaces.

352



353

Figure 7: (A) Level of perceptions of different indicators for Community based awareness measures, (B) The level of perceived implementation within the neighbourhood

354

355 **5.2. Factors of the motivation for implementing mitigation measures**

356 Only the aggregated indicators for motivation to implement preventive mazuku emission
 357 measures and adaptive strategies showed statistically significant variation across demographic
 358 groups (Appendix 1). No significant differences were found in overall motivation levels based on
 359 local awareness measures. Financial conditions—specifically household monthly income and the
 360 number of rooms in a dwelling—were positively associated with motivation to implement both
 361 types of measures (Fig.8).




Figure 8: The level of perceptions of the aggregated indicator according to significant determining factors. Perceptions are expressed on a numerical scale from 1 (very low) to 5 (very high). In each boxplot, the horizontal bold line represents the median, the red dot indicates the mean, and the small circles represent outliers. The letter on top on boxplots represents the post-hoc test results between groups of the same aggregate indicator not between the same group between two indicators.

363

364 Motivation levels for both preventive and adaptive measure increased with age and length of
365 residence, but only up to a certain point. Beyond approximately 46 years of age, or after more
366 than 17 years living in the area, motivation declined and then plateaued. Men exhibited higher
367 motivation to implement these measures than women. Furthermore, individuals who had not
368 previously experienced volcanic risk showed lower implementation willingness; however, their
369 willingness increased with the number of personal experiences of Nyiragongo eruption risk.

370 **5.3. Correlations**

371 Pairwise Spearman's rank-order correlations indicate that perceived efficacy is a stronger driver
372 of motivation than cost perceptions, although cost can either reinforce or hinder motivation
373 depending on the type of measure. Figure 9A shows that most measures have a strong and
374 statistically significant positive correlation between efficacy and motivation, particularly for
375 measures such as blocking gas with waste materials or raising beds to adapt to gas emissions.
376 This suggests that higher perceived effectiveness is consistently associated with a stronger
377 willingness to implement these measures. However, there is no relationship between motivation
378 and perceived efficacy for the measure of installing warning panels, which may be due to the fact
379 that this intervention depends on disaster risk authorities rather than the community.

380 Figure 9B also shows that there are mostly positive, though generally weak, relationships
381 between perceived efficacy and cost. Notably, for the awareness measures of avoiding high gas
382 areas in the early morning or after rainfall, there is no association between perceived efficacy and
383 cost. Figure 9C reveals a more mixed pattern between cost and motivation: while certain
384 adaptive and awareness measures (Measures 5, 6, 7, and 12) display a significant positive
385 association, some blocking measures (e.g., Measure 2) are negatively correlated, indicating that
386 higher perceived costs may discourage willingness to implement those interventions.



387


Figure 9: Pairwise Spearman's rank-order correlations. *** p value<0.001, ** pvalue<0.01 and * p value<0.1.

388

389 5.4. Spatial variation

390 The figure 10 presents the variation in the population's perceptions of efficacy across the
 391 sampling zones. It shows that aggregated efficacy is perceived very differently across the three
 392 sampling zones, with statistically significant differences. Zone 2 hosts a large proportion of the
 393 population who consider both awareness measures and those limiting mazuku emissions to be
 394 effective or even very effective. In contrast, zone 3 is home to the majority of people who regard
 395 emission-limiting or adaptation measures as ineffective. When grouping together those who
 396 perceive the measures as effective and those who consider them very effective, we find almost
 397 the same proportion of the population in Zone 3 regardless of the type of measure.

398



399

Figure 10: Spatial variation of perceived efficacy across different sampling zones⁴⁰⁰ We also
⁴⁰¹ assessed the
⁴⁰² variation in the perceived level of implementation for each measure within each sampling zone
⁴⁰³ (Annex B). It is evident that measures requiring substantial resources, regardless of their
⁴⁰⁴ category, are perceived as not implemented by a large proportion of the population in Zone 3
⁴⁰⁵ (over 65% to 85%). This is the case, for example, for heating or cementing courtyards, living on
⁴⁰⁶ upper floors or raising bed heights. In contrast, for the measure involving the use of waste
⁴⁰⁷ materials to limit mazuku emissions, only 24% of the population in Zone 3 perceive it as not
⁴⁰⁸ implemented. Awareness measures, such as identifying mazuku-prone areas for avoiding them
⁴⁰⁹ during cold periods (in the morning or after rainfall), are the most widely perceived as
⁴¹⁰ implemented across all three zones, although the proportions of the population in their perception
⁴¹¹ category vary by zone.

412

413 6. Discussions

414 6.1. Passive Risk Acceptance: Motivation and Efficacy Constrained by Limited Living 415 Options and resources scarcity

416 By 2019, one billion people were already living within 100 km of active volcanoes, with the
417 density of human activities continuing to increase (Brown et al., 2015; Freire et al., 2019). In
418 CO₂ diffused degassing zones not restricted as parks or reserves (Williams-Jones & Rymer,
419 2015), people may choose to reside in areas with CO₂ high-concentrations (Edmonds et al.,
420 2017; Hansell & Oppenheimer, 2004a, 2004b), as in the present case study. This may reflect a
421 risk acceptance. However, our findings indicate a more specific form of *passive* risk
422 acceptance (Wachinger et al., 2013b, 2018). Indeed, people are well aware of the risk posed by
423 mazuku and claim to know where they are located, yet many still choose to live close to, or even
424 on them. This could suggest that they have no other options left. Indeed, in Goma—a city
425 already extremely densely populated (Pech et al., 2018; Pech & Lakes, 2017) —people often
426 settle in these risky areas because, despite the volcanic hazards, Goma is perceived as safer than
427 the conflict-affected surrounding regions (Mafuko Nyandwi, Kervyn, Habiyaremye, et al., 2023;
428 Mafuko Nyandwi, Kervyn, Muhashy Habiyaremye, et al., 2023) Therefore, people have
429 developed local mitigation measures to compensate for the insufficiency of the official advice to
430 simply avoiding the area, as indicated on warning panels.

431 Wachinger et al. (2013a) describe this as the risk-mitigation paradox—a situation in which
432 people consciously choose to live exposed to hazards, and the choices of mitigation measures
433 being controlled by resource availability. In such contexts, most participants report being
434 motivated to identify high-concentration areas in order to avoid them during critical times, such
435 as early mornings or after rainfall, when mazuku concentration is high. Being less resource-
436 intensive, awareness-based measures were widely considered effective by the majority,
437 particularly among low-income households, who also felt these measures had been largely
438 implemented. Therefore, mazuku are perceived as a daily threat that can be controlled — through
439 preventive measures, awareness of high-risk zones and times of day, or by adapting the
440 environment (for example improved ventilation) to reduce its magnitude. It suggests that living
441 in a zone prone to mazuku gives rise to a widespread, yet often unrecognised, acceptance of risk.
442 Inhabitants develop everyday routines and coping practices in response to repeated exposure

443 (Walshe et al., 2023), and over time these behaviours become internalised and incorporated into
444 the community’s habitus—defined by (Bourdieu, 1990) as the set of structured dispositions
445 through which individuals perceive the hazardous environment and act in it. In effect, what
446 begins as a mitigation strategy gradually solidifies into a socialised readiness to ‘live with’ the
447 hazard rather than to challenge or transform it (Vergara-Pinto & Marín, 2023; Walshe et al.,
448 2023). This suggests that mazuku becomes embedded in the routines of everyday life, gradually
449 normalised, and that the mitigation practices sustaining a perceived sense of “safe exposure” are
450 reproduced through habit rather than being critically questioned or scrutinised.

451 However, (Paton, 2008) caution that if people overestimate the effectiveness of some mitigation
452 measures or their ability to respond to a hazard, they may be less inclined to recognise the need
453 for additional mitigation measures and less receptive to new awareness-raising initiatives. This is
454 evident here: residents are less motivated to comply with mazuku warning panels at all times of a
455 day because they believe they already know the “critical periods” (early mornings and after
456 rainfall). Yet, in this region, it has been demonstrated that concentration levels can change
457 suddenly following abrupt magmatic activities or volcanic events or due to diurnal–nocturnal
458 fluctuations (Balagizi et al., 2018b; M. Kasereka, 2017; Smets et al., 2010). Therefore, locally
459 contextualised awareness initiatives that build upon people’s risk experiences, knowledge, and
460 available resources may prove more effective. (Lutete Landu et al., 2023; Mafuko-Nyandwi et
461 al., 2024).

462 **6.2. The Influence of Risk Experience on Mazuku Mitigation**

463 The literature indicates that risk experience influences the perceptions of people living in hazard-
464 prone areas, whether in terms of risk perception or views on the implementation of mitigation
465 measures (Mafuko Nyandwi, Kervyn, Habiyaremye, et al., 2023; Sattler et al., 2000; Townshend
466 et al., 2015). In this perspective, our results show that the number of times an individual has
467 experienced the risk of a volcanic eruption positively influences both the motivation to
468 implement, and the perceived effectiveness of local mazuku mitigation measures. Moreover,
469 there is evidence of spatial variation in perceptions of efficacy of mitigation measures, despite no
470 comprehensive knowledge of how mazuku concentrations vary across different zones. Instead,

471 variation in perception aligns more closely with historical patterns of land occupation and
472 settlements.

473 This suggests that these patterns are more reflective of community-level perceptions and shared
474 risk experiences than of an objective individual evaluation of risk mitigation (Becker et al.,
475 2017). Before, the 2021 Nyiragongo eruption, we have observed already a spatial
476 homogenisation in people's perception of volcanic risk across different neighbourhoods of Goma
477 between old residents and newcomers (Mafuko Nyandwi, Kervyn, Habiyaemye, et al., 2023).
478 This was partly because a long time had passed since the last eruption, and partly because
479 Nyiragongo is an "open volcano" with a persistent reddish gas plume at its summit (Barrière et
480 al., 2022), serving over the years as a continual reminder of the volcanic threat. Meanwhile, the
481 mazuku hazard is silent, permanent, colourless and odourless (Smets et al., 2010). Thus, spatial
482 homogeneity in how people perceive the implementation of *mazuku* mitigation measures appears
483 to depend heavily on demographic factors, especially monthly income, which segregate
484 populations into different settlement zones. Interviews in the affected area have already revealed
485 three distinct settlement zones: high-income zone, transitional zone with middle-income
486 households, and low-income household zone with high proportion of IDPs.

487 The spatial homogenisation of risk perception is had been also documented in others context. In
488 an editorial review, Gaillard & Dibben, (2008) demonstrated that the spatial dimension of risk
489 perception is closely linked to the memory of past events or previous experiences of fatalities in
490 a given area. This collective memory can shape entire communities residing in hazardous areas,
491 fostering a strong attachment to their environment—as observed among populations in the
492 Southern Andes that have experienced seven eruptions in less than a century. (Vergara-Pinto &
493 Marín, 2023; Walshe et al., 2023) This means that it is not individual experience that matters
494 most, but rather the shared history of a community, in which the impacts of past fatalities remain
495 visible (such as the skeletons of animals asphyxiated by mazuku) or are passed down orally from
496 generation to generation, or from long-term residents to newcomers, or even from a neighbour to
497 another one (Gaillard & Dibben, 2008). Moreover, within the same zone, households tend to
498 implement only those measures that are affordable for them. This is the case with cementing
499 house yards or septic pits, which are widely perceived implemented in zone 1, where high-

500 income households live. Thus, the effective implementation of mitigation measures requires
501 empowering local communities through a co-creation approach.

502 **6.3. The Need for Co-Creation with Local Communities and Empowering Them**

503 In a systematic review, (Viveiros & Silva, (2024) discuss both the environmental and health
504 impacts of volcanic gases and highlight that mitigation strategies vary significantly between
505 volcanic regions. In our study, we also identified mitigation measures that are specific to the
506 Goma context, such as heating fires in courtyards to foster the dispersion of mazuku or using
507 waste materials to block its emission. This highlights the importance of co-creating knowledge
508 and mitigation measures with local communities (Pardo et al., 2015), rather than importing
509 solutions that may not be suited to the local context (Bird et al., 2011). Therefore, understanding
510 the incentives that drive these communities to mitigate mazuku-related risks is essential for
511 effective risk management (Barclay et al., 2008, 2015).

512 In this perspective, our findings support (Barclay et al., 2008), who noted that in many cases the
513 risk is well known to the exposed population, yet they may fail to act due to competing life
514 pressures such as resource constraints, rather than a lack of knowledge. We observed that both
515 the perceived efficacy of risk mitigation measures and their perceived level of implementation
516 vary across zones not because of differences in mazuku concentrations but because of resource
517 limitations. People report being motivated to adopt a mitigation measure if they perceive it as
518 effective and if it is affordable. In other words, even when a measure could be effective—such as
519 cementing courtyards or septic pits—motivation to implement it declines sharply if resources are
520 lacking and, paradoxically, our results indicate that the measure is then judged less effective.
521 Therefore, mitigation measures that align with capacity of specific social groups are likely to be
522 more effective than collective, one-size-fits-all solutions, like the installation of warning panels
523 that are now the only official mitigation measures implemented in Goma. Achieving this requires
524 researchers, decision-makers, and all other stakeholders involved in risk management to learn
525 from local communities practices and collaborate with them in designing mitigation strategies
526 that are locally contextualised.

527 **7. Limitations**

528 This study did not assess the actual physical effectiveness of the 12 risk mitigation measures.
529 Furthermore, data collection did not evaluate whether households had already been directly
530 affected by Mazuku, given that the main impact—loss of human life—could raise ethical
531 sensitivities. In addition, we did not assess whether households had individually implemented a
532 given measure but rather enquired about the level of implementation within the neighbourhood
533 as a whole. This approach was taken because, as highlighted during the interviews, the
534 implementation of such measures was considered more as a collective matter at community
535 level, since the sources of CO₂ emissions were dispersed across different locations.

536 **8. Conclusion**

537 This study employed a mixed-methods approach, combining qualitative and quantitative
538 techniques, to assess perceptions of the implementation of risk mitigation measures related to
539 emissions of magmatic dry gases—primarily carbon dioxide—locally known in the study area as
540 mazuku. Research of this kind is essential, given that the number of people living in active
541 volcanic zones has continued to rise over the centuries, and that cases of human fatalities and
542 livestock asphyxiation are regularly recorded in such areas.

543 The study identified three categories of risk mitigation measures implemented in the western part
544 of Goma, within the Virunga volcanic province: (1) measures aimed at limiting mazuku
545 emissions; (2) adaptive measures to reduce exposure to mazuku; and (3) awareness-related
546 measures based on local knowledge, transmitted orally from generation to generation or from
547 long-term residents to newcomers. Financial resources, along with risk experience—often linked
548 to length of residence—were found to positively influence both motivation and the perceived
549 effectiveness of the first two categories of measures. Perceptions of awareness-related measures
550 showed no significant variations. Moreover, the study highlights spatial variation in both the
551 level of implementation and the perceived effectiveness of these measures, not necessarily based
552 on individual evaluation but rather on community-level knowledge of the local environment.

553 This study offers novel insights into the implementation of risk mitigation practices addressing
554 volcanic gas emissions in active volcanic zones—such as heating courtyards or blocking gas
555 with household waste—examined through a Global South perspective characterised by rapid and
556 largely uncontrolled urbanisation. It reinforces the call, made by other scholars, for the co-

557 creation of mitigation strategies with local communities, rather than the imposition of externally
 558 derived solutions that may not be effective in the local context. Future research could
 559 complement these findings by assessing the actual effectiveness of such mitigation measures
 560 through physical measurements of mazuku concentrations—not only in public spaces but also
 561 within buildings—and by further examining local risk perception. Moreover, volcano monitoring
 562 programmes in Goma and the surrounding areas should diversify their focus to include
 563 systematic monitoring of mazuku and recognise it as a significant public risk requiring sustained
 564 attention.

565 Appendices

566 Appendix A

567 **Table A1: Results of test of variations of motivations according to demographic**
 568 **characteristics**
 569

1. Blocking gas measures			
Variable	Test	Statistic	P_Value
Gender	Wilcoxon	29341	0.0000
Age	Kruskal-Wallis	36.26726631	0.0000
Income	Kruskal-Wallis	117.044502	0.0000
Household size	Kruskal-Wallis	1.642291024	0.8012
Room number	Kruskal-Wallis	130.0287962	0.0000
Eruption experience	Kruskal-Wallis	86.4399316	0.0000
Residence duration	Kruskal-Wallis	28.48813659	0.0000
2. Adaptative mitigation measures			
Variable	Test	Statistic	P_Value
Age	Wilcoxon	28238	0.00000
Income	Kruskal-Wallis	33.48868	0.00000
Household size	Kruskal-Wallis	49.02454	0.00000
Room number	Kruskal-Wallis	2.09096	0.71903
Eruption experience	Kruskal-Wallis	76.40373	0.00000
Residence duration	Kruskal-Wallis	51.00693	0.00000
3. Community based awareness measures			
Variable	Test	Statistic	P_Value
Age	Wilcoxon	35057.5	0.063461708
Income	Kruskal-Wallis	1.733625	0.78460089
Household size	Kruskal-Wallis	14.45435	0.059776304
Room number	Kruskal-Wallis	1.374521	0.848611036
Eruption experience	Kruskal-Wallis	8.608284	0.071672068
Residence duration	Kruskal-Wallis	3.911153	0.418163549

570 **Appendix B: The spatial variations of level of implementation per sampling zones**
 571






Figure B1: Variation of level of implementation of blocking gas measures



573

Figure B2: Variation of level of implementation of adaptive mitigation measures



574

Figure B3: Variation of level of implementation of community based mitigation measures

575

576 Acknowledgement:

577 Research discussed in this publication has been supported by the Global Development Network
578 (GDN) and L'Agence Française de Développement (AFD). The views expressed in this article
579 are not necessarily those of GDN or AFD.

580 The author gratefully acknowledges the invaluable work of the enumerators who collected the
581 field data for this study, as well as the dedicated support of the University of Goma team
582 involved in the GDN project.

583 Data availability

584 Raw and processed data are available on Zotero (Mafuko Nyandwi, 2025). Research design and
585 the French questionnaire are available upon request from the corresponding author.

586 Ethical statement

587 The survey questionnaire and protocol were approved by the academic office of the University of
588 Goma and local authorities at the municipality and neighbourhood levels in Goma. Verbal
589 informed consent was obtained from the survey participants for their anonymized information to
590 be published in this article.

591 Competing interests

592 The author declares no conflict of interest

593 References

594 Balagizi, C. M., Kies, A., Kasereka, M. M., Tedesco, D., Yalire, M. M., & McCausland, W. A.
595 (2018). Natural hazards in Goma and the surrounding villages, East African Rift System.
596 *Natural Hazards*, 93(1), 31–66. <https://doi.org/10.1007/s11069-018-3288-x>

597 Barclay, J., Haynes, K., Houghton, B., & Johnston, D. (2015). Social Processes and Volcanic
598 Risk Reduction. In *The Encyclopedia of Volcanoes* (Second Edi). Dr Jenni Barclay,
599 Copyright © 2015. <https://doi.org/10.1016/b978-0-12-385938-9.00069-9>

600 Barclay, J., Haynes, K., Mitchell, T., Solana, C., Teeuw, R., Darnell, A., Croweller, H. S., Cole,
601 P., Pyle, D., Lowe, C., Fearnley, C., & Kelman, I. (2008). Framing volcanic risk

602 communication within disaster risk reduction: Finding ways for the social and physical
603 sciences to work together. *Geological Society Special Publication*, 305, 163–177.
604 <https://doi.org/10.1144/SP305.14>

605 Barrière, J., d'Oreye, N., Smets, B., Oth, A., Delhaye, L., Subira, J., Mashagiro, N., Derauw, D.,
606 Smittarello, D., & Syavulisembo, A. M. (2022). Intra-crater eruption dynamics at
607 Nyiragongo (DR Congo), 2002–2021. *Journal of Geophysical Research: Solid Earth*,
608 127(4), e2021JB023858.

609 Bird, D. K., Gísladóttir, G., & Dominey-Howes, D. (2011). Different communities, different
610 perspectives: Issues affecting residents' response to a volcanic eruption in southern Iceland.
611 *Bulletin of Volcanology*, 73(9), 1209–1227. <https://doi.org/10.1007/s00445-011-0464-1>

612 Bourdieu, P. (1990). *The logic of practice*. Stanford university press.

613 Brown, S. . K., & Jenkins, S. F. (2017). Global distribution of volcanic threat. *Global Volcanic*
614 *Hazards and Risk*, 2015, 359–369.

615 Brown, S. K., Auken, M. R., & Sparks, R. S. J. (2015). Populations around holocene volcanoes
616 and development of a population exposure index. *Global Volcanic Hazards and Risk*, 2015,
617 223–232. <https://doi.org/10.1017/CBO9781316276273.006>

618 Büscher, K., & Vlassenroot, K. (2010). Humanitarian presence and urban development: New
619 opportunities and contrasts in Goma, DRC. *Disasters*, 34(SUPPL. 2), 256–273.
620 <https://doi.org/10.1111/j.1467-7717.2010.01157.x>

621 Edmonds, M., Grattan, J., & Michnowicz, S. (2017). Volcanic gases: silent killers. In *Observing*
622 *the volcano world: volcano crisis communication* (pp. 65–83). Springer.

623 Freire, S., Florczyk, A. J., Pesaresi, M., & Sliuzas, R. (2019). An improved global analysis of
624 population distribution in proximity to active volcanoes, 1975–2015. *ISPRS International*
625 *Journal of Geo-Information*, 8(8), 341.

626 Gaillard, J. C., & Dibben, C. J. L. (2008). Volcanic risk perception and beyond. *Journal of*
627 *Volcanology and Geothermal Research*, 172(3–4), 163–169.
628 <https://doi.org/10.1016/j.jvolgeores.2007.12.015>

- 629 Hansell, A., & Oppenheimer, C. (2004). Health Hazards from Volcanic Gases: A Systematic
630 Literature Review. *Archives of Environmental Health*, 59(12), 628–639.
631 <https://doi.org/10.1080/00039890409602947>
- 632 Kasereka, M. M., Yalire, M. M., Minani, A. S., Samba, C. V, Bisusa, A. K., & Kamate, E. K.
633 (2017). Les risques liés aux mazuku dans la région de Goma , République démocratique du
634 Congo (rift est-africain. *Journal of Water Environnement Sciences*, 1, 164–174.
- 635 Loughlin, S. C., Vye-Brown, C., Sparks, R. S. J., Brown, S. K., Barclay, J., Calder, E., Cottrell,
636 E., Jolly, G., Komorowski, J. C., Mandeville, C., Newhall, C., Palma, J., Potter, S., &
637 Valentine, G. (2015). An introduction to global volcanic hazard and risk. In *Global*
638 *Volcanic Hazards and Risk* (Issue 2015). <https://doi.org/10.1017/CBO9781316276273.003>
- 639 Lutete Landu, E., Ilombe Mawe, G., Makanzu Imwangana, F., Bielders, C., Dewitte, O., Poesen,
640 J., Hubert, A., & Vanmaercke, M. (2023). Effectiveness of measures aiming to stabilize
641 urban gullies in tropical cities: Results from field surveys across D.R. Congo. *International*
642 *Soil and Water Conservation Research*, 11(1), 14–29.
643 <https://doi.org/https://doi.org/10.1016/j.iswcr.2022.10.003>
- 644 Mafuko Nyandwi, B. (2025). *Data on perception of the implementation and effectiveness of local*
645 *dry-gas degassing measures in the Goma area (East DRC)* . Zenodo.
646 <https://doi.org/10.5281/zenodo.17628103>
- 647 Mafuko Nyandwi, B., Kervyn, M., Habiyaremye, F. M., Kervyn, F., & Michellier, C. (2023).
648 Differences in volcanic risk perception among Goma’s population before the Nyiragongo
649 eruption of May 2021, Virunga volcanic province (DR Congo). *Natural Hazards and Earth*
650 *System Sciences*, 23(2), 933–953. <https://doi.org/10.5194/nhess-23-933-2023>
- 651 Mafuko Nyandwi, B., Kervyn, M., Muhashy Habiyaremye, F., Kervyn, F., & Michellier, C.
652 (2023). To go or not to go when the lava flow is coming? Understanding evacuation
653 decisions of Goma inhabitants during the 2021 Nyiragongo eruption crisis. In *Journal of*
654 *Volcanology and Geothermal Research* (Vol. 444).
655 <https://doi.org/10.1016/j.jvolgeores.2023.107947>

- 656 Mafuko-Nyandwi, B., Kervyn, M., Habiyaremye, F. M., Vanwing, T., Kervyn, F., Jacquet, W.,
657 Mitengezo, V., & Michellier, C. (2024). Building a prepared community to volcanic risk in
658 the global south: Assessment of awareness raising tools for high school students in Goma,
659 (East DR Congo). *Progress in Disaster Science*, 24, 100370.
660 <https://doi.org/https://doi.org/10.1016/j.pdisas.2024.100370>
- 661 Morgan, K. (1970). Sample size determination using Krejcie and Morgan table. *Kenya Projects*
662 *Organization (KENPRO)*, 38(1970), 607–610.
- 663 Pardo, N., Wilson, H., Procter, J. N., Lattughi, E., & Black, T. (2015). Bridging Māori
664 indigenous knowledge and western geosciences to reduce social vulnerability in active
665 volcanic regions. *Journal of Applied Volcanology*, 4(1), 5. [https://doi.org/10.1186/s13617-](https://doi.org/10.1186/s13617-014-0019-1)
666 [014-0019-1](https://doi.org/10.1186/s13617-014-0019-1)
- 667 Paton, D. (2008). Risk communication and natural hazard mitigation: How trust influences its
668 effectiveness. *International Journal of Global Environmental Issues*, 8(1–2), 2–16.
669 <https://doi.org/10.1504/IJGENVI.2008.017256>
- 670 Pech, L., Büscher, K., & Lakes, T. (2018). Intraurban development in a city under protracted
671 armed conflict: Patterns and actors in Goma, DR Congo. *Political Geography*, 66(August),
672 98–112. <https://doi.org/10.1016/j.polgeo.2018.08.006>
- 673 Pech, L., & Lakes, T. (2017). The impact of armed conflict and forced migration on urban
674 expansion in Goma: Introduction to a simple method of satellite-imagery analysis as a
675 complement to field research. *Applied Geography*, 88, 161–173.
676 <https://doi.org/10.1016/j.apgeog.2017.07.008>
- 677 Quesada-Román, A. (2022). Disaster risk assessment of informal settlements in the Global
678 South. *Sustainability*, 14(16), 10261.
- 679 Sattler, D. N., Kaiser, C. F., & Hittner, J. B. (2000). Disaster preparedness: Relationships among
680 prior experience, personal characteristics, and distress. *Journal of Applied Social*
681 *Psychology*, 30(7), 1396–1420. <https://doi.org/10.1111/j.1559-1816.2000.tb02527.x>
- 682 Smets, B., Tedesco, D., Kervyn, F., Kies, A., Vaselli, O., & Yalire, M. M. (2010). Dry gas vents
683 (“mazuku”) in Goma region (North-Kivu, Democratic Republic of Congo): Formation and

- 684 risk assessment. *Journal of African Earth Sciences*, 58(5), 787–798.
685 <https://doi.org/10.1016/j.jafrearsci.2010.04.008>
- 686 Townshend, I., Awosoga, O., Kulig, J., & Fan, H. Y. (2015). Social cohesion and resilience
687 across communities that have experienced a disaster. *Natural Hazards*, 76(2), 913–938.
688 <https://doi.org/10.1007/s11069-014-1526-4>
- 689 Vaselli, O., Capaccioni, B., Tedesco, D., Tassi, F., & Yalire, M. (2003). The “evil winds”
690 (mazukus) at Nyiragongo Volcano (Democratic Republic of Congo). *Acta Vulcanologica*,
691 *Journal of the National Volcanic Group of Italy*, 14–15, 123–128.
- 692 Vergara-Pinto, F., & Marín, A. (2023). Stratigraphy of volcanic memory: Sociocultural
693 dimensions of volcanic risk in the Southern Andes, Chile. *Journal of Contingencies and*
694 *Crisis Management*, 31(4), 1018–1033. <https://doi.org/10.1111/1468-5973.12474>
- 695 Viveiros, F., Gaspar, J. L., Ferreira, T., & Silva, C. (2016). Hazardous indoor CO₂
696 concentrations in volcanic environments. *Environmental Pollution*, 214, 776–786.
- 697 Viveiros, F., & Silva, C. (2024). Hazardous volcanic CO₂ diffuse degassing areas—A systematic
698 review on environmental impacts, health, and mitigation strategies. *Isience*, 27(10).
- 699 Vlassenroot, K., & Büscher, K. (2009). The City as Frontier: Urban Development and Identity
700 Processes in Goma. *Crisis*, 1797(61), 1–7.
- 701 Wachinger, G., Keilholz, P., & O’Brian, C. (2018). The Difficult Path from Perception to
702 Precautionary Action—Participatory Modeling as a Practical Tool to Overcome the Risk
703 Perception Paradox in Flood Preparedness. *International Journal of Disaster Risk Science*,
704 9(4), 472–485. <https://doi.org/10.1007/s13753-018-0203-8>
- 705 Wachinger, G., Renn, O., Begg, C., & Kuhlicke, C. (2013a). *The Risk Perception Paradox —*
706 *Implications for Governance and Communication of Natural Hazards*. 33(6).
707 <https://doi.org/10.1111/j.1539-6924.2012.01942.x>
- 708 Wachinger, G., Renn, O., Begg, C., & Kuhlicke, C. (2013b). The risk perception paradox-
709 implications for governance and communication of natural hazards. *Risk Analysis*, 33(6),
710 1049–1065. <https://doi.org/10.1111/j.1539-6924.2012.01942.x>

- 711 Walshe, R., Morin, J., Donovan, A., Vergara-Pinto, F., & Smith, C. (2023). Contrasting
712 memories and imaginaries of Lonquimay volcano, Chile. *International Journal of Disaster*
713 *Risk Reduction*, 97, 104003. <https://doi.org/https://doi.org/10.1016/j.ijdr.2023.104003>
- 714 Wauthier, C., Smets, B., Hooper, A., Kervyn, F., & D'Oreye, N. (2018). Identification of
715 subsiding areas undergoing significant magmatic carbon dioxide degassing, along the
716 northern shore of Lake Kivu, East African Rift. *Journal of Volcanology and Geothermal*
717 *Research*, 363, 40–49. <https://doi.org/https://doi.org/10.1016/j.jvolgeores.2018.08.018>
- 718 Williams-Jones, G., & Rymer, H. (2015). Hazards of volcanic gases. In *The encyclopedia of*
719 *volcanoes* (pp. 985–992). Elsevier.
- 720