

# Climatology of aerosol pH and its controlling factors at the Melpitz continental background site in central Europe

Vikram Pratap<sup>1</sup>, Christopher J. Hennigan<sup>1</sup>, Bastian Stieger<sup>2,3</sup>, Andreas Tilgner<sup>2</sup>, Laurent Poulain<sup>2</sup>,  
5 Dominik van Pinxteren<sup>2</sup>, Gerald Spindler<sup>2</sup>, and Hartmut Herrmann<sup>2</sup>

<sup>1</sup> Department of Chemical, Biochemical and Environmental Engineering, University of Maryland, Baltimore County, Baltimore, 21250, USA

<sup>2</sup> Atmospheric Chemistry Department (ACD), Leibniz Institute for Tropospheric Research (TROPOS),  
10 Permoserstr. 15, 04318 Leipzig, Germany

<sup>3</sup>now at: SKW Stickstoffwerke Piesteritz GmbH, Möllendorfer Straße 13, 06886 Lutherstadt Wittenberg

*Correspondence to:* Christopher Hennigan ([hennigan@umbc.edu](mailto:hennigan@umbc.edu)) or Hartmut Herrmann ([herrmann@tropos.de](mailto:herrmann@tropos.de))

15

**Abstract.** Aerosol acidity has importance for the chemical and physical properties of atmospheric aerosol particles and for many processes that affect their transformations and fate. Here, we characterize trends in PM<sub>10</sub> pH and its controlling factors over the period of 2010 – 2019 at the Melpitz research station in eastern Germany, a continental background site in central Europe. Aerosol liquid water content (ALWC) associated with inorganic species decreased by 3.4  $\mu\text{g m}^{-3} \text{ a}^{-1}$ , corresponding to a 50% decrease during the analyzed time period, in response to decreasing sulfate and nitrate. Aerosol pH exhibited an increase of 0.06 units per year, a trend that was distinct from other regions. The factors controlling aerosol pH varied by season. Temperature, the most important factor driving pH variability overall, was most important in summer (responsible for 51% of pH variability) and less important during spring and fall (22% and 27%, respectively). NH<sub>3</sub>, the second most important factor contributing to pH variability overall (29%), was most important during winter (38%) and far less important during summer (15%). Aerosol chemistry in Melpitz is influenced by the high buffering capacity contributed by NH<sub>4</sub><sup>+</sup>/NH<sub>3</sub> and, to a lesser degree, NO<sub>3</sub><sup>-</sup>/HNO<sub>3</sub>. Thermodynamic analysis of the aerosol system shows that secondary inorganic aerosol formation is most frequently HNO<sub>3</sub> limited, suggesting that factors that control NO<sub>x</sub>

20

25

30 would be more effective than  $\text{NH}_3$  controls in reducing PM mass concentrations. However, the non-linear response of gas-phase  $\text{HNO}_3$  and aerosol  $\text{NO}_3^-$  to  $\text{NO}_x$  emissions in the region, likely due to VOC controls on oxidant formation and subsequent impacts on  $\text{NO}_x$  conversion to  $\text{HNO}_3$ , highlights the challenge associated with PM reductions needed to attain new air quality standards in this region.

## 1 Introduction

35 Aerosol acidity (pH) varies greatly in the atmosphere, from highly acidic sub-micron particles with  $\text{pH} < 0$  to nascent sea spray particles with  $\text{pH} \sim 8$  (Pye et al., 2020). pH is tremendously important because it affects so many atmospheric processes, including the gas/particle partitioning of weakly acidic and basic semi-volatile compounds (Ahrens et al., 2012; Nah et al., 2018; Stieger et al., 2021), aqueous-phase reaction rates, including uptake and formation of secondary organic aerosols (Tilgner et al., 2021),  
40 solubility of metals in PM (Fang et al., 2017b), optical properties of light-absorbing brown carbon (Phillips et al., 2017), nutrient deposition and bioavailability (Meskhidze et al., 2003; Baker et al., 2021), and dry deposition rates and atmospheric lifetime of reactive nitrogen species (Nenes et al., 2021). The effects of pH have direct implications for human health through their impacts on PM mass concentrations and toxicity (Song et al., 2024; Dockery et al., 1996). Aerosol pH also has direct implications for Earth's  
45 climate through the effects on PM size and number distributions, light-absorbing properties of organics, and effects on global biogeochemical cycles (Mahowald, 2011; Mahowald et al., 2018).

The pH of atmospheric particles is affected by many factors. Historically, inorganic aerosol composition – focused on the most abundant compounds – was viewed as the driving factor of pH. Sulfate, nitrate, and ammonium are certainly important contributors to aerosol pH (Pye et al., 2020); however,  
50 minor contributors to aerosol mass, such as non-volatile cations (NVCs), can also have important influences on pH (Guo et al., 2018a). Organic acids can contribute to particle acidity in clean environments (Trebs et al., 2005), but their effect on pH is likely small under more polluted conditions (Battaglia et al., 2019). The impact of relative humidity (RH) on aerosol pH has been well known because of its direct effect on aerosol liquid water content (ALWC) (Guo et al., 2015). More recently, temperature  
55 has been identified as a key factor affecting pH due to its effects on acid-base and phase equilibria (Stieger et al., 2021; Battaglia et al., 2017; Tao and Murphy, 2019; Zheng et al., 2020; Campbell et al., 2024).

Together, this indicates that a confluence of factors including regulatory issues, land use, and climate change can produce changes in regional and local aerosol pH.

Few studies have examined aerosol pH in Europe to-date. Cabauw, the Netherlands, a site near  
60 the Atlantic coast surrounded by agricultural land, had mean pH values  $\sim 3.5$  across a 14-month study  
(Guo et al., 2018b). A remote site on the island of Crete, located in the Mediterranean Sea, typically had  
highly acidic aerosol pH values ( $\sim 1 - 1.5$ ), except when the biomass burning influence was high and pH  
was  $\sim 2.5 - 3$  (Bougiatioti et al., 2016). Several studies have characterized pH at a polluted site in the Po  
Valley, Italy, where pH is typically  $\sim 2 - 2.5$  in summer and between  $3 - 4$  (seasonal means) during other  
65 times of the year (Squizzato et al., 2013; Paglione et al., 2021).

Compared with the body of studies that have characterized trends in aerosol mass concentration  
and composition at locations globally, very few studies have characterized trends in aerosol pH through  
time series measurements. The only locations, to our knowledge, where trends in pH have been examined  
over extended time periods ( $> 5$  years) are the southeastern USA (1998 - 2013) (Weber et al., 2016),  
70 contiguous USA (2011 - 2020) (Pan et al., 2024), continental sites in Canada (2007 - 2016) (Tao and  
Murphy, 2019), the Po Valley, Italy (1993 - 2018) (Paglione et al., 2021), and eastern China (2011 - 2019)  
(Zhou et al., 2022). Characterizing trends in aerosol pH is important because many industrialized  
countries have implemented regulations that have reduced emissions of  $\text{SO}_2$  and  $\text{NO}_x$ , two key precursors  
to acidic inorganic aerosol components. Further, aerosol pH may also change in response to warmer  
75 temperatures and changing humidity conditions associated with climate change. Therefore, characterizing  
temporal trends in pH – and the factors contributing to pH – is required to better understand aerosol  
chemistry in these locations. Such analyses can also provide guidance for regions that are transitioning to  
more stringent air pollution controls – e.g., China – and regions projected to implement controls in the  
future – e.g., India and Pakistan. Within the present study, we characterize aerosol pH and its controlling  
80 factors over a decade at the continental background site Melpitz in central Europe. We focus on annual  
trends and seasonal characteristics of gas- and particle-phase composition, partitioning, meteorology, and  
their combined effects on aerosol pH.

## 2 Materials and Methods

### 2.1 Site description

85           The data presented in this analysis were collected between January 2010 – August 2019 at Melpitz, Germany (12°56'E, 51°32'N, 86 m a.s.l.), a rural background site in Saxony situated ~50 km north-east of Leipzig (Fig. S1). The research station at Melpitz is operated by the Leibniz-Institute for Tropospheric Research (TROPOS), where aerosol and gas-phase composition measurements have been conducted for more than three decades (Spindler et al., 2004; Spindler et al., 2013; Spindler et al., 2010). The site is a  
90   part of the European Monitoring and Evaluation Programme (EMEP) network and the Aerosol, Clouds and Trace Gases Research Infrastructure (ACTRIS). The sampling site is in a rural area surrounded by flat agricultural land with no major anthropogenic sources nearby (Atabakhsh et al., 2023). The prevailing wind directions are from the west, southwest, and east (Fig. S2). The southwest wind often has a maritime origin that crosses western Europe before reaching Melpitz, while easterly winds are largely dry  
95   continental air masses with strong anthropogenic influence from emissions in several central and Eastern European countries (Spindler et al., 2013; Stieger et al., 2018). Long-term measurements at the site show the highest PM<sub>2.5</sub> concentrations in air masses from east-southeast (28 µg m<sup>-3</sup>) and lowest in those from the northwest (19 µg m<sup>-3</sup>) (Spindler et al., 2013).

### 100   2.2 Gas and aerosol measurements

          A Monitor for AeRosols and Gases in ambient Air (MARGA, Metrohm-Applikon, The Netherlands) system was deployed to conduct hourly measurements of water-soluble gases and inorganic ionic components in aerosols (Stieger et al., 2019; Rumsey et al., 2014; Ten Brink et al., 2007). The instrument uses a wet-rotating denuder (WRD) and a steam-jet-aerosol-collector (SJAC) to capture trace  
105   gases (HCl, SO<sub>2</sub>, NH<sub>3</sub>, HNO<sub>3</sub>, HNO<sub>2</sub>) and water-soluble ions in particles (Cl<sup>-</sup>, NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>, NH<sub>4</sub><sup>+</sup>, Na<sup>+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>, Ca<sup>2+</sup>, and organic acid ions), respectively, at a time resolution of one hour. The sample air passes through a Teflon-coated PM<sub>10</sub> inlet (URG Inc., Chapel Hill, USA) and enters the WRD wetted by de-ionized water and captures water-soluble gases. The sample air then goes to a SJAC where the sample air is mixed with steam, creating a supersaturated environment that causes particles to grow rapidly into  
110   droplets, which are subsequently collected into a liquid sample stream. The liquid solutions from the

WRD and SJAC are analyzed online using a dual anion-cation ion chromatograph. Details of the ion chromatography protocols are provided by Stieger et al. (2018). A detailed performance characterization of the MARGA system has been discussed previously (ten Brink et al., 2007; Stieger et al., 2018).

## 2.3 Aerosol pH modelling and sensitivity

115 Aerosol pH was calculated using the MARGA measurements, along with temperature and relative humidity (RH), as inputs to the E-AIM (Extended – Aerosol Inorganics Model) thermodynamic equilibrium model (Clegg et al., 1998; Wexler and Clegg, 2002). E-AIM was run using Model IV in batch mode, with solids formation prevented (i.e., ‘metastable’ mode) according to Pye et al. (2020).  $H^+$  mole fraction and mole-fraction-based activity coefficients output from E-AIM were used to calculate pH using  
120 the conversion described by Pye et al. (2020). Inputs to the model were particulate inorganic ionic compounds ( $Na^+$ ,  $NH_4^+$ ,  $Cl^-$ ,  $NO_3^-$ , and  $SO_4^{2-}$ ), temperature, and RH. The  $Cl^-$  and  $NO_3^-$  inputs represented the total gas + aerosol concentrations ( $HCl + Cl^-$  and  $HNO_3 + NO_3^-$ , respectively), though aerosol  $NH_4^+$  and  $NH_3(g)$  were separate inputs to the model. E-AIM does not include the non-volatile cations (NVCs)  $K^+$ ,  $Mg^{2+}$ , and  $Ca^{2+}$ . Therefore,  $K^+$ ,  $Mg^{2+}$ , and  $Ca^{2+}$  were input as  $Na^+$  equivalents. The measurement  
125 values of NVCs that were below the method detection limit (MDL) were assigned the value of the MDL for model runs. For  $Na^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$  and  $K^+$ , those values were 0.02, 0.01, 0.01 and 0.01  $\mu g\ m^{-3}$ , respectively. Guo et al. (2018a) showed that this assumption affects our pH calculations by  $< 0.1$  pH unit (i.e., compared to simulations using zero for NVC inputs when measurements were below the MDL), in agreement with other studies that have shown a minor effect of NVCs on pH when present in low  
130 concentrations (Battaglia Jr et al., 2021; Tao and Murphy, 2021; Zheng et al., 2020).  $H^+$  and  $OH^-$  ions are required inputs to the model and were estimated using the total ion charge balance fulfilling the requirement of electroneutrality. Thermodynamic calculations of pH are challenged at low RH (Pye et al., 2020). Similarly, the operating temperature range for E-AIM model IV with  $NH_4$  and  $Cl^-$  or  $Na^+$  and other ions is 263.15 – 330 K (<https://www.aim.env.uea.ac.uk/aim/aim.php>). Therefore, samples with both  
135  $T < 264\ K$  and  $RH < 60\%$  were excluded from the present analysis (1% and 16% of observations, respectively). Further, data with modeled pH outside the range of -0.5 – 6.0 and ALWC outside the range of 0.34 – 500  $\mu g\ m^{-3}$  were outliers and were also excluded (3% of observations). Finally, the model was

not run if there were any data missing from the aerosol, gas-phase, or meteorology measurements (e.g., for instrument maintenance, calibrations, QA/QC checks, 20% of observations). Overall, with these conditions applied, we modeled 54614 out of a possible 86185 hourly data points (~63%) across the 10-year study, with  $n = 13130$  in spring (March, April, and May),  $n = 11350$  in summer (June, July, and August),  $n = 14200$  in autumn (September, October, and November), and  $n = 15934$  in winter (December, January, and February).

The E-AIM model was run without organics as inputs, a good assumption that has minimal effect on the predicted pH under the imposed RH limitation (Battaglia et al., 2019). Organic aerosol components can take up water and contribute substantially to ALWC, so the results presented herein do not account for the fraction of ALWC associated with those organic species. Based on AMS measurements of  $\text{PM}_{10}$  composition at the Melpitz research station, it is likely that water content associated with organics was ~20% of the total ALWC assuming an average kappa value of 0.15 for the organics (Atabakhsh et al., 2025). Although the model output represents ALWC associated with  $\text{PM}_{10}$ , Kakavas et al. (2021) showed that ALWC was many times higher in fine particles than coarse mode particles across Europe. The approximations and assumptions are evaluated through the measured and predicted  $\text{NH}_3$  partitioning, a key metric used to assess thermodynamic model calculations of aerosol pH.  $\text{NH}_3$  (slope = 1.006, intercept = 0.15,  $R^2 = 0.99$ ,  $n = 54617$ ) and  $\text{NH}_4^+$  (slope = 0.933, intercept = 0.22,  $R^2 = 0.928$ ,  $n = 54617$ ) showed excellent agreement between the predictions and measurements (Fig. S3), indicating the model assumptions employed were valid for this dataset.

The approach of Tao and Murphy (2021) was used to investigate the contribution of different factors to pH variability in Melpitz. The method uses pH calculated according to  $\text{NH}_3$  partitioning theory (Eq. 1) to identify the factors responsible for the difference in pH between two systems, whether that be different locations or different times at the same location. The framework of Tao and Murphy (2021) assesses the following factors for their influence on  $\text{NH}_3/\text{NH}_4^+$  phase partitioning, and hence pH: (1) temperature, (2) gas-phase  $\text{NH}_3$  concentration, (3) RH, (4) particle properties, and (5) an error term associated with simplifying assumptions in the derivation. Temperature affects equilibrium constants ( $K_H$ ,  $K_a$ , and  $K_w$ ) that affect  $\text{NH}_3/\text{NH}_4^+$  phase partitioning. Duan et al. (2025) demonstrated that T effects on activity coefficients and semi-volatile vapor pressures also contribute to its overarching influence on pH.

The  $\text{NH}_3$  (g) abundance clearly affects  $\text{NH}_3$  partitioning, as a  $\text{NH}_3/\text{NH}_4^+$  system at equilibrium will shift towards the gas phase in response to a reduction in  $\text{NH}_3$  and towards the particle phase in response to an  $\text{NH}_3$  increase.  $RH$  directly affects  $ALWC$ , which impacts the particle-phase activities of aerosol species, including  $\text{NH}_4^+$  and  $\text{H}^+$ , both of which directly factor into the computation of  $\text{NH}_3/\text{NH}_4^+$  phase partitioning. Finally, particle properties account for the particle hygroscopicity, which also influences  $ALWC$ , and the ammonium activity coefficient, which affects the ammonium activity.

In the present analysis, we used the average study conditions in Melpitz to calculate the factors responsible for seasonal variability in  $pH$ . The approach decomposes the  $pH$  difference between two points into the difference between key factors that control or influence  $pH$ , Equations 2 and 3 (taken from Eq. 8 and Eq. 4 in Tao and Murphy (2021)):

$$pH = \log_{10} \left[ \frac{n(\text{NH}_3) \cdot K_H \cdot T \cdot R}{\chi(\text{NH}_4^+) \cdot \gamma(\text{NH}_4^+) \cdot 55.509 \cdot P_{atm}} \right] \quad (1)$$

$$pH_2 - pH_1 = \Delta \log_{10}[n(\text{NH}_3)] + \Delta \log_{10} \left[ \frac{\kappa M(\text{NH}_4^+)_{avg}}{\rho(\text{NH}_4^+)_{avg} \gamma(\text{NH}_4^+)} \right] + \Delta \log_{10} \left[ \frac{RH}{1 - RH} \right] + \Delta \log_{10}[K_H \cdot T] + \Delta(\varepsilon) \quad (2)$$

$$pH_2 - pH_1 = \Delta f(\text{NH}_3) + \Delta f(\text{particle properties}) + \Delta f(RH) + \Delta f(T) + \Delta(\varepsilon) \quad (3)$$

where  $n(\text{NH}_3)$  is gas-phase molar concentration of ammonia ( $\text{mol m}^{-3}$ ),  $\kappa$  is the particle hygroscopicity parameter,  $M(\text{NH}_4^+)_{avg}$  is equivalent molecular weight of ammonium in the aerosol phase ( $\text{g mol}^{-1}$ ),  $\rho(\text{NH}_4^+)_{avg}$  is the density of particulate ammonium salt ( $\text{kg m}^{-3}$ ),  $\chi(\text{NH}_4^+)$  is the mole fraction of ammonium,  $\gamma(\text{NH}_4^+)$  is the mole fraction-based activity coefficient of ammonium,  $K_H$  is Henry's constant for  $\text{NH}_4^+$  salt (See equation 4, Tao and Murphy (2021)),  $RH$  is the relative humidity,  $T$  is the temperature (K),  $PP$  represents particle properties (a parameter that combines the influence of particle composition and concentration), and an error term ( $\Delta\varepsilon$ ) that accounts for simplifying assumptions used in

the calculation of pH (the term increases under low RH conditions (RH <50%)). The *PP* term includes the measured ratios of ammonium to sulfate and nitrate, as well as the E-AIM modeled  $\gamma(NH_4^+)$  and  $\kappa$ . The values for the parameters in equation (1) were obtained from the results of the E-AIM thermodynamic model. In this case,  $pH_I$  is the average pH for the entire 10-year (2010 – 2019) study in Melpitz, while  
 195  $pH_2$  is the modelled aerosol pH at a given time. The difference in pH between a single observation and the overall study average can then be attributed to the different factors given in Eq. 3. The error term was low ( $\Delta\epsilon \sim 1\text{-}3\%$  overall and within each season), suggesting the simplifying assumptions used in the definition of pH for this analysis were appropriate (Tao and Murphy, 2021). The pH calculated by  $NH_3$  partitioning showed excellent agreement with the E-AIM pH (slope = 1.02,  $R^2 = 0.994$ , Fig. S4),  
 200 suggesting that the method provides useful insight into the factors that affect pH in Melpitz.

We also applied the multiphase buffer theory developed by Zheng et al. (2020) to quantify the buffering capacity in Melpitz attributed to different compounds. The buffering capacity,  $\beta$ , represents the ratio between the mass of acid (base) added to a system and the resulting decrease (increase) in aerosol pH. Higher values of  $\beta$  indicate systems that will have relatively smaller changes in pH for a given amount  
 205 of acid or base added. Multiple compounds present in the atmosphere contribute to the aerosol buffering capacity, including the pairs  $NH_3\text{-}NH_4^+$ ,  $HNO_3\text{-}NO_3^-$ , and  $HSO_4^-\text{-}SO_4^{2-}$  as well as NVCs and  $H_2O$ . The concentrations of each buffering species or pair, as well as meteorological conditions, determine the overall  $\beta$  as well as the contribution of each acid-base pair to the buffering capacity. The calculation, based on Zheng et al. (2020) is:

$$210 \quad \beta = \frac{dn_{base}}{dpH} = 2.303 \left\{ \frac{K_w}{[H^+]} + [H^+] + \sum_i \frac{K_{a,i}^*[H^+]}{(K_{a,i}^* + [H^+])^2} [X_i]_{tot}^* \right\} \quad (4)$$

Where  $n_{base}$  is the molal concentration ( $\text{mol kg}^{-1}$ ), pH is the aerosol pH,  $K_w$  is the temperature-dependent water dissociation constant,  $K_{a,i}^*$  is the effective acid dissociation constant for species  $i$ ,  $[H^+]$  is the molality based aqueous concentration of  $H^+$  ( $\text{mol kg}^{-1}$ ), and  $[X_i]_{tot}^*$  is the total equivalent molality-based concentration of conjugate acid-base pair  $i$  (e.g.,  $[X_i]_{tot}^*$  for  $NH_4^+/NH_3$  would include  $[NH_4^+(\text{aq})]$ ,  
 215  $[NH_3(\text{aq})]$ , and  $[NH_3(\text{g})]$ , with all concentrations expressed as moles per kg water).



## 3 Results

### 3.1 Annual trends of aerosol pH

There were notable trends in several major species measured at Melpitz (Fig. 1, Table 1). The trend in annual sulfate concentrations was  $-0.15 \mu\text{g m}^{-3} \text{ a}^{-1}$ , a statistically significant decrease that corresponded to a  $\sim 60\%$  reduction from 2010 to 2019 (Fig. 1a). The trend in annual Tot-NO<sub>3</sub> concentrations was also  $-0.15 \mu\text{g m}^{-3} \text{ a}^{-1}$ , a statistically significant decrease that corresponded to a  $\sim 50\%$  reduction from 2010 to 2019 (Fig. 1b). These trends are the result of ongoing reductions in precursor SO<sub>2</sub> and NO<sub>x</sub> emissions across continental Europe (Turnock et al., 2015; Vestreng et al., 2009; Hamed et al., 2010; Jonson et al., 2022). The observed trends in aerosol sulfate are consistent with the broader trends of decreasing sulfate across central Europe that have occurred since the 1980s (Vestreng et al., 2007). In response to the decreases in SO<sub>4</sub><sup>2-</sup> and Tot-NO<sub>3</sub>, aerosol NH<sub>4</sub><sup>+</sup> concentrations exhibited a similar decrease of  $\sim 60\%$  from 2010 to 2019. Sulfate, nitrate, and ammonium (SNA) are the most abundant inorganic aerosol components in Melpitz (Spindler et al., 2013), and thus, they control ALWC abundance. Concentrations of ALWC decreased by  $3.4 \mu\text{g m}^{-3} \text{ a}^{-1}$  (Table 1), representing a  $\sim 50\%$  decrease from 2010 to 2019 (Fig. 1c) because of the 50 – 60% decreases in SNA.

Although the concentration of NH<sub>4</sub><sup>+</sup> decreased, the average Tot-NH<sub>3</sub> concentration remained relatively flat from 2010 – 2019 (Fig. 1d), indicating a shift in NH<sub>3</sub> partitioning towards the gas phase. The annual trend in the Tot-NH<sub>3</sub> concentration was not statistically significant at the 95% confidence level (Table 1). A moderate increase in total column NH<sub>3</sub> was observed over Germany for the period roughly corresponding to the present study, 2008 – 2018, although most of the observed increase occurred in 2017 and 2018 (Van Damme et al., 2021). NVCs showed a slight positive trend from 2010 – 2019 (Fig. 1e),  $+0.01 \mu\text{g m}^{-3} \text{ a}^{-1}$ , but the increase was not statistically significant at the 95% confidence level.

The annual average aerosol pH varied between 2.76 – 3.45 and followed a gradual linear increase of  $\sim 0.06$  pH units per year, leading to a 0.6 pH unit rise over the past ten years (Fig. 1f). The increasing trend in pH is statistically significant at the 95% confidence level (Table 1), demonstrating a decrease in particle acidity in Melpitz. The increasing pH trend in Melpitz is interesting to contrast with trends in other locations because it is the only site where particle acidity has shown a significant decrease (pH increase). Weber et al. (2016) found a slight decrease in pH in the southeastern United States over 1998

– 2013, though the trend appears flat from 2006 – 2013. Tao and Murphy (2019) observed no significant  
245 changes in pH in five Canadian cities from 2007 – 2016, though acidity in Ottawa increased (pH  
decreased). Paglione et al. (2021) used fog water composition measurements to estimate annual trends in  
aerosol pH in the Po Valley, Italy. While fog water pH has steadily increased since 1993, rising by 0.04  
units per year on average as a result of decreasing emissions of SO<sub>2</sub> and NO<sub>x</sub>, calculated aerosol pH  
declined by ~1.5 units from 1993 – 2018 (Paglione et al., 2021). Although the trends in aerosol pH in  
250 Melpitz appear contrary to those in North America and the Po Valley, the results are highly consistent  
with trends in fog water and cloud water pH observed over central Europe (Pye et al., 2020). Using  
composite results from dozens of sites across Europe, cloud and fog water pH has increased by 0.56 pH  
units per decade since 1980 (~2.2 pH unit increase from 1980 to 2020) (Pye et al., 2020). This closely  
matches the 0.6 pH unit increase in Melpitz aerosol for 2010 - 2019. Further, pH measured in cloud water  
255 at Whiteface Mountain, NY, has shown a marked increase over a similar timescale, suggesting key  
differences in the thermodynamic regime of the aerosol in North America (Lawrence et al., 2023).

### 3.2 Seasonal trends in aerosol pH, ALWC, and composition

Aerosol composition in Melpitz is characterized by large gradients in the seasonal concentrations  
260 of major components. Median SO<sub>4</sub><sup>2-</sup> concentrations were relatively consistent between seasons, with a  
range of only 1.14 – 1.37 µg m<sup>-3</sup>. However, the quartile and 90<sup>th</sup> percentile values indicate relatively  
frequent enhancements in SO<sub>4</sub><sup>2-</sup> concentrations during winter and similarly infrequent enhancements  
during summer (Fig. 2a). Under the coldest conditions, Melpitz was under heavy influence from emissions  
to the east and southeast (Poland, Czech Republic, and Austria) (Stieger et al., 2018). Sulfate showed an  
265 inverse relationship with temperature during winter (Fig. S5) due to higher energy demand and increased  
coal combustion in eastern Europe for domestic heating (Atabakhsh et al., 2023; Hamed et al., 2010).  
Tot-NO<sub>3</sub> concentrations exhibited a similar seasonal pattern as SO<sub>4</sub><sup>2-</sup>, with the highest median  
concentrations in winter (3.41 µg m<sup>-3</sup>), followed by spring (2.95 µg m<sup>-3</sup>), fall (2.53 µg m<sup>-3</sup>), and then  
summer (1.84 µg m<sup>-3</sup>, Fig. 2b). Average Tot-NO<sub>3</sub> concentrations in winter and spring were approximately  
270 double the summer average, while differences in the upper quartile and 90<sup>th</sup> percentile values were even  
larger. The seasonal Tot-NO<sub>3</sub> pattern is also apparent in Fig. 2b, where maxima in winter and minima in

summer occurred almost annually. The seasonal Tot-NO<sub>3</sub> results reflect higher energy demand and increased coal combustion under colder conditions in the region, as well as differences in the air mass source regions and boundary layer structure.

275 In contrast to SO<sub>4</sub><sup>2-</sup> and Tot-NO<sub>3</sub>, mean and median concentrations of Tot-NH<sub>3</sub> were ~1.5 times higher in the summer and fall compared to winter, and 2.3 times higher in spring (Fig. 2c). Tot-NH<sub>3</sub> concentrations in the spring regularly exceeded 10-15 µg m<sup>-3</sup>, an infrequent occurrence in other seasons (Fig. 2c). Melpitz is located in eastern Saxony, a region dense with farmland. The elevated Tot-NH<sub>3</sub> observed in spring is attributed to fertilizer applications at the beginning of the growing season, as this is  
280 a major source of NH<sub>3</sub> emissions in the region (Viatte et al., 2022). The observations in Melpitz are consistent with those in other parts of Europe, where NH<sub>3</sub> emissions peak in March, April, and May due to fertilizer spreading (Van Der Graaf et al., 2022). The region surrounding Melpitz shows large model-measurement differences in NH<sub>3</sub> emissions, and predictions are worse during the spring (van der Graaf et al., 2022).

285 Tot-NH<sub>3</sub> concentrations showed a clear relationship with temperature in all seasons except winter (Fig. 3a). Due to the strong effect of temperature on NH<sub>3</sub> partitioning (Fig. 3d), mean NH<sub>3</sub>(g) concentrations increased with temperature in each season (Fig. 3b). Mean gas-phase fraction  $\epsilon_{\text{NH}_3}$  ( $\epsilon_{\text{NH}_3} = \text{NH}_3(\text{g})/(\text{NH}_3(\text{g}) + \text{NH}_4^+)$ ) values showed highly consistent behavior with temperature across seasons despite changing aerosol composition. The results in Fig. 3 suggest that agricultural emissions represent  
290 the major source of Tot-NH<sub>3</sub> during most seasons in Melpitz, consistent with broader studies across Europe (Sutton et al., 2013; Backes et al., 2016a). Very strong correlations ( $r^2 \sim 0.8$ ) between NH<sub>3</sub>(g) and skin temperature (physical temperature of the earth's surface) – both remotely sensed – are observed in this area (Viatte et al., 2022). We observed similar temperature-NH<sub>3</sub> relationships within each season, but the annual results showed less consistency. We observed stronger seasonal gradients, especially much  
295 higher spring concentrations, than the adjacent area (lon [11, 12], lat [50, 52]). However, it should be noted that the IASI satellite results are based on twice-daily total column measurements, at 09:30 and 21:30 local time, and it is not straightforward to convert between total column NH<sub>3</sub> and local, in situ concentrations: this relationship varies by region (van der Graaf, ACP, 2022; Viatte et al., 2022).

The seasonal profile of ALWC was distinct from the other species plotted in Fig. 2. Mean values  
300 of ALWC were a factor of 3-4 higher than the median each season, reflecting skewed distributions.  
Median RH values were 74.7% (summer), 79.0% (spring), 89.2% (fall), and 89.6% (winter). ALWC  
increases exponentially with RH, so RH values above 90%, and especially above 95%, cause this  
observation. Approximately 28% of observations in the winter and 32% of observations in fall had an  
RH > 95% (Fig. S6), reflecting the humid conditions experienced in Melpitz during this time of year.  
305 Conversely, approximately 17% of observations in summer and 16% of observations in spring had an  
RH > 95% (Fig. S6). This explains why the mean, upper quartile, and 90<sup>th</sup> percentile ALWC  
concentrations were so much higher in winter and fall than in spring and summer. Similar skewed seasonal  
distributions in ALWC, with mean seasonal values near or exceeding upper quartile values, were also  
observed in a decadal study in Shanghai, China (Zhou et al., 2022).

310 Aerosol pH showed seasonal trends that reflect the combined influence of inorganic composition  
and meteorology (Fig. 2d). Aerosol pH was highest in spring with mean and median pH of 3.55,  
coinciding with the maxima in Tot-NH<sub>3</sub>. Lowest mean and medial pH values were observed during  
summer (2.83 and 2.89, respectively), when highest temperatures and lowest ALWC levels were  
observed. It is likely that our imposed RH limit (> 60%) for pH calculation removed a subset of the  
315 summer data with very low pH values (< 2) because of the strong effect of ALWC on pH. Therefore, it is  
more likely that the summer data exhibit lower quartile and 10<sup>th</sup> percentile values than Fig. 2d shows.  
Winter had mean and median pH values (3.04 and 3.21, respectively) that reflect the high sulfate and low  
Tot-NH<sub>3</sub> concentrations. Aerosol pH in fall exhibited the second highest season mean and median values  
(3.30 and 3.33, respectively). The pH in Melpitz showed a very different seasonal pattern from that  
320 observed in six Canadian cities, where pH was a minimum in summer and maximum in winter, with  
spring and fall in between as pH transitioned between the extremes (Tao and Murphy, 2019). As discussed  
above in Section 3.1, the thermodynamic drivers of pH Canada and the associated trends in pH appear to  
be different from those in Melpitz. Detailed analysis and discussion of the factors contributing to pH  
variability in Melpitz are presented in Section 3.4.

325

### 3.3 Diurnal trends in aerosol pH

Figure 4 shows the mean diurnal profiles of key species and parameters separated by season. Aerosol pH exhibited a maximum in the early morning hours (04:00-06:00, Fig. 4g) that coincided with the peak ALWC (Fig. 4a). As temperature increased throughout the morning and into the afternoon (Fig. 4b), ALWC and pH decreased. Daily minima in ALWC and pH occurred in the afternoon, coincident with the peak temperature, but rose in the afternoon and evening as temperature decreased. The diurnal profiles of ALWC and pH are qualitatively consistent with those from diverse locations, including the southeastern U.S. (Guo et al., 2015), Baltimore, MD (Battaglia et al., 2017), California (Guo et al., 2017), and Beijing (Ding et al., 2019). This reflects the strong response of pH to meteorological inputs, namely temperature and RH (Battaglia et al., 2017; Zheng et al., 2020; Tao and Murphy, ES&T, 2021). The pH profiles in Fig. 6 are consistent with those Fig. 3d, which shows mean pH was lowest in summer, followed by winter, fall, and spring. Likewise, the ALWC profiles in Fig. 4 are consistent with mean values shown in Fig. 2e, with summer having the lowest ALWC followed by spring, winter, and fall.

The diurnal profiles of Tot-NH<sub>3</sub>, Tot-NO<sub>3</sub>, and sulfate show different processes and sources contributing to their seasonal concentrations (Fig. 4d, e, and f, respectively). Sulfate peaked between 12:00 – 15:00 daily (local), with minima at night, reflecting secondary formation processes. The midday peak during winter is weaker than during other seasons, reflecting the increased importance of regional transport during this time (Stieger et al., 2018). Tot-NO<sub>3</sub> increased in the morning and peaked between 07:00 – 10:00 daily due to photochemical HNO<sub>3</sub> formation. The diurnal profiles of Tot-NO<sub>3</sub> were qualitatively similar during spring, summer, and fall, though the concentrations were quite different, consistent with Fig. 2b. Tot-NH<sub>3</sub> exhibited very different behavior than sulfate or Tot-NO<sub>3</sub>. During spring, summer, and fall, Tot-NH<sub>3</sub> concentrations strongly increased in the morning with the onset of sunrise and the daily temperature increase, approximately doubling between 04:00 and 09:00 each day. Dew is an important nighttime sink and morning reservoir for HONO in Melpitz (Ren et al., 2020). Therefore, in addition to fresh emissions, dew evaporation may be an important source of morning NH<sub>3</sub> (Wentworth et al., 2016). RH peaks at night and in the early morning and the frequency of high RH periods suggests dew regularly forms around Melpitz (Fig S6). While Tot-NH<sub>3</sub> emissions vary considerably by season, Fig. 3d shows that NH<sub>3</sub> partitioning is consistent across seasons and is mostly regulated by temperature.

ALWC shows similar diurnal profiles between seasons, with maxima around 04:00 – 06:00 and  
355 minima around 14:00 – 16:00 (Fig. 4a). The seasonal ALWC profiles are qualitatively consistent with  
those observed in other locations, including North America, China, and other locations in Europe (Pye et  
al., 2020). Aerosol pH profiles are also qualitative similar each season, with maxima at night and early  
morning and minima in the afternoon, due to the concentrating effects of the reduced ALWC (Fig. 4g).  
Although the seasonal profiles also follow the general behavior observed in many other locations, the  
360 amplitude was somewhat less, as the range in diurnal profiles was only ~0.5 pH units in summer and  
spring and only ~0.3 units in winter and fall. This suggests a higher aerosol pH buffering capacity. The  
calculated aerosol pH values are consistent with a modeling study of size-resolved aerosol pH over Europe  
(Kakavas et al., 2021). In May, aerosol pH predictions around Melpitz were ~3 for PM<sub>1</sub> and ~4 for PM<sub>1-2.5</sub>,  
consistent with our observations. The simulations capture well the diurnal trends; however, they  
365 overestimate the range in daily pH, predicting ~2-2.5 unit change in pH throughout the day (Kakavas et  
al., 2021).

Fig. 5 shows the aerosol buffering capacity calculated according to Zheng et al. (2020). Overall,  
aerosols in Melpitz have a high buffering capacity compared to other locations like the southeast USA  
(Zheng et al., 2020). NH<sub>4</sub><sup>+</sup>/NH<sub>3</sub> contributes most of the buffering capacity in Melpitz, followed by NO<sub>3</sub><sup>-</sup>  
370 /HNO<sub>3</sub>. The relative order of importance of buffering pairs was generally consistent across seasons, as  
well (Fig. S7). Other species were relatively minor contributors to the buffering capacity because of the  
observed aerosol pH range. The high buffering capacity leads to the relatively modest amplitude in diurnal  
pH values observed in Fig. 6a. By contrast, the eastern USA has a much smaller buffering capacity than  
central Europe, contributing to diurnal averages of aerosol pH that vary by 1 – 2 pH units throughout the  
375 day (Nah et al., 2018; Guo et al., 2015). Pye et al. (2020) found that diurnal profiles of aerosol pH in  
Melpitz show lower amplitude than similar profiles in a comparison to diverse environments. Overall, the  
high buffering capacity of aerosols in Melpitz suggests that pH may undergo more modest changes in  
response to climate change because of the high Tot-NH<sub>3</sub> and its contribution to buffering capacity.  
Further, analysis of pH across diverse regions of the globe revealed that NH<sub>3</sub>-NH<sub>4</sub><sup>+</sup> is the dominant  
380 buffering pair in the atmosphere, highlighting the important role of agricultural emissions in affecting  
aerosol chemistry (Zheng et al., 2020).

It should be noted that the present study characterizes the pH of  $\text{PM}_{10}$  but small particles are systematically more acidic than large particles. Kakavas et al. (2021) modeled aerosol pH over continental Europe during summer in four size bins:  $\text{PM}_1$ ,  $\text{PM}_{1-2.5}$ ,  $\text{PM}_{2.5-5}$ , and  $\text{PM}_{5-10}$ . They found that smaller size bins are systematically more acidic than larger size bins due to the higher presence of non-volatile cations from sea salt and dust in larger particles, consistent with prior studies (Keene et al., 2004; Fridlind and Jacobson, 2000; Fang et al., 2017a). The average pH of  $\text{PM}_1$  was 1.3 pH units lower than the largest coarse bin ( $\text{PM}_{5-10}$ ), however, this difference exhibited a strong spatial dependence. Coastal areas with high sea salt influence, such as Denmark and northwestern France, showed the largest pH differences between different size fractions. Kakavas et al. (2021) specifically analyzed conditions in Melpitz and found the pH differences between size fractions to vary diurnally. Throughout the day,  $\text{PM}_{5-10}$  was  $\sim 1$  pH unit higher than  $\text{PM}_1$ ; however, the differences between  $\text{PM}_1$  and  $\text{PM}_{1-2.5}$  were as large as  $\sim 1$  pH unit at night but as small as  $\sim 0.25$  pH unit in the daytime. Very similar differences and diurnal patterns were found for the  $\text{PM}_1$  vs.  $\text{PM}_{2.5-5}$  fractions. We expect the simulations of Kakavas et al. (2021) to represent an upper bound on pH differences between fine and coarse fractions in Melpitz because their simulations were specific to summer: Spindler et al. (2013) showed through analysis of long-term aerosol composition data that dust concentrations in Melpitz are highest during summer. Therefore, we expect the present results to inform the trends of aerosol pH in smaller size fractions ( $\text{PM}_1$  and  $\text{PM}_{2.5}$ ), with pH values  $\sim 0.25$ -1 unit lower, depending on the season and time of day.

### 3.4 Drivers of pH variability

The factors contributing to pH variability in Melpitz are shown in Fig. 6. Overall, temperature and RH together accounted for 45% of pH variability, relative to the study average. This is qualitatively

consistent with prior studies that have identified the important role of meteorology in controlling pH (Battaglia et al., 2017; Zheng et al., 2020; Tao and Murphy, 2021).  $\text{NH}_3$  was the second most important factor, responsible for 29% of pH variability over the entire study, followed by particle properties, which contributed 23% of the variability in pH. The role of  $\text{NH}_3$  in Melpitz was similar to its influence on pH variation in Toronto (25%, Tao and Murphy, 2021). Note that this analysis does not capture the directional change in pH resulting from each factor. In fact, a given factor can cause both positive and negative deviations from the average, depending on conditions. Because the method quantifies changes relative to the mean study conditions, lower temperatures observed in winter contribute to increases in pH while higher temperatures in summer contribute to decreases in pH. This can be seen in Equations 1 and 3, as well as the effects of temperature on  $\text{NH}_3$  partitioning (Fig. 3d). In this analysis, these opposing effects do not cancel out in quantifying the factor's contribution to pH variability.

Consistent with the results in Fig. 2, the factors contributing to pH variability also showed distinct seasonal differences (Fig. 6). Temperature was the most significant contributor to pH variability overall, but its influence ranged from 51% in summer to only 22% in spring. Likewise,  $\text{NH}_3$  – the second most important factor overall – ranged from 38% in winter to 15% in summer. It is important to note that the influence of  $\text{NH}_3$  on pH variability does not correlate with the Tot- $\text{NH}_3$  concentration. For example, summer and fall Tot- $\text{NH}_3$  concentrations were quite similar (Fig. 2c and Fig. 4d), yet  $\text{NH}_3$  had more influence on pH variability during fall than summer (22% vs. 15%, respectively). Particle properties, while the third most important factor overall, had the most important effect on pH variability in fall (32%). RH contributed 10% – 16% of pH variability in all seasons, far less than temperature during winter and summer, but closer to the effect on temperature during spring and fall.

It should be noted that the various factors contributing to pH variability are not completely independent. For example, temperature and  $\text{NH}_3$  often influence pH in opposite directions, but  $\text{NH}_3$  emissions and abundance have a clear dependence on temperature (Fig. 3). Similarly, temperature and RH are typically inversely related at ground level. RH affects ALWC, which changes solute concentrations in aqueous particles and gas-particle partitioning of semi-volatile compounds (Ansari and Pandis, 1999). Temperature is a master variable that affects most processes in the environment – with regards to aerosol pH, it affects compound vapor pressures, equilibrium constants, activity coefficients,



solubilities (i.e., overall partitioning), and reaction rates (Duan et al., 2025; Pye et al., 2020; Tilgner et al., 2021). Therefore, while their effects on aerosol pH can be separately quantified, these factors can somewhat offset each other in the ambient atmosphere relative to conditions where one factor changes independently of the others. A similar phenomenon was observed and noted for the Canadian data set  
435 analyzed by Tao and Murphy (2021).

The results in Fig. 6 also highlight interesting trends that provide additional context to the seasonal and diurnal data (Figs. 2 and 4). For example, while the pH values in winter and summer were similar (mean values differ by only 0.2 units), the factors contributing to pH variability were distinctly different. During winter,  $\text{NH}_3$  was a more important driver of pH variability than temperature (38% and 32%,  
440 respectively), but temperature was far more important during summer (51% to 15%). Conversely, pH values in spring and fall were also similar (mean values differ by 0.2 units), but the drivers of pH variability were more similar than summer-winter.

The present results are consistent with a study investigating the drivers of pH variability in Canada, where temperature was the most significant factor, followed by  $\text{NH}_3$  (Tao and Murphy, 2019;  
445 2021). The observations in Canada also found temperature to be a much more important factor in summer and winter than it was in spring and fall, consistent with the results in Melpitz. However, particle properties had a much greater influence on pH variability in Melpitz, ranging from 19% in winter to 32% in fall, than Canada, where it contributed  $< 10\%$  to pH variability in each season. Tao and Murphy (2021) showed that  $\gamma(\text{NH}_4^+)$  and  $\kappa$  were the two most important individual particle property factors contributing  
450 to pH variability. The results in Melpitz and Canada also differ significantly from a study that characterized pH variability at three coastal sites in China, where temperature was only a minor contributor (8 – 12%) to pH variability (Wang et al., 2022). There, ammonia was a major factor in driving pH variability (23 – 42%), similar to Melpitz and Canada, but RH exhibited a much stronger influence (25 – 39%) than it did in Melpitz and Canada. There are several likely reasons for the observed differences  
455 between Melpitz and Canada and the Chinese sites. The Chinese measurements only spanned 3 – 5 weeks at each site whereas the studies in Melpitz and Canada each included  $\sim 10$  years of data. It is likely that similarly brief periods of time in either location would also yield factor contributions that significantly differ from seasonal and annual averages. The coastal Chinese sites also experienced less daily

temperature variation – only  $\sim 2 - 3$  °C in daily temperature range – than either the Canadian or German  
460 studies, likely contributing to the much smaller effect of temperature on pH variability (Wang et al.,  
2022).

### 3.5 Sensitivity of PM mass to precursor availability

Aerosol concentrations always respond to emissions of non-volatile components and precursors,  
465 such as NVCs and sulfate. For semi-volatile components, especially  $\text{NH}_3$  and  $\text{HNO}_3$ , thermodynamic  
conditions dictate whether aerosol mass responds to changing concentrations. Following the approach of  
Nenes et al. (2020), we calculated the sensitivity of  $\text{PM}_{10}$  mass concentrations to  $\text{HNO}_3$  and  $\text{NH}_3$   
availability. The approach defines regimes where the aerosol mass is limited by  $\text{HNO}_3$  and  $\text{NH}_3$   
availability, mapped in pH and ALWC space. Aerosol mass can be sensitive to  $\text{NH}_3$ ,  $\text{HNO}_3$ , both species,  
470 or neither: temperature, ALWC, and pH are the key parameters that determine the thermodynamic regime  
for a given set of conditions (Nenes et al., 2020). Fig. 7 shows the different thermodynamic regimes by  
season calculated using the mean temperature within each season (275 K in winter, 281 K in spring, 291 K  
in summer, and 282 K in fall).  $\text{NH}_4\text{NO}_3$  concentrations in Melpitz are typically high, accounting for  $\sim 30\%$   
of annual  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  mass in the region (Spindler et al., 2013). Therefore, if a sensitivity to  $\text{NH}_3$  or  
475  $\text{HNO}_3$  (or both) is identified, this suggests the aerosol concentration would respond to changes in these  
semi-volatile precursors. In the present analysis, we only considered ALWC associated with inorganic  
aerosol components. While water-soluble OA likely exerts minor effects on pH (Battaglia et al., 2019), it  
can contribute to the ALWC. Consideration of the ALWC contributed by inorganic and organic  
compounds would result in a minor shift of the individual data points in Fig. 7, although the logarithmic  
480 scale indicates very few points would shift to a different thermodynamic regime (e.g., from insensitive to  
 $\text{HNO}_3$ -sensitive).

Aerosol concentrations in Melpitz were most often sensitive to  $\text{HNO}_3$ , ranging from 47% in winter  
to 83% in spring (Figs. 7 and 8). Very few observations showed sensitivity to  $\text{NH}_3$ , alone, with only 0.2%  
at a minimum (spring) and 2.6% maximum (summer). Many observations showed sensitivity to both  $\text{NH}_3$   
485 and  $\text{HNO}_3$ , from 16% in spring to 51% in winter. Therefore, the fraction of observations sensitive to  
 $\text{HNO}_3$  (either exclusively or with  $\text{NH}_3$ ) was 94% in summer and  $> 99\%$  in spring and fall. Less than 1%

of observations showed insensitivity to  $\text{NH}_3$  and  $\text{HNO}_3$  in winter, spring, and fall, and only 3.3% of observations in summer showed insensitivity to both species (Figs. 7 and 8). Temperature exerts a controlling effect on the partitioning coefficients of  $\text{NH}_3$  and  $\text{HNO}_3$  (affecting the intercept of the red and blue lines in Fig. 7), with higher temperatures increasing the region insensitive to both  $\text{NH}_3$  and  $\text{HNO}_3$  (white region in Fig. 7). This explains why the summer had more than an order of magnitude more observations in the insensitive regime, although this was still a very minor fraction of the overall observations in this season. For a given combination of temperature and pH, higher ALWC increases the condensed-phase fraction of  $\text{NH}_3$  and  $\text{HNO}_3$ , leading to conditions where the aerosol is sensitive to one or both of  $\text{NH}_3$  and  $\text{HNO}_3$ . Lower ALWC leads to the opposite effect, decreasing the sensitivity of aerosol mass to either  $\text{NH}_3$  or  $\text{HNO}_3$ . Thus, the generally high RH conditions in Melpitz (Fig. S6) and high ALWC levels push a lot of the observations into thermodynamic regimes sensitive to one or both of  $\text{NH}_3$  and  $\text{HNO}_3$ . This analysis offers insight into changes that may occur under a changing climate, as well. Higher temperatures will result in more acidic particles and lower ALWC (Battaglia et al., 2017). As shown in Fig. 7, this will make the aerosol mass less sensitive to changes in  $\text{NH}_3$  and  $\text{HNO}_3$ , offering fewer mitigation options for controlling PM levels.

The results in Figs. 7 and 8 suggest strategies to reduce  $\text{PM}_{10}$  concentrations in the region should prioritize  $\text{NO}_x$  emissions reductions over  $\text{NH}_3$  controls because it is the key  $\text{HNO}_3$  precursor. This result is consistent with a one-year study in Cabauw, Netherlands, which also found  $\text{NO}_x$  controls likely to reduce aerosol  $\text{NO}_3^-$  more than  $\text{NH}_3$  controls (Guo et al., 2018b; Nenes et al., 2020). Pay et al. (2012) conducted a modeling study over Europe and found results consistent with the present analysis – more frequent aerosol regimes limited by  $\text{HNO}_3$  availability, suggesting  $\text{NO}_x$  controls may be more effective at reducing  $\text{PM}_{2.5}$ . Another study modeled inorganic aerosol formation over Germany and found the aerosols insensitive to  $\text{NH}_3$  reductions, especially during the spring when  $\text{NH}_3$  emissions peak (Renner and Wolke, 2010). Both studies are consistent with the thermodynamic framework shown in Figs. 7 and 8. Backes et al. (2016b) modeled the response of aerosol concentrations over western Europe to changes in  $\text{NH}_3$  emissions and found that decreases in  $\text{NH}_3$  emissions led to decreases in  $\text{PM}_{2.5}$  and  $\text{PM}_{10}$  mass due to decreases in  $\text{NH}_4\text{NO}_3$ . In strong agreement with our thermodynamic analysis, aerosol mass showed the largest sensitivities to  $\text{NH}_3$  reductions in winter (Backes et al., 2016b).

515 Aerosol  $\text{NO}_3^-$  formation from  $\text{NO}_x$  is highly non-linear, suggesting limitations to the effectiveness of  $\text{NO}_x$  controls on secondary aerosol formation. For example, if a location falls under the  $\text{HNO}_3$  limited thermodynamic regime (or both  $\text{HNO}_3$  and  $\text{NH}_3$  limited), reductions in  $\text{NO}_x$  do not necessarily translate into PM reductions the way that  $\text{SO}_2$  reductions lead directly and almost linearly to reductions in aerosol sulfate. In the eastern US, there was a very weak response of aerosol nitrate to  $\text{NO}_x$  reductions due to  
520 simultaneous changes in  $\text{SO}_2$  emissions and changes in aerosol pH shifted the  $\text{NO}_3^-/\text{HNO}_3$  partitioning towards the particle phase (Shah et al., 2018). So, although  $\text{NO}_x$  reductions may decrease Tot- $\text{NO}_3$ , even minor changes in  $\text{NO}_3^-/\text{HNO}_3$  partitioning can keep  $\text{NO}_3^-$  relatively constant, or even lead to an increase (Ansari and Pandis, 1998). A similar phenomenon was also observed in the Salt Lake Valley, Utah, where wintertime ammonium nitrate formation did not respond to  $\text{NO}_x$  controls, but was instead sensitive to  
525 VOC emissions (Womack et al., 2019). This occurs because  $\text{NO}_x$  conversion to  $\text{HNO}_3$  is oxidant-limited in some environments. In such cases, control of VOC emissions, rather than  $\text{NO}_x$  or  $\text{NH}_3$ , can provide the most direct effect on aerosol  $\text{NH}_4\text{NO}_3$  (Dang et al., 2023; Womack et al., 2019). Changes in  $\text{NO}_3^-/\text{HNO}_3$  partitioning can also reduce the  $\text{HNO}_3$  dry deposition sink, increasing the atmospheric lifetime of Tot- $\text{NO}_3$  (Zhai et al., 2021). This complex chemistry is consistent with a recent study that found that a  
530 23% reduction in  $\text{NO}_x$  emissions during the COVID-19 lockdown period was associated with only a ~5% decrease in  $\text{PM}_{2.5}$  mass in Germany (Balamurugan et al., 2022). Further, actual emissions control measures need to account for costs and analysis by Liu et al. (2023) indicates  $\text{NH}_3$  controls are 5-10 times more cost effective than  $\text{NO}_x$  controls at reducing PM in Europe.

Therefore, although the thermodynamic predictions indicate that the aerosol system in Melpitz is  
535 most frequently sensitive to  $\text{HNO}_3$ , VOC controls on the conversion of  $\text{NO}_x$  to  $\text{HNO}_3$  suggest that  $\text{NO}_x$  reductions may not be as effective in reducing inorganic aerosol concentrations, at least at first. This helps explain the results of a modeling study of the sulfate-nitrate-ammonium system over Germany that predicted  $\text{NH}_3$  reductions will be more effective reducing PM than  $\text{NO}_x$  reductions due to the non-linear response described above (Banzhaf et al., 2013). Such predictions are difficult to test with observations,  
540 especially because the ‘accidental experiment’ offered by the COVID-19 lockdown actually showed higher atmospheric  $\text{NH}_3$  emissions in northwest Germany, likely due to interannual variability in meteorological parameters (Balamurugan et al., 2022).

## 4 Conclusions

Central Europe has seen dramatic changes in secondary aerosol precursor emissions, which have  
545 decreased  $\text{NO}_3^-$  and  $\text{SO}_4^{2-}$  concentrations (Nec, 2024).  $\text{NH}_3$  emissions have remained steady, or increased,  
over the same time period, potentially changing aerosol pH, as well. In Melpitz, the annual average pH  
has increased by 0.06 units per year over the 10-year period 2010 – 2019. This is similar to trends of fog  
water pH in the Po Valley, Italy, where an annual increase of 0.04 units has been observed since 1993  
(Paglione et al., 2021). However, inferred values of aerosol pH in the Po Valley have actually decreased  
550 by 0.03 units per year during the same time period due to the combination of changing aerosol  
composition and changing meteorology (warmer and drier). Similarly, aerosols in the southeastern US  
and Canada have become more acidic despite dramatic decreases in acidic aerosol precursors,  $\text{SO}_2$  and  
 $\text{NO}_x$  (Weber et al., 2016; Tao and Murphy, 2019), contrary to the observed trends in Melpitz. Compared  
with the large body of studies on aerosol composition and trends, few studies have characterized trends  
555 in aerosol pH and its controlling factors. These results underscore the need for more characterizations of  
aerosol pH trends across Europe, and beyond.

The present results highlight the critical factors that contribute to aerosol pH in central Europe.  
Similar to other locations, temperature is the most important factor driving pH variability. There is a  
strong seasonal cycle to the factors affecting pH, with  $\text{NH}_3$  and particle properties (composition and  
560 concentration) also strongly affecting pH variability. The factors contributing to pH variability do not  
scale with concentration or the absolute temperature or RH level. This analysis suggests that aerosol pH  
will continue changing in the future with regulations and climate; however, our results suggest that  
predictions of future changes in aerosol pH require a full treatment of the coupled emissions-chemistry-  
meteorology system that together determine particle acidity. Although the results were derived from  
565 measurements at a background site, similarities in other areas may also be expected. For example, in the  
megacity of Paris, similarities of  $\text{NH}_3$  partitioning, strong agricultural influences on secondary aerosol  
formation, and  $\text{NH}_4\text{NO}_3$  formation regime suggest that the observations in Melpitz may translate to urban  
environments across Europe, as well (Petetin et al., 2016).

Although it is widely observed that  $\text{NH}_3$  emissions in Europe peak during spring due to fertilizer  
570 application, variability in farming practices makes this quite challenging to accurately capture in models.

For example, the region surrounding Melpitz shows large model-measurement discrepancies in  $\text{NH}_3$  emissions, and predictions are worse during the spring (van der Graaf et al., 2022). This is problematic because  $\text{NH}_3$  emissions in Germany contribute significantly to episodic events of elevated  $\text{PM}_{2.5}$ , especially in spring, which are also challenging to model (Fortems-Cheiney et al., 2016). Although  $\text{NH}_3$  emissions represent a large source of uncertainty in model predictions of  $\text{NH}_4\text{NO}_3$  formation, errors come from numerous sources, including  $\text{NO}_x$  emissions, dry deposition rates,  $\text{HNO}_3$  formation from  $\text{NO}_x$  (uncertainty in OH), uncertainty in the thermodynamic partitioning, and model treatment of bidirectional flux (Petetin et al., 2016). Therefore, improved understanding of the aforementioned factors is required to advance predictive capabilities for springtime haze formation events in Europe.

Finally, these results provide additional context to the strategies that may be most effective reducing  $\text{PM}_{2.5}$  mass concentrations. We applied a recently developed framework to identify the thermodynamic regimes of  $\text{NH}_4\text{NO}_3$  aerosol formation through aerosol pH and ALWC (Nenes et al., 2020). Very few observations ( $< 1\%$  total) were insensitive to  $\text{NH}_3$  or  $\text{HNO}_3$ . Thermodynamic conditions indicate that  $\text{HNO}_3$  was more often limiting than  $\text{NH}_3$ , suggesting that  $\text{PM}_{2.5}$  control strategies in this region should target  $\text{NO}_x$  emissions reductions, next to ammonia reductions, especially in winter. However,  $\text{NO}_x$  reaction to form  $\text{HNO}_3$ , with subsequent partitioning to aerosol  $\text{NO}_3^-$ , is highly non-linear. Reductions in  $\text{NO}_x$  decrease Tot- $\text{NO}_3$ , but the system can respond through changes in  $\text{NO}_3^-/\text{HNO}_3$  partitioning such that aerosol concentrations remain relatively stable (e.g., Womack et al., 2019). Changes in  $\text{NO}_3^-/\text{HNO}_3$  partitioning can change  $\text{HNO}_3$  dry deposition rates, but partitioning and dry deposition are usually large sources of model uncertainty (e.g., Zakoura and Pandis, 2019). Therefore, future studies are needed to complement the thermodynamic calculations in order to address such complexities.

## Data Availability

All data presented in this work are available upon request.

## Author Contributions

V.P. – formal analysis, investigation, visualization, writing – original draft, writing – review and editing.

C.H. – conceptualization, supervision, formal analysis, funding acquisition, visualization, writing –

original draft, writing – review and editing. B.S. –investigation, validation, data curation, writing – review  
600 and editing. A.T. – formal analysis, investigation, data curation, writing – review and editing. L.P. - formal  
analysis, investigation, data curation, writing – review and editing. D.vP. - investigation, data curation,  
writing – review and editing. G.S. –investigation, validation, data curation, writing – review and editing.  
H.H. – conceptualization, supervision, resources, funding acquisition, writing – review and editing.

## 605 **Competing Interests**

The authors declare no competing financial interests.

## **Acknowledgements**

The authors acknowledge support in instruments operation in Melpitz by TROPOS ACD technicians  
610 Achim Grüner and René Rabe.

## **Financial Support**

C.H. and V.P. acknowledge support from the National Science Foundation through grants AGS-1719252  
and AGS-2430366. The TROPOS co-authors acknowledge financial support for the deployment of the  
615 MARGA system by the German federal environment protection agency Umweltbundesamt (UBA) under  
contract no. 52436 as well as from the European Regional Development fund by the European Union  
under contract no. 100188826. Further support by the European infrastructure projects ACTRIS (EU FP7;  
grant no. 262254) and ACTRIS-2 (grant no. 654109) and the RI-URBANS project (grant no. 101036245)  
are gratefully acknowledged.

620

625

**Table 1:** Summary of annual trend analysis for species shown in Fig. 1.

	Sulfate	Tot-NO <sub>3</sub> <sup>d</sup>	ALWC <sup>e</sup>	Tot-NH <sub>3</sub> <sup>f</sup>	NVCs <sup>g</sup>	pH
Slope <sup>a</sup>	-0.15	-0.15	-3.39	0.06	0.01	0.06
Statistically significant <sup>b</sup>	Yes	Yes	Yes	No	No	Yes
Intercept <sup>c</sup>	2.60	4.20	69.5	4.69	0.21	2.87
R <sup>2</sup>	0.73	0.80	0.71	0.05	0.10	0.75

<sup>a</sup>Linear least squares method; Units are  $\mu\text{g m}^{-3} \text{ a}^{-1}$ , except pH: pH units  $\text{a}^{-1}$

<sup>b</sup>Indicates whether the slope is statistically significant at the 95% confidence level

<sup>c</sup>Intercept relative to 2010


<sup>d</sup>Tot-NO<sub>3</sub> = NO<sub>3</sub><sup>-</sup> + HNO<sub>3</sub>

<sup>e</sup>Aerosol liquid water content


<sup>f</sup>Tot-NH<sub>3</sub> = NH<sub>4</sub><sup>+</sup> + NH<sub>3</sub>

<sup>g</sup>Non-volatile cations







645 **Figure 1:** Average monthly (dotted line) and annual (diamonds) trends in a) sulfate, b) total nitrate (Tot- $\text{NO}_3 = \text{HNO}_3 + \text{NO}_3^-$ ), c) aerosol liquid water content (ALWC), d) total  $\text{NH}_3$  (Tot- $\text{NH}_3 = \text{NH}_3 + \text{NH}_4^+$ ), e) non-volatile cations (NVCs), and f) aerosol pH in Melpitz over 2010 – 2019. The solid line in each panel is the linear least squares regression fit for the annual averages.




**Figure 2:** Box and whisker plots showing seasonal a) sulfate concentrations, b) total nitrate (Tot- $\text{NO}_3 = \text{HNO}_3 + \text{NO}_3^-$ ) concentrations, c) total  $\text{NH}_3$  (Tot- $\text{NH}_3 = \text{NH}_3 + \text{NH}_4^+$ ) concentrations, d) aerosol pH, e) aerosol liquid water content (ALWC), and temperatures. Data shown are the 10<sup>th</sup> and 90<sup>th</sup> percentiles (whiskers), quartiles (upper and lower box), median (solid line), and mean values (circles).



**Figure 3:** Seasonal characteristics of a) total NH<sub>3</sub> concentrations (Tot-NH<sub>3</sub> = NH<sub>3</sub> + NH<sub>4</sub><sup>+</sup>), b) NH<sub>3</sub>(g) concentrations, c) NH<sub>4</sub><sup>+</sup> concentrations, and NH<sub>3</sub> partitioning ( $\epsilon_{\text{NH}_3} = \text{NH}_3(\text{g})/(\text{NH}_3(\text{g}) + \text{NH}_4^+)$ ), all presented as a function of ambient temperature.




**Figure 4:** Average diurnal profiles of a) aerosol liquid water content, b) temperature, c) relative humidity, d) total  $\text{NH}_3$  ( $\text{NH}_3(\text{g}) + \text{NH}_4^+$ ), e) total nitrate ( $\text{HNO}_3(\text{g}) + \text{NO}_3^-$ ), f) sulfate concentrations, and g) aerosol pH within each season.







**Figure 5:** Buffering capacity calculated according to Zheng et al. (2020) evaluated for various species.

675 Vertical green lines represent the region enclosed by mean annual pH range of this work.




680 **Figure 6:** Contribution of various factors to changes in aerosol pH at Melpitz in winter, spring, summer, fall, and overall, calculated according to Tao and Murphy (2021).





**Figure 7:** Chemical regimes in each season showing when Melpitz aerosol mass concentrations are sensitive to  $\text{HNO}_3$  availability,  $\text{NH}_3$  availability, both, or neither, calculated according to Nenes et al. (2020).



**Figure 8:** Pie charts showing the chemical regimes in Melpitz by season with the fraction of observations in which  $\text{NH}_4\text{NO}_3$  aerosol mass concentrations are sensitive to  $\text{HNO}_3$  availability,  $\text{NH}_3$  availability, both, or neither, calculated according to Nenes et al. (2020).



## 695 References

- Ahrens, L., Harner, T., Shoeib, M., Lane, D. A., and Murphy, J. G.: Improved Characterization of Gas-Particle Partitioning for Per- and Polyfluoroalkyl Substances in the Atmosphere Using Annular Diffusion Denuder Samplers, *Environmental Science & Technology*, 46, 7199-7206, 10.1021/es300898s, 2012.
- 700 Ansari, A. S. and Pandis, S. N.: Response of Inorganic PM to Precursor Concentrations, *Environmental Science & Technology*, 32, 2706-2714, 10.1021/es971130j, 1998.
- Ansari, A. S. and Pandis, S. N.: An analysis of four models predicting the partitioning of semivolatile inorganic aerosol components, *Aerosol Science and Technology*, 31, 129-153, Doi 10.1080/027868299304200, 1999.
- Atabakhsh, S., Poulain, L., Bigi, A., Coen, M. C., Pöhlker, M., and Herrmann, H.: Trends of PM1 aerosol chemical composition, carbonaceous aerosol, and source over the last 10 years at Melpitz (Germany), *Atmospheric Environment*, 346, 121075, <https://doi.org/10.1016/j.atmosenv.2025.121075>, 2025.
- 705 Atabakhsh, S., Poulain, L., Chen, G., Canonaco, F., Prévôt, A. S. H., Pöhlker, M., Wiedensohler, A., and Herrmann, H.: A 1-year aerosol chemical speciation monitor (ACSM) source analysis of organic aerosol particle contributions from anthropogenic sources after long-range transport at the TROPOS research station Melpitz, *Atmospheric Chemistry and Physics*, 23, 6963-6988, 10.5194/acp-23-6963-2023, 2023.
- 710 Backes, A., Aulinger, A., Bieser, J., Matthias, V., and Quante, M.: Ammonia emissions in Europe, part I: Development of a dynamical ammonia emission inventory, *Atmospheric Environment*, 131, 55-66, 10.1016/j.atmosenv.2016.01.041, 2016a.
- Backes, A. M., Aulinger, A., Bieser, J., Matthias, V., and Quante, M.: Ammonia emissions in Europe, part II: How ammonia emission abatement strategies affect secondary aerosols, *Atmospheric Environment*, 126, 153-161, <https://doi.org/10.1016/j.atmosenv.2015.11.039>, 2016b.
- 715 Baker, A. R., Kanakidou, M., Nenes, A., Myriokefalitakis, S., Croot, P. L., Duce, R. A., Gao, Y., Guieu, C., Ito, A., Jickells, T. D., Mahowald, N. M., Middag, R., Perron, M. M. G., Sarin, M. M., Shelley, R., and Turner, D. R.: Changing atmospheric acidity as a modulator of nutrient deposition and ocean biogeochemistry, *Science Advances*, 7, doi:10.1126/sciadv.abd8800, 2021.
- 720 Balamurugan, V., Chen, J., Qu, Z., Bi, X., and Keutsch, F. N.: Secondary PM2.5 decreases significantly less than NO2 emission reductions during COVID lockdown in Germany, *Atmos. Chem. Phys.*, 22, 7105-7129, 10.5194/acp-22-7105-2022, 2022.
- Banzhaf, S., Schaap, M., Wichink Kruit, R. J., Denier van der Gon, H. A. C., Stern, R., and Builtjes, P. J. H.: Impact of emission changes on secondary inorganic aerosol episodes across Germany, *Atmos. Chem. Phys.*, 13, 11675-11693, 10.5194/acp-13-11675-2013, 2013.
- 725 Battaglia Jr, M. A., Balasus, N., Ball, K., Caicedo, V., Delgado, R., Carlton, A. G., and Hennigan, C. J.: Urban aerosol chemistry at a land-water transition site during summer – Part 2: Aerosol pH and liquid water content, *Atmos. Chem. Phys. Discuss.*, 2021, 1-30, 10.5194/acp-2021-368, 2021.
- Battaglia, M. A., Douglas, S., and Hennigan, C. J.: Effect of the Urban Heat Island on Aerosol pH, *Environmental Science & Technology*, 51, 13095-13103, 10.1021/acs.est.7b02786, 2017.

- 730 Battaglia, M. A., Weber, R. J., Nenes, A., and Hennigan, C. J.: Effects of water-soluble organic carbon on aerosol pH, *Atmospheric Chemistry and Physics*, 19, 14607-14620, 10.5194/acp-19-14607-2019, 2019.
- Bougiatioti, A., Nikolaou, P., Stavroulas, I., Kouvarakis, G., Weber, R., Nenes, A., Kanakidou, M., and Mihalopoulos, N.: Particle water and pH in the eastern Mediterranean: source variability and implications for nutrient availability, *Atmos. Chem. Phys.*, 16, 4579-4591, 10.5194/acp-16-4579-2016, 2016.
- 735 Campbell, J. R., Battaglia, M., Dingilian, K. K., Cesler-Maloney, M., Simpson, W. R., Robinson, E. S., DeCarlo, P. F., Temime-Roussel, B., D'Anna, B., Holen, A. L., Wu, J., Pratt, K. A., Dibb, J. E., Nenes, A., Weber, R. J., and Mao, J.: Enhanced aqueous formation and neutralization of fine atmospheric particles driven by extreme cold, *Science Advances*, 10, eado4373, doi:10.1126/sciadv.ado4373, 2024.
- 740 Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system  $\text{H}^+\text{-NH}_4^+\text{-SO}_4^{2-}\text{-NO}_3^-\text{-H}_2\text{O}$  at tropospheric temperatures, *Journal of Physical Chemistry A*, 102, 2137-2154, DOI 10.1021/jp973042r, 1998.
- Dang, R., Jacob, D. J., Zhai, S., Coheur, P., Clarisse, L., Van Damme, M., Pendergrass, D. C., Choi, J.-s., Park, J.-s., Liu, Z., and Liao, H.: Diagnosing the Sensitivity of Particulate Nitrate to Precursor Emissions Using Satellite Observations of Ammonia and Nitrogen Dioxide, *Geophysical Research Letters*, 50, e2023GL105761, <https://doi.org/10.1029/2023GL105761>, 2023.
- 745 Ding, J., Zhao, P. S., Su, J., Dong, Q., Du, X., and Zhang, Y. F.: Aerosol pH and its driving factors in Beijing, *Atmospheric Chemistry and Physics*, 19, 7939-7954, 10.5194/acp-19-7939-2019, 2019.
- Dockery, D. W., Cunningham, J., Damokosh, A. I., Neas, L. M., Spengler, J. D., Koutrakis, P., Ware, J. H., Raizenne, M., and Speizer, F. E.: Health effects of acid aerosols on North American children: respiratory symptoms, *Environ Health Persp*, 104, 500-505, doi:10.1289/ehp.96104500, 1996.
- 750 Duan, X., Zheng, G., Chen, C., Zhang, Q., and He, K.: Driving factors of aerosol acidity: a new hierarchical quantitative analysis framework and its application in Changzhou, China, *Atmos. Chem. Phys.*, 25, 3919-3928, 10.5194/acp-25-3919-2025, 2025.
- Fang, T., Guo, H., Zeng, L., Verma, V., Nenes, A., and Weber, R. J.: Highly Acidic Ambient Particles, Soluble Metals, and Oxidative Potential: A Link between Sulfate and Aerosol Toxicity, *Environmental Science & Technology*, 51, 2611-2620, 10.1021/acs.est.6b06151, 2017a.
- 755 Fang, T., Guo, H. Y., Zeng, L. H., Verma, V., Nenes, A., and Weber, R. J.: Highly Acidic Ambient Particles, Soluble Metals, and Oxidative Potential: A Link between Sulfate and Aerosol Toxicity, *Environmental Science & Technology*, 51, 2611-2620, 10.1021/acs.est.6b06151, 2017b.
- Fortems-Cheiney, A., Dufour, G., Hamaoui-Laguel, L., Foret, G., Siour, G., Van Damme, M., Meleux, F., Coheur, P.-F., Clerbaux, C., Clarisse, L., Favez, O., Wallasch, M., and Beekmann, M.: Unaccounted variability in  $\text{NH}_3$  agricultural sources detected by IASI contributing to European spring haze episode, *Geophysical Research Letters*, 43, 5475-5482, <https://doi.org/10.1002/2016GL069361>, 2016.
- 760 Fridlind, A. M. and Jacobson, M. Z.: A study of gas-aerosol equilibrium and aerosol pH in the remote marine boundary layer during the First Aerosol Characterization Experiment (ACE 1), *Journal of Geophysical Research: Atmospheres*, 105, 17325-17340, <https://doi.org/10.1029/2000JD900209>, 2000.

- Guo, H., Xu, L., Bougiatioti, A., Cerully, K. M., Capps, S. L., Hite, J. R., Carlton, A. G., Lee, S. H., Bergin, M. H., Ng, N. L., Nenes, A., and Weber, R. J.: Fine-particle water and pH in the southeastern United States, *Atmospheric Chemistry and Physics*, 15, 5211-5228, 10.5194/acp-15-5211-2015, 2015.
- 770 Guo, H. Y., Nenes, A., and Weber, R. J.: The underappreciated role of nonvolatile cations in aerosol ammonium-sulfate molar ratios, *Atmospheric Chemistry and Physics*, 18, 17307-17323, 10.5194/acp-18-17307-2018, 2018a.
- Guo, H. Y., Otjes, R., Schlag, P., Kiendler-Scharr, A., Nenes, A., and Weber, R. J.: Effectiveness of ammonia reduction on control of fine particle nitrate, *Atmospheric Chemistry and Physics*, 18, 12241-12256, 10.5194/acp-18-12241-2018, 2018b.
- 775 Guo, H. Y., Liu, J. M., Froyd, K. D., Roberts, J. M., Veres, P. R., Hayes, P. L., Jimenez, J. L., Nenes, A., and Weber, R. J.: Fine particle pH and gas-particle phase partitioning of inorganic species in Pasadena, California, during the 2010 CalNex campaign, *Atmospheric Chemistry and Physics*, 17, 5703-5719, 10.5194/acp-17-5703-2017, 2017.
- Hamed, A., Birmili, W., Joutsensaari, J., Mikkonen, S., Asmi, A., Wehner, B., Spindler, G., Jaatinen, A., Wiedensohler, A., Korhonen, H., Lehtinen, K. E. J., and Laaksonen, A.: Changes in the production rate of secondary aerosol particles in Central Europe in view of decreasing SO<sub>2</sub> emissions between 1996 and 2006, *Atmospheric Chemistry and Physics*, 10, 1071-1091, DOI 10.5194/acp-10-1071-2010, 2010.
- 780 Jonson, J. E., Fagerli, H., Scheuschner, T., and Tsyro, S.: Modelling changes in secondary inorganic aerosol formation and nitrogen deposition in Europe from 2005 to 2030, *Atmospheric Chemistry and Physics*, 22, 1311-1331, 10.5194/acp-22-1311-2022, 2022.
- Kakavas, S., Patoulias, D., Zakoura, M., Nenes, A., and Pandis, S. N.: Size-resolved aerosol pH over Europe during summer, *Atmospheric Chemistry and Physics*, 21, 799-811, 10.5194/acp-21-799-2021, 2021.
- 785 Keene, W. C., Pszenny, A. A. P., Maben, J. R., Stevenson, E., and Wall, A.: Closure evaluation of size-resolved aerosol pH in the New England coastal atmosphere during summer, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2004JD004801>, 2004.
- Lawrence, C. E., Casson, P., Brandt, R., Schwab, J. J., Dukett, J. E., Snyder, P., Yerger, E., Kelting, D., VandenBoer, T. C., and Lance, S.: Long-term monitoring of cloud water chemistry at Whiteface Mountain: the emergence of a new chemical regime, *Atmospheric Chemistry and Physics*, 23, 1619-1639, 10.5194/acp-23-1619-2023, 2023.
- 790 Liu, Z., Rieder, H. E., Schmidt, C., Mayer, M., Guo, Y., Winiwarter, W., and Zhang, L.: Optimal reactive nitrogen control pathways identified for cost-effective PM<sub>2.5</sub> mitigation in Europe, *Nature Communications*, 14, 4246, 10.1038/s41467-023-39900-9, 2023.
- Mahowald, N.: Aerosol Indirect Effect on Biogeochemical Cycles and Climate, *Science*, 334, 794-796, doi:10.1126/science.1207374, 2011.
- 795 Mahowald, N. M., Hamilton, D. S., Mackey, K. R. M., Moore, J. K., Baker, A. R., Scanza, R. A., and Zhang, Y.: Aerosol trace metal leaching and impacts on marine microorganisms, *Nature Communications*, 9, doi:10.1038/s41467-018-04970-7, 2018.
- Meskhidze, N., Chameides, W. L., Nenes, A., and Chen, G.: Iron mobilization in mineral dust: Can anthropogenic SO<sub>2</sub> emissions affect ocean productivity?: *Geophysical Research Letters*, 30, doi:10.1029/2003gl018035, 2003.
- 800 Nah, T., Guo, H., Sullivan, A. P., Chen, Y., Tanner, D. J., Nenes, A., Russell, A., Ng, N. L., Huey, L. G., and Weber, R. J.: Characterization of aerosol composition, aerosol acidity, and organic acid partitioning at an agriculturally intensive rural southeastern US site, *Atmos. Chem. Phys.*, 18, 11471-11491, 10.5194/acp-18-11471-2018, 2018.

NEC: Air pollution in Europe: 2024 reporting status under the National Emission reduction Commitments Directive, doi:10.2800/977119, 2024.

- 805 Nenes, A., Pandis, S. N., Weber, R. J., and Russell, A.: Aerosol pH and liquid water content determine when particulate matter is sensitive to ammonia and nitrate availability, *Atmos. Chem. Phys.*, 20, 3249-3258, 10.5194/acp-20-3249-2020, 2020.
- Nenes, A., Pandis, S. N., Kanakidou, M., Russell, A. G., Song, S., Vasilakos, P., and Weber, R. J.: Aerosol acidity and liquid water content regulate the dry deposition of inorganic reactive nitrogen, *Atmos. Chem. Phys.*, 21, 6023-6033, 10.5194/acp-21-6023-2021, 2021.
- 810 Paglione, M., Decesari, S., Rinaldi, M., Tarozzi, L., Manarini, F., Gilardoni, S., Facchini, M. C., Fuzzi, S., Bacco, D., Trentini, A., Pandis, S. N., and Nenes, A.: Historical Changes in Seasonal Aerosol Acidity in the Po Valley (Italy) as Inferred from Fog Water and Aerosol Measurements, *Environmental Science & Technology*, 55, 7307-7315, 10.1021/acs.est.1c00651, 2021.
- Pan, D., Mauzerall, D. L., Wang, R., Guo, X., Puchalski, M., Guo, Y., Song, S., Tong, D., Sullivan, A. P., Schichtel, B. A., Collett, J. L., and Zondlo, M. A.: Regime shift in secondary inorganic aerosol formation and nitrogen deposition in the rural  
815 United States, *Nature Geoscience*, 17, 617-623, 10.1038/s41561-024-01455-9, 2024.
- Pay, M. T., Jiménez-Guerrero, P., and Baldasano, J. M.: Assessing sensitivity regimes of secondary inorganic aerosol formation in Europe with the CALIOPE-EU modeling system, *Atmospheric Environment*, 51, 146-164, <https://doi.org/10.1016/j.atmosenv.2012.01.027>, 2012.
- Petetin, H., Sciare, J., Bressi, M., Gros, V., Rosso, A., Sanchez, O., Sarda-Estève, R., Petit, J. E., and Beekmann, M.: Assessing  
820 the ammonium nitrate formation regime in the Paris megacity and its representation in the CHIMERE model, *Atmos. Chem. Phys.*, 16, 10419-10440, 10.5194/acp-16-10419-2016, 2016.
- Phillips, S. M., Bellcross, A. D., and Smith, G. D.: Light Absorption by Brown Carbon in the Southeastern United States is pH-dependent, *Environmental Science & Technology*, 51, 6782-6790, 10.1021/acs.est.7b01116, 2017.
- Pye, H. O. T., Nenes, A., Alexander, B., Ault, A. P., Barth, M. C., Clegg, S. L., Collett, J. L., Fahey, K. M., Hennigan, C. J.,  
825 Herrmann, H., Kanakidou, M., Kelly, J. T., Ku, I. T., McNeill, V. F., Riemer, N., Schaefer, T., Shi, G. L., Tilgner, A., Walker, J. T., Wang, T., Weber, R., Xing, J., Zaveri, R. A., and Zuend, A.: The acidity of atmospheric particles and clouds, *Atmospheric Chemistry and Physics*, 20, 4809-4888, 10.5194/acp-20-4809-2020, 2020.
- Ren, Y. G., Stieger, B., Spindler, G., Grosselin, B., Mellouki, A., Tuch, T., Wiedensohler, A., and Herrmann, H.: Role of the  
830 dew water on the ground surface in HONO distribution: a case measurement in Melpitz, *Atmospheric Chemistry and Physics*, 20, 13069-13089, 10.5194/acp-20-13069-2020, 2020.
- Renner, E. and Wolke, R.: Modelling the formation and atmospheric transport of secondary inorganic aerosols with special attention to regions with high ammonia emissions, *Atmospheric Environment*, 44, 1904-1912, <https://doi.org/10.1016/j.atmosenv.2010.02.018>, 2010.
- Rumsey, I. C., Cowen, K. A., Walker, J. T., Kelly, T. J., Hanft, E. A., Mishoe, K., Rogers, C., Proost, R., Beachley, G. M.,  
835 Lear, G., Frelink, T., and Otjes, R. P.: An assessment of the performance of the Monitor for AeRosols and GAs in ambient air (MARGA): a semi-continuous method for soluble compounds, *Atmospheric Chemistry and Physics*, 14, 5639-5658, 10.5194/acp-14-5639-2014, 2014.
- Shah, V., Jaeglé, L., Thornton, J. A., Lopez-Hilfiker, F. D., Lee, B. H., Schroder, J. C., Campuzano-Jost, P., Jimenez, J. L.,  
840 Guo, H., Sullivan, A. P., Weber, R. J., Green, J. R., Fiddler, M. N., Bililign, S., Campos, T. L., Stell, M., Weinheimer, A. J., Montzka, D. D., and Brown, S. S.: Chemical feedbacks weaken the wintertime response of particulate sulfate and nitrate to

emissions reductions over the eastern United States, *Proceedings of the National Academy of Sciences*, 115, 8110-8115, doi:10.1073/pnas.1803295115, 2018.

Song, X., Wu, D., Su, Y., Li, Y., and Li, Q.: Review of health effects driven by aerosol acidity: Occurrence and implications for air pollution control, *Sci Total Environ*, 955, 176839, <https://doi.org/10.1016/j.scitotenv.2024.176839>, 2024.

- 845 Spindler, G., Grüner, A., Müller, K., Schlimper, S., and Herrmann, H.: Long-term size-segregated particle (PM<sub>10</sub>, PM<sub>2.5</sub>, PM<sub>1</sub>) characterization study at Melpitz - influence of air mass inflow, weather conditions and season, *Journal of Atmospheric Chemistry*, 70, 165-195, 10.1007/s10874-013-9263-8, 2013.

- Spindler, G., Müller, K., Brüggemann, E., Gnauk, T., and Herrmann, H.: Long-term size-segregated characterization of PM<sub>10</sub>, PM<sub>2.5</sub>, and PM<sub>1</sub> at the IfT research station Melpitz downwind of Leipzig (Germany) using high and low-volume filter  
850 samplers, *Atmospheric Environment*, 38, 5333-5347, 10.1016/j.atmosenv.2003.12.047, 2004.

Spindler, G., Brüggemann, E., Gnauk, T., Grüner, A., Müller, K., and Herrmann, H.: A four-year size-segregated characterization study of particles PM<sub>10</sub>, PM<sub>2.5</sub> and PM<sub>1</sub> depending on air mass origin at Melpitz, *Atmospheric Environment*, 44, 164-173, 10.1016/j.atmosenv.2009.10.015, 2010.

- Squizzato, S., Masiol, M., Brunelli, A., Pistollato, S., Tarabotti, E., Rampazzo, G., and Pavoni, B.: Factors determining the  
855 formation of secondary inorganic aerosol: a case study in the Po Valley (Italy), *Atmospheric Chemistry and Physics*, 13, 1927-1939, 10.5194/acp-13-1927-2013, 2013.

Stieger, B., Spindler, G., van Pinxteren, D., Grüner, A., Wallasch, M., and Herrmann, H.: Development of an online-coupled MARGA upgrade for the 2 h interval quantification of low-molecular-weight organic acids in the gas and particle phases, *Atmos Meas Tech*, 12, 281-298, 10.5194/amt-12-281-2019, 2019.

- 860 Stieger, B., van Pinxteren, D., Tilgner, A., Spindler, G., Poulain, L., Grüner, A., Wallasch, M., and Herrmann, H.: Strong Deviations from Thermodynamically Expected Phase Partitioning of Low-Molecular-Weight Organic Acids during One Year of Rural Measurements, *ACS Earth and Space Chemistry*, 5, 500-515, 10.1021/acsearthspacechem.0c00297, 2021.

- Stieger, B., Spindler, G., Fahlbusch, B., Müller, K., Grüner, A., Poulain, L., Thöni, L., Seitler, E., Wallasch, M., and Herrmann, H.: Measurements of PM<sub>10</sub> ions and trace gases with the online system MARGA at the research station Melpitz in Germany  
865 - A five-year study, *Journal of Atmospheric Chemistry*, 75, 33-70, 10.1007/s10874-017-9361-0, 2018.

- Sutton, M. A., Reis, S., Riddick, S. N., Dragosits, U., Nemitz, E., Theobald, M. R., Tang, Y. S., Braban, C. F., Vieno, M., Dore, A. J., Mitchell, R. F., Wanless, S., Daunt, F., Fowler, D., Blackall, T. D., Milford, C., Flechard, C. R., Loubet, B., Massad, R., Cellier, P., Personne, E., Coheur, P. F., Clarisse, L., Van Damme, M., Ngadi, Y., Clerbaux, C., Skjoth, C. A., Geels, C., Hertel, O., Kruit, R. J. W., Pinder, R. W., Bash, J. O., Walker, J. T., Simpson, D., Horváth, L., Misselbrook, T. H.,  
870 Bleeker, A., Dentener, F., and de Vries, W.: Towards a climate-dependent paradigm of ammonia emission and deposition, *Philos T R Soc B*, 368, doi:10.1098/rstb.2013.0166, 2013.

Tao, Y. and Murphy, J. G.: The sensitivity of PM<sub>2.5</sub> acidity to meteorological parameters and chemical composition changes: 10-year records from six Canadian monitoring sites, *Atmospheric Chemistry and Physics*, 19, 9309-9320, 10.5194/acp-19-9309-2019, 2019.

- 875 Tao, Y. and Murphy, J. G.: Simple Framework to Quantify the Contributions from Different Factors Influencing Aerosol pH Based on NH<sub>x</sub> Phase-Partitioning Equilibrium, *Environmental Science & Technology*, 55, 10310-10319, 10.1021/acs.est.1c03103, 2021.

- ten Brink, H., Otjes, R., Jongejan, P., and Slanina, S.: An instrument for semi-continuous monitoring of the size-distribution of nitrate, ammonium, sulphate and chloride in aerosol, *Atmospheric Environment*, 41, 2768-2779, <https://doi.org/10.1016/j.atmosenv.2006.11.041>, 2007.
- Tilgner, A., Schaefer, T., Alexander, B., Barth, M., Collett Jr, J. L., Fahey, K. M., Nenes, A., Pye, H. O. T., Herrmann, H., and McNeill, V. F.: Acidity and the multiphase chemistry of atmospheric aqueous particles and clouds, *Atmos. Chem. Phys.*, 21, 13483-13536, 10.5194/acp-21-13483-2021, 2021.
- Trebs, I., Metzger, S., Meixner, F. X., Helas, G. N., Hoffer, A., Rudich, Y., Falkovich, A. H., Moura, M. A. L., da Silva, R. S., Artaxo, P., Slanina, J., and Andreae, M. O.: The  $\text{NH}_4^+ \text{-NO}_3^- \text{-Cl}^- \text{-SO}_4^{2-} \text{-H}_2\text{O}$  aerosol system and its gas phase precursors at a pasture site in the Amazon Basin:: How relevant are mineral cations and soluble organic acids? -: art. no. D07303, *Journal of Geophysical Research-Atmospheres*, 110, doi:10.1029/2004jd005478, 2005.
- Turnock, S. T., Spracklen, D. V., Carslaw, K. S., Mann, G. W., Woodhouse, M. T., Forster, P. M., Haywood, J., Johnson, C. E., Dalvi, M., Bellouin, N., and Sanchez-Lorenzo, A.: Modelled and observed changes in aerosols and surface solar radiation over Europe between 1960 and 2009, *Atmospheric Chemistry and Physics*, 15, 9477-9500, 10.5194/acp-15-9477-2015, 2015.
- Van Damme, M., Clarisse, L., Franco, B., Sutton, M. A., Erismann, J. W., Kruit, R. W., van Zanten, M., Whitburn, S., Hadji-Lazaro, J., Hurtmans, D., Clerbaux, C., and Coheur, P. F.: Global, regional and national trends of atmospheric ammonia derived from a decadal (2008-2018) satellite record, *Environ Res Lett*, 16, doi:10.1088/1748-9326/abd5e0, 2021.
- van der Graaf, S., Dammers, E., Segers, A., Kranenburg, R., Schaap, M., Shephard, M. W., and Erismann, J. W.: Data assimilation of CrIS  $\text{NH}_3$  satellite observations for improving spatiotemporal  $\text{NH}_3$  distributions in LOTOS-EUROS, *Atmospheric Chemistry and Physics*, 22, 951-972, 10.5194/acp-22-951-2022, 2022.
- Vestreng, V., Myhre, G., Fagerli, H., Reis, S., and Tarrasón, L.: Twenty-five years of continuous sulphur dioxide emission reduction in Europe, *Atmospheric Chemistry and Physics*, 7, 3663-3681, DOI 10.5194/acp-7-3663-2007, 2007.
- Vestreng, V., Ntziachristos, L., Semb, A., Reis, S., Isaksen, I. S. A., and Tarrasón, L.: Evolution of  $\text{NO}_x$  emissions in Europe with focus on road transport control measures, *Atmospheric Chemistry and Physics*, 9, 1503-1520, 10.5194/acp-9-1503-2009, 2009.
- Viatte, C., Abeed, R., Yamanouchi, S., Porter, W. C., Safieddine, S., Van Damme, M., Clarisse, L., Herrera, B., Grutter, M., Coheur, P. F., Strong, K., and Clerbaux, C.:  $\text{NH}_3$  spatiotemporal variability over Paris, Mexico City, and Toronto, and its link to  $\text{PM}_{2.5}$  during pollution events, *Atmospheric Chemistry and Physics*, 22, 12907-12922, doi:10.5194/acp-22-12907-2022, 2022.
- Wang, G., Tao, Y., Chen, J., Liu, C., Qin, X., Li, H., Yun, L., Zhang, M., Zheng, H., Gui, H., Liu, J., Huo, J., Fu, Q., Deng, C., and Huang, K.: Quantitative Decomposition of Influencing Factors to Aerosol pH Variation over the Coasts of the South China Sea, East China Sea, and Bohai Sea, *Environmental Science & Technology Letters*, 9, 815-821, 10.1021/acs.estlett.2c00527, 2022.
- Weber, R. J., Guo, H. Y., Russell, A. G., and Nenes, A.: High aerosol acidity despite declining atmospheric sulfate concentrations over the past 15 years, *Nature Geoscience*, 9, 282-+, 10.1038/ngeo2665, 2016.
- Wentworth, G. R., Murphy, J. G., Benedict, K. B., Bangs, E. J., and Collett, J.: The role of dew as a night-time reservoir and morning source for atmospheric ammonia, *Atmospheric Chemistry and Physics*, 16, 7435-7449, 10.5194/acp-16-7435-2016, 2016.

- 915 Wexler, A. S. and Clegg, S. L.: Atmospheric aerosol models for systems including the ions  $\text{H}^+$ ,  $\text{NH}_4^+$ ,  $\text{Na}^+$ ,  $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Cl}^-$ ,  $\text{Br}^-$ , and  $\text{H}_2\text{O}$ : *Journal of Geophysical Research-Atmospheres*, 107, doi:10.1029/2001jd000451, 2002.
- Womack, C. C., McDuffie, E. E., Edwards, P. M., Bares, R., de Gouw, J. A., Docherty, K. S., Dubé, W. P., Fibiger, D. L., Franchin, A., Gilman, J. B., Goldberger, L., Lee, B. H., Lin, J. C., Long, R., Middlebrook, A. M., Millet, D. B., Moravek, A., Murphy, J. G., Quinn, P. K., Riedel, T. P., Roberts, J. M., Thornton, J. A., Valin, L. C., Veres, P. R., Whitehill, A. R., Wild, R. J., Warneke, C., Yuan, B., Baasandorj, M., and Brown, S. S.: An Odd Oxygen Framework for Wintertime Ammonium Nitrate Aerosol Pollution in Urban Areas: NO<sub>x</sub> and VOC Control as Mitigation Strategies, *Geophysical Research Letters*, 46, 4971-4979, <https://doi.org/10.1029/2019GL082028>, 2019.
- 920 Zakoura, M. and Pandis, S. N.: Improving fine aerosol nitrate predictions using a Plume-in-Grid modeling approach, *Atmospheric Environment*, 215, doi:10.1016/j.atmosenv.2019.116887, 2019.
- 925 Zhai, S., Jacob, D. J., Wang, X., Liu, Z., Wen, T., Shah, V., Li, K., Moch, J. M., Bates, K. H., Song, S., Shen, L., Zhang, Y., Luo, G., Yu, F., Sun, Y., Wang, L., Qi, M., Tao, J., Gui, K., Xu, H., Zhang, Q., Zhao, T., Wang, Y., Lee, H. C., Choi, H., and Liao, H.: Control of particulate nitrate air pollution in China, *Nature Geoscience*, 14, 389-395, 10.1038/s41561-021-00726-z, 2021.
- 930 Zheng, G. J., Su, H., Wang, S. W., Andreae, M. O., Poschl, U., and Cheng, Y. F.: Multiphase buffer theory explains contrasts in atmospheric aerosol acidity, *Science*, 369, 1374-+, 10.1126/science.aba3719, 2020.
- Zhou, M., Zheng, G., Wang, H., Qiao, L., Zhu, S., Huang, D., An, J., Lou, S., Tao, S., Wang, Q., Yan, R., Ma, Y., Chen, C., Cheng, Y., Su, H., and Huang, C.: Long-term trends and drivers of aerosol pH in eastern China, *Atmos. Chem. Phys.*, 22, 13833-13844, 10.5194/acp-22-13833-2022, 2022.