

Retrieving root-zone soil moisture from land surface modelling and GRACE/-FO and validating its dynamics with in-situ data over West Africa

Loudi Yap^{1,2}, Jürgen Kusche², Bamidele Oloruntoba^{3,4}, Helena Gerdener², and Harrie-Jan
Hendricks Franssen^{3,4}

¹Research Laboratory in Geodesy, National Institute of Cartography, Cameroon

²Institute of Geodesy and Geoinformation (IGG), University of Bonn, Germany

³Institute of Bio- and Geosciences: Agrosphere (IBG-3), Forschungszentrum Jülich GmbH,
52425 Jülich, Germany

⁴Centre for High-Performance Scientific Computing in Terrestrial Systems, Geoverbund
ABC/J, 52425 Jülich, Germany.

Corresponding author: L. Yap. E-mail addresses: loudiyap@uni-bonn.de, loudiyap@yahoo.fr.

June 29, 2026

Abstract

Rainfall variability in West Africa, driven by the West African Monsoon, poses significant challenges to agricultural productivity and livelihoods. In this context, understanding root-zone soil moisture (RZSM) dynamics is crucial since it serves as the primary water source for crops. While surface soil moisture (SSM) has been widely studied, research on RZSM remains limited. This study investigates RZSM dynamics across West Africa from 2003 to 2019 using multiple satellite-derived and model-based datasets, including ESA CCI v0.81, GLWS2.0, WaterGAP, CLM5.0, and in-situ observations. Results indicate that ESA CCI exhibits the strongest temporal and spatial alignment with ground measurements, whereas CLM5.0 and GLWS2.0 effectively capture latitudinal soil moisture ([SM](#)) gradients associated with climatic zones. A novel application of an analytical solution to Richards' equation was employed to translate surface moisture signals to deeper soil layers, demonstrating GLWS2.0's superior ability to reproduce seasonal patterns at various depths, notably in Benin and Niger. Despite challenges posed by sparse in-situ data and vegetation-induced signal attenuation, the study highlights the significant benefits of GRACE/-FO data assimilation in

enhancing model accuracy. The proposed depth-projection methodology improves the vertical representation of ~~soil-moisture~~SM, offering new insights into the dynamics of surface and subsurface water storage. These findings have important implications for agricultural forecasting, sustainable water resource management, and climate adaptation strategies in regions where accurate ~~soil~~moisture-SM data are essential for resilience planning.

Keywords

West Africa, Root-zone soil moisture, Satellite-derived soil moisture, Richards' equation, GRACE/-FO data assimilation

Highlights

- Comprehensive assessment of ~~root-zone-soil-moisture (RZSM-)~~RZSM dynamics across West Africa (2003–2019) using multiple satellite- and model-based datasets.
- Novel depth-projection approach based on Richards' equation translates ~~surface-soil-moisture~~SM signals to deeper layers, enhancing geodetic representation of subsurface water storage.
- ESA CCI shows the strongest temporal alignment with in-situ observations, while GLWS2.0 and CLM5.0 capture latitudinal and seasonal ~~soil-moisture (SM-)~~SM patterns effectively.
- Integration of GRACE/-FO satellite gravimetry improves GLWS2.0 accuracy, supporting geodetic-based monitoring of water resources and hydrological forecasting.
- Methodological framework advances understanding of surface and subsurface water mass redistribution, contributing to geodesy-informed sustainable water management and climate adaptation strategies.

1 Introduction

Rainfall in West Africa is largely driven by the West African Monsoon (WAM), characterized by significant spatial and temporal variability (?). This variability, often sporadic and unpredictable, increases the region’s vulnerability to droughts and floods, severely impacting agricultural productivity and leading to crop failures in rainfed systems (???). As a result, smallholder farmers, reliant on rainfed agriculture and constrained by financial limitations, face heightened risks to their livelihoods (?). These challenges significantly hinder economic development and exacerbate poverty in this already vulnerable region, which relies on agriculture for the livelihoods of around 70% of its estimated 420 million people, with a rapidly growing population at an annual rate of 2.2 – 2.8% (?). ~~Root-zone soil moisture (RZSM)~~ RZSM which refers to the amount of water stored in the soil within the root zone of vegetation, typically the top 1-2 meters serve as the primary water source for crops (?). RZSM directly influences plant growth, agricultural productivity, and water availability for ecosystems (????). Unlike ~~surface soil moisture (SSM)~~ SSM, which can quickly change due to weather conditions, RZSM represents the longer-term water storage available to plants, playing a key role in determining drought resilience and crop yields (?). Given that approximately 75% of the total crop area harvested globally consists of non-irrigated crops (??), the importance of RZSM in global food production and food security becomes even more pronounced. Monitoring RZSM is essential for understanding the water balance (?), drought and flood warning (??), managing irrigation (??), and modeling climate impacts on agriculture and natural vegetation (?). It potentially enhances forecasts and climate projections, guides water resources management, and supports precision agriculture by optimizing water usage.

Currently, ~~soil moisture SM~~ SM products (SSM or RZSM) can be generated using three different key approaches: in situ observations, remote sensing, and modelling (?). In situ observations involve ground-based sensors that measure ~~soil moisture SM~~ SM directly at specific points using gravimetric, tensiometric and nuclear methods (?). These measurements are reasonably accurate and can capture ~~soil moisture SM~~ SM at different depths. However, while in situ observations offer precision, they are labor-intensive and expensive to maintain, and their limited spatial coverage makes them difficult to scale across large regions (?). This poses challenges in using them for extensive, long-term monitoring over large areas. Nevertheless, the point-scale ground observations are often used as a benchmark for calibrating and validating ~~soil moisture SM~~ SM estimates from remote sensing and model simulations (???). In recent years, significant efforts have been made to establish in situ ~~soil moisture SM~~ SM monitoring networks across Africa, particularly exemplified by the AMMA-CATCH (African Monsoon Multi-disciplinary Analysis–Couplage de l’Atmosphère Tropicale et du Cycle Hydrologique) observa-

tory (?). Some of the data collected from these networks have been integrated into the International
 Soil Moisture Network (ISMN)(<https://ismn.geo.tuwien.ac.at/>) (?). These sparse in situ monitoring
 networks have played a crucial role in validating remotely sensed and simulated ~~soil-moisture~~ SM
 estimates over extended periods in various African regions (?). Remote sensing utilizes various meth-
 ods, such as microwave, optical, and thermal satellite sensors, to estimate ~~surface-soil-moisture~~ SSM
 across large areas (??). It is an effective technique for detecting the dynamic patterns of ~~soil-moisture~~
SM on regional and global scales. Various satellite instruments, including the Soil Moisture Active
 Passive (SMAP), Soil Moisture and Ocean Salinity (SMOS), METOP-A/B Advanced Scatterometer
 (ASCAT), Advanced Microwave Scanning Radiometer–EOS (AMSR-E), and products from the Euro-
 pean Space Agency’s Climate Change Initiative (ESA CCI), have been successfully utilized to retrieve
 SSM at a global scale with a temporal resolution of 2 to 3 days (???????). While the remote sensing
 approach offers broad spatial coverage and frequent updates, it primarily measures ~~soil-moisture~~ SM in
 the uppermost few centimeters (0–5 cm), often missing the deeper root zone dynamics. Additionally,
 its accuracy can be impacted by factors such as vegetation, weather conditions, radio frequency inter-
 ference (RFI), and topography, which can reduce measurement reliability. In contrast, the GRACE
 (Gravity Recovery and Climate Experiment) mission captures changes in total water storage by map-
 ping variations in Earth’s gravity field (?). It has been demonstrated that GRACE-observed Total
 Water Storage Anomalies (TWSA) can be translated into ~~surface-or-root-zone-soil-moisture~~ (SSM or
 RZSM) using a physically based approach (??). Although GRACE-based ~~soil-moisture~~ SM data have
 lower spatial resolution $\sim 3^\circ$ vs. 40 km for microwave data) and less frequent temporal sampling
 (monthly vs. daily), data assimilation and downscaling algorithms can be applied to make these two
 approaches comparable (?). Additionally, GRACE data are not affected at all by vegetation density
 and RFI. This study ~~will incorporate this~~ incorporates this advantage by combining GRACE-based
observations with a modelled data approach. The modelled data approach uses simulations that inte-
 grate climatic, soil, and vegetation information to estimate both ~~surface-and-root-zone-soil-moisture~~
SSM and RZSM across various spatial and temporal scales. Whether based on hydrological or land
 surface models, both approaches rely on similar equations to simulate ~~soil-moisture~~ SM according a
 water balance approach (?). Modelling allows for ~~soil-moisture~~ SM estimates to be generated at high
 spatial and temporal resolution, offering detailed insights into ~~soil-moisture~~ SM dynamics. However,
 while models provide comprehensive coverage and long-term predictions, their accuracy is highly de-
 pendent on the quality of meteorological input data and the assumptions made in parameterization,
 which can introduce significant uncertainties. As a result, each modelling approach has its strengths
 and limitations, and combining multiple methods can provide a more robust and reliable understand-

ing of ~~soil-moisture~~ SM dynamics. While numerous studies have focused on monitoring remotely sensed and modeled ~~surface-soil-moisture~~ SSM data over West Africa (???????) using in situ data, there has been little to no research, to our knowledge, specifically examining ~~root-zone-soil-moisture~~ RZSM in this region. Furthermore, while the remote sensing products monitored in this area primarily involve AMSR-E satellite data (???), the SMOS satellite mission (??), and ASCAT satellite data (?), ~~soil-moisture~~ SM products based on ESA CCI, CLM5.0, and GRACE/-FO assimilated data have not been comprehensively validated in this region. However, while many physically-based land surface models (e.g., CLM5.0) simulate the ~~soil-moisture~~ SM patterns at different depths by numerically solving Richard's equation, this remains a challenge for conceptual hydrological models. This study aims to retrieve the ~~root-zone-soil-moisture~~ RZSM from a conceptual model, WaterGAP, as well as the GRACE/-FO-based global assimilation model GLWS2.0 which is based on WaterGAP. The dynamics of these estimates will be validated against in-situ measurements, while additional ~~soil-moisture~~ SM products, including ESA CCI and CLM5.0, will be used for comparative analysis across the West Africa region. This is the first study, to our knowledge, that uses Richard's equation to consistently compare WaterGAP and GLWS2.0 (i.e., GRACE-derived ~~root-zone-soil-moisture~~ RZSM) to in-situ data sampled at different depths.

Our main research questions ~~are:~~ addressed in this study are: What is the correlation between the ~~soil-moisture (SM)~~ SM from the GRACE/-FO-based global assimilation model (GLWS2.0) and ESA CCI, in-situ data, and other land surface models in the region? How can the water content in the single ~~soil-moisture~~ SM reservoir from the conceptual hydrological models be translated to a ~~soil moisture~~ SM vertical profile? How does the ~~root-zone-soil-moisture (RZSM)~~ RZSM changes at each retrieval depth over 2003–2019 in this region? and What is the correlation of its dynamics with the physically-based model (CLM5.0), ESA CCI products, and in-situ data?

The analytical solution of Richards' equation (?) will be used to translate water content from the single ~~soil-moisture~~ SM reservoir in GLWS2.0 and WaterGAP to different depths. Our approach offers a distinct advantage over that of ?, who assumes that GRACE TWSA (~~Total Water Storage Anomaly~~) data have minimal contributions from sources such as groundwater, surface water, or lateral groundwater flow. In their case, any TWSA variation not physically attributable to ~~soil-moisture~~ SM and incompatible with their model is classified as error. By contrast, we work directly with the ~~soil moisture~~ SM anomaly derived from GLWS2.0 (or WaterGAP), where ~~non-soil-moisture~~ non-SM components have already been filtered out, offering a cleaner signal for analysis. However, it is important to note that complete separation of these additional hydrological signals from ~~soil-moisture~~ SM may still be imperfect within the assimilation or WaterGAP model. WaterGAP (and thus GLWS2.0) rep-

resents ~~soil-moisture~~ SM via a single layer that extends to the root zone. The model simulates varying surface water storages (lakes, wetlands, rivers, and reservoirs) and includes a conceptual groundwater representation. Human water use, i.e. surface and groundwater abstractions, are included in Water-GAP. This approach provides a promising solution to not only expand the shallow vertical support of the microwave satellites with good spatial resolution but also to better isolate the groundwater storage dynamics from the GRACE/-FO-based assimilated signal. This is particularly relevant in West Africa, where ~~soil-moisture~~ SM has been shown to be the dominant component of Total Water Storage (TWS) (???).

2 Study area, datasets and models

2.1 Study area

The study area encompasses West Africa, spanning from 17°W to 17°E and 2°N to 21°N (Fig. 1). The terrain in this region is predominantly low and flat (?). The area is characterized by a latitudinal gradient that includes three bioclimatic regions, progressing from north to south: the Sahelian, Sudanian, and Guinean zones (?). This gradient results in varying vegetation patterns (Fig.2), with the arid north experiencing a single rainy season and sparse vegetation, while the south has two rainy seasons and dense vegetation (?). Three meso-scale sites located at different latitudes in Benin, Niger, and Senegal, where AMMA SM observations have undergone advanced quality control procedures by the ISMN, have been selected for analysis in this study (Fig. 1). The Benin site, situated in the Sudanian climate zone, features sandy clay loam soils, woody savanna vegetation, and gently undulating topography (630–225 m asl), with about 1200 mm of annual rainfall concentrated in a single rainy season from April to October (?). Both the Niger and Senegal sites are located in the Sahel region characterized by a single rainy season between June and October. The Niger site experiences a semi-arid tropical climate, characterized by a long dry season from October to May, with an average yearly temperature of 29.2°C and 520 mm of annual rainfall (1990–2007)(?). The landscape features flat lateritic plateaus and sandy valleys within the Iullemeden sedimentary basin, which has endorheic hydrology and a continental terminal aquifer. Soils are sandy and weakly structured, contributing to erosion. Additionally, the original woody savannah has transformed into a mosaic of rainfed millet fields and shrubby savannah, mixed with degraded tiger bush vegetation (?). The Senegal site, situated in the Dahra region (15.432°W – 15.403°N), has a Sahelian climate with a mean yearly temperature of 29°C, peaking in May, and an annual precipitation of approximately 420 mm. The area features herbaceous vegetation dominated by annual grasses and a tree cover of about 3%,

178 with most water bodies being temporary, except for a few permanent ponds (??).




Figure 1: Location of the three meso-scale study sites in West Africa where in situ **soil moisture (SM)** stations from the **International Soil Moisture Network (ISMN)** are installed: Benin, Niger, and Senegal. Green dots indicate the locations of the individual **soil moisture SM** stations (four in Benin, three in Niger, and one in Senegal). Purple squares represent the 0.25° satellite grid cells used to compare satellite and model **soil moisture SM** products with the in situ observations. The red rectangle delineates the regional domain used for the gridded validation between ESA CCI **soil moisture SM** and the land surface models (CLM5.0, GLWS2.0, and WaterGAP). Photos used are adapted from (?)

179 2.2 Datasets and models

180 This section describes the hydrological and land surface models used in this study, and the observa-
 181 tional datasets employed for evaluation. Model simulations are forced with homogenized atmospheric
 182 reanalysis datasets, including GSWP3-W5E5, CRUNCEP (CRU (Climate Research unit Time Series)
 183 (?) and National Centre for Environmental Protection (NCEP) reanalysis (?), and WFDE5 ([WATCH](#)
 184 ([WATER](#) and global [CHange\(WATCH\)](#) Forcing Data [methodology applied to ERA5 reanalysis data](#)).
 185 These datasets offer more spatially sufficient information than direct measurements which are often
 186 sparse in Africa.

187 2.2.1 Hydrological models: WaterGAP and GLWS2.0

188 This study utilizes version 2.2e of the WaterGAP global hydrology model (?), which simulates daily
 189 water fluxes and storage on a 0.5° grid by solving water balance equations across ten water compart-
 190 ments. These compartments represent different storage components within the hydrological cycle,
 191 including surface water (rivers, lakes, reservoirs, wetlands), **soil moisture SM**, groundwater, snow,

glaciers, canopy water, and river floodplains. The model’s vertical water balance includes components such as the canopy, snow, and ~~soil-moisture~~[SM](#), while the lateral water balance accounts for storage in groundwater, lakes, artificial reservoirs, wetlands, and rivers. The vertical water balance is expressed in terms of water height (measured in millimeters), whereas the lateral water balance is calculated using volumetric units (in cubic meters) (?). Unlike many land surface models, WaterGAP incorporates human water use for various purposes, including irrigation, livestock, industry, domestic consumption, and cooling of thermal power plants, and is calibrated against long-term annual river discharge. A significant update in this version is the enhanced algorithm for surface and groundwater abstraction. The model employs forcing data from the homogenized GSWP3-W5E5 reanalysis dataset, which includes precipitation, temperature, long-wave radiation, and shortwave radiation (?), as well as information on the characteristics of surface water bodies (lakes, reservoirs, and wetlands), land cover, soil type, topography, and irrigated areas. Since its inception in 1996, WaterGAP has been instrumental in assessing the dynamic development of the human-water system, both historically and into the future, particularly in the context of climate change. The model has significantly improved our understanding of changes in continental water storage, with a particular emphasis on the overuse and depletion of water resources (?). GLWS 2.0, or the Global Land Water Storage dataset version 2.0, is developed by assimilating monthly ~~Total Water Storage Anomaly (TWSA)~~[TWSA](#) maps from GRACE and GRACE-FO into the WaterGAP global hydrological model. The Ensemble Kalman Filter (EnKF) (?) was used for assimilation, which is implemented through the Parallel Data Assimilation Framework (PDAF) (?). The assimilation process includes vertical disaggregation to optimally combine GRACE/GRACE-FO data with inputs from the hydrological model, resulting in ten distinct water compartments. GLWS 2.0 covers global land areas, excluding Greenland and Antarctica, with a spatial resolution of 0.5° and spans the period from 2003 to 2019, ensuring no gaps in the data. It also incorporates monthly uncertainty quantification at the grid cell level. Key improvements in GLWS 2.0 compared to its predecessor, GLWS 1.0, include the integration of the updated WaterGAP version 2.2e and minor bug fixes in the assimilation process. Comprehensive details about the development of GLWS 2.0 can be found in ?.

2.2.2 Land Surface Model: CLM5.0

In this study, we use CLM5.0, the latest version of the Community Land Model (CLM), which operates in land-only mode over the Coordinated Regional Climate Downscaling Experiment (CORDEX) Africa domain (??). This configuration uses atmospheric reanalysis datasets including GSWP3-W5E5, CRUNCEP, and WFDE5 as external forcings rather than coupling CLM5.0 with an atmospheric

model. CLM5.0 simulates key biophysical and biogeochemical processes, such as the interaction between incoming radiation and the canopy/soil, and the exchange of sensible heat, latent heat, and carbon with the atmosphere (?). Additionally, the model incorporates snow accumulation and melting, along with water and energy transport in the soil. It captures processes such as infiltration, surface runoff, deep percolation, stomatal physiology, and photosynthesis. To account for land surface variability, CLM5.0 divides each grid cell into multiple land units with unique soil or snow columns and plant functional types (PFTs), allowing for a more nuanced representation of surface heterogeneity (?). Compared to earlier versions like CLM4.5, CLM5.0 provides enhanced accuracy in simulating hydrological and ecological processes and introduces a more explicit representation of human land management, making it a powerful tool for analyzing land-atmosphere interactions and land-use impacts (?). The model relies on a comprehensive set of atmospheric forcing data, including precipitation, air temperature, shortwave and longwave radiation, specific humidity, surface air pressure, and wind speed. This data is available at different temporal resolutions: every 6 hours for CRUNCEP, every 3 hours for GSWP, and hourly for WFDE5. It operates at a high horizontal resolution of approximately 0.027° (around 3 km) with a 30-minute time step, and output data are further aggregated to a monthly scale for the analysis. ~~Soil moisture~~ SM in CLM5.0 is expressed in volumetric units (cm³/cm³) for each soil layer, structured through a 25-layer soil model extending to a depth of 42 meters. Of these, 20 layers are hydrologically and biogeochemically active, providing the simulation of vertical soil ~~moisture transport~~ water movement through the numerical solution of the Richards equation (?). This layered approach improves the model's capacity to capture the vertical distribution of water in the soil, critical for understanding root-zone hydrological processes. For further details on CLM5.0's methods for simulating processes, surface characterization, and vertical soil discretization, refer to ? and ?.

2.2.3 In-situ ~~soil moisture~~ SM data

~~Soil moisture~~ SM observations from three meso-scale sites in Benin, Niger, and Senegal (Figure 1) spanning from 2003 to 2019 were sourced from the ~~International Soil Moisture Network (ISMN)~~ ISMN (?) to evaluate the accuracy of model-simulated ~~soil moisture in both surface soil moisture (SSM) and root zone soil moisture (RZSM)~~ SM in both SSM and RZSM. Table 1 details the geographical coordinates of the ~~soil moisture~~ SM stations, land-cover types, and the depths of the available ~~soil moisture~~ SM probes. The locations of the ~~soil moisture~~ SM stations within the 0.25° satellite pixels are illustrated in Figure 1. Specifically, there are four stations in Benin (Belefoungou-top, Belefoungou-middle, Nalohou-top, and Nalohou-middle), three stations in Niger (Banizoumbou, Tondikiboro, and

Wankama), and one station in Senegal (Dahra). The stations in Benin and Niger are part of the AMMA-CATCH observatory, which monitors land-atmosphere interactions in West Africa (?), while the Dahra station in Senegal belongs to the Dahra super-site monitoring ecosystem dynamics in the Sahel (?). Each ~~soil-moisture~~SM dataset includes hourly observations in volumetric units (cm^3/cm^3) at various depths, as outlined in Table 1. Only observations that have undergone rigorous quality control procedures and were flagged as "Good" by the ISMN, were selected for analysis (?). This ensures that the in-situ data used is of the highest reliability, having passed through stringent checks for accuracy, consistency, and completeness. Additional information regarding the instruments used and the data quality control procedures for the observations can be found in the network reports and the associated references, which are accessible at <https://ismn.geo.tuwien.ac.at>. While additional ISMN sites exist in the region (e.g., in Ghana, Nigeria, and Côte d'Ivoire), many were installed relatively recently, primarily within the framework of the Trans-African Hydro-Meteorological Observatory (TAHMO), with most deployments occurring from 2019 onwards. Consequently, these stations do not provide sufficient temporal overlap with our analysis period (2003–2019) and do not meet the temporal coverage and quality requirements for inclusion in this study. By focusing exclusively on the high-quality observations, we aim to minimize errors and uncertainties in the results, leading to more robust and credible findings.

Table 1: Geographic coordinates, stations grouped by site, probe depths, and land-cover types where the probes are installed.

Sites	Stations	Latitude	Longitude	Probes depths (cm)	Land-cover
Benin	Nalohou (top)	9.743° N	1.606° E	5, 10, 20, 40, 60, 100	Mixed crops
	Nalohou (middle)	9.745° N	1.605° E	5, 10, 20, 40, 120	Mixed crops
	Belefoungou (top)	9.790° N	1.710° E	5, 10, 20, 40, 60, 100	Forest
	Belefoungou (middle)	9.795° N	1.715° E	5, 10, 20, 40, 50, 100	Forest
Niger	Wankama	13.646° N	2.632° E	5, 10–40, 40–70, 70–100, 100–130	Millet
	Banizoumbou	13.532° N	2.660° E	5	Fallow
	Tondikiboro	13.548° N	2.696° E	5, 10–40, 40–70, 70–100, 100–130	Fallow
Senegal	DAHRA	15.403° N	15.432° W	5, 10, 30, 50, 100	shrubland

2.2.4 ESA-CCI ~~soil-moisture~~SM

The ESA-CCI is a global satellite-observed ~~soil-moisture~~ (SM-)SM dataset developed under the European Space Agency's Climate Change Initiative (CCI). The ESA-CCI SM v0.81 is the latest version data used in this study, providing daily estimates of global ~~surface soil-moisture~~SSM, covering the top 2–5 cm of soil, over a long term (1978–2022) at a spatial resolution of 0.25° . The ESA-CCI SSM v0.81 dataset is generated by merging SM data from 12 single-sensor active and 5 passive microwave

sensors. For detailed information on the key features of these active and passive microwave sensors, please refer to ?. Despite some limitations associated with the chosen merging algorithm and the quality of individual data sources, the v0.81 version has shown significant promise for assessing model performance (??). In this study, the combined product, which integrates ~~soil-moisture~~ SM retrievals from both active and passive microwave sensors, is selected, as it benefits from the strengths of both types of observations and generally outperforms products that rely solely on single-sensor input (??). The dataset is available free of charge from the ESA website and other platforms, provided in volumetric units (cm^3/cm^3) and in NetCDF format, making it accessible for long-term climate and hydrological studies. A detailed description of the ESA-CCI SSM product can be found at: <http://www.esa-soilmoisture-cci.org/node/139>.

2.2.5 Land cover

The land cover data used in this study is sourced from versions v2.0.7cds and v2.1.1 of the ~~European Space Agency's Climate Change Initiative~~ ESA CCI Land Cover (~~ESA-CCI-LCLC~~) dataset, accessed from <https://www.esa-landcover-cci.org> (ESA, 2024; last access: 31 August 2024). Version v2.0.7cds covers the period 1992–2015, while v2.1.1 provides data for 2016–2019. However, for this research, the version v2.0.7cds is used for the period 2003–2015, and v2.1.1 for 2016–2019.

The ESA CCI-LC dataset offers global annual land cover maps at a 300-meter spatial resolution in NetCDF format, spanning 1992 to 2019. These maps are produced by integrating observations from multiple satellite sensors (e.g., MERIS, SPOT-VGT, PROBA-V) using time-series analysis and supervised machine learning classification based on the Land Cover Classification System (LCCS) (?). The dataset's accuracy has been validated by several studies (??). The land cover map for this study (Figure 2) was generated by processing NetCDF files, clipping the data to the West Africa region, and calculating the dominant land cover class per pixel for the period 2007–2019, with key land cover types including cropland, tree cover, shrubland, grassland, urban areas, and bare areas.

3 Methods

The methods are organized into four main steps. The detailed procedure for retrieving ~~root-zone-soil~~ moisture RZSM from conceptual models (GLWS2.0 and WaterGap) is outlined in Section 3.1, while the approach for assessing spatial footprint is discussed in Section 3.2. Data pre-processing procedure is covered in Section 3.3, and the performance validation approach is presented in Section 3.4.




Figure 2: Land cover map over the study area.

3.1 Projecting ~~root-zone-soil-moisture~~ RZSM to specific depth

An approach based on the analytical solution of Richards' equation is used to translate the water content from the single ~~soil-moisture~~ SM reservoir in the conceptual hydrological models to different depths. Equation (1) below was applied to retrieve the ~~soil-moisture~~ (~~SM~~) SM signal at any given depth, following the procedure outlined in (?) :

$$\frac{\partial \theta}{\partial t} = D \frac{\partial^2 \theta}{\partial z^2} - k \frac{\partial \theta}{\partial z} \quad (1)$$

Where θ is the volumetric ~~soil-moisture~~ SM content (cm^3/cm^3), t is the time (month), z is the soil depth (positive downward) (cm), and D is the effective soil water diffusivity (an average value over the entire saturation range) (cm^2/month). k represents the average slope of the soil hydraulic conductivity function, which describes the relationship between unsaturated hydraulic conductivity K (cm/month) and volumetric ~~soil-moisture~~ SM θ . The soil hydraulic parameters D and K vary with soil type and are calibrated at each site according to the methodology outlined in ?. The equation (1) illustrates

320 how the ~~soil-moisture~~ SM content, represented by prescribed boundary conditions, changes over time
 321 and depth due to soil water diffusion and conductivity. The closed-form solution for determining ~~soil~~
 322 ~~moisture~~ SM at any arbitrary depth during the N^{th} time step with time intervals of ΔT (as shown
 323 in Equation (2)) is derived by utilizing ? approach to solve Equation (1). This solution is developed
 324 while following the boundary and initial conditions specified in ?. This equation (2) is given by:

$$\theta_N(Z) = \begin{cases} \exp(-0.5Z + 0.25T)F_1U(Z, T) & \text{for } N = 1 \\ \exp(-0.5Z + 0.25T) \left\{ F_1U(Z, T) + \sum_{i=2}^N (F_i - F_{i-1})U(Z, [N - i + 1]\Delta T) \right\} & \text{for } N > 1 \end{cases} \quad (2a)$$

$$U(Z, T) = -0.5 \exp(Z) (Z + T + 1) \operatorname{erfc} \left[0.5 \left(\frac{Z}{\sqrt{T}} + \sqrt{T} \right) \right] + \sqrt{\frac{T}{\pi}} \exp \left(- \left[0.5 \left(\frac{Z}{\sqrt{T}} - \sqrt{T} \right) \right]^2 \right) + 0.5 \operatorname{erfc} \left[0.5 \left(\frac{Z}{\sqrt{T}} - \sqrt{T} \right) \right] \quad (2b)$$

329 where erfc is the complementary error function, N is the total number of discrete time steps, and T ,
 330 Z and F are dimensionless representations of time t , soil depth z , and net water flux f given by:

$$\begin{aligned} Z &= \frac{z \cdot k}{D}, \\ T &= \frac{t \cdot k^2}{D}, \\ F &= \frac{(f - k \cdot \theta_{\infty})}{(k \cdot \theta_{\infty})}. \end{aligned} \quad (3)$$

332 A closed-form solution is obtained by assuming a stepwise surface flux input F defined as follows:

$$F(T) = \begin{cases} F_1 & \text{for } 0 < T < \Delta T \\ F_2 & \text{for } \Delta T < T < 2\Delta T \\ \vdots & \\ F_N & \text{for } (N - 1)\Delta T < T < N\Delta T \end{cases} \quad (4)$$

334 θ_{∞} as defined in F , represents the long-term temporal mean of relative ~~soil-moisture~~ SM at a given
 335 site. The implementation of this approach in our research assumes that the time derivative of the
 336 ~~soil-moisture~~ SM component, derived from either the GLWS 2.0 or WaterGAP models, approximates
 337 the net water flux f as shown in Eq. (3). Central differencing is employed to avoid introducing
 338 a phase lag. This approximation yields the dimensionless flux F , which is used in Eq. (2a). The
 339 model is subsequently executed as a sequential computational workflow: first, the monthly single-layer

SM time series from GLWS2.0 or WaterGAP is used to compute temporal variations and derive f ; second, f is transformed into F (Eq. 3), representing the normalized forcing driving vertical moisture redistribution; third, the analytical solution in Eq. (2) is evaluated for each grid cell, time step, and predefined soil depth to propagate surface moisture variability into deeper layers; finally, the resulting depth-resolved SM profiles are used to compute RZSM by integrating over the corresponding soil layers. The analytical formulation of Eq. (2) enables the computation of the ~~soil-moisture-SM~~ profile from GLWS 2.0 or WaterGAP at monthly intervals, eliminating the risk of “truncation error” that can arise in numerical solutions of Richards’ equation (?). Using this formulation, depth-resolved ~~soil-moisture~~ SM can be derived iteratively for any layer of interest, starting from the surface and proceeding downward through the root zone. Representative values for the soil hydraulic parameters D (diffusion coefficient) and K (slope of the hydraulic conductivity function) are initially assumed, while θ_∞ is set as the long-term average of ~~surface-soil-moisture~~SSM. These parameters can later be calibrated against in situ observations, using the observed minimum and maximum ~~soil-moisture-SM~~ to constrain physically realistic values (?). This approach provides a robust and computationally efficient method to translate single-layer ~~soil-moisture-SM~~ estimates into full root-zone profiles, facilitating comparison across models, satellite products, and in situ measurements. The ~~soil-moisture-SM~~ profile at any desired depth, aligned with the node depths of CLM5 for subsequent comparison, is calculated using Eq. (2).

3.2 Spatial footprint

Spatial footprint, or spatial representativeness, refers to the area surrounding a ~~soil-moisture-(SM)-SM~~ station within which temporal ~~soil-moisture-SM~~ dynamics closely align with those observed in nearby regions, as captured in model outputs or remote sensing data. This metric is critical for evaluating a model’s ability to accurately reflect local ~~soil-moisture-SM~~ dynamics around each station, thereby supporting robust model validation. To understand spatial patterns of ~~soil-moisture-SM~~ dynamics in the study area, an insightful approach based on the spatial representativeness is used (???). This approach quantifies the area surrounding a SM station of interest for which its temporal dynamics are representative, i.e., its spatial footprint. This area is determined by first calculating Spearman’s rank-based correlation coefficient between the time-series of the station under investigation and the surrounding pixels of the studied models and then iteratively removing the furthest pixels away from the station of interest, focusing on keeping only those pixels that have a correlation above a predefined threshold (denoted as r_{cut}). The spatial representativeness is ultimately defined by the area covered by the convex hull surrounding these stations that exhibit a correlation above r_{cut} threshold (?). A

convex hull is the smallest polygon that can enclose all the stations that meet the correlation criteria.

3.3 Pre-processing and performance validation approach of SM products

The ~~soil-moisture~~SM products used in this study vary in spatial and temporal resolution, layer depths, and units. Consequently, these datasets undergo preprocessing to standardize their specifications for effective comparison through the following steps:

3.3.1 Scaling

To ensure consistency across the different ~~soil-moisture~~SM products used in this study, a few scaling procedures are applied. First, the ~~soil-moisture~~SM products have varying spatial resolutions, such as 0.5° for WaterGAP and GLWS2.0, 0.25° for ESA CCI, and 0.2° for CLM5.0. To enable consistent analysis, the outputs from CLM5.0 and ESA CCI are aggregated to match the 0.5° spatial resolution of WaterGAP and GLWS2.0. In addition to this spatial scaling, all datasets are also aggregated to a monthly temporal resolution to align with the standard resolution of GLWS2.0. Second, ~~soil-moisture~~SM is known to exhibit significant spatial variability, which poses challenges when using single station measurements to accurately represent model simulations across an entire 0.5° ~~soil-moisture (SM)~~SM grid. Direct comparisons between grid-based simulations and point-based observations can introduce substantial errors (?). To mitigate this, the study utilizes multiple in situ measurements at each site, which are grouped within the 0.5°x0.5° model pixel resolution area, covering three sites (Benin, Niger, and Senegal). The average of all SM measurements within each site is computed to provide the best possible approximation of the ~~soil-moisture~~SM at ground level. This approach, adopted by many authors such as ?, ?, and ?, ensures a more reliable comparison. To further refine the process, model values are interpolated to the corresponding in situ points using the nearest neighbor method. Third, another challenge in the validation process is the mismatch in depths between the model-based ~~soil-moisture (SM)~~SM estimates and the SM observations. The ESA CCI captures only ~~near-surface-soil-moisture~~near-SSM, whereas CLM5.0 and in situ data provide ~~soil-moisture~~SM measurements at various depths, including those within the root zone. In contrast, WaterGAP and GLWS2.0 represent the water content within a single ~~soil-moisture~~SM reservoir spanning the entire root zone. Consequently, these products are not directly comparable in terms of magnitude (?). This issue is addressed in ~~the first subsection~~ subsection 3.3.2 (Normalization)

3.3.2 Normalization

The soil water simulated by the GLWS2.0 and WaterGAP models, expressed as water depth (in millimeters), cannot be directly compared to in-situ measurements, ESA CCI data, or CLM5.0 outputs, which are reported in volumetric fraction. For validation purposes, numerous previous studies have used a range of normalization techniques to address the inconsistencies among SM estimates derived from global hydrological models, satellite-based retrievals, and in situ observations (??). Common methods include percentile-based normalization, moving-window anomalies, and statistical rescaling to standardize datasets, improving alignment across diverse sources and enhancing the accuracy of validation analyses. In this study, we employed a percentile-based normalization method using the 2nd and 98th percentiles (Eq. 4).

$$w_t = \frac{\theta_t - \theta_{wt}}{\theta_{fc} - \theta_{wt}} \quad (4)$$

In this approach, each data point in the time series θ_t is normalized by subtracting the 2nd percentile of the entire time series from individual monthly values (θ_{wt}), followed by dividing the resulting series by the difference between the 98th (θ_{fc}) and 2nd percentiles of the original time series (θ_{wt}). θ_{wt} and θ_{fc} denote the ~~soil moisture~~ SM at the wilting point (dry conditions) and at field capacity (wet conditions), respectively. w_t commonly referred to as the Soil Wetness Index (SWI), is widely used in land surface modeling and drought analysis (???). Its primary purpose is to highlight relative moisture anomalies and drought signals, rather than to represent absolute ~~soil moisture~~ SM values controlled by soil texture. Values of w_t close to 0 indicating ~~soil moisture~~ SM conditions approaching the wilting point (dry state) and values close to 1 indicating conditions near field capacity (wet state). This method, commonly referred to as "relative wetness" by ? and applied in recent studies (e.g., ?? and references therein), is robust and less sensitive to outliers, making it particularly suitable for our analysis. By normalizing in this way, the approach eliminates information about the original scale, enabling comparisons to focus exclusively on the relative seasonal patterns within each dataset. This avoids biases introduced by absolute magnitudes or imposed statistical alignments, as seen in methods like ?, ensuring meaningful comparisons across datasets.

3.3.3 Performance validation approach for SM products

After converting ~~soil moisture~~ SM simulated in the model and derived from satellites into relative wetness, quantitative evaluations were performed on the refined datasets. The seasonality of different SSM time series is analyzed by applying a 12-month rolling boxcar filter. A boxcar filter is an unweighted moving average filter with a fixed-width temporal window, used to smooth time series

by suppressing short-term variability while preserving long-term signals. This approach smooths out short-term fluctuations, effectively isolating long-term variations with periods longer than one year defined here as the seasonal component. Subtracting this smoothed, long-term component from the original time series reveals the residual signal, which captures the episodic variations, defined as short-term ~~soil-moisture~~-SM fluctuations and event-scale anomalies (wet and dry events) occurring on timescales shorter than one year.

For the validation of ~~surface-soil-moisture (SSM)~~-SSM and the ESA CCI grid, only ~~soil-moisture~~ SM probes located at depths ≤ 5 cm were considered to ensure consistency. It was assumed that the microwave retrievals from ESA CCI represent moisture content within the top 0–5 cm of the soil column. Three performance metrics were applied in this study: the Pearson correlation coefficient (R), bias, and root-mean-square error (RMSE) for validating with the *in situ* data, as well as R and RMSE for grid-based validation using the ESA CCI ~~soil-moisture~~-SM data. R is a key statistical metric in this study, as it assesses how well the temporal variability of *in situ* and ~~soil-moisture~~-SM product time series align. It remains unaffected by differences in mean or variance, which can arise from varying soil properties or scale discrepancies between in situ data and model or satellite footprints (???).


4 Results and Discussion

4.1 Performance validation of SSM products


In this study, we employed two validation approaches. First, we validated ~~surface-soil-moisture (SSM)~~-SSM products specifically, ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 by comparing ~~model-soil~~ ~~moisture-model~~-SM with in situ measurements taken at a depth of 5 cm across different study sites in Benin, Niger, and Senegal. Our analysis emphasized the temporal correspondence between the datasets (e.g., time/phase lags, dry and wet spell event periods) rather than the magnitudes. Dry and wet spells are defined as persistent periods of anomalous ~~soil-moisture~~-SM conditions, corresponding to sequences of consecutive time steps with ~~soil-moisture~~-SM values persistently below or above a defined threshold, respectively. This definition is conceptually consistent with the classical climatological definition of spells as continuous sequences of days with rainfall below or above a given threshold. Additionally, deseasonalized and episodic time series were compared with the corresponding in situ time series to further assess consistency and alignment. Second, we compared the GLWS2.0, WaterGAP, and CLM5 models against the ESA CCI gridded ~~soil-moisture~~-SM data.

4.1.1 In situ-based data validation of different SSM products


The normalized ~~soil moisture~~ SM time series from ESA CCI, GLWS2.0, WaterGAP, and CLM5.0, along with the in situ time series at each study site are overlaid and presented in Figure 3.



(a) Niger Study Site



(b) Benin Study Site



(c) Senegal Study Site


Figure 3: Normalized ~~soil moisture~~ SM time series for in situ measurements, ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 at three distinct study sites at a depth of 5 cm.

The ESA CCI, GLWS2.0 and CLM5.0 SM products effectively replicate the seasonal dynamics of


monthly ~~surface soil moisture (SSM)~~ SSM observed in ground-based measurements across different sites, demonstrating their robustness in capturing the effects of varying land cover and climate conditions in the region. The seasonal changes in ~~soil moisture~~ SM at a depth of 5 cm show marked variability, likely driven by the influence of the monsoon across the West African region. In situ observations in Niger and Senegal reveal a typical Sahelian seasonal cycle, marked by a distinct wet season from June to October, while the Benin site, located in the Sudanian climate zone, experiences a rainy season extending from April to October, along with a longer monsoon period. More broadly, the dominant role of monsoon systems in controlling seasonal ~~soil moisture~~ SM dynamics is not unique to West Africa. Similar large-scale monsoon-driven ~~soil moisture~~ SM cycles have been documented in other monsoon regions, such as the Tibetan Plateau, where ~~soil moisture~~ SM variability is strongly shaped by the South Asian summer monsoon (?). This underscores the importance of regional monsoon systems in determining ~~soil moisture~~ SM fluctuations across distinct climatic zones. The statistics from the normalized SSM signals, as presented in Table 2, highlight the performance of various datasets in terms of synchronization and correlation with in-situ measurements. In Benin, ESA, GLWS2.0, and WaterGap generally exhibit strong synchronization, with time lags close to 0 months. These datasets also demonstrate high correlations (ESA: 0.894, WaterGap: 0.856, GLWS2.0: 0.752). In Senegal, these same datasets show minimal time lag (ranging from 0 to 1 month), though the correlations are somewhat weaker (ESA: 0.646, WaterGap: 0.637, GLWS2.0: 0.574). In Niger, ESA remains well-aligned with in-situ measurements (0-month lag and a 0.864 correlation), while both GLWS2.0 and WaterGap experience slight delays (1-month lag), accompanied by weaker correlations (0.655 and 0.515, respectively).

4.1.2 SSM seasonality


To better highlight events in the time series, a deseasonalization method is applied. The seasonal maps derived from this process are presented in Fig. 4.



(a) Niger study site



(b) Benin study site



(c) Senegal study site

Figure 4: Episodic and deseasonalized soil-moisture-SM time series for the three study sites: (a) Niger, (b) Benin, and (c) Senegal.

The analysis of episodic and deseasonalized time series for soil-moisture-SM across Senegal, Niger, and Benin reveals distinct regional patterns influenced by the West African Monsoon (WAM). In the Sahelian zones of Niger and Senegal, typical seasonal cycles show sharp increases in soil-moisture-SM during the monsoon (June to October) and declines in the dry season (November to May). Deseasonalized data highlight anomalies, such as unseasonal rainfall or dry spells, which suggest shifts in the timing or intensity of the monsoon and potential impacts of climate variability, such as droughts or flooding. In Benin, located in the Sudanian climate zone, a longer rainy season (April to October) leads to a more extended period of increased soil-moisture-SM, with anomalous events like flooding or unexpected dry spells reflecting changes in rainfall distribution. Overall, deseasonalizing the time series reveals how soil-moisture-SM is sensitive to shifts in the monsoon, with delayed rains in the Sahel potentially causing droughts, while early or heavy rains in Benin could lead to flooding or soil saturation. The correlation between the seasonality of the ESA CCI product and the in situ data is generally high, exceeding 0.9 for episodic time series across all sites. At the Benin site, ESA CCI demonstrates the strongest performance, followed by the assimilation-based model GLWS2.0. Mean-

while, at the Niger and Senegal sites, WaterGAP also performs well, ranking closely behind ESA CCI.

Table 2: Statistics from the normalized SSM time series.

	Benin				Niger				Senegal			
	Lag (month)	R	bias	RMSE	Lag (month)	R	bias	RMSE	Lag (month)	R	bias	RMSE
Insitu – ESA	0	0.90	-0.12	0.18	0	0.86	-0.07	0.16	0	0.65	0.05	0.21
Insitu – GLWS2.0	0	0.75	0.16	0.28	1	0.66	0.12	0.27	0	0.57	0.22	0.36
Insitu – WaterGap	0	0.86	0.06	0.21	1	0.52	0.16	0.31	1	0.64	0.24	0.35
Insitu – CLM5	6	0.62	-0.17	0.55	-18	0.50	-0.18	0.46	17	0.44	-0.11	0.50

Table 3: Statistics of the episodic and deseasonalized components of the normalized ~~soil-moisture~~ SM time series across the Senegal, Niger, and Benin study sites.

	Benin			Niger			Senegal		
	R	bias	RMSE	R	bias	RMSE	R	bias	RMSE
Episodic time series									
Insitu – ESA	0.93	-0.00	0.13	0.93	-0.01	0.12	0.90	-0.03	0.16
Insitu – GLWS2.0	0.76	0.00	0.23	0.62	0.00	0.23	0.66	-0.01	0.21
Insitu – WaterGap	0.86	0.00	0.20	0.51	0.00	0.25	0.70	-0.01	0.20
Insitu – CLM5	0.71	-0.01	0.24	0.50	-0.00	0.22	0.34	-0.02	0.26
Deseasonalized time series									
Insitu – ESA	0.76	-0.12	0.12	-0.18	-0.07	0.11	0.34	0.08	0.18
Insitu – GLWS2.0	0.40	0.16	0.17	0.04	0.12	0.14	-0.47	0.24	0.31
Insitu – WaterGap	0.58	0.06	0.08	-0.28	0.16	0.18	-0.41	0.25	0.31
Insitu – CLM5	0.09	-0.17	0.21	0.05	-0.17	0.18	-0.35	-0.17	0.28

4.1.3 ESA CCI grid based evaluation of the different models

To evaluate the accuracy and performance of the global models used in this study (GLWS2.0, WaterGAP, and CLM5.0) a grid-based comparison was conducted using remotely sensed ~~surface-soil-moisture~~ (SSM)-SM data from ESA CCI. Figure 5 displays the Pearson correlation coefficient (R) and root-mean-square error (RMSE) metrics, providing insights into how well the GLWS2.0, WaterGAP, and CLM5.0 models perform relative to the ESA CCI grid. In this figure, gaps in the data particularly noticeable in forested and densely vegetated areas (shown as white regions near the coastline) high-

light areas where ~~soil-moisture~~SM estimates are challenging. This data masking in the ESA CCI SSM datasets occurs because microwave-based observations typically exclude areas with moderate to dense vegetation due to the strong attenuation of soil signals by the vegetation canopy (?).

The figure also reveals that the patterns in RMSE and correlation metrics vary with land cover type, demonstrating that land cover exerts a clear influence on the performance of these models. GLWS2.0 shows a notably lower RMSE than the other models across the region and exhibits strong spatial consistency across various land cover types. The lowest RMSE values for GLWS2.0 are observed in the semi-arid Sahel regions, likely due to its effective assimilation process that better captures ~~soil~~moistureSM dynamics in these environments.





Figure 5: Pearson correlation coefficient (R) and RMSE metrics illustrating the performance of GLWS2.0, WaterGAP, and CLM5 models relative to the ESA CCI grid.

The figure underscores the complexities of ~~soil-moisture~~SM modeling across the diverse climates and land covers of the West African region, where model performance and error metrics vary sig-


nificantly. While GLWS2.0 demonstrates spatial consistency across different land-cover types with relatively low RMSE values, this consistency is not as evident in other models. RMSE values are generally lower in the northern part of West Africa but higher in the southern regions. This variation may be attributed to the sporadic nature of ~~soil-moisture~~ SM in these areas, where conditions are highly dependent on occasional rainfall events. In these regions, ~~soil-moisture~~ SM levels can shift rapidly, with dry conditions prevailing and only brief, intense increases in moisture following rainfall. Both the satellite data (ESA CCI) and the hydrological models encounter challenges in accurately capturing these short-lived, extreme moisture events. However, because arid and semi-arid regions experience prolonged dry spells, the overall RMSE between model predictions and satellite observations remains relatively low, despite the missed short-term fluctuations. This pattern underscores the inherent challenges of modeling ~~soil-moisture~~ SM in arid and semi-arid regions, where infrequent but intense moisture changes add complexity to the dynamics. Additionally, the figures show that GLWS2.0 demonstrates a stronger correlation with ESA CCI data compared to the WaterGAP model, which does not exhibit a clear dependency on land-cover types. Although the correlation between ESA CCI and CLM5.0 appears to be the strongest overall, the RMSE values for GLWS2.0 are slightly more favorable in arid and semi-arid regions. This correlation is generally higher in southern West Africa and weaker in the north, where low ~~soil-moisture~~ SM values make it difficult to discern clear patterns. In the southern regions, more frequent and consistent rainfall leads to stable ~~soil-moisture~~ SM levels, allowing moisture dynamics to be easier to predict. Consequently, the spatial correlation pattern between GLWS2.0 and ESA CCI ~~soil-moisture~~ SM data reveals a clear latitudinal gradient aligned with annual rainfall and climate zones. This pattern is particularly evident in semi-arid regions that serve as transition zones between wet and dry climates, where the highest correlation values are observed (i.e. area between 9.5°N and 15°N). Such transitional regions, including the Indian subcontinent, the North American Great Plains, and southeastern Brazil, also exhibit increased correlation values due to pronounced seasonal and inter-annual variability in ~~soil-moisture-(SM)-~~ SM measurements (?). Similar findings were reported by ? over West Africa, where an evaluation of spatial correlations between GRACE-based assimilation, SSM simulations and satellite-derived SSM products (ASCAT, SMOS, and SMAP) indicated strong correlation patterns in transitional zones. This consistency suggests that SM measurements in these regions are particularly sensitive to seasonal dynamics and the distinct cycles of wet and dry phases. Moreover, GLWS2.0 effectively captures the general pattern of ~~soil~~ moisture SM in the south, where soils retain moisture better and vegetation moderates fluctuations.

4.2 Spatial Representativeness


Figure 6 illustrates the Spatial Representativeness (SR) of ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 using a cutoff Spearman correlation of 0.6, which effectively highlights how well each ~~soil moisture~~ SM product reflects observed ~~soil moisture~~ SM dynamics at each study site. This similarity threshold distinguishes between datasets that demonstrate moderate to strong agreement with in situ observations, categorizing them as performing well (correlation > 0.6) or less effectively (correlation ≤ 0.6) in capturing real-world variations. Notably, only ESA CCI and GLWS2.0 exhibit Spearman correlations greater than 0.6 with in situ data, indicating that these datasets accurately capture the spatial and temporal dynamics of ~~soil moisture~~ SM within their respective spatial footprints. This alignment suggests that their ~~soil moisture~~ SM estimates correlate quite well with the observed time series. Furthermore, the efficacy of the GLWS2.0 assimilation process likely enhances its ability to reflect real-world ~~soil moisture~~ SM conditions, particularly in arid and semi-arid regions characterized by sporadic moisture changes. In contrast, the lower correlations observed for WaterGAP and CLM5.0 point to limitations in these models' capacity to represent ~~soil moisture~~ SM variability at finer spatial scales. Overall, these findings underscore the importance of integrating observational data to enhance model accuracy in representing high-resolution ~~soil moisture~~ SM dynamics, suggesting that ESA CCI and GLWS2.0 are better suited for regional applications requiring high spatial sensitivity. Although the SR approach has not previously been applied to the satellite/model SM products examined in this study, it has proven effective for validating other SM products across various regions worldwide. For example, in the Little Washita watershed in the United States and the Yanco area in Australia, spatial representativeness has been used to explore the connections between SM spatial scales and timescales within the 50 km satellite footprint of SMOS, AMSR2, and ECMWF SM products (?). These authors demonstrate that the spatial representativeness of ~~surface soil moisture~~ SSM increases with longer timescales, but with greater variability in these regions. ? showcased the robustness and effectiveness of this approach for selecting appropriate ~~soil moisture~~ SM products. By applying it to analyze the temporal dynamics of absolute ~~soil moisture~~ SM across North America, they compared in situ observations with the European Space Agency's ECV-SM and ERA-Land datasets. ? demonstrating the robustness of this parameter-free method in climatology to quantify the spatial footprint of weather stations across Europe. Their findings show that temperature data generally exhibit greater representativeness than precipitation, with significant seasonal changes influenced by atmospheric circulation patterns, particularly in boreal winter.



(a) Niger site



(b) Benin site



(c) Senegal site

Figure 6: Spatial footprint of ~~soil moisture~~ SM products represented by a blue circle with a radius of 5° around each site, indicated in red. The convex hulls in light and dark blue represent the areas for which ESA CCI and GLWS2.0 exceed the specified cutoff threshold of 0.6, respectively.

4.3 Retrieving the ~~root-zone soil moisture~~ RZSM from GLWS2.0 and WaterGap and validating its dynamics using in-situ and CLM5

An analytical solution to Richards' equation is used to convert water content from GLWS2.0 and WaterGAP to different depths, enabling comparison of model-derived ~~root-zone soil moisture~~ RZSM with in-situ observations across depths (Figures 7 to 9). At shallow depth (5 cm), GLWS2.0, which incorporates GRACE/FO data, demonstrates superior performance than WaterGAP and yields results comparable to those of CLM5.0 in capturing the monthly ~~root-zone soil moisture (RZSM)~~ RZSM dynamics at the Niger and Benin sites (Figure 7). However, the temporal dynamics at the Senegal site are less consistent, particularly during the initial two years (2004–2006) of the in situ observation system. This discrepancy can be partially attributed to the Senegal site being represented by a single station, unlike Niger and Benin, which have at least three stations at different locations. The limited data from a single station poses challenges in accurately representing model simulations over a broader 0.5° ~~soil moisture~~ SM grid study (?). Furthermore, the WaterGAP model struggles to capture long-term (2006–2018) seasonal dynamics at all sites as recorded by in situ sensors. At this depth, a comparison of model performances in capturing seasonal ~~soil moisture~~ SM dynamics shows that the ESA CCI model aligns most closely with in situ measurements. ESA CCI achieves the lowest RMSE and the highest R^2 values across all study sites: Benin (RMSE = 0.19, R^2 = 0.71), Niger (RMSE = 0.17, R^2 = 0.66), and Senegal (RMSE = 0.21, R^2 = 0.53). The GLWS2.0 assimilation-based model ranks second in accuracy, with corresponding values in Benin (RMSE = 0.33, R^2 = 0.22), Niger (RMSE = 0.32, R^2 = -0.03), and Senegal (RMSE = 0.31, R^2 = 0.40). CLM5.0 provides performance metrics similar to those of GLWS2.0, with values in Benin (RMSE = 0.34, R^2 = 0.24), Niger (RMSE = 0.34, R^2 = -0.35), and Senegal (RMSE = 0.386, R^2 = 0.26). These results indicate that ESA CCI demonstrates the highest accuracy in reflecting seasonal ~~soil moisture~~ SM patterns, while GLWS2.0 and CLM5.0 show comparable but lower precision in fitting observed in situ measurements across all sites.

Extending the analysis to deeper layers (10, 40, and 100 cm) as illustrated in ~~figures~~ Figures 8, 9, and 10, which represent RZSM at these depths, both GLWS2.0 and CLM5.0 capture reasonably the seasonal dynamics of ~~soil moisture~~ SM at the Niger and Benin sites. These models show good alignment with in situ measurements across different depths, demonstrating their ability to reflect seasonal moisture changes as recorded by the local sensors. However, the WaterGAP model does not reflect this seasonality effectively. At the Senegal site, these seasonal patterns are notably absent, likely due to previously mentioned issues with the limited data from a single station, which may not fully represent local ~~soil moisture~~ SM variability at the model's grid scale (0.5° resolution). Furthermore,

seasonal consistency in GLWS2.0 projections across depths is less reliable at the Senegal site, situated near the coastline. This discrepancy could stem from coastal regions' unique characteristics, such as tidal influences and potential signal interference from the nearby ocean, which may impact the accuracy of GRACE/-FO-based observations used in the model's assimilation process. In comparing RZSM estimates retrieved from GLWS2.0 and WaterGAP to in situ observations and the physically based model CLM5.0 at depths of 10 cm and 40 cm, CLM5.0 demonstrates slightly better performance. This is reflected in CLM5.0's smaller RMSE and higher R^2 values across the study sites, indicating a closer alignment with observed moisture dynamics. However, at a depth of 100 cm, GLWS2.0 performs marginally better than CLM5.0, with slightly lower RMSE and higher R^2 metrics, suggesting a potential advantage of GLWS2.0 in capturing RZSM dynamics at this depth. Overall, while GLWS2.0 exhibits solid performance, particularly in comparison to WaterGAP, WaterGAP's RZSM estimates show more significant discrepancies from both in situ measurements and the results provided by CLM5.0, especially at all depths observed. In addition, CLM5.0 performed quite poor at ~~5cm~~5 cm, but relatively good at greater depths. This might be related to inclusion of measurement data by ESA CCI at 5 cm~~+~~, while CLM5.0 did not have data assimilation at this depth. The influence of 5 cm ~~soil-moisture~~SM measurements diminishes at greater depths, while apparently the model for vertical ~~soil-moisture~~SM transport scheme that CLM5.0 uses is better than for the other models. GLWS2.0, on the other hand, benefits from GRACE-based data assimilation at depth, which likely explains its comparatively stronger performance at 100 cm.

5 Conclusions and implications

This study assessed ~~root-zone-soil-moisture (RZSM)~~RZSM dynamics across West Africa (2003–2019) using GLWS2.0, WaterGAP, CLM5.0, ESA CCI v0.81, and in-situ measurements. Overall, ESA CCI shows the strongest temporal agreement with in situ data, characterized by near-zero time lags and consistently high correlations across all regions. CLM5.0 and GLWS2.0 show moderate to good performance, whereas WaterGAP performs less reliably. A grid-based validation reveals that CLM5.0 and GLWS2.0 correlate more strongly with ESA CCI across West Africa, displaying a distinct latitudinal gradient aligned with annual rainfall and climate zones. This gradient is particularly strong in the transition zones between wet and dry climates (9.5°N to 15°N). In contrast, WaterGAP lacks this dependency on land-cover types, suggesting its limitations in capturing regional SM variations. To evaluate how well each product captures spatially coherent ~~soil-moisture (SM)~~SM dynamics, a Spearman correlation threshold of 0.6 was applied. The results show that ESA CCI, GLWS2.0, and CLM5.0 consistently meet this threshold around the studied sites, indicating that these models ac-

curately capture the spatial and temporal dynamics of SM within their respective spatial footprints. However, it is important to acknowledge several limitations that may affect both the in-situ and grid-based validation, as well as the spatial footprint analysis. First, the limited number of in-situ probes at each study site may not adequately capture the local variability in ~~soil moisture (SM)~~SM, especially when compared to the coarser 0.5° spatial resolution of the models. Additionally, the availability of ESA CCI data is often restricted in forested and densely vegetated regions due to the strong attenuation of microwave signals by vegetation canopies, as noted by ?. Another source of uncertainty lies in the mismatch between the spatial representativeness of point-based observations and the model grid size, which can introduce discrepancies in the validation process. Moreover, the normalization applied to the datasets, while useful for comparative purposes, may have masked true differences in SM magnitudes across products. Finally, temporal gaps in data coverage —whether in in-situ records or satellite-derived products —can affect the consistency and reliability of the validation outcomes.

The analytical solution of Richards’ equation used to translate water content from a single ~~soil moisture~~SM reservoir to various depths provides additional insights into vertical ~~soil moisture~~SM dynamics. GLWS2.0 performs better than WaterGAP at depth, largely due to GRACE/-FO data assimilation, while CLM5.0 also shows robust performance despite not assimilating GRACE/-FO data. The novel application of this depth-translation approach, based on analytical solutions of Richards’ equation, enabled the projection of water content from a single ~~soil moisture~~SM reservoir to various depths across West Africa, providing insights that go beyond traditional SM analysis in the region. This methodology not only extends the shallow vertical range typically captured by microwave satellite sensors, but it also offers a way to differentiate surface and groundwater storage variations within the GRACE/-FO data, potentially expanding our understanding of water storage dynamics in complex hydrological settings. We acknowledge that the analytical solution of the Richards’ equation represents a simplification of reality. Implementing a more refined (numerical) version, together with more detailed representations of soil types, plant root water uptake, and soil hydraulics, would actually ~~mean~~require taking steps towards assimilating GRACE data into more refined land surface hydrology models. We believe this should be explored in future studies (?). By enhancing the representation of SM across different depths, this framework can improve agricultural forecasting and deepen our understanding of water cycle interactions across diverse landscapes, especially in regions where accurate SM data are essential for sustainable water management.

679 **Competing interests**


680 Harrie-Jan Hendricks Franssen is a member of the editorial board of Hydrology and Earth System
681 Sciences. The authors declare that they have no other competing interests.

682 **Author contributions**


683 LY: Data curation, methodology, software, formal analysis, visualization, interpretation, validation,
684 writing (original draft, review and editing). JK: Conceptualization, methodology, writing (review
685 and editing), supervision. BO: Methodology, model running, writing (review and editing). HG:
686 Interpretation, writing (review and editing). HJHF: Conceptualization, supervision, writing (review
687 and editing).

688 **Acknowledgments**


689 We acknowledge funding through University of Bonn's SDG fellowship program, and from Deutsche
690 Forschungsgemeinschaft (DFG, German Research Foundation), SFB 1502/1-2022 (Projektnummer:
691 450058266). We would like to thank Dr. Morteza Sadeghi for his valuable assistance and guidance
692 during the implementation of the analytical solution of Richards' equation. We are also grateful to
693 the anonymous reviewers for their constructive comments and suggestions.



(a) Niger study site




(b) Benin study site




(c) Senegal study site

Figure 7: ~~Soil moisture~~ SM time series for in situ measurements, ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 at three distinct study sites, projected at a depth of 5 cm.


- Node depth = 10 cm



(a) Niger study site




(b) Benin study site




(c) Senegal study site

Figure 8: Soil moisture time series for in situ measurements, ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 at three distinct study sites, projected at a depth of 10 cm.

- Node depth = 40 cm




(a) Niger study site




(b) Benin study site

Figure 9: Soil moisture SM time series for in situ measurements, ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 at three distinct study sites, projected at a depth of 40 cm.


- Node depth = 100 cm



(a) Niger study site



(b) Benin study site



(c) Senegal study site

Figure 10: Soil moisture SM time series for in situ measurements, ESA CCI, GLWS2.0, WaterGAP, and CLM5.0 at three distinct study sites, projected at a depth of 100 cm.