



Recent history and future demise of Jostedalsbreen, the largest ice cap in mainland Europe

Henning Åkesson¹, Kamilla Hauknes Sjørksen², Thomas Vikhamar Schuler¹, Thorben Dunse², Liss Marie Andreassen³, Mette Kusk Gillespie², Benjamin Aubrey Robson⁴, Thomas Schellenberger¹, and Jacob Clement Yde²

¹Department of Geosciences, University of Oslo, Oslo, Norway

²Department of Civil Engineering and Environmental Sciences, Western Norway University of Applied Sciences, Sogndal, Norway

³Norwegian Water Resources and Energy Directorate (NVE), Oslo, Norway

⁴Department of Earth Science, University of Bergen, Bergen, Norway

Correspondence: Henning Åkesson (henning.akesson@geo.uio.no)

Abstract. Glaciers and ice caps worldwide are in strong decline, and models project this trend to continue with future warming, with strong environmental and socio-economic implications. The Jostedalsbreen ice cap is the largest ice cap on the European continent (458 km² in 2019) and occupies 20% of the total glacier area of mainland Norway. Here we simulate the evolution of Jostedalsbreen since 1960, and its fate in a changing climate in the 21st-century and beyond (2300). This ice cap consists of glacier units with a great diversity in shape, steepness, hypsometry, and flow speed. We employ a coupled model system with higher-order 3-d ice dynamics forced by simulated surface mass balance that fully accounts for the mass-balance elevation feedback. We find that Jostedalsbreen may lose 12–74% of its present-day volume until 2100, depending on future greenhouse gas emissions. Regardless of emission scenario, the ice cap is likely to split into three parts during the second half of the 21st century. Our results suggest that Jostedalsbreen will likely be more resilient than many smaller glaciers and ice caps in Scandinavia. However, we show that by the year 2100, the ice cap may be committed to a complete disappearance during the 22nd century, under high emissions (RCP8.5). Under medium 21st-century emissions (RCP4.5), the ice cap is bound to shrink by 90% until 2300. Further simulations indicate that substantial mass losses undergone until 2100 are irreversible. Our study demonstrates a model approach for complex ice masses with numerous outlet glaciers such as ice caps, and how tightly linked future mass loss is to future greenhouse-gas emissions. Finally, uncertainties in future climate conditions, particularly precipitation, appear to be the largest source of uncertainty in future projections of maritime ice masses like Jostedalsbreen.

1 Introduction

Melting glaciers are powerful and concrete symbols of climate change, and can raise awareness and spur climate action among people who visit them (Dannevig and Rusdal, 2023). Since the year 2000, glaciers and ice caps have contributed nearly as much to sea-level rise as the combined input from the Greenland and Antarctic ice sheets (Hugonnet et al., 2021). In addition, the response of glaciers and ice caps to global warming has widespread societal implications at regional to local scales. This



includes consequences for tourism, hydropower production, agriculture and local ecosystems as a result of changes to glacier extent and the magnitude and timing of meltwater runoff (e.g. Milner et al., 2017; Huss et al., 2017). Such implications are particularly strong at Jostedalsbreen ice cap, the largest ice mass in mainland Norway and Europe, and the focus of this study. Meltwater from this ice cap feeds several hydropower stations, and Jostedalsbreen attracts over 600,000 visitors every year (Jostedalsbreen nasjonalparkstyre, 2021), with three glacier visitor centres, a dedicated museum and several curated glacier viewpoints. The ice cap is extensively used for recreation and exploration, including skiing and glacier hiking. Visitors play an important role for the livelihood of many local settlements in the vicinity of the ice cap.

The primary objective of this study is to assess the short- and long-term fate of Jostedalsbreen under future climate change. To achieve this, we demonstrate a comprehensive modelling approach of a topographically complex ice cap, consisting of 81 connected glacier units that are exceptionally diverse in shape, steepness, hypsometry and flow speed. This setting ranks among the most challenging real-world cases outside the polar ice sheets. The wide range of geometries and flow speeds found at Jostedalsbreen are representative of glaciers found in many glacierised regions of the world. Modelling the evolution of Jostedalsbreen can be viewed as an application to a small mountain range with many connected glaciers, an intermediary step between glacier-specific studies and large-scale regional or global applications. The treatment of so many connected glaciers introduces a number of challenges, which are less prominent or absent for studies of single glaciers or idealised geometries. This includes the need for detailed historical and contemporary input data, spatially constrained model parameters embedded with advanced ice-flow physics and surface mass balance, and climate projections suitable for a mountainous region.

Detailed modelling studies of single glaciers or ice caps have improved the physical realism in glacier models, evaluated the importance of physical processes and climate forcing, and projected future evolution and associated impacts on nature and society (e.g. Oerlemans, 1997; Aðalgeirsdóttir et al., 2005; Jouvett et al., 2009; Giesen and Oerlemans, 2010; Ziemen et al., 2016; Ekblom Johansson et al., 2022). Therefore, the rationale for in-depth studies of individual glaciers and ice caps remains strong (Zekollari et al., 2022). Such studies often resolve ice dynamics in two dimensions with simplified physics (e.g. Meur and Vincent, 2003; Giesen and Oerlemans, 2010; Ziemen et al., 2016; Åkesson et al., 2017; Schmidt et al., 2020). This is usually viable for simple glacier geometries, slow-flowing ice, and long time scales (Leysinger Vieli and Gudmundsson, 2004; Meur et al., 2004; Adhikari and Marshall, 2012). In contrast, ice caps may consist of numerous connected glaciers with variable geometry and dynamics, controlled to some extent by the underlying topography, often with fast-flowing outlet glaciers. This complex and dynamic setting may require 3-d modelling efforts with higher-order physics (Adhikari and Marshall, 2012; Zekollari et al., 2017), which only occurs as exceptions in the literature (e.g. Adhikari and Marshall, 2012; Gilbert et al., 2016; Zekollari et al., 2017).

While ice caps present challenges for detailed studies, these issues are exacerbated in regional or global glacier-evolution models. In such models, interactions between ice dynamics, geometry evolution, and glacier surface mass balance are typically represented in even simpler terms (Huss et al., 2012; Huss and Hock, 2015; Maussion et al., 2019; Rounce et al., 2023). In a warming climate, this means that the positive feedback between ice thinning and stronger surface melt at increasingly lower surface elevations, and associated dynamic adjustments, is not necessarily well captured. This *mass-balance elevation feedback* (Harrison et al., 2001) is particularly important for ice caps (Åkesson et al., 2017), where minor changes in elevation in their flat



interior regions affect the surface mass balance over large areas. Moreover, large-scale models generally simulate each glacier unit individually, often using simple 1-d flowline dynamics (MauSSION et al., 2019; Rounce et al., 2023). Individual treatment of glacier units has significant shortcomings when applied to ice-cap evolution over timescales of decades to centuries. As a result of strongly altered geometries, ice caps may split up into separate glacier units, ice divides may migrate, or the underlying topography can emerge in the middle of a glacier unit due to ice thinning. Such changes will heavily alter the flow regime, sometimes in the form of ‘ice piracy’ (Huiban et al., 2024), and will, in other situations, render some ice dynamically dead. These geometric changes are likely to occur in many glacierised areas of the world in the future. Here, the glacier-by-glacier approach will suffer, and the width-averaged 1d-flowline dynamics will no longer be valid. As such, large-scale glacier evolution models struggle to represent some of the most dynamic ice masses on the planet outside of the polar ice sheets (Millan et al., 2022). More detailed knowledge of the evolution and dynamics of ice caps, and how to best represent these changes in models, is therefore urgently needed (Zekollari et al., 2022). There is also a need to bridge the two end-members of glacier-scale simulations and regional-to-global-scale approaches.

A major challenge when simulating glacier dynamics is how to account for friction between ice and the underlying bedrock. Basal friction strongly influences glacier flow speeds (e.g. Iken, 1981), but is difficult to observe directly. Subglacial bulk properties are therefore usually inferred from surface velocities in the form of a spatially variable basal friction parameter (Morlighem et al., 2010; Gillet-Chaulet et al., 2012). This approach is a cornerstone in ice-sheet projections, supported by remote sensing datasets of ice-sheet wide surface speeds. In contrast, modelling efforts for glaciers and ice caps have not been able to fully take advantage of these spatially continuous velocity datasets. This is because the spatial resolution and/or temporal coverage of remote-sensing products often have been too poor to properly resolve the details of ice flow of glaciers and ice caps. Therefore, glacier modelling studies have often imposed some ad-hoc relationship for the friction parameter (Zekollari et al., 2022). The advent of detailed datasets of velocity for every glacier in the world (Friedl et al., 2021; Millan et al., 2022) presents new opportunities for friction inversions in detailed modelling studies. These datasets can however not be uncritically applied ‘off-the-shelf’, especially not for complex ice masses such as ice caps. For example, the accuracy of the underlying method may be smaller than the actual ice movement, or the data may represent snapshots or subsets of shorter periods, and hence not representative for annual mean velocities.

Another common challenge for modelling studies is poorly known ice thickness. The vast majority of Earth’s over 200,000 glaciers do not have measurements of ice thickness (Welty et al., 2020). This means that model applications must revert to global and regional thickness products (Farinotti et al., 2019; Millan et al., 2022; Frank and Pelt, 2024), which may not be accurate for detailed applications (Gillespie et al., 2024). Some of these thickness models struggle to represent connected ice complexes, because each glacier unit is treated individually. This usually leads to issues with thickness along ice divides. Poor representation of the subglacial topography may bias modelled glacier dynamics, which makes future glacier projections and associated local impacts uncertain at best.

A key conceptual and practical challenge is that the climate system is inert. Glaciers and ice caps are excellent examples of this. It can take years to decades for a glacier to respond significantly to a change in climate (e.g. Jóhannesson et al., 1989), and centuries to millennia before ice mass loss and retreat is manifested in a new steady-state (e.g. Marshall, 2005). To simulate the



historical evolution of a glacier, however, a realistic starting point is essential (Aschwanden et al., 2013). A biased initial model state will strongly affect historical and future model evolution (Aðalgeirsdóttir et al., 2014), yet obtaining this initial state with the correct climatic and dynamic memory imprinted is notoriously difficult. For example, the input data and climate forcing may be lacking, patchy or uncertain, or the model physics may be incomplete. These challenges are hard enough to resolve for a single glacier, and becomes increasingly difficult for the great number of glaciers involved in comprehensive applications like the one presented here.

In terms of future change, climate science has been largely preoccupied with the 21st century, not least reflected and fuelled by the landmark establishment of the Paris agreement in 2015 (Meinshausen et al., 2020), with the goal of limiting global warming to 1.5–2 °C above pre-industrial temperatures. Meeting the Paris goals is, at the time of writing, still technically feasible but extremely challenging, while the stakes are high for both nature and people worldwide (United Nations Environment Programme, 2024). To understand the full implications of current and near-future climate change, we however also need to consider future inertia and longer timescales. The concept of ‘committed glacier mass loss’ is around in the literature (e.g. Goldberg et al., 2015), sometimes referred to as mass loss ‘in the pipeline’, and yet deserves to be assessed in greater detail. Committed ice mass loss in a glaciological context refers to what will happen to a glacier with no further emissions, that is, a continuation of recent climate forcing into the future. This is a pertinent concept as we are heading towards the post-Paris agreement world beyond year 2100, where carbon levels in the atmosphere and global temperature rise may stabilise, or eventually even be reversed.

Some studies consider the fate of glaciers and ice caps beyond the year 2100 (e.g. Schmidt et al., 2020), but these are exceptions. There is certainly a need to consider these time scales in the glaciological context and in climate science at large (Meinshausen et al., 2020). In doing so, local and global policymakers can be informed about expected ice mass loss or glacier retreat, now and in the future.

In this study, we make progress from the modelling of individual glacier units towards detailed modelling studies of entire regions. For Jostedalsbreen, previous studies have focused on the evolution of single outlet glaciers (Nigardsbreen and Briksdalsbreen; Fig. 1) using 1-d flowline models (Oerlemans, 1997; Laumann and Nesje, 2009), or surface mass balance modelling (Jóhannesson et al., 1995; Sjurset et al., 2025). However, no study has previously attempted to simulate the historical or future evolution of the entire ice cap. We employ a coupled model system with 3-d ice dynamics, and surface mass balance simulated using a temperature-index model with spatially constrained parameters calibrated with a Bayesian approach. A newly acquired detailed dataset on ice thickness and bed topography (Gillespie et al., 2024) allows for realistic 3-d simulations of historical and future changes of the complex ice cap.

First, we simulate historical ice-cap changes since the 1960s. These simulations generate a plausible initial state for the future projections, and is used to validate our model setup against observations of ice thickness, ice margins and ice velocities. We then launch coupled simulations to the year 2100, using two scenarios of future greenhouse-gas emissions. Finally, we assess committed mass loss towards year 2300, the potential for ice-cap recovery after year 2100, as well as the possibility for glacier regrowth from scratch in the current and future climates.



125 2 Study area

Jostedalsbreen (61° 40'N; 7° 00'E) is a northeast-southwest-oriented ice cap located c. 100 km inland from the west coast of Norway (Fig. 1). It is the largest coherent ice mass in mainland Europe with an area of 458 km² in 2019 (Andreassen et al., 2022), covering an elevation range between 381 and c. 2001 m a.s.l. (Andreassen et al., 2022; Kjølmoen et al., 2024). The maximum ice thickness is about 630 m, and the ice volume is calculated to 70.6 ± 10.2 km³ (in 2018-2023; Gillespie et al., 2024). The ice cap is divided into 81 glacier units with great diversity in size, shape, steepness, and orientation (Fig. 1b; Andreassen et al., 2022). The larger glacier units are outlet glaciers terminating in glacial valleys. Glacier surface catchments are mainly determined by the bed and surrounding topography, although some discrepancies exist (Gillespie et al., 2024). Ice velocities reach up to a few hundred meters per year for the fastest outlet glaciers Nigardsbreen and Tunsbergdalsbreen, while most outlet glaciers typically move a few tens of meters annually (Wangensteen et al., 2006; Nagy and Andreassen, 2019).

135 Radar measurements suggest that the ice is predominantly temperate, but with isolated patches of cold ice (Gillespie et al., 2024). The ice cap is situated directly on Proterozoic gneiss bedrock belonging to the Western Gneiss Region of Norway (Hacker et al., 2010), except for the lower part of the outlet glacier Austerdalsbreen, which likely rests on unconsolidated glaciofluvial and till sediments (Seier et al., 2024).

Since the maximum extent of Jostedalsbreen during the Little Ice Age (c. 1740 to 1860 Gjerde et al., 2023), the ice cap has lost about 20% in area and volume (Carrivick et al., 2022). This recession has occurred progressively with decadal-scale interruptions of glacier advances or stillstands, as evidenced by sequences of moraine ridges deposited in front of most outlet glaciers (e.g. Erikstad and Sollid, 1986; Bickerton and Matthews, 1993). The latest advance of Jostedalsbreen's outlet glaciers occurred in the 1990s to early 2000s, and was a response to high winter snowfall in the preceding years (Andreassen et al., 2005; Nesje and Matthews, 2012). This illustrates that large-scale atmospheric circulation patterns, such as the North Atlantic Oscillation, influence the surface mass balance of Jostedalsbreen (Nesje et al., 2000). The current climate is characterized by mean annual air temperatures between -3 and +5°C and precipitation amounts between c. 1000 and c. 2000 mm a⁻¹, where most precipitation falls in the western part of the ice cap (Carrivick et al., 2022; Sjørnsen et al., 2025).

Jostedalsbreen has been extensively studied for more than a century. Long-term records of annual glacier front variation exist for six outlet glaciers (Austerdalsbreen, Brenndalsbreen, Briksdalsbreen [stopped in 2015], Fåbergstølsbreen, Nigardsbreen, Stigaholtbreen) dating back to 1899 (Andreassen and Elvehøy, 2021). Surface mass balance has been monitored for several outlet glaciers, but only the two records at Nigardsbreen (back to 1962) and Austdalsbreen (back to 1988) are maintained today (Andreassen et al., 2020; Kjølmoen et al., 2024).

3 Glacier input data

3.1 Glacier geometry

155 To simulate the evolution of Jostedalsbreen, we need historical and contemporary geometries of the ice cap. Ice-surface topography of 10 m resolution for the present day (2020 Digital Terrain Model; DTM) is used in calibration of the ice-flow




Figure 1. Ice-surface topography of Jostedalsgreen ice cap. Key outlet glaciers are named. Observed ice-cap extents in 1966 and 2019 are shown (Andreassen et al., 2023), as well as individual glacier catchments (blue). The inset map shows the location of Jostedalsgreen in southern Norway. Coordinates in UTM 33N, datum ETRS89 on main map and in geographical coordinates on inset.



model dynamics, as well as a final target in historical simulations (Sections 5.1 and 5.2). The source of the surface DTM on Jostedalsbreen is a lidar survey from 2020, except for the lower tongue of Tunsbergdalsbreen where the survey year is 2017 (Andreassen et al., 2023). Ice thickness has been extensively measured at Jostedalsbreen over the years 2018–2023 using ground- and helicopter-based ice radar. A distributed gridded dataset of ice thickness has been produced from these point measurements, by applying an ice-thickness model based on the inversion of surface topography (Gillespie et al., 2024). We hereafter refer to this dataset as the *observation-based* ice thickness. To derive the subglacial topography of Jostedalsbreen, this observation-based gridded ice thickness (Gillespie et al., 2024) was subtracted from the present-day ice-surface DTM. To produce a seamless bed topography map of the ice cap and nearby ice-free areas, we used the software QGIS to merge the subglacial topography with a present-day 10 m DTM of areas without glacier cover.

Meanwhile, another DTM is needed as a starting point for the historical simulations 1960–2020 outlined in Sect. 5.2. Andreassen et al. (2023) presented a DTM for 1966 based on photogrammetric reconstruction of historical aerial photographs. However, this DTM does not cover southern Jostedalsbreen. To produce a DTM for 1966 covering the entire ice cap, we extended the DTM from Andreassen et al. (2023) using another 10 m DTM produced by the company Hexagon. This DTM is based on the same 1966 aerial photographs and has a greater spatial extent, but is more noisy. The resulting DTM covered the entire ice cap, but contained data gaps over steep terrain and areas with low image contrast. We therefore georeferenced the N50 1966 topographic map from the Norwegian Mapping Authority (map sheet 1318-2; Paul et al., 2011), manually digitised the 20 m interval contour lines over the data voids, and interpolated the elevation using the TopotoRaster function within ArcGIS Pro. This interpolation tool is based on the ANUDEM algorithm, designed to produce a hydrologically accurate DTM that preserves ridgelines and stream networks (Hutchinson et al., 2011). These interpolated elevations were then used to fill the voids in the DTM. In the southern part of the ice cap, the resulting DTM had some artefacts at the boundaries between the high-resolution DTM from Andreassen et al. (2023) and the 1966 topographic contour map. These were smoothed using a 5x5 low-pass filter. The remaining traces of these artefacts quickly dissipated during the spinup simulation of the ice cap (Sect. 5.1). Between 1966 and 2020, there is no data of ice-surface topography available that covers the entire ice cap.

3.2 Ice velocity

To construct a representative present-day velocity dataset for calibration and validation of the ice-flow model (Sect. 5.1), we use a global velocity product as a baseline (Millan et al., 2022). Upon close inspection of this product, which is based on optical and synthetic aperture radar (SAR) imagery from 2017–2018, we discovered several artifacts. Mainly, we found spurious velocity magnitudes for flat areas high on the ice-cap plateau where image contrast or amplitude patterns are typically low. For example, ice along the ice divide of Nigardsbreen supposedly flows more than 100 m a^{-1} , while in reality, surface velocities near ice divides are typically limited to a few meters per year. To mediate unrealistic velocities close to ice divides, velocity maps were derived using InSAR from European Remote Sensing satellite (ERS) from 1996. Velocity maps with partial coverage of Jostedalsbreen were acquired by differential InSAR performed on three ERS-1/ERS-2 tandem pairs. The ascending pairs were acquired on 20/21 January 1996 and 11/12 February 1996, the descending on 22/23 March 1996. From those pairs, we derived line-of-sight velocities, which were then combined to a 2d-velocity map. The differential InSAR processing was performed



using the GAMMA Remote Sensing software. The 1996-velocity map was used to correct the global data from Millan et al. (2022) in areas with surface slopes lower than 0.08 and/or above 1775 m a.s.l. Judging from velocity data from complementary sources (Table A1), ice flow at the ice divides displays little year-to-year variability. Moreover, elevation changes at the ice-cap plateau since 1966 have been very small (less than ± 5 m; Andreassen et al., 2023), which suggests a negligible velocity change around the ice divides since 1996. We therefore judge our composite velocity dataset to be robustly constructed.

Overall, the global dataset from Millan et al. (2022) appears to be more representative for fast-flowing regions at lower elevations of Jostedalsbreen, where it is in close agreement with other velocity datasets that cover parts of the same time period (e.g. Nagy and Andreassen, 2019, see Table A1). Velocities from the global dataset are c. 100 to 500 m a⁻¹ for fast-flowing outlet glaciers, while differences among various data sources are on the order of 50–100 m a⁻¹ for a given glacier. However, this is not a straightforward comparison. Visual inspection of other velocity products covering parts of the 2010s (Table A1) suggests that velocities in the global dataset may be too high also in some fast-flowing areas. This is possibly because the 2017–2018 global data are biased towards high summer velocities and thus do not represent annual averages. The effect of this bias will vary from glacier to glacier, depending on the magnitude of seasonal velocity variations.

4 Model description

4.1 Ice-flow model

Ice flow and evolution of ice-cap geometry is simulated using the Ice-Sheet and Sea-level System Model (ISSM; Larour et al., 2012) considering 3-d higher order physics (Blatter, 1995; Pattyn, 2003). The model domain comprises the 1966 ice-cap extent and a c. 5 km buffer zone. This ensures an accurate representation of surface mass balance (SMB) around the margins (Sect. 4.3), allows for frontal fluctuations of outlet glaciers, and resolves interactions with nearby glaciers separate from the main ice cap. The finite-element model mesh consists of c. 87,000 horizontal elements spread over four vertical layers. To capture ice flow in steep terrain, we vary the mesh resolution from 100 to 300 m based on gradients in bedrock topography. In addition, we enforce a high resolution (100 m) for several key catchments (Fig. A1). This improves the representation of the two largest and fastest outlet glaciers Nigardsbreen and Tunsbergdalsbreen (area 41.7 and 46.2 km², respectively; Andreassen et al. (2022); ice velocities up to 300–500 m/a), and several other glaciers of particular interest to local hydropower and tourism (Austerdalsbreen and glaciers draining into Oldedalen; see Fig. 1). A time step of 0.02–0.05 years (7.3–18.25 days) is used depending on the experiment (Sect. 5), which ensures numerical stability.

Two of the 81 glacier units currently terminate in lakes (Austdalsbreen and Sygneskarsbreen in the north). Since calving is not included in the model, the lake surface is considered the bedrock topography.

Jostedalsbreen is considered a temperate ice cap (Sect. 2) and we assume a uniform ice temperature of 0 °C to compute the associated ice viscosity using an Arrhenius law (p. 75; Cuffey and Paterson, 2010). Glen's flow law is used to compute the strain rate in response to stress at any point in the glacier (p. 55; Cuffey and Paterson, 2010).



4.2 Basal friction

We use a linear viscous friction law (Budd et al., 1979) to compute basal drag τ_b as

$$\tau_b = -\alpha^2 N \mathbf{u}_b, \quad (1)$$

225 where α is a friction parameter, $\mathbf{u}_b$ is basal velocity and $N = \rho_i g H$, where ρ_i , g , H are ice density, gravitational acceleration and ice thickness, respectively. The friction parameter is allowed to vary from 10 to $400 \text{ s}^{(1/2)} \text{ m}^{(-1/2)}$. We do not account for input of surface melt and associated seasonal variations of N and $\mathbf{u}_b$.

We constrain the local values of the basal friction parameter α by inversion using an adjoint method that minimizes the misfit between the modelled u and observed ice velocities u^{obs} (e.g. Morlighem et al., 2010):

$$230 \quad \mathcal{J}(\mathbf{u}, \alpha) = \gamma_1 \frac{1}{2} \int_S \left((u_x - u_x^{obs})^2 + (u_y - u_y^{obs})^2 \right) dS + \gamma_2 \int_S \left(\log \left(\frac{\|\mathbf{u}\| + \epsilon}{\|\mathbf{u}^{obs}\| + \epsilon} \right) \right)^2 dS + \gamma_3 \frac{1}{2} \int_B \|\nabla \alpha\|^2 dB \quad (2)$$

The first and second terms are the absolute and logarithmic velocity misfits, respectively. The third term is a regularization term to prevent singularities and penalises inferred parameters to avoid overfitting, and $\gamma_1 = 2000$, $\gamma_2 = 1$, and $\gamma_3 = 3.2 \times 10^{-6}$ are respective weights for each of the terms in the cost function Eq. 2. S and B refer to the surface and bed, respectively.

Model velocities in Eq. 2 are computed using a fixed-geometry stress-balance calculation with the present-day data described
 235 in Sect. 3 (experiment ‘Calibration’ in Table 2). We use this inversion method to infer the spatially variable friction parameter α for the two largest and fastest outlet glaciers Tunsbergdalsbreen and Nigardsbreen, where we judge our mosaic velocity dataset described above to be the most representative. For the rest of the ice cap and for ice-free areas, we use a spatially variable friction parameter proportional to bedrock altitude z_b , following Åkesson et al. (2018):

$$\alpha = \beta_{max} \times \frac{\min[\max(0, z_b + z'), z_b]}{\max(z_b)}, \quad (3)$$

240 where $\beta_{max} = 400 \text{ s}^{(1/2)} \text{ m}^{(-1/2)}$ and $z' = 500 \text{ m a.s.l.}$ This simple parametrisation imposes a higher basal friction at higher elevations and more slippery conditions downglacier.

4.3 Surface mass balance model

To simulate SMB, we use the temperature-index model of Sjurset al. (2023, 2025). The model uses 1 km gridded daily mean temperature and daily total precipitation from seNorge_2018 (Lussana et al., 2019; Lussana, 2021), which is largely based on
 245 interpolation of measurements from a large network of weather stations across Norway (see Sjurset al. (2023) for details). The SMB model has been calibrated to Jostedalbreen with a Bayesian approach that uses seasonal glaciological observations to constrain accumulation and melt over the period 1962–2020, and decadal satellite-derived geodetic observations for 2000–2019 from Hugonnet et al. (2021) to derive spatially-distributed bias-corrections of temperature and precipitation for each glacier unit (Sjurset al., 2025). The SMB model domain covers Jostedalbreen and a 5 km buffer zone around the current
 250 ice-cap margins. This makes SMB available as forcing in case of glacier advance, which occurred for some outlet glaciers in



Table 1. Constants and parameter values used in this study.

Parameter	Symbol	Value	Unit
Ice density	ρ_i	917	kg m^{-3}
Gravitational acceleration	g	9.81	m s^{-2}
Rate factor	A	2.4×10^{-24}	$\text{s}^{-1} \text{Pa}^{-3}$
Basal friction parameter	α	10–400	$\text{s}^{(1/2)} \text{m}^{(-1/2)}$
Glen’s law exponent	n	3	
Mesh resolution	Δx	100–300	m
Time step	Δt	0.02–0.05	a

the 1990s and early 2000s. We employ SMB simulated using median values of the posterior distributions of model parameters from Sjrursen et al. (2025). For consistency, we propagate the glacier-specific temperature and precipitation bias-corrections outside of the glacier margins to fill the remainder of the SMB-model domain.

5 Experiment setup

255 5.1 Calibration and spinup of ice-flow model

We perform a dual ice-flow model optimisation to (i) obtain the best possible representation of ice velocities (‘Calibration’; Table 2), and (ii) obtain the most realistic initial topography for the historical simulation (‘Spinup’; Table 2). The success of (i) is measured by the root mean square error (RMSE) against present-day velocities in a fixed-geometry stress balance simulation. The performance for objective (ii) essentially depends on the accuracy of the entire suite of model choices, assumptions and input data (Sect. 3 and 4; see also Sect. 7.3). In addition, we assume that the ice cap was in steady-state in the 1960s. This assumption may be more or less accurate for different parts of the ice cap. By the 1960s, some parts of the ice cap may still have been adjusting to climatic changes from the Little Ice Age maximum in the 1700s and onwards into the 20th century. Although we deem this to have minor importance, this may introduce some localised bias, as discussed in Sect. 7.3.

To minimise both (i) the RMSE present-day velocity during the calibration, and (ii) the RMSE thickness at end-of-spinup, we varied the allowed maximum for the basal friction parameter $\beta_{max} = [200, 500] \text{ s}^{(1/2)} \text{m}^{(-1/2)}$ in the inversion. This was done for the large outlet glaciers Nigardsbreen and Tunsbergdalsbreen. Meanwhile, for areas using the friction parameterisation in Eq. 3, we tested different combinations of $\beta_{max} = [200, 500] \text{ s}^{(1/2)} \text{m}^{(-1/2)}$ and $z' = [200, 800] \text{ m a.s.l.}$ This process was done iteratively until the lowest possible RMSE for velocity (i) and thickness (ii) was found.

Once the optimal model dynamics is obtained, we perform a 250-year spinup with a fixed SMB forcing, using the mean annual SMB over the period 1960–1989 (Table 2). During this period, many outlet glacier positions remained relatively unchanged (<https://glacier.nve.no/viewer/CI/en/>; Andreassen and Elvehøy, 2021), suggesting that the ice cap was in approximate steady-state. The SMB during this period is therefore considered suitable as forcing for spinup of the 1960s ice cap. We initialise



Table 2. Overview of experiments performed in this study. Time period refers to year AD for which the experiment is run for, except for Spinup and Regrow-experiments, where no specific time period is modelled. The climate data ‘seNorge’ refers to the version seNorge_2018 (Lussana et al., 2019). Where SMB is given with years in brackets [yearX–yearY], experiments are performed using the mean annual SMB over the given period. Initial states refer to either the ice-surface data (Sect. 3.1) or modelled ice geometry at the final year of the stated simulations. Note that the ‘Calibration’ experiment is a fixed-geometry simulation and thus does not involve any climate data or SMB (Sect. 5.1).

Experiment	Time period	Climate data	SMB	Initial state
Calibration	2019	-	-	DTM2019
Spinup	250 yrs	seNorge	seNorge[1960-1989]	DTM1966
Historical	1960–2020	seNorge	Sjursen et al. (2025)	Spinup
Future4.5-ECC	2021–2100	ECEARTH/CCLM	RCP4.5	Historical
Future4.5-ECH	2021–2100	ECEARTH/HIRHAM	RCP4.5	Historical
Future4.5-CNC	2021–2100	CNRM/CCLM	RCP4.5	Historical
Future4.5-MPC	2021–2100	MPI/CCLM	RCP4.5	Historical
Future8.5-ECC	2021–2100	ECEARTH/CCLM	RCP8.5	Historical
Future8.5-ECH	2021–2100	ECEARTH/HIRHAM	RCP8.5	Historical
Future8.5-CNC	2021–2100	CNRM/CCLM	RCP8.5	Historical
Future8.5-MPC	2021–2100	MPI/CCLM	RCP8.5	Historical
CommitNow	2021–2300	seNorge	seNorge[2001–2020]	Historical
Commit4.5	2101–2300	ECEARTH/CCLM	RCP4.5[2081–2100]	Future4.5-ECC
Commit8.5	2101–2300	ECEARTH/CCLM	RCP8.5[2081–2100]	Future8.5-ECC
Reverse4.5	2101–2300	seNorge	seNorge[2001–2020]	Future4.5-ECC
Reverse8.5	2101–2300	seNorge	seNorge[2001–2020]	Future8.5-ECC
RegrowNow	1000 yrs	seNorge	seNorge[2001–2020]	no ice
Regrow4.5	1000 yrs	ECEARTH/CCLM	RCP4.5[2081–2100]	no ice
Regrow8.5	1000 yrs	ECEARTH/CCLM	RCP8.5[2081–2100]	no ice

the spinup using the 1966 ice-surface DTM (Sect. 3), and let the ice cap evolve freely until a steady-state is reached, which takes c. 250 years. The 250-year spinup prevents initial model drift when the historical simulation is launched. To measure the performance of the spinup, we calculate RMSE between simulated thickness at the end of spinup, and observation-based thickness in 1966. The latter was derived by subtracting bedrock elevation from the 1966 DTM (Sect. 3.1).

5.2 Historical ice-cap evolution

After the calibration and spinup, we launch historical simulations of Jostedalsbreen for the period 1960–2020 (‘Historical’ in Table 2). This period was chosen because a high-resolution climate reanalysis is available (Sect. 4.3). The 60-year period is



also long enough for SMB to have an effect on ice-cap dynamics and outlet glacier frontal variations. It also ensures that recent SMB forcing is imprinted ‘in the pipeline’ for ice-cap evolution in future simulations (Sect. 5.3.3).

For historical SMB simulations (1960–2020), we employ daily mean temperature and daily total precipitation from the seNorge_2018 reanalysis dataset. Meanwhile, we force the ice-flow model with annual SMB averaged from the simulated monthly SMB, since we are not interested in seasonal variations in historical simulations. Regardless, the mesh resolution of 100–300 m is too coarse to capture seasonal frontal variations.

For historical simulations, we forced the ice-flow model with simulated historical SMB (Sect. 4.3) in a one-way fashion (cf. Sect. 5.3.1), as we found that a two-way coupling was not needed, since elevation changes 1960–2020 are minor (Andreassen et al., 2023). The historical runs were therefore initiated with SMB downscaled to the 100 m 1966 DTM and then interpolated onto the mesh of the ice-flow model.

We use SMB model output with precipitation correction factors adjusted to each individual glacier, as outlined in Sect. 4.3. For both the spinup and historical simulations, we apply a positive SMB correction of 0.75 m w.e. in the northern part of the ice cap (cf. Fig. 1). Without this manual correction, the ice cap thins heavily during spinup, which is not realistic. Modelled SMB suggests that the northern part of the ice cap is the area with the largest mass loss over the spinup period 1960–1989 (Sjursen et al., 2025). In particular, the 1960s display considerable mass loss due to a strong negative anomaly in precipitation amounts. This could be a result of bias in the climate data and/or in the estimated precipitation correction factors for this region. Meanwhile, the assumption of a steady-state ice cap 1960–1989 is possibly less accurate in this area than others. In fact, geodetic mass balance reveals the largest thinning in the northern part of the ice cap over the 1960–2020 period (Andreassen et al., 2023),

The modelled ice-cap state at the end of the historical simulation is validated against present-day surface topography from 2020, and observed ice margins in year 2019. We also compare the model’s ability to reproduce present-day ice velocities (Sect. 3.2). A good model performance entails low RMSEs for both ice thickness and velocities. Note that the observed ice velocities have already been used in the calibration of basal friction parameters (Sect. 5.1), and should therefore not be viewed as a completely independent dataset for validation.

5.3 Future change

5.3.1 Coupling of SMB and ice-flow model

SMB-elevation feedback. In future simulations, the models for ice dynamics (Sect. 4.1) and SMB (Sect. 4.3) are two-way coupled at yearly intervals. Thereby, we account for the feedback between SMB and the changing glacier-surface elevation over time. At the end of each year, the modelled ice-surface topography is used as input for the downscaling routine described below. SMB is downscaled to the updated ice-surface topography on a monthly basis for the subsequent year, in 100 m resolution. The obtained 100 m SMB is then interpolated onto the mesh of the ice-flow model, which then simulates ice-cap evolution over the following year.



We distinguish between different representations of SMB, to isolate the effects and feedbacks with SMB involved. The reference-surface mass balance, SMB_{static} , refers to the SMB on an unchanging reference geometry (no elevation or area changes) and serves as a direct indicator of climatic variations (Elsberg et al., 2001; Huss et al., 2012). The conventional surface mass balance, $SMB_{transGeom}$, is produced with the coupling procedure described above and accounts for the dynamic adjustment of both ice-surface elevation and glacier area to the SMB forcing. To isolate the SMB-elevation feedback from the effect of glacier area changes, we also evaluate $SMB_{transArea}$, which is SMB for an evolving area, but without elevation changes.

High-resolution downscaling of SMB. To provide high-resolution distributed SMB to the ice-flow model we downscale the modelled SMB from the 1 km DTM of the seNorge_2018 dataset to 100 m resolution using the elevation-dependent downscaling procedure of Noël et al. (2016). This algorithm downscales SMB from a coarse to a high-resolution DTM grid based on linear regression to determine local SMB gradients. We apply the algorithm on a monthly time scale to retrieve monthly fields of SMB at 100 m resolution. First, for each grid cell of the 1 km DTM the local SMB gradient (slope of the regression) is computed using the 1 km SMB and elevation of the given cell and the adjacent cells. The intercept of the regression in the given cell is then found using the 1 km SMB and elevation and the local SMB gradient. Secondly, the 1 km regression coefficients (slope and intercept) are bilinearly interpolated to the 100 m DTM to retrieve 100 m regression coefficients. Finally, SMB is computed in each 100 m grid cell using the 100 m regression coefficients and elevation in the given cell.

5.3.2 Future SMB

To simulate SMB over the period 2021–2100, we construct time series of daily mean temperature and precipitation using a combination of daily variability in historical data (seNorge_2018) and future trends from regional and global climate model (RCM/GCM) projections (Table 2). The general procedure is based on the approach of van Pelt et al. (2021) and can be outlined as follows: (1) we detrend daily temperature and precipitation in seNorge_2018 for a 20-year historical reference period (2001–2020) by subtracting monthly linear trend models from the historical data, (2) for each combination of climate model and future emission scenario (eight in total) we compute monthly linear trends for 20-year periods (2021–2040, 2041–2060, 2061–2080 and 2081–2100), and establish piece-wise linear functions for future changes in temperature and precipitation, and (3) we superimpose the piece-wise linear functions on the detrended reference data, repeated for each future 20-year future period.

The basis for future trends in temperature and precipitation are 1 km downscaled and bias-corrected daily resolution temperature and precipitation fields (Wong et al., 2016) based on four EURO-CORDEX GCM/RCM combinations (Jacob et al., 2014) and two Representative Concentration Pathway (RCP) emission scenarios that have been used to assess climate change impacts in Norway (Hanssen-Bauer et al., 2017, see Table 2). The original 12.5 km resolution GCM/RCM fields have been re-gridded to 1 km resolution and bias-corrected (see Wong et al. (2016) for details) using 1 km resolution temperature and precipitation fields from the seNorge dataset (version 1.1; Mohr, 2008).

When detrending precipitation, the monthly trends and intercepts are distributed evenly over each day of the month. This results in some days having negative precipitation values, even after the addition of future monthly trends. Where this occurs, daily precipitation is set to zero. This means that the number of wet/dry days in future periods is not necessarily consistent with



the reference period. We judge this to be of minor importance since repeating past variability may not be fully representative of future conditions. Capping negative values at zero may also result in deviations between future precipitation sums and computed trends. However, since precipitation generally increases in future projections, large differences are unlikely to occur.

5.3.3 Future ice-cap evolution

350 After ensuring that the model can satisfactorily reproduce recent glacier changes and dynamics, we conduct a suite of experiments of future evolution of the ice cap (Table 2). These simulations are grouped into four categories: ‘Future’, ‘Commit’, ‘Reverse’ and ‘Regrow’. Below we describe each group of experiments in more detail.

‘Future’ simulations (Sect. 6.2.2) are run from 2021–2100, using SMB based on climate forcing from the four GCM/RCM combinations, and for RCP4.5 and RCP8.5 emission scenarios. Each of these simulations, eight in total, start from the transient
 355 model state at the end of the ‘Historical’ simulation. This ensures consistent internal model dynamics and, crucially, that future simulations account for the time-lagged ice-cap response to recent climate forcing. We specifically designed an experiment called ‘CommitNow’ to assess this aspect, where we let the historical simulation continue until 2300, using the mean annual SMB of 2001–2020 as a constant forcing.

The modelled ice cap in 2100 produced using the ECEARTH/CCLM model combination is used as the initial state for
 360 the ‘Commit’- and ‘Reverse’-experiments (Sections 6.2.3; 6.2.4). This model combination was chosen because it renders a mid-range volume evolution across the four climate model combinations (Sect. 6.2.2). The Commit experiments quantify mass losses ‘in the pipeline’ by 2100, without any further greenhouse gas emissions. In these simulations, the ice cap evolves for another 200 years until 2300, using mean annual SMB from 2081–2100 as a constant forcing. Meanwhile, Reverse experiments explore the potential to backpedal the modelled 21st-century evolution of the ice cap (Reverse4.5 and 8.5). To this end, mean
 365 annual SMB of 2001–2020 is used as forcing.

Finally, we assess whether Jostedalsgreen ice cap can regrow from scratch in the present or future climates (Sect. 7.2). These experiments run for 1000 years and use mean annual SMB for the present-day (2001–2020, RegrowNow) or end-of-the-century climate (2081–2100, Regrow4.5 and 8.5). SMB and ice dynamics are coupled every 20 years in these simulations, as opposed to every year in the other future runs. The longer coupling interval for ‘Regrow’ experiments saves computation time
 370 and has here negligible effect compared to coupling every year, due to the relatively slow dynamics involved when growing Jostedalsgreen from no ice.

6 Results

In this section we show the results of our historical (Sect. 6.1) and future (Sect. 6.2) simulations. The historical simulations provide insight in their own right; no study has previously simulated the recent evolution of Jostedalsgreen as a whole, including
 375 ice dynamics. These experiments also (1) generate the initial geometry and volume for future simulations and (2) provide benchmarks that give confidence in the modelled ice dynamics and SMB, and their ability to collectively produce an ice cap that resembles reality. These conditions need to be fulfilled before embarking on future projections.



6.1 Historical ice-cap change 1960–2020

Our simulations show that the modelled ice volume of Jostedalsbreen ice cap has changed very little since the 1960s. The simulated present-day volume is in excellent agreement with the observed (Fig. 2a). Meanwhile, the modelled volume in 1966 is a few km³ lower than observed, yet well within observational uncertainty.

Partitioning the volume change 1960–2020 reveals that modelled and observed volumes agree very well for the South, Central and North parts of the ice cap (Fig. 2a; cf. Fig. 1). Of these, the central ice cap amounts to c. two-thirds of the total ice volume of 70.6 ± 10.2 km³.

The modelled SMB and ice-volume evolution exhibit three distinct variations during the historical period: approximate balance 1960–1989, apart from slightly negative SMB in the early 1960s; positive SMB and ice-cap growth in the 1990s; and mass loss after the year 2000 (Fig. 2c). The reader is referred to Sjurset et al. (2025) for further details on the historical SMB.

The overall agreement between modelled and observed ice margins is good (Fig. 3), especially considering the complex topography and great diversity in glacier characteristics. This agreement includes the large outlets Austerdalsbreen, Tunsbergdalsbreen and Nigardsbreen on the eastern side of the ice cap (cf. Fig. 1). However, the model overestimates the total area by 23% (modelled area 566 km² in 2020; Fig. 9b), mainly due to too advanced margins in the north, where some areas with relatively coarse model elements contribute to an apparent too large areal extent. These areas generally host very thin ice (less than 50 m; Fig. 3a), which means that they contribute little to total ice volume (Fig. 2a). The modelled area shown in Fig. 2b includes all areas with a modelled ice thickness of 10 m or more. If instead areas with more than 20 m thick ice is considered as glacierised, the modelled area of 2020 becomes 520 km², which reduces the overestimation of areal extent from 23% to 13% (Fig. A2). This means that the present-day modelled ice cap hosts large areas (46 km²) of ice only 10–20 m thick, some which are even disconnected with the main ice cap. This sensitivity analysis, together with a visual inspection of the modelled ice-cap margins in Fig. 3b, illustrate that both the modelled volume and area should be considered satisfactory.

Overall, the modelled thickness agrees well with observation-based gridded thickness (RMSE = 24 m; Fig. 3b). Tunsbergdalsbreen, the longest (28 km) and thickest (maximum 626 m; Gillespie et al., 2024) glacier in Norway, sees excellent agreement for ice thickness (RMSE c. ± 30 m), with deviations within the typical 30–80 m uncertainty of the observation-based ice-thickness data for this glacier (Gillespie et al., 2024). This glacier alone contains roughly as much ice (10.8 km³) as the entire South (12.3 km³) and North (8.5 km³) parts of the ice cap, respectively (Gillespie et al., 2024). For the lower part of Nigardsbreen, the present-day ice surface and glacier front are reproduced within c. ± 15 m, while the upper part is somewhat thinner than the observation-based thickness, with thickness misfit similar to the upper limit of uncertainty for the observation-based dataset (c. 20–50 m, Fig. 3b; Gillespie et al., 2024).

Conversely, the model overestimates ice thickness by c. 20–100 m for a few very steep, narrow outlet glaciers, for example Briksdalsbreen and lower Austdalsbreen (Fig. 3b; cf. Fig. 1). For some of these outlet glaciers (e.g. Briksdalsbreen, Brenndalsbreen, Lodalsbreen), this is associated with frontal positions c. 0.5–1 km downvalley of the present-day observed termini (Fig. 3a). On the other hand, there are also some steep and/or narrow glaciers that are very well represented both in terms of terminus position, thickness and flow speeds (e.g. Fåbergstølsbreen, and several glacier units in the South; Fig. 3)




Figure 2. Simulated historical change (a–c) and future (d–f) evolution with experiment Future4.5-ECC (cf. Table 2). Recent historical periods of little change (gray), ice-cap growth (blue) and decay (red) are shaded in (a–c). Observed volumes and areas are shown as filled circles in (a) and (b). In (c), historical SMB with evolving model area is shown ($SMB_{transArea}$), but without elevation changes, since these make little difference (Sect. 5.2). Shown in (f) is SMB with transiently evolving geometry ($SMB_{transGeom}$), including the SMB–elevation feedback and area changes.




Figure 3. Comparison between model simulations in this study, and observation-based ice thickness (‘obs.’) and observed velocity at the end of the historical simulation (2020). (a) Modelled ice thickness; (b) thickness misfit, model - observation-based (Gillespie et al., 2024); (c) modelled annual glacier velocity in 2020; (d) velocity misfit, model - observed (corrected dataset from Millan et al. (2022)). Glacier outlines and divides from 2019 are shown with thin black lines (Andreassen et al., 2022). Ice-free areas within the model domain are shown in grayscale in (a) and (c).



Velocities are overall well reproduced ($\text{RMSE} = 23 \text{ m a}^{-1}$), with some spatial variability. For example, it appears that several modelled outlet glaciers are too slow in their steepest regions (e.g. ice falls of Nigardsbreen and Tunsbergsdalsbreen). This is true when comparing modelled velocities against the global velocity product (Millan et al., 2022). As mentioned in Sect. 3.2, velocities in this global dataset may be too high in some locations, which would mean that the modelled velocities in fact are more accurate than they appear. Ice thickness and ice margins are indeed well represented in many areas despite too slow apparent velocities. This indicates that the combined SMB–dynamics model system performs well regardless of the discrepancies between modelled and observed velocities.

6.2 Modelled future evolution 2021–2300

6.2.1 Reference surface mass balance

Future SMB simulations reveal that substantial changes in SMB forcing ($\text{SMB}_{\text{static}}$) on Jostedalsbreen can be expected over the coming century (Figs. 4 and 5). Moreover, differences between climate model combinations and emission scenarios are striking (Fig. 4c). The majority of climate models result in cumulative mass losses of around -25 m w.e. over the next 30–40 years, with relatively low variation between the RCP4.5 and 8.5 scenarios. After around year 2050, models show a larger spread with cumulative mass losses up to around -150 m w.e. by the end of the century. Note that this mass loss is for an SMB over a fixed geometry ($\text{SMB}_{\text{static}}$), that is, before considering the effects of future area changes and the SMB–elevation feedback (see Sect. 6.2.2 and Fig. 11). One model combination stands out from the rest: MPI/CCLM shows a large mass increase under the high-emission scenario (Fig. 4c). This is a result of rapid and extreme (unrealistic; Sect. 7.1) gains in future precipitation, along with little change in summer temperature, which renders a positive cumulative $\text{SMB}_{\text{static}}$ over the course of the 21st-century (Fig. 4). The near balance with ECEARTH/HIRHAM under a medium-emission scenario is a result of similar, but less extreme, mechanisms.

Our simulations indicate an increase in the frequency and magnitude of negative SMB throughout the century, under both medium- and high-emission scenarios (Fig. 4a and b, respectively). Under high-emission scenarios, ice-cap wide mass losses can reach SMB magnitudes similar to those seen only for the low-lying tongues of Jostedalsbreen today (e.g. around -6 to -7.5 m w.e. on the lower tongue of Nigardsbreen in 2023; Kjølmoen et al., 2024). Our results also indicate that, under both emission scenarios, years with positive ice-cap wide SMB can occur far into the 21st century, although this occurs significantly less often during the second half of the century. Uncertainties in future SMB are greater for high-emission scenarios, mainly due to the substantial spread between climate model projections (see Sect. 7.1).

Considering the mid-range climate forcing ECEARTH/CCLM, before considering the SMB–elevation feedback (Fig. A4), the ice cap will retain a substantial accumulation area at high elevations in the South and Central parts for RCP4.5 by the end of the 21st century (Fig. 5a), while annual SMB rates in the North will be mainly negative. For RCP8.5, SMB rates remain positive only for smaller parts on the central plateau (Fig. 5c), with a large reduction in SMB rates over the entire ice cap compared to present day (Fig. 5d). The greatest reduction in SMB occurs on low-lying glacier tongues and along ice-cap



margins (Fig. 5b and d). However, the South part, which has shown mostly positive SMB rates over the historical period
 445 (1960–2020; Sjurset al., 2025), also displays large negative changes in SMB by the end of the 21st century for this scenario.




Figure 4. Modelled glacier-wide annual reference surface mass balance (SMB_{static}, fixed DTM 2020 geometry and 2019 area) for Jostedalsgreen over the period 2021–2100. Results are presented as median annual mass balance across climate model combinations for (a) RCP4.5 and (b) RCP8.5, and (c) cumulative mass balance for each model and RCP combination. Bars in (a) and (b) show median modelled mass balance and whiskers show spread of models (minimum and maximum values).




Figure 5. Modelled annual reference surface mass balance rates (SMB_{static} , fixed DTM 2020 geometry and 2019 area) for Jostedalsgreen by the end of the 21st century (2081–2100 mean) for (a) RCP4.5 and (c) RCP8.5, with ECEARTH/CCLM climate forcing. Difference between end-of-century and present-day (2001–2020 mean) SMB_{static} rates for (b) RCP4.5 and (d) RCP8.5.

6.2.2 Ice-cap evolution in the 21st century

Our dynamical model simulations reveal that Jostedalsgreen stays in near balance until year 2040. Thereafter, the ice cap is projected to lose half (two-thirds) of its ice volume under medium (high) emissions until the year 2100 (Figs. 2d and 9b). These results are obtained with the ECEARTH/CCLM climate forcing, which gives an ice-volume loss that is mid-range across the four climate model combinations (Fig. 11; cf. Table 2). Model simulations forced by the full ensemble of climate models suggest that, until the year 2100, Jostedalsgreen will lose 10–75 % of its present-day volume under medium emissions (Fig. 11f). Meanwhile, for high emissions, future volume evolution ranges from 75% loss (CNRM/CCLM) to 17% gain (MPI/CCLM) by 2100. The latter unexpected future increase of volume with MPI/CCLM under high emissions is a result of positive SMB. While we include these results for completeness, we deem this climate projection unrealistic (Sect. 6.2.1). If we exclude the



455 MPI/CCLM results, the projected future mass loss is 12–74% for medium emissions, and 53–74% under high emissions (Fig. 11f).

In our experiments of future evolution until 2100, Jostedalsbreen splits up into three separate ice caps both under medium and high emissions (ECEARTH/CCLM, RCP4.5 and 8.5; Fig. 6). With medium emissions, North separates from the rest of the ice cap in the 2070s, while South becomes a separate ice cap in the 2090s (see Supplementary video 1). Meanwhile, 460 under high emissions, these breakups occur around 10–20 years earlier; in the 2050s for North, and in the 2080s for South (see Supplementary video 2). However, it is important to note that these separations and other small-scale details include a high degree of uncertainty. They can vary by several decades due to differences in the future climate projections (Sect. 7.1), underlying model limitations (Sect. 7.3) and local mismatches between the modelled and real-world geometry. For example, the model mesh is too coarse (Fig. A1) to resolve a few small ice-free patches (100–200 m across) which currently partly 465 separate the southern and central parts (cf. Fig. 1). Contemporary modelled ice thickness is 20–50 m in this area, while the measured ice thickness is less than 25 m (Gillespie et al., 2024). This suggests that a separation may occur several decades before the 2080s.

The North part of Jostedalsbreen, with its plateau situated c. 100–300 m lower than the Central ice cap (Fig. 1), is simulated to decline most strongly. The modelled future thinning here is 20–100 %, and total volume loss 74–93%, depending on emissions 470 (Fig. 6). This is already the area of the ice cap where current mass loss is greatest (Andreassen et al., 2023; Sjørnsen et al., 2025).

Regardless of emission pathway, most of the volume loss is driven by strong thinning of 20–200 m at the ice-cap margins (Fig. 6), with less pronounced thinning of 20–100 m on the plateau. This results in frontal retreat and an overall steepening of the ice cap’s outlet glaciers (Fig. 7ab). These patterns can be explained by a $SMB_{transGeom}$ that, over the course of the century, 475 becomes increasingly negative in ablation zones, while net annual SMB on the plateau remains positive for many of the years in the coming decades (Fig. 7cdf). Annual SMB on the plateau is in fact rather similar between the medium and high emissions, which is likely due to similar precipitation amounts (see Sect. 7.1). Meanwhile, SMB along the margins of key outlet glaciers such as Nigardsbreen and Tunsbergdalsbreen is 3–8 m w.e. more negative than at present (Fig. 7cd). This corresponds to typical annual SMB values of -12 to -18 m w.e. at the frontal regions, which presently are located at 300–700 m a.s.l.

480 Depending on the emission scenario, terminus retreat is simulated to be c. 2.5–5 km for Nigardsbreen and c. 5–10 km for Tunsbergdalsbreen, respectively (ECEARTH/CCLM; Fig. 7ab). Because mass loss at the high-lying plateau here is rather subdued, these glaciers still retain around half of their volume by the year 2100, even under high emissions (Fig. 8). Similarly, Austerdalsbreen loses the entire glacier tongue below its ice fall under RCP4.5, but two-thirds of the ice volume remains by 2100. To reiterate this point: Tunsbergdalsbreen retreats 10 km under high emissions (Fig. 7b), yet by the end of the century 485 still hosts large upper areas with ice more than 500 m thick (Fig. 6d). As we will see in Sect. 6.2.3, this is the most resilient part of the ice cap.




Figure 6. Future model thickness and thickness change under medium (a–c) and high (d–f) emissions, using ECEARTH/CCLM climate forcing. (a,d) Modelled ice thickness in year 2100; (b,e) Total ice thickness change (future–present); (c,f) Relative thickness change. In (e–f), yellows means a slight surface lowering, while dark red implies strong thinning. In (c,f), -100 % thickness change means complete deglaciation




Figure 7. Modelled ice-surface (a, b and e) and SMB evolution (c, d and f) 2021–2100 under RCP4.5 (ECEARTH/CCLM) for three transects on Jostedalsgreen. The three transects are shown in the inset map, which is slightly rotated for visibility.




Figure 8. Future mass loss for Southern, Central and Northern parts (delineated by dashed lines) and selected glacier basins (thick white outlines) of Jostedalsgreen ice cap. The discs show relative mass loss (% of 2020 volume) and remaining volume (km³) in year 2100 (RCP4.5, RCP8.5) and 2300 (Commit4.5, Commit8.5). The background map shows simulated thickness in year 2100 under scenario RCP4.5, and glacier outlines in 2019 (black lines Andreassen et al., 2022). Climate forcing is based on ECEARTH/CCLM (complete list in Table 2).



6.2.3 High committed mass loss 2100–2300

Our simulations show that if greenhouse-gas emissions were capped at present-day levels, half of the ice cap would still disappear (CommitNow; Figs. 9b and 10a). Under the medium-emission scenario RCP4.5, Jostedalsbreen loses around half
 490 of its present-day volume by 2100. Similarly, about one-third of the ice volume remains by 2100 under the high-emission scenario RCP8.5 (Fig. 9b). However, this is only half of the story, since Jostedalsbreen will not have reached a steady-state in 2100. Further mass loss is ‘in the pipeline’, as the ice cap undergoes additional dynamic adjustments and continues to thin due to the effective SMB-elevation feedback. In the high-emission case, our simulations show that the ice cap is committed to vanish completely by around 2200, even without any post-2100 climate warming (Commit8.5; Figs. 8 and 9b). Similarly, with
 495 moderate 21st-century emissions (RCP4.5), the ice cap is bound to lose 90% of its present-day volume by 2300 (Commit4.5).

Upper Tunsbergsdalsbreen and the nearby high-elevation areas to the north (upper Brenndalsbreen and southwest corner of upper Nigardsbreen) is the part of the ice cap that is most likely to survive the longest. Another resilient high-lying ice complex appears to be upper Fåbergstølsbreen and nearby areas along the local ice divides (Fig. 10bc).

6.2.4 Failure to recover

Our Reverse experiments test whether the ice cap can recover if future greenhouse-gas emissions are rolled back immediately
 500 at the year 2100. That is, resetting the climate forcing to an SMB associated with the prevailing climate of 2000–2020. We find that such hypothetical time travel leads to barely any recovery (Fig. 9b). The inability to recover holds regardless of whether emissions are moderate or high before the reversal. The main reason for the failure to recover is that the ice cap is currently strongly out of balance with the prevailing climate. In other words, turning back the clock from year 2100 to the
 505 climate of 2000–2020 means aiming for an ice-cap that is considerably smaller than that of the present-day. The CommitNow experiment illustrates this, where Jostedalsbreen’s volume is halved by the year 2300 (Fig. 9b). Indeed, the Reverse4.5 and Reverse8.5 simulations are on their way to steady-state volumes similar to that of the CommitNow experiment (c. 30–35 km³; Fig. 9b). Another aspect preventing recovery may be that by 2100, the ice cap is on the course to an even lower ice volume (cf. Commit4.5 and Commit8.5 experiments; Fig. 9b). Therefore, any ice-cap growth due to a more favourable SMB, will in fact
 510 compete with ongoing dynamic adjustments and associated SMB-elevation feedback working towards a smaller ice cap.

7 Discussion

7.1 Future surface mass balance

Sensitivity to climate models. The majority of our simulations suggest that Jostedalsbreen will experience a substantial mass loss over the course of the 21st century (Fig. 11). At the same time, modelled SMB based on the ensemble of climate models
 515 leads to a large spread of possible future ice-cap evolution. This suggests both a strong sensitivity to choice of climate model and scenario, and that there are substantial variations between projected temperature and precipitation changes between climate models (Fig. A5). Medium and high emission scenarios show a mean annual temperature increase over Jostedalsbreen




Figure 9. Historical and future evolution of (a) reference SMB (SMB_{static}), (b) ice volume, and (c) area, using future ECEARTH/CCLM climate forcing. Historical observations (gray circles) and uncertainties (whiskers) are shown. See Table 2 for a full experiment overview.




Figure 10. Maps of modelled ice thickness in 2300, after accounting for committed mass loss due to (a) SMB 2000–2020, (b) medium 21st-century emissions; (c) high 21st-century emissions. These ‘Commit’ simulations have temporally fixed SMB, and run over (a) 2021–2300; (b) 2101–2300; and (c) 2101–2300, see Table 2. The SMB in (b) and (c) is derived from ECEARTH/CCLM climate forcing. Present-day glacier outlines (2019) are shown with black lines (Andreassen et al., 2022).

(2019 area) of 1.8 °C and 3.4°C, respectively, by the end of the century. The mean increase in annual precipitation sums for the respective scenarios is 27% and 65%, with an increase of 30% and 67% in winter (Fig. A5c). However, for a given
 520 emission scenario there are considerable differences between the models. For example, projected change in summer (May–Sep) temperature varies from -0.3°C to +2.7°C under RCP4.5, and +0.2°C to 3.0°C under RCP8.5 (Fig. A5b). The spread in winter (Oct–Apr) temperature is smaller, around 1.2°C and 1.9°C for RCP4.5 and 8.5, respectively (Fig. A5a). Differences in precipitation between climate models and scenarios are even more striking, with models projecting winter precipitation changes between -14% and +72% for RCP4.5 and between +39% and +80% for RCP8.5 (Fig. A5d). Moreover, the timing of
 525 precipitation changes also varies across the climate models, both in terms of the time of year and the evolution throughout the century.

Seasonality. Our simulations allow us to assess future changes in seasonality and magnitudes of ablation and accumulation for Jostedalsgreen. With the ECEARTH/CCLM projections, the largest changes in the SMB for the transiently evolving Jostedalsgreen can be expected in autumn and summer under medium and high emissions, respectively (Fig. 12). Under an
 530 RCP4.5 pathway, the ablation season is prolonged by approximately a month, from mid-September to mid-October, with an equivalent shortening of the winter accumulation period (Fig. 12a). Meanwhile, ablation magnitudes at the peak of summer change relatively little. This contrasts with high emissions, where strong summer melt intensifies and is more prolonged



between June to September (Fig. 12b). These differences reflect both differing warming and precipitation trends between climate projections (Fig. A5), and variations in which seasons and months such changes are projected to occur. While both ECEARTH/CCLM scenarios render relatively strong winter warming (Fig. A5a), temperature increase in summer is smaller (Fig. A5b). Moreover, there are variations between the projections within seasons. For example, warming is stronger in October for RCP4.5 than RCP8.5, which could explain why autumn SMB changes are less pronounced for the latter. However, it should be noted that the future monthly SMB in Fig. 12 not only reflects climatic changes but also accounts for the feedback of glacier retreat and surface-elevation lowering ($SMB_{transGeom}$).

Precipitation. Most climate models show that precipitation gains increase with elevation. This explains why the models indicate larger precipitation increases over Jostedalsbreen than for western Norway in general (Hanssen-Bauer et al., 2017). This may be a natural consequence of increasing and intensifying precipitation; precipitation amounts on Jostedalsbreen are likely substantially affected by orographic uplift. However, the large precipitation increase over Jostedalsbreen may also be a result of how climate models are downscaled and bias-corrected to the seNorge version 1.1. dataset (Wong et al., 2016), which relies on precipitation lapse rates to extrapolate precipitation to higher elevations (Mohr, 2008). Regardless, our results highlight the sensitivity of glacier projections to climate model input and considerable uncertainty in future temperature and precipitation in mountainous regions. For glaciers with a substantial mass turnover and likely large future accumulation area, such as Jostedalsbreen, our results highlight that future glacier evolution strongly depends on future precipitation changes, and that projections of future glacier change remain uncertain due to the (in)ability of climate models to capture these changes.

SMB feedbacks. In future SMB simulations there are two competing feedbacks between SMB and glacier geometry changes at play (Elsberg et al., 2001; Huss et al., 2012). The first is the retreat feedback, which contributes to a more positive $SMB_{transArea}$, relative to the reference SMB_{static} , as the glacier retreats to higher elevations where ablation is smaller. The elevation feedback acts in the opposite direction, contributing to a more negative $SMB_{transGeom}$ as the glacier surface is lowered to elevations where ablation is higher (Huss et al., 2012). To quantify the relative and net impacts, we computed the transient SMB in Megatonnes (Mt) across the entire ensemble of climate model combinations (Fig. 11d). This analysis shows that, for most climate models, the total mass balance SMB_{static} becomes very negative towards the end of the century, since it ignores the two feedbacks. Meanwhile, the positive retreat feedback, which includes area changes, makes the $SMB_{transArea}$ c. 0.5–2.5 Mt less negative than SMB_{static} . Note that the glacier is still losing mass overall for all scenarios with this feedback included. Including also the elevation feedback, where glacier geometry and SMB is fully coupled, shifts the total mass balance 0.1–0.6 Mt more negative compared to SMB with the retreat feedback only. Simulation of long-term glacier evolution thus requires careful consideration of the mass-balance elevation feedback. If not, a strong positive bias in modelled SMB can be expected (Fig. 11).

7.2 Jostedalsbreen's fate compared to other glaciers in Norway

Potential for regrowth. A pertinent question is whether Jostedalsbreen can regrow from no ice to its current size under contemporary or future climate conditions. The answer is no, neither in the current climate, nor in a warmer future one, at least not using the chosen main climate model combination ECEARTH/CCLM. If Jostedalsbreen disappeared tomorrow, the




Figure 11. Future evolution of SMB and ice volume 2021–2100 across the ensemble of climate model forcings, cf. Table 2. (a) SMB with fixed present-day ice geometry (SMB_{static}); (b) SMB over the modelled area evolving over time, without accounting for the SMB-elevation feedback ($SMB_{transArea}$; see Sect. 4.3); (c) SMB over a transiently updated area and elevation, ($SMB_{transGeom}$); (d) total SMB of the representations in (a–c) in Megatonnes ($10^6 \text{ m}^3 \text{ w.e.}$), using a 5-year moving mean; (e) Cumulative $SMB_{transGeom}$; (f) Modelled ice volume with $SMB_{transGeom}$. Line styles shown in the legend in (f) are consistent in (a–c and e–f), representing climate model combinations ECEARTH/CCLM (ECC), ECEARTH/HIRHAM (ECH), CNRM/CCLM (CNC), and MPI/CCLM (MPC).




Figure 12. Mean monthly surface mass balance (SMB) for historical (SMB_{static} , grayscale; fixed DTM 2020 geometry and modelled 2020 area) and future ($SMB_{transGeom}$; colours) decades for ECEARTH/CCLM (a) RCP4.5 and (b) RCP8.5. The 2090s mean refers to the 11-year period 2090–2100.

present-day climate (2000–2020) would render a much smaller ice cap than the present-day Jostedalsgreen, with a 59% smaller ice volume (29 km^3), 30% smaller area (317 km^2 and 60% thinner ice [mean ice thickness of only 91 m]; Fig. A3). This implies 60% thinner ice on average than for the present-day (Gillespie et al., 2024). The modelled volume of this regenerated ice cap is similar to present-day Søndre Folgefonna (30 km^3 ; Ekblom Johansson et al., 2022), Norway’s third largest ice cap by area, and around three times larger than the present-day Hardangerjøkulen (ice volume of 9.3 km^3 , area of 71 km^2 , max ice thickness c. 350 m), one of the most well-studied ice caps in Norway (Andreassen et al., 2015). Meanwhile, we barely produce any glacier ice when trying to grow Jostedalsgreen from scratch using an SMB (2081–2100) associated with medium 21st-century emissions, and no ice at all with a high-emission climate (experiments Regrow4.5 and Regrow8.5). For Hardangerjøkulen, Åkesson et al. (2017) similarly found that a small negative SMB anomaly relative to the present-day would render a regrowth from no ice impossible. These findings collectively imply that if two of the largest Norwegian ice caps would melt away, a recovery appears very difficult, and would require a colder and/or wetter climate than the present one.

Large-scale studies. Regional studies of Scandinavian glaciers have shown that Jostedalsgreen will lose 35% (RCP4.5) to 50% (RCP8.5) of its area and volume by 2100 (Huss and Hock, 2018). This is slightly lower mass loss than estimated by the current study (49% for RCP4.5, 63% for RCP8.5; Fig. 8). Meanwhile, Compagno et al. (2021) suggested that Nigardsbreen would lose 64% (RCP4.5) to 85% (RCP8.5) of its present-day volume, which is roughly 50% more mass loss than our mid-range results for Nigardsbreen from ECEARTH/CCLM (Figs. 8). More broadly, estimates for 21st-century mass loss for Scandinavian glaciers and ice caps as a whole range from 40–80% (RCP4.5) and 70–100% (RCP8.5; e.g. Marzeion et al., 2012; Compagno et al., 2021; Rounce et al., 2023). These large-scale studies may perform well on regional scales, but their



estimates for individual ice masses are not directly comparable with those of a detailed study like ours. As emphasised in Sect. 1, regional studies treat connected glacier units separately, use crude ice-flow dynamics, and have not benefitted from such detailed ice-thickness data and surface mass balance simulations as we have in the current study. The methods, scales and implementations thus differ greatly compared to the detailed, higher-order, high-resolution, spatially continuous modelling approach presented here.

Detailed studies. For high-emission scenarios, some studies of other ice caps in Norway suggest almost complete disappearance by the year 2100 or soon thereafter, including Hardangerjøkulen (Giesen and Oerlemans, 2010; Åkesson et al., 2017), and Spørteggreen ice cap, c. 15 km east of Jostedalsbreen (Laumann and Nesje, 2014). Our findings suggest that Jostedalsbreen will be more long-lasting, and retain one-third to half of its present-day ice volume at the end of the 21st-century, depending on emissions (Fig. 9b). Not only does Jostedalsbreen contain more ice – its particular resilience is likely also related to the vast plateau being at high elevations, and relatively large precipitation amounts, which will increase even more in the future. Even so, an extrapolation of the high-emission modelled rate of mass loss by 2100 suggests a complete disappearance in the late 22nd-century, if emissions are not curtailed. While our Commit-experiments do not explicitly test this, they illustrate that a complete disappearance of Jostedalsbreen may be in the pipeline already by 2100.

Oerlemans (1997) simulated future evolution of Nigardsbreen (cf. Fig. 1) using a 1d-flowline model. For a warming of 1–2 °C and +10–20% precipitation, Nigardsbreen was projected to lose 24–42 % of its present-day (AD 1950) volume. This is somewhat lower than our modelled 40–56% volume loss for Nigardsbreen for RCP4.5 and RCP8.5, respectively (Fig. 8). However, the thickness data, models and future climate projections used by the current study and Oerlemans (1997) differ significantly.

The impact of future warming on Briksdalsbreen has previously been modelled using a flowline model together with a degree-day model for surface mass balance (Laumann and Nesje, 2009). While they did not provide volume projections, they estimate 21st-century frontal retreat of 2.5 to 5 km under scenarios similar to RCP4.5 and RCP8.5 used here. A similar retreat is simulated in the current study, along with a volume loss of 33% (RCP4.5) and 46% (RCP8.5) for the glaciers draining Oldedalen as a whole, using the ECEARTH/CCLM climate (Fig. 8).

Results from Søndre Folgefonna indicate that for RCP4.5 and RCP8.5 about 40% and 80%, respectively, of the ice cap will have melted away by the year 2120 (Ekblom Johansson et al., 2022). The northern parts of the ice cap are more vulnerable than the southern parts. Søndre Folgefonna's current ice volume is 30 km³ (Ekblom Johansson et al., 2022), around 40% smaller than the ice volume of Jostedalsbreen. Considering its smaller size, Søndre Folgefonna is surprisingly resilient to future warming. The knowledge of the bed topography underneath Søndre Folgefonna (Ekblom Johansson et al., 2022) is perhaps the only one in Norway that matches the level of detail in the newly-mapped topography under Jostedalsbreen (Gillespie et al., 2024), and maximum ice thicknesses are similar, with areas of ice more than 500 m thick existing for both ice caps. Some outlet glaciers from Folgefonna are known to have reached their Little Ice Age maximum extent as late as in the 1940s (Tvede, 1973; Tvede and Liestøl, 1976), and has not retreated much since, which may explain the relative resilience to warming. Folgefonna is also a more maritime ice cap than Jostedalsbreen, with its surface mass balance heavily controlled by high winter snowfall (Tvede, 1989). This may buffer Folgefonna from future warming, and perhaps also the western outlets of Jostedalsbreen. Such snowfall



620 buffering however has its limits. As climate warms, more precipitation falls as rain rather than snow, which, for example, has been found to diminish the buffering of Patagonian glaciers (Troch et al., 2024) located in a similar maritime climate (Tvede, 1973).

7.3 Limitations and future improvements

The performance of our model system for ice dynamics and SMB depends on a multitude of choices and assumptions. These
 625 includes uncertainty in historical and contemporary ice-surface data and bedrock topography (Sect. 3.1), uncertainty in velocity data (Sect. 3.2), ice-flow physics, basal friction, ice rheology, and mesh construction (Sect. 4.1), surface mass balance model and its parameters (Sect. 4.3), as well as uncertainties in climate projections (Sect. 7.1). Most of these factors also vary spatially across the 81 glacier units.

Input data. The gridded observation-based ice-thickness dataset is based on measured ice-thickness point data combined
 630 with an ice-thickness model based on inversion (Sect. 3.1). Despite the unprecedented level of detail in this dataset, the uncertainties can in theory be up to 50–100 m in areas without thickness observations, but in practice likely smaller, particularly close to radar survey lines (Gillespie et al., 2024). An overestimated (underestimated) observation-based thickness will affect modelled ice-flow and render a too slow (too fast) future mass loss and retreat. Meanwhile, present-day thickness and ice-cap margins are overall well reproduced by the historical simulations, yet there are some mismatches at the end-of-historical state
 635 (Fig. 3). These differences for individual glaciers will propagate into the future, affecting the timing of retreat. For example, the steep Briksdalsbreen and Brenndalsbreen outlets are too advanced for the present-day (cf. Figs 1 and 3a). This means that future modelled frontal positions may be located further downvalley compared to reality.

Initial state and spinup. The future projections presented here rely on the model geometry and dynamics of the historical simulations 1960–2020. In turn, the historical experiments are influenced by the initial ice-cap state of the 1960s, obtained
 640 through model spinup (Sect. 5.1). The spinup uses the 1966 ice-surface topography as a target, and assumes that the ice cap was in steady-state in the 1960s. In reality, the ice cap will have experienced ongoing adjustments to climate variations occurring between the Little Ice Age maximum in the 1800s up until the 1950s. Indeed, several records of frontal variations of Jostedalsbreen’s outlet glaciers show retreat in the 1930s through 1960s (<https://glacier.nve.no/viewer/CI/en/>; Andreassen and Elvehøy, 2021). Such adjustments are not accounted for in the historical simulations. This may partly explain why the model
 645 underestimates the reconstructed ice volume in 1966 (Fig. 2a), and partly why a relatively modest historical retreat and volume loss is simulated – although we emphasise that there are also underlying uncertainties in the observation-based ice thickness dataset (Gillespie et al., 2024). Regardless, not accounting for the real-world ongoing decline could have resulted in chosen model parameters being biased towards too positive SMB.

SMB and climate. Historical and future ice cap evolution is heavily influenced by the SMB forcing (Figs. 2 and 9), which is
 650 subject to its own limitations, in addition to the uncertainty in climate forcing. There are several features that our SMB model does not explicitly account for, including small-scale processes such as redistribution of snow by wind and avalanching, and differences in incoming solar radiation with different aspects and topographic shading. However, some of these characteristics could be accounted for by the spatially corrected model parameters (see Sjørnsen et al. (2025) for a detailed discussion). We



have also assumed that parameter values in the SMB model are constant over time, including in a changing future climate.
 655 Some studies indicate that this may cause an oversensitivity to temperature change (Gabbi et al., 2014; Ismail et al., 2023),
 although it is unclear if more physically-based approaches improve the performance when detailed meteorological data is
 lacking (Réveillet et al., 2018). Nevertheless, the above discussion in Sect. 7.1 illustrates that modelled future SMB, and thus
 ice-cap evolution, strongly depends on the choice of climate model, where especially precipitation is uncertain. Thus, the
 uncertainty in future climate forcing, illustrated by the large spread in future projected temperature and precipitation (Fig. A5),
 660 likely outweighs the uncertainty associated with SMB model parameter values.

Ice dynamics. Modelled ice dynamics is heavily influenced by the representation of basal friction (e.g. Brondex et al., 2019;
 Åkesson et al., 2021). Here we have taken advantage of a global velocity dataset for the fastest outlet glaciers. Comparison
 against this dataset suggests that several model glaciers are less dynamic compared to reality (Fig. 3d). If the velocity data is
 correct, this could partly explain why historical outlet-glacier retreat is underestimated in a few cases (e.g. steep glaciers such
 665 as Briksdalsbreen and Brenndalsbreen; Fig. 1). It follows that more dynamic outlet glaciers would render faster frontal retreat
 than what we have projected here. Like virtually all previous model studies, we also keep friction fixed in time, ignoring any
 feedbacks between increased meltwater supply and friction in a future warmer climate, which may accelerate mass loss further
 (e.g. de Fleurian et al., 2022).

The global velocity dataset used does not come without issues, as highlighted in Sect. 3.2. We illustrated a hybrid approach
 670 to mediate this, with inversion for the largest and most dynamic glaciers, where we judge the velocity data quality to be most
 representative. Several alternative approaches to basal friction were tested. We tried a transient inversion of the friction field,
 where ice thickness was allowed to evolve using a constant SMB, and the friction parameter adjusted at regular intervals
 (10–50 years) depending on local thickening or thinning, following Pollard and DeConto (2012); Santos et al. (2023). Despite
 theoretical promise, this approach resulted in large unrealistic drift of the ice-cap geometry during spinup, and was therefore
 675 abandoned. Moreover, we also tested a spatially uniform friction parameter, an elevation-dependent friction parameter every-
 where (Eq. 3), as well as imposing the friction inversion everywhere. All of these approaches rendered poorer results (higher
 RMSEs) than our approach outlined in Sect. 3.2.

Summary. All this illustrates the challenges of tackling a complex ice cap. Indeed, we expect and accept that the modelled
 and observation-based thickness diverges to some extent, and that not all modelled glacier fronts will reproduce their historical
 680 or present-day positions.

7.4 Global implications

Glacierised regions worldwide are in decline and are on track for severe mass loss over the 21st-century. If emissions are not
 strongly reduced, many glaciers are bound to recede or disappear, including in the European Alps (e.g. Jouvet et al., 2011),
 High-Mountain Asia (Van Tricht and Huybrechts, 2023); the sub-Antarctic (Verfaillie et al., 2021) and Scandinavia (Giesen
 685 and Oerlemans, 2010). This may have consequences for local societies (Carey et al., 2017) and impact proglacial ecosystems
 (Bosson et al., 2023). Numerous smaller mountain glaciers may disappear even under the 1.5°C goal of the Paris Agreement
 (e.g. Rounce et al., 2023). Large ice caps like Jostedalsbreen contain more ice and will therefore take longer to disappear.



Ice masses with flat-topped geometry and flow governed by topography are usually called ice caps in Scandinavia, Iceland, Greenland, and Canada, while referred to as ‘ice fields’ in Alaska and Patagonia. Studies from Alaska (Ziemen et al., 2016), Arctic Canada (Schaffer et al., 2023), Iceland (Schmidt et al., 2020), Greenland (Zekollari et al., 2017) and Patagonia (Troch et al., 2024) illustrate that large ice caps and ice fields may last several hundred years into the future even under strong warming. This does not mean that ice caps are necessarily more resilient in a warming climate – rather the opposite. Because of their flat interior plateaus, ice loss will accelerate greatly as the ELA rises to and above the elevation of vast plateau areas (Harrison et al., 2001; Åkesson et al., 2017). This switch to a negative SMB over large areas drastically reduces the accumulation area – a recipe of strong mass wastage and eventual disappearance. In the case of Jostedalsbreen, our simulations suggest that this situation occurs towards the end of the 21st-century (Fig. A4cf). Under high emissions, the associated accumulation-area ratio for 2081–2100 is approximately 0.1, meaning that 90% of the ice-cap area experiences net annual melt (Fig. A4ef). The strong long-term impact is illustrated with the Commit8.5 experiment, which leads to a 99% volume reduction of Jostedalsbreen by the year 2300 (Figs. 8 and 11c). Under the more moderate emission pathway RCP4.5, areas with net positive SMB by the end of the 21st century are somewhat larger (Fig. A4), yet Jostedalsbreen is at that stage committed to a mass loss of 90% of its present-day volume (Figs. 8 and 11c).

8 Conclusions and outlook

Jostedalsbreen is the largest ice mass on the European continent. The state and fate of this and similar ice masses worldwide is of great interest to local residents and stakeholders because of widespread natural and socio-economic implications.

To simulate the future evolution of Jostedalsbreen, we use a 3-d ice-flow model coupled with a surface mass balance model until 2100 and beyond to 2300, allowing us to investigate rapid geometric changes in complex terrain. Historical simulations with the ice-flow model from 1960 to 2020 serve to reproduce present-day conditions in terms of ice geometry and dynamics. This current-state ice cap then provides a vantage point for embarking on annually coupled ice-dynamical and mass balance simulations, fully accounting for the mass-balance elevation feedback.

- We find that Jostedalsbreen is subject to substantial mass loss until 2100, losing about half to two-thirds of its 2020 volume with the medium and high greenhouse-gas emission scenarios RCP4.5 and RCP8.5, respectively. There exists a large spread in simulated mass loss ranging from 12–74% and 53–74%, respectively, depending on the employed climate model, while excluding results from one model which generates unrealistic future winter accumulation. Focusing on results from the mid-range results obtained using the model ECEARTH/CCLM, Jostedalsbreen is projected to lose 49% (RCP4.5) and 63% (RCP8.5) of its present-day ice volume by 2100.
- Most simulations suggest that Jostedalsbreen will split into three separate ice caps over the course of the 21st century. The ice cap is however currently *not* (yet) committed to this breakup. This implies that if greenhouse-gas emissions are reduced over the next decades, the ice cap may evade future breakup and complete disappearance.



- Simulations until the year 2300 show that Jostedalsbreen is currently committed to losing almost half of its current volume by the year 2200. Similarly, prolonging the surface mass balance representative for medium- or high emissions of the end of the 21st century (2081–2100) suggest that the ice cap is bound to shrink by 90% until the year 2300, or disappear by 2200, respectively.
- Tunsbergdalsbreen, the largest glacier basin, appears to be most resilient to future climate change. Its upper reaches are likely to persist well beyond 2100, with 500 m thick ice still present by 2100.
- Experiments beyond 2100 with the climate reversed to that of the present-day indicate that large parts of mass losses until 2100 are irreversible. This is due to significant ice-surface lowering and the associated strong mass-balance elevation feedback, which results in SMB being about 2 m w.e. more negative than that observed in 2001–2020.
- We have demonstrated the importance of accounting for the mass-balance elevation feedback when assessing ice-cap evolution over long timescales. This feedback partly explains why Jostedalsbreen would be difficult to regrow if it disappeared – the current climate of the 2000s is not enough to regenerate the contemporary ice cap.

Jostedalsbreen is likely to survive longer under future warming than many smaller glaciers and ice caps in Scandinavia and worldwide. Future climate conditions determine the SMB over the ice cap, and appears to be the largest source of uncertainty in our projections. Our findings indicate that climate models particularly disagree on future precipitation changes, which affects glacier projections of maritime ice caps like Jostedalsbreen. This calls for further improvement of regional climate models and the representation of precipitation in mountainous areas.

Our detailed, spatially continuous model approach for an ice mass with 81 glacier units, with challenging topography and ice-flow regime, should serve as inspiration for studies of glaciers and ice caps worldwide. Numerical simulations of complex ice masses like Jostedalsbreen will also serve as critical training data for more robust and widely applicable emulators based on machine learning. The diverse characteristics make ice caps ideal yardsticks for such emulators in future studies. Finally, our findings on strong committed mass loss illustrate the need to consider time scales beyond the 21st century, and call for intensified research on irreversible change and committed mass loss.

Code and data availability. ISSM is open-source and can be downloaded from <https://issm.jpl.nasa.gov>. Model scripts will be made available in a public repository upon publication. Climate model projections for 2021–2100 can be downloaded from the Norwegian Center for Climate Services (NCCS) website (<https://klimaservicesenter.no/>; Jacob et al., 2014; Wong et al., 2016).

Video supplement. The following animations are supplied along with the article, and can be downloaded from <https://doi.org/10.5281/zenodo.14764904>.

- Supplementary video 1: Modelled ice-thickness evolution 2021–2100 (ECEARTH/CCLM, RCP4.5).
- Supplementary video 2: Modelled ice-thickness evolution 2021–2100 (ECEARTH/CCLM, RCP8.5).



- Supplementary video 3: Modelled ice-thickness evolution 2101–2300 (Commit4.5).
- Supplementary video 4: Modelled ice-thickness evolution 2101–2300 (Commit8.5).

750 Appendix A: Additional figures




Figure A1. Model mesh on underlying bed topography, as described in Sect. 4.

Author contributions. The study was conceptualised and formulated by HÅ, KHS, TVS and TD. HÅ carried out pre-processing of most input data, conducted dynamical model simulations, analysed model results, wrote most of the manuscript, and made all figures except Fig. 1 (made by BAR) and Figs. 5, 12, A5 (made by KHS). Model development and simulations of surface mass balance were done by KHS, who also pre-processed and analysed climate model data, and wrote substantial parts of the manuscript. HÅ and KHS jointly coded the coupling for dynamics and surface mass balance. TVS and TD contributed with advice on methodology, model strategy, visualisation, and




Figure A2. Simulated historical change (a–c) and future (d–f) evolution with experiment Future4.5-ECC (cf. Table 2), using a 20 m ice thickness threshold (cf. Fig. 2, which uses a 10 m threshold). Recent historical periods of little change (gray), ice-cap growth (blue) and decay (red) are shaded in (a–c). In (c), historical SMB with evolving model area is shown ($\text{SMB}_{\text{transArea}}$), but without elevation changes, since these make little difference (Sect. 5.2). Shown in (f) is SMB with transiently evolving geometry ($\text{SMB}_{\text{transGeom}}$), including the SMB–elevation feedback and area changes.




Figure A3. Model evolution of (a) SMB_{static} , (b) ice volume, and (c) area in experiments aiming to grow Jostedalsbreen from scratch. See Table 2 for experiment details.

comments on scientific writing. MKG and LMA prepared the ice-thickness data. LMA also provided guidance with data on glacier outlines and front positions. BAR and LMA prepared historical data for ice geometry. TS contributed with processing and advice on ice-velocity data. JCY contributed with funding acquisition, project management, and writing/editing. All authors contributed to writing and revising the manuscript, and approved the final version for submission.

760 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This work is a contribution to the JOSTICE project funded by the Research Council of Norway (RCN grant #302458) and ERC-2022-ADG grant agreement No 01096057 GLACMASS from the European Research Council. TS was funded under MASSIVE




Figure A4. Future simulated mean annual SMB_{transGeom}, ELA and accumulation-area ratio (AAR) under (a–c) RCP4.5 and (d–f) RCP8.5 emission pathways, using the climate model ECEARTH/CCLM. Yearly AAR and 10-year running means are shown in (b) and (e). Present-day glacier outlines (2019) are shown with black lines (Andreassen et al., 2022) on the SMB maps.

(RCN #315971). We would like to thank Wai Kwok Wong at the Norwegian Water Resources and Energy Directorate (NVE) for providing input on the selection of climate model and emission scenarios used in this study.




Figure A5. Future changes in (a, b) mean temperature and (c, d) total precipitation in (a, c) winter (Oct–Apr) and (b, d) summer (May–Sep) months for the ensemble of climate model forcing.

765 **References**

Adhikari, S. and Marshall, S. J.: Glacier volume-area relation for high-order mechanics and transient glacier states, *Geophysical Research Letters*, 39, n/a–n/a, <https://doi.org/10.1029/2012GL052712>, 2012.

Andreassen, L. and Elvehøy, H.: Norwegian glacier reference dataset for climate change studies, Tech. Rep. NVE report 33/2021, 2021.

Andreassen, L., Huss, M., Melvold, K., Elvehøy, H., and Winsvold, S.: Ice thickness measurements and volume estimates for glaciers in Norway, *Journal of Glaciology*, 61, 763–775, <https://doi.org/10.3189/2015JoG14J161>, 2015.

Andreassen, L. M., Elvehøy, H., Kjølmoen, B., Engeset, R. V., and Haakensen, N.: Glacier mass-balance and length variation in Norway, *Annals of Glaciology*, 42, 317–325, <https://doi.org/10.3189/172756405781812826>, 2005.

Andreassen, L. M., Elvehøy, H., Kjølmoen, B., and Belart, J. M.: Glacier change in Norway since the 1960s – an overview of mass balance, area, length and surface elevation changes, *Journal of Glaciology*, 66, 313–328, <https://doi.org/10.1017/jog.2020.10>, 2020.

Andreassen, L. M., Nagy, T., Kjølmoen, B., and Leigh, J. R.: An inventory of Norway’s glaciers and ice-marginal lakes from 2018–19 Sentinel-2 data, *Journal of Glaciology*, pp. 1–22, <https://doi.org/10.1017/jog.2022.20>, 2022.



Table A1. Overview of data sources used in this study.

Data type	Aquisition method	Time period	Resolution (m)	Usage	Reference
Subglacial topography	ice radar	2019–2023	10	Input	Gillespie et al. (2024)
Land-surface topography	aerial imagery	2019	10	Input	Andreassen et al. (2023)
Ice-surface topography	aerial imagery	1966	10	Input	Andreassen et al. (2023)
Ice-surface topography	topographic map	1966	50	Input	Paul et al. (2011)
Ice-surface topography	aerial imagery	2019	10	Validation	Andreassen et al. (2023)
Ice-surface velocity	radar+optical satellite imagery	2017–2018	50	Calibration	Millan et al. (2022)
Ice-surface velocity	radar satellite imagery	1996	20	Calibration	this study (Sect. 3.2)
Ice-surface velocity	optical satellite imagery	2015–2018	80	Validation	Nagy and Andreassen (2019)
Glacier outlines	optical satellite imagery	1966, 2019	10	Validation	Winsvold et al. (2014) Andreassen et al. (2022)
Glacier front positions	in-situ	various	1	Validation	Andreassen and Elvehøy (2021) Andreassen et al. (2023) Kjöllmoen et al. (2024)

Andreassen, L. M., Robson, B. A., Sjrursen, K. H., Elvehøy, H., Kjöllmoen, B., and Carrivick, J. L.: Spatio-temporal variability in geometry and geodetic mass balance of Jostedalsbreen ice cap, Norway, *Annals of Glaciology*, pp. 1–18, <https://doi.org/10.1017/aog.2023.70>, publisher: Cambridge University Press, 2023.

780 Aschwanden, A., Aðalgeirsdóttir, G., and Khroulev, C.: Hindcasting to measure ice sheet model sensitivity to initial states, *The Cryosphere*, 7, 1083–1093, <https://doi.org/10.5194/tc-7-1083-2013>, publisher: Copernicus GmbH, 2013.

Aðalgeirsdóttir, G., Gudmundsson, G. H., and Björnsson, H.: Volume sensitivity of Vatnajökull Ice Cap, Iceland, to perturbations in equilibrium line altitude, *Journal of Geophysical Research: Earth Surface*, 110, n/a–n/a, <https://doi.org/10.1029/2005JF000289>, 2005.

785 Aðalgeirsdóttir, G., Aschwanden, A., Khroulev, C., Boberg, F., Mottram, R., Lucas-Picher, P., and Christensen, J.: Role of model initialization for projections of 21st-century Greenland ice sheet mass loss, *Journal of Glaciology*, 60, 782–794, <https://doi.org/10.3189/2014JoG13J202>, 2014.

Bickerton, R. W. and Matthews, J. A.: ‘Little ice age’ variations of outlet glaciers from the jostedalsbreen ice-cap, Southern Norway: A regional lichenometric-dating study of ice-marginal moraine sequences and their climatic significance, *Journal of Quaternary Science*, 8, 45–66, <https://doi.org/10.1002/jqs.3390080105>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/jqs.3390080105>, 1993.

790 Blatter, H.: Velocity and stress fields in grounded glaciers: a simple algorithm for including deviatoric stress gradients, *Journal of Glaciology*, 41, 333–344, <https://doi.org/10.3189/S002214300001621X>, 1995.

Bosson, J. B., Huss, M., Cauvy-Fraunié, S., Clément, J. C., Costes, G., Fischer, M., Poulénard, J., and Arthaud, F.: Future emergence of new ecosystems caused by glacial retreat, *Nature*, 620, 562–569, <https://doi.org/10.1038/s41586-023-06302-2>, publisher: Nature Publishing Group, 2023.

795 Brondex, J., Gillet-Chaulet, F., and Gagliardini, O.: Sensitivity of centennial mass loss projections of the Amundsen basin to the friction law, *The Cryosphere*, 13, 177–195, <https://doi.org/10.5194/tc-13-177-2019>, 2019.



- Budd, W. F., Keage, P. L., and Blundy, N. A.: Empirical Studies of Ice Sliding, *Journal of Glaciology*, 23, 157–170, <https://doi.org/10.3189/S0022143000029804>, 1979.
- Carey, M., Molden, O. C., Rasmussen, M. B., Jackson, M., Nolin, A. W., and Mark, B. G.: Impacts of Glacier Recession and Declining Meltwater on Mountain Societies, *Annals of the American Association of Geographers*, 107, 350–359, <https://doi.org/10.1080/24694452.2016.1243039>, publisher: Taylor & Francis _eprint: <https://doi.org/10.1080/24694452.2016.1243039>, 2017.
- Carrivick, J. L., Andreassen, L. M., Nesje, A., and Yde, J. C.: A reconstruction of Jostedalsbreen during the Little Ice Age and geometric changes to outlet glaciers since then, *Quaternary Science Reviews*, 284, 107 501, <https://doi.org/10.1016/j.quascirev.2022.107501>, 2022.
- Compagno, L., Zekollari, H., Huss, M., and Farinotti, D.: Limited impact of climate forcing products on future glacier evolution in Scandinavia and Iceland, *Journal of Glaciology*, 67, 727–743, <https://doi.org/10.1017/jog.2021.24>, 2021.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Academic Press, 2010.
- Dannevig, H. and Rusdal, T.: Caring for melting glaciers, *Tourism Geographies*, 25, 1679–1695, <https://doi.org/10.1080/14616688.2023.2278762>, 2023.
- de Fleurian, B., Davy, R., and Langebroek, P. M.: Impact of runoff temporal distribution on ice dynamics, *The Cryosphere*, 16, 2265–2283, <https://doi.org/10.5194/tc-16-2265-2022>, publisher: Copernicus GmbH, 2022.
- Eklblom Johansson, F., Bakke, J., Støren, E. N., Gillespie, M. K., and Laumann, T.: Mapping of the Subglacial Topography of Folgefonna Ice Cap in Western Norway—Consequences for Ice Retreat Patterns and Hydrological Changes, *Frontiers in Earth Science*, 10, <https://www.frontiersin.org/article/10.3389/feart.2022.886361>, 2022.
- Elsberg, D. H., Harrison, W. D., Echelmeyer, K. A., and Krimmel, R. M.: Quantifying the effects of climate and surface change on glacier mass balance, *Journal of Glaciology*, 47, 649–658, <https://doi.org/10.3189/172756501781831783>, 2001.
- Erikstad, L. and Sollid, J. L.: Neoglaciation in South Norway using lichenometric methods, *Norsk Geografisk Tidsskrift - Norwegian Journal of Geography*, 40, 85–105, <https://doi.org/10.1080/00291958608552159>, 1986.
- Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus estimate for the ice thickness distribution of all glaciers on Earth, *Nature Geoscience*, 12, 168–173, <https://doi.org/10.1038/s41561-019-0300-3>, 2019.
- Frank, T. and Pelt, W. J. J. v.: Ice volume and thickness of all Scandinavian glaciers and ice caps, *Journal of Glaciology*, pp. 1–14, <https://doi.org/10.1017/jog.2024.25>, 2024.
- Friedl, P., Seehaus, T., and Braun, M.: Global time series and temporal mosaics of glacier surface velocities derived from Sentinel-1 data, *Earth System Science Data*, 13, 4653–4675, <https://doi.org/10.5194/essd-13-4653-2021>, publisher: Copernicus GmbH, 2021.
- Gabbi, J., Carenzo, M., Pellicciotti, F., Bauder, A., and Funk, M.: A comparison of empirical and physically based glacier surface melt models for long-term simulations of glacier response, *Journal of Glaciology*, 60, 1140–1154, <https://doi.org/10.3189/2014JoG14J011>, 2014.
- Giesen, R. H. and Oerlemans, J.: Response of the ice cap Hardangerjøkulen in southern Norway to the 20th and 21st century climates, *The Cryosphere*, 4, 191–213, <https://doi.org/10.5194/tc-4-191-2010>, 2010.
- Gilbert, A., Flowers, G. E., Miller, G. H., Rabus, B. T., Van Wychen, W., Gardner, A. S., and Copland, L.: Sensitivity of Barnes Ice Cap, Baffin Island, Canada, to climate state and internal dynamics: Barnes Ice Cap Stability, *Journal of Geophysical Research: Earth Surface*, 121, 1516–1539, <https://doi.org/10.1002/2016JF003839>, 2016.
- Gillespie, M. K., Andreassen, L. M., Huss, M., de Villiers, S., Sjørnsen, K. H., Aasen, J., Bakke, J., Cederstrøm, J. M., Elvehøy, H., Kjølmoen, B., Loe, E., Meland, M., Melvold, K., Nerhus, S. D., Røthe, T. O., Støren, E. W. N., Øst, K., and Yde, J. C.: Ice thickness and bed



- topography of Jostedalsbreen ice cap, Norway, *Earth System Science Data*, 16, 5799–5825, <https://doi.org/10.5194/essd-16-5799-2024>,
 835 publisher: Copernicus GmbH, 2024.
- Gillet-Chaulet, F., Gagliardini, O., Seddik, H., Nodet, M., Durand, G., Ritz, C., Zwinger, T., Greve, R., and Vaughan, D. G.: Greenland ice
 sheet contribution to sea-level rise from a new-generation ice-sheet model, *The Cryosphere*, 6, 1561–1576, <https://doi.org/10.5194/tc-6-1561-2012>, 2012.
- Gjerde, M., Hoel, O. L., and Nesje, A.: The ‘Little Ice Age’ advance of Nigardsbreen, Norway: A cross-disciplinary revision of the chrono-
 840 logical framework, *The Holocene*, p. 09596836231185830, <https://doi.org/10.1177/09596836231185830>, publisher: SAGE Publications
 Ltd, 2023.
- Goldberg, D. N., Heimbach, P., Joughin, I., and Smith, B.: Committed retreat of Smith, Pope, and Kohler Glaciers over the next 30 years
 inferred by transient model calibration, *The Cryosphere*, 9, 2429–2446, <https://doi.org/10.5194/tc-9-2429-2015>, 2015.
- Hacker, B. R., Andersen, T. B., Johnston, S., Kylander-Clark, A. R. C., Peterman, E. M., Walsh, E. O., and Young, D.: High-temperature
 845 deformation during continental-margin subduction & exhumation: The ultrahigh-pressure Western Gneiss Region of Norway, *Tectono-
 physics*, 480, 149–171, <https://doi.org/10.1016/j.tecto.2009.08.012>, 2010.
- Hanssen-Bauer, I., Førland, E., Haddeland, I., Hisdal, H., Mayer, S., Nesje, A., Nilsen, J., Sandven, S. A., Sandø, A., Sorteberg, A., and
 Ådlandsvi, B.: Climate in Norway 2100, Norwegian Center for Climate Services (NCCS) Report 1/2017, Norwegian Environment Agency
 (Miljødirektoratet), 2017.
- 850 Harrison, W. D., Elsberg, D. H., Echelmeyer, K. A., and Krimmel, R. M.: On the characterization of glacier response by a single time-scale,
Journal of Glaciology, 47, 659–664, <https://doi.org/10.3189/172756501781831837>, 2001.
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I., Brun, F., and Käab, A.:
 Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>,
 2021.
- 855 Huiban, F., Millan, R., Kjeldsen, K. K., Andresen, C. S., Dømgård, M., Dehecq, A., Brunt, S., Khan, S. A., Mouginot, J., and Bjørk,
 A. A.: Regional ice flow piracy following the collapse of Midgaard Glacier in Southeast Greenland, *Nature Communications*, 15, 9976,
<https://doi.org/10.1038/s41467-024-54045-z>, publisher: Nature Publishing Group, 2024.
- Huss, M. and Hock, R.: A new model for global glacier change and sea-level rise, *Frontiers in Earth Sciences*, 3,
<https://doi.org/10.3389/feart.2015.00054>, 2015.
- 860 Huss, M. and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8, 135–140,
<https://doi.org/10.1038/s41558-017-0049-x>, 2018.
- Huss, M., Hock, R., Bauder, A., and Funk, M.: Conventional versus reference-surface mass balance, *Journal of Glaciology*, 58, 278–286,
<https://doi.org/10.3189/2012JoG11J216>, 2012.
- Huss, M., Bookhagen, B., Huggel, C., Jacobsen, D., Bradley, R., Clague, J., Vuille, M., Buytaert, W., Cayan, D., Greenwood, G., Mark,
 865 B., Milner, A., Weingartner, R., and Winder, M.: Toward mountains without permanent snow and ice, *Earth’s Future*, 5, 418–435,
<https://doi.org/10.1002/2016EF000514>, 2017.
- Hutchinson, M. F., Xu, T., Stein, J. A., et al.: Recent progress in the ANUDEM elevation gridding procedure, *Geomorphometry*, 2011, 19–22,
 2011.
- Iken, A.: The Effect of the Subglacial Water Pressure on the Sliding Velocity of a Glacier in an Idealized Numerical Model, *Journal of*
 870 *Glaciology*, 27, 407–421, <https://doi.org/10.3189/S0022143000011448>, 1981.



- Ismail, M. F., Bogacki, W., Disse, M., Schäfer, M., and Kirschbauer, L.: Estimating degree-day factors of snow based on energy flux components, *The Cryosphere*, 17, 211–231, <https://doi.org/10.5194/tc-17-211-2023>, 2023.
- Jacob, D., Petersen, J., Eggert, B., Alias, A., Christensen, O., Bouwer, L., Braun, A., Colette, A., Déqué, M., Georgievski, G., Georgopoulou, E., Gobiet, A., Menut, L., Nikulin, G., Haensler, A., Hempelmann, N., Jones, C., Keuler, K., Kovats, S., and Yiou, P.: EURO-CORDEX: New high-resolution climate change projections for European impact research, *Regional Environmental Change*, 14, <https://doi.org/10.1007/s10113-013-0499-2>, 2014.
- Jostedalsbreen nasjonalparkstyre: Besøksstrategi Jostedalsbreen nasjonalpark 2021-2027, Tech. rep., https://www.nasjonalparkstyre.no/uploads/files_jostedalsbreen/Besoksstrategi-Jostedalsbreen-2021-2027-godkjent.pdf, 2021.
- Jouvet, G., Huss, M., Blatter, H., Picasso, M., and Rappaz, J.: Numerical simulation of Rhonegletscher from 1874 to 2100, *Journal of Computational Physics*, 228, 6426–6439, <https://doi.org/10.1016/j.jcp.2009.05.033>, 2009.
- Jouvet, G., Huss, M., Funk, M., and Blatter, H.: Modelling the retreat of Grosser Aletschgletscher, Switzerland, in a changing climate, *Journal of Glaciology*, 57, 1033–1045, <https://doi.org/10.3189/002214311798843359>, 2011.
- Jóhannesson, T., Raymond, C., and Waddington, E.: Time-Scale for Adjustment of Glaciers to Changes in Mass Balance, *Journal of Glaciology*, 35, 355–369, <https://doi.org/10.3189/S002214300000928X>, 1989.
- Jóhannesson, T., Sigurdsson, O., Laumann, T., and Kennett, M.: Degree-day glacier mass-balance modelling with applications to glaciers in Iceland, Norway and Greenland, *Journal of Glaciology*, 41, 345–358, <https://doi.org/10.3189/S0022143000016221>, 1995.
- Kjøllmoen, B., Andreassen, L. M., and Elvehøy, H.: Glaciological investigations in Norway 2023, NVE Rapport 22-2024, Tech. Rep. 22-2024, http://publikasjoner.nve.no/rapport/2024/rapport2024_22.pdf, 2024.
- Larour, E., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM): ICE SHEET SYSTEM MODEL, *Journal of Geophysical Research: Earth Surface*, 117, n/a–n/a, <https://doi.org/10.1029/2011JF002140>, 2012.
- Laumann, T. and Nesje, A.: The impact of climate change on future frontal variations of Briksdalsbreen, western Norway, *Journal of Glaciology*, 55, 789–796, <https://doi.org/10.3189/002214309790152366>, 2009.
- Laumann, T. and Nesje, A.: Spørteggbreen, western Norway, in the past, present and future: Simulations with a two-dimensional dynamical glacier model, *The Holocene*, 24, 842–852, <https://doi.org/10.1177/0959683614530446>, 2014.
- Leysinger Vieli, G. J.-M. C. and Gudmundsson, G. H.: On estimating length fluctuations of glaciers caused by changes in climatic forcing, *Journal of Geophysical Research: Earth Surface*, 109, <https://doi.org/10.1029/2003JF000027>, 2004.
- Lussana, C.: seNorge observational gridded datasets, seNorge_2018, versions 21.09 and 21.10, METreport 07/2021, The Norwegian Meteorological Institute (MET Norway), Oslo, Norway, 2021.
- Lussana, C., Tveito, O., Dobler, A., and Tunheim, K.: seNorge_2018, daily precipitation, and temperature datasets over Norway, *Earth System Science Data*, 11, 1531–1551, <https://doi.org/10.5194/essd-11-1531-2019>, 2019.
- Marshall, S. J.: Simulation of Vatnajökull ice cap dynamics, *Journal of Geophysical Research*, 110, F03009, <https://doi.org/10.1029/2004JF000262>, 2005.
- Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers, *The Cryosphere*, 6, 1295–1322, <https://doi.org/10.5194/tc-6-1295-2012>, publisher: Copernicus GmbH, 2012.
- Maussion, F., Butenko, A., Champollion, N., Dusch, M., Eis, J., Fourteau, K., Gregor, P., Jarosch, A. H., Landmann, J., Oesterle, F., Recinos, B., Rothenpieler, T., Vlug, A., Wild, C. T., and Marzeion, B.: The Open Global Glacier Model (OGGM) v1.1, *Geoscientific Model Development*, 12, 909–931, <https://doi.org/10.5194/gmd-12-909-2019>, publisher: Copernicus GmbH, 2019.



- Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500, *Geoscientific Model Development*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, publisher: Copernicus GmbH, 2020.
- Meur, E. L. and Vincent, C.: A two-dimensional shallow ice-flow model of Glacier de Saint-Sorlin, France, *Journal of Glaciology*, 49, 527–538, <https://doi.org/10.3189/172756503781830421>, 2003.
- Meur, E. L., Gagliardini, O., Zwinger, T., and Ruokolainen, J.: Glacier flow modelling: a comparison of the Shallow Ice Approximation and the full-Stokes solution, *Comptes Rendus. Physique*, 5, 709–722, <https://doi.org/10.1016/j.crh.2004.10.001>, 2004.
- Millan, R., Mouginot, J., Rabatel, A., and Morlighem, M.: Ice velocity and thickness of the world’s glaciers, *Nature Geoscience*, 15, 124–129, <https://doi.org/10.1038/s41561-021-00885-z>, number: 2 Publisher: Nature Publishing Group, 2022.
- Milner, A. M., Khamis, K., Battin, T. J., Brittain, J. E., Barrand, N. E., Füreder, L., Cauvy-Fraunié, S., Gíslason, G. M., Jacobsen, D., Hannah, D. M., Hodson, A. J., Hood, E., Lencioni, V., Ólafsson, J. S., Robinson, C. T., Tranter, M., and Brown, L. E.: Glacier shrinkage driving global changes in downstream systems, *Proceedings of the National Academy of Sciences*, 114, 9770–9778, <https://doi.org/10.1073/pnas.1619807114>, publisher: Proceedings of the National Academy of Sciences, 2017.
- Mohr, M.: New Routines for Gridding of Temperature and Precipitation Observations for “seNorge. no”, Tech. Rep. 08/2008, Norwegian Meteorological Institute, Oslo, Norway, 2008.
- Morlighem, M., Rignot, E., Seroussi, H., Larour, E., Ben Dhia, H., and Aubry, D.: Spatial patterns of basal drag inferred using control methods from a full-Stokes and simpler models for Pine Island Glacier, West Antarctica, *Geophysical Research Letters*, 37, n/a–n/a, <https://doi.org/10.1029/2010GL043853>, 2010.
- Nagy, T. and Andreassen, L.: Glacier surface velocity mapping with Sentinel-2 imagery in Norway. NVE Report 37, Tech. rep., 2019.
- Nesje, A. and Matthews, J. A.: The Briksdalsbre Event: A winter precipitation-induced decadal-scale glacial advance in southern Norway in the 1990s and its implications, *The Holocene*, 22, 249–261, <https://doi.org/10.1177/0959683611414938>, 2012.
- Nesje, A., Lie, O., and Dahl, S. O.: Is the North Atlantic Oscillation reflected in Scandinavian glacier mass balance records?, *Journal of Quaternary Science*, 15, 587–601, [https://doi.org/10.1002/1099-1417\(200009\)15:6<587::AID-JQS533>3.0.CO;2-2](https://doi.org/10.1002/1099-1417(200009)15:6<587::AID-JQS533>3.0.CO;2-2), 2000.
- Noël, B., van de Berg, W. J., Machguth, H., Lhermitte, S., Howat, I., Fettweis, X., and van den Broeke, M. R.: A daily, 1 km resolution data set of downscaled Greenland ice sheet surface mass balance (1958–2015), *The Cryosphere*, 10, 2361–2377, <https://doi.org/10.5194/tc-10-2361-2016>, 2016.
- Oerlemans, J.: A flowline model for Nigardsbreen, Norway: projection of future glacier length based on dynamic calibration with the historic record, *Annals of Glaciology*, 24, 382–389, <https://doi.org/10.3189/S0260305500012489>, 1997.
- Pattyn, F.: A new three-dimensional higher-order thermomechanical ice sheet model: Basic sensitivity, ice stream development, and ice flow across subglacial lakes, *Journal of Geophysical Research: Solid Earth*, 108, <https://doi.org/10.1029/2002JB002329>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2002JB002329>, 2003.
- Paul, F., Andreassen, L. M., and Winsvold, S. H.: A new glacier inventory for the Jostedalsgreen region, Norway, from Landsat TM scenes of 2006 and changes since 1966, *Annals of Glaciology*, 52, 153–162, <https://doi.org/10.3189/172756411799096169>, 2011.
- Pollard, D. and DeConto, R. M.: A simple inverse method for the distribution of basal sliding coefficients under ice sheets, applied to Antarctica, *The Cryosphere*, 6, 953–971, <https://doi.org/10.5194/tc-6-953-2012>, 2012.



- Réveillet, M., Six, D., Vincent, C., Rabatel, A., Dumont, M., Lafaysse, M., Morin, S., Vionnet, V., and Litt, M.: Relative performance of empirical and physical models in assessing the seasonal and annual glacier surface mass balance of Saint-Sorlin Glacier (French Alps), *The Cryosphere*, 12, 1367–1386, <https://doi.org/10.5194/tc-12-1367-2018>, 2018.
- Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., Farinotti, D., Menounos, B., and McNabb, R. W.: Global glacier change in the 21st century: Every increase in temperature matters, *Science*, 379, 78–83, <https://doi.org/10.1126/science.abo1324>, publisher: American Association for the Advancement of Science, 2023.
- Santos, T. D. D., Morlighem, M., Simões, J. C., and Devloo, P. R. B.: Sensitivity analysis of a King George Island outlet glacier, South Shetlands, Antarctica, *Anais da Academia Brasileira de Ciências*, 95, e20210560, <https://doi.org/10.1590/0001-3765202320210560>, publisher: Academia Brasileira de Ciências, 2023.
- Schaffer, N., Copland, L., Zdanowicz, C., and Hock, R.: Modeling the surface mass balance of Penny Ice Cap, Baffin Island, 1959–2099, *Annals of Glaciology*, 64, 330–342, <https://doi.org/10.1017/aog.2023.68>, 2023.
- Schmidt, L. S., Aðalgeirsdóttir, G., Pálsson, F., Langen, P. L., Guðmundsson, S., and Björnsson, H.: Dynamic simulations of Vatnajökull ice cap from 1980 to 2300, *Journal of Glaciology*, 66, 97–112, <https://doi.org/10.1017/jog.2019.90>, 2020.
- Seier, G., Abermann, J., Andreassen, L. M., Carrivick, J. L., Kielland, P. H., Löffler, K., Nesje, A., Robson, B. A., Røthe, T. O., Scheiber, T., Winkler, S., and Yde, J. C.: Glacier thinning, recession and advance, and the associated evolution of a glacial lake between 1966 and 2021 at Austerdalsbreen, western Norway, *Land Degradation & Development*, 35, 394–414, <https://doi.org/10.1002/ldr.4923>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/ldr.4923>, 2024.
- Sjursen, K. H., Dunse, T., Tambue, A., Schuler, T. V., and Andreassen, L. M.: Bayesian parameter estimation in glacier mass-balance modelling using observations with distinct temporal resolutions and uncertainties, *Journal of Glaciology*, p. 1–20, <https://doi.org/10.1017/jog.2023.62>, 2023.
- Sjursen, K. H., Dunse, T., Schuler, T. V., Andreassen, L. M., and Åkesson, H.: Spatiotemporal mass balance variability of Jostedal-breen Ice Cap, Norway, revealed by a temperature-index model using Bayesian inference, *Annals of Glaciology*, 66, 1–18, <https://doi.org/10.1017/aog.2024.41>, 2025.
- Troch, M., Åkesson, H., Cuzzzone, J. K., and Bertrand, S.: Precipitation drives western Patagonian glacier variability and may curb future ice mass loss, *Scientific Reports*, 14, 26744, <https://doi.org/10.1038/s41598-024-77486-4>, publisher: Nature Publishing Group, 2024.
- Tvede, A.: Folgefonni, en glasiologisk avvik, *Naturen*, 97, 11–16, 1973.
- Tvede, A. and Liestøl, O.: Blomsterskardbreen, Folgefonni, mass balance and recent fluctuations, *Nor. Polarinst. Arbok*, pp. 225–233, 1976.
- Tvede, A. M.: Floods Caused by a Glacier-Dammed Lake at the Folgefonni Ice Cap, Norway, *Annals of Glaciology*, 13, 262–264, <https://doi.org/10.3189/S0260305500008016>, 1989.
- United Nations Environment Programme: Emissions Gap Report 2024: No more hot air ... please! With a massive gap between rhetoric and reality, countries draft new climate commitments., Tech. rep., Nairobi, <https://doi.org/10.59117/20.500.11822/46404>, section: publications, 2024.
- van Pelt, W. J. J., Schuler, T. V., Pohjola, V. A., and Pettersson, R.: Accelerating future mass loss of Svalbard glaciers from a multi-model ensemble, *Journal of Glaciology*, 67, 485–499, <https://doi.org/10.1017/jog.2021.2>, 2021.
- Van Tricht, L. and Huybrechts, P.: Modelling the historical and future evolution of six ice masses in the Tien Shan, Central Asia, using a 3D ice-flow model, *The Cryosphere*, 17, 4463–4485, <https://doi.org/10.5194/tc-17-4463-2023>, publisher: Copernicus GmbH, 2023.
- Verfaillie, D., Charton, J., Schimmelpfennig, I., Stroebele, Z., Jomelli, V., Bétard, F., Favier, V., Caverio, J., Berthier, E., Goosse, H., Rinterknecht, V., Legentil, C., Charrassin, R., Aumaître, G., Bourlès, D. L., and Keddadouche, K.: Evolution of the Cook Ice Cap (Kerguelen



- Islands) between the last centuries and 2100 ce based on cosmogenic dating and glacio-climatic modelling, *Antarctic Science*, 33, 301–317, <https://doi.org/10.1017/S0954102021000080>, 2021.
- 985 Wangenstein, B., Tønsberg, O. M., Kääb, A., Eiken, T., and Hagen, J. O.: Surface Elevation Change and High Resolution Surface Velocities for Advancing Outlets of Jostedalsbreen, *Geografiska Annaler: Series A, Physical Geography*, 88, 55–74, <https://doi.org/10.1111/j.0435-3676.2006.00283.x>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.0435-3676.2006.00283.x>, 2006.
- Welty, E., Zemp, M., Navarro, F., Huss, M., Fürst, J. J., Gärtner-Roer, I., Landmann, J., Machguth, H., Naegeli, K., Andreassen, L. M., 990 Farinotti, D., Li, H., and Contributors, G.: Worldwide version-controlled database of glacier thickness observations, *Earth System Science Data*, 12, 3039–3055, <https://doi.org/10.5194/essd-12-3039-2020>, publisher: Copernicus GmbH, 2020.
- Winsvold, S. H., Andreassen, L. M., and Kienholz, C.: Glacier area and length changes in Norway from repeat inventories, *The Cryosphere*, 8, 1885–1903, 2014.
- Wong, W. K., Haddeland, I., Lawrence, D., and Beldring, S.: Gridded 1 x 1 km climate and hydrological projections for Norway, NVE report 995 59/2016, Norwegian Water Resources and Energy Directorate (NVE), Oslo, Norway, 2016.
- Zekollari, H., Huybrechts, P., Noël, B., van de Berg, W. J., and van den Broeke, M. R.: Sensitivity, stability and future evolution of the world’s northernmost ice cap, Hans Tausen Iskappe (Greenland), *The Cryosphere*, 11, 805–825, <https://doi.org/10.5194/tc-11-805-2017>, publisher: Copernicus GmbH, 2017.
- Zekollari, H., Huss, M., Farinotti, D., and Lhermitte, S.: Ice-Dynamical Glacier Evolution Modeling—A Review, *Reviews of Geophysics*, 1000 60, e2021RG000754, <https://doi.org/10.1029/2021RG000754>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021RG000754>, 2022.
- Ziemen, F. A., Hock, R., Aschwanden, A., Khroulev, C., Kienholz, C., Melkonian, A., and Zhang, J.: Modeling the evolution of the Juneau Icefield between 1971 and 2100 using the Parallel Ice Sheet Model (PISM), *Journal of Glaciology*, 62, 199–214, <https://doi.org/10.1017/jog.2016.13>, 2016.
- 1005 Åkesson, H., Nisancioglu, K. H., Giesen, R. H., and Morlighem, M.: Simulating the evolution of Hardangerjøkulen ice cap in southern Norway since the mid-Holocene and its sensitivity to climate change, *The Cryosphere*, 11, 281–302, <https://doi.org/10.5194/tc-11-281-2017>, 2017.
- Åkesson, H., Morlighem, M., Nisancioglu, K. H., Svendsen, J. I., and Mangerud, J.: Atmosphere-driven ice sheet mass loss paced by topography: Insights from modelling the south-western Scandinavian Ice Sheet, *Quaternary Science Reviews*, 195, 32–47, 1010 <https://doi.org/10.1016/j.quascirev.2018.07.004>, 2018.
- Åkesson, H., Morlighem, M., O’Regan, M., and Jakobsson, M.: Future Projections of Petermann Glacier Under Ocean Warming Depend Strongly on Friction Law, *Journal of Geophysical Research: Earth Surface*, 126, <https://doi.org/10.1029/2020JF005921>, 2021.