

Bottom topography effects on the internal wave climate in the Ionian Sea

Florian Kokoszka^{1,2}, Stefania Sparnocchia³, Davide Cavaliere⁴, Vincenzo Artale^{4,5}, Mireno Borghini⁶, Beatrice Giambenedetti⁵, Federico Falcini^{4*}

¹Istituto di Scienze Marine, Consiglio Nazionale delle Ricerche, Naples 34149, Italy

²Stazione Zoologica Anton Dohrn, Villa Comunale, Naples 80121, Italy

³Istituto di Scienze Marine, Consiglio Nazionale delle Ricerche, Trieste 34149, Italy

⁴Istituto di Scienze Marine, Consiglio Nazionale delle Ricerche, Roma 00133, Italy

⁵Istituto Nazionale di Geofisica e Vulcanologia, Roma 00143, Italy

⁶Istituto di Scienze Marine, Consiglio Nazionale delle Ricerche, Lerici 19032, Italy

* Correspondence:

Federico Falcini

federico.falcini@cnr.it

Keywords: Ionian Sea, internal waves, shear and strain ratio, seafloor topography, finescale parameterization.

Abstract

The abyssal Ionian Sea is a deep region of interest for the entire ocean circulation of the Mediterranean Sea, since it plays an important role in the ventilation processes of the whole basin. Here we investigate spatial patterns of internal wave climate over the bottom of the Ionian sub-basin. To identify regional features of the internal wave field in terms of vertical shear and strain, we analyze LADCP and CTD profiles, measured across the basin in 2007, covering various seafloor morphologies (shelf, shelf break, and abyssal plain). Our results show that increasing seafloor roughness reduces the absolute values of shear-to-strain ratio, a pattern also influenced by correlations between slope and roughness. Roughness appears to constrain waves toward higher frequencies, with high shear-to-strain ratios associated with lower frequencies and flatter propagation angles, and low ratios linked to higher frequencies and steeper beams. Spectral analyses indicate that rougher regions enhance strain variance at small vertical scales while reducing shear variance at larger scales, leading to flatter shear spectra in the low-wavenumber band. Together, these findings suggest that roughness redistributes energy from large-scale (low-mode, low-vertical-wavenumber internal waves with vertical scales $O(10^2\text{--}10^3\text{ m})$) toward small-scale (high-vertical-wavenumber internal waves with vertical scales $O(10\text{--}10^1\text{ m})$), fundamentally altering the balance of internal wave energy across scales. These results provide useful knowledge for ad hoc finescale parameterization based on seafloor topography.

1 Introduction

The abyssal circulation, when particularly constrained by seabed topography, can bring to the generation of internal waves (IW) that enhance finescale shear (i.e., vertical variation of the horizontal velocity) and strain (i.e., vertical gradient of isopycnal displacements) over rough bathymetry (Polzin et al., 1997). The resulting small-scale mixing due to breaking of IWs is, therefore, likely responsible for vertical energy transport, which contributes to ocean ventilation (Munk, 1966; Osborne and Burch, 1980; Walin, 1982; Wunsch and Ferrari, 2004; Garrett and Kunze, 2007; Nikurashin and Ferrari, 2013; Ferrari, 2014; Ferrari et al., 2016; Cavaliere et al., 2021; La Forgia et al., 2021).

Despite its thorough implications in the ocean circulation, the relationship between the intensity of overturning circulation and deep, IW-induced mixing rates is not yet fully understood (Garrett and Laurent, 2002; Ferrari, 2014). Paucity of deep ocean measurements hampers our knowledge on abyssal mixing processes and, consequently, ocean heat content variability (Munk and Wunsch, 1998; Ferrari and Wunsch, 2009; Nikurashin and Ferrari, 2011; Waterhouse et al., 2014; de Lavergne et al., 2016; MacKinnon et al., 2017; Ferron et al., 2017; Artale et al., 2018). On the other hand, it is quite established by now that rough bathymetry leads to enhanced IW generation, scattering, and breaking, and thus increased levels of turbulent dissipation (Polzin et al., 1997; Ledwell et al., 2000; St. Laurent et al., 2012; Ferrari et al., 2016; Mashayek et al., 2017). The analysis of IW-field characteristics from microstructure and fine-structure observations suggests that bottom-generated IWs play a major role in determining the spatial distribution of turbulent dissipation (Sheen et al., 2013; Kunze et al., 2006; Ferron et al., 2014; Ferron et al., 2016).

Recent observations indicate that IW field characteristics, shaped by local topography and forcing, show significant variability not only in amplitude but also in frequency composition, affecting the structure and energy distribution of the wave packets and the resulting turbulent mixing (Chinn et al., 2016). In particular, Chinn et al. (2016) explicitly investigates spatial and temporal variability of the shear-to-strain ratio using moored profilers. These authors show that the ratio varies over more than an order of magnitude (≈ 0.8 – 10), it is often controlled by either shear or strain variability (site-dependent), and is strongly influenced by local forcing and topography. These considerations inspired us to investigate the potential role of IWs in mixing processes within the eastern part of the Mediterranean basin, where synchronous Lowered Acoustic Doppler Current Profiler (LADCP) and Conductivity, Temperature, Depth (CTD) data were available.

1.1 The regional oceanographic context

The Mediterranean Sea is often considered as a “miniature ocean”, due to its morpho-bathymetric characteristics and its thermohaline cell, where cold and/or salty waters sink during winter at specific regions and, subsequently, spread to intermediate and deep, near-bottom layers (Wüst, 1961; Bergamasco and Malanotte-Rizzoli, 2010; Millot and Taupier-Letage, 2005; Somot et al., 2006; Tsimplis et al., 2006; Schroeder et al., 2016; Artale et al., 2018). Moreover, among the Mediterranean-type climates, i.e., those defined by temperate, wet winters and hot/warm, dry summers over the western edges of five continents, the Mediterranean is the only one in which the atmospheric variability produces an intense deep-water formation (Seager et al., 2019).

Several analyses of the Mediterranean overturning circulation showed that this marginal basin is well ventilated, in comparison to the world ocean (Wüst, 1961; Malanotte-Rizzoli and Hecht, 1988; Pinardi and Masetti, 2000; Theocharis et al., 2002; Artale et al., 2006; Malanotte-Rizzoli et al., 2014; Schneider et al., 2014). Negative hydrological balance and cold-dry winds, blowing during the winter season, make the Mediterranean Sea an extraordinary place of intense ocean water mass transformation (Pinardi and Masetti, 2000), and thus, an ocean laboratory to study the main processes involved in the global ocean circulation.

Surface water from the Atlantic Ocean enters the Mediterranean basin through the Strait of Gibraltar (Figure 1a), forming a shallow overturning circulation at the upper layer. Flowing eastward, along the African coasts, this Atlantic water increases its salinity and density, then sinking in the Levantine basin in the eastern Mediterranean and forming the Levantine Intermediate Water (LIW). The LIW flows back into the opposite direction, mixing with the surrounding water mass found along its travel back and leaving the Mediterranean Sea through the Strait of Gibraltar, which works as a salt-valve (Artale et al., 2006). Deep water exchanges between the western and the eastern Mediterranean basins occur within the shallow Sicily Channel (Figure 1a), which makes the deep-water circulation of the two sub-basins rather independent from each other (Sparnocchia et al., 1999; Napolitano et al., 2003; Béranger et al., 2004; Schroeder et al., 2006). In the eastern basin, the Eastern Mediterranean Deep Water (EMDW) is formed, alternately, in separate regions of the Eastern Mediterranean Sea (Figure 1a), i.e., the southern Adriatic Sea and the Aegean Sea (Gačić et al., 2010). This cold and high salinity winter water, after deep convection, flows into the Ionian Sea (Wüst, 1961; Roether and Schlitzer, 1991; Schlitzer et al., 1991; Roether et al., 1996; Bensi et al., 2013a; Bensi et al., 2013b; Bellacicco et al., 2016), i.e., the deepest sub-basin of the Mediterranean Sea, which has a maximum depth close to the mean depth of the global ocean (Figure 1a). Such a thermohaline circulation is, therefore, characterized by a superposition of intermediate and deep, zonal and meridional overturning cells (Bergamasco and Malanotte-Rizzoli, 2010).

In the Ionian Sea boundary currents show intense downwelling due to their interaction with the topographic constraints while the deepest part of the basin is characterized by extremely weak stratification (Waldman et al., 2018), which induces a deepening and barotropization of the flow (Holte and Straneo, 2017; Send and Testor, 2017). These deep waters are characterized by IW-turbulence and mixing (van Haren and Gostiaux, 2011). Moreover, the Ionian Sea shows a semi-diurnal lunar tidal (M2) and a dominant motion that is around the inertial frequency, mainly as freely propagating IWs (Gerkema and Zimmerman, 2008). For this reason, the Ionian Sea constitutes a unique example for studying the fundamental dynamics and processes of the abyssal layers in the Mediterranean basin, also in the light of turbulent mixing due to IW breaking (Rubino et al., 2012; Artale et al., 2018; Giambenedetti et al., 2024).

1.2 Internal waves and internal tides in the Eastern Mediterranean Sea

Internal waves constitute a fundamental component of the Mediterranean Sea dynamics and play a key role in vertical energy transfer and in the long-term redistribution of heat and tracers. The semi-enclosed geometry of the Mediterranean, its mid-latitude setting, and the presence of steep and irregular bathymetry strongly shape the internal wave field, distinguishing it from that of the open ocean (Millot and Taupier-Letage, 2005; Schroeder et al., 2016).

In particular, the Eastern Mediterranean is characterized by weak deep stratification, quasi-homogeneous abyssal layers, and pronounced topographic gradients, conditions that make this sub-basin highly sensitive to internal wave processes and their variability (van Haren and Gostiaux, 2011). Despite the very low buoyancy frequencies typical of its deep layers, long-term high-resolution temperature measurements revealed intense internal wave activity in the deep Ionian Sea: vertical displacements reached several hundreds of meters, with significant vertical velocities occurring at near-inertial frequencies (van Haren and Gostiaux, 2011). These observations further showed that energetic internal waves advected through abyssal waters can generate strong vertical shear and episodic overturning, indicating that internal wave-induced mixing remains effective even under extremely weak stratification. Weak stratification does not inhibit internal wave dynamics; rather, it modifies their geometry and frequency content, allowing motions with substantial vertical components to develop in the deep basin (van Haren and Millot, 2004; van Haren and Millot, 2005).

Evidence of energetic internal wave variability has also been reported in other Mediterranean sub-basins, such as the Western Mediterranean, where studies in the Tyrrhenian Sea documented active internal wave fields associated with mesoscale circulation and boundary processes (van Haren, 2004; Buffett et al., 2017). Although these studies focus on different hydrographic and dynamical settings, they support the broader view that internal waves remain dynamically relevant throughout the Mediterranean Sea, including deep environments.

In addition to broadband internal wave activity, recent studies have highlighted the importance of internal tides as a persistent and organized contribution to the Eastern and central Mediterranean internal wave climate. Although barotropic tides are relatively weak in the Mediterranean Sea, their interaction with steep bathymetric features, particularly in straits and along major topographic gradients, can generate energetic baroclinic internal tides (Artale et al., 1989; Oddo et al., 2023). Observational and modeling studies in the central Mediterranean have shown that diurnal internal tides, often with Kelvin-like characteristics, are generated at the transition between shallow regions of the Sicily Channel and the deep Ionian basin, propagate away from their generation sites, and contribute significantly to the internal wave energy budget while inducing strong vertical shear (Oddo et al., 2023). Direct observations in the deep Ionian Sea further indicate that baroclinic internal tides penetrate into abyssal layers, modulating temperature and velocity fields even under weak stratification (Giambenedetti et al., 2023).

This body of evidence defines a Mediterranean internal wave climate that is both spatially heterogeneous and dynamically rich, shaped by the interplay between stratification, bathymetry, and tidal forcing. Within this framework, the Ionian Sea emerges as a key region where internal wave activity intersects with large-scale circulation variability. The hydrographic structure of Atlantic Water, Levantine Intermediate Water, and deeper water masses controls internal wave propagation paths and vertical structure (Budillon et al., 2010), while variability in this structure directly affects the efficiency and vertical reach of internal wave-induced mixing, particularly along continental slopes and over irregular bathymetry. On longer timescales, decadal changes in water mass properties further modulate the background conditions for internal wave dynamics and internal redistribution of momentum and tracers (Gačić et al., 2010).

When interacting with rough topography, internal waves and internal tides have been linked to the redistribution of heat and tracers in abyssal layers, contributing to the observed variability of heat content and influencing long-term basin-scale circulation (Artale et al., 2018). This mechanism becomes particularly relevant during periods of reduced deep-water formation, when internal mixing plays an increasingly important role in controlling abyssal ventilation. Internal tides, in particular, represent a continuous source of energy that sustains mixing in deep layers, complementing episodic convective events and reinforcing the coupling between the deep Ionian interior and boundary regions (Giambenedetti et al., 2023; Oddo et al., 2023). Overall, this internal wave climate provides the dynamical background against which internal wave-driven mixing processes operate and contribute to the thermohaline structure and long-term evolution of the Ionian Sea and the broader Eastern Mediterranean basins.

1.3 The KM3NeT project and a preliminary study on seafloor-induced enhancement of isopycnal vertical strain

An intense observational activity was conducted in the Ionian Sea in the framework of the Cubic Kilometre Neutrino Telescope (KM3NeT) project, from April 2006 to May 2009 (Kats et al., 2006). During the KM3Net cruise (July 2007), hydrographic measurements were spread over a large area, including a section that almost synoptically spanned the central Ionian Sea, from Sicily to Greece. On the western side and up to the central abyssal plane, a marked stratification was found in the deep layers, with a warmer and saltier water on the bottom, below a fresher and slightly colder layer (located approximately at about 2750 m), probably due to deep water formation processes at different times (Sparnocchia et al., 2011). Deep current meter measurements collected in the westernmost part in the period 2007–2008 showed a quite energetic and impulsive abyssal circulation pattern, strongly affected by the bathymetric constraints, with maximum velocities up to 15 cm/s and cyclonic and anti-cyclonic mesoscale structures, with a period of 5-to-11 days, superimposed on the background geostrophic flow (Rubino et al., 2012; Meccia et al., 2015).

Artale et al. (2018), by analyzing water column characteristics of the Ionian Sea, observed during the last three decades, focused on the hydrological processes occurring at the bottom layer. They found that the ocean circulation in this abyssal plain is strongly affected by an interplay between advection and diffusion. The progressive warming and salinification of this sub-basin produced warmer near-bottom waters, causing an anomalous heat storage of $\sim 1.6 \text{ W/m}^2$, i.e., a value three times larger than the equivalent climate trend occurring in the same period at global scale (Bindoff et al., 2007). The analysis of Artale et al. (2018) investigated the triggering of a diapycnal mixing due to rough bathymetry. To explore topographic-induced mixing, these authors estimated dissipation rates from the "CTD strain-based" parameterization (Kunze et al., 2006), showing the role of the sea bottom in enhancing isopycnal vertical strain. In particular, they highlighted that the turbulent kinetic energy generated by breaking internal waves is large within a few hundred meters of rough-bottom topography and decays to weaker values farther up in the water column; the bottom enhancement of turbulence reflects the generation of energetic waves, impinging over topography and breaking locally.

From the synergetic use of LADCP and CTD data, collected in the Ionian Sea during the KM3Net cruise in 2007, here we expand the analysis of Artale et al. (2018) to investigate shear variance in addition to strain variance, also exploring the role of topographic characteristics, such as bottom slope and roughness. In particular, we explore different seafloor characteristics, in terms of bathymetric gradient and bottom roughness. LADCP profiles, synchronous with CTD casts, allowed estimation of regional values of shear-to-strain ratio R_ω (Kunze et al., 2006). In the frame of the Garrett-Munk parametrization (Polzin & Lvov, 2011; See Data and methods), we explore the relation between R_ω

and topographic features of the Ionian Sea. This approach employs CTD profiles to determine the isopycnal vertical strain ($\langle \xi_z^2 \rangle_{in situ}$), and LADCP profiles for the vertical shear of horizontal velocities ($\langle V_z^2 \rangle_{in situ}$), assuming that the variance is due to the presence of IWs (See Data and methods). These variances are used to estimate the local energy's level of the IW field, to modulate a canonical value of the dissipation rate of turbulent kinetic energy ϵ_0 (Garrett and Munk, 1975).

2 Data and methods

2.1 CTD-LADCP casts

Our analysis is based on CTD and LADCP profiles acquired in July 2007 during a research cruise aboard the R/V Urania, conducted within the framework of the KM3NeT project (<https://www.km3net.org/>; Borghini et al., 2025). CTD-LADCP stations were distributed over a wide area of the northwestern Ionian Sea (Figure 1a), including strategic sites for the KM3NeT (Neutrino Telescope) infrastructure that had been identified by the particle physics community (Katz, 2006).

CTD data were acquired from the sea surface to the bottom by using a SBE911-plus calibrated before the cruise at the NATO center in La Spezia and were processed by the SBE Sea Soft program, following the standard procedure suggested by the manufacturer. LADCP profiles were collected at a subset of stations using a dual-head system installed on the rosette made of two RDI BB/WH 300 kHz instruments looking upward and downward respectively. The instruments were set to acquire 20 bins with size 10 m and the LADCP data were processed using the LDEO LADCP software Version IX.13 (Visbeck, 2002), using CTD and respective GPS fixes as auxiliary data.

2.2 Finescale parameterization

The small spatial scales (order of centimeters) and intermittent temporal nature (from minutes to hours) involved in internal-wave-driven turbulent mixing, imply that direct measurements are difficult to achieve. Indeed, measurements resolving small-scale turbulence rely on ship-based microstructure profilers, not yet in wide use and, in any case, not on board during the 2007 KM3NeT research cruise. To fill this gap, finescale parameterizations are being used to estimate the turbulent kinetic energy dissipation rate and diapycnal diffusivity from more common instruments, such as CTDs, ADCPs, and Argo profiles (Whalen et al., 2020; Polzin et al., 2014). Finescale parameterizations aim to infer the centimeter-scale turbulent energy dissipation rate, operating on the intermediate [O(10-100) m] vertical wavelengths that are assumed to mediate energy transfer between large and small scales in the ocean (Polzin et al., 2014). Finescale parameterizations are formulated by reference to the Garrett-Munk 75 model (hereafter, GM) (Garrett & Munk, 1975; Cairns & Williams, 1976; Polzin & Lvov, 2011), which provides an empirical expression for the internal wave energy density in the spectral domain. Strain- and shear-based parameterizations allow to estimate the in-situ internal wave energy level, which serves to establish a deviation from the GM model, and to adjust its canonical value for the turbulent dissipation rate ϵ_0 to a more realistic estimate. In particular, the turbulent dissipation rate can be expressed through the form of the finescale parameterization proposed by Kunze et al. (2006) as:

$$\epsilon_{iw} = \epsilon_0 \frac{N^2}{N_0^2} \left(\frac{E}{E_{GM}} \right)^2 F(R_\omega) L(f, N) \quad (1)$$

where E is the in situ internal wavefield variance and N the in-situ buoyancy frequency, while E_{GM} , $\varepsilon_0 = 7 \times 10^{-10} W/kg$ and $N_0 = 5.2 \times 10^{-3} rad/s$ are the GM internal wavefield variance, dissipation rate and Brünt-Väisälä frequency, respectively (Garrett and Munk, 1975; Munk, 1981). Depending on whether the parameterization is based on shear or strain, $(E_{in-situ}/E_{GM})^2$ in Eq. (1) can be estimated from the shear variance as $\langle V_z^2 \rangle_{in-situ}^2 / \langle V_z^2 \rangle_{GM}^2$, or from the strain variance as $\langle \xi_z^2 \rangle_{in-situ}^2 / \langle \xi_z^2 \rangle_{GM}^2$.

The factor $L(f, N)$ in (1) represents the latitude effect, calculated as $L(f, N) = f \cosh^{-1}(N/f) / f_0 \cosh^{-1}(N_0/f_0)$ with f the Coriolis parameter and f_0 the Coriolis's frequency at 30° . The term R_ω represents the internal wavefield's aspect ratio and the bulk frequency content, and it is defined as the ratio of the buoyancy-normalized shear variance to the strain variance:

254

$$R_\omega = (\langle V_z^2 \rangle / N^2) / \langle \xi_z^2 \rangle, \quad (2)$$

256

whose effect on dissipation rates ε_{iw} is modeled by the term $F(R_\omega)$:

258

$$F(R_\omega) = h_1(R_\omega) = 3/4(1 + 1/R_\omega)\sqrt{2/(R_\omega - 1)} \quad (\text{for shear-based parameterization}) \quad (3a)$$

$$F(R_\omega) = h_2(R_\omega) = R_\omega(R_\omega + 1)/(6\sqrt{2R_\omega - 1}) \quad (\text{for strain-based parameterization}) \quad (3b)$$

261

Equations (3a,b) point out that the error associated with incorrectly assuming R_ω is not symmetric (Chinn et al. 2016). For $R_\omega = 3$ (i.e., the GM value), the correction functions are equal to 1 and do not affect ε_{iw} . When $R_\omega > 3$, shear variance dominates, low-frequency internal waves contribute more strongly, and h_1 decreases (underestimates ε_{iw}), while h_2 increases (overestimates ε_{iw}). When $R_\omega < 3$, strain variance dominates, high frequency internal waves contribute more to the ratio: h_1 increases (overestimates ε_{iw}), h_2 decreases (underestimates ε_{iw}). This makes the shear-based parameterizations more reliable when $R_\omega > 3$, and conversely the strain-based parameterizations are more reliable when $R_\omega < 3$ (Chinn et al. 2016).

The determination of R_ω requires shear and strain variance measurements through LADCP and CTD observations, respectively. The strain ξ_z is defined as the vertical derivative of the isopycnal displacement and is generally calculated as $\xi_z = (N^2 - N_{fit}^2)/\overline{N^2}$, where N^2 is the Brünt-Väisälä frequency, $\overline{N^2}$ a mean value over the vertical segment of water column, and N_{fit}^2 a polynomial quadratic fit: $N^2 - N_{fit}^2$ represents the IW's density perturbation, normalized by $\overline{N^2}$ to obtain the vertical derivative of the isopycnal displacements.

Instead of determining ξ_z , we follow the approach of Ferron et al. (2014) and we determine ξ from the density fluctuations as $\xi = (g/\rho_0)(\rho_{HP}/\overline{N^2})$, with g/ρ_0 the standard gravity on the reference

density, and ρ_{HP} density filtered by a high-pass Butterworth filter with a cutoff wavelength of 160m to remove the density fluctuations associated with the background stratification, whose signal would contaminate the integration of the strain spectra (Kunze et al., 2006). This advantageously avoids fitting a polynomial to N^2 (e.g., quadratic fit in Kunze et al., 2006), that could be inconsistent with the shape of the density profile in case of strong vertical gradients. This method is a variant from Kunze et al. (2006), and follows Ferron et al. (2014). Strain variance is then calculated as the integral of the isopycnal displacement spectra between k_{min} and k_{max} :

$$\langle \xi_z^2 \rangle = \int_{k_{min}}^{k_{max}} k^2 S(\xi) dk. \quad (4)$$

Similarly, the shear variance $\langle V_z^2 \rangle$ is obtained from the velocity spectra as

$$\langle V_z^2 \rangle = \int_{k_{min}}^{k_{max}} k^2 S(u, v) dk. \quad (5)$$

To integrate the spectra, k_{min} and k_{max} must be carefully determined, as shear and strain rely on different instrument limitations and can be sensitive to distinct phenomenological contributions. In general k_{min} is fixed by the choice of the data segment length $k_{min} = 2\pi/L \text{ rad m}^{-1}$, where L is generally of 200-to-500 m, i.e., a compromise between the larger vertical wavelengths to be considered (the spectral resolution to be employed) and the desired vertical resolution. For strain, in particular, the integration is performed from $k_{min} = 2\pi/O(150m)$ (Kunze, 2006, Pollman, 2017; 2020), to exclude contributions from larger-scale background stratification, and continues up to $k_{max} = 2\pi/10m$ (the GM upper limit).

For shear, a velocity noise is expected to dominate at high wavenumber, especially in weakly stratified layers lacking turbulent microstructures or suspended material (“water-column reflectors”) (Kunze et al., 2006), which tend to reflect the acoustic signal emitted by Doppler profilers like the LADCP. Their absence reduces the quality of the acoustic return, leading to higher measurement uncertainty and potentially spurious velocity variance, especially at small vertical scales. Thus, wavenumbers larger than $k_{max} = 2\pi/O(100 - 50m)$ represent the upper limit range of usable vertical wavenumbers for integration.

An additional saturation criterion is applied to limit the integration of spectra (Gargett, 1990, Kunze et al., 2006). Specifically, the integration is stopped when reaching the saturation value of $0.66 N^2$ for the shear variance and 0.22 for strain. This threshold represents the onset of internal wave breaking or turbulence, where the linear internal wave regime breaks down. The corresponding vertical wavenumber k_c at which this value is reached is retained as the upper limit for the integration. This approach constrains the finescale parameterization to physically realistic shear and strain levels, avoiding overestimation of internal wave energy due to instrument noise or unresolved scales, particularly in regions of enhanced energy.

In our case, strain and shear are derived from CTD and LADCP measurements from the bottommost 640 m of each vertical profile. Signals are then linearly detrended and Fourier transformed with a Hamming tapering window. Strain and shear variance estimates are obtained by integrating the 640 m segments, overlapping every 10 m from the bottom to the surface, to account for non-homogeneity in the statistical distribution of the water column properties and to have a statistically robust representation of the fine-scale variability. Final variances are obtained by averaging collections of estimates over finite-sized segments (here 80 m).

The shear integration upper limit is identified from the LADCP velocity uncertainty. The LDEO LADCP processing (velocity-inversion method; Visbeck, 2002) provides, for each station, a depth-dependent single-bin velocity error profile, i.e. the estimated uncertainty associated with each vertical velocity bin, which is used to constrain the usable shear wavenumber range. The average $\overline{V_{ERR}} = 0.075 \text{ m.s}^{-1}$ is used to estimate a velocity error by segment as $V_{NOISE} = \overline{V_{ERR}}/\sqrt{640\text{m}/10\text{m}}$. Its associated variance V_{NOISE}^2 is assumed to arise from white velocity noise, uncorrelated in space and therefore uniformly distributed across vertical wavenumbers. In the spectral domain, this corresponds to a flat velocity noise spectrum which, once differentiated, produces a shear noise spectrum used to identify the noise-dominated upper wavenumber limit..

The resulting spectra are then averaged by stratification bins, where we highlight their overall shape and the noise-free bandwidths we retain for integration, from 640 to 107 m. This choice is conservative, as our lowest stratification levels are close to the $N_{ERR} = 5 \cdot 10^{-4} \text{ s}^{-1}$ (discussed in Kunze, 2006), and consistent with the integration set up of Ferron (2014) that stopped the integration at 107 m in the most limitative case. Additionally, this treatment combined both data from downward and upward profiles, in relatively small bin size (10 m), limiting the expected variance loss at short scales due to the various steps of data processing (Polzin, 2002; Thurnherr, 2012). For strain, integration is led from 128 to 10 m (the GM upper limit).

Since shear and strain are integrated on their respective bandwidths, instead of using Eq. (2) directly, the shear-to-strain ratio is calculated as:

$$R_{\omega} = R_{\omega}^{GM} (< V_z^2 > / < V_z^2 >_{GM}) / (< \xi_z^2 > / < \xi_z^2 >_{GM}), \quad (6)$$

where $R_{\omega}^{GM} = 3$, both shear and strain factors are divided by their associated value in the GM model, evaluated on the same spectral bandwidth, respectively.

2.3 Internal wave beams and impact of bathymetry on R_{ω}

Traditional finescale parameterizations of turbulent kinetic energy dissipation rely on the assumption that the internal wave field follows a GM-like structure. In this framework, a fixed value of the shear-to-strain ratio, typically $R_{\omega} \approx 3$, is used to represent the relative contributions of vertical shear and isopycnal strain to the finescale energy budget. This fixed value implicitly assumes a broadband, statistically stationary internal wave field, with no significant spatial or temporal variability in spectral composition. However, these models sometimes fail to capture the true dynamics, for

instance, near mixing hotspots where the internal wave spectra can be significantly distorted, showing bias toward higher or lower frequencies (Chinn et al., 2016; Polzin et al., 2014; Ferron et al., 2014)..

Accumulating evidence from both observations and modeling (Ijichi and Hibiya, 2015, 2017; Takahashi, et al. 2021; Dematteis et al., 2024) indicates that this assumption is often violated, especially in regions influenced by complex bathymetry, variable stratification, or energetic boundary currents. In these settings, the internal wave spectrum is distorted, leading to departures from the canonical balance between shear and strain. As a result, R_ω can vary significantly in space and time, reflecting changes in the dominant frequency content of the wave field.

Ijichi and Hibiya (2017) used 3D eikonal equations to simulate how internal waves transfer energy across distorted spectra, including variations in the wave energy level, the local buoyancy frequency, and the inertial frequency. Their results confirm that energy transfer rates are consistent with the Henyey et al. (1986) model, which predicts dissipation from internal wave-wave interactions and highlights the importance of the spectral shear-to-strain ratio R_ω . The study questions the accuracy of existing finescale parameterization models, particularly in environments with distorted spectra near boundaries or topographic features. To address these limitations, Ijichi and Hibiya (2015, 2017) propose a revised parameterization that explicitly considers both broadband and narrowband spectra. This leads to more accurate dissipation estimates by incorporating a dynamically varying R_ω instead of assuming a constant value based on idealized spectral shapes. Their approach was further developed by Takahashi et al. (2021), who showed how vertical wavenumber spectra distortions (such as spectral humps) can bias traditional estimates, emphasizing the need to resolve the spatial variability of R_ω .

From a physical point of view, R_ω represents the ratio between the buoyancy-normalized vertical shear variance and the isopycnal strain variance. The spectral structure of the internal wave field determines whether energy is primarily stored in the shear (associated with horizontal motion) or in the strain (associated with vertical isopycnal displacement). The theoretical foundation for this lies in the internal wave dispersion relation:

$$\omega^2 = \frac{N^2 k_h^2 + f^2 k_z^2}{k_h^2 + k_z^2}, \quad (7)$$

where ω is the wave frequency, $\mathbf{k}_h = (k_x, k_y)$ and k_z the horizontal and vertical wavenumbers, respectively.

Equation (7) shows the intrinsic relation between horizontal (vertical) wavenumber and the shear-to-strain ratio. When $\omega \approx f$, it follows that $k_z \gg k_h$ and that the wavefield is dominated by horizontal motion, yielding high shear $\partial u / \partial z$, i.e., the wavefield varies rapidly with depth and more slowly in the horizontal direction. This configuration results in particle motions that are predominantly horizontal but vary significantly over short vertical distances.

In observed internal wave spectra, low-frequency components (near the Coriolis frequency f) tend to concentrate energy in horizontal motions that change rapidly with depth, making shear-dominated fields a hallmark of low-frequency wave regimes (Kunze et al., 2006; Musgrave et al., 2022).

Conversely, when $\omega \approx N$, $k_h \gg k_z$ and the particle motion becomes primarily vertical, enhancing isopycnal displacement and thus the strain $\partial \xi / \partial z$ (i.e., significant vertical displacements of isopycnals). It results that the wave propagates with a nearly vertical beam and the wave field varies slowly with depth. For this condition, although the vertical structure of the wave is broad (i.e., fewer oscillations along z), the amplitude of vertical displacements is large. As a result, the isopycnals experience stronger stretching and compression, leading to enhanced strain energy. This explains why internal waves at high frequencies, despite their low vertical wavenumber, contribute disproportionately to the strain variance. In observed internal wave spectra, higher-frequency components tend to be associated with enhanced isopycnal displacement relative to velocity shear, leading to strain-dominated regimes (Garrett and Munk, 1975; Kunze et al., 2006).

A propagation angle of an internal wave ray can be determined using the dispersion relation:

$$\frac{k_z^2}{k_x^2 + k_y^2} = \frac{N^2 - \omega^2}{\omega^2 - f^2}, \quad (8)$$

thus,

$$\tan^2 \alpha = \frac{N^2 - \omega^2}{\omega^2 - f^2}. \quad (9)$$

From Equation (9), the angle of the wave trajectory (i.e., the beam angle α) relative to the horizontal can be determined. The angle α describes a cone of propagation that supports the wave:

$$\alpha = \tan^{-1} \sqrt{\frac{N^2 - \omega^2}{\omega^2 - f^2}}. \quad (10)$$

The wave energy propagates with the group velocity $\mathbf{c}_g$, perpendicular to the wave direction $\mathbf{k}$, at an angle $\beta = 90^\circ - \alpha$. For R_ω it is possible to derive the following expression (see Kunze et al., 2006; Ijichi and Hibiya, 2015):

$$R_\omega = \frac{\langle V_z^2 \rangle}{\langle \xi_z^2 \rangle N^2} = \frac{\omega^2 + f^2}{\omega^2 - f^2} \simeq 1 + \frac{2f^2 k_z^2}{N^2 k_h^2}, \quad (11)$$

from which we estimate a local value of ω in the water-column, given $N(z)$ and the local f , that we assume representative of a certain regional extent in the horizontal. From ω we can estimate β perpendicular to α . The relation between R_ω , ω and β will be exploited later in our study to associate the beam angle with R_ω value and compare it with the bathymetrical slope to infer propagation insights. Equation (10), along with the physical interpretation of Equation (7), highlights that the wave geometry (i.e., the ratio k_z^2/k_h^2) dominates the shear-strain dynamics: for $k_z \gg k_h$ we have high shear and R_ω values, and for $k_h \gg k_z$ high strain values and low R_ω . In other words, high R_ω values correspond to low-frequency, shear-dominated conditions, while low R_ω indicates high-frequency, strain-dominated fields.

A growing body of evidence shows that turbulent mixing is significantly enhanced in regions of rough bottom topography and sloping boundaries (Polzin et al., 1997; Ledwell et al., 2000; Kunze et al., 2006; Ferrari et al., 2016; Mashayek et al., 2017; Naveira Garabato et al., 2025). Internal wave interactions with these features can lead to reflection, scattering, and energy transfer toward higher vertical wavenumbers, increasing the likelihood of wave breaking and dissipation (Nash et al., 2004; Legg & Adcroft, 2003; Musgrave et al., 2022). In addition to tidal processes, geostrophic flows impinging on small-scale topography generate broadband internal waves that can radiate and dissipate energy locally, contributing to abyssal mixing (Nikurashin & Ferrari, 2010a, 2010b). These mechanisms, collectively, explain much of the spatial variability in observed mixing rates and highlight the importance of accurately representing topographic interactions in ocean mixing parameterizations.

When an internal wave encounters a rough topography, its propagation is disturbed, promoting scattering, reflection, or the generation of higher-frequency waves. This tends to break down low-frequency, large-scale wave structures and shift spectral energy toward higher vertical wavenumbers.

Since strain is associated with large vertical displacements, characteristic of high-frequency waves, this shift leads to a systematic decrease of the shear-to-strain variance ratio R_ω in rougher regions, as documented in Kunze et al. (2006), where departures from the canonical Garrett–Munk spectrum were observed in high-roughness environments. As a result, areas of high roughness typically exhibit lower R_ω , indicating increased strain. In contrast, smoother topographic regions preserve the lower-frequency shear-dominated character, yielding higher R_ω .

In our investigation of the impact of bathymetry on R_ω , we analyze the KM3NeT hydrological stations in relation to bathymetric features such as slope and topographic roughness, calculated using the GEBCO gridded data (<https://www.gebco.net>) over the Ionian Sea area, at 15 arc-second intervals. In particular, slopes are estimated between the hydrological cast position and the points located apart in the eight cardinal directions around, then the average is retained. A distance of 10 km between the two points is chosen to avoid overestimation of the topographic slope, due to the presence of small bathymetric features. Topographic roughness is calculated from the variance of the bathymetry over the same neighboring area (i.e., 20-by-20 square kilometers). R_ω values are then compared with the local bathymetric roughness, i.e., the logarithm of the variance of seafloor elevation (Figures 1c,d).

3 Results

To investigate the influence of bottom topography on internal wave shear-to-strain ratio from Equation (6), our dataset is bin-averaged every 80 m, from bottom towards the surface. From the first upward segment of 640 m we obtain values at the center of the segment (i.e. 320 m from bottom), and

the associated first binned value is available in bin 4, centered at 360 m from bottom. We stop at bin 12 (centered 920 m from the bottom). By doing this, we consider 8 bins of 80 m.

Results show that most of the near-bottom values of R_ω are larger than the canonical value of 3, given by the GM model (Figure 1; Supplementary Table S1). Small values, from 1.01 to 10, are generally observed in the shelf break regions, as well as at some deep stations (Figure 1c; Supplementary Table S1), or close to the Malta escarpment (Figure 1a,b,c). For deep, offshore casts, estimations show large values. High values of shear-to-strain ratio are evident in the western part of the abyssal plain, between 2000 and 3000 m depth, i.e., along the offshore portion of the four cross-shore transects T1, T2, T3, and T4 (Figure 1b,c). Here, R_ω shows values between ~ 10 and ~ 30 (Figures 1c; Supplementary Table S1).

In general, regions with higher roughness show lower R_ω (Figure 1d,e), suggesting, for those regions, a shift toward higher-frequency energy and enhanced vertical displacement (i.e., strain). High values of bottom roughness are distributed along the eastern and northern shelf; a smoother bathymetry is recognized in the abyssal plain (Figure 1e). By plotting R_ω against bathymetric slope and roughness (Figure 1d), we find that although a linear relationship is difficult to establish, an interesting pattern emerges: increased roughness appears to reduce R_ω . This effect could also be attributed to the bathymetric slope, as slope and roughness are correlated, with steeper zones leading to greater variance in elevation. We hypothesize that this pattern indicates waves being constrained toward higher frequency bands by the roughness.

Indeed, the increasing R_ω corresponds to decreasing frequency and flatter beam angles (i.e., more horizontal propagation), while low R_ω indicates steeper beams and higher frequencies (Figure 2a). The grey vertical line and shaded area in Figure 2a highlight the mean and observed range of R_ω in our dataset, contextualizing the frequencies and angles involved. This framework sets the stage for comparing beam angles with local bathymetric slopes. A Δ -slope is inferred as the difference between the beam angle in Equation 10 and the bathymetry slope. This angle controls the reflection regime: near-critical reflection (Δ -slope $\approx 0^\circ$) concentrates energy in horizontal shear, while supercritical conditions (Δ -slope $< 0^\circ$) promote upward-propagating waves and stronger vertical displacements (strain) (Musgrave et al., 2022). R_ω is clearly related to Δ -slope (Figure 2b): higher R_ω values correspond to smaller angle differences, while lower R_ω values are associated with steeper conditions. This indicates a shift in frequency content, illustrated by the associated ω -distribution (Figure 2c), with frequencies increasing as Δ -slope grows. Additionally, roughness appears to constrain the dynamical range of R_ω dispersion. This suggests a critical threshold where high frequencies and smaller scales begin to dominate over shear.

By organizing the shear and strain spectra into bins of roughness and Δ -slope to identify whether specific bandwidths are affected, one might expect enhanced signals at larger vertical scales (≈ 300 m) or smaller scales (≈ 100 m), depending on the preferential frequency content potentially enhanced by the bathymetric features. The resulting spectra (Figure 3b) show that strain variance increases at high vertical wavenumbers (small vertical scales) in those regions characterized by the highest roughness (bin 5.1). A tendency toward reduced shear variance at low wavenumbers is observed in rougher regions, accompanied by a flattening of the shear spectrum. Given the limited number of realizations, these differences are interpreted as systematic patterns rather than statistically significant changes. Indeed, shear spectra under rougher conditions (blue spectra in Figure 3b) appear flatter in the lower wavenumber band compared to smoother regions. Intermediate roughness levels (green spectra in Figure 3b) show increased energy around vertical scales of 64 m, potentially indicating enhanced

energy distribution at shorter scales. Strain spectra exhibit a distinct pattern for the highest roughness, with higher energy levels compared to other bins. These observations suggest that increasing roughness reduces energy distribution at larger scales (shear lower band) while enhancing strain variance contributions at smaller scales (upper band).

To isolate the influence of slope geometry, we use the beam-to-slope angle difference Δ -slope. The shear spectra in Figure 3c shows a clear dependence on Δ -slope. Indeed, higher R_ω values correspond to smaller angle differences, while lower R_ω values are associated with steeper conditions. This indicates a shift in frequency content, with frequencies increasing as Δ -slope grows. Spectra in Figure 3c are coherent with what we observed in Figure 2 and suggest a critical threshold where high frequencies and smaller scales begin to dominate over shear: the range of R_ω appears to be limited when the roughness increases. However we do not have enough data to establish if there exist an actual "threshold value" of roughness above which the spread of R_ω is limited. This interpretation is further confirmed by the patterns observed in both Figures 3 and 4: as Δ -slope approaches 0, lower frequencies become dominant, indicating more horizontal motion. This is associated with increased shear and a higher prevalence of critical Richardson instability events (Figure 4a,b), where the Richardson number is defined as $R_i = \frac{N^2}{\left[\left(\frac{\partial u}{\partial z}\right)^2 + \left(\frac{\partial v}{\partial z}\right)^2\right]}$: R_i values show their minimum values tending toward criticality with low values of delta slope (i.e., flat angles). Yet, the deviation from the GM model is evident in the shear ratio, with values exceeding 8–10 times the model (Figure 4c,d), while the strain consistently remains below the GM model's predictions (Figure 4e,f).

4 Discussions and conclusions

Our analysis in the Ionian Sea contributes to the investigation of the IW mixing, as described by Kunze et al. (2006). We focused on hydrological stations where we took advantage of synchronous LADCP and CTD profiles from which we infer regional values of the near-bottom ratio between shear and strain variances (R_ω). Values for R_ω are necessary in the finescale parameterization for deriving both dissipation rate of turbulent kinetic energy and diapycnal diffusivity (i.e., ϵ_{iw} and K_{iw} , respectively). Our goal, in particular, was to quantitatively assess the spatial patterns of shear-to-strain ratio to highlight the role of morpho-bathymetric features in enhancing dissipation and diffusion rates, in terms of depth, slope, and roughness.

We found a general relation between bottom depth and shear-to-strain variance ratio (R_ω), which tends to increase from the shelf break to the abyss: large (small) values of R_ω occur over areas with gentle (steep) topographic slopes and/or low (high) seafloor roughness. This supports the hypothesis that gentler slopes and smoother seafloor generate lower-frequency internal waves, whereas steeper and rougher topography favors higher-frequency waves, leading to a corresponding increase in the shear (strain) contribution. In particular, for the northwest group of stations (i.e., the shelf-to-plain transects T1, T2, and T3; Figure 1b), we found coherent trends of R_ω , which co-varies with depth, slope, and roughness.

Our observations suggest that, in rougher conditions, the redistribution of internal wave energy toward higher frequencies and smaller vertical scales enhances strain variance relative to shear. The resulting decrease in R_ω indicates a wave field that departs from low-frequency, shear-dominated regimes and becomes increasingly strain-dominated, consistent with enhanced spectral distortion over rough topography (Chinn et al., 2016). In abyssal waters (i.e., deeper than 3000 m), specific stations of transects T4 and T5 (e.g., from Station NK6 to Station 11; Figure 1b) show a behavior similar to the one observed in the shelf-to-plain transects: R_ω increases with distance from the shelf break, starting from $R_\omega \sim 10$, although we are in abyssal waters. We notice, however, that those abyssal stations of

Transect 4 that show low values of R_ω (i.e., values we expect in rough areas) are close to the topographic constraints of the Malta Escarpment (Figure 1a), where energetic mesoscale meandering features and vortical structures at the bottom, were observed (Rubino et al., 2012; Meccia et al., 2015). These low values of shear-to-strain ratio may be due to the eventual mesoscale eddy–internal wave coupling that could represent a sink of eddy energy and then a source of IW energy (Polzin, 2010; Takahashi et al., 2021). These abyssal stations in Transect 4 may therefore represent those particular, “nontraditional” cases (Gerkema and Shrira, 2005) where deep mesoscale activity is able to trigger diapycnal mixing by breaking of IWs. Finally, we notice that in Ferron et al. (2014) a 160m- high-pass filter is explicitly applied on density to remove largescale density variations, that can lead to spectral shape differing from Kunze et al. (2006) for scale larger than 160m. The overall level of our strain spectra is also closer to GM, which is consistent with the fact that our observed internal-wave field is close to GM in amplitude.

Kunze et al. (2006) suggested that the lack of reflectors in the water column at deep locations could lead to an overestimation of the shear variance, reflected ultimately in strong values of R_ω . However, within the limits of our analysis, the shear spectra do not display clear indications of dominant noise contamination, given the conservative choices adopted for the integration bands.

In addition to this potential instrumental bias, we also examined whether thermohaline structure could affect the interpretation of R_ω . In particular, we observe that the bottom layer of several eastern abyssal stations exhibits thermohaline gradients that could be consistent with double-diffusive regimes (Miles, 1961; Ruddick, 1983; Thorpe, 2005). However, an additional analysis of Turner angles (Tu) indicates that only a few thin layers fall within active values $|Tu| > 75^\circ$ (13.6% for salt Fingering, 1.8% for diffusive convection). These few layers do not display particularly significant density structures. We therefore consider that double-diffusive layers are only minimally represented in our dataset and are unlikely to significantly influence the overall analysis of R_ω .

Our analysis on shear-to-strain ratio is based on CTD-LADCP casts that were carried out during a single oceanographic cruise in 2007. Therefore, we cannot perform any analysis on the temporal variability of the spatial patterns we observe. However, we can frame our results within the interannual analysis (1977-2011) presented by Artale et al. (2018). These authors highlighted the presence of a “new” bottom Ionian water, from the 2003, resulting from the continuous entrainment of the warmer, upper waters, a process that would cause a loss of kinetic energy and gain of potential energy in the deep layer (Marotzke and Scott, 1999). Based on this analysis we can argue that the spatial pattern of shear-to-strain ratio we observe is likely depicting the particular barotropization of the bottom layer found in Artale et al. (2018), characterized by a homogeneous bottom layer between 3000 and 4000 m depth and weak horizontal gradients of potential temperature over the Ionian basin (Artale et al., 2018). Our results are consistent with previous studies (i.e., Chinn et al., 2016; Polzin et al., 2014; Ferron et al., 2014) in showing that shear–strain balance is not universal but systematically modulated by environmental conditions, here specifically seafloor roughness and slope in the Ionian Sea.

Finally, our analysis aims to provide useful insights for ocean circulation models, which are often too sensitive to vertical eddy diffusivity and are largely affected by inaccuracy at deep layers (Gargett and Holloway, 1982; Wright and Stocker, 1992). In particular, the large variability of the shear-to-strain ratio we found, may help deriving both dissipation rate of turbulent kinetic energy and diapycnal diffusivity in the finescale parameterization.

594 **5 Conflict of Interest**

595 The authors declare that the research was conducted in the absence of any commercial or financial
596 relationships that could be construed as a potential conflict of interest.
597

598 **6 Author Contributions**

599 FF, VA, and FK delineated, supervised the study. SS, MB, and FF collected the data. SS and FK
600 processed and analyzed the CTD and LADCP data. FK, SS, VA, DC, and FF contributed with
601 resources, analyzed the data and wrote the manuscript.
602

603 **7 Funding**

604 This work was partially supported by EMODnet (European Marine Observation and Data Network
605 Physics), the Copernicus Climate Change Service (C3S) project, and the Flagship Project RITMARE
606 (The Italian Research for the Sea), coordinated by the Italian National Research Council and funded
607 by the Italian Ministry of Education, University and Research.
608

609 **8 Acknowledgments**

610 Data were collected in the framework of the project “Specific Support Action for the Design Study
611 for a Deep-Sea Facility in the Mediterranean for Neutrino Astronomy and Associated Sciences”
612 (KM3NeT), EU contract no. 011937. We thank Daniele Iudicone, Manuel Bensi, and Enrico
613 Zambianchi for the insightful discussion at the initial stage of the investigation.
614

615 **9 Data and Code Availability**

616 All functions used for obtaining Equation (6) are in Kokoszka, F. (2025).

617 Data are available on Sea Scientific Open Data Publication (SEANOE),
618 <https://www.seanoe.org/preview/108742?token=NkpdOxO5966RjZ--Rqdw1c0dCZ-6iGSW>
619 Codes that were used for this analysis are available at
620 <https://zenodo.org/records/17170422> (DOI: [10.5281/zenodo.17170421](https://doi.org/10.5281/zenodo.17170421))
621
622
623
624
625
626
627
628
629
630
631
632
633

Figures




Figure 1. a) The Mediterranean Sea bathymetry and the main regional features that are discussed within the text; the red box indicates the study area. b) The Ionian Sea bathymetry and the main regional features that are discussed within the text; LADCP-CTD casts are grouped in five main hydrological transects: four shelf-to-plain transects, i.e., T1 ($\blacktriangle$), T2 ($\blacksquare$), T3 ($\blacklozenge$), T4 ($\bullet$), and one abyssal transect T5 ($\times$). c) Shear-to-strain ratio R_ω at the bottom of the LADCP-CTD profiles, averaged from bins of 80m in the layer 360-920m distant from the bottom. d) Bottom roughness ($\log_{10}$ of $\text{var}(z)$) at the cast sites. e) R_ω as a function of bathymetric slope (x-axis), and roughness (in color).




Figure 2. a) Correspondence between R_ω and ω (black), and the beam angle associated with R_ω (orange), computed at the average latitude of the study (36.75°N). Typical values, from high to low R_ω , are reported. The average R_ω value of our study is indicated with the vertical line marker, and its distribution range is indicated by the gray area. **b)** Distributions of R_ω and **c)** ω in the layer 360-920 m distant from bottom, in function of the difference between the associated beam angle and the bathymetry slope (Δ -slope). Roughness is indicated in color. Higher values on the x-axis indicate a beam angle steeper, relative to the bathymetry slope, while lower values indicate a beam aligning with the slope. The critical range where the alignment is close to be parallel to the slope is arbitrarily marked by the shaded red area from -1 to 1 degrees, out of which a supercritical range is indicated in orange.




Figure 3. a) Buoyancy-normalized shear (red) and strain (blue) wavenumber spectra, with their associated GM model (upper black line: shear; lower black line: strain). Spectra are averaged by bins of stratification (thick to thin, from weak to higher stratification). Number of spectra by bins, with the average N of the bin are reported in legend. Shear noise spectra associated with a noise velocity of 0.075 m/s are indicated in gray (thickness indicates its association to the N bins). Instrumental wavenumber limits are indicated with the vertical dashed lines: upper limit of $2\pi/128$ rad $\cdot$ m $^{-1}$ for shear due to noise contamination; lower limit of $2\pi/160$ rad $\cdot$ m $^{-1}$ for strain to filter the large-scale background stratification. Retained wavenumber bandwidths for integration are indicated with the shaded ranges on top and bottom (640-107 m for shear; 128-10 m for strain). **b)** Associated estimates of $R\omega$ are shown by station and by vertical bins of 80 m, within the same bottom 1000 m layer (lower panel). **c,d)** Buoyancy-normalized shear (**top**) and strain (**bottom**) wavenumber spectra, with their associated GM model (upper black: shear; lower black: strain). Spectra are averaged by bins of roughness (**c**), and by bins of the difference between the beam angle and the bathymetry slope angle (**d**). For both (c) and (d) number of spectra found by bins and the average bin value are reported in legend. Shear noise spectra associated with a noise velocity of 0.075 m/s are indicated in gray (thickness indicates its association to the N bins). Instrumental wavenumber limits are indicated with the vertical dotted lines: upper limit of $2\pi/128$ rad $\cdot$ m $^{-1}$ for shear due to noise contamination; lower limit of $2\pi/160$ rad $\cdot$ m $^{-1}$ for strain to filter the large-scale background stratification. Retained wavenumber bandwidths for integration are indicated with the shaded ranges on top and bottom (640-107 m for shear; 128-10 m for strain).




Figure 4. Distributions of R_ω (left) and ω (right) in the layer 360-920m distant from the bottom, in function of the difference between the associated beam angle and the bathymetry slope angle. Colours are used to indicate (a,b) a bulk Richardson number (in logarithmic scale), (c,d) the shear-to-shear-GM ratio, (e,f) the strain-to-strain-GM ratio. Higher values on the x-axis indicate a beam angle steeper, relative to the bathymetry slope, while lower values indicate a beam aligning with the slope. The critical range where the alignment is close to be parallel to the slope is arbitrarily marked by the shaded red area from -1 to 1 degrees, out of which a supercritical range is indicated in orange.

724 10 References

- 725 Artale, V., Provenzale, A., & Santoleri, R. (1989). Analysis of internal temperature oscillations of tidal
726 period on the Sicilian continental shelf. *Continental Shelf Research*, 9(10), 867-888.
- 727 Artale, V., Calmanti, S., Malanotte-Rizzoli, P., Pisacane, G., Rupolo, V., & Tsimplis, M. (2006). The
728 Atlantic and Mediterranean Sea as connected systems. In *Developments in Earth and Environmental*
729 *Sciences* (Vol. 4, pp. 283-323). Elsevier.
- 730 Artale, V., Falcini, F., Marullo, S., Bensi, M., Kokoszka, F., Iudicone, D., & Rubino, A. (2018).
731 Linking mixing processes and climate variability to the heat content distribution of the Eastern
732 Mediterranean abyss. *Scientific reports*, 8(1), 1-10.
- 733 Bellacicco, M., Anagnostou, C., Falcini, F., Rinaldi, E., Tripsanas, K., & Salusti, E. (2016). The 1987
734 Aegean dense water formation: A streamtube investigation by comparing theoretical model results,
735 satellite, field, and numerical data with contourite distribution. *Marine Geology*, 375, 120-133.
- 736 Bensi, M., Cardin, V., Rubino, A., Notarstefano, G., & Poulain, P. M. (2013b). Effects of winter
737 convection on the deep layer of the Southern Adriatic Sea in 2012. *Journal of Geophysical Research:*
738 *Oceans*, 118(11), 6064-6075.
- 739 Bensi, M., Rubino, A., Cardin, V., Hainbucher, D., & Mancero-Mosquera, I. (2013a). Structure and
740 variability of the abyssal water masses in the Ionian Sea in the period 2003-2010. *Journal of*
741 *Geophysical Research: Oceans*, 118(2), 931-943.
- 742 Béranger, K., Mortier, L., Gasparini, G. P., Gervasio, L., Astraldi, M., & Crépon, M. (2004). The
743 dynamics of the Sicily Strait: a comprehensive study from observations and models. *Deep Sea*
744 *Research Part II: Topical Studies in Oceanography*, 51(4-5), 411-440.
- 745 Bergamasco, A., & Malanotte-Rizzoli, P. (2010). The circulation of the Mediterranean Sea: a historical
746 review of experimental investigations. *Advances in Oceanography and Limnology*, 1(1), 11-28.
- 747 Bindoff, N. L., Willebrand, J., Artale, V., Cazenave, A., Gregory, J. M., Gulev, S., ... & Woodworth,
748 P. (2007). Observations: oceanic climate change and sea level. *Climate Change 2007: The Physical*
749 *Science Basis*, eds Solomon S, et al. (Cambridge Univ Press, Cambridge, UK), pp 747–845.
- 750 Borghini, M., Falcini, F., Kokoszka, F., Sparnocchia S. (2025). CTD and LADCP data from the
751 KM3NET cruise, Ionian Sea, July 2007. *SEANOE*
752 (<https://www.seanoe.org/preview/108742?token=NkpdOxO5966RjZ--Rqdw1c0dCZ-6iGSW>).
- 753
- 754 Cairns, J. L., & Williams, G. O. (1976). Internal wave observations from a midwater float, 2. *Journal*
755 *of Geophysical Research*, 81(12), 1943-1950.
- 756 Cavaliere, D., La Forgia, G., Adduce, C., Alpers, W., Martorelli, E., & Falcini, F. (2021). Breaking
757 location of Internal Solitary Waves over a sloping seabed. *Journal of Geophysical Research: Oceans*,
758 126(2), e2020JC016669.

Chinn, B. S., Girton, J. B., & Alford, M. H. (2016). The impact of observed variations in the shear-to-strain ratio of internal waves on inferred turbulent diffusivities. *Journal of Physical Oceanography*, 46(11), 3299-3320.

De Lavergne, C., Madec, G., Le Sommer, J., Nurser, A. G., & Naveira Garabato, A. C. (2016). The impact of a variable mixing efficiency on the abyssal overturning. *Journal of Physical Oceanography*, 46(2), 663-681.

Dematteis, G., Le Boyer, A., Pollmann, F., Polzin, K. L., Alford, M. H., Whalen, C. B., & Lvov, Y. V. (2024). Interacting internal waves explain global patterns of interior ocean mixing. *Nature Communications*, 15(1), 7468.

Egbert, G. D., & Ray, R. D. (2000). Significant dissipation of tidal energy in the deep ocean inferred from satellite altimeter data. *Nature*, 405(6788), 775-778.

Ferrari, R. (2014). What goes down must come up. *Nature*, 513(7517), 179-180.

Ferrari, R., and Wunsch, C. (2009) Ocean Circulation Kinetic Energy: Reservoirs, Sources, and Sinks. *Annual Review of Fluid Mechanics*, 41, 253–282.
<https://doi.org/10.1146/annurev.fluid.40.111406.102139>

Ferrari, R., Mashayek, A., McDougall, T. J., Nikurashin, M., & Campin, J. M. (2016). Turning ocean mixing upside down. *Journal of Physical Oceanography*, 46(7), 2239-2261.

Ferron, B., Kokoszka, F., Mercier, H., & Lherminier, P. (2014). Dissipation rate estimates from microstructure and finescale internal wave observations along the A25 Greenland–Portugal OVIDE line. *Journal of Atmospheric and Oceanic Technology*, 31(11), 2530-2543.

Ferron, B., Kokoszka, F., Mercier, H., Lherminier, P., Huck, T., Rios, A., & Thierry, V. (2016). Variability of the turbulent kinetic energy dissipation along the A25 Greenland–Portugal transect repeated from 2002 to 2012. *Journal of Physical Oceanography*, 46(7), 1989-2003.

Ferron, B., Bouruet Aubertot, P., Cuypers, Y., Schroeder, K., & Borghini, M. (2017). How important are diapycnal mixing and geothermal heating for the deep circulation of the Western Mediterranean?. *Geophysical Research Letters*, 44(15), 7845-7854.

Gačić, M., Borzelli, G. E., Civitarese, G., Cardin, V., & Yari, S. (2010). Can internal processes sustain reversals of the ocean upper circulation? The Ionian Sea example. *Geophysical research letters*, 37(9).

Gargett, A. E. (1990). Do we really know how to scale the turbulent kinetic energy dissipation rate ε due to breaking of oceanic internal waves?. *Journal of Geophysical Research: Oceans*, 95(C9), 15971-15974.

Gargett, A. E., & Holloway, G. (1992). Sensitivity of the GFDL ocean model to different diffusivities for heat and salt. *Journal of physical oceanography*, 22(10), 1158-1177.

Garrett, C., & Kunze, E. (2007). Internal tide generation in the deep ocean. *Annu. Rev. Fluid Mech.*, 39, 57-87.

795 Garrett, C., & Laurent, L. S. (2002). Aspects of deep ocean mixing. *Journal of oceanography*, 58(1),
796 11-24.

797 Garrett, C., & Munk, W. (1975). Space-time scales of internal waves: A progress report. *Journal of*
798 *Geophysical Research*, 80(3), 291-297.

799 Gerkema, T., & Shrira, V. I. (2005). Near-inertial waves on the “nontraditional” β plane. *Journal of*
800 *Geophysical Research: Oceans*, 110(C1).

801 Gerkema, T., & Zimmerman, J. T. F. (2008). *An introduction to internal waves*. Lecture Notes, Royal
802 NIOZ, Texel, 207.

803 Henyey, F. S., Wright, J., & Flatté, S. M. (1986). Energy and action flow through the internal wave
804 field: An eikonal approach. *Journal of Geophysical Research: Oceans*, 91(C7), 8487-8495.

805 Holte, J., & Straneo, F. (2017). Seasonal overturning of the Labrador Sea as observed by Argo floats.
806 *Journal of Physical Oceanography*, 47(10), 2531-2543.

807 Ijichi, T., & Hibiya, T. (2015). Frequency-based correction of finescale parameterization of turbulent
808 dissipation in the deep ocean. *Journal of Atmospheric and Oceanic Technology*, 32(8), 1526-1535.

809 Ijichi, T., & Hibiya, T. (2017). Eikonal calculations for energy transfer in the deep-ocean internal wave
810 field near mixing hotspots. *Journal of Physical Oceanography*, 47(1), 199-210.

811 Jones, C. S., & Abernathey, R. P. (2019). Isopycnal mixing controls deep ocean ventilation.
812 *Geophysical Research Letters*, 46(22), 13144-13151.

813 Katz, U. F. (2006). KM3NeT: Towards a km³ Mediterranean neutrino telescope. *Nuclear Instruments*
814 *and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated*
815 *Equipment*, 567(2), 457-461.

816 Kokoszka, F. (2025). Supporting code, fszk/internal-waves-finescale: Finescale parameterization
817 shear/strain GM (v1.0.0). *Zenodo*. <https://doi.org/10.5281/zenodo.17170422>

818 Kunze, E., & Toole, J. M. (1997). Tidally driven vorticity, diurnal shear, and turbulence atop Fieberling
819 Seamount. *Journal of Physical Oceanography*, 27(12), 2663-2693.

820 Kunze, E., Firing, E., Hummon, J. M., Chereskin, T. K., & Thurnherr, A. M. (2006). Global abyssal
821 mixing inferred from lowered ADCP shear and CTD strain profiles. *Journal of Physical*
822 *Oceanography*, 36(8), 1553-1576.

823 La Forgia, G., Cavaliere, D., Adduce, C., Falcini, F. (2021). Mixing efficiency for breaking internal
824 solitary waves. *Journal of Geophysical Research: Oceans*, 2021JC017275-TR (in revision).

825 Ledwell, J. R., Montgomery, E. T., Polzin, K. L., Laurent, L. S., Schmitt, R. W., & Toole, J. M. (2000).
826 Evidence for enhanced mixing over rough topography in the abyssal ocean. *Nature*, 403(6766), 179-
827 182.

828 Legg, S., & Adcroft, A. (2003). Internal wave breaking at concave and convex continental
829 slopes. *Journal of Physical Oceanography*, 33(11), 2224-2246.

- 830 Levitus, S., Antonov, J. I., Boyer, T. P., & Stephens, C. (2000). Warming of the world ocean. *Science*,
831 287(5461), 2225-2229.
- 832 MacKinnon, J. A., et al. (2017). Climate Process Team on Internal Wave–Driven Ocean Mixing.
833 Bulletin of the American Meteorological Society, 98(11), 2429–2454.
834 <https://doi.org/10.1175/BAMS-D-16-0030.1>
- 835 Malanotte-Rizzoli, P., & Hecht, A. (1988). Large-scale properties of the eastern mediterranean-A
836 review. *Oceanologica Acta*, 11(4), 323-335.
- 837 Malanotte-Rizzoli, P., Artale, V., Borzelli-Eusebi, G. L., Brenner, S., Crise, A., Gacic, M., ... &
838 Triantafyllou, G. (2014). Physical forcing and physical/biochemical variability of the Mediterranean
839 Sea: a review of unresolved issues and directions for future research. *Ocean Science*, 10(3), 281-322.
- 840 Marotzke, J., & Scott, J. R. (1999). Convective mixing and the thermohaline circulation. *Journal of*
841 *Physical Oceanography*, 29(11), 2962-2970.
- 842 Mashayek, A., Ferrari, R., Merrifield, S., Ledwell, J. R., St Laurent, L., & Garabato, A. N. (2017).
843 Topographic enhancement of vertical turbulent mixing in the Southern Ocean. *Nature*
844 *communications*, 8(1), 1-12.
- 845 Meccia, V. L., Borghini, M., & Sparnocchia, S. (2015). Abyssal circulation and hydrographic
846 conditions in the Western Ionian Sea during Spring–Summer 2007 and Autumn–Winter 2007–2008.
847 *Deep Sea Research Part I: Oceanographic Research Papers*, 104, 26-40.
- 848 Miles, J. W. (1961). On the stability of heterogeneous shear flows. *Journal of Fluid Mechanics*, 10(4),
849 496-508.
- 850 Millot, C., & Taupier-Letage, I. (2005). Circulation in the Mediterranean sea. *The Mediterranean Sea*,
851 29-66.
- 852 Munk, W. H. (1966). Abyssal recipes. In *Deep Sea Research and Oceanographic Abstracts* (Vol. 13,
853 No. 4, pp. 707-730). Elsevier.
- 854 Munk, W. H. (1981). Internal waves and small-scale processes. *Evolution of physical oceanography*,
855 264-291.
- 856 Munk, W., & Wunsch, C. (1998). Abyssal recipes II: Energetics of tidal and wind mixing. *Deep Sea*
857 *Research Part I: Oceanographic Research Papers*, 45(12), 1977-2010.
- 858 Musgrave, R., Pollmann, F., Kelly, S., & Nikurashin, M. (2022). The lifecycle of topographically-
859 generated internal waves. In *Ocean mixing* (pp. 117-144). Elsevier.
- 860 Napolitano, E., Sannino, G., Artale, V., & Marullo, S. (2003). Modeling the baroclinic circulation in
861 the area of the Sicily channel: The role of stratification and energy diagnostics. *Journal of*
862 *Geophysical Research: Oceans*, 108(C7).
- 863 Nash, J. D., Kunze, E., Toole, J. M., & Schmitt, R. W. (2004). Internal tide reflection and turbulent
864 mixing on the continental slope. *Journal of Physical Oceanography*, 34(5), 1117-1134.

- 865 Naveira Garabato, A. C., Spingys, C. P., Castro, B. F., Couto, N., Drake, H. F., Forryan, A., ... &
866 Alford, M. H. (2025). Connecting mixing to upwelling along the ocean's sloping
867 boundary. *Geophysical Research Letters*, 52(22), e2025GL119186.
- 868 Nikurashin, M., & Ferrari, R. (2010a). Radiation and dissipation of internal waves generated by
869 geostrophic motions impinging on small-scale topography: Theory. *Journal of Physical*
870 *Oceanography*, 40(5), 1055-1074.
- 871 Nikurashin, M., & Ferrari, R. (2010b). Radiation and dissipation of internal waves generated by
872 geostrophic motions impinging on small-scale topography: Application to the Southern
873 Ocean. *Journal of Physical Oceanography*, 40(9), 2025-2042.
- 874 Nikurashin, M., & Ferrari, R. (2013). Overturning circulation driven by breaking internal waves in the
875 deep ocean. *Geophysical Research Letters*, 40(12), 3133-3137.
- 876 Nikurashin, M., and Ferrari, R. (2011). Global energy conversion rate from geostrophic flows into
877 internal lee waves in the deep ocean. *Geophysical Research Letters*, 38, L08610.
878 <https://doi.org/10.1029/2011GL046576>
- 879 Oddo, P., Poulain, P. M., Falchetti, S., Storto, A., & Zappa, G. (2023). Internal tides in the central
880 Mediterranean Sea: observational evidence and numerical studies. *Ocean Dynamics*, 73(3), 145-
881 163.
- 882 Osborn, T. R., & Cox, C. S. (1972). Oceanic fine structure. *Geophysical Fluid Dynamics*, 3(4), 321-
883 345.
- 884 Osborne, A. R., & Burch, T. L. (1980). Internal solitons in the Andaman Sea. *Science*, 208(4443), 451-
885 460.
- 886 Pinardi, N., & Masetti, E. (2000). Variability of the large scale general circulation of the Mediterranean
887 Sea from observations and modelling: a review. *Palaeogeography, Palaeoclimatology,*
888 *Palaeoecology*, 158(3-4), 153-173.
- 889 Pollmann, F. (2020). Global characterization of the ocean's internal wave spectrum. *Journal of*
890 *Physical Oceanography*, 50(7), 1871-1891.
- 891 Pollmann, F., Eden, C., & Olbers, D. (2017). Evaluating the global internal wave model IDEMIX
892 using finestructure methods. *Journal of Physical Oceanography*, 47(9), 2267-2289.
- 893 Polzin, K. L. (2010). Mesoscale eddy–internal wave coupling. Part II: Energetics and results from
894 PolyMode. *Journal of physical oceanography*, 40(4), 789-801.
- 895 Polzin, K. L., & Lvov, Y. V. (2011). Toward regional characterizations of the oceanic internal
896 wavefield. *Reviews of geophysics*, 49(4).
- 897 Polzin, K. L., Garabato, A. C. N., Huussen, T. N., Sloyan, B. M., & Waterman, S. (2014). Finescale
898 parameterizations of turbulent dissipation. *Journal of Geophysical Research: Oceans*, 119(2), 1383-
899 1419.

900 Polzin, K. L., Naveira Garabato, A. C., Huussen, T. N., Sloyan, B. M., & Waterman, S. (2014).
 901 Finescale parameterizations of turbulent dissipation. *Journal of Geophysical Research:*
 902 *Oceans*, 119(2), 1383-1419.

903 Polzin, K. L., Toole, J. M., Ledwell, J. R., & Schmitt, R. W. (1997). Spatial variability of turbulent
 904 mixing in the abyssal ocean. *Science*, 276(5309), 93-96.

905 Polzin, K., Kunze, E., Hummon, J., & Firing, E. (2002). The finescale response of lowered ADCP
 906 velocity profiles. *Journal of Atmospheric and Oceanic Technology*, 19(2), 205-224.

907 Roether, W., & Schlitzer, R. (1991). Eastern Mediterranean deep water renewal on the basis of
 908 chlorofluoromethane and tritium data. *Dynamics of Atmospheres and Oceans*, 15(3-5), 333-354.

909 Roether, W., Manca, B. B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., ... & Luchetta, A.
 910 (1996). Recent changes in eastern Mediterranean deep waters. *Science*, 271(5247), 333-335.

911 Rubino, A., Falcini, F., Zanchettin, D., Bouche, V., Salusti, E., Bensi, M., ... & Capone, A. (2012).
 912 Abyssal undular vortices in the Eastern Mediterranean basin. *Nature communications*, 3(1), 1-6.

913 Ruddick, B. (1983). A practical indicator of the stability of the water column to double-diffusive
 914 activity. *Deep Sea Research Part A. Oceanographic Research Papers*, 30(10), 1105-1107.

915 Schlitzer, R., Roether, W., Oster, H., Junghans, H. G., Hausmann, M., Johannsen, H., & Michelato, A.
 916 (1991). Chlorofluoromethane and oxygen in the Eastern Mediterranean. *Deep Sea Research Part A.*
 917 *Oceanographic Research Papers*, 38(12), 1531-1551.

918 Schneider, A., Tanhua, T., Roether, W., & Steinfeldt, R. (2014). Changes in ventilation of the
 919 Mediterranean Sea during the past 25 year. *Ocean Science*, 10(1), 1-16.

920 Schroeder, K., Chiggiato, J., Bryden, H. L., Borghini, M., & Ben Ismail, S. (2016). Abrupt climate
 921 shift in the Western Mediterranean Sea. *Scientific reports*, 6(1), 23009.

922 Schroeder, K., Gasparini, G. P., Tangherlini, M., & Astraldi, M. (2006). Deep and intermediate water
 923 in the western Mediterranean under the influence of the Eastern Mediterranean Transient.
 924 *Geophysical Research Letters*, 33(21).

925 Seager, R., Osborn, T. J., Kushnir, Y., Simpson, I. R., Nakamura, J., & Liu, H. (2019). Climate
 926 variability and change of Mediterranean-type climates. *Journal of Climate*, 32(10), 2887-2915.

927 Send, U., & Testor, P. (2017). Direct observations reveal the deep circulation of the western
 928 Mediterranean Sea. *Journal of Geophysical Research: Oceans*, 122(12), 10091-10098.

929 Sheen, K. L., Brearley, J. A., Naveira Garabato, A. C., Smeed, D. A., Waterman, S., Ledwell, J. R., ...
 930 & Watson, A. J. (2013). Rates and mechanisms of turbulent dissipation and mixing in the Southern
 931 Ocean: Results from the Diapycnal and Isopycnal Mixing Experiment in the Southern Ocean
 932 (DIMES). *Journal of Geophysical Research: Oceans*, 118(6), 2774-2792.

933 Somot, S., Sevault, F., & Déqué, M. (2006). Transient climate change scenario simulation of the
 934 Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model.
 935 *Climate Dynamics*, 27(7), 851-879.

- 936 Sparnocchia, S., Gasparini, G. P., Astraldi, M., Borghini, M., & Pistek, P. (1999). Dynamics and
937 mixing of the Eastern Mediterranean outflow in the Tyrrhenian basin. *Journal of Marine*
938 *Systems*, 20(1-4), 301-317.
- 939 Sparnocchia, S., Gasparini, G.P., Schroeder, K., Borghini, M. (2011). Oceanographic conditions in the
940 NEMO region during the KM3NeT project (April2006–May2009). *Nucl. Instrum. Methods Phys.*
941 *Res. A*, S87–S90. <http://dx.doi.org/10.1016/j.nima.2010.06.231> 626-627.
- 942 St. Laurent, L., Naveira Garabato, A. C., Ledwell, J. R., Thurnherr, A. M., Toole, J. M., & Watson, A.
943 J. (2012). Turbulence and diapycnal mixing in Drake Passage. *Journal of Physical Oceanography*,
944 42(12), 2143-2152.
- 945 Takahashi, A., Hibiya, T., & Naveira Garabato, A. C. (2021). Influence of the distortion of vertical
946 wavenumber spectra on estimates of turbulent dissipation using the finescale parameterization:
947 Eikonal calculations. *Journal of Physical Oceanography*, 51(5), 1723-1733.
- 948 Theocharis, A., Klein, B., Nittis, K., & Roether, W. (2002). Evolution and status of the Eastern
949 Mediterranean Transient (1997–1999). *Journal of Marine Systems*, 33, 91-116.
- 950 Thorpe, S. A. (2005). *The turbulent ocean*. Cambridge University Press.
- 951 Tsimplis, M. N., Zervakis, V., Josey, S. A., Peneva, E. L., Struglia, M. V., Stanev, E. V., ... & Oguz,
952 T. (2006). Changes in the oceanography of the Mediterranean Sea and their link to climate variability.
953 In *Developments in earth and environmental sciences* (Vol. 4, pp. 227-282). Elsevier.
- 954 Thurnherr 2012: The Finescale Response of Lowered ADCP Velocity Measurements Processed with
955 Different Methods. https://journals.ametsoc.org/view/journals/atot/29/4/jtech-d-11-00158_1.xml.
956 <https://doi.org/10.1175/JTECH-D-11-00158.1>
- 957 van Haren, H., & Gostiaux, L. (2011). Large internal waves advection in very weakly stratified deep
958 Mediterranean waters. *Geophysical Research Letters*, 38(22).
- 959 Visbeck, M. (2002). Deep velocity profiling using lowered acoustic Doppler current profilers: Bottom
960 track and inverse solutions. *Journal of atmospheric and oceanic technology*, 19(5), 794-807.
- 961 Waldman, R., Brüggemann, N., Bosse, A., Spall, M., Somot, S., & Sevault, F. (2018). Overturning the
962 Mediterranean thermohaline circulation. *Geophysical Research Letters*, 45(16), 8407-8415.
- 963 Walin, G. (1982). On the relation between sea-surface heat flow and thermal circulation in the ocean.
964 *Tellus*, 34(2), 187-195.
- 965 Waterhouse, A. F., et al. (2014). Global patterns of diapycnal mixing from measurements and models.
966 *Journal of Physical Oceanography*, 44(7), 1854–1872. <https://doi.org/10.1175/JPO-D-13-0133.1>
- 967 Whalen, C. B., De Lavergne, C., Naveira Garabato, A. C., Klymak, J. M., MacKinnon, J. A., &
968 Sheen, K. L. (2020). Internal wave-driven mixing: Governing processes and consequences for
969 climate. *Nature Reviews Earth & Environment*, 1(11), 606-621.
- 970 Wright, D. G., & Stocker, T. F. (1992). Sensitivities of a zonally averaged global ocean circulation
971 model. *Journal of Geophysical Research: Oceans*, 97(C8), 12707-12730.

972 Wüst, G. (1961). On the vertical circulation of the Mediterranean Sea. *Journal of Geophysical*
973 *Research*, 66(10), 3261-3271.

974