Dear Dr. Ian Delaney,

We thank you, Dr. Herreid, and Dr. Wild for the feedback provided and for the opportunity to further improve our manuscript. We revised the manuscript in response to the reviewers' suggestions.

Below is our updated point-by-point response to the comments (in blue), specifying the corrections made in the manuscript (in quotation marks).

The revised manuscript and a marked-up version of the manuscript have been uploaded.

We believe that these revisions have substantially strengthened the manuscript and enhanced its relevance to the broad readership of The Cryosphere.

Thank you and best regards,

On behalf of all coauthors,
Yulia Antropova
Carleton University

Response to “RC1: 'Comment on egusphere-2025-4771', Sam Herreid, 13 Jan 2026”

Dear Dr. Herreid,

Thank you for reviewing our manuscript and for your thoughtful and valuable feedback, which have helped us improve the manuscript’s quality and scope.

In response to your major comments, we have revised our discussion of IBE and removed the figure panels demonstrating the correlation as well as the reconstructed surface displacements without IBE. We have also added an IBE correction based solely on GEM, which may be useful for the broader glaciological community. In addition, we revised our terminology and the Discussion section based on the comments from both reviewers.

Our detailed responses to all comments are provided below in blue font.

Review for **Intra-annual grounding line migration and retreat: insights from high resolution satellite and in-situ observations over Milne Glacier in the Canadian High Arctic**

The article “Intra-annual grounding line migration and retreat: insights from high resolution satellite and in-situ observations over Milne Glacier in the Canadian High Arctic” by Antropova et al. presents a well organized narrative of both methodologies and and local results exploring the intra-annual changes of the flexure zone of a marine termination glacier in the Canadian High Arctic. Ground and airborne data are combined with a dense time series of SAR images to unpack what is happening not just at a high resolution throughout a year but also looking along three longitudinal transects that exhibit variability that the authors do a nice job of unpacking and quantifying. Finally, they put this work into a wider context by looking at the bed topography for where future pinning points might be and how they might affect subglacial melt.

Below are my main comments followed by in line minor comments.

The authors spend some time exploring the importance of including the inverse barometer effect (IBE), but I think this as been well established by Padman et al., 2003, Padman et al, 2008 and e.g. Chen et al., 2023. While quantifying and partitioning the relative terms is very useful, I’m not sure if IBE needs to be demonstrated as important against a simulation excluding it.

We agree that IBE is well documented already and have reduced the IBE discussion accordingly. We retained only three panels showing the correlation between “SSH” (i.e., IBE included) and “DInSAR displacement” (Fig. 4 d-f in the initial manuscript) and moved “DTide” (i.e., without IBE) vs. “DInSAR displacement” panels (Fig. 4 a-c in the initial manuscript) to the Appendix (Fig. A3 in the revised manuscript). We also removed panels related to reconstructed ice surface height based on simulated tides only from Fig. 6 in the initial version of the manuscript. We revised the text accordingly, removing the discussion of IBE from the Abstract and reducing it in the Results and Discussion sections.

The Discussion section was revised to highlight the differences between our study and previously published DInSAR studies that explored the IBE, as follows:

"The intra-annual vertical displacement of the Milne Glacier ice tongue is governed by changes in SSH, which, in turn, are driven by variations in tides and atmospheric pressure. SSH has been identified as a major driving mechanism of grounding-line migration for Thwaites Glacier in West Antarctica (Rignot et al., 2024), Fisher, Mellor, and Lambert glaciers in East Antarctica (Chen et al., 2023), and Jakobshavn Isbræ in Greenland (Kim et al., 2024). These locations have relatively large (i.e. ≥ 1 m) tidal ranges, therefore atmospheric pressure only plays a minor role in the displacement of floating ice shelves and ice tongues. For marine-terminating glacier systems similar to Milne Glacier, where tidal range is much smaller than in the aforementioned studies, atmospheric pressure variability plays a more substantial role in controlling SSH, and, consequently, the displacement of the floating ice tongue."

The authors use GEM to fill data gaps in the pressure record, this is somewhat of a footnote. I'm not sure if there are other studies that do this, but since the authors have a dense timeseries of supporting data it could be a useful result for the community to use only GEM data and see if the 15 x 15 km grid 9 km away can reproduce the same tidal model.

Model-based IBE corrections have been discussed previously in the literature, for example through the use of ECMWF atmospheric reanalyses (ERA5) by Chen et al. (2023) and Rignot et al. (2024); however, we are not aware of any direct examples using GEM model. In response to this suggestion, we added a figure demonstrating a solely GEM-based IBE correction in the Appendix (Fig. 1AR middle panels and Fig. A3 d-f in the revised manuscript) to facilitate comparison with the predominantly in situ-based IBE corrections and to make our analysis more useful to the glaciological community. Given that, in general, the in-situ-based IBE corrections tend to yield higher correlations (Fig. 1AR) and, in our view, add value to this case-based study we use our primary analysis based on in situ observations for analysis.




Figure 1AR: Correlation results calculated for the three transects. Upper panels (moved to the Appendix): DInSAR displacement and T_TIDE-simulated tides. Middle panels (added to the Appendix): DInSAR displacement and T_TIDE-simulated tides + GEM-based IBE. Lower panels (retained in the main text): DInSAR displacement and T_TIDE-simulated tides + in situ–based IBE.

There seems to be some inconsistencies throughout with regard to FZ and GZ, both in the terms used and what was calculated and how. I would ask the authors to read through this carefully and make sure terms are consistent throughout and clearly derived if used later in the paper.

We revised L52-54 and the caption of Fig. 1 to explain how the FZ and GZ terms are defined and calculated. As per request from the second reviewer, we adopted "tidal flexure zone" (equivalent to "flexure zone") and "grounding zone" terms and updated them consistently throughout the manuscript.

We note that there are terminology discrepancies in the literature stemming from differences between oceanographic usage and that of the DInSAR community, particularly for terms such as "grounding zone," "grounding line," and "hinge line." Numerous studies (e.g., Rignot, 1996; Rignot, 1998; Fricker & Padman, 2006; Fricker et al., 2009; Sykes et al., 2009; Hogg et al., 2016; Dawson & Bamber, 2017; Freer et al., 2023) define the "grounding zone" as a feature spanning from the hinge line, i.e., the inland limit of tidal flexure, past the grounding line to the line of hydrostatic equilibrium.

In contrast, more recent DInSAR-focused studies (e.g., Mohajerani et al., 2021; Chen et al., 2023; Ciracì et al., 2023) tend to refer to the DInSAR-derived landward limit of tidal ice flexure as the "grounding line" (rather than the hinge line), and to the region over which this landward limit migrates with the tides as the "grounding zone." In addition, the size of this zone is commonly reported using the term "grounding zone width," even though it is measured along the glacier flow direction.

In this manuscript, we adopt the following terminology to remain consistent with recent literature and to address the reviewers' comments:

Hinge line: the DInSAR-derived inland limit of tidal flexure.
Tidal-flexure zone or flexure zone, initially referred to as the "ice flexure zone": the zone of ice flexure due to tides, spanning from the hinge line to the line of hydrostatic equilibrium. The width of the flexure zone (initially referred to as the "FZ span") is measured along the three red transects (Fig. 1).
Grounding zone: the region over which the hinge line migrates between its most landward and most seaward positions in response to tides. The width of the grounding zone (initially referred to as the "GZ span") is measured along the three red transects (Fig. 1). The following text from the manuscript describes how it was calculated:

"The Milne Glacier grounding-zone width for the entire period of observations (2023 - 2024) was calculated as the difference between the maximum and minimum distances from point A to the most seaward and landward DInSAR-derived hinge lines, respectively, along the three transects (e.g., max $|A\ B_W|$ – min $|A\ B_W|$ for the western transect)."

We also note that the terms "hinge line" and "grounding line" can be used interchangeably when discussing DInSAR-derived short-term migration or retreat processes, as hinge-line and grounding-line migrations are directly correlated (Rignot, 1996; Hogg et al., 2016), as clarified in the Introduction.

There is a use of stable and unstable which I found to be imperfect terms for an intra-annual time series. In Section 4.2 the authors relax this language and talk in terms of season. I think the stable/unstable delineation might be more easy to follow and possibly more accurate to label in terms of seasons. A possible title could be "Seasonal grounding line migration (I'm not sure if there is a long enough time series here to identify permanent retreat?). Also in the title is "high resolution" high temporal resolution? I think this is really the key factor that make the study unique.

Thank you for the suggestions. We adopted the terms "steady" and "unsteady" periods to describe the analyzed velocity regimes and to address comments from both reviewers.
We would like to avoid the use of "seasonal" grounding line migration because we do not have data spanning multiple seasons to reliably identify seasonal regimes. Additionally, our division of periods is based primarily on changes in velocity and SAR data availability rather than on seasons (e.g., the summer season is not represented, and the fall and winter seasons overlap).

In the title, "high resolution" refers to both spatial and temporal resolutions. We would like to retain the title in its initial short form, but we have a revised a clarifying sentence in the abstract as follows:

"We explore flexure zone changes on Milne Glacier, Nunavut, Canada from 2023 to 2024 at a high spatial resolution of 10 m using 4-day RADARSAT Constellation Mission (RCM) data acquisitions from four different orbits."

The paper is largely a collection of substories, one of which is the notch which I had the hardest time following. Maybe there's a figure that could help the reader along.

We added Fig. 2AR (i.e., Fig. A2 in Appendix of the revised manuscript), which demonstrates examples of differential interferograms with and without the new eastern notch and pinning points.




Figure 2AR: DInSAR image examples from orbit 103 showing: a) the flexure zone without the eastern notch, b) the flexure zone with the newly formed notch around the eastern pinning point.

We also modified the caption of Fig. 3 to better clarify the connection between the new pinning point and the notch:

"A pronounced retreat associated with the formation of a new notch around the pinning point (Fig. A2) at the eastern margin was discovered in October 2023, after the March-April 2023 hinge line delineations by Antropova et al. (2024)."

Finally, as an intra-annual study, I think a nice addition to the discussion section could be a recommendation to future researchers, who might be limited to a more sparse dataset, on how they might be confident or cautious comparing data from different instances in the seasonal cycle.

We revised Section 4.2 of Discussion with respect to SAR-line-of sight errors due to velocity contributions and due to ice stretching and compression caused by bending (as per request from the second reviewer).

We added the following text regarding the reliability of DInSAR results, considering the quantified errors associated with the seasonal cycle and SAR viewing geometry.

"This uncertainty would increase if there was an even greater velocity variability. Additionally, larger tidal displacements and thicker ice can enhance errors associated with ice extension or compression caused by bending. For instance, the error due to ice bending can be on the order of one ice thickness based on a 1-D model (Rack et al., 2017). This effect can be inhibited within embayments and enhanced around protrusions of the grounding line, as suggested by a 2-D model (Wild et al., 2018). Thus, flexure-zone delineations from orbits with high incidence angles, which are sensitive to horizontal surface displacements, should be interpreted with consideration of the associated uncertainties when a glacier experiences substantial velocity variability and/or pronounced tidal bending."

L14-15: The language isn't quite correct here, either something like "Knowledge of changes at the boundary…is critical for…" or "Changes…the grounding line control glacier stability…"

Thank you. We have revised the sentence as follows:
"Observing and quantifying changes of the boundary where marine-terminating glaciers transition from grounded to floating, known as the grounding line, is critical for advancing our understanding of glacier stability."

L15: Is there a more succinct way to build directly off of the grounding line established in the first sentence to flexure zone rather than seemingly developing a new term and then at the end of the sentence connecting it back to grounding line? Within a system with a flexure zone, isn't there not an "actual grounding line" location but rather a moving will occupy throughout an observed tidal maximum and minimum?

We revised the text as follows:

"The grounding line is difficult to observe directly but can be inferred from the ice surface tidal-flexure zone. This zone spans from the inland limit of tidal flexure — the hinge line, across the grounding line to the hydrostatic equilibrium line, where the ice tongue becomes freely floating."

We note that flexure zone is the zone where ice experiences flexure with tides, and the grounding zone, where the DInSAR-derived hinge line moves with tides (in line with the DInSAR nomenclature).

L20-L21: measured or modeled tides? Was atmospheric pressure implicit or explicit?

Tides were simulated using the T_TIDE algorithm based on our 10-month in-situ pressure records in 2016-2017 (Antropova et al., 2024). The atmospheric pressure was implicit (i.e., it becomes "unexplained variation" not accounted for by the tidal constituents) in the simulated tides, but then it was added through the inverse barometric effect (IBE) correction to get sea surface height (SSH).

The following clarifications were added to Section 2.2.2:

"The vertical component of DInSAR-derived surface displacement of the floating part of the glacier was analysed with respect to the changes in sea surface height (SSH) due to tides and atmospheric pressure. We used tidal constituents derived from in situ tidal observations recorded every 6 seconds with an RBR Duet datalogger deployed at ~10 m depth in a lateral bay on the western side of Milne Fiord (i.e., tide gauge location in Fig.1 a) between September 3, 2016 and July 4, 2017 to simulate tides at a 1-minute interval for the period of SAR observations using the T_TIDE toolbox (Pawlowicz et al., 2002). Simulated tide heights were corrected by adding the inverse barometer effect (IBE) (Wunsch & Stammer, 1997; Padman et al., 2003), while ocean tide loading on the Earth's crust was neglected, as it has a relatively minor impact (Wild et al., 2025)."

L22: 0.81, 0.94 m?

These numbers represent Pearson's correlation coefficients; therefore, they are dimensionless. This information was removed from the revised Abstract.

L22-25: Is the vertical displacement a key result of your project? I would think the key point to get across is sensitivity in z to determine a meaningful x,y. The following sentence is unnecessary detail for the an abstract in how it's presented here. As it reads, here it should just be +/- error bounds for 0.81 and 0.94, but I would argue that these numbers aren't so fundamental to the study and the sentence of interest here is z accuracy as it relates to x,y accuracy.

We revised the Abstract to emphasize horizontal (x,y) uncertainty due to SAR line-of-sight (z) error, as follows:

"The most reliable flexure zone delineation (horizontal uncertainty of ±8 m) was achieved with ascending RCM tracks at low incidence angles during periods when the ice velocity was steady and bending was not substantial, resulting in a combined SAR line-of-sight error of 0.13 cm. A worst-case flexure-zone delineation uncertainty of ±222 m was noted when a descending track with a high incidence angle was used and glacier velocity became more variable, increasing the combined SAR line-of-sight error to 3.26 cm."

L25: Maybe you mean the downstream edge of the GZ, presumably there is a gradient somewhere in the upper boundary of the GZ.

This statement was removed from the abstract. It was based on Fig. 11 and its discussion:

"The grounded ice moved downstream at about 100 m yr^{-1} and then accelerated to 200 m yr^{-1} at ~3 km up-glacier of the FZ, reaching a velocity of 270 m yr^{-1} in the FZ and then up to 320 m yr^{-1} down-glacier of it during summer months. Hence, Milne Glacier FZ zone was associated with ~2.7-fold increase of glacier velocity compared against its upglacier grounded part shown in Fig. 11."

L25: How should a reader interpret this result, is this abnormal? A change? Did this seasonal cycle drive the retreat in the following sentence?

We replaced this sentence with information about line-of-sight errors and the corresponding horizontal errors due to velocity in the DInSAR-derived bounds (see the L22–25 comment above).

L25: new meaning previously unreported or newly developed within some amount of time?

Corrected to "newly developed" as this grounding line retreat was observed from our DInSAR images within our 2023-2024 period of observations.

L26: Why grounding line here and not GZ?

We believe that both GZ and grounding line would be appropriate in this context; however, we prefer to use only the terms 'grounding line' and flexure zone already introduced in the Abstract to avoid reader confusion arising from the terminology discrepancies mentioned above and to save space in the Abstract.

L27: "...shows how bed features can alter the grounding…"

The sentence was revised as follows:

"Low-error DInSAR images enabled the detection of a newly developed ~2.4 km grounding-line retreat associated with water intrusion behind a pinning point at the eastern margin of the glacier. Our results underscore that care must be taken to ensure that the low-error DInSAR results are used and that pinning points alter the grounding-line short-term migration and its long-term retreat."

L31: Add citation(s)

Three citations: "(Black & Joughin, 2022; Kochtitzky & Copland, 2022; Rantanen et al., 2022)" were added.

L31: Establish this connection: Rising atmospheric and ocean temperatures can cause marine-terminating glaciers to speed up, which then leads to …

The sentence was revised as follows:

"Rising atmospheric and oceanic temperatures can cause these glaciers to accelerate, leading to increased sea level rise and discharge of icebergs or ice islands into adjacent waters (Morlighem et al., 2019; Wood et al., 2021; Carnahan et al., 2022; Holmes et al., 2025)."

L32: Check formatting of IOCC ref. Is it 2021 or 2023?

The "IPCC (2021)" citation was replaced with original research articles from peer-reviewed journals, as requested by the second reviewer: "(Morlighem et al., 2019; Wood et al., 2021; Carnahan et al., 2022; Holmes et al., 2025)."

L33, try to establish the location for the reader, something like, "In the Arctic, …"

We modified the first sentence of the corresponding paragraph to emphasize that we are narrowing its focus to Arctic-related research, which is more relevant to our area of interest:

"Marine-terminating glaciers, which drain ice masses from land to the ocean, have experienced dramatic changes in recent decades due to a warming climate, particularly in the Arctic, where temperatures are rising faster than the global average (Black & Joughin, 2022; Kochtitzky & Copland, 2022; Rantanen et al., 2022)."

L33: What was the change? Faster and smaller?

We revised the sentence to clarify the documented changes as follows:

"Increased ice discharge and ice mass losses associated with Arctic glacier volume decrease, retreat, and deterioration were recently documented by Millan et al. (2017, 2023) and Van Wychen et al. (2020)."

L38-39: Suggest: …exhibit dynamic behaviour at two key scales, ysub-day variability from tides and [insert some time scale, e.g. years to decadal] changes as a response to climate and glacier dynamics.

Thank you, but we prefer to retain the original text, as distinguishing between timescales in this context is uncertain due to tidal variations spanning sub-daily/daily cycles (e.g., semi-diurnal and diurnal) to longer-term cycles (e.g., spring-neap-spring ~14 - 15 days, perigean-apogean ~27 days, seasonal equinox events, lunar nodal ~18.6 years) and sea-level rise. Similarly, defining an appropriate timescale for grounding line retreat due to long-term glacier changes is problematic, as it varies greatly with environmental and glacier-specific factors, which interact and can result in a rapid and pronounced grounding line retreat, as documented in this study.

L41: maybe effective rather than efficient

Thank you - we have modified it to "effective," as it is more appropriate in this context.

L45: Yes, Fricker at al, uses grounding zone for flexure zone, Freer et al, in their Fig. 1 do differentiate their points, F, H as the flexure zone proxy but also F_max and F_min, the limits of the GL. I would encourage the authors to affirm the language flexure zone for what DInSAR measures, the ice grounding zone for the true max and min of tidal migration of the GL (ice used here to differentiate from an oceanographers definition of the grounding zone which I believe is slightly different from a glaciologist's perspective), finally the grounding line referred to not a fixed location but moving both diurnally with tide and over years with glacier-wide and climate changes. I think the authors are mostly holding to this, but it's not perfectly concrete. (your L48 and 50-54 are very good and clear). I think you could just define FZ and GZ, I'm not sure what span adds.

Our study assumes that a single DInSAR image can be used to delineate the "tidal flexure zone" that occurs between three or four SAR data acquisitions. This zone spans from the inland limit of

the fringe pattern (i.e., hinge line) to its seaward limit (i.e., hydrostatic equilibrium line), whereas multiple DInSAR image patterns can be used to monitor the landward and seaward bounds of hinge line migration referred to as "grounding zone" in the most recent glaciological studies (e.g., Mohajerani et al., 2021; Chen et al., 2023; Ciracì et al., 2023) and in our study. Given the availability of frequent DInSAR images from different orbits with different acquisition times in our study, we were able to quantify a reliable DInSAR-derived grounding zone width, which serves as an adequate proxy for the true "Fmax-to-Fmin" distance, occurring between maximum and minimum tides modulated by atmospheric pressure. We hope that this explanation, together with our response to the terminology question above, fully clarifies our use of terminology in the manuscript.

L47: This "however" is a little misplaced I think, make sure there is clear conceptual contrast.

We replace it with "Nevertheless" to emphasize the coexistence of definitions.

L49, 58, Chen et al., 2023 is missing in References

Thank you, we added the citation to the list.

L59 GZ spans for consistency.

We prefer to retain "grounding zones" rather than "grounding zone widths" here, as in this context we discuss melt across the entire grounding zone.

L61: I think the objective is to understand the controls and responses of a system that happens to be deteriorating. We hope these observations will work in a phase of glacier growth also.

Thank you, we revised the sentence to refine our objectives:

"To improve these models, a detailed investigation of the drivers and glacier-system responses governing grounding-line migration over short-term cycles and its long-term advance or retreat is required. The reliability of satellite-based observations used to monitor these processes should also be thoroughly evaluated."

L65: Generally would say model results or inputs not model datasets

Thank you; this has been revised as follows:
"Our study explores intra-annual changes in the Milne Glacier flexure zone based on a combination of RCM DInSAR-derived results, in situ and simulated datasets, and a numerical weather prediction model output."

L64,66 if using FZ "span", use throughout

FZ (changed to "flexure zone") refers to the entire zone of ice flexure delineated based on the DInSAR fringe pattern from one DInSAR image/result here. "Span" was changed to "width" in the current version of the manuscript.

L70: what does marked mean? Could just say acceleration and no acceleration for a and b. Is it acceleration and deceleration that makes an unstable regime?

We revised these sentences to clarify the details as follows:

"We further assess this correlation across the four SAR orbits with respect to the compressional and tensional strains caused by ice-surface bending in the flexure zone during two velocity periods:
a) a steady period, when the in situ-measured Milne Glacier velocity did not change appreciably during winter and the DInSAR assumption of invariant velocities was met; and
b) an unsteady period, when velocity changes were evident, the glacier decelerated after summer peak velocity, and the DInSAR assumption of invariant velocities was violated."

Please note that "steady/unsteady period" terminology was adopted instead of "stable/unstable regime".

L71: In 64-66 you establish intra-annual as a short term, is this still the same short term? Or do you mean 1d or 4d changes?

We removed "short-term" and revised the first objective to avoid giving the impression of establishing full equivalence between the intra-annual and short-term timescales, and to provide further details about the time scale:

"Delineate the Milne Glacier flexure zone using RCM data acquisitions from four different orbits, with a 4-day repeat interval within each orbit and about a 1-day interval between orbits, spanning from February 17, 2023 to September 17, 2024;"

We note that, although changes in flexure zone occur in response to SSH (i.e., tides plus IBE), our definition of a "short-term" period is limited to DInSAR observations generally with a 4-day repeat interval within each orbit and about 1-day interval between orbits; for example, in January of 2024, the coverage was: orbit 103: Jan. 5, 9, 13; orbit 43: Jan. 9, 13, 17; orbit 58: Jan. 10, 14, 18; orbit 09: Jan. 11, 15, 19. The RCM acquisition timeline throughout 2023-2024 is shown in Fig. 2a.

L93-95: Does Van Wychen et al, provide a mechanism for these velocity changes? Obviously velocity context is useful, but will the speed up slow down cycle come back into the story here?

Thank you, we added a clarifying transitional sentence to show the connection between changes in velocity and grounding line retreat and discuss mechanisms for these velocity changes:

"Van Wychen et al. (2020) suggested that this cyclical behaviour may result from prominent bedrock features that modulate velocity where the glacier rests below sea level. In contrast, Antropova et al. (2024) noted that the post-2016 acceleration was preceded by the highest grounding-line retreat rate along the glacier's centerline between 2011 and 2018, suggesting the presence of feedback mechanisms between grounding-line retreat and upstream glacier acceleration."

We are working on a follow-up paper, which confirms glacier thinning above the grounding zone. We are testing the hypothesis that dynamic thinning, associated with a velocity increase in this zone and likely triggered by grounding line retreat, is involved. However, we consider this outside of the scope of this current paper.

L96-98: There isn't a citation here, is this your result? Add citation or move to results.

These results are from Antropova et al. (2024). Since this work is already cited in the previous and following sentences, we prefer not to add another citation here.

L123: automated or manually delineated?

"manually" was added.

L129: It's possible resampling the DEM to a lower resolution can help maintain coherence with the crevasse pattern in the interferograms, just a suggestion for the future.

Thank you. The ArcticDEM resampling could be considered in the future, but based on our experience with 30 m ASTER and GLO-30 Copernicus DEMs, these areas were also affected by noise, when these lower-resolution DEMs were used.

L134: It's not clear which dataset actually defined the three transect lines. I think figure 1 could benefit from an inset with the surface and bed airborne and in situ radar transects and the IceBridge flightlines.

We added the IPR and IceBridge transects to Fig. 1 and the following text: (Fig. 1 a and Fig. 10 in the Results section)" to direct readers to the figure presenting surface and bed airborne and in situ radar transects.

L138-L138, Isn't the distance from Bw1 to Bw2 a change over time, your migration term not the GZ?

Yes, we used the distance from B_{W1} to B_{W2} as a simplified example based on only two DInSAR images to illustrate the concept behind the term. However, the DInSAR-derived grounding-zone width reported in this study was calculated using all DInSAR images over the entire observation period.

We revised the sentence as follows to better explain this idea:

"- grounding-zone width (i.e., the distance along the glacier between the most landward and most seaward hinge-line delineations based on all DInSAR images over the entire observation period). For illustrative purposes, the grounding zone is shown by the distance between BW1 and BW2 in Fig. 1, based on DInSAR image 1 and 2 associated with data acquisitions in March-April 2024 and September 2024."

L141: Was the unwrapped phase calibrated to the surrounding bedrock? What is a phase discontinuity? A phase jump? How was this corrected?

We calibrated the unwrapped phase with respect to the mean value of the grounded ice surface as follows: for orbits 43, 58, and 103, the grounded ice surface means were negligible; therefore, correction for the incidence angle alone was sufficient (i.e., dividing the derived displacement by the cosine of the incidence angle). For orbit 09, an additional calibration relative to the grounded ice surface mean was applied.

Yes, this sentence refers to phase jumps, also referred to as discontinuous shifts in the measured phase. These were corrected using a MATLAB script that identified and removed phase jumps exceeding 2 cm along the three transects. This threshold was chosen to be slightly more conservative than half of the RCM C-band wavelength (i.e., 5.5 cm / 2).

We revised the sentence to better clarify these processing steps as follows:

"The final calibration of the SAR line-of-sight displacement was conducted along three transects (Fig. 1, described below) using an automated script. Values were divided by the cosine of the incidence angle and additionally calibrated to the mean values of the grounded portions of the transects for each DInSAR image from orbits 09 and 103. Phase jumps exceeding 2 cm (i.e., a more conservative threshold than half of the RCM C-band wavelength) were corrected for all orbits."

L148: change and to are. In this study are…

We revised the text to incorporate this change and improve readability:

"Flexure-zone variables explored in this study are computed along the western, central, and eastern transects (subscript W, C, and E) shown in red (through points A, B and C). **These variables include:** 1) distance to hinge lines (A to BW1, A to BC1, A to BE1 for flexure zone in DInSAR image 1, blue polygon, and A to BW2, A to BC2, A to BE2 for flexure zone in DInSAR image 2, orange polygon), 2) width of the two flexure zones (BW1 to CW1, BC1 to CC1, BE1 to CE1 and BW2 to CW2, BC2 to CC2, BE2 to CE2), 3) Grounding-zone width (BW1 to BW2, BC1 to BC2, BE1 to BE2), and 4) surface displacement height of the floating ice."

L174: The notch isn't perfectly clear to the reader at this point in the article.

We added the following information to Section "2.2.2 Analysis of tides and reconstruction of ice-surface height timeseries":

"To explore the origin of the newly developed notch along the eastern margin, we tested the hypothesis that it appears in DInSAR images associated with SAR acquisitions during periods of high SSH, suggesting that seawater intrudes farther inland and results in ice ungrounding around a pinning point. We calculated the minimum and maximum SSH values, as well as the standard deviation, for each DInSAR result associated with three or four SAR data acquisitions. These values were then divided into two groups corresponding to: (1) DInSAR results without the notch

and (2) DInSAR results with the newly developed notch present, to test whether SSH differs significantly between the groups."

We also added a new figure demonstrating DInSAR results with/without the notch, as stated above, to the Appendix

L180: It looks like the unstable period is the white block between sable-blue and unstable-red. And why isn't all of the flat velocity before the beginning of the first blue block considered stable? It seems to me these are environmental classifications that are actually instrumental classifications, where data was available and where there was coherence. I would encourage the authors to say something like "winter observation block" and late-summer/fall observation block" otherwise this is very confusing for the reader. Layer L191 admits this.

Yes, this breakdown was based on SAR data availability and glacier velocity, as clarified in the text (previously L191).

We revised the caption of Fig. 2 to better explain this:

"**The winter SAR observation block** associated with a steady glacier period, highlighted in blue, corresponds to relatively steady and low velocity values of 0.4 m d^{-1}. **The fall SAR observation block** associated with an unsteady period, highlighted in orange, corresponds to velocity values decreasing from 0.6 m d-1 to 0.4 m d^{-1}."

L195: It is technically "not stable" because there is change in velocity, but I wouldn't call a predictable seasonal slowdown as "unstable"

We adopted "steady/unsteady" terminology in the current version of the manuscript.

L227: A height range converted to a spatial extent using ArcticDEM?

L227 refers to the DInSAR-derived grounding zone located between the most landward and most seaward hinge-line positions. Its width was calculated as the distance between points with XY coordinates at the intersections with the three red transects: first, the distance from point A to the most seaward hinge-line position was measured; next, the distance from point A to the most landward position was measured; finally, the difference between these two distances was computed to characterize the grounding zone width.

We revised the sentence to better clarify this:
"The Milne Glacier grounding-zone width for the entire period of observations (2023 - 2024) was calculated as the difference between the maximum and minimum distances from point A to the most seaward and landward DInSAR-derived hinge lines, respectively, along the three transects (e.g., max |A BW| – min |A B W| for the western transect)."

L246: Don't you have bed geometry and ice thickness (L220), why not explore this rather than leave as speculative "likely due to a distinctive bed geometry and ice thickness"

This sentence was removed from this section, as the corresponding clarifications are provided in the Discussion section.

L283: Can this or a plot like this include the pinning point(s)? It would be cool to see how the Eastern transect changes

We have added this figure to the Appendix (as noted above) to better illustrate the DInSAR results with and without the notch. This figure demonstrates that the primary changes at the ice surface occur farther east of the eastern transect.

As shown in Figure 3b, the changes observed from orbit 43 at the ice surface along the eastern transect are not pronounced. The hinge line migrates only approximately 400 m landward (from km 39.4 to km 39 along the IceBridge transect). Therefore, we retain the current version of the figure, as adding vertical lines to indicate the most landward hinge-line positions, with and without the notch, would not substantially improve its clarity.

We also refer to Figure 9b to highlight the bed rises, suggesting that the glacier remains grounded near these features and farther east (between km 39.5 and 40.5, labeled as the "pinning point region"), when high tide allows water to intrude farther inland.

L294: Maybe I missed it, what is the reference height SSH is based on?

SSH is derived from T_TIDE-simulated tides referenced to mean sea level and subsequently corrected for the inverse barometer effect (IBE), assuming that a 1 hPa increase (decrease) in atmospheric pressure results in a 1 cm drop (rise) in sea level (Wunsch & Stammer, 1997; Padman et al., 2003). In the T_TIDE MATLAB package (Pawlowicz et al., 2002), mean sea level is determined as the mean level of the input data series, recorded by our pressure sensor. Then T_TIDE computes amplitudes and Greenwich phase lags based on the deviation from this mean. Therefore, our simulated tides were reconstructed relative to this in situ mean baseline (considered as mean sea level) recorded by our pressure sensor.

L303: Table 3, It's essentially a new fringe or two that you're saying is a persistent feature and thus not error. From this table it seems possibly within error but if it's spatially consistent this is maybe a better argument. Why no middle value along with SD and max min, and the first column says averaged across all observations, but really divided between sets of notch and no notch?

Yes, the surface changes are minimal (shown in Fig. 5c and Fig. A2 in the Appendix); however, they are spatially consistent and persistent across different orbits, as shown in Figure 3.
The purpose of this table was to clarify the origin of the newly developed notch in the eastern margin, specifically to test our assumption that the notch (i.e., ungrounding around the pinning point) occurs during high SSH (tides+IBE). For this reason, we examined two categories of DInSAR images: with and without the notch. The occurrence of the notch during periods of higher SSH maxima (0.21 m vs. 0.13 m for observations with and without the notch, respectively) supports our interpretation that it caused by seawater intrusion.

We revised the sentence to better explain this:

"Our hypothesis that the newly observed notch formed during periods of high SSH, leading to seawater intrusion and ice ungrounding around the eastern pinning point, was evaluated using statistical characteristics of SSH (i.e., sea surface heights at the time of SAR data acquisitions rather than their displacement values of DSSH) along the eastern transect. The calculated characteristics for the DInSAR results were divided into two groups 1) cases without the notch, and 2) cases when the notch was observed (Table 3)."

L318-319: Is the loss of correlation a suggestion to future scientists that acceleration/decelation lowers the confidence of DDInSAR measurements? Is there also something we're learning about this particular glacier? It's read like a result for the particular glacier but I think the key result is methodological.

Thank you, we added the following clarifying sentence to better explain the associated methodological issues:

"This example underscores that pronounced changes in velocity violate the DInSAR assumption of invariant velocities between SAR data acquisitions and reduce the correlation between DSSH and DInSAR displacement. Therefore, the DInSAR images should be interpreted with consideration of errors arising from variable velocity."

L380-384: I'm not sure if I recall the authors giving their method to derive GZ from the data.

The following sentence (also discussed above) clarifies this:

"The Milne Glacier grounding-zone width for the entire period of observations (2023 - 2024) was calculated as the difference between the maximum and minimum distances from point A to the most seaward and landward DInSAR-derived hinge lines, respectively, along the three transects (e.g., max |A BW| – min |A B W| for the western transect)."

L385: East and West is reversed

Thank you. We have revised the figure references to direct readers to the appropriate panels corresponding to the West and East sections.

L386: Did you calculate this flux or is it just assumed based on a roughly parallel bed profile to cross-sectional ice surface?

This text has been revised to provide additional information:

"The widest grounding zone along the western transect of the glacier was associated with the highest velocity of 166.4 m yr^{-1}, averaged over a year and across the observed grounding zone width (Fig. 10 c), and the thickest ice (mean value of 225.6 m, Fig. 9 d). This section discharged approximately 37 540 m3 yr^{-1} of ice along a 1 m-wide flow line. The narrower and thinner grounding zones along the eastern and central transects, with mean thicknesses of 210.0 m and

200.5 m (Fig. 9 b, c), were associated with slower average velocities of 144.6 m yr^{-1} and 159.8 m yr^{-1} (Fig. 10 a, b), discharged approximately 30 366 m3 yr^{-1} and 32 040 m3 yr^{-1} along a 1 m-wide flow line, respectively."

L393-401: I think the role of IBE is well established in Padman et al., 2003 (cited but missing in references), also Padman et al., 2008, and as you cite Chen et al., 2023. Is/was there an assumption that the tidal effect would be different for marine terminating glaciers in the Canadian Arctica?

Thank you. As discussed above, we reduced our discussion of the IBE. While we acknowledge the fact that the IBE has been well explained in the literature with respect to SSH and is expected to be more pronounced where tidal ranges are small, our aim was to demonstrate that IBE plays a critical role in DInSAR-derived surface elevation changes for glacier tongues flowing into fjords with small tidal amplitudes, such as Milne Fiord.
DInSAR-related studies include IBE corrections, however, most of them focus on ice shelves and glacier tongues in Antarctica or Greenland, where the modeled tidal range is larger (approximately ±1 m) than in our study; consequently, the relative contribution of IBE is smaller in these cases. For instance, Rignot et al. (2024) demonstrated for Thwaites Glacier that the modeled tidal range (–80.8 cm to 96.5 cm) changed by only about 6% after including IBE (–93.5 cm to 95.1 cm). Similarly, Chen et al. reported a tidal range (IBE included) of approximately –90 to 112 cm and showed that applying the IBE correction improved the model fit (reducing the root-mean-square error from 0.932 to 0.780). Wild et al., 2025 reported modeled tides of ±0.4 m and the IBE, of ±0.15 m at Priestley Glacier.
In contrast, for Milne Fiord, the simulated tidal range spanned from –20 cm to 21 cm over the observation period. After correcting for IBE, the range increased 2.3-fold, spanning from –44 cm to 50 cm, which substantially improved the correlation between the SSH and DInSAR derived surface displacements shown in Fig. 4 of the original version of the manuscript.

L410-412: The grounding line migration of 800m wasn't explained or reported earlier. This should be a result and I'm not sure I know how the grounding line analysis was done.

In the Introduction we stated: "We also assume that the relative movements (e.g., short-term migration and long-term retreat) of the DInSAR-derived 'hinge line' and the actual 'grounding line' are equivalent, as their positions are strongly correlated (Rignot, 1996)." Therefore, our grounding line analysis with respect to its migration was based on the DInSAR-derived hinge line positions.

We moved L410-412 to the Results section and provided additional details (800 m was based on observations from orbit 09 only):

"The grounding line shifted up-glacier by 1 km compared to the most landward position observed during March 27-April 4 2023 in our previous study (i.e., Antropova et al., 2024), and exhibited a shift of approximately 690 m along the eastern transect between the most landward hinge-line positions derived from all DInSAR images with and without the notch in this study. This shift was even more pronounced farther east, where the notch advanced landward by 2.5 km compared to the last reported position and by 2.4 km (measured parallel to the eastern profile) for observations with and without the notch during the period covered by this study."

The sentence in the Discussion part was revised as follows:

"Milne Glacier's topography within its grounding zone controlled the grounding zone width over the period of observations: the newly detected notch, associated with a pinning point, caused a 30% increase in the grounding-zone width along the eastern transect."

L420: Do you mean for future predictions of pinning points? Because for a general understanding of active pinning points we don't need detailed information of bed geometry, just good DDInSAR.

Yes, thank you, we revised the sentence to better communicate this idea:

"Therefore, detailed information on bed geometry up-glacier of the grounding zone is essential for improving predictions of grounding-line retreat, particularly for identifying where potential pinning points may form due to water intrusion and ice ungrounding around bed rises."

Citation: https://doi.org/10.5194/egusphere-2025-4771-RC1

Response to “RC2: 'Comment on egusphere-2025-4771', Christian Wild, 21 Jan 2026”

Dear Dr. Wild,

We thank you for reviewing our manuscript and for providing a thoughtful, detailed, and insightful critique of our approach to exploring glaciological processes. In response to your major comments, we conducted: 1) additional ice-bending analysis, 2) improved ice surface displacement reconstruction, and 3) revision to our analysis and discussion of flexure-zone and grounding-zone widths and their landward positions. We hope that these changes, along with minor revisions, have improved both the quality and scope of the manuscript, and, in particular, its relevance to the cryospheric modeling community.

Our detailed responses to all comments are provided below in blue font.

Review of Antropova et al., submitted to The Cryosphere Discussion
Intra-annual grounding line migration and retreat: insights from high-resolution satellite and in-situ observations over Milne Glacier in the Canadian High Arctic
By Christian T. Wild
January 2026

Summary:
Antropova et al. present a detailed differential interferometric SAR (DInSAR) analysis of the tidal-flexure zone of Milne Glacier. Using a high-temporal-resolution RADARSAT dataset with 4-day repeat intervals, combined with in situ observations, they test the commonly assumed steady-flow condition applied in many DInSAR grounding-line studies. They also show that intra-annual grounding-line retreat occurs in a spatially and temporally heterogeneous pattern. They link this variability to subglacial topography and the development of localized pinning points, and offer speculations about the implications for basal melt rates and the potential for upstream ocean-water intrusion beneath the ice.

Strengths:
The authors demonstrate strong command of the DInSAR technique from a remote-sensing perspective and reveal clear, instructive examples of well-known processes associated with tidal flexure of ice shelves, which I elaborate on below. I have rarely seen such a thorough quantification of the steady-flow assumption, nor such careful consideration of how different satellite tracks (and associated variations in incidence angle and azimuth) capture the tidal-flexure signal. For this, I commend the authors. I particularly appreciate their use of available GNSS data for validation. However, because the GNSS measurements may be temporally biased (limited to a three-hour acquisition window per day, L183, which may also explain why they yield substantially higher ice velocities than the SAR-derived estimates in Fig. 2a), I wonder whether the analysis should be extended to include SAR-derived velocities alone. This would make the approach more transferable to settings where GNSS data are unavailable and could provide a clearer spatial picture of how the steady-flow assumption performs across the tidal-flexure zone, where velocity gradients are most pronounced towards their lateral margins.

Thank you for your feedback on our study and for recognizing its strengths in the remote sensing component and the use of in situ observations. We consider the use of in situ observations, including GNSS-derived velocities, to be an essential component of this work. We assume the GNSS-derived velocity records to be representative of grounding-zone velocities along the glacier centreline. We have clarified our use of GNSS records in our response below as well as in the manuscript. In addition, we improved our discussion of temporal biases with respect to SAR data acquisition times and quantified uncertainties associated with the derived velocities and tidal bending.
Although SAR-derived velocities provide promising research avenues, we consider this analysis to be outside the scope of the current manuscript. Therefore, SAR-derived velocities are used mainly to support our GNSS-derived velocities, which have much better temporal resolution and are well suited to our DInSAR analysis.

Weaknesses:
The study lacks a clearly defined glaciological hypothesis and instead reads largely as an enumeration of observations that have been explored extensively in the past.. Similar datasets have already been used to address, among other things: (1) improvements to tidal models (Wild et al., 2019), (2) the depth of the neutral layer in a deforming beam, (3) effective ice stiffness (Wild et al., 2018), (4) shear-margin strength (Wild et al., 2025), (5) two-dimensional effects on flexural patterns (Rack et al., 2017; Wild et al., 2018). Many of these applications have been published in this journal, yet the authors appear to be insufficiently engaged with this literature. This is reflected in an incomplete reference list, shortcomings in the introduction, and at times imprecise or incorrect terminology.
Given these issues, and considering that most recent glaciological DInSAR studies in *The Cryosphere* now combine DInSAR with multi-sensor approaches (Li et al., 2023), machine-learning methods (Ramanath et al., 2025), broad spatial (Picton et al., 2023) or temporal coverage (Chien et al., 2026), or numerical modeling of tidal flexure (Ross et al., 2024; Wild et al., 2025), I recommend that this manuscript may be more suitably placed in a more method-focused journal such as IEEE or a similar outlet. In that context, it would serve as an excellent technical reference that showcases the paper's considerable methodological strengths. If the manuscript were to be framed more explicitly within a glaciological scope, it could be strengthened by a robust investigation of short-term ice-flow dynamics and their relationship to changing buttressing as pinning points emerge and evolve.

Thank you for the constructive critique and suggestions. We agree that our study has an observational remote-sensing focus. It provides a thorough investigation and quantification of SAR-detected changes in a tidal-flexure zone from four different orbits at a high spatial and temporal resolution, which have not previously been explored in the literature. Our comprehensive dataset and approach contribute to a better understanding of the impact of variable glacier velocity with respect to different SAR viewing geometries on the delineation of SAR-derived tidal flexure-zones. In addition, we explore changes in the tidal flexure zone with respect to environmental and glacier-specific factors. Our advances in these regards should be of interest to the broader glaciological community and well-aligned with The Cryosphere's scope. Therefore, we believe that our study is a better fit for The Cryosphere than for IEEE or a similar outlet.
We acknowledge that the initial version of the manuscript did not explicitly discuss the processes examined in detail by Wild et al. (2018, 2019, 2025) and Rack et al. (2017). To address comments

regarding the processes acting in the glacier flexure zone and to situate our study more clearly within a glaciological context, we have:

- incorporated an analysis of ice compression and extension effects due to bending (i.e., Major Comment 1 below),
- included all sinusoids in the unraveling/reconstruction (i.e., Major Comment 2 below), and
- revised our analysis and discussion of flexure-zone width and hinge-line landward migration, referred to as the "distance to hinge line" (i.e., Major Comments 3 and 4 below).

We also recognize the recent trend in DInSAR-based grounding-line studies toward automated machine learning methods, the use of multiple sensors and/or broader spatial coverage, and integration with modeling, particularly in The Cryosphere. However, our high-temporal-resolution observations of the glacier flexure zone, which explain the short-term drivers of grounding-line migration and suggest the mechanisms of its long-term retreat due to ice deterioration around pinning points, are also of interest to the cryospheric community. In particular, our integrates multiple data types, presents high-spatiotemporal-resolution observations of glaciological processes from a SAR platform, offers insights about glaciological application of DInSAR methodology and its steady flow assumption with respect to SAR viewing geometry, and has implications for glaciers well beyond our study site.

In addition, we considered machine learning methods for automated flexure-zone delineation introduced by Mohajerani et al. (2021). However, this approach has limitations and can be problematic in areas with non-standard fringe patterns, such as pinning points, particularly in high-noise areas that are close to decorrelation (e.g., Fig. 9 in Ramanath et al., 2025). Therefore, we opted for a manual delineation approach to accurately document the emergence of a pinning point on the eastern margin of Milne Glacier.

We also note that many recently published grounding-line studies, including those cited above (Li et al., 2023; Ramanath et al., 2025; Picton et al., 2023; Chien et al., 2026; Ross et al., 2024; Wild et al., 2025) focus on Antarctica and/or Greenland. Our work contributes to improved representation of Arctic glaciers outside of Greenland, that are relatively understudied. These glaciers may serve as analogues for the future evolution of marine-terminating glaciers in Greenland and Antarctica, given their rapid retreat and disintegration, as discussed in our Introduction.

Regarding the comment that our terminology was at times "imprecise or incorrect," we attribute this primarily to inconsistencies in terminology across the literature (i.e., discrepancies between the oceanographic grounding zone and DInSAR grounding zone as detailed in our responses to terminology-related comments from both reviewers), rather than to insufficient engagement with studies published in The Cryosphere. We have revised the terminology throughout the manuscript to address these concerns and to clarify our usage.

The additional analyses addressing the major comments, together with the revisions to the manuscript text addressing the minor comments, have substantially strengthened the paper and enhanced its relevance to the broad readership of The Cryosphere.

Major comments:

1) <u>The bending effect introducing spurious signals in the tidal-flexure zone</u>




Figure R1: (left) from Wild et al., 2018, (right) from Rack et al., 2017. Red illustrates compression, blue horizontal extension. Arrows indicate magnitude and direction of geometric displacement around a neutral-layer.

The interpretation of individual DInSAR observations warrants more careful consideration of the vertical structure of tidal flexure. These measurements are inherently limited to the ice surface, whereas flexural deformation occurs about a neutral layer located approximately midway through the ice thickness (as illustrated in Fig. R1). In that schematic, the upper part of the beam moves to the left (red) while the lower part moves to the right (blue), highlighting that the ice deforms in opposite directions above and below the neutral layer. Vice versa for low tide as also illustrated. As a result, the surface experiences a harmonic back-and-forth motion during
the tidal cycle, which can generate an apparent, spurious grounding-line migration on the order of one local ice thickness (Rack et al., 2017), which is within the y-range of the scatter presented in Fig. 9a.

This effect is further complicated in two dimensions by grounding-line sinuosity (Wild et al., 2018), which modifies the surface flexure pattern relative to a simple one-dimensional beam (it's dampened within embayments, and propelled around protrusions of the grounding line). In principle, this "bending effect" can be quantified and subtracted using a simple elastic model (with the temporal mean grounding line position, an ice thickness map and the DSSH). The tidal-flexure slopes in both the x and y directions are then combined with assumptions about the distance to the neutral layer. The corresponding x and y components of the bending effect can then be rotated into the radar line-of-sight using straightforward trigonometric relationships.

$$\int_x \delta_x \, dx = Be_x = \frac{H}{2} w'_x$$
$$\int_y \delta_y \, dy = Be_y = \frac{H}{2} w'_y.$$

Given the importance of this effect for grounding-line interpretation, the manuscript would benefit from a more explicit treatment of how surface flexure, the position of the neutral layer, and two-dimensional grounding-line geometry influence the inferred grounding-line signal.

In particular, I am concerned that the delineation of the tidal-flexure zone may be partly biased. Support for this interpretation is provided by the observation that DInSAR-derived displacements acquired at higher incidence angles show improved correlation during the unstable regime (Fig. 8a/b and L319–321). As the radar incidence angle increases, the sensor becomes more sensitive to horizontal surface motion and its temporal variability, which amplifies the contribution of geometric surface deformation associated with flexure, compared to a lower incidence angle.

This also has implications for the authors' conclusion that "a larger grounding-zone width promotes increased ice flow" (L443-445). Thicker ice is expected to exhibit a more pronounced bending effect than thinner ice, and the associated geometric surface motion should be greatest at the grounding line itself, where curvature is most pronounced. This could, at least in part, create or exaggerate the appearance of a wider grounding zone independent of dynamic ice flow.

For this reason, the multi-incidence-angle DInSAR dataset presented in the manuscript offers a valuable opportunity. The differing sensitivity of the observations to horizontal versus vertical motion provides a promising, and untapped, means of constraining the depth of the neutral layer beneath the ice surface. Such an analysis would also represent a significant contribution to ongoing glaciological efforts to infer basal melt rates from phase-sensitive radar measurements in the tidal-flexure zone (Vaňková et al., 2020).

Thank you for suggesting an analysis of the effect of bending on DInSAR-derived surface displacement. We have implemented the elastic model described by Rack et al. (2017), using known thickness values along the three transects. An example of horizontal surface displacement along the central transect, associated with the highest and lowest tides during the observation period, is shown in Fig. 1AR.




Figure 1AR: Horizontal surface displacements assuming an ice thickness of 210 m (i.e., the mean thickness of the grounding zone along the central transect) and an elastic Young's modulus of 1.5 GPa for (a) the highest tide of 0.50 m and (b) the lowest tide of −0.44 m during the observation period.

Using this simple elastic model, we assessed the contribution from differences in ice extension or compression caused by tides at the times of SAR data acquisition. We then quantified these

contributions for all orbits ("MeanLOSbend error" in Fig. 2AR). As expected, the most pronounced contribution from horizontal displacement due to velocity variability and bending was associated with orbit 09 (high incidence angle, descending pass).




Figure 2AR (updated Fig. 8 from the initial manuscript): DInSAR-derived vertical displacement versus differential sea surface height (DSSH) along the western and central transects with respect to the glacier velocity periods as defined in Fig. 2. Steady period with minimal changes in velocities and low line-of-sight errors for four orbits: a), c), e), g); and unsteady period with more pronounced changes in velocities and high line-of-sight errors for four orbits: b), d), f), h). LOShV represents error due to variable velocity based on in situ GNSS records, and LOShB represents error due to ice surface extension or compression from tidal bending along the central (western) transect.

We added information on the calculation of uncertainties associated with line-of-sight errors in Methods "2.2.3 Assessment of uncertainties in flexure-zone delineations due to velocity variability, ice bending, and SAR viewing geometry":

"The uncertainty of the delineated hinge-line positions was assessed based on the sum of the computed line-of-sight errors arising from variations in ice velocity and ice bending. This combined error was expressed as a fraction of a SAR fringe, representing a vertical displacement of half the wavelength (i.e., 2.75 cm for RCM). To estimate the horizontal extent corresponding to a fraction of a SAR fringe, the typical fringe width along the central transect was calculated from the DInSAR images used for flexure-zone delineation. A Laplacian filter was applied to the DInSAR images to enhance fringe edges, after which fringe widths were computed for all images and the mean, median, and mode values were determined. Horizontal uncertainties were then calculated as fractions of the typical fringe width along the transect, represented by the median value."

The uncertainty results were added in Results "3.3 Uncertainty analysis results" as follows:

"The three ascending orbits with low incidence angles yielded the most reliable results, characterized by high correlations between DSSH and DInSAR-derived displacements and the lowest mean LOShV and LOShB errors during the steady period. In contrast, descending orbit 09, acquired at a high incidence angle, exhibited the highest mean LOShV and LOShB errors during the unsteady period. Along the central transect during the steady period, the combined mean LOShV and LOShB errors for orbits 43 and 103 were 0.12 cm and 0.13 cm, respectively. These combined line-of-sight errors represented 4.4% and 4.7% of the vertical displacement detected within a single RCM DInSAR fringe for orbits 43 and 103, respectively. Given the calculated median horizontal fringe width of 168.7 m (mean: 236.5 m; modal bin range: 100–120 m, representing 15% of total observations), the corresponding horizontal uncertainty during the steady period was approximately ±7–8 m, which is less than one pixel.
During the unsteady period for orbit 09, the maximum combined line-of-sight error along the central transect was 3.26 cm. Given the median horizontal fringe width of 187.1 m (mean: 212.5 m; modal bin range: 160–180 m, representing 16% of total observations), this corresponds to a horizontal uncertainty of ±221.8 m (i.e., approximately 22 pixels) for the flexure-zone delineations. Thus, flexure-zone delineations from orbits 43 and 103 during the steady period represent the most reliable results, with the lowest uncertainty, whereas delineations from orbit 09 during the unsteady period exhibit the greatest uncertainty."

2) Unraveling (here called reconstruction) of DInSAR to single tides

The authors correctly note that DInSAR images represent a double difference of three or four consecutive SAR acquisitions, each taken at different phases of the tidal cycle and therefore containing contributions from multiple tidal constituents as well as atmospheric effects. Consequently, a direct comparison between tide models and DInSAR observations requires either that the same SAR differencing procedure be applied to the modeled tides (as in Fig. 4), or that the DInSAR signal be decomposed into individual tidal components (as attempted in Fig. 6).

The authors pursue the latter approach by fitting a single harmonic with a prescribed angular frequency corresponding to one dominant tidal constituent, both with and without the inclusion of the inverse barometer effect (IBE). While the fit does improve when the IBE is included (Fig. 6), this improvement is expected and has been demonstrated in numerous previous work (e.g., Rignot et al., 2000; Padman et al., 2003; Wild et al., 2022). Importantly, the remaining root-mean-square error (~13 cm; Fig. 6b/c) remains large relative to the total displacement range of approximately -40 to +60 cm in the corresponding double-difference interferograms (Fig. 5a). In my view, this relative magnitude of residual error limits the physical interpretability of the harmonic decomposition using a single sinusoid. I would also like to draw the author's attention to (Minchew et al., 2017), who underscore the importance of using all sinusoids for the unraveling/reconstruction (with former knowledge of their phase from GNSS).

In previous work, we developed and applied an alternative approach for unraveling DInSAR to single tides for Darwin Glacier (Wild et al., 2019) and subsequently for Priestley Glacier (Wild et al., 2025), both published in *The Cryosphere*. This method uses the mean tide-deflection ratio (the so-called alpha map first introduced by Han and Lee, 2015) to separate DInSAR observations into individual tidal components, reducing the residual RMSE to a level comparable to interferometric noise (on the order of 1 cm for Sentinel-1). Based on the flexure curves shown in Fig. 5, the Milne Glacier dataset appears well suited for this type of analysis, and its application here could substantially strengthen the study and its glaciological scope. In this context, relying solely on a single-harmonic fit, despite recognizing the importance of IBE forcing, seems a missed opportunity to fully exploit the dataset. I would be very willing to discuss or advise on such an analysis if helpful, while recognizing that this would extend beyond the minimum expectations for a reviewer.

Two additional benefits of the alpha-map approach are worth noting. First, it naturally averages over the bending effect associated with the position of the neutral layer, because DInSAR observations sample all phases of the tidal cycle (as illustrated in Fig. 5). Second, it mitigates the influence of viscoelastic effects, which appear to be present in several flexure curves along the West and Central transects, through temporal averaging (discussed further below), so it avoids the implementation of a numerical model.

Our manuscript focuses on the "standard" DInSAR method and its assessment with respect to horizontal displacement components caused by variable velocity and bending effects (added in the revised version of the manuscript). Therefore, we consider the alpha-map approach to be outside the scope of this manuscript.
However, based on the feedback above, we improved our initial least-squares fitting approach by including all tidal constituents (i.e., 59) computed using the T_TIDE algorithm (Pawlowicz et al., 2002). The coefficients for each harmonic component were estimated using a regularized least-squares approach. To minimize the effect of horizontal displacement in our DInSAR results, the reconstruction was performed only for the stable season (Fig. 3AR). The associated figure (Fig. 6) and text in Methods, Results, and Appendix sections were revised accordingly.
As a result, the remaining RMSE decreased from ~13 cm to ~5 cm (Fig. 3AR, lower panels), which is a substantial improvement. Moreover, given that T_TIDE shows a correlation of 0.96 and an RMSE of 3.4 cm with our in situ tidal observations (details provided in the corresponding comment below), the remaining RMSE of ~5 cm in our reconstruction analysis is reasonable.




Figure 3AR (updated Fig. 6 from the initial manuscript): Reconstructed ice surface height. a) Time-series of ice surface heights and sea surface heights (SSH, i.e., tides and the inverse barometer effect, IBE). Relationship between the differential reconstructed ice surface displacement and DInSAR-based displacement calculated along the: b) western and c) central transects. Number of observations is 55.

3) Tidal forcing and flexure-zone width

I would appreciate further clarification regarding the physical basis for the reported relationship between flexure-zone width and the magnitude of the tidal forcing. From my understanding, for a given tidal forcing, the width of the flexure zone should primarily depend on ice stiffness, which is itself governed by ice thickness and the effective Young's modulus (Fig. R2). Under this framework, the magnitude of the tidal forcing should affect displacement amplitudes, but not the spatial extent of the flexural response. Similar for ice viscosity, which affects the timing and shape of the flexure, but not its horizontal extent.




Fig. R2: from Wild et al., 2017. Both panels show that for a given tidal forcing (here -0.6 m) and ice thickness, the flexure-zone width is only dependent on the Young's Modulus. Ice viscosity affects the shape and timing of the flexural wave.

For this reason, I am concerned about the interpretation presented in Fig. 9b, as well as the conclusions drawn from it, may not be physically well founded. In addition, it is unclear whether the linear relationships shown are statistically significant. While some structure is visible in the scatterplot, this pattern may alternatively reflect temporal aliasing in the SAR-data acquisitions, potentially related to tidal phase or inverse barometer effects at the time of SAR-data collection.

To explore this possibility, it would be helpful if the authors could reproduce the classification used in Fig. 2a (i.e., the different triangle symbols) in Fig. 6a and explicitly assess whether systematic biases arise from the timing of the SAR-data acquisitions relative to tidal forcing and IBE. Such an analysis would help to clarify whether the observed relationship reflects a physical signal or an acquisition-related artifact.

We agree that the width of the flexure zone depends on ice stiffness, which is governed by ice thickness and the effective Young's modulus, in turn defined by tensile and compressive stresses. We also think that glacier geometry (i.e., slope steepness and/or direction, i.e., retrograde or prograde, and the presence of pinning points) affects the distribution of tensile and compressive stresses in the flexure zone, as well as its resulting width under varying tides, which is likely nonlinear. Our aim was not to fit a linear function but to demonstrate the presence of an appreciable trend with a moderate correlation (~ 0.4).

We added the requested figure (Fig. 4AR) showing SAR data acquisition times to assess potential temporal aliasing. Additionally, we show the association between DInSAR surface displacement and flexure zone width (Fig. 5AR) for different orbits. We find no obvious evidence of pronounced

temporal aliasing due to acquisition timing, except that orbit 09 acquisitions were associated with slightly higher tides and orbit 43 with slightly lower tides than the other orbits during the unstable season (red dashed lines). We removed Figure 9 to avoid potential misinterpretation regarding linear fitting, and keep only Table 2 summarizing the flexure zone span across the transects, and revised our discussion to take the suggested comments into account.




Figure 4AR: SAR data acquisition times with respect to sea surface heights during the 2023-2024 stable season.




Figure 5AR: Association between DInSAR surface displacement and flexure zone width. Orbit 09 is shown with diamonds, orbit 43 with circles, orbit 58 with squares, and orbit 103 with triangles.

The following text was added to the Discussion:

"These distinct slices of the flexure zones along the three transects differ due to variations in bed geometry, ice thickness, and likely ice stiffness. While the width of the flexure zone is governed by ice stiffness, which is determined by ice thickness and the effective Young's modulus (Wild

et al., 2017), pinning points can modify these relationships. Pinning points act to physically prop up the floating ice tongue, at least at certain water levels, thereby influencing local ice stiffness, ice flexure patterns, and the extent of inland water intrusion with changing tides. Previous studies have shown that the presence of pinning points affects lateral and basal shear stresses, generating upstream compression and downstream tension, which, in turn, influences flow variability and can locally enhance the stiffness of the glacier system (Robel et al., 2017; Still & Hulbe, 2021). In this study, both of the observed pinning points are located near the lateral shear zones, which are associated with reduced ice stiffness (Wild et al., 2025). Therefore, the complex interplay between local ice stiffness and glacier bed geometry results in the observed flexure-zone widths, which tend to be larger near the glacier margins where the pinning points are present."

4) Tidal forcing and grounding-zone width

Furthermore, the manuscript presents a similar analysis relating tidal forcing to grounding-zone width (referred to as the "Distance to HL" in Fig. 9a). This type of correlation implicitly assumes that tidal flexure responds instantaneously to the forcing, i.e., that the ice behaves purely elastically. However, at tidal frequencies, glacier ice is well known to exhibit viscoelastic behavior (e.g., Reeh et al., 2003; Wild et al., 2017). Under such conditions, a time lag between the tidal forcing and the flexural response is expected. Importantly, this delay is not constant but is anticipated to increase nonlinearly from the freely floating portion of the ice shelf toward the grounding (hinge) line.

This viscoelastic response provides a plausible alternative explanation for the observed negative correlations. Time delays on the order of tens of minutes to several hours have been reported in similar settings and should be carefully accounted for when interpreting correlations between tidal forcing and grounding-zone metrics. In this context, applying a simple linear fit across all observations may not be physically appropriate.

A useful first step toward quantifying this time delay would be to correlate the available GNSS observations acquired within the tidal-flexure zone with an independent tide model representative of the freely floating ice shelf. For example, the Gr1kmTM model by Howard and Padman (2021) could be used for this purpose. Introducing an artificial time lag and identifying the lag that maximizes correlation between the modeled tides and the GNSS data, following an approach similar to that described by Wild et al., 2025, would help to assess the magnitude and spatial variability of the viscoelastic response. Such an analysis could substantially strengthen the physical interpretation of the results presented in Fig. 9a.

Similarly to the correlation discussed above (Fig. 5AR), we do not aim to fit a linear function but only to demonstrate the presence of a trend and a moderate correlation. In addition, our analysis of SSH with respect to DInSAR results, divided into two groups (i.e., "notch present" and "no notch") confirms that the grounding line migrates farther inland (i.e., when a "notch" is present and, thus, the distance to the hinge line is shorter) under high SSH.

We also note that our simulated tides are representative of a freely floating ice shelf based on pressure records from an RBR Duet datalogger moored in a lateral bay. Furthermore, the T_TIDE

simulated tides showed better agreement with in situ observations than the Gr1kmTM model (details are provided in our response to the corresponding question below).

To quantify the time delay between the simulated tides and the ice flexure response, we calculated the correlation between the simulated tides and GNSS records using different time shifts (Fig. 6AR). The maximum correlation achieved is about 0.2 with a time shift of 15 minutes (Fig. 6AR c). This low correlation is influenced by the limited GNSS observations and their own oscillations (Fig. 6AR b). Since our GNSS records were collected for only three hours, they unfortunately did not capture a full tidal period, preventing reliable quantification of the correlation and the time shift. Therefore, we prefer to remove Figure 9 to avoid potential misinterpretation.




Figure 6AR: Correlation analysis between SSH and GNSS records in the flexure zone: a) Time series of SSH and GNSS-derived surface heights; b) Zoom-in on the October records to demonstrate oscillations in the GNSS data; c) Correlation between SSH and GNSS (running mean) with respect to time shift.

We added Table 4 "Results for the distances from point A to the hinge line, grounding-zone widths, and the associated SSH standard deviations" and Figure 8 "Timing of RCM SAR data acquisitions and the associated SSH values when the maximum (upward-pointing triangles) and minimum (downward-pointing triangles) distances from point A to the hinge line were detected in the DInSAR images…", along with the associated text in "3.4 Grounding-line migration and grounding-zone width results".

Minor comments:

L14: Reword from "Changes at the boundary" to "Changes of the boundary" as the grounding line location generally serves as a proxy for stability considerations in numerical models.

We revised the sentence as follows:
"Observing and quantifying changes of the boundary where marine-terminating glaciers transition from grounded to floating, known as the grounding line, is critical for advancing our understanding of glacier stability."

L15: Terminology. Reword here and throughout from "ice-flexure zone" to "tidal-flexure zone"

Thank you. We have revised the terminology and replaced "ice-flexure zone" with "tidal-flexure zone," since "tidal flexure" or "flexure" are more commonly used in combination with "zone" in the literature. While terms such as "ice flexure" or "tidal ice flexure" are also used, they generally appear without "zone" (e.g., Brunt et al., 2010; Freer et al., 2023; Wallis et al., 2024).

We also clarified our use of this term as follows:

"The DInSAR method detects the differential tidal-flexure zone, hereafter referred to as the flexure zone, where the glacier surface flexure with tides is captured at SAR data acquisition times."

L19: Terminology. Reword here and throughout from "Double Difference Interferometric SAR (DDInSAR)" to "Differential Interferometric SAR (DInSAR)". The word interferometric implies a (phase) difference between SAR images, the differential implies that two interferograms are being differenced. I know that DDInSAR can be found in the literature, but DInSAR is the modern terminology and the community strives to get rid of DDInSAR.

Thank you for the detailed clarifications. Yes, both versions of this term can be found in the literature. We have revised the terminology to adopt "DInSAR".

L23/24: Terminology. Reword here and throughout from "stable/unstable regime" to "steady/unsteady period". Stability of a glacier system is carefully defined in the ice-dynamics modeling community and we shouldn't use the term for other processes than MISI or MICI. Here, even if Milne Glacier experiences MISI, the analysis is concerned with the assumption of steady flow in DInSAR processing. In line with this, also reword "changes in glacier velocity were minimal" to "glacier flow was steady" and "more variable" to "unsteady". I also wonder if a single observation of an unsteady period warrants the classification as a "regime" or "phase". I suggest rewording these terms to "period".

We have revised the terminology to adopt "steady/unsteady period."
This text was revised as follows:

"The most reliable flexure zone delineation (horizontal uncertainty of ±8 m) was achieved with ascending RCM tracks at low incidence angles during periods when the ice velocity was steady and bending was not substantial, resulting in a combined SAR line-of-sight error of 0.13 cm. A

worst-case flexure-zone delineation uncertainty of ±222 m was noted when a descending track with a high incidence angle was used and glacier velocity became more variable, increasing the combined SAR line-of-sight error to 3.26 cm."

L32: Please replace the citation of the IPCC with original research in a discipline-specific journal like *The Cryosphere*.

The sentence and the citation were revised as follows:
"Rising atmospheric and oceanic temperatures can cause these glaciers to accelerate, leading to increased sea level rise and discharge of icebergs or ice islands into adjacent waters (Morlighem et al., 2019; Wood et al., 2021; Carnahan et al., 2022; Holmes et al., 2025)."

L38: Change "migrating with tides" to "migrating with tides over short time scales (Freer et al., 2023)"

The sentence was revised as follows:
"The grounding line exhibits dynamic behaviour, migrating with tides over short time scales (Freer et al., 2023) and advancing or retreating in response to changes in glacier dynamics over the long term (Milillo et al., 2022; Chartrand et al., 2024)."

L41: Change "DDInSAR" here and throughout to "DInSAR". This means that "DSSH" can later be more intuitively compared to "DInSAR".

The term and its abbreviation have been updated throughout the manuscript.

L43: "4 days" please compare to other often used SAR satellites like Sentinel-1, TerraSAR-X, COSMO, etc. Is this a dedicated mission, if so, please add this piece of information.

We revised the sentence and added additional information to compare RCM with other operational missions explored in the literature:

"The satellite-based Differential Interferometric Synthetic Aperture Radar (DInSAR) method has been proven to be highly effective for accurate monitoring of grounding-line migration, using SAR imagery with high temporal (1-6 days) and spatial (1 m to a few metres) resolution, along with consistent multi-temporal coverage over large areas ranging from tens to several hundreds of kilometres (Rignot et al., 2011; Milillo et al., 2017). Among the currently operational SAR missions explored for DInSAR grounding-line applications, the RADARSAT Constellation Mission (RCM, C-band) has a 4-day acquisition cycle (Antropova et al., 2024; Gadi et al., 2025). This is comparable to COSMO-SkyMed (X-band) allowing 1 to 4 day revisits (Milillo et al., 2022; Rignot et al., 2024), and shorter than Sentinel-1 (C-band) with 6 to 12 day revisits (shorter with the A+B or A+C constellations) (Ramanath et al., 2025; Wild et al., 2025), TerraSAR-X (X-band) providing 11-day revisit (shorter with TanDEM-X) (Wild et al., 2018; Wild et al., 2025), and ALOS PALSAR (L-band) with a 46-day revisit (Rignot et al., 2011). The relatively short revisit time of RCM enables high phase coherence, while its C-band wavelength provides sensitivity to small vertical displacements, allowing reliable DInSAR measurements for grounding-line monitoring."

L43: Kim et al. 2024 use a terrestrial radar system, while the present study is about a space-borne system. Please clarify the difference or replace this reference with other DInSAR based studies of which there are many other relevant ones.

Thank you. Yes, this terrestrial radar-based DInSAR study was cited in the context of sub-daily acquisitions. We decided to focus on satellite-based SAR missions only and revised the text accordingly (as stated in our response above); the citation "Kim et al., 2024" was removed.

L46: Define "Hydrostatic line" here, as it is used later in the paper but I can't remember to have seen its definition.

The sentence was revised as follows:

"The DInSAR method detects the differential tidal-flexure zone, hereafter referred to as the flexure zone, where the glacier surface flexure with tides is captured at SAR data acquisition times. This zone, also referred to as the grounding zone in some studies (e.g., Fricker et al., 2009; Freer et al., 2023), spans from the inland limit of tidal flexure (i.e., the hinge line) past the grounding line to the seaward limit of flexure (i.e., hydrostatic equilibrium line), where hydrostatic equilibrium is reached and the ice tongue becomes completely buoyant."

Also, a clarifying sentence was added to the Abstract:

"This zone spans from the inland limit of tidal flexure — the hinge line, across the grounding line to the hydrostatic equilibrium line, where the ice tongue becomes freely floating."

L50: Terminology. Reword here and throughout "FZ span" to "tidal-flexure zone", and "GZ span" to "grounding zone" and refer to their width.

We replaced the abbreviations FZ and GZ with their full terms, although abbreviations seem to be acceptable in The Cryosphere (e.g., use of 'GL' for grounding line by Le Meur et al., 2014). We note that the "tidal-flexure zone" is equivalent to the "flexure zone" in the context of this manuscript, as defined in the text and consistent with recent studies.
We acknowledge that "grounding zone width" is more commonly used in the literature; however, some argue that this terminology can be misleading, since the distance is measured along the glacier flow rather than across it. We believe that terms such as along glacier "extent" or "span" could be more precise. Nevertheless, we adopt the term "width" for both the tidal-flexure zone and the grounding zone, to remain consistent with its frequent usage in the literature. We also note that "FZ span" is equivalent to FZ width measured along the glacier from a DInSAR result, while GZ width is calculated based on all available DInSAR images.

L55: Causes of complexity. The most relevant citations for the present study are Rack et al., 2017 and Wild et al., 2018.

Thank you, we added these citations:

"Recent DInSAR-based studies highlighted the complexity of grounding-line migration with tidal cycles and explored the causes of this complexity, including glacier geometry, underlying bed type, subglacial hydrology, and sea water intrusions (e.g., Mohajerani et al., 2021; Ciracì et al., 2023, Rignot et al., 2024; Kim et al., 2024), as well as the contributions of compressional and tensional strain components induced by ice-surface bending to the DInSAR signal (Rack et al., 2017; Wild et al., 2018)."

L65: What "model datasets" ? It sets up the expectation of the reader to see some form of tidal-flexure modeling. Coming back to this after reading the paper, I think this is only GEM, right ? I think to warrant the "model datasets" here requires to perform the analysis at least with and without the GEM-derived IBE, in comparison with the GNSS analysis.

We revised the following sentences in the Introduction section to better clarify our datasets:

"Our study explores intra-annual changes in the Milne Glacier flexure zone based on a combination of RCM DInSAR-derived results, in situ and simulated datasets, and a numerical weather prediction model output. The objectives of this study are to:
- Delineate the Milne Glacier flexure zone using RCM data acquisitions from four different orbits, with a 4-day repeat interval within each orbit and about a 1-day interval between orbits, spanning from February 17, 2023 to September 17, 2024;
- Examine the relationship between DInSAR-derived ice-surface vertical displacement and displacement of sea surface height (DSSH), computed from simulated tides and in situ atmospheric pressure records together with modeled atmospheric pressure."

We added the GEM model-based IBE correction to the Appendix (Fig. A3) for comparison with in situ-based IBE correction, as requested by another reviewer. GEM-derived IBE correction is also shown Fig. A1."

L67: Perfect location to add a statement about the quantification of the steady-flow assumption during times when it is clearly violated. I believe this should be the real focus of the manuscript.

Thank you, we refined this objective in the Introduction section as follows:

"We further assess this correlation across the four SAR orbits with respect to the compressional and tensional strains caused by ice-surface bending in the flexure zone during two velocity periods:
a) a steady period, when the in situ-measured Milne Glacier velocity did not change appreciably during winter and the DInSAR assumption of invariant velocities was met; and
b) an unsteady period, when velocity changes were evident, the glacier decelerated after summer peak velocity, and the DInSAR assumption of invariant velocities was violated;"

L78: Add information about the tidal range and tidal regime. Is it a diurnal or semi-diurnal tide ? How long is the spring-neap-spring tidal cycle ? The introduction of the epishelf lake is of course interesting, but the space should be used for more relevant information to the paper, such as what is known about Milne Glacier's velocity variability, and how it motivates a hypothesis (I suggest to use the known velocity variability as a motivation to analyse how different viewing geometries are affected by horizontal flow variability).

We revised the "Study Site" section, removing some details about the epishelf lake formation and adding information related to tidal processes:

"The present-day intact floating ice tongue, located deep within Milne Fiord, is protected from drifting ice along the northern coast of Ellesmere Island (Mortimer et al., 2012). The floating ice tongue experiences moderate stresses due to tidal bending in the flexure zone. Milne Fiord is characterized by a semi-diurnal tidal regime with a relatively small tidal range, spanning from –20 cm to 21 cm over the observation period of this study, and a 14.8-day spring-neap-spring tidal cycle. However, atmospheric pressure also plays an important role in modulating tides in the fjord (Wunsch & Stammer, 1997; Padman et al., 2003), further affecting the displacement of the ice tongue and increasing the stresses it experiences.
The glacier's transition zone between grounded ice and the floating ice tongue exhibited notable velocity fluctuations over the past decades. Ice velocity decreased from ~190 m yr−1 to ~100 m yr−1 between the 1990s and 2005 (Millan et al., 2017), increased to 120 - 140 m yr−1 in 2010, slowed down again to 80 - 100 m yr−1 from 2011 to 2015, and then accelerated to 140 - 160 m yr−1 between 2016 and 2020 (Van Wychen et al., 2020). Van Wychen et al. (2020) suggested that this cyclical behaviour may result from prominent bedrock features that modulate velocity where the glacier rests below sea level. In contrast, Antropova et al. (2024) noted that the post-2016 acceleration was preceded by the highest grounding-line retreat rate along the glacier's centerline between 2011 and 2018, suggesting the presence of feedback mechanisms between grounding-line retreat and upstream glacier acceleration."

L106: How many km upstream is the mentioned prograde bed slope ? How many years of average grounding-line retreat will it take to be reached ?

We added the suggested details as follows:

"This positive feedback loop may be underway currently and is expected to persist until the grounding line reaches a prograde bed slope roughly 1 km up-glacier. Based on the glacier's 2011-2023 average retreat rate, this would occur in 2035. However, grounding-line retreat is inherently non-linear and governed by the complex interplay of glacier-specific and environmental controls. Reaching this slope will help stabilize the glacier and inhibit further recession of its grounding line."

L108: Data and methods. Please add sub-section titles such as "Differential InSAR processing", "In-situ data", "Tide modeling", "Uncertainty analysis", "Grounding-zone width modeling", etc…

We distinguished the following subsections and revised their order based on our analysis:

2.2 Data and methods
2.2.1 Processing of SAR data
2.2.2 Analysis of tides and reconstruction of ice-surface height timeseries
2.2.3 Assessment of uncertainties in flexure-zone delineations due to velocity variability, ice bending, and SAR viewing geometry
2.2.4 Analysis of grounding-line migration and grounding-zone width

The Results section was also revised accordingly.

L111: A 10m spatial resolution is stated in the abstract, here 5m native resolution and the table shows about 2.5m raw pixel size. I know how these are affected by multi-looking and the topographic correction, so please describe these processing steps in more detail and state the chosen variables, which is also a great motivation to acquire a higher-resolution DEM to not lose any information in RCM.

We provided additional details on the RCM SAR products and our choice of spatial resolution for the final DInSAR results:

"We used RCM single look complex (SLC) images with HH polarization, acquired in 'High Resolution 5m Mode' (MDA, 2021) from different orbits during 2023–2024 (Table 1). This mode has a nominal resolution (i.e., theoretical resolution, which is based on radar system parameters) of 5 m and a swath width of 30 km. The native SLC pixel spacing is 1.4–2.8 m in the range direction and 2.4–2.8 m in the azimuth direction depending on the incidence angle (Table 1)."

"Interferograms were computed for each pair of RCM data acquisitions with a 4-day interval. During processing, we applied 3 × 3 multilooking to reduce speckle, improve the signal-to-noise ratio and interferometric coherence, and reduce file sizes to facilitate phase unwrapping. After multilooking, the effective ground pixel spacing was approximately 9–10 m. Topographic phase removal and terrain correction were performed using the 10 m ArcticDEM dataset (Porter et al., 2023). Interferograms were unwrapped using the minimum-cost flow phase unwrapping algorithm described by Costantini (1998). During the initial unwrapping, the processing software automatically selected an arbitrary stable reference area. Then, differential interferograms were calculated at 10 m resolution (i.e., the final spatial resolution of DInSAR images used for flexure zone delineations) and the corresponding unwrapped results represented by three consecutive data acquisitions (over 8 days) or four data acquisitions (over ≥12 days) if there was a gap between data acquisitions."

RCM modes descriptions by MDA, (2021) can be found at:
https://download-telecharger.services.geo.ca/pub/csa_asc/Space-technology_Technologie-spatiale/radarsat_constellation_mission_plan/RCM-SP-52-9092_Product_Spec_1-15_Public.pdf

L115: What seems to me the most important information is that orbit 9 has twice as large incidence angle as the other orbits. This should be better visible in the table and the description of the RCM data, which sets the paper up nicely for developing the unsteady-flow quantification or even the bending-effect hypotheses outlined in the major comments.

We highlighted this information in the table by adding the following footnote:

"[2] The incidence angle of orbit 09 is 1.4 - 2 times greater than that of orbits 43, 58, and 103."

L121: reword to "effect of steady glacier flow".

We added "steady" to the sentence:

"This method reveals the surface displacement of the floating ice tongue along the SAR line-of-sight, by removing the effect of steady glacier flow, by differencing two consecutive interferograms, under the assumption that the glacier velocity remains constant between SAR data acquisitions."

L122: hydrostatic equilibrium line is not defined yet. Terminology. Also, please avoid confusion with the abbreviation of hinge line. Commonly HL denotes the hydrostatic line, and hinge line is spelled out.

The hydrostatic equilibrium line is now defined in the Introduction (as per the comment above). The "hinge line" term is spelled out in the manuscript text. Fig. 9 from the initial version of the manuscript with the "HL" notation was removed.

L123: Terminology. Reword to "landward and seaward limits of the tidal-flexure pattern". The boundaries normally refer to already delineated GL and HL. Also, reword "glacier flexure" to "tidal flexure". I understand that the main contributor to "glacier flexure" here is the IBE, but the geophysical process is commonly called tidal flexure, and the IBE is one contributor (aside ocean tides and tidal loading on the Earth's crust, which is neglected here, right ?).

We revised the sentence as follows:
"The hinge line and the hydrostatic equilibrium line are then manually or automatically delineated by examining the landward and seaward limits of the fringe pattern in the DInSAR images representing the flexure zone."

We replaced "glacier flexure" with "flexure zone." We consider the term "flexure zone," equivalent to "tidal-flexure zone," to be appropriate, as it is defined in the Introduction and is widely accepted in the literature. Indeed, we believe that the use of "flexure zone" is more intuitive, as it allows encompassing multiple driving factors (e.g., atmospheric pressure, ice thickness and stiffness, glacier geometry, and the presence of pinning points that can redistribute stresses and shape the flexure zone), rather than referring solely to tidal forcing.

We also added the clarification about ocean tide load in "2.2.2 Analysis of tides and reconstruction of ice-surface height timeseries" section:

"Simulated tide heights were corrected by adding the inverse barometer effect (IBE) (Wunsch & Stammer, 1997; Padman et al., 2003), while ocean tide loading on the Earth's crust was neglected, as it has a relatively minor impact (Wild et al., 2025)."

L126: To better distinguish between DInSAR images (the double-difference result) and times of single tide snapshots, please reword here and throughout to "three or four consecutive SAR data acquisitions" or "SAR scenes" to underline the snapshot character of the SAR data.

We replaced the corresponding entries with "SAR data acquisitions."

L129/130: First and only mention of phase-loss in lateral margins. Please add how ice-flow rotation leads to loss of coherence required for interferometry. Also, please read Wild et al., 2025 how weakening of ice properties in these zones affect stiffness and thus the flexural response of the entire glacier. This signal provides an exciting avenue for further research on bulk ice stiffness and buttressing. This section also lacks information of phase unwrapping and the coordinate used as a reference, which is also not shown in the figures.

The suggested information was added to the text as follows:

"The western and eastern flexure-zone boundaries were often affected by noise. These areas of low interferometric coherence typically occur due to strong velocity gradients, shear, ice deformation, and rotation as glacier velocities decrease from the centre toward the lateral margins as a result of lateral drag (Forster et al., 1999; Mannerfelt et al., 2025). These processes contribute to ice weakening and a reduction in stiffness in the lateral shear zone (e.g., a fivefold reduction in Young's modulus was reported by Wild et al., 2025). This interpretation is supported by our in situ observations and high-resolution imagery demonstrating extensive crevassing in this zone, particularly along the western margin of Milne Glacier, where a tributary glacier further enhances these processes. The resulting loss of coherence in the flexure-zone's western and eastern boundaries prevented their reliable delineation and the use of the entire area of the flexure zone for our analysis."

We added the following information about phase unwrapping and corrections:

"Interferograms were unwrapped using the minimum-cost flow phase unwrapping algorithm described by Costantini (1998). During the initial unwrappig, the processing software automatically selected an arbitrary stable reference area."

L142: How were phase discontinuities corrected ? Only along the transects ? This and the mention of division by cosine of the incidence angle to get displacement should be moved to a dedicated DInSAR processing subsection in the methods.

Yes, the phase discontinuities were corrected along the transects with known bed geometry. We added clarifying details to the text and moved it to "2.2.1 Processing of SAR data" section:

"The final calibration of the SAR line-of-sight displacement was conducted along three transects (Fig. 1, described below) using an automated script. Values were divided by the cosine of the incidence angle and additionally calibrated to the mean values of the grounded portions of the transects for each DInSAR image from orbits 09 and 103. Phase jumps exceeding 2 cm (i.e., a more conservative threshold than half of the RCM C-band wavelength) were corrected for all orbits."

L154: The correct references for the importance of atmospheric pressure are Rignot et al., 2000 and Padman et al., 2003.

Thank you. We have replaced the citation with these two "(Wunsch & Stammer, 1997; Padman et al., 2003)" earlier studies.

L156: t-tide analysis. Please quantify how accurately the t-tide prediction replicates the input data, and how well it fits to an independent tidal model (such as Gr1kmTM). How do these uncertainties affect the results ? Is there a time delay between modeled tides and GNSS displacement ?

Our T_TIDE simulations were conducted based on 'a 10-month in situ pressure record from an RBR Duet datalogger moored at ~10 m depth in a lateral bay off the west side of Milne Fiord (Antropova et al., 2024)', rather than on GNSS ice surface displacement data measured on a freely floating ice tongue. The manuscript text was revised to explain this directly, rather than referring to Antropova et al. (2024). In this study, GNSS records were primarily used to investigate displacement due to ice flow in the grounding zone, where the ice was not freely floating; therefore, these records were not suitable for tidal simulations. We computed the requested statistics using our pressure-record observations and consider that the question of a time delay is not relevant to our dataset.
The comparison of detrended 2016-2017 in situ pressure-based tide records with tides computed using T_TIDE and Gr1kmTM (Fig. 7AR) shows correlation of 0.83 vs. 0.73 and RMSE values of 4.7 cm vs. 5.9 cm, for T_TIDE vs. Gr1kmTM, respectively. Therefore, T_TIDE is more accurate than Gr1kmTM as it shows better agreement with the observed tides than Gr1kmTM.




Figure 7AR: In-situ pressure-based tide records and simulated tides using T-tide algorithm and Gr1kmTM model.

We also added to the manuscript a comparison of an independent tide data sample (i.e., not used as an input for the tide simulations) collected in July 2023 against the simulated tides, with GEM-based IBE included, to the Appendix (Fig. A1). Based on this comparison, which represents the observed conditions better than the detrending approach above and falls within our observation period, the correlation was 0.96 vs. 0.93 and the RMSE was 3.4 cm vs. 4.6 cm for the T_TIDE- and Gr1kmTM -based outputs, respectively. Thus, T_TIDE demonstrated better agreement with the observed tides than GR1kmTM.

The following text was added to the manuscript:

"The SSH computed using T_TIDE showed a higher correlation (0.96 vs. 0.93) and lower RMSE (3.4 cm vs. 4.6 cm) than the independent tidal model GR1kmTM (Howard and Padman, 2021), when compared with repeated in situ tidal observations collected in July 2023 (Fig. A1). This supports the choice of the T_TIDE algorithm for our analysis."

The following figure was added to the Appendix (i.e., Fig. A1):




Figure 8AR: Observed tide records measured in situ in July 2023 (black) and computed sea surface heights (SSH) using the T-tide algorithm and the Gr1kmTM model, with IBE added (red and blue, respectively). The correlation between the observed and computed SSH based on the T_TIDE and Gr1kmTM is 0.96 vs. 0.93 and the RMSE is 3.4 cm vs. 4.6 cm, respectively.

L157: Reword "corrected" to "added to" the IBE, since atmospheric pressure variations are a contributor to the tidal forcing (sum of ocean tides, load tides, and the IBE). Also, please add the conversion factor, was it 1cm drop for every positive anomaly of 1hPa ? Were pressure anomalies calculated over the entire time span, or over shorter chunks ?

Yes, we used a conversion factor of a 1 cm drop for every positive 1 hPa pressure anomaly, and vice versa.
The IBE was calculated over the entire study period.

The corresponding sentences were revised as follows:

"Simulated tide heights were corrected by adding the inverse barometer effect (IBE) (Wunsch & Stammer, 1997; Padman et al., 2003), while ocean tide loading on the Earth's crust was neglected, as it has a relatively minor impact (Wild et al., 2025)."

"To calculate the IBE over the entire observation period, we assumed that a 1 hPa change in atmospheric pressure results in a corresponding -1 cm change in sea level (Wunsch & Stammer, 1997; Padman et al., 2003). This correction was linearly interpolated to a 1-minute interval and then added to the simulated tides to compute the SSH."

L160: What's the temporal resolution of GEM ? Fig. 2c indicates it is lower resolution than the AWS data. Was the AWS pressure record low-pass filtered to a common cutoff frequency before any gaps were filled to account for any differences in temporal resolution ?

The temporal resolution of GEM is 1 hour, whereas the temporal resolution of AWS is 30 minutes. We added information about time intervals to the corresponding section. Our tidal time series were simulated at a 1-minute interval in order to better match the acquisition times of the SAR data. The pressure records from AWS and GEM were used to calculate the IBE values, which were then linearly interpolated to a 1-minute interval and added to the tidal time series. The following clarifications were added to the manuscript:

We also note that a gap in the AWS records occurred between July 9 and July 21, 2023, during which no coherent DInSAR results were available for our analysis, and after July 22, 2024, when only one SAR triplet was acquired in September. Therefore, the GEM-based correction corresponds to only one DInSAR result in our analysis.

L162: Please quantify "good agreement" between GEM and AWS data. For a direct comparison, the AWS records will need to be low-pass filtered with the cutoff frequency of GEM before any correlations and RMSE can be calculated.

The following information was added to the manuscript:
"In particular, the correlation between the GEM and in situ atmospheric pressure records used for IBE calculations was 0.96, with the RMSE of 3.6 hPa."

L163: Reword "DDInSAR result" to "DInSAR image" and also change to "SAR data acquisition" or "SAR scenes".

"SAR acquisition(s)" entries were replaced with "SAR data acquisition," as stated above.

L164: Reword "DDInSAR subtraction", which implies a triple difference to "DInSAR combinations", which I think is not meant here.

"DDInSAR" entries were replaced with "DInSAR," as stated above.

L165: Where along each profile is DInSAR-derived surface height taken ? Normally a "freely-floating area" is averaged over, but this reads as if the far left side of each profile was taken ? To minimize the effect of interferogram noise, I suggest averaging over everything larger than 47 km distance, or even better any DInSAR displacement beyond the delineated hydrostatic lines.

Yes, the DInSAR-derived surface heights was calculated as the mean values. We added the following clarifying information to the manuscript:

"We assessed the relationship between DSSH and DInSAR-derived surface height displacement of the freely floating ice tongue (i.e., mean values calculated from 44 km downfjord along the three transects in Fig. 1) based on the correlation coefficient, the root mean squared error (RMSE), and bias."

L171: DInSAR images contain information about the net sum of all tidal constituents (plus IBE and tidal loading) so fitting with a single constituent is physically invalid. Please refer to the methodology presented by Minchew et al., 2016 how to deal with apriori information on each sinusoid before the inversion, or employ an alpha map based approach for the unraveling to single tides.

We improved our initial least-squares fitting approach by including all tidal constituents (i.e., 59) computed using the T_TIDE algorithm and IBE (Fig. 6). The corresponding method description was revised accordingly.

The following text was added to Section "3.2 Tidal analysis and ice-surface height reconstruction results":

"The ice-surface height time series, reconstructed using a regularized least-squares approach based on 59 tidal constituents and DInSAR displacement, is shown in Fig. 6 a. The differential reconstructed ice surface displacement and DInSAR displacement are highly correlated (r= 0.97) and yield an RMSE of approximately 5 cm for both the western and central transects during the steady period, which is reasonable given that our SSH showed a correlation of 0.96 and an RMSE of 3.4 cm with in situ tidal observations (Fig. A1)."

L172: Please reword here and throughout from "SAR acquisitions" to "SAR data acquisitions" or "SAR scenes".

"SAR acquisition(s)" entries were replaced with "SAR data acquisition," as stated above.

L174-176: "We used SSH…" I do not follow this sentence, please rephrase and say what hypothesis is investigated and how it is done.

L174-176 were revised as follows:

"To explore the origin of the newly developed notch along the eastern margin, we tested the hypothesis that it appears in DInSAR images associated with SAR acquisitions during periods of high SSH, suggesting that seawater intrudes farther inland and results in ice ungrounding around a pinning point. We calculated the minimum and maximum SSH values, as well as the standard deviation, for each DInSAR result associated with three or four SAR data acquisitions. These values were then divided into two groups corresponding to: (1) DInSAR results without the notch and (2) DInSAR results with the newly developed notch present, to test whether SSH differs significantly between the groups."

L176: Reword "grounding-line delineations" with "tidal flexure zone delineations".

This sentence was revised as per the comment above, and the term "grounding-line delineations" was removed.

L180: Terminology: Reword “SAR observations” to “SAR data acquisitions” or “SAR scenes”. The word observation normally refers to processed DInSAR images.

“SAR observations” was replaced with “SAR data acquisitions”.

L183: Do the three hour GNSS acquisition period introduce a tidal bias in the derived mean daily velocity ?Is the error of 1.5 cm considered in the uncertainty analysis ?

The GNSS device was installed in the flexure zone; therefore, GNSS measurements (daily mean values for the stabilized GNSS position between 18:00 and 19:00 UTC) were also affected by tidal bending. SAR data for orbits 43, 58, and 103 were acquired approximately 1 hour after the GNSS records. Orbit 09 had a much larger timing lag relative to the GNSS observations and therefore captured a tidal signal different from that recorded by the GNSS.

SAR data acquisition times (UTC):
Orbit 43: 19:58:00;
Orbit 09: 13:19:00;
Orbit 58: 20:06:00;
Orbit 103: 20:30:00;

The following information was added to better explain this:

“The final GNSS position was represented by the mean latitude and longitude values stabilized between 18:00 and 19:00 UTC, which was close to the SAR acquisition times for orbits 43, 58, and 103, but involved a larger time gap for orbit 09. Thus, the GNSS measurements likely captured a more representative tidal signal for orbits 43, 58, and 103 than for orbit 09.”

We did not explicitly account for an error of 1.5 cm. Velocity was calculated through the change in distance relative to the previous GNSS position; assuming a similar positional error for each record, the error largely cancels out.

L190: Quantify “agreed well” with an RMSE. Looking at Fig. 2a this is only true for the steady periods but the agreement isn’t that good during summer and into the unsteady period.

We note that SAR velocities were computed to support our GNSS-based velocities during the period of missing records, which corresponds to the period of good agreement. The following information was added:

“The missing GNSS records (10.5% of the total records within the analysed periods) at the end of the unsteady period and beginning of the steady period in 2024 were linearly interpolated from neighbouring observations and compared against SAR-derived velocity. …
The interpolated values for the missing GNSS records agreed within ±0.01 m d^{-1} with SAR-derived velocities.”

In addition, SAR velocity is affected by speckle noise. While values tend to be more consistent and reliable when analyzed and spatially averaged over larger areas, point-based comparison, as

in our case, is much more complicated due to noise, especially when melt processes affect the SAR signal. The figure demonstrates that the agreement is very good during the stable (cold) season. It is not surprising that summertime results demonstrate weaker agreement, as SAR data acquisitions are much sparser during this period, the SAR signal is affected by melt processes, and, likely, by much more pronounced glacier deformation due to ice flow acceleration.

L192: Reword from "effect of velocity" to "effect of velocity variability"

We added the word "variability".

L201: Please label "Purple Valley" in Fig. 1

The label was added.

L204: Reword "DDInSAR observations" to "SAR data acquisition times for four different orbits".

We made the requested replacement.

L209: Please add the percentage of missing GNSS records of the total period of investigation.

"(10.5% of the total records within the analysed periods)" as per the comment above.

L211-218: Very nice. It would have been beneficial to also conduct this part of the error analysis with the SAR-derived velocity fields and analyze any spatial patterns of the steady flow assumption.

We believe that our approach, based on in situ GNSS measurements, is of interest to the broader community of The Cryosphere. Therefore, we retain the GNSS-based velocities in our analysis and consider a detailed analysis using SAR-derived velocities to be beyond the scope of this manuscript.

L236: Do the delineations of grounding-zone width need to be corrected for the "bending effect" to be comparable to the theoretical width determined by Tsai and Gudmundsson ? In other words, is this a model for neutral layer displacement ?

We infer that Tsai & Gudmundsson (2015) assumed negligible bending stresses for "sufficiently long" ice shelves. Likewise, Chen et al. (2023) and Mohajerani et al. (2021) neglected bending stresses, although they compared their DInSAR-derived results with the theoretical width predicted by Tsai & Gudmundsson (2015).

We added the following statements to better clarify that bending stresses were not considered here:

"This equation was derived by Tsai and Gudmundsson (2015) based on prograde bed geometry and was later applied to both retrograde and prograde bed geometries (e.g., Chen et al., 2023), assuming negligible bending stresses at the ice surface."

L240: reword “RCM images” to “SAR-data acquisitions" or “SAR scenes”.

We replaced “RCM images” with “RCM SAR scenes” here.

L245: This is not surprising, given that ice stiffness controls the width of the tidal-flexure zone.

This sentence was removed, as this part is covered in detail in the Discussion section

L254: Reword “during the last part of our intensive observation period” to “ towards the end of the observation period”.

The “toward the end of the current observation period” replacement was done.

L263: Reword “atmospheric pressure” to “IBE” or “atmospheric pressure variability”, which is what is causing this signal.

The “atmospheric pressure” was replaced with “IBE”.

L267-269: “These correlations…” move to Discussion.

These sentences have been moved to the Discussion section.

L269-272: I think that one must add that the result of a double difference can have a larger absolute value than any of the individual values. So a 2.3 fold increase is essentially a measure of the non-linearity of the investigated process (a tiny change at the times of the SAR-data acquisitions will result in a much larger value in the DInSAR combination). So adding the IBE, which is known to dominate in areas with small tidal range, will greatly improve any correlation with DInSAR measurements on the freely-floating part. Therefore, I think that it is useful to report on how the IBE improves the correlation at Milne Glacier, but it shouldn’t be a main finding with a dedicated Figure in the main text. I suggest moving Fig.4 to the Appendix or SI, and cut back on the interpretation in the main text.

Thank you. We retained only panels for correlation between SSH and DInSAR displacement in the manuscript and moved the other panels to the Appendix. We also revised the text of the manuscript to reduce the discussion of IBE and fully removed it from the Abstract.

L272: Great that correlations improved after the IBE, but the RMSE is still huge when compared to the signal, about 15 cm of 50 cm DSSH. Please refer to the “tide-deflection ratio” approach outlined above on how to further minimize the RMSE to be within interferogram noise (<1cm).

Thank you for your suggestion. However, our study aims to explore the “standard” DInSAR method and its assumption of steady flow velocity. Therefore, we consider reporting these results to be a necessary step in our analysis. As reported above, the correlation between the observed and computed SSH based on T_TIDE was 0.96, and the RMSE was 3.4 cm. We also note that our RMSE is comparable to values reported in recent studies using the same method (e.g., inferred

from Fig. 2c in Chen et al., 2023, and Fig. 3b in Rignot et al., 2024). Therefore, we consider our RMSE values to be reasonable for the method used in this study.
In addition, we emphasize that our GNSS records do not provide reliable information on ice surface displacement for the freely floating ice tongue, and thus we do not have the opportunity to use them for validation as done by Wild et al. (2019). We prefer to retain our "standard" DInSAR approach and consider the "tide-deflection ratio" approach to be outside the scope of the current manuscript. However, we would be happy to share our interferograms and other available processed datasets with interested members of the glaciological community for further exploration of these processes using the "tide-deflection ratio" and other approaches.

L301: I don't quite agree with this interpretation (which also needs to be moved to the Discussion). The width of the flexure zone is independent of DSSH and only determined by ice stiffness. Also, the higher the DSSH, the more pronounced the spurious surface movement by the bending effect and thus the grounding-zone width.

We agree that ice stiffness controls the width of the flexure zone; however, pinning points can influence stress distribution, flow variability, and the apparent stiffness of the glacier system.
We moved the revised clarifications in the Discussion section:

"Previous studies have shown that the presence of pinning points affects lateral and basal shear stresses, generating upstream compression and downstream tension, which, in turn, influences flow variability and can locally enhance the stiffness of the glacier system (Robel et al., 2017; Still & Hulbe, 2021). In this study, both of the observed pinning points are located near the lateral shear zones, which are associated with reduced ice stiffness (Wild et al., 2025). Therefore, the complex interplay between local ice stiffness and glacier bed geometry results in the observed flexure-zone widths, which tend to be larger near the glacier margins where the pinning points are present."

L304-309: Nice quantification of the steady-flow assumption, which is the strength of the paper.

Thank you.

L314/315: Is this surprising ? Rather start this paragraph with why one would want to look at individual orbits and say that different incidence angles are more or less sensitive to horizontal flow variability.

No, it is not. We added the following sentence for a smoother transition:

"These results are more informative when examined with respect to each of the four orbits, as the sensitivity of DInSAR measurements to vertical and horizontal motion components depends on the incidence angle and look geometry (e.g., Rocca et al., 2010; Hu et al., 2014)."

L320: The different sensitivity of different orbits is interesting and additional evidence that the bending effect is at play. This signal should ideally be explored for a more glaciological research question.

We added the additional analysis of contributions due to bending to our analysis (Fig. 2AR).

L330: Is this a typo ? Orbit 43 here should be orbit 09, right ?

Thank you; this was a typo and has been corrected to orbit 09.

L331: Replace “different viewing geometry” to “higher sensitivity to horizontal displacement and variability thereof.”

We revised the sentence as follows:

“Descending orbit 09, with a high incidence angle, exhibited the opposite pattern: a higher correlation during the unsteady period than during the steady period (0.93 vs. 0.83 in Fig.7 a, b), which we attribute to its higher sensitivity to horizontal displacement and, consequently to velocity variability as well as ice extension and compression due to tidal bending.”

L333-346: I think this paragraph is speculative and the argumentation is invalid for the reasons outlined above and should therefore be removed. At a minimum, it should be moved from the results section to the Discussion section. The part talking about grounding-zone width (here distance between HL and point A) requires a dedicated investigation of the tidal forcing at the times of SAR data acquisition, which are here presented in their DInSAR combinations.
Similar to the width of the tidal-flexure zone (here Flexure zone span) and differential tidal displacement, where a linear regression over all the data seems physically unmotivated. The trends look statistically insignificant. I think this also presents either biases during the times of SAR-data acquisition or a temporal signal (which then can be linked to the emergence of new pinning points).

As per our response to the major comments, Figure 9 was removed, and information about flexure zone width and distance to the hinge line is presented in a table. Our discussion was revised with respect to ice thickness, stiffness, viscoelastic response to tidal flexure, and emerging pinning points:

“Based on the explored flexure-zone variables, such as distance to the hinge line and flexure-zone width, we found that the grounding line migrates differently along each of the three transects, in spite of experiencing the same water level changes. These distinct slices of the flexure zones along the three transects differ due to variations in bed geometry, ice thickness, and likely ice stiffness. While the width of the flexure zone is governed by ice stiffness, which is determined by ice thickness and the effective Young’s modulus (Wild et al., 2017), pinning points can modify these relationships. Pinning points act to physically prop up the floating ice tongue, at least at certain water levels, thereby influencing local ice stiffness, ice flexure patterns, and the extent of inland water intrusion with changing tides. Previous studies have shown that the presence of pinning points affects lateral and basal shear stresses, generating upstream compression and downstream tension, which, in turn, influences flow variability and can locally enhance the stiffness of the glacier system (Robel et al., 2017; Still & Hulbe, 2021). In this study, both of the observed pinning points are located near the lateral shear zones, which are associated with reduced ice stiffness (Wild et al., 2025). Therefore, the complex interplay between local ice stiffness and glacier bed geometry results in the observed flexure-zone widths, which tend to be larger near the glacier margins where the pinning points are present.”

L351-367: This reads all like Discussion and should be moved out of the results section.

This section was revised. Text providing interpretations and discussion of future grounding-line retreat along the eastern transect with respect to bed rises was moved to the Discussion section. The manuscript text reporting values for the grounding zone width, the calculated theoretical grounding zone width, and the present-day pinning points along the transects was retained here. This text presents our results based on Fig. 9 of the revised manuscript and, therefore, we believe it should remain close to the figure.

L355: Wait, I thought the observations are not aligned with the theory by Tsai and Gudmundsson ? What did I miss ?

We never claimed that the observations are inconsistent with Tsai's theory regarding the idea that the widest grounding zone occurs over the shallowest bed slope, and the steepest bed slope corresponds to the narrowest grounding zone (i.e., the expected direction of the relationship). The only discrepancy between the theory and the observations lies in the magnitude of the observed versus predicted values.

We added revised the sentence as follows:

"The western transect (Fig. 9 d), associated with the shallowest bed slope, had the largest grounding-zone width, whereas the central transect (Fig. 9 c), associated with the steepest slope, had the smallest grounding-zone width, which aligns with theory (Tsai & Gudmundsson, 2015) and observations from elsewhere (e.g., Rignot et al., 2024)."

L374: Here we switch from m/d to m/yr. Please choose one unit throughout the manuscript (text and figures) so velocities can be easier compared between satellite and GNSS.

Thank you for the suggestion, but we prefer to use the "m yr^{-1}" notation in this section because it facilitates comparison with previous studies cited in the Study site description as well as provides a context for the grounding-zone width calculated over the entire SAR data acquisition period. As clarified above, the daily velocities shown in Fig. 2 were used solely to support our daily GNSS observations. We added GNSS data as a scatter plot using "m yr^{-1}" notation to facilitate the comparison.

We added the following clarifying sentence:

"To provide context for the observed grounding-zone and flexure-zone widths with respect to changes in glacier velocities, and to allow direct comparison with previously reported annual velocities (Millan et al., 2017; Van Wychen et al., 2020), we present velocities in annual notation in Fig. 10."

L377: A 2.7 fold increase between flexure-zone and grounded glacier speeds. This is more than the temporal acceleration observed in the GNSS data (eye-balled from Fig. 2a about 2.1 fold). Taken together, these observations indicate temporal variability of longitudinal strain rates. Was

this signal investigated with respect to the emergence of new pinning points and short-term variability in buttressing strength? This provides another exciting avenue for this manuscript as spatial patterns could also be investigated with the data at hand.

We do not think so, as the GNSS data show a similar acceleration: the highest velocity peak was 1.09 m d^{-1}, while the steady-season velocity was about 0.4 m d^{-1}, corresponding to the same ~2.7-fold increase. Based on our GNSS records from 2019, 2024, and 2025, acceleration to 1.0 – 1.2 m d^{-1} is typical of the July velocity peak.

L386: Typo. "Fig. 11a" should be "Fig. 11c", right ? Also, the claim about larger ice discharge in the west requires calculation using the ice thicknesses shown in Fig. 10 and surface velocities.

Yes, thank you, the typo was corrected.
The following calculations were added to support the statement:

"The grounding zone, where the grounding line migrated during the observation period, exhibited differences in velocity along the three transects. The widest grounding zone along the western transect of the glacier was associated with the highest velocity of 166.4 m yr^{-1}, averaged over a year and across the observed grounding zone width (Fig. 10 c), and the thickest ice (mean value of 225.6 m, Fig. 9 d). This section discharged approximately 37 540 m^3 yr^{-1} of ice along a 1 m-wide flow line. The narrower and thinner grounding zones along the eastern and central transects, with mean thicknesses of 210.0 m and 200.5 m (Fig. 9 b, c), were associated with slower average velocities of 144.6 m yr^{-1} and 159.8 m yr^{-1} (Fig. 10 a, b), discharged approximately 30 366 m^3 yr^{-1} and 32 040 m^3 yr^{-1} along a 1 m-wide flow line, respectively."

L.387: Typo. "Fig. 11 b,c" should be "Fig. a,b", right ?

Yes, thank you, the typo was corrected.

Discussion: I found this section underwhelming as it's largely a summary of the presented results, without a clear evaluation of a glaciological hypothesis. I provided a few different avenues above and hope that depending on the target journal, this section will be reworked with either a remote-sensing focus or glaciological focus.

We revised the Discussion section, in particular, added discussion of the suggested bending effect to provide more detail on the remote-sensing aspects of the explored glaciological processes.

L400: Reword to "atmospheric pressure variability"

"Variability" was added to the phrase "atmospheric pressure."

L410: Reword "pinning points drastically affected" to "pinning points control"

The suggested replacement was done.

L421: I suggest starting this paragraph with how DInSAR assumes steady flow to isolate the differential vertical component commonly referred to as a fringe pattern delineating the tidal-flexure zone. Then state how the present study quantifies this assumption with available GNSS data. The difference for different orbits then depends on the sensitivity to horizontal flow variability, which should be investigated with the available SAR-derived velocity fields.

Thank you, this part was revised as follows:

"This study is based on a commonly used DInSAR approach, which assumes steady glacier flow between SAR data acquisitions so that it can be isolated through interferogram differencing to reveal the vertical component of displacement due to tidal flexure. We examined the validity of this assumption by quantifying the SAR line-of-sight error caused by velocity variability using available in situ GNSS records. We also assessed SAR line-of-sight error resulting from ice extension and compression due to tidal bending. Analysis of these errors with respect to four RCM orbits with different incidence angles, and both ascending and descending passes - thus different sensitivities to horizontal flow variability - provides insights about the robustness of our DInSAR flexure-zone delineations and the associated uncertainties."

L429: Typo. "39.6 deg" is "39.5 deg" in Tab. 1

Thank you, the typo was corrected.

L431: Replace "viewing geometry" to "sensitivity to horizontal flow variability"

The suggested replacement was done as follows: "its higher sensitivity to horizontal displacement".

L435: Regarding the loss of coherence. Did you try to mask only ice-covered areas when matching slcs (the step before calculating interferograms). ? I found that this often greatly increases coherence compared to offset fitting the entire scene.

We did not apply masking of only ice-covered areas during SLC matching. Our interferograms have very short temporal baselines (a few days), and overall coherence across the scenes is generally high. The main areas of coherence loss correspond to zones of very rapid ice motion/deformation, where phase decorrelation occurs due to physical displacement between acquisitions. In such cases, the loss of coherence is primarily driven by glacier dynamics and cannot be significantly improved through alternative coregistration strategies such as masking during offset estimation.

L437: Why are more in-situ velocity observations required ? I understood that the beginning of this paragraph that even though the velocity variability is large, the spurious signal in DInSAR issmall compared to the tidal forcing. I would have appreciated a statement how well only using SAR-derived velocity fields from speckle tracking can be used for estimating the spatial patterns of the steady flow assumption.

Thank you. We revised the sentence (please see below) to highlight the importance of using SAR-derived 3D/2D velocity fields. However, as mentioned above, we would like to retain our in situ GNSS analysis and consider the suggested only SAR-derived velocity analysis outside the scope of the current manuscript.

"With the increasing number of SAR observations, larger DInSAR datasets during the unsteady season can be used along with SAR-derived 2-D velocities (e.g., Samsonov et al., 2021), to further investigate and quantify how the flexure-zone delineation is affected by the SAR viewing geometry with respect to changes in glacier velocity and ice bending across the entire glacier grounding zone."

L443: I think this is the bending effect, it's just more pronounced if the background flow is elevated. Another motivation for using the 2D velocity fields for a detailed analysis.

We agree that 2D or 3D velocity fields can provide additional information that may be useful for a more detailed analysis in the future. However, the precision of SAR offset tracking is typically on the order of about 1/10–1/20 of the pixel size. For our data, with a pixel spacing of ~2.5 m, this corresponds to approximately 0.1–0.25 m, which is about an order of magnitude lower than the precision achievable with InSAR.

L450: This sentence is quite speculative as the observations only show that the grounding line has retreated and acceleration is non-linear, but not that it's a runaway effect (which would require a numerical model of viscous ice-dynamics).

We revised the sentence as follows:

"This study documents further glacier acceleration in the grounding zone, accompanied by the formation of the eastern notch, which was preceded by grounding-line retreat (Antropova et al., 2024). These findings support the idea that Milne Glacier, currently resting on a retrograde slope in its grounding zone, is likely susceptible to marine ice-sheet instability."

L494: I want to encourage the authors to perform the suggested further investigation of viewing geometry.

The suggested bending analysis has been added to the manuscript.

Data availability:
I understand that raw data are often subject to license agreements and can therefore not be published, but how about the processed InSAR and DInSAR images ? I want to encourage the authors to provide them (together with the datetimes of the corresponding SAR scenes) to the scientific community.

We revised the text as follows to better clarify the data availability:

"The authors do not have permission to share raw RCM data, but it can be accessed by registering a vetted user account (https://www.asc-csa.gc.ca/eng/satellites/radarsat/access-to-data/about.asp).

InSAR results, as well as other field, simulated, and modeled data used in this study, are available upon request. The IceBridge airborne radar data (IceBridge MCoRDS L2 Ice Thickness, Version 1 [IRMCR2_20140401_03], https://doi.org/10.5067/GDQ0CUCVTE2Q) are freely available at https://nsidc.org/data/irmcr2/versions/1"

Where is the IceBridge data available. Please provide a link and DOI.

We added details as per the comment above. This information is also included in the Acknowledgements and References sections.

The modern standard is to provide field-data sets to the scientific community through an online provider (PANGEA, USAP-DC, NSIDC, etc) about 2 years after they've been acquired. And not by request to the corresponding author.

Thank you for your suggestion. As of now we will be happy to share our datasets by request to provide additional clarifications. In the future, we will consider preparing our datasets for publishing online.

References:
Unify the formatting. Schoof, 2007 seems missing.

Thank you, Schoof (2007) was added to the list. We revised the formatting of references.

Figures

Figure 1:
I suggest adding two panels with the mean velocity fields over the steady and unsteady periods. Or the steady one, with the second showing the spatial variability of percentage acceleration during the unsteady phase. Also, add the extent of panel b to panel a. And the location of GNSS and the phase unwrapping point to panel b. Two arrows with flight direction and azimuth of the different orbits might also be useful to estimate the sensitivity to horizontal flow variability.

Thank you for the suggestions. We prefer not to include SAR-derived velocity fields, as our error analysis of DInSAR results is interpreted with respect to in situ GNSS observations. The extent of panel b has been added to panel a. We also added GNSS locations throughout the year to panel b; these locations were already present on panel a. We did not mark the phase unwrapping point in the figure, as it is represented by an arbitrarily stable reference area automatically defined by the algorithm. Details about the SAR viewing geometry (i.e., pass, incidence angle, and track angle represented as azimuth clockwise from True North) are presented in Table 1; therefore, we prefer not to repeat them here to avoid overcrowding of the introductory figure.

Figure 2:
Panel (a): What is going on with the SAR-derived velocity spike in 2024-01-02 ? It seems to overlap with an increase in air temperature - is this a real signal ?

We believe this is likely a real signal because: 1) it coincided with the air temperature rise, as you mentioned; 2) the spike was observed at other nearby locations (i.e., 3×3 pixel windows) and also associated with acceleration in Fig.11; 3) our second GNSS, located ~15 km up-glacier from the grounding zone, also recorded a small jump at this time (i.e., circled in Fig. 9AR below). Unfortunately, our GNSS located in the grounding zone stopped recording at this time, and we do not have any other solid confirmations;




Fig. 9AR: SAR-derived velocity (green) and GNSS-derived velocity in the grounding zone (black), and ~15 km up glacier(blue) of the grounding zone.

We added the following information to the manuscript:

"These SAR-derived velocities demonstrated a surprising spike in January–February 2024 (Fig. 2 a). This spike coincided with a rise in air temperature and likely represented real glacier acceleration. However, because the GNSS receiver ran out of battery and did not record in situ measurements during this period, these records were excluded from the velocity variability analysis."

Panel (c): Please change the pressure unit from "kPa" to hPa", which is the standard practical unit

We have changed the units to "hPa" on the plot.

Figure 3:
Panel (d): Please add a unit to the colorbar or in the caption. I think these are radians. It looks like there is a nice freely-floating area towards the end of the profiles, so please average the DInSAR displacement over some distance before the comparison to DSSH.

Yes, thank you, these are radians, which are dimensionless. We changed the label to "$-\pi$, π".
The reported DInSAR displacement values were averaged over the freely floating part of the transects.

Figure 4:
Move out of main text.

We retain panels d-f (i.e., correlation between DDInSAR displacement and DSSH) as we consider these results to be essential part of our analysis. Panels a-c (i.e., correlation between DDInSAR displacement and simulated tides, without IBE) were moved to the Appendix.
Please note that the final statistics changed slightly due to experimentation with corrections based on the cosine of the incidence angle and/or calibration based on the mean values. The final results are based on the approach described in the manuscript and cited above.

Figure 5:
Correcting the flexure curves for the "bending effect" would eliminate the spurious grounding-line migration within one local ice thickness (about plus/minus 300 m). Also, some flexure curves show distinct bumps upstream of the grounding line, which are indicative of viscoelasticity or grounding line migration between the SAR-data acquisitions. Why are the different shapes of the flexure curves not being discussed with respect to the timing (or tide, IBE) at the times of their underlying SAR scenes ?

Our additional bending analysis demonstrates that the mean error due to the bending effect is low for orbit 43 (i.e., mean absolute value of 0.06 cm) during both steady and unsteady periods. Therefore, we prefer to retain the DInSAR-derived displacement results without applying a bending correction.

The following statement was added in the figure caption:

"The line-of-sight errors due to tidal bending and velocity variability for orbit 43 are relatively low (Figure 7 c, d). During the steady period, the combined line-of-sight mean error is 0.12 cm, whereas during the unsteady period it is 0.3 cm. These errors result in uncertainties of ±7.4 m (i.e., less than one pixel) and ±18.4 m (i.e., less than two pixels), respectively, for the hinge-line delineations.

The following discussion of the surface displacement curves was added to the manuscript:

"The shape of the surface displacement curves suggests that ice ungrounding mostly occurs between 38 and 40 km down-glacier from point A. Two prominent surface displacement bumps, approximately 3 cm in height, located between 32 and 35 km, were derived from DInSAR images based on SAR acquisitions with relatively large temporal gaps. Specifically, the first DInSAR image was generated from RCM data acquired on September 15 (SSH: −8.3 cm), September 19 (SSH: −7.9 cm), October 17 (SSH: 4.4 cm), and October 21 (SSH: 9.8 cm) 2023. The second DInSAR image was computed from acquisitions on October 21 (SSH: 9.8 cm), October 25 (SSH: 13.8 cm), November 2 (SSH: −8.6 cm), and November 6 (SSH: 6.3 cm) 2023. Our uncertainty analysis, discussed in detail below, confirmed that errors due to ice bending were relatively low, whereas velocity-related errors exceeded the steady-period mean value for both DInSAR images. Therefore, we attribute these bumps to velocity variability and the associated surface changes occurring within the acquisition intervals.
Both of these curves also illustrate an additional noticeable displacement trend of diverging from zero at approximately 36 km and subsequently increasing or decreasing up to 7 cm at 41 km. The same trend was observed in two other DInSAR images based on SAR acquisitions from: 1)

November 6, 2023 (SSH: 6.4 cm), November 10, 2023 (SSH: 4.9 cm), and November 14, 2023 (SSH: −0.8 cm); and 2) March 1, 2024 (SSH: 4.5 cm), March 5, 2024 (SSH: 19.4 cm), February 10, 2024 (SSH: −3.3 cm), and February 14, 2024 (SSH: −12.1 cm). Errors due to velocity variability and ice bending were small (only a bending error of 0.1 cm in the March–February DInSAR image exceeded the mean error value) and well below the observed displacement. We attribute this displacement pattern to the combined effect of low and high tides, which likely induced a viscoelastic ice response (Wild et al., 2018)."

Figure 6:
Again, the main take-away is that including the IBE improves the correlation but the large RMSE are kept. Does this Figure need to be shown in the main text ?
Panel (a): cyan label: Reword to "Reconstructed SSH (tide-based only)" and black label to "Reconstructed SSH (tides plus IBE)"
Panels (b-e): Are unraveled (or single-tide) or DInSAR combination results plotted against each other ? I think it should be single tides, so the x-values would correspond to the red curve in panel a ? Is the x-label incorrect ?
Please swap the order of panel b/c and d/e, to match the description in L173 and logic of Fig.4

We removed panels related to the tide-based only analysis and change the primary focus of the figure to reconstruction analysis rather then the IBE. The updated figure is based on Fig. RA3 above.
The x-label is correct. The figure shows differential displacements at SAR data acquisition times based on the reconstructed time series vs. DInSAR-derived displacements.
Panels d and e were removed, and the order of panels b and c was swapped.

Figure 7:
Is this separating parts of Fig. 4 d/e into steady/unsteady periods as defined by GNSS data as colors ? Another strength of the presented analysis, and I think the whole story about including the IBE to improve the statistics can easily be moved here by including two additional panels and moving Figs. 4 and 6 to the Appendix/SI.

Yes, this figure separates parts of Fig. 4d/e from the original version of the manuscript into steady and unsteady periods. We prefer to move this figure, which generalizes the results for all orbits, to the Appendix and keep Figure 7 (i.e., correlation by orbits and by steady/unsteady) of the revised manuscript in the main text to better demonstrate horizontal errors due to velocity variability and bending across the orbits.

Figure 8:
I don't think this figure adds to the storyline and can be summarized when presenting Fig. 7 in the main text. It should therefore also be moved to the Appendix/SI.

We prefer to retain this figure and remove Figure 7 in the initial version of the manuscript, as explained in our response regarding Figure 7.

Figure 9:

For the reasons outlined above, temporarily integrating the data points by fitting a linear regression isn't physically motivated in the manuscript's current form. I think the panel (a) shows a signal that can be further explored, but since it doesn't add to the story Fig. 9 should therefore be removed from the manuscript.
We know from Schmeltz et al., 2002, and others that it's non-linear.

We removed this figure from the manuscript. Information about the flexure zone width and the distance to the hinge line is presented in table form instead (i.e., Table 2, 4).

Figure 10:
I was a bit confused if flexure zones or grounding zones are displayed in panel a and b-d ? I think it's grounding zone limits, please clarify.

Panel (a) is an example of flexure zone, (b)-(d) are grounding zones over observation period.
To improve figure clarity, we replaced the flexure-zone example in panel (a) with multiple flexure-zone delineations associated with low line-of-sight errors due to tidal bending and velocity variability (i.e., orbits 43 and 103) during the steady period and thus considered reliable. We also added DInSAR-derived grounding zones based on both "reliable" images and all DInSAR images for better comparison.

Figure 11:
Add "Hofmöller diagrams" at the start of the caption. Unify the labels to either m/d or m/yr. Also, add the location of the GNSS to one of the transects (consider a colored scatter with the same colormap, or its difference to the SAR-derived velocities through time). Interestingly, the acceleration in Jan/Feb 2024 occurs through the tidal-flexure zone. Was the signal investigated ?

We added "Hovmöller diagram of …" to the caption and included a colored scatter plot using the same colormap to show GNSS-derived locations along the central transect and velocities in the figure.

The units are shown in m yr^{-1} in this figure for both GNSS-derived and SAR-derived velocities, as explained above.

The question about acceleration was addressed above (i.e., under the "Figure 2 comment"): this signal coincided with a temperature increase. We believe it is a real signal, but we do not have any other reliable confirmation.

References

Chien, Y., Zhou, C., Sun, S., Chen, Y., Wang, T., Zhang, B., 2026. Ephemeral grounding on the Pine Island Ice Shelf, West Antarctica, from 2014–2023. The Cryosphere 20, 245–263. https://doi.org/10.5194/tc-20-245-2026

Freer, B.I.D., Marsh, O.J., Hogg, A.E., Fricker, H.A., Padman, L., 2023. Modes of Antarctic tidal grounding line migration revealed by Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) laser altimetry. The Cryosphere 17, 4079–4101. https://doi.org/10.5194/tc-17-4079-2023

Han, H., Lee, H., 2015. Tide-corrected flow velocity and mass balance of Campbell Glacier Tongue, East Antarctica, derived from interferometric SAR. Remote Sens. Environ. 160, 180–192. https://doi.org/10.1016/j.rse.2015.01.014

Li, T., Dawson, G.J., Chuter, S.J., Bamber, J.L., 2023. Grounding line retreat and tide-modulated ocean channels at Moscow University and Totten Glacier ice shelves, East Antarctica. The Cryosphere 17, 1003–1022. https://doi.org/10.5194/tc-17-1003-2023

Minchew, B.M., Simons, M., Riel, B., Milillo, P., 2017. Tidally induced variations in vertical and horizontal motion on Rutford Ice Stream, West Antarctica, inferred from remotely sensed observations. J. Geophys. Res. Earth Surf. 122, 167–190. https://doi.org/10.1002/2016JF003971

Padman, L., Erofeeva, S., Joughin, I., 2003. Tides of the Ross Sea and Ross Ice Shelf cavity. Antarct. Sci. 15, 31–40. https://doi.org/10.1017/S0954102003001032

Picton, H.J., Stokes, C.R., Jamieson, S.S.R., Floricioiu, D., Krieger, L., 2023. Extensive and anomalous grounding line retreat at Vanderford Glacier, Vincennes Bay, Wilkes Land, East Antarctica. The Cryosphere 17, 3593–3616. https://doi.org/10.5194/tc-17-3593-2023

Rack, W., King, M.A., Marsh, O.J., Wild, C.T., Floricioiu, D., 2017. Analysis of ice shelf flexure and its InSAR representation in the grounding zone of the southern McMurdo Ice Shelf. The Cryosphere 11, 2481–2490. https://doi.org/10.5194/tc-11-2481-2017

Ramanath, S., Krieger, L., Floricioiu, D., Diaconu, C.-A., Heidler, K., 2025. Automatic grounding line delineation of DInSAR interferograms using deep learning. The Cryosphere 19, 2431–2455. https://doi.org/10.5194/tc-19-2431-2025

Reeh, N., Christensen, E.L., Mayer, C., Olesen, O.B., 2003. Tidal bending of glaciers: a linear viscoelastic approach. Ann. Glaciol. 37, 83–89. https://doi.org/10.3189/172756403781815663

Rignot, E., Padman, L., MacAyeal, D.R., Schmeltz, M., 2000. Observation of ocean tides below the Filchner and Ronne Ice Shelves, Antarctica, using synthetic aperture radar interferometry: Comparison with tide model predictions. J. Geophys. Res. Oceans 105, 19615–19630. https://doi.org/10.1029/1999JC000011

Ross, N., Milillo, P., Nakshatrala, K.B., Ballarini, R., Stubblefield, A., Dini, L., 2024. Importance of ice elasticity in simulating tide-induced grounding line variations along prograde bed slopes. https://doi.org/10.5194/egusphere-2024-875

Schmeltz, M., Rignot, E., MacAyeal, D., 2002. Tidal flexure along ice-sheet margins: comparison of InSAR with an elastic-plate model. Ann. Glaciol. 34, 202–208. https://doi.org/10.3189/172756402781818049

Vaňková, I., Nicholls, K.W., Corr, H.F.J., Makinson, K., Brennan, P.V., 2020. Observations of Tidal Melt and Vertical Strain at the Filchner-Ronne Ice Shelf, Antarctica. J. Geophys. Res. Earth Surf. 125, e2019JF005280. https://doi.org/10.1029/2019JF005280

Wild, C.T., Alley, K.E., Muto, A., Truffer, M., Scambos, T.A., Pettit, E.C., 2022. Weakening of the pinning point buttressing Thwaites Glacier, West Antarctica. The Cryosphere 16, 397–417. https://doi.org/10.5194/tc-16-397-2022

Wild, C.T., Drews, R., Neckel, N., Lee, J., Kim, S., Han, H., Lee, W.S., Helm, V., Rosier, S.H.R., Marsh, O.J., Rack, W., 2025. Monitoring shear-zone weakening in East Antarctic outlet glaciers through differential InSAR measurements. The Cryosphere 19, 4533–4554. https://doi.org/10.5194/tc-19-4533-2025
Wild, C.T., Marsh, O.J., Rack, W., 2019. Differential interferometric synthetic aperture radar for tide modelling in Antarctic ice-shelf grounding zones. The Cryosphere 13, 3171–3191. https://doi.org/10.5194/tc-13-3171-2019
Wild, C.T., Marsh, O.J., Rack, W., 2018. Unraveling InSAR Observed Antarctic Ice-Shelf Flexure Using 2-D Elastic and Viscoelastic Modeling. Front. Earth Sci. 6, 28. https://doi.org/10.3389/feart.2018.00028
Wild, C.T., Marsh, O.J., Rack, W., 2017. Viscosity and elasticity: a model intercomparison of ice-shelf bending in an Antarctic grounding zone. J. Glaciol. 63, 573–580. https://doi.org/10.1017/jog.2017.15

References (RC1)

Antropova, Y. K., Mueller, D., Samsonov, S. V., Komarov, A. S., Bonneau, J., & Crawford, A. J. (2024). Grounding-line retreat of Milne Glacier, Ellesmere Island, Canada over 1966–2023 from satellite, airborne, and ground radar data. Remote Sensing of Environment, 315, 114478. https://doi.org/10.1016/j.rse.2024.114478

Black, T. E., & Joughin, I. (2022). Multi-decadal retreat of marine-terminating outlet glaciers in northwest and central-west Greenland. The Cryosphere, 16(3), 807–824. https://doi.org/10.5194/tc-16-807-2022

Carnahan, E., Catania, G., & Bartholomaus, T. C. (2022). Observed mechanism for sustained glacier retreat and acceleration in response to ocean warming around Greenland. The Cryosphere, 16(10), 4305–4317. https://doi.org/10.5194/tc-16-4305-2022

Chen, H., Rignot, E., Scheuchl, B., & Ehrenfeucht, S. (2023). Grounding Zone of Amery Ice Shelf, Antarctica, From Differential Synthetic-Aperture Radar Interferometry. Geophysical Research Letters, 50(6), e2022GL102430. https://doi.org/10.1029/2022GL102430

Ciracì, E., Rignot, E., Scheuchl, B., Tolpekin, V., Wollersheim, M., An, L., Milillo, P., Bueso-Bello, J.-L., Rizzoli, P., & Dini, L. (2023). Melt rates in the kilometer-size grounding zone of Petermann Glacier, Greenland, before and during a retreat. Proceedings of the National Academy of Sciences, 120(20), e2220924120. https://doi.org/10.1073/pnas.2220924120

Dawson, G. J., & Bamber, J. L. (2017). Antarctic Grounding Line Mapping From CryoSat-2 Radar Altimetry. Geophysical Research Letters, 44(23), 11,886-11,893. https://doi.org/10.1002/2017GL075589

Freer, B. I. D., Marsh, O. J., Hogg, A. E., Fricker, H. A., & Padman, L. (2023). Modes of Antarctic tidal grounding line migration revealed by Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) laser altimetry. The Cryosphere, 17(9), 4079–4101. https://doi.org/10.5194/tc-17-4079-2023

Fricker, H. A., & Padman, L. (2006). Ice shelf grounding zone structure from ICESat laser altimetry. Geophysical Research Letters, 33(15). https://doi.org/10.1029/2006GL026907

Fricker, H. A., Coleman, R., Padman, L., Scambos, T. A., Bohlander, J., & Brunt, K. M. (2009). Mapping the grounding zone of the Amery Ice Shelf, East Antarctica using InSAR, MODIS and ICESat. Antarctic Science, 21(5), 515–532. https://doi.org/10.1017/S095410200999023X

Hogg, A. E., Shepherd, A., Gourmelen, N., & Engdahl, M. (2016). Grounding line migration from 1992 to 2011 on Petermann Glacier, North-West Greenland. Journal of Glaciology, 62(236), 1104–1114. doi: 10.1017/jog.2016.83

Holmes, F. A., Barnett, J., Åkesson, H., Morlighem, M., Nilsson, J., Kirchner, N., & Jakobsson, M. (2025). Sea level rise contribution from Ryder Glacier in northern Greenland varies by an order of magnitude by 2300 depending on future emissions. The Cryosphere, 19(7), 2695–2714. https://doi.org/10.5194/tc-19-2695-2025

Kochtitzky, W., & Copland, L. (2022). Retreat of Northern Hemisphere Marine-Terminating Glaciers, 2000–2020. Geophysical Research Letters, 49(3), e2021GL096501. https://doi.org/10.1029/2021GL096501

Millan, R., Jager, E., Mouginot, J., Wood, M. H., Larsen, S. H., Mathiot, P., Jourdain, N. C., & Bjørk, A. (2023). Rapid disintegration and weakening of ice shelves in North Greenland. Nature Communications, 14(1), 6914. https://doi.org/10.1038/s41467-023-42198-2

Millan, R., Mouginot, J., & Rignot, E. (2017). Mass budget of the glaciers and ice caps of the Queen Elizabeth Islands, Canada, from 1991 to 2015. Environmental Research Letters, 12(2), 024016. https://doi.org/10.1088/1748-9326/aa5b04

Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J. L., Prats-Iraola, P., & Dini, L. (2022). Rapid glacier retreat rates observed in West Antarctica. Nature Geoscience, 15(1), Article 1. https://doi.org/10.1038/s41561-021-00877-z

Mohajerani, Y., Jeong, S., Scheuchl, B., Velicogna, I., Rignot, E., & Milillo, P. (2021). Automatic delineation of glacier grounding lines in differential interferometric synthetic-aperture radar data using deep learning. Scientific Reports, 11(1), Article 1. https://doi.org/10.1038/s41598-021-84309-3

Morlighem, M., Wood, M., Seroussi, H., Choi, Y., & Rignot, E. (2019). Modeling the response of northwest Greenland to enhanced ocean thermal forcing and subglacial discharge. The Cryosphere, 13(2), 723–734. https://doi.org/10.5194/tc-13-723-2019

Padman, L., King, M., Goring, D., Corr, H., & Coleman, R. (2003). Ice-shelf elevation changes due to atmospheric pressure variations. Journal of Glaciology, 49(167), 521–526. https://doi.org/10.3189/172756503781830386

Pawlowicz, R., Beardsley, B., Lentz, S., 2002. Classical tidal harmonic analysis including error estimates in MATLAB using T_TIDE. Comput. Geosci. 28 (8), 929–937. https://doi.org/10.1016/S0098-3004(02)00013-4.

Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., & Laaksonen, A. (2022). The Arctic has warmed nearly four times faster than the globe since 1979. Communications Earth & Environment, 3(1), 168. https://doi.org/10.1038/s43247-022-00498-3

Rignot, E., Ciracì, E., Scheuchl, B., Tolpekin, V., Wollersheim, M., & Dow, C. (2024). Widespread seawater intrusions beneath the grounded ice of Thwaites Glacier, West Antarctica. Proceedings of the National Academy of Sciences, 121(22), e2404766121. https://doi.org/10.1073/pnas.2404766121

Rignot, E., & MacAyeal, D. R. (1998). Ice-shelf dynamics near the front of the Filchner—Ronne Ice Shelf, Antarctica, revealed by SAR interferometry. Journal of Glaciology, 44(147), 405–418. https://doi.org/10.3189/S0022143000002732

Rignot, E. (1996). Tidal motion, ice velocity and melt rate of Petermann Gletscher, Greenland, measured from radar interferometry. Journal of Glaciology, 42(142), 476–485. https://doi.org/10.3189/S0022143000003464

Sykes, H. J., Murray, T., & Luckman, A. (2009). The location of the grounding zone of Evans Ice Stream, Antarctica, investigated using SAR interferometry and modelling. Annals of Glaciology, 50(52), 35–40. doi: 10.3189/172756409789624292

Van Wychen, W. V., Burgess, D., Kochtitzky, W., Nikolic, N., Copland, L., & Gray, L. (2020). RADARSAT-2 Derived Glacier Velocities and Dynamic Discharge Estimates for the Canadian High Arctic: 2015–2020. Canadian Journal of Remote Sensing, 46(6), 695–714. https://doi.org/10.1080/07038992.2020.1859359

Wild, C. T., Drews, R., Neckel, N., Lee, J., Kim, S., Han, H., Lee, W. S., Helm, V., Rosier, S. H. R., Marsh, O. J., & Rack, W. (2025). Monitoring shear-zone weakening in East Antarctic outlet glaciers through differential InSAR measurements. The Cryosphere, 19(10), 4533–4554. https://doi.org/10.5194/tc-19-4533-2025

Wood, M., Rignot, E., Fenty, I., An, L., Bjørk, A., van den Broeke, M., Cai, C., Kane, E., Menemenlis, D., Millan, R., Morlighem, M., Mouginot, J., Noël, B., Scheuchl, B., Velicogna, I., Willis, J. K., & Zhang, H. (2021). Ocean forcing drives glacier retreat in Greenland. Science Advances, 7(1), eaba7282. https://doi.org/10.1126/sciadv.aba7282

Wunsch, C., & Stammer, D. (1997). Atmospheric loading and the oceanic “inverted barometer” effect. Reviews of Geophysics, 35(1), 79–107. https://doi.org/10.1029/96RG03037

References (RC2)

Antropova, Y. K., Mueller, D., Samsonov, S. V., Komarov, A. S., Bonneau, J., & Crawford, A. J. (2024). Grounding-line retreat of Milne Glacier, Ellesmere Island, Canada over 1966–2023 from satellite, airborne, and ground radar data. Remote Sensing of Environment, 315, 114478. https://doi.org/10.1016/j.rse.2024.114478

Brunt, K. M., Fricker, H. A., Padman, L., Scambos, T. A., & O'Neel, S. (2010). Mapping the grounding zone of the Ross Ice Shelf, Antarctica, using ICESat laser altimetry. Annals of Glaciology, 51(55), 71–79. https://doi.org/10.3189/172756410791392790

Carnahan, E., Catania, G., & Bartholomaus, T. C. (2022). Observed mechanism for sustained glacier retreat and acceleration in response to ocean warming around Greenland. The Cryosphere, 16(10), 4305–4317. https://doi.org/10.5194/tc-16-4305-2022

Chartrand, A. M., Howat, I. M., Joughin, I. R., & Smith, B. E. (2024). Thwaites Glacier thins and retreats fastest where ice-shelf channels intersect its grounding zone. The Cryosphere, 18(11), 4971–4992. https://doi.org/10.5194/tc-18-4971-2024

Chen, H., Rignot, E., Scheuchl, B., & Ehrenfeucht, S. (2023). Grounding Zone of Amery Ice Shelf, Antarctica, From Differential Synthetic-Aperture Radar Interferometry. Geophysical Research Letters, 50(6), e2022GL102430. https://doi.org/10.1029/2022GL102430

Chien, Y., Zhou, C., Sun, S., Chen, Y., Wang, T., & Zhang, B. (2026). Ephemeral grounding on the Pine Island Ice Shelf, West Antarctica, from 2014–2023. The Cryosphere, 20(1), 245–263. https://doi.org/10.5194/tc-20-245-2026

Ciracì, E., Rignot, E., Scheuchl, B., Tolpekin, V., Wollersheim, M., An, L., Milillo, P., Bueso-Bello, J.-L., Rizzoli, P., & Dini, L. (2023). Melt rates in the kilometer-size grounding zone of Petermann Glacier, Greenland, before and during a retreat. Proceedings of the National Academy of Sciences, 120(20), e2220924120. https://doi.org/10.1073/pnas.2220924120

Costantini, M. (1998). A novel phase unwrapping method based on network programming. IEEE Transactions on Geoscience and Remote Sensing, 36(3), 813–821. https://doi.org/10.1109/36.673674

Forster, R. R., Rignot, E., Isacks, B. L., & Jezek, K. C. (1999). Interferometric radar observations of Glaciares Europa and Penguin, Hielo Patagónico Sur, Chile. Journal of Glaciology, 45(150), 325–337. https://doi.org/10.3189/S0022143000001829

Freer, B. I. D., Marsh, O. J., Hogg, A. E., Fricker, H. A., & Padman, L. (2023). Modes of Antarctic tidal grounding line migration revealed by Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) laser altimetry. The Cryosphere, 17(9), 4079–4101. https://doi.org/10.5194/tc-17-4079-2023

Fricker, H. A., Coleman, R., Padman, L., Scambos, T. A., Bohlander, J., & Brunt, K. M. (2009). Mapping the grounding zone of the Amery Ice Shelf, East Antarctica using InSAR, MODIS and ICESat. Antarctic Science, 21(5), 515–532. https://doi.org/10.1017/S095410200999023X

Gadi, R., Rignot, E., Menemenlis, D., & Scheuchl, B. (2025). Contrasting melt regime in the Ice Grounding Zone of Thwaites Glacier, West Antarctica. Proceedings of the National Academy of Sciences, 122(48), e2512626122. https://doi.org/10.1073/pnas.2512626122

Holmes, F. A., Barnett, J., Åkesson, H., Morlighem, M., Nilsson, J., Kirchner, N., & Jakobsson, M. (2025). Sea level rise contribution from Ryder Glacier in northern Greenland varies by an order of magnitude by 2300 depending on future emissions. The Cryosphere, 19(7), 2695–2714. https://doi.org/10.5194/tc-19-2695-2025

Howard, S. and Padman, L. (2021). Gr1kmTM: Greenland 1 kilometer Tide Model, Arctic Data Center, https://doi.org/10.18739/A2251FM3S

Hu, J., Li, Z. W., Ding, X. L., Zhu, J. J., Zhang, L., & Sun, Q. (2014). Resolving three-dimensional surface displacements from InSAR measurements: A review. Earth-Science Reviews, 133, 1–17. https://doi.org/10.1016/j.earscirev.2014.02.005

Kim, J. H., Rignot, E., Holland, D., & Holland, D. (2024). Seawater Intrusion at the Grounding Line of Jakobshavn Isbræ, Greenland, From Terrestrial Radar Interferometry. Geophysical Research Letters, 51(6), e2023GL106181. https://doi.org/10.1029/2023GL106181

Le Meur, E., Sacchettini, M., Garambois, S., Berthier, E., Drouet, A. S., Durand, G., Young, D., Greenbaum, J. S., Holt, J. W., Blankenship, D. D., Rignot, E., Mouginot, J., Gim, Y., Kirchner, D., de Fleurian, B., Gagliardini, O., & Gillet-Chaulet, F. (2014). Two independent methods for mapping the grounding line of an outlet glacier – an example from the Astrolabe Glacier, Terre Adélie, Antarctica. The Cryosphere, 8(4), 1331–1346. https://doi.org/10.5194/tc-8-1331-2014

Li, T., Dawson, G. J., Chuter, S. J., & Bamber, J. L. (2023). Grounding line retreat and tide-modulated ocean channels at Moscow University and Totten Glacier ice shelves, East Antarctica. The Cryosphere, 17(2), 1003–1022. https://doi.org/10.5194/tc-17-1003-2023

Mannerfelt, E. S., Schellenberger, T., & Kääb, A. M. (2025). Tracking glacier surge evolution using interferometric SAR coherence—Examples from Svalbard. Journal of Glaciology, 71, e43. https://doi.org/10.1017/jog.2025.27

MDA Space (MDA). (2021). RADARSAT Constellation Mission (RCM) product specification. Retrieved from: https://download-telecharger.services.geo.ca/pub/csa_asc/Space-technology_Technologie-spatiale/radarsat_constellation_mission_plan/RCM-SP-52-9092_Product_Spec_1-15_Public.pdf

Millan, R., Mouginot, J., & Rignot, E. (2017). Mass budget of the glaciers and ice caps of the Queen Elizabeth Islands, Canada, from 1991 to 2015. Environmental Research Letters, 12(2), 024016. https://doi.org/10.1088/1748-9326/aa5b04

Milillo, P., Rignot, E., Mouginot, J., Scheuchl, B., Morlighem, M., Li, X., & Salzer, J. T. (2017). On the Short-term Grounding Zone Dynamics of Pine Island Glacier, West Antarctica, Observed With COSMO-SkyMed Interferometric Data. Geophysical Research Letters, 44(20), 10,436-10,444. https://doi.org/10.1002/2017GL074320

Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J. L., Prats-Iraola, P., & Dini, L. (2022). Rapid glacier retreat rates observed in West Antarctica. Nature Geoscience, 15(1), Article 1. https://doi.org/10.1038/s41561-021-00877-z

Mohajerani, Y., Jeong, S., Scheuchl, B., Velicogna, I., Rignot, E., & Milillo, P. (2021). Automatic delineation of glacier grounding lines in differential interferometric synthetic-aperture radar data using deep learning. Scientific Reports, 11(1), Article 1. https://doi.org/10.1038/s41598-021-84309-3

Morlighem, M., Wood, M., Seroussi, H., Choi, Y., & Rignot, E. (2019). Modeling the response of northwest Greenland to enhanced ocean thermal forcing and subglacial discharge. The Cryosphere, 13(2), 723–734. https://doi.org/10.5194/tc-13-723-2019

Mortimer, C. A., Copland, L., & Mueller, D. R. (2012). Volume and area changes of the Milne Ice Shelf, Ellesmere Island, Nunavut, Canada, since 1950. Journal of Geophysical Research. Earth Surface; Washington, 117(4). http://dx.doi.org.proxy.library.carleton.ca/10.1029/2011JF002074

Padman, L., King, M., Goring, D., Corr, H., & Coleman, R. (2003). Ice-shelf elevation changes due to atmospheric pressure variations. Journal of Glaciology, 49(167), 521–526. https://doi.org/10.3189/172756503781830386

Pawlowicz, R., Beardsley, B., Lentz, S., 2002. Classical tidal harmonic analysis including error estimates in MATLAB using T_TIDE. Comput. Geosci. 28 (8), 929–937. https://doi.org/10.1016/S0098-3004(02)00013-4.

Picton, H. J., Stokes, C. R., Jamieson, S. S. R., Floricioiu, D., & Krieger, L. (2023). Extensive and anomalous grounding line retreat at Vanderford Glacier, Vincennes Bay, Wilkes Land, East Antarctica. The Cryosphere, 17(8), 3593–3616. https://doi.org/10.5194/tc-17-3593-2023

Porter, Claire, et al., 2023, "ArcticDEM, Version 4.1", https://doi.org/10.7910/DVN/3VDC4W, Harvard Dataverse, V1.

Rack, W., King, M. A., Marsh, O. J., Wild, C. T., & Floricioiu, D. (2017). Analysis of ice shelf flexure and its InSAR representation in the grounding zone of the southern McMurdo Ice Shelf. The Cryosphere, 11(6), 2481–2490. https://doi.org/10.5194/tc-11-2481-2017

Ramanath, S., Krieger, L., Floricioiu, D., Diaconu, C.-A., & Heidler, K. (2025). Automatic grounding line delineation of DInSAR interferograms using deep learning. The Cryosphere, 19(7), 2431–2455. https://doi.org/10.5194/tc-19-2431-2025

Rignot, E., Ciracì, E., Scheuchl, B., Tolpekin, V., Wollersheim, M., & Dow, C. (2024). Widespread seawater intrusions beneath the grounded ice of Thwaites Glacier, West Antarctica. Proceedings of the National Academy of Sciences, 121(22), e2404766121. https://doi.org/10.1073/pnas.2404766121

Rignot, E., Mouginot, J., & Scheuchl, B. (2011). Antarctic grounding line mapping from differential satellite radar interferometry. Geophysical Research Letters, 38(10). https://doi.org/10.1029/2011GL047109

Rignot, E., Padman, L., MacAyeal, D. R., & Schmeltz, M. (2000). Observation of ocean tides below the Filchner and Ronne Ice Shelves, Antarctica, using synthetic aperture radar interferometry: Comparison with tide model predictions. Journal of Geophysical Research: Oceans, 105(C8), 19615–19630. https://doi.org/10.1029/1999JC000011

Robel, A. A., Tsai, V. C., Minchew, B., & Simons, M. (2017). Tidal modulation of ice shelf buttressing stresses. Annals of Glaciology, 58(74), 12–20. https://doi.org/10.1017/aog.2017.22

Rocca, F., Prati, C., & Ferretti, A. (2010). Space-borne SARs: Impact of wavelengths and scan modes on ground motion studies. Annals of GIS, 16(2), 69–79. https://doi.org/10.1080/19475683.2010.492124

Ross, N., Milillo, P., Nakshatrala, K., Ballarini, R., Stubblefield, A., & Dini, L. (2025). Importance of ice elasticity in simulating tide-induced grounding line variations along prograde bed slopes. The Cryosphere, 19(6), 1995–2015. https://doi.org/10.5194/tc-19-1995-2025

Schoof, C. (2007). Ice sheet grounding line dynamics: Steady states, stability, and hysteresis. Journal of Geophysical Research: Earth Surface, 112(F3). https://doi.org/10.1029/2006JF000664

Still, H., & Hulbe, C. (2021). Mechanics and dynamics of pinning points on the Shirase Coast, West Antarctica. The Cryosphere, 15(6), 2647–2665. https://doi.org/10.5194/tc-15-2647-2021

Tsai, V. C., & Gudmundsson, G. H. (2015). An improved model for tidally modulated grounding-line migration. Journal of Glaciology, 61(226), 216–222. https://doi.org/10.3189/2015JoG14J152

Van Wychen, W. V., Burgess, D., Kochtitzky, W., Nikolic, N., Copland, L., & Gray, L. (2020). RADARSAT-2 Derived Glacier Velocities and Dynamic Discharge Estimates for the Canadian High Arctic: 2015–2020. Canadian Journal of Remote Sensing, 46(6), 695–714. https://doi.org/10.1080/07038992.2020.1859359

Wallis, B. J., Hogg, A. E., Zhu, Y., & Hooper, A. (2024). Change in grounding line location on the Antarctic Peninsula measured using a tidal motion offset correlation method. The Cryosphere, 18(10), 4723–4742. https://doi.org/10.5194/tc-18-4723-2024

Wild, C. T., Drews, R., Neckel, N., Lee, J., Kim, S., Han, H., Lee, W. S., Helm, V., Rosier, S. H. R., Marsh, O. J., & Rack, W. (2025). Monitoring shear-zone weakening in East Antarctic outlet glaciers through differential InSAR measurements. The Cryosphere, 19(10), 4533–4554. https://doi.org/10.5194/tc-19-4533-2025

Wild, C. T., Marsh, O. J., & Rack, W. (2019). Differential interferometric synthetic aperture radar for tide modelling in Antarctic ice-shelf grounding zones. The Cryosphere, 13(12), 3171–3191. https://doi.org/10.5194/tc-13-3171-2019

Wild, C. T., Marsh, O. J., & Rack, W. (2018). Unraveling InSAR Observed Antarctic Ice-Shelf Flexure Using 2-D Elastic and Viscoelastic Modeling. Frontiers in Earth Science, 6. https://doi.org/10.3389/feart.2018.00028

Wood, M., Rignot, E., Fenty, I., An, L., Bjørk, A., van den Broeke, M., Cai, C., Kane, E., Menemenlis, D., Millan, R., Morlighem, M., Mouginot, J., Noël, B., Scheuchl, B., Velicogna, I., Willis, J. K., & Zhang, H. (2021). Ocean forcing drives glacier retreat in Greenland. Science Advances, 7(1), eaba7282. https://doi.org/10.1126/sciadv.aba7282

Wunsch, C., & Stammer, D. (1997). Atmospheric loading and the oceanic “inverted barometer” effect. Reviews of Geophysics, 35(1), 79–107. https://doi.org/10.1029/96RG03037