

Continuing the MLS water vapor record with OMPS LP using neural networks

Michael D. Himes^{1,2}, Natalya A. Kramarova², Krzysztof Wargan^{3,2}, Sean M. Davis⁴, and Glen Jaross²

¹Morgan State University, Baltimore, MD, USA

²NASA Goddard Space Flight Center, Greenbelt, MD, USA

³Science Systems Association, Inc., Lanham, MD, USA

⁴NOAA Chemical Sciences Laboratory, Boulder, CO, USA

Correspondence: Michael D. Himes (michael.d.himes@nasa.gov)

Abstract. Stratospheric water vapor (SWV) plays an important role in atmospheric chemistry, dynamics, and radiative forcing. Satellite measurements by the Aura Microwave Limb Sounder (MLS), SciSat-1 Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS), and Stratospheric Aerosol and Gas Experiment III on the International Space Station (SAGE III/ISS) have provided key constraints on SWV for the past decades. MLS provides the best geographical coverage among these instruments, but it approaches the end of its life cycle in the coming years, which will result in a data desert for satellite-based SWV measurements given that ACE-FTS and SAGE III/ISS only measure at a few dozen geolocations per day. The Ozone Mapping and Profiler Suite Limb Profiler (OMPS LP) is flying aboard the Suomi National Polar-orbiting Partnership (SNPP) and NOAA-21 satellites and is planned for additional platforms in the coming years. While not designed to measure SWV, it shows weak sensitivity to it, particularly in the wake of the Hunga eruption’s significant injection of water vapor into the stratosphere. By utilizing the frequent co-locations between OMPS LP and MLS measurements, we developed a neural network-based retrieval algorithm to estimate SWV from SNPP OMPS LP radiances between 11.5–40.5 km. We find that the LP SWV profiles typically agree with MLS within 5%, and agreement with ACE-FTS and SAGE III/ISS is typically within 10%. We show that the SNPP-trained model is applicable to NOAA-21 OMPS LP without retraining, though minor differences in radiances between the instruments results in a $\sim 5\%$ bias under most conditions. Our results suggest that OMPS LP can continue the global water vapor record in the lower stratosphere into the 2030s, though continued independent measurements by satellite and balloon-borne instruments will be key to verifying the stability of our approach for quantifying decadal-scale SWV variability.

1 Introduction

Stratospheric water vapor (SWV) influences atmospheric dynamics, chemistry, and radiative forcing (e.g., Ramanathan and Inamdar, 2006; Charlesworth et al., 2023; Niemeier et al., 2023; Fleming et al., 2024). While SWV is typically 3–6 parts per million by volume (ppmv), water vapor concentrations in the upper troposphere can reach up to 1000 ppmv (e.g., Read et al., 2022). Deep convective systems and tropical upwelling via the Brewer-Dobson circulation can transport tropospheric air into the lower stratosphere, which comprises an important contribution to SWV (e.g., Fueglistaler et al., 2009; Khaykin et al., 2009;

Randel and Jensen, 2013; Dauhut et al., 2016). Rising atmospheric temperature due to climate change increases the amount of water vapor held in tropospheric air, which in turn increases the amount of water vapor transported into the stratosphere (Yue et al., 2019; Nowack et al., 2023). Long-term satellite measurements of SWV provide key constraints on the SWV budget and serve as important sources for data assimilation and validation of reanalysis frameworks (e.g., Davis et al., 2017; Hersbach et al., 2020; Wargan et al., 2023; Knowland et al., 2025).

Presently, satellite retrievals of SWV profiles are performed by the Aura Microwave Limb Sounder (MLS; Waters et al., 2006; Livesey et al., 2021), SciSat-1 Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS; Bernath et al., 2005; Boone et al., 2023), Stratospheric Aerosol and Gas Experiment III aboard the International Space Station (SAGE III/ISS Cisewski et al., 2014), and the Thermosphere Ionosphere Mesosphere Energetics Dynamics (TIMED) satellite's Sounding of the Atmosphere using Broadband Emission Radiometry (SABER; Russell et al., 1999) instruments. These instruments provide well validated H₂O products that typically show median differences within ~10% among each other as well as with ground-based and in-situ measurements (e.g., Carleer et al., 2008; Hurst et al., 2014; Rong et al., 2019; Davis et al., 2021; Park et al., 2021; De Los Ríos et al., 2024). Since May 2024, the MLS receiver used for the H₂O retrievals now only operates around 6 days per month to preserve measurement lifetime and will continue to do so until the end of the Aura mission, which significantly limits the spatiotemporal coverage of the MLS H₂O product. Following the decommissioning of Aura, ACE-FTS and SAGE III/ISS will continue to provide their H₂O products, but their geographical coverage is limited given they are solar occultation instruments. SABER takes around 1400 scans per day and, depending on time of year, views between 52°S–83°N or 83°S–52°N; the local times of the measurements change by up to 12 hours over TIMED's two-month yaw cycle. The Canadian High-altitude Aerosols, Water vapour, and Clouds mission (HAWC; Langille et al., 2025) is planned to launch early next decade, presenting a gap in global geographical coverage of SWV between Aura's decommissioning and HAWC's launch (Salawitch et al., 2025).

The Ozone Mapping and Profiler Suite (OMPS) Limb Profiler (LP; Dittman et al., 2002; Leitch et al., 2003) is currently flying aboard the Suomi National Polar-orbiting Partnership (SNPP; launched October 28, 2011) and NOAA-21 (launched November 10, 2022) satellites, and it is planned for the Joint Polar Satellite System (JPSS) 4 and 3 satellites, which are estimated to launch in 2027 and 2032, respectively. Using 3 slits, the instrument measures limb-scattered radiances between 290–1000 nm with a 1 km vertical sampling. Each satellite completes 14–15 Sun-synchronous orbits per day. SNPP's OMPS LP takes measurements every ~1° latitude, while NOAA-21's LP takes measurements every ~0.4° latitude, resulting in around 7000 and 17,500 measured radiance profiles per day, respectively. However, OMPS LP is only weakly sensitive to H₂O, which has challenged the application of traditional radiative transfer-based retrieval methods.

Our solution to retrieve water vapor profiles from OMPS LP measurements is deep learning (Goodfellow et al., 2016). Neural networks (NNs) learn to model complex, nonlinear processes in a data-driven manner. Given a set of corresponding inputs and outputs, the NN weights are tuned to approximate the underlying process, without explicit knowledge about it. Given OMPS LP's weak sensitivity to H₂O and the well validated MLS H₂O product, NNs could learn to accurately predict H₂O profiles from OMPS LP measurements at altitudes with sufficient sensitivity by using co-located MLS H₂O profiles as

the target outputs. This would result in an MLS-like H₂O product, thereby continuing the MLS global SWV record following the end of the Aura mission.

60 Here we present an OMPS LP water vapor product between 11.5–40.5 km produced by a NN trained on co-located LP-MLS measurements. In Section 2 we investigate the sensitivity of LP to water vapor under conditions before and after the Hunga Tonga–Hunga Ha’apai (hereafter Hunga) eruption, which injected more than 50 Tg of water into the stratosphere (e.g., Millán et al., 2022; Vömel et al., 2022; Khaykin et al., 2022; APARC, 2025). In Section 3 we discuss our methodology to train the NN as well as validate its predictions using other instruments. We present and discuss our results in Section 4, including the
65 limitations of our approach. Finally, we present our conclusions in Section 5.

2 OMPS LP sensitivity to H₂O

We investigate OMPS LP’s H₂O sensitivity using the Gauss-Seidel limb scattering (GSLs) radiative transfer model (RTM) (see Herman et al., 1994; Loughman et al., 2004, 2005, 2015). To calculate H₂O cross sections, we use the HITRAN 2020 database (Gordon et al., 2022) via the HITRAN Application Programming Interface (Kochanov et al., 2016). We then convolve those
70 high-resolution cross sections with the OMPS LP bandpasses such that RTM calculations at a given wavelength will be more consistent with what LP would measure for the assumed conditions. Using two selected co-located MLS H₂O profiles from before and after the Hunga eruption, we simulate radiances between 550–1025 nm in 5 nm intervals at altitudes of 3.5–50.5 km in 1 km intervals. For the two RTM simulations, all parameters are kept identical except for the H₂O profile ensuring that any differences in the Jacobians are due solely to variations in H₂O.

75 Based on the Jacobians output by the RTM, we select 12 wavelengths measured by OMPS LP (554, 596, 654, 720, 728, 824, 917, 929, 943, 956, 970, and 983 nm) that show the highest sensitivity to H₂O in their spectral region. Figure 1 shows an example of these Jacobians at 945 nm for the selected H₂O profile after the Hunga eruption. The H₂O enhancement between 20–30 km attributable to the Hunga eruption results in as much as a four times increase in LP’s sensitivity to H₂O. However, LP is very weakly sensitive to H₂O, and the sensitivity becomes negligible above 30 km. For more details on LP’s sensitivity
80 to H₂O, see Appendix A.

3 Deep learning methods

3.1 Data curation

Since there is currently no water vapor product derived from OMPS LP measurements, we instead use the MLS version 5 water vapor product as our ground truth. To prioritize times where SNPP and Aura have closely aligned orbits, we select dates that
85 have

- at least one orbit with 60 consecutive co-locations that are within 30 minutes and within 100 km, and
- at least 250 total co-locations on that day that satisfy the above co-location criteria.




Figure 1. Example of OMPS LP’s sensitivity to H_2O at 945 nm in the tropics. Panels (a) and (b) show the Jacobians for selected MLS H_2O profiles from before and after the Hunga eruption, respectively. Panel (c) shows the selected water vapor profiles, which show an enhancement in water vapor around 24 km attributable to the Hunga eruption. Panel (d) shows the ratio of (b) to (a). The increased water vapor concentration between 21–27 km due to Hunga results in LP’s sensitivity increasing by up to $4\times$ in this altitude range.

These criteria are satisfied every couple days due to their similar equatorial crossing times around 1:30 in the afternoon. On dates between February 2014 and December 2024 that satisfy the above criteria, we co-locate OMPS LP and MLS measurements within 6 hours and 100 km to build a data set of OMPS LP radiances and the corresponding MLS water vapor profiles. This results in 2,074,101 co-locations, with almost half occurring at high latitudes. For context, SNPP OMPS LP measures over 2.5 million vertical profiles per year.

We apply the recommended MLS data screening criteria except for the period of January 15 – February 12, 2022, as it has been shown that the recommended screening criteria filter out many valid profiles affected by Hunga (Millán et al., 2024). We limit the MLS water vapor profiles to ≤ 261 hPa. We log-linearly interpolate the water vapor profiles from the MLS pressure grid to OMPS LP’s geometric height (11.5–40.5 km in 1 km steps) using the NASA Global Earth Observing System Forward Processing for Instrument Teams (GEOS FP-IT; Lucchesi, 2015) pressures interpolated in time and space to the LP measurements. We limit the altitude range to 11.5–40.5 km because 10.5 km can exceed 261 hPa and the H₂O sensitivity of OMPS LP becomes ~ 0 above 40.5 km.

We also consider a similar methodology for ACE-FTS and SAGE III/ISS data with recommended screening criteria applied to both datasets. We utilize co-location criteria of within 1 day, within 2° latitude, and within 1113 km longitude (equal to 10° longitude at the equator), consistent with the criteria used in Davis et al. (2021). These data sets are used to investigate whether it is viable to train exclusively on ACE-FTS or SAGE III/ISS data and whether training on a combination of MLS, ACE-FTS, and/or SAGE III/ISS data offers benefits over only training on MLS data.

3.2 Neural network methods

For each co-located measurement, we construct an input–output pair to be used during NN training. The inputs are comprised of

- LP radiances at 554, 596, 654, 720, 728, 824, 917, 929, 943, 956, 970, and 983 nm,
- GEOS FP-IT pressures and temperatures interpolated in time and space to the LP measurements, and
- the solar zenith angle of the LP measurement.

These inputs are formatted as 2-D “images” (wavelength \times altitude) with four channels (radiance, pressure, temperature, solar zenith angle), similar to a standard RGBA image. The radiances vary at each point in the 2-D image, the pressures and temperatures vary only with respect to altitude, and the solar zenith angle is constant throughout. For each input image, the corresponding outputs are the co-located MLS H₂O profile that has been interpolated to the LP altitude grid. See Appendix B for a discussion of the relative importance of these inputs.

To address the latitudinal sampling bias inherent in the co-located data set, we first select a subset of the data such that there are roughly the same number of samples in each 5° latitudinal bin, resulting in 1,137,100 input–output pairs. We ensure that extrema for each latitudinal bin are included in this subset. We then split these data into training (used to update NN weights), validation (monitors for overfitting during training), and testing (tests model generality on unseen data after training

120 is complete) sets in a proportion of roughly 75%, 15%, and 10%, respectively. For context, this results in the training and validation sets containing a total of $\sim 3.7\%$ of all available SNPP OMPS LP data over the 2014–2024 period considered.

To investigate how the training data impacts the algorithm’s ability to generalize to unseen data, we run additional experiments where certain years are omitted from the training data set. One test omits 2015–2016, which exhibits typical SWV, while the other test omits 2024, which contains SWV perturbations from Hunga.

125 For data pre-processing and NN training, we utilize the open-source Python package MARGE (Himes et al., 2022), which uses TensorFlow (Abadi et al., 2016) via the Keras API. We pre-process the data by taking the base-10 logarithm of the OMPS LP radiances, GEOS FP-IT pressures, and MLS H₂O profiles, then scale each input and output parameter to be within the closed interval $[-1, 1]$ based on their training set extrema at each altitude.

To determine a neural network architecture well suited to solving this problem, we perform a Bayesian hyperparameter optimization (Akiba et al., 2019). The selected architecture is similar to the landmark AlexNet architecture (Krizhevsky et al., 2012); for details on our optimization procedure, the selected architecture, and the training details, see Appendix C. Using the chosen architecture, we train an ensemble of 10 neural networks using a mean-squared-error (MSE) loss function; members are only differentiated by their random initialization. The ensemble size is held constant for all retrievals; it was determined by adding ensemble members until the mean and standard deviation of the ensemble’s predictions did not significantly change. 135 The ensemble’s mean prediction provides the retrieved H₂O profile. The standard deviation among the ensemble provides an uncertainty estimate for the NNs’ prediction of the MLS H₂O profile. MLS profiles also have some uncertainty; we factor this into the LP uncertainty estimate by combining these sources in quadrature, assuming that they are uncorrelated:

$$\sigma_{LP}^2 = \sigma_{NN}^2 + \sigma_{MLS}^2 \quad (1)$$

We assume these terms are uncorrelated given that LP’s errors at the wavelengths considered are dominated by systematics, and these should be unrelated to any random or systematic errors in the MLS data; this assumption has a negligible effect on the 140 resulting LP uncertainty, as discussed in Section 4.1. Typically, σ_{NN} is small in the stratosphere due to close agreement among the NNs, leading the total LP uncertainties to be dominated by σ_{MLS} in this regime. In the troposphere, where strong scattering limits the NNs’ accuracy, the LP uncertainties are dominated by σ_{NN} . We estimate the MLS uncertainty by computing 5° monthly zonal means at each pressure level considered for the reported MLS uncertainties over the 2015–2019 period. The 145 chosen time period is a minor consideration, as the MLS data quality and description document reports that other time periods are similar to their reported January–March 2009 period (Livesey et al., 2022). These seasonal uncertainties are then log-linearly interpolated in time and pressure to the LP altitude grid using the GEOS-IT pressure data.

Additionally, we apply these methods to permutations of the co-located MLS, ACE-FTS, and/or SAGE III/ISS data sets. When combining more than one data set, we consider two approaches, one where we use the data as is, and another where we 150 bias correct the ACE-FTS and SAGE III/ISS data such that they have a global median difference of 0% at all altitudes with respect to MLS.

3.3 Evaluation and validation

To evaluate the NN’s typical accuracy and how well it generalizes to unseen data, we calculate the root mean square error (RMSE) and coefficient of determination (R^2) for the validation and test sets. We validate our LP water vapor product by comparing with satellite measurements, balloon-borne measurements, and an assimilation/reanalysis product.

For satellite measurements, we consider the MLS version 5 (Lambert et al., 2020), SAGE III/ISS version 6 (NASA/LARC/SD/ASDC, 2025), and ACE-FTS version 5.3 (Boone et al., 2023) products. For MLS, we consider co-locations within 6 hours, while for SAGE III/ISS and ACE-FTS we consider co-locations within 24 hours. For all three, we only consider the co-location if it is separated by less than 1000 km and within 2° latitude. When multiple co-locations satisfy these criteria, we use the co-location with the shortest distance. For each of these instruments, we compute both a global median percent difference as well as zonal median percent differences in 5° latitude bins. We found anomalous values in the SAGE III/ISS and ACE-FTS data sets that differ by a factor of up to 10,000 compared with the layers above and below it, even after applying each product’s recommended screening criteria. To screen out the most extreme of these unrealistic values, we apply a very conservative 20σ median rejection routine to the data set of percent differences.

We investigate whether our product shows similar properties as the MLS product by performing a multiple linear regression (MLR) on monthly means for a $5^\circ \times 5^\circ$ grid, with proxies for a linear trend, seasonal cycle, quasi-biennial oscillation (QBO), and El Niño Southern Oscillation (ENSO), as these terms explain the majority of stratospheric H_2O variability. For the seasonal cycle term, we also include a phase offset term for lag in months, given that it takes time for the effects to propagate upwards through the stratosphere. For the ENSO term, we regress using the sea surface temperature anomaly with a fitted lag in months, as previous work showed this is necessary to maximize correlation (e.g., Garcia et al., 2007; Calvo et al., 2010; Yu et al., 2022). For the QBO term, we use coefficients for the two leading empirical orthogonal functions for the QBO wind time series between January 1956–February 2025.

To compare with balloon-borne measurements, we consider NOAA Frost Point Hygrometer (Hurst et al., 2011) and Cryogenic Frost point Hygrometer (Vömel et al., 2007a, b) soundings from Boulder, USA; Hilo, USA; Lauder, New Zealand; San José, Costa Rica; Lindenburg, Germany; and Biak, Indonesia. For each sounding, we co-locate satellite measurements within 24 hours, 2° latitude, and 1113 km longitude (equal to 10° at the equator). If multiple co-locations meet these criteria, we select the profile that minimizes the distance from the sounding. We calculate the median absolute difference and percent difference over the data set of co-locations between each satellite instrument and the frost point measurements.

We additionally compare with the Modern-Era Retrospective analysis for Research and Applications version 2 (MERRA-2) Stratospheric Composition Reanalysis of Aura MLS (M2-SCREAM) reanalysis product (Wargan et al., 2023), which assimilates MLS products including H_2O , as this guarantees a co-location for all OMPS LP measurements. To assess whether our methodology blindly memorizes the days it sees during training (where co-locations with MLS are frequent) or if it learns to generalize to days it does not see during training (where co-locations with MLS are less frequent), we compute the mean differences and standard deviation of the differences between the LP product and M2-SCREAM for two subsets of 2021 data:




Figure 2. Performance summary for the ensemble of neural networks when applied to the test set. The relative RMSE (RMSE divided by the mean H_2O profile in the training set) is used to better highlight how the errors compare to the average H_2O VMRs at each altitude. Panel (a) shows the performance metrics calculated over the full test set. Error bars denote the standard deviation among the NNs’ performances. Panels (b) and (c) show the performance metrics calculated as a function of latitude. Below the tropopause, the RMSE is on the order of or larger than the average H_2O VMR, while in the stratosphere the RMSE is typically $<10\%$ of the average H_2O VMR.

185 one that contains the days where training data were drawn from, and another that contains the days where no data were used during training.

NOAA-21 OMPS LP has an insufficient period overlapping with MLS measurements (2023–present, with MLS only taking measurements for ~ 6 days per month since May 2024), inhibiting the use of NOAA-21 OMPS LP data for our NN training methodology. Given the imminent termination of Aura MLS and that the SNPP satellite will presumably cease operations before the end of NOAA-21 or the subsequent JPSS satellites, we test the application of our SNPP-trained model to NOAA-21
 190 OMPS LP measurements to determine whether our model can generalize to future iterations of the same instrument, as Himes et al. (2025b) found this to be the case for OMPS LP aerosol retrievals.

4 Results and Discussion

4.1 Training on MLS data

195 Figure 2 shows the global mean and standard deviation of the R^2 and RMSE metrics as well as the mean R^2 and RMSE as a function of latitude for the NN ensemble when applied to the test set of data not seen during training. Globally, R^2 is >0.7 except between 15.5–18.5 km. Figure 2c shows this reduction in R^2 is due to the upper troposphere in the tropics as

well as localized reductions in the mid-latitudes, discussed in detail below. When considering only events outside the tropics, the global R^2 for 15.5–16.5 km increases to ~ 0.7 , while considering only events within the tropics results in R^2 reducing to ~ 0.55 for these altitudes. In the stratosphere, the global RMSE is $< 10\%$ of the H_2O VMR. Errors increase below 18.5 km as measurements increasingly occur in the troposphere where water vapor VMR increases substantially. The discontinuity in RMSE at 32.5 km is related to the discontinuity in MLS v5 a priori profiles (Millán et al., 2024).

Figure 2c exhibits local minima in R^2 in the mid-latitudes. This reduction in R^2 is a numerical effect associated with how the NN models are trained combined with the variability of MLS water vapor. The equation for R^2 involves the ratio of the residual sum of squares (RSS) to the total sum of squares (TSS). The combination of the MSE loss function and data normalization procedure optimizes the NN models to make predictions such that the errors (RSS) are relatively evenly distributed (homoscedasticity). TSS, on the other hand, is proportional to the variance of the MLS water vapor values. When MLS water vapor variance is reduced, the R^2 must also reduce if the agreement between LP and MLS remains constant. These mid-latitude regions are local minima in MLS water vapor variance, and thus they are also local minima in R^2 given the expected homoscedasticity of the NN models. However, the absolute minimum in water vapor variance occurs in the SH mid-latitudes, while the absolute minimum in R^2 occurs in the NH mid-latitudes. In the NH, there is an additional effect due to reduced water vapor sensitivity at certain combinations of altitudes, SZA, single scattering angle, aerosol, and H_2O . Based on RTM simulations, the spectral features of water vapor at 930–970 nm shift from reductions in radiances in the SH and tropics to increases in radiances in the NH. At the transition between these regimes, water vapor sensitivity is at its minimum, corresponding to the minimum in R^2 in the NH mid-latitudes. For more details on this, see Appendix A.

Table 1 summarizes the OMPS LP H_2O product characteristics. The along-track resolution is $\sim 1^\circ$ for SNPP OMPS LP and $\sim 0.4^\circ$ for NOAA-21 OMPS LP. Given that the MLS H_2O product has a vertical resolution of 2.8–3.8 km over the pressure range we consider and that the LP vertical field of view is 1.2–1.5 km, the LP H_2O product’s vertical resolution may be between 1.2 and 3.8 km. However, more analysis is needed to better estimate the LP H_2O vertical resolution and how it changes as a function of altitude and under different conditions. At present, we conservatively report a vertical resolution ~ 3.8 km.

Figure 3 shows a comparison between the NN, MLS, and LP uncertainties for a co-located event in the tropics in January 2019. The NN uncertainty is smaller than the MLS uncertainty by a factor of ~ 2.5 in the stratosphere, while in the troposphere the NN uncertainty is larger than the MLS uncertainty by a factor of ~ 3 . Consequently, the combined LP uncertainty is dominated by the NN in the troposphere and by MLS in the stratosphere. We show two cases for this combined LP uncertainty, comprising the lower (uncorrelated) and upper (correlated) bounds for the combined NN-MLS uncertainty. They differ by ~ 0.1 ppm, which is negligible relative to the volume mixing ratios of typical water vapor profiles. The uncertainties reported within the LP H_2O product are calculated assuming no correlation (see Eq. 1), as the random and systematic errors of MLS are unlikely to be correlated with LP’s systematic errors at the wavelengths considered.

The results of our experiments that omit certain years from the training set reveal that some years are more important than others for achieving a well-generalized model. When omitting 2015–2016, we find that the model is generally unaffected and still performs well during those years. However, when omitting 2024, we find that the model begins producing severely inaccurate predictions by March 2024. This behavior is likely explained by the difference between these considered periods:

Table 1. OMPS LP H₂O product characteristics

Parameter	Instrument/Altitude	Value
Along-track resolution	SNPP OMPS LP	~1° latitude
	NOAA-21 OMPS LP	~0.4° latitude
Vertical resolution		~3.8 km
Precision, 0°–5°N	11.5 km	11.6 ppmv
	16.5 km	0.324 ppmv
	21.5 km	0.292 ppmv
	26.5 km	0.297 ppmv
	31.5 km	0.311 ppmv
	36.5 km	0.346 ppmv
Precision, 55°–60°N	11.5 km	0.754 ppmv
	16.5 km	0.322 ppmv
	21.5 km	0.316 ppmv
	26.5 km	0.325 ppmv
	31.5 km	0.344 ppmv
	36.5 km	0.377 ppmv

Notes. The reported vertical resolution is a conservative estimate based on the reported vertical resolution for MLS; there is not currently a per-altitude estimate. The reported precision values are the average uncertainties in the specified latitudinal bands.

while 2015–2016 were ordinary years in terms of SWV, the continued evolution of elevated SWV from the Hunga eruption into 2024 is atypical and not represented by the data available during 2014–2023. By including a small fraction of 2024 data during training, we find that the model continues performing accurately up to the present time of writing this manuscript. As the stratosphere returns to pre-Hunga conditions, we expect that the NNs will continue producing accurate retrievals of H₂O, but continued comparisons with other instruments designed to measure water vapor, such as ACE-FTS, will be critical to ensuring that accuracy.

Since the input radiances are at wavelengths affected by aerosols, we analyze the model errors as a function of aerosol extinction reported in the OMPS LP aerosol product. We note that a weak anti-correlation exists between the predicted H₂O VMR and the aerosol extinction at 675 nm in the lower stratosphere, though this is a real phenomenon rather than an artifact of our model. Stratospheric aerosol extinction generally peaks immediately above the tropopause and decreases over the few kilometers above it, while stratospheric H₂O VMR typically is at a minimum immediately above the tropopause due to the cold trap and increases over the few kilometers above it. We find that the NNs’ percentage error as a function of aerosol extinction at 675 nm is uncorrelated, indicating that the success of our approach is not dependent on aerosol conditions. See Appendix D for additional details.




Figure 3. Comparison of uncertainties for a co-located event in the tropics in January 2019. In the troposphere, the NN uncertainty is larger than the corresponding MLS uncertainty by a factor of ~ 3 , while in the stratosphere the MLS uncertainty is larger than the NN uncertainty by a factor of ~ 2.5 . The LP uncertainties are a combination of these NN and MLS uncertainties; the uncorrelated and correlated cases differ by ~ 0.1 ppm.

In general, the differences with respect to MLS are independent of the presence of upper tropospheric clouds identified by the OMPS LP measurements (Chen et al., 2016). However, events affected by polar stratospheric clouds (PSCs) identified by OMPS LP show a median bias around -2% at most altitudes. When considering the error as a function of distance below the PSC, the median bias can exceed -20% at 17 km below the PSC, which only occurs for PSCs at ≥ 28.5 km. However, the standard deviation of these errors can be substantial, where it averages around 33% between 15.5–22.5 km, with a maximum of 66% at 22.5 km. Given this behavior, we recommend that events contaminated by PSCs should not be used for scientific studies; the data product includes quality flags for these events.

4.2 Training on ACE-FTS and SAGE III/ISS data

We also considered training on co-located ACE-FTS and/or SAGE III/ISS data as an alternative to the co-located MLS data set. Results for these experiments were not satisfactory.

When training exclusively on ACE-FTS or SAGE III/ISS data, we find that the resulting NN models do not properly generalize to unseen data, indicating that these data sets are not sufficient to solve this problem. Part of this may be explained by the

typically significant measurement time differences between LP and the other instruments; LP measures in the early afternoon,
260 while ACE-FTS and SAGE III/ISS measure at sunrise/sunset, and these times only coincide for select geolocations depending
on the time of year. However, the number of co-locations seems to be a more limiting factor. When restricting the MLS data set
to similar sizes as the ACE-FTS and SAGE III/ISS data sets, we find that the resulting performance is poor. Given the success
when using the MLS-LP data set of ~ 1 million co-locations but the failure when considering tens of thousands of co-locations,
these results emphasize that our methodology relies on a large data set of co-located profiles.

265 Additionally, when including ACE-FTS and/or SAGE III/ISS data alongside the MLS data, we find significantly degraded
performance, regardless of whether or not the ACE-FTS and SAGE III/ISS data were de-biased with respect to MLS. This is
likely attributable to the variances between MLS, ACE-FTS, and SAGE III/ISS being on the order of the natural variability of
water vapor, which inhibits NN learning.

Despite our negative results when training on ACE-FTS and/or SAGE III/ISS data, we cannot rule out that alternative ML
270 approaches not considered here could utilize ACE-FTS and/or SAGE III/ISS data to derive water vapor profiles from LP
radiances.

4.3 Comparisons with satellite measurements

Figure 4 summarizes the global median percent differences between LP stratospheric H_2O profiles and co-located MLS, ACE-
FTS, and SAGE III/ISS profiles. For this comparison, we filter all LP tropospheric measurements by using the nearest co-
275 located tropopause altitude reported in the GEOS FP-IT product. The error bars show the standard error of the median, which
is generally negligible except at low altitudes for ACE-FTS and at high altitudes for SAGE III/ISS. Where LP detects a cloud,
we exclude any measurements at or below the cloud top, though we find that this criterion does not significantly alter the
results. Note that for the MLS comparisons we include all co-located data, including those used during training; this choice
does not bias the results, as discussed later in Section 4.5.

280 Differences with respect to MLS are less than 2% at all altitudes ≥ 14.5 km, with a maximum difference of 4.1% at 11.5 km.
When considering only 2025 data, the most extreme difference is 7.7% at 13.5 km, with a typical difference of $\sim 5\%$ below
22 km and $< 2\%$ above 25 km. Given that we trained on MLS data, this close agreement is expected and shows the model has
learned a good approximation to retrieve MLS-like water vapor profiles.

When comparing with ACE-FTS, we find agreement within 10% except between 11.5–13.5 km, where differences can reach
285 up to 19.3%. In general, the differences increase with altitude from -19.3% at 11.5 km up to 8.8% at 40.5 km.

LP's differences with respect to SAGE III/ISS are generally around 6% or less. Between 11.5–13.5 km, differences can reach
up to 16.9%. Given that a similar increase in differences at these altitudes is seen when comparing with both ACE-FTS and
SAGE III/ISS but not with MLS, this suggests that either ACE-FTS and SAGE III/ISS are biased high in this regime, or MLS
is biased low in this regime and the LP product has inherited this bias.

290 Figure 5 and Figure 6 show the median percent differences and standard error of the median, respectively, in 5° latitudinal
bins for the comparisons with MLS, ACE-FTS, and SAGE III/ISS. The results are generally consistent with those shown in
Figure 4. The notable exceptions occur at the lowest altitudes. For the comparisons with MLS, differences can reach up to 8%




Figure 4. Summary of global median percent differences between the LP stratospheric H₂O profiles and co-located (a) MLS, (b) ACE-FTS, and (c) SAGE III/ISS profiles. Horizontal uncertainty bars indicate the standard error of the median.

between 11.5 - 13.5 km just outside of the tropics. For ACE-FTS, the deviations at these altitudes can exceed 21%, but notably the standard error of the median is typically $\sim 11\%$ in this region, suggesting that this low bias may not be as substantial as it appears. However, comparisons with SAGE III/ISS also show this low bias at these altitudes, where the standard error is negligible. Together, this suggests that the LP product has a slight systematic low bias at 11.5–13.5 km just outside the tropics, but at latitudes $\geq 45^\circ$, the agreement with MLS suggests the biases when comparing with ACE-FTS and SAGE III/ISS are related to statistical differences between those products and MLS.

Figure 7 shows the “tape recorder” of alternating positive and negative anomalies in H₂O VMR, primarily attributable to seasonal changes in H₂O. For this plot, we subtract the pre-Hunga mean profile from each daily zonal mean to produce the daily anomaly. Before 2025, the OMPS LP and MLS tape recorders show excellent agreement throughout the stratosphere, with OMPS LP correctly capturing the increase in H₂O due to the Hunga eruption. For ~ 3 months in early 2025, OMPS LP shows a positive bias >1.0 ppm above 30 km that is not seen in the corresponding MLS data (Figure 7c). This bias is likely due to the weak H₂O sensitivity at these altitudes, which inhibits the NNs’ ability to reliably infer the H₂O VMR at these altitudes. Since the NNs correctly infer the H₂O at these altitudes before 2025, it suggests that the pre-2025 data were successfully predicted based on the shape of profiles at the lower altitudes that have sensitivity. With an absence of 2025 data in the training




Figure 5. Summary of median percent differences between the LP stratospheric H₂O profiles and co-located (a) MLS, (b) ACE-FTS, and (c) SAGE III/ISS profiles in 5° latitudinal bins. The pink line indicates the median tropopause altitude for a given latitudinal bin. Gray stippling indicates where there are no data, whether due to lack of statistical significance (high latitudes) or due to being in the troposphere (below the tropopause).

set, the NN model relies on profiles from the training data, primarily those influenced by elevated H₂O associated with the Hunga eruption. In 2025, the NNs perform reasonably well below 30 km, suggesting that there is sufficient sensitivity for the determined approximation to remain accurate when applied to unseen data, though there is a slight overestimation (~0.25 ppm) in mid-2025 between 25–30 km. We therefore advise that users exercise caution when using the OMPS LP H₂O product above 30 km in the tropics. Additionally, as the excess SWV from Hunga continues to evolve, it is possible that those conditions will be different enough from the training data that the NNs’ predictions become inaccurate; we will continue to monitor their performance to ensure the LP NN-based algorithm continues to provide reasonable H₂O profiles.

The NNs are largely trained on normal conditions. While Hunga is an anomaly, its impact has lasted years and can be seen globally, and thus the NNs are exposed to various Hunga-affected conditions. Since our data selection process does not consider H₂O concentration or profile states, rare short-lived events are severely undersampled relative to nominal conditions. Despite this, we find that the NN models can still retrieve reasonably accurate profiles for rare cases not seen during training. Figure 8 shows the H₂O profiles retrieved by LP and MLS for an event over the Pacific Ocean on 8 January 2020. This event exhibits a water vapor enhancement in the lower stratosphere due to the intense wildfires in Australia during December 2019 – January 2020. Pyrocumulonimbus clouds carried moist, tropospheric air into the lower stratosphere, where it was then transported eastward by winds. Despite that these H₂O concentrations were seen in <0.045% of the training data set, the NNs are able




Figure 6. Like Figure 5, but for the standard error of the median.

to identify this enhancement. The LP and MLS VMRs of this enhancement differ by up to a factor of two, but it is unclear whether this is due to LP underestimating the VMR, MLS overestimating the VMR, a combination of both, or differences in measured air masses.

325 4.4 Multiple linear regression analysis

In general, the results of our MLR analysis show similar behavior between the LP and MLS products, but the fitted coefficients for the LP data tend to be less than the corresponding MLS coefficients. The main exception to this is the seasonal phase offset, where both products closely agree. Figure 9 shows an example of the fitted coefficients for the seasonal amplitude at 14.5 km. The South Asian monsoon stands out clearly in both panels, though the LP product's fitted amplitudes for this region are around 1 ppm less than the corresponding fits for MLS.

Regarding water vapor trends, the LP product generally shows weaker trends in the stratosphere when compared with MLS. In some locations (particularly south and southeast Asia, central Africa, and central America), the LP product shows greater trends in the upper troposphere. Where both products show a trend of increasing H₂O, LP tends to show a lesser trend than MLS (Figure 10).




Figure 7. Parts-per-million anomaly in H₂O VMRs for the daily zonal means within 2.5° of the Equator for the (a) OMPS LP and (b) MLS water vapor products. Panel (c) shows the difference between the LP and MLS H₂O anomalies. The anomaly is determined by subtracting the pre-Hunga mean profile from each daily zonal mean profile. Beginning in May 2024, the MLS data become more sparse due to only taking measurements around 6 days each month.




Figure 8. LP and MLS water vapor profiles from 8 January 2020, which feature a lower stratospheric enhancement caused by the Australian wildfires, are plotted over a 2D histogram of the water vapor profiles outside the tropics included in the training data set. Each altitude is normalized such that the probabilities sum to 1. White regions indicate no data. Despite the low frequency of the VMRs of this enhancement in the training data set (<0.045%), the ML model successfully identifies this rare situation.

335 4.5 Comparisons with M2-SCREAM

Figure 11 compares the OMPS LP and M2-SCREAM H₂O products for the year 2021. Note that M2-SCREAM assimilates MLS v4.2, which is biased high for water vapor, while LP is trained on MLS v5, resulting in a persistent ~0.5 ppmv bias between the products. Days in which data were (Fig. 11a) or were not (Fig. 11b) included during training are shown separately but look almost identical, highlighting that the NNs' predictions are equally accurate whether or not they saw data from that day
 340 during training. Additionally, the standard deviation of the differences between OMPS LP and M2-SCREAM are consistently less than the standard deviation among OMPS LP or M2-SCREAM profiles, indicating that the OMPS LP product is more accurate than the natural variability of H₂O. Overall, our results suggest that the NN predictions are in good agreement with M2-SCREAM for data not seen during training.




Figure 9. Results for the seasonal amplitude at 14.5 km fitted via MLR for the (a) LP and (b) MLS water vapor products. The large amplitude over South Asia is attributable to the annual monsoon’s strong seasonal impact on water vapor. Panel (c) shows the difference between the fitted seasonal amplitude for the LP and MLS products.




Figure 10. Results for the linear trend in H₂O at 18.5 km fitted via MLR for the (a) LP and (b) MLS water vapor products. In general, LP shows a weaker trend of increasing H₂O than MLS.




Figure 11. Comparison between OMPS LP and M2-SCREAM for days in 2021 where (a) some data were included during training and (b) no data were included during training. Note that for days in which data were included during training, only a small percentage (1–4%) of data on those days were used in the training data set. For each panel, the left subpanel shows the probability density function of the differences between OMPS LP and M2-SCREAM as horizontal colored bars, one standard deviation of the differences as the vertical marks, the mean difference as plus signs, the mean difference \pm the standard deviation of OMPS LP H₂O as dashed lines, and the mean difference \pm the standard deviation of M2-SCREAM as dotted lines. The right subpanel shows the mean H₂O profile for each product. Panels (a) and (b) look nearly identical, indicating that the model is retrieving H₂O from the information content embedded in LP radiances rather than blindly memorizing the training data. The ~ 0.5 ppmv offset between the products is due to differences in the MLS version used by each product.

4.6 Comparisons with balloon-borne measurements

345 Figure 12 shows the median differences between satellite instruments (MLS, SAGE III/ISS, ACE-FTS, and OMPS LP) and the
 frost point hygrometer soundings from the six stations considered (see Section 3.3). The LP product agrees with the frost point
 measurements within 0.3 ppmv and within 10% between 16.5–27.5 km; this is in close agreement with the MLS results, which
 is expected given that we trained on MLS data. The only notable difference in agreement between MLS and LP is that the LP
 product shows a slightly reduced bias between 16.5–21.5 km. Like in Davis et al. (2021), we find that the satellite instruments
 350 show a dry bias in the upper troposphere compared to the frost point measurements, which may be due to spatiotemporal
 variability between the co-located measurements and/or reduced data quality in this regime.




Figure 12. Comparisons between satellite instruments and frost point balloon measurements. Panel (a) shows the median difference in ppmv limited to altitudes where the mean water vapor is <10 ppmv, and panel (b) shows the median percent difference. In both panels, the shaded regions show the median ± 2 standard errors of the mean.

4.7 Application to NOAA-21 OMPS LP

Paralleling the SNPP comparisons in Section 4.3, Figures 13 and 14 show respectively the global median percent differences and the 5° zonal median percent differences between NOAA-21 OMPS LP and MLS, ACE-FTS, and SAGE III/ISS. For global
 355 comparisons, NOAA-21 OMPS LP shows a persistent $\sim 5\%$ offset with respect to the corresponding SNPP comparisons at all altitudes. A similar low bias is seen in the NOAA-21 OMPS LP aerosol data, suggesting that this bias is attributable to differences in radiances between OMPS LP on SNPP and NOAA-21. Given our methodology, it is unclear whether the NNs have learned to implicitly account for a bias in the SNPP radiances or if the problem is related to the calibration of NOAA-21 radiances. However, this bias is not strictly a -5% shift for all conditions; the zonal comparisons show that the tropics exhibit
 360 a positive bias not seen in the corresponding SNPP comparisons. Further investigation is necessary to understand the cause of these biases. If the origin of these biases is not able to be determined, they could be addressed via a soft calibration approach.

5 Conclusions

We presented a water vapor retrieval product derived from SNPP OMPS LP measurements via a neural network (NN) trained on co-located MLS version 5 water vapor profiles. In general, the LP H_2O product is consistent with other water vapor products




Figure 13. Like Figure 4, but for NOAA-21 OMPS LP.

365 considered here. We find that our method typically agrees with MLS within 5% at all altitudes considered. The results of our multiple linear regression analysis show good correspondence between LP and MLS for seasonal water vapor variations, including for the south Asian monsoon. LP’s tape recorder in the tropics also shows close agreement with MLS, capturing both the alternating positive and negative seasonal anomalies as well as the large water vapor injection from the Hunga eruption. Agreement with SAGE III/ISS version 6 and ACE-FTS version 5.3 water vapor profiles is typically within 10% above 15 km and within 20% below 15 km. When compared with frost point balloon measurements, OMPS LP generally agrees within 10% in the stratosphere, closely mirroring comparisons between those frost point measurements and MLS. Comparisons with the M2-SCREAM reanalysis product show similar behavior between days included in training and days omitted from training, indicating that our method is retrieving H₂O from the LP radiances rather than memorizing the training data. Overall, we find that the LP product generally performs comparably to MLS over the 11.5–40.5 km altitude range considered, enabling the continuation of the MLS water vapor record for these altitudes. The exception is in the tropics above 30 km, where a period in 2025 shows larger errors relative to MLS than those seen during the 2014–2024 training period; we advise that users exercise caution when using LP H₂O data in this regime.

375




Figure 14. Like Figure 5, but for NOAA-21 OMPS LP.

When applying the same methodology but using SAGE III/ISS and/or ACE-FTS data for training, we find significantly reduced performance. We similarly find poor performance when limiting the MLS-LP data set to the same size as the SAGE
 380 III/ISS and ACE-FTS data sets, which suggests that the success of our approach relies on a large training data set of co-located profiles. When including SAGE III/ISS and/or ACE-FTS data alongside MLS data, we also find poor performance, regardless of whether or not the SAGE III/ISS and ACE-FTS data are bias corrected to have a 0% median difference with MLS at all altitudes. This suggests that the variances between the three satellite data products inhibit NN learning.

Despite insufficient co-located data to train a well generalized model specific for NOAA-21 OMPS LP, we find that the
 385 SNPP-trained NN is applicable to NOAA-21 OMPS LP measurements without retraining. For NOAA-21 data, we find a persistent negative bias of $\sim 5\%$ under most conditions when compared with the corresponding SNPP results; this pattern is also seen in comparisons between the SNPP and NOAA-21 OMPS LP aerosol products, suggesting that it is due to differences in the radiances rather than poor generalization of the SNPP-trained NN. However, the source of this bias is unclear at the time of writing; future work should explore approaches to identify the origin of this bias, characterize it, and correct it, if possible.
 390 Assuming that the NN model continues performing well over the coming years, our results suggest that this SNPP-trained model will be applicable to OMPS LP onboard JPSS-4 and 3, which are planned to launch in 2027 and 2032, respectively, thereby extending the MLS water vapor record into the 2030s, albeit at a reduced altitude range. Continued satellite and balloon-borne measurements from instruments with physics-based stratospheric H_2O products, such as ACE-FTS, SAGE III/ISS, and

frost point hygrometer soundings, will be integral to ensuring that our NN-based retrievals continue to perform well in the
395 coming years.

Code and data availability. The MARGE software is available on GitHub at <https://github.com/exosports/MARGE> (Himes, 2022). All data and results related to the MARGE software for this work are publicly available under the Reproducible Research Software License at <https://doi.org/10.5281/zenodo.17237404> (Himes et al., 2025a). The SNPP OMPS LP version 1.0 H₂O data product is available at <https://doi.org/10.5067/C1BD8BLEBH04> (Himes, 2025a). The NOAA-21 OMPS LP version 1.0 H₂O data product is available at <https://doi.org/10.5067/XNK38X2VQGZ0> (Himes, 2025b). The SNPP OMPS LP version 2.6 L1G data product is available at <https://doi.org/10.5067/YVE3FSNJ59RQ> (Jaross, 2023).
400

Appendix A: Additional details on OMPS LP sensitivity to H₂O

To better understand OMPS LP's sensitivity to H₂O, we use the GSLS RTM (Herman et al., 1994; Loughman et al., 2004, 2005, 2015) to simulate viewing conditions and atmospheric states measured by OMPS LP. Results can be grouped into two regimes, based
405 on the impact of aerosols: the southern latitudes and tropics, and the northern latitudes. In the southern latitudes and tropics, aerosol scattering does not significantly impact the H₂O signal. In the northern latitudes, LP's viewing conditions measure forward scattering from aerosols, resulting in significant changes in radiances that reduce the relative strength of the H₂O signal.

Figure A1 and Figure A2 show the RTM simulations and OMPS LP measurements, respectively, for an event in the tropics on
410 3 May 2019. A decrease in radiances attributable to H₂O absorption can be seen between 930–970 nm. The RTM simulations also show weak absorption around 724 and 824 nm, but the LP measurements do not have any obvious absorption features there. It is important to highlight that the RTM contains a number of simplifying assumptions that are not representative of reality (e.g., it assumes a simple reflective surface at the bottom of the atmosphere and ignores water vapor absorption within clouds) and therefore may be missing some important contributions in this regime necessary to model real measurements.

415 In the northern hemisphere, the H₂O signal become less clear. The interaction between aerosol scattering and H₂O causes the H₂O signal to weaken under these conditions, leading to no notable changes or even a small increase in radiances, rather than decrease as seen in the tropics and southern latitudes. Figure A3 shows RTM simulations for the viewing conditions and atmospheric state of a measurement by OMPS LP in the northern mid-latitudes on 3 May 2019. The simulation with H₂O shows no water vapor absorption but rather shows a small increase in radiances. When neglecting aerosols, this setup produces
420 a water vapor absorption signature between 930–970 nm as expected. Figure A4 shows the OMPS LP measurement for those conditions; there are no obvious spectral features attributable to H₂O, though the small positive bump at 956 nm may be related. Given the RTM results, there must be some transition region where the H₂O absorption feature vanishes before the radiances begin to increase as seen in Figure A3. That transition region likely occurs for specific combinations of SZA, single scattering angle, aerosol, and H₂O. Given the larger than expected reduction in R^2 around 40° N in Figure 2, we believe that feature is
425 in part explained by reduced H₂O sensitivity in this transition region. Measured radiances at higher latitudes begin to show a




Figure A1. Radiative transfer simulations with and without H₂O based on a measurement by OMPS LP on 3 May 2019 at 5.4° N latitude, 106.7° E longitude, where the solar zenith angle is 23.2° and the single scattering angle is 103.6°. Temperature and pressure profiles are from GEOS IT; O₃, surface reflectivity, and the aerosol extinction profile at 675 nm (extrapolated to other wavelengths assuming a gamma-function size distribution) are from the OMPS LP ozone retrieval algorithm; and the H₂O profile is from a co-located MLS measurement. Radiances are altitude normalized at 40.5 km. Black dashed vertical lines mark LP channels used in the ML model for the H₂O retrievals. The LP bandpasses are shown as black curves, which are vertically shifted to avoid overlaps. Original RTM simulations with water vapor absorption are performed at a 0.1 nm spectral resolution (blue curve). Simulated radiances convolved with LP bandpasses are shown in green. Convolved radiances simulated without accounting for water vapor absorption are shown in orange. The lower panel shows the differences between convolved radiances simulated with and without water vapor (differences between orange and green lines), highlighting spectral regions where water vapor absorption is expected.




Figure A2. Altitude-normalized radiances measured by OMPS LP for the event modeled in Figure A1. At these altitudes, including in the lower stratosphere, a small decrease in radiances is observed around 930–970 nm, which we believe is attributed to water vapor absorption. However, the measured radiances do not show reduction around 720 and 824 nm as predicted by the RTM simulations, possibly due to the RTM’s simplifying assumptions.

negative spectral slope in the 930–970 nm range, rather than monotonically increasing as would be expected under conditions with a strong aerosol signal, which we believe may be indicative of H₂O.

Rozanov et al. (2011) found that doubling tropospheric H₂O causes significantly larger changes to visible and near-infrared stratospheric radiances than doubling stratospheric H₂O, except in the 1353–1410 nm range where stratospheric perturbations result in larger changes to stratospheric radiances. We selected an OMPS LP measurement from 21 September 2015 to carry out a similar test using the GSLS RTM. The tropopause height and the temperature and pressure profiles are from GEOS-IT, the O₃ profile is from climatology, the surface reflectivity is set to 0.28, and the H₂O profile is from a co-located MLS measurement. For these tests, we neglect aerosols. Figure A5 shows the change in convolved radiances at various OMPS LP wavelengths when doubling H₂O at each altitude (a) in the troposphere and (b) in the stratosphere. Consistent with Rozanov et al. (2011), we find that doubling tropospheric H₂O has a more significant impact on stratospheric radiances than doubling stratospheric H₂O. However, when applying altitude normalization, we find the reverse. Figure A6 shows the change in convolved radiances




Figure A3. Like Figure A1, but for a measurement at 61.6° N latitude, 87.0° E longitude, solar zenith angle of 46.2° , and a single scattering angle of 49.5° . Simulations without aerosol are shown as dashed orange and green lines. Unlike in Figure A1, the lower panel here shows almost no differences between the simulations with and without H_2O in 930 – 970 nm spectral range, and even indicate a small increase in radiances simulated with H_2O compared with the simulation without H_2O . When omitting aerosol, radiances decrease due to H_2O absorption, confirming the increase in radiances for these viewing conditions is due to aerosol.

when applying altitude normalization at 40.5 km, demonstrating that doubling stratospheric H_2O has a more significant impact on altitude-normalized stratospheric radiances than doubling tropospheric H_2O . Whether altitude normalization is applied or not, Figures A5 and A6 show that the change in radiances is $<3\%$ when doubling H_2O concentrations for these conditions.

440 Despite that high surface reflectivities boost the impact of tropospheric perturbations and reduce the impact of stratospheric perturbations, it is not significant enough to alter these conclusions. However, given the RTM limitations mentioned above, it is possible that altitude normalization of real measurements does not cancel out contributions from tropospheric H_2O as much as suggested by these simulations.

445 Based on these analyses of simulated and measured radiances, we believe that the ML model derives water vapor from the spectral dependence of measured radiances and is able to distinguish different conditions by being exposed to a large range of cases during the training. Given the ML model's agreement with MLS, ACE-FTS, and SAGE III/ISS when applied to LP measurements in 2025 and 2026, the model can infer H_2O at northern high latitudes, because otherwise a lack of information content would manifest as reduced performance in NH regions compared to other regions, which are not seen in our validation




Figure A4. Like Figure A2, but for the event simulated in Figure A3. Unlike in Figure A2, there are no obvious spectral features attributable to H₂O in the 930–970 nm spectral band.

450 results. In an experiment where we added single scattering angle as additional input parameter (note that single scattering angle is consistently different between southern and northern hemispheres for the LP viewing geometry and can serve as a proxy for aerosol contamination), we found no change in results. This test, coupled with the tests in Appendix B, confirms that the model relies on the spectral signature of measured radiances to infer the information about the water vapor concentration under different conditions.

Appendix B: Investigation of relative importance of neural network input parameters

455 The NNs presented in the manuscript utilize a combination of LP radiances and solar zenith angles as well as GEOS-IT data for temperatures and pressures. To roughly assess the relative importance of these inputs, we conducted experiments where NNs were trained on (1) only the meteorological GEOS-IT data, (2) only the LP radiances, and (3) using pressure/temperature climatology in place of GEOS-IT data.




Figure A5. Change in convolved radiances at LP's resolution when doubling H₂O at each altitude (a) in the troposphere and (b) in the stratosphere. Tropospheric perturbations in H₂O have a larger impact than stratospheric perturbations.




Figure A6. Like Figure A5, but for convolved radiances altitude-normalized at 40.5 km. Altitude normalization cancels out much of the diffuse light coming from the surface and troposphere, leading to stratospheric perturbations having a larger impact than tropospheric perturbations on stratospheric radiances.




Figure B1. Like Figure 2, but also including results for NNs trained on (1) only GEOS-IT meteorology and (2) only LP radiances. The performance gap between the model trained on only GEOS-IT data and those trained with LP radiances indicates that the LP radiances offer additional information that enables a more accurate model.

Figure B1 shows the NN ensemble’s RMSE and R^2 values for the model presented in the manuscript alongside those performance metrics for NNs trained on (1) only the GEOS-IT meteorological data and (2) only LP radiances. The reduced performance of the NN trained on only GEOS-IT data indicates that the LP radiances enable the determination of a better solution for retrieving SWV. This performance difference can also be seen in Figure B2, which shows the tape recorder plot for the NN trained only on the GEOS-IT data. Notably, the GEOS-IT-only NN model features positive anomaly artifacts (especially in mid 2018), misses the SWV enhancement in January 2022 attributable to the Hunga eruption, and consistently shows larger deviations from MLS in 2025 than those seen in Figure 7. In contrast, the NN trained on only LP radiances and climatological meteorological data result in almost the same performance as the NN ensemble presented in the main text of the manuscript. Given these results, we conclude that the LP radiance data are the primary source of information content for this problem.

Appendix C: Neural network optimization, architecture, and training

To optimize the neural network architecture for this problem, we performed a Bayesian hyperparameter optimization over the number and types of layers, number of nodes per layer, and activation functions. We considered fully connected NNs, convolutional NNs, and architectures that utilize both fully connected and convolutional layers. In addition to standard fully




Figure B2. Like Figure 7, but for the NN trained on only the meteorological GEOS-IT data. The absence of LP radiance and solar zenith angle data results in reduced performance.

connected layers, we also considered Concrete Dropout layers (Gal et al., 2017), which include a trainable parameter for the layer’s dropout rate. For convolutional architectures, we considered architectures with and without pooling layers.

The selected architecture is similar to the landmark AlexNet architecture (Krizhevsky et al., 2012), with hidden layers and activation functions consisting of Conv2D(32)–ReLU–MaxPool2D–Conv2D(64)–ReLU–MaxPool2D–CD(256)–ReLU–CD(256)–ReLU, where Conv2D(m) indicates a two-dimensional convolutional layer with m feature maps using a kernel size of 5, ReLU indicates the rectified linear unit activation function, MaxPool2D indicates a two-dimensional pooling layer that selects the maximum value within a 2×2 window, and CD(n) indicates a Concrete Dropout layer with n nodes. This is followed by a fully-connected output layer of 30 nodes, corresponding to the H₂O VMR at the 30 altitudes spanning 11.5–40.5 km. Note that other architectures performed similarly to the selected architecture; we found that the training data set played a more significant role in model performance.

We optimized the learning rate policy according to the method described by Himes et al. (2025b) and trained each NN using the mean-squared-error loss over the validation set until early stopping engaged after a patience of 60 epochs. On average, models trained for 638 epochs, which required an average of almost 7 hours to train using an Nvidia V100 graphics processing unit.

On our processing system, running our retrieval algorithm using the central processing unit requires around 12 and 16 seconds to process one SNPP and NOAA-21 orbit, respectively; specialized graphics processing units would reduce this runtime.

Appendix D: Evaluation of sensitivity to aerosols

The LP wavelengths used in this study are affected by aerosols. However, as discussed in Section 4.1, the NN model errors are generally uncorrelated with aerosol extinction. Figure D1 shows a 2-D histogram of LP-MLS differences vs. aerosol extinction at 675 nm. The vast majority of the probability density is found clustered around a difference of $\sim 0\%$. At the largest extinctions (>0.01), there appears to be a slight negative bias, but this is generally related to the negative bias associated with PSCs discussed in Section 4.1. Figure D2 shows a time series of mean orbital LP-MLS differences, with three major eruptions over this time period marked for reference. The LP-MLS differences do not feature any notable deviations in the time following these eruptions, suggesting that variations in aerosol loading do not negatively impact the model performance.




Figure D1. Histogram of the LP-MLS percent differences vs. OMPS LP aerosol extinction at 675 nm at 11.5 km. Errors are clustered around 0%, indicating aerosols do not significantly impact model performance.




Figure D2. Time series plot of the average percentage difference between LP-MLS co-locations per LP orbit. Three major eruptions are denoted on the plot for context. Errors appear unrelated to those major eruptions.

Author contributions. MDH curated the data set used for the NNs; trained the NNs; developed the software that produces the data product; performed comparisons with MLS, SAGE III/ISS, and ACE-FTS; applied MLR to separate trend, seasonal, QBO, and ENSO contributions; and wrote the initial draft of the manuscript. NAK oversaw the project, provided guidance on the validation efforts, and provided advice on figures. KW performed the comparisons between the LP H₂O product and M2-SCREAM. SMD performed the comparisons between
500 satellite instruments and balloon-borne measurements. GJ suggested the initial idea for the project and provided advice based on previous efforts to retrieve H₂O from OMPS LP. All co-authors advised ways to investigate and improve the performance of the method, reviewed the manuscript, and provided advice on the text and figures.

Competing interests. At least one of the authors is a member of the editorial board of *Atmospheric Measurement Techniques*.

Acknowledgements. The authors would like to thank the OMPS LP characterization team for producing the Level-1 gridded data used in this
505 work. We are grateful for Alyn Lambert, William Read, Luis Millan, Nathaniel Livesey, and all members of the Aura MLS water vapor team for their work on the valuable data product that made this work possible. We also thank the science teams responsible for the ACE-FTS, SAGE III/ISS, and frost point hygrometer data used for validation in this work. Additionally, we thank James Johnson and Daniel Kahn for feedback on the data product formatting and assisting with releasing the data product. We appreciate the thoughtful comments provided by the anonymous reviewers as they have improved the quality of this manuscript. We also recognize contributors to NumPy, SciPy, Matplotlib,
510 TensorFlow, Keras, Optuna, Dask, the Python Programming Language, and the free and open-source community.

Financial support. This work was supported in part by the NASA Goddard Space Flight Center Earth Science Division's Strategic Science program and NASA grant 80NSSC26K0049.

References

- Abadi, M., Barham, P., Chen, J., Chen, Z., Davis, A., Dean, J., Devin, M., Ghemawat, S., Irving, G., Isard, M., et al.: Tensorflow: a system
515 for large-scale machine learning., in: OSDI, vol. 16, pp. 265–283, 2016.
- Akiba, T., Sano, S., Yanase, T., Ohta, T., and Koyama, M.: Optuna: A Next-Generation Hyperparameter Optimization Framework, in:
Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining, KDD '19, p. 2623–2631,
Association for Computing Machinery, New York, NY, USA, ISBN 9781450362016, <https://doi.org/10.1145/3292500.3330701>, 2019.
- APARC: The Hunga Volcanic Eruption Atmospheric Impacts Report, Yunqian Zhu, Graham Mann, Paul A. Newman, and William Randel
520 (Eds.), APARC Report No. 11, WCRP-10/2025, <https://doi.org/10.34734/FZJ-2025-05237>, 2025.
- Bernath, P. F., McElroy, C. T., Abrams, M. C., Boone, C. D., Butler, M., Camy-Peyret, C., Carleer, M., Clerbaux, C., Coheur, P.-F., Colin, R.,
DeCola, P., DeMazière, M., Drummond, J. R., Dufour, D., Evans, W. F. J., Fast, H., Fussen, D., Gilbert, K., Jennings, D. E., Llewellyn,
E. J., Lowe, R. P., Mahieu, E., McConnell, J. C., McHugh, M., McLeod, S. D., Michaud, R., Midwinter, C., Nassar, R., Nichitiu, F.,
Nowlan, C., Rinsland, C. P., Rochon, Y. J., Rowlands, N., Semeniuk, K., Simon, P., Skelton, R., Sloan, J. J., Soucy, M.-A., Strong,
525 K., Tremblay, P., Turnbull, D., Walker, K. A., Walkty, I., Wardle, D. A., Wehrle, V., Zander, R., and Zou, J.: Atmospheric Chemistry
Experiment (ACE): Mission overview, *Geophysical Research Letters*, 32, <https://doi.org/https://doi.org/10.1029/2005GL022386>, 2005.
- Boone, C., Bernath, P., and Lecours, M.: Version 5 retrievals for ACE-FTS and ACE-imagers, *Journal of Quantitative Spectroscopy and
Radiative Transfer*, 310, 108 749, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2023.108749>, 2023.
- Calvo, N., Garcia, R. R., Randel, W. J., and Marsh, D. R.: Dynamical Mechanism for the Increase in Tropical Upwelling in
530 the Lowermost Tropical Stratosphere during Warm ENSO Events, *Journal of the Atmospheric Sciences*, 67, 2331 – 2340,
<https://doi.org/10.1175/2010JAS3433.1>, 2010.
- Carleer, M. R., Boone, C. D., Walker, K. A., Bernath, P. F., Strong, K., Sica, R. J., Randall, C. E., Vömel, H., Kar, J., Höpfner, M., Milz, M.,
von Clarmann, T., Kivi, R., Valverde-Canossa, J., Sioris, C. E., Izawa, M. R. M., Dupuy, E., McElroy, C. T., Drummond, J. R., Nowlan,
C. R., Zou, J., Nichitiu, F., Lossow, S., Urban, J., Murtagh, D., and Dufour, D. G.: Validation of water vapour profiles from the Atmospheric
535 Chemistry Experiment (ACE), *Atmospheric Chemistry and Physics Discussions*, 8, 4499–4559, <https://doi.org/10.5194/acpd-8-4499-2008>, 2008.
- Charlesworth, E., Plöger, F., Birner, T., Baikhadzhaev, R., Abalos, M., Abraham, N. L., Akiyoshi, H., Bekki, S., Dennison, F., Jöckel, P.,
Keeble, J., Kinnison, D., Morgenstern, O., Plummer, D., Rozanov, E., Strode, S., Zeng, G., Egorova, T., and Riese, M.: Stratospheric water
vapor affecting atmospheric circulation, *Nature Communications*, 14, 3925, <https://doi.org/10.1038/s41467-023-39559-2>, 2023.
- 540 Chen, Z., DeLand, M., and Bhartia, P. K.: A new algorithm for detecting cloud height using OMPS/LP measurements, *Atmospheric Mea-
surement Techniques*, 9, 1239–1246, <https://doi.org/10.5194/amt-9-1239-2016>, 2016.
- Cisewski, M., Zawodny, J., Gasbarre, J., Eckman, R., Topiwala, N., Rodriguez-Alvarez, O., Cheek, D., and Hall, S.: The Stratospheric
Aerosol and Gas Experiment (SAGE III) on the International Space Station (ISS) Mission, in: *Sensors, Systems, and Next-Generation
Satellites XVIII*, edited by Meynart, R., Neeck, S. P., and Shimoda, H., vol. 9241 of *Society of Photo-Optical Instrumentation Engineers
(SPIE) Conference Series*, p. 924107, <https://doi.org/10.1117/12.2073131>, 2014.
- 545 Dauhut, T., Chaboureaud, J.-P., Escobar, J., and Mascart, P.: Giga-LES of Hector the Convectoid and Its Two Tallest Updrafts up to the
Stratosphere, *Journal of the Atmospheric Sciences*, 73, 5041 – 5060, <https://doi.org/10.1175/JAS-D-16-0083.1>, 2016.

- Davis, S. M., Hegglin, M. I., Fujiwara, M., Dragani, R., Harada, Y., Kobayashi, C., Long, C., Manney, G. L., Nash, E. R., Potter, G. L., Tegtmeier, S., Wang, T., Wargan, K., and Wright, J. S.: Assessment of upper tropospheric and stratospheric water vapor and ozone in reanalyses as part of S-RIP, *Atmospheric Chemistry and Physics*, 17, 12 743–12 778, <https://doi.org/10.5194/acp-17-12743-2017>, 2017.
- 550 Davis, S. M., Damadeo, R., Flittner, D., Rosenlof, K. H., Park, M., Randel, W. J., Hall, E. G., Huber, D., Hurst, D. F., Jordan, A. F., Kizer, S., Millan, L. F., Selkirk, H., Taha, G., Walker, K. A., and Vömel, H.: Validation of SAGE III/ISS Solar Water Vapor Data With Correlative Satellite and Balloon-Borne Measurements, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 803, <https://doi.org/https://doi.org/10.1029/2020JD033803>, e2020JD033803 2020JD033803, 2021.
- 555 De Los Ríos, K., Ordoñez, P., Stiller, G. P., Raspollini, P., Gai, M., Walker, K. A., Peña Ortiz, C., and Acosta, L.: Comparison of the H₂O, HDO and δ D stratospheric climatologies between the MIPAS-ESA V8, MIPAS-IMK V5 and ACE-FTS V4.1/4.2 satellite datasets, *Atmospheric Measurement Techniques*, 17, 3401–3418, <https://doi.org/10.5194/amt-17-3401-2024>, 2024.
- Dittman, M. G., Leitch, J. W., Chrisp, M., Rodriguez, J. V., Sparks, A. L., McComas, B. K., Zaun, N. H., Frazier, D., Dixon, T., Philbrick, R. H., and Wasinger, D.: Limb broadband imaging spectrometer for the NPOESS Ozone Mapping and Profiler Suite (OMPS), in: *Earth Observing Systems VII*, edited by Barnes, W. L., vol. 4814, pp. 120 – 130, International Society for Optics and Photonics, SPIE, <https://doi.org/10.1117/12.453751>, 2002.
- 560 Fleming, E. L., Newman, P. A., Liang, Q., and Oman, L. D.: Stratospheric Temperature and Ozone Impacts of the Hunga Tonga-Hunga Ha’apai Water Vapor Injection, *Journal of Geophysical Research (Atmospheres)*, 129, e2023JD039298, <https://doi.org/10.1029/2023JD039298>, 2024.
- 565 Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Reviews of Geophysics*, 47, <https://doi.org/https://doi.org/10.1029/2008RG000267>, 2009.
- Gal, Y., Hron, J., and Kendall, A.: Concrete dropout, in: *Proceedings of the 31st International Conference on Neural Information Processing Systems, NIPS’17*, p. 3584–3593, Curran Associates Inc., Red Hook, NY, USA, ISBN 9781510860964, 2017.
- Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., and Sassi, F.: Simulation of secular trends in the middle atmosphere, 1950–2003, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2006JD007485>, 2007.
- 570 Goodfellow, I., Bengio, Y., and Courville, A.: *Deep Learning*, MIT Press, 2016.
- Gordon, I., Rothman, L., Hargreaves, R., Hashemi, R., Karlovets, E., Skinner, F., Conway, E., Hill, C., Kochanov, R., Tan, Y., Wcislo, P., Finenko, A., Nelson, K., Bernath, P., Birk, M., Boudon, V., Campargue, A., Chance, K., Coustenis, A., Drouin, B., Flaud, J., Gamache, R., Hodges, J., Jacquemart, D., Mlawer, E., Nikitin, A., Perevalov, V., Rotger, M., Tennyson, J., Toon, G., Tran, H., Tyuterev, V., Adkins, E., Baker, A., Barbe, A., Canè, E., Császár, A., Dudaryonok, A., Egorov, O., Fleisher, A., Fleurbaey, H., Foltynowicz, A., Furtenbacher, T., Harrison, J., Hartmann, J., Horneman, V., Huang, X., Karman, T., Karns, J., Kass, S., Kleiner, I., Kofman, V., Kwabia-Tchana, F., Lavrentieva, N., Lee, T., Long, D., Lukashovskaya, A., Lyulin, O., Makhnev, V., Matt, W., Massie, S., Melosso, M., Mikhailenko, S., Mondelain, D., Müller, H., Naumenko, O., Perrin, A., Polyansky, O., Raddaoui, E., Raston, P., Reed, Z., Rey, M., Richard, C., Tóbiás, R., Sadiq, I., Schwenke, D., Starikova, E., Sung, K., Tamassia, F., Tashkun, S., Vander Auwera, J., Vasilenko, I., Viganin, A., Villanueva, G., 580 Vispoel, B., Wagner, G., Yachmenev, A., and Yurchenko, S.: The HITRAN2020 molecular spectroscopic database, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 277, 107 949, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2021.107949>, 2022.
- Herman, B. M., Ben-David, A., and Thome, K. J.: Numerical technique for solving the radiative transfer equation for a spherical shell atmosphere, *Appl. Opt.*, 33, 1760–1770, <https://doi.org/10.1364/AO.33.001760>, 1994.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, 585

- P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 590 Himes, M. D.: MARGE, GitHub, <https://github.com/exosports/MARGE> (last access: 15 August 2025), 2022.
- Himes, M. D.: OMPS-NPP LP L2 Water Vapor (H₂O) Vertical Profile swath orbital 3slit V1.0, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], <https://doi.org/10.5067/C1BD8BLEBH04>, 2025a.
- Himes, M. D.: OMPS-N21 LP L2 Water Vapor (H₂O) Vertical Profile swath orbital 3slit V1.0, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], <https://doi.org/10.5067/XNK38X2VQGZ0>, 2025b.
- 595 Himes, M. D., Harrington, J., Cobb, A. D., Güneş Baydin, A., Soboczenski, F., O’Beirne, M. D., Zorzan, S., Wright, D. C., Scheffer, Z., Domagal-Goldman, S. D., and Arney, G. N.: Accurate Machine-learning Atmospheric Retrieval via a Neural-network Surrogate Model for Radiative Transfer, *Planetary Science Journal*, 3, 91, <https://doi.org/10.3847/PSJ/abe3fd>, 2022.
- Himes, M. D., Kramarova, N. A., Wargan, K., Davis, S. M., and Jaross, G.: Research Compendium for Himes et al. (2026): "Continuing the MLS water vapor record with OMPS LP using neural networks", Zenodo [data set], <https://doi.org/10.5281/zenodo.17237404>, 2025a.
- 600 Himes, M. D., Taha, G., Kahn, D., Zhu, T., and Kramarova, N. A.: Using neural networks for near-real-time aerosol retrievals from OMPS Limb Profiler measurements, *Atmospheric Measurement Techniques*, 18, 2523–2536, <https://doi.org/10.5194/amt-18-2523-2025>, 2025b.
- Hurst, D. F., Oltmans, S. J., Vömel, H., Rosenlof, K. H., Davis, S. M., Ray, E. A., Hall, E. G., and Jordan, A. F.: Stratospheric water vapor trends over Boulder, Colorado: Analysis of the 30 year Boulder record, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/https://doi.org/10.1029/2010JD015065>, 2011.
- 605 Hurst, D. F., Lambert, A., Read, W. G., Davis, S. M., Rosenlof, K. H., Hall, E. G., Jordan, A. F., and Oltmans, S. J.: Validation of Aura Microwave Limb Sounder stratospheric water vapor measurements by the NOAA frost point hygrometer, *Journal of Geophysical Research: Atmospheres*, 119, 1612–1625, <https://doi.org/https://doi.org/10.1002/2013JD020757>, 2014.
- Jaross, G.: OMPS-NPP L1G LP Radiance EV Wavelength-Altitude Grid swath orbital 3slit V2.6, Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], <https://doi.org/10.5067/YVE3FSNJ59RQ>, 2023.
- 610 Khaykin, S., Pommereau, J.-P., Korshunov, L., Yushkov, V., Nielsen, J., Larsen, N., Christensen, T., Garnier, A., Lukyanov, A., and Williams, E.: Hydration of the lower stratosphere by ice crystal geysers over land convective systems, *Atmospheric Chemistry and Physics*, 9, 2275–2287, <https://doi.org/10.5194/acp-9-2275-2009>, 2009.
- Khaykin, S., Podglajen, A., Ploeger, F., Grooß, J.-U., Tence, F., Bekki, S., Khlopenkov, K., Bedka, K., Rieger, L., Baron, A., Godin-Beekmann, S., Legras, B., Sellitto, P., Sakai, T., Barnes, J., Uchino, O., Morino, I., Nagai, T., Wing, R., Baumgarten, G., Gerding,
- 615 M., Dufлот, V., Payen, G., Jumelet, J., Querel, R., Liley, B., Bourassa, A., Clouser, B., Feofilov, A., Hauchecorne, A., and Ravetta, F.: Global perturbation of stratospheric water and aerosol burden by Hunga eruption, *Communications Earth and Environment*, 3, 316, <https://doi.org/10.1038/s43247-022-00652-x>, 2022.
- Knowland, K. E., Wales, P. A., Wargan, K., Weir, B., Pawson, S., Damadeo, R., and Flittner, D.: Stratospheric Water Vapor Beyond NASA’s Aura MLS: Assimilating SAGE III/ISS Profiles for a Continued Climate Record, *Geophysical Research Letters*, 52, e2024GL112610, <https://doi.org/https://doi.org/10.1029/2024GL112610>, e2024GL112610 2024GL112610, 2025.
- 620 Kochanov, R. V., Gordon, I. E., Rothman, L. S., Wcisło, P., Hill, C., and Wilzewski, J. S.: HITRAN Application Programming Interface (HAPI): A comprehensive approach to working with spectroscopic data, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 177, 15–30, <https://doi.org/10.1016/j.jqsrt.2016.03.005>, 2016.

- Krizhevsky, A., Sutskever, I., and Hinton, G. E.: Imagenet classification with deep convolutional neural networks, in: *Advances in neural information processing systems*, pp. 1097–1105, 2012.
- Lambert, A., Read, W., and Livesey, N.: MLS/Aura Level 2 water vapor (H₂O) mixing ratio V005, Goddard Earth Sciences Data and Information Services Center (GES DISC) [data set], <https://doi.org/10.5067/Aura/MLS/DATA2508>, 2020.
- Langille, J., Rieger, L. A., Blanchard, Y., Blanchet, J.-P., Bourassa, A., Degenstein, D., Huang, Y., Strong, K., Walker, K., Zawada, D., Braun, S., Cole, J., Mariani, Z., McLinden, C., Paquin-Ricard, D., Sioris, C., Qu, Z., Wolde, M., Wang, X., Al-Abadleh, H. A., Ariya, P., Beltrami, H., Chang, R., Fletcher, C., Goldblatt, C., Grenier, P., Gyakum, J., Kushner, P., Luca, A. D., MacDougall, A. H., O'Neill, N., Pausata, F., Sica, R., Tan, I., Thériault, J. M., Tegtmeier, S., Toohey, M., Ward, W., and Wiacek, A.: The High-altitude Aerosols, Water vapour and Clouds mission: concept, scientific objectives and data products, *Bulletin of the American Meteorological Society*, pp. BAMS-D-23-0309.1, <https://doi.org/10.1175/BAMS-D-23-0309.1>, 2025.
- Leitch, J. W., Rodriguez, J. V., Dittman, M. G., Frazier, D., McComas, B. K., Philbrick, R. H., Wasinger, D., Valle, T. J., Dixon, T., Dooley, D., Munzer, R., and Larsen, J. C.: Limb scatter ozone profiling sensor for the NPOESS ozone mapping and profiler suite (OMPS), in: *Optical Remote Sensing of the Atmosphere and Clouds III*, edited by Huang, H.-L., Lu, D., and Sasano, Y., vol. 4891 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, pp. 13–21, <https://doi.org/10.1117/12.467527>, 2003.
- Livesey, N. J., Read, W. G., Froidevaux, L., Lambert, A., Santee, M. L., Schwartz, M. J., Millán, L. F., Jarnot, R. F., Wagner, P. A., Hurst, D. F., Walker, K. A., Sheese, P. E., and Nedoluha, G. E.: Investigation and amelioration of long-term instrumental drifts in water vapor and nitrous oxide measurements from the Aura Microwave Limb Sounder (MLS) and their implications for studies of variability and trends, *Atmospheric Chemistry and Physics*, 21, 15 409–15 430, <https://doi.org/10.5194/acp-21-15409-2021>, 2021.
- Livesey, N. J., Read, W. G., Wagner, P. A., Froidevaux, L., Santee, M. L., Schwartz, M. J., Lambert, A., Millan Valle, L. F., Pumphrey, H. C., Manney, G. L., Fuller, R. A., Jarnot, R. F., Knosp, B. W., and Lay, R. R.: Earth Observing System Aura Microwave Limb Sounder Version 5.0x Level 2 and 3 data quality and description document, Tech. Rep. JPL D-105336 Rev. B, Jet Propulsion Laboratory, https://mls.jpl.nasa.gov/data/v5-0_data_quality_document.pdf, 2022.
- Loughman, R., Flittner, D., Nyaku, E., and Bhartia, P. K.: Gauss–Seidel limb scattering (GSLs) radiative transfer model development in support of the Ozone Mapping and Profiler Suite (OMPS) limb profiler mission, *Atmospheric Chemistry and Physics*, 15, 3007–3020, <https://doi.org/10.5194/acp-15-3007-2015>, 2015.
- Loughman, R. P., Griffioen, E., Oikarinen, L., Postolyakov, O. V., Rozanov, A., Flittner, D. E., and Rault, D. F.: Comparison of radiative transfer models for limb-viewing scattered sunlight measurements, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD003854>, 2004.
- Loughman, R. P., Flittner, D. E., Herman, B. M., Bhartia, P. K., Hilsenrath, E., and McPeters, R. D.: Description and sensitivity analysis of a limb scattering ozone retrieval algorithm, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2004JD005429>, 2005.
- Lucchesi, R.: File Specification for GEOS-5 FP-IT, Tech. Rep. GMAO Office Note No. 2, NASA Global Modeling and Assimilation Office, <https://gmao.gsfc.nasa.gov/pubs/docs/Lucchesi865.pdf>, 2015.
- Millán, L., Santee, M. L., Lambert, A., Livesey, N. J., Werner, F., Schwartz, M. J., Pumphrey, H. C., Manney, G. L., Wang, Y., Su, H., Wu, L., Read, W. G., and Froidevaux, L.: The Hunga Tonga–Hunga Ha’apai Hydration of the Stratosphere, *Geophysical Research Letters*, 49, e2022GL099 381, <https://doi.org/10.1029/2022GL099381>, e2022GL099381 2022GL099381, 2022.

- 660 Millán, L., Read, W. G., Santee, M. L., Lambert, A., Manney, G. L., Neu, J. L., Pitts, M. C., Werner, F., Livesey, N. J., and Schwartz, M. J.: The Evolution of the Hunga Hydration in a Moistening Stratosphere, *Geophysical Research Letters*, 51, e2024GL110841, <https://doi.org/https://doi.org/10.1029/2024GL110841>, e2024GL110841 2024GL110841, 2024.
NASA/LARC/SD/ASDC: SAGE III/ISS L2 Solar Event Species Profiles (HDF5) V006, https://doi.org/10.5067/ISS/SAGEIII/SOLAR_HDF5_L2-V6.0, 2025.
- 665 Niemeier, U., Wallis, S., Timmreck, C., van Pham, T., and von Savigny, C.: How the Hunga Tonga—Hunga Ha’apai Water Vapor Cloud Impacts Its Transport Through the Stratosphere: Dynamical and Radiative Effects, *Geophysical Research Letters*, 50, e2023GL106482, <https://doi.org/https://doi.org/10.1029/2023GL106482>, e2023GL106482 2023GL106482, 2023.
Nowack, P., Ceppi, P., Davis, S. M., Chiodo, G., Ball, W., Diallo, M. A., Hassler, B., Jia, Y., Keeble, J., and Joshi, M.: Response of stratospheric water vapour to warming constrained by satellite observations, *Nature Geoscience*, 16, 577–583, 2023.
- 670 Park, M., Randel, W. J., Damadeo, R. P., Flittner, D. E., Davis, S. M., Rosenlof, K. H., Livesey, N., Lambert, A., and Read, W.: Near-Global Variability of Stratospheric Water Vapor Observed by SAGE III/ISS, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034274, <https://doi.org/https://doi.org/10.1029/2020JD034274>, e2020JD034274 2020JD034274, 2021.
Ramanathan, V. and Inamdar, A.: The radiative forcing due to clouds and water vapor, p. 119–151, Cambridge University Press, 2006.
Randel, W. J. and Jensen, E. J.: Physical processes in the tropical tropopause layer and their roles in a changing climate, *Nature Geoscience*, 6, 169–176, 2013.
- Read, W. G., Stiller, G., Lossow, S., Kiefer, M., Khosrawi, F., Hurst, D., Vömel, H., Rosenlof, K., Dinelli, B. M., Raspollini, P., Nedoluha, G. E., Gille, J. C., Kasai, Y., Eriksson, P., Sioris, C. E., Walker, K. A., Weigel, K., Burrows, J. P., and Rozanov, A.: The SPARC Water Vapor Assessment II: assessment of satellite measurements of upper tropospheric humidity, *Atmospheric Measurement Techniques*, 15, 3377–3400, <https://doi.org/10.5194/amt-15-3377-2022>, 2022.
- 680 Rong, P., Russell, J. M., Marshall, B. T., Gordley, L. L., Mlynczak, M. G., and Walker, K. A.: Validation of water vapor measured by SABER on the TIMED satellite, *Journal of Atmospheric and Solar-Terrestrial Physics*, 194, 105 099, <https://doi.org/https://doi.org/10.1016/j.jastp.2019.105099>, 2019.
Rozanov, A., Weigel, K., Bovensmann, H., Dhomse, S., Eichmann, K.-U., Kivi, R., Rozanov, V., Vömel, H., Weber, M., and Burrows, J. P.: Retrieval of water vapor vertical distributions in the upper troposphere and the lower stratosphere from SCIAMACHY limb measurements, *Atmospheric Measurement Techniques*, 4, 933–954, <https://doi.org/10.5194/amt-4-933-2011>, 2011.
- Russell, III, J. M., Mlynczak, M. G., Gordley, L. L., Jr., J. J. T., and Esplin, R. W.: Overview of the SABER experiment and preliminary calibration results, in: *Optical Spectroscopic Techniques and Instrumentation for Atmospheric and Space Research III*, edited by Larar, A. M., vol. 3756, pp. 277 – 288, International Society for Optics and Photonics, SPIE, <https://doi.org/10.1117/12.366382>, 1999.
- Salawitch, R. J., Smith, J. B., Selkirk, H., Wargan, K., Chipperfield, M. P., Hossaini, R., Levelt, P. F., Livesey, N. J., McBride, L. A.,
690 Millán, L. F., Moyer, E., Santee, M. L., Schoeberl, M. R., Solomon, S., Stone, K., and Worden, H. M.: The Imminent Data Desert: The Future of Stratospheric Monitoring in a Rapidly Changing World, *Bulletin of the American Meteorological Society*, 106, E540 – E563, <https://doi.org/10.1175/BAMS-D-23-0281.1>, 2025.
- Vömel, H., Barnes, J. E., Forno, R. N., Fujiwara, M., Hasebe, F., Iwasaki, S., Kivi, R., Komala, N., Kyrö, E., Leblanc, T., Morel, B., Ogino, S.-Y., Read, W. G., Ryan, S. C., Saraspriya, S., Selkirk, H., Shiotani, M., Valverde Canossa, J., and Whiteman, D. N.: Validation of
695 Aura Microwave Limb Sounder water vapor by balloon-borne Cryogenic Frost point Hygrometer measurements, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2007JD008698>, 2007a.

- Vömel, H., David, D. E., and Smith, K.: Accuracy of tropospheric and stratospheric water vapor measurements by the cryogenic frost point hygrometer: Instrumental details and observations, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/https://doi.org/10.1029/2006JD007224>, 2007b.
- 700 Vömel, H., Evan, S., and Tully, M.: Water vapor injection into the stratosphere by Hunga Tonga-Hunga Ha'apai, *Science*, 377, 1444–1447, <https://doi.org/10.1126/science.abq2299>, 2022.
- Wargan, K., Weir, B., Manney, G. L., Cohn, S. E., Knowland, K. E., Wales, P. A., and Livesey, N. J.: M2-SCREAM: A Stratospheric Composition Reanalysis of Aura MLS Data With MERRA-2 Transport, *Earth and Space Science*, 10, e2022EA002632, <https://doi.org/https://doi.org/10.1029/2022EA002632>, e2022EA002632 2022EA002632, 2023.
- 705 Waters, J., Froidevaux, L., Harwood, R., Jarnot, R., Pickett, H., Read, W., Siegel, P., Cofield, R., Filipiak, M., Flower, D., Holden, J., Lau, G., Livesey, N., Manney, G., Pumphrey, H., Santee, M., Wu, D., Cuddy, D., Lay, R., Loo, M., Perun, V., Schwartz, M., Stek, P., Thurstans, R., Boyles, M., Chandra, K., Chavez, M., Chen, G.-S., Chudasama, B., Dodge, R., Fuller, R., Girard, M., Jiang, J., Jiang, Y., Knosp, B., LaBelle, R., Lam, J., Lee, K., Miller, D., Oswald, J., Patel, N., Pukala, D., Quintero, O., Scaff, D., Van Snyder, W., Tope, M., Wagner, P., and Walch, M.: The Earth observing system microwave limb sounder (EOS MLS) on the aura Satellite, *IEEE Transactions on Geoscience and Remote Sensing*, 44, 1075–1092, <https://doi.org/10.1109/TGRS.2006.873771>, 2006.
- 710 Yu, W., Garcia, R., Yue, J., Russell III, J., and Mlynczak, M.: Variability of Water Vapor in the Tropical Middle Atmosphere Observed From Satellites and Interpreted Using SD-WACCM Simulations, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036714, <https://doi.org/https://doi.org/10.1029/2022JD036714>, e2022JD036714 2022JD036714, 2022.
- Yue, J., Russell III, J., Gan, Q., Wang, T., Rong, P., Garcia, R., and Mlynczak, M.: Increasing Water Vapor in the Stratosphere and Mesosphere After 2002, *Geophysical Research Letters*, 46, 13 452–13 460, <https://doi.org/https://doi.org/10.1029/2019GL084973>, 2019.
- 715