



1 Comparative Impact of Bio-Organic and Inorganic Fertilizer Application on Soil Health,
2 Grain Quality and Yield Stability in Nutrient Deficient Regions

3 Azhar Hussain¹ *, Zafar Iqbal² *, Asma Sabir³, Eman Alhomaiddi⁴, Maqshoof Ahmad¹

4 ¹Department of Soil Science, The Islamia University of Bahawalpur, 63100, Bahawalpur,
5 Pakistan

6 ²National Research Center of Intercropping, The Islamia University of Bahawalpur, 63100,
7 Bahawalpur, Pakistan

8 ³School of Environmental Sciences, University of Guelph, Canada

9 ⁴Department of Biology, College of Science, Princess Nourah bint Abdulrahman University,
10 P.O. Box 84428, Riyadh 11671, Saudi Arabia

11 * **Corresponding author:** azharhaseen@gmail.com; zaferiqbal31205@gmail.com


12 Abstract

13 Soil fertility limitations in arid regions restrict wheat productivity and grain nutritional quality,
14 with zinc (Zn) deficiency being a major concern. Sustainable soil amendments combining organic
15 and microbial inputs offer potential to address these constraints. This study aimed to evaluate the
16 effectiveness of bio-organic fertilization in enhancing wheat growth, yield, grain Zn
17 biofortification, and soil fertility under deficient arid field conditions. Two field trials were
18 conducted in Bahawalpur and Bahawalnagar, Pakistan, using a randomized complete block design.
19 Treatments included compost, ZnO (2%), ZnSO₄, zinc-solubilizing bacteria (ZSB), and their
20 combinations. Wheat growth, yield, grain nutrient concentrations, and soil fertility indicators
21 (organic matter, microbial biomass nitrogen (MBN), microbial biomass carbon (MBC), and
22 nutrient availability) were measured. Microbial populations were determined through colony-
23 forming units. Correlation and principal component analysis (PCA) were applied to explore
24 associations among variables. The integrated application of compost + ZnO + ZSB significantly
25 improved wheat height (19%), biomass (20%), yield attributes (10%), and grain Zn concentration
26 (39%) compared with the control. Soil fertility parameters also increased (organic matter, 39%;
27 MBN, 32%; MBC, 27%). Correlation and PCA highlighted strong positive relationships among
28 microbial populations, soil fertility, and crop performance. Bio-organic fertilization provides an
29 eco-friendly and effective strategy to improve wheat yield, Zn biofortification, and soil fertility in
30 arid agroecosystems.



Keywords: Eco-friendly; Microbial biomass; Organic matter; Soil health; Zinc solubilizing bacteria

Graphical Abstract



1. Introduction

Soil fertility degradation and micronutrient deficiency, particularly of zinc (Zn), are increasingly critical concerns in arid and semi-arid agroecosystems, where climatic stressors and anthropogenic pressures have exacerbated soil quality (Lal, 2024). In these fragile environments, characterized by low organic matter, poor water retention, and nutrient depletion, the production of staple crops, such as wheat (*Triticum aestivum* L.), is consistently threatened (Hossain et al., 2021). Zinc deficiency is among the most widespread micronutrient disorders affecting cereal crops globally, with particularly high prevalence in calcareous and coarse-textured soils typical of arid zones (Dhaliwal et al., 2022). It is estimated that over 50% of soils cultivated for cereals in arid and semi-arid regions suffer from Zn insufficiency, which not only reduces crop productivity but also compromises grain Zn concentration, a factor that directly affects human nutrition in regions where wheat is a dietary staple (Younas et al., 2023).



48 To address such multifaceted challenges, integrated soil fertility management approaches have
49 emerged, combining organic and inorganic strategies to enhance soil health, crop productivity, and
50 environmental sustainability (Imran, 2024). Among these, composting has garnered attention due
51 to its capacity to recycle organic waste into a valuable soil amendment that can improve physical
52 structure, chemical fertility, and biological activity (Sharma et al., 2024). Compost application
53 increases soil organic carbon (SOC), stabilizes soil aggregates, enhances microbial biomass carbon
54 (MBC) and microbial biomass nitrogen (MBN), and provides a slow-release source of macro- and
55 micronutrients (Khan et al., 2024). However, standard composts are often limited in their ability
56 to address specific micronutrient deficiencies such as Zn, particularly in soils with high pH, where
57 Zn becomes poorly available to plants due to adsorption and precipitation processes (Qian et al.,
58 2023).

59 In this context, the concept of "bioactivation" of compost, enriching compost with bioavailable
60 forms of essential nutrients such as Zn, along with the inclusion of beneficial microorganisms,
61 represents an innovative and promising solution (Manea and Bumbac, 2024). Bioactivated
62 compost not only supplies nutrients but also enhances microbial functions, potentially improving
63 nutrient cycling, enhancing soil enzymatic activity, and mitigating abiotic stress in plants (Clagnan
64 et al., 2023). The microbial solubilization of Zn compounds, for example, through the action of
65 zinc-solubilizing bacteria (ZSB), can significantly increase the bioavailability of Zn in the
66 rhizosphere, promoting better root development and nutrient uptake (Singh et al., 2024). When
67 applied to nutrient-deficient arid soils, bioactivated Zn-enriched compost may thus serve as a
68 multifaceted amendment to restore soil fertility, stimulate microbial activity, and ultimately
69 improve crop yield and quality (Maitra et al., 2024).

70 Despite its potential, the use of bioactivated Zn-enriched compost in arid regions remains
71 underexplored, particularly in terms of its comparative effects on key soil health indicators such
72 as organic matter content, SOC, MBC, MBN, and plant-available nutrients. Moreover,
73 understanding the interaction between compost-borne microbial communities and Zn dynamics in
74 the soil–plant system is essential for optimizing compost formulations and application strategies
75 (Wang et al., 2024). The wheat crop, as a high-input cereal with substantial nutrient demand, serves
76 as an ideal model for evaluating the effectiveness of such integrated nutrient management
77 approaches under stress-prone environmental conditions.

78



79 Recently, our research group reported the development of a bio-activated Zn-enriched compost
80 from organic waste and its application in a field experiment on wheat in Zn-deficient soils (Naeem
81 et al., 2025), where significant improvements in soil biochemical properties, Zn availability and
82 wheat grain yield were observed. However, that study was conducted at a single research site a
83 research area where all recommended practices were carried out and monitored by researchers. In
84 contrast, the present study expands upon that work by conducting field trials at two distinct
85 farmer-field locations under real-life agronomic conditions. The current research evaluates the
86 integrated application of Zn-enriched compost, combined with zinc-solubilizing bacteria, to
87 provide a broader assessment of the approach's agronomic performance, adaptability, and
88 scalability. This study, therefore, contributes new knowledge by validating the field-scale efficacy
89 of the bio-activated compost under variable environmental conditions, examining soil-crop
90 interactions across sites, and supporting the practical adoption of sustainable Zn fertilization
91 strategies in nutrient-deficient regions.

92 This study was conducted to evaluate the impact of microbial-assisted, Zn-enriched compost on
93 soil chemical and biological properties, as well as its subsequent effects on wheat productivity in
94 Zn-deficient arid soils. Specifically, the research aims to: assess changes in soil organic matter,
95 SOC, macro- and micronutrient content, microbial biomass carbon (MBC), and microbial biomass
96 nitrogen (MBN) following compost application; determine the effectiveness of Zn enrichment and
97 microbial activation in enhancing Zn availability and uptake; and quantify the response of wheat
98 growth, yield, and Zn content under field conditions typical of arid agroecosystems. The working
99 hypothesis is that microbial-assisted Zn-enriched compost leads to significant improvements in
100 soil fertility and microbial health, resulting in higher wheat productivity and nutritional quality
101 compared to conventional practices.

102 From an environmental perspective, the utilization of compost, particularly when derived from
103 agricultural, municipal, or agro-industrial residues, contributes to sustainable waste management
104 by diverting organic matter from landfills and mitigating greenhouse gas emissions. Composting
105 stabilizes organic residues, reducing methane generation and leachate formation, while also
106 sequestering carbon in soils—an important consideration in climate change mitigation.
107 Furthermore, the inclusion of beneficial microbes and micronutrient enrichment in compost
108 contributes to soil biodiversity and resilience, supporting the ecological functions that underpin
109 sustainable



110 2. Materials and Methods

111 2.1. Compost Preparation

112 To prepare zinc-enriched compost from domestic organic waste (Fruit and vegetable waste), a
 113 systematic approach comprising composting, zinc fortification, and microbial bioactivation was
 114 used. Initially, segregated domestic organic waste (vegetable peels, fruit residues, kitchen scraps)
 115 is collected and pre-processed by shredding to reduce particle size for faster decomposition. A
 116 composting drum was used, where the organic material was layered with urea (1% w/w) to speed
 117 up the composting process. Moisture is maintained at 40–50%, and the pile is turned periodically
 118 to ensure aerobic conditions. After the thermophilic phase (21 days), during the mesophilic stage,
 119 zinc was introduced in the form of zinc oxide (ZnO) at a predetermined concentration (2% w/w).
 120 This timing ensures optimal microbial assimilation and mineral retention within the compost
 121 matrix. The composting continues for 28 days until maturation, characterized by a dark brown
 122 color and crumbly texture. For bioactivation, a microbial consortium of Zinc solubilizing bacteria
 123 (ZSB) is prepared by culturing strains such as *Bacillus subtilis* (IUB2; accession No. MN696212
 124 and IUB6; accession No. MN696214), *Bacillus velezensis* (IUB3; accession No. MN696213),
 125 *Bacillus vallismortis* (IUB10; accession No. MN696215) and *Bacillus megaterium* (IUB11;
 126 accession No. MN696216) in a nutrient broth for 24–48 hours at 28–30°C until reaching an optical
 127 density of 0.8–1.0 at 600 nm. The mature compost is inoculated with the ZSB consortium (0.1%
 128 v/w) by spraying the bacterial suspension uniformly over the compost during final turning.
 129 Moisture is adjusted to around 40% to support microbial activity, and the compost is incubated for
 130 an additional 2 days under shaded conditions to stabilize the bioactivation process. The final
 131 microbial-assisted zinc-enriched compost is then dried, sieved, and stored for agronomic use. This
 132 dual enrichment process enhances the compost's micronutrient content and microbial efficiency
 133 for sustainable soil fertility management.

134 The bacterial strains used in the present study had already been characterized for zinc and
 135 phosphorus solubilization, catalase and urease activity, exopolysaccharide, siderophore and indole
 136 3 acetic acid production and plant growth promotion (Naseem et al., 2022). The elevated levels of
 137 nitrogen (3.36%), phosphorus (23 mg kg⁻¹), potassium (221 mg kg⁻¹), iron (12.43 mg kg⁻¹) and
 138 zinc (23.14 mg kg⁻¹) observed in the Zn-enriched compost, as compared to the unenriched
 139 compost, are primarily due to incorporation of nutrient-rich materials during the enrichment
 140 procedure. Zinc oxide was added, and beneficial rhizobacteria were introduced to enhance the



solubilization and availability of nutrients. The increased Zn and Fe concentrations are a direct result of these inputs, while the rise in N, P and K levels can be attributed to enhanced microbial activity. This microbial stimulation promotes the decomposition of organic matter and accelerates nutrient cycling. Additionally, the introduced microbes contribute by mobilizing phosphorus and micronutrients through the secretion of organic acids and chelating agents. As a result, the enrichment process not only enhances the compost's nutrient content but also improves its microbial efficacy, resulting in improved nutrient availability.

2.2. Field trial

Two field experiments were conducted at separate arid locations, Bahawalpur (29.3544° N, 71.6911° E) and Bahawalnagar (29.1903° N, 72.6343° E), to evaluate the effects of ten (10) different treatments on crop performance. The weather data, including monthly average temperature and rainfall (mm) during the cropping season (2021-22), were presented in fig. 1. The treatments included the sole and combined application of a market source of zinc (ZnSO₄), a cheap source of zinc (ZnO), compost, zinc solubilizing rhizobacteria, and microbial-assisted zinc-enriched compost. The experiment consists of ten treatments; T0: Control; T1: ZnSO₄; T2: ZnO; T3: Compost; T4: ZSB; T5: compost + 2% ZnO; T6: ZnSO₄ + ZSB; T7: ZnO + ZSB; T8: compost + ZSB; T9: compost + 2% ZnO + ZSB. A randomized complete block design (RCBD) was used with three replications per treatment at each site to ensure statistical reliability.

Before sowing, composite soil samples were collected from the 0–15 cm depth across each site and analyzed for baseline fertility status. The soils at both locations were characterized as deficient arid soils with the average properties: pH (7.69 and 8.1), electrical conductivity (EC) (1.27 and 1.42 dS m⁻¹), organic matter (0.43 and 0.51%), total nitrogen (0.032 and 0.035%), available phosphorus (8.8 and 9.8 mg kg⁻¹), available potassium (73 and 85 mg kg⁻¹), DTPA-extracted zinc (0.63 and 0.71 mg kg⁻¹) and DTPA extracted iron (0.56 and 0.51 mg kg⁻¹) in trial I (Bahawalpur) and II (Bahawalnagar), respectively. Based on soil test results and crop nutrient requirements, recommended doses of fertilizers (e.g., 120 kg of N ha⁻¹, 90 kg of P ha⁻¹, and 60 kg of K ha⁻¹) were applied. Phosphorus and potassium were incorporated at the time of seedbed preparation using diammonium phosphate (DAP) and sulfate of potash (SOP), respectively, while nitrogen was split-applied: one-third as a basal dose using urea at sowing, and the remaining two-thirds in two equal splits at tiller formation and early flowering stages to minimize losses and enhance nutrient use efficiency. A basal dose of ZnSO₄ (33% Zn) and microbial-assisted zinc-enriched



172 compost were applied @ 12 kg ha⁻¹ and 250 kg ha⁻¹, respectively. Chemical fertilizer application
173 was reduced by 5% in the treatment where Microbial-assisted zinc-enriched compost was used.
174 All agronomic practices were kept uniform across treatments.

175 **2.3. Soil and plant samples collection and analysis**

176 The growth parameters, such as plant height, number of tillers and biomass, were recorded at
177 harvest. Yield attributes, including 1000-grain weight, grain yield per hectare, and harvest index,
178 were measured post-harvest. For nutrient analysis, plant tissues (e.g., grain) were collected at
179 harvest, dried, ground, and analyzed for macro- and micronutrient content using standard
180 procedures. Rhizosphere soil samples were carefully collected at harvest by gently shaking the soil
181 adhered to the root zone, then analyzed for microbial population (CFU g⁻¹ soil), total organic
182 carbon, available NPK, DTPA extracted Fe and Zn, and microbial biomass carbon and nitrogen.

183 **2.4. Plant analysis**

184 The determination of nitrogen (N), phosphorus (P), and potassium (K) in grain samples was
185 conducted using a wet digestion method involving sulfuric acid (H₂SO₄) and hydrogen peroxide
186 (H₂O₂). Oven-dried grain samples (0.5 g) were placed in digestion tubes and initially treated with
187 concentrated H₂SO₄ (6 mL) to break down organic matter. Hydrogen peroxide (2 mL) was added
188 dropwise to enhance oxidation and complete the digestion. The mixture was heated until the
189 solution became clear, indicating complete digestion. The digested samples were allowed to cool,
190 diluted with distilled water, and filtered. The clear filtrate was analyzed for total N using the
191 Kjeldahl method, P content was determined colorimetrically using the molybdenum blue method,
192 and K was quantified using a flame photometer. This method ensures efficient mineralization of
193 organic components and accurate estimation of macronutrients essential for grain quality
194 assessment (Ryan et al., 2001).

195 The analysis of zinc (Zn) and iron (Fe) concentrations in grain samples was conducted using the
196 diacid digestion method, employing a mixture of nitric acid (HNO₃) and perchloric acid (HClO₄).
197 Oven-dried and ground grain sample (0.5 g) was accurately weighed into a digestion flask. A 10
198 mL aliquot of concentrated HNO₃ was added, and the mixture was allowed to pre-digest overnight
199 at room temperature to initiate breakdown of organic matter. The following day, 4 mL of
200 concentrated HClO₄ was added, and the sample was subjected to controlled heating on a digestion
201 block, gradually increasing the temperature up to 350 °C until white fumes appeared, indicating
202 the completion of organic matrix oxidation. The digestion was continued until a clear solution was



203 obtained. After cooling, the digested sample was diluted with deionized water and filtered. The
204 final volume was made up to a known volume (50 mL) with deionized water. The concentrations
205 of Zn and Fe in the digest were then determined using an atomic absorption spectrophotometer
206 (AAS). All glassware was acid-washed to minimize contamination, and analytical-grade reagents
207 were used throughout the procedure. The instrument was standardized by using respective standard
208 solutions of Fe and Zn (Antreich, 2012).

209 **2.5. Soil chemical analysis**

210 For the soil biochemical analysis, soil samples were systematically collected from designated study
211 sites using a clean stainless-steel auger to minimize contamination. Samples were air-dried, sieved
212 through a 2 mm mesh to remove debris and stones, and stored in polythene bags for laboratory
213 analysis. Chemical properties were assessed by measuring soil organic matter content through the
214 Walkley-Black method (Jha et al., 2014). Available nutrients, including nitrogen (N), phosphorus
215 (P), and potassium (K), were quantified using Kjeldahl digestion for N (Estefan et al., 2013), Olsen
216 methods for P (Olsen, 1954), and flame photometry for K after ammonium acetate extraction
217 (Shuman and Duncan, 1990).

218 **2.6. Soil biochemical analysis**

219 For the analysis of soil microbial biomass carbon (MBC) and microbial biomass nitrogen (MBN),
220 soil samples were collected from the study area and air-dried before sieving through a 2 mm mesh.
221 The fumigation-extraction method was employed, where one set of samples was fumigated with
222 chloroform vapor for 24 hours to lyse microbial cells, while the other set served as non-fumigated
223 controls (Brookes et al., 1985). Following fumigation, both sets were extracted with 0.5 M K₂SO₄
224 solution, and the extracts were analyzed for dissolved organic carbon and nitrogen using a total
225 organic carbon analysis and the Kjeldahl method, respectively. Microbial biomass carbon and
226 nitrogen were calculated by subtracting the values of non-fumigated samples from fumigated ones.
227 All analyses were conducted in triplicate to ensure accuracy and reproducibility, and standard
228 quality control procedures were followed throughout the process.

229 For the analysis of soil ammonium (NH₄⁺) and nitrate (NO₃⁻) nitrogen, soil extracts were prepared
230 by shaking 10 g of soil with 100 mL of 2 M potassium chloride (KCl) solution for 30 minutes to
231 displace exchangeable ammonium and nitrate ions. The suspension was then filtered through a
232 Whatman No. 42 filter paper. Ammonium nitrogen concentration in the filtrate was determined
233 colorimetrically using the indophenol blue method, while nitrate nitrogen was measured by the



phenol disulfonic acid method and readings were taken on UV spectrophotometer at 220 nm (Kachurina et al., 2000). All analyses were performed in triplicate to ensure accuracy, and results were expressed in mg kg^{-1} of soil. Quality control included calibration curves using standard NH_4^+ and NO_3^- solutions, reagent blanks, and periodic checks with known reference materials.

2.7. Economic analysis

The economic analysis was estimated, and subsequently, the costs and returns of various items used in this study for wheat cultivation, as normally practiced by farmers (control), and the application of a market source of zinc and microbial-assisted zinc-enriched compost (Khan et al., 2012) were apportioned. Gross margin was calculated by the following formula to make comparisons.

$$\text{Gross Margin (GM)} = \text{Total Revenue (TR)} - \text{Total Variable Cost (TVC)}$$

2.8. Statistical Analysis

The statistical analysis in this study was conducted using a combination of multivariate techniques. Analysis of Variance (ANOVA) was employed to test for significant differences between group means. Principal Component Analysis (PCA) was done to reduce dimensionality and identify the gradients of variation in the dataset, facilitating visualization of patterns and clustering. Pearson correlation coefficients were calculated to quantify the strength and direction of linear associations between pairs of continuous variables. All analyses were performed using OriginPro 2021b and Statistix 8.1 software, with significance levels set at $\alpha = 0.05$.

3. Results

The results from both field trials consistently demonstrated the positive role of microbial-assisted zinc-enriched compost in improving wheat growth, nutrient accumulation, and soil health. Treatments integrating compost, zinc sources, and zinc-solubilizing bacteria (ZSB) performed better than sole or dual applications, highlighting the synergistic effect of organic matter and microbial activity. The results demonstrated the influence of microbial-assisted zinc-enriched compost on plant performance, grain nutritional quality, and soil fertility parameters.

3.1. Comparative effect of bio-organic and organic fertilizers on soil biochemical properties

In both Trial I (Bahawalpur) and Trial II (Bahawalnagar), the application of microbial-assisted zinc-enriched compost significantly improved soil fertility parameters compared to the control and sole application of inorganic fertilizer (Table 1). The highest enhancement in organic matter (OM), total organic carbon (TOC), and macro- and micronutrient contents (N, P, K, Fe, and Zn) was



observed in the treatment combining compost with 2% ZnO and zinc-solubilizing bacteria (ZSB). In Trial I, the treatment Compost + 2% ZnO + ZSB (T9) resulted in a remarkable increase in soil OM and TOC by 38.6% over the control. Nitrogen (N) content increased by 17.1%, phosphorus (P) by 23.2%, and potassium (K) by 17.9%. Micronutrient concentrations were also significantly enhanced, with iron (Fe) increasing by 22.6% and zinc (Zn) by an impressive 22.0% relative to the control. Similarly, in Trial II, the same treatment (Compost + 2% ZnO + ZSB) demonstrated the most effective results, improving OM by 38.3%, TOC by 31.8%, N by 20.4%, P by 20.2%, and K by 22.0%. Increases in Fe and Zn were recorded at 25.8% and 20.6%, respectively, compared to untreated soil.

In Trial I, the treatment (compost + 2% ZnO + ZSB) led to a maximum CFU count (34.1×10^6 CFU g⁻¹ soil), showing an increase of 50.3% over the control. Microbial biomass carbon (MBC) and microbial biomass nitrogen (MBN) were significantly elevated, with increases of 21% and 26%, respectively, compared to the untreated control. Additionally, ammonium-N and nitrate-N concentrations increased by 25% and 23%, respectively. Similarly, in Trial II, the Compost + 2% ZnO + ZSB treatment again resulted in the highest values, with CFU increasing by 39%, MBC by 27%, MBN by 32%, ammonium-N by 28.2%, and nitrate-N by 27.7% relative to the control.

Among individual and dual component treatments, Compost + ZSB and compost + 2% ZnO showed superior performance compared to sole applications. Compost + ZSB increased MBC by 16% in Trial I and 22% in Trial II, whereas compost + 2% ZnO enhanced nitrate-N by 12% in both trials. The integration of compost, zinc (in the form of ZnO), and zinc-solubilizing bacteria (ZSB) synergistically improved soil biochemical properties, underscoring the potential of microbial-assisted zinc-enriched compost as a sustainable soil amendment in enhancing soil fertility and microbial function.

3.2. Multivariate analysis of the studied parameters compares the effectiveness of bio-organic and inorganic fertilizers to improve soil health

The significance of the results was further confirmed through multivariate analysis. Principal component analysis (PCA) revealed clear separation among treatments, indicating distinct influences of different zinc and organic amendments on soil biochemical properties (Fig. 2). In trial I, the two principal components (PC1 and PC2) accounted for 95.58% of the total variance, with PC1 alone contributing 92.55%. Treatments involving combinations of compost and zinc sources, especially Compost + 2% ZnO + ZSB and Compost + ZSB, were closely associated with



296 higher scores on PC1, primarily driven by elevated levels of soil organic matter (OM), total organic
 297 carbon (TOC), nitrogen (N), and available phosphorus (P). These treatments clustered in the
 298 positive quadrant of the biplot, indicating strong synergistic effects on soil fertility indicators. On
 299 the other hand, the control and mineral ZnSO₄-alone treatment showed negative associations with
 300 PC1 and PC2, reflecting minimal improvements in soil nutrient status. Zinc oxide in combination
 301 with ZSB (ZnO + ZSB) showed a moderate effect, clustering near the origin, suggesting limited
 302 but positive contributions to zinc availability and other nutrients.

303 In Trial II, PCA highlighted similar patterns with the two principal components explaining 96.12%
 304 of the total variance, where PC1 accounted for 91.52% and PC2 for 4.60%. Compost-based
 305 treatments again formed a distinct cluster, particularly Compost + 2% ZnO + ZSB and Compost +
 306 ZSB, indicating their significant influence on soil OM, TOC, and micronutrient (Fe and Zn)
 307 concentrations. Control and ZnSO₄ alone consistently showed the lowest contributions across both
 308 principal components, reinforcing their limited efficacy in improving soil fertility. Overall, the
 309 PCA confirmed that treatments integrating organic matter (compost) and biological agents (ZSB)
 310 with zinc sources (ZnO) led to more comprehensive improvements in soil nutrient status compared
 311 to inorganic treatments.

312 **3.3. Effectiveness of bio-organic fertilizer to improve plant growth and yield**

313 In both Trial I (Bahawalpur) and Trial II (Bahawalnagar), the application of microbial-assisted
 314 zinc-enriched compost demonstrated a significant improvement in wheat growth, particularly in
 315 terms of plant height, shoot dry biomass, 1000-grain weight and grain yield (Fig. 3). Among the
 316 treatments, the combined application of Compost + 2% ZnO + ZSB (T9) yielded the most
 317 pronounced effects. In Trial I, the Compost + 2% ZnO + ZSB treatment increased plant height by
 318 16.7% and shoot dry biomass by 17.6% compared to the control. Similarly, in Trial II, this
 319 treatment resulted in a 19.4% increase in plant height and a 19.8% increase in shoot dry biomass
 320 over the control. The next most effective treatments were Compost + ZSB (T8) and Compost +
 321 ZnO (T5). Compost + ZSB enhanced plant height by 12.5% and 12.1% and shoot biomass by
 322 15.1% and 12.8% in Trials I and II, respectively. Compost + ZnO led to increases of 9.7% and
 323 13.6% in plant height, and 4.7% and 10.5% in biomass for Trials I and II, respectively.

324 In Trial I, the highest 1000-grain weight was recorded in the compost + 2% ZnO + ZSB (T9)
 325 treatment, showing an 8.7% increase over the control. This was followed by compost + ZSB
 326 (8.2%) and ZnO + ZSB (7.7%). In terms of grain yield per hectare, the compost + 2% ZnO + ZSB



327 treatment led to a significant 9.5% increase compared to the control. Other notable increases in
328 yield included compost + ZSB (7.3%), ZnSO₄ + ZSB (6.3%), and ZnO + ZSB (7.7%). Trial II
329 confirmed the trends observed in Trial I. The compost + 2% ZnO + ZSB treatment again yielded
330 the highest 1000-grain weight, with a 9.6% increase over the control. Compost + ZSB and ZnSO₄
331 + ZSB treatments improved 1000-grain weight by 7.3% and 5.9%, respectively. Grain yield was
332 similarly enhanced, with the compost + 2% ZnO + ZSB treatment resulting in a 8.5% increase
333 over the control. The compost + ZSB (6.7%), ZnSO₄ + ZSB (5.0%), and ZnO + ZSB (6.6%)
334 treatments also showed considerable improvements.

335 In contrast, sole applications of ZnO, ZnSO₄, or compost alone had moderate effects on both
336 parameters, with increases ranging between 1%–3% depending on the trial and variable measured.
337 The ZSB treatment alone showed a limited but positive effect, suggesting that microbial inoculants
338 are more effective when combined with zinc sources and organic amendments.

339 **3.4. Comparative improvement in grain quality by the application of bioorganic and** 340 **inorganic fertilizers**

341 Based on the results obtained from both Trial I (Bahawalpur) and Trial II (Bahawalnagar), the
342 application of microbial-assisted zinc-enriched compost significantly improved the concentration
343 of macronutrients (N, P, K) and micronutrients (Fe and Zn) in wheat grains compared to the control
344 and individual application of inorganic fertilizers (Fig. 4). In Trial I, the combined treatment of
345 Compost + 2% ZnO + ZSB (T9) demonstrated the highest nutrient enhancement across all
346 measured parameters. This treatment led to an increase of 16.5% in nitrogen (N), 13.9% in
347 phosphorus (P), and 11.3% in potassium (K) contents compared to the control. In terms of
348 micronutrients, grain iron (Fe) content (30.4 mg kg⁻¹) and zinc (Zn) content (38.5 mg kg⁻¹) were
349 recorded, which were increased by 27.6% and 37.3%, respectively, relative to the control.
350 Similarly, in Trial II, the Compost + 2% ZnO + ZSB treatment consistently outperformed all other
351 treatments. Compared to the control, it increased grain nitrogen by 15.9%, phosphorus by 15.0%,
352 potassium by 13.7%, iron by 27.9%, and zinc by 39.4%. The treatments ZnO + ZSB and ZnSO₄ +
353 ZSB also showed considerable improvements over the sole applications of ZnO or ZnSO₄,
354 suggesting the synergistic effect of zinc-solubilizing bacteria (ZSB) in mobilizing native and
355 supplemented zinc sources.

356 The Compost + ZSB treatment also enhanced nutrient accumulation in grains, albeit to a lesser
357 extent than the Compost + 2% ZnO + ZSB. Sole compost and ZSB treatments moderately



increased nutrient content compared to the control, while the sole application of ZnSO_4 and ZnO resulted in lower improvements than when these sources were bioactivated. Overall, the combined application of compost, ZnO , and ZSB (Compost + 2% ZnO + ZSB) was the most effective strategy for promoting the biofortification of wheat grains with essential nutrients in both trials.

3.5. Correlation analysis demonstrates the interactive effect of improved soil fertility parameters on yield

Correlation analysis revealed significant positive relationships among soil biochemical parameters, bacterial population in term of colony forming units (CFU g^{-1} soil), microbial biomass carbon (MBC), and microbial biomass nitrogen (MBN) and key soil nutrients (N, P, K, Fe, and Zn) with crop yield (Fig. 5). Specifically, CFU exhibited a strong positive correlation with MBC ($r = 0.82$, $p < 0.01$) and MBN ($r = 0.76$, $p < 0.01$), indicating that enhanced microbial abundance supports higher microbial biomass. MBC and MBN were significantly correlated with available nitrogen ($r = 0.79$ and 0.74 , respectively; $p < 0.01$), suggesting active microbial involvement in nutrient mineralization and cycling. Available phosphorus and potassium were also positively correlated with microbial indicators (MBC–P: $r = 0.68$; MBN–K: $r = 0.63$; $p < 0.05$), underscoring the role of microbial processes in improving nutrient availability. Moreover, micronutrients such as Fe and Zn showed moderate but significant correlations with MBC ($r = 0.59$ and 0.57 , respectively; $p < 0.05$), suggesting microbial-mediated enhancement of micronutrient solubility. Importantly, crop yield demonstrated strong positive correlations with MBC ($r = 0.85$), MBN ($r = 0.80$), and CFU ($r = 0.78$), as well as with macronutrients N ($r = 0.83$), P ($r = 0.77$), and K ($r = 0.75$). These findings collectively indicate that microbial activity and biomass are closely linked to nutrient availability and are reliable predictors of soil fertility and crop productivity.

3.6. Comparison of bio-organic and inorganic (ZnSO_4) fertilizers for cost of production and net returns for one hectare of wheat

In the study area, the total cost of cultivating one hectare of control (untreated), ZnSO_4 (market source of zinc) and microbial-assisted zinc-enriched compost (bio-organic fertilizer) of wheat comprised several components, including land preparation, sowing, seed, fertilizer application, plant protection (weedicide and pesticide), manual labor, irrigation, and harvesting and threshing operations (Table 2). The production cost per hectare was calculated as Rs. 230,371 for control wheat, Rs. 235,353 for ZnSO_4 -treated wheat, and Rs. 239,438 for the microbial-assisted zinc-enriched compost-treated wheat. The treated wheat incurred higher costs than the control



(untreated) wheat, mainly due to the application of zinc fertilizers. For wheat cultivation, the total production cost also accounted for opportunity costs, calculated as a 12% markup from land preparation to fertilizer application. These amounted to Rs. 14,529 for control, Rs. 15,063 for ZnSO_4 , and Rs. 15,446 for the microbial-assisted zinc-enriched compost. Manual labor costs were Rs. 2965, Rs. 3,033, and Rs. 3,478, respectively. As detailed in Table 3, the total cost (TC) was broken down into Total Variable Cost (TVC) and Total Fixed Cost (TFC). The TVC included expenses related to land preparation, seed, fertilizers, labor, markup, and harvesting and threshing. The estimated TVC per hectare was Rs. 167,484 for control, Rs. 172,534 for ZnSO_4 , and Rs. 176,551 for the microbial-assisted zinc-enriched compost. TFC, on the other hand, included fixed expenses such as land rent and *abiana* (government water charges), which collectively amounted to Rs. 62,887 per hectare. Land rent was treated as an opportunity cost, as most farmers in the region owned their land. The total cost of production (TC) was derived by summing the TVC and TFC.

The gross margin, net return, and revenue-to-cost (R/C) ratio were calculated for the cultivation of one hectare of wheat treated as control, ZnSO_4 , and microbial-assisted zinc-enriched compost (Table 3). Based on Equation 1, the gross margins were estimated at Rs. 111,997 for control, Rs. 112,304 for ZnSO_4 and Rs. 130,578 for the microbial-assisted zinc-enriched compost. Net returns were determined by deducting the total costs (TC) from total revenue (TR). When land rent (LR) was included, the net returns amounted to Rs. 49,110 for control, Rs. 49,417 for ZnSO_4 , and Rs. 67,691 for the microbial-assisted zinc-enriched compost. Excluding land rent, the net returns increased to Rs. 110,885, Rs. 111,192, and Rs. 129,466, respectively. Likewise, the revenue-to-cost ratios, including land rent, were 1.21 for control and ZnSO_4 , and 1.28 for the microbial-assisted zinc-enriched compost. Without considering land rent, these ratios improved to 1.67, 1.65, and 1.74, respectively. Overall, the economic analysis revealed that the application of the market source of zinc (ZnSO_4) has a negligible impact on farmers' net income. Whereas the application of microbial-assisted zinc-enriched compost showed a significant increase in net income.

4. Discussion

The results of both trials distinctly highlight the agronomic and soil-enhancing potential of bioorganic fertilizer, particularly when integrated with zinc oxide (ZnO) and zinc-solubilizing bacteria (ZSB). The treatment combination of Compost + 2% ZnO + ZSB consistently outperformed all other treatments across all growth, yield, nutrient uptake, and soil fertility



420 parameters. These findings underscore the efficacy of an integrated nutrient management approach
421 combining organic, inorganic, and biological amendments in enhancing wheat productivity and
422 soil health.

423 Soil chemical properties were significantly influenced by the bio-organic fertilizer application.
424 The observed increase in zinc (Zn) availability can be attributed to two complementary
425 mechanisms: the gradual and sustainable release of Zn from ZnO particles and the active microbial
426 solubilization of Zn compounds facilitated by Zn-solubilizing bacteria (ZSB) (Huang et al., 2022).
427 These processes ensure a continuous supply of bioavailable Zn, reducing the risk of nutrient
428 fixation in soil. In addition, the enhanced availability of nitrogen (N), phosphorus (P), and
429 potassium (K) is primarily linked to accelerated microbial mineralization of organic matter,
430 enzymatic hydrolysis of complex organic compounds, and the decomposition of compost materials
431 (Reimer et al., 2023). Collectively, these mechanisms improve soil nutrient cycling and ensure
432 synchronized nutrient release that matches plant demand.

433 Biological indicators of soil fertility, including colony-forming units (CFU), microbial biomass
434 carbon (MBC), and microbial biomass nitrogen (MBN), were improved by the Compost + 2%
435 ZnO + ZSB treatment. This demonstrates not only the proliferative effect of compost as a microbial
436 substrate but also the synergistic interactions between microbial inoculants and native soil
437 microbiota, which enhance microbial colonization and community stability (Dincă et al., 2022).
438 ZSB play a critical mechanistic role by releasing organic acids (Sethi et al., 2025), siderophores
439 (Zhu et al., 2025), and hydrolytic enzymes that mobilize nutrients (Mujumdar et al., 2024),
440 improve rhizosphere activity, and stimulate beneficial plant–microbe interactions (Jalal et al.,
441 2024). Furthermore, the observed increases in nitrate-N and ammonium-N concentrations indicate
442 enhanced nitrogen transformation pathways, including ammonification, nitrification, and
443 mineralization processes, actively mediated by microbial communities (Duan et al., 2023). These
444 transformations improve nitrogen turnover and ensure steady nutrient availability to crops, thereby
445 supporting higher productivity and nutrient use efficiency.

446 Plant growth, measured in terms of plant height and shoot dry biomass, was significantly improved
447 by the Compost + 2% ZnO + ZSB treatment. These improvements are attributable to enhanced
448 nutrient availability, particularly zinc and nitrogen, resulting from the synergistic effect of the
449 organic matrix, micronutrient enrichment, and microbial solubilization. Zinc plays a critical role
450 in auxin synthesis and enzyme activation (Wang et al., 2023), and its increased availability in a



451 bioavailable form likely stimulated vegetative growth. Moreover, the compost component
452 improved soil physical structure and provided a steady nutrient supply (Kelbesa, 2021), while ZSB
453 enhanced micronutrient solubility through the production of organic acids and siderophores
454 (Asghar et al., 2024).

455 The sole application of ZnSO_4 or ZnO resulted in comparatively modest gains, indicating the
456 limitations of mineral zinc sources in calcareous or alkaline soils where Zn availability is
457 inherently low due to fixation (Gupta et al., 2024). In contrast, microbial inoculation with ZSB in
458 conjunction with ZnO improved plant growth, suggesting the pivotal role of ZSB in solubilizing
459 ZnO particles (Sethi et al., 2025). Compost + ZSB treatment also performed well, reaffirming the
460 benefits of incorporating biological inputs into organic nutrient management systems.

461 In terms of yield, the Compost + 2% ZnO + ZSB treatment consistently showed the highest grain
462 yield. Enhanced grain filling and development can be attributed to the continuous availability of
463 zinc, phosphorus, and nitrogen, facilitated by microbial mineralization and compost-mediated
464 nutrient retention (Campana et al., 2025). Furthermore, the observed increase in 1000-grain weight
465 in treatments involving ZSB aligns with previous research, which reported that ZSB not only
466 solubilizes native zinc but also stimulates root growth and enhances nutrient uptake efficiency
467 (Singh et al., 2024).

468 Nutrient accumulation in grains, particularly of nitrogen (N), phosphorus (P), potassium (K), iron
469 (Fe), and zinc (Zn), was significantly higher in the Compost + 2% ZnO + ZSB treatment. The
470 improvement in nutrient uptake is primarily linked to enhanced microbial activity, which mobilizes
471 native and added micronutrients and the organic matter that prevents nutrient leaching and
472 enhances cation exchange capacity (Dhaliwal et al., 2024). These results are crucial in the context
473 of human nutrition, particularly in zinc-deficient regions, and support the promotion of agronomic
474 biofortification strategies.

475 The findings of the study validate the hypothesis that integrating organic matter (compost),
476 micronutrient supplementation (ZnO), and microbial inoculants (ZSB) offers a sustainable,
477 efficient, and environmentally friendly strategy to improve crop growth, yield, nutrient density,
478 and soil health. This integrated approach is promising in zinc-deficient soils and can serve as a
479 cornerstone of sustainable agricultural and biofortification efforts in developing regions. Future
480 studies should explore the scalability of the strategy under different agroecological zones and
481 cropping systems, as well as its long-term impacts on soil microbial ecology and plant health.



482 The economic analysis of wheat cultivation under different zinc fertilization treatments
483 demonstrated that although input costs were higher for treated plots, especially for microbial-
484 assisted zinc-enriched compost, the overall profitability was enhanced, particularly under bio-
485 organic fertilization. This aligns with research, which reported that the addition of external nutrient
486 sources in wheat production generally increases variable costs, especially when micronutrient
487 fortification is involved (Ali and Tsou, 1997). Despite the increased costs, the economic benefits
488 in terms of gross margins and net returns were significantly higher for the microbial-assisted zinc-
489 compost treatment. This improvement may be attributed to enhanced nutrient use efficiency and
490 improved plant growth parameters due to the synergistic effects of compost and beneficial
491 microbes. It is evident from the literature that microbial inoculants can promote nutrient
492 solubilization and uptake, translating into higher yields (Sammauria et al., 2020) and better
493 financial returns. The higher net income in the microbial-assisted zinc-enriched compost treatment
494 relative to the control suggests that bio-organic-assisted biofortification not only improves soil
495 health and crop yield but also enhances farm-level economic resilience. This strategy offers a
496 sustainable solution for nutrient-deficient soils, supporting the transition towards low-input, high-
497 efficiency farming systems (Akbar et al., 2020).

498 Although initial input costs were higher with microbial-assisted zinc-enriched compost, the
499 significant gains in yield and net return justify its adoption. The data strongly support integrating
500 microbial-assisted composts into conventional fertilization regimes to improve agronomic
501 performance and economic returns under resource-constrained conditions. The present study was
502 conducted under specific soil and climatic conditions and over limited cropping seasons, which
503 may restrict the generalization of results across diverse agroecological zones. Long-term effects
504 on soil microbial diversity, nutrient cycling, and micronutrient buildup were not assessed, and the
505 findings were restricted to wheat without testing in other cropping systems. Future research should
506 therefore focus on multi-location and multi-season trials, long-term soil health monitoring, and
507 extending the approach to different crops. Additionally, developing cost-effective microbial
508 inoculant formulations and delivery mechanisms will be crucial to enhance adoption, particularly
509 by resource-constrained farmers.

510 **5. Conclusion**

511 The integrated application of microbial-assisted zinc-enriched compost (Compost + 2% ZnO +
512 ZSB) enhances wheat growth, yield, grain nutrient biofortification, and soil fertility compared to



513 sole amendments. This combined treatment consistently outperformed others across both trials,
514 markedly increasing plant height, biomass, grain weight, and yield, while significantly elevating
515 macronutrient (N, P, K) and micronutrient (Fe, Zn) concentrations in grains. Soil chemical and
516 biological parameters also showed pronounced improvements, with enhanced organic matter,
517 microbial biomass, and nutrient availability, underscoring the synergistic effects of organic,
518 microbial, and zinc amendments. These findings highlight the efficacy of integrated nutrient
519 management strategies involving microbial-assisted zinc-enriched compost for sustainable wheat
520 production and soil health restoration, offering promising avenues for addressing micronutrient
521 deficiencies in cereal crops.

522 **Data Availability Statement**

523 The authors confirm that the data, figures, and tables included in this manuscript are original and
524 have not been previously published. The primary data are accessible and can be provided upon
525 request. Relevant data is submitted on the NCBI website, and accession numbers MN696212,
526 MN696213, MN696214, MN696215 and MN696216 are included in the text.

527 **Author Contributions**

528 All authors contributed to research and manuscript preparation. **Conceptualization:** Azhar
529 Hussain, Zafar Iqbal. **Methodology:** Azhar Hussain, Maqshoof Ahmad, Asma Sabir.
530 **Investigation:** Azhar Hussain, Asma Sabir, Maqshoof Ahmad. **Data Curation:** Azhar Hussain,
531 Maqshoof Ahmad, Eman Alhomaidi. **Formal Analysis:** Azhar Hussain, Zafar Iqbal, Asma Sabir.
532 **Writing – Original Draft:** Azhar Hussain, Zafar Iqbal. **Writing – Review & Editing:** Asma
533 Sabir, Eman Alhomaidi, Maqshoof Ahmad. **Supervision:** Maqshoof Ahmad. **Funding**
534 **Acquisition:** Eman Alhomaidi, Maqshoof Ahmad. All authors reviewed and commented on the
535 previous version of the manuscript and approved the final manuscript for publication.

536 **Competing interest**

537 The authors declared no potential conflict/competing interest relevant to this work.

538

539 **Acknowledgments**



540 This work was supported by Princess Nourah bint Abdulrahman University Researchers
 541 Supporting Project number (PNURSP2025R317), Princess Nourah bint Abdulrahman University,
 542 Riyadh, Saudi Arabia. The authors are grateful to the Department of Soil Science, The Islamia
 543 University of Bahawalpur, Pakistan, and the National Research Program for Universities (NRPU)
 544 project number 6292/Punjab/NRPU/R&D/HEC/2016 for providing research facilities.

545 **Funding**

546 This work was Supported by Princess Nourah bint Abdulrahman University Researchers
 547 Supporting Project number (PNURSP2025R317), Princess Nourah bint Abdulrahman University,
 548 Riyadh, Saudi Arabia.

549

550 **References**

- 551 Akbar, F., Rahman, A. U., and Rehman, A.: Genetic Engineering of Rice to Survive in Nutrient-Deficient Soil,
 552 Rice Research for Quality Improvement: Genomics and Genetic Engineering: Volume 1: Breeding
 553 Techniques and Abiotic Stress Tolerance, 437-464, 2020.
- 554 Ali, M. and Tsou, S. C.: Combating micronutrient deficiencies through vegetables—a neglected food
 555 frontier in Asia, *Food policy*, 22, 17-38, 1997.
- 556 Antreich, S.: Heavy metal stress in plants-a closer look, Protocol of the project practicum “Heavy metal
 557 stress in plants, University of Vienna, 1-13, 2012.
- 558 Asghar, M. U., Hussain, A., Anwar, H., Dar, A., Ahmad, H. T., Nazir, Q., Tariq, N., and Jamshaid, M. U.:
 559 Investigating the Efficacy of Dry Region Zinc Solubilizing Bacteria for Growth Promotion of Maize and
 560 Wheat under Axenic Conditions, *Plant Health*, 3, 01-18, 2024.
- 561 Brookes, P. C., Landman, A., Pruden, G., and Jenkinson, D.: Chloroform fumigation and the release of soil
 562 nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil, *Soil biology and*
 563 *biochemistry*, 17, 837-842, 1985.
- 564 Campana, E., Ciriello, M., Lentini, M., Rouphael, Y., and De Pascale, S.: Sustainable Agriculture Through
 565 Compost Tea: Production, Application, and Impact on Horticultural Crops, *Horticulturae*, 11, 433, 2025.
- 566 Clagnan, E., Cucina, M., De Nisi, P., Dell’Orto, M., D’Imporzano, G., Kron-Morelli, R., Llenas-Argelaguet, L.,
 567 and Adani, F.: Effects of the application of microbiologically activated bio-based fertilizers derived from
 568 manures on tomato plants and their rhizospheric communities, *Scientific Reports*, 13, 22478, 2023.
- 569 Dhaliwal, S. S., Sharma, V., and Shukla, A. K.: Impact of micronutrients in mitigation of abiotic stresses in
 570 soils and plants—A progressive step toward crop security and nutritional quality, *Advances in Agronomy*,
 571 173, 1-78, 2022.
- 572 Dhaliwal, S. S., Dubey, S. K., Kumar, D., Toor, A. S., Walia, S. S., Randhawa, M. K., Kaur, G., Brar, S. K.,
 573 Khambalkar, P. A., and Shivey, Y. S.: Enhanced Organic Carbon Triggers Transformations of
 574 Macronutrients, Micronutrients, and Secondary Plant Nutrients and Their Dynamics in the Soil under
 575 Different Cropping Systems-A Review, *Journal of Soil Science and Plant Nutrition*, 24, 5272-5292, 2024.
- 576 Dincă, L. C., Grenni, P., Onet, C., and Onet, A.: Fertilization and soil microbial community: a review, *Applied*
 577 *Sciences*, 12, 1198, 2022.



- 578 Duan, F., Peng, P., Yang, K., Shu, Y., and Wang, J.: Straw return of maize and soybean enhances soil
579 biological nitrogen fixation by altering the N-cycling microbial community, *Applied Soil Ecology*, 192,
580 105094, 2023.
- 581 Estefan, G., Sommer, R., and Ryan, J.: Methods of soil, plant, and water analysis, A manual for the West
582 Asia and North Africa region, 3, 65-119, 2013.
- 583 Gupta, N., Gupta, A., Sharma, V., Kaur, T., Rajan, R., Mishra, D., Singh, J., and Pandey, K.: Biofortification
584 of Legumes: Enhancing Protein and Micronutrient Content, in: *Harnessing Crop Biofortification for*
585 *Sustainable Agriculture*, Springer, 225-253, 2024.
- 586 Hossain, A., Skalicky, M., Brestic, M., Maitra, S., Ashraful Alam, M., Syed, M. A., Hossain, J., Sarkar, S., Saha,
587 S., and Bhadra, P.: Consequences and mitigation strategies of abiotic stresses in wheat (*Triticum aestivum*
588 L.) under the changing climate, *Agronomy*, 11, 241, 2021.
- 589 Huang, H., Chen, J., Liu, S., and Pu, S.: Impact of ZnO nanoparticles on soil lead bioavailability and microbial
590 properties, *Science of The Total Environment*, 806, 150299, 2022.
- 591 Imran: Integration of organic, inorganic and bio fertilizer, improve maize-wheat system productivity and
592 soil nutrients, *Journal of Plant Nutrition*, 47, 2494-2510, 2024.
- 593 Jalal, A., Júnior, E. F., and Teixeira Filho, M. C. M.: Interaction of zinc mineral nutrition and plant growth-
594 promoting bacteria in tropical agricultural systems: a review, *Plants*, 13, 571, 2024.
- 595 Jha, P., Biswas, A., Lakaria, B. L., Saha, R., Singh, M., and Rao, A. S.: Predicting total organic carbon content
596 of soils from Walkley and Black analysis, *Communications in soil science and plant analysis*, 45, 713-725,
597 2014.
- 598 Kachurina, O., Zhang, H., Raun, W., and Krenzer, E.: Simultaneous determination of soil aluminum,
599 ammonium-and nitrate-nitrogen using 1 M potassium chloride extraction, *Communications in soil science*
600 *and plant analysis*, 31, 893-903, 2000.
- 601 Kelbesa, W. A.: Effect of compost in improving soil properties and its consequent effect on crop
602 production—A review, *Journal of Natural Sciences Research*, 12, 15-25, 2021.
- 603 Khan, M. T., Aleinikovienė, J., and Butkevicienė, L.-M.: Innovative organic fertilizers and cover crops:
604 Perspectives for sustainable agriculture in the Era of climate change and organic agriculture, *Agronomy*,
605 14, 2871, 2024.
- 606 Lal, R.: Soil Degradation Effects on Human Malnutrition and Under-Nutrition, *Medical Research Archives*,
607 12, 2024.
- 608 Maitra, S., Shankar, T., Hossain, A., Sairam, M., Sagar, L., Sahoo, U., Gaikwad, D. J., Pramanick, B., Mandal,
609 T. K., and Sarkar, S.: Climate-smart millets production in future for food and nutritional security, in:
610 *Adapting to climate change in agriculture-theories and practices: Approaches for adapting to climate*
611 *change in agriculture in India*, Springer, 11-41, 2024.
- 612 Manea, E. E. and Bumbac, C.: Sludge Composting—Is This a Viable Solution for Wastewater Sludge
613 Management?, *Water* (20734441), 16, 2024.
- 614 Mujumdar, S. S., Tikekar, S., and Tapkir, S.: Role of Bacteria in Plant Nutrient Mobilization: A Review, *Open*
615 *Access Journal of Microbiology and Biotechnology*, 9, 1-13, 2024.
- 616 Naeem, M., Iqbal, Z., Hussain, A., Jamil, M., Ahmad, H. T., Ismail, A. M., El-Mogy, M. M., El Ganainy, S. M.,
617 El-Beltagi, H. S., and Hadid, M. L.: Conversion of organic waste into bio-activated zinc-enriched compost
618 to improve the health of deficient soils and wheat yield, 2025.
- 619 Naseem, S., Hussain, A., Iqbal, Z., Mustafa, A., Mumtaz, M. Z., Manzoor, A., Jamil, M., and Ahmad, M.:
620 Exopolysaccharide and Siderophore Production Ability of Zn Solubilizing Bacterial Strains Improve Growth,
621 *Physiology and Antioxidant Status of Maize and Wheat*, *Polish Journal of Environmental Studies*, 31, 2022.
- 622 Olsen, S. R.: Estimation of available phosphorus in soils by extraction with sodium bicarbonate, 939, US
623 Department of Agriculture 1954.



- 624 Qian, S., Zhou, X., Fu, Y., Song, B., Yan, H., Chen, Z., Sun, Q., Ye, H., Qin, L., and Lai, C.: Biochar-compost as
625 a new option for soil improvement: Application in various problem soils, *Science of the total environment*,
626 870, 162024, 2023.
- 627 Reimer, M., Kopp, C., Hartmann, T., Zimmermann, H., Ruser, R., Schulz, R., Müller, T., and Möller, K.:
628 Assessing long term effects of compost fertilization on soil fertility and nitrogen mineralization rate,
629 *Journal of Plant Nutrition and Soil Science*, 186, 217-233, 2023.
- 630 Ryan, J., Estefan, G., and Rashid, A.: Soil and plant analysis laboratory manual, ICARDA2001.
- 631 Sammauria, R., Kumawat, S., Kumawat, P., Singh, J., and Jatwa, T. K.: Microbial inoculants: potential tool
632 for sustainability of agricultural production systems, *Archives of microbiology*, 202, 677-693, 2020.
- 633 Sethi, G., Behera, K. K., Sayyed, R., Adarsh, V., Sipra, B., Singh, L., Alamro, A. A., and Behera, M.: Enhancing
634 soil health and crop productivity: the role of zinc-solubilizing bacteria in sustainable agriculture, *Plant*
635 *Growth Regulation*, 1-17, 2025.
- 636 Sharma, A., Soni, R., and Soni, S. K.: From waste to wealth: exploring modern composting innovations and
637 compost valorization, *Journal of Material Cycles and Waste Management*, 26, 20-48, 2024.
- 638 Shuman, L. and Duncan, R.: Soil exchangeable cations and aluminum measured by ammonium chloride,
639 potassium chloride, and ammonium acetate, *Communications in Soil Science and Plant Analysis*, 21, 1217-
640 1228, 1990.
- 641 Singh, S., Chhabra, R., Sharma, A., and Bisht, A.: Harnessing the power of zinc-solubilizing bacteria: a
642 catalyst for a sustainable agrosystem, *Bacteria*, 3, 15-29, 2024.
- 643 Wang, Z., Fu, X., and Kuramae, E. E.: Insight into farming native microbiome by bioinoculant in soil-plant
644 system, *Microbiological Research*, 127776, 2024.
- 645 Wang, Z., Wang, Y., Du, Q., Yan, P., Yu, B., Li, W.-X., and Zou, C.-Q.: The auxin signaling pathway contributes
646 to phosphorus-mediated zinc homeostasis in maize, *BMC Plant Biology*, 23, 20, 2023.
- 647 Younas, N., Fatima, I., Ahmad, I. A., and Ayyaz, M. K.: Alleviation of zinc deficiency in plants and humans
648 through an effective technique; biofortification: A detailed review, *Acta Ecologica Sinica*, 43, 419-425,
649 2023.
- 650 Zhu, X.-X., Shi, L.-N., Shi, H.-M., and Ye, J.-R.: Characterization of the *Priestia megaterium* ZS-3 siderophore
651 and studies on its growth-promoting effects, *BMC microbiology*, 25, 133, 2025.

652



Table 1: Effect of Microbial-assisted zinc-enriched compost on soil biochemical properties

Treatment	Organic matter (%)		Total organic carbon (g kg ⁻¹)		Bacterial population (CFU × 10 ⁶)		Total nitrogen (%)	
	Trial I	Trial II	Trial I	Trial II	Trial I	Trial II	Trial I	Trial II
T0	0.47 ± 0.01 d	0.54 ± 0.01 e	5.8 ± 0.3 d	6.3 ± 0.3 c	22.7 ± 0.7 f	13.1 ± 0.3 f	0.029 ± 0.0004 f	0.033 ± 0.0004 f
T1	0.47 ± 0.00 d	0.56 ± 0.00 de	6.0 ± 0.4 cd	6.5 ± 0.3 bc	23.7 ± 0.5 ef	14.4 ± 0.7 def	0.029 ± 0.0002 ef	0.033 ± 0.0001 ef
T2	0.48 ± 0.00 d	0.55 ± 0.01 de	6.0 ± 0.2 cd	6.5 ± 0.2 c	25.4 ± 0.7 de	13.5 ± 0.3 ef	0.029 ± 0.0001 ef	0.033 ± 0.0004 ef
T3	0.61 ± 0.01 b	0.70 ± 0.01 b	7.0 ± 0.2 bc	7.5 ± 0.3 abc	29.1 ± 0.8 b	14.9 ± 0.4 cde	0.031 ± 0.0004 cd	0.036 ± 0.0004 cd
T4	0.48 ± 0.01 cd	0.57 ± 0.00 cd	6.2 ± 0.2 cd	7.0 ± 0.4 bc	26.8 ± 1.0 cd	14.0 ± 0.2 def	0.029 ± 0.0003 ef	0.034 ± 0.0002 e
T5	0.62 ± 0.01 b	0.71 ± 0.00 b	6.9 ± 0.5 bc	7.3 ± 0.6 abc	29.9 ± 0.7 b	16.3 ± 0.7 bc	0.032 ± 0.0006 bc	0.036 ± 0.0002 bc
T6	0.49 ± 0.00 cd	0.57 ± 0.00 cd	6.4 ± 0.2 cd	6.9 ± 0.5 bc	28.0 ± 0.3 bc	15.4 ± 0.4 cd	0.030 ± 0.0002 de	0.034 ± 0.0001 e
T7	0.50 ± 0.01 c	0.58 ± 0.00 c	6.4 ± 0.2 bcd	6.9 ± 0.5 bc	26.9 ± 0.2 cd	16.2 ± 0.7 bc	0.031 ± 0.0003 cd	0.035 ± 0.0002 d
T8	0.63 ± 0.01 ab	0.72 ± 0.01 b	7.5 ± 0.7 ab	7.9 ± 0.4 ab	29.5 ± 0.6 b	17.1 ± 0.2 ab	0.032 ± 0.0006 ab	0.037 ± 0.0003 bc
T9	0.65 ± 0.01 a	0.75 ± 0.01 a	8.0 ± 0.6 a	8.3 ± 0.7 a	34.1 ± 0.9 a	18.2 ± 0.3 a	0.034 ± 0.0006 a	0.039 ± 0.0007 a
LSD ($p \leq 0.05$)	0.0249 ****	0.0207 ****	1.0147 **	1.3130 ns	2.0018 ****	1.4673 ****	1.244 ****	1.034 ****
Treatment	Ammonium nitrogen (mg kg ⁻¹)		Nitrate nitrogen (mg kg ⁻¹)		Microbial biomass nitrogen (mg kg ⁻¹)		Microbial biomass carbon (mg kg ⁻¹)	
	Trial I	Trial II	Trial I	Trial II	Trial I	Trial II	Trial I	Trial II
T0	9.9 ± 0.2 e	10.2 ± 0.2 f	13.6 ± 0.5 c	14.7 ± 0.5 d	133 ± 4.6 c	134 ± 2.0 f	651 ± 3.8 e	654 ± 3.8 e
T1	10.2 ± 0.2 de	10.4 ± 0.2 ef	14.0 ± 0.9 bc	15.4 ± 0.3 cd	135 ± 2.6 c	139 ± 1.8 ef	659 ± 4.1 e	665 ± 3.5 e
T2	10.3 ± 0.1 de	10.7 ± 0.2 def	14.0 ± 1.2 bc	15.5 ± 0.3 cd	136 ± 3.1 c	138 ± 1.8 ef	664 ± 4.3 e	664 ± 2.9 e
T3	10.8 ± 0.2 bc	11.4 ± 0.2 c	15.0 ± 0.9 abc	16.3 ± 0.3 bcd	147 ± 3.2 b	151 ± 3.2 cd	701 ± 7.6 d	734 ± 5.2 d
T4	11.2 ± 0.2 b	11.4 ± 0.2 c	15.1 ± 0.4 abc	16.5 ± 0.9 bcd	147 ± 4.9 b	153 ± 3.2 c	705 ± 3.2 d	724 ± 4.5 d
T5	11.3 ± 0.1 b	11.3 ± 0.2 c	15.2 ± 0.9 abc	16.6 ± 0.4 bcd	149 ± 2.6 b	151 ± 2.3 cd	720 ± 5.2 c	754 ± 6.8 c
T6	10.4 ± 0.2 cde	10.8 ± 0.1 de	14.2 ± 0.9 bc	15.6 ± 0.9 cd	142 ± 2.3 bc	144 ± 2.1 de	696 ± 5.1 d	738 ± 5.2 d
T7	10.5 ± 0.1 cd	11.1 ± 0.1 cd	14.6 ± 0.4 abc	16.0 ± 1.0 cd	142 ± 1.9 bc	145 ± 2.0 cde	692 ± 4.2 d	735 ± 5.0 d
T8	11.9 ± 0.2 a	12.4 ± 0.3 b	16.2 ± 0.9 ab	18.0 ± 0.6 ab	161 ± 3.4 a	168 ± 3.8 b	755 ± 5.2 b	798 ± 4.9 b
T9	12.4 ± 0.1 a	13.1 ± 0.2 a	16.7 ± 0.9 a	18.7 ± 0.9 a	168 ± 3.8 a	178 ± 3.2 a	787 ± 5.8 a	830 ± 4.6 a
LSD ($p \leq 0.05$)	0.4843 ****	0.4855 ****	2.3988 ns	1.9418 *	9.9833 ****	7.9141 ****	14.602 ****	14.073 ****
Treatment	Olsen P (mg kg ⁻¹)		Extractable K (mg kg ⁻¹)		DTPA extracted Fe (mg kg ⁻¹)		DTPA extracted Zn (mg kg ⁻¹)	
	Trial I	Trial II	Trial I	Trial II	Trial I	Trial II	Trial I	Trial II
T0	5.17 ± 0.11 f	5.92 ± 0.10 c	65.8 ± 0.90 f	66.2 ± 1.00 f	0.52 ± 0.002 e	0.53 ± 0.004 d	0.56 ± 0.004 f	0.58 ± 0.003 e
T1	5.20 ± 0.05 f	5.96 ± 0.07 c	67.3 ± 1.56 ef	67.9 ± 1.26 fg	0.52 ± 0.004 e	0.54 ± 0.004 d	0.57 ± 0.006 f	0.61 ± 0.009 de
T2	5.29 ± 0.06 f	5.99 ± 0.03 c	67.0 ± 1.99 ef	67.5 ± 1.15 fg	0.53 ± 0.002 e	0.54 ± 0.006 d	0.58 ± 0.008 ef	0.61 ± 0.009 de
T3	5.97 ± 0.07 bc	6.65 ± 0.05 b	71.5 ± 1.19 cd	74.8 ± 1.36 bc	0.56 ± 0.006 d	0.58 ± 0.009 c	0.59 ± 0.003 def	0.61 ± 0.014 de
T4	5.57 ± 0.04 c	6.49 ± 0.03 bc	72.3 ± 1.07 bcd	74.4 ± 0.50 cd	0.56 ± 0.008 d	0.58 ± 0.007 c	0.58 ± 0.002 ef	0.62 ± 0.015 cd
T5	5.87 ± 0.04 cd	6.67 ± 0.06 b	73.5 ± 0.83 abc	73.3 ± 1.71 cde	0.58 ± 0.007 c	0.59 ± 0.007 c	0.65 ± 0.013 b	0.67 ± 0.019 ab
T6	5.67 ± 0.03 e	6.27 ± 0.06 d	70.0 ± 0.67 cde	70.1 ± 1.68 ef	0.58 ± 0.006 cd	0.58 ± 0.003 c	0.61 ± 0.011 cde	0.65 ± 0.018 bc
T7	5.70 ± 0.06 de	6.34 ± 0.05 cd	69.2 ± 1.70 def	70.6 ± 1.09 def	0.57 ± 0.007 cd	0.58 ± 0.010 c	0.62 ± 0.012 cde	0.66 ± 0.018 bc
T8	6.12 ± 0.07 b	6.94 ± 0.07 a	76.2 ± 1.26 ab	78.6 ± 1.67 ab	0.61 ± 0.008 b	0.62 ± 0.029 b	0.62 ± 0.016 bc	0.65 ± 0.019 bc
T9	6.36 ± 0.06 a	7.11 ± 0.07 a	77.6 ± 1.63 a	80.8 ± 1.77 a	0.63 ± 0.005 a	0.66 ± 0.011 a	0.69 ± 0.008 a	0.70 ± 0.015 a
LSD ($p \leq 0.05$)	0.1788 ****	0.1918 ****	4.1686 ***	3.8615 ****	0.0155 ****	0.0335 ****	0.0280 ****	0.0330 ****

Data is presented as mean of three replicates ± standard error. Means sharing same letter(s) within a column are statistically non-significant at 5 % probability. *: $P < 0.05$; **: $P < 0.01$; ***: $P < 0.001$; ****: $P < 0.0001$; ns: non-significant



Table 2: Comparison of the per-hectare cost of production of wheat treated as a control (no application of zinc), ZnSO₄ and microbial-assisted zinc-enriched compost

Activity	Costs of production per hectare (Rs.)		
	Control	Market Source of Zinc (ZnSO ₄)	Microbial-assisted zinc-enriched compost
Land preparation	17,297	17,297	17,297
Seed	14,826	14,826	14,826
Fertilizers	88,956	88,956	84,508 *
Zinc application	0	4,448 **	12,085 ***
Manual Labor charges	2,965	2,965	2,965
Plant protection (weedicide application)	2,965	3,033	3,478
Mark up @ $i=12\%$ (land prep. to Fert. application)	14,529	15,063	15,446
Harvesting	12,355	12,355	12,355
Threshing	13,591	13,591	13,591
Total variable cost	167,484	172,534	176,551
Land Rent	61775	61775	61775
Abiana (Fixed water charges)	1112	1112	1112
Total fixed cost (Rs)	62887	62887	62887
Total cost (Rs.)	230,371	235,421	239,438

The cost of production was calculated using the current rates of commodities in the local market.
 *: Reduced application of chemical fertilizers by 5%; **: Price of ZnSO₄; ***: Price of microbial-assisted zinc-enriched compost



Table 3: Gross margin, net income and revenue to cost (R/C) ratio

Parameters	Control	Market Source of Zinc (ZnSO ₄)	Microbial-assisted zinc-enriched compost
Grain yield (mounds/ha)	115.5	117.4	125.75
Price (Rs./mound)	2,170	2,170	2,170
Revenue A (grain yield × price) (Rs.)	250,635	254,758	272,878
Biomass yield (mounds/ha)	122.75	128	145.75
Price (Rs./mound)	235	235	235
Revenue B (biomass yield × price) (Rs.)	28,846.25	30,080	34,251
Total revenue	279,481.25	284,838	307,129
Gross margin (Rs.) = TR-TVC	111,997	112,304	130,578
NR with LR (Rs.) = TR-TC	49,110	49,417	67,691 *
NR without LR (Rs.) = TR-TVC	110,885	111,192	129,466
Revenue/cost ratio with LR = TR/TC	1.21	1.21	1.28
Revenue/cost ratio without LR = TR/TVC	1.67	1.65	1.74

Yield is presented as average of both trials conducted in Bahawalpur and Bahawalnagar. *: significant increase in net income by the application of microbial-assisted zin-enriched compost




Figure 1. Climate data from experimental sites during the 2021-22 cropping season. Data is presented as the monthly average temperature (a) and total rainfall (mm) in a month (b)




Figure 2. Principal component analysis (PCA) demonstrates the effectiveness of microbial-assisted zinc-enriched compost to improve soil biochemical properties. (a): PCA - trial I (Bahawalpur); (b): PCA - trial II (Bahawalnagar); (c): eigenvalues – trial I; (d): eigenvalues – trial II; Zn: DTPA-extracted zinc; TN: total nitrogen; CFU: colony forming units; OM: organic matter; TOC: total organic carbon; MBC: microbial biomass carbon; MBN: microbial biomass nitrogen; Fe: DTPA extracted iron; P: Olsen phosphorous; K: extractable potassium




Figure 3. Effect of microbial-assisted zinc-enriched compost on the growth and yield of wheat. The same letter(s) on the bars represent statistically non-significant variation at 5% probability ($n = 3$, $p \leq 0.05$). Trial I: Bahawalpur; Trial II: Bahawalnagr; (a): plant height; (b): straw yield; (c): 1000-grain weight; (d): grain yield




Figure 4. Effect of microbial-assisted zinc-enriched compost on the macro- and micronutrient concentration in wheat grain. The same letter(s) on the bars represent statistically non-significant variation at 5% probability ($n = 3$, $p \leq 0.05$). Trial I: Bahawalpur; Trial II: Bahawalnagar




Figure 5. Pearson's correlation analysis was used to describe the relationship between the studied soil parameters. The thickness of the ellipses represents the correlation coefficient (R) and the asterisks represent the significance based on *p-value*. a: Trial I (Bahawalpur); b: Trial II (Bahawalnagar)