

A fault-based application to model seismicity rates for seismic hazard assessment in the southern Apennines (Italy)

Giulia Alessandrini¹, Octavi Gómez-Novell², Silvia Castellaro³, Michela Giustiniani⁴, Umberta Tinivella⁴
¹ Istituto Nazionale di Geofisica e Vulcanologia (INGV), Sezione di Bologna, Bologna, 40127, Italy
² Instituto Geológico y Minero de España (IGME-CSIC), Madrid, 28003, Spain
³ Department of Physics and Astronomy, Alma Mater Studiorum Università di Bologna, Bologna, 40127, Italy
⁴ Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS), Borgo Grotta Gigante, Sgonico, 34010, Italy
Correspondence to: Giulia Alessandrini (giulia.alessandrini@ingv.it)

Abstract. Although fault-based approaches to seismic hazard assessment have been increasingly adopted worldwide, the official Italian hazard model, on which the national building code is based, still relies on a catalogue-based framework, with well-known limitations in capturing the long-term recurrences of large-magnitude events. In this study, we present a fault-based application to model seismicity rates for the southern Apennines (Italy) that incorporates a multi-fault rupture assumption. This area is of particular interest due to its active seismicity and the presence of large dams, for which robust long-term hazard estimates are essential. We use the SHERIFS code to model seismicity rates at the fault system-level, which allow us to explore epistemic uncertainties of fault and seismicity parameters (rupture scenarios, scaling laws, *b*-values and background seismicity). Our results highlight the key role of rupture models: scenarios allowing multi-fault ruptures outperform single-fault rupture models in terms of agreement with the regional seismicity and paleoseismic rates. Our findings support the inclusion of multi-fault rupture models in PSHA logic trees for the region and emphasize the need for improved fault behaviour characterization in southern Italy.

1 Introduction

In recent decades, fault-based approaches to seismic hazard assessment have gained increasing attention worldwide as an alternative to traditional methods relying exclusively on earthquake catalogues. These latter approaches, based on the original formulation of Probabilistic Seismic Hazard Analyses (PSHA) by Cornell (1968), have completeness limitations related to the combined effect of short earthquake catalogues (usually no more than a thousand years) and the long recurrence intervals of moderate-to large-magnitude earthquakes. These effects hamper the earthquake forecasting capabilities of PSHA, especially in regions with lower deformation rates. Moreover, catalogue-based approaches classically employ distributed seismicity models (area sources, grid sources, etc.) which, although suitable for representing background seismicity, i.e. earthquakes occurring on unmapped or poorly constrained faults, tend to unify or smooth out the seismic hazard over larger regions rather

Definizione stile	... [35]
Definizione stile	... [34]
Definizione stile	... [33]
Definizione stile	... [32]
Definizione stile	... [31]
Definizione stile	... [30]
Definizione stile	... [29]
Definizione stile	... [28]
Definizione stile	... [27]
Definizione stile	... [26]
Definizione stile	... [25]
Definizione stile	... [24]
Definizione stile	... [23]
Definizione stile	... [22]
Definizione stile	... [21]
Definizione stile	... [20]
Definizione stile	... [19]
Definizione stile	... [18]
Definizione stile	... [17]
Definizione stile	... [16]
Definizione stile	... [15]
Definizione stile	... [14]
Definizione stile	... [13]
Definizione stile	... [12]
Definizione stile	... [11]
Definizione stile	... [10]
Definizione stile	... [9]
Definizione stile	... [8]
Definizione stile	... [7]
Definizione stile	... [6]
Definizione stile	... [5]
Definizione stile	... [4]
Definizione stile	... [3]
ha formattato	... [36]
Formattato	... [37]
ha formattato	... [39]
Formattato	... [38]
ha formattato	... [40]
ha eliminato: ³ ,	
ha formattato	... [41]
ha eliminato: Giustiniani ¹	
ha formattato	... [42]
ha eliminato: Tinivella ¹	
ha formattato	... [43]
ha eliminato: ¹ Istituto	
ha formattato	... [44]
ha eliminato: Oceanografia e di	
ha formattato	... [45]
ha eliminato: Sperimentale (OGS), Borgo Grotta Gigante	... [46]
ha formattato	... [47]
ha eliminato:	
ha formattato	... [48]
ha eliminato: ³ Department	
ha formattato	[49]

52 than localizing ~~it~~ along main active fault structures. Conversely, fault-based PSHA allows also ~~the incorporation of~~ fault-
53 specific geological data (e.g., paleoseismology, geomorphology) into the source modelling, ~~enabling~~ a time-extended and more
54 reliable characterization and localization of the seismic activity of a region. ~~This approach provides improved spatial resolution~~
55 ~~of seismic hazard associated with moderate-to-large magnitude earthquakes ($M_w \geq 6.5-7.0$), whose occurrence is strongly~~
56 ~~controlled by the geometry, slip rates and segmentation of major active faults (e.g., Field et al., 2009; Gerstenberger et al.,~~
57 ~~2024).~~
58 Quantitative comparisons between the two approaches have shown that incorporating geological fault information may
59 significantly modify both the spatial variability and the amplitude of ground-motion estimates. For example, fault-based
60 models have been shown to produce increases in peak ground acceleration (PGA, 10% probability of exceedance in 50 years)
61 of up to 0.3 g compared to catalogue-based models in regions characterized by low deformation rates and limited instrumental
62 seismicity (e.g., Williams et al., 2023), while differences exceeding 100% have been observed for sites located close to major
63 active faults (Gómez-Novell et al., 2020b). In California, the transition from area-source models to the fault-based UCERF
64 framework reduced discrepancies between observed and modelled rates of intermediate-magnitude earthquakes ($6.5 \leq M \leq$
65 7.0), improving consistency with historical catalogues (Field et al., 2009).
66 Despite the evolution and implementation of fault-based PSHA approaches in many countries worldwide (e.g., New Zealand,
67 Gerstenberger et al., 2024; California, Petersen et al., 2024), catalogue-based PSHA remains the standard practice, especially
68 for official seismic hazard assessment and regulatory applications such as the building codes (e.g., Spain, NCSE-02, 2002;
69 Italy, Stucchi et al., 2011). In Italy, for example, the national building code still relies on an area-source Earthquake Rupture
70 Forecast (ERF; NTC, 2008, 2018).
71 However, the reliance of catalogue-based hazard models on historical earthquakes can lead to the exclusion of faults with
72 paleoseismic evidence of rupture -and thus capable of generating significant earthquakes- that have not produced events within
73 the historical record. A relevant example, in Italy, is the Mount Vettore Fault, which ruptured during the 2016 Central
74 Apennines earthquake even though it was previously classified as silent (Galadini & Galli, 2003; Galli et al., 2019).
75 Beyond these limitations, catalogue-based approaches do not account for the complexity of earthquake ruptures observed in
76 nature. Seismological, geological and paleoseismological data show that earthquake ruptures can be very complex, involving
77 the simultaneous activation of faults with different characteristics. Recent complex coseismic ruptures highlighted the need to
78 include the multi-fault earthquakes in PSHA, going thus beyond strict fault segmentation assumptions (e.g., M_w 6.9 Irpinia in
79 1980, Bernard & Zollo, 1989; M_w 7.1 El Mayor-Cucapah in 2010, Wei et al., 2011; M_w 8.6 Sumatra in 2012, Zhang et al.,
80 2012; M_w 6.5 central Italy in 2016, Peruzza et al., 2016; M_w 7.8 Kaikoura in 2016, Hamling et al., 2017; ~~Mw 7.8 and Mw 7.7~~
81 ~~Türkiye-Syria in 2023, Liu et al., 2023).~~ Capturing such multi-fault ruptures is a key challenge in fault-based PSHA:
82 implementations in California with UCERF3 (Field et al., 2009) or New Zealand (Stirling et al., 2012) pioneered the inclusion
83 of such fault rupture complexities in PSHA.

ha eliminato: them
ha eliminato: to incorporate
ha eliminato: allowing for

ha eliminato: Gerstenberger et al., 2024; California, Petersen et al., 2024),
ha formattato: Inglese (Regno Unito)
ha formattato: Inglese (Regno Unito)

ha eliminato: (Galadini & Galli, 2003; Galli et al., 2019).

ha formattato: Inglese (Regno Unito)

ha eliminato:).

ha eliminato: 2
ha formattato: Colore carattere: Nero
Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

91 However, incorporating faults into seismic hazard assessment in low-strain regions, such as Europe, is not yet a widespread
92 practice and remains a topic of debate in many countries. A regulated implementation of such fault-based models in Europe
93 remains a challenge mainly due to the larger recurrence intervals of the active faults and therefore to the scarcity of data to
94 characterize such activity. In recent years, some initiatives such as the Fault2SHA Working Group of the European
95 Seismological Commission have been founded to discuss the implementation of fault-based PSHA in Europe (central Italy,
96 Peruzza et al., 2011; Valentini et al., 2018, 2019; France, Scotti et al., 2014; Greece, Deligiannakis et al., 2018; and Spain,
97 Gómez-Novell et al., 2020b). Along this line, in the recent MPS19 Seismic Hazard Model of Italy, two fault-based models
98 were included in the earthquake rupture forecast, marking a significant step forward in Europe (Meletti et al., 2021; Visini et
99 al., 2021).

100 In this study we model fault seismicity rates in the southern Apennines using a relaxed segmentation framework, i.e.,
101 considering multi-fault rupture scenarios. To our knowledge, this is the first study in the area to adopt such an approach for
102 seismic hazard assessment. The Irpinia region, severely affected by the 1980 Mw 6.9 earthquake, the third deadliest in Italy
103 (Peruzza, 2018), hosts several large earth dams located near active but poorly characterized faults, highlighting the need for
104 accurate estimates of fault activity rates.

105 The study is designed as a comprehensive sensitivity test, aimed at observing how various model configurations of fault
106 ruptures, fault and seismicity parameters impact fault system-level seismicity rates for seismic hazard assessment.

107 **1.1 Geological and seismological setting**


108 The study area, covering about 50,000 km², is centred around the Irpinia district, one of the most seismically active regions of
109 the southern Apennines (southern Italy; Fig. 1). The Southern Apennines are part of an east-verging thrust belt related to the
110 west-dipping subduction of the Apulian lithosphere beneath the Tyrrhenian domain (Doglioni et al., 1996). Since the Late
111 Quaternary, slab rollback and trench retreat have promoted extensional tectonics along the Apennine chain, resulting in a
112 system of large, segmented NW-SE-striking normal faults accommodating crustal stretching (Doglioni et al., 1996). At the
113 same time, strike-slip to oblique-slip faulting is observed in the outer sectors of the belt and within the Adriatic foreland. These
114 structures are commonly interpreted as the reactivation of inherited crustal discontinuities within the Apulian foreland,
115 occurring at depth (typically ~15-30 km; Latorre et al., 2023) and partially decoupled from the shallow extensional regime
116 affecting the chain (generally confined within the upper ~10-15 km of the crust; e.g., Boncio et al., 2007; Vannoli & Burrato,
117 2018). Recent studies (e.g., Adinolfi et al., 2015) have proposed that the coexistence of extensional and strike-slip faulting
118 may reflect a vertically partitioned seismotectonic system, with shallow normal faulting and deeper strike-slip deformation.
119 However, the relationship between these deformation regimes, and whether they are mechanically coupled or spatially
120 overlapping, remains debated.

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: The study area, covering about 50,000 km², is centred around the Irpinia district, in southern Italy (Fig. 1). This area is located in the southern Apennines, an east-verging thrust belt related to the west-dipping subduction of the Apulian lithosphere (Doglioni et al., 1996).



ha eliminato: 3

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra




Figure 1. Geographical distribution of the $M_w \geq 4.0$ earthquakes occurred in the southern Apennines between 1019 and 2017 ($4.0 \leq M_w \leq 7.2$; reference catalogue: CPTI15 v4.0, Rovida et al., 2022) and the fault traces of the 35 seismogenic sources used for this study (references: Valentini et al., 2017; DISS Working Group, 2025). Earthquakes are colour-coded by magnitude class (M_w bin = 1.0), with larger symbols representing stronger events; major historical earthquakes are labelled by year. Faults are colour-coded by tectonic domain. Latitude and longitude grid shown in WGS84 coordinates.

This tectonic framework explains the subdivision of the study area into two main domains or fault systems. The strongest seismic events of the region (Fig. 1) mostly occurred in one of the domains, a segmented belt of large normal NW–SE striking faults, running along the chain axis, hereafter “SubArea 1” (DISS Working Group, 2025). In the other system, i.e., the foreland, faulting develops along E–W right-lateral strike-slip to oblique-slip faults, hereafter “SubArea 2” (DISS Working Group, 2025).

The strongest events (Fig. 1) reported in the Parametric Catalogue of Italian Earthquakes - CPTI15 v4.0 (Rovida et al., 2022) for this region are the 1456 Sannio-Irpinia sequence (M_w 7.2), which is one of the most destructive events that took place in the Italian peninsula, the 1688 (M_w 7.1) and the 1857 (M_w 7.1) earthquakes. The probably most famous destructive earthquake,

ha eliminato: 1.

ha eliminato: M_w

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato: M_w

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Colore carattere: Nero

ha eliminato: The regions surrounding the Irpinia district lie in one of the most seismically active regions of the southern Apennines. Tectonically, the area is divided into two main domains or fault systems. The strongest seismic events of the region (Fig. 1) mostly occurred in one of the domains, a segmented belt of large normal NW–SE striking faults, running along the chain axis, hereafter “SubArea 1” (DISS Working Group, 2025). In the other system, i.e., the foreland, faulting develops along E–W right-lateral strike-slip to oblique-slip faults related to the roll-back of the Adriatic foreland, hereafter “SubArea 2” (DISS Working Group, 2025). In this domain, instrumental earthquakes characterized by strike-slip kinematics generally involve a deeper seismogenic thickness (≈ 15 –30 km deep; Latorre et al., 2023) than earthquakes with extensional kinematics, that exclusively characterize the first kilometres of crustal thickness (≈ 0 –15 km deep; Vannoli & Burrato, 2018).

ha eliminato: 4

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

160 the Mw 6.9 that hit the Irpinia district in 1980, had devastating effects in a large area of the southern Apennines, causing
161 thousands of casualties and high damage to buildings. Latorre et al. (2023) observe that the highest earthquake concentration
162 in the southern Apennines consistently occurs along the faults responsible for the 1980, Mw 6.9 Irpinia earthquake. Close to
163 this epicentre, it is also worth to mention the 1694 (Mw 6.7) Irpinia-Basilicata earthquake, considered the ancestor of the 1980
164 (DISS Working Group, 2025), the 1851 (Mw 6.5), and the 1930 (Mw 6.7) Irpinia earthquake.

165 **2 Methods, data and model parameters**

166 **2.1 The SHERIFS method**

167 In this study, we employ SHERIFS V1.1 (Seismic Hazard and Earthquake Rate In Fault Systems), an approach and
168 accompanying code developed by [Chartier et al. \(2019\)](#), [designed to model seismicity rates in active fault systems using](#)
169 [geological fault data as inputs. The approach treats fault systems as a whole, explicitly accounting for multi-fault rupture](#)
170 [scenarios in the estimation of seismicity rates. So far, SHERIFS has been tested in several regions worldwide including SE](#)
171 [Spain \(Gómez-Novell, et al., 2020a, Gómez-Novell, et al., 2020b\), the Marmara Region in Turkey \(Chartier et al., 2021\), the](#)
172 [Tibetan Plateau \(Cheng et al., 2021\) and the Levant Fault \(El Kadri et al., 2025\).](#)

173 To derive seismicity rates from geological slip-rate data, SHERIFS requires several inputs: *(i)* the 3D geometry and slip rates
174 of the fault system, *(ii)* a set of multi-fault rupture scenarios based on user-defined criteria (e.g., inter-fault distance), and *(iii)*
175 a target Magnitude-Frequency Distribution (MFD) specified at the fault-system level (e.g., a Gutenberg-Richter law). The
176 epistemic uncertainties of the input parameters in SHERIFS are explored through an iterative process that performs the
177 computation *n* times exploring *n* different values of the input parameters within their uncertainty ranges. The result is the
178 average and uncertainty dispersion cloud of all samples explored.

179 One of the main characteristics of SHERIFS is that the approach treats the prescribed slip rate of the fault system as a budget.
180 Through an iterative procedure, this slip rate budget is distributed among faults according to the magnitudes they can
181 accommodate, following analytical relationships that link seismic moment rate to seismicity rates. The slip rate budget is spent
182 until the pre-imposed MFD target is reached and the slip rate budgets are consumed. In some cases, the target is reached before
183 the budget of all faults is exhausted and therefore there is remaining slip rate budget that is not converted into seismicity rates.
184 This remaining budget, namely Non-Main-Shock slip (hereafter NMS) in SHERIFS, is indeed an artifact of the modelling.
185 However, according to the authors, small percentages of NMS (generally below 30-40% of the total budget of the fault), can
186 be interpreted as a natural phenomenon (e.g., post-seismic slip or creep) as geological slip rates might not be exclusively
187 resulting from the coseismic phase, but also from other post-seismic relaxation processes. Conversely, NMS percentages above
188 30-40% are less easily attributable to the natural phenomena described. Instead, they most likely indicate that the combination
189 of input hypotheses used (fault parameters, target MFD and rupture hypotheses) is not optimal in the SHERIFS framework
190 and that they should be reconsidered (Chartier et al., 2019).

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

Formattato: Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 0,76 cm

ha eliminato: Chartier et al. (2019), designed to model seismicity rates in active fault systems using geological fault data as inputs. The approach treats fault systems as a whole, explicitly accounting for multi-fault rupture scenarios in the estimation of seismicity rates. So far, SHERIFS has been tested in several regions worldwide including SE Spain (Gómez-Novell, et al., 2020a, Gómez-Novell, et al., 2020b), the Marmara Region in Turkey (Chartier et al., 2021), the Tibetan Plateau (Cheng et al., 2021) and the Levant Fault (El Kadri et al., 2024)

ha formattato: Inglese (Regno Unito)

ha formattato: Tipo di carattere: Corsivo, Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Tipo di carattere: Corsivo, Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha formattato: Tipo di carattere: Corsivo, Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: -

ha eliminato: .

ha formattato: Inglese (Regno Unito)

ha formattato: Inglese (Regno Unito)

ha eliminato: coseismic

ha eliminato: 5

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra




Figure 2. Structure of the main model branches (blue) used to explore epistemic uncertainty in the fault-based model, and of the consistency check criteria (violet) adopted to assess their performance.

In our model, the epistemic uncertainty is explored in two ways:

1. **Model branches:** we define 13 modelling branches (Fig. 2), covering variables such as rupture scenarios, scaling relationships, b -values, and background seismicity and buffer areas.
2. **Random samples:** we run 10 random samples per branch that explore different values of slip rate on faults within their uncertainty ranges.


The inputs used in our modelling, which are discussed in detail in the following sections, are archived and openly accessible in a Zenodo repository (Alessandrini et al., 2026).

2.2 Input data and model parameters

2.2.1 Fault sources

We use 35 faults of the southern Apennines for our study. All fault identification names (thereafter IDs), and their main characteristics are listed in Table 1, while their geometries are shown in Fig. 3. Fault traces, 3D geometry (dip and seismogenic depth), slip rates and kinematics are mainly extracted from the Individual Seismogenic Sources (ISS) within the Database of Individual Seismogenic Sources - DISS 3.3.1 (DISS Working Group, 2025), a geo-referenced archive of information on seismogenic sources and of their seismogenic potential (Basili et al., 2008). This database is a reference in Italy and many recent seismic hazard studies have employed this database for seismic source modelling (e.g., Stucchi et al., 2011; Woessner et al., 2015). We include some additional active faults reported by Valentini et al. (2017), that are not part of the DISS (respectively faults 11, 15, 16, 26, 27, 28, 29, 30, 32, 33, 34, 35).

Model



ha eliminato:

ha eliminato: 2.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: 10

Formattato: Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm

ha eliminato: 2025

ha eliminato: <#>Input data and model parameters

Fault sources

We use 35 faults of the southern Apennines for our study. We extract these fault sources, namely their traces, 3D geometry (dip, seismogenic depth), slip rates and kinematics mainly from the Individual Seismogenic Sources (ISS) within the Database of Individual Seismogenic Sources - DISS 3.3.1 (DISS Working Group, 2025), a geo-referenced archive of information on seismogenic sources and of their seismogenic potential (Basili et al., 2008). This database is a reference in Italy and many recent seismic hazard studies have employed this database for seismic source modelling (e.g., Stucchi et al., 2011; Woessner et al., 2015). Additionally, we include some additional active faults reported by Valentini et al. (2017), that are not part of the DISS (respe... [55])

ha eliminato: 6

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra




Figure 3. Representation of the 35 fault sections, coloured based on their mean slip rates (references: Valentini et al., 2017; DISS Working Group, 2025).

Fault geometries and segmentation are directly adopted from these datasets, which define individual seismogenic sources based on geological, geophysical and seismotectonic criteria, including geological mapping, seismicity distribution, geomorphic evidence and geodetic observations. Potential discontinuities in fault traces, particularly within the strike-slip systems, may reflect the presence of unmapped or blind structures. For example, additional faults are identified in the ITHACA catalogue (ITaly HAZards from CApable faults; ITHACA Working Group, 2019). However, these structures generally lack the quantitative parameters required for earthquake rate modelling, such as slip rates and seismogenic depths. For this reason, only faults with sufficiently constrained geometric and kinematic parameters are included in the present modelling framework. The adopted segmentation therefore reflects the current state of knowledge on active faults for seismic hazard applications in the region.

The slip rates we assigned to each fault come from the DISS 3.3.1 database and by Valentini et al. (2017) and are constrained by geological and geodynamic observations. Variability in slip rates across adjacent faults is consistent with the heterogeneous accommodation of strain within fault systems, which depends on factors such as fault geometry, orientation and mechanical properties. This behaviour has been documented in several tectonic settings, including the Apennines, and is reflected in regional fault databases (e.g., Faure Walker et al., 2021).

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: The slip rates we assigned to each fault come from the DISS 3.3.1 database and by Valentini et al. (2017), and are based on different methods: mainly geodynamic and geological constraints.




Figure 3. Representation of the 35 fault sections, colored

ha eliminato: 7

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

285
286
287
288

Table 1. List of faults employed in the models. Faults highlighted by a star (*) are the integration of Valentini et al. (2017) while the other faults come from the DISS 3.3.1 database (DISS Working Group, 2025). For each of them, we list here the name, the fault ID used in the model, the dip angle, the kinematics (S = strike-slip, N = normal), the seismogenic depth range (km), length and the slip rate range (mm/yr). Paleoseismic data are available for faults reported in the DISS database and here marked by a double star (**).

Fault Name	Fault ID	Dip (°)	Kinematics	Seismogenic depth range (km)	Length	Min slip rate (mm/yr)	Mean slip rate (mm/yr)	Max slip rate (mm/yr)
Frosolone	1	70	S	11 - 25	36.0	0.1	0.55	1.0
Ripabottoni	2	86	S	12 - 20	9.4	0.1	0.3	0.5
San Giuliano di Puglia	3	82	S	12 - 20	10.5	0.1	0.3	0.5
San Severo	4	80	S	6 - 21	34.0	0.1	0.3	0.5
San Marco Lamis	5	80	S	0 - 12	10.0	0.1	0.55	1.0
San Giovanni Rotondo	6	80	S	0 - 12	11.0	0.1	0.55	1.0
Monte Sant' Angelo	7	80	S	0 - 12	20.0	0.5	0.85	1.2
Ascoli Satriano	8	80	S	13 - 21	12.6	0.1	0.3	0.5
Cerignola	9	80	S	11 - 22	18.6	0.1	0.3	0.5
Bisceglie	10	80	S	13 - 19	8.6	0.1	0.3	0.5
*Matese	11	60	N	0 - 13	48.3	0.2	1.05	1.9
Carpino - Le Piane	12	60	N	1 - 7	8.9	0.1	0.55	1.0
Boiano Basin	13	55	N	1 - 12	24.0	0.1	0.55	1.0
Tammaro Basin	14	60	N	1 - 13	25.0	0.1	0.55	1.0
*Ailano-Piedimonte	15	60	N	0 - 12	17.6	0.15	0.25	0.35
*Benevento	16	55	N	0 - 10	25.0	0.35	0.65	0.95
Ariano Irpino	17	70	N	11 - 25	30.0	0.1	0.55	1.0
Ufita Valley	18	64	N	1 - 14	25.6	0.1	0.55	1.0
Bisaccia	19	64	N	1 - 15	31.0	0.1	0.55	1.0
Melfi	20	80	S	12 - 23	17.2	0.1	0.3	0.5
**Pescopagano	21	70	N	1 - 10	15.0	0.4	0.5	0.6
**Colliano	22	60	N	1 - 14	28.0	0.4	0.5	0.6
**San Gregorio Magno	23	60	N	1 - 14	9.0	0.4	0.5	0.6
Melandro - Pergola	24	60	N	1 - 11	17.9	0.1	0.55	1.0
Agri Valley	25	60	N	1 - 13	23.0	0.1	0.55	1.0
*Monte Alpi	26	60	N	0 - 13	10.9	0.35	0.63	0.9
*Caggiano-Diano	27	60	N	0 - 12	46.0	0.35	0.78	1.2
*Maratea	28	60	N	0 - 13	21.6	0.46	0.58	0.7
*Agri	29	50	N	0 - 13	34.9	0.8	1.05	1.3
*Alburni	30	60	N	0 - 8	20.4	0.35	0.53	0.7
Potenza	31	88	S	15 - 21	7.9	0.1	0.3	0.5
*Palagianello	32	90	S	0 - 22	73.3	0.1	0.3	0.5
*Vulturno	33	60	N	0 - 13	15.7	0.23	0.4	0.57
*Avella	34	55	N	0 - 13	20.5	0.2	0.45	0.7
*Vulturara	35	60	N	0 - 13	23.7	0.2	0.28	0.35

289

ha eliminato: ¶

ha eliminato: 1.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Colore carattere: Nero

Tabella formattata

ha eliminato: 8¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

293 2.2.2 Segmentation rules and rupture scenarios

294 We explore fault-rupture scenarios to compute system-level seismicity rates under different hypotheses of fault connectivity
295 within the fault system. We define three fault rupture scenarios for the whole fault system (Table 2), each representing an
296 incremental increase in fault connectivity and, consequently, in multi-fault rupture length. These scenarios are exploratory
297 hypotheses, and as such, other configurations could be tested.

299 Table 2. List of the three fault rupture scenarios, characterized by incremental multi-fault ruptures. For each scenario the
300 combinations of the maximum ruptures are listed, and the expected maximum magnitude range is provided. Fault sections (from 1
301 to 35) are shown in Fig. 3.

Rupture scenario	Fault/multi-fault ruptures	Maximum magnitude range (M_w)
Set_0	Only single fault section ruptures allowed	7.0 – 7.4
Set_1	F2 F3	7.1 – 7.4
	F4 F5 F6 F7	
	F11 F14	
	F18 F19	
	F22 F23	
Set_2	F24 F27 F30	7.1 – 7.6
	F1 F2 F3	
	F4 F5 F6 F7	
	F8 F9	
	F11 F12 F13 F14 F15 F16 F17	
	F18 F19	
	F21 F22 F23 F24 F25 F27 F29 F30 F35	

- 302
- 303 • **Set_0.** In this hypothesis, only single fault section ruptures are allowed. The segment lengths reported in the literature
304 for each fault define the maximum rupture length, and multi-segment ruptures involving neighbouring faults are not
305 allowed.
 - 306 • **Set_1.** In this scenario a subset of multi-fault ruptures is allowed based on geometric compatibility rules (i.e., faults
307 with similar strike, dip direction, seismogenic depth), kinematics compatibility, as well as a fault distance threshold.
308 In terms of kinematic compatibility, we follow the cumulative rake change criterion suggested by Milner et al. (2013),
309 which limits the total rake variation within a rupture to 180 degrees to prevent kinematic incompatibility. This means
310 that only faults with the same kinematics are allowed to rupture together. In terms of distance for rupture propagation,
311 we select 5 km as the threshold for rupture propagation, consistent with previous studies on multi-fault rupture
312 connectivity, including the JCERF-3 framework (e.g., Milner et al., 2013; Field et al., 2014; Scotti et al., 2020),
313 where similar inter-fault jump distances are commonly used. Based on the above criteria, Set_1 permits ruptures
314 between neighbouring faults up to a maximum of 4 faults rupturing together (Fig. 4) with rupture lengths up to 75
315 km, which falls well within the typical rupture lengths observed in Gerstenberger et al. (2024), usually below a few
316 hundred kilometres.

Formattato: Struttura + Livello:3 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,27 cm

ha eliminato: (Table 2),

ha eliminato: 2.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: M_w

Tabella formattata

ha eliminato: F6F 7

ha formattato: Colore carattere: Nero

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

Formattato: Normale, Rientro: Sinistro: 0,63 cm, Sporgente 0,63 cm, SpazioDopo: 0 pt, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato: neighboring

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

ha formattato: Colore carattere: Nero

ha formattato: Colore carattere: Nero

ha eliminato: excessive

ha formattato: Colore carattere: Nero

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

ha eliminato: values reported in

ha eliminato: literature

ha eliminato:).

ha formattato: Colore carattere: Nero, Inglese (Regno Unito)

ha formattato: Colore carattere: Nero

ha formattato: Colore carattere: Nero

ha formattato: Colore carattere: Nero

ha formattato ... [57]


ha formattato: Colore carattere: Nero

ha eliminato: 9

ha formattato: Colore carattere: Nero

Formattato ... [56]

326 • **Set 2.** The distinct feature of Set 2 is the adoption of a larger distance threshold of 10 km for rupture propagation, while maintaining the requirement of compatible faulting mechanism. This configuration is intentionally more speculative and is introduced to explore an upper-bound range of rupture behaviours and their impact on seismicity rates. The choice of extending the distance threshold beyond 5 km is motivated by observations of complex multi-fault ruptures in the study area, such as the 1980 M_W 6.9 Irpinia earthquake, which involved multiple adjacent fault segments (faults 21–22–23) separated by distances exceeding 5 km. Under this configuration, up to 8 faults can rupture together (Fig. 4) resulting in rupture lengths up to 184 km.



334 **Figure 4.** Maximum extent (number of faults) of the multi-fault rupture scenarios in Set 1 and Set 2 explored in this study. Faults with the same colour are allowed to rupture together in a same rupture within that particular scenario set.

337 In order to study the relationship between magnitude and rupture area, we employ the empirical scaling relationships of Wells and Coppersmith (1994), Leonard (2010), and Thingbaijam et al. (2017). These scaling laws, widely used in PSHA applications, have broad applicability across tectonic settings and are implemented in the current version of the SHERIFS code. Leonard (2010) has been employed in regional studies in Italy and is used for the parametrization of national databases (e.g., DISS Working Group, 2025), supporting its suitability for the study area. Wells and Coppersmith (1994) provide a robust and widely adopted model that does not require specific adjustments between continental extensional and compressional regimes. In addition, Thingbaijam et al. (2017) provides coefficients specifically derived for normal faults, which represent the dominant faulting style in the Apennines, allowing a better representation of this tectonic regime. The maximum magnitude (M_{max}) range is set by the scaling relationship, and it is shown for each scenario in Table 2. A comparative evaluation of their performance within the study area is presented in Sect. 4.3, where their implications for the modelling results are discussed.

- ha eliminato: application
- ha eliminato: 10 km
- ha eliminato: ruptures, with faults allowed to rupture together as long as they share
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha eliminato: same
- ha eliminato: The decision to extend the distance threshold in this set was made
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha eliminato: a wider
- ha eliminato: to evaluate
- ha eliminato: onto
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha eliminato: This assumption allows for
- ha eliminato: rupturing
- ha formattato: Colore carattere: Nero
- ha formattato: Colore carattere: Nero
- ha eliminato: with
- ha formattato: Colore carattere: Nero
- ha eliminato: In order to study the relationship between magnitude and rupture area, the Wells & Coppersmith (1994), Leonard (2010) and Thingbaijam et al. (2017) empirical laws have been employed, as they represent the ones implemented in the current version of the code. The maximum magnitude (M_{max}) range is set by the scaling relationship, and it is shown for each scenario in Table 2. A comparative evaluation of their performance within the study area is presented in Sect. 4.3, where their implications for the modelling results are discussed.
- ha eliminato: 4.
- ha formattato ... [60]
- Formattato ... [61]
- ha formattato ... [62]
- ha formattato: Colore carattere: Automatico
- ha eliminato: 10
- ha formattato: Colore carattere: Nero
- Formattato ... [58]

374
375
376
377
378
379

380
381
382
383
384
385
386
387
388
389
390
391

392
393
394
395
396
397
398
399
400
401

2.2.3 Model MFD shape

To model the magnitude-frequency distribution (MFD) of seismicity on individual faults, we adopt a Gutenberg-Richter (G-R) relationship, which is widely used in PSHA and is consistent with the Italian national seismic catalogue (e.g., CPTI; Rovida et al., 2022). To explore the influence of epistemic uncertainty, we test two different *b*-value ranges: a broader interval (0.9-1.1) based on the national CPTI catalogue (Rovida et al., 2022), and a narrower one (0.93-0.96) supported by previous studies focused on this region (Gulia & Meletti, 2007; Meletti et al., 2008).

2.2.4 Background seismicity

One common challenge in fault-based PSHA is selecting a magnitude threshold for a confident assignation of seismicity to the known active faults, as small-to-moderate earthquakes rarely cause surface faulting and therefore may not be related to the known active faults. Because the classical selection of a cut-off threshold can be somewhat arbitrary, SHERIFS deals with this issue by incorporating magnitude-dependent ratios that determine the proportion of the seismicity that will be assigned to the faults, usually increasing with magnitude. That is, for each magnitude a user-defined proportion -ratio- of the seismicity rates of the faults is subtracted and assigned to the background as a distributed source. This approach allows for a smoother distribution of seismicity between faults and background, and is consistent with real observations.

We explore two alternative scenarios for background seismicity:

- **BG_1**, a fault-only model where all seismicity is attributed to the known active faults.
- **BG_2**, a hybrid model where a magnitude-dependent fraction of seismicity is assigned to the background buffer (Fig. 5), outside the mapped faults.

To define the BG_2 scenario, we set up a buffer area of 10 km around the faults (Fig. 5), where background seismicity is assigned. This 10-km distance is consistent with background zone sizing proposed by Chartier et al. (2019) (e.g., 10 or 20 km from faults). In order to evaluate the sensitivity of the results to this modelling choice, we also tested alternative buffer distances of 5 km and 15 km. The comparison between buffer sizes is further discussed in Sect. 4.4, and the corresponding models and figures are provided in the Zenodo repository (Alessandrini et al., 2026).

The ratios for each magnitude bin (Table 3) are custom set up using a magnitude-dependent criterion, based on the assumption that larger earthquakes are more likely to be associated with known, geomorphologically recognized faults, whereas smaller events may occur in secondary unmapped or buried faults. Consequently, BG_2 assigns higher background ratios to lower magnitudes.

Formattato: Struttura + Livello:3 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,27 cm

ha eliminato: –

ha eliminato: –

Formattato: Struttura + Livello:3 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,27 cm

ha eliminato: ,


ha eliminato: ,

ha formattato: Colore carattere: Nero

Formattato

... [64]

ha formattato: Colore carattere: Nero



ha eliminato:

... [65]

ha formattato: Colore carattere: Nero

ha eliminato: (Table 3)

Formattato: SpazioDopo: 8 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha eliminato: 11†

ha formattato: Colore carattere: Nero

Formattato

... [63]




Figure 5. Buffer area (10 km around the faults, dashed line) used in the BG 2 scenario to define the background seismicity and to extract the catalogue for the consistency check. Earthquakes from the CPTI15 v4.0 catalogue (Rovida et al., 2022) are shown for reference. Internal buffer areas are also displayed for the two structural domains (SubArea 1, blue shaded buffer and SubArea 2, pink shaded buffer).

Table 3. Ratio of seismicity occurring on known faults for each background scenario considered in the model (M_w bin = 0.5).

Magnitude (M_W)	4.0	4.5	5.0	5.5	6.0	6.5	7.0	7.5	8.0
Background 1 (BG_1)	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Background 2 (BG_2)	0.4	0.4	0.4	0.5	0.6	0.7	0.8	0.9	1.0

422 **2.2.5 Model setup**

423 Each branch of the model is computed 10 times using different random values (samples) within the uncertainty ranges of the
424 different input parameters (e.g., fault slip rates, *b*-values) to explore the epistemic variability of the seismicity rates linked to
425 the input parameter uncertainties. This results in the computation of 1080 different models, 360 for each rupture scenario
426 branch.

427 **2.3 Comparison with observations**

428 **2.3.1 Regional historical and instrumental seismicity**

429 We adopt the Parametric Catalogue of Italian Earthquakes - CPTI15 v4.0 (Rovida et al., 2022) as a regional seismic catalogue
430 to compare the synthetic MFDs modelled in this study and to evaluate the models that best fit such seismicity rates. CPTI is
431 nowadays the main tool for seismic hazard models in Italy, as it homogenises the magnitudes of pre-instrumental (historical)
432 and instrumental events (Rovida et al., 2020). The catalogue contains all known Italian earthquakes from year 1000 AD. All
433 the available events for this work have been selected within a radius of about 150 km encompassing the study area in the
434 southern Apennines. The resulting selection (Fig. 1) spans a time interval from year 1019 to 2017, and a moment magnitude
435 range of $2.9 \leq M_w \leq 7.2$. The minimum magnitude for our analysis is set at $M_w 4.0$, and the completeness magnitudes to obtain
436 the GR distributions for the CPTI catalogue are reported in Rovida et al. (2020). For the comparison with the models, only the
437 events falling within the buffer area in Fig. 5, which is regarded as the area of influence of the active faults, are considered.

438 **2.3.2 Paleoearthquake rates**

439 In faults with available paleoseismological studies and, as such, estimates of paleoearthquake recurrence intervals, we compare
440 the modelled participation rates of that fault with the estimated paleoearthquake rates from paleoseismological studies
441 (paleoearthquake rate = $1/\text{paleoearthquake recurrence interval}$). We considered that the paleoseismological earthquake rates
442 (paleorates) data represent earthquakes of $M_w \geq 6.0$ - 6.5 , based on the statistical fact that most surface rupturing earthquakes
443 (i.e., those observed in paleoseismology) have a magnitude equal or higher than the said threshold (Baize et al., 2020;
444 Nurminen et al., 2022).

445 Paleoseismic data, including annual paleoearthquake rates, are available for three faults activated during the 1980 Mw 6.9
446 Irpinia earthquake: Pescopagano (21), Colliano (22), and San Gregorio Magno (23), thanks to several studies (e.g., Pantosti &
447 Valensise, 1990; D'Addezio et al., 1991; Pantosti et al., 1993).

448 **3 Results**

449 Our results show that the choice of rupture scenario is the primary factor driving differences in the modelled seismicity rates,
450 whereas the other explored branches have a much more limited impact. For this reason, to enable a direct comparison of the

Formattato: Struttura + Livello:3 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,27 cm

ha eliminato: 360

ha eliminato: 120

Formattato: Struttura + Livello:3 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,27 cm

ha eliminato: Rovida et al. (2020).

Formattato: Struttura + Livello:3 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,27 cm

ha eliminato: M_w

Formattato: Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 0,76 cm

ha formattato: Inglese (Regno Unito)

ha eliminato: 134

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

different rupture scenarios, we display our results for each rupture scenario showing a single branch for the scaling relationship, b -value, background seismicity and buffer area ("Leonard (2010)", " b -value = 0.9 – 1.1", "BG 1", "buffer area = 10 km", respectively). The results for all the other branches are available in the Zenodo repository of the paper (Alessandrini et al., 2026). The quality of our models is tested through the evaluation of their fit with the regional earthquake catalogue MFD, the overall model performance based on the SHERIFS NMS proportion, and the agreement with paleoseismic slip rates.

3.1 Modelled MFDs: fit with the earthquake catalogue

In the upper three panels of Fig. 6, we show the fit between the synthetic MFDs, i.e., the modelled seismicity rates, and the regional seismic catalogue for the different rupture scenarios. In each scenario, the synthetic MFD is formed by combining 10 different MFDs, generated from the 10 random samples explored in the modelling. Results show that the choice of the rupture configuration directly influences the shape of the synthetic MFDs. The intermediate multi-fault rupture scenario Set_1 and the large multi-fault rupture scenario Set_2 exhibit the best fit with the catalogue, as the modelled rates with SHERIFS overlap with the catalogue-derived MFD. In particular, Set_1 displays a strong agreement with the catalogue for magnitudes up to M_w 6.0, while Set_2 tends to slightly underestimate the MFD across this magnitude range, but the modelled and observed rates still remain in good agreement when accounting for the respective variability. Conversely, for Set_0 there is no match between the modelled and catalogue MFDs: the synthetic MFD lies consistently below the catalogue-derived curve across the entire magnitude range, even when considering their respective uncertainty bounds.


3.2 Non-Main-Shock slip

The choice of a rupture configuration also influences how the slip rate budget is consumed in the different SHERIFS iterations. When part of this budget remains unspent at the end of the calculation, it is referred to as non-mainshock slip. The NMS is expressed as a percentage and represented through histograms for each of the 10 random samples in each scenario (Fig. 6, lower panels). Following Chartier et al. (2019), NMS percentages above 30-40% in a model indicate poor model performance. Accordingly, we use the NMS calculated by SHERIFS as an indicator of model quality: lower NMS correspond to better performance, as the models are able to spend the prescribed slip rate in the computation. The NMS is also useful to differentiate between models with otherwise similar fits with the catalogue, as in the case of Set_1 and Set_2 (Fig. 6, upper panels). In our results, Set_1 exhibits the lowest NMS values among all models (mean value of 15.5%), with 90% of its samples falling below the 40% threshold. Set_2 shows slightly higher values, but its mean value (23.9%) remains well within the acceptable range. Conversely, Set_0 displays consistently high NMS values, ranging from (85%-97%). These results confirm that Set_1 and Set_2 yield the most robust performances, in line with their closer agreement with the catalogue-derived seismicity rates (Fig. 6).

ha eliminato: and ...background seismicity and buffer area ("Leonard (2010),"..., " b -value = 0.9 – 1.1,"..., "BG 1", "buffer = 10 km", respectively). The results for all the other branches are available in the Zenodo repository of the paper (Alessandrini et al., 2025) ... [105]

ha formattato: Colore carattere: Colore personalizzato(79;129;189))

ha eliminato: M_w ...w 6.0, while it slightly underestimates the seismicity rates at higher magnitudes....Set_2 tends to slightly underestimate the MFD across the entire ... [106]



ha eliminato: Figure 6.

ha formattato: ... [107]

ha eliminato: 2... exhibits the lowest NMS values among all models,...(mean value of 15.5%), with 80...0% of its samples falling below the 40% threshold. Set_1... shows slightly higher values, but its mean value (40.1...3.9%) remains well within the acceptable range. Conversely, Set_0 displays consistently high NMS v... [108]

ha eliminato: 14

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra




Figure 6. Columns show different rupture scenarios. Above: comparison between the modelled MFDs (green) and the seismicity rates from the historical and instrumental regional catalogues within the study area (red). Solid green line is the mean MFD and green patches represent the uncertainty (16–84 percentiles). Below: NMS (expressed as a percentage) for each random sample explored in the model.

3.3 Paleoearthquake rates on faults

Here we compare the participation rate curves modelled with SHERIFS in each of the three fault sections with the paleorates documented (faults 21, 22, 23; Fig. 3), for each rupture scenario, including their magnitude and rate uncertainty ranges (Fig. 7). The participation rates are derived from all ruptures involving these fault sections.

While all three rupture scenarios tend to underestimate the paleoseismic rates for fault 21, Set 1 and Set 2 closely match the paleoearthquake rate mean values and their uncertainty ranges for faults 22 and 23. Set 1 only slightly overestimates the central value for fault 22, but remains within the confidence intervals. In contrast, Set 0 consistently and significantly underestimates the measured rates for all three faults, confirming its weaker performance in line with the previous quality tests.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero


Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero, Motivo: Trasparente

ha spostato (inserimento) [4]

ha spostato (inserimento) [5]

ha formattato: Motivo: Trasparente (Colore personalizzato(183;221;232)))



ha eliminato:

ha eliminato: 15

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra




Figure 7. Comparison of the modelled paleoearthquake rates on faults and paleoseismic data. Green solid curves are the modelled mean cumulative participation rates rupturing the specific fault section; dashed lines are the dispersions of individual samples; purple points are the paleorates measured on the fault trench; purple boxes are the paleodata uncertainty ranges. The location of each fault (rows) is highlighted in red within the fault maps, for the three different scenarios (columns).

4 Discussion

4.1 Segmentation models in the southern Apennines

The results of our analysis show that the impact of different rupture scenarios leads to markedly different behaviours in terms of seismicity rates, MFD shape, and expected maximum magnitudes. These results have implications for the segmentation models to be considered for PSHA in the southern Apennines.

Both Set_1 and Set_2, which allow ruptures involving multiple adjacent faults, fit well the catalogue. Set_1, in particular, stands out as the most balanced configuration, showing strong consistency with the observed seismicity and producing a maximum expected magnitude (M_w 7.4) closer to historical records with respect to Set_2 (M_w 7.6). Conversely, individual segment ruptures show lower consistency with the observations for the study region.

The good agreement of multi-fault rupture earthquake rates with the seismic and paleoseismic records in the region, indicates that such multi-fault rupture scenarios are statistically consistent with observations in the southern Apennines, given the

ha eliminato: 7.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero, Motivo: Trasparente (Colore personalizzato(RGB(183;221;232)))

ha eliminato: ¶

ha formattato: Motivo: Trasparente (Colore personalizzato(RGB(183;221;232)))

ha eliminato: less feasibility

ha formattato: Colore carattere: Colore personalizzato(RGB(191;191;191))

ha eliminato: feasible

ha eliminato: 16¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

578 modelling set up and fault input data used. These results are also in line with historical observations, such as the 1980 Irpinia
579 earthquake, which involved the rupture of multiple fault segments (21-22-23; Fig. 3, Table 1).
580 Considering these multi-fault scenarios also has important implications for the maximum expected magnitude in the region,
581 as they can reach values up to M_w 7.6 (Set_2) that are unseen in the historical records of the region (CPTI; Rovida et al., 2022).
582 However, it is important to remark that SHERIFS is not a physics-based model (e.g., it does not account for kinematic rupture
583 compatibility, stress transfer, etc.) to evaluate whether the fault system can accommodate such large magnitudes. Instead, it
584 relies on statistical metrics derived from observations, which might not fully capture the specific characteristics of the southern
585 Apennines fault system.
586 Despite this and as observed in other tectonic regions (e.g., California, Field et al., 2014; New Zealand, Quigley et al., 2017;
587 Taiwan, Chang et al., 2023), relaxing segmentation can give enhanced performance of seismicity models for PSHA. While the
588 rupture scenario models explored in this study are a sample of the possible rupture scenario models to be explored, our results
589 give valuable insight for future seismic hazard evaluations in the region.

590 **4.2 Tectonic domains impact**


591 The modelled faults in the study area span two tectonically distinct domains: SubArea 1, characterised by extensional normal
592 faulting, and SubArea 2, dominated by strike-slip structures (Fig. 1). These tectonic domains not only differ in fault kinematics
593 but also in their seismic expression: SubArea 1 shows a denser clustering of seismicity, while SubArea 2 appears notably more
594 quiescent. To investigate how these differences may influence model behaviour, we analysed the two sub-areas independently,
595 that is, we computed the models considering only the faults and seismicity catalogue within each sub-area (the background
596 seismicity buffers specific to each sub-area are shown in Fig. 5).
597 The results show that the tectonic domain influences the statistical performance of the tested rupture hypotheses. In particular,
598 we observe that the modelled rates of SubArea 1 faults (Fig. 8) return considerably better fits with the catalogue rates than
599 SubArea 2 (Fig. 9), which systematically underestimates the catalogue. In SubArea 1, Set_1 is the one with better overall
600 performance for its agreement with the catalogue for $M_w > 6$ and considerably low NMS percentages. The other sets
601 underperform Set_1: Set_0 has higher NMS percentages, although holding mean values within the 40% threshold; Set_2
602 slightly underestimates the catalogue rates. These results, especially the agreement of Set_1 and the NMS values across models,
603 are consistent with the full model analysis (both sub-areas combined) and suggest that the extensional domain of the Southern
604 Apennines (SubArea 1) contributes substantially to the overall seismicity rates reproduced by the model. Conversely, the
605 results of SubArea 2 indicate that the currently available fault data do not fully capture the seismicity rates observed in the
606 catalogue in this area.
607 This can be due to two principal reasons. First, the faults in this region generally have lower slip rates (Fig. 3), and in some
608 cases, relatively long lengths. Since SHERIFS treats slip rate as a budget that is spent on larger magnitudes first (due to its
609 moment rate formulation), these faults tend to release most of their seismic moment through fewer, higher-magnitude events.
610 As a consequence, the earthquake rates across the full MFD are reduced, particularly at lower magnitudes.

ha eliminato: consistent... Iso in line with historical observations, such as the 1980 Irpinia earthquake that simultaneously ruptured 3 faults... which involved the rupture of multiple fault segments (21-22-23; Fig. 3, Table 1), highlighting the potential of the region to generate such events. ... [109]

ha eliminato: ... [110]

ha eliminato: clearly ...how that the tectonic domain has a strong influence on how different...influences the statistical performance of the tested rupture hypotheses perform. We... In particular, we observe that the modelled rates of SubArea 1 faults (Fig. 8) return considerably better fits with the catalogue rates than SubArea 2 (Fig. 9), which systematically underestimates the catalogue. In SubArea 1, Set_1 is the one with better overall performance for its agreement with the catalogue for the M_w 4.5-...w > 6 range ...nd ... [111]

ha eliminato: plays a dominant role in controlling the regional seismicity.



ha eliminato: ... [112]

ha eliminato: Conversely, the results of SubArea 2 show that the active faults considered for the modelling are not fully able to explain the seismicity rates of the catalogue in this area. ...his can be due to two principal reasons. One contributing factor is that... ... [114]

ha formattato ... [113]

ha eliminato: 17

ha formattato: Colore carattere: Nero
Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra





Figure 8, SubArea_1 models: columns show different rupture scenarios. Above: comparison between the modelled MFDs (green) and the seismicity rates from the historical and instrumental regional catalogues within the study area (red). Below: NMS (expressed as a percentage) for each random sample explored in the model.

A second factor may lie in the limited geological constraints available for several faults in SubArea 2. Many slip rates in this region come from regional geodynamic models rather than direct geological constraints (see DISS Working Group, 2025), and paleoseismic data are notably scarce. Although some structures, particularly in the Gargano Promontory, are well studied and associated with significant seismicity, the overall quantity and resolution of geological data in SubArea 2 remain more limited than in SubArea 1, likely due to the lower number of large earthquakes historically recorded in the region. Additionally, certain faults in SubArea 2 are modelled with long, simplified traces that may not fully represent their internal segmentation. This generalisation could lead to an overestimation of rupture length in the models, ultimately affecting seismicity rate calculations. Similar effects have been observed in previous SHERIFS applications (e.g., Gómez-Novell et al., 2020a), where simplified fault geometries impacted the agreement between modelled and observed seismicity.

Overall, the observed differences between the two sub-areas highlight how variations in the level of geological constraint may influence model performance. While the lower seismic activity in SubArea 2 could suggest a role of tectonic behaviour, the modelling framework adopted in this study is not intended to resolve tectonic controls on seismicity. Rather, it evaluates the



ha eliminato: Figure 9.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Colore carattere: Nero

ha eliminato: —


ha eliminato: —

ha eliminato: 18

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

704 statistical consistency of alternative rupture hypotheses with available observations, which are inherently influenced by the
705 quality and completeness of input data.
706



707
708 **Figure 9.** SubArea 2 models: columns show different rupture scenarios. Above: comparison between the modelled MFDs (green)
709 and the seismicity rates from the historical and instrumental regional catalogues within the study area (red). Below: NMS (expressed
710 as a percentage) for each random sample explored in the model.
711

712 4.3 Scaling relationship impact

713 Among the three scaling relationships tested, the differences between Leonard (2010) and Wells & Coppersmith (1994) are
714 minor, with both yielding consistent and reliable results across all rupture scenarios. The Leonard (2010) scaling relationship
715 was ultimately selected as the reference scaling relationship in this study (Fig. 6), as it delivers a slightly better fit to the MFDs,
716 lower NMS percentages, being also a more recent formulation applicable to a wider range of tectonic settings.
717 In contrast, the scaling law by Thingbaijam et al. (2017), while producing a closer match to the regional MFD in the single-
718 fault scenario (Set_0), results in only minor differences across the three rupture sets (Fig. 10), making it less effective in

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: 19

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

distinguishing between different rupture behaviours. This, together with a generally broader dispersion and a systematically higher proportion of non-main-shock slip, limits its applicability in our modelling framework.





Figure 10. Rows correspond to different scaling laws. Columns 1–3 show the comparison between modelled MFDs (green) and seismicity rates from the historical and instrumental regional catalogues within the study area (red), for the three rupture scenarios. Column 4 displays the NMS (expressed as a percentage) for each random sample explored under all rupture scenarios.

This behaviour may be attributed to the nature of the Thingbaijam et al. (2017) scaling law, which instead of geological data is based on source modelling inversion datasets. This scaling law tends to estimate smaller magnitudes for a given rupture area compared to Wells & Coppersmith (1994), as it can be seen by the maximum magnitudes of the models depending on the scaling relationship. Because SHERIFS operates under the preservation of the seismic moment assumption (budget), the reduction of M_{max} in the Thingbaijam et al. (2017)-based models is compensated in the whole MFD, which leads to more similar earthquake rates across all models. This formulation seems less effective for our fault system and therefore has not adopted as the reference model.

4.4 Impact of MFD shape and background seismicity

Concerning the MFD shape assumption, the choice of a Gutenberg–Richter distribution fits the observed seismicity well and proves to be a valid and robust statistical representation. This choice is also consistent with long-standing applications of the GR law to the instrumental seismicity in Italy (e.g., Stucchi et al., 2011; Rovida et al., 2020; Meletti et al., 2021). Exploring a wider or a narrower b -value range has also a limited impact in our results (Fig. 11). Generally, the wider b -value range of 0.9–1.1 describes slightly better the regional seismicity and shows slightly lower NMS percentages. This is because the MFD shape is less restrictive in these models and allows the modelling to better capture the rate variability of the catalogue.



ha eliminato:

ha eliminato: 10.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: 20

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra





Figure 11. Rows correspond to different b-values. Columns 1–3 show the comparison between modelled MFDs (green) and seismicity rates from the historical and instrumental regional catalogues within the study area (red), for the three rupture scenarios. Column 4 displays the NMS (expressed as a percentage) for each random sample explored under all rupture scenarios.

Consistent with previous studies that acknowledge the relevance of distributed background seismicity models to account for events not clearly linked to known faults (e.g., Stirling et al., 2002; Chartier et al., 2019; Gerstenberger et al., 2024), we configured BG_2 to include a magnitude-dependent fraction of seismicity within a 10 km buffer zone surrounding the mapped faults. Set 0 is the one showing the larger differences related to background seismicity, allowing to partially compensate the model underestimation of the earthquake catalogue rates (Fig. 12). Conversely, the effect of BG_2 on multi-fault rupture models (Set_1 and Set_2) is minimal, and in some cases slightly worsens the fit to the observed MFD. Additionally, BG_1 yields slightly lower NMS values (Fig. 12), with only marginal differences. This indicates that introducing background seismicity does not produce any significant improvement in our modelling results and, as such, the relative performance between rupture scenario sets remains invariant.

To evaluate the sensitivity of these results to the definition of the background area, we additionally tested buffer distances of 5 km and 15 km (available in the Zenodo repository; Alessandrini et al., 2026). The results obtained with these alternative configurations are consistent with those derived for the 10 km buffer, confirming that the overall outcomes are not significantly influenced by the selected buffer size. Moreover, a 5-km buffer from fault traces is not optimal because it does not always capture the whole surface projection area of the faults and, as such, might exclude some seismicity related to the faults at study, while the 15 km buffer might include seismicity out of the known fault systems.

Although a distributed seismicity component is widely considered essential in modern PSHA frameworks, especially to account for unmapped faults, the similarity between fault seismicity rates and earthquake catalogue indicates that most of the



ha eliminato:

ha eliminato: 11.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: . This approach results in slightly higher seismicity rates at lower magnitudes compared to the fault-only configuration BG_1 (Fig. 12).

ha eliminato: 21

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

recorded seismicity within the buffer is likely linked to the active fault structures considered. Nonetheless, the introduction of BG_2 also adds an additional layer of epistemic uncertainty, as the magnitude-dependent ratios adopted are based on expert judgment and remain inherently arbitrary. Given that our tests indicate a limited impact on overall model performance, we prefer to adopt the simpler BG_1 configuration, which avoids unnecessary complexity without compromising the quality of the results.




Figure 12. Rows correspond to different background seismicity scenarios. Columns 1–3 show the comparison between modelled MFDs (green) and seismicity rates from the historical and instrumental regional catalogues within the study area (red), for the three rupture scenarios. Column 4 displays the NMS (expressed as a percentage) for each random sample explored under all rupture scenarios.

4.5 Model weighting and implications for probabilistic seismic hazard

The main challenges in fault-based PSHA applications adopting relaxed segmentation frameworks concern: (i) the definition of possible earthquake rupture scenarios and (ii) the estimation of their associated rates of occurrence. While the approach employed in this study does not aim to resolve the physical plausibility of multi-fault ruptures in the southern Apennines which would require physics-based investigations- it provides a systematic framework to structure and explore epistemic uncertainties related to rupture configurations and earthquake rates within a PSHA context. In this framework, the approach does allow to narrow some of these uncertainties through consistency analyses with available observational data (instrumental seismicity and paleoseismic rates). At the same time, our hypothesis-based setup allows preserving the probabilistic nature of PSHA, as the results are intended to be implemented within a logic-tree scheme. It is important to note that this study represents only a subset of the many rupture scenarios and seismicity parameters that could be explored for the southern Apennines and is therefore not intended to confirm or rule out other configurations. We

ha eliminato: Necessarily, enlarging the buffer area to encompass a larger portion of the seismicity of the region, would likely widen the differences between BG_1 and BG_2 as the chances of including seismicity out of the known fault would increase accordingly.

ha eliminato: ... [115]

ha eliminato: 12.

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

ha formattato: Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Formattato: Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

ha formattato: Colore carattere: Nero

ha eliminato: Our study suggests that

ha eliminato: are feasible, based on their agreement

ha eliminato: catalogue

ha eliminato: observations. However,

ha eliminato: .

ha eliminato: specific fault rupture

ha eliminato: 22

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

also acknowledge the limitations in model quality check, as the available datasets, namely the regional seismic catalogue and the sparse paleoseismic records, may be insufficient to fully assess model performance.

In this sense, the main outcome of this study is to provide a modelling framework that enables the use of geological-based criteria to assign weights to rupture hypotheses within a PSHA logic tree. The modelling framework we propose follows a logic-tree structure that can be directly implemented within PSHA engines such as OpenQuake, as SHERIFS is explicitly designed to explore epistemic uncertainties in fault parameters and rupture configurations and to propagate them into probabilistic hazard calculations. The weights derived for the different model branches remain region-specific, as they are based on comparisons with local observational constraints that are not directly transferable to other tectonic settings. However, the proposed weighting rationale and the general methodological framework can be used in other active regions worldwide.

Based on the performance of the different models across all consistency tests (summarized in Table 4), Set 1 emerges as the most robust configuration and should be assigned the highest weight in a PSHA logic tree, given its strong agreement with the catalogue MFD. Set 2, while slightly less aligned with the catalogue MFD, provides the best results in terms of NMS and paleoseismic fit, and should therefore receive a comparable weight. In contrast, Set 0 shows consistently poorer performance and limited agreement with observational data, and should be assigned a lower weight.

Table 4. Summary table of the main outcomes discussed for each branch explored.

Features	Branches Explored	Key Findings	Preferred Branch
Rupture Scenarios	• Set 0	Superior performance of Set 1 and Set 2 over Set 0 against catalogue and paleorates	Set 1
	• Set 1		
	• Set 2		
Tectonic Domains	• Extensional (SubArea 1)	SubArea 1 drives regional seismicity	Not included as branching model
	• Strike-Slip (SubArea 2)		
Scaling Laws	• Wells & Coppersmith (1994)	Superior equal performance of Wells & Coppersmith (1994) and Leonard (2010) over Thingbaijam et al. (2017) against catalogue	Leonard (2010)
	• Leonard (2010)		
	• Thingbaijam et al. (2017)		
MFD Shape (b-value)	• 0.9 – 1.1	Minor differences, but broader b-value range fits better the catalogue	0.9–1.1
	• 0.93 – 0.96		
Background Seismicity	• BG 1	Minor differences, but BG 2 ratios are more speculative	BG 1
	• BG 2		
Buffer Area	• 5 km	Limited sensitivity to buffer size; 10 km adopted as reference configuration	10 km
	• 10 km		
	• 15 km		

ha eliminato: —

ha eliminato: —

ha eliminato: Instead, our aim

ha eliminato: Table 4),

ha spostato in basso [8]: As for the other modelling branches explored, while their influence is less pronounced than that of the rupture scenarios, they can still serve as additional criteria for developing hazard models. Among the scaling relationships tested, Leonard (2010) is our preferred choice due to its consistent performance across scenarios. Wells & Coppersmith (1994) yields

ha spostato in basso [9]: 1.1) results in better agreement

ha eliminato: Regarding the b-value, we find that adopti (... [117]

ha eliminato: 1

ha eliminato: 4.

ha formattato (... [118]

ha formattato (... [120]

ha formattato: Colore carattere: Nero

Formattato (... [119]

ha formattato: Colore carattere: Automatico

ha formattato: Colore carattere: Nero

Formattato (... [121]

Formattato: Tabulazioni: Non a 1,27 cm

ha formattato: Colore carattere: Automatico

ha formattato: Colore carattere: Nero

Formattato (... [122]

ha formattato: Colore carattere: Automatico

Formattato: Tabulazioni: Non a 1,27 cm

ha formattato: Colore carattere: Nero

Formattato (... [123]

Formattato: Tabulazioni: Non a 1,27 cm

ha formattato: Colore carattere: Automatico

ha formattato: Colore carattere: Automatico

ha formattato: Colore carattere: Nero

Formattato (... [124]

Formattato: Tabulazioni: Non a 1,27 cm

ha formattato: Colore carattere: Automatico

ha formattato: Colore carattere: Nero

Formattato (... [125]

ha eliminato: 231

ha formattato: Colore carattere: Nero

Formattato (... [116]

855
856
857
858
859
860
861
862
863
864
865
866
867

868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886

As for the other modelling branches explored, while their influence is less pronounced than that of the rupture scenarios, they can still serve as additional criteria for developing hazard models. Among the scaling relationships tested, Leonard (2010) is our preferred choice due to its consistent performance across scenarios. Wells & Coppersmith (1994) yields very similar results and can therefore be considered a valid alternative. In contrast, the relationship proposed by Thingbaijam et al. (2017) generally results in poorer model performance, with greater dispersion in the MFDs and higher NMS percentages.

Regarding the b -value, we find that adopting a broader range (0.9–1.1) results in better agreement between the modelled and observed seismicity, making it our preferred option. As for background seismicity, BG 1 (where all seismicity is attributed to the faults) is favoured, as it explains the observed seismicity well and yields model performances comparable to BG 2 for logic tree weighting purposes. Additionally, BG 1 avoids the added layer of arbitrariness required by BG 2, thus reducing unnecessary complexity in the modelling framework. Sensitivity tests performed using alternative background buffer distances (5 km, 10 km and 15 km) show only minor variations in model performance, with the 10 km configuration representing the preferred compromise between spatial association of seismicity with mapped faults and the obtained results.

5 Conclusions

This study explores the impact of different variables on fault-based seismicity rate modelling in the southern Apennines, using a relaxed segmentation approach that allows multi-fault ruptures. The region, affected by the 1980 Mw 6.9 Irpinia earthquake, includes several critical infrastructures, highlighting the need for accurate seismic hazard assessment.

We computed seismicity rates from geological data in 35 seismogenic sources employing the SHERIFS algorithm and performing an exploration tree that combines different rupture scenarios and seismic parameter assumptions (e.g., scaling relationships, b -value, background seismicity, buffer area) to explore intrinsic epistemic uncertainties in fault data. We evaluated model performance by comparing them against the regional earthquake catalogue data and paleoearthquake rates from paleoseismic studies, as well as quantifying computational model performance.

Our results show that the scenarios allowing earthquake ruptures involving multiple adjacent faults consistently show better agreement with observations on regional seismicity and paleoseismic data than single segment rupture scenarios. These findings highlight the importance of including fault rupture complexity in fault-based PSHA models, especially in regions with complex fault systems like the southern Apennines. Our results also establish criteria to weight seismicity models for future PSHA implementations in the region.

Nevertheless, this modelling has some limitations. The limited availability and quality of fault data, along with the necessary simplification of fault geometries, result in poorly constrained input parameters; at the same time, the regional seismic catalogue and the limited number of paleoseismic constraints may be insufficient to fully evaluate model performance, introducing non-negligible uncertainty into the results. Moreover, the definition of background seismicity through magnitude-dependent ratios, particularly in BG 2, is expert-based and therefore open to alternative assumptions; additionally, the

ha spostato (inserimento) [8]

ha spostato (inserimento) [9]

ha eliminato: ¶

Formattato: Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 0,76 cm

ha eliminato: major

ha eliminato:).

ha eliminato: artifacts

ha eliminato: These results support the feasibility of multi-fault ruptures in the southern Apennines, and

ha eliminato: such ruptures

ha eliminato: Italy. These findings

ha eliminato: ¶
Input data: the

ha eliminato: in the region

Formattato: Normale, SpazioDopo: 0 pt, Nessun elenco puntato o numerato

ha eliminato: results in key input parameters that are often

ha eliminato: and introduce a

ha eliminato: our outcomes. ¶
Non-physics...

ha eliminato: approach: SHERIFS does not include

ha eliminato: 24¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

903 modelling framework itself relies on statistical consistency with observations rather than physics-based rupture modelling,
904 which could further refine both rupture scenario selection and the evaluation of seismicity rates.
905 Addressing these aspects will be essential for the next generation of fault-based seismic hazard models in southern Italy, where
906 the high seismicity and the presence of large population and critical infrastructure demand accurate estimates of large-
907 magnitude events, including those with long recurrence intervals.

908 **Code availability**

909 The SHERIFS code (Seismic Hazard and Earthquake Rates in Fault Systems; Chartier et al., 2019) is publicly available at
910 <https://github.com/tomchartier/SHERIFS>. The version used in this study (v1.1) is no longer available in the public repository
911 but can be obtained upon request from the original authors.

912 **Data availability**

913 All SHERIFS input and output files used in this study are publicly available through Zenodo:
914 <https://doi.org/10.5281/zenodo.17183318>.

915 **Author contribution**

916 Conceptualization and software: G.A. and O.G.-N.; methodology and writing—original draft preparation: G.A.; writing—
917 review and editing: G.A., O.G.-N., and S.C.; supervision: S.C., M.G., and U.T.

918 **Competing Interests**

919 The authors declare that they have no conflict of interest.

920 **Acknowledgments**

921 The authors are sincerely grateful to Oona Scotti (Institut de Radioprotection et de Sûreté Nucléaire, France) for her helpful
922 discussions on the model. The first author also wishes to thank the Universitat de Barcelona and the Istituto Nazionale di
923 Oceanografia e di Geofisica Sperimentale (OGS) for hosting and supporting the development of this work.

ha eliminato: modellèd

ha eliminato: ¶

Consistency check: the available datasets may be insufficient to fully assess model performance. The regional seismic catalogue includes large uncertainties, especially for historical earthquakes, which may affect spatial attribution (e.g., events located near but outside the modelled buffer zone). Paleoseismic data are available for only a small number of faults and may not be representative of the whole system.¶

Effect of the epistemic uncertainty: the selection of magnitude-dependent ratios, especially in BG_2, is inherently arbitrary and might influence model outcomes, although our tests indicate that the overall impact remains limited.

ha eliminato: both

ha eliminato: dams

ha eliminato: Team list¶

Giulia Alessandrini (G.A.)¶

Octavi Gómez-Novell (O.G.-N.)¶

Silvia Castellaro (S.C.)¶

Michela Giustiniani (M.G.)¶

Umberto Tinivella (U.T.)¶

ha eliminato: 25¶

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

945 **Financial support**

946 This research was conducted in the framework of activities made possible by the Consultancy Contract Commissioned between
947 the Consorzio per la Bonifica della Capitanata and the Department of Physics and Astronomy of the University of Bologna.
948 Octavi Gómez-Novell acknowledges support from the Generation D initiative, promoted by Red.es, an organisation attached
949 to the Ministry for Digital Transformation and the Civil Service, and financed by the Recovery, Transformation and Resilience
950 Plan through the European Union's NextGenerationEU.

951 **References**

952 [Adinolfi, G. M., De Matteis, R., Orefice, A., Festa, G., Zollo, A., De Nardis, R., and Lavecchia, G.: The September 27, 2012,](#)
953 [ML 4.1, Benevento earthquake: A case of strike-slip faulting in the southern Apennines \(Italy\), Tectonophysics,](#)
954 <https://doi.org/10.1016/j.tecto.2015.06.036>, 2015.
955 Alessandrini, G., Gómez-Novell, O., Castellaro, S., Giustiniani, M., and Tinivella, U.: Dataset for fault-based seismicity rates
956 in the southern Apennines (Italy) - Version 2, Zenodo [data set], <https://doi.org/10.5281/zenodo.17183317>, 2026.
957 Baize, S., Nurminen, F., Sarmiento, A., Dawson, T., Takao, M., Scotti, O., Azuma, T., Boncio, P., Champenois, J., Cinti, F.
958 R., Civico, R., Costa, C., Guerrieri, L., Marti, E., McCalpin, J., Okumura, K., and Villamor, P.: A worldwide and unified
959 database of surface ruptures (SURE) for fault displacement hazard analyses, Seismol. Res. Lett., 91, 499–520,
960 <https://doi.org/10.1785/0220190144>, 2020.
961 Basili, R., Valensise, G., Vannoli, P., Burrato, P., Fracassi, U., Mariano, S., Tiberti, M. M., and Boschi, E.: The Database of
962 Individual Seismogenic Sources (DISS), version 3: summarizing 20 years of research on Italy’s earthquake geology,
963 Tectonophysics, 453, 20–43, <https://doi.org/10.1016/j.tecto.2007.04.014>, 2008.
964 Bernard, P. and Zollo, A.: The Irpinia (Italy) 1980 earthquake: detailed analysis of a complex normal faulting, J. Geophys.
965 Res.-Sol. Ea., 94, 1631–1647, <https://doi.org/10.1029/JB094iB02p01631>, 1989.
966 Boncio, P., Lavecchia, G., and Pace, B.: Defining a model of 3D seismogenic sources for seismic hazard assessment
967 applications: the case of central Apennines (Italy), J. Seismol., 8, 407–425,
968 <https://doi.org/10.1023/B:JOSE.0000038449.78801.05>, 2004.
969 [Boncio, P., Mancini, T., Lavecchia, G., and Selvaggi, G.: Seismotectonics of strike-slip earthquakes within the deep crust of](#)
970 [southern Italy: Geometry, kinematics, stress field and crustal rheology of the Potenza 1990–1991 seismic sequences \(Mmax](#)
971 [5.7\), Tectonophysics, 445, 281–300, https://doi.org/10.1016/j.tecto.2007.08.001](#), 2007.
972 Chang, C. C., Chang, C. Y., Gao, J. C., and Chan, C. H.: Quantifying the probability and uncertainty of multiple-structure
973 rupture for Taiwan, Terr. Atmos. Ocean. Sci., 34, 7, <https://doi.org/10.1007/s44195-023-00040-8>, 2023.
974 Chartier, T., Scotti, O., Lyon-Caen, H., and Boiselet, A.: Methodology for earthquake rupture rate estimates of fault networks:
975 example for the western Corinth rift, Greece, Nat. Hazards Earth Syst. Sci., 17, 1857–1869, [https://doi.org/10.5194/nhess-17-](https://doi.org/10.5194/nhess-17-1857-2017)
976 [1857-2017](https://doi.org/10.5194/nhess-17-1857-2017), 2017.
977 Chartier, T., Scotti, O., and Lyon-Caen, H.: SHERIFS: open-source code for computing earthquake rates in fault systems and
978 constructing hazard models, Seismol. Res. Lett., 90, 1–10, <https://doi.org/10.1785/0220180332>, 2019.
979 Chartier, T., Scotti, O., Lyon-Caen, H., Richard-Dinger, K., Dieterich, J. H., and Shaw, B. E.: Modelling earthquake rates and
980 associated uncertainties in the Marmara Region, Turkey, Nat. Hazards Earth Syst. Sci., 21, 2733–2751,
981 <https://doi.org/10.5194/nhess-21-2733-2021>, 2021.

ha eliminato:),
ha eliminato: 17183318, 2025

ha eliminato: 26
ha formattato: Colore carattere: Nero
Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

984 Cheng, J., Chartier, T., and Xu, X.: Multisegment rupture hazard modeling along the Xianshuihe Fault Zone, southeastern
985 Tibetan Plateau, Seismol. Res. Lett., 92, 951–964, <https://doi.org/10.1785/0220200117>, 2021.

986 Cornell, C. A.: Engineering seismic risk analysis, Bull. Seismol. Soc. Am., 58, 1583–1606, 1968.

987 D'Addezio, G., Pantosti, D., and Valensise, G.: Paleoearthquakes along the Irpinia fault at Pantano di S. Gregorio Magno
988 (southern Italy), Il Quaternario, 4, 121–136, 1991.

989 Deligiannakis, G., Papanikolaou, I. D., and Roberts, G.: Fault specific GIS based seismic hazard maps for the Attica region,
990 Greece, Geomorphology, 306, 264–282, <https://doi.org/10.1016/j.geomorph.2016.12.005>, 2018.

991 DISS Working Group: Database of Individual Seismogenic Sources (DISS), Version 3.3.1: a compilation of potential sources
992 for earthquakes larger than M 5.5 in Italy and surrounding areas, Istituto Nazionale di Geofisica e Vulcanologia (INGV),
993 [dataset], <https://doi.org/10.13127/diss3.3.1>, 2025.

994 Doglioni, C., Harabaglia, P., Martinelli, G., Mongelli, F., and Zito, G.: A geodynamic model of the Southern Apennines
995 accretionary prism, Terra Nova, 8, 540–547, <https://doi.org/10.1111/j.1365-3121.1996.tb00783.x>, 1996.

996 El Kadri, S., Beauval, C., Brax, M., and Klinger, Y.: Implementation of an interconnected fault system in PSHA: example on
997 the Levant fault, Nat. Hazards Earth Syst. Sci., 25, 3397–3417, <https://doi.org/10.5194/nhess-25-3397-2025>, 2025.

998 Faure Walker, J., Boncio, P., Pace, B., Roberts, G., Benedetti, L., Scotti, O., Visini, F., and Peruzza, L.: Fault2SHA Central
999 Apennines database and structuring active fault data for seismic hazard assessment, Sci. Data, 8, 87,
1000 <https://doi.org/10.1038/s41597-021-00868-0>, 2021.

1001 Field, E. H., Dawson, T. E., Felzer, K. R., Frankel, A. D., Gupta, V., Jordan, T. H., et al.: Uniform California Earthquake
1002 Rupture Forecast, Version 2 (UCERF 2), Bull. Seismol. Soc. Am., 99, 2053–2107, <https://doi.org/10.1785/0120080049>, 2009.

1003 Field, E. H., Arrowsmith, R. J., Biasi, G. P., Bird, P., Dawson, T. E., Felzer, K. R., Jackson, D. D., Johnson, K. M., Jordan, T.
1004 H., Madden, C., Michael, A. J., Milner, K. R., Page, M. T., Parsons, T., Powers, P. M., Shaw, B. E., Thatcher, W. R., Weldon,
1005 R. J., and Zeng, Y.: Uniform California Earthquake Rupture Forecast, Version 3 (UCERF3)—The time-independent model,
1006 Bull. Seismol. Soc. Am., 104, 1122–1180, <https://doi.org/10.1785/0120130164>, 2014.

1007 Galadini, F. and Galli, P.: Paleoseismology of silent faults in the Central Apennines (Italy): the Mt. Vettore and Laga Mts.
1008 faults, Ann. Geophys., 46, <https://doi.org/10.4401/ag-3457>, 2003.

1009 Galli, P., Galderisi, A., Peronace, E., Giaccio, B., Hajdas, I., Messina, P., and Polpetta, F.: The awakening of the dormant
1010 Mount Vettore fault (2016 central Italy earthquake, Mw 6.6): paleoseismic clues on its millennial silences, Tectonics, 38, 687–
1011 705, <https://doi.org/10.1029/2018TC005326>, 2019.

1012 Gerstenberger, M. C., Van Dissen, R., Rollins, C., DiCaprio, C., Thingbaijam, K. K., Bora, S., and Williams, C.: The seismicity
1013 rate model for the 2022 Aotearoa New Zealand national seismic hazard model, Bull. Seismol. Soc. Am., 114, 182–216,
1014 <https://doi.org/10.1785/0120230165>, 2024.

1015 Gómez-Novell, O., Chartier, T., García-Mayordomo, J., Ortuño, M., Masana, E., Insua-Arévalo, J. M., and Scotti, O.:
1016 Modelling earthquake rupture rates in fault systems for seismic hazard assessment: the Eastern Betics Shear Zone, Eng. Geol.,
1017 265, 105452, <https://doi.org/10.1016/j.enggeo.2019.105452>, 2020a.

1018 Gómez-Novell, O., García-Mayordomo, J., Ortuño, M., Masana, E., and Chartier, T.: Fault system-based probabilistic seismic
1019 hazard assessment of a moderate seismicity region: the Eastern Betics Shear Zone (SE Spain), Front. Earth Sci., 8, 579398,
1020 <https://doi.org/10.3389/feart.2020.579398>, 2020b.

1021 Gulia, L. and Meletti, C.: Testing the b-value variability in Italy and its influence on Italian PSHA, Boll. Geof. Teor. Appl.,
1022 49, 59–76, 2007.

1023 Gutenberg, B. and Richter, C. F.: Frequency of earthquakes in California, Bull. Seismol. Soc. Am., 34, 185–188, 1944.

ha eliminato: ,

ha eliminato: . Discuss. [preprint],

ha eliminato: 2024-184, in review, 2024

ha eliminato: 27

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

1027 Hamling, I. J., Hreinsdóttir, S., Clark, K., Elliott, J., Liang, C., Fielding, E., and Stirling, M.: Complex multifault rupture during
1028 the 2016 Mw 7.8 Kaikōura earthquake, New Zealand, Science, 356, eaam7194, <https://doi.org/10.1126/science.aam7194>,
1029 2017.

1030 [ITHACA Working Group. ITHACA \(ITaLY HAZard from CApable faulting\), A database of active capable faults of the Italian](#)
1031 [territory. Version December 2019. ISPRA Geological Survey of Italy. \[dataset\]. Web Portal](#)
1032 <http://sgi2.isprambiente.it/ithacaweb/Mappatura.aspx>

1033 Latorre, D., Di Stefano, R., Castello, B., Michele, M., and Chiaraluce, L.: An updated view of the Italian seismicity from
1034 probabilistic location in 3D velocity models: the 1981–2018 Italian catalog of absolute earthquake locations (CLASS),
1035 Tectonophysics, 846, 229664, <https://doi.org/10.1016/j.tecto.2022.229664>, 2023.

1036 Leonard, M.: Earthquake fault scaling: Self-consistent relating of rupture length, width, average displacement, and moment
1037 release, Bull. Seismol. Soc. Am., 100, 1971–1988, <https://doi.org/10.1785/0120090189>, 2010.

1038 [Liu, C., Lay, T., Wang, R., Taymaz, T., Xie, Z., Xiong, X., Irmak, T. S., Kaharam, M. and Erman, C.: Complex multi-fault](#)
1039 [rupture and triggering during the 2023 earthquake doublet in southeastern Türkiye, Nat. Commun., 14, 5564,](#)
1040 <https://doi.org/10.1038/s41467-023-41404-5>, 2023.

1041 Meletti, C., Galadini, F., Valensise, G., Stucchi, M., Basili, R., Barba, S., Vannucci, G., and Boschi, E.: A seismic source zone
1042 model for the seismic hazard assessment of the Italian territory, Tectonophysics, 450, 85–108,
1043 <https://doi.org/10.1016/j.tecto.2008.01.003>, 2008.

1044 Meletti, C., Marzocchi, W., D'Amico, V., Lanzano, G., Luzi, L., Martinelli, F., Pace, B., Rovida, A., Taroni, M., Visini, F. and
1045 the MPS19 Working Group: The new Italian seismic hazard model (MPS19), Ann. Geophys., 64, [https://doi.org/10.4401/ag-](https://doi.org/10.4401/ag-8579)
1046 8579, 2021.

1047 Milner, K. R., Page, M. T., Field, E. H., Parsons, T., Biasi, G. P., and Shaw, B. E.: Appendix T—Defining the inversion rupture
1048 set using plausibility filters, US Geol. Surv. Open-File Rept., 2013–1165, 2013.

1049 NCSE (Norma de la Construcción Sismorresistente Española): Real Decreto 997/2002, de 27 de septiembre, por el que se
1050 aprueba la norma de construcción sismorresistente: parte general y edificación (NCSR-02), Bol. Of. Estado, 244, 35898–
1051 35967, 2002.

1052 NTC (Norme Tecniche sulle Costruzioni): Decreto Ministeriale 14/01/2008, Ministry of Infrastructures and Transportations,
1053 G.U. S.O. n.29 on 4/2/2008, 2008.

1054 NTC (Norme Tecniche sulle Costruzioni): Decreto Ministeriale 17/01/2018, Ministry of Infrastructures and Transportations,
1055 G.U. S.O. n.8 on 20/2/2018, 2018.

1056 Nurminen, F., Baize, S., Boncio, P., Blumetti, A. M., Cinti, F. R., Civico, R., and Guerrieri, L.: SURE 2.0—New release of the
1057 worldwide database of surface ruptures for fault displacement hazard analyses, Sci. Data, 9, 729,
1058 <https://doi.org/10.1038/s41597-022-01835-z>, 2022.

1059 Pace, B., Visini, F., Scotti, O., and Peruzza, L.: Preface: Linking faults to seismic hazard assessment in Europe, Nat. Hazards
1060 Earth Syst. Sci., 18, 1349–1350, <https://doi.org/10.5194/nhess-18-1349-2018>, 2018.

1061 Pantosti, D. and Valensise, G.: Faulting mechanism and complexity of the November 23, 1980, Campania-Lucania Earthquake,
1062 inferred from surface observations, J. Geophys. Res.-Sol. Ea., 95, 15319–15341, <https://doi.org/10.1029/JB095iB10p15319>,
1063 1990.

1064 Pantosti, D., Schwartz, D. P., and Valensise, G.: Paleoseismology along the 1980 surface rupture of the Irpinia Fault:
1065 Implications for earthquake recurrence in the southern Apennines, Italy, J. Geophys. Res.-Sol. Ea., 98, 6561–6577,
1066 <https://doi.org/10.1029/92JB02277>, 1993.

1067 Peruzza, L.: Se i terremoti fossero lavatrici, Sapere, 3, 12–16, <https://doi.org/10.12919/sapere.2018.03.1>, 2018.

ha eliminato: 28

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

1068 Peruzza, L., Pace, B., and Visini, F.: Fault-Based Earthquake Rupture Forecast in Central Italy: Remarks after the L'Aquila
1069 Mw 6.3 Event, Bull. Seismol. Soc. Am., 101, 404–412, <https://doi.org/10.1785/0120090276>, 2011.

1070 Peruzza, R., Gee, B., Pace, B., Roberts, G., Scotti, O., Visini, F., Benedetti, L., and Pagani, M.: PSHA after a strong earthquake:
1071 Hints for the recovery, Ann. Geophys., 59, 35, <https://doi.org/10.4401/ag-7257>, 2016.

1072 Petersen, M. D., Shumway, A. M., Powers, P. M., Field, E. H., Moschetti, M. P., Jaiswal, K. S., and others: The 2023 US 50-
1073 state national seismic hazard model: Overview and implications, Earthq. Spectra, 40, 5–88,
1074 <https://doi.org/10.1177/87552930231215428>, 2024.

1075 Quigley, M., Mohammadi, H., Jiménez, A., and Duffy, B.: Multi-fault earthquakes with kinematic and geometric rupture
1076 complexity: how common?, 8th International INQUA Meeting on Paleoseismology, Active Tectonics and Archeoseismology
1077 (PATA), New Zealand, 13–16 November 2017, p. 316–318, 2017.

1078 Rovida, A., Locati, M., Camassi, R., Lolli, B., and Gasperini, P.: The Italian earthquake catalogue CPTI15, Bull. Earthq. Eng.,
1079 18, 2953–2984, <https://doi.org/10.1007/s10518-020-00818-y>, 2020.

1080 Rovida, A., Locati, M., Camassi, R., Lolli, B., Gasperini, P., and Antonucci, A.: Catalogo Parametrico dei Terremoti Italiani
1081 (CPTI15), versione 4.0, Istituto Nazionale di Geofisica e Vulcanologia (INGV) [data set],
1082 <https://doi.org/10.13127/cpti/cpti15.4>, 2022.

1083 Scotti, O., Clément, C., and Baumont, D.: Seismic Hazard for Design and Verification of Nuclear Installations in France:
1084 Regulatory context, debated issues and ongoing developments, Boll. Geof. Teor. Appl., <https://doi.org/10.4430/bgta0080>,
1085 2014.

1086 Scotti, O., Visini, F., Faure Walker, J., Peruzza, L., Pace, B., Benedetti, L., Boncio, P., and Roberts, G.: Which Fault Threatens
1087 Me Most? Bridging the Gap Between Geologic Data-Providers and Seismic Risk Practitioners, Front. Earth Sci., 8, 626401,
1088 <https://doi.org/10.3389/feart.2020.626401>, 2020.

1089 Stirling, M. W., Verry, G. H. M., and Berryman, K. R.: A new seismic hazard model for New Zealand, Bull. Seismol. Soc.
1090 Am., 92, 1878–1903, <https://doi.org/10.1785/0120010156>, 2002.

1091 Stirling, M., McVerry, G., Gerstenberger, M., Litchfield, N., Van Dissen, R., Berryman, K., Barnes, P., Wallace, L., Villamor,
1092 P., Langridge, R., Lamarche, G., Nodder, S., Reyners, M., Bradley, B., Rhoades, D., Smith, W., Nicol, A., Pettinga, J., Clark,
1093 K., and Jacobs, K.: National Seismic Hazard Model for New Zealand: 2010 Update, Bull. Seismol. Soc. Am., 102, 1514–1542,
1094 <https://doi.org/10.1785/0120110170>, 2012.

1095 Stucchi, M., Meletti, C., Montaldo, V., Crowley, H., Calvi, G. M., and Boschi, E.: Seismic Hazard Assessment (2003–2009)
1096 for the Italian Building Code, Bull. Seismol. Soc. Am., 101, 1885–1911, <https://doi.org/10.1785/0120100130>, 2011.

1097 Thingbaijam, K. K. S., Mai, P. M., and Goda, K.: New empirical earthquake source-scaling laws, Bull. Seismol. Soc. Am.,
1098 107, 2225–2246, <https://doi.org/10.1785/0120170017>, 2017.

1099 Valentini, A.: Allowing multi-fault earthquakes and relaxing fault segmentation in Central Apennines (Italy): Hints for fault-
1100 based PSHA, Boll. Geofis. Teor. Appl., DOI 10.4430/bgta0339, 2020.

1101 Valentini, A., Visini, F., and Pace, B.: Integrating faults and past earthquakes into a probabilistic seismic hazard model for
1102 peninsular Italy, Nat. Hazards Earth Syst. Sci., 17, 2017–2039, <https://doi.org/10.5194/nhess-17-2017-2017>, 2017.

1103 Valentini, A., Pace, B., Boncio, P., Visini, F., Pagliaroli, A., and Pergalani, F.: Analisi probabilistica fault-based della
1104 pericolosità sismica volta alla definizione di input sismici: osservazioni dopo la sequenza sismica del 2016 in Italia Centrale,
1105 GNGTS 2018 Sessione 2.1 (Atti del 37° Convegno Nazionale del Gruppo Nazionale di Geofisica della Terra Solida), Italy,
1106 2018.

1107 Valentini, A., Pace, B., Boncio, P., Visini, F., Pagliaroli, A., and Pergalani, F.: Definition of Seismic Input From Fault-Based
1108 PSHA: Remarks After the 2016 Central Italy Earthquake Sequence, Tectonics, 38, 595–620,
1109 <https://doi.org/10.1029/2018TC005086>, 2019.

ha eliminato: 29

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

1110 Vannoli, P. and Burrato, P.: La sismicità del Veneto tra eventi storici e sorgenti sismogenetiche, in: Rischio sismico in Italia:
1111 analisi e prospettive per una geologia del rischio, Supplemento al n. 1/2018, Società Italiana di Geologia Ambientale (SIGEA),
1112 pp. 139–147, 2018.

1113 Visini, F., Pace, B., Meletti, C., Marzocchi, W., Akinci, A., Azzaro, R., Barani, S., Barberi, G., Barreca, G., Basili, R., Bird,
1114 P., Bonini, M., Burrato, P., Busetti, M., Carafa, M., Cocina, O., Console, R., Corti, G., D’Agostino, N., ... Zuccolo, E.:
1115 Earthquake Rupture Forecasts for the MPS19 Seismic Hazard Model of Italy, *Ann. Geophys.*, 64, 3, [https://doi.org/10.4401/ag-](https://doi.org/10.4401/ag-8608)
1116 8608, 2021.

1117 Wei, S., Fielding, E., Leprince, S., Sladen, A., Avouac, J. P., Helmberger, D., Hauksson, E., Chu, R., Simons, M., Hudnut, K.,
1118 Herring, T., and Briggs, R.: Superficial simplicity of the 2010 El Mayorg-Cucapah earthquake of Baja California in Mexico,
1119 *Nat. Geosci.*, 4, 615–618, <https://doi.org/10.1038/ngeo1213>, 2011.

1120 Wells, D. L. and Coppersmith, K. J.: New Empirical Relationships among Magnitude, Rupture Length, Rupture Width,
1121 Rupture Area, and Surface Displacement, *Bull. Seismol. Soc. Am.*, 84, 4, 974–1002,
1122 <https://doi.org/10.1785/BSSA0840040974>, 1994.

1123 ~~Williams, J. N., Werner, M. J., Goda, K., Wedmore, L. N., De Risi, R., Biggs, J., Mdala, H., Dulanya, Z., Fagereng, A.,~~
1124 ~~Mphemo, F., and Chindandali, P.: Fault-based probabilistic seismic hazard analysis in regions with low strain rates and a thick~~
1125 ~~seismogenic layer: A case study from Malawi, *Geophys. J. Int.*, 233, 1557–1579, <https://doi.org/10.1093/gji/ggad038>,~~
1126 ~~2023.Woessner, J., Laurentiu, D., Giardini, D., Crowley, H., Cotton, F., Grünthal, G., Valensise, G., Arvidsson, R., Basili, R.,~~
1127 ~~Demircioglu, M. B., Hiemer, S., Meletti, C., Musson, R. W., Rovida, A. N., Sesetyan, K., and Stucchi, M.: The 2013 European~~
1128 ~~Seismic Hazard Model: Key components and results, *Bull. Earthq. Eng.*, 13, 12, 3553–3596, [https://doi.org/10.1007/s10518-](https://doi.org/10.1007/s10518-015-9795-1)~~
1129 ~~015-9795-1~~, 2015.

1130 Zhang, H., Chen, J., and Ge, Z.: Multi-fault rupture and successive triggering during the 2012 Mw 8.6 Sumatra offshore
1131 earthquake, *Geophys. Res. Lett.*, 39, 22, <https://doi.org/10.1029/2012GL054207>, 2012.

ha eliminato: Woessner, J.,

ha formattato: Inglese (Regno Unito)

ha eliminato: 30

ha formattato: Colore carattere: Nero

Formattato: Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

Pagina 1: [1] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Colore carattere: Nero		
Pagina 1: [2] Formattato	Giulia Alessandrini	09/04/26 16:53:00
Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra		
Pagina 1: [3] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Normale (Web)		
Pagina 1: [4] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Nessuna spaziatura: Interlinea: 1,5 righe		
Pagina 1: [5] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Testo nota a piè di pagina		
Pagina 1: [6] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Bibliografia		
Pagina 1: [7] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Testo commento		
Pagina 1: [8] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Acknowledgments		
Pagina 1: [9] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
History		
Pagina 1: [10] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Authornames		
Pagina 1: [11] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Authors		
Pagina 1: [12] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Correspondence		
Pagina 1: [13] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Piè di pagina		
Pagina 1: [14] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Didascalia		
Pagina 1: [15] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Equation		
Pagina 1: [16] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00

Testo fumetto

Pagina 1: [17] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Affiliation		
Pagina 1: [18] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Paragrafo elenco		
Pagina 1: [19] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
MS title		
Pagina 1: [20] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Copernicus_Word_template		
Pagina 1: [21] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Name		
Pagina 1: [22] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Kontakt		
Pagina 1: [23] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Intestazione		
Pagina 1: [24] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Bullets: Rientro: Sinistro: 0 cm, Prima riga: 0 cm, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Tabulazioni: 0,63 cm, Tabulazione elenco		
Pagina 1: [25] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Betreff		
Pagina 1: [26] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Titolo 9: Struttura + Livello:9 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 10,16 cm + Tabulazione dopo: 11,43 cm + Imposta un rientro di: 11,43 cm		
Pagina 1: [27] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Titolo 8: Struttura + Livello:8 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 8,89 cm + Tabulazione dopo: 10,16 cm + Imposta un rientro di: 10,16 cm		
Pagina 1: [28] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Titolo 7: Struttura + Livello:7 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 7,62 cm + Tabulazione dopo: 8,89 cm + Imposta un rientro di: 8,89 cm		
Pagina 1: [29] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
Titolo 6: Tipo di carattere: (Predefinito) Times New Roman, Colore carattere: Colore personalizzato(RGB(36;63;97)), Inglese (Regno Unito), Rientro: Sinistro: 0 cm, Sporgente 2,03 cm		
Pagina 1: [30] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00

Titolo 5: Tipo di carattere: (Predefinito) Times New Roman, Colore carattere: Colore personalizzato(RGB(54;96;145)), Inglese (Regno Unito), Rientro: Sinistro: 0 cm, Sporgente 1,78 cm

Pagina 1: [31] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
---	----------------------------	--------------------------

Titolo 4: Rientro: Sinistro: 0 cm, Sporgente 1,52 cm

Pagina 1: [32] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
---	----------------------------	--------------------------

Titolo 3: Rientro: Sinistro: 0 cm, Sporgente 1,27 cm

Pagina 1: [33] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
---	----------------------------	--------------------------

Titolo 2: Struttura + Livello:2 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 1,02 cm

Pagina 1: [34] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
---	----------------------------	--------------------------

Titolo 1

Pagina 1: [35] Definizione stile	Giulia Alessandrini	09/04/26 16:53:00
---	----------------------------	--------------------------

Normale

Pagina 1: [36] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Tipo di carattere: 17 pt, Grassetto, Colore carattere: Nero

Pagina 1: [37] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Normale, SpazioPrima: 18 pt, Interlinea: multipla 1,83 ri, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Pagina 1: [38] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Distanza dal testo: 0 cm

Pagina 1: [39] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [40] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Tipo di carattere: 12 pt, Colore carattere: Nero

Pagina 1: [40] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Tipo di carattere: 12 pt, Colore carattere: Nero

Pagina 1: [41] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Tipo di carattere: 12 pt, Colore carattere: Nero

Pagina 1: [42] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Tipo di carattere: 12 pt, Colore carattere: Nero

Pagina 1: [43] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [44] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [45] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [46] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

▼

Pagina 1: [47] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [48] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [49] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero, Italiano

Pagina 1: [50] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero, Italiano

Pagina 1: [51] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Normale, SpazioPrima: 6 pt, Dopo: 18 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Pagina 1: [52] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero, Italiano

Pagina 1: [53] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Struttura + Livello:1 + Stile numerazione: 1, 2, 3, ... + Comincia da:1 + Allineamento: A sinistra + Allinea a: 0 cm + Imposta un rientro di: 0,76 cm

Pagina 1: [54] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Inglese (Regno Unito)

Pagina 6: [55] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

▼

Pagina 1: [56] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

Pagina 9: [57] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Colore personalizzato(RGB(79;129;189))

Pagina 1: [58] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

Pagina 10: [59] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Pagina 10: [60] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 10: [61] Formattato	Giulia Alessandrini	09/04/26 16:53:00
-----------------------------------	----------------------------	--------------------------

Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Pagina 10: [62] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 1: [63] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

Pagina 11: [64] Formattato	Giulia Alessandrini	09/04/26 16:53:00
-----------------------------------	----------------------------	--------------------------

Normale, SpazioDopo: 0 pt, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo),

Pagina 11: [65] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Pagina 1: [66] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
-------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 1: [67] Formattato	Giulia Alessandrini	09/04/26 16:53:00
----------------------------------	----------------------------	--------------------------

Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

Pagina 12: [68] Formattato	Giulia Alessandrini	09/04/26 16:53:00
-----------------------------------	----------------------------	--------------------------

Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Pagina 12: [69] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 12: [69] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 12: [70] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 12: [71] Formattato	Giulia Alessandrini	09/04/26 16:53:00
-----------------------------------	----------------------------	--------------------------

Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Pagina 12: [72] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero		
Pagina 12: [73] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero		
Pagina 12: [73] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero		
Pagina 12: [74] Tabella formattata	Giulia Alessandrini	09/04/26 16:53:00
Tabella formattata		
Pagina 12: [75] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [76] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [77] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [78] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [79] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [80] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [81] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [82] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [83] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [84] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [85] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [86] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		

Pagina 12: [87] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [88] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [89] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [90] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [91] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [92] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [93] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [94] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [95] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [96] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [97] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [98] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [99] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [100] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [101] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		
Pagina 12: [102] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
Tipo di carattere: 10 pt		

Pagina 12: [103] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Tipo di carattere: 10 pt

Pagina 12: [104] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Tipo di carattere: 10 pt

Pagina 14: [105] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [105] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [105] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [105] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [105] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [106] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [106] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [106] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [107] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 14: [107] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼		
---	--	--

Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 14: [108] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [109] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [109] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [109] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [110] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [110] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [110] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------



▼

Pagina 17: [111] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼

Pagina 17: [112] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼

Pagina 17: [113] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 17: [113] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Colore carattere: Nero

Pagina 17: [114] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼

Pagina 17: [114] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼

Pagina 22: [115] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼

Pagina 1: [116] Formattato	Giulia Alessandrini	09/04/26 16:53:00
-----------------------------------	----------------------------	--------------------------

Normale, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo), Tabulazioni: 7,96 cm, Allineato al centro + 15,92 cm, Allineato a destra

Pagina 23: [117] ha eliminato	Giulia Alessandrini	09/04/26 16:53:00
--------------------------------------	----------------------------	--------------------------

▼

Pagina 23: [118] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 23: [119] Formattato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

Normale, SpazioDopo: 10 pt, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Pagina 23: [120] ha formattato	Giulia Alessandrini	09/04/26 16:53:00
---------------------------------------	----------------------------	--------------------------

Tipo di carattere: 9 pt, Grassetto, Colore carattere: Nero

Pagina 23: [121] Formattato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

Normale, Rientro: Sinistro: 0,21 cm, Sporgente 1,27 cm, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bord

Pagina 23: [122] Formattato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

Normale, Rientro: Sinistro: 0,21 cm, Sporgente 1,25 cm, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bord

Pagina 23: [123] Formattato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

Normale, Rientro: Sinistro: 0,21 cm, Sporgente 1,25 cm, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bord

Pagina 23: [124] Formattato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

Normale, Rientro: Sinistro: 0,21 cm, Sporgente 1,25 cm, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bord

Pagina 23: [125] Formattato	Giulia Alessandrini	09/04/26 16:53:00
------------------------------------	----------------------------	--------------------------

Normale, Rientro: Sinistro: 0,21 cm, Sporgente 1,25 cm, Struttura + Livello:1 + Stile numerazione: Punto elenco + Allinea a: 0,63 cm + Imposta un rientro di: 1,27 cm, Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bord